

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
All races 3/, both sexes										
2004.....	816.5	685.2	29.9	14.7	18.7	66.1	94.0	96.0	107.9	151.7
2003.....	841.9	700.0	31.5	14.7	19.1	66.4	96.4	95.2	111.3	159.1
2002.....	847.3	695.0	31.2	15.2	19.5	67.8	95.2	94.7	111.6	161.3
2001.....	848.5	683.4	33.3	15.3	19.2	66.9	95.0	96.2	113.5	165.9
2000.....	854.0	736.7	32.4	15.8	20.3	67.1	93.6	91.2	111.0	159.2
1999.....	857.0	736.0	34.2	16.9	20.4	68.6	90.8	90.4	113.4	160.8
All races 3/, male										
2004.....	817.6	753.7	32.4	16.4	21.7	91.0	138.0	135.6	143.2	193.8
2003.....	840.3	777.4	35.1	16.2	23.1	92.3	140.3	133.5	148.9	202.7
2002.....	846.6	761.5	35.2	16.7	23.1	94.0	140.8	134.4	149.4	206.6
2001.....	846.4	749.8	37.0	16.7	22.8	93.7	141.1	135.6	151.2	213.7
2000.....	853.0	806.5	35.9	17.6	24.2	93.3	138.0	128.8	147.9	205.4
1999.....	859.2	805.0	37.9	18.6	24.4	94.7	133.1	127.0	152.0	207.6
All races 3/, female										
2004.....	815.4	613.4	27.3	13.0	15.4	39.9	47.3	54.5	72.0	109.3
2003.....	843.4	619.1	27.8	13.0	15.0	39.0	49.8	55.4	73.0	115.3
2002.....	848.0	625.3	27.0	13.5	15.9	40.1	47.3	53.6	73.3	116.1
2001.....	850.4	613.9	29.5	13.9	15.3	38.5	46.8	55.8	75.2	118.2
2000.....	855.0	663.4	28.7	14.0	16.1	39.3	47.1	52.8	73.6	113.2
1999.....	854.9	663.6	30.3	15.0	16.2	41.0	46.6	53.2	74.4	114.4
White, both sexes										
2004.....	863.2	574.0	27.0	13.4	17.4	64.4	88.3	87.3	98.3	140.6
2003.....	890.1	591.3	28.5	13.7	17.8	64.7	89.4	86.1	101.9	145.5
2002.....	895.7	586.7	28.1	14.0	18.1	66.3	88.0	85.0	101.1	146.8
2001.....	895.1	570.9	30.7	14.4	17.9	64.2	87.1	87.0	102.7	151.2
2000.....	900.2	610.6	29.2	14.8	19.2	64.5	85.1	81.0	99.0	144.0
1999.....	901.4	606.1	30.8	15.3	19.2	65.3	81.6	80.8	101.4	144.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
White, male										
2004.....	854.2	631.6	29.4	15.1	20.6	87.1	129.1	123.2	130.8	180.9
2003.....	877.6	658.7	31.5	15.3	21.4	88.3	129.1	120.0	137.0	187.6
2002.....	884.0	650.9	31.5	15.4	21.2	90.5	128.9	119.7	136.0	188.7
2001.....	881.9	627.6	34.2	15.3	21.3	88.3	128.2	122.4	137.4	196.7
2000.....	887.8	667.6	32.6	16.5	23.1	88.1	124.6	113.9	133.6	188.2
1999.....	892.1	667.0	33.9	16.8	22.7	88.3	118.5	113.5	136.5	188.8
White, female										
2004.....	871.9	513.6	24.4	11.6	14.2	40.4	44.3	48.8	64.3	98.9
2003.....	902.3	520.7	25.5	12.1	14.0	39.6	46.7	50.0	65.2	102.2
2002.....	907.0	519.4	24.5	12.6	14.8	40.4	44.4	48.1	64.8	103.9
2001.....	907.9	511.5	27.1	13.3	14.4	38.5	43.2	49.6	66.6	104.9
2000.....	912.3	550.5	25.5	13.0	15.2	39.3	43.0	46.5	63.0	99.2
1999.....	910.4	542.0	27.5	13.8	15.4	40.8	42.3	46.7	65.0	99.5
Black, both sexes										
2004.....	744.3	1,284.7	44.8	21.6	25.3	81.4	138.8	167.2	191.3	254.4
2003.....	763.6	1,273.8	46.8	19.7	25.7	80.0	147.3	168.6	194.1	275.2
2002.....	768.4	1,263.6	47.1	21.3	27.5	82.0	146.8	172.1	200.7	284.0
2001.....	773.5	1,304.5	47.5	20.7	25.8	86.3	151.3	168.8	200.5	286.0
2000.....	781.1	1,426.1	49.9	21.8	26.6	87.4	155.0	168.4	206.9	283.1
1999.....	788.1	1,457.3	52.6	25.2	28.0	91.9	154.4	163.5	204.9	293.5
Black, male										
2004.....	792.6	1,414.2	48.6	23.1	28.6	121.0	210.1	244.7	259.9	323.6
2003.....	813.7	1,410.0	53.7	21.6	31.4	120.9	224.5	249.6	262.5	344.8
2002.....	816.7	1,351.5	54.4	24.1	33.4	121.7	227.6	255.7	272.9	366.5
2001.....	823.9	1,426.4	53.2	23.8	31.6	130.4	236.3	247.0	271.0	364.1
2000.....	834.1	1,567.6	54.5	24.5	32.1	130.1	240.6	246.8	275.0	361.6
1999.....	847.4	1,592.8	59.0	29.3	35.0	137.7	239.9	238.0	278.9	372.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All causes

Black, female

2004.....	700.3	1,149.9	40.9	20.1	21.9	40.7	67.0	94.7	129.5	192.6
2003.....	717.9	1,132.2	39.7	17.8	19.8	37.8	70.6	93.6	132.5	213.1
2002.....	724.4	1,172.0	39.5	18.5	21.3	41.0	68.0	95.7	135.7	210.6
2001.....	727.7	1,176.3	41.7	17.5	19.8	40.8	69.4	97.5	137.3	216.6
2000.....	733.0	1,279.8	45.3	19.0	21.0	43.7	73.9	97.3	145.6	213.4
1999.....	734.3	1,317.4	46.1	21.0	20.7	45.2	73.2	96.1	138.6	223.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

I. Certain infectious and parasitic diseases (A00-B99)

All races 3/, both sexes

2004.....	22.0	11.8	1.2	0.4	0.4	0.7	1.6	3.3	6.8	12.5
2003.....	22.2	12.3	1.2	0.5	0.5	0.8	1.5	3.4	7.2	14.4
2002.....	22.2	14.4	1.1	0.5	0.6	0.8	1.5	3.9	8.2	15.3
2001.....	21.2	14.0	1.5	0.5	0.5	1.0	1.7	4.1	9.0	15.5
2000.....	21.0	13.4	1.4	0.5	0.5	0.8	1.6	4.4	10.4	16.0
1999.....	21.5	14.8	1.5	0.7	0.6	0.7	1.8	5.1	11.6	16.5

All races 3/, male

2004.....	23.7	13.8	1.1	0.4	0.4	0.7	1.8	3.6	8.4	16.8
2003.....	24.0	12.6	1.4	0.4	0.4	0.9	1.6	3.9	9.3	19.7
2002.....	24.2	14.8	1.2	0.5	0.6	0.7	1.6	4.4	10.5	21.6
2001.....	23.3	14.7	1.6	0.5	0.6	1.0	1.7	4.7	12.0	22.2
2000.....	23.0	14.1	1.5	0.5	0.5	0.9	1.8	4.9	13.8	23.1
1999.....	23.7	17.0	1.8	0.7	0.6	0.7	1.9	6.3	15.9	23.7

All races 3/, female

2004.....	20.3	9.6	1.2	0.4	0.4	0.7	1.3	3.1	5.2	8.2
2003.....	20.6	12.1	1.1	0.5	0.5	0.8	1.5	3.0	5.0	9.0
2002.....	20.2	14.0	0.9	0.4	0.6	0.9	1.4	3.5	5.8	8.9
2001.....	19.2	13.2	1.3	0.5	0.5	0.9	1.7	3.5	6.0	8.9
2000.....	19.0	12.7	1.4	0.4	0.5	0.7	1.4	3.8	6.9	9.0
1999.....	19.3	12.5	1.2	0.7	0.5	0.7	1.8	3.9	7.2	9.3

White, both sexes

2004.....	19.9	9.9	1.1	0.4	0.3	0.5	0.9	1.8	3.7	8.2
2003.....	20.0	9.0	1.0	0.4	0.4	0.7	1.0	1.8	4.3	8.9
2002.....	19.7	12.4	1.0	0.4	0.5	0.7	0.9	2.3	4.5	9.6
2001.....	18.6	10.6	1.3	0.5	0.4	0.7	1.0	2.1	5.1	9.8
2000.....	18.2	10.6	1.2	0.4	0.4	0.6	1.0	2.3	5.8	10.3
1999.....	18.6	11.9	1.3	0.6	0.4	0.6	1.1	2.9	7.1	10.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

I. Certain infectious and parasitic diseases (A00-B99)

White, male

2004.....	20.9	11.6	1.1	0.4	0.3	0.5	1.2	2.0	5.2	11.6
2003.....	20.9	8.6	1.2	0.3	0.3	0.8	1.0	2.3	5.9	13.1
2002.....	20.9	12.7	1.2	0.5	0.6	0.6	1.1	2.9	6.3	14.5
2001.....	19.8	11.3	1.5	0.5	0.5	0.7	1.1	2.7	7.3	15.0
2000.....	19.3	11.2	1.2	0.5	0.4	0.8	1.2	2.9	8.5	15.9
1999.....	19.8	14.3	1.6	0.6	0.5	0.6	1.3	3.8	10.6	16.7

White, female

2004.....	19.0	8.1	1.1	0.4	0.3	0.5	0.7	1.5	2.2	4.7
2003.....	19.1	9.4	0.9	0.4	0.4	0.5	1.0	1.4	2.5	4.5
2002.....	18.6	12.1	0.8	0.4	0.5	0.8	0.8	1.6	2.6	4.6
2001.....	17.4	9.9	1.2	0.5	0.4	0.6	0.9	1.5	2.8	4.5
2000.....	17.2	10.0	1.2	0.4	0.5	0.5	0.7	1.8	2.9	4.6
1999.....	17.4	9.4	1.1	0.6	0.4	0.6	0.8	1.9	3.5	4.8

Black, both sexes

2004.....	40.4	21.8	1.5	*	0.8	1.6	5.3	13.2	27.2	42.5
2003.....	41.7	28.2	2.3	0.9	0.9	1.6	4.5	13.5	26.6	51.9
2002.....	42.9	24.6	1.7	*	1.2	1.5	4.6	14.8	32.8	54.9
2001.....	42.3	31.2	2.2	0.6	1.1	2.4	5.9	16.6	35.7	55.4
2000.....	43.2	29.9	2.4	0.8	0.7	1.6	5.4	17.6	41.6	56.7
1999.....	44.5	31.1	2.5	1.3	1.1	1.3	6.0	19.6	42.4	57.5

Black, male

2004.....	47.8	24.3	*	*	*	1.7	5.6	13.9	31.5	55.4
2003.....	50.0	31.8	2.5	*	*	1.3	4.6	14.8	33.6	68.5
2002.....	51.7	25.3	1.6	*	*	1.4	4.6	14.6	40.9	75.9
2001.....	51.2	33.2	2.6	*	*	2.4	5.6	18.6	46.4	76.4
2000.....	52.7	31.3	2.8	*	*	1.7	5.4	19.4	52.7	79.0
1999.....	54.6	33.2	3.2	1.3	1.4	1.3	5.1	23.8	56.2	78.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ years	year 2/ years	years	years	years	years	years	years	years	years

I. Certain infectious and parasitic diseases (A00-B99)

Black, female

2004.....	33.7	19.1	*	*	*	1.5	5.1	12.5	23.4	31.0
2003.....	34.2	24.4	2.0	*	*	2.0	4.5	12.3	20.4	37.2
2002.....	35.0	23.9	1.7	*	1.4	1.7	4.6	14.9	25.5	36.4
2001.....	34.2	29.0	1.8	*	*	2.3	6.2	14.8	26.2	36.6
2000.....	34.5	28.4	2.1	*	*	1.4	5.4	16.1	31.7	36.9
1999.....	35.3	28.9	1.9	1.3	*	*	6.9	15.8	30.1	39.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Zoonotic and other bacterial diseases (A20-A49)

All races 3/, both sexes

2004.....	12.0	7.5	0.7	0.2	0.2	0.3	0.5	0.7	1.1	1.6
2003.....	12.3	7.9	0.7	0.2	0.2	0.4	0.6	0.7	1.0	1.6
2002.....	12.3	8.6	0.6	0.3	0.3	0.4	0.4	0.7	1.0	1.6
2001.....	11.9	8.9	0.9	0.2	0.3	0.5	0.5	0.6	1.0	1.5
2000.....	11.6	8.6	0.8	0.3	0.2	0.4	0.5	0.6	0.9	1.6
1999.....	11.7	9.1	0.8	0.3	0.2	0.3	0.6	0.7	1.0	1.6

All races 3/, male

2004.....	11.0	8.9	0.7	0.2	*	0.3	0.5	0.6	1.2	1.7
2003.....	11.1	8.4	0.8	*	0.2	0.4	0.5	0.6	1.1	1.7
2002.....	11.1	8.6	0.7	0.3	0.4	0.4	0.5	0.8	1.0	1.6
2001.....	10.8	9.5	0.9	0.2	0.3	0.5	0.5	0.6	1.0	1.7
2000.....	10.4	8.9	0.9	0.3	*	0.5	0.5	0.7	1.0	1.6
1999.....	10.5	10.4	0.9	0.3	0.2	0.3	0.7	0.7	1.1	1.7

All races 3/, female

2004.....	12.9	6.1	0.7	0.3	0.2	0.3	0.5	0.7	1.1	1.5
2003.....	13.5	7.5	0.5	0.3	0.3	0.4	0.6	0.8	0.8	1.6
2002.....	13.5	8.5	0.6	0.2	0.3	0.4	0.4	0.7	0.9	1.6
2001.....	12.9	8.3	0.8	*	0.3	0.5	0.4	0.6	1.0	1.4
2000.....	12.9	8.3	0.7	0.2	*	0.3	0.4	0.5	0.9	1.5
1999.....	12.9	7.7	0.7	0.3	0.3	0.3	0.5	0.6	0.9	1.5

White, both sexes

2004.....	11.8	6.1	0.6	0.3	0.2	0.3	0.4	0.6	0.9	1.4
2003.....	12.1	5.6	0.5	0.2	0.2	0.3	0.5	0.6	0.8	1.3
2002.....	12.1	7.3	0.6	0.3	0.3	0.4	0.4	0.6	0.8	1.3
2001.....	11.7	6.8	0.8	0.2	0.3	0.5	0.3	0.5	0.7	1.2
2000.....	11.3	7.1	0.7	0.2	0.2	0.4	0.4	0.5	0.7	1.2
1999.....	11.3	7.2	0.8	0.3	0.2	0.3	0.5	0.5	0.8	1.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Zoonotic and other bacterial diseases (A20-A49)

White, male

2004.....	10.8	7.3	0.6	0.3	*	0.3	0.4	0.5	0.9	1.5
2003.....	10.8	5.6	0.7	*	*	0.4	0.5	0.5	0.8	1.4
2002.....	10.9	7.3	0.7	0.4	0.3	0.3	0.4	0.7	0.8	1.3
2001.....	10.5	7.5	0.9	*	0.3	0.5	0.4	0.5	0.7	1.4
2000.....	10.0	7.2	0.8	0.3	*	0.5	0.5	0.5	0.8	1.3
1999.....	10.0	8.5	0.8	0.3	*	0.3	0.6	0.6	0.9	1.4

White, female

2004.....	12.8	4.9	0.6	*	*	0.3	0.4	0.6	0.8	1.3
2003.....	13.4	5.6	0.4	*	*	0.3	0.6	0.6	0.7	1.2
2002.....	13.4	7.4	0.5	*	*	0.4	0.4	0.5	0.8	1.3
2001.....	12.8	6.1	0.7	*	*	0.4	0.3	0.5	0.8	1.0
2000.....	12.6	7.0	0.7	*	*	0.3	0.3	0.5	0.6	1.1
1999.....	12.6	5.9	0.7	0.3	*	0.3	0.4	0.5	0.8	1.1

Black, both sexes

2004.....	16.2	15.1	0.9	*	*	*	1.2	1.4	2.7	2.8
2003.....	16.9	19.6	1.2	*	*	0.7	0.9	1.7	2.3	3.9
2002.....	16.9	14.8	0.8	*	0.6	*	0.7	1.8	2.2	3.8
2001.....	16.5	20.3	1.1	*	*	0.9	1.0	1.2	2.8	3.9
2000.....	16.8	18.4	1.2	*	*	*	0.9	1.6	2.7	4.2
1999.....	17.0	19.0	0.9	*	*	*	1.2	1.7	2.2	4.2

Black, male

2004.....	15.3	16.9	*	*	*	*	*	*	3.1	2.6
2003.....	16.0	22.0	*	*	*	*	*	1.6	2.9	3.9
2002.....	15.7	15.7	*	*	*	*	*	*	2.6	4.1
2001.....	15.6	20.7	*	*	*	*	1.4	*	2.8	4.0
2000.....	15.6	19.8	*	*	*	*	*	2.1	2.8	4.2
1999.....	16.3	20.5	*	*	*	*	*	*	2.8	4.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Zoonotic and other bacterial diseases (A20-A49)

Black, female

2004.....	17.1	13.3	*	*	*	*	*	1.5	2.3	3.0
2003.....	17.7	17.0	*	*	*	*	*	1.9	1.6	3.9
2002.....	18.0	13.9	*	*	*	*	*	2.2	2.0	3.5
2001.....	17.3	19.8	*	*	*	*	*	*	2.8	3.8
2000.....	18.0	16.9	*	*	*	*	*	*	2.6	4.2
1999.....	17.7	17.5	*	*	*	*	*	1.9	1.6	3.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Whooping cough (A37)

White, male

Table with 11 columns (years 2004-1999) and 11 rows (age groups) for White, male. All cells contain an asterisk (*).

White, female

Table with 11 columns (years 2004-1999) and 11 rows (age groups) for White, female. All cells contain an asterisk (*).

Black, both sexes

Table with 11 columns (years 2004-1999) and 11 rows (age groups) for Black, both sexes. All cells contain an asterisk (*).

Black, male

Table with 11 columns (years 2004-1999) and 11 rows (age groups) for Black, male. All cells contain an asterisk (*).

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Scarlet fever and erysipelas (A38,A46)

All races 3/, both sexes

Table with 11 columns (years 2004-1999) and 11 rows (age groups) showing asterisks for all data points.

All races 3/, male

Table with 11 columns (years 2004-1999) and 11 rows (age groups) showing asterisks for all data points.

All races 3/, female

Table with 11 columns (years 2004-1999) and 11 rows (age groups) showing asterisks for all data points.

White, both sexes

Table with 11 columns (years 2004-1999) and 11 rows (age groups) showing asterisks for all data points.

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Meningococcal infection (A39)

All races 3/, both sexes

2004.....	0.0	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	0.1	*	*	*
2002.....	0.1	*	*	*	*	0.1	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	*	*	*	*
2000.....	0.1	0.6	*	*	*	0.2	0.1	*	*	*
1999.....	0.1	0.8	0.2	*	*	0.1	*	*	*	*

All races 3/, male

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.0	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	*	*	*	*
1999.....	0.1	1.0	0.2	*	*	0.1	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Septicemia (A40-A41)

All races 3/, both sexes

2004.....	11.4	6.6	0.5	0.2	0.1	0.2	0.4	0.6	1.0	1.5
2003.....	11.7	6.9	0.5	0.2	0.2	0.3	0.5	0.6	0.9	1.5
2002.....	11.7	7.3	0.5	0.2	0.3	0.2	0.3	0.6	0.9	1.5
2001.....	11.3	7.7	0.7	0.1	0.2	0.3	0.4	0.5	0.9	1.4
2000.....	11.1	7.2	0.6	0.2	0.1	0.2	0.3	0.5	0.9	1.4
1999.....	11.0	7.5	0.6	0.2	0.1	0.2	0.5	0.5	0.8	1.4

All races 3/, male

2004.....	10.4	8.0	0.5	*	*	0.2	0.4	0.5	1.1	1.5
2003.....	10.5	7.4	0.7	*	*	0.3	0.4	0.6	1.0	1.6
2002.....	10.6	7.7	0.5	0.3	0.2	0.2	0.4	0.7	1.0	1.5
2001.....	10.2	8.4	0.8	*	0.2	0.3	0.4	0.5	0.9	1.5
2000.....	9.8	7.1	0.7	0.2	*	0.2	0.4	0.6	0.9	1.4
1999.....	9.8	9.0	0.6	0.2	*	*	0.5	0.5	0.9	1.5

All races 3/, female

2004.....	12.3	5.2	0.6	0.2	*	0.2	0.4	0.7	1.0	1.4
2003.....	12.9	6.5	0.4	0.2	*	0.2	0.5	0.7	0.8	1.4
2002.....	12.9	7.0	0.5	*	0.3	0.3	0.3	0.6	0.9	1.5
2001.....	12.4	7.0	0.6	*	*	0.3	0.3	0.5	0.9	1.2
2000.....	12.3	7.3	0.6	*	*	*	0.2	0.4	0.8	1.4
1999.....	12.2	5.8	0.5	0.2	*	*	0.4	0.5	0.7	1.3

White, both sexes

2004.....	11.2	5.2	0.5	0.2	*	0.2	0.3	0.5	0.8	1.3
2003.....	11.5	4.6	0.4	0.2	0.2	0.2	0.4	0.5	0.7	1.2
2002.....	11.5	6.1	0.5	0.2	0.2	0.2	0.3	0.5	0.8	1.2
2001.....	11.1	5.4	0.7	0.2	0.2	0.2	0.3	0.4	0.7	1.1
2000.....	10.8	5.7	0.6	0.2	0.1	0.2	0.2	0.4	0.6	1.1
1999.....	10.6	5.5	0.5	0.2	0.1	0.2	0.4	0.4	0.7	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Septicemia (A40-A41)

White, male

2004.....	10.2	6.3	0.5	*	*	*	0.3	0.4	0.8	1.4
2003.....	10.2	4.6	0.6	*	*	0.3	0.4	0.4	0.8	1.3
2002.....	10.3	6.2	0.5	0.3	0.2	*	0.4	0.6	0.8	1.2
2001.....	9.9	6.2	0.8	*	*	0.3	0.3	0.4	0.6	1.3
2000.....	9.4	5.8	0.6	*	*	*	0.3	0.4	0.7	1.2
1999.....	9.3	6.9	0.5	*	*	*	0.5	0.5	0.8	1.2

White, female

2004.....	12.2	4.0	0.5	*	*	*	0.3	0.6	0.8	1.2
2003.....	12.8	4.5	*	*	*	*	0.4	0.6	0.7	1.1
2002.....	12.8	6.0	0.4	*	*	0.3	*	0.4	0.8	1.2
2001.....	12.3	4.6	0.6	*	*	*	0.3	0.4	0.7	0.9
2000.....	12.1	5.6	0.5	*	*	*	*	0.4	0.5	1.0
1999.....	11.9	4.0	0.5	*	*	*	*	0.4	0.6	0.9

Black, both sexes

2004.....	15.6	14.5	*	*	*	*	1.0	1.3	2.5	2.7
2003.....	16.3	18.6	0.9	*	*	*	0.7	1.6	2.2	3.6
2002.....	16.3	13.5	*	*	*	*	*	1.6	2.2	3.3
2001.....	15.8	19.3	0.9	*	*	*	0.7	1.1	2.5	3.7
2000.....	16.2	16.4	1.0	*	*	*	0.8	1.5	2.5	3.7
1999.....	16.1	17.7	*	*	*	*	1.0	1.4	1.7	3.7

Black, male

2004.....	14.6	16.3	*	*	*	*	*	*	2.8	2.5
2003.....	15.3	21.1	*	*	*	*	*	1.6	2.7	3.6
2002.....	15.0	14.8	*	*	*	*	*	*	2.5	3.7
2001.....	14.9	20.1	*	*	*	*	*	*	2.5	3.7
2000.....	14.9	16.0	*	*	*	*	*	2.1	2.5	3.5
1999.....	15.3	19.8	*	*	*	*	*	*	2.1	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Septicemia (A40-A41)

Black, female

2004.....	16.4	12.7	*	*	*	*	*	1.4	2.1	2.9
2003.....	17.2	16.1	*	*	*	*	*	1.6	1.6	3.5
2002.....	17.4	12.1	*	*	*	*	*	1.9	2.0	3.1
2001.....	16.7	18.6	*	*	*	*	*	*	2.5	3.7
2000.....	17.4	16.9	*	*	*	*	*	*	2.5	3.9
1999.....	16.9	15.4	*	*	*	*	*	1.6	1.4	3.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Viral diseases (A80-B34)

All races 3/, both sexes

2004.....	6.8	2.9	0.3	0.1	0.2	0.3	0.9	2.4	5.3	10.3
2003.....	7.1	2.9	0.4	0.2	0.2	0.3	0.8	2.5	5.8	12.1
2002.....	7.4	4.1	0.3	0.1	0.2	0.3	0.9	2.9	6.8	13.1
2001.....	7.4	3.0	0.4	0.2	0.2	0.4	1.1	3.3	7.7	13.4
2000.....	7.5	2.9	0.4	0.2	0.2	0.3	1.0	3.6	9.1	13.9
1999.....	7.6	3.5	0.5	0.3	0.2	0.3	1.0	4.1	10.0	14.1

All races 3/, male

2004.....	9.5	3.2	0.4	*	*	0.3	1.0	2.7	6.7	14.3
2003.....	10.1	2.5	0.4	*	*	0.3	0.9	3.0	7.7	17.2
2002.....	10.6	4.6	0.4	*	0.2	0.3	1.0	3.3	9.1	19.3
2001.....	10.5	3.1	0.5	0.2	*	0.4	1.0	3.8	10.7	19.7
2000.....	10.8	2.9	0.4	*	0.2	0.3	1.1	4.1	12.3	20.7
1999.....	10.9	4.0	0.6	0.3	0.3	0.3	1.0	5.3	14.2	21.0

All races 3/, female

2004.....	4.1	2.6	0.3	*	*	0.3	0.8	2.1	3.9	6.3
2003.....	4.2	3.3	0.4	*	*	0.3	0.8	2.0	3.9	7.0
2002.....	4.4	3.6	*	*	0.2	0.4	0.8	2.6	4.5	6.9
2001.....	4.3	2.8	0.3	0.2	*	0.3	1.2	2.7	4.6	7.1
2000.....	4.4	2.9	0.4	*	0.3	0.3	0.9	3.1	5.8	7.1
1999.....	4.3	3.1	0.4	0.3	*	0.3	1.0	2.9	5.8	7.2

White, both sexes

2004.....	4.6	2.6	0.3	*	*	0.2	0.4	1.0	2.6	6.2
2003.....	4.9	2.3	0.4	*	*	0.2	0.4	1.2	3.2	7.1
2002.....	5.1	3.7	0.3	*	0.2	0.2	0.4	1.5	3.4	7.9
2001.....	5.0	2.3	0.3	0.2	*	0.2	0.5	1.4	4.1	8.1
2000.....	5.1	2.1	0.3	0.2	0.2	0.2	0.4	1.7	4.8	8.7
1999.....	5.2	2.8	0.4	0.2	0.1	0.2	0.5	2.1	5.9	8.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Viral diseases (A80-B34)

White, male

2004.....	6.8	2.8	0.3	*	*	*	0.5	1.3	4.0	9.4
2003.....	7.2	1.9	0.3	*	*	0.3	0.5	1.6	4.8	11.0
2002.....	7.5	4.1	0.4	*	*	*	0.5	2.0	5.2	12.8
2001.....	7.4	2.4	0.4	*	*	*	0.6	1.9	6.3	13.0
2000.....	7.5	2.2	*	*	*	0.2	0.6	2.2	7.4	14.0
1999.....	7.6	3.4	0.6	*	*	*	0.6	2.9	9.2	14.5

White, female

2004.....	2.5	2.4	*	*	*	*	*	0.6	1.2	3.0
2003.....	2.7	2.7	0.4	*	*	*	0.3	0.7	1.6	3.0
2002.....	2.8	3.3	*	*	*	0.3	0.3	0.9	1.6	2.9
2001.....	2.7	2.2	*	*	*	*	0.4	0.9	1.8	3.2
2000.....	2.8	2.1	*	*	*	*	*	1.2	2.2	3.3
1999.....	2.8	2.1	*	*	*	*	0.4	1.2	2.4	3.3

Black, both sexes

2004.....	21.5	4.2	*	*	0.6	0.9	4.0	11.3	23.6	38.5
2003.....	22.3	5.0	*	*	*	0.8	3.4	11.3	23.6	46.9
2002.....	23.5	5.8	*	*	*	1.0	3.5	12.3	29.5	49.7
2001.....	23.7	6.6	0.8	*	0.6	1.3	4.7	15.0	31.8	50.1
2000.....	24.1	7.3	0.9	*	*	0.9	4.2	15.7	37.9	50.9
1999.....	24.3	7.3	1.1	0.9	0.6	0.8	4.3	17.0	38.3	51.0

Black, male

2004.....	29.4	*	*	*	*	*	4.1	12.2	27.1	51.1
2003.....	31.1	*	*	*	*	*	3.6	12.8	29.3	63.3
2002.....	33.0	6.4	*	*	*	*	3.7	12.4	37.1	69.8
2001.....	33.1	7.5	*	*	*	1.4	3.9	17.1	42.7	70.6
2000.....	34.3	7.0	*	*	*	*	4.0	16.9	48.7	72.9
1999.....	35.0	7.5	*	*	*	*	3.8	21.5	51.2	71.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Viral diseases (A80-B34)

Black, female

2004.....	14.2	*	*	*	*	*	3.9	10.4	20.6	27.2
2003.....	14.2	*	*	*	*	*	3.3	9.9	18.4	32.3
2002.....	14.9	*	*	*	*	*	3.3	12.2	22.7	31.9
2001.....	15.0	*	*	*	*	*	5.4	13.0	22.1	31.9
2000.....	14.8	7.6	*	*	*	*	4.3	14.6	28.3	31.5
1999.....	14.6	7.0	*	*	*	*	4.9	13.0	26.7	33.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Rabies (A82)

White, male

Table with 11 columns (years 1999-2004) and 11 rows (age groups) for White, male. All cells contain an asterisk (*).

White, female

Table with 11 columns (years 1999-2004) and 11 rows (age groups) for White, female. All cells contain an asterisk (*).

Black, both sexes

Table with 11 columns (years 1999-2004) and 11 rows (age groups) for Black, both sexes. All cells contain an asterisk (*).

Black, male

Table with 11 columns (years 1999-2004) and 11 rows (age groups) for Black, male. All cells contain an asterisk (*).

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Viral hepatitis (B15-B19)

All races 3/, both sexes

2004.....	1.8	*	*	*	*	*	*	0.1	0.2	0.7
2003.....	1.9	*	*	*	*	*	*	0.1	0.2	0.8
2002.....	2.0	*	*	*	*	*	*	0.2	0.3	1.0
2001.....	2.0	*	*	*	*	*	*	*	0.3	0.9
2000.....	1.9	*	*	*	*	*	*	0.1	0.3	1.0
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9

All races 3/, male

2004.....	2.4	*	*	*	*	*	*	*	0.2	0.8
2003.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2002.....	2.7	*	*	*	*	*	*	0.2	0.4	1.4
2001.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.3
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.2

All races 3/, female

2004.....	1.3	*	*	*	*	*	*	*	*	0.5
2003.....	1.3	*	*	*	*	*	*	*	*	0.4
2002.....	1.4	*	*	*	*	*	*	*	*	0.6
2001.....	1.4	*	*	*	*	*	*	*	0.2	0.6
2000.....	1.4	*	*	*	*	*	*	*	0.2	0.6
1999.....	1.3	*	*	*	*	*	*	*	0.2	0.6

White, both sexes

2004.....	1.8	*	*	*	*	*	*	*	0.2	0.6
2003.....	1.8	*	*	*	*	*	*	*	0.2	0.8
2002.....	2.0	*	*	*	*	*	*	0.1	0.3	1.0
2001.....	1.9	*	*	*	*	*	*	*	0.2	0.9
2000.....	1.8	*	*	*	*	*	*	*	0.2	1.0
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All races 3/, both sexes

2004.....	4.4	*	*	*	*	0.1	0.8	2.2	5.1	9.5
2003.....	4.7	*	*	*	0.1	0.2	0.7	2.3	5.5	11.2
2002.....	4.9	*	*	*	0.1	0.2	0.7	2.6	6.4	11.9
2001.....	5.0	*	*	*	0.1	0.2	0.9	3.1	7.3	12.3
2000.....	5.1	*	*	*	0.1	0.2	0.8	3.3	8.7	12.8
1999.....	5.3	*	0.2	0.2	0.1	0.2	0.9	3.9	9.5	13.0

All races 3/, male

2004.....	6.6	*	*	*	*	*	0.9	2.6	6.4	13.3
2003.....	7.1	*	*	*	*	*	0.8	2.8	7.3	15.9
2002.....	7.4	*	*	*	*	*	0.7	3.0	8.5	17.7
2001.....	7.6	*	*	*	*	0.2	0.8	3.7	10.2	18.4
2000.....	7.9	*	*	*	*	*	0.8	3.8	11.9	19.2
1999.....	8.2	*	*	0.2	*	*	0.8	5.1	13.6	19.5

All races 3/, female

2004.....	2.4	*	*	*	*	*	0.7	1.8	3.7	5.6
2003.....	2.4	*	*	*	*	*	0.7	1.8	3.7	6.5
2002.....	2.5	*	*	*	*	0.2	0.6	2.3	4.2	6.2
2001.....	2.5	*	*	*	*	0.2	1.0	2.5	4.4	6.3
2000.....	2.5	*	*	*	*	*	0.7	2.9	5.5	6.4
1999.....	2.5	*	*	*	*	*	1.0	2.7	5.4	6.6

White, both sexes

2004.....	2.4	*	*	*	*	*	0.2	0.8	2.3	5.4
2003.....	2.5	*	*	*	*	*	0.3	0.9	2.9	6.1
2002.....	2.6	*	*	*	*	*	0.3	1.2	3.0	6.8
2001.....	2.7	*	*	*	*	*	0.3	1.3	3.8	7.1
2000.....	2.8	*	*	*	*	*	0.3	1.5	4.5	7.6
1999.....	3.0	*	*	*	*	*	0.4	2.0	5.4	7.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease (B20-B24)

White, male

2004.....	3.9	*	*	*	*	*	0.4	1.2	3.6	8.4
2003.....	4.2	*	*	*	*	*	0.3	1.4	4.5	9.7
2002.....	4.3	*	*	*	*	*	0.3	1.8	4.7	11.1
2001.....	4.5	*	*	*	*	*	0.4	1.8	5.8	11.6
2000.....	4.7	*	*	*	*	*	0.4	2.0	7.1	12.5
1999.....	5.0	*	*	*	*	*	0.4	2.8	8.6	13.0

White, female

2004.....	0.8	*	*	*	*	*	*	0.4	1.0	2.4
2003.....	0.9	*	*	*	*	*	*	0.5	1.3	2.4
2002.....	0.9	*	*	*	*	*	*	0.7	1.3	2.3
2001.....	0.9	*	*	*	*	*	0.3	0.8	1.6	2.4
2000.....	1.0	*	*	*	*	*	*	1.0	1.9	2.6
1999.....	1.0	*	*	*	*	*	0.3	1.0	2.1	2.7

Black, both sexes

2004.....	18.8	*	*	*	*	0.7	3.7	10.8	23.2	37.4
2003.....	19.6	*	*	*	*	0.6	3.2	11.2	23.2	46.0
2002.....	20.8	*	*	*	*	0.8	3.3	11.7	29.1	48.2
2001.....	21.1	*	*	*	*	1.1	4.4	14.5	31.5	48.8
2000.....	21.4	*	*	*	*	0.7	3.7	15.2	37.1	49.8
1999.....	21.8	*	*	0.7	*	0.7	4.1	16.6	37.8	49.9

Black, male

2004.....	25.9	*	*	*	*	*	3.8	11.9	26.5	50.1
2003.....	27.7	*	*	*	*	*	3.2	12.7	28.6	61.9
2002.....	29.5	*	*	*	*	*	3.3	11.7	36.6	68.4
2001.....	30.1	*	*	*	*	*	3.5	16.8	42.3	69.2
2000.....	31.0	*	*	*	*	*	3.6	16.4	47.9	71.2
1999.....	31.9	*	*	*	*	*	3.5	21.1	50.6	69.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Black, female

2004.....	12.4	*	*	*	*	*	3.7	9.8	20.3	25.9
2003.....	12.3	*	*	*	*	*	3.1	9.8	18.3	31.9
2002.....	12.8	*	*	*	*	*	3.3	11.7	22.2	30.3
2001.....	12.9	*	*	*	*	*	5.2	12.4	21.8	30.7
2000.....	12.8	*	*	*	*	*	3.8	14.1	27.4	30.8
1999.....	12.6	*	*	*	*	*	4.7	12.5	26.3	32.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

All races 3/, both sexes

2004.....	1.7	*	*	*	*	*	0.3	0.9	2.1	3.7
2003.....	1.8	*	*	*	*	*	0.3	1.0	2.2	4.3
2002.....	1.9	*	*	*	*	*	0.3	1.1	2.5	4.9
2001.....	1.9	*	*	*	*	*	0.4	1.3	2.9	4.8
2000.....	2.0	*	*	*	*	*	0.4	1.4	3.4	5.2
1999.....	2.1	*	*	*	*	*	0.4	1.7	3.8	5.4

All races 3/, male

2004.....	2.6	*	*	*	*	*	0.4	1.1	2.7	5.2
2003.....	2.7	*	*	*	*	*	0.3	1.1	3.0	5.9
2002.....	2.9	*	*	*	*	*	0.4	1.2	3.3	7.1
2001.....	2.9	*	*	*	*	*	0.4	1.6	3.9	7.1
2000.....	3.1	*	*	*	*	*	0.4	1.7	4.7	7.8
1999.....	3.3	*	*	*	*	*	0.4	2.2	5.3	8.0

All races 3/, female

2004.....	1.0	*	*	*	*	*	0.3	0.8	1.6	2.1
2003.....	0.9	*	*	*	*	*	0.2	0.8	1.4	2.6
2002.....	1.0	*	*	*	*	*	0.2	1.0	1.8	2.7
2001.....	1.0	*	*	*	*	*	0.4	1.0	1.9	2.5
2000.....	1.0	*	*	*	*	*	0.4	1.1	2.1	2.6
1999.....	1.0	*	*	*	*	*	0.5	1.2	2.4	2.8

White, both sexes

2004.....	1.0	*	*	*	*	*	*	0.4	1.0	2.3
2003.....	1.0	*	*	*	*	*	*	0.4	1.2	2.5
2002.....	1.1	*	*	*	*	*	0.1	0.5	1.2	2.9
2001.....	1.1	*	*	*	*	*	0.2	0.6	1.6	2.9
2000.....	1.1	*	*	*	*	*	0.2	0.7	1.8	3.1
1999.....	1.2	*	*	*	*	*	0.1	0.9	2.2	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
White, male										
2004.....	1.6	*	*	*	*	*	*	0.6	1.5	3.5
2003.....	1.7	*	*	*	*	*	*	0.6	1.9	3.8
2002.....	1.7	*	*	*	*	*	*	0.8	1.7	4.6
2001.....	1.8	*	*	*	*	*	*	0.9	2.4	4.6
2000.....	1.9	*	*	*	*	*	*	0.9	2.9	5.0
1999.....	2.0	*	*	*	*	*	*	1.3	3.3	5.3
White, female										
2004.....	0.4	*	*	*	*	*	*	*	0.5	1.0
2003.....	0.4	*	*	*	*	*	*	*	0.5	1.1
2002.....	0.4	*	*	*	*	*	*	0.3	0.7	1.2
2001.....	0.4	*	*	*	*	*	*	0.3	0.7	1.2
2000.....	0.4	*	*	*	*	*	*	0.5	0.8	1.2
1999.....	0.4	*	*	*	*	*	*	0.5	1.0	1.3
Black, both sexes										
2004.....	7.2	*	*	*	*	*	1.8	4.1	9.8	13.5
2003.....	7.1	*	*	*	*	*	1.1	4.4	8.9	16.5
2002.....	8.0	*	*	*	*	*	1.4	4.5	11.4	18.6
2001.....	8.0	*	*	*	*	*	1.8	5.6	12.0	18.1
2000.....	8.3	*	*	*	*	*	1.6	6.1	14.2	20.2
1999.....	8.6	*	*	*	*	*	2.1	6.9	15.1	20.1
Black, male										
2004.....	9.7	*	*	*	*	*	1.7	4.0	11.3	18.1
2003.....	10.2	*	*	*	*	*	*	4.9	11.4	21.5
2002.....	11.4	*	*	*	*	*	1.6	4.0	14.8	25.8
2001.....	11.3	*	*	*	*	*	1.9	6.1	15.2	26.2
2000.....	11.9	*	*	*	*	*	1.5	7.1	18.7	29.5
1999.....	12.6	*	*	*	*	*	1.7	8.3	19.7	28.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

Black, female

2004.....	4.9	*	*	*	*	*	1.8	4.3	8.4	9.4
2003.....	4.3	*	*	*	*	*	*	3.9	6.6	12.0
2002.....	4.9	*	*	*	*	*	*	5.0	8.3	12.3
2001.....	4.9	*	*	*	*	*	1.7	5.2	9.1	11.0
2000.....	5.0	*	*	*	*	*	1.8	5.2	10.2	11.9
1999.....	5.0	*	*	*	*	*	2.5	5.6	11.0	12.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)

All races 3/, both sexes

2004.....	0.3	*	*	*	*	*	*	*	0.3	0.7
2003.....	0.3	*	*	*	*	*	*	0.1	0.3	0.7
2002.....	0.3	*	*	*	*	*	*	0.1	0.3	0.7
2001.....	0.3	*	*	*	*	*	*	0.2	0.5	0.7
2000.....	0.4	*	*	*	*	*	*	0.2	0.5	0.9
1999.....	0.4	*	*	*	*	*	*	0.2	0.8	1.0

All races 3/, male

2004.....	0.5	*	*	*	*	*	*	*	0.5	1.1
2003.....	0.5	*	*	*	*	*	*	*	0.4	1.1
2002.....	0.5	*	*	*	*	*	*	*	0.5	1.2
2001.....	0.6	*	*	*	*	*	*	0.2	0.7	1.3
2000.....	0.6	*	*	*	*	*	*	0.2	0.8	1.5
1999.....	0.7	*	*	*	*	*	*	0.3	1.3	1.7

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	0.2
2003.....	0.1	*	*	*	*	*	*	*	*	0.2
2002.....	0.1	*	*	*	*	*	*	*	*	0.2
2001.....	0.1	*	*	*	*	*	*	*	0.2	*
2000.....	0.1	*	*	*	*	*	*	*	0.2	0.2
1999.....	0.1	*	*	*	*	*	*	*	0.2	0.3

White, both sexes

2004.....	0.2	*	*	*	*	*	*	*	0.2	0.5
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.5
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.5
2001.....	0.2	*	*	*	*	*	*	*	0.3	0.6
2000.....	0.3	*	*	*	*	*	*	*	0.4	0.6
1999.....	0.3	*	*	*	*	*	*	*	0.6	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)

White, male

2004.....	0.4	*	*	*	*	*	*	*	0.3	0.9
2003.....	0.4	*	*	*	*	*	*	*	0.3	0.9
2002.....	0.4	*	*	*	*	*	*	*	0.4	1.0
2001.....	0.4	*	*	*	*	*	*	*	0.5	1.1
2000.....	0.5	*	*	*	*	*	*	*	0.7	1.2
1999.....	0.5	*	*	*	*	*	*	*	1.1	1.5

White, female

2004.....	0.0	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

Black, both sexes

2004.....	1.0	*	*	*	*	*	*	*	1.0	1.9
2003.....	1.0	*	*	*	*	*	*	*	0.9	1.9
2002.....	1.1	*	*	*	*	*	*	*	1.0	2.1
2001.....	1.1	*	*	*	*	*	*	*	1.6	2.1
2000.....	1.2	*	*	*	*	*	*	0.7	1.5	2.6
1999.....	1.2	*	*	*	*	*	*	0.8	2.1	2.5

Black, male

2004.....	1.6	*	*	*	*	*	*	*	*	3.1
2003.....	1.4	*	*	*	*	*	*	*	*	3.0
2002.....	1.6	*	*	*	*	*	*	*	*	3.4
2001.....	1.7	*	*	*	*	*	*	*	2.7	3.4
2000.....	1.9	*	*	*	*	*	*	*	2.2	4.3
1999.....	1.9	*	*	*	*	*	*	*	3.2	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)

All races 3/, both sexes

2004.....	0.4	*	*	*	*	*	*	0.2	0.5	0.9
2003.....	0.5	*	*	*	*	*	*	0.2	0.5	1.2
2002.....	0.5	*	*	*	*	*	*	0.3	0.7	1.0
2001.....	0.5	*	*	*	*	*	*	0.3	0.8	1.3
2000.....	0.6	*	*	*	*	*	*	0.3	1.1	1.5
1999.....	0.6	*	*	*	*	*	*	0.4	1.1	1.5

All races 3/, male

2004.....	0.7	*	*	*	*	*	*	0.3	0.6	1.3
2003.....	0.7	*	*	*	*	*	*	0.3	0.6	1.8
2002.....	0.8	*	*	*	*	*	*	0.4	1.1	1.7
2001.....	0.8	*	*	*	*	*	*	0.4	1.2	2.2
2000.....	0.9	*	*	*	*	*	*	0.4	1.6	2.3
1999.....	0.9	*	*	*	*	*	*	0.6	1.6	2.5

All races 3/, female

2004.....	0.2	*	*	*	*	*	*	*	0.4	0.4
2003.....	0.2	*	*	*	*	*	*	*	0.3	0.6
2002.....	0.2	*	*	*	*	*	*	0.3	0.4	0.4
2001.....	0.2	*	*	*	*	*	*	*	0.4	0.5
2000.....	0.2	*	*	*	*	*	*	0.2	0.6	0.7
1999.....	0.2	*	*	*	*	*	*	0.2	0.5	0.6

White, both sexes

2004.....	0.3	*	*	*	*	*	*	*	0.3	0.6
2003.....	0.3	*	*	*	*	*	*	*	0.3	0.8
2002.....	0.3	*	*	*	*	*	*	0.1	0.4	0.6
2001.....	0.3	*	*	*	*	*	*	0.1	0.5	0.9
2000.....	0.4	*	*	*	*	*	*	0.2	0.7	1.0
1999.....	0.4	*	*	*	*	*	*	0.2	0.7	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)

	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
White, male										
2004.....	0.5	*	*	*	*	*	*	*	0.4	1.0
2003.....	0.5	*	*	*	*	*	*	*	0.4	1.4
2002.....	0.5	*	*	*	*	*	*	*	0.8	1.2
2001.....	0.5	*	*	*	*	*	*	*	0.8	1.5
2000.....	0.6	*	*	*	*	*	*	0.3	1.1	1.8
1999.....	0.6	*	*	*	*	*	*	0.3	1.1	1.9
White, female										
2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	0.3
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	0.2
1999.....	0.1	*	*	*	*	*	*	*	*	0.3
Black, both sexes										
2004.....	1.7	*	*	*	*	*	*	1.2	1.8	3.1
2003.....	1.7	*	*	*	*	*	*	1.0	1.6	3.7
2002.....	1.7	*	*	*	*	*	*	1.4	2.6	3.6
2001.....	1.9	*	*	*	*	*	*	1.3	3.1	4.7
2000.....	2.1	*	*	*	*	*	*	1.6	4.0	4.7
1999.....	2.1	*	*	*	*	*	*	1.5	3.8	5.1
Black, male										
2004.....	2.3	*	*	*	*	*	*	*	1.6	4.0
2003.....	2.5	*	*	*	*	*	*	*	2.1	4.9
2002.....	2.5	*	*	*	*	*	*	*	3.5	5.5
2001.....	2.8	*	*	*	*	*	*	1.7	4.1	7.3
2000.....	3.2	*	*	*	*	*	*	1.8	5.4	6.2
1999.....	3.2	*	*	*	*	*	*	2.1	5.7	7.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ year 1/	year 2/ year 1/	years	years	years	years	years	years	years	years

Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)

Black, female

2004.....	1.1	*	*	*	*	*	*	*	1.9	2.2
2003.....	0.9	*	*	*	*	*	*	*	*	2.6
2002.....	1.0	*	*	*	*	*	*	1.4	1.9	1.8
2001.....	1.1	*	*	*	*	*	*	*	2.1	2.5
2000.....	1.2	*	*	*	*	*	*	*	2.8	3.5
1999.....	1.1	*	*	*	*	*	*	*	2.2	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

All races 3/, both sexes

2004.....	0.7	*	*	*	*	*	*	0.3	0.8	1.4
2003.....	1.1	*	*	*	*	*	0.2	0.5	1.2	2.4
2002.....	1.2	*	*	*	*	*	0.1	0.6	1.4	2.6
2001.....	1.2	*	*	*	*	*	0.3	0.8	1.6	2.9
2000.....	1.2	*	*	*	*	*	0.1	0.7	1.9	2.9
1999.....	1.2	*	*	*	*	*	0.2	0.8	2.0	2.8

All races 3/, male

2004.....	1.1	*	*	*	*	*	*	0.4	1.0	1.9
2003.....	1.6	*	*	*	*	*	*	0.5	1.5	3.3
2002.....	1.7	*	*	*	*	*	*	0.6	1.9	3.9
2001.....	1.8	*	*	*	*	*	*	0.8	2.2	4.2
2000.....	1.8	*	*	*	*	*	*	0.7	2.4	4.1
1999.....	1.8	*	*	*	*	*	*	1.1	2.7	3.9

All races 3/, female

2004.....	0.4	*	*	*	*	*	*	0.3	0.5	0.9
2003.....	0.6	*	*	*	*	*	0.2	0.5	0.9	1.5
2002.....	0.7	*	*	*	*	*	*	0.5	0.9	1.4
2001.....	0.7	*	*	*	*	*	0.4	0.7	1.0	1.7
2000.....	0.7	*	*	*	*	*	*	0.7	1.4	1.6
1999.....	0.7	*	*	*	*	*	0.2	0.6	1.3	1.7

White, both sexes

2004.....	0.3	*	*	*	*	*	*	*	0.3	0.6
2003.....	0.5	*	*	*	*	*	*	0.1	0.5	1.1
2002.....	0.5	*	*	*	*	*	*	0.2	0.5	1.3
2001.....	0.6	*	*	*	*	*	*	0.2	0.7	1.5
2000.....	0.6	*	*	*	*	*	*	0.3	0.8	1.5
1999.....	0.6	*	*	*	*	*	*	0.3	0.9	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

White, male

2004.....	0.5	*	*	*	*	*	*	*	0.4	0.9
2003.....	0.8	*	*	*	*	*	*	*	0.7	1.7
2002.....	0.9	*	*	*	*	*	*	*	0.8	2.2
2001.....	0.9	*	*	*	*	*	*	*	1.1	2.4
2000.....	0.9	*	*	*	*	*	*	0.3	1.2	2.5
1999.....	0.9	*	*	*	*	*	*	0.5	1.4	2.2

White, female

2004.....	0.1	*	*	*	*	*	*	*	*	0.3
2003.....	0.2	*	*	*	*	*	*	*	0.3	0.5
2002.....	0.2	*	*	*	*	*	*	*	*	0.5
2001.....	0.2	*	*	*	*	*	*	*	0.4	0.6
2000.....	0.2	*	*	*	*	*	*	*	0.5	0.6
1999.....	0.2	*	*	*	*	*	*	*	0.4	0.6

Black, both sexes

2004.....	3.5	*	*	*	*	*	*	1.8	3.9	6.9
2003.....	5.3	*	*	*	*	*	1.0	2.8	6.0	11.5
2002.....	5.7	*	*	*	*	*	*	3.0	7.1	11.7
2001.....	5.7	*	*	*	*	*	1.2	4.1	7.6	12.8
2000.....	5.5	*	*	*	*	*	*	3.9	9.1	12.3
1999.....	5.8	*	*	*	*	*	0.9	4.1	9.2	12.7

Black, male

2004.....	5.0	*	*	*	*	*	*	1.9	5.2	9.3
2003.....	7.3	*	*	*	*	*	*	2.5	7.2	15.1
2002.....	8.0	*	*	*	*	*	*	3.3	9.2	17.0
2001.....	8.0	*	*	*	*	*	*	4.7	10.4	17.9
2000.....	7.8	*	*	*	*	*	*	4.1	11.0	16.7
1999.....	8.3	*	*	*	*	*	*	5.4	11.9	17.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

Black, female

2004.....	2.2	*	*	*	*	*	*	1.8	2.7	4.8
2003.....	3.6	*	*	*	*	*	*	3.0	4.9	8.2
2002.....	3.6	*	*	*	*	*	*	2.6	5.2	7.0
2001.....	3.6	*	*	*	*	*	1.9	3.6	5.1	8.2
2000.....	3.5	*	*	*	*	*	*	3.7	7.3	8.4
1999.....	3.6	*	*	*	*	*	*	2.9	6.7	8.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Unspecified human immunodeficiency virus (HIV) disease (B24)

All races 3/, both sexes

2004.....	1.2	*	*	*	*	*	0.2	0.6	1.4	2.8
2003.....	1.0	*	*	*	*	*	0.2	0.5	1.4	2.7
2002.....	1.0	*	*	*	*	*	0.2	0.6	1.4	2.7
2001.....	1.0	*	*	*	*	*	0.2	0.6	1.5	2.5
2000.....	1.0	*	*	*	*	*	0.1	0.7	1.8	2.4
1999.....	1.0	*	*	*	*	*	0.2	0.8	1.9	2.3

All races 3/, male

2004.....	1.7	*	*	*	*	*	0.3	0.8	1.7	3.8
2003.....	1.5	*	*	*	*	*	*	0.7	1.8	3.8
2002.....	1.5	*	*	*	*	*	*	0.7	1.7	3.9
2001.....	1.5	*	*	*	*	*	*	0.7	2.2	3.6
2000.....	1.5	*	*	*	*	*	*	0.7	2.4	3.6
1999.....	1.5	*	*	*	*	*	*	0.9	2.7	3.5

All races 3/, female

2004.....	0.7	*	*	*	*	*	*	0.5	1.2	1.9
2003.....	0.5	*	*	*	*	*	*	0.4	1.0	1.5
2002.....	0.5	*	*	*	*	*	*	0.5	1.1	1.5
2001.....	0.5	*	*	*	*	*	0.2	0.4	0.9	1.4
2000.....	0.5	*	*	*	*	*	*	0.6	1.2	1.2
1999.....	0.5	*	*	*	*	*	*	0.6	1.1	1.2

White, both sexes

2004.....	0.6	*	*	*	*	*	*	0.2	0.6	1.5
2003.....	0.5	*	*	*	*	*	*	0.2	0.7	1.2
2002.....	0.5	*	*	*	*	*	*	0.3	0.6	1.3
2001.....	0.5	*	*	*	*	*	*	0.2	0.7	1.2
2000.....	0.5	*	*	*	*	*	*	0.3	0.8	1.3
1999.....	0.5	*	*	*	*	*	*	0.4	1.1	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Unspecified human immunodeficiency virus (HIV) disease (B24)

White, male

2004.....	1.0	*	*	*	*	*	*	*	0.9	2.2
2003.....	0.9	*	*	*	*	*	*	0.3	1.1	2.0
2002.....	0.8	*	*	*	*	*	*	0.4	0.9	2.2
2001.....	0.8	*	*	*	*	*	*	0.3	1.1	2.1
2000.....	0.8	*	*	*	*	*	*	0.5	1.3	2.1
1999.....	0.9	*	*	*	*	*	*	0.5	1.7	2.2

White, female

2004.....	0.2	*	*	*	*	*	*	*	0.3	0.8
2003.....	0.2	*	*	*	*	*	*	*	0.3	0.5
2002.....	0.2	*	*	*	*	*	*	*	0.3	0.4
2001.....	0.1	*	*	*	*	*	*	*	0.2	0.3
2000.....	0.2	*	*	*	*	*	*	*	0.4	0.4
1999.....	0.2	*	*	*	*	*	*	*	0.4	0.4

Black, both sexes

2004.....	5.4	*	*	*	*	*	1.1	3.4	6.8	12.0
2003.....	4.5	*	*	*	*	*	0.7	2.5	5.8	12.6
2002.....	4.3	*	*	*	*	*	0.8	2.4	6.9	12.2
2001.....	4.4	*	*	*	*	*	0.9	2.8	7.2	11.1
2000.....	4.3	*	*	*	*	*	0.8	2.9	8.4	10.1
1999.....	4.1	*	*	*	*	*	0.7	3.3	7.5	9.5

Black, male

2004.....	7.3	*	*	*	*	*	*	4.2	7.1	15.6
2003.....	6.3	*	*	*	*	*	*	3.2	6.8	17.5
2002.....	6.0	*	*	*	*	*	*	2.8	7.8	16.7
2001.....	6.3	*	*	*	*	*	*	3.5	9.9	14.5
2000.....	6.3	*	*	*	*	*	*	2.5	10.6	14.6
1999.....	5.9	*	*	*	*	*	*	4.0	10.1	13.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Unspecified human immunodeficiency virus (HIV) disease (B24)

Black, female

2004.....	3.7	*	*	*	*	*	*	2.7	6.5	8.8
2003.....	2.8	*	*	*	*	*	*	1.9	4.8	8.2
2002.....	2.8	*	*	*	*	*	*	2.1	6.2	8.2
2001.....	2.7	*	*	*	*	*	*	2.2	4.8	8.1
2000.....	2.6	*	*	*	*	*	*	3.3	6.3	6.1
1999.....	2.4	*	*	*	*	*	*	2.7	5.2	6.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)

All races 3/, both sexes

2004.....	0.3	2.3	0.3	*	*	0.1	0.1	*	0.1	0.1
2003.....	0.3	2.2	0.4	*	*	0.1	*	*	*	0.1
2002.....	0.4	3.2	0.3	*	*	*	0.1	*	0.1	0.2
2001.....	0.3	2.1	0.3	0.1	*	0.1	*	*	*	0.1
2000.....	0.4	1.8	0.3	*	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.4	2.7	0.3	0.1	*	*	*	*	0.1	0.2

All races 3/, male

2004.....	0.3	2.7	0.3	*	*	*	*	*	*	*
2003.....	0.3	1.8	0.4	*	*	*	*	*	*	*
2002.....	0.4	3.6	0.4	*	*	*	*	*	*	*
2001.....	0.3	2.1	0.4	*	*	*	*	*	*	*
2000.....	0.4	1.9	0.3	*	*	*	*	*	*	*
1999.....	0.4	2.8	0.4	*	*	*	*	*	*	0.2

All races 3/, female

2004.....	0.3	1.9	0.3	*	*	*	*	*	*	*
2003.....	0.3	2.6	0.4	*	*	*	*	*	*	*
2002.....	0.4	2.8	*	*	*	*	*	*	*	*
2001.....	0.3	2.0	0.3	*	*	*	*	*	*	*
2000.....	0.4	1.7	*	*	*	*	*	*	*	*
1999.....	0.4	2.5	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.3	2.1	0.2	*	*	*	*	*	*	0.1
2003.....	0.4	1.8	0.3	*	*	*	*	*	*	*
2002.....	0.4	3.1	0.3	*	*	*	*	*	0.1	0.1
2001.....	0.3	1.7	0.3	*	*	*	*	*	*	*
2000.....	0.4	1.5	0.3	*	*	*	*	*	*	0.1
1999.....	0.4	2.3	0.3	*	*	*	*	*	*	0.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

All races 3/, both sexes

2004.....	1.5	1.0	*	*	*	*	0.1	0.2	0.2	0.5
2003.....	1.5	1.1	*	*	*	*	*	0.2	0.3	0.5
2002.....	1.3	1.3	*	*	*	*	0.1	0.2	0.3	0.4
2001.....	1.1	1.6	*	*	*	*	0.1	0.2	0.3	0.5
2000.....	1.0	1.3	*	*	*	*	*	0.1	0.3	0.4
1999.....	1.4	1.6	0.1	*	*	*	0.1	0.3	0.5	0.7

All races 3/, male

2004.....	1.8	1.3	*	*	*	*	0.2	*	0.3	0.6
2003.....	1.7	1.3	*	*	*	*	*	0.2	0.3	0.6
2002.....	1.5	1.5	*	*	*	*	*	0.2	0.4	0.6
2001.....	1.2	1.6	*	*	*	*	*	0.2	0.3	0.6
2000.....	1.1	1.5	*	*	*	*	*	*	0.3	0.6
1999.....	1.6	1.9	*	*	*	*	*	0.4	0.5	0.8

All races 3/, female

2004.....	1.3	*	*	*	*	*	*	*	*	0.4
2003.....	1.3	*	*	*	*	*	*	*	0.2	0.4
2002.....	1.0	1.2	*	*	*	*	*	*	0.3	0.3
2001.....	1.0	1.6	*	*	*	*	*	*	0.2	0.4
2000.....	0.9	1.1	*	*	*	*	*	*	0.2	0.3
1999.....	1.3	1.2	*	*	*	*	*	0.3	0.4	0.5

White, both sexes

2004.....	1.6	0.8	*	*	*	*	*	0.1	0.2	0.4
2003.....	1.5	0.9	*	*	*	*	*	*	0.2	0.4
2002.....	1.3	1.1	*	*	*	*	*	0.2	0.2	0.3
2001.....	1.1	1.2	*	*	*	*	*	0.2	0.2	0.4
2000.....	1.0	1.0	*	*	*	*	*	*	0.2	0.3
1999.....	1.3	1.2	*	*	*	*	*	0.2	0.3	0.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

White, male

2004.....	1.8	*	*	*	*	*	*	*	*	0.5
2003.....	1.7	*	*	*	*	*	*	*	*	0.6
2002.....	1.5	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.2	*	*	*	*	*	*	*	*	0.5
2000.....	1.1	1.3	*	*	*	*	*	*	*	0.5
1999.....	1.5	1.6	*	*	*	*	*	0.3	0.3	0.7

White, female

2004.....	1.3	*	*	*	*	*	*	*	*	0.4
2003.....	1.3	*	*	*	*	*	*	*	*	0.3
2002.....	1.0	*	*	*	*	*	*	*	*	0.2
2001.....	1.0	1.4	*	*	*	*	*	*	*	0.3
2000.....	0.9	*	*	*	*	*	*	*	*	*
1999.....	1.2	*	*	*	*	*	*	*	0.3	0.3

Black, both sexes

2004.....	1.5	*	*	*	*	*	*	*	*	0.8
2003.....	1.5	*	*	*	*	*	*	*	*	0.9
2002.....	1.5	*	*	*	*	*	*	*	0.8	1.0
2001.....	1.3	3.7	*	*	*	*	*	*	0.8	1.0
2000.....	1.3	3.3	*	*	*	*	*	*	0.8	1.2
1999.....	2.2	3.8	*	*	*	*	*	0.9	1.7	1.9

Black, male

2004.....	1.8	*	*	*	*	*	*	*	*	*
2003.....	1.8	*	*	*	*	*	*	*	*	*
2002.....	1.8	*	*	*	*	*	*	*	*	*
2001.....	1.5	*	*	*	*	*	*	*	*	*
2000.....	1.5	*	*	*	*	*	*	*	*	1.7
1999.....	2.3	*	*	*	*	*	*	*	1.9	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ year 1/	year 2/ year 1/	years	years	years	years	years	years	years	years

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

Black, female

2004.....	1.3	*	*	*	*	*	*	*	*	*
2003.....	1.4	*	*	*	*	*	*	*	*	*
2002.....	1.1	*	*	*	*	*	*	*	*	*
2001.....	1.1	*	*	*	*	*	*	*	*	*
2000.....	1.1	*	*	*	*	*	*	*	*	*
1999.....	2.2	*	*	*	*	*	*	*	1.4	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Mycoses (B35-B49)

All races 3/, both sexes

2004.....	0.4	0.9	*	*	*	*	*	*	0.1	0.1
2003.....	0.4	1.0	*	*	*	*	*	*	0.1	0.1
2002.....	0.4	1.2	*	*	*	*	*	*	0.2	0.2
2001.....	0.4	1.4	*	*	*	*	*	*	*	0.1
2000.....	0.4	1.3	*	*	*	*	*	*	0.1	0.2
1999.....	0.7	1.4	*	*	*	*	*	0.2	0.2	0.3

All races 3/, male

2004.....	0.4	1.0	*	*	*	*	*	*	*	*
2003.....	0.4	1.1	*	*	*	*	*	*	*	*
2002.....	0.5	1.2	*	*	*	*	*	*	0.2	0.2
2001.....	0.4	1.4	*	*	*	*	*	*	*	*
2000.....	0.5	1.5	*	*	*	*	*	*	*	0.3
1999.....	0.8	1.6	*	*	*	*	*	0.2	0.3	0.4

All races 3/, female

2004.....	0.3	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.3	1.1	*	*	*	*	*	*	*	*
2001.....	0.4	1.5	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*
1999.....	0.6	1.1	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.3	0.7	*	*	*	*	*	*	*	0.1
2003.....	0.4	0.7	*	*	*	*	*	*	*	0.1
2002.....	0.4	0.9	*	*	*	*	*	*	*	0.1
2001.....	0.4	1.0	*	*	*	*	*	*	*	*
2000.....	0.4	1.0	*	*	*	*	*	*	*	0.2
1999.....	0.6	1.0	*	*	*	*	*	0.1	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Sequelae of tuberculosis (B90)

All races 3/, both sexes

2004.....	0.0	...	*	*	*	*	*	*	*
2003.....	0.0	...	*	*	*	*	*	*	*
2002.....	0.0	...	*	*	*	*	*	*	*
2001.....	0.0	...	*	*	*	*	*	*	*
2000.....	0.0	...	*	*	*	*	*	*	*
1999.....	0.1	...	*	*	*	*	*	*	*

All races 3/, male

2004.....	0.0	...	*	*	*	*	*	*	*
2003.....	0.0	...	*	*	*	*	*	*	*
2002.....	0.0	...	*	*	*	*	*	*	*
2001.....	0.0	...	*	*	*	*	*	*	*
2000.....	0.0	...	*	*	*	*	*	*	*
1999.....	0.1	...	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.0	...	*	*	*	*	*	*	*
2003.....	0.0	...	*	*	*	*	*	*	*
2002.....	0.0	...	*	*	*	*	*	*	*
2001.....	0.0	...	*	*	*	*	*	*	*
2000.....	0.0	...	*	*	*	*	*	*	*
1999.....	0.1	...	*	*	*	*	*	*	*

White, both sexes

2004.....	0.0	...	*	*	*	*	*	*	*
2003.....	0.0	...	*	*	*	*	*	*	*
2002.....	0.0	...	*	*	*	*	*	*	*
2001.....	0.0	...	*	*	*	*	*	*	*
2000.....	0.0	...	*	*	*	*	*	*	*
1999.....	0.1	...	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Sequelae of tuberculosis (B90)

Black, female

2004.....	*	...	*	*	*	*	*	*	*
2003.....	*	...	*	*	*	*	*	*	*
2002.....	*	...	*	*	*	*	*	*	*
2001.....	*	...	*	*	*	*	*	*	*
2000.....	*	...	*	*	*	*	*	*	*
1999.....	*	...	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Sequelae of poliomyelitis (B91)

All races 3/, both sexes

2004.....	0.1	...	*	*	*	*	*	*	*
2003.....	0.1	...	*	*	*	*	*	*	*
2002.....	0.1	...	*	*	*	*	*	*	*
2001.....	0.1	...	*	*	*	*	*	*	*
2000.....	0.1	...	*	*	*	*	*	*	*
1999.....	0.1	...	*	*	*	*	*	*	*

All races 3/, male

2004.....	0.1	...	*	*	*	*	*	*	*
2003.....	0.1	...	*	*	*	*	*	*	*
2002.....	0.1	...	*	*	*	*	*	*	*
2001.....	0.1	...	*	*	*	*	*	*	*
2000.....	0.1	...	*	*	*	*	*	*	*
1999.....	0.1	...	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.1	...	*	*	*	*	*	*	*
2003.....	0.1	...	*	*	*	*	*	*	*
2002.....	0.1	...	*	*	*	*	*	*	*
2001.....	0.1	...	*	*	*	*	*	*	*
2000.....	0.1	...	*	*	*	*	*	*	*
1999.....	0.1	...	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	...	*	*	*	*	*	*	*
2003.....	0.1	...	*	*	*	*	*	*	*
2002.....	0.1	...	*	*	*	*	*	*	*
2001.....	0.1	...	*	*	*	*	*	*	*
2000.....	0.1	...	*	*	*	*	*	*	*
1999.....	0.1	...	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Sequelae of poliomyelitis (B91)

White, male

2004.....	0.1	...	*	*	*	*	*	*	*
2003.....	0.1	...	*	*	*	*	*	*	*
2002.....	0.1	...	*	*	*	*	*	*	*
2001.....	0.1	...	*	*	*	*	*	*	*
2000.....	0.1	...	*	*	*	*	*	*	*
1999.....	0.1	...	*	*	*	*	*	*	*

White, female

2004.....	0.1	...	*	*	*	*	*	*	*
2003.....	0.1	...	*	*	*	*	*	*	*
2002.....	0.1	...	*	*	*	*	*	*	*
2001.....	0.1	...	*	*	*	*	*	*	*
2000.....	0.1	...	*	*	*	*	*	*	*
1999.....	0.1	...	*	*	*	*	*	*	*

Black, both sexes

2004.....	*	...	*	*	*	*	*	*	*
2003.....	*	...	*	*	*	*	*	*	*
2002.....	*	...	*	*	*	*	*	*	*
2001.....	*	...	*	*	*	*	*	*	*
2000.....	*	...	*	*	*	*	*	*	*
1999.....	*	...	*	*	*	*	*	*	*

Black, male

2004.....	*	...	*	*	*	*	*	*	*
2003.....	*	...	*	*	*	*	*	*	*
2002.....	*	...	*	*	*	*	*	*	*
2001.....	*	...	*	*	*	*	*	*	*
2000.....	*	...	*	*	*	*	*	*	*
1999.....	*	...	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Sequelae of poliomyelitis (B91)

Black, female

2004.....	*	...	*	*	*	*	*	*	*
2003.....	*	...	*	*	*	*	*	*	*
2002.....	*	...	*	*	*	*	*	*	*
2001.....	*	...	*	*	*	*	*	*	*
2000.....	*	...	*	*	*	*	*	*	*
1999.....	*	...	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

II. Neoplasms (C00-D48)

All races 3/, both sexes

2004.....	193.2	3.6	2.8	2.9	2.5	3.8	4.9	6.9	12.0	22.1
2003.....	196.2	3.4	2.8	2.8	2.8	3.6	4.9	7.2	12.3	23.6
2002.....	197.9	3.5	3.0	2.9	2.7	3.7	5.2	7.0	13.0	24.1
2001.....	199.2	2.9	3.1	2.7	2.7	3.8	5.2	7.7	13.0	25.3
2000.....	201.3	4.0	3.1	2.7	2.7	3.9	5.4	7.4	12.9	25.2
1999.....	201.8	3.3	3.1	2.8	2.7	4.0	5.5	7.6	13.1	25.8

All races 3/, male

2004.....	203.1	3.5	2.9	3.2	2.7	4.3	5.8	7.1	10.9	18.7
2003.....	206.0	3.2	3.2	3.1	2.9	4.4	5.4	7.2	11.5	20.0
2002.....	208.5	3.5	3.1	3.1	3.0	4.2	6.0	7.4	11.7	20.5
2001.....	210.1	2.8	3.3	2.7	2.9	4.5	6.1	7.6	11.6	21.7
2000.....	212.0	4.2	3.4	2.9	3.0	4.6	6.3	7.4	11.8	22.0
1999.....	213.7	3.4	3.3	2.9	2.8	4.6	6.4	8.0	12.2	22.3

All races 3/, female

2004.....	183.7	3.7	2.7	2.5	2.3	3.2	4.0	6.6	13.1	25.5
2003.....	186.6	3.6	2.4	2.5	2.7	2.7	4.4	7.2	13.1	27.3
2002.....	187.6	3.6	2.8	2.7	2.4	3.2	4.4	6.5	14.2	27.8
2001.....	188.7	3.0	2.9	2.7	2.6	3.2	4.3	7.7	14.5	28.8
2000.....	191.1	3.8	2.8	2.5	2.4	3.2	4.4	7.4	14.1	28.4
1999.....	190.3	3.2	2.9	2.6	2.5	3.3	4.6	7.2	13.9	29.3

White, both sexes

2004.....	205.7	3.8	2.8	2.9	2.5	3.8	4.8	6.7	11.6	21.4
2003.....	208.9	3.6	2.9	2.9	2.8	3.7	5.0	7.1	11.9	22.8
2002.....	210.6	3.7	2.9	3.0	2.7	3.8	5.4	6.8	12.6	23.3
2001.....	211.7	2.9	3.4	2.7	2.7	3.9	5.2	7.5	12.6	24.4
2000.....	213.9	4.1	3.1	2.8	2.8	3.8	5.3	7.4	12.6	24.0
1999.....	213.9	3.2	3.3	2.8	2.6	3.8	5.4	7.6	12.6	24.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

II. Neoplasms (C00-D48)

White, male

2004.....	215.0	3.8	2.9	3.3	2.7	4.4	5.6	7.2	11.0	18.6
2003.....	218.3	3.4	3.2	3.3	2.9	4.6	5.4	7.4	11.2	20.0
2002.....	220.6	3.9	3.0	3.1	3.0	4.4	6.2	7.2	11.6	19.9
2001.....	221.7	2.7	3.5	2.7	2.6	4.5	6.1	7.5	11.6	21.2
2000.....	223.3	4.4	3.5	2.9	3.1	4.4	6.3	7.4	11.8	21.2
1999.....	224.8	3.4	3.5	3.0	2.8	4.3	6.2	7.9	11.9	21.7

White, female

2004.....	196.7	3.9	2.8	2.5	2.2	3.1	3.9	6.2	12.4	24.3
2003.....	199.7	3.8	2.5	2.6	2.7	2.8	4.4	6.8	12.6	25.7
2002.....	200.9	3.5	2.7	2.9	2.5	3.2	4.6	6.4	13.6	26.9
2001.....	201.9	3.1	3.2	2.8	2.7	3.2	4.3	7.6	13.7	27.8
2000.....	204.8	3.9	2.7	2.6	2.5	3.1	4.4	7.3	13.4	26.9
1999.....	203.3	2.9	3.2	2.6	2.4	3.3	4.5	7.2	13.4	27.5

Black, both sexes

2004.....	165.0	*	2.8	2.8	2.9	4.2	6.1	9.4	16.1	29.4
2003.....	167.2	*	2.6	2.6	3.0	3.2	5.2	9.1	16.2	32.3
2002.....	168.9	3.1	3.5	2.8	2.8	3.8	4.8	8.5	17.0	32.4
2001.....	170.2	*	2.1	2.8	3.2	4.0	5.4	9.4	16.8	33.6
2000.....	172.4	3.3	3.2	2.4	2.5	4.6	6.2	8.6	16.3	35.3
1999.....	174.5	4.3	2.5	2.8	3.2	4.9	7.0	8.9	16.6	36.9

Black, male

2004.....	179.2	*	3.2	2.7	2.9	4.4	7.2	8.3	12.9	21.0
2003.....	181.2	*	3.0	2.8	3.3	3.7	5.8	7.7	14.2	22.8
2002.....	184.3	*	3.3	3.6	3.0	4.3	5.4	8.6	14.3	26.2
2001.....	187.5	*	2.6	2.9	3.8	4.7	6.3	8.7	13.0	27.8
2000.....	191.5	*	3.3	2.7	2.8	5.7	7.3	7.7	13.3	28.6
1999.....	194.1	*	2.8	2.6	3.5	6.1	7.8	9.4	14.9	28.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

II. Neoplasms (C00-D48)

Black, female

2004.....	152.1	*	2.4	2.8	2.9	3.9	5.1	10.4	18.9	37.0
2003.....	154.4	*	2.1	2.3	2.8	2.6	4.6	10.3	18.1	40.6
2002.....	155.0	*	3.7	1.9	2.5	3.3	4.2	8.5	19.3	37.9
2001.....	154.6	*	1.6	2.6	2.5	3.2	4.5	10.0	20.3	38.8
2000.....	155.0	*	3.0	2.2	2.3	3.6	5.2	9.5	19.0	41.2
1999.....	156.8	*	2.1	2.9	2.8	3.7	6.3	8.4	18.1	44.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

All races 3/, both sexes

2004.....	188.6	1.8	2.5	2.7	2.3	3.5	4.7	6.6	11.5	21.5
2003.....	191.5	1.9	2.5	2.6	2.6	3.4	4.6	6.8	11.8	23.0
2002.....	193.2	1.8	2.6	2.7	2.5	3.5	5.0	6.6	12.5	23.5
2001.....	194.4	1.6	2.7	2.4	2.5	3.6	4.9	7.3	12.6	24.6
2000.....	196.5	2.4	2.7	2.4	2.6	3.7	5.1	7.0	12.5	24.5
1999.....	197.0	1.8	2.7	2.5	2.5	3.7	5.3	7.3	12.5	25.2

All races 3/, male

2004.....	198.4	1.8	2.6	3.0	2.5	4.1	5.5	6.8	10.4	18.1
2003.....	201.3	1.7	2.8	2.9	2.7	4.2	5.1	6.8	10.9	19.3
2002.....	203.8	2.0	2.7	2.9	2.8	4.0	5.7	7.0	11.3	19.8
2001.....	205.3	1.5	2.9	2.4	2.6	4.3	5.7	7.2	11.2	21.0
2000.....	207.2	2.6	3.0	2.6	2.9	4.3	5.9	7.1	11.3	21.2
1999.....	208.9	1.9	2.9	2.6	2.7	4.3	6.1	7.6	11.6	21.6

All races 3/, female

2004.....	179.1	1.9	2.4	2.3	2.2	2.9	3.8	6.3	12.6	24.9
2003.....	182.0	2.1	2.1	2.3	2.5	2.5	4.2	6.8	12.6	26.7
2002.....	183.0	1.6	2.4	2.5	2.2	3.0	4.2	6.2	13.7	27.2
2001.....	183.9	1.8	2.5	2.4	2.3	2.9	4.1	7.4	14.0	28.1
2000.....	186.2	2.3	2.5	2.2	2.2	3.0	4.2	7.0	13.7	27.8
1999.....	185.6	1.7	2.5	2.3	2.3	3.0	4.4	7.0	13.4	28.7

White, both sexes

2004.....	200.7	2.0	2.5	2.7	2.3	3.6	4.5	6.4	11.1	20.8
2003.....	203.7	2.1	2.6	2.8	2.6	3.5	4.7	6.7	11.3	22.2
2002.....	205.5	1.8	2.5	2.8	2.5	3.6	5.2	6.5	12.1	22.7
2001.....	206.4	1.7	3.0	2.5	2.4	3.7	5.0	7.2	12.2	23.7
2000.....	208.6	2.5	2.8	2.5	2.6	3.6	5.1	7.0	12.1	23.3
1999.....	208.7	1.8	2.9	2.5	2.4	3.6	5.2	7.3	12.1	24.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

White, male

2004.....	209.9	1.9	2.5	3.1	2.5	4.2	5.3	6.8	10.4	18.0
2003.....	213.1	1.8	2.9	3.1	2.7	4.3	5.1	7.0	10.6	19.3
2002.....	215.5	2.2	2.7	2.9	2.7	4.2	5.9	6.9	11.1	19.2
2001.....	216.4	1.5	3.1	2.4	2.4	4.4	5.8	7.2	11.2	20.5
2000.....	218.1	2.8	3.1	2.6	2.9	4.2	5.9	7.1	11.2	20.5
1999.....	219.5	2.0	3.2	2.7	2.6	4.1	5.9	7.6	11.3	21.1

White, female

2004.....	191.7	2.1	2.5	2.3	2.1	2.9	3.7	6.0	11.9	23.7
2003.....	194.6	2.3	2.3	2.4	2.5	2.6	4.2	6.5	12.0	25.2
2002.....	195.8	1.4	2.3	2.6	2.4	3.0	4.4	6.1	13.2	26.3
2001.....	196.7	2.0	2.8	2.5	2.4	2.9	4.1	7.2	13.3	27.1
2000.....	199.4	2.3	2.4	2.3	2.3	3.0	4.2	7.0	13.0	26.3
1999.....	198.2	1.5	2.7	2.2	2.2	3.0	4.3	7.0	12.9	27.1

Black, both sexes

2004.....	161.9	*	2.5	2.5	2.6	3.8	5.9	8.9	15.1	28.4
2003.....	164.3	*	2.2	2.3	2.8	2.9	4.9	8.5	15.7	31.0
2002.....	165.9	*	2.9	2.6	2.5	3.6	4.5	8.0	16.3	31.7
2001.....	167.1	*	1.9	2.4	2.8	3.6	5.1	9.1	16.4	32.7
2000.....	169.3	*	2.8	2.1	2.3	4.4	5.9	8.0	15.8	34.5
1999.....	171.3	*	2.0	2.4	3.0	4.5	6.6	8.6	15.8	35.8

Black, male

2004.....	176.2	*	2.8	2.5	2.4	4.1	6.8	7.9	12.2	19.7
2003.....	178.3	*	2.4	2.4	3.1	3.5	5.6	7.0	13.5	21.7
2002.....	181.5	*	2.6	3.3	2.9	4.1	5.0	8.2	14.1	25.6
2001.....	184.5	*	2.4	2.5	3.4	4.3	5.9	8.5	12.4	26.8
2000.....	188.5	*	2.7	2.4	2.7	5.2	7.1	7.3	12.9	27.6
1999.....	191.0	*	2.1	2.3	3.3	5.6	7.3	8.8	14.0	27.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

Black, female

2004.....	148.9	*	2.2	2.5	2.7	3.5	5.0	9.9	17.8	36.2
2003.....	151.4	*	2.1	2.1	2.5	2.3	4.3	10.0	17.7	39.2
2002.....	151.7	*	3.2	1.9	2.1	3.1	4.0	7.9	18.3	37.1
2001.....	151.3	*	*	2.2	2.2	2.9	4.3	9.7	20.0	37.9
2000.....	151.8	*	2.9	1.8	2.0	3.5	4.7	8.5	18.3	40.5
1999.....	153.3	*	1.9	2.6	2.7	3.4	5.9	8.3	17.5	43.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races 3/, both sexes

2004.....	2.7	*	*	*	*	*	*	*	0.1	0.3
2003.....	2.7	*	*	*	*	*	*	*	0.2	0.3
2002.....	2.7	*	*	*	*	*	*	0.1	0.2	0.4
2001.....	2.7	*	*	*	*	*	*	*	0.2	0.4
2000.....	2.7	*	*	*	*	*	*	*	0.2	0.4
1999.....	2.7	*	*	*	*	*	*	*	0.1	0.4

All races 3/, male

2004.....	3.7	*	*	*	*	*	*	*	0.2	0.3
2003.....	3.7	*	*	*	*	*	*	*	0.3	0.4
2002.....	3.7	*	*	*	*	*	*	*	0.2	0.4
2001.....	3.7	*	*	*	*	*	*	*	0.2	0.5
2000.....	3.6	*	*	*	*	*	*	*	*	0.5
1999.....	3.6	*	*	*	*	*	*	*	*	0.5

All races 3/, female

2004.....	1.7	*	*	*	*	*	*	*	*	0.2
2003.....	1.7	*	*	*	*	*	*	*	*	0.2
2002.....	1.7	*	*	*	*	*	*	*	*	0.3
2001.....	1.8	*	*	*	*	*	*	*	*	0.3
2000.....	1.8	*	*	*	*	*	*	*	*	0.2
1999.....	1.8	*	*	*	*	*	*	*	*	0.3

White, both sexes

2004.....	2.7	*	*	*	*	*	*	*	0.1	0.2
2003.....	2.7	*	*	*	*	*	*	*	0.2	0.3
2002.....	2.7	*	*	*	*	*	*	*	0.1	0.3
2001.....	2.7	*	*	*	*	*	*	*	0.1	0.3
2000.....	2.7	*	*	*	*	*	*	*	0.1	0.3
1999.....	2.7	*	*	*	*	*	*	*	0.1	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of digestive organs (C15-C26)

All races 3/, both sexes

2004.....	44.7	*	0.1	*	*	0.2	0.5	0.9	2.1	4.3
2003.....	45.5	*	0.1	*	*	0.1	0.4	1.0	2.1	4.6
2002.....	45.7	*	0.2	*	*	0.1	0.5	1.0	2.0	4.6
2001.....	46.0	*	*	*	*	0.2	0.4	0.9	2.1	5.0
2000.....	46.4	*	0.2	*	*	0.2	0.3	0.9	2.1	4.5
1999.....	46.4	*	0.2	*	*	0.2	0.4	1.0	2.0	4.6

All races 3/, male

2004.....	49.8	*	*	*	*	0.2	0.5	1.1	2.4	5.0
2003.....	50.4	*	*	*	*	*	0.4	1.1	2.3	5.5
2002.....	50.7	*	0.3	*	*	*	0.5	1.2	2.3	5.3
2001.....	50.5	*	*	*	*	*	0.4	0.9	2.4	6.0
2000.....	50.8	*	*	*	*	0.2	0.4	1.1	2.5	5.5
1999.....	50.7	*	0.3	*	*	0.2	0.4	1.2	2.3	5.3

All races 3/, female

2004.....	39.8	*	*	*	*	*	0.4	0.7	1.8	3.6
2003.....	40.7	*	*	*	*	*	0.4	0.9	1.9	3.8
2002.....	40.8	*	*	*	*	*	0.4	0.8	1.7	3.8
2001.....	41.7	*	*	*	*	*	0.3	0.8	1.8	4.0
2000.....	42.3	*	*	*	*	*	0.3	0.8	1.6	3.4
1999.....	42.3	*	*	*	*	*	0.4	0.8	1.7	3.8

White, both sexes

2004.....	46.3	*	*	*	*	0.1	0.4	0.8	1.9	4.0
2003.....	47.0	*	*	*	*	0.1	0.4	0.9	1.8	4.3
2002.....	47.3	*	0.2	*	*	0.1	0.4	0.9	1.8	4.3
2001.....	47.6	*	*	*	*	0.1	0.3	0.8	1.9	4.5
2000.....	48.0	*	*	*	*	0.2	0.3	0.8	1.8	4.0
1999.....	48.0	*	0.2	*	*	0.1	0.4	0.8	1.7	4.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of digestive organs (C15-C26)

White, male

2004.....	51.3	*	*	*	*	*	0.4	0.9	2.2	4.8
2003.....	52.0	*	*	*	*	*	0.4	1.0	1.9	5.1
2002.....	52.3	*	*	*	*	*	0.4	1.0	2.0	4.8
2001.....	52.1	*	*	*	*	*	0.3	0.8	2.1	5.4
2000.....	52.3	*	*	*	*	*	0.3	0.9	2.2	5.0
1999.....	52.2	*	*	*	*	*	0.4	1.0	2.0	4.8

White, female

2004.....	41.3	*	*	*	*	*	0.4	0.7	1.6	3.2
2003.....	42.2	*	*	*	*	*	0.3	0.8	1.7	3.4
2002.....	42.4	*	*	*	*	*	0.5	0.8	1.6	3.7
2001.....	43.2	*	*	*	*	*	0.4	0.8	1.6	3.5
2000.....	43.9	*	*	*	*	*	*	0.6	1.4	3.1
1999.....	43.9	*	*	*	*	*	0.4	0.6	1.5	3.4

Black, both sexes

2004.....	42.7	*	*	*	*	*	0.9	1.4	3.1	5.4
2003.....	43.7	*	*	*	*	*	*	1.2	3.1	6.8
2002.....	43.6	*	*	*	*	*	*	1.4	2.7	5.9
2001.....	44.1	*	*	*	*	*	*	1.2	3.1	7.7
2000.....	44.5	*	*	*	*	*	*	1.3	3.3	6.9
1999.....	44.6	*	*	*	*	*	*	1.8	3.1	7.2

Black, male

2004.....	47.7	*	*	*	*	*	*	1.6	3.4	5.7
2003.....	48.4	*	*	*	*	*	*	*	3.6	7.4
2002.....	48.5	*	*	*	*	*	*	1.9	3.5	7.6
2001.....	48.1	*	*	*	*	*	*	*	3.7	9.5
2000.....	49.2	*	*	*	*	*	*	*	4.1	8.5
1999.....	49.4	*	*	*	*	*	*	2.3	3.7	7.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of digestive organs (C15-C26)

Black, female

2004.....	38.2	*	*	*	*	*	*	*	2.8	5.2
2003.....	39.3	*	*	*	*	*	*	*	2.6	6.3
2002.....	39.2	*	*	*	*	*	*	*	2.0	4.4
2001.....	40.5	*	*	*	*	*	*	*	2.6	6.1
2000.....	40.3	*	*	*	*	*	*	1.4	2.5	5.5
1999.....	40.4	*	*	*	*	*	*	*	2.5	6.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of esophagus (C15)

All races 3/, both sexes

2004.....	4.4	*	*	*	*	*	*	*	0.1	0.3
2003.....	4.4	*	*	*	*	*	*	*	*	0.4
2002.....	4.4	*	*	*	*	*	*	*	*	0.4
2001.....	4.4	*	*	*	*	*	*	*	0.1	0.3
2000.....	4.3	*	*	*	*	*	*	*	0.1	0.3
1999.....	4.3	*	*	*	*	*	*	*	0.1	0.3

All races 3/, male

2004.....	7.0	*	*	*	*	*	*	*	0.2	0.5
2003.....	6.9	*	*	*	*	*	*	*	*	0.6
2002.....	6.9	*	*	*	*	*	*	*	*	0.6
2001.....	6.9	*	*	*	*	*	*	*	*	0.5
2000.....	6.7	*	*	*	*	*	*	*	0.2	0.6
1999.....	6.6	*	*	*	*	*	*	*	0.2	0.6

All races 3/, female

2004.....	2.0	*	*	*	*	*	*	*	*	*
2003.....	2.0	*	*	*	*	*	*	*	*	*
2002.....	2.0	*	*	*	*	*	*	*	*	*
2001.....	2.0	*	*	*	*	*	*	*	*	*
2000.....	2.1	*	*	*	*	*	*	*	*	*
1999.....	2.0	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	4.7	*	*	*	*	*	*	*	*	0.4
2003.....	4.7	*	*	*	*	*	*	*	*	0.4
2002.....	4.6	*	*	*	*	*	*	*	*	0.4
2001.....	4.6	*	*	*	*	*	*	*	*	0.3
2000.....	4.5	*	*	*	*	*	*	*	0.2	0.4
1999.....	4.4	*	*	*	*	*	*	*	0.1	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of stomach (C16)

All races 3/, both sexes

2004.....	4.0	*	*	*	*	*	*	0.1	0.5	0.7
2003.....	4.2	*	*	*	*	*	*	0.2	0.4	0.8
2002.....	4.2	*	*	*	*	*	0.1	0.3	0.4	0.7
2001.....	4.3	*	*	*	*	*	*	0.2	0.4	0.8
2000.....	4.5	*	*	*	*	*	*	0.2	0.4	0.7
1999.....	4.6	*	*	*	*	*	*	0.2	0.4	0.8

All races 3/, male

2004.....	4.8	*	*	*	*	*	*	*	0.4	0.7
2003.....	4.9	*	*	*	*	*	*	*	0.4	0.8
2002.....	5.0	*	*	*	*	*	*	0.3	0.5	0.7
2001.....	5.1	*	*	*	*	*	*	0.2	0.5	0.9
2000.....	5.3	*	*	*	*	*	*	0.2	0.5	0.9
1999.....	5.5	*	*	*	*	*	*	0.2	0.5	0.9

All races 3/, female

2004.....	3.3	*	*	*	*	*	*	*	0.5	0.6
2003.....	3.4	*	*	*	*	*	*	*	0.4	0.8
2002.....	3.5	*	*	*	*	*	*	*	0.4	0.7
2001.....	3.5	*	*	*	*	*	*	*	0.4	0.7
2000.....	3.7	*	*	*	*	*	*	*	0.4	0.6
1999.....	3.7	*	*	*	*	*	*	0.3	0.3	0.8

White, both sexes

2004.....	3.8	*	*	*	*	*	*	*	0.4	0.7
2003.....	3.9	*	*	*	*	*	*	0.2	0.4	0.7
2002.....	4.0	*	*	*	*	*	*	0.2	0.4	0.7
2001.....	4.1	*	*	*	*	*	*	0.2	0.4	0.6
2000.....	4.2	*	*	*	*	*	*	0.1	0.3	0.6
1999.....	4.3	*	*	*	*	*	*	0.2	0.4	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of colon, rectum and anus (C18-C21)

All races 3/, both sexes

2004.....	18.3	*	*	*	*	*	0.2	0.4	0.9	2.0
2003.....	19.2	*	*	*	*	*	0.2	0.5	0.9	2.1
2002.....	19.7	*	*	*	*	*	0.2	0.5	1.0	2.0
2001.....	20.0	*	*	*	*	*	0.2	0.4	1.1	2.3
2000.....	20.4	*	*	*	*	*	0.1	0.4	0.9	2.1
1999.....	20.5	*	*	*	*	*	0.2	0.4	1.0	1.9

All races 3/, male

2004.....	18.6	*	*	*	*	*	0.2	0.5	1.0	2.1
2003.....	19.6	*	*	*	*	*	*	0.6	0.9	2.2
2002.....	20.1	*	*	*	*	*	0.3	0.5	1.1	2.1
2001.....	20.2	*	*	*	*	*	*	0.4	1.1	2.6
2000.....	20.6	*	*	*	*	*	*	0.4	1.1	2.4
1999.....	20.6	*	*	*	*	*	0.2	0.6	1.0	2.0

All races 3/, female

2004.....	18.0	*	*	*	*	*	*	0.4	0.9	2.0
2003.....	18.9	*	*	*	*	*	*	0.4	0.9	2.0
2002.....	19.2	*	*	*	*	*	*	0.4	0.9	1.9
2001.....	19.8	*	*	*	*	*	*	0.5	1.0	1.9
2000.....	20.2	*	*	*	*	*	*	0.4	0.8	1.8
1999.....	20.3	*	*	*	*	*	*	0.3	0.9	1.9

White, both sexes

2004.....	19.2	*	*	*	*	*	0.2	0.4	0.9	1.9
2003.....	20.2	*	*	*	*	*	0.1	0.5	0.9	1.9
2002.....	20.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2001.....	21.0	*	*	*	*	*	0.2	0.4	1.0	2.1
2000.....	21.5	*	*	*	*	*	*	0.4	0.9	1.9
1999.....	21.6	*	*	*	*	*	0.2	0.4	0.9	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Colon (C18)

All races 3/, both sexes

2004.....	15.2	*	*	*	*	*	0.1	0.3	0.7	1.4
2003.....	16.1	*	*	*	*	*	0.1	0.4	0.7	1.7
2002.....	16.5	*	*	*	*	*	0.2	0.3	0.7	1.5
2001.....	16.8	*	*	*	*	*	0.1	0.3	0.8	1.7
2000.....	17.3	*	*	*	*	*	*	0.3	0.8	1.6
1999.....	17.4	*	*	*	*	*	0.2	0.3	0.7	1.5

All races 3/, male

2004.....	15.1	*	*	*	*	*	*	0.4	0.6	1.4
2003.....	16.1	*	*	*	*	*	*	0.5	0.7	1.7
2002.....	16.6	*	*	*	*	*	0.2	0.3	0.8	1.5
2001.....	16.7	*	*	*	*	*	*	0.3	0.9	1.9
2000.....	17.1	*	*	*	*	*	*	0.3	0.9	1.7
1999.....	17.2	*	*	*	*	*	*	0.4	0.8	1.5

All races 3/, female

2004.....	15.2	*	*	*	*	*	*	0.3	0.7	1.4
2003.....	16.1	*	*	*	*	*	*	0.3	0.7	1.7
2002.....	16.4	*	*	*	*	*	*	0.3	0.7	1.4
2001.....	16.9	*	*	*	*	*	*	0.3	0.8	1.5
2000.....	17.4	*	*	*	*	*	*	0.3	0.6	1.4
1999.....	17.5	*	*	*	*	*	*	0.3	0.7	1.5

White, both sexes

2004.....	15.9	*	*	*	*	*	*	0.3	0.6	1.3
2003.....	16.9	*	*	*	*	*	*	0.4	0.6	1.6
2002.....	17.3	*	*	*	*	*	0.2	0.3	0.7	1.4
2001.....	17.6	*	*	*	*	*	0.1	0.3	0.8	1.6
2000.....	18.1	*	*	*	*	*	*	0.3	0.7	1.4
1999.....	18.2	*	*	*	*	*	*	0.3	0.7	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Rectosigmoid junction and rectum (C19-C20)

All races 3/, both sexes

2004.....	2.9	*	*	*	*	*	*	0.1	0.3	0.5
2003.....	2.9	*	*	*	*	*	*	*	0.2	0.4
2002.....	3.0	*	*	*	*	*	*	0.1	0.2	0.5
2001.....	3.0	*	*	*	*	*	*	0.1	0.2	0.5
2000.....	3.0	*	*	*	*	*	*	0.1	0.2	0.5
1999.....	3.0	*	*	*	*	*	*	0.1	0.2	0.3

All races 3/, male

2004.....	3.3	*	*	*	*	*	*	*	0.3	0.6
2003.....	3.3	*	*	*	*	*	*	*	0.3	0.4
2002.....	3.4	*	*	*	*	*	*	*	0.3	0.5
2001.....	3.4	*	*	*	*	*	*	*	0.2	0.6
2000.....	3.4	*	*	*	*	*	*	*	*	0.6
1999.....	3.3	*	*	*	*	*	*	0.2	0.2	0.4

All races 3/, female

2004.....	2.5	*	*	*	*	*	*	*	*	0.5
2003.....	2.6	*	*	*	*	*	*	*	*	0.3
2002.....	2.6	*	*	*	*	*	*	*	*	0.4
2001.....	2.6	*	*	*	*	*	*	*	*	0.3
2000.....	2.6	*	*	*	*	*	*	*	*	0.3
1999.....	2.7	*	*	*	*	*	*	*	*	0.3

White, both sexes

2004.....	3.1	*	*	*	*	*	*	0.1	0.2	0.5
2003.....	3.1	*	*	*	*	*	*	*	0.2	0.3
2002.....	3.2	*	*	*	*	*	*	0.1	0.2	0.4
2001.....	3.2	*	*	*	*	*	*	0.1	0.2	0.4
2000.....	3.2	*	*	*	*	*	*	*	0.1	0.4
1999.....	3.2	*	*	*	*	*	*	0.1	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Of liver and intrahepatic bile ducts (C22)

All races 3/, both sexes

2004.....	5.2	*	0.1	*	*	*	0.1	0.1	0.3	0.5
2003.....	5.1	*	0.1	*	*	*	0.1	0.1	0.3	0.5
2002.....	4.9	*	0.2	*	*	*	*	0.2	0.3	0.6
2001.....	4.7	*	*	*	*	*	*	0.2	0.2	0.6
2000.....	4.6	*	0.1	*	*	*	*	0.1	0.3	0.5
1999.....	4.4	*	0.2	*	*	*	0.1	0.2	0.2	0.6

All races 3/, male

2004.....	6.8	*	*	*	*	*	*	0.2	0.3	0.8
2003.....	6.6	*	*	*	*	*	*	*	0.4	0.8
2002.....	6.4	*	*	*	*	*	*	0.2	0.4	0.8
2001.....	6.1	*	*	*	*	*	*	*	0.3	0.9
2000.....	5.9	*	*	*	*	*	*	*	0.4	0.7
1999.....	5.6	*	0.3	*	*	*	*	0.2	0.3	0.8

All races 3/, female

2004.....	3.7	*	*	*	*	*	*	*	0.2	0.2
2003.....	3.5	*	*	*	*	*	*	*	0.2	0.3
2002.....	3.4	*	*	*	*	*	*	*	0.3	0.3
2001.....	3.4	*	*	*	*	*	*	*	*	0.3
2000.....	3.3	*	*	*	*	*	*	*	*	0.3
1999.....	3.3	*	*	*	*	*	*	*	*	0.3

White, both sexes

2004.....	5.1	*	*	*	*	*	*	*	0.2	0.4
2003.....	4.9	*	*	*	*	*	*	*	0.2	0.4
2002.....	4.8	*	0.2	*	*	*	*	*	0.2	0.4
2001.....	4.6	*	*	*	*	*	*	*	0.2	0.5
2000.....	4.5	*	*	*	*	*	*	*	0.2	0.4
1999.....	4.4	*	0.2	*	*	*	*	*	*	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Liver (C22.0,C22.2-C22.9)

All races 3/, both sexes

2004.....	4.0	*	0.1	*	*	*	0.1	0.1	0.2	0.3
2003.....	3.9	*	0.1	*	*	*	0.1	0.1	0.2	0.4
2002.....	3.8	*	0.2	*	*	*	*	0.2	0.2	0.4
2001.....	3.7	*	*	*	*	*	*	0.1	0.2	0.5
2000.....	3.6	*	0.1	*	*	*	*	0.1	0.2	0.4
1999.....	3.5	*	0.2	*	*	*	0.1	0.1	0.2	0.4

All races 3/, male

2004.....	5.6	*	*	*	*	*	*	*	0.3	0.5
2003.....	5.5	*	*	*	*	*	*	*	0.3	0.6
2002.....	5.3	*	*	*	*	*	*	0.2	0.3	0.6
2001.....	5.1	*	*	*	*	*	*	*	0.2	0.8
2000.....	5.0	*	*	*	*	*	*	*	0.3	0.6
1999.....	4.7	*	0.3	*	*	*	*	*	0.2	0.6

All races 3/, female

2004.....	2.5	*	*	*	*	*	*	*	*	*
2003.....	2.4	*	*	*	*	*	*	*	*	*
2002.....	2.4	*	*	*	*	*	*	*	*	0.2
2001.....	2.4	*	*	*	*	*	*	*	*	0.2
2000.....	2.3	*	*	*	*	*	*	*	*	0.3
1999.....	2.4	*	*	*	*	*	*	*	*	0.2

White, both sexes

2004.....	3.8	*	*	*	*	*	*	*	*	0.2
2003.....	3.7	*	*	*	*	*	*	*	*	0.3
2002.....	3.7	*	0.2	*	*	*	*	*	0.1	0.3
2001.....	3.5	*	*	*	*	*	*	*	0.2	0.3
2000.....	3.4	*	*	*	*	*	*	*	0.1	0.3
1999.....	3.4	*	0.2	*	*	*	*	*	*	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of pancreas (C25)

All races 3/, both sexes

2004.....	10.8	*	*	*	*	*	*	*	0.2	0.6
2003.....	10.6	*	*	*	*	*	*	0.1	0.2	0.7
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.7
2001.....	10.5	*	*	*	*	*	*	*	0.2	0.8
2000.....	10.4	*	*	*	*	*	*	*	0.2	0.6
1999.....	10.4	*	*	*	*	*	*	*	0.2	0.8

All races 3/, male

2004.....	10.9	*	*	*	*	*	*	*	0.3	0.7
2003.....	10.5	*	*	*	*	*	*	*	0.2	0.9
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.8
2001.....	10.3	*	*	*	*	*	*	*	0.2	0.9
2000.....	10.3	*	*	*	*	*	*	*	0.2	0.8
1999.....	10.4	*	*	*	*	*	*	*	0.3	0.9

All races 3/, female

2004.....	10.7	*	*	*	*	*	*	*	*	0.5
2003.....	10.6	*	*	*	*	*	*	*	*	0.5
2002.....	10.5	*	*	*	*	*	*	*	*	0.6
2001.....	10.6	*	*	*	*	*	*	*	*	0.6
2000.....	10.5	*	*	*	*	*	*	*	*	0.5
1999.....	10.5	*	*	*	*	*	*	*	0.2	0.6

White, both sexes

2004.....	11.4	*	*	*	*	*	*	*	0.2	0.6
2003.....	11.2	*	*	*	*	*	*	*	0.2	0.7
2002.....	11.1	*	*	*	*	*	*	*	0.1	0.7
2001.....	11.1	*	*	*	*	*	*	*	0.2	0.7
2000.....	11.0	*	*	*	*	*	*	*	0.2	0.6
1999.....	11.0	*	*	*	*	*	*	*	0.2	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)

All races 3/, both sexes

2004.....	55.4	*	*	*	*	*	0.1	0.3	0.6	2.4
2003.....	56.0	*	*	*	*	*	0.1	0.2	0.7	2.6
2002.....	56.3	*	*	*	*	*	0.2	0.2	0.7	2.9
2001.....	56.5	*	*	*	*	*	0.1	0.2	0.7	3.1
2000.....	57.0	*	*	*	*	*	0.1	0.2	0.9	3.1
1999.....	56.2	*	*	*	*	*	0.1	0.3	0.8	3.1

All races 3/, male

2004.....	64.4	*	*	*	*	*	*	0.4	0.7	2.4
2003.....	65.4	*	*	*	*	*	*	0.2	0.7	2.6
2002.....	66.1	*	*	*	*	*	0.3	0.3	0.8	3.0
2001.....	67.2	*	*	*	*	*	*	0.2	0.7	3.1
2000.....	68.1	*	*	*	*	*	*	0.4	0.8	3.5
1999.....	68.0	*	*	*	*	*	*	0.3	0.8	3.2

All races 3/, female

2004.....	46.7	*	*	*	*	*	*	*	0.5	2.4
2003.....	46.9	*	*	*	*	*	*	0.3	0.6	2.6
2002.....	46.8	*	*	*	*	*	*	*	0.7	2.8
2001.....	46.1	*	*	*	*	*	*	0.3	0.7	3.1
2000.....	46.2	*	*	*	*	*	*	*	0.9	2.7
1999.....	45.0	*	*	*	*	*	*	0.2	0.7	3.1

White, both sexes

2004.....	59.7	*	*	*	*	*	*	0.3	0.6	2.4
2003.....	60.3	*	*	*	*	*	0.1	0.2	0.7	2.5
2002.....	60.7	*	*	*	*	*	0.1	0.2	0.7	2.8
2001.....	60.8	*	*	*	*	*	0.1	0.2	0.7	3.1
2000.....	61.2	*	*	*	*	*	0.1	0.2	0.9	2.9
1999.....	60.2	*	*	*	*	*	*	0.3	0.8	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of trachea, bronchus and lung (C33-C34)

All races 3/, both sexes

2004.....	53.8	*	*	*	*	*	*	0.2	0.5	2.2
2003.....	54.4	*	*	*	*	*	*	0.2	0.6	2.4
2002.....	54.7	*	*	*	*	*	*	0.2	0.6	2.7
2001.....	54.8	*	*	*	*	*	*	0.2	0.6	2.9
2000.....	55.3	*	*	*	*	*	*	0.2	0.8	2.9
1999.....	54.5	*	*	*	*	*	*	0.2	0.7	2.9

All races 3/, male

2004.....	62.0	*	*	*	*	*	*	0.3	0.6	2.2
2003.....	62.9	*	*	*	*	*	*	*	0.7	2.4
2002.....	63.7	*	*	*	*	*	*	*	0.7	2.6
2001.....	64.7	*	*	*	*	*	*	*	0.6	2.8
2000.....	65.5	*	*	*	*	*	*	0.3	0.7	3.3
1999.....	65.4	*	*	*	*	*	*	0.2	0.7	2.8

All races 3/, female

2004.....	45.9	*	*	*	*	*	*	*	0.5	2.2
2003.....	46.1	*	*	*	*	*	*	0.2	0.5	2.5
2002.....	46.0	*	*	*	*	*	*	*	0.6	2.7
2001.....	45.3	*	*	*	*	*	*	*	0.6	2.9
2000.....	45.4	*	*	*	*	*	*	*	0.8	2.6
1999.....	44.1	*	*	*	*	*	*	*	0.6	3.0

White, both sexes

2004.....	58.1	*	*	*	*	*	*	0.2	0.5	2.2
2003.....	58.7	*	*	*	*	*	*	0.2	0.6	2.4
2002.....	59.1	*	*	*	*	*	*	0.2	0.6	2.6
2001.....	59.1	*	*	*	*	*	*	0.2	0.6	2.9
2000.....	59.5	*	*	*	*	*	*	0.2	0.8	2.7
1999.....	58.5	*	*	*	*	*	*	0.2	0.7	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of bone and articular cartilage (C40-C41)

All races 3/, both sexes

2004.....	0.4	*	*	*	0.3	0.5	0.4	0.3	0.2	0.2
2003.....	0.4	*	*	*	0.3	0.5	0.4	0.3	0.2	0.2
2002.....	0.4	*	*	0.1	0.3	0.5	0.4	0.3	0.2	0.2
2001.....	0.5	*	*	*	0.3	0.5	0.5	0.3	0.2	0.2
2000.....	0.4	*	*	*	0.3	0.5	0.4	0.2	0.2	0.2
1999.....	0.4	*	*	0.1	0.3	0.6	0.4	0.2	0.2	0.2

All races 3/, male

2004.....	0.5	*	*	*	0.3	0.7	0.5	0.3	0.2	0.2
2003.....	0.5	*	*	*	0.3	0.7	0.5	0.3	0.2	0.3
2002.....	0.5	*	*	*	0.3	0.5	0.5	0.4	0.2	0.2
2001.....	0.5	*	*	*	0.4	0.6	0.7	0.3	0.2	0.2
2000.....	0.5	*	*	*	0.4	0.6	0.5	0.3	0.3	0.2
1999.....	0.5	*	*	*	0.3	0.7	0.5	0.3	0.2	0.2

All races 3/, female

2004.....	0.4	*	*	*	0.3	0.4	0.3	0.2	*	*
2003.....	0.4	*	*	*	0.3	0.3	0.3	0.3	*	*
2002.....	0.4	*	*	*	0.3	0.4	0.3	*	*	*
2001.....	0.4	*	*	*	0.3	0.4	0.3	0.2	0.2	0.2
2000.....	0.4	*	*	*	0.2	0.4	0.3	*	*	*
1999.....	0.4	*	*	*	0.3	0.5	0.3	*	0.2	*

White, both sexes

2004.....	0.5	*	*	*	0.3	0.5	0.4	0.3	0.2	0.2
2003.....	0.5	*	*	*	0.3	0.6	0.4	0.3	0.2	0.2
2002.....	0.4	*	*	*	0.3	0.5	0.4	0.3	0.2	0.2
2001.....	0.5	*	*	*	0.4	0.6	0.5	0.3	0.2	0.2
2000.....	0.5	*	*	*	0.3	0.5	0.4	0.2	0.2	0.2
1999.....	0.5	*	*	*	0.3	0.7	0.4	0.2	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Melanoma and other malignant neoplasms of skin (C43-C44)

All races 3/, both sexes

2004.....	3.5	*	*	*	*	*	0.1	0.4	0.6	1.0
2003.....	3.5	*	*	*	*	*	0.1	0.4	0.7	0.9
2002.....	3.4	*	*	*	*	*	0.2	0.4	0.6	1.1
2001.....	3.5	*	*	*	*	*	0.2	0.4	0.6	1.0
2000.....	3.4	*	*	*	*	*	0.2	0.3	0.6	1.1
1999.....	3.4	*	*	*	*	*	0.2	0.3	0.7	1.2

All races 3/, male

2004.....	4.7	*	*	*	*	*	*	0.4	0.7	1.1
2003.....	4.6	*	*	*	*	*	*	0.5	0.8	1.2
2002.....	4.5	*	*	*	*	*	*	0.4	0.8	1.2
2001.....	4.6	*	*	*	*	*	*	0.5	0.6	1.3
2000.....	4.5	*	*	*	*	*	*	0.4	0.7	1.1
1999.....	4.5	*	*	*	*	*	*	0.4	0.8	1.5

All races 3/, female

2004.....	2.4	*	*	*	*	*	*	0.3	0.5	0.8
2003.....	2.4	*	*	*	*	*	*	0.3	0.5	0.7
2002.....	2.4	*	*	*	*	*	*	0.4	0.4	0.9
2001.....	2.5	*	*	*	*	*	*	0.3	0.5	0.8
2000.....	2.5	*	*	*	*	*	*	0.3	0.5	1.1
1999.....	2.4	*	*	*	*	*	*	0.2	0.6	0.8

White, both sexes

2004.....	4.2	*	*	*	*	*	0.2	0.4	0.8	1.2
2003.....	4.2	*	*	*	*	*	0.2	0.4	0.8	1.1
2002.....	4.1	*	*	*	*	*	0.2	0.5	0.8	1.3
2001.....	4.2	*	*	*	*	*	0.2	0.5	0.7	1.2
2000.....	4.0	*	*	*	*	*	0.2	0.4	0.8	1.3
1999.....	4.0	*	*	*	*	*	0.2	0.4	0.8	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant melanoma of skin (C43)

All races 3/, both sexes

2004.....	2.7	*	*	*	*	*	0.1	0.3	0.6	0.9
2003.....	2.7	*	*	*	*	*	0.1	0.3	0.6	0.9
2002.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2001.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2000.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0
1999.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0

All races 3/, male

2004.....	3.5	*	*	*	*	*	*	0.4	0.7	1.0
2003.....	3.5	*	*	*	*	*	*	0.4	0.8	1.1
2002.....	3.3	*	*	*	*	*	*	0.4	0.8	1.1
2001.....	3.4	*	*	*	*	*	*	0.4	0.6	1.2
2000.....	3.3	*	*	*	*	*	*	0.3	0.7	1.0
1999.....	3.3	*	*	*	*	*	*	0.4	0.7	1.3

All races 3/, female

2004.....	1.9	*	*	*	*	*	*	0.3	0.5	0.8
2003.....	1.9	*	*	*	*	*	*	0.2	0.5	0.7
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.9
2001.....	1.9	*	*	*	*	*	*	0.3	0.5	0.7
2000.....	2.0	*	*	*	*	*	*	0.3	0.5	1.0
1999.....	1.9	*	*	*	*	*	*	0.2	0.6	0.8

White, both sexes

2004.....	3.3	*	*	*	*	*	0.2	0.4	0.8	1.1
2003.....	3.2	*	*	*	*	*	0.2	0.4	0.8	1.1
2002.....	3.1	*	*	*	*	*	0.2	0.5	0.7	1.2
2001.....	3.2	*	*	*	*	*	0.2	0.5	0.7	1.2
2000.....	3.1	*	*	*	*	*	0.2	0.4	0.8	1.2
1999.....	3.1	*	*	*	*	*	0.2	0.4	0.8	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of mesothelial and soft tissue (C45-C49)

All races 3/, both sexes

2004.....	2.4	*	0.2	0.1	0.2	0.3	0.4	0.4	0.5	0.7
2003.....	2.4	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.6
2002.....	2.4	*	0.1	0.1	0.2	0.3	0.4	0.4	0.4	0.7
2001.....	2.4	*	*	0.1	0.2	0.3	0.3	0.4	0.6	0.6
2000.....	2.4	*	0.1	*	0.1	0.4	0.4	0.5	0.6	0.7
1999.....	2.4	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.7

All races 3/, male

2004.....	2.8	*	*	*	0.2	0.3	0.4	0.5	0.6	0.8
2003.....	2.8	*	*	*	*	0.4	0.4	0.5	0.6	0.7
2002.....	2.8	*	*	*	0.2	0.3	0.4	0.5	0.5	0.8
2001.....	2.8	*	*	*	*	0.4	0.4	0.5	0.7	0.6
2000.....	2.9	*	*	*	*	0.4	0.5	0.5	0.6	0.8
1999.....	2.8	*	*	*	0.2	0.4	0.4	0.5	0.7	0.9

All races 3/, female

2004.....	2.0	*	*	*	0.2	0.3	0.3	0.2	0.4	0.5
2003.....	2.0	*	*	*	0.2	0.3	0.3	0.4	0.4	0.6
2002.....	1.9	*	*	*	0.2	0.3	0.3	0.4	0.3	0.5
2001.....	2.0	*	*	*	*	0.3	0.2	0.4	0.4	0.6
2000.....	2.0	*	*	*	*	0.4	0.3	0.4	0.6	0.6
1999.....	1.9	*	*	*	*	*	0.4	0.4	0.4	0.6

White, both sexes

2004.....	2.7	*	*	0.2	0.2	0.3	0.3	0.4	0.5	0.6
2003.....	2.6	*	*	0.1	0.2	0.4	0.4	0.4	0.5	0.7
2002.....	2.6	*	*	0.1	0.2	0.3	0.4	0.4	0.4	0.7
2001.....	2.6	*	*	*	0.2	0.3	0.3	0.4	0.5	0.6
2000.....	2.7	*	*	*	0.1	0.4	0.3	0.5	0.6	0.6
1999.....	2.6	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of other mesothelial and soft tissue (C47-C49)

All races 3/, both sexes

2004.....	1.6	*	0.2	0.1	0.2	0.3	0.4	0.3	0.5	0.6
2003.....	1.5	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.6
2002.....	1.5	*	0.1	0.1	0.2	0.3	0.4	0.4	0.4	0.6
2001.....	1.5	*	*	0.1	0.2	0.3	0.3	0.4	0.5	0.5
2000.....	1.6	*	0.1	*	0.1	0.4	0.4	0.4	0.5	0.6
1999.....	1.5	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.7

All races 3/, male

2004.....	1.4	*	*	*	0.2	0.3	0.4	0.5	0.6	0.7
2003.....	1.4	*	*	*	*	0.4	0.4	0.4	0.6	0.6
2002.....	1.4	*	*	*	0.2	0.3	0.4	0.5	0.5	0.7
2001.....	1.4	*	*	*	*	0.4	0.4	0.4	0.6	0.5
2000.....	1.4	*	*	*	*	0.4	0.5	0.5	0.5	0.7
1999.....	1.5	*	*	*	0.2	0.4	0.4	0.5	0.6	0.7

All races 3/, female

2004.....	1.7	*	*	*	0.2	0.3	0.3	0.2	0.4	0.4
2003.....	1.6	*	*	*	0.2	0.3	0.3	0.3	0.4	0.5
2002.....	1.6	*	*	*	0.2	0.3	0.3	0.4	0.3	0.5
2001.....	1.6	*	*	*	*	0.3	0.2	0.4	0.4	0.5
2000.....	1.7	*	*	*	*	0.4	0.3	0.4	0.5	0.5
1999.....	1.6	*	*	*	*	*	0.4	0.4	0.4	0.6

White, both sexes

2004.....	1.7	*	*	0.2	0.2	0.3	0.3	0.4	0.5	0.6
2003.....	1.6	*	*	0.1	0.2	0.4	0.4	0.4	0.4	0.6
2002.....	1.6	*	*	0.1	0.2	0.3	0.4	0.4	0.4	0.6
2001.....	1.6	*	*	*	0.2	0.3	0.3	0.4	0.5	0.6
2000.....	1.7	*	*	*	0.1	0.3	0.3	0.5	0.6	0.6
1999.....	1.6	*	*	0.1	0.2	0.3	0.3	0.4	0.4	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of breast (C50)

All races 3/, both sexes

2004.....	14.1	*	*	*	*	*	*	0.4	1.6	3.9
2003.....	14.4	*	*	*	*	*	*	0.3	1.7	4.3
2002.....	14.5	*	*	*	*	*	*	0.3	1.7	4.1
2001.....	14.7	*	*	*	*	*	*	0.5	1.8	4.3
2000.....	15.0	*	*	*	*	*	*	0.4	1.9	4.3
1999.....	14.9	*	*	*	*	*	*	0.4	1.7	4.3

All races 3/, male

2004.....	0.3	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	27.5	*	*	*	*	*	*	0.8	3.2	7.8
2003.....	28.2	*	*	*	*	*	*	0.6	3.4	8.5
2002.....	28.3	*	*	*	*	*	*	0.7	3.5	8.2
2001.....	28.6	*	*	*	*	*	*	0.9	3.6	8.6
2000.....	29.2	*	*	*	*	*	*	0.8	3.8	8.5
1999.....	28.9	*	*	*	*	*	*	0.9	3.4	8.5

White, both sexes

2004.....	14.6	*	*	*	*	*	*	0.3	1.3	3.4
2003.....	15.0	*	*	*	*	*	*	0.2	1.5	3.8
2002.....	15.1	*	*	*	*	*	*	0.3	1.5	3.6
2001.....	15.2	*	*	*	*	*	*	0.4	1.5	3.9
2000.....	15.7	*	*	*	*	*	*	0.4	1.6	3.9
1999.....	15.5	*	*	*	*	*	*	0.4	1.5	3.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of breast (C50)

Black, female

2004.....	27.5	*	*	*	*	*	*	1.8	6.2	15.1
2003.....	28.5	*	*	*	*	*	*	1.6	6.5	15.1
2002.....	28.2	*	*	*	*	*	*	1.6	6.9	15.1
2001.....	28.3	*	*	*	*	*	*	2.2	7.6	14.2
2000.....	27.9	*	*	*	*	*	*	*	7.1	14.3
1999.....	28.2	*	*	*	*	*	*	1.8	6.9	16.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of female genital organs (C51-C58)

All races 3/, both sexes

2004.....	9.2	*	*	*	*	*	0.2	0.6	1.1	2.0
2003.....	9.3	*	*	*	*	*	0.2	0.5	1.1	2.0
2002.....	9.4	*	*	*	*	*	0.2	0.4	1.2	2.2
2001.....	9.4	*	*	*	*	*	0.2	0.5	1.2	2.3
2000.....	9.4	*	*	*	*	*	0.2	0.6	1.2	2.2
1999.....	9.3	*	*	*	*	*	0.2	0.5	1.1	2.2

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	18.1	*	*	*	*	*	0.4	1.1	2.2	4.1
2003.....	18.3	*	*	*	*	*	0.4	1.1	2.1	4.1
2002.....	18.5	*	*	*	*	*	0.5	0.9	2.5	4.5
2001.....	18.5	*	*	*	*	*	0.3	1.1	2.5	4.5
2000.....	18.4	*	*	*	*	*	0.4	1.2	2.4	4.4
1999.....	18.2	*	*	*	*	*	0.4	1.1	2.2	4.4

White, both sexes

2004.....	9.7	*	*	*	*	*	0.2	0.5	1.1	2.0
2003.....	9.8	*	*	*	*	*	0.2	0.5	1.0	1.9
2002.....	9.9	*	*	*	*	*	0.2	0.4	1.2	2.2
2001.....	9.9	*	*	*	*	*	0.1	0.5	1.2	2.2
2000.....	9.9	*	*	*	*	*	0.2	0.7	1.1	2.1
1999.....	9.8	*	*	*	*	*	0.2	0.5	1.1	2.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of female genital organs (C51-C58)

Black, female

2004.....	16.4	*	*	*	*	*	*	2.2	2.7	5.4
2003.....	16.3	*	*	*	*	*	*	2.0	3.1	6.0
2002.....	16.4	*	*	*	*	*	*	*	3.3	6.3
2001.....	16.2	*	*	*	*	*	*	1.7	3.2	5.6
2000.....	16.4	*	*	*	*	*	*	*	3.7	5.8
1999.....	16.1	*	*	*	*	*	*	*	2.5	5.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of cervix uteri (C53)

All races 3/, both sexes

2004.....	1.3	*	*	*	*	*	*	0.4	0.7	1.1
2003.....	1.3	*	*	*	*	*	*	0.3	0.7	1.2
2002.....	1.4	*	*	*	*	*	*	0.2	0.8	1.3
2001.....	1.4	*	*	*	*	*	*	0.3	0.8	1.3
2000.....	1.5	*	*	*	*	*	*	0.4	0.7	1.3
1999.....	1.5	*	*	*	*	*	*	0.3	0.8	1.3

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	2.6	*	*	*	*	*	*	0.7	1.4	2.2
2003.....	2.7	*	*	*	*	*	*	0.6	1.4	2.3
2002.....	2.7	*	*	*	*	*	*	0.5	1.5	2.5
2001.....	2.8	*	*	*	*	*	*	0.5	1.5	2.6
2000.....	2.9	*	*	*	*	*	*	0.8	1.5	2.6
1999.....	3.0	*	*	*	*	*	*	0.6	1.5	2.5

White, both sexes

2004.....	1.2	*	*	*	*	*	*	0.3	0.7	1.1
2003.....	1.2	*	*	*	*	*	*	0.3	0.6	1.1
2002.....	1.3	*	*	*	*	*	*	0.2	0.8	1.2
2001.....	1.3	*	*	*	*	*	*	0.3	0.7	1.3
2000.....	1.4	*	*	*	*	*	*	0.4	0.7	1.2
1999.....	1.4	*	*	*	*	*	*	0.3	0.7	1.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of cervix uteri (C53)

Black, female

2004.....	3.9	*	*	*	*	*	*	1.4	1.7	3.0
2003.....	4.0	*	*	*	*	*	*	*	2.2	3.8
2002.....	4.2	*	*	*	*	*	*	*	2.0	3.9
2001.....	4.0	*	*	*	*	*	*	*	2.1	3.5
2000.....	4.5	*	*	*	*	*	*	*	2.7	4.0
1999.....	4.4	*	*	*	*	*	*	*	1.9	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of ovary (C56)

All races 3/, both sexes

2004.....	5.0	*	*	*	*	*	0.1	0.2	0.3	0.6
2003.....	5.0	*	*	*	*	*	*	0.2	0.3	0.6
2002.....	5.1	*	*	*	*	*	0.1	0.2	0.3	0.7
2001.....	5.1	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	5.0	*	*	*	*	*	*	0.2	0.3	0.6
1999.....	4.9	*	*	*	*	*	*	0.1	0.2	0.7

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	9.9	*	*	*	*	*	0.2	0.3	0.6	1.2
2003.....	9.9	*	*	*	*	*	*	0.3	0.5	1.3
2002.....	10.0	*	*	*	*	*	0.3	0.4	0.6	1.4
2001.....	9.9	*	*	*	*	*	*	0.4	0.6	1.4
2000.....	9.8	*	*	*	*	*	*	0.4	0.6	1.3
1999.....	9.6	*	*	*	*	*	*	0.3	0.5	1.4

White, both sexes

2004.....	5.5	*	*	*	*	*	*	0.1	0.3	0.6
2003.....	5.6	*	*	*	*	*	*	0.2	0.3	0.6
2002.....	5.6	*	*	*	*	*	0.1	0.2	0.3	0.7
2001.....	5.6	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	5.5	*	*	*	*	*	*	0.2	0.3	0.6
1999.....	5.4	*	*	*	*	*	*	0.1	0.2	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of male genital organs (C60-C63)

All races 3/, both sexes

2004.....	10.1	*	*	*	*	*	0.2	0.2	0.2	0.2
2003.....	10.4	*	*	*	*	*	0.1	0.2	0.2	0.3
2002.....	10.8	*	*	*	*	*	0.2	0.2	0.3	0.3
2001.....	11.0	*	*	*	*	*	0.1	0.2	0.2	0.2
2000.....	11.3	*	*	*	*	*	0.2	0.2	0.2	0.3
1999.....	11.6	*	*	*	*	*	0.2	0.3	0.3	0.3

All races 3/, male

2004.....	20.5	*	*	*	*	*	0.4	0.4	0.4	0.5
2003.....	21.1	*	*	*	*	*	0.2	0.4	0.4	0.6
2002.....	21.9	*	*	*	*	*	0.3	0.4	0.5	0.5
2001.....	22.4	*	*	*	*	*	0.3	0.5	0.4	0.5
2000.....	22.9	*	*	*	*	*	0.4	0.4	0.4	0.6
1999.....	23.6	*	*	*	*	*	0.3	0.5	0.5	0.6

All races 3/, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	10.2	*	*	*	*	*	0.2	0.2	0.2	0.3
2003.....	10.5	*	*	*	*	*	0.1	0.2	0.3	0.3
2002.....	10.9	*	*	*	*	*	0.2	0.2	0.3	0.3
2001.....	11.0	*	*	*	*	*	0.2	0.3	0.2	0.3
2000.....	11.2	*	*	*	*	*	0.2	0.3	0.2	0.3
1999.....	11.6	*	*	*	*	*	0.2	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of testis (C62)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
2003.....	0.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2002.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
2001.....	0.1	*	*	*	*	*	0.1	0.2	0.2	0.2
2000.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
1999.....	0.1	*	*	*	*	*	0.1	0.2	0.3	0.2

All races 3/, male

2004.....	0.2	*	*	*	*	*	0.3	0.4	0.4	0.4
2003.....	0.2	*	*	*	*	*	0.2	0.4	0.4	0.5
2002.....	0.3	*	*	*	*	*	0.3	0.4	0.5	0.4
2001.....	0.2	*	*	*	*	*	0.3	0.4	0.4	0.4
2000.....	0.2	*	*	*	*	*	0.3	0.4	0.4	0.5
1999.....	0.3	*	*	*	*	*	0.3	0.5	0.5	0.5

All races 3/, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.3
2003.....	0.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2002.....	0.2	*	*	*	*	*	0.2	0.2	0.3	0.2
2001.....	0.1	*	*	*	*	*	0.2	0.3	0.2	0.2
2000.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.3
1999.....	0.2	*	*	*	*	*	0.2	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of urinary tract (C64-C68)

All races 3/, both sexes

2004.....	8.8	*	*	0.1	*	*	*	0.1	0.2	0.5
2003.....	8.7	*	*	0.1	*	*	*	0.1	0.2	0.5
2002.....	8.8	*	*	0.1	*	*	*	*	0.2	0.6
2001.....	8.7	*	0.1	0.2	*	*	*	0.1	0.2	0.5
2000.....	8.7	*	*	0.1	*	*	*	*	0.2	0.6
1999.....	8.5	*	*	0.1	*	*	*	0.2	0.2	0.6

All races 3/, male

2004.....	11.7	*	*	*	*	*	*	*	0.2	0.7
2003.....	11.6	*	*	*	*	*	*	*	0.3	0.7
2002.....	11.6	*	*	*	*	*	*	*	0.3	0.8
2001.....	11.6	*	*	*	*	*	*	*	0.2	0.6
2000.....	11.4	*	*	*	*	*	*	*	0.2	0.7
1999.....	11.2	*	*	*	*	*	*	*	0.3	0.8

All races 3/, female

2004.....	6.0	*	*	*	*	*	*	*	*	0.4
2003.....	6.0	*	*	*	*	*	*	*	*	0.3
2002.....	6.1	*	*	*	*	*	*	*	*	0.4
2001.....	6.0	*	*	0.2	*	*	*	*	*	0.3
2000.....	6.0	*	*	*	*	*	*	*	*	0.4
1999.....	5.9	*	*	*	*	*	*	*	*	0.4

White, both sexes

2004.....	9.9	*	*	*	*	*	*	*	0.1	0.5
2003.....	9.7	*	*	0.2	*	*	*	*	0.2	0.4
2002.....	9.8	*	*	0.1	*	*	*	*	0.2	0.5
2001.....	9.7	*	*	0.1	*	*	*	*	0.2	0.4
2000.....	9.6	*	*	*	*	*	*	*	0.2	0.5
1999.....	9.3	*	*	*	*	*	*	*	0.2	0.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes

2004.....	4.4	*	0.6	0.9	0.7	0.5	0.5	0.7	1.1	1.7
2003.....	4.4	*	0.7	0.9	0.8	0.6	0.6	0.9	1.2	1.9
2002.....	4.4	*	0.7	1.0	0.6	0.5	0.5	0.7	1.2	1.7
2001.....	4.4	*	0.7	0.8	0.6	0.5	0.6	0.7	1.4	1.8
2000.....	4.5	0.5	0.7	0.8	0.7	0.5	0.6	0.8	1.2	2.0
1999.....	4.6	*	0.7	0.9	0.6	0.6	0.6	0.9	1.3	1.9

All races 3/, male

2004.....	4.9	*	0.7	1.0	0.7	0.6	0.7	0.8	1.3	2.1
2003.....	5.0	*	0.7	0.9	0.8	0.6	0.6	0.9	1.4	2.3
2002.....	5.0	*	0.8	1.1	0.7	0.6	0.7	0.7	1.5	2.0
2001.....	5.0	*	0.7	0.8	0.6	0.6	0.7	0.9	1.7	2.2
2000.....	5.1	*	0.7	0.8	0.8	0.6	0.7	1.0	1.6	2.6
1999.....	5.1	*	0.7	0.9	0.6	0.7	0.7	1.1	1.7	2.3

All races 3/, female

2004.....	3.9	*	0.6	0.9	0.7	0.5	0.4	0.6	0.8	1.2
2003.....	3.9	*	0.6	0.9	0.7	0.5	0.5	0.9	0.9	1.5
2002.....	3.9	*	0.7	0.9	0.5	0.4	0.3	0.7	1.0	1.4
2001.....	3.9	*	0.6	0.9	0.6	0.4	0.6	0.5	1.1	1.4
2000.....	3.9	*	0.6	0.8	0.6	0.4	0.5	0.7	0.8	1.4
1999.....	4.1	*	0.8	0.9	0.7	0.5	0.5	0.8	0.9	1.5

White, both sexes

2004.....	4.9	*	0.7	1.0	0.6	0.6	0.6	0.7	1.2	1.9
2003.....	5.0	*	0.7	0.9	0.8	0.6	0.6	1.0	1.3	2.1
2002.....	5.0	*	0.7	1.0	0.6	0.5	0.5	0.7	1.4	1.9
2001.....	5.0	*	0.7	0.9	0.6	0.6	0.6	0.8	1.5	2.1
2000.....	5.1	*	0.7	0.8	0.7	0.5	0.7	0.9	1.3	2.2
1999.....	5.2	*	0.8	0.9	0.6	0.5	0.6	1.1	1.4	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of brain (C71)

All races 3/, both sexes

2004.....	4.3	*	0.6	0.9	0.6	0.5	0.5	0.7	1.1	1.6
2003.....	4.4	*	0.6	0.9	0.7	0.6	0.5	0.8	1.1	1.9
2002.....	4.4	*	0.7	1.0	0.6	0.5	0.5	0.7	1.2	1.7
2001.....	4.3	*	0.6	0.8	0.6	0.5	0.6	0.7	1.3	1.8
2000.....	4.4	0.5	0.7	0.8	0.7	0.5	0.6	0.8	1.2	2.0
1999.....	4.5	*	0.7	0.9	0.6	0.5	0.5	0.9	1.3	1.9

All races 3/, male

2004.....	4.8	*	0.7	1.0	0.6	0.6	0.6	0.8	1.3	2.1
2003.....	4.9	*	0.7	0.8	0.8	0.6	0.6	0.9	1.4	2.3
2002.....	4.9	*	0.8	1.1	0.7	0.6	0.7	0.7	1.4	2.0
2001.....	4.9	*	0.6	0.8	0.6	0.6	0.6	0.9	1.6	2.2
2000.....	5.0	*	0.7	0.8	0.8	0.6	0.7	0.9	1.6	2.6
1999.....	5.0	*	0.7	0.9	0.6	0.6	0.6	1.0	1.6	2.2

All races 3/, female

2004.....	3.8	*	0.5	0.9	0.6	0.4	0.4	0.6	0.8	1.2
2003.....	3.8	*	0.6	0.9	0.7	0.5	0.5	0.8	0.9	1.5
2002.....	3.8	*	0.6	0.9	0.5	0.4	0.3	0.7	1.0	1.4
2001.....	3.8	*	0.6	0.9	0.6	0.4	0.6	0.5	1.0	1.4
2000.....	3.9	*	0.6	0.7	0.6	0.4	0.5	0.7	0.7	1.4
1999.....	4.0	*	0.7	0.9	0.6	0.4	0.5	0.8	0.9	1.5

White, both sexes

2004.....	4.9	*	0.6	0.9	0.6	0.5	0.6	0.7	1.2	1.9
2003.....	4.9	*	0.7	0.9	0.8	0.6	0.6	0.9	1.3	2.1
2002.....	4.9	*	0.6	1.0	0.6	0.5	0.5	0.7	1.3	1.9
2001.....	4.9	*	0.7	0.9	0.6	0.5	0.6	0.8	1.5	2.1
2000.....	5.0	*	0.7	0.8	0.7	0.5	0.6	0.9	1.3	2.2
1999.....	5.1	*	0.8	0.8	0.6	0.5	0.6	1.0	1.4	2.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of thyroid and other endocrine glands (C73-C75)

All races 3/, both sexes

2004.....	0.7	*	0.5	0.4	0.1	*	0.1	*	*	0.2
2003.....	0.7	*	0.4	0.4	0.1	*	*	*	0.1	0.2
2002.....	0.7	*	0.5	0.4	0.1	*	*	*	0.1	0.1
2001.....	0.7	*	0.5	0.3	0.1	*	*	0.1	*	0.2
2000.....	0.7	*	0.5	0.3	0.1	*	*	*	0.1	0.2
1999.....	0.7	*	0.5	0.4	0.1	*	*	*	0.1	0.2

All races 3/, male

2004.....	0.6	*	0.5	0.4	*	*	*	*	*	*
2003.....	0.6	*	0.5	0.5	*	*	*	*	*	0.2
2002.....	0.6	*	0.4	0.4	*	*	*	*	*	*
2001.....	0.7	*	0.5	0.3	*	*	*	*	*	*
2000.....	0.7	*	0.5	0.3	*	*	*	*	*	*
1999.....	0.7	*	0.5	0.4	*	*	*	*	*	0.2

All races 3/, female

2004.....	0.8	*	0.5	0.3	*	*	*	*	*	*
2003.....	0.7	*	0.4	0.4	*	*	*	*	*	0.2
2002.....	0.8	*	0.5	0.3	*	*	*	*	*	*
2001.....	0.8	*	0.4	0.2	*	*	*	*	*	*
2000.....	0.8	*	0.5	0.3	*	*	*	*	*	0.2
1999.....	0.8	*	0.4	0.4	*	*	*	*	*	0.2

White, both sexes

2004.....	0.8	*	0.5	0.3	*	*	*	*	*	0.2
2003.....	0.7	*	0.4	0.4	*	*	*	*	*	0.2
2002.....	0.8	*	0.4	0.3	0.1	*	*	*	*	0.2
2001.....	0.8	*	0.5	0.3	*	*	*	*	*	0.1
2000.....	0.8	*	0.5	0.4	*	*	*	*	*	0.2
1999.....	0.8	*	0.5	0.4	*	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

All races 3/, both sexes

2004.....	13.5	*	*	*	*	0.2	0.3	0.5	0.8	1.4
2003.....	13.8	*	*	*	0.1	0.2	0.3	0.4	0.6	1.4
2002.....	14.1	*	0.1	*	*	0.3	0.3	0.4	0.9	1.5
2001.....	14.1	*	*	*	0.1	0.2	0.3	0.5	0.8	1.6
2000.....	14.5	*	0.1	*	0.1	0.2	0.3	0.4	0.7	1.6
1999.....	15.7	*	*	*	0.1	0.2	0.4	0.5	0.9	1.7

All races 3/, male

2004.....	14.1	*	*	*	*	0.3	0.4	0.5	0.8	1.5
2003.....	14.3	*	*	*	*	0.3	0.3	0.4	0.7	1.3
2002.....	14.7	*	*	*	*	0.3	0.4	0.4	0.9	1.6
2001.....	14.5	*	*	*	*	0.3	0.3	0.5	0.9	1.5
2000.....	14.8	*	*	*	*	0.2	0.3	0.5	0.7	1.5
1999.....	16.2	*	*	*	*	0.2	0.5	0.6	1.1	1.7

All races 3/, female

2004.....	13.0	*	*	*	*	*	*	0.5	0.8	1.3
2003.....	13.4	*	*	*	*	*	0.3	0.4	0.6	1.5
2002.....	13.5	*	*	*	*	0.2	0.2	0.3	0.8	1.5
2001.....	13.6	*	*	*	*	*	0.2	0.5	0.6	1.6
2000.....	14.1	*	*	*	*	0.2	0.3	0.4	0.7	1.7
1999.....	15.3	*	*	*	*	*	0.3	0.4	0.8	1.7

White, both sexes

2004.....	14.4	*	*	*	*	0.2	0.3	0.5	0.8	1.4
2003.....	14.7	*	*	*	*	0.2	0.3	0.4	0.6	1.3
2002.....	14.9	*	*	*	*	0.3	0.3	0.3	0.9	1.4
2001.....	14.8	*	*	*	*	0.2	0.3	0.4	0.7	1.5
2000.....	15.3	*	*	*	*	0.2	0.3	0.5	0.7	1.5
1999.....	16.6	*	*	*	*	0.2	0.4	0.5	0.9	1.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes

2004.....	18.6	0.6	0.8	0.9	0.9	1.4	1.7	1.8	2.3	2.7
2003.....	19.1	0.8	0.9	0.9	1.0	1.2	1.7	1.9	2.4	3.1
2002.....	19.5	0.7	0.8	0.8	1.0	1.5	1.9	1.9	2.6	3.2
2001.....	19.8	0.7	1.0	0.8	1.0	1.5	2.0	2.3	2.7	3.5
2000.....	20.1	1.0	0.9	0.8	1.0	1.6	2.1	2.1	2.5	3.4
1999.....	20.2	0.8	0.9	0.7	1.0	1.5	2.2	2.2	2.6	3.7

All races 3/, male

2004.....	20.4	*	0.9	1.1	1.0	1.6	2.1	2.1	2.7	3.2
2003.....	21.0	*	1.0	1.0	1.1	1.6	2.0	2.3	3.0	3.5
2002.....	21.4	*	0.8	0.9	1.2	1.8	2.2	2.3	3.0	3.7
2001.....	21.5	*	1.0	0.9	1.1	1.8	2.3	2.6	3.1	4.2
2000.....	21.6	1.1	1.0	0.9	1.2	1.9	2.6	2.3	3.0	3.9
1999.....	21.7	*	1.0	0.8	1.2	1.7	2.7	2.4	3.1	4.4

All races 3/, female

2004.....	16.8	*	0.6	0.7	0.7	1.1	1.3	1.5	1.9	2.2
2003.....	17.3	1.1	0.8	0.7	0.9	0.9	1.4	1.6	1.7	2.6
2002.....	17.7	*	0.7	0.8	0.8	1.2	1.7	1.5	2.2	2.6
2001.....	18.1	*	1.0	0.7	0.8	1.2	1.7	2.0	2.2	2.7
2000.....	18.6	*	0.9	0.7	0.8	1.2	1.5	1.9	1.9	3.0
1999.....	18.7	*	0.8	0.6	0.9	1.3	1.8	1.9	2.2	3.0

White, both sexes

2004.....	20.2	0.6	0.8	1.0	0.9	1.4	1.7	1.7	2.2	2.6
2003.....	20.9	0.9	0.9	0.9	1.0	1.3	1.7	1.9	2.3	3.0
2002.....	21.3	*	0.8	0.9	1.0	1.5	2.1	1.9	2.5	3.1
2001.....	21.5	0.7	1.1	0.8	1.0	1.6	2.0	2.1	2.6	3.4
2000.....	21.9	1.2	1.0	0.9	1.1	1.6	2.0	2.1	2.4	3.3
1999.....	21.9	0.8	1.0	0.7	1.0	1.5	2.3	2.1	2.5	3.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
White, male										
2004.....	22.2	*	0.8	1.2	1.0	1.7	2.1	2.1	2.6	3.1
2003.....	23.1	*	1.1	1.2	1.1	1.6	2.0	2.2	2.9	3.4
2002.....	23.4	*	0.9	0.9	1.2	1.9	2.4	2.3	2.9	3.5
2001.....	23.3	*	1.1	0.9	1.0	1.9	2.3	2.4	3.2	4.0
2000.....	23.6	*	1.0	0.9	1.2	1.9	2.5	2.3	2.9	3.8
1999.....	23.6	*	1.0	0.9	1.1	1.7	2.8	2.3	2.9	4.3
White, female										
2004.....	18.3	*	0.7	0.7	0.8	1.1	1.3	1.3	1.8	2.1
2003.....	18.8	1.4	0.8	0.7	0.9	0.8	1.5	1.6	1.7	2.5
2002.....	19.2	*	0.8	0.9	0.8	1.1	1.7	1.5	2.2	2.6
2001.....	19.7	*	1.1	0.7	0.9	1.2	1.6	1.9	2.0	2.8
2000.....	20.3	*	1.0	0.8	0.9	1.2	1.4	1.8	1.9	2.9
1999.....	20.4	*	0.9	0.6	0.9	1.3	1.7	1.9	2.1	2.9
Black, both sexes										
2004.....	13.6	*	*	0.7	0.8	1.4	1.9	2.8	3.1	3.7
2003.....	13.4	*	*	0.6	1.0	1.1	1.7	2.4	3.3	4.1
2002.....	13.6	*	*	*	1.0	1.4	1.5	2.3	3.4	3.8
2001.....	14.1	*	*	0.9	1.0	1.5	2.1	3.2	3.5	4.5
2000.....	13.9	*	0.8	0.6	0.7	1.9	2.6	2.8	3.1	4.4
1999.....	14.1	*	*	0.6	1.3	1.7	2.9	2.6	3.2	4.6
Black, male										
2004.....	14.4	*	*	*	*	1.4	2.5	2.9	3.6	4.3
2003.....	13.9	*	*	*	1.2	*	2.1	2.5	4.2	4.7
2002.....	14.6	*	*	*	1.1	1.6	1.3	2.8	4.4	5.1
2001.....	15.1	*	*	*	1.7	1.8	2.2	3.8	4.0	6.5
2000.....	14.7	*	*	*	*	2.3	3.2	2.6	4.4	4.8
1999.....	15.1	*	*	*	1.7	2.1	3.1	3.3	4.2	5.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Black, female

2004.....	12.8	*	*	*	*	1.4	1.3	2.7	2.7	3.1
2003.....	12.9	*	*	*	*	*	1.3	2.4	2.4	3.6
2002.....	12.8	*	*	*	*	1.3	1.6	2.0	2.5	2.7
2001.....	13.1	*	*	*	*	*	2.1	2.7	3.1	2.8
2000.....	13.1	*	*	*	*	1.4	2.1	3.0	1.9	4.1
1999.....	13.2	*	*	*	*	1.4	2.8	2.0	2.3	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Hodgkin's disease (C81)

All races 3/, both sexes

2004.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.3
1999.....	0.5	*	*	*	*	0.2	0.3	0.4	0.3	0.4

All races 3/, male

2004.....	0.5	*	*	*	*	*	0.3	0.3	0.4	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.5
2000.....	0.5	*	*	*	*	*	0.3	0.3	0.5	0.4
1999.....	0.6	*	*	*	*	0.2	0.4	0.4	0.4	0.5

All races 3/, female

2004.....	0.4	*	*	*	*	*	*	0.3	0.3	0.2
2003.....	0.4	*	*	*	*	*	0.2	0.4	0.4	0.3
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3
2001.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.3
2000.....	0.4	*	*	*	*	*	0.2	0.3	0.4	0.3
1999.....	0.4	*	*	*	*	*	0.3	0.4	0.3	0.3

White, both sexes

2004.....	0.5	*	*	*	*	*	0.2	0.3	0.3	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4
1999.....	0.6	*	*	*	*	0.2	0.3	0.4	0.4	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

All races 3/, both sexes

2004.....	7.2	*	*	0.1	*	0.3	0.4	0.6	0.7	0.9
2003.....	7.4	*	*	*	0.1	0.3	0.4	0.5	0.8	1.1
2002.....	7.6	*	*	*	0.2	0.2	0.5	0.6	0.8	1.2
2001.....	7.9	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4
2000.....	8.1	*	*	0.1	0.1	0.3	0.6	0.6	0.8	1.3
1999.....	8.2	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4

All races 3/, male

2004.....	7.7	*	*	*	*	0.3	0.5	0.9	0.8	1.2
2003.....	8.0	*	*	*	*	0.3	0.5	0.6	1.0	1.4
2002.....	8.2	*	*	*	0.3	0.3	0.6	0.7	1.0	1.5
2001.....	8.4	*	*	*	0.2	0.4	0.5	0.8	1.0	1.8
2000.....	8.6	*	*	*	*	0.4	0.8	0.7	1.0	1.7
1999.....	8.6	*	*	*	0.2	0.3	0.6	0.7	1.1	1.8

All races 3/, female

2004.....	6.7	*	*	*	*	*	0.3	0.4	0.6	0.6
2003.....	6.8	*	*	*	*	*	0.3	0.4	0.5	0.7
2002.....	7.1	*	*	*	*	*	0.4	0.4	0.6	0.8
2001.....	7.3	*	*	*	*	*	0.4	0.6	0.6	1.0
2000.....	7.6	*	*	*	*	*	0.4	0.6	0.6	0.9
1999.....	7.8	*	*	*	*	0.2	0.4	0.4	0.7	1.1

White, both sexes

2004.....	8.0	*	*	*	*	0.3	0.4	0.6	0.6	0.9
2003.....	8.3	*	*	*	*	0.2	0.4	0.4	0.7	1.1
2002.....	8.5	*	*	*	0.2	0.2	0.5	0.6	0.7	1.1
2001.....	8.8	*	*	*	0.1	0.3	0.4	0.6	0.8	1.3
2000.....	9.1	*	*	*	0.1	0.3	0.6	0.6	0.8	1.2
1999.....	9.2	*	*	*	*	0.2	0.5	0.5	0.8	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

White, male

2004.....	8.5	*	*	*	*	0.4	0.5	0.8	0.7	1.3
2003.....	9.0	*	*	*	*	0.3	0.5	0.6	0.9	1.4
2002.....	9.1	*	*	*	0.3	0.3	0.6	0.8	0.9	1.4
2001.....	9.3	*	*	*	*	0.4	0.5	0.7	1.0	1.6
2000.....	9.6	*	*	*	*	0.5	0.8	0.6	1.0	1.6
1999.....	9.6	*	*	*	*	0.3	0.6	0.6	1.0	1.7

White, female

2004.....	7.5	*	*	*	*	*	0.3	0.3	0.5	0.6
2003.....	7.7	*	*	*	*	*	0.4	0.3	0.4	0.7
2002.....	8.0	*	*	*	*	*	0.3	0.4	0.6	0.8
2001.....	8.3	*	*	*	*	*	0.4	0.5	0.5	1.0
2000.....	8.7	*	*	*	*	*	0.4	0.5	0.6	0.8
1999.....	8.8	*	*	*	*	*	0.4	0.4	0.6	1.0

Black, both sexes

2004.....	3.7	*	*	*	*	*	0.7	1.1	1.0	1.2
2003.....	3.6	*	*	*	*	*	*	0.9	1.2	1.5
2002.....	3.9	*	*	*	*	*	*	0.8	1.5	1.4
2001.....	4.1	*	*	*	*	*	*	1.1	1.3	2.0
2000.....	3.8	*	*	*	*	*	1.0	1.1	0.9	1.9
1999.....	4.0	*	*	*	*	*	*	0.9	1.2	2.1

Black, male

2004.....	4.0	*	*	*	*	*	*	*	*	*
2003.....	3.8	*	*	*	*	*	*	*	1.8	1.6
2002.....	4.3	*	*	*	*	*	*	*	2.2	2.0
2001.....	4.6	*	*	*	*	*	*	*	*	3.1
2000.....	4.2	*	*	*	*	*	*	*	*	2.7
1999.....	4.6	*	*	*	*	*	*	*	1.7	2.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Non-Hodgkin's lymphoma (C82-C85)

All races 3/, both sexes

2004.....	7.1	*	*	0.1	*	0.3	0.4	0.6	0.7	0.9
2003.....	7.4	*	*	*	0.1	0.3	0.4	0.5	0.8	1.1
2002.....	7.6	*	*	*	0.2	0.2	0.5	0.6	0.8	1.2
2001.....	7.8	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4
2000.....	8.1	*	*	0.1	0.1	0.3	0.6	0.6	0.8	1.3
1999.....	8.2	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4

All races 3/, male

2004.....	7.6	*	*	*	*	0.3	0.5	0.8	0.8	1.2
2003.....	8.0	*	*	*	*	0.3	0.5	0.6	1.0	1.4
2002.....	8.1	*	*	*	0.3	0.3	0.6	0.7	1.0	1.5
2001.....	8.4	*	*	*	0.2	0.4	0.5	0.8	1.0	1.8
2000.....	8.6	*	*	*	*	0.4	0.8	0.7	1.0	1.7
1999.....	8.6	*	*	*	0.2	0.3	0.6	0.7	1.1	1.8

All races 3/, female

2004.....	6.6	*	*	*	*	*	0.3	0.4	0.6	0.6
2003.....	6.8	*	*	*	*	*	0.3	0.4	0.5	0.7
2002.....	7.1	*	*	*	*	*	0.4	0.4	0.6	0.8
2001.....	7.3	*	*	*	*	*	0.4	0.6	0.6	1.0
2000.....	7.6	*	*	*	*	*	0.4	0.6	0.6	0.9
1999.....	7.7	*	*	*	*	0.2	0.4	0.4	0.7	1.0

White, both sexes

2004.....	8.0	*	*	*	*	0.3	0.4	0.5	0.6	0.9
2003.....	8.3	*	*	*	*	0.2	0.4	0.4	0.7	1.1
2002.....	8.5	*	*	*	0.2	0.2	0.5	0.6	0.7	1.1
2001.....	8.8	*	*	*	0.1	0.3	0.4	0.6	0.8	1.3
2000.....	9.1	*	*	*	0.1	0.3	0.6	0.6	0.8	1.2
1999.....	9.2	*	*	*	*	0.2	0.5	0.5	0.8	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Non-Hodgkin's lymphoma (C82-C85)

White, male

2004.....	8.5	*	*	*	*	0.4	0.5	0.8	0.7	1.2
2003.....	8.9	*	*	*	*	0.3	0.5	0.6	0.9	1.4
2002.....	9.1	*	*	*	0.3	0.3	0.6	0.8	0.9	1.4
2001.....	9.3	*	*	*	*	0.4	0.5	0.7	1.0	1.6
2000.....	9.6	*	*	*	*	0.5	0.8	0.6	1.0	1.6
1999.....	9.6	*	*	*	*	0.3	0.6	0.6	1.0	1.7

White, female

2004.....	7.5	*	*	*	*	*	0.3	0.3	0.5	0.6
2003.....	7.7	*	*	*	*	*	0.4	0.3	0.4	0.7
2002.....	8.0	*	*	*	*	*	0.3	0.4	0.6	0.8
2001.....	8.2	*	*	*	*	*	0.4	0.5	0.5	1.0
2000.....	8.6	*	*	*	*	*	0.4	0.5	0.6	0.8
1999.....	8.8	*	*	*	*	*	0.4	0.4	0.6	1.0

Black, both sexes

2004.....	3.7	*	*	*	*	*	0.7	1.1	1.0	1.2
2003.....	3.6	*	*	*	*	*	*	0.9	1.2	1.5
2002.....	3.9	*	*	*	*	*	*	0.8	1.5	1.4
2001.....	4.0	*	*	*	*	*	*	1.1	1.2	2.0
2000.....	3.7	*	*	*	*	*	1.0	1.1	0.9	1.9
1999.....	4.0	*	*	*	*	*	*	0.9	1.2	2.1

Black, male

2004.....	4.0	*	*	*	*	*	*	*	*	*
2003.....	3.8	*	*	*	*	*	*	*	1.8	1.6
2002.....	4.3	*	*	*	*	*	*	*	2.2	2.0
2001.....	4.5	*	*	*	*	*	*	*	*	3.0
2000.....	4.2	*	*	*	*	*	*	*	*	2.7
1999.....	4.6	*	*	*	*	*	*	*	1.7	2.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Leukemia (C91-C95)

All races 3/, both sexes

2004.....	7.3	0.6	0.8	0.8	0.7	1.0	1.1	0.9	1.2	1.4
2003.....	7.4	0.7	0.8	0.8	0.9	0.9	1.1	1.1	1.1	1.5
2002.....	7.5	0.5	0.7	0.7	0.8	1.2	1.2	1.0	1.3	1.5
2001.....	7.5	0.6	1.0	0.8	0.8	1.1	1.3	1.2	1.4	1.6
2000.....	7.6	1.0	0.9	0.7	0.8	1.2	1.2	1.2	1.2	1.6
1999.....	7.5	0.7	0.8	0.6	0.9	1.1	1.4	1.2	1.3	1.6

All races 3/, male

2004.....	8.3	*	0.9	1.0	0.8	1.2	1.3	1.0	1.4	1.5
2003.....	8.4	*	0.9	0.9	1.0	1.2	1.2	1.3	1.4	1.6
2002.....	8.5	*	0.7	0.7	0.9	1.4	1.4	1.2	1.5	1.6
2001.....	8.5	*	1.0	0.8	0.9	1.3	1.5	1.4	1.6	1.8
2000.....	8.5	1.1	0.9	0.8	1.0	1.4	1.5	1.3	1.5	1.6
1999.....	8.4	*	0.8	0.7	0.9	1.3	1.7	1.3	1.4	1.8

All races 3/, female

2004.....	6.3	*	0.6	0.6	0.7	0.8	0.8	0.8	0.9	1.2
2003.....	6.4	1.1	0.8	0.7	0.8	0.6	0.9	0.9	0.8	1.4
2002.....	6.5	*	0.6	0.8	0.7	1.0	1.0	0.8	1.1	1.3
2001.....	6.6	*	1.0	0.7	0.7	0.9	1.0	1.0	1.2	1.3
2000.....	6.7	*	0.9	0.6	0.6	0.9	0.9	1.0	0.9	1.6
1999.....	6.7	*	0.7	0.5	0.8	0.9	1.1	1.1	1.2	1.5

White, both sexes

2004.....	8.0	*	0.7	0.8	0.8	1.0	1.1	0.9	1.2	1.3
2003.....	8.2	0.9	0.9	0.9	0.9	0.9	1.1	1.1	1.1	1.4
2002.....	8.2	*	0.7	0.8	0.8	1.2	1.3	1.0	1.3	1.5
2001.....	8.3	0.6	1.1	0.7	0.8	1.2	1.3	1.1	1.4	1.5
2000.....	8.3	1.2	1.0	0.7	0.9	1.2	1.2	1.1	1.1	1.6
1999.....	8.3	0.8	0.9	0.7	0.8	1.1	1.4	1.2	1.3	1.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Leukemia (C91-C95)

White, male

2004.....	9.1	*	0.8	1.0	0.8	1.2	1.4	1.0	1.4	1.4
2003.....	9.4	*	1.0	1.0	1.0	1.2	1.2	1.3	1.4	1.4
2002.....	9.3	*	0.8	0.7	0.9	1.4	1.5	1.2	1.5	1.6
2001.....	9.3	*	1.1	0.8	0.8	1.4	1.6	1.3	1.6	1.7
2000.....	9.4	*	0.9	0.8	1.1	1.4	1.5	1.3	1.4	1.7
1999.....	9.2	*	0.9	0.8	0.9	1.2	1.8	1.3	1.4	1.8

White, female

2004.....	6.9	*	0.7	0.7	0.7	0.8	0.9	0.8	1.0	1.2
2003.....	7.0	1.4	0.8	0.7	0.8	0.6	0.9	0.9	0.8	1.4
2002.....	7.1	*	0.6	0.8	0.7	1.0	1.1	0.8	1.1	1.4
2001.....	7.3	*	1.1	0.7	0.8	0.9	1.0	0.9	1.2	1.4
2000.....	7.4	*	1.0	0.7	0.7	0.9	0.9	0.9	0.9	1.6
1999.....	7.3	*	0.8	0.6	0.8	1.0	1.0	1.1	1.2	1.5

Black, both sexes

2004.....	4.8	*	*	*	0.6	1.1	0.9	1.1	1.5	1.8
2003.....	4.6	*	*	*	0.8	0.7	1.0	1.0	1.4	2.0
2002.....	4.7	*	*	*	0.7	0.9	0.8	1.1	1.4	1.5
2001.....	4.7	*	*	0.8	0.9	0.9	1.2	1.4	1.4	1.8
2000.....	4.8	*	*	*	0.6	1.5	1.3	1.4	1.8	1.7
1999.....	4.7	*	*	*	1.1	1.2	1.7	1.1	1.4	1.8

Black, male

2004.....	5.3	*	*	*	*	*	*	*	1.8	2.1
2003.....	5.0	*	*	*	*	*	1.3	*	1.8	2.5
2002.....	5.1	*	*	*	*	*	*	*	1.6	1.9
2001.....	5.2	*	*	*	1.5	*	*	1.7	1.8	2.3
2000.....	5.3	*	*	*	*	2.0	1.8	*	2.5	1.4
1999.....	5.2	*	*	*	1.5	1.5	1.7	*	1.7	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Leukemia (C91-C95)

Black, female

2004.....	4.4	*	*	*	*	*	*	*	1.5
2003.....	4.2	*	*	*	*	*	*	*	1.7
2002.....	4.3	*	*	*	*	*	*	*	*
2001.....	4.3	*	*	*	*	*	*	*	1.4
2000.....	4.2	*	*	*	*	*	1.5	*	1.9
1999.....	4.3	*	*	*	*	*	1.8	*	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Lymphoid leukemia (C91)

All races 3/, both sexes

2004.....	2.1	*	0.3	0.5	0.3	0.5	0.4	0.3	0.3	0.3
2003.....	2.2	*	0.4	0.4	0.4	0.4	0.5	0.3	0.3	0.4
2002.....	2.2	*	0.2	0.4	0.4	0.5	0.5	0.3	0.4	0.4
2001.....	2.2	*	0.3	0.4	0.4	0.5	0.5	0.4	0.4	0.4
2000.....	2.2	*	0.3	0.4	0.5	0.5	0.4	0.3	0.2	0.3
1999.....	2.2	*	0.3	0.3	0.4	0.5	0.5	0.4	0.3	0.3

All races 3/, male

2004.....	2.5	*	0.4	0.5	0.4	0.6	0.6	0.4	0.4	0.4
2003.....	2.6	*	0.4	0.5	0.6	0.6	0.6	0.4	0.4	0.4
2002.....	2.6	*	*	0.4	0.5	0.7	0.6	0.4	0.4	0.5
2001.....	2.5	*	0.3	0.4	0.5	0.7	0.7	0.4	0.5	0.5
2000.....	2.6	*	0.3	0.4	0.6	0.7	0.7	0.4	0.3	0.4
1999.....	2.6	*	0.3	0.4	0.4	0.6	0.7	0.4	0.4	0.4

All races 3/, female

2004.....	1.7	*	*	0.4	0.3	0.4	0.3	0.2	0.2	0.3
2003.....	1.8	*	0.4	0.3	0.3	0.3	0.3	0.3	*	0.3
2002.....	1.8	*	*	0.4	0.4	0.4	0.4	*	0.3	0.3
2001.....	1.8	*	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.3
2000.....	1.8	*	0.3	0.3	0.3	0.4	0.2	0.2	*	0.2
1999.....	1.9	*	0.3	0.2	0.3	0.3	0.4	0.3	0.2	0.3

White, both sexes

2004.....	2.3	*	0.3	0.5	0.4	0.5	0.4	0.3	0.3	0.3
2003.....	2.4	*	0.4	0.5	0.4	0.5	0.5	0.4	0.3	0.3
2002.....	2.4	*	0.2	0.4	0.5	0.5	0.6	0.3	0.4	0.4
2001.....	2.4	*	0.3	0.4	0.4	0.6	0.6	0.4	0.4	0.3
2000.....	2.4	*	0.4	0.4	0.5	0.5	0.5	0.3	0.2	0.3
1999.....	2.5	*	0.4	0.4	0.4	0.5	0.6	0.4	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Myeloid leukemia (C92)

All races 3/, both sexes

2004.....	3.4	*	0.4	0.2	0.3	0.4	0.5	0.5	0.6	0.8
2003.....	3.4	*	0.3	0.2	0.3	0.4	0.4	0.5	0.6	0.8
2002.....	3.4	*	0.3	0.2	0.2	0.4	0.4	0.5	0.7	0.8
2001.....	3.5	*	0.5	0.2	0.2	0.4	0.5	0.6	0.8	1.0
2000.....	3.5	*	0.4	0.2	0.3	0.5	0.5	0.6	0.8	1.1
1999.....	3.3	*	0.4	0.2	0.3	0.4	0.6	0.6	0.7	1.0

All races 3/, male

2004.....	3.8	*	0.4	0.3	0.2	0.4	0.5	0.4	0.7	0.8
2003.....	3.8	*	0.3	0.2	0.3	0.5	0.4	0.6	0.7	0.8
2002.....	3.9	*	0.4	*	0.2	0.4	0.4	0.6	0.8	0.9
2001.....	3.9	*	0.5	0.2	0.2	0.4	0.5	0.7	0.8	1.0
2000.....	3.9	*	0.4	*	0.3	0.5	0.6	0.6	0.9	0.9
1999.....	3.7	*	0.4	*	0.3	0.4	0.7	0.6	0.7	1.0

All races 3/, female

2004.....	3.0	*	0.3	*	0.3	0.3	0.5	0.5	0.6	0.7
2003.....	3.0	*	*	*	0.4	0.3	0.4	0.4	0.5	0.8
2002.....	3.0	*	*	0.2	0.2	0.4	0.4	0.4	0.6	0.8
2001.....	3.1	*	0.5	*	0.3	0.4	0.5	0.5	0.7	0.9
2000.....	3.1	*	0.3	*	0.2	0.4	0.4	0.6	0.6	1.2
1999.....	3.0	*	0.3	*	0.3	0.4	0.5	0.6	0.6	0.9

White, both sexes

2004.....	3.7	*	0.3	0.2	0.2	0.3	0.5	0.5	0.6	0.7
2003.....	3.7	*	0.3	0.2	0.3	0.3	0.4	0.6	0.6	0.8
2002.....	3.8	*	0.3	0.2	0.2	0.4	0.4	0.5	0.7	0.8
2001.....	3.8	*	0.5	0.2	0.2	0.4	0.5	0.6	0.8	0.9
2000.....	3.8	*	0.4	0.2	0.3	0.4	0.5	0.5	0.7	1.1
1999.....	3.6	*	0.4	0.1	0.3	0.3	0.6	0.6	0.7	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified leukemia (C94-C95)

All races 3/, both sexes

2004.....	1.8	*	0.1	0.1	0.1	0.2	0.2	0.1	0.2	0.2
2003.....	1.8	*	0.2	0.1	0.1	0.1	0.2	0.2	0.2	0.3
2002.....	1.8	*	*	0.2	0.1	0.2	0.2	0.2	0.2	0.2
2001.....	1.8	*	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2
2000.....	1.9	*	0.2	0.1	0.1	0.2	0.2	0.3	0.2	0.3
1999.....	1.9	*	*	0.2	0.2	0.2	0.3	0.2	0.3	0.3

All races 3/, male

2004.....	2.0	*	*	*	*	0.2	0.2	*	0.3	0.3
2003.....	1.9	*	*	*	*	*	0.2	0.2	0.2	0.3
2002.....	2.0	*	*	*	*	0.3	0.3	0.2	0.3	0.2
2001.....	2.0	*	0.3	*	0.2	*	0.3	0.3	0.3	0.3
2000.....	2.0	*	*	*	*	0.2	0.2	0.4	0.2	0.3
1999.....	2.1	*	*	0.2	*	0.3	0.3	0.2	0.3	0.3

All races 3/, female

2004.....	1.6	*	*	*	*	*	*	*	*	0.2
2003.....	1.6	*	*	*	*	*	*	*	*	0.3
2002.....	1.6	*	*	*	*	*	*	*	0.2	0.2
2001.....	1.6	*	*	*	*	0.2	*	*	*	0.2
2000.....	1.7	*	*	*	*	0.2	0.2	0.2	*	0.2
1999.....	1.8	*	*	*	*	0.2	*	*	0.3	0.3

White, both sexes

2004.....	1.9	*	*	0.1	0.1	0.2	0.2	*	0.2	0.2
2003.....	2.0	*	0.2	0.1	0.1	0.1	0.2	0.2	0.2	0.3
2002.....	2.0	*	*	0.1	*	0.2	0.2	0.2	0.2	0.2
2001.....	2.0	*	0.3	0.1	0.2	0.2	0.2	0.2	0.2	0.3
2000.....	2.1	*	0.2	0.1	*	0.2	0.2	0.3	0.2	0.3
1999.....	2.1	*	*	0.2	0.1	0.3	0.3	0.2	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

All races 3/, both sexes

2004.....	3.8	0.8	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.4
2003.....	3.9	0.9	0.2	0.2	0.1	0.1	0.2	0.2	0.4	0.5
2002.....	3.8	0.9	0.3	0.2	0.2	0.1	0.2	0.2	0.3	0.4
2001.....	4.0	0.5	0.3	0.2	0.2	0.1	0.2	0.2	0.3	0.5
2000.....	4.0	0.7	0.3	0.3	0.2	0.1	0.2	0.2	0.3	0.5
1999.....	4.0	0.8	0.4	0.3	0.2	0.2	0.2	0.2	0.4	0.4

All races 3/, male

2004.....	4.1	*	0.3	*	*	*	0.2	0.3	0.4	0.4
2003.....	4.1	*	0.3	*	*	*	*	0.2	0.4	0.6
2002.....	4.0	*	0.3	*	0.2	*	0.2	0.2	0.3	0.4
2001.....	4.2	*	0.3	0.2	0.2	*	*	0.2	0.3	0.6
2000.....	4.2	*	0.4	0.3	*	*	0.3	0.3	0.3	0.6
1999.....	4.2	*	0.4	0.2	*	*	0.3	0.2	0.5	0.5

All races 3/, female

2004.....	3.6	*	*	*	*	*	*	*	0.3	0.4
2003.....	3.8	*	*	*	*	*	*	*	0.3	0.3
2002.....	3.7	*	0.3	*	*	*	*	*	0.3	0.3
2001.....	3.9	*	0.3	0.2	0.2	*	*	*	0.3	0.5
2000.....	3.9	*	*	0.2	*	*	*	0.2	0.3	0.4
1999.....	3.8	*	0.3	0.3	*	0.2	*	*	0.3	0.4

White, both sexes

2004.....	4.3	0.8	0.3	0.2	0.1	0.1	0.2	0.3	0.3	0.4
2003.....	4.4	0.8	0.2	0.1	0.1	0.1	0.2	0.2	0.4	0.4
2002.....	4.3	1.0	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.4
2001.....	4.5	*	0.3	0.2	0.2	0.1	0.1	0.2	0.3	0.5
2000.....	4.5	0.7	0.3	0.3	0.2	*	0.2	0.2	0.4	0.5
1999.....	4.4	0.7	0.3	0.3	0.2	0.2	0.2	0.2	0.4	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

All races 3/, both sexes

2004.....	3.2	2.3	0.4	0.3	0.2	0.4	0.5	0.7	0.8	1.1
2003.....	3.3	2.6	0.5	0.2	0.3	0.3	0.5	0.6	0.9	1.2
2002.....	3.4	2.0	0.5	0.2	0.2	0.4	0.5	0.7	0.9	1.2
2001.....	3.3	2.6	0.5	0.2	0.2	0.4	0.5	0.6	1.1	1.2
2000.....	3.3	1.8	0.5	0.2	0.2	0.3	0.5	0.6	0.9	1.3
1999.....	3.2	2.5	0.5	0.2	0.2	0.4	0.6	0.6	1.0	1.2

All races 3/, male

2004.....	2.8	2.8	0.5	0.2	0.2	0.4	0.5	0.7	0.9	1.2
2003.....	2.9	2.9	0.5	*	0.3	0.3	0.6	0.6	0.9	1.2
2002.....	2.9	2.5	0.5	*	0.2	0.4	0.6	0.9	1.0	1.2
2001.....	2.8	2.9	0.6	0.2	0.3	0.4	0.5	0.7	1.2	1.3
2000.....	2.8	2.1	0.5	0.3	0.2	0.3	0.5	0.6	0.9	1.2
1999.....	2.8	3.0	0.6	0.2	0.3	0.4	0.7	0.7	1.1	1.4

All races 3/, female

2004.....	3.6	1.8	0.4	0.3	0.2	0.4	0.5	0.8	0.7	1.0
2003.....	3.7	2.3	0.4	*	0.3	0.3	0.5	0.7	0.9	1.2
2002.....	3.8	1.5	0.5	0.2	*	0.3	0.5	0.6	0.8	1.2
2001.....	3.9	2.4	0.3	0.2	*	0.4	0.5	0.5	1.0	1.1
2000.....	3.8	1.6	0.6	*	0.2	0.3	0.5	0.6	1.0	1.3
1999.....	3.7	1.9	0.4	0.3	*	0.4	0.6	0.5	1.0	1.1

White, both sexes

2004.....	3.0	2.0	0.3	0.2	0.2	0.3	0.2	0.4	0.3	0.5
2003.....	3.0	2.3	0.4	*	0.2	0.2	0.3	0.3	0.4	0.6
2002.....	3.1	1.9	0.4	0.2	*	0.2	0.3	0.3	0.5	0.6
2001.....	3.1	2.3	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.5
2000.....	3.1	1.7	0.4	0.1	0.2	0.2	0.2	0.2	0.4	0.6
1999.....	3.0	1.8	0.4	0.2	0.2	0.3	0.2	0.2	0.4	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

White, male

2004.....	2.6	2.2	0.3	*	*	0.3	0.3	0.4	0.3	0.6
2003.....	2.6	2.8	0.3	*	0.2	*	0.3	*	0.5	0.7
2002.....	2.6	2.5	0.5	*	*	0.3	0.3	0.4	0.6	0.7
2001.....	2.6	2.4	0.6	*	0.3	*	0.2	0.4	0.5	0.6
2000.....	2.5	1.7	0.4	*	*	*	*	*	0.4	0.6
1999.....	2.5	2.2	0.5	*	*	0.3	*	0.3	0.4	0.7

White, female

2004.....	3.3	1.7	*	*	*	0.3	*	0.4	0.3	0.4
2003.....	3.5	1.7	0.4	*	*	*	*	0.3	0.4	0.5
2002.....	3.6	*	0.4	*	*	*	0.3	0.3	0.4	0.5
2001.....	3.6	2.1	*	*	*	*	*	*	0.5	0.4
2000.....	3.6	1.7	0.4	*	*	*	*	*	0.3	0.6
1999.....	3.4	1.4	*	*	*	0.3	*	*	0.4	0.4

Black, both sexes

2004.....	5.7	3.5	0.9	*	*	1.1	2.2	2.9	4.1	5.1
2003.....	6.0	3.0	0.9	*	0.6	0.7	2.3	2.8	4.0	5.2
2002.....	6.0	*	0.8	*	*	0.9	2.1	3.2	3.8	5.5
2001.....	5.9	3.5	*	*	*	1.1	2.0	2.4	4.7	5.5
2000.....	5.8	*	1.2	0.6	0.6	0.7	2.1	3.0	4.7	5.7
1999.....	5.7	5.5	1.4	0.6	0.6	0.9	2.8	2.9	5.1	5.8

Black, male

2004.....	5.3	*	*	*	*	1.4	2.0	3.0	4.7	5.6
2003.....	5.6	*	*	*	*	*	2.2	2.7	4.1	5.3
2002.....	5.5	*	*	*	*	*	2.7	4.1	4.3	5.2
2001.....	5.2	*	*	*	*	1.3	2.2	2.9	5.6	5.9
2000.....	5.1	*	*	*	*	*	2.1	3.0	4.4	5.8
1999.....	5.3	6.5	*	*	*	*	3.3	3.3	5.4	6.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ years	year 2/ years	years	years	years	years	years	years	years	years

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

Black, female

2004.....	6.0	*	*	*	*	*	2.4	2.9	3.5	4.7
2003.....	6.3	*	*	*	*	*	2.4	2.9	3.9	5.2
2002.....	6.4	*	*	*	*	*	1.6	2.3	3.3	5.7
2001.....	6.4	*	*	*	*	*	1.8	1.9	3.8	5.0
2000.....	6.4	*	*	*	*	*	2.2	3.0	5.0	5.6
1999.....	6.2	*	*	*	*	*	2.3	2.5	4.7	5.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Anemias (D50-D64)

White, male

2004.....	1.1	*	*	*	*	*	*	*	*	*
2003.....	1.1	*	*	*	*	*	*	*	*	*
2002.....	1.1	*	*	*	*	*	*	*	*	*
2001.....	1.1	*	*	*	*	*	*	*	*	*
2000.....	1.1	*	*	*	*	*	*	*	*	*
1999.....	1.1	*	*	*	*	*	*	*	*	*

White, female

2004.....	1.8	*	*	*	*	*	*	*	*	*
2003.....	1.8	*	*	*	*	*	*	*	*	*
2002.....	1.8	*	*	*	*	*	*	*	*	*
2001.....	1.9	*	*	*	*	*	*	*	*	*
2000.....	1.9	*	*	*	*	*	*	*	*	*
1999.....	1.9	*	*	*	*	*	*	*	*	*

Black, both sexes

2004.....	2.7	*	*	*	*	0.9	2.0	2.2	2.4	2.4
2003.....	2.7	*	*	*	*	*	1.9	1.9	2.2	2.2
2002.....	2.8	*	*	*	*	0.9	1.7	2.5	2.3	2.6
2001.....	2.7	*	*	*	*	0.9	1.5	1.9	2.5	2.1
2000.....	2.6	*	*	*	*	*	1.6	1.7	2.8	2.2
1999.....	2.8	*	0.8	*	*	0.7	2.2	2.1	3.0	2.2

Black, male

2004.....	2.6	*	*	*	*	1.2	1.9	2.3	3.0	3.0
2003.....	2.7	*	*	*	*	*	1.8	2.2	2.3	2.2
2002.....	2.7	*	*	*	*	*	2.3	3.2	2.5	2.7
2001.....	2.5	*	*	*	*	*	1.6	2.5	3.3	2.2
2000.....	2.3	*	*	*	*	*	1.7	1.6	2.7	1.9
1999.....	2.7	*	*	*	*	*	2.8	2.3	3.3	2.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Anemias (D50-D64)

Black, female

2004.....	2.8	*	*	*	*	*	2.1	2.2	1.9	1.9
2003.....	2.8	*	*	*	*	*	2.1	1.7	2.2	2.2
2002.....	3.0	*	*	*	*	*	*	1.8	2.2	2.4
2001.....	2.8	*	*	*	*	*	1.5	*	1.8	2.1
2000.....	3.0	*	*	*	*	*	1.6	1.9	2.9	2.5
1999.....	3.0	*	*	*	*	*	1.7	1.8	2.8	2.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Hemolytic anemias (D55-D59)

White, male

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, female

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*

Black, both sexes

2004.....	1.5	*	*	*	*	0.8	1.9	2.2	2.3	2.1
2003.....	1.3	*	*	*	*	*	1.7	1.9	2.0	1.9
2002.....	1.4	*	*	*	*	0.7	1.6	2.2	2.3	2.3
2001.....	1.4	*	*	*	*	0.7	1.4	1.6	2.3	1.8
2000.....	1.4	*	*	*	*	*	1.5	1.6	2.4	2.0
1999.....	1.5	*	0.8	*	*	0.6	1.9	1.9	2.8	2.1

Black, male

2004.....	1.5	*	*	*	*	*	1.8	2.2	3.0	2.5
2003.....	1.4	*	*	*	*	*	1.5	2.1	2.0	1.9
2002.....	1.4	*	*	*	*	*	2.0	2.9	2.4	2.5
2001.....	1.4	*	*	*	*	*	1.5	2.0	3.1	1.9
2000.....	1.3	*	*	*	*	*	1.5	1.6	2.1	1.9
1999.....	1.7	*	*	*	*	*	2.6	2.2	3.1	2.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Hemolytic anemias (D55-D59)

Black, female

2004.....	1.4	*	*	*	*	*	2.0	2.1	1.8	1.8
2003.....	1.3	*	*	*	*	*	2.0	1.7	2.0	2.0
2002.....	1.4	*	*	*	*	*	*	1.7	2.2	2.1
2001.....	1.3	*	*	*	*	*	*	*	1.5	1.8
2000.....	1.4	*	*	*	*	*	1.5	1.6	2.6	2.0
1999.....	1.3	*	*	*	*	*	*	1.8	2.5	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)

All races 3/, both sexes

2004.....	0.9	0.9	*	*	*	*	*	0.1	0.2	0.3
2003.....	0.9	1.0	*	*	*	*	0.1	0.1	0.2	0.2
2002.....	0.9	0.6	*	*	*	*	0.1	0.1	0.2	0.2
2001.....	0.9	0.8	*	*	*	*	0.1	0.2	0.3	
2000.....	0.9	0.6	*	*	*	*	0.1	0.1	0.1	0.3
1999.....	0.9	1.1	*	*	*	*	0.1	0.1	0.2	0.3

All races 3/, male

2004.....	0.8	1.2	*	*	*	*	*	*	*	0.3
2003.....	0.8	1.1	*	*	*	*	*	*	*	0.2
2002.....	0.8	*	*	*	*	*	*	*	0.3	0.2
2001.....	0.8	*	*	*	*	*	*	*	*	0.3
2000.....	0.8	*	*	*	*	*	*	*	*	0.2
1999.....	0.8	1.4	*	*	*	*	*	*	*	0.3

All races 3/, female

2004.....	1.0	*	*	*	*	*	*	*	0.2	0.3
2003.....	1.0	*	*	*	*	*	*	*	*	0.3
2002.....	1.0	*	*	*	*	*	*	*	*	0.3
2001.....	1.0	*	*	*	*	*	*	*	0.4	0.2
2000.....	1.0	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.0	*	*	*	*	*	*	*	0.2	0.3

White, both sexes

2004.....	0.9	0.8	*	*	*	*	*	0.1	0.1	0.2
2003.....	0.9	0.9	*	*	*	*	0.1	*	0.1	0.2
2002.....	1.0	*	*	*	*	*	*	0.1	0.2	0.2
2001.....	0.9	0.7	*	*	*	*	*	*	0.2	0.2
2000.....	0.9	*	*	*	*	*	*	*	*	0.3
1999.....	0.9	0.9	*	*	*	*	*	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Certain disorders involving the immune mechanism (D80-D89)

All races 3/, both sexes

2004.....	0.5	*	*	*	*	*	*	0.1	0.2	0.4
2003.....	0.5	0.5	*	*	*	*	*	0.1	0.3	0.5
2002.....	0.5	0.5	*	*	*	*	*	*	0.2	0.5
2001.....	0.5	0.8	*	*	*	*	*	*	0.3	0.5
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	0.5
1999.....	0.4	*	0.2	*	*	*	*	0.1	0.3	0.5

All races 3/, male

2004.....	0.4	*	*	*	*	*	*	*	0.2	0.4
2003.....	0.5	*	*	*	*	*	*	*	0.3	0.6
2002.....	0.5	*	*	*	*	*	*	*	0.3	0.5
2001.....	0.4	*	*	*	*	*	*	*	0.4	0.6
2000.....	0.5	*	*	*	*	*	*	*	0.3	0.6
1999.....	0.4	*	*	*	*	*	*	*	0.4	0.5

All races 3/, female

2004.....	0.5	*	*	*	*	*	*	*	*	0.4
2003.....	0.6	*	*	*	*	*	*	*	0.3	0.4
2002.....	0.6	*	*	*	*	*	*	*	*	0.5
2001.....	0.6	*	*	*	*	*	*	*	*	0.4
2000.....	0.5	*	*	*	*	*	*	*	0.2	0.5
1999.....	0.4	*	*	*	*	*	*	*	0.3	0.4

White, both sexes

2004.....	0.3	*	*	*	*	*	*	*	*	0.1
2003.....	0.3	*	*	*	*	*	*	*	*	0.2
2002.....	0.3	*	*	*	*	*	*	*	*	0.2
2001.....	0.3	0.8	*	*	*	*	*	*	0.2	0.2
2000.....	0.3	*	*	*	*	*	*	*	0.1	0.2
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Certain disorders involving the immune mechanism (D80-D89)

White, male

2004.....	0.3	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	0.3
2002.....	0.3	*	*	*	*	*	*	*	0.3
2001.....	0.3	*	*	*	*	*	*	0.2	0.3
2000.....	0.3	*	*	*	*	*	*	*	0.2
1999.....	0.3	*	*	*	*	*	*	*	0.2

White, female

2004.....	0.3	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*

Black, both sexes

2004.....	1.7	*	*	*	*	*	*	*	1.2	2.0
2003.....	1.9	*	*	*	*	*	*	*	1.3	2.6
2002.....	1.9	*	*	*	*	*	*	*	1.0	2.5
2001.....	1.8	*	*	*	*	*	*	*	1.4	2.6
2000.....	1.8	*	*	*	*	*	*	0.7	1.3	2.8
1999.....	1.5	*	*	*	*	*	*	*	1.5	2.5

Black, male

2004.....	1.6	*	*	*	*	*	*	*	*	1.9
2003.....	1.7	*	*	*	*	*	*	*	*	3.1
2002.....	1.7	*	*	*	*	*	*	*	*	2.2
2001.....	1.6	*	*	*	*	*	*	*	1.9	3.0
2000.....	1.6	*	*	*	*	*	*	*	*	3.2
1999.....	1.3	*	*	*	*	*	*	*	1.7	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

IV. Endocrine, nutritional and metabolic diseases (E00-E88)

All races 3/, both sexes

2004.....	33.7	5.7	0.9	0.5	0.6	0.9	1.6	2.2	3.6	5.5
2003.....	34.5	6.2	1.2	0.5	0.6	1.0	1.6	2.6	3.6	5.5
2002.....	34.5	6.6	0.9	0.5	0.6	0.9	1.7	2.5	3.7	5.6
2001.....	34.0	7.2	1.2	0.5	0.6	1.0	1.5	2.3	3.2	5.3
2000.....	33.5	6.8	1.0	0.4	0.7	1.0	1.6	2.3	3.4	4.9
1999.....	33.3	6.9	1.0	0.5	0.6	1.1	1.6	2.2	3.1	4.9

All races 3/, male

2004.....	32.2	7.1	1.0	0.5	0.6	0.9	1.7	2.4	3.9	6.4
2003.....	32.5	6.8	1.4	0.6	0.6	1.1	1.7	2.8	4.3	6.0
2002.....	31.9	7.4	0.8	0.6	0.7	0.9	1.9	2.7	4.4	6.3
2001.....	30.9	8.0	1.2	0.4	0.7	1.0	1.5	2.4	3.6	6.1
2000.....	30.2	7.4	1.1	0.4	0.7	1.0	1.7	2.6	4.1	5.6
1999.....	29.8	7.3	1.1	0.5	0.6	1.1	1.5	2.2	3.6	5.8

All races 3/, female

2004.....	35.2	4.2	0.8	0.5	0.5	0.8	1.6	2.0	3.3	4.5
2003.....	36.5	5.5	1.1	0.3	0.5	0.9	1.5	2.4	2.9	4.9
2002.....	37.0	5.8	1.0	0.3	0.5	0.9	1.6	2.3	2.9	4.8
2001.....	37.0	6.4	1.2	0.5	0.5	1.0	1.6	2.2	2.8	4.5
2000.....	36.7	6.2	1.0	0.3	0.6	1.0	1.4	2.0	2.7	4.3
1999.....	36.6	6.5	0.9	0.5	0.6	1.1	1.7	2.1	2.7	4.1

White, both sexes

2004.....	33.7	5.8	0.9	0.5	0.6	0.8	1.6	2.0	3.2	4.7
2003.....	34.5	5.4	1.2	0.5	0.6	1.0	1.6	2.2	3.2	4.8
2002.....	34.4	6.2	1.0	0.5	0.6	0.9	1.6	2.3	3.2	5.0
2001.....	34.0	6.8	1.2	0.5	0.6	0.9	1.4	2.0	2.8	4.7
2000.....	33.5	6.3	1.1	0.4	0.7	1.0	1.5	1.9	3.0	4.4
1999.....	33.1	6.0	1.1	0.4	0.6	1.1	1.6	2.0	2.7	4.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

IV. Endocrine, nutritional and metabolic diseases (E00-E88)

White, male

2004.....	32.5	7.2	1.0	0.5	0.6	0.8	1.7	2.2	3.4	5.5
2003.....	32.9	6.3	1.4	0.7	0.7	1.1	1.7	2.3	3.9	5.5
2002.....	32.2	7.6	0.8	0.7	0.7	0.9	1.6	2.5	3.9	5.7
2001.....	31.2	7.2	1.3	0.5	0.7	1.0	1.4	2.0	3.2	5.3
2000.....	30.6	6.8	1.2	0.4	0.8	1.0	1.7	2.3	3.7	5.0
1999.....	29.9	6.3	1.2	0.5	0.6	1.2	1.4	2.1	3.0	5.1

White, female

2004.....	34.9	4.3	0.9	0.6	0.5	0.9	1.5	1.9	2.9	3.8
2003.....	36.1	4.5	1.1	0.4	0.5	0.9	1.5	2.2	2.5	4.1
2002.....	36.6	4.7	1.1	0.3	0.5	1.0	1.5	2.1	2.6	4.2
2001.....	36.8	6.3	1.2	0.5	0.5	0.9	1.5	2.0	2.5	4.0
2000.....	36.3	5.7	1.0	0.4	0.6	1.0	1.4	1.6	2.3	3.8
1999.....	36.2	5.7	0.9	0.4	0.6	1.1	1.7	2.0	2.4	3.4

Black, both sexes

2004.....	41.8	5.9	*	*	0.6	1.1	2.4	4.2	7.7	12.2
2003.....	42.3	9.9	1.0	*	*	1.2	2.2	5.5	7.1	11.0
2002.....	42.6	8.3	0.8	*	0.7	1.0	2.9	4.9	7.5	10.7
2001.....	41.6	10.3	0.8	*	0.6	1.1	2.5	4.5	6.4	10.5
2000.....	41.4	8.9	0.8	*	*	1.1	2.5	5.3	6.8	9.6
1999.....	41.4	11.4	0.8	0.6	0.6	1.1	2.2	3.6	6.4	10.2

Black, male

2004.....	37.7	7.7	*	*	*	1.4	2.2	4.8	8.8	14.2
2003.....	37.1	8.9	*	*	*	*	2.1	6.7	8.5	11.3
2002.....	36.6	7.0	*	*	*	1.2	3.5	5.5	9.3	12.0
2001.....	35.9	11.7	*	*	*	*	2.4	5.2	7.8	12.8
2000.....	34.6	9.9	*	*	*	*	2.7	5.5	8.0	10.7
1999.....	35.2	12.7	*	*	*	*	2.3	3.4	7.8	11.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

IV. Endocrine, nutritional and metabolic diseases (E00-E88)

Black, female

2004.....	45.5	*	*	*	*	2.5	3.6	6.7	10.3
2003.....	47.0	10.8	*	*	*	1.3	4.4	5.9	10.6
2002.....	48.0	9.7	*	*	*	2.4	4.3	5.8	9.6
2001.....	46.8	8.8	*	*	*	2.5	3.9	5.2	8.6
2000.....	47.6	7.9	*	*	*	2.2	5.1	5.7	8.6
1999.....	47.1	10.1	*	*	*	2.1	3.9	5.1	8.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Endocrine diseases (E00-E34)

All races 3/, both sexes

2004.....	25.8	0.6	0.1	*	0.2	0.3	0.7	1.1	2.2	3.5
2003.....	26.5	0.7	0.1	*	0.1	0.3	0.7	1.3	2.2	3.5
2002.....	26.4	1.0	*	*	0.1	0.3	0.7	1.3	2.1	3.7
2001.....	26.1	1.1	*	*	0.2	0.3	0.6	1.2	2.0	3.5
2000.....	25.6	1.2	0.1	*	0.1	0.3	0.6	1.2	2.2	3.2
1999.....	25.5	1.0	0.1	*	0.1	0.3	0.7	1.1	2.0	3.4

All races 3/, male

2004.....	25.0	*	*	*	0.2	0.3	0.6	1.3	2.4	4.4
2003.....	25.3	*	*	*	*	0.3	0.8	1.4	2.6	4.0
2002.....	24.8	1.1	*	*	0.2	0.3	0.8	1.4	2.6	4.2
2001.....	24.1	1.3	*	*	0.2	0.3	0.7	1.2	2.3	4.1
2000.....	23.4	1.3	*	*	*	0.3	0.7	1.3	2.7	3.7
1999.....	23.3	*	*	*	*	0.3	0.7	1.1	2.3	4.0

All races 3/, female

2004.....	26.7	*	*	*	*	0.2	0.7	0.9	1.9	2.7
2003.....	27.7	*	*	*	*	0.2	0.6	1.3	1.8	2.9
2002.....	28.0	*	*	*	*	0.3	0.6	1.2	1.7	3.1
2001.....	28.0	*	*	*	*	0.3	0.5	1.2	1.6	2.9
2000.....	27.7	1.1	*	*	*	0.2	0.6	1.1	1.7	2.8
1999.....	27.5	1.3	*	*	*	0.3	0.7	1.1	1.8	2.8

White, both sexes

2004.....	25.3	0.7	*	*	*	0.2	0.5	0.9	1.8	3.0
2003.....	26.1	*	*	*	*	0.2	0.6	1.0	1.9	3.0
2002.....	26.0	1.1	*	*	*	0.3	0.6	1.1	1.9	3.2
2001.....	25.7	1.1	*	*	0.1	0.2	0.4	0.9	1.7	3.0
2000.....	25.1	1.0	*	*	*	0.3	0.5	0.9	1.9	2.8
1999.....	24.9	1.0	*	*	*	0.3	0.6	1.0	1.7	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Endocrine diseases (E00-E34)

White, male

2004.....	24.9	*	*	*	*	*	0.6	1.0	2.0	3.7
2003.....	25.3	*	*	*	*	0.3	0.7	0.9	2.3	3.4
2002.....	24.8	1.3	*	*	*	0.3	0.6	1.2	2.2	3.7
2001.....	24.1	*	*	*	*	0.3	0.5	0.9	1.9	3.5
2000.....	23.5	*	*	*	*	0.3	0.6	1.0	2.4	3.2
1999.....	23.2	*	*	*	*	0.3	0.6	1.0	1.9	3.5

White, female

2004.....	25.8	*	*	*	*	*	0.5	0.8	1.6	2.3
2003.....	26.8	*	*	*	*	*	0.5	1.1	1.5	2.5
2002.....	27.1	*	*	*	*	0.3	0.5	1.0	1.5	2.7
2001.....	27.2	*	*	*	*	*	0.4	0.9	1.5	2.5
2000.....	26.8	*	*	*	*	*	0.4	0.8	1.3	2.5
1999.....	26.5	1.4	*	*	*	0.3	0.5	1.0	1.5	2.3

Black, both sexes

2004.....	34.2	*	*	*	*	0.7	1.5	2.3	4.9	8.0
2003.....	34.7	*	*	*	*	*	1.3	3.5	4.7	7.5
2002.....	34.5	*	*	*	*	*	1.6	3.1	4.6	7.6
2001.....	34.0	*	*	*	*	0.6	1.4	3.4	4.4	7.3
2000.....	33.8	*	*	*	*	*	1.5	3.3	4.5	6.7
1999.....	34.0	*	*	*	*	*	1.2	2.1	4.4	7.1

Black, male

2004.....	31.0	*	*	*	*	*	1.3	2.8	5.8	9.9
2003.....	30.5	*	*	*	*	*	1.4	4.4	5.4	8.7
2002.....	29.5	*	*	*	*	*	1.8	3.3	6.0	8.7
2001.....	29.2	*	*	*	*	*	1.5	4.0	5.8	9.3
2000.....	28.1	*	*	*	*	*	1.5	3.5	5.0	7.6
1999.....	28.5	*	*	*	*	*	*	1.8	5.2	8.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Endocrine diseases (E00-E34)

Black, female

2004.....	37.2	*	*	*	*	*	1.8	1.8	4.2	6.2
2003.....	38.6	*	*	*	*	*	*	2.7	4.1	6.5
2002.....	39.0	*	*	*	*	*	1.4	2.8	3.4	6.6
2001.....	38.3	*	*	*	*	*	1.3	2.8	3.1	5.5
2000.....	39.1	*	*	*	*	*	1.5	3.2	4.1	5.8
1999.....	39.0	*	*	*	*	*	*	2.5	3.8	6.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Diabetes mellitus (E10-E14)

All races 3/, both sexes

2004.....	24.9	*	*	*	0.1	0.2	0.5	0.9	2.0	3.4
2003.....	25.5	*	*	*	0.1	0.2	0.5	1.2	2.1	3.3
2002.....	25.4	*	*	*	0.1	0.3	0.6	1.2	2.0	3.5
2001.....	25.1	*	*	*	0.2	0.2	0.5	1.1	1.8	3.3
2000.....	24.6	*	*	*	*	0.2	0.6	1.1	2.0	3.0
1999.....	24.5	*	*	*	0.1	0.3	0.6	1.0	1.9	3.2

All races 3/, male

2004.....	24.4	*	*	*	*	0.3	0.5	1.0	2.3	4.2
2003.....	24.8	*	*	*	*	0.3	0.6	1.2	2.5	3.9
2002.....	24.2	*	*	*	*	0.3	0.7	1.3	2.4	4.1
2001.....	23.5	*	*	*	0.2	0.3	0.6	1.2	2.2	3.9
2000.....	22.9	*	*	*	*	0.3	0.7	1.2	2.5	3.5
1999.....	22.8	*	*	*	*	0.2	0.6	1.0	2.2	3.8

All races 3/, female

2004.....	25.4	*	*	*	*	*	0.6	0.8	1.8	2.5
2003.....	26.2	*	*	*	*	*	0.5	1.1	1.7	2.8
2002.....	26.5	*	*	*	*	0.3	0.5	1.1	1.5	3.0
2001.....	26.6	*	*	*	*	0.2	0.5	1.1	1.5	2.7
2000.....	26.3	*	*	*	*	*	0.5	0.9	1.6	2.5
1999.....	26.2	*	*	*	*	0.3	0.5	1.0	1.6	2.6

White, both sexes

2004.....	24.4	*	*	*	*	0.2	0.4	0.8	1.7	2.9
2003.....	25.0	*	*	*	*	0.2	0.5	0.9	1.8	2.8
2002.....	24.9	*	*	*	*	0.2	0.5	1.0	1.7	3.1
2001.....	24.6	*	*	*	*	0.2	0.4	0.8	1.6	2.8
2000.....	24.1	*	*	*	*	0.2	0.5	0.8	1.8	2.7
1999.....	23.9	*	*	*	*	0.2	0.5	0.9	1.6	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Diabetes mellitus (E10-E14)

White, male

2004.....	24.3	*	*	*	*	*	0.4	0.8	2.0	3.6
2003.....	24.8	*	*	*	*	*	0.6	0.8	2.2	3.4
2002.....	24.2	*	*	*	*	*	0.5	1.1	2.1	3.7
2001.....	23.5	*	*	*	*	*	0.4	0.8	1.8	3.3
2000.....	22.9	*	*	*	*	0.3	0.5	1.0	2.3	3.1
1999.....	22.7	*	*	*	*	0.3	0.5	0.9	1.8	3.4

White, female

2004.....	24.5	*	*	*	*	*	0.4	0.7	1.5	2.1
2003.....	25.2	*	*	*	*	*	0.4	1.0	1.4	2.3
2002.....	25.6	*	*	*	*	*	0.4	0.9	1.3	2.5
2001.....	25.7	*	*	*	*	*	0.3	0.8	1.4	2.3
2000.....	25.3	*	*	*	*	*	0.4	0.7	1.2	2.3
1999.....	25.0	*	*	*	*	*	0.4	0.9	1.4	2.1

Black, both sexes

2004.....	33.2	*	*	*	*	0.6	1.3	2.0	4.6	7.5
2003.....	33.8	*	*	*	*	*	1.0	3.2	4.4	7.3
2002.....	33.6	*	*	*	*	*	1.4	2.8	4.3	7.3
2001.....	33.1	*	*	*	*	*	1.4	3.2	4.2	6.9
2000.....	32.8	*	*	*	*	*	1.5	2.8	4.1	6.0
1999.....	33.0	*	*	*	*	*	1.1	1.8	4.1	6.6

Black, male

2004.....	30.2	*	*	*	*	*	*	2.5	5.4	9.5
2003.....	29.8	*	*	*	*	*	*	4.1	5.2	8.4
2002.....	29.0	*	*	*	*	*	1.7	3.0	5.7	8.5
2001.....	28.5	*	*	*	*	*	1.5	3.9	5.6	8.7
2000.....	27.4	*	*	*	*	*	1.5	3.1	4.5	7.1
1999.....	27.7	*	*	*	*	*	*	1.6	5.0	7.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Diabetes mellitus (E10-E14)

Black, female

2004.....	36.0	*	*	*	*	*	1.6	1.6	3.8	5.6
2003.....	37.4	*	*	*	*	*	*	2.3	3.7	6.3
2002.....	37.8	*	*	*	*	*	*	2.6	3.1	6.3
2001.....	37.2	*	*	*	*	*	1.3	2.7	2.9	5.3
2000.....	37.8	*	*	*	*	*	1.5	2.5	3.8	5.0
1999.....	37.8	*	*	*	*	*	*	2.0	3.3	5.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Nutritional diseases (E40-E68)

All races 3/, both sexes

2004.....	2.5	*	*	*	*	*	0.2	0.6	0.9	1.2
2003.....	2.5	*	*	*	*	0.1	0.3	0.6	0.8	1.3
2002.....	2.6	*	*	*	*	*	0.3	0.6	0.9	1.2
2001.....	2.4	*	*	*	*	*	0.3	0.4	0.7	1.2
2000.....	2.6	*	*	*	*	*	0.2	0.5	0.7	1.1
1999.....	2.5	*	*	*	*	0.1	0.2	0.4	0.6	0.9

All races 3/, male

2004.....	2.1	*	*	*	*	*	0.3	0.6	0.9	1.3
2003.....	2.1	*	*	*	*	*	0.3	0.8	1.1	1.2
2002.....	2.2	*	*	*	*	*	0.4	0.6	1.1	1.4
2001.....	1.9	*	*	*	*	*	0.3	0.6	0.8	1.2
2000.....	2.0	*	*	*	*	*	0.2	0.6	0.9	1.2
1999.....	2.0	*	*	*	*	*	*	0.4	0.8	1.1

All races 3/, female

2004.....	2.8	*	*	*	*	*	*	0.6	0.9	1.2
2003.....	2.9	*	*	*	*	*	0.2	0.5	0.6	1.3
2002.....	3.0	*	*	*	*	*	0.3	0.5	0.8	1.1
2001.....	2.9	*	*	*	*	*	0.3	0.3	0.7	1.2
2000.....	3.1	*	*	*	*	*	*	0.4	0.5	0.9
1999.....	2.9	*	*	*	*	*	0.3	0.4	0.4	0.8

White, both sexes

2004.....	2.4	*	*	*	*	*	0.2	0.5	0.7	1.0
2003.....	2.6	*	*	*	*	*	0.2	0.5	0.7	1.1
2002.....	2.6	*	*	*	*	*	0.2	0.5	0.7	1.1
2001.....	2.4	*	*	*	*	*	0.2	0.4	0.6	1.0
2000.....	2.6	*	*	*	*	*	0.1	0.3	0.5	0.9
1999.....	2.5	*	*	*	*	*	0.2	0.3	0.5	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Nutritional diseases (E40-E68)

White, male

2004.....	2.1	*	*	*	*	*	*	0.5	0.7	1.1
2003.....	2.2	*	*	*	*	*	0.3	0.6	1.0	1.2
2002.....	2.2	*	*	*	*	*	0.3	0.5	0.8	1.3
2001.....	1.9	*	*	*	*	*	*	0.5	0.7	1.0
2000.....	2.1	*	*	*	*	*	*	0.4	0.7	1.2
1999.....	2.0	*	*	*	*	*	*	0.3	0.7	0.9

White, female

2004.....	2.8	*	*	*	*	*	*	0.5	0.7	0.8
2003.....	3.0	*	*	*	*	*	*	0.4	0.4	1.0
2002.....	3.0	*	*	*	*	*	*	0.4	0.6	0.9
2001.....	2.9	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	3.1	*	*	*	*	*	*	*	0.3	0.7
1999.....	3.0	*	*	*	*	*	*	0.3	0.3	0.6

Black, both sexes

2004.....	3.3	*	*	*	*	*	*	1.6	2.2	3.3
2003.....	3.0	*	*	*	*	*	*	1.5	1.9	2.5
2002.....	3.3	*	*	*	*	*	0.7	1.4	2.4	2.3
2001.....	3.0	*	*	*	*	*	0.8	0.8	1.7	2.8
2000.....	3.1	*	*	*	*	*	*	1.3	1.6	2.1
1999.....	3.0	*	*	*	*	*	*	1.2	1.3	2.1

Black, male

2004.....	2.6	*	*	*	*	*	*	1.6	2.4	3.2
2003.....	2.4	*	*	*	*	*	*	1.8	2.3	1.9
2002.....	2.6	*	*	*	*	*	*	1.6	2.9	2.3
2001.....	2.5	*	*	*	*	*	*	*	1.7	2.8
2000.....	2.5	*	*	*	*	*	*	*	2.1	2.2
1999.....	2.6	*	*	*	*	*	*	*	1.8	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Obesity and other hyperalimntation (E65-E68)

All races 3/, both sexes

2004.....	1.4	*	*	*	*	0.2	0.6	0.9	1.2
2003.....	1.4	*	*	*	*	0.1	0.2	0.6	0.8
2002.....	1.3	*	*	*	*	0.3	0.5	0.9	1.1
2001.....	1.1	*	*	*	*	0.3	0.4	0.7	1.1
2000.....	1.1	*	*	*	*	0.2	0.4	0.6	0.9
1999.....	0.9	*	*	*	*	0.2	0.4	0.6	0.8

All races 3/, male

2004.....	1.4	*	*	*	*	0.2	0.6	0.9	1.2
2003.....	1.3	*	*	*	*	0.2	0.7	1.0	1.2
2002.....	1.2	*	*	*	*	0.3	0.5	1.1	1.3
2001.....	1.0	*	*	*	*	0.2	0.5	0.7	1.1
2000.....	1.0	*	*	*	*	0.2	0.5	0.8	1.1
1999.....	0.9	*	*	*	*	*	0.4	0.8	1.0

All races 3/, female

2004.....	1.5	*	*	*	*	*	0.6	0.9	1.1
2003.....	1.5	*	*	*	*	0.2	0.5	0.6	1.2
2002.....	1.3	*	*	*	*	0.2	0.5	0.7	0.9
2001.....	1.2	*	*	*	*	0.3	0.3	0.6	1.1
2000.....	1.1	*	*	*	*	*	0.3	0.4	0.7
1999.....	1.0	*	*	*	*	0.2	0.4	0.4	0.7

White, both sexes

2004.....	1.4	*	*	*	*	0.1	0.5	0.7	0.9
2003.....	1.4	*	*	*	*	0.2	0.5	0.7	1.0
2002.....	1.2	*	*	*	*	0.2	0.4	0.7	1.0
2001.....	1.0	*	*	*	*	0.2	0.4	0.6	0.9
2000.....	1.0	*	*	*	*	*	0.3	0.5	0.8
1999.....	0.9	*	*	*	*	0.1	0.3	0.5	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Obesity and other hyperalimentation (E65-E68)

White, male

2004.....	1.4	*	*	*	*	*	*	0.5	0.7	1.1
2003.....	1.4	*	*	*	*	*	*	0.6	0.9	1.1
2002.....	1.2	*	*	*	*	*	*	0.4	0.8	1.2
2001.....	1.0	*	*	*	*	*	*	0.5	0.7	1.0
2000.....	1.0	*	*	*	*	*	*	0.4	0.7	1.1
1999.....	0.9	*	*	*	*	*	*	0.3	0.6	0.8

White, female

2004.....	1.4	*	*	*	*	*	*	0.4	0.7	0.8
2003.....	1.4	*	*	*	*	*	*	0.4	0.4	1.0
2002.....	1.2	*	*	*	*	*	*	0.4	0.5	0.8
2001.....	1.1	*	*	*	*	*	*	*	0.5	0.9
2000.....	1.0	*	*	*	*	*	*	*	0.3	0.6
1999.....	0.9	*	*	*	*	*	*	*	0.3	0.5

Black, both sexes

2004.....	2.2	*	*	*	*	*	*	1.5	2.2	3.0
2003.....	1.9	*	*	*	*	*	*	1.5	1.9	2.5
2002.....	2.0	*	*	*	*	*	0.7	1.4	2.3	2.0
2001.....	1.8	*	*	*	*	*	0.7	0.8	1.6	2.6
2000.....	1.6	*	*	*	*	*	*	1.2	1.5	1.8
1999.....	1.4	*	*	*	*	*	*	1.2	1.3	1.9

Black, male

2004.....	1.7	*	*	*	*	*	*	1.6	2.4	2.8
2003.....	1.4	*	*	*	*	*	*	1.8	2.2	1.9
2002.....	1.5	*	*	*	*	*	*	1.6	2.9	2.1
2001.....	1.3	*	*	*	*	*	*	*	1.6	2.6
2000.....	1.3	*	*	*	*	*	*	*	1.8	2.1
1999.....	1.1	*	*	*	*	*	*	*	1.8	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Metabolic disorders (E70-E88)

All races 3/, both sexes

2004.....	5.4	4.9	0.8	0.4	0.4	0.5	0.8	0.5	0.6	0.7
2003.....	5.5	5.2	1.1	0.4	0.4	0.6	0.7	0.7	0.5	0.7
2002.....	5.5	5.5	0.8	0.4	0.4	0.5	0.7	0.6	0.6	0.6
2001.....	5.5	6.0	1.0	0.4	0.4	0.6	0.7	0.6	0.5	0.7
2000.....	5.4	5.3	0.9	0.3	0.5	0.6	0.8	0.6	0.6	0.7
1999.....	5.3	5.6	0.8	0.4	0.5	0.7	0.7	0.6	0.5	0.6

All races 3/, male

2004.....	5.1	6.0	0.8	0.4	0.4	0.5	0.8	0.5	0.7	0.7
2003.....	5.1	5.8	1.1	0.5	0.5	0.7	0.7	0.7	0.6	0.8
2002.....	4.9	6.1	0.8	0.5	0.5	0.5	0.7	0.7	0.7	0.7
2001.....	4.9	6.5	1.1	0.4	0.4	0.6	0.6	0.6	0.6	0.8
2000.....	4.8	5.5	0.9	0.3	0.6	0.6	0.8	0.8	0.6	0.7
1999.....	4.5	6.3	0.9	0.4	0.5	0.7	0.7	0.7	0.5	0.7

All races 3/, female

2004.....	5.8	3.8	0.7	0.4	0.3	0.5	0.8	0.5	0.5	0.7
2003.....	5.9	4.5	1.0	0.3	0.3	0.6	0.7	0.7	0.4	0.7
2002.....	6.0	4.9	0.9	0.3	0.4	0.6	0.7	0.6	0.5	0.5
2001.....	6.1	5.5	1.0	0.4	0.4	0.6	0.7	0.6	0.5	0.5
2000.....	6.0	5.0	0.8	0.3	0.4	0.7	0.7	0.5	0.5	0.6
1999.....	6.1	4.9	0.7	0.4	0.4	0.7	0.8	0.6	0.5	0.5

White, both sexes

2004.....	5.9	5.0	0.8	0.4	0.4	0.6	0.9	0.6	0.6	0.7
2003.....	5.9	4.6	1.1	0.4	0.5	0.7	0.8	0.7	0.6	0.7
2002.....	5.9	5.0	0.9	0.4	0.5	0.6	0.8	0.7	0.7	0.7
2001.....	5.9	5.6	1.1	0.4	0.4	0.7	0.8	0.7	0.6	0.7
2000.....	5.8	5.0	0.9	0.3	0.6	0.7	0.9	0.7	0.6	0.7
1999.....	5.7	4.9	0.9	0.4	0.5	0.8	0.9	0.8	0.5	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Metabolic disorders (E70-E88)

Black, female

2004.....	4.4	*	*	*	*	*	*	*	*
2003.....	4.9	9.0	*	*	*	*	*	*	*
2002.....	5.1	8.5	*	*	*	*	*	*	*
2001.....	4.9	8.5	*	*	*	*	*	*	*
2000.....	4.9	*	*	*	*	*	*	*	*
1999.....	4.6	8.1	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Cystic fibrosis (E84)

All races 3/, both sexes

2004.....	0.2	*	*	*	0.1	0.3	0.5	0.3	0.3	0.2
2003.....	0.2	*	*	*	0.1	0.3	0.4	0.4	0.2	0.2
2002.....	0.2	*	*	*	0.1	0.3	0.5	0.4	0.3	0.2
2001.....	0.2	*	*	*	0.1	0.4	0.5	0.4	0.2	0.2
2000.....	0.2	*	*	*	0.2	0.4	0.5	0.4	0.3	0.3
1999.....	0.2	*	*	*	0.2	0.4	0.5	0.4	0.3	0.2

All races 3/, male

2004.....	0.2	*	*	*	*	0.3	0.5	0.3	0.3	0.2
2003.....	0.2	*	*	*	*	0.3	0.4	0.4	0.3	0.2
2002.....	0.2	*	*	*	*	0.2	0.4	0.4	0.4	0.2
2001.....	0.2	*	*	*	*	0.3	0.4	0.4	0.3	0.2
2000.....	0.2	*	*	*	*	0.3	0.5	0.4	0.3	0.2
1999.....	0.2	*	*	*	*	0.3	0.5	0.4	0.3	*

All races 3/, female

2004.....	0.2	*	*	*	*	0.3	0.6	0.3	0.3	0.2
2003.....	0.2	*	*	*	*	0.3	0.5	0.4	*	*
2002.....	0.2	*	*	*	*	0.4	0.5	0.3	0.2	0.2
2001.....	0.2	*	*	*	*	0.4	0.6	0.4	0.2	*
2000.....	0.2	*	*	*	0.2	0.4	0.5	0.3	0.4	0.3
1999.....	0.2	*	*	*	*	0.5	0.6	0.4	0.2	0.2

White, both sexes

2004.....	0.2	*	*	*	0.2	0.3	0.6	0.4	0.3	0.3
2003.....	0.2	*	*	*	*	0.4	0.5	0.5	0.3	0.2
2002.....	0.2	*	*	*	0.2	0.4	0.5	0.5	0.4	0.2
2001.....	0.2	*	*	*	0.2	0.5	0.6	0.5	0.3	0.2
2000.....	0.2	*	*	*	0.2	0.4	0.6	0.4	0.4	0.3
1999.....	0.2	*	*	*	0.2	0.5	0.7	0.5	0.3	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

All races 3/, both sexes

2004.....	2.2	1.6	*	*	*	*	*	0.1	0.2
2003.....	2.4	1.9	0.2	*	*	*	*	0.1	0.2
2002.....	2.5	1.8	*	*	*	*	*	0.1	0.2
2001.....	2.7	2.2	0.2	*	*	*	*	0.1	0.2
2000.....	2.8	1.9	*	*	2*	*	*	*	0.1
1999.....	2.9	1.9	0.1	*	*	*	*	0.1	0.2

All races 3/, male

2004.....	1.6	1.9	*	*	*	*	*	*	*
2003.....	1.7	2.2	*	*	*	*	*	*	*
2002.....	1.7	2.1	*	*	*	*	*	*	*
2001.....	1.8	2.2	*	*	*	*	*	*	0.2
2000.....	2.0	1.8	*	*	*	*	*	*	*
1999.....	2.0	2.3	*	*	*	*	*	*	0.2

All races 3/, female

2004.....	2.8	1.3	*	*	*	*	*	*	*
2003.....	3.1	1.7	*	*	*	*	*	*	0.3
2002.....	3.3	1.5	*	*	*	*	*	*	*
2001.....	3.6	2.2	*	*	*	*	*	*	*
2000.....	3.5	2.0	*	*	*	*	*	*	*
1999.....	3.8	1.5	*	*	*	*	*	*	0.2

White, both sexes

2004.....	2.4	1.4	*	*	*	*	*	*	*
2003.....	2.5	1.4	*	*	*	*	*	*	0.2
2002.....	2.7	1.4	*	*	*	*	*	*	0.1
2001.....	2.9	1.7	0.2	*	*	*	*	*	0.2
2000.....	2.9	1.5	*	*	*	*	*	*	0.1
1999.....	3.1	1.3	*	*	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other metabolic disorders (E70-E83,E85,E88)

All races 3/, both sexes

2004.....	3.0	3.1	0.7	0.3	0.2	0.2	0.2	0.1	0.2	0.3
2003.....	2.9	3.1	0.8	0.3	0.3	0.3	0.2	0.2	0.2	0.3
2002.....	2.8	3.4	0.7	0.3	0.3	0.2	0.2	0.2	0.2	0.3
2001.....	2.6	3.7	0.8	0.3	0.2	0.2	0.1	0.1	0.1	0.3
2000.....	2.4	3.2	0.8	0.2	0.3	0.2	0.2	0.2	0.2	0.2
1999.....	2.2	3.4	0.7	0.3	0.3	0.2	0.1	0.1	0.2	0.3

All races 3/, male

2004.....	3.3	3.8	0.7	0.3	0.3	*	0.2	*	0.2	0.4
2003.....	3.2	3.5	1.0	0.4	0.4	0.3	0.2	*	*	0.4
2002.....	3.0	3.6	0.7	0.4	0.4	0.3	0.3	0.2	0.2	0.4
2001.....	2.9	4.1	0.9	0.3	0.3	0.3	*	*	*	0.4
2000.....	2.6	3.6	0.8	0.3	0.4	0.2	0.3	0.3	0.3	0.3
1999.....	2.4	3.7	0.8	0.3	0.3	0.3	*	*	*	0.3

All races 3/, female

2004.....	2.8	2.4	0.6	0.3	*	*	*	*	*	0.2
2003.....	2.6	2.8	0.7	0.2	*	0.2	*	*	*	0.2
2002.....	2.5	3.2	0.7	*	0.2	*	*	*	*	*
2001.....	2.3	3.2	0.7	0.3	*	*	*	*	*	0.2
2000.....	2.3	2.8	0.7	0.2	*	*	*	*	*	0.2
1999.....	2.1	3.1	0.6	0.3	0.2	*	*	*	*	0.2

White, both sexes

2004.....	3.3	3.4	0.7	0.3	0.3	0.2	0.2	0.2	0.2	0.3
2003.....	3.2	3.1	0.9	0.4	0.3	0.3	0.2	0.1	0.2	0.3
2002.....	3.0	3.3	0.7	0.3	0.3	0.2	0.2	0.2	0.2	0.3
2001.....	2.8	3.7	0.8	0.4	0.3	0.2	0.1	0.1	0.1	0.3
2000.....	2.7	3.4	0.8	0.3	0.3	0.2	0.2	0.2	0.1	0.3
1999.....	2.4	3.1	0.8	0.3	0.3	0.2	0.1	0.1	0.1	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

V. Mental and behavioral disorders (F01-F99)

All races 3/, both sexes

2004.....	21.0	*	*	*	0.1	0.4	1.0	1.6	2.1	3.8
2003.....	20.7	*	*	*	*	0.4	1.0	1.5	2.1	4.0
2002.....	19.3	*	*	*	0.1	0.3	0.8	1.3	2.0	4.0
2001.....	17.8	*	*	*	*	0.4	0.8	1.4	2.2	3.7
2000.....	16.4	*	*	*	*	0.4	0.9	1.3	2.3	3.9
1999.....	15.1	*	*	*	0.1	0.4	0.8	1.3	2.3	4.1

All races 3/, male

2004.....	16.2	*	*	*	*	0.5	1.4	2.4	3.0	5.5
2003.....	15.7	*	*	*	*	0.5	1.4	2.3	2.9	5.6
2002.....	14.8	*	*	*	*	0.4	1.1	1.8	2.7	5.6
2001.....	13.8	*	*	*	*	0.5	1.2	2.1	3.2	5.2
2000.....	12.8	*	*	*	*	0.5	1.3	1.9	3.3	6.0
1999.....	12.3	*	*	*	*	0.5	1.1	2.0	3.3	6.0

All races 3/, female

2004.....	25.6	*	*	*	*	0.4	0.7	0.8	1.3	2.1
2003.....	25.5	*	*	*	*	0.3	0.5	0.7	1.2	2.3
2002.....	23.7	*	*	*	*	0.2	0.5	0.7	1.3	2.4
2001.....	21.7	*	*	*	*	0.3	0.3	0.8	1.2	2.2
2000.....	19.8	*	*	*	*	0.3	0.5	0.7	1.3	1.8
1999.....	17.8	*	*	*	*	*	0.5	0.6	1.4	2.2

White, both sexes

2004.....	23.1	*	*	*	*	0.5	1.1	1.7	2.1	3.8
2003.....	22.8	*	*	*	*	0.4	1.0	1.6	2.1	3.9
2002.....	21.3	*	*	*	*	0.4	0.9	1.3	2.1	3.9
2001.....	19.6	*	*	*	*	0.4	0.8	1.4	2.2	3.6
2000.....	17.9	*	*	*	*	0.4	0.9	1.4	2.2	3.7
1999.....	16.5	*	*	*	*	0.4	0.8	1.4	2.4	3.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

V. Mental and behavioral disorders (F01-F99)

White, male

2004.....	17.2	*	*	*	*	0.5	1.6	2.4	2.9	5.5
2003.....	16.7	*	*	*	*	0.5	1.5	2.4	2.9	5.6
2002.....	15.7	*	*	*	*	0.5	1.2	1.9	2.8	5.5
2001.....	14.5	*	*	*	*	0.5	1.2	2.0	3.2	5.1
2000.....	13.4	*	*	*	*	0.5	1.3	2.0	3.1	5.7
1999.....	12.8	*	*	*	*	0.6	1.1	2.1	3.4	5.6

White, female

2004.....	28.8	*	*	*	*	0.4	0.7	0.8	1.3	2.1
2003.....	28.7	*	*	*	*	0.4	0.5	0.7	1.2	2.2
2002.....	26.7	*	*	*	*	0.3	0.5	0.7	1.3	2.3
2001.....	24.4	*	*	*	*	0.3	0.4	0.8	1.2	2.1
2000.....	22.2	*	*	*	*	0.3	0.6	0.8	1.2	1.7
1999.....	20.1	*	*	*	*	*	0.4	0.6	1.3	1.8

Black, both sexes

2004.....	14.4	*	*	*	*	*	*	1.7	2.5	4.1
2003.....	14.0	*	*	*	*	*	0.7	1.6	2.2	5.2
2002.....	13.2	*	*	*	*	*	0.7	1.2	2.3	4.9
2001.....	12.6	*	*	*	*	*	0.8	1.4	2.3	4.9
2000.....	11.8	*	*	*	*	*	1.1	1.4	3.3	5.3
1999.....	10.7	*	*	*	*	*	0.9	1.3	2.4	6.8

Black, male

2004.....	14.0	*	*	*	*	*	*	3.0	4.1	5.8
2003.....	13.4	*	*	*	*	*	*	2.2	2.7	7.4
2002.....	13.1	*	*	*	*	*	*	1.8	2.9	6.9
2001.....	13.0	*	*	*	*	*	1.4	2.5	3.4	7.1
2000.....	12.5	*	*	*	*	*	1.8	2.3	4.6	8.0
1999.....	11.9	*	*	*	*	*	*	2.0	3.1	9.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

V. Mental and behavioral disorders (F01-F99)

Black, female

2004.....	14.8	*	*	*	*	*	*	*	2.6
2003.....	14.5	*	*	*	*	*	*	1.8	3.3
2002.....	13.3	*	*	*	*	*	*	1.8	3.1
2001.....	12.1	*	*	*	*	*	*	1.4	3.0
2000.....	11.2	*	*	*	*	*	*	2.2	2.8
1999.....	9.6	*	*	*	*	*	*	1.8	4.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	All ages 1/ 2	Under 1 year 2/ 3	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------	-------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

All races 3/, both sexes

2004.....	3.6	*	*	*	*	0.3	0.9	1.4	2.0	3.5
2003.....	3.5	*	*	*	*	0.2	0.8	1.3	1.9	3.7
2002.....	3.3	*	*	*	*	0.2	0.7	1.1	1.8	3.7
2001.....	3.2	*	*	*	*	0.3	0.6	1.2	2.0	3.4
2000.....	3.1	*	*	*	*	0.2	0.7	1.1	2.1	3.6
1999.....	3.1	*	*	*	*	0.3	0.6	1.1	2.1	3.8

All races 3/, male

2004.....	5.6	*	*	*	*	0.4	1.3	2.2	2.8	5.2
2003.....	5.3	*	*	*	*	0.3	1.2	2.0	2.7	5.3
2002.....	5.1	*	*	*	*	0.3	1.0	1.7	2.5	5.3
2001.....	4.9	*	*	*	*	0.4	1.1	1.8	2.9	4.9
2000.....	4.8	*	*	*	*	0.3	1.2	1.7	3.0	5.7
1999.....	4.8	*	*	*	*	0.4	1.0	1.7	3.0	5.6

All races 3/, female

2004.....	1.8	*	*	*	*	*	0.4	0.6	1.1	1.8
2003.....	1.7	*	*	*	*	*	0.3	0.5	1.1	2.1
2002.....	1.6	*	*	*	*	*	0.3	0.5	1.0	2.2
2001.....	1.5	*	*	*	*	*	0.2	0.6	1.0	1.9
2000.....	1.5	*	*	*	*	*	0.3	0.6	1.1	1.6
1999.....	1.4	*	*	*	*	*	0.3	0.4	1.2	1.9

White, both sexes

2004.....	3.6	*	*	*	*	0.3	1.0	1.5	1.9	3.5
2003.....	3.4	*	*	*	*	0.3	0.8	1.4	1.9	3.6
2002.....	3.2	*	*	*	*	0.3	0.7	1.1	1.8	3.6
2001.....	3.0	*	*	*	*	0.3	0.7	1.2	1.9	3.3
2000.....	3.0	*	*	*	*	0.3	0.8	1.2	2.0	3.5
1999.....	2.9	*	*	*	*	0.3	0.7	1.1	2.1	3.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

White, male

2004.....	5.5	*	*	*	*	0.5	1.4	2.3	2.8	5.2
2003.....	5.2	*	*	*	*	0.3	1.3	2.2	2.7	5.2
2002.....	4.9	*	*	*	*	0.4	1.1	1.7	2.7	5.2
2001.....	4.6	*	*	*	*	0.5	1.1	1.8	2.9	4.7
2000.....	4.6	*	*	*	*	0.4	1.2	1.8	2.9	5.5
1999.....	4.6	*	*	*	*	0.4	1.0	1.7	3.1	5.2

White, female

2004.....	1.8	*	*	*	*	*	0.5	0.6	1.1	1.8
2003.....	1.7	*	*	*	*	*	0.3	0.5	1.0	1.9
2002.....	1.5	*	*	*	*	*	0.3	0.5	1.0	2.0
2001.....	1.5	*	*	*	*	*	0.3	0.6	1.0	1.8
2000.....	1.4	*	*	*	*	*	0.3	0.6	1.0	1.5
1999.....	1.3	*	*	*	*	*	0.3	0.4	1.1	1.6

Black, both sexes

2004.....	4.3	*	*	*	*	*	*	1.2	2.2	3.8
2003.....	4.3	*	*	*	*	*	*	1.4	2.0	4.8
2002.....	4.4	*	*	*	*	*	*	1.1	2.0	4.7
2001.....	4.6	*	*	*	*	*	*	1.2	2.1	4.6
2000.....	4.6	*	*	*	*	*	0.9	1.1	3.1	4.8
1999.....	4.7	*	*	*	*	*	*	1.1	2.2	6.2

Black, male

2004.....	6.9	*	*	*	*	*	*	2.1	3.6	5.5
2003.....	6.7	*	*	*	*	*	*	1.9	2.5	6.8
2002.....	7.0	*	*	*	*	*	*	1.6	2.7	6.6
2001.....	7.4	*	*	*	*	*	*	2.0	3.0	6.7
2000.....	7.3	*	*	*	*	*	1.5	2.0	4.3	7.2
1999.....	7.4	*	*	*	*	*	*	1.8	2.8	9.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

Black, female

2004.....	2.0	*	*	*	*	*	*	*	2.3
2003.....	2.0	*	*	*	*	*	*	1.5	3.1
2002.....	2.1	*	*	*	*	*	*	1.4	3.0
2001.....	2.1	*	*	*	*	*	*	*	2.7
2000.....	2.1	*	*	*	*	*	*	2.0	2.6
1999.....	2.3	*	*	*	*	*	*	1.7	3.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Mental and behavioral disorders due to use of alcohol (F10)

All races 3/, both sexes

2004.....	2.5	*	*	*	*	0.1	0.3	0.5	1.0	2.0
2003.....	2.4	*	*	*	*	0.1	0.3	0.4	1.0	2.0
2002.....	2.4	*	*	*	*	0.1	0.2	0.5	0.8	2.2
2001.....	2.3	*	*	*	*	0.1	0.2	0.5	1.0	2.2
2000.....	2.2	*	*	*	*	*	0.2	0.4	1.0	2.2
1999.....	2.2	*	*	*	*	*	0.2	0.4	1.0	2.3

All races 3/, male

2004.....	3.9	*	*	*	*	*	0.5	0.8	1.4	3.0
2003.....	3.8	*	*	*	*	*	0.5	0.7	1.5	3.0
2002.....	3.8	*	*	*	*	*	0.3	0.8	1.3	3.3
2001.....	3.7	*	*	*	*	0.2	0.4	0.7	1.5	3.3
2000.....	3.6	*	*	*	*	*	0.3	0.7	1.5	3.6
1999.....	3.6	*	*	*	*	*	0.4	0.7	1.4	3.4

All races 3/, female

2004.....	1.0	*	*	*	*	*	*	*	0.5	1.0
2003.....	1.1	*	*	*	*	*	*	*	0.4	1.0
2002.....	1.0	*	*	*	*	*	*	*	0.4	1.1
2001.....	1.0	*	*	*	*	*	*	0.2	0.5	1.0
2000.....	0.9	*	*	*	*	*	*	*	0.4	0.9
1999.....	0.9	*	*	*	*	*	*	*	0.6	1.1

White, both sexes

2004.....	2.5	*	*	*	*	*	0.3	0.5	0.9	2.0
2003.....	2.4	*	*	*	*	0.1	0.2	0.5	1.0	2.1
2002.....	2.4	*	*	*	*	0.1	0.2	0.5	0.8	2.2
2001.....	2.3	*	*	*	*	0.1	0.2	0.5	1.0	2.2
2000.....	2.2	*	*	*	*	*	0.2	0.4	0.9	2.2
1999.....	2.1	*	*	*	*	*	0.3	0.4	1.0	2.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

All races 3/, both sexes

2004.....	1.2	*	*	*	*	0.2	0.6	0.9	1.0	1.5
2003.....	1.1	*	*	*	*	0.1	0.5	0.9	0.9	1.6
2002.....	0.9	*	*	*	*	0.1	0.5	0.6	1.0	1.5
2001.....	0.9	*	*	*	*	0.1	0.4	0.7	0.9	1.3
2000.....	0.9	*	*	*	*	0.1	0.5	0.7	1.1	1.4
1999.....	0.9	*	*	*	*	0.2	0.4	0.6	1.1	1.5

All races 3/, male

2004.....	1.6	*	*	*	*	0.3	0.8	1.4	1.4	2.2
2003.....	1.5	*	*	*	*	*	0.8	1.3	1.2	2.2
2002.....	1.3	*	*	*	*	*	0.7	0.9	1.3	1.9
2001.....	1.2	*	*	*	*	0.2	0.6	1.1	1.4	1.6
2000.....	1.2	*	*	*	*	*	0.8	1.0	1.5	2.1
1999.....	1.3	*	*	*	*	0.3	0.6	1.0	1.5	2.2

All races 3/, female

2004.....	0.7	*	*	*	*	*	0.3	0.4	0.6	0.9
2003.....	0.7	*	*	*	*	*	0.2	0.4	0.6	1.0
2002.....	0.6	*	*	*	*	*	0.3	0.4	0.6	1.0
2001.....	0.5	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	0.6	*	*	*	*	*	0.2	0.4	0.7	0.8
1999.....	0.5	*	*	*	*	*	*	0.3	0.6	0.8

White, both sexes

2004.....	1.1	*	*	*	*	0.2	0.7	1.0	1.0	1.5
2003.....	1.0	*	*	*	*	0.1	0.6	0.9	0.9	1.5
2002.....	0.9	*	*	*	*	0.1	0.5	0.7	1.0	1.4
2001.....	0.8	*	*	*	*	0.2	0.5	0.7	1.0	1.1
2000.....	0.8	*	*	*	*	0.2	0.6	0.7	1.0	1.3
1999.....	0.8	*	*	*	*	0.2	0.4	0.6	1.1	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

White, male

2004.....	1.5	*	*	*	*	0.3	0.9	1.5	1.4	2.2
2003.....	1.4	*	*	*	*	*	0.9	1.4	1.3	2.1
2002.....	1.2	*	*	*	*	*	0.7	1.0	1.4	1.8
2001.....	1.1	*	*	*	*	0.3	0.7	1.1	1.4	1.5
2000.....	1.1	*	*	*	*	*	0.8	1.0	1.4	1.9
1999.....	1.1	*	*	*	*	0.3	0.6	1.0	1.6	2.0

White, female

2004.....	0.7	*	*	*	*	*	0.4	0.5	0.6	0.8
2003.....	0.6	*	*	*	*	*	0.3	0.3	0.6	0.8
2002.....	0.6	*	*	*	*	*	*	0.3	0.6	1.0
2001.....	0.5	*	*	*	*	*	*	0.4	0.5	0.8
2000.....	0.5	*	*	*	*	*	*	0.5	0.6	0.7
1999.....	0.4	*	*	*	*	*	*	0.3	0.5	0.7

Black, both sexes

2004.....	1.8	*	*	*	*	*	*	0.8	1.2	2.2
2003.....	1.7	*	*	*	*	*	*	1.0	1.3	2.9
2002.....	1.7	*	*	*	*	*	*	0.8	1.1	2.5
2001.....	1.6	*	*	*	*	*	*	0.9	1.2	2.5
2000.....	1.6	*	*	*	*	*	*	*	1.9	3.0
1999.....	1.7	*	*	*	*	*	*	0.8	1.4	3.0

Black, male

2004.....	2.6	*	*	*	*	*	*	*	2.1	2.9
2003.....	2.4	*	*	*	*	*	*	*	*	3.9
2002.....	2.4	*	*	*	*	*	*	*	*	3.2
2001.....	2.3	*	*	*	*	*	*	*	1.6	3.0
2000.....	2.3	*	*	*	*	*	*	*	2.3	4.6
1999.....	2.5	*	*	*	*	*	*	*	1.6	4.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Mental retardation (F70-F79)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	*	0.1	*	0.1
2003.....	0.2	*	*	*	*	*	*	0.1	*	0.1
2002.....	0.2	*	*	*	*	*	*	*	*	0.1
2001.....	0.1	*	*	*	*	*	*	*	0.1	0.1
2000.....	0.1	*	*	*	*	*	*	*	*	0.1
1999.....	0.2	*	*	*	*	*	*	0.1	0.1	0.2

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	*	*	*	*	*	*	*	*	0.1
2003.....	0.2	*	*	*	*	*	*	*	*	0.1
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	0.1	0.1
2000.....	0.2	*	*	*	*	*	*	*	*	0.1
1999.....	0.2	*	*	*	*	*	*	0.1	0.1	0.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

VI. Diseases of the nervous system (G00-G98)

All races 3/, both sexes

2004.....	38.1	9.2	2.0	1.1	1.4	1.9	2.1	2.2	2.4	3.2
2003.....	37.7	10.0	1.8	1.3	1.4	1.9	2.4	2.3	2.4	3.3
2002.....	35.7	9.5	2.1	1.1	1.3	1.8	2.2	2.1	2.3	3.3
2001.....	33.9	8.9	2.1	1.1	1.3	2.0	2.2	2.1	2.3	3.2
2000.....	32.3	10.2	2.1	1.1	1.4	1.8	2.0	1.9	2.3	3.2
1999.....	30.1	11.6	2.4	1.3	1.3	1.9	2.2	1.9	2.2	3.2

All races 3/, male

2004.....	30.0	10.5	2.0	1.2	1.7	2.6	2.6	2.6	2.7	3.6
2003.....	29.9	11.0	1.8	1.4	1.6	2.6	3.0	2.7	2.7	3.6
2002.....	28.5	11.0	2.4	1.2	1.5	2.3	2.8	2.6	2.8	4.0
2001.....	27.4	10.0	2.2	1.1	1.4	2.7	2.9	2.5	2.3	3.8
2000.....	26.2	11.5	2.1	1.1	1.6	2.2	2.6	2.3	2.6	3.5
1999.....	25.0	12.2	2.6	1.4	1.4	2.5	3.0	2.3	2.5	3.7

All races 3/, female

2004.....	45.9	8.0	2.0	1.0	1.2	1.1	1.6	1.7	2.1	2.9
2003.....	45.3	9.0	1.8	1.1	1.1	1.3	1.7	1.9	2.1	3.0
2002.....	42.8	7.8	1.7	1.0	1.1	1.2	1.6	1.7	1.8	2.5
2001.....	40.2	7.8	1.9	1.2	1.1	1.2	1.5	1.6	2.3	2.5
2000.....	38.2	8.8	2.0	1.1	1.2	1.3	1.3	1.5	2.1	2.8
1999.....	35.1	11.0	2.1	1.2	1.1	1.2	1.4	1.6	1.8	2.7

White, both sexes

2004.....	42.7	9.1	1.8	1.0	1.4	1.8	2.0	2.1	2.2	3.1
2003.....	42.4	9.2	1.6	1.2	1.2	1.9	2.4	2.2	2.2	3.0
2002.....	40.2	9.3	1.8	1.1	1.2	1.7	2.0	2.0	2.2	3.1
2001.....	38.0	7.9	1.8	1.1	1.2	1.9	2.1	1.8	2.1	3.0
2000.....	36.3	9.4	1.8	1.1	1.3	1.7	2.0	1.8	2.2	3.0
1999.....	33.7	10.4	2.2	1.2	1.1	1.8	2.1	1.9	2.0	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

VI. Diseases of the nervous system (G00-G98)

White, male

2004.....	33.4	10.6	1.8	1.2	1.6	2.4	2.5	2.7	2.5	3.5
2003.....	33.3	11.1	1.5	1.4	1.4	2.6	3.1	2.6	2.4	3.3
2002.....	31.7	11.0	2.1	1.2	1.3	2.4	2.5	2.5	2.6	3.8
2001.....	30.5	8.7	2.1	1.1	1.4	2.6	2.8	2.2	2.1	3.7
2000.....	29.1	10.3	2.0	1.1	1.5	2.3	2.6	2.3	2.5	3.3
1999.....	27.7	11.8	2.5	1.2	1.2	2.5	2.8	2.3	2.3	3.6

White, female

2004.....	51.8	7.5	1.7	0.9	1.1	1.1	1.5	1.4	1.8	2.7
2003.....	51.4	7.1	1.7	1.0	1.0	1.2	1.6	1.8	2.0	2.7
2002.....	48.6	7.5	1.5	0.9	1.0	1.0	1.5	1.5	1.7	2.3
2001.....	45.4	7.1	1.6	1.1	1.0	1.1	1.3	1.4	2.1	2.3
2000.....	43.2	8.4	1.6	1.0	1.1	1.1	1.3	1.3	1.9	2.6
1999.....	39.5	9.0	1.8	1.1	1.1	1.0	1.3	1.5	1.7	2.5

Black, both sexes

2004.....	21.9	10.4	3.2	1.7	1.9	2.8	3.2	3.3	4.8	5.0
2003.....	20.8	14.6	2.8	1.7	2.2	2.3	3.0	3.6	4.2	6.1
2002.....	19.5	11.6	3.3	1.3	2.0	2.3	3.5	3.6	4.1	5.5
2001.....	19.0	14.4	3.1	1.5	1.8	3.0	3.0	3.9	4.2	5.4
2000.....	17.9	14.3	3.4	1.4	2.2	2.5	2.5	3.2	3.8	5.4
1999.....	16.7	17.8	3.3	1.9	1.8	2.5	3.5	2.8	3.5	5.3

Black, male

2004.....	17.3	10.4	3.2	1.2	2.1	3.9	3.5	2.8	4.9	5.0
2003.....	17.5	12.5	3.2	1.3	2.8	2.8	3.3	4.0	5.2	6.6
2002.....	16.7	11.9	4.0	1.3	2.4	2.5	4.8	4.3	4.8	6.7
2001.....	16.1	17.1	2.9	1.2	1.7	4.0	3.5	5.0	4.5	6.5
2000.....	15.4	17.9	3.1	1.1	2.7	2.3	3.0	3.5	3.8	5.8
1999.....	14.6	15.9	3.2	2.3	2.4	2.9	5.0	3.3	3.9	5.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

VI. Diseases of the nervous system (G00-G98)

Black, female

2004.....	26.1	10.5	3.3	2.2	1.7	1.7	2.9	3.8	4.7	5.0
2003.....	23.9	16.7	2.3	2.1	1.6	1.8	2.6	3.2	3.3	5.7
2002.....	22.2	11.2	2.7	1.3	1.6	2.1	2.4	2.9	3.5	4.5
2001.....	21.5	11.7	3.4	1.8	1.8	1.9	2.5	2.9	4.0	4.4
2000.....	20.2	10.6	3.6	1.8	1.8	2.7	2.1	2.8	3.8	4.9
1999.....	18.5	19.8	3.4	1.5	1.2	2.0	2.1	2.3	3.1	5.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Meningitis (G00,G03)

All races 3/, both sexes

2004.....	0.2	1.6	0.2	*	*	*	*	*	0.1	0.1
2003.....	0.3	1.9	0.2	*	*	*	0.1	*	0.1	0.1
2002.....	0.2	1.8	*	*	0.1	*	*	*	0.1	0.1
2001.....	0.3	1.9	0.2	*	*	*	0.1	*	0.1	0.1
2000.....	0.3	2.4	0.2	*	*	*	*	*	0.1	0.2
1999.....	0.3	3.1	0.3	*	*	*	0.1	0.1	0.1	0.1

All races 3/, male

2004.....	0.2	1.5	*	*	*	*	*	*	*	*
2003.....	0.3	1.9	*	*	*	*	*	*	*	0.2
2002.....	0.3	2.0	*	*	*	*	*	*	*	*
2001.....	0.3	1.9	*	*	*	*	*	*	*	0.2
2000.....	0.3	2.5	0.3	*	*	*	*	*	*	0.2
1999.....	0.3	3.1	0.3	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.2	1.7	*	*	*	*	*	*	*	*
2003.....	0.2	2.0	*	*	*	*	*	*	*	*
2002.....	0.2	1.7	*	*	*	*	*	*	*	*
2001.....	0.2	2.0	*	*	*	*	*	*	*	*
2000.....	0.2	2.3	*	*	*	*	*	*	*	*
1999.....	0.3	3.0	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	1.4	*	*	*	*	*	*	*	*
2003.....	0.2	1.7	*	*	*	*	*	*	*	*
2002.....	0.2	1.7	*	*	*	*	*	*	*	*
2001.....	0.2	1.5	*	*	*	*	*	*	0.1	*
2000.....	0.2	1.8	0.2	*	*	*	*	*	*	0.1
1999.....	0.3	2.3	0.3	*	*	*	*	*	*	0.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Meningitis (G00,G03)

Black, female

2004.....	0.4	*	*	*	*	*	*	*	*
2003.....	0.5	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*
1999.....	0.5	9.7	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Multiple sclerosis (G35)

All races 3/, both sexes

2004.....	1.1	*	*	*	*	*	*	*	0.2	0.4
2003.....	1.1	*	*	*	*	*	*	*	0.2	0.4
2002.....	1.1	*	*	*	*	*	*	*	0.1	0.4
2001.....	1.1	*	*	*	*	*	*	*	0.2	0.4
2000.....	1.0	*	*	*	*	*	*	0.1	0.2	0.4
1999.....	1.0	*	*	*	*	*	*	*	0.2	0.4

All races 3/, male

2004.....	0.8	*	*	*	*	*	*	*	*	0.3
2003.....	0.8	*	*	*	*	*	*	*	*	0.3
2002.....	0.8	*	*	*	*	*	*	*	*	0.4
2001.....	0.7	*	*	*	*	*	*	*	*	0.3
2000.....	0.7	*	*	*	*	*	*	*	*	0.3
1999.....	0.6	*	*	*	*	*	*	*	*	0.3

All races 3/, female

2004.....	1.5	*	*	*	*	*	*	*	0.2	0.5
2003.....	1.4	*	*	*	*	*	*	*	0.2	0.6
2002.....	1.4	*	*	*	*	*	*	*	*	0.5
2001.....	1.4	*	*	*	*	*	*	*	0.2	0.5
2000.....	1.3	*	*	*	*	*	*	*	0.3	0.5
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.5

White, both sexes

2004.....	1.2	*	*	*	*	*	*	*	*	0.4
2003.....	1.2	*	*	*	*	*	*	*	0.1	0.4
2002.....	1.2	*	*	*	*	*	*	*	*	0.4
2001.....	1.2	*	*	*	*	*	*	*	0.2	0.4
2000.....	1.1	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.0	*	*	*	*	*	*	*	0.2	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Epilepsy (G40-G41)

All races 3/, both sexes

2004.....	0.5	*	0.2	*	0.1	0.2	0.3	0.4	0.5	0.5
2003.....	0.5	*	0.2	*	0.2	0.2	0.4	0.4	0.4	0.5
2002.....	0.5	*	0.1	0.1	0.1	0.2	0.4	0.4	0.4	0.5
2001.....	0.5	*	*	0.1	*	0.2	0.3	0.4	0.4	0.5
2000.....	0.5	*	0.2	*	0.1	0.2	0.3	0.3	0.4	0.5
1999.....	0.5	*	0.2	0.1	0.1	0.2	0.3	0.4	0.4	0.5

All races 3/, male

2004.....	0.5	*	*	*	*	0.2	0.3	0.5	0.6	0.5
2003.....	0.5	*	*	*	0.2	0.2	0.4	0.5	0.5	0.6
2002.....	0.5	*	*	*	*	*	0.4	0.5	0.5	0.6
2001.....	0.5	*	*	*	*	0.2	0.4	0.5	0.4	0.7
2000.....	0.5	*	*	*	*	*	0.4	0.4	0.5	0.5
1999.....	0.5	*	0.3	*	*	0.3	0.4	0.6	0.5	0.7

All races 3/, female

2004.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.4
2003.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.4
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.4
2001.....	0.4	*	*	*	*	*	*	0.3	0.4	0.3
2000.....	0.4	*	*	*	*	0.2	*	0.2	0.4	0.4
1999.....	0.4	*	*	*	*	*	0.3	0.3	0.3	0.3

White, both sexes

2004.....	0.4	*	0.2	*	0.1	0.2	0.3	0.4	0.4	0.4
2003.....	0.4	*	0.2	*	0.1	0.2	0.4	0.4	0.4	0.5
2002.....	0.4	*	*	*	*	0.2	0.3	0.4	0.4	0.5
2001.....	0.4	*	*	0.1	*	0.2	0.2	0.3	0.4	0.4
2000.....	0.4	*	0.2	*	*	0.1	0.3	0.3	0.4	0.4
1999.....	0.4	*	0.2	*	*	0.2	0.3	0.4	0.4	0.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Infantile cerebral palsy (G80)

All races 3/, both sexes

2004.....	0.5	*	0.5	0.5	0.7	0.8	0.7	0.6	0.3	0.3
2003.....	0.5	*	0.4	0.6	0.6	0.7	0.7	0.6	0.4	0.3
2002.....	0.4	*	0.6	0.5	0.5	0.6	0.7	0.5	0.2	0.2
2001.....	0.4	*	0.5	0.4	0.5	0.7	0.6	0.4	0.3	0.2
2000.....	0.4	*	0.4	0.5	0.6	0.6	0.6	0.3	0.3	0.2
1999.....	0.4	*	0.5	0.4	0.5	0.5	0.5	0.4	0.3	0.3

All races 3/, male

2004.....	0.6	*	0.4	0.5	0.7	1.0	0.8	0.7	0.3	0.3
2003.....	0.5	*	0.5	0.7	0.7	0.8	0.7	0.6	0.4	0.3
2002.....	0.5	*	0.7	0.6	0.5	0.8	0.8	0.5	0.3	0.3
2001.....	0.5	*	0.5	0.4	0.6	0.8	0.7	0.4	0.4	0.2
2000.....	0.4	*	0.3	0.5	0.7	0.7	0.7	0.4	0.4	0.2
1999.....	0.4	*	0.6	0.5	0.5	0.6	0.6	0.4	0.3	0.2

All races 3/, female

2004.....	0.4	*	0.5	0.4	0.6	0.5	0.6	0.4	0.3	0.3
2003.....	0.4	*	0.4	0.5	0.5	0.6	0.6	0.5	0.3	0.3
2002.....	0.4	*	0.5	0.5	0.5	0.5	0.5	0.4	0.2	0.2
2001.....	0.3	*	0.4	0.5	0.5	0.6	0.5	0.3	0.3	0.2
2000.....	0.3	*	0.4	0.6	0.4	0.5	0.5	0.3	0.3	0.2
1999.....	0.3	*	0.4	0.4	0.4	0.4	0.3	0.4	0.3	0.3

White, both sexes

2004.....	0.5	*	0.3	0.4	0.6	0.7	0.7	0.6	0.3	0.3
2003.....	0.4	*	0.3	0.5	0.5	0.6	0.6	0.6	0.3	0.3
2002.....	0.4	*	0.5	0.5	0.4	0.6	0.5	0.4	0.2	0.2
2001.....	0.4	*	0.4	0.4	0.5	0.6	0.6	0.3	0.3	0.2
2000.....	0.4	*	0.3	0.5	0.5	0.6	0.5	0.3	0.3	0.2
1999.....	0.3	*	0.5	0.4	0.4	0.5	0.4	0.4	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

All races 3/, both sexes

2004.....	6.9	6.7	1.1	0.5	0.5	0.8	0.9	0.9	1.2	1.9
2003.....	7.1	7.3	0.9	0.5	0.6	0.9	1.2	1.1	1.2	1.8
2002.....	6.9	6.9	1.1	0.3	0.6	0.8	1.0	1.0	1.3	1.8
2001.....	6.7	6.1	1.2	0.5	0.6	0.9	1.1	1.0	1.1	1.8
2000.....	6.7	7.1	1.2	0.4	0.5	0.9	1.0	1.0	1.2	1.8
1999.....	6.5	7.7	1.2	0.5	0.6	0.9	1.2	0.9	1.1	1.7

All races 3/, male

2004.....	7.3	7.8	1.1	0.5	0.6	1.2	1.3	1.2	1.4	2.2
2003.....	7.5	8.3	0.9	0.5	0.7	1.3	1.7	1.4	1.4	2.1
2002.....	7.3	8.4	1.3	0.4	0.7	1.2	1.4	1.4	1.7	2.4
2001.....	7.1	7.1	1.2	0.5	0.7	1.5	1.6	1.4	1.1	2.3
2000.....	7.2	8.1	1.3	0.4	0.6	1.3	1.4	1.3	1.5	2.1
1999.....	7.0	8.6	1.2	0.6	0.6	1.5	1.7	1.1	1.3	2.2

All races 3/, female

2004.....	6.5	5.6	1.1	0.4	0.4	0.4	0.5	0.7	1.0	1.6
2003.....	6.6	6.2	1.0	0.5	0.4	0.5	0.6	0.7	1.0	1.4
2002.....	6.5	5.4	0.9	0.3	0.4	0.4	0.6	0.7	0.9	1.3
2001.....	6.3	5.0	1.2	0.5	0.5	0.3	0.7	0.7	1.1	1.2
2000.....	6.3	6.0	1.1	0.4	0.5	0.5	0.6	0.6	1.0	1.4
1999.....	6.1	6.8	1.2	0.5	0.5	0.4	0.6	0.6	0.8	1.3

White, both sexes

2004.....	7.3	6.8	1.0	0.5	0.5	0.8	0.9	0.9	1.1	1.9
2003.....	7.5	6.8	0.9	0.5	0.5	1.0	1.2	1.1	1.2	1.7
2002.....	7.3	6.9	1.0	0.3	0.6	0.8	1.0	1.0	1.3	1.8
2001.....	7.0	5.8	1.1	0.5	0.6	0.9	1.1	0.9	1.0	1.8
2000.....	7.1	6.9	1.1	0.4	0.5	0.9	1.0	0.9	1.2	1.7
1999.....	6.9	7.5	1.1	0.5	0.5	1.0	1.2	0.9	1.0	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

White, male

2004.....	7.8	8.0	1.0	0.5	0.7	1.3	1.3	1.2	1.3	2.2
2003.....	8.0	8.2	0.7	0.5	0.6	1.5	1.8	1.4	1.3	1.9
2002.....	7.8	8.5	1.1	0.4	0.7	1.3	1.4	1.4	1.7	2.4
2001.....	7.5	6.6	1.1	0.5	0.7	1.5	1.6	1.2	1.0	2.4
2000.....	7.6	7.4	1.2	0.4	0.5	1.4	1.5	1.3	1.5	2.1
1999.....	7.4	8.4	1.1	0.5	0.5	1.6	1.8	1.2	1.2	2.1

White, female

2004.....	6.8	5.6	1.0	0.4	0.4	0.3	0.4	0.5	0.9	1.5
2003.....	7.0	5.2	1.0	0.4	0.4	0.5	0.6	0.7	1.0	1.4
2002.....	6.9	5.2	0.8	0.3	0.4	0.3	0.5	0.6	0.8	1.3
2001.....	6.6	4.9	1.0	0.5	0.4	0.3	0.6	0.6	0.9	1.2
2000.....	6.6	6.3	1.0	0.5	0.4	0.4	0.6	0.5	0.9	1.3
1999.....	6.4	6.5	1.1	0.5	0.5	0.3	0.6	0.6	0.8	1.2

Black, both sexes

2004.....	6.2	6.8	1.7	*	0.6	0.9	1.0	1.4	2.1	2.6
2003.....	6.2	9.9	1.5	*	0.9	0.7	1.0	1.4	1.6	3.0
2002.....	6.0	7.9	1.7	*	0.7	0.9	1.3	1.4	1.8	2.7
2001.....	6.1	8.0	2.0	*	0.7	1.1	1.4	1.7	2.2	2.4
2000.....	5.9	7.5	1.6	*	0.6	0.9	1.0	1.7	1.6	2.7
1999.....	5.7	9.1	1.4	0.8	0.8	0.8	1.4	0.9	1.7	2.6

Black, male

2004.....	6.2	7.4	1.9	*	*	1.3	*	*	2.2	2.8
2003.....	6.3	9.8	1.8	*	1.2	*	*	1.8	1.9	3.7
2002.....	6.3	7.8	2.0	*	*	*	1.7	1.7	2.1	3.5
2001.....	6.3	10.2	1.9	*	*	1.6	1.5	2.5	2.2	2.9
2000.....	6.2	10.9	1.6	*	*	*	*	2.3	1.7	3.0
1999.....	5.7	10.4	*	*	1.2	*	2.3	*	2.1	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

IX. Diseases of the circulatory system (I00-I99)

All races 3/, both sexes

2004.....	294.9	14.5	1.5	0.6	1.0	2.2	4.3	7.5	13.0	24.8
2003.....	311.9	14.8	1.5	0.7	1.0	2.6	4.7	7.3	13.6	25.7
2002.....	320.2	16.5	1.5	0.6	1.1	2.5	4.0	7.3	13.1	25.9
2001.....	325.5	15.4	1.9	0.7	1.1	2.2	4.4	7.4	13.4	25.5
2000.....	334.6	17.4	1.5	0.7	1.1	2.5	4.4	7.0	12.5	24.7
1999.....	342.0	17.6	1.5	0.7	1.0	2.8	4.2	6.6	12.9	26.3

All races 3/, male

2004.....	282.7	15.6	1.5	0.5	1.1	2.8	5.4	9.5	16.7	32.6
2003.....	297.7	16.2	1.5	0.8	1.2	3.1	5.5	8.7	17.4	34.0
2002.....	304.7	17.3	1.5	0.6	1.3	3.2	5.1	9.1	16.6	33.5
2001.....	307.6	15.8	2.0	0.6	1.1	2.7	5.4	8.9	16.8	33.5
2000.....	317.2	18.3	1.7	0.8	1.2	2.9	5.1	8.3	15.7	31.9
1999.....	324.2	18.5	1.6	0.8	1.1	3.4	4.9	7.9	16.4	34.6

All races 3/, female

2004.....	306.7	13.5	1.5	0.7	1.0	1.6	3.2	5.5	9.3	16.9
2003.....	325.8	13.3	1.5	0.6	0.8	2.1	3.8	5.9	9.8	17.4
2002.....	335.2	15.7	1.4	0.7	0.9	1.7	2.9	5.4	9.5	18.2
2001.....	342.8	15.0	1.8	0.8	1.1	1.7	3.4	5.8	9.9	17.5
2000.....	351.3	16.5	1.4	0.6	1.0	2.1	3.7	5.6	9.4	17.6
1999.....	359.1	16.6	1.4	0.6	1.0	2.1	3.6	5.3	9.4	18.0

White, both sexes

2004.....	313.3	12.6	1.2	0.5	0.9	1.9	3.7	6.0	10.8	21.5
2003.....	331.7	11.9	1.2	0.6	0.9	2.3	3.8	5.9	11.3	21.8
2002.....	341.0	14.1	1.3	0.6	0.9	2.2	3.4	5.9	10.6	22.1
2001.....	346.4	12.7	1.6	0.6	1.0	2.0	3.5	6.1	11.2	21.8
2000.....	356.4	15.3	1.4	0.6	0.9	2.1	3.6	5.7	10.4	21.1
1999.....	364.2	15.2	1.3	0.7	0.9	2.4	3.5	5.5	10.9	22.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

IX. Diseases of the circulatory system (I00-I99)

White, male

2004.....	298.1	13.7	1.3	0.5	1.0	2.4	4.7	7.8	14.1	28.6
2003.....	314.0	12.5	1.1	0.6	1.1	2.7	4.5	7.1	14.7	29.9
2002.....	322.2	15.2	1.5	0.5	1.1	2.8	4.4	7.6	13.7	29.0
2001.....	324.9	13.7	1.6	0.6	1.0	2.3	4.4	7.5	14.3	29.2
2000.....	335.6	16.1	1.6	0.7	1.1	2.5	4.3	7.0	13.5	28.1
1999.....	342.5	15.9	1.5	0.8	0.9	2.9	3.9	6.6	14.2	30.2

White, female

2004.....	328.3	11.4	1.1	0.6	0.8	1.4	2.7	4.1	7.3	14.1
2003.....	348.9	11.2	1.4	0.5	0.7	1.8	3.0	4.7	7.7	13.6
2002.....	359.5	13.0	1.2	0.7	0.8	1.5	2.3	4.1	7.4	14.9
2001.....	367.4	11.7	1.5	0.7	0.9	1.6	2.6	4.6	7.9	14.2
2000.....	376.6	14.4	1.1	0.5	0.8	1.7	2.9	4.4	7.2	13.9
1999.....	385.2	14.4	1.2	0.6	0.9	1.9	3.0	4.3	7.4	14.4

Black, both sexes

2004.....	262.4	23.0	2.9	1.1	1.8	4.2	8.0	17.6	29.6	51.1
2003.....	275.3	27.1	2.8	1.2	1.6	4.5	10.0	16.8	30.5	54.2
2002.....	278.2	28.6	2.0	0.8	2.0	4.1	8.1	17.2	31.0	54.4
2001.....	282.4	29.3	3.6	1.0	1.6	4.0	9.7	17.0	29.5	52.7
2000.....	287.0	28.6	2.6	1.0	1.9	4.8	9.4	15.9	28.7	52.4
1999.....	291.1	29.9	2.5	1.0	1.8	5.0	9.3	14.8	27.5	56.2

Black, male

2004.....	259.1	24.6	2.8	*	1.8	5.6	10.2	21.1	36.5	66.9
2003.....	270.6	32.7	3.3	1.4	1.7	5.8	12.0	20.5	37.1	66.6
2002.....	270.5	28.8	2.0	*	2.3	5.3	9.8	21.1	38.5	68.3
2001.....	274.3	29.0	3.8	*	1.5	5.1	11.6	20.8	36.0	66.9
2000.....	277.6	29.7	2.2	*	1.8	5.3	10.5	18.7	33.7	63.2
1999.....	284.2	31.6	2.6	*	2.1	6.4	11.1	18.0	33.8	70.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

IX. Diseases of the circulatory system (I00-I99)

Black, female

2004.....	265.3	21.3	3.0	1.3	1.8	2.7	5.7	14.2	23.4	36.9
2003.....	279.5	21.3	2.4	*	1.5	3.3	8.0	13.3	24.5	43.2
2002.....	285.2	28.5	2.1	*	1.6	2.8	6.5	13.6	24.2	42.0
2001.....	289.9	29.6	3.4	1.3	1.6	2.9	8.0	13.4	23.5	40.1
2000.....	295.6	27.4	2.9	*	2.0	4.4	8.4	13.5	24.1	42.8
1999.....	297.4	28.2	2.4	*	1.5	3.5	7.6	11.8	21.9	43.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Hypertensive diseases (I10-I13)

All races 3/, both sexes

2004.....	18.6	*	*	*	*	0.1	0.2	0.7	1.4	2.7
2003.....	18.4	*	*	*	*	*	0.3	0.7	1.5	2.7
2002.....	17.2	*	*	*	*	*	0.2	0.7	1.3	2.5
2001.....	16.4	*	*	*	*	*	0.2	0.6	1.3	2.2
2000.....	15.9	*	*	*	*	*	0.2	0.5	1.0	2.0
1999.....	15.4	*	*	*	*	*	0.2	0.4	1.1	2.2

All races 3/, male

2004.....	16.0	*	*	*	*	*	0.3	1.0	1.9	3.9
2003.....	15.4	*	*	*	*	*	0.4	0.8	1.9	3.6
2002.....	14.5	*	*	*	*	*	0.3	1.0	1.8	3.5
2001.....	13.5	*	*	*	*	*	0.3	0.7	1.7	2.8
2000.....	13.0	*	*	*	*	*	0.3	0.6	1.3	2.5
1999.....	12.6	*	*	*	*	*	0.3	0.6	1.5	2.8

All races 3/, female

2004.....	21.2	*	*	*	*	*	*	0.5	0.9	1.5
2003.....	21.2	*	*	*	*	*	*	0.5	1.0	1.8
2002.....	19.9	*	*	*	*	*	*	0.4	0.9	1.6
2001.....	19.2	*	*	*	*	*	*	0.5	0.9	1.6
2000.....	18.6	*	*	*	*	*	*	0.4	0.7	1.5
1999.....	18.1	*	*	*	*	*	*	0.2	0.7	1.5

White, both sexes

2004.....	17.2	*	*	*	*	*	0.2	0.4	0.9	1.7
2003.....	16.9	*	*	*	*	*	0.2	0.4	0.9	1.7
2002.....	15.8	*	*	*	*	*	0.2	0.4	0.8	1.5
2001.....	15.0	*	*	*	*	*	*	0.3	0.8	1.4
2000.....	14.4	*	*	*	*	*	0.1	0.3	0.5	1.2
1999.....	13.9	*	*	*	*	*	*	0.2	0.7	1.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Hypertensive diseases (I10-I13)

White, male

2004.....	14.2	*	*	*	*	*	0.2	0.6	1.3	2.5
2003.....	13.6	*	*	*	*	*	0.3	0.6	1.2	2.6
2002.....	12.7	*	*	*	*	*	0.2	0.6	1.0	2.2
2001.....	11.8	*	*	*	*	*	*	0.4	1.1	2.0
2000.....	11.3	*	*	*	*	*	*	0.4	0.8	1.7
1999.....	10.8	*	*	*	*	*	*	0.4	1.1	1.7

White, female

2004.....	20.1	*	*	*	*	*	*	*	0.4	0.8
2003.....	20.1	*	*	*	*	*	*	*	0.6	0.8
2002.....	18.8	*	*	*	*	*	*	*	0.5	0.9
2001.....	18.2	*	*	*	*	*	*	*	0.4	0.8
2000.....	17.4	*	*	*	*	*	*	*	0.3	0.7
1999.....	16.9	*	*	*	*	*	*	*	0.3	0.6

Black, both sexes

2004.....	32.2	*	*	*	*	*	0.7	3.0	4.8	9.7
2003.....	32.1	*	*	*	*	*	1.0	2.2	5.4	9.1
2002.....	30.7	*	*	*	*	*	0.7	2.4	5.3	9.3
2001.....	29.1	*	*	*	*	*	0.7	2.0	4.9	7.9
2000.....	28.9	*	*	*	*	*	0.8	1.8	4.4	7.5
1999.....	28.6	*	*	*	*	*	1.0	1.5	3.9	8.9

Black, male

2004.....	31.3	*	*	*	*	*	*	3.7	6.0	14.1
2003.....	30.8	*	*	*	*	*	1.4	2.2	7.2	11.0
2002.....	29.3	*	*	*	*	*	*	3.6	7.0	13.1
2001.....	27.7	*	*	*	*	*	*	2.4	6.2	9.1
2000.....	26.8	*	*	*	*	*	*	2.0	5.2	8.5
1999.....	26.8	*	*	*	*	*	*	1.9	4.4	11.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Hypertensive diseases (I10-I13)

Black, female

2004.....	33.0	*	*	*	*	*	*	2.4	3.8	5.8
2003.....	33.3	*	*	*	*	*	*	2.2	3.8	7.5
2002.....	31.9	*	*	*	*	*	*	*	3.8	6.0
2001.....	30.5	*	*	*	*	*	*	1.7	3.8	6.8
2000.....	30.8	*	*	*	*	*	*	1.6	3.6	6.5
1999.....	30.3	*	*	*	*	*	*	*	3.6	6.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Hypertensive heart disease (I11)

All races 3/, both sexes

2004.....	9.7	*	*	*	*	*	0.1	0.5	1.0	1.9
2003.....	9.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2002.....	9.2	*	*	*	*	*	0.1	0.4	1.0	1.8
2001.....	8.7	*	*	*	*	*	*	0.3	0.9	1.6
2000.....	8.4	*	*	*	*	*	0.1	0.4	0.7	1.4
1999.....	8.1	*	*	*	*	*	0.1	0.3	0.8	1.5

All races 3/, male

2004.....	8.9	*	*	*	*	*	*	0.7	1.5	2.8
2003.....	8.8	*	*	*	*	*	0.3	0.7	1.4	2.7
2002.....	8.2	*	*	*	*	*	0.2	0.7	1.4	2.6
2001.....	7.5	*	*	*	*	*	*	0.4	1.2	2.1
2000.....	7.2	*	*	*	*	*	0.2	0.4	0.9	1.9
1999.....	6.8	*	*	*	*	*	*	0.4	1.0	2.0

All races 3/, female

2004.....	10.5	*	*	*	*	*	*	0.3	0.5	1.0
2003.....	10.7	*	*	*	*	*	*	0.3	0.7	1.2
2002.....	10.2	*	*	*	*	*	*	*	0.6	1.0
2001.....	9.8	*	*	*	*	*	*	0.3	0.5	1.1
2000.....	9.7	*	*	*	*	*	*	0.3	0.5	1.0
1999.....	9.4	*	*	*	*	*	*	*	0.5	1.0

White, both sexes

2004.....	8.9	*	*	*	*	*	*	0.3	0.6	1.3
2003.....	8.8	*	*	*	*	*	0.1	0.3	0.7	1.3
2002.....	8.3	*	*	*	*	*	*	0.3	0.6	1.3
2001.....	7.8	*	*	*	*	*	*	0.2	0.5	1.1
2000.....	7.6	*	*	*	*	*	*	0.2	0.4	0.9
1999.....	7.3	*	*	*	*	*	*	0.2	0.5	0.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Hypertensive heart disease (I11)

White, male

2004.....	7.8	*	*	*	*	*	*	0.5	1.0	2.0
2003.....	7.7	*	*	*	*	*	*	0.5	1.0	2.1
2002.....	7.1	*	*	*	*	*	*	0.4	0.9	1.8
2001.....	6.4	*	*	*	*	*	*	0.3	0.9	1.6
2000.....	6.1	*	*	*	*	*	*	0.3	0.6	1.4
1999.....	5.8	*	*	*	*	*	*	0.3	0.8	1.3

White, female

2004.....	9.9	*	*	*	*	*	*	*	0.3	0.6
2003.....	10.0	*	*	*	*	*	*	*	0.3	0.6
2002.....	9.5	*	*	*	*	*	*	*	0.3	0.7
2001.....	9.2	*	*	*	*	*	*	*	*	0.5
2000.....	9.0	*	*	*	*	*	*	*	*	0.5
1999.....	8.7	*	*	*	*	*	*	*	*	0.5

Black, both sexes

2004.....	17.6	*	*	*	*	*	*	2.1	3.2	6.1
2003.....	17.9	*	*	*	*	*	0.6	1.5	3.5	6.0
2002.....	17.2	*	*	*	*	*	*	1.5	3.9	5.8
2001.....	16.0	*	*	*	*	*	*	1.1	3.1	5.1
2000.....	16.1	*	*	*	*	*	*	1.4	2.8	4.9
1999.....	15.5	*	*	*	*	*	*	1.0	2.6	5.7

Black, male

2004.....	18.4	*	*	*	*	*	*	2.5	4.6	8.8
2003.....	18.1	*	*	*	*	*	*	1.7	4.3	7.3
2002.....	17.5	*	*	*	*	*	*	2.4	5.4	8.5
2001.....	15.9	*	*	*	*	*	*	*	4.0	5.9
2000.....	15.6	*	*	*	*	*	*	1.6	3.5	5.8
1999.....	15.2	*	*	*	*	*	*	*	2.8	7.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Hypertensive heart disease (I11)

Black, female

2004.....	16.9	*	*	*	*	*	*	1.8	2.1	3.6
2003.....	17.8	*	*	*	*	*	*	*	2.7	4.8
2002.....	16.9	*	*	*	*	*	*	*	2.6	3.4
2001.....	16.2	*	*	*	*	*	*	*	2.3	4.4
2000.....	16.5	*	*	*	*	*	*	*	2.1	4.1
1999.....	15.9	*	*	*	*	*	*	*	2.5	4.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Ischemic heart diseases (I20-I25)

All races 3/, both sexes

2004.....	153.7	*	*	*	*	0.1	0.5	1.5	3.9	9.4
2003.....	165.1	0.5	*	*	*	0.2	0.6	1.4	3.9	10.1
2002.....	171.4	0.6	*	*	*	0.1	0.5	1.3	3.6	10.1
2001.....	176.3	0.6	*	*	*	0.2	0.7	1.5	4.0	9.9
2000.....	183.1	0.8	*	*	*	0.2	0.5	1.2	3.4	9.8
1999.....	189.8	0.7	*	*	*	0.3	0.5	1.3	3.7	10.4

All races 3/, male

2004.....	161.6	*	*	*	*	0.2	0.7	2.1	6.0	13.9
2003.....	172.2	*	*	*	*	0.2	0.8	2.0	5.9	15.2
2002.....	178.4	*	*	*	*	*	0.7	2.0	5.2	15.2
2001.....	181.7	*	*	*	*	0.2	0.9	2.2	5.9	15.0
2000.....	188.7	*	*	*	*	0.2	0.6	1.8	5.1	14.7
1999.....	195.4	*	*	*	*	0.3	0.6	1.7	5.5	15.8

All races 3/, female

2004.....	146.1	*	*	*	*	*	0.2	0.9	1.8	4.9
2003.....	158.1	*	*	*	*	*	0.4	0.8	1.9	4.9
2002.....	164.7	*	*	*	*	*	0.3	0.6	1.9	4.9
2001.....	171.2	*	*	*	*	*	0.4	0.7	2.1	4.9
2000.....	177.6	*	*	*	*	*	0.3	0.6	1.7	5.0
1999.....	184.5	*	*	*	*	*	0.3	0.8	1.9	5.1

White, both sexes

2004.....	166.1	*	*	*	*	*	0.4	1.3	3.6	9.2
2003.....	178.3	*	*	*	*	0.2	0.5	1.3	3.7	9.8
2002.....	185.4	*	*	*	*	0.1	0.4	1.2	3.3	9.6
2001.....	190.5	*	*	*	*	0.1	0.5	1.3	3.8	9.5
2000.....	197.9	0.8	*	*	*	0.2	0.4	1.0	3.1	9.4
1999.....	205.1	*	*	*	*	0.2	0.3	1.2	3.5	9.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Ischemic heart diseases (I20-I25)

White, male

2004.....	174.3	*	*	*	*	*	0.5	1.9	5.6	13.6
2003.....	185.5	*	*	*	*	*	0.7	1.8	5.6	15.0
2002.....	192.5	*	*	*	*	*	0.6	1.9	5.0	14.6
2001.....	195.8	*	*	*	*	*	0.8	2.0	5.5	14.5
2000.....	203.6	*	*	*	*	*	0.6	1.6	4.8	14.3
1999.....	210.5	*	*	*	*	0.3	0.5	1.6	5.1	15.1

White, female

2004.....	158.1	*	*	*	*	*	*	0.7	1.6	4.7
2003.....	171.4	*	*	*	*	*	0.4	0.7	1.7	4.4
2002.....	178.4	*	*	*	*	*	*	0.5	1.6	4.5
2001.....	185.3	*	*	*	*	*	0.3	0.6	1.9	4.5
2000.....	192.4	*	*	*	*	*	0.3	0.5	1.4	4.5
1999.....	200.0	*	*	*	*	*	*	0.7	1.7	4.6

Black, both sexes

2004.....	121.0	*	*	*	*	*	1.2	2.9	6.5	12.7
2003.....	129.2	*	*	*	*	*	1.2	2.4	6.1	14.0
2002.....	132.9	*	*	*	*	*	0.9	2.1	5.6	15.1
2001.....	136.9	*	*	*	*	*	1.5	2.8	6.4	14.9
2000.....	140.1	*	*	*	*	*	0.7	2.4	6.0	14.6
1999.....	143.7	*	*	*	*	*	1.2	1.9	5.8	16.1

Black, male

2004.....	125.2	*	*	*	*	*	1.8	3.4	9.6	18.4
2003.....	133.4	*	*	*	*	*	1.6	3.1	9.2	19.6
2002.....	135.3	*	*	*	*	*	*	2.7	7.5	22.0
2001.....	138.9	*	*	*	*	*	1.9	3.9	9.3	21.6
2000.....	141.5	*	*	*	*	*	*	3.7	8.0	20.3
1999.....	146.0	*	*	*	*	*	*	2.9	8.4	23.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Ischemic heart diseases (I20-I25)

Black, female

2004.....	117.1	*	*	*	*	*	*	2.4	3.7	7.6
2003.....	125.3	*	*	*	*	*	*	1.7	3.3	9.1
2002.....	130.8	*	*	*	*	*	*	1.5	4.0	9.0
2001.....	135.2	*	*	*	*	*	*	1.7	3.8	9.0
2000.....	138.8	*	*	*	*	*	*	*	4.3	9.7
1999.....	141.7	*	*	*	*	*	*	*	3.4	9.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Acute myocardial infarction (I21-I22)

All races 3/, both sexes

2004.....	53.4	*	*	*	*	*	0.2	0.6	1.6	3.8
2003.....	58.7	*	*	*	*	*	0.3	0.5	1.5	4.1
2002.....	62.3	*	*	*	*	*	0.2	0.5	1.5	4.0
2001.....	64.9	*	*	*	*	*	0.2	0.6	1.8	4.2
2000.....	68.5	0.5	*	*	*	*	0.2	0.6	1.3	4.0
1999.....	71.5	*	*	*	*	0.1	0.2	0.5	1.6	4.1

All races 3/, male

2004.....	57.4	*	*	*	*	*	0.3	0.8	2.3	5.7
2003.....	62.6	*	*	*	*	*	0.3	0.8	2.2	6.2
2002.....	66.2	*	*	*	*	*	0.3	0.7	2.1	6.0
2001.....	68.6	*	*	*	*	*	0.3	0.7	2.5	6.2
2000.....	72.7	*	*	*	*	*	0.2	0.8	1.9	5.8
1999.....	76.1	*	*	*	*	*	0.3	0.7	2.3	6.2

All races 3/, female

2004.....	49.6	*	*	*	*	*	*	0.4	0.8	1.9
2003.....	54.8	*	*	*	*	*	0.2	0.3	0.8	2.0
2002.....	58.4	*	*	*	*	*	*	0.3	0.8	2.1
2001.....	61.3	*	*	*	*	*	*	0.4	1.0	2.1
2000.....	64.6	*	*	*	*	*	*	0.3	0.7	2.2
1999.....	67.0	*	*	*	*	*	*	0.3	0.9	2.0

White, both sexes

2004.....	57.8	*	*	*	*	*	0.1	0.5	1.6	3.8
2003.....	63.4	*	*	*	*	*	0.2	0.5	1.4	4.1
2002.....	67.3	*	*	*	*	*	0.2	0.4	1.4	4.0
2001.....	70.1	*	*	*	*	*	0.2	0.5	1.7	4.1
2000.....	74.1	*	*	*	*	*	0.2	0.5	1.2	4.0
1999.....	77.4	*	*	*	*	*	0.1	0.5	1.5	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Acute myocardial infarction (I21-I22)

White, male

2004.....	62.3	*	*	*	*	*	0.7	2.4	5.6
2003.....	67.7	*	*	*	*	0.3	0.7	2.0	6.3
2002.....	71.8	*	*	*	*	0.3	0.6	2.1	5.9
2001.....	74.4	*	*	*	*	0.3	0.6	2.5	6.1
2000.....	78.8	*	*	*	*	*	0.6	1.8	6.0
1999.....	82.7	*	*	*	*	*	0.6	2.1	6.2

White, female

2004.....	53.3	*	*	*	*	*	0.4	0.7	1.9
2003.....	59.2	*	*	*	*	*	*	0.8	1.9
2002.....	62.8	*	*	*	*	*	*	0.7	1.9
2001.....	66.0	*	*	*	*	*	0.3	0.9	2.0
2000.....	69.6	*	*	*	*	*	0.3	0.6	2.0
1999.....	72.3	*	*	*	*	*	0.3	0.9	1.9

Black, both sexes

2004.....	42.0	*	*	*	*	*	1.2	2.2	4.9
2003.....	45.7	*	*	*	*	*	0.9	2.3	5.6
2002.....	48.5	*	*	*	*	*	0.9	2.3	5.5
2001.....	50.0	*	*	*	*	*	1.1	2.5	5.8
2000.....	52.2	*	*	*	*	*	1.2	2.3	4.9
1999.....	53.3	*	*	*	*	*	0.8	2.4	5.5

Black, male

2004.....	42.4	*	*	*	*	*	*	2.7	7.7
2003.....	46.6	*	*	*	*	*	*	3.5	7.5
2002.....	48.3	*	*	*	*	*	*	3.0	7.6
2001.....	49.2	*	*	*	*	*	*	3.4	8.0
2000.....	52.0	*	*	*	*	*	1.6	2.7	6.5
1999.....	53.4	*	*	*	*	*	*	3.4	7.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Acute myocardial infarction (I21-I22)

Black, female

2004.....	41.6	*	*	*	*	*	*	*	1.8	2.4
2003.....	44.9	*	*	*	*	*	*	*	*	3.9
2002.....	48.8	*	*	*	*	*	*	*	1.6	3.6
2001.....	50.6	*	*	*	*	*	*	*	1.7	3.8
2000.....	52.5	*	*	*	*	*	*	*	1.9	3.6
1999.....	53.2	*	*	*	*	*	*	*	1.4	3.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other forms of chronic ischemic heart disease (I20,I25)

All races 3/, both sexes

2004.....	99.5	*	*	*	*	*	0.3	0.9	2.3	5.5
2003.....	105.3	*	*	*	*	*	0.3	0.9	2.4	5.8
2002.....	108.0	*	*	*	*	*	0.3	0.8	2.0	6.0
2001.....	110.3	*	*	*	*	*	0.4	0.9	2.2	5.6
2000.....	113.3	*	*	*	*	*	0.3	0.6	2.0	5.7
1999.....	117.2	*	*	*	*	0.1	0.3	0.7	2.1	6.2

All races 3/, male

2004.....	103.3	*	*	*	*	*	0.4	1.3	3.6	8.1
2003.....	108.5	*	*	*	*	*	0.4	1.2	3.7	8.8
2002.....	110.9	*	*	*	*	*	0.4	1.3	3.0	9.2
2001.....	111.9	*	*	*	*	*	0.5	1.5	3.3	8.6
2000.....	114.8	*	*	*	*	*	0.4	1.1	3.1	8.7
1999.....	118.0	*	*	*	*	0.2	0.3	1.0	3.2	9.4

All races 3/, female

2004.....	95.7	*	*	*	*	*	*	0.5	0.9	3.0
2003.....	102.3	*	*	*	*	*	0.2	0.5	1.0	2.8
2002.....	105.2	*	*	*	*	*	*	0.4	1.0	2.8
2001.....	108.8	*	*	*	*	*	0.2	0.3	1.1	2.7
2000.....	111.9	*	*	*	*	*	*	0.2	1.0	2.8
1999.....	116.4	*	*	*	*	*	*	0.4	1.0	3.1

White, both sexes

2004.....	107.5	*	*	*	*	*	0.2	0.8	2.0	5.4
2003.....	113.8	*	*	*	*	*	0.3	0.8	2.2	5.6
2002.....	116.9	*	*	*	*	*	0.2	0.8	1.8	5.6
2001.....	119.1	*	*	*	*	*	0.3	0.8	2.0	5.3
2000.....	122.6	*	*	*	*	*	0.2	0.6	1.9	5.3
1999.....	126.5	*	*	*	*	*	0.2	0.7	1.9	5.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other forms of chronic ischemic heart disease (I20,I25)

White, male

2004.....	111.1	*	*	*	*	*	0.3	1.2	3.1	7.9
2003.....	116.6	*	*	*	*	*	0.4	1.1	3.5	8.5
2002.....	119.4	*	*	*	*	*	0.4	1.3	2.8	8.6
2001.....	120.1	*	*	*	*	*	0.4	1.4	3.0	8.2
2000.....	123.5	*	*	*	*	*	0.3	0.9	2.9	8.2
1999.....	126.5	*	*	*	*	*	*	0.9	3.0	8.8

White, female

2004.....	103.9	*	*	*	*	*	*	0.3	0.9	2.8
2003.....	111.0	*	*	*	*	*	*	0.4	0.9	2.5
2002.....	114.4	*	*	*	*	*	*	0.3	0.8	2.5
2001.....	118.1	*	*	*	*	*	*	*	0.9	2.4
2000.....	121.6	*	*	*	*	*	*	*	0.8	2.4
1999.....	126.5	*	*	*	*	*	*	0.4	0.9	2.7

Black, both sexes

2004.....	78.0	*	*	*	*	*	*	1.6	4.1	7.7
2003.....	82.3	*	*	*	*	*	*	1.5	3.7	8.2
2002.....	83.2	*	*	*	*	*	*	1.1	3.2	9.6
2001.....	85.8	*	*	*	*	*	0.8	1.7	3.8	8.9
2000.....	86.6	*	*	*	*	*	*	1.3	3.6	9.5
1999.....	89.3	*	*	*	*	*	0.7	1.1	3.4	10.1

Black, male

2004.....	81.5	*	*	*	*	*	*	1.9	6.7	10.5
2003.....	85.5	*	*	*	*	*	*	2.1	5.7	11.7
2002.....	85.7	*	*	*	*	*	*	*	4.1	14.4
2001.....	88.2	*	*	*	*	*	*	2.4	5.9	13.3
2000.....	88.0	*	*	*	*	*	*	2.1	5.0	13.4
1999.....	91.1	*	*	*	*	*	*	1.7	5.0	14.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other forms of chronic ischemic heart disease (I20,I25)

Black, female

2004.....	74.7	*	*	*	*	*	*	*	1.7	5.2
2003.....	79.4	*	*	*	*	*	*	*	2.0	5.0
2002.....	81.0	*	*	*	*	*	*	*	2.3	5.3
2001.....	83.5	*	*	*	*	*	*	*	1.9	4.9
2000.....	85.3	*	*	*	*	*	*	*	2.4	5.9
1999.....	87.6	*	*	*	*	*	*	*	1.9	5.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All races 3/, both sexes

2004.....	21.7	*	*	*	*	*	*	0.2	0.8	1.7
2003.....	23.1	*	*	*	*	*	*	0.3	0.8	1.9
2002.....	23.6	*	*	*	*	*	*	0.2	0.7	1.9
2001.....	23.9	*	*	*	*	*	*	0.2	0.7	1.8
2000.....	24.7	*	*	*	*	*	*	0.2	0.5	1.7
1999.....	25.8	*	*	*	*	*	*	0.2	0.6	2.0

All races 3/, male

2004.....	23.9	*	*	*	*	*	*	0.3	1.2	2.5
2003.....	25.2	*	*	*	*	*	*	0.4	1.3	2.9
2002.....	25.4	*	*	*	*	*	*	0.3	1.1	3.0
2001.....	25.3	*	*	*	*	*	*	0.3	1.0	2.8
2000.....	26.0	*	*	*	*	*	*	0.3	0.9	2.8
1999.....	26.8	*	*	*	*	*	*	0.2	0.9	3.1

All races 3/, female

2004.....	19.5	*	*	*	*	*	*	*	0.3	0.8
2003.....	21.1	*	*	*	*	*	*	*	0.2	0.9
2002.....	21.9	*	*	*	*	*	*	*	0.3	0.8
2001.....	22.5	*	*	*	*	*	*	*	0.4	0.8
2000.....	23.5	*	*	*	*	*	*	*	0.2	0.7
1999.....	24.8	*	*	*	*	*	*	*	0.3	0.9

White, both sexes

2004.....	22.1	*	*	*	*	*	*	0.2	0.7	1.6
2003.....	23.6	*	*	*	*	*	*	0.2	0.7	1.9
2002.....	24.1	*	*	*	*	*	*	0.2	0.6	1.7
2001.....	24.3	*	*	*	*	*	*	0.2	0.6	1.7
2000.....	25.1	*	*	*	*	*	*	*	0.5	1.6
1999.....	26.1	*	*	*	*	*	*	0.1	0.5	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Atherosclerotic cardiovascular disease, so described (I25.0)

White, male

2004.....	24.1	*	*	*	*	*	*	0.3	1.0	2.5
2003.....	25.3	*	*	*	*	*	*	0.3	1.3	2.8
2002.....	25.5	*	*	*	*	*	*	0.3	1.0	2.7
2001.....	25.3	*	*	*	*	*	*	0.3	0.9	2.7
2000.....	26.0	*	*	*	*	*	*	*	0.8	2.5
1999.....	26.6	*	*	*	*	*	*	*	0.8	2.9

White, female

2004.....	20.1	*	*	*	*	*	*	*	0.3	0.8
2003.....	21.9	*	*	*	*	*	*	*	*	0.9
2002.....	22.7	*	*	*	*	*	*	*	*	0.7
2001.....	23.3	*	*	*	*	*	*	*	0.3	0.7
2000.....	24.3	*	*	*	*	*	*	*	*	0.6
1999.....	25.6	*	*	*	*	*	*	*	*	0.7

Black, both sexes

2004.....	25.0	*	*	*	*	*	*	*	1.5	2.0
2003.....	26.0	*	*	*	*	*	*	*	1.3	2.6
2002.....	27.0	*	*	*	*	*	*	*	1.3	3.5
2001.....	27.7	*	*	*	*	*	*	*	1.0	3.0
2000.....	28.5	*	*	*	*	*	*	*	0.9	2.9
1999.....	29.9	*	*	*	*	*	*	*	1.2	3.7

Black, male

2004.....	28.9	*	*	*	*	*	*	*	2.6	3.0
2003.....	30.5	*	*	*	*	*	*	*	2.3	3.9
2002.....	31.1	*	*	*	*	*	*	*	1.9	5.4
2001.....	31.5	*	*	*	*	*	*	*	*	4.6
2000.....	32.3	*	*	*	*	*	*	*	*	4.7
1999.....	33.8	*	*	*	*	*	*	*	1.7	5.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races 3/, both sexes

2004.....	77.8	*	*	*	*	*	0.2	0.7	1.5	3.9
2003.....	82.2	*	*	*	*	*	0.3	0.6	1.6	3.9
2002.....	84.4	*	*	*	*	*	0.2	0.6	1.3	4.1
2001.....	86.4	*	*	*	*	*	0.3	0.7	1.5	3.8
2000.....	88.6	*	*	*	*	*	0.2	0.5	1.5	4.0
1999.....	91.4	*	*	*	*	0.1	0.2	0.6	1.5	4.3

All races 3/, male

2004.....	79.4	*	*	*	*	*	0.3	1.0	2.3	5.6
2003.....	83.3	*	*	*	*	*	0.3	0.8	2.3	5.9
2002.....	85.5	*	*	*	*	*	0.3	0.9	1.9	6.2
2001.....	86.5	*	*	*	*	*	0.4	1.1	2.3	5.8
2000.....	88.8	*	*	*	*	*	0.2	0.8	2.2	5.9
1999.....	91.2	*	*	*	*	0.2	0.2	0.8	2.2	6.3

All races 3/, female

2004.....	76.2	*	*	*	*	*	*	0.4	0.7	2.1
2003.....	81.1	*	*	*	*	*	*	0.4	0.8	1.9
2002.....	83.3	*	*	*	*	*	*	0.3	0.8	2.0
2001.....	86.2	*	*	*	*	*	*	0.2	0.7	1.9
2000.....	88.4	*	*	*	*	*	*	*	0.8	2.1
1999.....	91.6	*	*	*	*	*	*	0.3	0.7	2.2

White, both sexes

2004.....	85.4	*	*	*	*	*	0.2	0.6	1.4	3.7
2003.....	90.2	*	*	*	*	*	0.2	0.5	1.5	3.7
2002.....	92.8	*	*	*	*	*	0.2	0.6	1.2	3.9
2001.....	94.8	*	*	*	*	*	0.2	0.6	1.4	3.6
2000.....	97.4	*	*	*	*	*	0.1	0.4	1.4	3.7
1999.....	100.4	*	*	*	*	*	0.1	0.5	1.4	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

White, male

2004.....	87.0	*	*	*	*	*	0.2	0.9	2.1	5.5
2003.....	91.3	*	*	*	*	*	0.3	0.7	2.2	5.7
2002.....	93.9	*	*	*	*	*	0.3	0.9	1.8	5.9
2001.....	94.8	*	*	*	*	*	0.4	1.1	2.1	5.5
2000.....	97.5	*	*	*	*	*	*	0.7	2.1	5.6
1999.....	99.9	*	*	*	*	*	*	0.7	2.1	6.0

White, female

2004.....	83.8	*	*	*	*	*	*	0.3	0.6	2.0
2003.....	89.1	*	*	*	*	*	*	0.3	0.7	1.6
2002.....	91.7	*	*	*	*	*	*	*	0.6	1.9
2001.....	94.9	*	*	*	*	*	*	*	0.6	1.7
2000.....	97.3	*	*	*	*	*	*	*	0.6	1.8
1999.....	100.8	*	*	*	*	*	*	0.3	0.6	2.0

Black, both sexes

2004.....	52.9	*	*	*	*	*	*	1.3	2.6	5.7
2003.....	56.3	*	*	*	*	*	*	1.0	2.5	5.6
2002.....	56.2	*	*	*	*	*	*	0.9	1.9	6.1
2001.....	58.0	*	*	*	*	*	*	1.2	2.8	5.8
2000.....	58.1	*	*	*	*	*	*	0.8	2.8	6.5
1999.....	59.3	*	*	*	*	*	*	0.8	2.2	6.4

Black, male

2004.....	52.6	*	*	*	*	*	*	1.6	4.1	7.5
2003.....	55.0	*	*	*	*	*	*	*	3.4	7.8
2002.....	54.5	*	*	*	*	*	*	*	2.3	9.0
2001.....	56.7	*	*	*	*	*	*	1.8	4.6	8.7
2000.....	55.8	*	*	*	*	*	*	*	3.8	8.7
1999.....	57.3	*	*	*	*	*	*	*	3.3	9.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other heart diseases (I26-I51)

All races 3/, both sexes

2004.....	56.6	10.0	1.1	0.4	0.7	1.5	2.5	3.6	5.0	7.8
2003.....	58.5	10.4	1.1	0.5	0.7	1.7	2.7	3.6	5.4	7.8
2002.....	58.8	11.7	1.0	0.4	0.7	1.8	2.3	3.8	5.2	8.1
2001.....	58.6	11.1	1.4	0.5	0.7	1.5	2.5	3.6	5.1	8.0
2000.....	58.8	12.0	1.1	0.5	0.7	1.8	2.6	3.5	5.2	7.5
1999.....	59.4	12.9	1.2	0.5	0.7	2.0	2.6	3.5	5.3	8.2

All races 3/, male

2004.....	50.6	10.5	1.1	0.3	0.8	1.9	3.2	4.5	6.1	9.4
2003.....	52.3	11.4	1.0	0.5	0.8	2.1	3.2	4.1	6.5	9.5
2002.....	52.4	12.1	1.0	0.4	0.8	2.4	3.0	4.6	6.6	9.6
2001.....	51.8	10.8	1.5	0.4	0.8	1.9	3.1	4.3	6.0	9.9
2000.....	52.2	12.2	1.2	0.6	0.9	2.1	3.1	4.2	6.2	9.1
1999.....	53.1	13.1	1.3	0.5	0.8	2.4	3.0	4.0	6.4	10.3

All races 3/, female

2004.....	62.4	9.4	1.1	0.5	0.6	1.2	1.8	2.6	3.9	6.1
2003.....	64.5	9.3	1.2	0.4	0.5	1.3	2.1	3.1	4.3	6.1
2002.....	65.0	11.4	1.0	0.5	0.5	1.1	1.7	3.0	3.8	6.6
2001.....	65.2	11.5	1.3	0.5	0.7	1.1	1.9	2.9	4.1	6.0
2000.....	65.1	11.8	0.9	0.4	0.6	1.4	2.1	2.9	4.1	5.9
1999.....	65.5	12.7	1.0	0.4	0.7	1.5	2.2	2.9	4.2	6.1

White, both sexes

2004.....	60.4	8.7	0.9	0.3	0.6	1.3	2.2	2.8	4.0	6.5
2003.....	62.2	8.2	0.9	0.4	0.6	1.5	2.2	2.8	4.4	6.4
2002.....	62.6	10.0	0.9	0.4	0.6	1.5	1.9	3.0	4.1	6.6
2001.....	62.3	9.3	1.1	0.4	0.7	1.3	2.0	2.9	4.1	6.5
2000.....	62.5	10.6	1.0	0.4	0.6	1.4	2.1	2.8	4.3	6.1
1999.....	62.9	11.0	1.0	0.4	0.6	1.8	2.1	2.7	4.4	6.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other heart diseases (I26-I51)

White, male

2004.....	53.0	9.1	0.9	*	0.7	1.6	2.9	3.6	5.0	8.0
2003.....	54.7	8.8	0.8	0.4	0.8	1.8	2.5	3.2	5.5	7.9
2002.....	54.9	10.5	1.0	0.4	0.7	2.1	2.5	3.6	5.2	7.9
2001.....	54.1	9.6	1.2	0.4	0.7	1.5	2.4	3.5	4.8	8.1
2000.....	54.6	10.8	1.2	0.5	0.8	1.7	2.5	3.4	5.2	7.5
1999.....	55.1	11.4	1.1	0.5	0.7	2.1	2.4	3.3	5.4	8.5

White, female

2004.....	67.6	8.3	0.8	0.4	0.5	1.0	1.5	2.0	3.1	4.9
2003.....	69.6	7.6	1.0	0.3	0.4	1.1	1.8	2.4	3.2	5.0
2002.....	70.2	9.6	0.8	0.4	0.5	1.0	1.3	2.3	2.9	5.3
2001.....	70.4	8.9	1.1	0.5	0.6	1.0	1.5	2.3	3.4	4.8
2000.....	70.2	10.4	0.7	0.3	0.4	1.2	1.6	2.2	3.3	4.6
1999.....	70.5	10.7	0.8	0.4	0.6	1.4	1.9	2.2	3.4	4.8

Black, both sexes

2004.....	50.8	14.8	2.3	0.8	1.3	3.0	4.5	8.0	11.9	17.9
2003.....	52.5	19.6	2.0	0.9	1.0	3.0	5.9	8.9	12.8	18.1
2002.....	52.5	20.2	1.5	*	1.2	3.1	5.1	9.5	13.0	18.9
2001.....	52.8	20.1	2.9	0.8	1.1	2.9	5.5	8.7	12.3	18.4
2000.....	52.7	19.0	1.9	0.8	1.4	3.6	6.0	8.8	11.9	17.8
1999.....	54.4	22.3	2.0	0.7	1.2	3.5	5.6	8.6	12.4	19.5

Black, male

2004.....	50.4	15.7	2.3	*	1.3	4.0	5.6	10.4	13.9	21.5
2003.....	51.5	23.2	2.2	*	1.2	3.9	7.5	10.5	14.3	22.5
2002.....	50.8	20.9	*	*	1.3	4.0	6.2	11.9	16.4	22.8
2001.....	51.7	17.7	3.2	*	1.2	3.8	7.0	10.7	14.9	23.1
2000.....	51.5	18.9	1.9	*	1.4	4.3	6.7	10.1	14.3	21.6
1999.....	54.3	21.8	2.1	*	1.5	4.8	6.8	10.2	15.2	24.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other heart diseases (I26-I51)

Black, female

2004.....	51.1	13.9	2.3	*	1.3	2.0	3.4	5.8	10.1	14.7
2003.....	53.4	15.8	1.8	*	*	2.1	4.2	7.3	11.4	14.3
2002.....	54.0	19.4	1.8	*	*	2.2	4.0	7.4	9.8	15.5
2001.....	53.8	22.7	2.5	*	*	2.0	4.1	6.9	10.0	14.1
2000.....	53.9	19.2	2.0	*	1.3	2.9	5.3	7.6	9.8	14.5
1999.....	54.5	22.8	1.8	*	*	2.2	4.5	7.1	9.9	15.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)

All races 3/, both sexes

2004.....	4.5	3.8	0.2	*	*	0.2	0.4	0.7	0.9	1.4
2003.....	4.8	3.3	0.2	*	*	0.2	0.4	0.7	1.1	1.5
2002.....	4.8	4.6	0.2	*	0.1	0.2	0.3	0.7	0.9	1.4
2001.....	4.7	4.2	0.1	*	*	0.2	0.4	0.8	1.0	1.4
2000.....	4.6	4.9	0.2	*	*	0.2	0.4	0.6	1.1	1.4
1999.....	4.7	5.7	*	0.1	*	0.3	0.5	0.7	1.0	1.5

All races 3/, male

2004.....	3.6	4.1	*	*	*	*	0.3	0.5	0.7	1.1
2003.....	3.9	3.7	*	*	*	0.2	0.3	0.4	0.8	1.3
2002.....	3.8	4.7	*	*	*	*	0.2	0.7	0.7	1.3
2001.....	3.8	3.9	*	*	*	*	0.3	0.6	0.7	1.1
2000.....	3.8	5.1	*	*	*	*	0.3	0.5	0.9	1.2
1999.....	3.9	5.9	*	*	*	0.3	0.3	0.5	0.8	1.3

All races 3/, female

2004.....	5.3	3.5	*	*	*	0.3	0.5	0.8	1.1	1.8
2003.....	5.7	2.9	*	*	*	0.3	0.6	0.9	1.3	1.8
2002.....	5.7	4.5	*	*	*	0.2	0.5	0.8	1.1	1.6
2001.....	5.5	4.6	*	*	*	*	0.6	0.9	1.3	1.7
2000.....	5.5	4.6	*	*	*	0.3	0.6	0.8	1.2	1.6
1999.....	5.5	5.6	*	*	*	0.3	0.7	0.9	1.2	1.7

White, both sexes

2004.....	4.5	3.2	*	*	*	0.2	0.3	0.5	0.7	1.2
2003.....	4.8	2.5	*	*	*	0.2	0.3	0.5	0.9	1.2
2002.....	4.8	3.9	*	*	*	0.2	0.3	0.5	0.7	1.3
2001.....	4.7	3.6	*	*	*	0.1	0.4	0.6	0.8	1.1
2000.....	4.7	4.1	0.2	*	*	0.2	0.4	0.5	0.9	1.2
1999.....	4.7	5.2	*	*	*	0.3	0.4	0.5	0.8	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)

White, male

2004.....	3.6	3.4	*	*	*	*	*	0.4	0.6	0.8
2003.....	3.9	2.9	*	*	*	*	*	0.3	0.7	1.0
2002.....	3.9	4.1	*	*	*	*	*	0.5	0.6	1.1
2001.....	3.8	3.4	*	*	*	*	0.3	0.5	0.6	0.9
2000.....	3.8	4.3	*	*	*	*	0.3	0.5	0.7	1.1
1999.....	3.9	5.4	*	*	*	0.2	*	0.4	0.6	1.2

White, female

2004.....	5.4	3.0	*	*	*	0.3	0.4	0.6	0.9	1.6
2003.....	5.7	2.0	*	*	*	0.3	0.5	0.8	1.1	1.4
2002.....	5.8	3.7	*	*	*	*	0.4	0.5	0.8	1.4
2001.....	5.6	3.9	*	*	*	*	0.5	0.7	1.1	1.4
2000.....	5.5	3.9	*	*	*	0.3	0.4	0.6	1.0	1.3
1999.....	5.5	4.9	*	*	*	0.4	0.6	0.6	1.0	1.3

Black, both sexes

2004.....	5.6	5.9	*	*	*	*	1.1	1.6	2.2	3.4
2003.....	6.1	6.5	*	*	*	*	1.2	1.4	2.5	3.8
2002.....	5.9	8.2	*	*	*	*	0.8	2.2	2.5	3.1
2001.....	5.8	6.8	*	*	*	*	0.8	1.9	2.3	3.2
2000.....	5.7	8.0	*	*	*	*	1.0	1.6	2.5	3.1
1999.....	6.2	9.4	*	*	*	*	1.0	1.8	2.6	3.3

Black, male

2004.....	4.6	6.5	*	*	*	*	*	1.6	1.5	3.3
2003.....	5.2	7.1	*	*	*	*	*	*	2.0	3.4
2002.....	4.5	9.0	*	*	*	*	*	2.2	1.8	2.9
2001.....	4.8	6.9	*	*	*	*	*	*	1.7	2.8
2000.....	4.6	7.7	*	*	*	*	*	*	2.5	2.5
1999.....	5.1	8.8	*	*	*	*	*	*	2.5	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)

Black, female

2004.....	6.5	*	*	*	*	*	*	1.5	2.9	3.5
2003.....	6.9	*	*	*	*	*	1.3	1.6	2.9	4.2
2002.....	7.2	7.3	*	*	*	*	*	2.2	3.1	3.3
2001.....	6.8	6.6	*	*	*	*	*	2.4	2.9	3.5
2000.....	6.8	8.3	*	*	*	*	1.5	2.2	2.6	3.6
1999.....	7.2	10.1	*	*	*	*	*	2.3	2.7	4.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pulmonary embolism (I26)

All races 3/, both sexes

2004.....	2.8	*	*	*	*	0.1	0.3	0.4	0.6	1.0
2003.....	3.0	*	*	*	*	0.1	0.3	0.4	0.7	1.1
2002.....	3.0	*	*	*	*	0.1	0.3	0.5	0.6	1.0
2001.....	3.0	*	*	*	*	*	0.3	0.5	0.7	0.9
2000.....	3.0	*	*	*	*	0.1	0.3	0.4	0.7	1.0
1999.....	3.2	*	*	*	*	0.2	0.3	0.4	0.7	0.9

All races 3/, male

2004.....	2.3	*	*	*	*	*	0.2	0.4	0.5	0.7
2003.....	2.6	*	*	*	*	*	0.2	0.3	0.6	0.9
2002.....	2.5	*	*	*	*	*	*	0.5	0.5	0.9
2001.....	2.6	*	*	*	*	*	0.2	0.4	0.5	0.8
2000.....	2.6	*	*	*	*	*	*	0.3	0.6	0.8
1999.....	2.7	*	*	*	*	*	0.3	0.3	0.6	0.9

All races 3/, female

2004.....	3.2	*	*	*	*	*	0.3	0.5	0.8	1.3
2003.....	3.4	*	*	*	*	*	0.4	0.5	0.8	1.2
2002.....	3.5	*	*	*	*	*	0.3	0.5	0.7	1.1
2001.....	3.5	*	*	*	*	*	0.4	0.6	0.9	1.1
2000.....	3.4	*	*	*	*	*	0.4	0.5	0.7	1.1
1999.....	3.7	*	*	*	*	*	0.5	0.6	0.8	1.0

White, both sexes

2004.....	2.8	*	*	*	*	*	0.2	0.4	0.6	0.8
2003.....	3.0	*	*	*	*	*	0.2	0.3	0.6	0.8
2002.....	3.0	*	*	*	*	*	0.2	0.3	0.5	0.9
2001.....	3.1	*	*	*	*	*	0.3	0.4	0.6	0.7
2000.....	3.0	*	*	*	*	*	0.2	0.4	0.6	0.8
1999.....	3.2	*	*	*	*	0.1	0.3	0.3	0.5	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pulmonary embolism (I26)

White, male

2004.....	2.3	*	*	*	*	*	*	0.3	0.5	0.5
2003.....	2.6	*	*	*	*	*	*	*	0.5	0.7
2002.....	2.5	*	*	*	*	*	*	0.3	0.4	0.7
2001.....	2.6	*	*	*	*	*	*	0.3	0.4	0.6
2000.....	2.6	*	*	*	*	*	*	0.4	0.5	0.7
1999.....	2.7	*	*	*	*	*	*	*	0.5	0.8

White, female

2004.....	3.2	*	*	*	*	*	0.3	0.4	0.7	1.2
2003.....	3.4	*	*	*	*	*	0.3	0.5	0.6	0.9
2002.....	3.5	*	*	*	*	*	0.3	0.4	0.6	1.0
2001.....	3.6	*	*	*	*	*	0.3	0.4	0.8	0.8
2000.....	3.4	*	*	*	*	*	0.3	0.3	0.6	0.8
1999.....	3.7	*	*	*	*	*	0.4	0.4	0.6	0.8

Black, both sexes

2004.....	3.6	*	*	*	*	*	*	0.9	1.4	2.4
2003.....	4.0	*	*	*	*	*	1.0	1.0	1.9	2.9
2002.....	3.8	*	*	*	*	*	*	1.6	1.5	2.1
2001.....	3.8	*	*	*	*	*	*	1.4	1.7	2.4
2000.....	3.9	*	*	*	*	*	0.9	0.9	1.6	2.3
1999.....	4.3	*	*	*	*	*	0.8	1.3	2.0	2.0

Black, male

2004.....	3.2	*	*	*	*	*	*	*	*	2.4
2003.....	3.6	*	*	*	*	*	*	*	1.6	2.7
2002.....	3.0	*	*	*	*	*	*	1.6	*	2.0
2001.....	3.3	*	*	*	*	*	*	*	*	2.1
2000.....	3.3	*	*	*	*	*	*	*	1.5	2.0
1999.....	3.8	*	*	*	*	*	*	*	2.0	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pulmonary embolism (I26)

Black, female

2004.....	3.9	*	*	*	*	*	*	*	1.6	2.4
2003.....	4.3	*	*	*	*	*	*	*	2.2	3.1
2002.....	4.4	*	*	*	*	*	*	1.5	1.6	2.2
2001.....	4.3	*	*	*	*	*	*	1.7	2.0	2.7
2000.....	4.4	*	*	*	*	*	*	1.4	1.6	2.6
1999.....	4.8	*	*	*	*	*	*	1.5	2.1	2.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

All races 3/, both sexes

2004.....	1.7	3.7	0.1	*	*	0.1	0.1	0.2	0.3	0.4
2003.....	1.8	3.2	0.1	*	*	0.1	0.1	0.2	0.4	0.5
2002.....	1.8	4.6	0.2	*	*	*	*	0.2	0.3	0.4
2001.....	1.6	4.2	0.1	*	*	*	0.2	0.3	0.3	0.5
2000.....	1.6	4.7	0.2	*	*	0.1	0.1	0.2	0.4	0.5
1999.....	1.5	5.7	*	0.1	*	0.1	0.2	0.2	0.3	0.5

All races 3/, male

2004.....	1.2	4.0	*	*	*	*	*	*	*	0.4
2003.....	1.3	3.6	*	*	*	*	*	*	0.2	0.4
2002.....	1.3	4.7	*	*	*	*	*	*	0.2	0.4
2001.....	1.2	3.9	*	*	*	*	*	*	0.2	0.3
2000.....	1.2	4.9	*	*	*	*	*	*	0.3	0.4
1999.....	1.2	5.9	*	*	*	*	*	*	*	0.4

All races 3/, female

2004.....	2.2	3.5	*	*	*	*	*	0.3	0.4	0.5
2003.....	2.3	2.8	*	*	*	0.2	*	0.4	0.5	0.6
2002.....	2.2	4.4	*	*	*	*	*	0.3	0.4	0.5
2001.....	2.0	4.5	*	*	*	*	0.2	0.4	0.4	0.6
2000.....	2.1	4.5	*	*	*	*	*	0.3	0.5	0.5
1999.....	1.8	5.5	*	*	*	*	0.3	0.3	0.5	0.7

White, both sexes

2004.....	1.7	3.2	*	*	*	*	*	0.2	0.2	0.4
2003.....	1.8	2.5	*	*	*	0.1	*	0.2	0.3	0.4
2002.....	1.8	3.8	*	*	*	*	*	0.1	0.2	0.4
2001.....	1.7	3.6	*	*	*	*	*	0.2	0.3	0.4
2000.....	1.7	4.0	0.2	*	*	*	*	0.2	0.3	0.4
1999.....	1.5	5.1	*	*	*	0.2	0.2	0.2	0.3	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

White, male

2004.....	1.2	3.3	*	*	*	*	*	*	0.3
2003.....	1.4	2.9	*	*	*	*	*	*	0.3
2002.....	1.3	4.0	*	*	*	*	*	*	0.3
2001.....	1.2	3.4	*	*	*	*	*	*	0.3
2000.....	1.2	4.2	*	*	*	*	*	*	0.4
1999.....	1.2	5.4	*	*	*	*	*	*	0.4

White, female

2004.....	2.2	3.0	*	*	*	*	*	*	0.4
2003.....	2.3	2.0	*	*	*	*	0.3	0.4	0.5
2002.....	2.2	3.6	*	*	*	*	*	0.2	0.4
2001.....	2.1	3.8	*	*	*	*	0.3	0.4	0.6
2000.....	2.1	3.8	*	*	*	*	*	0.4	0.4
1999.....	1.8	4.8	*	*	*	*	*	0.4	0.5

Black, both sexes

2004.....	2.0	5.7	*	*	*	*	*	0.8	1.0
2003.....	2.1	6.2	*	*	*	*	*	*	0.9
2002.....	2.2	8.2	*	*	*	*	*	1.0	1.1
2001.....	2.0	6.8	*	*	*	*	*	*	0.8
2000.....	1.9	7.6	*	*	*	*	*	1.0	0.8
1999.....	1.9	9.3	*	*	*	*	*	*	1.3

Black, male

2004.....	1.4	6.5	*	*	*	*	*	*	*
2003.....	1.5	6.8	*	*	*	*	*	*	*
2002.....	1.5	9.0	*	*	*	*	*	*	*
2001.....	1.5	6.9	*	*	*	*	*	*	*
2000.....	1.3	7.4	*	*	*	*	*	*	*
1999.....	1.3	8.8	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other non-pulmonary forms of heart disease (I30-I51)

All races 3/, both sexes

2004.....	52.2	6.2	0.9	0.3	0.6	1.3	2.2	2.9	4.1	6.3
2003.....	53.6	7.1	0.9	0.4	0.6	1.5	2.2	3.0	4.3	6.3
2002.....	54.0	7.1	0.8	0.4	0.6	1.6	2.0	3.1	4.3	6.6
2001.....	53.9	6.9	1.3	0.4	0.6	1.3	2.0	2.9	4.1	6.6
2000.....	54.1	7.1	0.9	0.4	0.7	1.5	2.2	2.9	4.1	6.1
1999.....	54.7	7.2	1.0	0.4	0.7	1.7	2.1	2.8	4.3	6.7

All races 3/, male

2004.....	47.1	6.4	1.0	0.2	0.7	1.7	2.9	3.9	5.4	8.3
2003.....	48.3	7.7	0.8	0.4	0.8	1.9	2.9	3.7	5.7	8.2
2002.....	48.6	7.3	0.9	0.3	0.7	2.2	2.7	4.0	5.9	8.3
2001.....	48.0	6.9	1.3	0.3	0.7	1.7	2.7	3.7	5.3	8.8
2000.....	48.5	7.1	1.0	0.4	0.8	2.0	2.8	3.6	5.3	7.9
1999.....	49.1	7.3	1.1	0.4	0.8	2.2	2.6	3.6	5.6	9.0

All races 3/, female

2004.....	57.1	5.9	0.9	0.4	0.5	0.9	1.3	1.8	2.8	4.4
2003.....	58.8	6.4	1.0	0.3	0.4	1.0	1.5	2.2	3.0	4.4
2002.....	59.3	6.9	0.8	0.4	0.4	0.9	1.2	2.2	2.7	4.9
2001.....	59.7	7.0	1.2	0.5	0.6	1.0	1.3	2.0	2.8	4.4
2000.....	59.6	7.2	0.7	0.3	0.5	1.0	1.5	2.1	2.9	4.3
1999.....	60.0	7.1	0.9	0.4	0.6	1.2	1.5	2.0	3.0	4.5

White, both sexes

2004.....	55.9	5.4	0.7	0.3	0.5	1.1	2.0	2.3	3.3	5.3
2003.....	57.4	5.7	0.8	0.3	0.5	1.2	1.9	2.3	3.5	5.2
2002.....	57.8	6.2	0.8	0.3	0.5	1.4	1.6	2.5	3.4	5.4
2001.....	57.6	5.7	1.0	0.4	0.6	1.1	1.6	2.3	3.3	5.3
2000.....	57.8	6.5	0.8	0.3	0.5	1.2	1.7	2.3	3.4	4.9
1999.....	58.1	5.9	0.9	0.3	0.6	1.5	1.7	2.2	3.6	5.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other non-pulmonary forms of heart disease (I30-I51)

White, male

2004.....	49.4	5.7	0.8	*	0.6	1.4	2.7	3.2	4.4	7.1
2003.....	50.8	5.8	0.7	0.4	0.7	1.6	2.4	3.0	4.8	6.8
2002.....	51.1	6.4	0.9	0.3	0.6	1.9	2.3	3.2	4.6	6.9
2001.....	50.3	6.3	1.0	0.3	0.6	1.4	2.1	3.0	4.3	7.2
2000.....	50.8	6.4	1.0	0.4	0.7	1.6	2.2	2.9	4.5	6.4
1999.....	51.1	6.0	1.0	0.4	0.6	1.8	2.2	2.9	4.8	7.4

White, female

2004.....	62.2	5.2	0.7	0.3	0.4	0.7	1.2	1.4	2.2	3.3
2003.....	63.9	5.6	0.9	0.3	0.3	0.8	1.3	1.5	2.2	3.6
2002.....	64.4	5.9	0.7	0.4	0.4	0.8	0.9	1.8	2.1	3.9
2001.....	64.8	5.0	1.0	0.4	0.5	0.8	1.0	1.6	2.2	3.4
2000.....	64.6	6.6	0.6	0.3	0.4	0.9	1.1	1.6	2.3	3.3
1999.....	64.9	5.8	0.8	0.3	0.5	1.1	1.2	1.6	2.4	3.5

Black, both sexes

2004.....	45.2	8.9	2.1	0.8	1.2	2.7	3.4	6.5	9.7	14.5
2003.....	46.4	13.0	1.7	0.7	1.0	2.7	4.7	7.5	10.3	14.3
2002.....	46.6	12.0	1.1	*	1.1	2.7	4.3	7.3	10.5	15.8
2001.....	47.0	13.4	2.7	0.7	1.0	2.5	4.7	6.8	10.0	15.2
2000.....	47.0	11.1	1.6	0.6	1.3	3.2	4.9	7.2	9.4	14.7
1999.....	48.2	12.9	1.8	0.7	1.2	3.2	4.6	6.8	9.8	16.2

Black, male

2004.....	45.8	9.2	2.2	*	1.2	3.8	4.6	8.8	12.4	18.1
2003.....	46.4	16.1	1.8	*	1.2	3.6	6.5	9.4	12.2	19.0
2002.....	46.3	11.9	*	*	1.3	3.7	5.6	9.7	14.6	19.9
2001.....	46.9	10.8	3.0	*	*	3.3	6.4	9.3	13.2	20.4
2000.....	46.9	11.2	*	*	1.4	4.1	6.2	9.2	11.8	19.1
1999.....	49.2	13.0	2.0	*	1.4	4.5	5.9	9.0	12.7	21.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other non-pulmonary forms of heart disease (I30-I51)

Black, female

2004.....	44.6	8.6	2.0	*	1.2	1.6	2.2	4.3	7.2	11.3
2003.....	46.5	9.9	1.6	*	*	1.8	3.0	5.7	8.5	10.1
2002.....	46.8	12.1	*	*	*	1.7	3.0	5.1	6.8	12.2
2001.....	47.1	16.1	2.3	*	*	1.8	3.1	4.5	7.1	10.6
2000.....	47.1	10.9	1.6	*	1.2	2.1	3.8	5.4	7.2	10.8
1999.....	47.3	12.8	1.6	*	*	1.8	3.4	4.8	7.2	11.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races 3/, both sexes

2004.....	0.3	0.5	0.2	*	*	*	*	*	0.1	0.1
2003.....	0.3	0.8	0.1	*	*	0.1	*	0.1	0.1	0.1
2002.....	0.3	0.6	0.1	*	*	*	*	*	0.1	0.2
2001.....	0.3	0.8	0.2	*	*	*	0.1	0.1	0.1	0.2
2000.....	0.3	0.5	0.2	*	*	*	0.1	0.1	0.1	0.2
1999.....	0.3	0.7	0.2	*	*	*	0.1	*	0.2	0.2

All races 3/, male

2004.....	0.3	*	*	*	*	*	*	*	*	0.2
2003.....	0.3	*	*	*	*	*	0.2	*	*	0.2
2002.....	0.3	*	*	*	*	*	*	*	0.2	*
2001.....	0.3	*	*	*	*	*	*	*	*	0.2
2000.....	0.3	*	*	*	*	*	*	*	*	0.3
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.3

All races 3/, female

2004.....	0.3	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	0.2

White, both sexes

2004.....	0.2	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	0.1	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	0.1
2001.....	0.3	*	*	*	*	*	*	*	0.1	0.2
2000.....	0.3	*	*	*	*	*	*	0.1	*	0.1
1999.....	0.3	*	*	*	*	*	*	*	0.1	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of endocardium (I34-I38)

All races 3/, both sexes

2004.....	6.9	*	*	*	*	0.1	0.1	0.2	0.4	0.5
2003.....	6.9	*	*	*	*	*	0.1	0.2	0.4	0.5
2002.....	6.9	*	*	*	*	*	0.1	0.3	0.3	0.6
2001.....	6.9	*	*	*	*	*	0.1	0.3	0.3	0.6
2000.....	7.0	*	*	*	*	*	0.2	0.2	0.3	0.6
1999.....	7.0	*	*	*	*	0.1	0.2	0.2	0.5	0.6

All races 3/, male

2004.....	5.5	*	*	*	*	*	*	*	0.4	0.6
2003.....	5.4	*	*	*	*	*	0.2	*	0.4	0.6
2002.....	5.4	*	*	*	*	*	*	0.2	0.4	0.7
2001.....	5.4	*	*	*	*	*	*	0.3	0.3	0.7
2000.....	5.5	*	*	*	*	*	*	0.3	0.3	0.7
1999.....	5.5	*	*	*	*	*	0.2	*	0.5	0.8

All races 3/, female

2004.....	8.3	*	*	*	*	*	*	*	0.3	0.5
2003.....	8.3	*	*	*	*	*	*	0.3	0.4	0.4
2002.....	8.4	*	*	*	*	*	*	0.4	0.2	0.5
2001.....	8.2	*	*	*	*	*	*	0.2	0.3	0.5
2000.....	8.5	*	*	*	*	*	*	0.2	0.4	0.6
1999.....	8.5	*	*	*	*	*	*	0.3	0.5	0.5

White, both sexes

2004.....	7.8	*	*	*	*	*	0.1	0.1	0.4	0.5
2003.....	7.8	*	*	*	*	*	*	0.2	0.3	0.5
2002.....	7.9	*	*	*	*	*	*	0.3	0.3	0.5
2001.....	7.8	*	*	*	*	*	*	0.2	0.3	0.5
2000.....	7.9	*	*	*	*	*	*	0.2	0.3	0.5
1999.....	7.9	*	*	*	*	0.1	0.2	0.2	0.5	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Nonrheumatic mitral valve disorders (I34)

All races 3/, both sexes

2004.....	0.9	*	*	*	*	*	*	*	0.1	0.2
2003.....	0.9	*	*	*	*	*	*	*	0.1	0.1
2002.....	1.0	*	*	*	*	*	*	*	0.1	0.2
2001.....	1.0	*	*	*	*	*	*	0.1	0.1	0.2
2000.....	1.0	*	*	*	*	*	*	*	0.1	0.2
1999.....	1.0	*	*	*	*	*	*	*	0.2	0.2

All races 3/, male

2004.....	0.6	*	*	*	*	*	*	*	*	*
2003.....	0.6	*	*	*	*	*	*	*	*	*
2002.....	0.7	*	*	*	*	*	*	*	*	*
2001.....	0.7	*	*	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*	*	*
1999.....	0.7	*	*	*	*	*	*	*	*	0.2

All races 3/, female

2004.....	1.1	*	*	*	*	*	*	*	*	0.2
2003.....	1.1	*	*	*	*	*	*	*	*	*
2002.....	1.2	*	*	*	*	*	*	*	*	0.2
2001.....	1.3	*	*	*	*	*	*	*	*	0.2
2000.....	1.3	*	*	*	*	*	*	*	0.2	0.2
1999.....	1.3	*	*	*	*	*	*	*	0.2	*

White, both sexes

2004.....	1.0	*	*	*	*	*	*	*	0.1	0.2
2003.....	1.0	*	*	*	*	*	*	*	*	0.1
2002.....	1.1	*	*	*	*	*	*	*	*	0.2
2001.....	1.1	*	*	*	*	*	*	0.1	0.2	0.2
2000.....	1.1	*	*	*	*	*	*	*	0.1	0.2
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases of endocardium (I36-I38)

All races 3/, both sexes

2004.....	1.7	*	*	*	*	*	*	*	0.1	0.2
2003.....	1.7	*	*	*	*	*	*	0.1	0.1	0.2
2002.....	1.7	*	*	*	*	*	*	0.1	0.1	0.2
2001.....	1.6	*	*	*	*	*	*	*	*	0.2
2000.....	1.6	*	*	*	*	*	*	*	0.1	0.3
1999.....	1.6	*	*	*	*	*	*	*	0.2	0.2

All races 3/, male

2004.....	1.4	*	*	*	*	*	*	*	*	0.3
2003.....	1.3	*	*	*	*	*	*	*	*	0.2
2002.....	1.3	*	*	*	*	*	*	*	*	0.3
2001.....	1.3	*	*	*	*	*	*	*	*	0.2
2000.....	1.2	*	*	*	*	*	*	*	*	0.3
1999.....	1.3	*	*	*	*	*	*	*	0.2	0.2

All races 3/, female

2004.....	2.0	*	*	*	*	*	*	*	*	0.2
2003.....	2.1	*	*	*	*	*	*	*	*	*
2002.....	2.0	*	*	*	*	*	*	*	*	0.2
2001.....	1.9	*	*	*	*	*	*	*	*	0.2
2000.....	1.9	*	*	*	*	*	*	*	*	0.2
1999.....	2.0	*	*	*	*	*	*	*	0.2	0.2

White, both sexes

2004.....	1.9	*	*	*	*	*	*	*	0.1	0.2
2003.....	1.8	*	*	*	*	*	*	*	0.1	0.2
2002.....	1.8	*	*	*	*	*	*	*	0.1	0.2
2001.....	1.7	*	*	*	*	*	*	*	*	0.2
2000.....	1.7	*	*	*	*	*	*	*	*	0.2
1999.....	1.7	*	*	*	*	*	*	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Cardiomyopathy (I42)

All races 3/, both sexes

2004.....	8.7	2.8	0.2	0.1	0.2	0.5	0.8	1.2	1.6	2.7
2003.....	9.2	3.1	0.4	0.1	0.2	0.5	0.8	1.3	1.7	2.6
2002.....	9.6	2.9	0.2	0.1	0.2	0.6	0.8	1.4	1.8	2.8
2001.....	9.4	3.4	0.4	0.1	0.2	0.6	0.8	1.1	1.7	2.6
2000.....	9.7	3.1	0.3	0.1	0.3	0.5	0.9	1.2	1.8	2.6
1999.....	9.8	3.5	0.2	0.1	0.2	0.6	0.7	1.2	1.8	3.0

All races 3/, male

2004.....	10.3	2.7	0.3	*	0.2	0.7	1.2	1.7	2.4	3.8
2003.....	10.8	3.4	0.3	*	0.2	0.6	1.1	1.8	2.4	3.7
2002.....	11.3	2.7	*	*	0.3	0.8	1.1	2.0	2.6	3.7
2001.....	11.0	2.9	0.5	*	0.2	0.7	1.1	1.5	2.4	3.8
2000.....	11.4	2.8	0.3	*	0.4	0.7	1.2	1.6	2.6	3.6
1999.....	11.5	3.1	*	*	0.3	0.9	0.9	1.6	2.4	4.3

All races 3/, female

2004.....	7.2	3.0	*	*	0.2	0.3	0.4	0.6	0.9	1.7
2003.....	7.7	2.9	0.4	*	*	0.4	0.5	0.7	1.1	1.5
2002.....	8.0	3.1	*	*	*	0.3	0.4	0.8	1.0	2.0
2001.....	8.0	3.9	0.3	*	*	0.4	0.4	0.7	0.9	1.5
2000.....	8.0	3.4	*	*	*	0.3	0.6	0.8	1.1	1.6
1999.....	8.1	3.8	*	*	*	0.3	0.5	0.8	1.2	1.7

White, both sexes

2004.....	8.7	2.7	0.2	*	0.2	0.4	0.7	0.9	1.2	2.1
2003.....	9.2	3.0	0.3	*	0.1	0.4	0.7	0.9	1.3	1.9
2002.....	9.6	2.8	0.2	*	0.2	0.4	0.6	1.0	1.3	2.1
2001.....	9.3	2.8	0.3	*	0.2	0.4	0.6	0.8	1.3	2.0
2000.....	9.5	3.0	0.2	*	0.2	0.3	0.6	0.8	1.4	1.9
1999.....	9.6	3.1	0.2	*	0.2	0.4	0.5	0.9	1.4	2.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Cardiomyopathy (I42)

White, male

2004.....	10.2	2.6	*	*	*	0.5	1.1	1.3	1.8	3.0
2003.....	10.7	2.9	*	*	*	0.5	0.9	1.3	1.9	2.8
2002.....	11.2	2.4	*	*	0.3	0.6	0.9	1.4	1.9	2.8
2001.....	10.8	2.7	0.4	*	*	0.5	0.8	1.2	1.8	2.9
2000.....	11.1	2.7	*	*	0.3	0.5	0.9	1.1	1.9	2.7
1999.....	11.1	2.6	*	*	*	0.6	0.7	1.2	2.0	3.2

White, female

2004.....	7.2	2.8	*	*	*	0.3	0.3	0.4	0.6	1.2
2003.....	7.8	3.1	*	*	*	*	0.4	0.4	0.7	1.1
2002.....	8.0	3.2	*	*	*	0.3	*	0.5	0.7	1.4
2001.....	7.9	2.9	*	*	*	0.3	0.3	0.5	0.7	1.1
2000.....	8.0	3.4	*	*	*	*	0.3	0.5	0.8	1.1
1999.....	8.1	3.5	*	*	*	0.3	0.3	0.5	0.8	1.3

Black, both sexes

2004.....	11.2	*	*	*	*	1.1	1.5	2.8	4.6	7.1
2003.....	11.9	3.5	*	*	*	1.1	1.9	3.8	4.7	7.0
2002.....	12.2	3.4	*	*	*	1.2	2.1	4.0	5.2	7.9
2001.....	12.4	6.8	0.9	*	*	1.3	2.1	3.2	4.6	7.1
2000.....	12.9	3.6	*	*	*	1.4	2.3	3.6	4.8	7.3
1999.....	13.2	5.1	*	*	0.6	1.7	1.9	3.2	4.4	8.0

Black, male

2004.....	13.7	*	*	*	*	1.8	2.2	3.8	6.8	9.7
2003.....	14.5	*	*	*	*	*	2.6	5.3	6.3	9.9
2002.....	14.9	*	*	*	*	1.7	2.6	5.9	7.6	10.3
2001.....	14.9	*	*	*	*	1.7	2.9	4.1	6.7	9.9
2000.....	15.8	*	*	*	*	2.0	2.9	4.7	7.0	10.0
1999.....	16.4	*	*	*	*	2.7	2.5	4.5	5.8	11.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Cardiomyopathy (I42)

Black, female

2004.....	8.9	*	*	*	*	*	*	1.8	2.7	4.7
2003.....	9.5	*	*	*	*	*	*	2.4	3.2	4.5
2002.....	9.8	*	*	*	*	*	1.6	2.2	3.1	5.8
2001.....	10.2	8.8	*	*	*	*	*	2.3	2.7	4.5
2000.....	10.3	*	*	*	*	*	1.7	2.7	2.9	4.9
1999.....	10.3	*	*	*	*	*	*	2.0	3.1	4.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Conduction disorders and cardiac dysrhythmias (I44-I49)

All races 3/, both sexes

2004.....	12.8	0.8	0.1	*	0.2	0.3	0.6	0.7	0.9	1.3
2003.....	13.4	0.8	0.1	*	0.2	0.4	0.5	0.6	1.0	1.4
2002.....	13.6	1.4	0.1	0.1	0.1	0.4	0.5	0.7	0.9	1.3
2001.....	13.5	0.9	0.2	*	0.2	0.3	0.5	0.6	0.8	1.3
2000.....	13.6	1.0	0.2	*	0.2	0.4	0.5	0.6	0.8	1.3
1999.....	14.3	1.0	0.3	*	0.2	0.4	0.5	0.7	0.9	1.3

All races 3/, male

2004.....	11.5	*	*	*	*	0.4	0.7	0.8	1.1	1.6
2003.....	12.0	*	*	*	0.2	0.5	0.6	0.6	1.1	1.7
2002.....	12.2	1.6	*	*	*	0.6	0.7	0.8	1.1	1.6
2001.....	12.0	*	*	*	0.2	0.4	0.7	0.7	1.0	1.6
2000.....	12.1	1.1	*	*	0.2	0.5	0.6	0.8	0.9	1.6
1999.....	12.9	1.2	0.3	*	0.2	0.5	0.6	0.8	1.1	1.5

All races 3/, female

2004.....	14.1	*	*	*	*	0.2	0.4	0.5	0.8	1.0
2003.....	14.8	*	*	*	*	0.3	0.4	0.5	0.8	1.2
2002.....	15.1	1.2	*	*	*	0.2	0.3	0.5	0.7	1.1
2001.....	15.1	*	*	*	0.2	*	0.3	0.4	0.7	1.0
2000.....	15.1	*	*	*	*	0.3	0.4	0.4	0.7	1.0
1999.....	15.7	*	*	*	*	0.4	0.4	0.5	0.7	1.0

White, both sexes

2004.....	13.5	0.7	*	*	0.1	0.3	0.5	0.5	0.8	1.1
2003.....	14.2	*	*	*	0.1	0.3	0.4	0.5	0.8	1.3
2002.....	14.4	1.1	*	*	0.1	0.4	0.4	0.6	0.7	1.1
2001.....	14.3	0.9	0.2	*	0.2	0.3	0.4	0.5	0.7	1.0
2000.....	14.4	1.0	0.2	*	0.1	0.4	0.5	0.5	0.7	1.2
1999.....	15.2	0.8	0.3	*	0.2	0.4	0.4	0.6	0.8	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Conduction disorders and cardiac dysrhythmias (I44-I49)

White, male

2004.....	11.9	*	*	*	*	0.4	0.6	0.6	0.9	1.3
2003.....	12.5	*	*	*	*	0.4	0.5	0.5	1.1	1.5
2002.....	12.7	1.6	*	*	*	0.7	0.6	0.7	0.9	1.3
2001.....	12.4	*	*	*	*	0.4	0.6	0.6	0.9	1.3
2000.....	12.7	*	*	*	*	0.5	0.6	0.7	0.8	1.5
1999.....	13.5	*	0.3	*	*	0.5	0.5	0.7	1.0	1.3

White, female

2004.....	15.0	*	*	*	*	*	0.4	0.4	0.7	0.9
2003.....	15.9	*	*	*	*	0.3	0.3	0.4	0.6	1.1
2002.....	16.1	*	*	*	*	*	*	0.4	0.6	0.9
2001.....	16.1	*	*	*	*	*	*	0.4	0.5	0.8
2000.....	16.1	*	*	*	*	*	0.4	0.3	0.5	0.9
1999.....	16.9	*	*	*	*	0.4	0.4	0.4	0.5	0.8

Black, both sexes

2004.....	12.1	*	*	*	*	*	0.8	1.4	1.9	2.9
2003.....	12.4	*	*	*	*	0.6	0.8	1.4	2.0	2.4
2002.....	12.5	*	*	*	*	*	0.9	1.3	2.0	2.9
2001.....	12.6	*	*	*	*	*	1.0	1.0	1.9	3.2
2000.....	12.3	*	*	*	*	*	0.8	1.1	1.7	2.5
1999.....	13.0	*	*	*	*	*	1.1	1.4	2.0	3.1

Black, male

2004.....	11.9	*	*	*	*	*	*	1.5	1.9	3.6
2003.....	12.0	*	*	*	*	*	*	1.6	1.7	2.6
2002.....	11.9	*	*	*	*	*	*	*	2.5	3.4
2001.....	12.2	*	*	*	*	*	1.5	*	2.1	3.8
2000.....	11.7	*	*	*	*	*	*	*	1.7	3.3
1999.....	12.9	*	*	*	*	*	*	1.6	2.2	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ year 2/	year 2/ year 2/	years	years	years	years	years	years	years	years

Conduction disorders and cardiac dysrhythmias (I44-I49)

Black, female

2004.....	12.3	*	*	*	*	*	*	*	1.9	2.4
2003.....	12.8	*	*	*	*	*	*	*	2.2	2.2
2002.....	13.1	*	*	*	*	*	*	*	1.5	2.4
2001.....	12.9	*	*	*	*	*	*	*	1.6	2.7
2000.....	12.8	*	*	*	*	*	*	*	1.7	1.8
1999.....	13.1	*	*	*	*	*	*	*	1.8	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Heart failure (I50)

All races 3/, both sexes

2004.....	19.5	0.7	*	*	*	*	0.1	0.2	0.3	0.5
2003.....	19.8	0.9	*	*	*	*	0.1	0.1	0.3	0.4
2002.....	19.6	0.6	*	*	*	*	*	0.2	0.3	0.5
2001.....	20.0	0.6	*	*	*	*	*	0.2	0.2	0.5
2000.....	19.8	0.8	*	*	*	0.1	*	0.1	0.2	0.3
1999.....	19.7	*	*	*	*	*	*	0.1	0.2	0.4

All races 3/, male

2004.....	15.4	*	*	*	*	*	*	0.3	0.3	0.7
2003.....	15.7	1.4	*	*	*	*	*	*	0.4	0.5
2002.....	15.3	*	*	*	*	*	*	0.2	0.4	0.5
2001.....	15.5	*	*	*	*	*	*	0.3	0.3	0.7
2000.....	15.3	1.1	*	*	*	*	*	*	0.2	0.4
1999.....	15.2	*	*	*	*	*	*	*	*	0.5

All races 3/, female

2004.....	23.4	*	*	*	*	*	*	*	0.2	0.4
2003.....	23.7	*	*	*	*	*	*	*	0.2	0.4
2002.....	23.7	*	*	*	*	*	*	*	0.3	0.4
2001.....	24.3	*	*	*	*	*	*	*	0.2	0.4
2000.....	24.1	*	*	*	*	*	*	*	*	0.3
1999.....	24.0	*	*	*	*	*	*	*	*	0.3

White, both sexes

2004.....	21.5	0.6	*	*	*	*	*	0.1	0.2	0.4
2003.....	21.8	0.6	*	*	*	*	*	*	0.2	0.3
2002.....	21.6	*	*	*	*	*	*	*	0.2	0.3
2001.....	22.1	*	*	*	*	*	*	0.2	0.2	0.4
2000.....	21.8	*	*	*	*	*	*	*	0.1	0.2
1999.....	21.6	*	*	*	*	*	*	*	0.1	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)

All races 3/, both sexes

2004.....	0.2	0.6	0.2	*	*	0.1	0.1	0.1	0.1	0.2
2003.....	0.2	0.7	0.2	*	0.1	0.1	0.1	0.2	0.2	0.2
2002.....	0.3	0.8	0.1	*	*	0.2	0.1	0.1	0.2	0.3
2001.....	0.3	0.6	0.3	*	*	0.1	0.1	0.2	0.2	0.3
2000.....	0.2	0.6	0.1	*	*	0.1	0.1	0.2	0.2	0.2
1999.....	0.3	0.7	0.2	*	*	0.2	0.2	0.2	0.2	0.3

All races 3/, male

2004.....	0.2	*	*	*	*	*	0.2	*	*	0.3
2003.....	0.2	*	*	*	*	0.2	*	*	0.2	0.3
2002.....	0.3	*	*	*	*	*	*	*	0.2	0.3
2001.....	0.3	*	0.3	*	*	*	*	*	*	0.4
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.3
1999.....	0.3	*	*	*	*	*	0.3	0.2	0.3	0.3

All races 3/, female

2004.....	0.2	*	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*	*	0.2
2002.....	0.3	*	*	*	*	*	*	*	*	0.2
2001.....	0.2	*	*	*	*	*	*	0.2	0.2	0.2
2000.....	0.2	*	*	*	*	*	*	*	*	0.2
1999.....	0.2	*	*	*	*	*	*	*	*	0.2

White, both sexes

2004.....	0.2	*	*	*	*	*	0.2	0.1	*	0.2
2003.....	0.2	*	*	*	*	0.1	*	0.2	0.1	0.2
2002.....	0.3	*	*	*	*	0.1	*	0.1	0.2	0.2
2001.....	0.2	*	0.2	*	*	*	*	0.1	0.1	0.3
2000.....	0.2	*	*	*	*	0.1	*	0.1	0.2	0.2
1999.....	0.2	*	*	*	*	0.2	0.2	0.1	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

All races 3/, both sexes

2004.....	3.4	0.6	*	*	*	0.1	0.3	0.5	0.5	0.7
2003.....	3.4	0.5	*	*	*	0.2	0.3	0.3	0.6	0.7
2002.....	3.3	0.6	*	*	*	0.2	0.3	0.3	0.5	0.8
2001.....	3.2	*	*	*	*	0.1	0.3	0.3	0.6	0.8
2000.....	3.1	0.8	*	*	*	0.2	0.3	0.3	0.5	0.6
1999.....	2.9	0.8	*	*	*	0.2	0.2	0.3	0.5	0.7

All races 3/, male

2004.....	3.5	*	*	*	*	0.2	0.5	0.7	0.8	1.0
2003.....	3.4	*	*	*	*	0.2	0.5	0.5	0.9	1.1
2002.....	3.3	*	*	*	*	0.3	0.4	0.4	0.7	1.2
2001.....	3.2	*	*	*	*	0.2	0.4	0.5	0.8	1.3
2000.....	3.1	*	*	*	*	0.3	0.5	0.5	0.8	0.9
1999.....	3.0	*	*	*	*	0.3	0.3	0.4	0.8	1.1

All races 3/, female

2004.....	3.3	*	*	*	*	*	*	*	0.2	0.4
2003.....	3.3	*	*	*	*	*	*	*	0.3	0.4
2002.....	3.2	*	*	*	*	*	*	*	*	0.5
2001.....	3.1	*	*	*	*	*	*	*	0.3	0.4
2000.....	3.1	*	*	*	*	*	*	*	*	0.4
1999.....	2.8	*	*	*	*	*	*	*	0.2	0.4

White, both sexes

2004.....	3.5	*	*	*	*	*	0.3	0.4	0.4	0.7
2003.....	3.5	*	*	*	*	0.1	0.3	0.2	0.5	0.7
2002.....	3.4	*	*	*	*	0.1	0.2	0.3	0.4	0.8
2001.....	3.2	*	*	*	*	0.1	0.2	0.3	0.5	0.7
2000.....	3.2	0.7	*	*	*	*	0.2	0.3	0.5	0.5
1999.....	3.0	0.7	*	*	*	0.2	0.2	0.2	0.4	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
White, male										
2004.....	3.6	*	*	*	*	*	0.4	0.6	0.7	1.0
2003.....	3.5	*	*	*	*	*	0.4	0.4	0.9	1.0
2002.....	3.4	*	*	*	*	0.2	0.4	0.4	0.6	1.2
2001.....	3.2	*	*	*	*	*	0.3	0.5	0.7	1.1
2000.....	3.2	*	*	*	*	*	0.3	0.3	0.8	0.8
1999.....	3.0	*	*	*	*	*	0.3	0.4	0.7	1.0
White, female										
2004.....	3.5	*	*	*	*	*	*	*	*	0.4
2003.....	3.5	*	*	*	*	*	*	*	*	0.3
2002.....	3.4	*	*	*	*	*	*	*	*	0.4
2001.....	3.3	*	*	*	*	*	*	*	0.3	0.3
2000.....	3.2	*	*	*	*	*	*	*	*	0.2
1999.....	2.9	*	*	*	*	*	*	*	*	0.3
Black, both sexes										
2004.....	3.6	*	*	*	*	*	*	0.9	1.1	1.1
2003.....	3.5	*	*	*	*	*	0.8	0.8	1.1	1.5
2002.....	3.4	*	*	*	*	*	*	*	1.1	1.4
2001.....	3.4	*	*	*	*	*	*	*	1.2	1.5
2000.....	3.5	*	*	*	*	*	0.8	0.9	0.9	1.4
1999.....	3.3	*	*	*	*	*	*	*	1.2	1.5
Black, male										
2004.....	4.0	*	*	*	*	*	*	1.7	1.5	*
2003.....	3.7	*	*	*	*	*	*	*	*	2.5
2002.....	3.8	*	*	*	*	*	*	*	1.5	1.8
2001.....	3.5	*	*	*	*	*	*	*	1.9	2.2
2000.....	3.9	*	*	*	*	*	*	*	*	1.7
1999.....	3.6	*	*	*	*	*	*	*	1.9	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Cerebrovascular diseases (I60-I69)

All races 3/, both sexes

2004.....	51.1	3.1	0.3	0.2	0.2	0.3	0.7	1.1	1.8	3.4
2003.....	54.2	2.5	0.3	0.1	0.2	0.4	0.6	1.0	1.9	3.6
2002.....	56.4	2.9	0.3	0.2	0.3	0.3	0.6	1.0	1.8	3.6
2001.....	57.4	2.7	0.4	0.2	0.2	0.3	0.7	1.1	1.9	3.8
2000.....	59.6	3.3	0.3	0.1	0.2	0.3	0.7	1.1	1.9	3.9
1999.....	60.0	2.7	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.0

All races 3/, male

2004.....	40.7	3.4	0.3	*	0.2	0.4	0.7	1.1	1.7	3.7
2003.....	42.9	2.8	0.3	*	0.2	0.5	0.6	1.0	2.0	3.8
2002.....	44.2	3.2	0.4	*	0.3	0.3	0.7	1.0	1.7	3.6
2001.....	45.2	3.1	0.3	0.2	0.2	0.3	0.7	1.1	2.0	3.9
2000.....	46.9	3.8	*	*	0.2	0.4	0.7	1.1	1.9	3.8
1999.....	47.1	3.3	0.3	*	*	0.3	0.6	1.0	1.9	3.9

All races 3/, female

2004.....	61.2	2.8	*	*	0.2	0.3	0.7	1.0	1.8	3.1
2003.....	65.1	2.2	0.3	*	*	0.4	0.7	0.9	1.8	3.3
2002.....	68.2	2.5	0.3	*	0.3	*	0.5	0.9	1.9	3.7
2001.....	69.2	2.3	0.4	*	0.2	0.3	0.6	1.0	1.9	3.6
2000.....	71.8	2.7	0.4	*	0.3	0.3	0.7	1.0	1.9	4.0
1999.....	72.3	2.1	0.3	*	0.2	0.3	0.6	1.0	1.8	4.1

White, both sexes

2004.....	53.7	2.5	0.2	0.2	0.2	0.3	0.6	0.9	1.4	2.9
2003.....	57.0	2.0	0.2	0.1	0.2	0.4	0.5	0.8	1.6	2.7
2002.....	59.5	2.3	0.3	0.2	0.2	0.2	0.6	0.8	1.5	2.9
2001.....	60.5	1.9	0.3	0.2	0.2	0.3	0.5	0.9	1.7	3.1
2000.....	62.8	2.5	0.3	0.1	0.2	0.3	0.6	1.0	1.6	3.0
1999.....	63.3	2.3	0.3	0.1	0.2	0.3	0.6	0.9	1.5	3.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Cerebrovascular diseases (I60-I69)

White, male

2004.....	41.8	2.8	*	*	*	0.3	0.6	1.1	1.4	3.1
2003.....	44.2	2.0	*	*	*	0.4	0.6	0.8	1.7	2.8
2002.....	45.7	2.6	0.4	*	*	0.3	0.7	1.0	1.4	2.8
2001.....	46.6	2.2	*	*	*	0.3	0.7	0.9	1.8	3.1
2000.....	48.4	3.0	*	*	*	0.4	0.7	1.1	1.6	3.0
1999.....	48.7	2.8	*	*	*	0.3	0.6	0.9	1.7	3.2

White, female

2004.....	65.3	2.2	*	*	*	0.3	0.6	0.7	1.5	2.7
2003.....	69.5	2.0	*	*	*	0.3	0.5	0.8	1.5	2.5
2002.....	73.0	2.0	*	*	0.3	*	0.5	0.7	1.6	2.9
2001.....	74.0	1.6	0.4	*	*	0.3	0.4	0.9	1.6	3.0
2000.....	76.9	2.0	*	*	0.3	*	0.6	0.9	1.5	3.0
1999.....	77.6	1.9	*	*	*	*	0.6	0.9	1.4	3.3

Black, both sexes

2004.....	46.9	6.0	*	*	*	*	1.0	2.2	4.1	7.0
2003.....	49.3	4.9	*	*	*	0.7	1.2	2.0	4.1	8.8
2002.....	50.0	5.5	*	*	*	*	0.8	1.9	4.2	8.2
2001.....	51.1	6.6	*	*	*	*	1.3	2.2	3.6	8.1
2000.....	52.5	7.5	*	*	*	*	1.1	1.5	4.3	9.6
1999.....	52.2	4.6	*	*	*	*	0.8	1.8	3.6	9.1

Black, male

2004.....	41.5	6.8	*	*	*	*	*	1.7	4.1	8.5
2003.....	43.2	5.9	*	*	*	*	*	2.6	4.1	9.7
2002.....	43.5	5.8	*	*	*	*	*	1.7	4.0	7.5
2001.....	44.6	8.1	*	*	*	*	*	2.8	3.4	8.8
2000.....	46.1	8.6	*	*	*	*	*	*	4.1	9.2
1999.....	45.9	*	*	*	*	*	*	1.9	3.7	8.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Cerebrovascular diseases (I60-I69)

Black, female

2004.....	51.9	*	*	*	*	*	*	2.6	4.0	5.7
2003.....	54.8	*	*	*	*	*	1.6	1.5	4.1	8.0
2002.....	55.8	*	*	*	*	*	*	2.2	4.3	8.9
2001.....	56.9	*	*	*	*	*	1.7	1.7	3.8	7.4
2000.....	58.3	*	*	*	*	*	*	1.6	4.6	10.0
1999.....	57.9	*	*	*	*	*	*	1.8	3.6	9.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Subarachnoid hemorrhage (I60)

All races 3/, both sexes

2004.....	2.1	*	*	*	*	*	0.2	0.3	0.6	1.2
2003.....	2.1	*	*	*	*	*	0.2	0.4	0.6	1.2
2002.....	2.1	*	*	*	*	*	0.2	0.3	0.7	1.3
2001.....	2.2	*	*	*	*	0.1	0.2	0.4	0.7	1.3
2000.....	2.3	*	*	*	*	*	0.3	0.4	0.7	1.3
1999.....	2.3	*	*	*	*	0.1	0.2	0.4	0.7	1.6

All races 3/, male

2004.....	1.5	*	*	*	*	*	0.2	0.3	0.6	1.0
2003.....	1.5	*	*	*	*	*	0.2	0.4	0.7	1.1
2002.....	1.5	*	*	*	*	*	0.3	0.3	0.6	1.1
2001.....	1.5	*	*	*	*	*	0.2	0.4	0.7	1.0
2000.....	1.6	*	*	*	*	*	0.3	0.4	0.7	1.0
1999.....	1.6	*	*	*	*	*	*	0.4	0.7	1.3

All races 3/, female

2004.....	2.6	*	*	*	*	*	0.3	0.3	0.6	1.4
2003.....	2.6	*	*	*	*	*	*	0.4	0.6	1.3
2002.....	2.8	*	*	*	*	*	0.2	0.4	0.8	1.5
2001.....	2.8	*	*	*	*	*	0.2	0.3	0.7	1.5
2000.....	3.0	*	*	*	*	*	0.3	0.4	0.8	1.6
1999.....	3.0	*	*	*	*	*	0.2	0.4	0.8	1.8

White, both sexes

2004.....	2.1	*	*	*	*	*	0.2	0.3	0.6	1.1
2003.....	2.0	*	*	*	*	*	0.2	0.3	0.6	1.0
2002.....	2.1	*	*	*	*	*	0.2	0.3	0.5	1.1
2001.....	2.1	*	*	*	*	*	0.2	0.3	0.6	1.1
2000.....	2.3	*	*	*	*	*	0.3	0.5	0.7	1.2
1999.....	2.3	*	*	*	*	*	0.2	0.4	0.7	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Subarachnoid hemorrhage (I60)

Black, female

2004.....	2.8	*	*	*	*	*	*	*	1.8
2003.....	2.9	*	*	*	*	*	*	*	2.9
2002.....	3.1	*	*	*	*	*	*	1.6	2.9
2001.....	3.2	*	*	*	*	*	*	*	3.0
2000.....	3.5	*	*	*	*	*	*	1.9	3.3
1999.....	3.3	*	*	*	*	*	*	1.4	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intracerebral and other intracranial hemorrhage (I61-I62)

All races 3/, both sexes

2004.....	9.4	1.3	*	*	*	0.1	0.3	0.4	0.6	1.2
2003.....	9.4	0.8	*	*	*	0.2	0.3	0.4	0.6	1.3
2002.....	9.4	1.0	*	*	*	0.1	0.2	0.3	0.6	1.3
2001.....	9.4	1.0	*	*	*	0.1	0.3	0.4	0.7	1.5
2000.....	9.7	1.1	*	*	*	0.1	0.3	0.3	0.7	1.5
1999.....	9.1	1.1	*	*	*	0.1	0.2	0.3	0.6	1.5

All races 3/, male

2004.....	8.9	1.4	*	*	*	*	0.3	0.5	0.6	1.6
2003.....	8.9	*	*	*	*	0.2	0.3	0.4	0.7	1.6
2002.....	8.7	1.1	*	*	*	*	0.3	0.4	0.7	1.3
2001.....	8.8	1.2	*	*	*	0.2	0.3	0.5	0.8	1.8
2000.....	9.1	1.3	*	*	*	*	0.3	0.4	0.8	1.7
1999.....	8.5	1.3	*	*	*	*	0.2	0.3	0.7	1.7

All races 3/, female

2004.....	9.8	1.3	*	*	*	*	0.2	0.4	0.6	0.8
2003.....	9.8	*	*	*	*	*	0.3	0.3	0.6	1.0
2002.....	10.0	1.0	*	*	*	*	*	0.3	0.6	1.3
2001.....	9.9	*	*	*	*	*	0.3	0.4	0.7	1.1
2000.....	10.4	*	*	*	*	*	0.3	0.2	0.6	1.3
1999.....	9.8	*	*	*	*	*	*	0.4	0.5	1.3

White, both sexes

2004.....	9.4	1.0	*	*	*	*	0.2	0.3	0.5	0.9
2003.....	9.3	*	*	*	*	0.1	0.2	0.3	0.5	0.9
2002.....	9.5	0.9	*	*	*	*	0.2	0.3	0.5	1.0
2001.....	9.4	0.7	*	*	*	0.1	0.3	0.3	0.6	1.1
2000.....	9.8	0.7	*	*	*	*	0.3	0.3	0.5	1.0
1999.....	9.2	0.8	*	*	*	*	0.2	0.3	0.4	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intracerebral and other intracranial hemorrhage (I61-I62)

White, male

2004.....	8.8	*	*	*	*	*	0.3	0.4	0.5	1.2
2003.....	8.6	*	*	*	*	*	0.2	0.3	0.5	1.1
2002.....	8.7	*	*	*	*	*	0.3	0.4	0.5	1.0
2001.....	8.6	*	*	*	*	*	0.3	0.4	0.7	1.4
2000.....	8.9	*	*	*	*	*	0.3	0.4	0.6	1.2
1999.....	8.3	*	*	*	*	*	*	0.3	0.6	1.2

White, female

2004.....	10.1	*	*	*	*	*	*	*	0.4	0.6
2003.....	10.0	*	*	*	*	*	*	*	0.5	0.8
2002.....	10.2	*	*	*	*	*	*	*	0.5	0.9
2001.....	10.1	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	10.7	*	*	*	*	*	*	*	0.4	0.9
1999.....	10.0	*	*	*	*	*	*	0.3	0.3	0.9

Black, both sexes

2004.....	10.2	*	*	*	*	*	*	1.1	1.6	3.0
2003.....	10.5	*	*	*	*	*	*	0.9	1.5	3.2
2002.....	9.7	*	*	*	*	*	*	*	1.7	3.1
2001.....	10.0	*	*	*	*	*	*	1.1	1.6	3.3
2000.....	10.3	*	*	*	*	*	*	*	2.0	4.4
1999.....	9.5	*	*	*	*	*	*	*	1.4	4.4

Black, male

2004.....	10.7	*	*	*	*	*	*	*	1.7	4.6
2003.....	11.0	*	*	*	*	*	*	*	1.6	4.0
2002.....	10.1	*	*	*	*	*	*	*	2.0	3.1
2001.....	10.3	*	*	*	*	*	*	*	1.5	4.2
2000.....	10.9	*	*	*	*	*	*	*	2.2	5.2
1999.....	9.8	*	*	*	*	*	*	*	1.5	5.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intracerebral and other intracranial hemorrhage (I61-I62)

Black, female

2004.....	9.8	*	*	*	*	*	*	1.4	1.6
2003.....	10.1	*	*	*	*	*	*	1.4	2.5
2002.....	9.4	*	*	*	*	*	*	1.4	3.2
2001.....	9.8	*	*	*	*	*	*	1.6	2.4
2000.....	9.8	*	*	*	*	*	*	1.7	3.8
1999.....	9.2	*	*	*	*	*	*	1.4	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Stroke, not specified as hemorrhage or infarction (I64)

All races 3/, both sexes

2004.....	26.0	*	*	*	*	*	*	0.1	0.3	0.6
2003.....	28.4	*	*	*	*	*	*	0.1	0.4	0.6
2002.....	30.1	*	*	*	*	*	*	0.1	0.3	0.6
2001.....	31.1	*	*	*	*	*	*	0.1	0.2	0.5
2000.....	32.1	*	*	*	*	*	*	0.2	0.2	0.5
1999.....	32.6	*	*	*	*	*	*	0.1	0.2	0.4

All races 3/, male

2004.....	19.7	*	*	*	*	*	*	*	0.2	0.6
2003.....	21.5	*	*	*	*	*	*	*	0.4	0.7
2002.....	22.6	*	*	*	*	*	*	*	*	0.7
2001.....	23.5	*	*	*	*	*	*	*	*	0.5
2000.....	24.2	*	*	*	*	*	*	*	0.3	0.6
1999.....	24.7	*	*	*	*	*	*	*	0.2	0.4

All races 3/, female

2004.....	32.2	*	*	*	*	*	*	*	0.4	0.5
2003.....	35.0	*	*	*	*	*	*	*	0.3	0.5
2002.....	37.4	*	*	*	*	*	*	*	0.4	0.5
2001.....	38.5	*	*	*	*	*	*	*	0.3	0.5
2000.....	39.7	*	*	*	*	*	*	*	*	0.5
1999.....	40.3	*	*	*	*	*	*	*	0.3	0.5

White, both sexes

2004.....	27.6	*	*	*	*	*	*	*	0.2	0.4
2003.....	30.2	*	*	*	*	*	*	*	0.2	0.4
2002.....	32.1	*	*	*	*	*	*	*	0.2	0.4
2001.....	33.1	*	*	*	*	*	*	*	0.2	0.4
2000.....	34.2	*	*	*	*	*	*	*	0.2	0.4
1999.....	34.8	*	*	*	*	*	*	*	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other cerebrovascular diseases and their sequelae (I67,I69)

All races 3/, both sexes

2004.....	9.9	1.4	0.2	*	*	*	*	*	0.1	0.2
2003.....	10.3	1.2	0.1	*	*	*	*	*	0.1	0.2
2002.....	10.5	1.4	0.2	*	*	*	*	*	0.1	0.3
2001.....	10.3	1.2	0.2	*	*	*	*	*	0.2	0.3
2000.....	10.5	1.6	0.2	*	*	*	*	*	0.1	0.3
1999.....	10.6	1.1	0.1	*	*	*	*	*	0.1	0.2

All races 3/, male

2004.....	7.6	1.5	*	*	*	*	*	*	*	0.2
2003.....	7.8	1.2	*	*	*	*	*	*	*	0.3
2002.....	7.9	1.7	0.3	*	*	*	*	*	*	0.3
2001.....	7.9	1.5	*	*	*	*	*	*	*	0.3
2000.....	8.1	1.8	*	*	*	*	*	*	*	0.2
1999.....	8.1	1.4	*	*	*	*	*	*	*	0.2

All races 3/, female

2004.....	12.1	1.4	*	*	*	*	*	*	*	0.2
2003.....	12.7	1.2	*	*	*	*	*	*	*	0.2
2002.....	13.0	1.1	*	*	*	*	*	*	*	0.3
2001.....	12.7	*	*	*	*	*	*	*	*	0.3
2000.....	12.8	1.4	*	*	*	*	*	*	*	0.3
1999.....	12.9	*	*	*	*	*	*	*	*	0.3

White, both sexes

2004.....	10.5	1.1	*	*	*	*	*	*	0.1	0.2
2003.....	11.0	1.0	*	*	*	*	*	*	*	0.2
2002.....	11.3	0.9	0.2	*	*	*	*	*	0.1	0.2
2001.....	11.1	0.9	0.2	*	*	*	*	*	0.2	0.3
2000.....	11.3	1.2	*	*	*	*	*	*	*	0.2
1999.....	11.4	0.9	*	*	*	*	*	*	0.1	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of arteries, arterioles and capillaries (I70-I78)

All races 3/, both sexes

2004.....	12.1	*	*	*	*	*	0.2	0.3	0.5	0.9
2003.....	13.0	*	*	*	*	0.2	0.2	0.3	0.5	0.8
2002.....	13.4	*	*	*	*	0.2	0.2	0.3	0.6	0.8
2001.....	13.8	*	*	*	*	0.1	0.2	0.4	0.5	0.9
2000.....	14.4	*	*	*	*	*	0.2	0.3	0.5	0.8
1999.....	14.6	*	*	*	*	*	0.2	0.3	0.5	0.8

All races 3/, male

2004.....	11.6	*	*	*	*	*	0.3	0.5	0.7	1.2
2003.....	12.6	*	*	*	*	*	0.3	0.4	0.6	1.2
2002.....	13.0	*	*	*	*	*	0.3	0.3	0.7	0.8
2001.....	13.3	*	*	*	*	*	0.2	0.4	0.7	1.3
2000.....	14.1	*	*	*	*	*	0.3	0.4	0.7	0.9
1999.....	14.1	*	*	*	*	*	0.2	0.4	0.7	1.1

All races 3/, female

2004.....	12.6	*	*	*	*	*	*	*	0.4	0.6
2003.....	13.3	*	*	*	*	*	*	*	0.3	0.5
2002.....	13.8	*	*	*	*	*	*	0.2	0.4	0.7
2001.....	14.4	*	*	*	*	*	*	0.3	0.3	0.6
2000.....	14.6	*	*	*	*	*	*	*	0.4	0.6
1999.....	15.2	*	*	*	*	*	*	*	0.3	0.6

White, both sexes

2004.....	13.2	*	*	*	*	*	0.2	0.3	0.4	0.7
2003.....	14.2	*	*	*	*	0.1	0.2	0.3	0.4	0.6
2002.....	14.7	*	*	*	*	0.2	0.2	0.2	0.5	0.7
2001.....	15.1	*	*	*	*	*	0.2	0.3	0.5	0.8
2000.....	15.7	*	*	*	*	*	0.2	0.2	0.5	0.7
1999.....	16.0	*	*	*	*	*	0.2	0.3	0.4	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Aortic aneurysm and dissection (I71)

All races 3/, both sexes

2004.....	4.7	*	*	*	*	*	0.2	0.3	0.4	0.6
2003.....	5.1	*	*	*	*	*	0.1	0.2	0.3	0.6
2002.....	5.1	*	*	*	*	0.1	*	0.2	0.4	0.5
2001.....	5.3	*	*	*	*	*	0.1	0.2	0.3	0.6
2000.....	5.6	*	*	*	*	*	0.1	0.2	0.3	0.5
1999.....	5.7	*	*	*	*	*	0.1	0.2	0.3	0.6

All races 3/, male

2004.....	5.6	*	*	*	*	*	0.3	0.4	0.5	0.9
2003.....	6.3	*	*	*	*	*	0.2	0.3	0.5	0.9
2002.....	6.4	*	*	*	*	*	*	0.3	0.5	0.6
2001.....	6.5	*	*	*	*	*	*	0.4	0.5	0.9
2000.....	7.0	*	*	*	*	*	*	0.3	0.5	0.6
1999.....	7.0	*	*	*	*	*	*	0.3	0.5	0.9

All races 3/, female

2004.....	3.8	*	*	*	*	*	*	*	0.2	0.3
2003.....	3.9	*	*	*	*	*	*	*	*	0.3
2002.....	4.0	*	*	*	*	*	*	*	0.2	0.3
2001.....	4.2	*	*	*	*	*	*	*	*	0.3
2000.....	4.3	*	*	*	*	*	*	*	0.2	0.3
1999.....	4.4	*	*	*	*	*	*	*	*	0.3

White, both sexes

2004.....	5.1	*	*	*	*	*	0.2	0.2	0.3	0.5
2003.....	5.6	*	*	*	*	*	*	0.2	0.3	0.4
2002.....	5.7	*	*	*	*	*	*	0.1	0.3	0.5
2001.....	5.9	*	*	*	*	*	0.1	0.2	0.3	0.5
2000.....	6.2	*	*	*	*	*	*	0.2	0.3	0.4
1999.....	6.3	*	*	*	*	*	*	0.2	0.3	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races 3/, both sexes

2004.....	3.4	*	*	*	*	*	*	*	0.1	0.2
2003.....	3.4	*	*	*	*	*	*	*	0.1	0.2
2002.....	3.5	*	*	*	*	*	*	0.1	0.1	0.2
2001.....	3.5	*	*	*	*	*	*	0.1	0.2	0.3
2000.....	3.6	*	*	*	*	*	*	*	0.2	0.2
1999.....	3.6	*	*	*	*	*	*	*	0.1	0.2

All races 3/, male

2004.....	2.9	*	*	*	*	*	*	*	*	*
2003.....	3.0	*	*	*	*	*	*	*	*	0.2
2002.....	3.0	*	*	*	*	*	*	*	*	*
2001.....	3.1	*	*	*	*	*	*	*	*	0.2
2000.....	3.2	*	*	*	*	*	*	*	*	0.2
1999.....	3.1	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	3.9	*	*	*	*	*	*	*	*	0.3
2003.....	3.8	*	*	*	*	*	*	*	*	0.2
2002.....	4.0	*	*	*	*	*	*	*	*	0.3
2001.....	4.0	*	*	*	*	*	*	*	*	0.3
2000.....	4.0	*	*	*	*	*	*	*	*	0.3
1999.....	4.1	*	*	*	*	*	*	*	*	0.2

White, both sexes

2004.....	3.5	*	*	*	*	*	*	*	*	0.2
2003.....	3.6	*	*	*	*	*	*	*	*	0.2
2002.....	3.7	*	*	*	*	*	*	*	0.1	0.2
2001.....	3.7	*	*	*	*	*	*	*	0.1	0.2
2000.....	3.8	*	*	*	*	*	*	*	0.2	0.2
1999.....	3.8	*	*	*	*	*	*	*	0.1	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes

2004.....	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.5
2003.....	1.6	0.9	*	*	*	*	0.2	0.3	0.4	0.6
2002.....	1.6	0.7	*	*	*	*	0.1	0.2	0.5	0.6
2001.....	1.6	0.5	*	*	*	*	0.1	0.2	0.4	0.4
2000.....	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.6
1999.....	1.4	0.6	*	*	*	*	*	0.2	0.3	0.4

All races 3/, male

2004.....	1.4	*	*	*	*	*	*	0.3	0.4	0.5
2003.....	1.4	1.0	*	*	*	*	*	0.2	0.4	0.5
2002.....	1.4	*	*	*	*	*	*	*	0.5	0.6
2001.....	1.4	*	*	*	*	*	*	*	0.3	0.5
2000.....	1.4	*	*	*	*	*	*	*	0.4	0.6
1999.....	1.3	*	*	*	*	*	*	*	0.3	0.5

All races 3/, female

2004.....	1.7	*	*	*	*	*	*	0.3	0.4	0.5
2003.....	1.9	*	*	*	*	*	0.2	0.3	0.5	0.6
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.6
2001.....	1.9	*	*	*	*	*	*	0.3	0.4	0.4
2000.....	1.8	*	*	*	*	*	*	0.4	0.4	0.5
1999.....	1.6	*	*	*	*	*	*	*	0.3	0.4

White, both sexes

2004.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.4
2003.....	1.6	0.9	*	*	*	*	0.1	0.2	0.3	0.5
2002.....	1.6	0.6	*	*	*	*	*	0.2	0.4	0.5
2001.....	1.7	*	*	*	*	*	*	0.2	0.3	0.3
2000.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.5
1999.....	1.5	*	*	*	*	*	*	0.1	0.3	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)

All races 3/, both sexes

2004.....	1.2	*	*	*	*	*	0.1	0.3	0.4	0.4
2003.....	1.2	*	*	*	*	*	0.2	0.3	0.4	0.5
2002.....	1.2	*	*	*	*	*	0.1	0.2	0.4	0.5
2001.....	1.2	*	*	*	*	*	0.1	0.2	0.3	0.4
2000.....	1.2	*	*	*	*	*	0.1	0.3	0.3	0.5
1999.....	0.9	*	*	*	*	*	*	0.1	0.3	0.3

All races 3/, male

2004.....	1.0	*	*	*	*	*	*	0.3	0.4	0.4
2003.....	1.0	*	*	*	*	*	*	0.2	0.3	0.5
2002.....	1.0	*	*	*	*	*	*	*	0.5	0.5
2001.....	1.0	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.0	*	*	*	*	*	*	*	0.3	0.5
1999.....	0.8	*	*	*	*	*	*	*	0.3	0.4

All races 3/, female

2004.....	1.4	*	*	*	*	*	*	0.2	0.4	0.5
2003.....	1.5	*	*	*	*	*	0.2	0.3	0.4	0.5
2002.....	1.5	*	*	*	*	*	*	0.3	0.4	0.5
2001.....	1.5	*	*	*	*	*	*	0.3	0.4	0.4
2000.....	1.4	*	*	*	*	*	*	0.4	0.3	0.4
1999.....	1.1	*	*	*	*	*	*	*	0.3	0.3

White, both sexes

2004.....	1.2	*	*	*	*	*	*	0.2	0.3	0.3
2003.....	1.2	*	*	*	*	*	*	0.2	0.3	0.4
2002.....	1.2	*	*	*	*	*	*	0.1	0.3	0.4
2001.....	1.2	*	*	*	*	*	*	0.2	0.3	0.3
2000.....	1.2	*	*	*	*	*	*	0.2	0.3	0.4
1999.....	0.9	*	*	*	*	*	*	*	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

X. Diseases of the respiratory system (J00-J98)

All races 3/, both sexes

2004.....	77.3	15.8	1.9	0.6	0.8	1.1	1.5	1.7	2.3	4.0
2003.....	81.1	17.5	2.2	0.7	1.0	1.2	1.8	1.9	2.7	4.4
2002.....	81.4	15.8	1.9	0.7	1.0	1.2	1.5	1.9	2.6	4.4
2001.....	80.8	17.3	1.8	0.6	0.8	1.0	1.6	1.9	2.8	4.7
2000.....	82.1	17.1	1.7	0.7	0.9	1.1	1.7	2.0	2.9	4.5
1999.....	82.4	18.1	2.1	0.7	0.9	1.3	1.6	2.1	2.9	4.6

All races 3/, male

2004.....	75.2	17.6	1.8	0.7	0.9	1.2	1.7	1.9	2.8	4.2
2003.....	78.2	20.6	2.5	0.8	1.0	1.3	2.0	2.3	3.1	4.6
2002.....	78.9	18.8	2.2	0.7	0.9	1.4	1.6	1.9	2.8	4.6
2001.....	78.2	19.7	1.9	0.7	0.8	1.1	1.9	1.9	2.9	4.9
2000.....	80.2	18.9	1.7	0.8	0.9	1.3	1.9	2.2	3.1	4.7
1999.....	81.3	19.6	2.2	0.7	1.1	1.4	1.7	2.2	3.0	4.9

All races 3/, female

2004.....	79.4	14.1	1.9	0.5	0.6	0.9	1.2	1.4	1.9	3.7
2003.....	84.0	14.2	1.8	0.6	0.9	1.0	1.6	1.6	2.4	4.2
2002.....	83.7	12.6	1.5	0.7	1.0	0.9	1.3	1.8	2.5	4.3
2001.....	83.2	14.7	1.8	0.6	0.7	0.9	1.4	1.9	2.6	4.4
2000.....	84.0	15.2	1.7	0.6	0.8	0.9	1.5	1.8	2.7	4.2
1999.....	83.4	16.6	1.9	0.6	0.8	1.1	1.4	2.1	2.8	4.4

White, both sexes

2004.....	86.4	12.2	1.5	0.4	0.6	0.9	1.3	1.4	2.1	3.4
2003.....	90.6	14.4	1.8	0.6	0.8	0.9	1.5	1.6	2.4	3.9
2002.....	90.8	12.0	1.4	0.6	0.7	0.9	1.3	1.7	2.3	3.8
2001.....	89.7	12.3	1.5	0.5	0.7	0.8	1.3	1.5	2.2	3.8
2000.....	91.2	13.3	1.5	0.6	0.7	0.9	1.5	1.7	2.5	3.8
1999.....	91.2	13.3	1.6	0.6	0.8	1.1	1.2	1.6	2.3	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

X. Diseases of the respiratory system (J00-J98)

White, male

2004.....	82.8	13.5	1.5	0.4	0.7	1.0	1.6	1.5	2.4	3.5
2003.....	85.8	16.9	2.0	0.6	0.8	0.9	1.7	1.9	2.6	4.1
2002.....	86.7	14.6	1.7	0.6	0.6	1.1	1.4	1.6	2.5	3.7
2001.....	85.7	14.3	1.5	0.6	0.7	0.8	1.6	1.4	2.3	4.1
2000.....	87.7	15.1	1.3	0.7	0.7	1.0	1.7	1.7	2.5	3.9
1999.....	88.8	14.7	1.5	0.6	0.9	1.2	1.4	1.7	2.5	4.1

White, female

2004.....	89.9	10.9	1.4	0.4	0.5	0.8	1.0	1.3	1.7	3.3
2003.....	95.2	11.8	1.6	0.5	0.8	0.9	1.3	1.3	2.1	3.6
2002.....	94.7	9.2	1.2	0.5	0.8	0.8	1.1	1.7	2.1	3.9
2001.....	93.7	10.2	1.6	0.4	0.6	0.8	1.0	1.6	2.2	3.5
2000.....	94.6	11.3	1.6	0.5	0.7	0.8	1.2	1.7	2.4	3.6
1999.....	93.6	11.8	1.7	0.6	0.6	1.0	1.1	1.5	2.2	3.7

Black, both sexes

2004.....	44.2	33.4	3.7	1.5	1.5	2.2	2.6	3.5	4.8	8.5
2003.....	45.9	32.7	3.5	1.4	1.7	2.4	3.4	4.4	5.7	8.5
2002.....	46.7	34.1	4.1	1.2	2.3	2.5	2.9	3.5	5.6	9.4
2001.....	47.2	39.6	3.2	1.2	1.5	2.0	3.6	4.3	6.8	11.1
2000.....	48.0	37.7	3.1	1.2	1.9	2.5	3.4	4.4	6.1	9.6
1999.....	49.2	41.6	4.0	1.1	2.1	2.0	3.6	5.7	6.9	10.2

Black, male

2004.....	47.9	37.1	3.5	2.2	1.8	2.6	2.6	4.9	6.1	9.9
2003.....	50.0	38.7	4.5	1.8	2.0	3.1	3.9	5.5	7.4	8.8
2002.....	50.2	39.2	4.7	*	2.4	3.4	3.2	4.0	6.3	10.7
2001.....	50.5	43.7	3.8	1.2	1.7	2.6	3.5	4.6	7.8	11.3
2000.....	52.4	39.3	3.9	1.4	2.2	3.1	3.7	5.9	7.7	10.7
1999.....	54.0	44.6	5.1	1.4	2.5	2.3	4.3	5.6	7.3	10.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ years	year 2/ years	years	years	years	years	years	years	years	years

X. Diseases of the respiratory system (J00-J98)

Black, female

2004.....	40.9	29.6	3.8	*	1.2	1.8	2.5	2.2	3.6	7.3
2003.....	42.1	26.6	2.4	*	1.3	1.7	2.9	3.3	4.2	8.2
2002.....	43.5	28.8	3.5	1.4	2.2	1.5	2.7	3.1	4.9	8.3
2001.....	44.1	35.3	2.5	*	1.4	1.4	3.6	4.0	5.9	10.9
2000.....	44.1	36.0	2.2	*	1.5	1.9	3.2	3.0	4.6	8.6
1999.....	44.8	38.6	2.9	*	1.6	1.8	2.9	5.7	6.5	9.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of the respiratory system (J10-J98)

All races 3/, both sexes

2004.....	77.3	15.7	1.8	0.6	0.8	1.0	1.5	1.6	2.3	4.0
2003.....	81.1	17.1	2.1	0.7	0.9	1.1	1.8	1.9	2.7	4.3
2002.....	81.3	15.5	1.8	0.7	0.9	1.1	1.4	1.9	2.6	4.4
2001.....	80.7	17.0	1.7	0.6	0.8	1.0	1.6	1.9	2.8	4.7
2000.....	82.0	16.9	1.7	0.7	0.9	1.1	1.7	2.0	2.9	4.5
1999.....	82.3	17.8	2.0	0.7	0.9	1.2	1.6	2.1	2.9	4.6

All races 3/, male

2004.....	75.1	17.4	1.8	0.7	0.9	1.2	1.7	1.9	2.7	4.2
2003.....	78.1	20.1	2.4	0.8	1.0	1.3	2.0	2.3	3.0	4.5
2002.....	78.9	18.6	2.1	0.7	0.9	1.4	1.6	1.9	2.8	4.5
2001.....	78.1	19.4	1.8	0.6	0.8	1.1	1.9	1.8	2.9	4.9
2000.....	80.1	18.7	1.7	0.8	0.9	1.3	1.9	2.2	3.1	4.7
1999.....	81.3	19.2	2.1	0.7	1.1	1.3	1.7	2.1	3.0	4.8

All races 3/, female

2004.....	79.4	14.0	1.8	0.4	0.6	0.9	1.2	1.4	1.9	3.7
2003.....	83.9	13.9	1.8	0.6	0.9	1.0	1.5	1.6	2.3	4.2
2002.....	83.7	12.4	1.4	0.6	1.0	0.9	1.3	1.8	2.5	4.3
2001.....	83.2	14.6	1.7	0.6	0.7	0.9	1.4	1.9	2.6	4.4
2000.....	83.9	15.1	1.6	0.6	0.8	0.9	1.5	1.8	2.7	4.2
1999.....	83.3	16.4	1.9	0.6	0.7	1.1	1.4	2.0	2.8	4.4

White, both sexes

2004.....	86.3	12.1	1.4	0.3	0.6	0.9	1.3	1.4	2.1	3.4
2003.....	90.5	14.1	1.7	0.6	0.8	0.9	1.5	1.6	2.3	3.9
2002.....	90.7	11.9	1.3	0.6	0.7	0.9	1.2	1.7	2.3	3.8
2001.....	89.7	12.1	1.4	0.5	0.7	0.8	1.3	1.5	2.2	3.8
2000.....	91.1	13.1	1.4	0.6	0.7	0.9	1.4	1.7	2.4	3.8
1999.....	91.2	13.1	1.5	0.6	0.7	1.1	1.2	1.6	2.3	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of the respiratory system (J10-J98)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
White, male										
2004.....	82.7	13.2	1.4	0.4	0.7	1.0	1.6	1.5	2.4	3.5
2003.....	85.8	16.4	1.9	0.6	0.8	0.9	1.7	1.8	2.5	4.1
2002.....	86.7	14.5	1.6	0.6	0.6	1.1	1.4	1.6	2.5	3.7
2001.....	85.6	14.1	1.4	0.5	0.7	0.8	1.6	1.4	2.3	4.1
2000.....	87.6	14.8	1.3	0.7	0.7	1.0	1.7	1.7	2.4	3.9
1999.....	88.7	14.5	1.4	0.6	0.9	1.2	1.4	1.7	2.5	4.1
White, female										
2004.....	89.9	10.8	1.3	0.3	0.5	0.7	1.0	1.3	1.7	3.3
2003.....	95.2	11.6	1.5	0.5	0.8	0.8	1.3	1.3	2.1	3.6
2002.....	94.7	9.0	1.1	0.5	0.8	0.8	1.1	1.7	2.1	3.9
2001.....	93.6	10.0	1.5	0.4	0.6	0.8	1.0	1.6	2.2	3.5
2000.....	94.5	11.3	1.6	0.5	0.7	0.7	1.2	1.7	2.4	3.6
1999.....	93.5	11.7	1.6	0.6	0.6	1.0	1.1	1.5	2.2	3.7
Black, both sexes										
2004.....	44.2	33.4	3.7	1.5	1.5	2.2	2.6	3.5	4.8	8.4
2003.....	45.8	32.3	3.4	1.4	1.7	2.4	3.4	4.4	5.7	8.4
2002.....	46.6	33.5	4.0	1.2	2.3	2.5	2.9	3.5	5.6	9.4
2001.....	47.1	39.1	3.0	1.2	1.5	1.9	3.6	4.3	6.8	11.1
2000.....	48.0	37.6	3.0	1.2	1.8	2.5	3.3	4.3	6.1	9.6
1999.....	49.1	41.0	3.9	1.0	2.1	2.0	3.6	5.7	6.7	10.1
Black, male										
2004.....	47.8	37.1	3.5	2.2	1.8	2.5	2.6	4.9	6.1	9.8
2003.....	49.9	38.4	4.4	1.8	2.0	3.1	3.8	5.5	7.4	8.7
2002.....	50.2	38.3	4.7	*	2.4	3.4	3.1	4.0	6.3	10.7
2001.....	50.4	42.8	3.6	1.2	1.7	2.5	3.5	4.6	7.8	11.2
2000.....	52.4	39.3	3.9	1.4	2.2	3.1	3.5	5.8	7.7	10.7
1999.....	53.9	43.9	4.8	1.3	2.5	2.2	4.3	5.6	7.0	10.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of the respiratory system (J10-J98)

Black, female

2004.....	40.9	29.6	3.8	*	1.2	1.8	2.5	2.2	3.6	7.2
2003.....	42.1	26.0	2.3	*	1.3	1.7	2.9	3.3	4.2	8.1
2002.....	43.4	28.5	3.4	1.4	2.2	1.5	2.6	3.1	4.9	8.3
2001.....	44.1	35.3	2.5	*	1.3	1.4	3.6	4.0	5.9	10.9
2000.....	44.0	35.7	2.2	*	1.5	1.9	3.1	3.0	4.6	8.6
1999.....	44.7	37.9	2.9	*	1.6	1.8	2.9	5.7	6.4	9.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
All races 3/, both sexes										
2004.....	19.9	6.3	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.4
2003.....	21.8	7.3	0.7	0.2	0.3	0.3	0.6	0.8	1.0	1.7
2002.....	22.5	6.3	0.7	0.2	0.2	0.3	0.4	0.7	1.0	1.7
2001.....	21.7	7.2	0.7	0.2	0.2	0.3	0.6	0.6	1.0	1.8
2000.....	22.6	7.4	0.6	0.2	0.2	0.3	0.6	0.7	1.1	1.7
1999.....	22.2	8.1	0.8	0.2	0.2	0.3	0.5	0.7	1.0	2.0
All races 3/, male										
2004.....	18.3	7.5	0.5	*	0.2	0.4	0.7	0.7	1.2	1.7
2003.....	19.6	8.5	0.7	0.2	0.2	0.4	0.7	0.9	1.3	1.9
2002.....	20.2	7.6	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8
2001.....	19.5	7.8	0.7	0.2	0.2	0.3	0.7	0.6	1.2	2.0
2000.....	20.3	8.1	0.6	0.2	*	0.3	0.7	0.8	1.2	1.8
1999.....	19.8	8.5	0.7	0.2	0.2	0.3	0.6	0.8	1.1	2.2
All races 3/, female										
2004.....	21.5	5.0	0.8	*	*	0.2	0.4	0.4	0.7	1.2
2003.....	23.9	6.0	0.6	0.2	0.3	0.2	0.6	0.6	0.8	1.5
2002.....	24.8	5.0	0.6	*	0.2	0.3	0.4	0.7	0.8	1.5
2001.....	23.8	6.7	0.7	*	0.2	0.3	0.5	0.6	0.8	1.6
2000.....	24.8	6.6	0.6	*	*	0.3	0.6	0.6	0.9	1.7
1999.....	24.6	7.7	0.8	*	*	0.3	0.5	0.6	0.8	1.8
White, both sexes										
2004.....	21.6	4.7	0.5	*	0.2	0.3	0.5	0.5	0.9	1.2
2003.....	23.7	6.0	0.6	0.2	0.3	0.3	0.7	0.6	0.9	1.5
2002.....	24.6	5.0	0.5	0.2	0.2	0.3	0.4	0.7	0.9	1.4
2001.....	23.5	5.1	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.5
2000.....	24.4	5.6	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.6
1999.....	24.1	5.9	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
White, male										
2004.....	19.5	5.8	0.4	*	*	0.4	0.7	0.6	1.1	1.4
2003.....	20.9	7.1	0.7	*	0.3	0.4	0.7	0.8	1.1	1.7
2002.....	21.7	6.4	0.5	*	*	0.4	0.5	0.6	1.0	1.5
2001.....	20.6	5.8	0.6	*	*	0.3	0.7	0.5	1.0	1.6
2000.....	21.5	6.1	0.4	0.3	*	0.3	0.6	0.6	1.0	1.7
1999.....	21.0	6.7	0.4	*	0.2	0.3	0.5	0.6	0.9	1.9
White, female										
2004.....	23.6	3.5	0.6	*	*	*	0.3	0.4	0.7	1.0
2003.....	26.5	4.8	0.5	*	0.3	*	0.6	0.5	0.7	1.3
2002.....	27.4	3.6	0.5	*	*	*	0.4	0.7	0.7	1.4
2001.....	26.2	4.4	0.6	*	*	0.3	0.3	0.6	0.7	1.4
2000.....	27.3	5.0	0.6	*	*	*	0.5	0.6	0.9	1.5
1999.....	27.1	5.2	0.7	*	*	0.3	0.4	0.4	0.6	1.6
Black, both sexes										
2004.....	14.6	13.5	1.2	*	*	*	0.8	0.9	1.5	3.0
2003.....	15.1	13.0	0.9	*	*	*	0.7	1.6	2.0	3.3
2002.....	15.5	12.7	1.5	*	*	*	0.7	1.0	2.0	3.5
2001.....	15.5	16.3	1.0	*	*	*	1.0	1.2	2.0	3.7
2000.....	16.1	16.4	1.2	*	*	*	1.0	1.4	2.0	2.9
1999.....	16.1	18.3	1.7	*	*	*	1.3	1.7	2.2	4.0
Black, male										
2004.....	14.9	15.1	*	*	*	*	*	*	1.9	3.7
2003.....	15.8	15.2	*	*	*	*	*	1.6	2.8	3.5
2002.....	15.3	13.7	1.6	*	*	*	*	*	2.4	4.0
2001.....	15.8	15.9	*	*	*	*	*	*	2.7	4.5
2000.....	16.5	17.9	1.7	*	*	*	*	2.0	2.5	3.0
1999.....	16.3	17.9	1.9	*	*	*	*	1.9	2.5	4.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumonia (J12-J18)

Black, female

2004.....	14.4	11.7	1.7	*	*	*	*	*	2.3
2003.....	14.6	10.8	*	*	*	*	1.5	1.4	3.1
2002.....	15.6	11.8	*	*	*	*	*	1.6	3.1
2001.....	15.1	16.7	*	*	*	*	*	1.4	3.0
2000.....	15.8	14.9	*	*	*	*	*	1.6	2.7
1999.....	15.9	18.8	*	*	*	*	1.5	1.9	3.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Viral pneumonia, not elsewhere classified (J12)

All races 3/, both sexes

2004.....	0.1	1.3	0.1	*	*	*	*	*	*
2003.....	0.1	1.6	*	*	*	*	*	*	*
2002.....	0.1	1.4	*	*	*	*	*	*	*
2001.....	0.1	1.9	*	*	*	*	*	*	*
2000.....	0.1	1.4	*	*	*	*	*	*	*
1999.....	0.1	2.1	0.1	*	*	*	*	*	*

All races 3/, male

2004.....	0.1	1.7	*	*	*	*	*	*	*
2003.....	0.1	1.8	*	*	*	*	*	*	*
2002.....	0.1	1.8	*	*	*	*	*	*	*
2001.....	0.1	2.0	*	*	*	*	*	*	*
2000.....	0.1	1.7	*	*	*	*	*	*	*
1999.....	0.1	2.2	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*
2003.....	0.1	1.3	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*
2001.....	0.1	1.7	*	*	*	*	*	*	*
2000.....	0.2	1.2	*	*	*	*	*	*	*
1999.....	0.2	1.9	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	1.0	*	*	*	*	*	*	*
2003.....	0.1	1.4	*	*	*	*	*	*	*
2002.....	0.1	1.2	*	*	*	*	*	*	*
2001.....	0.1	1.3	*	*	*	*	*	*	*
2000.....	0.1	1.1	*	*	*	*	*	*	*
1999.....	0.1	1.7	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Pneumonia due to other or unspecified organisms (J16,J18)

All races 3/, both sexes

2004.....	18.4	4.1	0.4	0.1	0.1	0.3	0.5	0.4	0.8	1.2
2003.....	20.0	4.6	0.4	0.1	0.2	0.2	0.6	0.7	0.9	1.4
2002.....	20.8	3.9	0.5	0.1	0.2	0.3	0.4	0.6	0.8	1.4
2001.....	20.0	4.6	0.5	0.2	0.2	0.3	0.5	0.5	0.9	1.5
2000.....	20.8	5.1	0.4	0.2	0.1	0.2	0.5	0.6	0.9	1.4
1999.....	20.7	4.7	0.5	0.1	0.2	0.3	0.5	0.6	0.9	1.6

All races 3/, male

2004.....	16.7	4.9	0.3	*	*	0.4	0.6	0.6	1.0	1.5
2003.....	17.8	5.5	0.4	*	0.2	0.3	0.6	0.8	1.1	1.6
2002.....	18.5	4.6	0.5	*	*	0.3	0.5	0.5	1.0	1.6
2001.....	17.8	4.9	0.5	*	0.2	0.3	0.6	0.5	1.0	1.7
2000.....	18.5	5.3	0.4	*	*	0.3	0.6	0.7	1.0	1.5
1999.....	18.3	5.0	0.5	*	*	0.3	0.5	0.6	1.0	1.8

All races 3/, female

2004.....	20.0	3.3	0.5	*	*	*	0.3	0.3	0.6	0.9
2003.....	22.2	3.8	0.5	*	*	*	0.5	0.5	0.7	1.3
2002.....	23.0	3.1	0.4	*	*	0.3	0.3	0.6	0.7	1.3
2001.....	22.2	4.2	0.5	*	*	0.3	0.4	0.5	0.7	1.3
2000.....	23.1	4.9	0.4	*	*	0.2	0.5	0.4	0.8	1.4
1999.....	23.0	4.3	0.5	*	*	0.3	0.4	0.5	0.7	1.4

White, both sexes

2004.....	19.9	2.9	0.3	*	*	0.3	0.5	0.4	0.8	1.0
2003.....	21.8	3.8	0.4	*	0.2	0.2	0.6	0.6	0.8	1.3
2002.....	22.7	2.9	0.4	0.1	0.2	0.3	0.4	0.5	0.7	1.2
2001.....	21.7	3.0	0.5	0.2	0.2	0.2	0.5	0.4	0.7	1.3
2000.....	22.6	3.9	0.3	0.2	0.1	0.2	0.5	0.5	0.8	1.3
1999.....	22.5	3.3	0.4	0.1	0.2	0.3	0.4	0.5	0.7	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumonia due to other or unspecified organisms (J16,J18)

White, male

2004.....	17.8	3.4	*	*	*	0.4	0.7	0.5	0.9	1.2
2003.....	19.0	4.5	0.4	*	*	0.3	0.6	0.7	0.9	1.4
2002.....	19.9	3.7	0.4	*	*	0.3	0.5	0.5	0.9	1.3
2001.....	18.9	3.5	0.5	*	*	*	0.6	0.4	0.8	1.4
2000.....	19.6	4.3	*	*	*	0.2	0.6	0.6	0.9	1.4
1999.....	19.5	3.8	*	*	*	0.3	0.4	0.5	0.8	1.6

White, female

2004.....	22.0	2.5	0.3	*	*	*	*	0.3	0.6	0.8
2003.....	24.6	3.0	0.4	*	*	*	0.5	0.4	0.6	1.1
2002.....	25.5	2.0	0.3	*	*	*	0.3	0.5	0.6	1.2
2001.....	24.5	2.5	0.4	*	*	0.3	0.3	0.5	0.6	1.1
2000.....	25.5	3.5	0.4	*	*	*	0.4	0.4	0.7	1.2
1999.....	25.4	2.8	0.4	*	*	0.3	0.3	0.4	0.6	1.2

Black, both sexes

2004.....	13.2	10.3	1.0	*	*	*	0.8	0.7	1.2	2.6
2003.....	13.7	8.8	*	*	*	*	0.7	1.3	1.9	2.9
2002.....	14.2	8.2	1.0	*	*	*	*	0.9	1.6	3.0
2001.....	14.1	11.8	0.8	*	*	*	1.0	1.1	1.7	3.3
2000.....	14.7	11.2	0.8	*	*	*	0.9	1.2	1.9	2.4
1999.....	14.7	11.4	1.1	*	*	*	1.1	1.4	1.9	3.4

Black, male

2004.....	13.4	12.5	*	*	*	*	*	*	*	3.5
2003.....	14.2	10.1	*	*	*	*	*	*	2.6	3.1
2002.....	13.9	8.1	*	*	*	*	*	*	2.2	3.3
2001.....	14.3	11.1	*	*	*	*	*	*	2.4	3.8
2000.....	14.9	10.6	*	*	*	*	*	1.8	2.3	2.5
1999.....	14.8	12.0	*	*	*	*	*	1.6	2.4	3.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumonia due to other or unspecified organisms (J16,J18)

Black, female

2004.....	13.0	8.0	*	*	*	*	*	*	1.8
2003.....	13.3	7.4	*	*	*	*	*	1.4	2.7
2002.....	14.4	8.2	*	*	*	*	*	*	2.8
2001.....	14.0	12.6	*	*	*	*	*	*	2.8
2000.....	14.6	11.9	*	*	*	*	*	1.5	2.4
1999.....	14.6	10.7	*	*	*	*	*	1.5	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic lower respiratory diseases (J40-J47)

All races 3/, both sexes

2004.....	41.5	0.9	0.3	0.2	0.3	0.4	0.4	0.6	0.7	1.3
2003.....	43.5	0.8	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
2002.....	43.3	1.0	0.4	0.2	0.4	0.5	0.5	0.6	0.9	1.6
2001.....	43.2	1.0	0.3	0.2	0.3	0.4	0.5	0.5	0.9	1.6
2000.....	43.4	0.9	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
1999.....	44.5	0.9	0.4	0.2	0.4	0.5	0.5	0.7	0.9	1.3

All races 3/, male

2004.....	40.6	1.2	0.4	0.3	0.4	0.5	0.5	0.6	0.9	1.2
2003.....	42.4	1.1	0.5	0.2	0.5	0.5	0.6	0.7	0.9	1.3
2002.....	42.9	1.1	0.6	0.2	0.5	0.6	0.5	0.7	0.9	1.5
2001.....	42.7	1.1	0.4	0.2	0.3	0.4	0.6	0.6	0.9	1.5
2000.....	43.5	1.2	0.4	0.2	0.5	0.6	0.6	0.7	0.9	1.3
1999.....	45.6	*	0.4	0.3	0.6	0.6	0.7	0.7	0.8	1.2

All races 3/, female

2004.....	42.5	*	*	*	0.3	0.3	0.4	0.5	0.5	1.4
2003.....	44.4	*	*	*	0.3	0.3	0.4	0.5	0.7	1.5
2002.....	43.7	*	0.3	*	0.4	0.3	0.5	0.5	0.9	1.7
2001.....	43.7	*	*	*	0.2	0.3	0.4	0.5	0.9	1.7
2000.....	43.2	*	0.3	0.2	0.4	0.3	0.5	0.6	0.8	1.4
1999.....	43.4	*	0.3	0.2	0.3	0.5	0.4	0.8	1.0	1.4

White, both sexes

2004.....	47.4	0.7	0.2	*	0.2	0.2	0.3	0.4	0.5	1.1
2003.....	49.5	0.7	0.2	*	0.2	0.2	0.4	0.5	0.6	1.2
2002.....	49.2	0.8	0.2	*	0.2	0.3	0.4	0.5	0.7	1.3
2001.....	49.0	0.7	0.2	*	0.2	0.3	0.3	0.4	0.6	1.2
2000.....	49.0	0.8	0.2	0.1	0.2	0.3	0.4	0.4	0.6	1.0
1999.....	50.2	*	0.3	0.2	0.3	0.4	0.4	0.5	0.6	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic lower respiratory diseases (J40-J47)

White, male

2004.....	45.5	*	*	*	0.3	0.3	0.4	0.4	0.6	1.0
2003.....	47.4	1.3	*	*	0.3	0.2	0.5	0.4	0.5	1.1
2002.....	47.8	*	0.3	*	*	0.4	0.4	0.5	0.8	1.1
2001.....	47.6	*	*	*	*	*	0.3	0.4	0.5	1.2
2000.....	48.3	*	*	*	0.3	0.3	0.5	0.5	0.6	0.9
1999.....	50.6	*	*	*	0.4	0.4	0.4	0.4	0.5	1.0

White, female

2004.....	49.3	*	*	*	*	*	0.3	0.4	0.4	1.2
2003.....	51.5	*	*	*	*	*	0.3	0.5	0.7	1.3
2002.....	50.5	*	*	*	*	*	0.4	0.4	0.7	1.5
2001.....	50.3	*	*	*	*	0.3	*	0.3	0.7	1.2
2000.....	49.7	*	*	*	*	*	0.3	0.4	0.6	1.2
1999.....	49.8	*	*	*	*	0.4	0.4	0.5	0.7	1.2

Black, both sexes

2004.....	19.2	*	0.8	0.9	1.0	1.3	1.2	1.7	1.9	3.0
2003.....	20.2	*	1.0	0.7	1.1	1.3	1.3	1.7	2.1	2.8
2002.....	20.7	*	1.2	0.6	1.6	1.4	1.2	1.6	2.2	3.8
2001.....	20.4	*	0.9	0.7	0.9	1.0	1.6	1.8	3.0	4.5
2000.....	20.8	*	0.8	0.7	1.4	1.4	1.3	1.9	2.4	3.5
1999.....	21.9	*	*	0.6	1.5	1.2	1.5	2.5	2.7	3.2

Black, male

2004.....	22.0	*	*	1.3	1.3	1.6	1.3	2.4	2.6	3.1
2003.....	23.3	*	1.6	*	1.5	1.6	1.5	2.4	3.1	2.8
2002.....	24.1	*	1.6	*	1.7	2.2	1.5	2.2	2.3	4.5
2001.....	23.6	*	*	*	*	1.4	1.8	2.1	3.4	4.3
2000.....	24.3	*	*	*	1.7	2.0	1.5	2.4	3.1	3.7
1999.....	26.2	*	*	*	2.1	1.5	2.2	2.6	2.7	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

Black, female

2004.....	16.6	*	*	*	*	*	*	1.4	2.9
2003.....	17.3	*	*	*	*	*	*	*	2.9
2002.....	17.7	*	*	*	1.4	*	*	2.1	3.2
2001.....	17.5	*	*	*	*	*	1.4	1.5	2.6
2000.....	17.6	*	*	*	*	*	*	1.4	1.7
1999.....	18.0	*	*	*	*	*	*	2.4	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

All races 3/, both sexes

2004.....	41.2	0.9	0.3	0.2	0.3	0.4	0.4	0.6	0.7	1.3
2003.....	43.1	0.8	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
2002.....	42.9	1.0	0.4	0.2	0.4	0.5	0.5	0.6	0.9	1.5
2001.....	42.9	1.0	0.3	0.2	0.3	0.4	0.5	0.5	0.9	1.6
2000.....	43.0	0.9	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.3
1999.....	44.2	0.9	0.3	0.2	0.4	0.5	0.5	0.7	0.9	1.3

All races 3/, male

2004.....	40.4	1.2	0.4	0.3	0.4	0.5	0.5	0.6	0.8	1.2
2003.....	42.3	1.1	0.5	0.2	0.5	0.4	0.6	0.7	0.8	1.3
2002.....	42.7	1.1	0.6	0.2	0.5	0.6	0.5	0.7	0.9	1.4
2001.....	42.5	1.1	0.4	0.2	0.3	0.4	0.5	0.6	0.9	1.5
2000.....	43.3	1.2	0.4	0.2	0.5	0.5	0.6	0.7	0.9	1.3
1999.....	45.4	*	0.4	0.3	0.6	0.6	0.6	0.7	0.8	1.2

All races 3/, female

2004.....	42.0	*	*	*	0.3	0.3	0.4	0.5	0.5	1.3
2003.....	44.0	*	*	*	0.3	0.3	0.4	0.5	0.7	1.5
2002.....	43.2	*	0.3	*	0.4	0.3	0.5	0.5	0.8	1.6
2001.....	43.2	*	*	*	0.2	0.3	0.4	0.5	0.9	1.6
2000.....	42.8	*	0.3	0.2	0.4	0.3	0.4	0.5	0.7	1.4
1999.....	43.0	*	0.3	0.2	0.3	0.5	0.4	0.7	1.0	1.4

White, both sexes

2004.....	47.0	0.7	0.2	*	0.2	0.2	0.3	0.4	0.5	1.1
2003.....	49.1	0.7	0.2	*	0.2	0.2	0.4	0.5	0.6	1.2
2002.....	48.8	0.8	0.2	*	0.2	0.3	0.4	0.4	0.7	1.3
2001.....	48.6	0.7	0.2	*	0.2	0.2	0.3	0.3	0.6	1.2
2000.....	48.7	0.8	0.2	0.1	0.2	0.2	0.4	0.4	0.6	1.0
1999.....	49.8	*	0.3	0.2	0.3	0.4	0.4	0.4	0.6	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

White, male

2004.....	45.3	*	*	*	0.3	0.3	0.3	0.4	0.6	1.0
2003.....	47.2	1.3	*	*	0.3	*	0.5	0.4	0.5	1.1
2002.....	47.6	*	0.3	*	*	0.4	0.3	0.5	0.7	1.1
2001.....	47.4	*	*	*	*	*	0.3	0.4	0.5	1.2
2000.....	48.1	*	*	*	0.3	0.3	0.5	0.4	0.6	0.9
1999.....	50.3	*	*	*	0.4	0.4	0.4	0.4	0.5	0.9

White, female

2004.....	48.7	*	*	*	*	*	0.3	0.4	0.4	1.2
2003.....	51.0	*	*	*	*	*	0.3	0.5	0.6	1.3
2002.....	49.9	*	*	*	*	*	0.4	0.4	0.7	1.5
2001.....	49.8	*	*	*	*	0.3	*	0.3	0.7	1.1
2000.....	49.2	*	*	*	*	*	0.3	0.4	0.6	1.2
1999.....	49.3	*	*	*	*	0.4	0.4	0.5	0.7	1.2

Black, both sexes

2004.....	19.1	*	0.8	0.9	1.0	1.3	1.2	1.7	1.9	2.9
2003.....	20.1	*	1.0	0.7	1.1	1.3	1.3	1.6	2.1	2.8
2002.....	20.7	*	1.2	0.6	1.6	1.4	1.2	1.6	2.2	3.8
2001.....	20.3	*	0.9	0.7	0.9	1.0	1.6	1.7	2.9	4.5
2000.....	20.7	*	0.8	0.7	1.4	1.4	1.3	1.9	2.3	3.5
1999.....	21.8	*	*	0.6	1.5	1.2	1.5	2.4	2.7	3.2

Black, male

2004.....	21.9	*	*	1.3	1.3	1.6	1.3	2.4	2.6	3.0
2003.....	23.3	*	1.6	*	1.5	1.6	1.5	2.3	3.1	2.8
2002.....	24.1	*	1.6	*	1.7	2.2	1.5	2.2	2.3	4.5
2001.....	23.6	*	*	*	*	1.4	1.8	2.1	3.4	4.3
2000.....	24.3	*	*	*	1.7	1.9	1.5	2.4	3.1	3.7
1999.....	26.1	*	*	*	2.1	1.5	2.2	2.6	2.7	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Black, female

2004.....	16.5	*	*	*	*	*	*	1.4	2.9
2003.....	17.3	*	*	*	*	*	*	*	2.9
2002.....	17.6	*	*	*	1.4	*	*	2.1	3.2
2001.....	17.4	*	*	*	*	*	1.4	2.5	4.6
2000.....	17.5	*	*	*	*	*	1.4	1.6	3.3
1999.....	17.9	*	*	*	*	*	2.2	2.8	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Asthma (J45-J46)

All races 3/, both sexes

2004.....	1.3	*	0.2	0.2	0.3	0.4	0.4	0.5	0.5	0.7
2003.....	1.4	*	0.2	0.2	0.4	0.3	0.4	0.5	0.6	0.7
2002.....	1.5	*	0.3	0.2	0.4	0.4	0.4	0.5	0.6	0.9
2001.....	1.5	*	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9
2000.....	1.6	*	0.2	0.2	0.4	0.4	0.5	0.5	0.7	0.8
1999.....	1.7	*	0.2	0.2	0.4	0.5	0.5	0.6	0.7	0.9

All races 3/, male

2004.....	1.0	*	*	0.2	0.4	0.5	0.5	0.5	0.6	0.6
2003.....	1.0	*	0.4	0.2	0.4	0.4	0.5	0.6	0.7	0.6
2002.....	1.1	*	0.4	0.2	0.5	0.6	0.5	0.7	0.6	0.8
2001.....	1.1	*	0.3	0.2	0.3	0.3	0.4	0.5	0.6	0.8
2000.....	1.2	*	0.3	0.2	0.5	0.5	0.5	0.7	0.8	0.7
1999.....	1.2	*	*	0.2	0.6	0.5	0.6	0.6	0.6	0.8

All races 3/, female

2004.....	1.6	*	*	*	0.2	0.3	0.3	0.5	0.4	0.9
2003.....	1.8	*	*	*	0.3	0.3	0.4	0.4	0.6	0.9
2002.....	1.8	*	*	*	0.4	0.2	0.4	0.4	0.6	1.0
2001.....	1.9	*	*	*	0.2	0.3	0.4	0.4	0.8	1.0
2000.....	2.0	*	*	0.2	0.3	0.3	0.4	0.4	0.6	0.9
1999.....	2.1	*	*	*	0.3	0.4	0.4	0.6	0.8	1.0

White, both sexes

2004.....	1.1	*	*	*	0.2	0.2	0.3	0.3	0.3	0.6
2003.....	1.2	*	*	*	0.2	0.2	0.3	0.4	0.4	0.5
2002.....	1.3	*	*	*	0.2	0.3	0.3	0.4	0.5	0.7
2001.....	1.3	*	*	*	0.2	0.2	0.2	0.3	0.4	0.5
2000.....	1.4	*	*	*	0.2	0.2	0.4	0.3	0.5	0.5
1999.....	1.5	*	*	0.1	0.2	0.4	0.3	0.4	0.4	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Asthma (J45-J46)

White, male

2004.....	0.7	*	*	*	*	0.3	0.3	0.3	0.4	0.4
2003.....	0.8	*	*	*	0.3	*	0.3	0.4	0.4	0.4
2002.....	0.9	*	*	*	*	0.3	0.3	0.4	0.5	0.5
2001.....	0.8	*	*	*	*	*	0.2	0.3	0.3	0.5
2000.....	0.9	*	*	*	0.2	0.3	0.4	0.4	0.5	0.4
1999.....	0.9	*	*	*	0.3	0.4	0.4	0.4	0.4	0.6

White, female

2004.....	1.5	*	*	*	*	*	*	0.4	0.3	0.7
2003.....	1.6	*	*	*	*	*	*	0.4	0.5	0.7
2002.....	1.7	*	*	*	*	*	0.3	0.4	0.6	0.9
2001.....	1.7	*	*	*	*	*	*	*	0.6	0.6
2000.....	1.8	*	*	*	*	*	0.3	0.3	0.5	0.6
1999.....	2.0	*	*	*	*	0.4	*	0.4	0.5	0.8

Black, both sexes

2004.....	2.6	*	*	0.7	1.0	1.2	1.1	1.6	1.7	2.0
2003.....	2.7	*	0.9	0.7	1.0	1.1	1.3	1.4	1.9	2.0
2002.....	2.9	*	0.9	*	1.5	1.3	1.2	1.6	1.6	2.9
2001.....	3.0	*	0.8	0.7	0.8	0.9	1.4	1.7	2.6	3.4
2000.....	3.2	*	*	0.7	1.4	1.3	1.2	1.8	2.0	2.7
1999.....	3.2	*	*	0.6	1.4	1.1	1.4	2.3	2.4	2.6

Black, male

2004.....	2.5	*	*	*	1.2	1.5	1.3	2.2	2.3	1.8
2003.....	2.4	*	1.5	*	1.4	1.3	1.4	2.1	2.7	1.9
2002.....	2.8	*	*	*	1.7	2.1	1.5	2.2	1.7	3.4
2001.....	2.6	*	*	*	*	*	1.5	1.9	3.0	3.0
2000.....	2.8	*	*	*	1.7	1.7	*	2.3	2.6	2.7
1999.....	2.8	*	*	*	2.0	1.3	2.0	2.6	2.6	2.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Asthma (J45-J46)

Black, female

2004.....	2.7	*	*	*	*	*	*	*	*	2.2
2003.....	2.9	*	*	*	*	*	*	*	*	2.1
2002.....	3.0	*	*	*	1.4	*	*	*	1.4	2.3
2001.....	3.3	*	*	*	*	*	*	1.4	2.3	3.7
2000.....	3.5	*	*	*	*	*	*	*	1.4	2.6
1999.....	3.5	*	*	*	*	*	*	2.0	2.3	2.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Lung diseases due to external agents (J60-J70)

All races 3/, both sexes

2004.....	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2003.....	6.4	*	*	*	*	*	0.2	0.1	0.2	0.3
2002.....	6.5	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.5	*	*	*	*	*	0.1	0.2	0.2	0.3
2000.....	6.3	*	*	*	*	*	0.1	0.1	0.2	0.3
1999.....	5.9	*	*	*	*	*	0.1	0.2	0.2	0.3

All races 3/, male

2004.....	6.7	*	*	*	*	*	*	0.2	*	0.3
2003.....	6.9	*	*	*	*	*	0.3	*	0.3	0.4
2002.....	7.0	*	*	*	*	*	*	*	0.2	0.4
2001.....	7.0	*	*	*	*	*	*	*	0.3	0.4
2000.....	6.9	*	*	*	*	*	*	*	0.3	0.4
1999.....	6.5	*	*	*	*	*	*	0.2	0.3	0.3

All races 3/, female

2004.....	5.5	*	*	*	*	*	*	*	*	0.2
2003.....	5.9	*	*	*	*	*	*	*	*	0.3
2002.....	6.0	*	*	*	*	*	*	*	*	0.3
2001.....	6.1	*	*	*	*	*	*	*	0.2	0.2
2000.....	5.8	*	*	*	*	*	*	*	*	0.2
1999.....	5.3	*	*	*	*	*	*	*	0.2	0.2

White, both sexes

2004.....	6.7	*	*	*	*	*	*	0.2	0.2	0.3
2003.....	7.0	*	*	*	*	*	0.1	*	0.2	0.3
2002.....	7.2	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	7.1	*	*	*	*	*	*	0.2	0.2	0.3
2000.....	7.0	*	*	*	*	*	*	*	0.2	0.2
1999.....	6.4	*	*	*	*	*	*	0.2	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumonitis due to solids and liquids (J69)

All races 3/, both sexes

2004.....	5.7	*	*	*	*	*	0.1	0.2	0.2	0.3
2003.....	6.0	*	*	*	*	*	0.2	0.1	0.2	0.3
2002.....	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2000.....	5.9	*	*	*	*	*	0.1	0.1	0.2	0.3
1999.....	5.5	*	*	*	*	*	0.1	0.2	0.2	0.3

All races 3/, male

2004.....	5.9	*	*	*	*	*	*	0.2	*	0.3
2003.....	6.1	*	*	*	*	*	0.2	*	0.2	0.4
2002.....	6.3	*	*	*	*	*	*	*	0.2	0.4
2001.....	6.1	*	*	*	*	*	*	*	0.2	0.4
2000.....	6.1	*	*	*	*	*	*	*	0.2	0.4
1999.....	5.7	*	*	*	*	*	*	0.2	0.3	0.3

All races 3/, female

2004.....	5.5	*	*	*	*	*	*	*	*	0.2
2003.....	5.8	*	*	*	*	*	*	*	*	0.3
2002.....	5.9	*	*	*	*	*	*	*	*	0.3
2001.....	6.0	*	*	*	*	*	*	*	0.2	*
2000.....	5.8	*	*	*	*	*	*	*	*	0.2
1999.....	5.3	*	*	*	*	*	*	*	*	0.2

White, both sexes

2004.....	6.3	*	*	*	*	*	*	0.2	0.2	0.3
2003.....	6.5	*	*	*	*	*	0.1	*	0.2	0.3
2002.....	6.7	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.6	*	*	*	*	*	*	0.2	0.2	0.2
2000.....	6.5	*	*	*	*	*	*	*	0.2	0.2
1999.....	5.9	*	*	*	*	*	*	0.1	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumonitis due to food and vomit (J69.0)

All races 3/, both sexes

2004.....	5.7	*	*	*	*	*	0.1	0.2	0.2	0.3
2003.....	6.0	*	*	*	*	*	0.2	0.1	0.2	0.3
2002.....	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2000.....	5.9	*	*	*	*	*	0.1	0.1	0.2	0.3
1999.....	5.5	*	*	*	*	*	0.1	0.2	0.2	0.3

All races 3/, male

2004.....	5.9	*	*	*	*	*	*	0.2	*	0.3
2003.....	6.1	*	*	*	*	*	0.2	*	0.2	0.4
2002.....	6.3	*	*	*	*	*	*	*	0.2	0.4
2001.....	6.1	*	*	*	*	*	*	*	0.2	0.4
2000.....	6.1	*	*	*	*	*	*	*	0.2	0.4
1999.....	5.7	*	*	*	*	*	*	0.2	0.3	0.3

All races 3/, female

2004.....	5.5	*	*	*	*	*	*	*	*	0.2
2003.....	5.8	*	*	*	*	*	*	*	*	0.3
2002.....	5.9	*	*	*	*	*	*	*	*	0.3
2001.....	6.0	*	*	*	*	*	*	*	0.2	*
2000.....	5.8	*	*	*	*	*	*	*	*	0.2
1999.....	5.3	*	*	*	*	*	*	*	*	0.2

White, both sexes

2004.....	6.3	*	*	*	*	*	*	0.2	0.2	0.3
2003.....	6.5	*	*	*	*	*	0.1	*	0.2	0.3
2002.....	6.7	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.6	*	*	*	*	*	*	0.2	0.2	0.2
2000.....	6.5	*	*	*	*	*	*	*	0.2	0.2
1999.....	5.9	*	*	*	*	*	*	0.1	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other diseases of respiratory system (J80-J84,J93-J98)

All races 3/, both sexes

2004.....	8.5	5.9	0.5	0.1	0.1	0.2	0.3	0.3	0.4	0.8
2003.....	8.0	6.2	0.4	*	0.1	0.2	0.3	0.3	0.5	0.7
2002.....	7.9	6.1	0.4	0.1	0.1	0.2	0.3	0.3	0.5	0.7
2001.....	8.3	6.5	0.4	0.1	0.2	0.2	0.4	0.4	0.5	0.8
2000.....	8.3	6.4	0.4	0.1	0.2	0.3	0.3	0.4	0.6	0.9
1999.....	8.1	5.8	0.4	0.2	0.1	0.2	0.2	0.4	0.6	0.8

All races 3/, male

2004.....	8.5	5.9	0.6	*	*	*	0.2	0.3	0.4	0.8
2003.....	7.8	7.1	0.4	*	*	0.2	0.3	0.3	0.5	0.7
2002.....	7.7	7.3	0.5	*	*	0.2	0.3	0.3	0.4	0.7
2001.....	8.1	8.1	0.4	*	0.2	0.2	0.4	0.3	0.4	0.8
2000.....	8.1	7.1	0.4	*	*	0.2	0.4	0.4	0.6	1.0
1999.....	7.9	6.6	0.4	*	*	0.3	0.3	0.3	0.6	0.8

All races 3/, female

2004.....	8.6	5.8	0.3	*	*	0.2	0.3	0.3	0.4	0.8
2003.....	8.1	5.3	0.4	*	*	*	*	0.3	0.6	0.7
2002.....	8.0	4.9	0.3	*	*	0.2	0.3	0.4	0.5	0.8
2001.....	8.6	4.9	0.4	*	*	*	0.3	0.5	0.5	0.9
2000.....	8.4	5.7	0.3	*	*	0.3	0.2	0.4	0.6	0.8
1999.....	8.3	4.9	0.3	*	*	*	*	0.4	0.7	0.8

White, both sexes

2004.....	9.3	4.2	0.4	*	*	0.2	0.3	0.3	0.3	0.6
2003.....	8.7	4.6	0.3	*	*	0.2	0.2	0.3	0.5	0.6
2002.....	8.6	4.2	0.3	*	*	0.2	0.2	0.3	0.4	0.6
2001.....	9.1	4.5	0.4	*	0.2	0.2	0.3	0.3	0.4	0.6
2000.....	9.0	4.9	0.4	0.1	0.2	0.2	0.3	0.4	0.5	0.7
1999.....	8.8	4.1	0.3	0.1	0.1	0.2	0.3	0.3	0.6	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ 2	Under 1 year 2/ 3	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------	-------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other diseases of respiratory system (J80-J84,J93-J98)

Black, female

2004.....	5.5	13.3	*	*	*	*	*	*	*	1.5
2003.....	5.0	10.5	*	*	*	*	*	*	*	1.6
2002.....	5.4	11.8	*	*	*	*	*	*	*	1.6
2001.....	6.2	11.7	*	*	*	*	*	*	*	2.6
2000.....	5.7	15.2	*	*	*	*	*	*	*	1.7
1999.....	5.9	12.1	*	*	*	*	*	*	1.5	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XI. Diseases of the digestive system (K00-K92)

All races 3/, both sexes

2004.....	29.4	14.2	0.8	0.2	0.3	0.4	0.9	1.3	2.9	7.0
2003.....	30.2	13.8	0.8	0.3	0.3	0.5	0.8	1.6	3.2	7.3
2002.....	30.1	12.6	0.8	0.2	0.3	0.4	0.9	1.4	3.2	7.3
2001.....	30.1	11.9	0.8	0.3	0.3	0.4	0.7	1.4	3.2	8.1
2000.....	29.9	13.1	0.7	0.3	0.3	0.4	1.0	1.4	3.3	8.0
1999.....	29.7	13.2	0.7	0.3	0.4	0.5	0.7	1.3	3.5	7.9

All races 3/, male

2004.....	30.0	15.6	0.9	0.3	0.2	0.4	0.9	1.6	3.9	9.0
2003.....	30.9	16.6	0.9	0.3	0.3	0.6	0.8	1.8	4.0	9.0
2002.....	30.7	15.1	0.9	0.3	0.3	0.4	0.9	1.5	4.1	9.1
2001.....	30.6	13.4	0.9	0.3	0.4	0.4	0.8	1.6	4.0	10.3
2000.....	30.3	14.8	0.9	0.3	0.3	0.4	1.1	1.8	4.1	10.2
1999.....	30.4	16.5	0.7	0.2	0.4	0.5	0.7	1.5	4.3	9.9

All races 3/, female

2004.....	28.8	12.8	0.6	*	0.3	0.5	0.8	1.0	2.0	5.0
2003.....	29.4	10.8	0.7	0.2	0.3	0.4	0.7	1.3	2.3	5.6
2002.....	29.5	10.0	0.7	*	0.4	0.4	0.8	1.3	2.3	5.6
2001.....	29.7	10.3	0.6	0.3	0.3	0.3	0.7	1.2	2.4	6.0
2000.....	29.4	11.4	0.5	0.3	0.4	0.3	0.8	1.0	2.5	5.8
1999.....	29.1	9.7	0.8	0.3	0.4	0.4	0.8	1.1	2.7	5.9

White, both sexes

2004.....	31.5	10.7	0.6	0.2	0.2	0.4	0.7	1.1	2.8	6.8
2003.....	32.3	11.0	0.7	0.3	0.3	0.5	0.6	1.3	3.1	7.1
2002.....	32.2	10.1	0.7	0.2	0.3	0.4	0.8	1.2	3.0	7.2
2001.....	32.1	8.7	0.7	0.3	0.3	0.3	0.6	1.2	3.0	7.9
2000.....	31.7	10.0	0.5	0.2	0.3	0.3	0.9	1.2	3.1	7.6
1999.....	31.5	10.1	0.7	0.2	0.4	0.4	0.6	1.2	3.3	7.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XI. Diseases of the digestive system (K00-K92)

White, male

2004.....	31.6	11.4	0.7	0.3	*	0.4	0.7	1.4	3.8	8.6
2003.....	32.6	13.9	0.9	0.3	0.3	0.6	0.7	1.5	4.0	8.9
2002.....	32.4	12.1	0.7	0.3	0.3	0.4	0.9	1.2	4.0	9.0
2001.....	31.9	9.3	0.8	0.3	0.3	0.4	0.6	1.4	3.8	9.9
2000.....	31.7	11.2	0.6	0.3	0.3	0.4	1.0	1.7	3.9	9.8
1999.....	31.7	12.5	0.5	*	0.3	0.5	0.6	1.4	4.0	9.7

White, female

2004.....	31.3	9.9	0.5	*	*	0.4	0.6	0.8	1.9	4.9
2003.....	32.0	8.0	0.6	*	0.3	0.3	0.5	1.1	2.2	5.3
2002.....	32.1	7.9	0.7	*	0.3	0.4	0.6	1.1	1.9	5.4
2001.....	32.2	8.1	0.5	0.3	0.3	0.3	0.5	1.1	2.3	5.9
2000.....	31.6	8.8	0.4	*	0.3	0.3	0.7	0.7	2.2	5.3
1999.....	31.2	7.5	0.8	0.3	0.4	0.4	0.6	1.0	2.5	5.5

Black, both sexes

2004.....	23.7	31.6	1.4	*	*	0.7	1.7	2.2	3.6	8.6
2003.....	24.2	29.0	1.2	*	*	0.7	1.4	3.1	3.6	8.9
2002.....	23.7	25.9	1.4	*	*	0.6	1.5	2.5	4.3	8.0
2001.....	25.1	28.6	1.2	*	0.6	0.6	1.5	2.6	4.3	10.1
2000.....	25.2	29.6	1.5	*	*	*	1.6	2.9	4.7	10.8
1999.....	25.1	29.6	1.1	*	*	0.7	1.8	2.1	4.6	10.1

Black, male

2004.....	26.4	36.2	1.7	*	*	*	1.6	2.5	5.0	11.4
2003.....	27.1	32.4	*	*	*	*	1.6	3.6	4.5	10.0
2002.....	26.0	31.4	1.7	*	*	*	*	2.9	4.9	9.1
2001.....	28.4	33.5	*	*	*	*	*	3.3	5.3	13.2
2000.....	27.7	33.9	2.0	*	*	*	2.1	3.0	5.1	13.6
1999.....	27.8	36.4	*	*	*	*	1.6	2.2	5.7	12.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ year 1/	year 2/ year 2/	years	years	years	years	years	years	years	years

XI. Diseases of the digestive system (K00-K92)

Black, female

2004.....	21.1	26.9	*	*	*	*	1.7	2.0	2.3	6.2
2003.....	21.6	25.4	*	*	*	*	1.3	2.7	2.9	7.9
2002.....	21.5	20.3	*	*	*	*	1.8	2.1	3.8	7.0
2001.....	22.1	23.3	*	*	*	*	1.7	2.0	3.4	7.3
2000.....	23.0	25.1	*	*	*	*	*	2.8	4.3	8.4
1999.....	22.7	22.5	*	*	*	*	1.9	2.0	3.6	8.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Diseases of other parts of digestive system (K20-K92)

All races 3/, both sexes

2004.....	29.3	14.1	0.8	0.2	0.3	0.4	0.9	1.3	2.9	7.0
2003.....	30.1	13.7	0.8	0.3	0.3	0.5	0.7	1.5	3.2	7.3
2002.....	30.0	12.5	0.8	0.2	0.3	0.4	0.9	1.4	3.2	7.3
2001.....	30.1	11.8	0.8	0.3	0.3	0.4	0.7	1.4	3.2	8.1
2000.....	29.8	13.0	0.7	0.3	0.3	0.4	1.0	1.4	3.3	8.0
1999.....	29.7	13.1	0.7	0.3	0.4	0.5	0.7	1.3	3.5	7.9

All races 3/, male

2004.....	29.9	15.5	0.9	0.3	0.2	0.4	0.9	1.6	3.9	8.9
2003.....	30.9	16.5	0.9	0.3	0.3	0.6	0.8	1.8	4.0	9.0
2002.....	30.6	15.0	0.9	0.3	0.3	0.4	0.9	1.5	4.1	9.1
2001.....	30.5	13.3	0.9	0.3	0.4	0.4	0.8	1.6	4.0	10.3
2000.....	30.3	14.8	0.9	0.3	0.3	0.4	1.1	1.8	4.1	10.2
1999.....	30.3	16.4	0.7	0.2	0.4	0.5	0.7	1.5	4.3	9.9

All races 3/, female

2004.....	28.7	12.6	0.6	*	0.3	0.5	0.8	1.0	2.0	5.0
2003.....	29.4	10.8	0.7	0.2	0.3	0.4	0.6	1.3	2.3	5.6
2002.....	29.4	9.9	0.7	*	0.4	0.4	0.8	1.3	2.3	5.6
2001.....	29.7	10.2	0.6	0.3	0.3	0.3	0.7	1.2	2.4	6.0
2000.....	29.4	11.2	0.5	0.3	0.4	0.3	0.8	1.0	2.5	5.8
1999.....	29.0	9.7	0.8	0.3	0.4	0.4	0.7	1.1	2.7	5.9

White, both sexes

2004.....	31.4	10.5	0.6	0.2	0.2	0.4	0.7	1.1	2.8	6.8
2003.....	32.2	11.0	0.7	0.3	0.3	0.5	0.6	1.3	3.1	7.1
2002.....	32.2	10.0	0.7	0.2	0.3	0.4	0.8	1.2	3.0	7.2
2001.....	32.0	8.6	0.7	0.3	0.3	0.3	0.5	1.2	3.0	7.9
2000.....	31.6	9.9	0.5	0.2	0.3	0.3	0.9	1.2	3.0	7.6
1999.....	31.4	10.0	0.7	0.2	0.4	0.4	0.6	1.2	3.3	7.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of other parts of digestive system (K20-K92)

White, male

2004.....	31.6	11.3	0.7	0.3	*	0.4	0.7	1.4	3.7	8.6
2003.....	32.6	13.8	0.9	0.3	0.3	0.6	0.7	1.5	4.0	8.9
2002.....	32.4	12.0	0.7	0.3	0.3	0.4	0.9	1.2	4.0	9.0
2001.....	31.9	9.3	0.8	0.3	0.3	0.4	0.6	1.4	3.8	9.9
2000.....	31.7	11.2	0.6	0.3	0.3	0.4	1.0	1.6	3.9	9.8
1999.....	31.7	12.4	0.5	*	0.3	0.5	0.6	1.4	4.0	9.7

White, female

2004.....	31.2	9.7	0.5	*	*	0.4	0.6	0.8	1.9	4.9
2003.....	31.9	8.0	0.6	*	0.3	0.3	0.5	1.1	2.2	5.3
2002.....	32.0	7.9	0.7	*	0.3	0.4	0.6	1.1	1.9	5.4
2001.....	32.1	8.0	0.5	0.3	0.3	0.3	0.5	1.0	2.3	5.9
2000.....	31.6	8.6	0.4	*	0.3	0.3	0.7	0.7	2.2	5.3
1999.....	31.2	7.5	0.8	0.3	0.4	0.4	0.6	1.0	2.5	5.5

Black, both sexes

2004.....	23.6	31.6	1.4	*	*	0.7	1.7	2.1	3.5	8.6
2003.....	24.2	28.8	1.2	*	*	0.7	1.4	3.1	3.6	8.9
2002.....	23.6	25.9	1.4	*	*	0.6	1.4	2.5	4.2	7.9
2001.....	25.0	28.6	1.2	*	0.6	0.6	1.5	2.6	4.3	10.1
2000.....	25.2	29.4	1.5	*	*	*	1.6	2.9	4.6	10.8
1999.....	25.1	29.6	1.1	*	*	0.7	1.8	2.1	4.6	10.1

Black, male

2004.....	26.4	36.2	1.7	*	*	*	1.6	2.5	4.9	11.3
2003.....	27.1	32.4	*	*	*	*	1.5	3.5	4.5	10.0
2002.....	25.9	31.4	1.7	*	*	*	*	2.9	4.8	9.0
2001.....	28.3	33.5	*	*	*	*	*	3.2	5.3	13.2
2000.....	27.7	33.9	2.0	*	*	*	2.1	3.0	5.1	13.6
1999.....	27.7	36.4	*	*	*	*	1.6	2.2	5.7	12.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Diseases of other parts of digestive system (K20-K92)

Black, female

2004.....	21.1	26.9	*	*	*	*	1.7	1.8	2.3	6.1
2003.....	21.5	25.0	*	*	*	*	1.3	2.7	2.9	7.9
2002.....	21.5	20.3	*	*	*	*	1.7	2.1	3.7	7.0
2001.....	22.0	23.3	*	*	*	*	1.7	2.0	3.4	7.3
2000.....	22.9	24.8	*	*	*	*	*	2.8	4.3	8.4
1999.....	22.6	22.5	*	*	*	*	1.9	2.0	3.6	8.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

All races 3/, both sexes

2004.....	4.9	6.9	0.2	*	*	*	*	*	0.2	0.3
2003.....	5.1	6.8	0.1	*	*	*	0.1	0.1	0.1	0.2
2002.....	5.0	5.8	0.1	*	*	*	0.1	0.1	0.1	0.3
2001.....	5.0	5.8	0.1	*	*	*	*	0.1	0.1	0.2
2000.....	5.1	6.0	0.1	*	*	*	0.1	0.1	0.1	0.3
1999.....	5.0	6.1	0.1	*	*	*	*	0.1	0.2	0.2

All races 3/, male

2004.....	3.5	7.7	0.2	*	*	*	*	*	*	0.3
2003.....	3.7	8.2	*	*	*	*	*	*	*	0.2
2002.....	3.6	6.5	*	*	*	*	*	*	0.2	0.3
2001.....	3.6	6.5	*	*	*	*	*	*	*	0.2
2000.....	3.6	6.3	*	*	*	*	*	*	*	0.3
1999.....	3.5	7.6	*	*	*	*	*	*	*	0.2

All races 3/, female

2004.....	6.3	6.0	*	*	*	*	*	*	0.2	0.2
2003.....	6.5	5.3	*	*	*	*	*	*	*	0.2
2002.....	6.4	5.0	*	*	*	*	*	*	*	0.2
2001.....	6.4	5.1	*	*	*	*	*	*	*	0.2
2000.....	6.5	5.8	*	*	*	*	*	*	*	0.2
1999.....	6.4	4.4	*	*	*	*	*	*	0.2	0.3

White, both sexes

2004.....	5.3	4.6	0.2	*	*	*	*	*	0.1	0.2
2003.....	5.5	4.8	*	*	*	*	*	*	*	0.2
2002.....	5.4	4.2	*	*	*	*	*	*	0.1	0.2
2001.....	5.4	4.1	*	*	*	*	*	*	*	0.2
2000.....	5.4	3.8	*	*	*	*	0.1	*	*	0.2
1999.....	5.4	4.2	*	*	*	*	*	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of intestines and peritoneum (K52,K58-K63)

All races 3/, both sexes

2004.....	1.7	2.0	0.2	*	*	*	*	*	*	0.2
2003.....	1.7	2.4	0.1	*	*	*	*	*	0.1	0.2
2002.....	1.7	2.0	0.2	*	*	*	*	*	*	0.2
2001.....	1.7	1.4	0.2	*	*	*	*	*	*	0.2
2000.....	1.6	1.7	*	*	*	*	*	*	0.1	0.1
1999.....	1.6	1.5	0.2	*	*	*	*	*	*	0.2

All races 3/, male

2004.....	1.3	2.3	*	*	*	*	*	*	*	*
2003.....	1.3	2.9	*	*	*	*	*	*	*	0.2
2002.....	1.3	2.4	*	*	*	*	*	*	*	*
2001.....	1.2	1.6	*	*	*	*	*	*	*	*
2000.....	1.2	1.8	*	*	*	*	*	*	*	*
1999.....	1.2	2.0	*	*	*	*	*	*	*	0.2

All races 3/, female

2004.....	2.1	1.7	*	*	*	*	*	*	*	0.2
2003.....	2.1	1.8	*	*	*	*	*	*	*	*
2002.....	2.0	1.5	*	*	*	*	*	*	*	0.2
2001.....	2.1	1.1	*	*	*	*	*	*	*	0.2
2000.....	2.0	1.6	*	*	*	*	*	*	*	*
1999.....	1.9	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	1.9	1.6	*	*	*	*	*	*	*	0.2
2003.....	1.8	2.1	*	*	*	*	*	*	*	0.1
2002.....	1.8	1.6	0.2	*	*	*	*	*	*	0.2
2001.....	1.8	0.8	*	*	*	*	*	*	*	0.1
2000.....	1.7	1.4	*	*	*	*	*	*	*	0.1
1999.....	1.7	1.2	0.2	*	*	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of liver (K70-K76)

All races 3/, both sexes

2004.....	12.3	2.5	0.2	*	*	0.1	0.3	0.6	1.8	4.9
2003.....	12.6	1.5	0.2	*	*	0.2	0.2	0.7	2.0	5.3
2002.....	12.4	2.2	0.2	*	*	0.1	0.3	0.7	1.9	5.3
2001.....	12.2	1.9	0.2	*	*	0.1	0.3	0.6	2.0	6.1
2000.....	12.0	2.0	0.2	*	*	*	0.4	0.6	2.1	6.0
1999.....	12.0	2.3	0.1	*	*	0.2	0.3	0.6	2.2	6.0

All races 3/, male

2004.....	15.7	2.5	*	*	*	*	0.3	0.7	2.5	6.3
2003.....	16.1	2.2	0.2	*	*	*	0.2	0.9	2.5	6.7
2002.....	15.8	3.1	*	*	*	*	0.3	0.8	2.4	6.8
2001.....	15.7	2.3	*	*	*	*	0.3	0.7	2.5	7.8
2000.....	15.5	2.7	*	*	*	*	0.4	0.8	2.6	7.8
1999.....	15.6	2.9	*	*	*	*	*	0.6	2.8	7.6

All races 3/, female

2004.....	9.0	2.4	*	*	*	*	0.3	0.4	1.2	3.5
2003.....	9.1	*	*	*	*	*	0.3	0.6	1.4	4.0
2002.....	9.0	1.3	*	*	*	*	0.3	0.6	1.4	3.9
2001.....	8.9	1.6	*	*	*	*	0.3	0.5	1.5	4.4
2000.....	8.7	1.3	*	*	*	*	0.3	0.4	1.6	4.2
1999.....	8.6	1.6	*	*	*	*	0.3	0.5	1.6	4.3

White, both sexes

2004.....	13.0	1.9	0.2	*	*	*	0.2	0.5	1.9	4.9
2003.....	13.3	1.2	0.2	*	*	*	0.2	0.6	2.0	5.5
2002.....	13.1	1.9	*	*	*	*	0.2	0.6	1.9	5.4
2001.....	12.9	1.6	0.2	*	*	*	0.2	0.5	2.0	6.1
2000.....	12.6	1.6	*	*	*	*	0.3	0.5	2.0	5.8
1999.....	12.5	1.7	*	*	*	0.1	0.2	0.5	2.1	5.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of liver (K70-K76)

White, male

2004.....	16.6	1.8	*	*	*	*	0.3	0.6	2.6	6.2
2003.....	17.1	1.8	*	*	*	*	*	0.7	2.6	6.9
2002.....	16.8	2.7	*	*	*	*	0.3	0.7	2.4	6.9
2001.....	16.4	1.9	*	*	*	*	*	0.6	2.5	7.7
2000.....	16.2	2.0	*	*	*	*	0.3	0.7	2.6	7.7
1999.....	16.3	2.0	*	*	*	*	*	0.7	2.8	7.6

White, female

2004.....	9.5	1.9	*	*	*	*	*	0.4	1.2	3.5
2003.....	9.7	*	*	*	*	*	*	0.5	1.3	4.0
2002.....	9.5	*	*	*	*	*	*	0.5	1.3	3.9
2001.....	9.4	1.3	*	*	*	*	*	0.4	1.5	4.5
2000.....	9.1	*	*	*	*	*	*	0.3	1.4	3.9
1999.....	8.9	1.4	*	*	*	*	*	0.4	1.5	4.1

Black, both sexes

2004.....	9.8	5.4	*	*	*	*	0.6	0.7	1.4	4.8
2003.....	10.0	3.0	*	*	*	*	0.6	1.4	1.8	5.0
2002.....	9.9	3.7	*	*	*	*	*	1.0	1.9	4.5
2001.....	10.4	3.4	*	*	*	*	*	0.9	1.9	5.8
2000.....	10.4	4.2	*	*	*	*	0.7	1.2	2.3	6.9
1999.....	10.8	5.0	*	*	*	*	*	0.9	2.5	6.3

Black, male

2004.....	12.9	*	*	*	*	*	*	*	1.8	6.1
2003.....	13.3	*	*	*	*	*	*	1.6	1.9	5.8
2002.....	12.6	*	*	*	*	*	*	*	2.2	4.8
2001.....	14.0	*	*	*	*	*	*	*	2.4	7.8
2000.....	13.5	6.4	*	*	*	*	*	*	2.4	8.9
1999.....	14.1	7.2	*	*	*	*	*	*	2.7	7.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of liver (K70-K76)

Black, female

2004.....	7.0	*	*	*	*	*	*	*	3.6
2003.....	7.1	*	*	*	*	*	*	1.7	4.3
2002.....	7.4	*	*	*	*	*	*	1.7	4.2
2001.....	7.2	*	*	*	*	*	*	1.5	4.0
2000.....	7.7	*	*	*	*	*	*	2.3	5.1
1999.....	7.9	*	*	*	*	*	*	2.3	5.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Alcoholic liver disease (K70)

All races 3/, both sexes

2004.....	4.3	*	*	*	*	*	*	0.2	0.9	2.5
2003.....	4.3	*	*	*	*	*	*	0.3	0.9	2.9
2002.....	4.2	*	*	*	*	*	*	0.2	0.9	2.8
2001.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3
2000.....	4.3	*	*	*	*	*	*	0.3	1.1	3.3
1999.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3

All races 3/, male

2004.....	6.3	*	*	*	*	*	*	0.3	1.3	3.3
2003.....	6.4	*	*	*	*	*	*	0.4	1.4	3.7
2002.....	6.2	*	*	*	*	*	*	0.3	1.2	3.6
2001.....	6.4	*	*	*	*	*	*	0.3	1.4	4.4
2000.....	6.5	*	*	*	*	*	*	0.5	1.5	4.3
1999.....	6.5	*	*	*	*	*	*	0.2	1.4	4.3

All races 3/, female

2004.....	2.3	*	*	*	*	*	*	*	0.5	1.8
2003.....	2.2	*	*	*	*	*	*	*	0.5	2.0
2002.....	2.3	*	*	*	*	*	*	*	0.7	2.0
2001.....	2.3	*	*	*	*	*	*	*	0.7	2.3
2000.....	2.2	*	*	*	*	*	*	*	0.6	2.2
1999.....	2.1	*	*	*	*	*	*	*	0.8	2.4

White, both sexes

2004.....	4.6	*	*	*	*	*	*	0.2	1.0	2.6
2003.....	4.5	*	*	*	*	*	*	0.2	1.0	2.9
2002.....	4.4	*	*	*	*	*	*	0.2	0.9	2.8
2001.....	4.5	*	*	*	*	*	*	0.2	1.1	3.4
2000.....	4.5	*	*	*	*	*	*	0.3	1.1	3.2
1999.....	4.4	*	*	*	*	*	*	0.2	1.0	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Fibrosis and cirrhosis of liver (K74)

All races 3/, both sexes

2004.....	4.9	*	*	*	*	*	*	0.3	1.2
2003.....	5.2	*	*	*	*	*	0.2	0.4	1.2
2002.....	5.1	*	*	*	*	*	0.1	0.4	1.4
2001.....	5.0	*	*	*	*	*	0.1	0.4	1.6
2000.....	5.0	*	*	*	*	*	0.1	0.5	1.5
1999.....	5.0	*	*	*	*	*	0.1	0.5	1.4

All races 3/, male

2004.....	5.8	*	*	*	*	*	*	0.5	1.7
2003.....	6.1	*	*	*	*	*	*	0.5	1.6
2002.....	6.0	*	*	*	*	*	*	0.5	1.9
2001.....	6.0	*	*	*	*	*	*	0.5	1.9
2000.....	5.9	*	*	*	*	*	*	0.6	2.0
1999.....	5.9	*	*	*	*	*	*	0.7	2.0

All races 3/, female

2004.....	4.0	*	*	*	*	*	*	0.2	0.7
2003.....	4.2	*	*	*	*	*	*	0.3	0.9
2002.....	4.2	*	*	*	*	*	*	0.3	0.9
2001.....	4.2	*	*	*	*	*	*	0.3	1.2
2000.....	4.1	*	*	*	*	*	*	0.4	0.9
1999.....	4.1	*	*	*	*	*	*	0.4	0.9

White, both sexes

2004.....	5.3	*	*	*	*	*	*	0.3	1.2
2003.....	5.6	*	*	*	*	*	*	0.4	1.3
2002.....	5.5	*	*	*	*	*	*	0.4	1.4
2001.....	5.4	*	*	*	*	*	*	0.4	1.6
2000.....	5.4	*	*	*	*	*	*	0.4	1.5
1999.....	5.3	*	*	*	*	*	*	0.6	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of liver (K71-K72,K75-K76)

All races 3/, both sexes

2004.....	3.1	2.3	0.2	*	*	0.1	0.2	0.3	0.6	1.2
2003.....	3.1	1.5	0.2	*	*	0.1	0.2	0.3	0.6	1.2
2002.....	2.9	2.1	0.2	*	*	*	0.2	0.3	0.5	1.1
2001.....	2.7	1.9	0.2	*	*	*	0.2	0.2	0.5	1.1
2000.....	2.6	1.9	0.2	*	*	*	0.2	0.2	0.5	1.2
1999.....	2.6	2.1	0.1	*	*	0.1	0.2	0.2	0.6	1.2

All races 3/, male

2004.....	3.5	2.3	*	*	*	*	0.2	0.4	0.7	1.3
2003.....	3.6	2.1	0.2	*	*	*	*	0.3	0.6	1.3
2002.....	3.5	2.8	*	*	*	*	*	0.3	0.6	1.2
2001.....	3.2	2.2	*	*	*	*	*	*	0.5	1.4
2000.....	3.0	2.5	*	*	*	*	0.2	*	0.5	1.5
1999.....	3.1	2.7	*	*	*	*	*	0.2	0.7	1.4

All races 3/, female

2004.....	2.6	2.3	*	*	*	*	*	0.3	0.5	1.0
2003.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.1
2002.....	2.3	1.3	*	*	*	*	*	0.3	0.4	1.0
2001.....	2.3	1.5	*	*	*	*	*	0.2	0.5	0.9
2000.....	2.2	1.3	*	*	*	*	*	*	0.5	0.9
1999.....	2.2	1.5	*	*	*	*	0.2	*	0.4	0.9

White, both sexes

2004.....	3.1	1.6	0.2	*	*	*	0.2	0.2	0.6	1.1
2003.....	3.2	1.2	0.2	*	*	*	0.1	0.3	0.6	1.2
2002.....	3.0	1.8	*	*	*	*	*	0.3	0.5	1.1
2001.....	2.8	1.5	0.2	*	*	*	0.1	0.2	0.4	1.0
2000.....	2.6	1.6	*	*	*	*	0.2	0.1	0.5	1.1
1999.....	2.6	1.5	*	*	*	*	0.1	0.2	0.5	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Disorders of biliary tract and pancreas (K83-K86)

All races 3/, both sexes

2004.....	1.7	*	*	*	*	*	0.1	0.3	0.4	0.7
2003.....	1.7	*	*	*	*	*	*	0.2	0.4	0.6
2002.....	1.7	*	*	*	*	*	0.2	0.2	0.4	0.6
2001.....	1.7	*	*	*	*	*	0.1	0.2	0.5	0.7
2000.....	1.7	*	*	*	*	*	0.2	0.3	0.4	0.7
1999.....	1.6	*	*	*	*	*	0.2	0.3	0.4	0.6

All races 3/, male

2004.....	1.8	*	*	*	*	*	*	0.4	0.6	1.0
2003.....	1.8	*	*	*	*	*	*	0.3	0.6	0.8
2002.....	1.9	*	*	*	*	*	0.2	0.3	0.6	0.8
2001.....	1.8	*	*	*	*	*	*	0.3	0.6	1.0
2000.....	1.8	*	*	*	*	*	0.2	0.4	0.6	0.9
1999.....	1.7	*	*	*	*	*	*	0.4	0.6	0.8

All races 3/, female

2004.....	1.6	*	*	*	*	*	*	*	*	0.3
2003.....	1.6	*	*	*	*	*	*	*	0.2	0.4
2002.....	1.5	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.7	*	*	*	*	*	*	*	0.3	0.5
2000.....	1.6	*	*	*	*	*	*	*	0.2	0.4
1999.....	1.5	*	*	*	*	*	*	*	0.3	0.4

White, both sexes

2004.....	1.7	*	*	*	*	*	*	0.2	0.3	0.6
2003.....	1.7	*	*	*	*	*	*	0.2	0.4	0.5
2002.....	1.8	*	*	*	*	*	0.1	0.2	0.4	0.6
2001.....	1.8	*	*	*	*	*	*	0.2	0.4	0.6
2000.....	1.7	*	*	*	*	*	0.2	0.2	0.4	0.6
1999.....	1.6	*	*	*	*	*	*	0.2	0.4	0.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pancreatitis (K85,K86.0-K86.1)

All races 3/, both sexes

2004.....	1.2	*	*	*	*	*	0.1	0.2	0.3	0.6
2003.....	1.2	*	*	*	*	*	*	0.2	0.3	0.6
2002.....	1.2	*	*	*	*	*	0.2	0.2	0.4	0.6
2001.....	1.3	*	*	*	*	*	0.1	0.2	0.4	0.6
2000.....	1.2	*	*	*	*	*	0.2	0.2	0.4	0.6
1999.....	1.2	*	*	*	*	*	0.1	0.2	0.4	0.5

All races 3/, male

2004.....	1.3	*	*	*	*	*	*	0.4	0.6	0.9
2003.....	1.3	*	*	*	*	*	*	0.2	0.5	0.8
2002.....	1.4	*	*	*	*	*	0.2	0.3	0.6	0.8
2001.....	1.4	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	1.3	*	*	*	*	*	0.2	0.3	0.6	0.9
1999.....	1.3	*	*	*	*	*	*	0.3	0.6	0.7

All races 3/, female

2004.....	1.0	*	*	*	*	*	*	*	*	0.2
2003.....	1.1	*	*	*	*	*	*	*	*	0.4
2002.....	1.1	*	*	*	*	*	*	*	*	0.4
2001.....	1.2	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.1	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.1	*	*	*	*	*	*	*	0.3	0.3

White, both sexes

2004.....	1.2	*	*	*	*	*	*	0.2	0.3	0.5
2003.....	1.2	*	*	*	*	*	*	0.1	0.3	0.5
2002.....	1.3	*	*	*	*	*	0.1	0.2	0.3	0.5
2001.....	1.3	*	*	*	*	*	*	0.2	0.3	0.6
2000.....	1.2	*	*	*	*	*	0.2	0.2	0.3	0.6
1999.....	1.2	*	*	*	*	*	*	0.2	0.4	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

All races 3/, both sexes

2004.....	4.9	*	*	*	0.2	0.3	0.5	0.7	0.7	1.1
2003.....	5.0	*	*	*	0.1	0.3	0.5	0.7	0.9	1.1
2002.....	5.0	*	*	*	0.1	0.2	0.5	0.7	0.7	1.2
2001.....	4.9	*	*	*	0.1	0.3	0.6	0.6	0.8	1.3
2000.....	4.9	*	*	*	0.1	0.2	0.6	0.7	0.8	1.1
1999.....	4.8	*	*	*	0.1	0.3	0.5	0.6	0.8	1.2

All races 3/, male

2004.....	2.9	*	*	*	*	0.2	0.3	0.3	0.4	0.7
2003.....	3.0	*	*	*	*	0.2	0.2	0.4	0.4	0.8
2002.....	3.0	*	*	*	*	*	0.2	0.4	0.5	0.7
2001.....	2.9	*	*	*	*	*	0.2	0.3	0.5	1.0
2000.....	2.9	*	*	*	*	*	0.4	0.4	0.5	0.6
1999.....	2.8	*	*	*	*	*	0.2	0.3	0.5	0.8

All races 3/, female

2004.....	6.7	*	*	*	0.2	0.3	0.7	1.1	1.0	1.6
2003.....	6.9	*	*	*	*	0.5	0.8	1.1	1.4	1.4
2002.....	7.0	*	*	*	*	0.4	0.8	1.0	1.0	1.6
2001.....	6.9	*	*	*	*	0.5	0.9	1.0	1.2	1.7
2000.....	6.8	*	*	*	*	0.3	0.8	1.1	1.2	1.6
1999.....	6.7	*	*	*	0.2	0.4	0.7	1.0	1.2	1.7

White, both sexes

2004.....	5.0	*	*	*	0.1	0.2	0.3	0.3	0.4	0.8
2003.....	5.2	*	*	*	*	0.3	0.3	0.4	0.6	0.7
2002.....	5.2	*	*	*	*	0.2	0.3	0.4	0.4	0.8
2001.....	5.1	*	*	*	*	0.3	0.4	0.3	0.6	1.0
2000.....	5.1	*	*	*	*	*	0.3	0.5	0.6	0.7
1999.....	5.0	*	*	*	*	0.2	0.3	0.4	0.6	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

Black, female

2004.....	6.5	*	*	*	*	*	2.3	3.7	3.2	5.1
2003.....	6.4	*	*	*	*	1.4	2.6	3.5	4.2	4.1
2002.....	6.3	*	*	*	*	*	3.0	3.4	3.4	5.6
2001.....	6.3	*	*	*	*	*	2.7	3.6	3.6	4.8
2000.....	6.2	*	*	*	*	*	2.5	3.2	3.4	4.9
1999.....	6.2	*	*	*	*	1.4	2.4	3.1	3.4	5.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Systemic lupus erythematosus (M32)

All races 3/, both sexes

2004.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.5
2003.....	0.5	*	*	*	*	0.2	0.3	0.3	0.4	0.4
2002.....	0.4	*	*	*	*	0.1	0.3	0.3	0.3	0.5
2001.....	0.5	*	*	*	*	0.2	0.4	0.4	0.4	0.5
2000.....	0.5	*	*	*	*	0.1	0.3	0.4	0.3	0.5
1999.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.5

All races 3/, male

2004.....	0.1	*	*	*	*	*	*	*	*	0.2
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.7	*	*	*	*	*	0.5	0.7	0.6	0.7
2003.....	0.8	*	*	*	*	0.3	0.5	0.6	0.7	0.7
2002.....	0.8	*	*	*	*	0.2	0.5	0.5	0.5	0.9
2001.....	0.8	*	*	*	*	0.3	0.6	0.7	0.6	0.8
2000.....	0.8	*	*	*	*	*	0.5	0.6	0.6	0.9
1999.....	0.8	*	*	*	*	0.2	0.5	0.6	0.6	0.9

White, both sexes

2004.....	0.3	*	*	*	*	*	0.1	0.2	0.2	0.3
2003.....	0.4	*	*	*	*	*	0.2	0.2	0.2	0.3
2002.....	0.3	*	*	*	*	*	*	*	0.1	0.2
2001.....	0.4	*	*	*	*	*	0.2	0.1	0.2	0.2
2000.....	0.4	*	*	*	*	*	0.2	0.2	0.2	0.3
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Systemic lupus erythematosus (M32)

Black, female

2004.....	2.0	*	*	*	*	*	1.7	2.4	2.4	2.4
2003.....	1.9	*	*	*	*	*	1.9	1.9	2.3	2.4
2002.....	2.1	*	*	*	*	*	2.1	2.3	2.0	3.8
2001.....	2.0	*	*	*	*	*	2.0	2.7	2.2	3.2
2000.....	2.0	*	*	*	*	*	1.8	2.5	2.0	2.8
1999.....	2.0	*	*	*	*	*	1.9	1.9	2.1	3.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)

All races 3/, both sexes

2004.....	1.6	*	*	*	*	*	0.1	0.2	0.2	0.4
2003.....	1.6	*	*	*	*	0.1	0.1	0.2	0.3	0.4
2002.....	1.7	*	*	*	*	*	0.1	0.2	0.3	0.5
2001.....	1.7	*	*	*	*	*	0.1	0.2	0.3	0.5
2000.....	1.7	*	*	*	*	*	0.2	0.2	0.4	0.4
1999.....	1.7	*	*	*	*	0.1	*	0.2	0.3	0.5

All races 3/, male

2004.....	1.0	*	*	*	*	*	*	*	*	0.2
2003.....	1.0	*	*	*	*	*	*	*	*	0.4
2002.....	1.0	*	*	*	*	*	*	*	0.2	0.4
2001.....	1.0	*	*	*	*	*	*	*	*	0.4
2000.....	1.1	*	*	*	*	*	*	0.2	0.3	0.2
1999.....	1.0	*	*	*	*	*	*	*	*	0.4

All races 3/, female

2004.....	2.2	*	*	*	*	*	*	0.3	0.2	0.6
2003.....	2.2	*	*	*	*	*	*	0.4	0.4	0.5
2002.....	2.3	*	*	*	*	*	0.2	0.3	0.4	0.5
2001.....	2.3	*	*	*	*	*	*	0.3	0.4	0.6
2000.....	2.3	*	*	*	*	*	*	0.3	0.5	0.6
1999.....	2.3	*	*	*	*	*	*	0.2	0.4	0.6

White, both sexes

2004.....	1.7	*	*	*	*	*	*	*	0.1	0.3
2003.....	1.7	*	*	*	*	*	*	*	0.1	0.3
2002.....	1.7	*	*	*	*	*	*	0.2	0.2	0.3
2001.....	1.8	*	*	*	*	*	*	0.1	0.2	0.4
2000.....	1.7	*	*	*	*	*	*	0.2	0.3	0.2
1999.....	1.7	*	*	*	*	*	*	*	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

XIV. Diseases of the genitourinary system (N00-N98)

All races 3/, both sexes

2004.....	20.5	4.9	0.2	*	*	0.2	0.4	0.6	1.0	1.7
2003.....	20.7	5.1	0.1	*	0.1	0.2	0.4	0.7	1.1	1.6
2002.....	20.2	4.6	*	*	*	0.2	0.4	0.7	1.0	1.6
2001.....	20.0	3.9	0.2	*	*	0.1	0.4	0.6	0.9	1.7
2000.....	19.4	5.0	0.1	*	*	0.2	0.4	0.5	1.0	1.5
1999.....	19.0	5.1	0.2	0.1	0.1	0.2	0.3	0.7	0.9	1.5

All races 3/, male

2004.....	18.5	5.8	*	*	*	*	0.4	0.6	0.8	1.8
2003.....	18.6	6.0	*	*	*	0.2	0.4	0.6	1.1	1.6
2002.....	18.1	5.6	*	*	*	*	0.4	0.6	1.0	1.6
2001.....	17.7	4.2	*	*	*	*	0.4	0.6	0.9	1.8
2000.....	17.2	6.1	*	*	*	0.2	0.3	0.5	1.0	1.5
1999.....	16.9	5.0	*	*	*	*	0.3	0.6	0.9	1.6

All races 3/, female

2004.....	22.5	4.0	*	*	*	*	0.3	0.7	1.1	1.7
2003.....	22.6	4.3	*	*	*	0.2	0.5	0.7	1.1	1.5
2002.....	22.3	3.6	*	*	*	*	0.4	0.9	1.0	1.6
2001.....	22.2	3.6	*	*	*	*	0.5	0.6	1.0	1.6
2000.....	21.5	3.9	*	*	*	*	0.5	0.5	1.0	1.5
1999.....	21.1	5.1	*	*	*	0.2	0.4	0.8	0.9	1.4

White, both sexes

2004.....	20.6	3.6	*	*	*	0.2	0.3	0.5	0.7	1.3
2003.....	20.7	4.1	*	*	*	0.2	0.3	0.5	0.8	1.1
2002.....	20.3	4.0	*	*	*	*	0.3	0.5	0.8	1.1
2001.....	20.0	3.0	*	*	*	*	0.4	0.4	0.7	1.2
2000.....	19.4	3.6	*	*	*	0.2	0.4	0.4	0.7	1.0
1999.....	18.9	4.0	*	*	*	0.1	0.2	0.5	0.7	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XIV. Diseases of the genitourinary system (N00-N98)

White, male

2004.....	18.5	4.2	*	*	*	*	0.3	0.4	0.4	1.2
2003.....	18.6	4.8	*	*	*	*	0.3	0.5	0.8	1.0
2002.....	18.1	4.7	*	*	*	*	0.3	0.4	0.8	1.1
2001.....	17.7	3.4	*	*	*	*	0.3	0.4	0.6	1.5
2000.....	17.2	4.6	*	*	*	*	0.3	0.4	0.7	1.0
1999.....	16.6	4.5	*	*	*	*	*	0.4	0.7	1.1

White, female

2004.....	22.7	2.8	*	*	*	*	0.3	0.5	0.9	1.3
2003.....	22.8	3.4	*	*	*	*	0.3	0.6	0.8	1.1
2002.....	22.5	3.2	*	*	*	*	0.3	0.6	0.7	1.1
2001.....	22.3	2.6	*	*	*	*	0.5	0.5	0.8	1.0
2000.....	21.5	2.6	*	*	*	*	0.4	0.5	0.7	1.0
1999.....	21.2	3.4	*	*	*	*	0.3	0.6	0.6	1.0

Black, both sexes

2004.....	25.7	11.6	*	*	*	*	0.9	1.6	2.8	4.9
2003.....	26.0	11.1	*	*	*	*	1.0	1.6	3.4	5.1
2002.....	25.2	8.3	*	*	*	*	1.1	2.3	2.6	4.8
2001.....	25.1	9.2	*	*	*	*	0.8	1.6	2.6	4.7
2000.....	24.4	12.7	*	*	*	*	*	1.2	2.9	4.9
1999.....	24.3	11.1	*	*	*	*	1.0	1.8	2.5	4.6

Black, male

2004.....	23.4	13.0	*	*	*	*	*	*	3.1	5.8
2003.....	24.0	12.5	*	*	*	*	*	*	3.9	6.0
2002.....	23.2	10.5	*	*	*	*	*	1.9	2.4	4.6
2001.....	22.2	9.3	*	*	*	*	*	2.1	3.1	3.8
2000.....	21.9	14.1	*	*	*	*	*	*	3.1	5.6
1999.....	22.4	8.1	*	*	*	*	*	*	2.4	5.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XIV. Diseases of the genitourinary system (N00-N98)

Black, female

2004.....	27.7	10.2	*	*	*	*	*	1.8	2.6	4.0
2003.....	27.7	9.6	*	*	*	*	*	1.8	2.9	4.3
2002.....	27.0	6.1	*	*	*	*	1.3	2.7	2.8	5.0
2001.....	27.8	9.1	*	*	*	*	*	*	2.1	5.5
2000.....	26.6	11.2	*	*	*	*	*	*	2.8	4.3
1999.....	26.1	14.1	*	*	*	*	*	2.2	2.5	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Diseases of urinary system (N00-N39)

All races 3/, both sexes

2004.....	20.2	4.8	0.2	*	*	0.2	0.4	0.6	0.9	1.7
2003.....	20.3	5.0	0.1	*	0.1	0.2	0.4	0.7	1.1	1.5
2002.....	19.9	4.6	*	*	*	0.1	0.4	0.7	0.9	1.5
2001.....	19.7	3.7	0.2	*	*	0.1	0.4	0.6	0.9	1.7
2000.....	19.1	4.8	0.1	*	*	0.2	0.4	0.5	0.9	1.4
1999.....	18.7	4.7	0.1	0.1	0.1	0.2	0.3	0.7	0.9	1.4

All races 3/, male

2004.....	18.0	5.8	*	*	*	*	0.4	0.6	0.8	1.8
2003.....	18.2	5.9	*	*	*	0.2	0.4	0.6	1.1	1.6
2002.....	17.7	5.6	*	*	*	*	0.4	0.6	1.0	1.5
2001.....	17.3	4.2	*	*	*	*	0.4	0.6	0.9	1.7
2000.....	16.8	6.0	*	*	*	0.2	0.3	0.5	1.0	1.5
1999.....	16.4	5.0	*	*	*	*	0.3	0.6	0.9	1.6

All races 3/, female

2004.....	22.2	3.8	*	*	*	*	0.3	0.7	1.1	1.6
2003.....	22.4	4.1	*	*	*	*	0.4	0.7	1.0	1.4
2002.....	22.0	3.5	*	*	*	*	0.4	0.9	0.9	1.4
2001.....	22.0	3.2	*	*	*	*	0.5	0.6	1.0	1.6
2000.....	21.2	3.6	*	*	*	*	0.4	0.5	0.9	1.3
1999.....	20.9	4.3	*	*	*	*	0.3	0.7	0.8	1.3

White, both sexes

2004.....	20.2	3.5	*	*	*	0.1	0.3	0.4	0.6	1.2
2003.....	20.4	4.1	*	*	*	0.2	0.3	0.5	0.8	1.0
2002.....	20.0	3.9	*	*	*	*	0.3	0.5	0.7	1.0
2001.....	19.7	2.8	*	*	*	*	0.4	0.4	0.7	1.2
2000.....	19.0	3.5	*	*	*	0.2	0.4	0.4	0.7	1.0
1999.....	18.6	3.7	*	*	*	0.1	0.2	0.5	0.6	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of urinary system (N00-N39)										
White, male										
2004.....	18.0	4.2	*	*	*	*	0.3	0.4	0.4	1.2
2003.....	18.1	4.8	*	*	*	*	0.3	0.5	0.8	1.0
2002.....	17.7	4.7	*	*	*	*	0.3	0.4	0.8	1.1
2001.....	17.3	3.3	*	*	*	*	0.3	0.4	0.6	1.5
2000.....	16.7	4.5	*	*	*	*	0.3	0.4	0.7	1.0
1999.....	16.2	4.5	*	*	*	*	*	0.4	0.7	1.1
White, female										
2004.....	22.4	2.8	*	*	*	*	0.3	0.5	0.8	1.3
2003.....	22.5	3.3	*	*	*	*	0.3	0.5	0.8	1.0
2002.....	22.2	3.0	*	*	*	*	*	0.6	0.6	1.0
2001.....	22.0	2.3	*	*	*	*	0.4	0.5	0.8	1.0
2000.....	21.3	2.4	*	*	*	*	0.4	0.4	0.7	0.9
1999.....	20.9	2.8	*	*	*	*	*	0.5	0.6	0.9
Black, both sexes										
2004.....	25.3	11.2	*	*	*	*	0.8	1.6	2.8	4.8
2003.....	25.7	10.6	*	*	*	*	0.9	1.5	3.3	5.0
2002.....	24.8	8.3	*	*	*	*	1.0	2.3	2.5	4.5
2001.....	24.8	8.9	*	*	*	*	0.8	1.6	2.6	4.4
2000.....	24.1	12.2	*	*	*	*	*	1.2	2.8	4.7
1999.....	23.9	9.9	*	*	*	*	0.9	1.7	2.4	4.5
Black, male										
2004.....	23.0	13.0	*	*	*	*	*	*	3.0	5.8
2003.....	23.7	12.2	*	*	*	*	*	*	3.9	5.9
2002.....	22.8	10.5	*	*	*	*	*	1.9	2.4	4.5
2001.....	21.8	9.3	*	*	*	*	*	2.1	3.1	3.6
2000.....	21.6	14.1	*	*	*	*	*	*	3.0	5.5
1999.....	22.0	8.1	*	*	*	*	*	*	2.4	5.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ 2	Under 1 year 2/ 3	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------	-------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Diseases of urinary system (N00-N39)

Black, female

2004.....	27.4	9.3	*	*	*	*	*	1.8	2.5	3.8
2003.....	27.4	9.0	*	*	*	*	*	1.6	2.7	4.1
2002.....	26.6	6.1	*	*	*	*	*	2.6	2.6	4.4
2001.....	27.5	8.5	*	*	*	*	*	*	2.1	5.2
2000.....	26.3	10.3	*	*	*	*	*	*	2.6	4.0
1999.....	25.7	11.7	*	*	*	*	*	2.0	2.4	3.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Renal failure (N17-N19)

All races 3/, both sexes

2004.....	14.2	4.0	*	*	*	0.1	0.3	0.4	0.7	1.3
2003.....	14.4	4.2	*	*	*	0.1	0.3	0.5	0.8	1.2
2002.....	13.9	4.1	*	*	*	*	0.3	0.5	0.7	1.2
2001.....	13.6	3.1	*	*	*	*	0.3	0.4	0.6	1.2
2000.....	13.0	4.0	*	*	*	0.1	0.3	0.4	0.7	1.2
1999.....	12.4	4.3	*	*	*	0.1	0.2	0.4	0.7	1.1

All races 3/, male

2004.....	13.8	5.1	*	*	*	*	0.3	0.4	0.6	1.4
2003.....	14.1	4.8	*	*	*	*	0.3	0.5	0.9	1.3
2002.....	13.7	5.1	*	*	*	*	0.3	0.4	0.7	1.3
2001.....	13.2	3.5	*	*	*	*	0.3	0.4	0.6	1.3
2000.....	12.6	5.1	*	*	*	*	0.2	0.4	0.7	1.3
1999.....	12.1	4.5	*	*	*	*	*	0.3	0.7	1.3

All races 3/, female

2004.....	14.6	3.0	*	*	*	*	0.2	0.4	0.8	1.2
2003.....	14.6	3.7	*	*	*	*	0.2	0.5	0.8	1.1
2002.....	14.2	3.0	*	*	*	*	0.3	0.6	0.7	1.1
2001.....	14.0	2.7	*	*	*	*	0.4	0.4	0.7	1.2
2000.....	13.3	2.8	*	*	*	*	0.3	0.3	0.7	1.0
1999.....	12.7	4.1	*	*	*	*	0.3	0.5	0.7	0.9

White, both sexes

2004.....	13.9	2.9	*	*	*	*	0.2	0.3	0.4	0.9
2003.....	14.0	3.5	*	*	*	*	0.2	0.4	0.5	0.8
2002.....	13.6	3.4	*	*	*	*	0.2	0.3	0.5	0.8
2001.....	13.2	2.2	*	*	*	*	0.3	0.3	0.4	0.8
2000.....	12.6	2.7	*	*	*	*	0.2	0.3	0.5	0.7
1999.....	12.0	3.3	*	*	*	*	0.2	0.3	0.5	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Renal failure (N17-N19)

White, male

2004.....	13.6	3.6	*	*	*	*	*	0.3	0.3	0.9
2003.....	13.8	4.1	*	*	*	*	*	0.4	0.5	0.8
2002.....	13.4	4.2	*	*	*	*	*	*	0.5	0.9
2001.....	13.0	2.6	*	*	*	*	*	0.3	0.4	1.0
2000.....	12.4	3.6	*	*	*	*	0.3	0.3	0.5	0.8
1999.....	11.8	3.9	*	*	*	*	*	*	0.5	0.8

White, female

2004.....	14.1	2.1	*	*	*	*	*	0.3	0.5	1.0
2003.....	14.2	2.8	*	*	*	*	*	0.4	0.5	0.7
2002.....	13.8	2.6	*	*	*	*	*	0.4	0.5	0.7
2001.....	13.4	1.8	*	*	*	*	0.3	0.3	0.5	0.7
2000.....	12.8	1.7	*	*	*	*	*	*	0.5	0.7
1999.....	12.2	2.8	*	*	*	*	*	0.3	0.5	0.6

Black, both sexes

2004.....	20.0	9.8	*	*	*	*	0.7	1.2	2.3	4.0
2003.....	20.3	8.9	*	*	*	*	0.6	1.3	2.8	4.4
2002.....	19.5	7.7	*	*	*	*	0.8	1.7	2.0	3.9
2001.....	19.3	8.1	*	*	*	*	0.7	1.2	2.1	3.9
2000.....	18.5	11.1	*	*	*	*	*	1.0	2.5	4.1
1999.....	18.2	9.6	*	*	*	*	*	1.4	2.2	3.8

Black, male

2004.....	18.9	11.9	*	*	*	*	*	*	2.7	5.1
2003.....	19.7	9.2	*	*	*	*	*	*	3.4	5.4
2002.....	18.8	9.6	*	*	*	*	*	*	2.0	4.4
2001.....	17.7	8.7	*	*	*	*	*	1.6	2.6	3.3
2000.....	17.3	13.4	*	*	*	*	*	*	2.7	4.9
1999.....	17.1	8.1	*	*	*	*	*	*	2.2	4.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Renal failure (N17-N19)

Black, female

2004.....	21.0	7.7	*	*	*	*	*	1.9	3.0
2003.....	20.9	8.7	*	*	*	*	*	2.3	3.5
2002.....	20.2	*	*	*	*	*	1.9	2.0	3.5
2001.....	20.7	7.6	*	*	*	*	*	1.7	4.4
2000.....	19.6	8.6	*	*	*	*	*	2.3	3.4
1999.....	19.1	11.1	*	*	*	*	1.6	2.2	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Noninflammatory disorders of female genital tract (N80-N98)

White, male

Table with 11 columns (Year, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) showing death rates for White males.

White, female

Table with 11 columns (Year, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) showing death rates for White females.

Black, both sexes

Table with 11 columns (Year, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) showing death rates for Black both sexes.

Black, male

Table with 11 columns (Year, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) showing death rates for Black males.

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

XV. Pregnancy, childbirth and the puerperium (000-099)

All races 3/, both sexes

2004.....	0.2	*	0.2	0.6	0.7	0.7	0.5
2003.....	0.2	*	0.1	0.4	0.6	0.6	0.5
2002.....	0.1	*	0.1	0.3	0.4	0.4	0.4
2001.....	0.1	*	0.2	0.4	0.5	0.5	0.3
2000.....	0.1	*	*	0.4	0.4	0.5	0.4
1999.....	0.1	*	0.2	0.3	0.5	0.4	0.4

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	0.5	*	0.4	1.3	1.5	1.4	0.9
2003.....	0.4	*	0.3	0.9	1.2	1.2	1.0
2002.....	0.3	*	0.3	0.6	0.9	0.9	0.7
2001.....	0.3	*	0.4	0.8	1.0	0.9	0.7
2000.....	0.3	*	*	0.8	0.9	0.9	0.8
1999.....	0.3	*	0.3	0.7	0.9	0.9	0.8

White, both sexes

2004.....	0.2	*	0.1	0.4	0.5	0.6	0.4
2003.....	0.1	*	*	0.3	0.5	0.4	0.3
2002.....	0.1	*	*	0.2	0.3	0.3	0.2
2001.....	0.1	*	0.1	0.3	0.3	0.4	0.2
2000.....	0.1	*	*	0.3	0.3	0.3	0.3
1999.....	0.1	*	*	0.2	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ year 1/	year 2/ year 1/	years	years	years	years	years	years	years	years

XV. Pregnancy, childbirth and the puerperium (000-099)

Black, female

2004.....	1.2	*	*	4.0	4.3	3.2	2.0
2003.....	1.0	*	*	2.1	2.3	3.5	2.7
2002.....	0.8	*	*	2.0	2.8	1.9	2.3
2001.....	0.8	*	*	2.1	2.6	2.1	1.9
2000.....	0.7	*	*	2.2	2.2	2.3	1.7
1999.....	0.9	*	*	2.1	2.7	2.3	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

All races 3/, both sexes

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

White, both sexes

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

Black, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Ectopic pregnancy (000)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, both sexes

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Ectopic pregnancy (000)

Black, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Spontaneous abortion (003)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, both sexes

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Spontaneous abortion (003)

Black, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Medical abortion (004)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, both sexes

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Medical abortion (004)

Black, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other abortion (005)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, both sexes

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other abortion (005)

Black, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, both sexes

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ years	year 2/ years	years	years	years	years	years	years	years	years

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

Black, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other direct obstetric deaths (O10-O92)

All races 3/, both sexes

2004.....	0.1	*	0.1	0.3	0.4	0.4	0.2
2003.....	0.1	*	*	0.3	0.3	0.3	0.3
2002.....	0.1	*	0.1	0.2	0.3	0.3	0.3
2001.....	0.1	*	0.1	0.3	0.4	0.4	0.2
2000.....	0.1	*	*	0.3	0.3	0.4	0.3
1999.....	0.1	*	0.1	0.2	0.3	0.3	0.3

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	0.2	*	0.2	0.6	0.8	0.8	0.5
2003.....	0.2	*	*	0.5	0.7	0.7	0.7
2002.....	0.2	*	0.2	0.4	0.7	0.7	0.6
2001.....	0.2	*	0.3	0.5	0.8	0.7	0.5
2000.....	0.2	*	*	0.6	0.6	0.8	0.6
1999.....	0.2	*	0.3	0.5	0.7	0.7	0.6

White, both sexes

2004.....	0.1	*	*	0.2	0.2	0.3	0.2
2003.....	0.1	*	*	0.2	0.3	0.2	0.2
2002.....	0.1	*	*	*	0.2	0.3	0.2
2001.....	0.1	*	*	0.2	0.3	0.3	0.1
2000.....	0.1	*	*	0.2	0.2	0.3	0.2
1999.....	0.1	*	*	0.2	0.2	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other direct obstetric deaths (O10-O92)

Black, female

2004.....	0.7	*	*	2.0	2.9	2.0	*
2003.....	0.7	*	*	1.3	1.7	2.5	2.1
2002.....	0.6	*	*	1.5	1.9	1.6	1.7
2001.....	0.6	*	*	1.5	2.0	1.6	*
2000.....	0.5	*	*	1.6	1.5	1.7	*
1999.....	0.6	*	*	*	2.0	1.7	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Eclampsia and pre-eclampsia (011,013-016)

All races 3/, both sexes

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	0.1	*
1999.....	0.0	*	*	*	0.1	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	*
1999.....	0.1	*	*	*	0.2	*	*

White, both sexes

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Eclampsia and pre-eclampsia (O11,O13-O16)

Black, female

2004.....	0.1	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Hemorrhage of pregnancy and childbirth and placenta previa (020,044-046,067,072)

All races 3/, both sexes

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

White, both sexes

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ years	year 2/ years	years	years	years	years	years	years	years	years

Hemorrhage of pregnancy and childbirth and placenta previa (020,044-046,067,072)

Black, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Complications predominately related to the puerperium (085-092)

All races 3/, both sexes

2004.....	0.0	*	*	*	0.1	0.1	0.1
2003.....	0.0	*	*	*	0.1	*	0.1
2002.....	0.0	*	*	*	*	0.1	*
2001.....	0.0	*	*	*	0.1	0.1	*
2000.....	0.0	*	*	0.1	*	0.1	*
1999.....	0.0	*	*	*	*	0.1	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	0.1	*	*	*	0.2	0.2	0.2
2003.....	0.1	*	*	*	0.2	*	0.2
2002.....	0.1	*	*	*	*	0.2	*
2001.....	0.1	*	*	*	0.3	0.2	*
2000.....	0.1	*	*	0.2	*	0.3	*
1999.....	0.1	*	*	*	*	0.2	*

White, both sexes

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ years	year 2/ years	years	years	years	years	years	years	years	years

Complications predominately related to the puerperium (085-092)

Black, female

2004.....	0.2	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Obstetric embolism (088)

All races 3/, both sexes

2004.....	0.0	*	*	*	*	*
2003.....	0.0	*	*	*	*	*
2002.....	0.0	*	*	*	*	*
2001.....	0.0	*	*	*	*	*
2000.....	0.0	*	*	*	0.1	*
1999.....	0.0	*	*	*	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	0.0	*	*	*	*	*
2003.....	0.0	*	*	*	*	*
2002.....	0.0	*	*	*	*	*
2001.....	0.0	*	*	*	*	*
2000.....	0.0	*	*	*	0.2	*
1999.....	0.0	*	*	*	*	*

White, both sexes

2004.....	0.0	*	*	*	*	*
2003.....	0.0	*	*	*	*	*
2002.....	0.0	*	*	*	*	*
2001.....	0.0	*	*	*	*	*
2000.....	0.0	*	*	*	*	*
1999.....	0.0	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Obstetric embolism (088)

Black, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other complications predominately related to the puerperium (085-087,089-092)

All races 3/, both sexes

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

White, both sexes

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications predominately related to the puerperium (085-087,089-092)

Black, female

2004.....	0.1	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other direct obstetric causes (010,012,021-043,047-066,068-071,073-075)

All races 3/, both sexes

2004.....	0.1	*	*	0.1	0.2	0.2	0.1
2003.....	0.0	*	*	0.1	*	0.2	0.1
2002.....	0.0	*	*	0.1	0.1	0.1	0.1
2001.....	0.0	*	*	*	0.1	0.1	*
2000.....	0.0	*	*	0.1	*	0.1	0.1
1999.....	0.0	*	*	*	0.1	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	0.1	*	*	0.3	0.4	0.3	0.2
2003.....	0.1	*	*	0.2	*	0.3	0.2
2002.....	0.1	*	*	0.2	0.3	0.3	0.3
2001.....	0.1	*	*	*	0.3	0.2	*
2000.....	0.1	*	*	0.2	*	0.2	0.2
1999.....	0.1	*	*	*	0.2	*	*

White, both sexes

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ years	year 2/ years	years	years	years	years	years	years	years	years

All other direct obstetric causes (010,012,021-043,047-066,068-071,073-075)

Black, female

2004.....	0.3	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Obstetric death of unspecified cause (095)

All races 3/, both sexes

Table with 11 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for 'All races 3/, both sexes'. Values are mostly 0.0 or *.

All races 3/, male

Table with 11 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for 'All races 3/, male'. Values are mostly

All races 3/, female

Table with 11 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for 'All races 3/, female'. Values are mostly 0.0 or *.

White, both sexes

Table with 11 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for 'White, both sexes'. Values are mostly *.

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Obstetric death of unspecified cause (095)

Black, female

2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

All races 3/, both sexes

2004.....	0.1	*	*	*	0.1	0.2	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	0.1	*	*	*	0.2	0.3	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, both sexes

2004.....	0.0	*	*	*	*	0.1	*
2003.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Black, female

2004.....	0.2	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Indirect obstetric deaths (O98-O99)

All races 3/, both sexes

2004.....	0.0	*	*	0.1	0.1	0.1	0.1
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	0.1	*	*	0.3	0.3	0.2	0.2
2003.....	0.1	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

White, both sexes

2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Indirect obstetric deaths (O98-O99)

White, male

Table with 11 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for White, male. All cells contain '...'.

White, female

Table with 11 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for White, female. Values range from 0.0 to 0.1, with asterisks in the 10-14, 15-19, 20-24, 25-29, 30-34, and 35-39 age groups.

Black, both sexes

Table with 11 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Black, both sexes. Values range from 0.1 to 0.1, with asterisks in the 10-14, 15-19, 20-24, 25-29, 30-34, and 35-39 age groups.

Black, male

Table with 11 columns (Year, All ages 1/, Under 1 year 2/, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years) and 6 rows (2004, 2003, 2002, 2001, 2000, 1999) for Black, male. All cells contain '...'.

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Indirect obstetric deaths (O98-O99)

Black, female

2004.....	0.2	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

XVI. Certain conditions originating in the perinatal period (P00-P96)

All races 3/, both sexes

2004.....	4.8	345.3	0.4	0.1	*	*	*	*	*	*
2003.....	4.9	356.0	0.5	*	*	*	*	*	*	*
2002.....	4.9	349.7	0.4	*	*	*	*	*	*	*
2001.....	4.9	340.5	0.5	0.1	*	*	*	*	*	*
2000.....	5.0	365.7	0.5	0.1	*	*	*	*	*	*
1999.....	5.1	371.0	0.6	0.1	*	*	*	*	*	*

All races 3/, male

2004.....	5.6	384.2	0.5	*	*	*	*	*	*	*
2003.....	5.7	397.4	0.5	*	*	*	*	*	*	*
2002.....	5.7	385.7	0.5	*	*	*	*	*	*	*
2001.....	5.6	376.6	0.6	*	*	*	*	*	*	*
2000.....	5.8	403.9	0.7	*	*	*	*	*	*	*
1999.....	5.9	408.6	0.7	*	*	*	*	*	*	*

All races 3/, female

2004.....	4.1	304.6	0.3	*	*	*	*	*	*	*
2003.....	4.2	312.8	0.5	*	*	*	*	*	*	*
2002.....	4.2	312.0	0.3	*	*	*	*	*	*	*
2001.....	4.2	302.6	0.4	*	*	*	*	*	*	*
2000.....	4.3	325.6	0.4	*	*	*	*	*	*	*
1999.....	4.4	331.7	0.5	*	*	*	*	*	*	*

White, both sexes

2004.....	3.7	272.1	0.3	*	*	*	*	*	*	*
2003.....	3.8	285.1	0.4	*	*	*	*	*	*	*
2002.....	3.7	277.2	0.3	*	*	*	*	*	*	*
2001.....	3.7	268.5	0.4	*	*	*	*	*	*	*
2000.....	3.7	283.7	0.4	*	*	*	*	*	*	*
1999.....	3.8	285.9	0.5	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Newborn affected by maternal factors and by complications of pregnancy, labor and delivery (P00-P04)

All races 3/, both sexes

2004.....	1.1	76.2	*	*	*	*	*	*	*
2003.....	1.1	79.1	*	*	*	*	*	*	*
2002.....	1.1	75.9	*	*	*	*	*	*	*
2001.....	1.0	70.2	*	*	*	*	*	*	*
2000.....	1.0	72.7	*	*	*	*	*	*	*
1999.....	1.0	72.1	*	*	*	*	*	*	*

All races 3/, male

2004.....	1.2	81.9	*	*	*	*	*	*	*
2003.....	1.2	85.6	*	*	*	*	*	*	*
2002.....	1.2	83.2	*	*	*	*	*	*	*
2001.....	1.2	77.8	*	*	*	*	*	*	*
2000.....	1.1	78.7	*	*	*	*	*	*	*
1999.....	1.1	76.7	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.9	70.3	*	*	*	*	*	*	*
2003.....	1.0	72.2	*	*	*	*	*	*	*
2002.....	0.9	68.3	*	*	*	*	*	*	*
2001.....	0.8	62.2	*	*	*	*	*	*	*
2000.....	0.9	66.5	*	*	*	*	*	*	*
1999.....	0.9	67.4	*	*	*	*	*	*	*

White, both sexes

2004.....	0.8	60.0	*	*	*	*	*	*	*
2003.....	0.9	65.5	*	*	*	*	*	*	*
2002.....	0.8	61.1	*	*	*	*	*	*	*
2001.....	0.8	57.4	*	*	*	*	*	*	*
2000.....	0.8	59.4	*	*	*	*	*	*	*
1999.....	0.7	56.5	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Disorders related to short gestation and low birth weight, not elsewhere classified (P07)

White, male

2004.....	1.3	93.6	*	*	*	*	*	*	*
2003.....	1.3	97.9	*	*	*	*	*	*	*
2002.....	1.2	89.3	*	*	*	*	*	*	*
2001.....	1.2	83.5	*	*	*	*	*	*	*
2000.....	1.1	84.0	*	*	*	*	*	*	*
1999.....	1.2	86.1	*	*	*	*	*	*	*

White, female

2004.....	0.9	73.2	*	*	*	*	*	*	*
2003.....	1.0	79.2	*	*	*	*	*	*	*
2002.....	1.0	74.6	*	*	*	*	*	*	*
2001.....	0.9	72.1	*	*	*	*	*	*	*
2000.....	0.9	76.1	*	*	*	*	*	*	*
1999.....	0.9	72.2	*	*	*	*	*	*	*

Black, both sexes

2004.....	4.7	273.6	*	*	*	*	*	*	*
2003.....	5.0	289.6	*	*	*	*	*	*	*
2002.....	5.0	279.4	*	*	*	*	*	*	*
2001.....	4.9	281.7	*	*	*	*	*	*	*
2000.....	5.1	304.0	*	*	*	*	*	*	*
1999.....	5.3	313.5	*	*	*	*	*	*	*

Black, male

2004.....	5.6	304.2	*	*	*	*	*	*	*
2003.....	6.0	323.6	*	*	*	*	*	*	*
2002.....	5.9	306.5	*	*	*	*	*	*	*
2001.....	5.8	304.2	*	*	*	*	*	*	*
2000.....	5.9	329.4	*	*	*	*	*	*	*
1999.....	6.2	347.4	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intrauterine hypoxia and birth asphyxia (P20-P21)

All races 3/, both sexes

2004.....	0.2	13.0	*	*	*	*	*	*	*
2003.....	0.2	13.9	*	*	*	*	*	*	*
2002.....	0.2	14.5	*	*	*	*	*	*	*
2001.....	0.2	13.2	*	*	*	*	*	*	*
2000.....	0.2	16.6	*	*	*	*	*	*	*
1999.....	0.2	16.1	0.1	*	*	*	*	*	*

All races 3/, male

2004.....	0.2	14.6	*	*	*	*	*	*	*
2003.....	0.2	14.7	*	*	*	*	*	*	*
2002.....	0.2	14.5	*	*	*	*	*	*	*
2001.....	0.2	14.3	*	*	*	*	*	*	*
2000.....	0.3	18.1	*	*	*	*	*	*	*
1999.....	0.3	17.2	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.2	11.3	*	*	*	*	*	*	*
2003.....	0.2	13.2	*	*	*	*	*	*	*
2002.....	0.2	14.4	*	*	*	*	*	*	*
2001.....	0.2	12.1	*	*	*	*	*	*	*
2000.....	0.2	15.0	*	*	*	*	*	*	*
1999.....	0.2	15.0	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	11.8	*	*	*	*	*	*	*
2003.....	0.2	12.5	*	*	*	*	*	*	*
2002.....	0.2	13.2	*	*	*	*	*	*	*
2001.....	0.2	12.2	*	*	*	*	*	*	*
2000.....	0.2	14.6	*	*	*	*	*	*	*
1999.....	0.2	14.4	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intrauterine hypoxia and birth asphyxia (P20-P21)

White, male

2004.....	0.2	13.0	*	*	*	*	*	*	*
2003.....	0.2	13.7	*	*	*	*	*	*	*
2002.....	0.2	13.4	*	*	*	*	*	*	*
2001.....	0.2	13.0	*	*	*	*	*	*	*
2000.....	0.2	15.5	*	*	*	*	*	*	*
1999.....	0.2	15.0	*	*	*	*	*	*	*

White, female

2004.....	0.1	10.6	*	*	*	*	*	*	*
2003.....	0.1	11.3	*	*	*	*	*	*	*
2002.....	0.2	13.0	*	*	*	*	*	*	*
2001.....	0.2	11.5	*	*	*	*	*	*	*
2000.....	0.2	13.7	*	*	*	*	*	*	*
1999.....	0.2	13.8	*	*	*	*	*	*	*

Black, both sexes

2004.....	0.4	20.0	*	*	*	*	*	*	*
2003.....	0.4	22.0	*	*	*	*	*	*	*
2002.....	0.4	22.1	*	*	*	*	*	*	*
2001.....	0.4	19.8	*	*	*	*	*	*	*
2000.....	0.5	26.2	*	*	*	*	*	*	*
1999.....	0.4	25.9	*	*	*	*	*	*	*

Black, male

2004.....	0.4	22.5	*	*	*	*	*	*	*
2003.....	0.4	22.6	*	*	*	*	*	*	*
2002.....	0.4	21.5	*	*	*	*	*	*	*
2001.....	0.4	22.2	*	*	*	*	*	*	*
2000.....	0.6	29.4	*	*	*	*	*	*	*
1999.....	0.5	29.3	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intrauterine hypoxia and birth asphyxia (P20-P21)

Black, female

2004.....	0.3	17.3	*	*	*	*	*	*	*
2003.....	0.4	21.3	*	*	*	*	*	*	*
2002.....	0.4	22.7	*	*	*	*	*	*	*
2001.....	0.3	17.3	*	*	*	*	*	*	*
2000.....	0.4	22.8	*	*	*	*	*	*	*
1999.....	0.4	22.5	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Respiratory distress of newborn (P22)

All races 3/, both sexes

2004.....	0.3	21.5	*	*	*	*	*	*	*
2003.....	0.3	20.8	*	*	*	*	*	*	*
2002.....	0.3	23.4	*	*	*	*	*	*	*
2001.....	0.4	25.1	*	*	*	*	*	*	*
2000.....	0.4	26.3	*	*	*	*	*	*	*
1999.....	0.4	29.2	*	*	*	*	*	*	*

All races 3/, male

2004.....	0.3	24.0	*	*	*	*	*	*	*
2003.....	0.3	24.1	*	*	*	*	*	*	*
2002.....	0.4	26.2	*	*	*	*	*	*	*
2001.....	0.4	27.8	*	*	*	*	*	*	*
2000.....	0.5	33.0	*	*	*	*	*	*	*
1999.....	0.5	33.7	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.3	18.8	*	*	*	*	*	*	*
2003.....	0.2	17.3	*	*	*	*	*	*	*
2002.....	0.3	20.4	*	*	*	*	*	*	*
2001.....	0.3	22.2	*	*	*	*	*	*	*
2000.....	0.2	19.2	*	*	*	*	*	*	*
1999.....	0.3	24.6	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	17.6	*	*	*	*	*	*	*
2003.....	0.2	17.4	*	*	*	*	*	*	*
2002.....	0.2	18.7	*	*	*	*	*	*	*
2001.....	0.3	20.0	*	*	*	*	*	*	*
2000.....	0.3	21.0	*	*	*	*	*	*	*
1999.....	0.3	23.6	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Respiratory distress of newborn (P22)

Black, female

2004.....	0.6	38.0	*	*	*	*	*	*	*
2003.....	0.5	33.7	*	*	*	*	*	*	*
2002.....	0.9	50.9	*	*	*	*	*	*	*
2001.....	0.7	44.7	*	*	*	*	*	*	*
2000.....	0.7	46.3	*	*	*	*	*	*	*
1999.....	0.7	47.7	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other respiratory conditions originating in the perinatal period (P23-P28)

White, male

2004.....	0.4	28.5	*	*	*	*	*	*	*
2003.....	0.4	30.8	*	*	*	*	*	*	*
2002.....	0.4	29.8	*	*	*	*	*	*	*
2001.....	0.5	34.1	*	*	*	*	*	*	*
2000.....	0.5	35.8	*	*	*	*	*	*	*
1999.....	0.6	45.1	*	*	*	*	*	*	*

White, female

2004.....	0.3	19.9	*	*	*	*	*	*	*
2003.....	0.3	20.4	*	*	*	*	*	*	*
2002.....	0.3	20.9	*	*	*	*	*	*	*
2001.....	0.3	25.5	*	*	*	*	*	*	*
2000.....	0.4	28.7	*	*	*	*	*	*	*
1999.....	0.4	31.8	*	*	*	*	*	*	*

Black, both sexes

2004.....	1.1	61.3	*	*	*	*	*	*	*
2003.....	1.0	56.4	*	*	*	*	*	*	*
2002.....	1.1	60.5	*	*	*	*	*	*	*
2001.....	1.4	78.4	*	*	*	*	*	*	*
2000.....	1.4	83.1	*	*	*	*	*	*	*
1999.....	1.5	87.2	*	*	*	*	*	*	*

Black, male

2004.....	1.2	64.1	*	*	*	*	*	*	*
2003.....	1.3	66.3	*	*	*	*	*	*	*
2002.....	1.3	67.4	*	*	*	*	*	*	*
2001.....	1.7	88.6	*	*	*	*	*	*	*
2000.....	1.8	96.6	*	*	*	*	*	*	*
1999.....	1.8	94.3	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)

All races 3/, both sexes

2004.....	0.9	64.9	*	*	*	*	*	*	*
2003.....	0.9	65.9	*	*	*	*	*	*	*
2002.....	0.8	58.1	*	*	*	*	*	*	*
2001.....	0.9	60.2	*	*	*	*	*	*	*
2000.....	0.9	69.3	*	*	*	*	*	*	*
1999.....	0.9	69.1	*	*	*	*	*	*	*

All races 3/, male

2004.....	1.1	73.7	*	*	*	*	*	*	*
2003.....	1.1	74.4	*	*	*	*	*	*	*
2002.....	1.0	65.1	*	*	*	*	*	*	*
2001.....	1.0	68.5	*	*	*	*	*	*	*
2000.....	1.1	78.0	*	*	*	*	*	*	*
1999.....	1.1	77.0	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.7	55.7	*	*	*	*	*	*	*
2003.....	0.8	57.1	*	*	*	*	*	*	*
2002.....	0.7	50.7	*	*	*	*	*	*	*
2001.....	0.7	51.6	*	*	*	*	*	*	*
2000.....	0.8	60.1	*	*	*	*	*	*	*
1999.....	0.8	60.7	*	*	*	*	*	*	*

White, both sexes

2004.....	0.7	53.8	*	*	*	*	*	*	*
2003.....	0.7	54.8	*	*	*	*	*	*	*
2002.....	0.7	49.0	*	*	*	*	*	*	*
2001.....	0.7	50.4	*	*	*	*	*	*	*
2000.....	0.7	56.1	*	*	*	*	*	*	*
1999.....	0.7	55.0	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes

2004.....	3.6	137.9	3.6	1.0	0.9	1.2	1.1	1.1	1.0	1.0
2003.....	3.6	140.4	3.4	0.9	1.0	1.1	1.1	1.0	1.1	1.1
2002.....	3.7	139.4	3.4	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001.....	3.7	136.7	3.6	0.9	0.9	1.3	1.3	1.2	1.1	1.2
2000.....	3.8	150.9	3.2	1.0	1.0	1.1	1.1	1.1	1.3	1.2
1999.....	3.7	144.2	3.6	1.0	1.1	1.1	1.1	1.1	1.2	1.2

All races 3/, male

2004.....	3.8	139.0	3.9	1.1	0.9	1.4	1.3	1.2	1.1	1.2
2003.....	3.8	147.6	3.4	0.9	1.0	1.4	1.2	1.2	1.1	1.2
2002.....	3.9	142.6	3.5	1.0	1.1	1.4	1.3	1.4	1.4	1.2
2001.....	3.9	143.1	3.6	1.0	1.0	1.5	1.4	1.3	1.2	1.2
2000.....	4.0	155.4	3.3	1.0	1.0	1.3	1.4	1.2	1.5	1.4
1999.....	3.9	147.9	4.0	1.0	1.1	1.3	1.3	1.1	1.3	1.3

All races 3/, female

2004.....	3.4	136.8	3.2	1.0	0.8	1.1	0.9	0.9	1.0	0.8
2003.....	3.4	132.8	3.5	0.9	1.0	0.8	1.0	0.8	1.1	1.1
2002.....	3.5	136.1	3.3	1.0	1.0	1.0	1.1	0.9	1.1	1.2
2001.....	3.4	129.9	3.7	0.8	0.8	1.0	1.1	1.1	1.0	1.1
2000.....	3.5	146.2	3.1	0.9	1.0	0.9	0.9	0.9	1.1	1.1
1999.....	3.5	140.3	3.2	1.0	1.1	0.9	1.0	1.1	1.2	1.1

White, both sexes

2004.....	3.5	134.5	3.2	1.0	0.9	1.1	1.0	1.1	1.1	1.0
2003.....	3.5	137.6	3.2	0.9	1.0	1.1	1.0	1.1	1.1	1.2
2002.....	3.6	138.3	3.3	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001.....	3.6	135.0	3.2	0.9	0.9	1.2	1.2	1.2	1.2	1.1
2000.....	3.6	147.1	2.9	0.9	0.9	1.0	1.2	1.1	1.4	1.2
1999.....	3.6	141.3	3.2	0.9	1.0	1.1	1.1	1.1	1.3	1.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, male

2004.....	3.6	134.3	3.5	1.1	0.9	1.2	1.1	1.3	1.1	1.1
2003.....	3.7	143.2	3.1	0.8	0.9	1.3	1.1	1.2	1.1	1.2
2002.....	3.8	143.3	3.4	0.9	1.0	1.3	1.3	1.4	1.5	1.2
2001.....	3.8	143.0	3.2	1.0	1.1	1.4	1.3	1.4	1.3	1.2
2000.....	3.9	151.9	2.9	0.9	0.9	1.1	1.4	1.2	1.6	1.4
1999.....	3.8	146.5	3.5	0.8	1.0	1.3	1.3	1.2	1.3	1.4

White, female

2004.....	3.4	134.7	2.9	0.8	0.8	1.0	0.9	1.0	1.1	0.8
2003.....	3.4	131.8	3.2	0.9	1.0	0.8	1.0	0.9	1.1	1.1
2002.....	3.5	133.1	3.1	1.1	1.0	1.0	1.1	0.9	1.1	1.2
2001.....	3.4	126.7	3.3	0.8	0.7	0.9	1.1	1.0	1.1	1.0
2000.....	3.5	142.0	2.9	1.0	0.9	0.9	0.9	0.9	1.1	1.0
1999.....	3.5	135.7	2.9	1.0	1.0	0.8	0.9	1.1	1.2	1.1

Black, both sexes

2004.....	4.5	162.9	5.1	1.5	1.0	1.8	1.7	1.1	1.1	1.2
2003.....	4.5	157.5	4.7	1.1	1.1	1.2	1.7	1.2	1.5	1.2
2002.....	4.7	155.2	4.5	1.2	1.2	1.7	1.5	1.3	1.4	1.4
2001.....	4.6	158.4	5.3	1.0	1.2	1.9	1.8	1.4	1.2	1.5
2000.....	4.9	179.2	5.0	1.3	1.4	1.8	1.3	1.4	1.4	1.6
1999.....	4.8	169.3	5.5	1.4	1.7	1.4	1.5	1.0	1.2	1.4

Black, male

2004.....	5.1	169.9	5.8	1.2	*	2.0	2.5	*	1.6	1.7
2003.....	5.1	170.7	4.8	1.2	*	1.7	1.8	*	*	*
2002.....	5.1	151.7	4.6	1.4	1.4	2.1	1.7	1.7	1.6	*
2001.....	5.0	160.8	5.5	*	*	2.6	2.2	*	*	*
2000.....	5.4	181.4	5.5	1.6	1.4	2.5	*	1.9	1.7	1.8
1999.....	5.2	172.7	6.2	1.4	1.8	1.7	1.8	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Black, female

2004.....	4.0	155.6	4.3	1.8	*	1.6	*	*	*	*
2003.....	3.9	143.8	4.6	*	*	*	1.5	*	1.6	*
2002.....	4.3	158.9	4.4	*	*	1.3	1.3	*	*	1.4
2001.....	4.3	155.9	5.1	*	1.4	1.3	1.5	1.7	*	1.7
2000.....	4.4	176.9	4.6	*	1.5	*	*	*	*	1.5
1999.....	4.3	165.8	4.8	1.5	1.7	*	*	*	*	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Anencephaly and similar malformations (Q00)

Black, female

2004.....	*	*	*	*	*	*	*	*	*
2003.....	0.1	7.1	*	*	*	*	*	*	*
2002.....	0.1	7.9	*	*	*	*	*	*	*
2001.....	0.1	7.6	*	*	*	*	*	*	*
2000.....	0.1	7.3	*	*	*	*	*	*	*
1999.....	0.1	8.1	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)

All races 3/, both sexes

2004.....	0.3	10.8	0.8	0.3	0.2	0.2	*	*	*	*
2003.....	0.3	10.5	0.6	0.3	0.2	0.2	0.1	*	0.1	*
2002.....	0.3	9.3	0.8	0.3	0.2	0.2	0.1	*	*	0.1
2001.....	0.3	9.7	0.8	0.2	0.2	0.2	0.1	0.1	*	*
2000.....	0.3	10.7	0.6	0.3	0.2	0.1	0.1	*	*	*
1999.....	0.3	9.4	0.7	0.3	0.2	0.1	*	0.1	0.1	*

All races 3/, male

2004.....	0.3	9.2	0.9	0.2	0.2	0.2	*	*	*	*
2003.....	0.3	10.8	0.7	0.2	*	0.2	*	*	*	*
2002.....	0.3	9.0	0.7	0.3	*	*	*	*	*	*
2001.....	0.3	9.4	0.8	0.2	0.2	*	*	*	*	*
2000.....	0.3	10.6	0.6	0.3	0.2	*	*	*	*	*
1999.....	0.3	8.6	0.7	0.3	0.2	*	*	*	*	*

All races 3/, female

2004.....	0.3	12.4	0.8	0.3	0.2	0.2	*	*	*	*
2003.....	0.3	10.2	0.6	0.3	*	*	*	*	*	*
2002.....	0.3	9.6	0.9	0.3	0.3	*	*	*	*	*
2001.....	0.3	10.0	0.9	*	*	*	*	*	*	*
2000.....	0.3	10.9	0.7	0.3	0.2	*	*	*	*	*
1999.....	0.3	10.2	0.7	0.3	0.2	*	*	*	*	*

White, both sexes

2004.....	0.3	10.7	0.8	0.3	0.2	0.2	*	*	*	*
2003.....	0.3	10.2	0.6	0.2	0.2	0.2	0.1	*	*	*
2002.....	0.3	9.4	0.7	0.3	0.2	0.2	0.2	*	*	*
2001.....	0.3	9.6	0.7	0.2	0.2	0.2	*	*	*	*
2000.....	0.3	10.5	0.6	0.3	0.2	*	*	*	*	*
1999.....	0.2	8.8	0.6	0.2	0.2	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Congenital malformations of heart (Q20-Q24)

All races 3/, both sexes

2004.....	1.0	34.1	1.2	0.4	0.3	0.6	0.5	0.5	0.5	0.4
2003.....	1.1	36.1	1.4	0.3	0.4	0.4	0.4	0.5	0.5	0.5
2002.....	1.1	37.4	1.2	0.3	0.4	0.6	0.5	0.5	0.6	0.5
2001.....	1.2	38.6	1.3	0.2	0.4	0.6	0.6	0.5	0.5	0.5
2000.....	1.2	42.9	1.2	0.3	0.4	0.6	0.5	0.5	0.7	0.5
1999.....	1.3	42.1	1.3	0.3	0.5	0.6	0.5	0.6	0.6	0.6

All races 3/, male

2004.....	1.1	36.7	1.4	0.4	0.3	0.7	0.6	0.6	0.6	0.5
2003.....	1.2	40.1	1.5	0.3	0.4	0.6	0.5	0.6	0.6	0.6
2002.....	1.2	40.4	1.5	0.3	0.5	0.7	0.6	0.7	0.8	0.6
2001.....	1.3	41.8	1.4	0.3	0.4	0.9	0.7	0.6	0.7	0.6
2000.....	1.3	44.5	1.3	0.3	0.4	0.7	0.6	0.6	0.9	0.6
1999.....	1.4	45.7	1.5	0.3	0.6	0.8	0.7	0.6	0.7	0.7

All races 3/, female

2004.....	0.9	31.4	1.0	0.3	0.3	0.5	0.4	0.4	0.4	0.3
2003.....	1.0	31.9	1.4	0.3	0.4	0.2	0.3	0.4	0.5	0.5
2002.....	1.0	34.4	1.0	0.3	0.3	0.4	0.3	0.4	0.5	0.5
2001.....	1.1	35.2	1.2	0.2	0.3	0.3	0.5	0.5	0.4	0.5
2000.....	1.1	41.3	1.2	0.3	0.3	0.4	0.4	0.4	0.5	0.4
1999.....	1.2	38.3	1.1	0.3	0.4	0.4	0.4	0.5	0.6	0.5

White, both sexes

2004.....	1.0	33.2	1.1	0.3	0.3	0.5	0.4	0.5	0.5	0.4
2003.....	1.0	35.0	1.3	0.3	0.4	0.4	0.3	0.5	0.5	0.5
2002.....	1.1	36.4	1.3	0.3	0.4	0.6	0.4	0.5	0.7	0.6
2001.....	1.1	37.4	1.2	0.2	0.3	0.5	0.6	0.5	0.6	0.5
2000.....	1.2	41.5	1.1	0.3	0.3	0.5	0.5	0.5	0.7	0.5
1999.....	1.3	42.0	1.1	0.3	0.4	0.6	0.5	0.6	0.7	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations of heart (Q20-Q24)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
White, male										
2004.....	1.1	35.9	1.3	0.4	0.4	0.6	0.4	0.7	0.6	0.5
2003.....	1.1	39.3	1.3	0.3	0.4	0.5	0.4	0.6	0.6	0.6
2002.....	1.2	40.0	1.5	0.3	0.4	0.7	0.6	0.6	0.8	0.6
2001.....	1.2	41.3	1.3	0.3	0.4	0.7	0.6	0.6	0.7	0.6
2000.....	1.3	43.6	1.2	0.3	0.4	0.6	0.6	0.6	0.9	0.6
1999.....	1.3	46.2	1.3	0.3	0.5	0.8	0.6	0.6	0.7	0.7
White, female										
2004.....	0.9	30.5	0.9	0.3	0.3	0.4	0.4	0.4	0.4	0.3
2003.....	1.0	30.5	1.3	0.3	0.4	*	*	0.4	0.5	0.5
2002.....	1.0	32.7	1.0	0.3	0.3	0.4	0.3	0.4	0.6	0.5
2001.....	1.0	33.2	1.0	*	0.3	0.3	0.5	0.4	0.4	0.5
2000.....	1.1	39.3	1.1	0.3	0.3	0.4	0.4	0.4	0.6	0.4
1999.....	1.2	37.6	1.0	0.3	0.3	0.4	0.4	0.6	0.6	0.5
Black, both sexes										
2004.....	1.3	42.2	1.7	0.7	*	0.9	0.7	*	*	*
2003.....	1.4	41.7	2.1	*	0.6	0.7	0.8	*	0.8	*
2002.....	1.4	44.3	1.3	*	*	0.8	0.7	*	0.7	*
2001.....	1.6	47.7	1.9	*	*	1.2	1.0	*	*	*
2000.....	1.6	51.5	1.8	*	0.7	1.1	0.7	*	*	0.7
1999.....	1.6	47.9	2.2	*	0.8	0.7	0.9	*	*	0.8
Black, male										
2004.....	1.5	44.8	2.0	*	*	*	*	*	*	*
2003.....	1.6	44.9	2.2	*	*	*	*	*	*	*
2002.....	1.6	43.0	*	*	*	*	*	*	*	*
2001.....	1.7	48.8	1.7	*	*	1.9	*	*	*	*
2000.....	1.8	49.9	2.1	*	*	1.6	*	*	*	*
1999.....	1.8	50.1	2.5	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations of heart (Q20-Q24)

Black, female

2004.....	1.1	39.5	*	*	*	*	*	*	*
2003.....	1.2	38.3	2.1	*	*	*	*	*	*
2002.....	1.3	45.7	*	*	*	*	*	*	*
2001.....	1.4	46.6	2.0	*	*	*	*	*	*
2000.....	1.4	53.2	*	*	*	*	*	*	*
1999.....	1.4	45.6	2.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other congenital malformations of circulatory system (Q25-Q28)

Black, female

2004.....	0.3	9.6	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.3	6.6	*	*	*	*	*	*	*
2000.....	0.4	7.6	*	*	*	*	*	*	*
1999.....	0.4	10.4	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other congenital malformations and deformations (Q10-Q18,Q35-Q89)

All races 3/, both sexes

2004.....	0.9	36.4	0.8	0.2	0.1	0.2	0.2	0.2	0.2	0.2
2003.....	0.9	37.5	0.7	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2002.....	0.9	36.2	0.7	0.2	0.2	0.3	0.3	0.2	0.2	0.2
2001.....	0.9	36.3	0.8	0.2	0.2	0.2	0.2	0.2	0.2	0.3
2000.....	0.9	37.9	0.7	0.2	0.2	0.2	0.2	0.2	0.2	0.3
1999.....	0.9	38.3	0.9	0.2	0.1	0.1	0.2	0.2	0.2	0.2

All races 3/, male

2004.....	1.0	38.7	0.9	*	*	0.3	0.3	0.2	0.2	0.3
2003.....	1.0	42.8	0.7	0.2	*	0.3	0.3	0.3	0.2	0.2
2002.....	1.0	39.2	0.6	0.2	0.2	0.3	0.3	0.2	*	0.2
2001.....	1.0	41.1	0.8	*	*	0.2	0.3	0.2	*	0.3
2000.....	1.0	42.1	0.7	0.2	*	0.2	0.2	0.2	0.3	0.3
1999.....	1.0	42.5	1.1	0.2	*	*	0.3	*	0.2	0.2

All races 3/, female

2004.....	0.9	34.0	0.7	*	*	0.2	0.2	*	0.2	*
2003.....	0.8	31.9	0.7	*	*	*	0.2	*	0.2	0.2
2002.....	0.8	33.1	0.8	*	*	0.2	0.3	*	*	*
2001.....	0.8	31.3	0.8	*	*	*	*	0.2	*	0.3
2000.....	0.8	33.6	0.6	*	0.2	*	0.2	*	*	0.2
1999.....	0.8	33.8	0.8	*	*	*	*	*	*	0.2

White, both sexes

2004.....	0.9	35.6	0.8	0.2	0.1	0.2	0.2	0.2	0.2	0.2
2003.....	0.9	36.4	0.7	0.1	0.1	0.2	0.2	0.2	0.2	0.3
2002.....	0.9	37.0	0.7	0.2	0.2	0.2	0.3	0.2	0.2	0.2
2001.....	0.9	36.0	0.7	0.2	0.1	0.2	0.2	0.2	0.2	0.3
2000.....	0.9	37.4	0.6	0.2	0.2	0.2	0.3	0.2	0.2	0.3
1999.....	0.9	36.8	0.9	0.2	0.1	0.2	0.2	0.2	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other congenital malformations and deformations (Q10-Q18,Q35-Q89)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
White, male										
2004.....	0.9	37.5	0.8	*	*	0.3	0.3	*	*	0.2
2003.....	1.0	41.0	0.7	*	*	0.3	0.3	0.3	0.3	0.2
2002.....	1.0	40.4	0.6	*	*	0.2	0.3	0.3	*	*
2001.....	1.0	40.6	0.6	*	*	0.3	0.3	*	*	0.3
2000.....	1.0	41.3	0.7	*	*	*	0.3	*	0.2	0.3
1999.....	1.0	42.1	1.0	*	*	*	0.3	*	*	0.2
White, female										
2004.....	0.9	33.7	0.7	*	*	*	*	*	0.3	*
2003.....	0.8	31.6	0.7	*	*	*	*	*	*	0.3
2002.....	0.8	33.4	0.7	*	*	*	0.3	*	*	*
2001.....	0.8	31.1	0.7	*	*	*	*	*	*	0.3
2000.....	0.8	33.2	0.6	*	*	*	*	*	*	0.2
1999.....	0.8	31.2	0.7	*	*	*	*	*	*	0.2
Black, both sexes										
2004.....	1.1	40.2	1.1	*	*	*	*	*	*	*
2003.....	1.2	44.4	0.9	*	*	*	*	*	*	*
2002.....	1.1	37.1	1.0	*	*	*	*	*	*	*
2001.....	1.2	41.9	0.9	*	*	*	*	*	*	*
2000.....	1.1	42.9	1.0	*	*	*	*	*	*	*
1999.....	1.2	46.3	1.4	*	*	*	*	*	*	*
Black, male										
2004.....	1.4	47.4	1.5	*	*	*	*	*	*	*
2003.....	1.4	52.1	*	*	*	*	*	*	*	*
2002.....	1.3	40.1	*	*	*	*	*	*	*	*
2001.....	1.4	47.6	*	*	*	*	*	*	*	*
2000.....	1.4	48.0	*	*	*	*	*	*	*	*
1999.....	1.3	49.8	1.6	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)

All races 3/, both sexes

2004.....	0.4	24.4	0.3	*	*	*	*	*	*	*
2003.....	0.4	23.5	0.3	*	*	*	*	*	*	*
2002.....	0.4	24.3	0.2	*	*	*	*	*	*	*
2001.....	0.4	22.7	0.3	*	*	*	*	*	*	*
2000.....	0.4	24.2	0.3	*	*	*	*	*	*	*
1999.....	0.3	22.5	0.3	*	*	*	*	*	*	*

All races 3/, male

2004.....	0.3	20.2	*	*	*	*	*	*	*	*
2003.....	0.3	20.0	*	*	*	*	*	*	*	*
2002.....	0.3	20.7	*	*	*	*	*	*	*	*
2001.....	0.3	20.1	*	*	*	*	*	*	*	*
2000.....	0.3	21.4	*	*	*	*	*	*	*	*
1999.....	0.3	18.7	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.4	28.8	0.4	*	*	*	*	*	*	*
2003.....	0.4	27.1	0.4	*	*	*	*	*	*	*
2002.....	0.4	28.0	*	*	*	*	*	*	*	*
2001.....	0.4	25.4	0.5	*	*	*	*	*	*	*
2000.....	0.4	27.2	0.3	*	*	*	*	*	*	*
1999.....	0.4	26.5	0.3	*	*	*	*	*	*	*

White, both sexes

2004.....	0.4	23.9	0.2	*	*	*	*	*	*	*
2003.....	0.3	23.0	0.3	*	*	*	*	*	*	*
2002.....	0.4	24.6	0.2	*	*	*	*	*	*	*
2001.....	0.3	22.9	0.3	*	*	*	*	*	*	*
2000.....	0.3	23.8	0.2	*	*	*	*	*	*	*
1999.....	0.3	22.7	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2004.....	10.3	82.4	1.2	0.3	0.3	1.0	2.1	2.7	2.8	3.6
2003.....	10.8	82.9	1.6	0.4	0.5	1.5	2.6	3.2	3.4	4.8
2002.....	10.4	85.7	1.6	0.3	0.4	1.3	2.4	3.0	3.6	4.6
2001.....	11.4	86.5	1.8	0.4	0.5	1.8	3.0	3.5	4.2	6.2
2000.....	11.3	95.6	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.7
1999.....	9.6	93.2	1.5	0.3	0.4	1.1	2.0	2.3	3.1	4.1

All races 3/, male

2004.....	9.5	96.9	1.2	0.3	0.2	1.3	3.0	3.6	3.6	4.6
2003.....	10.2	96.5	1.8	0.3	0.6	1.9	3.4	4.4	4.5	6.4
2002.....	10.1	98.5	1.8	0.4	0.4	1.6	3.2	4.1	4.9	6.1
2001.....	11.3	98.9	2.1	0.4	0.5	2.4	4.0	4.8	5.6	8.0
2000.....	11.5	110.6	1.7	0.5	0.5	1.9	4.0	4.7	5.9	7.6
1999.....	9.4	106.6	1.7	0.3	0.4	1.5	2.8	2.9	4.2	5.3

All races 3/, female

2004.....	11.0	67.3	1.1	0.2	0.4	0.7	1.3	1.7	2.0	2.7
2003.....	11.4	68.6	1.5	0.5	0.4	1.0	1.7	1.9	2.3	3.3
2002.....	10.7	72.3	1.4	0.2	0.3	0.8	1.6	1.8	2.3	3.1
2001.....	11.4	73.6	1.5	0.4	0.5	1.1	1.9	2.2	2.8	4.4
2000.....	11.2	79.9	1.5	0.4	0.4	1.2	1.8	2.2	2.8	3.7
1999.....	9.7	79.2	1.3	0.3	0.3	0.8	1.2	1.6	2.1	3.0

White, both sexes

2004.....	10.5	65.8	1.0	0.2	0.2	1.0	2.2	2.6	2.7	3.6
2003.....	11.1	68.1	1.4	0.4	0.4	1.5	2.5	3.1	3.4	4.7
2002.....	10.5	71.0	1.3	0.2	0.3	1.2	2.3	2.8	3.5	4.4
2001.....	11.6	70.9	1.5	0.3	0.4	1.8	3.0	3.5	4.1	6.0
2000.....	11.4	77.2	1.5	0.4	0.4	1.5	2.8	3.2	4.2	5.6
1999.....	9.4	74.7	1.3	0.3	0.3	1.0	1.8	2.0	2.9	3.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, male

2004.....	9.4	76.9	1.1	0.3	*	1.2	3.0	3.6	3.5	4.6
2003.....	10.1	82.4	1.4	0.3	0.4	2.0	3.4	4.2	4.5	6.1
2002.....	9.8	82.5	1.4	0.3	0.4	1.6	3.2	3.9	4.7	5.8
2001.....	11.2	82.0	1.7	0.3	0.5	2.4	4.0	4.9	5.5	7.7
2000.....	11.3	89.4	1.7	0.4	0.4	1.9	3.9	4.4	5.7	7.7
1999.....	9.0	86.4	1.6	0.3	0.3	1.3	2.6	2.7	3.8	4.9

White, female

2004.....	11.6	54.2	1.0	*	0.3	0.7	1.3	1.5	1.9	2.5
2003.....	12.0	53.3	1.3	0.4	0.3	1.0	1.6	1.9	2.1	3.2
2002.....	11.2	59.0	1.2	*	*	0.8	1.4	1.6	2.2	3.0
2001.....	11.9	59.3	1.4	0.4	0.4	1.2	1.9	1.9	2.7	4.3
2000.....	11.5	64.3	1.3	0.3	0.4	1.1	1.6	1.9	2.6	3.5
1999.....	9.9	62.4	1.1	0.3	*	0.7	1.0	1.3	2.1	2.5

Black, both sexes

2004.....	11.6	171.7	1.9	*	0.6	1.3	2.3	3.9	4.2	5.2
2003.....	12.3	159.3	3.1	*	1.0	1.5	3.4	4.9	5.0	7.1
2002.....	12.3	162.3	2.6	0.7	0.7	1.8	3.2	4.6	5.6	7.5
2001.....	12.6	169.5	2.8	0.6	0.6	1.8	3.8	4.6	5.3	8.2
2000.....	13.7	197.1	2.3	0.6	0.8	2.2	4.3	5.3	6.3	7.3
1999.....	12.7	192.8	2.7	*	*	1.7	3.7	4.3	5.4	7.8

Black, male

2004.....	12.6	204.3	2.1	*	*	1.6	3.1	4.8	5.5	6.3
2003.....	13.5	170.7	3.4	*	1.3	1.8	4.2	7.3	6.2	9.6
2002.....	14.1	183.0	3.5	*	*	2.4	3.6	5.8	7.8	10.3
2001.....	13.9	187.4	3.5	*	*	2.4	5.2	5.5	6.5	10.5
2000.....	15.5	227.1	2.3	*	*	2.0	5.6	7.1	8.2	8.3
1999.....	14.6	217.6	2.8	*	*	2.4	4.7	5.2	8.3	9.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, female

2004.....	10.8	137.7	1.7	*	*	*	1.5	3.0	2.9	4.3
2003.....	11.3	147.5	2.9	*	*	*	2.7	2.7	3.9	4.9
2002.....	10.6	140.8	1.8	*	*	*	2.7	3.5	3.7	5.1
2001.....	11.4	150.6	2.0	*	*	*	2.4	3.9	4.2	6.2
2000.....	12.0	166.0	2.3	*	*	2.3	3.0	3.6	4.6	6.3
1999.....	10.9	167.1	2.7	*	*	*	2.6	3.4	2.8	6.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Ill-defined and unknown causes of mortality (R95-R99)

All races 3/, both sexes

2004.....	5.3	80.7	1.0	0.2	0.2	0.8	1.7	2.1	2.2	3.0
2003.....	6.0	81.2	1.3	0.3	0.4	1.2	2.0	2.5	2.7	4.0
2002.....	6.0	83.8	1.3	0.2	0.3	1.0	1.9	2.3	2.8	3.7
2001.....	7.1	84.8	1.5	0.3	0.3	1.5	2.5	3.0	3.6	5.4
2000.....	7.3	93.9	1.2	0.3	0.3	1.3	2.4	2.7	3.7	4.8
1999.....	5.9	91.3	1.3	0.2	0.3	0.9	1.7	1.7	2.6	3.3

All races 3/, male

2004.....	6.0	94.9	1.1	0.2	*	1.0	2.3	2.9	2.9	3.8
2003.....	6.8	94.3	1.4	0.2	0.4	1.6	2.7	3.4	3.7	5.3
2002.....	6.8	96.5	1.5	0.3	0.3	1.3	2.6	3.2	3.8	5.1
2001.....	8.2	96.5	1.8	0.3	0.3	2.0	3.4	4.1	4.9	6.9
2000.....	8.5	108.9	1.4	0.3	0.4	1.6	3.3	3.7	5.1	6.5
1999.....	6.7	104.3	1.4	0.2	0.3	1.2	2.4	2.2	3.5	4.2

All races 3/, female

2004.....	4.6	65.8	0.9	*	0.2	0.6	1.0	1.2	1.5	2.2
2003.....	5.2	67.6	1.2	0.4	0.3	0.9	1.2	1.5	1.7	2.7
2002.....	5.1	70.4	1.1	*	0.2	0.6	1.2	1.4	1.8	2.4
2001.....	6.1	72.4	1.2	0.2	0.4	1.0	1.5	1.7	2.3	3.8
2000.....	6.2	78.0	1.1	0.2	0.3	1.0	1.4	1.7	2.2	3.2
1999.....	5.2	77.7	1.1	0.3	0.2	0.6	0.9	1.2	1.6	2.4

White, both sexes

2004.....	5.2	64.4	0.8	0.2	0.1	0.8	1.8	2.1	2.2	3.0
2003.....	5.9	66.4	1.1	0.3	0.3	1.3	2.0	2.4	2.7	4.0
2002.....	5.8	69.5	1.1	0.2	0.2	1.0	2.0	2.2	2.8	3.6
2001.....	7.1	69.3	1.3	0.2	0.3	1.5	2.5	3.0	3.6	5.4
2000.....	7.2	75.6	1.2	0.3	0.3	1.2	2.3	2.6	3.7	4.9
1999.....	5.6	73.1	1.1	0.2	0.2	0.8	1.6	1.6	2.4	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Ill-defined and unknown causes of mortality (R95-R99)

White, male

2004.....	5.8	75.7	0.9	*	*	1.0	2.5	3.1	3.0	3.9
2003.....	6.6	80.2	1.1	*	0.3	1.7	2.8	3.3	3.8	5.2
2002.....	6.5	81.1	1.1	*	0.2	1.3	2.7	3.2	3.8	4.9
2001.....	8.1	79.9	1.5	*	0.3	2.0	3.5	4.3	4.8	6.9
2000.....	8.2	87.8	1.4	0.3	0.3	1.6	3.3	3.6	5.2	6.6
1999.....	6.2	84.5	1.3	*	0.2	1.1	2.3	2.1	3.3	4.0

White, female

2004.....	4.6	52.7	0.7	*	*	0.6	1.0	1.1	1.5	2.1
2003.....	5.2	51.9	1.1	0.3	0.2	0.8	1.2	1.5	1.6	2.7
2002.....	5.1	57.4	1.0	*	*	0.6	1.2	1.2	1.8	2.4
2001.....	6.1	58.2	1.1	*	0.3	1.0	1.5	1.6	2.2	3.8
2000.....	6.2	62.7	0.9	*	0.3	0.9	1.3	1.6	2.1	3.1
1999.....	5.1	61.2	0.8	0.3	*	0.5	0.7	1.0	1.6	2.1

Black, both sexes

2004.....	7.3	168.5	1.7	*	*	0.9	1.4	2.5	2.6	3.8
2003.....	8.2	158.1	2.5	*	0.7	1.2	2.5	3.4	3.8	5.4
2002.....	8.4	158.0	2.3	*	0.6	1.1	2.0	3.4	3.6	5.6
2001.....	8.8	167.3	2.3	*	*	1.2	2.5	3.4	4.4	6.3
2000.....	9.9	194.0	1.8	*	0.7	1.7	3.3	3.8	4.6	5.8
1999.....	9.1	189.1	2.3	*	*	1.3	2.7	2.8	4.0	5.8

Black, male

2004.....	8.8	199.0	1.7	*	*	*	1.8	2.9	3.5	4.2
2003.....	9.9	168.7	2.8	*	*	1.4	3.2	5.2	4.7	7.2
2002.....	10.4	178.1	2.9	*	*	1.4	2.5	4.0	5.1	8.0
2001.....	10.4	184.7	2.9	*	*	1.7	3.5	4.2	5.3	7.9
2000.....	12.0	224.2	1.9	*	*	1.7	4.2	5.2	6.3	6.5
1999.....	11.2	212.7	2.3	*	*	1.7	3.6	3.7	6.2	7.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Ill-defined and unknown causes of mortality (R95-R99)

Black, female

2004.....	6.0	136.8	1.6	*	*	*	*	2.1	1.7	3.5
2003.....	6.7	147.2	2.1	*	*	*	1.9	1.7	2.9	3.8
2002.....	6.5	137.1	1.7	*	*	*	1.6	2.8	2.2	3.5
2001.....	7.4	149.0	1.6	*	*	*	1.5	2.6	3.5	4.9
2000.....	7.9	162.7	1.6	*	*	1.6	2.4	2.5	3.1	5.2
1999.....	7.2	164.8	2.4	*	*	*	1.9	2.0	2.1	4.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other ill-defined and unknown causes of mortality (R96-R99)

All races 3/, both sexes

2004.....	4.5	25.6	1.0	0.2	0.2	0.8	1.7	2.1	2.2	3.0
2003.....	5.2	27.2	1.3	0.3	0.4	1.2	2.0	2.5	2.7	4.0
2002.....	5.2	26.9	1.3	0.2	0.3	1.0	1.9	2.3	2.8	3.7
2001.....	6.3	29.4	1.5	0.3	0.3	1.5	2.5	3.0	3.6	5.4
2000.....	6.4	27.6	1.2	0.3	0.3	1.3	2.4	2.7	3.7	4.8
1999.....	5.0	21.6	1.3	0.2	0.3	0.9	1.7	1.7	2.6	3.3

All races 3/, male

2004.....	5.1	31.2	1.1	0.2	*	1.0	2.3	2.9	2.9	3.8
2003.....	5.9	31.7	1.4	0.2	0.4	1.6	2.7	3.4	3.7	5.3
2002.....	5.9	31.1	1.5	0.3	0.3	1.3	2.6	3.2	3.8	5.1
2001.....	7.3	32.7	1.8	0.3	0.3	2.0	3.4	4.1	4.9	6.9
2000.....	7.4	31.5	1.4	0.3	0.4	1.6	3.3	3.7	5.1	6.5
1999.....	5.5	24.4	1.4	0.2	0.3	1.2	2.4	2.2	3.5	4.2

All races 3/, female

2004.....	4.0	19.7	0.9	*	0.2	0.6	1.0	1.2	1.5	2.2
2003.....	4.6	22.5	1.2	0.4	0.3	0.9	1.2	1.5	1.7	2.7
2002.....	4.5	22.5	1.1	*	0.2	0.6	1.2	1.4	1.8	2.4
2001.....	5.5	25.8	1.2	0.2	0.4	1.0	1.5	1.7	2.3	3.8
2000.....	5.5	23.4	1.1	0.2	0.3	1.0	1.4	1.7	2.2	3.2
1999.....	4.5	18.6	1.1	0.3	0.2	0.6	0.9	1.2	1.6	2.4

White, both sexes

2004.....	4.6	19.1	0.8	0.2	0.1	0.8	1.8	2.1	2.2	3.0
2003.....	5.3	22.5	1.1	0.3	0.3	1.3	2.0	2.4	2.7	4.0
2002.....	5.2	21.8	1.1	0.2	0.2	1.0	2.0	2.2	2.8	3.6
2001.....	6.5	24.6	1.3	0.2	0.3	1.5	2.5	3.0	3.6	5.4
2000.....	6.5	21.5	1.2	0.3	0.3	1.2	2.3	2.6	3.7	4.9
1999.....	4.9	15.7	1.1	0.2	0.2	0.8	1.6	1.6	2.4	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other ill-defined and unknown causes of mortality (R96-R99)

White, male

2004.....	5.1	23.5	0.9	*	*	1.0	2.5	3.1	3.0	3.9
2003.....	5.9	27.3	1.1	*	0.3	1.7	2.8	3.3	3.8	5.2
2002.....	5.8	25.6	1.1	*	0.2	1.3	2.7	3.2	3.8	4.9
2001.....	7.3	27.6	1.5	*	0.3	2.0	3.5	4.3	4.8	6.9
2000.....	7.4	24.4	1.4	0.3	0.3	1.6	3.3	3.6	5.2	6.6
1999.....	5.3	18.2	1.3	*	0.2	1.1	2.3	2.1	3.3	4.0

White, female

2004.....	4.1	14.4	0.7	*	*	0.6	1.0	1.1	1.5	2.1
2003.....	4.7	17.6	1.1	0.3	0.2	0.8	1.2	1.5	1.6	2.7
2002.....	4.6	17.9	1.0	*	*	0.6	1.2	1.2	1.8	2.4
2001.....	5.6	21.4	1.1	*	0.3	1.0	1.5	1.6	2.2	3.8
2000.....	5.7	18.4	0.9	*	0.3	0.9	1.3	1.6	2.1	3.1
1999.....	4.5	13.1	0.8	0.3	*	0.5	0.7	1.0	1.6	2.1

Black, both sexes

2004.....	5.4	58.7	1.7	*	*	0.9	1.4	2.5	2.6	3.8
2003.....	6.4	53.4	2.5	*	0.7	1.2	2.5	3.4	3.8	5.4
2002.....	6.5	53.8	2.3	*	0.6	1.1	2.0	3.4	3.6	5.6
2001.....	6.8	53.0	2.3	*	*	1.2	2.5	3.4	4.4	6.3
2000.....	7.6	59.7	1.8	*	0.7	1.7	3.3	3.8	4.6	5.8
1999.....	6.8	50.5	2.3	*	*	1.3	2.7	2.8	4.0	5.8

Black, male

2004.....	6.4	69.7	1.7	*	*	*	1.8	2.9	3.5	4.2
2003.....	7.9	59.5	2.8	*	*	1.4	3.2	5.2	4.7	7.2
2002.....	8.2	60.7	2.9	*	*	1.4	2.5	4.0	5.1	8.0
2001.....	8.0	58.4	2.9	*	*	1.7	3.5	4.2	5.3	7.9
2000.....	9.3	70.0	1.9	*	*	1.7	4.2	5.2	6.3	6.5
1999.....	8.5	57.6	2.3	*	*	1.7	3.6	3.7	6.2	7.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other ill-defined and unknown causes of mortality (R96-R99)

Black, female

2004.....	4.6	47.2	1.6	*	*	*	*	2.1	1.7	3.5
2003.....	5.1	47.0	2.1	*	*	*	1.9	1.7	2.9	3.8
2002.....	5.0	46.6	1.7	*	*	*	1.6	2.8	2.2	3.5
2001.....	5.7	47.3	1.6	*	*	*	1.5	2.6	3.5	4.9
2000.....	6.1	48.9	1.6	*	*	1.6	2.4	2.5	3.1	5.2
1999.....	5.3	43.3	2.4	*	*	*	1.9	2.0	2.1	4.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

All races 3/, both sexes

2004.....	3.3	1.7	0.2	*	0.1	0.2	0.5	0.6	0.6	0.7
2003.....	3.2	1.6	0.3	0.1	0.1	0.2	0.6	0.7	0.7	0.8
2002.....	2.9	1.9	0.3	*	0.1	0.3	0.5	0.6	0.8	0.9
2001.....	2.7	1.8	0.3	0.1	0.1	0.3	0.5	0.5	0.6	0.8
2000.....	2.6	1.8	0.4	0.1	0.1	0.3	0.5	0.7	0.7	0.8
1999.....	2.5	1.9	0.3	*	*	0.3	0.4	0.6	0.6	0.8

All races 3/, male

2004.....	2.5	2.0	*	*	*	0.3	0.6	0.7	0.7	0.8
2003.....	2.5	2.2	0.4	*	*	0.3	0.7	1.0	0.8	1.1
2002.....	2.4	1.9	0.4	*	*	0.4	0.5	0.8	1.0	1.1
2001.....	2.3	2.3	0.3	*	0.2	0.4	0.6	0.7	0.7	1.0
2000.....	2.3	1.7	0.3	*	*	0.3	0.7	0.9	0.7	1.1
1999.....	2.1	2.3	0.3	*	*	0.3	0.4	0.7	0.7	1.0

All races 3/, female

2004.....	4.1	1.5	*	*	*	*	0.3	0.5	0.5	0.5
2003.....	3.9	1.1	0.3	*	*	*	0.5	0.4	0.6	0.6
2002.....	3.5	1.9	*	*	*	0.2	0.4	0.4	0.6	0.7
2001.....	3.2	1.2	0.3	*	*	*	0.4	0.4	0.5	0.6
2000.....	3.0	1.8	0.4	*	*	0.3	0.4	0.5	0.6	0.5
1999.....	2.9	1.5	*	*	*	*	0.3	0.5	0.5	0.6

White, both sexes

2004.....	3.5	1.4	0.2	*	*	0.2	0.4	0.5	0.5	0.6
2003.....	3.3	1.8	0.3	*	*	0.2	0.5	0.6	0.6	0.7
2002.....	3.0	1.5	0.3	*	*	0.2	0.4	0.6	0.7	0.8
2001.....	2.8	1.7	0.2	*	0.1	0.3	0.4	0.5	0.6	0.7
2000.....	2.7	1.6	0.4	*	0.1	0.3	0.5	0.6	0.5	0.7
1999.....	2.5	1.6	0.3	*	*	0.2	0.3	0.5	0.5	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
White, male										
2004.....	2.5	1.2	*	*	*	0.3	0.5	0.5	0.5	0.7
2003.....	2.5	2.2	0.4	*	*	0.3	0.7	0.9	0.7	0.9
2002.....	2.4	1.4	*	*	*	0.2	0.4	0.8	0.8	0.9
2001.....	2.2	2.2	*	*	*	0.4	0.5	0.6	0.6	0.8
2000.....	2.2	1.6	0.3	*	*	0.3	0.6	0.8	0.6	1.0
1999.....	2.1	1.9	*	*	*	0.3	0.3	0.6	0.5	0.9
White, female										
2004.....	4.4	1.5	*	*	*	*	0.3	0.4	0.4	0.4
2003.....	4.2	1.3	*	*	*	*	0.4	0.3	0.5	0.5
2002.....	3.7	1.6	*	*	*	*	*	0.4	0.5	0.6
2001.....	3.4	*	*	*	*	*	0.4	0.3	0.5	0.5
2000.....	3.2	1.6	0.4	*	*	*	0.3	0.4	0.4	0.4
1999.....	3.0	*	*	*	*	*	*	0.3	0.5	0.4
Black, both sexes										
2004.....	3.5	3.2	*	*	*	*	1.0	1.4	1.6	1.4
2003.....	3.4	*	*	*	*	*	0.9	1.5	1.2	1.7
2002.....	3.2	4.3	*	*	*	0.7	1.1	1.2	2.1	1.9
2001.....	3.1	*	*	*	*	*	1.3	1.3	0.9	1.9
2000.....	3.0	*	*	*	*	*	1.0	1.5	1.7	1.4
1999.....	3.0	3.6	*	*	*	*	1.0	1.4	1.4	2.0
Black, male										
2004.....	3.3	*	*	*	*	*	1.3	1.9	2.1	2.1
2003.....	3.2	*	*	*	*	*	*	2.1	1.5	2.4
2002.....	3.3	*	*	*	*	*	*	1.7	2.8	2.3
2001.....	3.1	*	*	*	*	*	1.7	*	*	2.6
2000.....	3.0	*	*	*	*	*	*	2.0	1.9	1.8
1999.....	3.0	*	*	*	*	*	*	*	2.1	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ year	year 2/ year	years	years	years	years	years	years	years	years

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

Black, female

2004.....	3.7	*	*	*	*	*	*	*	*	*
2003.....	3.6	*	*	*	*	*	*	*	*	*
2002.....	3.2	*	*	*	*	*	*	*	1.4	1.6
2001.....	3.0	*	*	*	*	*	*	*	*	1.3
2000.....	3.0	*	*	*	*	*	*	*	1.5	*
1999.....	3.0	*	*	*	*	*	*	1.4	*	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XX. External causes of mortality (*U01-*U03,V01-Y89)

All races 3/, both sexes

2004.....	57.9	36.7	13.0	6.5	9.8	51.4	70.7	62.5	55.5	59.5
2003.....	57.4	34.4	13.6	6.3	9.5	50.7	72.1	60.3	56.0	60.3
2002.....	56.9	33.8	13.7	6.8	9.9	52.6	71.9	60.6	55.5	61.1
2001 4/.....	56.2	35.1	14.3	7.2	9.9	51.0	70.6	60.9	55.6	62.4
2000.....	53.8	34.5	14.6	7.7	10.6	51.7	68.9	56.5	51.9	57.4
1999.....	54.2	33.9	15.3	8.1	10.7	52.7	67.2	56.4	54.0	57.6

All races 3/, male

2004.....	80.3	40.4	14.7	7.6	12.3	74.0	111.2	98.0	84.0	87.2
2003.....	80.2	39.2	15.9	7.3	12.7	73.8	113.1	94.6	85.5	88.7
2002.....	79.8	37.6	16.3	7.9	12.7	76.5	114.0	95.6	84.8	90.3
2001 4/.....	79.3	38.9	16.8	8.5	12.9	75.1	113.0	96.1	85.3	92.4
2000.....	75.8	38.1	17.0	8.8	14.0	75.5	109.5	89.6	79.6	86.0
1999.....	76.2	38.3	17.3	9.5	14.1	76.3	106.5	88.9	82.6	86.0

All races 3/, female

2004.....	36.2	32.9	11.3	5.4	7.2	27.6	27.7	25.5	26.4	31.5
2003.....	35.3	29.3	11.2	5.3	6.1	26.3	28.6	24.7	25.9	31.9
2002.....	34.8	29.8	11.1	5.7	6.9	27.5	27.6	24.4	25.7	31.8
2001 4/.....	34.0	31.2	11.8	5.8	6.7	25.4	26.2	24.7	25.4	32.5
2000.....	32.5	30.8	12.1	6.4	7.1	26.5	26.5	22.6	23.8	29.0
1999.....	33.0	29.2	13.3	6.6	7.2	27.8	26.2	23.3	25.2	29.4

White, both sexes

2004.....	59.0	31.3	11.9	5.8	9.3	50.9	67.7	59.0	54.1	59.9
2003.....	58.2	28.9	12.5	5.7	8.9	49.9	67.7	56.4	54.7	60.5
2002.....	57.4	26.1	12.3	5.9	9.3	52.1	67.3	56.1	53.8	60.3
2001 4/.....	56.5	28.8	13.3	6.5	9.3	49.5	65.6	57.1	53.9	62.0
2000.....	53.6	27.6	13.1	7.0	10.3	50.4	63.2	51.7	49.6	56.6
1999.....	53.7	27.9	13.6	7.1	10.5	50.6	61.4	52.1	51.7	56.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XX. External causes of mortality (*U01-*U03,V01-Y89)

White, male

2004.....	80.3	35.3	13.7	6.7	11.7	71.4	104.7	90.1	80.0	87.0
2003.....	79.7	32.8	14.7	6.5	11.9	70.6	104.0	86.0	82.1	87.5
2002.....	79.0	30.7	14.5	6.8	11.8	74.1	104.5	86.0	80.8	87.9
2001 4/.....	78.3	31.8	15.8	7.4	12.1	71.1	102.8	88.2	81.6	91.2
2000.....	74.1	31.8	15.4	8.1	13.4	71.7	98.4	80.1	75.1	84.0
1999.....	74.2	32.6	15.4	8.2	13.6	71.1	95.2	80.8	77.8	83.3

White, female

2004.....	38.1	27.0	10.2	4.9	6.7	29.2	27.9	25.7	26.9	32.0
2003.....	37.1	24.8	10.3	4.9	5.8	27.9	28.6	24.7	26.1	32.8
2002.....	36.4	21.3	10.1	5.0	6.6	28.9	27.7	24.3	25.6	32.2
2001 4/.....	35.2	25.7	10.5	5.6	6.5	26.5	26.0	24.3	25.2	32.2
2000.....	33.6	23.2	10.7	5.9	7.0	27.7	25.7	22.0	23.2	28.7
1999.....	33.9	23.1	11.7	5.9	7.1	28.9	25.4	22.2	24.6	29.0

Black, both sexes

2004.....	62.3	66.5	19.4	10.4	12.7	58.8	97.9	97.1	78.2	71.1
2003.....	62.8	63.5	20.2	9.2	12.3	58.5	105.7	96.5	78.6	72.6
2002.....	63.7	72.8	21.0	11.2	12.9	59.9	106.7	100.3	78.6	78.5
2001 4/.....	64.2	68.9	21.2	10.6	12.6	62.5	107.7	94.8	76.8	76.9
2000.....	63.9	70.6	22.8	11.3	13.0	63.3	111.4	94.0	76.1	73.4
1999.....	65.2	67.1	24.8	13.3	13.2	68.4	108.3	91.2	76.9	75.1

Black, male

2004.....	96.9	68.2	21.4	12.4	15.7	94.8	166.8	171.2	133.1	112.1
2003.....	98.2	72.0	23.9	11.0	16.5	96.9	180.8	170.4	132.1	117.2
2002.....	99.9	73.5	26.0	13.1	17.9	96.4	184.7	178.2	132.9	126.6
2001 4/.....	100.2	77.2	24.1	14.2	17.7	102.8	189.7	165.4	128.5	117.9
2000.....	100.2	72.3	25.5	13.3	18.2	103.1	193.6	165.9	126.7	117.9
1999.....	101.7	71.2	27.7	16.7	18.0	110.9	189.6	158.4	127.6	119.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/	year 2/	years	years	years	years	years	years	years	years

XX. External causes of mortality (*U01-*U03,V01-Y89)

Black, female

2004.....	30.8	64.8	17.4	8.3	9.5	21.8	28.6	27.8	28.7	34.5
2003.....	30.5	54.7	16.4	7.4	7.9	18.9	31.0	28.2	30.3	32.9
2002.....	30.9	72.0	15.8	9.1	7.9	22.2	30.7	29.1	29.9	35.7
2001 4/.....	31.5	60.2	18.3	6.9	7.4	20.9	28.8	30.4	30.3	40.5
2000.....	30.9	68.8	19.9	9.2	7.6	22.6	33.5	29.0	30.5	34.0
1999.....	32.2	62.8	21.9	9.9	8.2	25.0	31.1	30.4	31.3	35.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes

2004.....	38.1	25.8	10.3	5.7	7.3	32.9	41.1	34.6	30.6	34.8
2003.....	37.6	23.6	10.9	5.5	7.2	33.0	41.1	32.8	30.2	35.0
2002.....	37.0	23.5	10.5	5.9	7.3	35.0	40.9	32.3	30.8	34.9
2001.....	35.7	24.2	11.2	6.4	7.4	32.8	39.5	31.4	28.5	34.1
2000.....	34.8	23.1	11.9	6.8	7.7	33.4	38.8	30.4	28.6	32.9
1999.....	35.1	22.3	12.4	7.1	8.1	33.3	37.5	30.0	29.2	32.2

All races 3/, male

2004.....	49.8	29.0	11.8	6.8	9.0	44.6	61.8	52.2	45.0	49.7
2003.....	49.3	26.4	13.0	6.4	9.3	44.7	61.8	49.4	45.1	50.2
2002.....	48.9	26.8	12.9	7.1	9.2	47.6	62.4	49.5	46.1	50.5
2001.....	47.2	26.2	13.3	7.6	9.5	45.3	60.7	48.0	43.1	49.5
2000.....	46.2	25.7	14.0	7.8	9.9	45.8	58.9	47.0	43.1	48.9
1999.....	46.4	25.4	14.4	8.5	10.4	44.8	57.1	46.1	43.9	47.4

All races 3/, female

2004.....	26.8	22.5	8.7	4.6	5.4	20.6	19.1	16.2	15.9	19.7
2003.....	26.2	20.7	8.7	4.6	4.9	20.6	19.2	15.5	15.1	19.7
2002.....	25.6	20.0	8.1	4.7	5.3	21.8	18.4	14.5	15.2	19.2
2001.....	24.5	22.1	8.9	5.0	5.3	19.5	17.2	14.3	13.8	18.7
2000.....	23.8	20.5	9.7	5.7	5.5	20.3	17.8	13.5	13.9	17.0
1999.....	24.1	19.0	10.3	5.6	5.6	21.1	17.1	13.7	14.3	17.0

White, both sexes

2004.....	40.2	22.2	10.0	5.2	7.1	36.0	44.4	36.5	32.1	36.1
2003.....	39.5	20.0	10.6	5.2	7.0	35.9	44.3	34.6	31.6	36.2
2002.....	38.7	18.6	9.9	5.2	7.0	37.9	43.8	33.5	32.2	35.5
2001.....	37.0	20.6	11.0	5.8	7.1	34.9	41.7	32.7	29.3	34.8
2000.....	35.9	19.1	11.1	6.2	7.6	36.0	40.7	30.9	29.1	33.4
1999.....	36.0	18.9	11.5	6.3	7.9	35.6	38.9	30.8	29.3	32.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
White, male										
2004.....	52.0	25.5	11.5	6.1	8.9	48.5	67.0	54.6	46.5	51.4
2003.....	51.3	22.5	12.8	6.0	9.0	48.4	66.2	51.9	46.8	51.4
2002.....	50.4	22.3	11.8	6.3	8.8	51.0	66.4	50.7	47.9	50.9
2001.....	48.4	22.1	13.3	6.7	9.0	47.8	64.2	49.6	44.0	50.3
2000.....	47.0	21.7	13.2	7.2	9.6	48.8	61.7	47.2	43.7	49.5
1999.....	46.9	21.8	13.2	7.4	10.1	47.2	58.9	47.1	43.8	47.5
White, female										
2004.....	28.8	18.8	8.4	4.3	5.2	22.9	20.1	17.2	16.9	20.3
2003.....	28.0	17.4	8.3	4.4	4.8	22.7	20.7	16.1	15.7	20.6
2002.....	27.3	14.7	7.8	4.2	5.2	23.9	19.6	15.1	15.9	19.7
2001.....	25.9	18.9	8.5	4.8	5.1	21.1	17.8	14.8	14.0	19.0
2000.....	25.1	16.4	9.0	5.3	5.5	22.3	18.3	13.8	13.8	17.0
1999.....	25.3	15.8	9.6	5.1	5.6	23.3	17.6	13.8	14.3	17.0
Black, both sexes										
2004.....	32.8	45.8	12.5	8.5	8.5	21.6	30.1	32.2	30.6	34.9
2003.....	32.4	43.5	13.4	7.3	8.1	20.7	30.6	29.8	29.9	35.5
2002.....	33.1	48.9	14.1	9.3	9.0	23.9	32.0	32.8	30.1	38.1
2001.....	33.5	43.1	13.6	9.4	9.4	24.7	33.0	30.5	30.0	35.8
2000.....	33.5	44.5	16.2	9.7	8.8	23.7	34.3	33.1	31.9	35.9
1999.....	35.2	41.6	17.7	11.3	9.6	25.0	34.6	31.5	32.9	36.6
Black, male										
2004.....	46.7	48.3	14.2	10.4	10.1	31.2	44.8	51.7	48.9	51.5
2003.....	46.1	47.6	15.7	8.7	10.6	30.0	47.6	46.6	46.9	54.9
2002.....	47.9	51.1	18.5	11.1	12.1	34.6	49.5	53.6	47.7	58.9
2001.....	48.2	48.5	15.4	12.6	12.9	36.1	51.0	48.3	46.6	53.3
2000.....	49.0	47.7	18.8	11.9	12.1	34.5	52.9	54.1	49.6	55.7
1999.....	51.0	45.9	21.0	14.5	13.0	36.6	54.7	51.0	51.7	55.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ years	year 2/ years	years	years	years	years	years	years	years	years

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Black, female

2004.....	20.1	43.2	10.6	6.5	6.7	11.8	15.4	14.1	14.1	20.0
2003.....	19.9	39.3	11.0	5.9	5.5	11.2	13.7	14.3	14.7	18.3
2002.....	19.7	46.6	9.6	7.5	5.8	12.8	14.9	13.8	14.4	19.7
2001.....	20.1	37.5	11.9	6.2	5.8	12.9	15.6	14.2	15.1	20.2
2000.....	19.5	41.3	13.5	7.6	5.4	12.7	16.7	14.1	15.9	18.3
1999.....	20.8	37.3	14.3	8.0	6.1	13.1	15.6	13.8	16.0	19.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Transport accidents (V01-V99,Y85)

All races 3/, both sexes

2004.....	16.4	3.6	4.1	3.5	5.1	26.0	28.5	20.9	16.7	16.4
2003.....	16.5	3.7	4.1	3.5	5.1	26.7	28.8	20.1	16.5	16.7
2002.....	16.8	3.1	4.1	3.5	4.9	28.4	29.9	20.5	17.1	17.1
2001.....	16.6	3.7	4.4	3.8	5.0	26.6	29.2	20.9	16.9	17.4
2000.....	16.6	4.6	4.5	4.1	5.3	27.1	29.3	20.4	16.8	17.1
1999.....	16.6	5.0	4.6	4.4	5.3	27.1	28.2	20.0	17.0	16.9

All races 3/, male

2004.....	23.0	3.9	4.4	3.9	6.0	33.6	42.0	31.2	24.4	24.0
2003.....	23.3	4.0	4.5	4.0	6.3	34.7	42.4	29.8	24.8	24.1
2002.....	23.7	3.5	4.8	4.0	5.9	37.0	44.4	31.0	25.4	24.7
2001.....	23.5	3.3	5.0	4.3	6.2	35.4	43.9	31.4	25.4	24.7
2000.....	23.3	4.8	5.0	4.4	6.4	35.5	43.3	30.8	24.7	24.8
1999.....	23.1	5.2	5.1	5.0	6.5	34.8	41.8	29.9	24.7	24.0

All races 3/, female

2004.....	9.9	3.2	3.8	3.0	4.1	18.0	14.2	10.2	8.9	8.8
2003.....	10.0	3.4	3.7	3.0	3.8	18.3	14.3	10.0	8.1	9.3
2002.....	10.1	2.8	3.4	3.1	3.9	19.3	14.6	9.7	8.7	9.6
2001.....	10.0	4.2	3.8	3.3	3.8	17.3	13.8	10.1	8.3	10.2
2000.....	10.2	4.3	4.0	3.8	4.1	18.1	14.6	9.8	8.7	9.5
1999.....	10.4	4.8	4.2	3.6	4.1	19.0	14.0	10.0	9.3	9.8

White, both sexes

2004.....	16.9	3.3	4.0	3.4	5.2	28.6	30.3	21.4	17.0	16.7
2003.....	17.0	3.6	4.0	3.5	5.2	29.2	30.4	20.7	16.8	17.0
2002.....	17.3	2.7	3.9	3.4	5.0	31.0	31.5	20.7	17.6	17.2
2001.....	17.0	3.7	4.3	3.6	5.0	28.6	30.4	21.6	17.1	17.6
2000.....	17.0	4.1	4.3	4.1	5.4	29.6	30.5	20.3	16.8	17.3
1999.....	17.0	4.6	4.1	4.1	5.5	29.4	29.2	20.4	17.2	16.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Transport accidents (V01-V99,Y85)

White, male

2004.....	23.7	3.3	4.2	3.8	6.3	36.7	44.8	31.5	24.3	24.3
2003.....	23.9	4.0	4.5	3.9	6.5	37.6	44.4	30.6	25.1	24.2
2002.....	24.2	3.1	4.5	3.8	6.0	40.2	46.7	31.0	25.9	24.5
2001.....	23.9	3.2	4.9	4.0	6.2	37.8	45.8	32.3	25.4	24.7
2000.....	23.6	4.5	4.7	4.4	6.5	38.4	45.0	30.3	24.6	24.8
1999.....	23.4	5.2	4.4	4.7	6.7	37.3	43.1	30.2	24.6	23.7

White, female

2004.....	10.3	3.2	3.8	2.9	4.1	20.0	14.7	10.5	9.3	8.9
2003.....	10.3	3.1	3.4	3.1	3.9	20.2	15.3	10.2	8.2	9.6
2002.....	10.5	2.2	3.4	2.9	3.9	21.3	15.3	9.8	9.0	9.7
2001.....	10.3	4.1	3.6	3.1	3.8	18.9	14.0	10.4	8.4	10.4
2000.....	10.5	3.7	3.8	3.7	4.3	20.2	15.1	9.9	8.7	9.8
1999.....	10.7	3.9	3.8	3.5	4.3	21.0	14.4	10.1	9.6	9.8

Black, both sexes

2004.....	15.2	4.8	5.1	4.0	4.8	16.6	22.7	22.4	19.3	17.6
2003.....	15.1	4.2	5.0	3.6	4.5	16.2	23.2	20.3	18.3	17.9
2002.....	15.4	5.3	5.0	4.6	4.6	17.9	24.6	23.0	18.0	19.2
2001.....	15.9	*	5.2	5.3	5.3	18.8	25.8	21.0	18.9	18.1
2000.....	16.0	6.5	5.6	4.5	4.8	17.3	26.6	23.5	19.4	17.9
1999.....	16.3	7.3	7.1	5.8	5.0	18.7	25.8	21.3	18.5	19.9

Black, male

2004.....	22.8	5.9	5.4	4.4	5.3	22.9	33.2	36.1	31.5	27.9
2003.....	22.6	*	5.1	4.5	5.7	22.7	36.3	31.8	29.4	28.4
2002.....	23.1	*	6.4	5.0	5.6	24.5	37.1	37.5	28.9	29.8
2001.....	23.8	*	5.7	6.6	7.3	26.5	39.1	33.2	30.0	27.2
2000.....	24.1	6.7	6.2	4.9	6.5	24.2	40.3	38.2	30.7	28.6
1999.....	24.2	*	8.0	7.2	6.4	25.8	39.2	34.3	29.4	29.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Transport accidents (V01-V99,Y85)

Black, female

2004.....	8.2	*	4.8	3.6	4.2	10.0	12.1	9.6	8.3	8.5
2003.....	8.3	*	5.0	2.8	3.4	9.5	10.2	9.8	8.2	8.4
2002.....	8.3	*	3.4	4.2	3.5	11.2	12.5	9.8	8.2	9.8
2001.....	8.7	*	4.7	4.0	3.4	10.8	13.0	9.9	8.9	10.0
2000.....	8.5	*	5.0	4.1	3.0	10.3	13.7	10.2	9.2	8.4
1999.....	9.2	8.7	6.2	4.4	3.6	11.4	13.1	9.5	8.6	11.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Railway accidents (V05,V15,V80.6,V81.2-V81.9)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.2
2003.....	0.1	*	*	*	*	0.1	0.2	0.2	0.2	0.3
2002.....	0.2	*	*	*	*	0.2	0.3	0.2	0.2	0.2
2001.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.3
2000.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.3
1999.....	0.2	*	*	*	*	0.2	0.3	0.2	0.2	0.3

All races 3/, male

2004.....	0.3	*	*	*	*	0.3	0.3	0.4	0.4	0.3
2003.....	0.3	*	*	*	*	0.3	0.3	0.3	0.4	0.5
2002.....	0.3	*	*	*	*	0.3	0.5	0.3	0.4	0.4
2001.....	0.3	*	*	*	*	0.3	0.3	0.4	0.4	0.5
2000.....	0.3	*	*	*	*	0.4	0.4	0.4	0.3	0.5
1999.....	0.3	*	*	*	*	0.3	0.6	0.4	0.4	0.6

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	*	0.1	0.2	0.2	0.2	0.2
2003.....	0.1	*	*	*	*	0.1	0.2	0.2	0.2	0.3
2002.....	0.2	*	*	*	*	0.2	0.3	0.2	0.2	0.2
2001.....	0.2	*	*	*	*	0.2	0.2	0.3	0.2	0.2
2000.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.3
1999.....	0.2	*	*	*	*	0.2	0.3	0.2	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes

2004.....	15.3	3.5	4.0	3.3	4.8	25.2	27.5	19.7	15.6	15.2
2003.....	15.4	3.6	3.9	3.3	4.7	25.7	27.5	19.0	15.4	15.4
2002.....	15.7	3.0	3.9	3.3	4.5	27.6	28.8	19.7	16.0	15.8
2001.....	15.4	3.6	4.1	3.6	4.6	25.7	28.0	19.6	15.6	15.9
2000.....	15.4	4.4	4.2	3.8	4.8	26.0	28.0	19.2	15.5	15.6
1999.....	15.2	4.8	4.2	3.9	4.8	25.9	26.5	18.4	15.4	15.1

All races 3/, male

2004.....	21.3	3.8	4.2	3.7	5.6	32.5	40.4	29.3	22.4	21.9
2003.....	21.4	3.9	4.4	3.7	5.9	33.3	40.4	28.2	23.0	21.9
2002.....	21.9	3.3	4.5	3.8	5.3	36.0	42.6	29.6	23.6	22.4
2001.....	21.5	3.2	4.5	4.0	5.7	34.0	42.1	29.5	23.1	22.3
2000.....	21.3	4.6	4.7	4.1	5.8	33.9	41.2	28.6	22.6	22.1
1999.....	20.9	5.0	4.7	4.4	5.7	33.1	39.1	27.2	22.2	21.2

All races 3/, female

2004.....	9.5	3.2	3.7	2.8	3.9	17.5	13.8	9.7	8.5	8.3
2003.....	9.5	3.3	3.5	2.8	3.5	17.8	13.9	9.5	7.7	8.8
2002.....	9.8	2.8	3.3	2.9	3.6	18.9	14.3	9.4	8.3	9.2
2001.....	9.5	4.0	3.6	3.0	3.5	16.9	13.4	9.5	8.0	9.6
2000.....	9.7	4.2	3.8	3.5	3.8	17.5	14.2	9.5	8.2	9.1
1999.....	9.7	4.7	3.8	3.3	3.8	18.3	13.4	9.4	8.5	9.0

White, both sexes

2004.....	15.8	3.2	3.9	3.2	4.9	27.8	29.2	20.0	15.7	15.3
2003.....	15.8	3.5	3.8	3.3	4.8	28.1	29.1	19.5	15.7	15.6
2002.....	16.2	2.6	3.7	3.2	4.6	30.2	30.4	19.8	16.4	15.8
2001.....	15.7	3.5	3.9	3.3	4.6	27.6	29.1	20.2	15.7	16.1
2000.....	15.7	3.9	4.0	3.7	5.0	28.4	29.1	19.0	15.4	15.8
1999.....	15.5	4.5	3.8	3.7	5.0	28.1	27.4	18.6	15.5	14.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, male

2004.....	21.9	3.3	4.0	3.6	5.9	35.6	43.2	29.5	22.2	22.1
2003.....	22.0	3.9	4.3	3.6	6.0	36.1	42.3	28.8	23.3	21.9
2002.....	22.4	2.9	4.2	3.6	5.5	39.0	44.8	29.5	23.9	22.2
2001.....	21.9	3.1	4.4	3.7	5.6	36.2	43.8	30.1	23.1	22.3
2000.....	21.6	4.2	4.5	4.0	5.9	36.8	42.6	27.9	22.4	22.1
1999.....	21.1	5.0	4.1	4.2	5.9	35.5	40.4	27.4	22.1	20.8

White, female

2004.....	9.8	3.2	3.7	2.7	3.9	19.5	14.1	9.9	8.9	8.3
2003.....	9.8	3.0	3.2	2.9	3.6	19.6	14.8	9.6	7.7	9.0
2002.....	10.1	2.2	3.2	2.7	3.7	20.9	15.0	9.5	8.6	9.3
2001.....	9.8	3.8	3.4	2.9	3.5	18.5	13.5	9.7	8.0	9.8
2000.....	10.0	3.5	3.6	3.4	4.0	19.5	14.6	9.6	8.2	9.3
1999.....	10.0	3.9	3.5	3.2	4.0	20.2	13.7	9.5	8.7	8.9

Black, both sexes

2004.....	14.3	4.7	4.9	3.8	4.5	15.8	21.8	21.2	18.4	16.6
2003.....	14.2	4.2	4.9	3.4	4.3	15.5	22.3	19.5	17.3	16.7
2002.....	14.5	5.2	4.8	4.3	4.2	17.4	24.0	22.4	17.2	17.9
2001.....	14.8	*	5.0	4.9	4.8	18.1	25.2	20.1	17.7	16.7
2000.....	15.0	6.5	5.4	4.3	4.6	16.5	25.8	22.8	18.0	16.4
1999.....	15.1	7.1	6.4	5.3	4.6	17.8	24.8	20.1	17.0	18.2

Black, male

2004.....	21.2	*	5.2	4.2	5.0	21.6	31.5	33.8	29.8	26.0
2003.....	21.1	*	5.0	4.1	5.2	21.6	34.6	30.5	27.7	26.3
2002.....	21.5	*	6.3	4.8	5.0	23.7	36.1	36.2	27.4	27.6
2001.....	22.0	*	5.3	6.2	6.5	25.5	38.0	31.9	28.1	24.9
2000.....	22.5	6.7	5.8	4.5	6.2	22.8	38.8	36.7	28.5	26.0
1999.....	22.2	*	7.0	6.3	5.7	24.5	37.7	32.4	27.0	27.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, female

2004.....	7.9	*	4.7	3.5	4.0	9.8	12.0	9.4	8.1	8.2
2003.....	8.0	*	4.8	2.7	3.4	9.2	10.1	9.4	8.0	8.2
2002.....	8.0	*	3.4	3.8	3.3	11.0	12.2	9.7	7.9	9.3
2001.....	8.2	*	4.6	3.6	3.1	10.5	12.9	9.5	8.4	9.5
2000.....	8.2	*	4.9	4.0	2.9	10.1	13.5	10.2	8.6	7.9
1999.....	8.7	8.7	5.8	4.2	3.5	11.0	12.6	9.0	8.1	10.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pedestrian involved in collision with motor vehicle (V02-V04)

All races 3/, both sexes

2004.....	1.1	*	1.0	0.5	0.5	0.8	1.0	1.0	0.9	1.1
2003.....	1.2	*	1.1	0.5	0.5	1.0	1.1	1.1	0.9	1.2
2002.....	1.3	*	1.0	0.6	0.5	1.0	1.1	1.0	1.0	1.3
2001.....	1.2	*	1.1	0.6	0.7	0.9	1.1	1.0	1.0	1.4
2000.....	1.2	*	1.2	0.7	0.6	0.8	1.0	1.0	1.0	1.2
1999.....	1.2	*	1.1	0.8	0.5	0.8	1.1	1.0	1.0	1.3

All races 3/, male

2004.....	1.6	*	1.2	0.6	0.6	1.1	1.5	1.4	1.4	1.6
2003.....	1.7	*	1.4	0.7	0.7	1.3	1.6	1.6	1.4	1.7
2002.....	1.7	*	1.3	0.7	0.6	1.4	1.7	1.6	1.6	1.8
2001.....	1.8	*	1.4	0.8	1.0	1.2	1.8	1.6	1.5	1.9
2000.....	1.7	*	1.4	0.8	0.7	1.0	1.5	1.5	1.5	1.7
1999.....	1.7	*	1.2	0.9	0.7	1.2	1.6	1.5	1.5	1.9

All races 3/, female

2004.....	0.7	*	0.8	0.4	0.3	0.4	0.5	0.5	0.5	0.7
2003.....	0.7	*	0.8	0.4	0.4	0.6	0.6	0.5	0.4	0.6
2002.....	0.8	*	0.6	0.5	0.4	0.6	0.5	0.5	0.5	0.8
2001.....	0.8	*	0.8	0.4	0.4	0.5	0.5	0.3	0.5	0.8
2000.....	0.8	*	0.9	0.6	0.4	0.6	0.4	0.4	0.5	0.7
1999.....	0.7	*	1.0	0.7	0.4	0.5	0.5	0.5	0.4	0.6

White, both sexes

2004.....	1.1	*	1.0	0.5	0.5	0.8	1.0	1.0	0.8	1.1
2003.....	1.2	*	1.1	0.4	0.5	1.0	1.1	1.0	0.9	1.1
2002.....	1.2	*	0.9	0.5	0.5	0.9	1.1	1.0	1.0	1.1
2001.....	1.2	*	1.0	0.5	0.7	0.8	1.2	1.0	1.0	1.2
2000.....	1.1	*	1.0	0.6	0.6	0.8	1.0	0.9	0.9	1.1
1999.....	1.1	*	1.0	0.6	0.5	0.8	1.0	0.9	0.9	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pedestrian involved in collision with motor vehicle (V02-V04)

White, male										
2004.....	1.5	*	1.2	0.5	0.6	1.2	1.5	1.5	1.3	1.5
2003.....	1.6	*	1.4	0.6	0.6	1.4	1.7	1.7	1.3	1.6
2002.....	1.6	*	1.2	0.7	0.6	1.3	1.8	1.5	1.5	1.5
2001.....	1.7	*	1.4	0.6	1.0	1.2	1.8	1.7	1.4	1.8
2000.....	1.6	*	1.2	0.7	0.7	1.0	1.6	1.3	1.4	1.5
1999.....	1.5	*	1.1	0.7	0.7	1.2	1.6	1.3	1.4	1.6
White, female										
2004.....	0.7	*	0.8	0.4	0.3	0.4	0.5	0.4	0.4	0.7
2003.....	0.7	*	0.8	0.3	0.3	0.6	0.5	0.4	0.5	0.6
2002.....	0.7	*	0.6	0.4	0.4	0.5	0.5	0.4	0.5	0.6
2001.....	0.7	*	0.7	0.3	0.4	0.5	0.5	0.3	0.5	0.7
2000.....	0.7	*	0.8	0.5	0.4	0.6	0.5	0.4	0.4	0.6
1999.....	0.7	*	1.0	0.5	0.4	0.5	0.5	0.4	0.4	0.5
Black, both sexes										
2004.....	1.4	*	1.1	0.9	*	*	1.1	1.0	1.2	1.3
2003.....	1.5	*	1.5	1.0	0.8	1.0	1.0	1.1	1.3	1.4
2002.....	1.6	*	1.3	1.0	0.6	1.3	1.0	1.4	1.4	2.4
2001.....	1.7	*	1.5	1.2	0.9	1.1	1.0	1.0	1.2	2.2
2000.....	1.7	*	1.9	1.2	0.7	0.8	0.8	1.3	1.1	2.1
1999.....	1.8	*	1.6	1.6	0.7	0.8	1.1	1.7	1.5	2.4
Black, male										
2004.....	2.1	*	*	*	*	*	1.6	*	2.1	1.9
2003.....	2.3	*	1.9	1.3	*	1.2	1.4	*	2.3	2.5
2002.....	2.4	*	1.7	*	*	1.7	1.5	2.1	2.2	3.6
2001.....	2.4	*	1.7	1.8	1.4	1.3	1.6	1.6	1.7	2.8
2000.....	2.5	*	2.3	1.5	*	*	*	2.1	1.6	3.4
1999.....	2.8	*	2.0	1.8	*	*	1.8	2.6	2.7	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pedestrian involved in collision with motor vehicle (V02-V04)

Black, female

2004.....	0.8	*	*	*	*	*	*	*	*
2003.....	0.8	*	*	*	*	*	*	*	*
2002.....	0.9	*	*	*	*	*	*	*	1.4
2001.....	1.0	*	*	*	*	*	*	*	1.7
2000.....	0.8	*	*	*	*	*	*	*	*
1999.....	0.9	*	*	1.3	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)

All races 3/, both sexes

2004.....	0.2	*	*	0.2	0.3	0.3	0.2	0.1	0.1	0.3
2003.....	0.2	*	*	0.2	0.4	0.3	0.1	0.1	0.1	0.2
2002.....	0.2	*	*	0.2	0.3	0.2	0.1	0.1	0.2	0.2
2001.....	0.2	*	*	0.2	0.3	0.3	0.2	0.2	0.2	0.2
2000.....	0.2	*	*	0.2	0.5	0.2	0.1	0.1	0.2	0.2
1999.....	0.2	*	*	0.3	0.4	0.3	0.2	0.2	0.2	0.2

All races 3/, male

2004.....	0.4	*	*	0.3	0.5	0.4	0.3	0.3	0.2	0.5
2003.....	0.4	*	*	0.2	0.6	0.4	0.3	0.2	*	0.4
2002.....	0.4	*	*	0.3	0.6	0.4	0.2	0.2	0.4	0.3
2001.....	0.4	*	*	0.3	0.5	0.5	0.3	0.4	0.3	0.4
2000.....	0.4	*	*	0.4	0.8	0.4	0.3	0.2	0.3	0.3
1999.....	0.4	*	*	0.5	0.6	0.6	0.4	0.3	0.4	0.4

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	0.2	*	*	*	*	*

White, both sexes

2004.....	0.2	*	*	0.2	0.3	0.2	0.2	0.2	0.1	0.3
2003.....	0.2	*	*	0.2	0.4	0.3	0.2	0.2	*	0.2
2002.....	0.2	*	*	0.1	0.3	0.2	0.2	0.2	0.2	0.2
2001.....	0.2	*	*	0.2	0.3	0.3	0.2	0.2	0.2	0.2
2000.....	0.2	*	*	0.2	0.4	0.3	0.2	*	0.2	0.2
1999.....	0.2	*	*	0.3	0.4	0.4	0.2	0.2	0.3	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)

All races 3/, both sexes

2004.....	1.4	*	*	*	0.2	1.1	2.7	2.3	2.1	2.0
2003.....	1.3	*	*	*	0.2	1.0	2.5	2.1	2.0	1.9
2002.....	1.1	*	*	*	0.1	0.8	2.3	2.0	1.8	1.7
2001.....	1.1	*	*	*	0.1	0.8	2.3	2.2	1.9	1.7
2000.....	1.0	*	*	*	0.1	0.8	2.1	2.1	1.6	1.5
1999.....	0.8	*	*	*	0.1	0.6	2.0	1.8	1.3	1.3

All races 3/, male

2004.....	2.5	*	*	*	0.4	2.0	5.0	4.2	3.9	3.5
2003.....	2.3	*	*	*	0.3	1.7	4.4	3.9	3.5	3.4
2002.....	2.1	*	*	*	*	1.5	4.1	3.7	3.3	3.0
2001.....	2.0	*	*	*	0.3	1.5	4.2	4.1	3.4	2.9
2000.....	1.8	*	*	*	0.2	1.5	3.8	3.9	2.9	2.7
1999.....	1.6	*	*	*	0.2	1.1	3.6	3.3	2.4	2.4

All races 3/, female

2004.....	0.3	*	*	*	*	0.2	0.4	0.3	0.3	0.4
2003.....	0.3	*	*	*	*	0.2	0.4	0.3	0.4	0.4
2002.....	0.2	*	*	*	*	*	0.3	0.3	0.3	0.5
2001.....	0.2	*	*	*	*	*	0.3	0.3	0.4	0.5
2000.....	0.2	*	*	*	*	*	*	0.2	0.2	0.4
1999.....	0.2	*	*	*	*	*	0.2	0.2	*	0.2

White, both sexes

2004.....	1.5	*	*	*	0.2	1.2	3.0	2.3	2.1	2.0
2003.....	1.4	*	*	*	0.2	1.1	2.7	2.1	1.9	1.9
2002.....	1.2	*	*	*	*	0.9	2.3	2.0	1.8	1.9
2001.....	1.1	*	*	*	0.2	1.0	2.4	2.2	1.8	1.8
2000.....	1.0	*	*	*	0.1	0.9	2.1	2.0	1.5	1.6
1999.....	0.9	*	*	*	0.2	0.7	2.0	1.6	1.3	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)

White, male

2004.....	2.7	*	*	*	0.4	2.2	5.4	4.1	3.7	3.5
2003.....	2.4	*	*	*	0.3	2.0	4.8	3.7	3.3	3.2
2002.....	2.2	*	*	*	*	1.7	4.2	3.6	3.1	3.1
2001.....	2.1	*	*	*	0.3	1.8	4.4	4.0	3.2	3.0
2000.....	1.9	*	*	*	*	1.6	3.9	3.6	2.6	2.8
1999.....	1.6	*	*	*	0.2	1.2	3.7	2.9	2.4	2.5

White, female

2004.....	0.3	*	*	*	*	*	0.4	0.4	0.3	0.5
2003.....	0.3	*	*	*	*	*	0.5	0.3	0.4	0.4
2002.....	0.2	*	*	*	*	*	0.3	0.3	0.4	0.6
2001.....	0.2	*	*	*	*	*	0.3	0.3	0.4	0.6
2000.....	0.2	*	*	*	*	*	*	0.3	0.3	0.4
1999.....	0.2	*	*	*	*	*	*	0.3	*	0.2

Black, both sexes

2004.....	1.1	*	*	*	*	0.8	1.8	3.0	3.2	2.4
2003.....	1.1	*	*	*	*	*	1.7	3.4	3.4	2.7
2002.....	0.9	*	*	*	*	*	2.2	2.8	2.4	1.5
2001.....	0.9	*	*	*	*	*	2.1	2.9	2.7	1.3
2000.....	0.9	*	*	*	*	*	2.2	3.2	2.9	1.5
1999.....	0.8	*	*	*	*	*	2.3	3.1	1.7	1.2

Black, male

2004.....	2.2	*	*	*	*	1.4	3.5	6.0	6.4	4.9
2003.....	2.2	*	*	*	*	*	3.2	7.0	6.7	5.7
2002.....	1.8	*	*	*	*	*	4.3	5.5	5.1	3.2
2001.....	1.7	*	*	*	*	*	4.0	5.9	5.4	2.7
2000.....	1.9	*	*	*	*	*	4.2	6.6	5.8	3.1
1999.....	1.6	*	*	*	*	*	4.5	6.4	3.5	2.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accident involving collision with railway train (V25,V35,V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	0.1	0.1	*	0.1	*
2003.....	0.1	*	*	*	*	0.1	0.1	*	*	*
2002.....	0.1	*	*	*	*	0.1	0.1	*	*	0.1
2001.....	0.1	*	*	*	*	*	0.2	0.1	*	0.1
2000.....	0.1	*	*	*	*	0.2	0.1	0.1	0.1	*
1999.....	0.1	*	*	*	*	0.2	0.2	*	*	*

All races 3/, male

2004.....	0.1	*	*	*	*	0.2	*	*	0.2	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.2	*	*	*
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	*
2000.....	0.1	*	*	*	*	0.2	0.2	*	0.2	*
1999.....	0.1	*	*	*	*	0.2	0.2	*	*	*

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	*	0.2	*	*	0.1	*
2003.....	0.1	*	*	*	*	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	0.1	0.2	*	*	*
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	0.1
2000.....	0.1	*	*	*	*	0.2	0.1	0.1	0.2	*
1999.....	0.1	*	*	*	*	0.2	0.2	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	0.1	0.1	*	0.1	*
2003.....	0.1	*	*	*	*	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	0.1	0.1	*	*	*
2001.....	0.1	*	*	*	*	*	0.2	0.1	*	0.1
2000.....	0.1	*	*	*	*	0.2	0.1	0.1	0.1	*
1999.....	0.1	*	*	*	*	0.2	0.2	*	*	*

All races 3/, male

2004.....	0.1	*	*	*	*	*	*	*	0.2	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.2	*	*	*
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	*
2000.....	0.1	*	*	*	*	0.2	*	*	0.2	*
1999.....	0.1	*	*	*	*	0.2	*	*	*	*

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	*	0.2	*	*	0.1	*
2003.....	0.1	*	*	*	*	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	0.1	0.2	*	*	*
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	0.1
2000.....	0.1	*	*	*	*	0.2	0.1	0.1	0.2	*
1999.....	0.1	*	*	*	*	0.2	0.2	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

All races 3/, both sexes

2004.....	1.6	*	0.2	0.2	0.3	3.9	4.4	2.5	1.7	1.6
2003.....	1.7	*	0.2	0.2	0.4	4.0	4.4	2.8	1.9	1.6
2002.....	1.8	*	0.2	0.2	0.4	4.5	4.6	2.8	1.9	1.8
2001.....	1.5	*	0.1	0.2	0.2	3.7	4.0	2.2	1.8	1.5
2000.....	1.4	*	0.2	0.2	0.3	3.4	3.5	2.0	1.5	1.6
1999.....	1.4	*	0.1	0.1	0.3	3.3	3.3	2.0	1.6	1.4

All races 3/, male

2004.....	2.4	*	*	0.2	0.3	5.3	6.6	3.8	2.5	2.6
2003.....	2.6	*	*	0.2	0.6	5.7	6.8	4.3	2.9	2.4
2002.....	2.6	*	*	0.2	0.4	6.3	7.0	4.4	2.9	2.7
2001.....	2.2	*	*	*	0.3	5.2	6.1	3.4	2.7	2.1
2000.....	2.0	*	*	*	0.3	4.8	5.7	3.2	2.2	2.3
1999.....	2.0	*	*	*	0.4	4.4	5.3	3.2	2.5	2.1

All races 3/, female

2004.....	0.8	*	*	*	0.3	2.3	1.9	1.1	0.8	0.7
2003.....	0.9	*	0.3	*	0.3	2.2	1.8	1.3	0.9	0.8
2002.....	0.9	*	*	*	0.3	2.6	1.9	1.1	0.8	0.9
2001.....	0.8	*	*	*	0.2	2.1	1.7	1.0	0.8	0.8
2000.....	0.8	*	*	*	0.3	1.9	1.3	0.8	0.8	1.0
1999.....	0.7	*	*	*	0.2	2.2	1.3	0.8	0.7	0.7

White, both sexes

2004.....	1.6	*	*	0.2	0.3	4.3	4.6	2.5	1.6	1.7
2003.....	1.8	*	0.2	0.2	0.4	4.4	4.8	2.8	2.0	1.7
2002.....	1.8	*	0.2	0.2	0.3	4.9	4.8	2.8	1.9	1.8
2001.....	1.5	*	*	0.1	0.3	3.8	3.9	2.2	1.7	1.5
2000.....	1.4	*	*	0.2	0.3	3.6	3.5	1.9	1.5	1.7
1999.....	1.3	*	*	0.1	0.3	3.6	3.3	2.0	1.6	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

White, male

2004.....	2.4	*	*	*	0.3	6.0	7.1	3.8	2.3	2.7
2003.....	2.6	*	*	0.3	0.6	6.2	7.3	4.2	3.0	2.5
2002.....	2.6	*	*	*	0.3	6.7	7.3	4.4	3.0	2.5
2001.....	2.2	*	*	*	0.3	5.3	6.1	3.4	2.6	2.1
2000.....	2.0	*	*	*	0.3	5.0	5.6	3.0	2.2	2.3
1999.....	2.0	*	*	*	0.4	4.7	5.3	3.1	2.4	2.0

White, female

2004.....	0.9	*	*	*	0.3	2.6	1.9	1.2	0.9	0.7
2003.....	0.9	*	*	*	0.3	2.5	2.0	1.4	0.9	0.9
2002.....	1.0	*	*	*	0.3	2.9	2.0	1.2	0.8	1.0
2001.....	0.8	*	*	*	*	2.3	1.5	1.0	0.8	0.8
2000.....	0.8	*	*	*	0.3	2.1	1.3	0.8	0.9	1.0
1999.....	0.7	*	*	*	*	2.4	1.2	0.8	0.7	0.6

Black, both sexes

2004.....	1.7	*	*	*	*	2.2	3.8	3.0	2.3	1.8
2003.....	1.6	*	*	*	*	2.8	3.4	3.2	1.7	1.3
2002.....	1.9	*	*	*	*	3.3	4.3	3.1	2.1	2.4
2001.....	1.8	*	*	*	*	3.6	4.7	2.6	2.3	1.7
2000.....	1.6	*	*	*	*	2.6	3.7	2.7	1.4	1.7
1999.....	1.6	*	*	*	*	2.8	4.1	2.5	2.1	1.8

Black, male

2004.....	2.5	*	*	*	*	3.1	5.3	4.9	4.1	2.6
2003.....	2.6	*	*	*	*	4.0	5.9	5.7	2.7	2.3
2002.....	3.0	*	*	*	*	5.1	6.7	5.8	3.3	4.0
2001.....	2.8	*	*	*	*	5.3	7.4	4.2	3.7	2.6
2000.....	2.5	*	*	*	*	3.9	5.9	4.8	2.6	2.7
1999.....	2.6	*	*	*	*	4.0	6.4	4.3	3.6	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ year 1/	year 2/ year 1/	years	years	years	years	years	years	years	years

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

Black, female

2004.....	0.9	*	*	*	*	1.2	2.4	*	*	*
2003.....	0.7	*	*	*	*	1.6	*	*	*	*
2002.....	0.9	*	*	*	*	1.5	1.9	*	*	*
2001.....	0.8	*	*	*	*	1.7	2.1	*	*	*
2000.....	0.7	*	*	*	*	*	1.7	*	*	*
1999.....	0.8	*	*	*	*	1.6	1.9	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

All races 3/, both sexes

2004.....	3.6	1.5	0.8	0.8	1.0	6.2	5.7	4.0	3.2	3.0
2003.....	3.7	1.5	0.9	0.8	0.8	6.2	6.0	4.1	3.4	3.2
2002.....	3.9	1.1	1.0	0.8	0.9	6.9	6.8	4.2	3.4	3.4
2001.....	3.7	1.6	1.0	0.8	1.0	6.3	5.9	4.2	3.1	3.3
2000.....	3.7	1.5	0.9	0.9	0.9	6.2	6.0	4.1	3.2	3.3
1999.....	3.7	1.6	0.9	0.9	1.0	6.0	5.7	4.0	3.5	3.0

All races 3/, male

2004.....	4.4	1.9	0.8	0.8	1.1	7.3	7.5	5.4	4.1	4.0
2003.....	4.5	1.4	0.9	0.7	0.9	7.0	7.9	5.4	4.5	4.1
2002.....	4.8	1.5	1.0	0.9	0.9	8.3	9.1	6.0	4.3	4.2
2001.....	4.5	1.6	0.9	0.8	1.0	7.6	8.1	5.7	4.3	4.2
2000.....	4.5	1.5	0.9	0.9	0.9	7.1	8.0	5.3	4.3	4.1
1999.....	4.4	1.7	1.0	0.9	1.0	7.1	7.5	5.2	4.6	3.5

All races 3/, female

2004.....	2.9	1.1	0.9	0.8	1.0	5.0	3.9	2.6	2.2	2.1
2003.....	2.9	1.5	0.9	0.9	0.7	5.3	4.1	2.7	2.3	2.3
2002.....	3.0	*	0.9	0.8	0.9	5.3	4.4	2.4	2.4	2.5
2001.....	2.9	1.6	1.1	0.9	1.0	4.9	3.6	2.7	2.0	2.4
2000.....	2.9	1.5	0.9	0.9	0.8	5.2	3.8	3.0	2.2	2.4
1999.....	2.9	1.4	0.7	0.9	1.1	4.8	3.8	2.7	2.4	2.4

White, both sexes

2004.....	3.8	1.5	0.8	0.8	1.1	6.8	6.2	4.1	3.2	3.1
2003.....	3.9	1.4	0.9	0.8	0.8	6.9	6.3	4.3	3.6	3.2
2002.....	4.1	1.0	0.9	0.8	1.0	7.6	7.2	4.2	3.4	3.4
2001.....	3.8	1.5	1.0	0.8	1.1	6.8	6.3	4.4	3.2	3.3
2000.....	3.8	1.5	0.9	0.9	0.9	6.8	6.2	4.2	3.2	3.3
1999.....	3.8	1.5	0.8	0.9	1.1	6.6	5.9	4.1	3.5	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

White, male

2004.....	4.5	1.9	0.8	0.9	1.2	7.9	8.2	5.5	3.9	4.0
2003.....	4.6	1.5	0.9	0.7	1.0	7.8	8.2	5.7	4.7	4.2
2002.....	5.0	1.4	1.0	0.9	1.0	9.0	9.7	5.9	4.3	4.2
2001.....	4.7	1.6	0.9	0.7	1.1	8.1	8.5	5.9	4.3	4.1
2000.....	4.6	1.5	0.9	1.0	0.9	7.8	8.1	5.3	4.2	4.1
1999.....	4.5	1.8	0.9	0.9	1.0	7.6	7.7	5.4	4.6	3.2

White, female

2004.....	3.0	*	0.8	0.8	1.0	5.7	4.1	2.6	2.3	2.1
2003.....	3.1	1.4	0.9	0.9	0.7	6.0	4.3	2.9	2.4	2.2
2002.....	3.2	*	0.9	0.7	1.0	6.1	4.6	2.3	2.4	2.6
2001.....	3.0	1.5	1.1	0.9	1.0	5.3	3.9	2.9	2.1	2.4
2000.....	3.1	1.5	0.9	0.9	0.9	5.7	4.1	2.9	2.3	2.4
1999.....	3.1	*	0.7	0.9	1.2	5.4	3.9	2.7	2.4	2.5

Black, both sexes

2004.....	3.1	*	0.9	0.8	0.8	3.7	4.2	4.5	3.9	3.1
2003.....	3.2	*	0.9	0.8	0.7	3.5	5.2	3.8	3.4	3.3
2002.....	3.3	*	1.2	0.8	0.7	4.1	5.5	5.5	4.1	3.4
2001.....	3.3	*	1.1	0.9	0.8	4.4	4.9	4.3	3.5	3.7
2000.....	3.3	*	1.0	0.7	0.8	3.7	5.5	4.7	3.7	3.3
1999.....	3.4	*	1.1	0.9	1.0	3.6	5.1	4.2	4.0	4.2

Black, male

2004.....	4.0	*	*	*	*	5.0	5.1	6.3	6.2	4.2
2003.....	4.1	*	*	*	*	4.6	7.2	5.2	4.8	3.7
2002.....	4.4	*	1.6	*	*	5.7	7.0	7.8	5.7	4.5
2001.....	4.4	*	*	*	*	5.6	6.9	5.9	5.2	4.8
2000.....	4.4	*	*	*	*	4.5	8.0	6.0	5.7	4.0
1999.....	4.5	*	*	*	*	5.0	6.9	5.7	5.5	5.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

Black, female

2004.....	2.2	*	*	*	*	2.3	3.3	2.8	1.7	2.1
2003.....	2.4	*	*	*	*	2.3	3.2	2.6	2.2	3.1
2002.....	2.4	*	*	*	*	2.4	4.1	3.4	2.6	2.5
2001.....	2.3	*	*	*	*	3.2	3.1	2.8	2.1	2.8
2000.....	2.2	*	*	*	*	2.9	3.2	3.5	1.9	2.6
1999.....	2.3	*	*	*	*	2.2	3.4	2.9	2.7	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

All races 3/, both sexes

2004.....	1.4	*	0.3	0.4	0.4	2.9	3.2	1.9	1.5	1.6
2003.....	1.5	*	0.3	0.3	0.5	3.4	3.2	2.2	1.6	1.6
2002.....	1.5	*	0.3	0.3	0.5	3.7	3.4	2.3	1.8	1.7
2001.....	1.4	*	0.2	0.2	0.5	2.8	3.3	1.9	1.7	1.6
2000.....	1.3	0.6	0.2	0.2	0.5	2.9	3.2	2.2	1.5	1.5
1999.....	1.3	*	0.3	0.3	0.3	3.0	3.1	2.0	1.5	1.5

All races 3/, male

2004.....	1.9	*	0.3	0.4	0.4	3.6	4.7	2.9	2.1	2.3
2003.....	2.1	*	0.3	0.3	0.6	4.3	4.8	3.2	2.4	2.2
2002.....	2.2	*	0.4	0.3	0.5	4.6	5.0	3.4	2.6	2.4
2001.....	2.0	*	*	0.3	0.5	3.4	5.0	2.8	2.6	2.3
2000.....	1.9	*	*	0.2	0.6	3.9	4.7	3.4	2.2	2.1
1999.....	1.9	*	0.3	0.3	0.3	3.9	4.6	3.0	2.1	2.1

All races 3/, female

2004.....	0.8	*	0.3	0.3	0.4	2.3	1.6	0.9	0.9	0.8
2003.....	0.9	*	0.4	0.3	0.4	2.5	1.6	1.1	0.8	1.0
2002.....	0.9	*	*	0.3	0.4	2.7	1.7	1.1	1.0	1.1
2001.....	0.8	*	*	0.2	0.5	2.1	1.5	1.0	0.8	0.9
2000.....	0.7	*	*	0.3	0.4	1.9	1.6	0.9	0.8	0.9
1999.....	0.8	*	0.3	0.3	0.3	2.2	1.5	0.9	0.8	0.9

White, both sexes

2004.....	1.4	*	0.2	0.4	0.4	3.3	3.3	2.0	1.6	1.6
2003.....	1.5	*	0.3	0.3	0.5	3.8	3.5	2.3	1.7	1.6
2002.....	1.6	*	0.3	0.3	0.5	4.0	3.6	2.5	1.9	1.8
2001.....	1.4	*	0.2	0.2	0.5	3.1	3.5	1.9	1.7	1.7
2000.....	1.4	*	0.2	0.2	0.4	3.2	3.4	2.3	1.6	1.5
1999.....	1.3	*	0.3	0.3	0.3	3.3	3.3	2.0	1.5	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

White, male

2004.....	2.0	*	*	0.4	0.4	4.0	5.0	3.0	2.2	2.4
2003.....	2.1	*	*	*	0.5	4.8	5.1	3.5	2.5	2.1
2002.....	2.2	*	0.3	0.3	0.6	5.1	5.3	3.5	2.7	2.4
2001.....	2.0	*	*	*	0.5	3.7	5.5	2.8	2.6	2.4
2000.....	2.0	*	*	*	0.5	4.3	5.0	3.5	2.4	2.2
1999.....	1.9	*	*	0.3	0.3	4.2	4.9	3.1	2.2	2.1

White, female

2004.....	0.8	*	*	0.3	0.4	2.6	1.5	0.9	0.8	0.8
2003.....	0.9	*	0.3	0.4	0.4	2.8	1.7	1.1	0.7	1.2
2002.....	0.9	*	*	0.3	0.4	2.8	1.7	1.3	1.1	1.1
2001.....	0.8	*	*	*	0.5	2.4	1.5	0.9	0.8	1.0
2000.....	0.7	*	*	0.3	0.3	2.1	1.6	1.0	0.7	0.9
1999.....	0.8	*	*	0.3	0.3	2.4	1.5	1.0	0.8	0.9

Black, both sexes

2004.....	1.1	*	*	*	*	1.4	2.2	1.5	1.7	1.3
2003.....	1.2	*	*	*	*	1.2	2.1	1.6	1.8	1.5
2002.....	1.2	*	*	*	*	1.8	2.2	1.7	1.2	1.4
2001.....	1.1	*	*	*	*	1.3	2.0	1.8	1.7	1.0
2000.....	1.1	*	*	*	0.6	1.4	2.5	1.6	1.1	1.3
1999.....	1.2	*	*	*	*	1.7	2.2	1.6	1.4	1.6

Black, male

2004.....	1.7	*	*	*	*	2.0	3.0	2.4	2.0	1.9
2003.....	1.7	*	*	*	*	1.7	3.1	2.0	2.3	2.6
2002.....	1.8	*	*	*	*	2.2	3.2	3.0	2.2	1.8
2001.....	1.6	*	*	*	*	1.8	2.4	2.5	3.2	1.5
2000.....	1.6	*	*	*	*	1.8	3.5	3.0	1.7	2.3
1999.....	1.7	*	*	*	*	2.2	2.9	2.8	2.0	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ year 1/	year 2/ year 1/	years	years	years	years	years	years	years	years

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

Black, female

2004.....	0.6	*	*	*	*	*	1.3	*	1.4	*
2003.....	0.7	*	*	*	*	*	*	*	*	*
2002.....	0.6	*	*	*	*	1.4	*	*	*	*
2001.....	0.6	*	*	*	*	*	1.6	*	*	*
2000.....	0.6	*	*	*	*	*	1.5	*	*	*
1999.....	0.7	*	*	*	*	*	1.5	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of special-use motor vehicle involved in any accident (V83-V86)

All races 3/, both sexes

2004.....	0.4	*	*	0.2	0.6	0.7	0.6	0.5	0.4	0.4
2003.....	0.4	*	*	0.2	0.5	0.6	0.6	0.4	0.4	0.3
2002.....	0.3	*	*	0.2	0.5	0.6	0.5	0.4	0.3	0.4
2001.....	0.4	*	*	0.2	0.4	0.6	0.5	0.4	0.4	0.3
2000.....	0.4	*	*	0.2	0.4	0.7	0.5	0.4	0.3	0.3
1999.....	0.4	*	*	0.2	0.4	0.4	0.5	0.4	0.4	0.3

All races 3/, male

2004.....	0.7	*	*	0.3	0.9	1.0	1.0	0.9	0.6	0.7
2003.....	0.7	*	*	0.4	0.7	1.0	1.0	0.7	0.7	0.6
2002.....	0.6	*	*	0.3	0.7	0.9	0.8	0.7	0.5	0.7
2001.....	0.7	*	*	0.3	0.7	1.1	0.8	0.7	0.7	0.6
2000.....	0.7	*	*	0.3	0.6	1.1	0.9	0.7	0.6	0.6
1999.....	0.7	*	*	0.3	0.6	0.7	0.8	0.6	0.7	0.6

All races 3/, female

2004.....	0.1	*	*	*	0.3	0.3	*	*	*	*
2003.....	0.1	*	*	*	0.3	*	*	*	*	*
2002.....	0.1	*	*	*	0.2	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	0.2	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.4	*	*	0.2	0.7	0.8	0.6	0.6	0.5	0.4
2003.....	0.4	*	*	0.3	0.6	0.6	0.7	0.5	0.5	0.4
2002.....	0.4	*	*	0.2	0.6	0.7	0.6	0.5	0.4	0.5
2001.....	0.4	*	*	0.3	0.5	0.8	0.5	0.5	0.5	0.4
2000.....	0.4	*	*	0.3	0.5	0.7	0.6	0.4	0.4	0.4
1999.....	0.4	*	*	0.2	0.4	0.5	0.6	0.4	0.5	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

All races 3/, both sexes

2004.....	5.5	1.2	1.6	1.0	1.4	9.3	9.6	7.2	5.5	5.0
2003.....	5.3	1.3	1.3	1.1	1.4	9.2	9.4	6.0	4.9	5.2
2002.....	5.5	1.1	1.3	1.0	1.3	9.9	9.9	6.6	5.5	5.1
2001.....	5.8	1.3	1.5	1.2	1.3	10.2	10.6	7.2	5.4	5.8
2000.....	6.0	1.8	1.6	1.3	1.6	10.7	11.4	7.2	5.9	5.8
1999.....	6.0	2.3	1.6	1.3	1.7	11.1	10.5	7.0	5.7	6.0

All races 3/, male

2004.....	7.3	1.2	1.6	1.0	1.4	11.6	13.6	10.2	7.3	6.7
2003.....	7.0	1.6	1.5	1.2	1.5	11.6	13.3	8.6	7.1	7.0
2002.....	7.3	1.1	1.4	1.1	1.4	12.4	14.3	9.3	7.7	7.1
2001.....	7.7	1.1	1.6	1.3	1.4	13.3	15.5	10.4	7.5	7.7
2000.....	8.0	1.9	1.8	1.3	1.7	13.7	16.0	10.2	8.2	8.0
1999.....	7.9	2.1	1.8	1.4	1.8	13.8	15.0	9.9	7.6	8.0

All races 3/, female

2004.....	3.7	1.3	1.5	1.0	1.5	6.8	5.3	4.0	3.6	3.4
2003.....	3.6	*	1.1	0.9	1.2	6.6	5.2	3.3	2.6	3.4
2002.....	3.7	1.1	1.3	0.9	1.1	7.3	5.4	3.8	3.1	3.2
2001.....	3.9	1.5	1.3	1.1	1.1	6.9	5.5	4.0	3.4	4.0
2000.....	4.1	1.7	1.4	1.3	1.4	7.5	6.6	4.0	3.6	3.6
1999.....	4.2	2.5	1.5	1.2	1.5	8.2	5.8	4.1	3.8	4.0

White, both sexes

2004.....	5.6	1.1	1.5	0.9	1.3	10.0	10.1	7.2	5.6	5.0
2003.....	5.3	1.2	1.2	1.1	1.4	9.8	9.7	6.1	4.9	5.2
2002.....	5.6	0.8	1.3	0.9	1.3	10.9	10.3	6.6	5.6	5.1
2001.....	5.9	1.1	1.4	1.1	1.2	11.0	10.9	7.4	5.5	5.8
2000.....	6.2	1.5	1.6	1.3	1.6	11.8	11.9	7.1	5.8	5.9
1999.....	6.2	2.2	1.4	1.2	1.7	12.0	11.0	7.3	5.7	6.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

White, male

2004.....	7.3	*	1.5	0.9	1.2	12.4	14.4	10.1	7.4	6.5
2003.....	7.1	1.6	1.4	1.2	1.5	12.3	13.6	8.9	7.0	6.9
2002.....	7.4	*	1.3	0.9	1.4	13.5	14.7	9.2	7.8	7.0
2001.....	7.8	*	1.5	1.2	1.2	14.2	15.9	10.6	7.6	7.4
2000.....	8.2	1.6	1.8	1.2	1.8	15.0	16.8	10.0	8.2	8.1
1999.....	8.0	2.2	1.4	1.3	1.8	14.8	15.6	10.3	7.5	8.1

White, female

2004.....	3.9	1.4	1.5	0.9	1.4	7.4	5.5	4.1	3.8	3.4
2003.....	3.6	*	1.0	0.9	1.3	7.2	5.5	3.2	2.6	3.5
2002.....	3.9	*	1.3	0.9	1.1	8.0	5.6	3.8	3.2	3.1
2001.....	4.0	1.3	1.3	1.0	1.1	7.6	5.5	4.1	3.3	4.2
2000.....	4.2	1.4	1.4	1.3	1.5	8.4	6.6	4.0	3.4	3.7
1999.....	4.3	2.3	1.4	1.2	1.5	9.0	6.0	4.1	3.9	4.1

Black, both sexes

2004.....	5.4	*	2.1	1.3	2.0	6.7	8.1	7.6	5.7	6.1
2003.....	5.2	*	1.6	1.0	1.3	6.0	8.6	6.1	5.5	5.8
2002.....	5.1	*	1.5	1.4	1.2	6.3	8.7	7.7	5.6	5.9
2001.....	5.6	*	1.7	1.8	1.7	6.7	10.0	7.4	5.9	6.4
2000.....	5.9	3.6	1.9	1.5	1.3	7.0	10.8	8.7	7.4	6.0
1999.....	5.8	*	2.6	1.7	1.6	7.9	9.4	6.9	6.0	6.4

Black, male

2004.....	7.8	*	2.3	1.2	2.3	8.6	12.0	11.7	8.3	9.3
2003.....	7.4	*	1.6	*	1.3	8.3	13.4	9.1	8.4	8.6
2002.....	7.3	*	1.8	1.9	1.4	7.9	13.1	11.7	8.3	8.8
2001.....	8.0	*	1.8	2.2	2.2	9.3	14.9	11.3	8.2	9.7
2000.....	8.5	*	2.0	1.4	1.5	9.4	15.3	13.0	10.2	9.2
1999.....	8.2	*	3.1	2.0	1.9	10.5	14.1	10.2	9.0	8.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

Black, female

2004.....	3.2	*	1.9	1.3	1.7	4.8	4.1	3.9	3.4	3.2
2003.....	3.2	*	*	*	1.2	3.6	3.9	3.4	2.9	3.4
2002.....	3.1	*	*	*	*	4.7	4.4	4.0	3.3	3.4
2001.....	3.4	*	*	1.3	1.2	3.9	5.4	3.8	3.8	3.5
2000.....	3.6	*	1.8	1.6	*	4.5	6.6	4.8	5.0	3.2
1999.....	3.7	*	2.1	1.4	1.4	5.4	4.9	3.9	3.2	4.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2004.....	0.3	*	*	0.1	0.1	0.3	0.3	0.2	0.2	0.2
2003.....	0.3	*	*	0.1	0.1	0.5	0.5	0.3	0.3	0.2
2002.....	0.2	*	*	*	0.2	0.1	0.2	0.1	0.2	0.2
2001.....	0.3	*	0.2	0.1	0.2	0.3	0.3	0.2	0.2	0.4
2000.....	0.4	*	0.2	0.2	0.2	0.4	0.4	0.3	0.4	0.3
1999.....	0.6	*	0.3	0.3	0.2	0.7	0.7	0.5	0.6	0.6

All races 3/, male

2004.....	0.4	*	*	*	*	0.4	0.4	0.4	0.3	0.4
2003.....	0.5	*	*	*	*	0.6	0.7	0.5	0.4	0.3
2002.....	0.3	*	*	*	0.2	0.2	0.4	*	*	0.4
2001.....	0.5	*	*	*	0.2	0.4	0.5	0.3	0.4	0.6
2000.....	0.5	*	*	*	0.3	0.5	0.6	0.5	0.5	0.5
1999.....	0.8	*	0.3	0.3	0.3	0.9	1.1	0.9	0.7	0.8

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	0.3	0.2	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	0.2	*	*	0.2	*
1999.....	0.3	*	0.3	*	*	0.5	0.3	*	0.4	0.3

White, both sexes

2004.....	0.3	*	*	*	*	0.3	0.3	0.2	0.2	0.2
2003.....	0.3	*	*	*	0.1	0.5	0.5	0.4	0.3	0.2
2002.....	0.2	*	*	*	0.2	0.1	0.2	*	0.2	0.2
2001.....	0.3	*	0.2	*	0.2	0.3	0.3	0.2	0.2	0.3
2000.....	0.4	*	*	0.2	0.2	0.4	0.4	0.3	0.4	0.3
1999.....	0.6	*	0.3	0.2	0.3	0.7	0.8	0.6	0.6	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Water transport accidents (V90-V94)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	0.2	0.2	0.3	0.3	0.2
2003.....	0.2	*	*	*	0.1	0.2	0.3	0.2	0.2	0.3
2002.....	0.2	*	*	*	*	0.2	0.3	0.2	0.3	0.3
2001.....	0.2	*	*	*	*	0.3	0.3	0.3	0.2	0.2
2000.....	0.2	*	*	*	*	0.3	0.3	0.3	0.3	0.3
1999.....	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.3

All races 3/, male

2004.....	0.3	*	*	*	*	0.3	0.4	0.4	0.4	0.4
2003.....	0.4	*	*	*	*	0.4	0.5	0.3	0.4	0.5
2002.....	0.4	*	*	*	*	0.3	0.4	0.4	0.6	0.6
2001.....	0.4	*	*	*	*	0.5	0.5	0.5	0.4	0.4
2000.....	0.4	*	*	*	*	0.5	0.5	0.6	0.4	0.6
1999.....	0.4	*	*	*	0.2	0.4	0.6	0.5	0.5	0.5

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	*	*	*	*	0.2	0.3	0.3	0.3	0.2
2003.....	0.2	*	*	*	*	0.2	0.3	0.2	0.2	0.3
2002.....	0.2	*	*	*	*	0.2	0.3	0.2	0.4	0.4
2001.....	0.2	*	*	*	*	0.3	0.3	0.3	0.2	0.2
2000.....	0.2	*	*	*	*	0.3	0.3	0.3	0.3	0.3
1999.....	0.2	*	*	*	0.1	0.3	0.4	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Air and space transport accidents (V95-V97)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	0.2	0.2	0.3	0.3
2003.....	0.3	*	*	*	*	*	0.2	0.2	0.2	0.3
2002.....	0.2	*	*	*	*	*	0.2	0.1	0.2	0.3
2001.....	0.3	*	*	*	*	*	0.2	0.4	0.4	0.4
2000.....	0.3	*	*	*	*	0.1	0.3	0.3	0.3	0.3
1999.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.3

All races 3/, male

2004.....	0.4	*	*	*	*	*	0.2	0.4	0.5	0.5
2003.....	0.4	*	*	*	*	*	0.3	0.3	0.3	0.5
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.3	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.5	0.7	0.7
2000.....	0.5	*	*	*	*	*	0.5	0.5	0.6	0.6
1999.....	0.4	*	*	*	*	*	0.3	0.5	0.5	0.5

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	0.2	*	0.2
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.4
2003.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.4
2002.....	0.3	*	*	*	*	*	0.2	0.2	0.2	0.3
2001.....	0.4	*	*	*	*	*	0.3	0.4	0.5	0.5
2000.....	0.3	*	*	*	*	*	0.3	0.4	0.4	0.4
1999.....	0.3	*	*	*	*	*	0.2	0.4	0.4	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified transport accidents and their sequelae (V98-V99,Y85)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	0.2	0.2	0.2	0.3
2003.....	0.2	*	*	*	*	*	0.2	0.2	0.2	0.3
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	0.2	*	*	*	*	0.1	0.1	0.2	0.2	0.2
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.3
1999.....	0.2	*	*	*	*	0.1	0.1	0.2	0.2	0.3

All races 3/, male

2004.....	0.3	*	*	*	*	*	0.2	0.4	0.4	0.4
2003.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.4
2002.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.4
2001.....	0.3	*	*	*	*	*	0.2	0.2	0.4	0.3
2000.....	0.3	*	*	*	*	*	*	0.3	0.3	0.4
1999.....	0.3	*	*	*	*	*	*	0.3	0.3	0.4

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	*	*	*	*	*	0.2	0.2	0.2	0.3
2003.....	0.2	*	*	*	*	*	0.2	0.2	0.2	0.3
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	0.2	*	*	*	*	*	*	0.2	0.2	0.2
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.3
1999.....	0.2	*	*	*	*	*	*	0.2	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Falls (W00-W19)

All races 3/, both sexes

2004.....	6.4	0.6	0.3	*	0.1	0.4	0.7	0.8	0.8	1.1
2003.....	5.9	*	0.3	*	0.1	0.4	0.7	0.7	0.7	1.2
2002.....	5.6	*	0.2	*	0.1	0.4	0.8	0.7	0.8	1.2
2001.....	5.3	0.6	0.2	0.2	0.2	0.4	0.9	0.8	0.9	1.2
2000.....	4.7	*	0.2	*	0.1	0.5	0.7	0.7	0.8	1.0
1999.....	4.7	*	0.4	0.1	0.1	0.6	0.7	0.8	0.9	1.2

All races 3/, male

2004.....	6.8	*	0.4	*	*	0.7	1.3	1.3	1.4	1.8
2003.....	6.2	*	0.5	*	0.2	0.7	1.1	1.3	1.1	2.0
2002.....	6.0	*	0.4	*	*	0.6	1.4	1.2	1.4	1.7
2001.....	5.8	*	0.3	0.2	0.2	0.8	1.5	1.5	1.5	2.1
2000.....	5.2	*	0.3	*	*	0.8	1.2	1.2	1.5	1.8
1999.....	5.2	*	0.5	*	0.2	0.9	1.2	1.4	1.5	2.0

All races 3/, female

2004.....	6.0	*	*	*	*	*	*	0.2	0.2	0.5
2003.....	5.6	*	*	*	*	*	0.2	*	0.3	0.3
2002.....	5.3	*	*	*	*	*	*	*	0.3	0.6
2001.....	4.8	*	*	*	*	*	*	*	0.3	0.4
2000.....	4.3	*	*	*	*	*	0.2	*	*	0.3
1999.....	4.3	*	*	*	*	*	*	*	0.3	0.4

White, both sexes

2004.....	7.3	*	0.3	*	0.1	0.5	0.8	0.8	0.9	1.2
2003.....	6.7	*	0.3	*	0.2	0.5	0.8	0.8	0.8	1.2
2002.....	6.4	*	0.2	*	*	0.4	0.9	0.8	0.9	1.2
2001.....	6.0	*	0.2	0.2	0.2	0.5	1.0	0.9	0.9	1.3
2000.....	5.3	*	0.2	*	*	0.6	0.8	0.8	0.9	1.0
1999.....	5.2	*	0.4	*	0.2	0.6	0.8	0.9	0.9	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Falls (W00-W19)

White, male

2004.....	7.6	*	0.4	*	*	0.8	1.4	1.4	1.5	2.0
2003.....	6.9	*	0.5	*	*	0.8	1.3	1.4	1.2	2.1
2002.....	6.6	*	*	*	*	0.7	1.6	1.4	1.5	1.8
2001.....	6.4	*	0.3	0.2	0.2	0.9	1.8	1.6	1.6	2.1
2000.....	5.7	*	*	*	*	1.0	1.4	1.4	1.5	1.7
1999.....	5.6	*	0.5	*	*	1.1	1.3	1.5	1.5	2.0

White, female

2004.....	7.0	*	*	*	*	*	*	*	0.3	0.4
2003.....	6.5	*	*	*	*	*	0.3	*	0.3	0.3
2002.....	6.2	*	*	*	*	*	*	*	0.3	0.7
2001.....	5.5	*	*	*	*	*	*	*	0.3	0.4
2000.....	5.0	*	*	*	*	*	0.3	*	*	0.3
1999.....	4.9	*	*	*	*	*	*	*	0.3	0.5

Black, both sexes

2004.....	2.6	*	*	*	*	*	*	*	0.7	1.1
2003.....	2.4	*	*	*	*	*	*	*	*	1.1
2002.....	2.2	*	*	*	*	*	*	*	*	1.1
2001.....	2.3	*	*	*	*	*	*	*	*	1.1
2000.....	2.1	*	*	*	*	*	*	*	*	1.0
1999.....	2.4	*	*	*	*	*	*	*	1.1	0.9

Black, male

2004.....	3.4	*	*	*	*	*	*	*	*	1.5
2003.....	3.0	*	*	*	*	*	*	*	*	1.7
2002.....	3.0	*	*	*	*	*	*	*	*	1.8
2001.....	3.1	*	*	*	*	*	*	*	*	2.0
2000.....	2.8	*	*	*	*	*	*	*	*	1.7
1999.....	3.3	*	*	*	*	*	*	*	1.8	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Fall from one level to another (W10-W17)

All races 3/, both sexes

2004.....	1.1	*	0.1	*	*	0.2	0.4	0.4	0.4	0.6
2003.....	1.1	*	0.2	*	*	0.3	0.5	0.5	0.4	0.6
2002.....	1.2	*	0.1	*	*	0.3	0.6	0.5	0.5	0.7
2001.....	1.2	*	0.1	*	*	0.3	0.6	0.6	0.5	0.7
2000.....	1.0	*	0.2	*	*	0.4	0.5	0.5	0.6	0.6
1999.....	1.1	*	0.2	*	*	0.4	0.4	0.6	0.6	0.6

All races 3/, male

2004.....	1.7	*	*	*	*	0.4	0.8	0.7	0.8	1.0
2003.....	1.6	*	0.3	*	*	0.5	0.9	0.9	0.7	1.1
2002.....	1.7	*	*	*	*	0.4	1.0	0.9	0.9	1.1
2001.....	1.7	*	*	*	*	0.5	1.1	1.1	0.9	1.2
2000.....	1.6	*	*	*	*	0.6	0.9	0.9	1.1	1.0
1999.....	1.7	*	0.3	*	*	0.6	0.8	1.0	1.0	1.1

All races 3/, female

2004.....	0.6	*	*	*	*	*	*	*	*	*
2003.....	0.7	*	*	*	*	*	*	*	*	*
2002.....	0.6	*	*	*	*	*	*	*	*	0.2
2001.....	0.6	*	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	1.3	*	*	*	*	0.2	0.5	0.4	0.5	0.6
2003.....	1.2	*	0.2	*	*	0.3	0.6	0.6	0.4	0.7
2002.....	1.3	*	*	*	*	0.3	0.7	0.6	0.6	0.7
2001.....	1.3	*	*	*	*	0.4	0.7	0.7	0.6	0.7
2000.....	1.1	*	*	*	*	0.4	0.6	0.6	0.7	0.6
1999.....	1.2	*	0.2	*	*	0.4	0.5	0.6	0.6	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Unspecified fall (W19)

All races 3/, both sexes

2004.....	3.2	*	*	*	*	0.1	0.2	0.2	0.2	0.3
2003.....	2.9	*	*	*	*	*	*	*	0.2	0.3
2002.....	2.7	*	*	*	*	*	*	*	0.1	0.3
2001.....	2.7	*	*	*	*	*	0.1	0.1	0.2	0.3
2000.....	2.5	*	*	*	*	*	*	0.2	0.1	0.3
1999.....	2.8	*	*	*	*	0.1	0.2	0.1	0.3	0.5

All races 3/, male

2004.....	3.1	*	*	*	*	*	0.3	0.3	0.3	0.5
2003.....	2.7	*	*	*	*	*	*	*	0.2	0.6
2002.....	2.5	*	*	*	*	*	*	*	0.3	0.4
2001.....	2.6	*	*	*	*	*	*	*	0.3	0.5
2000.....	2.5	*	*	*	*	*	*	0.3	0.2	0.5
1999.....	2.8	*	*	*	*	*	0.3	0.2	0.4	0.7

All races 3/, female

2004.....	3.2	*	*	*	*	*	*	*	*	*
2003.....	3.0	*	*	*	*	*	*	*	*	*
2002.....	2.8	*	*	*	*	*	*	*	*	0.2
2001.....	2.7	*	*	*	*	*	*	*	*	*
2000.....	2.6	*	*	*	*	*	*	*	*	*
1999.....	2.8	*	*	*	*	*	*	*	*	0.2

White, both sexes

2004.....	3.6	*	*	*	*	0.1	0.2	0.2	0.2	0.4
2003.....	3.2	*	*	*	*	*	*	*	0.2	0.3
2002.....	3.0	*	*	*	*	*	*	*	*	0.3
2001.....	3.1	*	*	*	*	*	*	*	0.2	0.3
2000.....	2.9	*	*	*	*	*	*	0.2	0.1	0.3
1999.....	3.1	*	*	*	*	*	0.2	0.2	0.3	0.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental discharge of firearms (W32-W34)

All races 3/, both sexes

2004.....	0.2	*	*	*	0.2	0.4	0.4	0.3	0.2	0.2
2003.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.2	0.3
2002.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.2
2001.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.2	0.3
2000.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.3
1999.....	0.3	*	*	*	0.3	0.6	0.7	0.4	0.3	0.3

All races 3/, male

2004.....	0.4	*	*	*	0.3	0.7	0.8	0.5	0.4	0.4
2003.....	0.5	*	*	*	0.3	0.8	0.9	0.7	0.4	0.5
2002.....	0.5	*	*	*	0.3	1.0	0.9	0.7	0.6	0.4
2001.....	0.5	*	*	*	0.3	1.0	0.9	0.7	0.4	0.5
2000.....	0.5	*	*	*	0.4	0.9	0.9	0.7	0.4	0.5
1999.....	0.5	*	*	*	0.5	1.1	1.2	0.6	0.6	0.5

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	*	*	*	0.2	0.4	0.4	0.3	0.2	0.2
2003.....	0.2	*	*	*	0.2	0.4	0.5	0.3	0.2	0.3
2002.....	0.3	*	*	*	0.2	0.4	0.5	0.3	0.3	0.2
2001.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.4
2000.....	0.3	*	*	*	0.3	0.5	0.4	0.3	0.3	0.3
1999.....	0.3	*	*	*	0.3	0.6	0.6	0.4	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental discharge of firearms (W32-W34)

White, male

2004.....	0.4	*	*	*	0.3	0.6	0.7	0.4	0.3	0.4
2003.....	0.4	*	*	*	0.3	0.8	0.8	0.6	0.4	0.5
2002.....	0.4	*	*	*	0.3	0.8	0.8	0.6	0.5	0.4
2001.....	0.5	*	*	*	0.4	0.9	0.9	0.6	0.5	0.6
2000.....	0.5	*	*	*	0.5	0.9	0.8	0.6	0.4	0.5
1999.....	0.5	*	*	*	0.5	1.0	1.0	0.6	0.6	0.5

White, female

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

Black, both sexes

2004.....	0.3	*	*	*	*	*	0.9	*	*	*
2003.....	0.3	*	*	*	*	0.7	0.9	1.0	*	*
2002.....	0.4	*	*	*	*	1.1	0.8	0.9	*	*
2001.....	0.3	*	*	*	*	0.8	0.8	*	*	*
2000.....	0.4	*	*	*	*	0.9	0.9	*	*	*
1999.....	0.4	*	*	*	*	1.2	1.1	*	*	*

Black, male

2004.....	0.6	*	*	*	*	*	1.6	*	*	*
2003.....	0.6	*	*	*	*	1.3	1.7	1.9	*	*
2002.....	0.8	*	*	*	*	2.2	1.5	1.8	*	*
2001.....	0.6	*	*	*	*	1.6	1.5	*	*	*
2000.....	0.7	*	*	*	*	1.6	1.5	*	*	*
1999.....	0.8	*	*	*	*	2.1	2.1	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental drowning and submersion (W65-W74)

All races 3/, both sexes

2004.....	1.1	1.5	2.7	0.7	0.7	1.5	1.3	1.1	0.8	0.9
2003.....	1.1	1.4	2.9	0.6	0.7	1.4	1.4	1.0	0.8	1.0
2002.....	1.2	1.6	2.9	0.8	0.8	1.6	1.5	1.2	1.0	1.2
2001.....	1.2	1.7	3.0	0.8	0.8	1.6	1.4	1.1	0.8	1.0
2000.....	1.2	2.0	3.2	1.0	0.8	1.8	1.5	1.1	1.0	1.1
1999.....	1.3	1.8	3.2	0.9	0.9	1.8	1.5	1.1	1.1	1.1

All races 3/, male

2004.....	1.8	1.8	3.2	0.9	1.0	2.6	2.3	1.9	1.4	1.6
2003.....	1.8	1.7	3.7	0.9	1.0	2.5	2.5	1.8	1.4	1.7
2002.....	1.9	1.9	3.9	1.2	1.1	2.8	2.8	2.1	1.7	2.0
2001.....	1.8	1.5	3.7	1.2	1.1	2.8	2.5	1.9	1.4	1.6
2000.....	2.0	2.2	4.0	1.3	1.2	3.3	2.5	1.9	1.7	1.9
1999.....	2.0	2.0	4.0	1.3	1.3	3.2	2.8	2.0	2.0	1.9

All races 3/, female

2004.....	0.5	1.2	2.1	0.4	0.3	0.3	0.2	0.3	0.2	0.2
2003.....	0.5	1.2	2.0	0.4	0.3	*	0.2	*	*	0.3
2002.....	0.5	1.2	1.9	0.3	0.4	0.3	*	0.3	0.3	0.4
2001.....	0.5	1.9	2.2	0.5	0.5	0.3	0.3	0.3	0.2	0.4
2000.....	0.5	1.7	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3
1999.....	0.5	1.6	2.4	0.5	0.4	0.3	0.3	*	0.3	0.3

White, both sexes

2004.....	1.1	1.6	3.0	0.5	0.4	1.2	1.3	1.1	0.8	0.9
2003.....	1.1	1.4	3.1	0.6	0.5	1.3	1.3	1.0	0.8	1.0
2002.....	1.1	1.6	3.0	0.6	0.5	1.4	1.5	1.1	1.0	1.1
2001.....	1.1	1.7	3.3	0.6	0.5	1.3	1.4	1.0	0.7	1.0
2000.....	1.2	1.6	3.5	0.8	0.6	1.6	1.4	1.1	1.0	1.1
1999.....	1.2	1.5	3.5	0.7	0.5	1.6	1.5	1.0	1.1	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental drowning and submersion (W65-W74)

White, male

2004.....	1.7	1.8	3.5	0.7	0.6	2.1	2.2	1.8	1.3	1.5
2003.....	1.8	1.5	4.0	0.8	0.7	2.3	2.4	1.7	1.3	1.7
2002.....	1.8	1.9	4.0	0.9	0.6	2.4	2.7	1.9	1.7	1.9
2001.....	1.7	1.6	4.2	0.8	0.6	2.3	2.4	1.7	1.2	1.5
2000.....	1.9	1.8	4.4	1.0	0.8	2.8	2.4	1.8	1.7	1.8
1999.....	1.9	1.7	4.3	0.9	0.7	2.8	2.5	1.7	1.9	1.8

White, female

2004.....	0.5	1.4	2.4	0.3	*	*	0.3	0.3	0.3	0.3
2003.....	0.5	1.3	2.2	0.3	*	*	*	*	*	0.3
2002.....	0.4	*	2.0	0.3	0.3	0.3	*	0.3	*	0.4
2001.....	0.5	1.8	2.4	0.5	0.3	*	0.3	0.3	*	0.4
2000.....	0.5	*	2.5	0.5	0.3	0.3	0.3	0.3	*	0.4
1999.....	0.5	1.4	2.6	0.4	*	0.3	0.3	*	0.3	0.4

Black, both sexes

2004.....	1.4	*	1.9	1.3	1.8	2.4	1.2	1.3	0.9	1.1
2003.....	1.3	*	2.1	1.0	1.6	1.8	1.4	1.1	1.1	0.8
2002.....	1.6	*	2.5	1.8	2.2	2.3	1.9	1.6	1.2	1.5
2001.....	1.6	*	1.9	1.7	2.3	2.9	1.5	1.2	1.1	1.2
2000.....	1.6	4.2	2.2	1.9	2.1	2.8	1.7	1.3	0.9	1.2
1999.....	1.8	3.5	2.4	2.2	2.6	2.7	1.9	1.8	1.4	1.1

Black, male

2004.....	2.3	*	2.8	1.6	2.7	4.3	2.3	2.2	1.6	2.2
2003.....	2.2	*	3.0	1.4	2.4	3.4	2.6	2.1	2.2	1.7
2002.....	2.8	*	3.4	2.8	3.5	4.3	3.7	3.1	2.0	2.8
2001.....	2.7	*	2.2	2.7	3.2	5.2	3.1	2.6	2.2	2.2
2000.....	2.7	*	2.8	2.7	3.2	5.2	3.3	2.3	1.8	2.3
1999.....	3.1	*	3.2	3.4	3.9	5.0	3.8	3.5	2.7	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental drowning and submersion (W65-W74)

Black, female

2004.....	0.5	*	*	*	*	*	*	*	*
2003.....	0.4	*	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*
2001.....	0.5	*	1.6	*	1.3	*	*	*	*
2000.....	0.6	*	1.7	*	*	*	*	*	*
1999.....	0.5	*	1.6	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

All races 3/, both sexes

2004.....	1.5	1.4	0.4	*	*	0.1	0.2	0.2	0.2	0.3
2003.....	1.5	1.5	0.6	*	*	0.1	0.2	0.2	0.2	0.3
2002.....	1.4	1.5	0.5	*	*	*	0.2	0.2	0.2	0.3
2001.....	1.5	1.5	0.5	*	*	*	0.2	0.1	0.2	0.3
2000.....	1.5	2.0	0.6	*	*	0.1	0.1	0.2	0.3	0.3
1999.....	1.4	1.7	0.6	0.1	0.1	*	0.1	0.1	0.3	0.3

All races 3/, male

2004.....	1.6	2.3	0.6	*	*	0.2	0.2	0.3	0.3	0.3
2003.....	1.6	1.6	0.7	*	*	*	0.2	0.3	0.3	0.5
2002.....	1.5	2.0	0.6	*	*	*	0.2	0.3	0.3	0.4
2001.....	1.5	2.0	0.6	*	*	*	0.2	*	0.2	0.4
2000.....	1.6	2.3	0.7	*	*	*	*	0.2	0.4	0.5
1999.....	1.4	1.9	0.8	*	*	*	*	*	0.3	0.3

All races 3/, female

2004.....	1.5	*	0.3	*	*	*	*	*	0.2	0.3
2003.....	1.4	1.5	0.4	*	*	*	*	*	*	0.2
2002.....	1.4	1.0	0.3	*	*	*	*	*	*	0.3
2001.....	1.4	1.0	0.4	*	*	*	*	*	*	0.3
2000.....	1.5	1.8	0.5	*	*	*	*	*	0.2	0.2
1999.....	1.4	1.4	0.4	*	*	*	*	*	*	0.2

White, both sexes

2004.....	1.6	1.4	0.3	*	*	0.2	0.2	0.2	0.2	0.3
2003.....	1.6	1.3	0.5	*	*	*	0.2	0.2	0.2	0.3
2002.....	1.5	1.4	0.4	*	*	*	0.1	0.2	0.2	0.3
2001.....	1.5	1.3	0.5	*	*	*	0.2	*	0.2	0.3
2000.....	1.6	1.7	0.5	*	*	*	*	0.2	0.3	0.4
1999.....	1.4	1.4	0.5	*	*	*	0.1	*	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

All races 3/, both sexes

2004.....	0.5	16.4	0.3	0.1	0.3	0.2	0.2	0.2	0.3	0.3
2003.....	0.4	13.9	0.5	0.1	0.2	0.1	0.2	0.3	0.2	0.4
2002.....	0.5	14.3	0.4	0.1	0.2	0.2	0.3	0.2	0.3	0.3
2001.....	0.5	13.7	0.4	0.1	0.2	0.2	0.2	0.2	0.3	0.4
2000.....	0.5	11.8	0.4	0.2	0.3	0.2	0.3	0.2	0.3	0.4
1999.....	0.6	10.8	0.5	0.1	0.3	0.2	0.3	0.3	0.2	0.3

All races 3/, male

2004.....	0.7	17.7	0.4	0.2	0.5	0.3	0.4	0.3	0.4	0.5
2003.....	0.6	16.3	0.5	*	0.3	0.3	0.4	0.5	0.4	0.6
2002.....	0.7	15.7	0.5	*	0.4	0.4	0.5	0.4	0.5	0.5
2001.....	0.7	15.3	0.5	0.2	0.4	0.4	0.4	0.4	0.5	0.6
2000.....	0.7	13.2	0.5	*	0.5	0.4	0.5	0.4	0.5	0.7
1999.....	0.8	12.6	0.5	0.2	0.5	0.4	0.5	0.5	0.4	0.5

All races 3/, female

2004.....	0.3	14.9	*	*	*	*	*	*	*	*
2003.....	0.3	11.4	0.4	*	*	*	*	*	*	*
2002.....	0.3	12.8	0.4	*	*	*	*	*	*	*
2001.....	0.3	12.0	0.3	*	*	*	*	*	*	*
2000.....	0.3	10.3	0.3	*	*	*	*	*	*	*
1999.....	0.4	8.9	0.5	*	*	*	*	*	*	*

White, both sexes

2004.....	0.5	13.2	0.4	0.1	0.3	0.2	0.3	0.2	0.3	0.4
2003.....	0.4	11.1	0.5	*	0.2	0.2	0.3	0.3	0.3	0.4
2002.....	0.4	10.7	0.4	*	0.3	0.3	0.3	0.3	0.3	0.4
2001.....	0.5	10.8	0.4	0.2	0.3	0.2	0.3	0.3	0.3	0.4
2000.....	0.5	9.4	0.4	0.2	0.3	0.3	0.3	0.3	0.3	0.4
1999.....	0.6	8.7	0.5	0.1	0.3	0.3	0.3	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental exposure to electric current (W85-W87)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
2003.....	0.1	*	*	*	*	0.1	0.2	0.3	0.2	0.2
2002.....	0.1	*	*	*	*	0.2	0.2	0.3	0.2	0.2
2001.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.3
2000.....	0.1	*	*	*	*	0.1	0.2	0.2	0.2	0.2
1999.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.3

All races 3/, male

2004.....	0.2	*	*	*	*	*	0.4	0.4	0.5	0.3
2003.....	0.3	*	*	*	*	*	0.4	0.5	0.3	0.4
2002.....	0.3	*	*	*	*	0.3	0.4	0.5	0.4	0.5
2001.....	0.3	*	*	*	*	*	0.4	0.4	0.5	0.5
2000.....	0.3	*	*	*	*	0.2	0.5	0.4	0.4	0.5
1999.....	0.3	*	*	*	*	0.3	0.5	0.4	0.4	0.5

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	*	*	0.3	0.2	0.3	0.2
2003.....	0.1	*	*	*	*	0.1	0.3	0.3	0.2	0.2
2002.....	0.2	*	*	*	*	0.2	0.3	0.3	0.3	0.3
2001.....	0.2	*	*	*	*	*	0.2	0.2	0.3	0.3
2000.....	0.2	*	*	*	*	0.2	0.3	0.3	0.3	0.3
1999.....	0.2	*	*	*	*	0.2	0.3	0.3	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

All races 3/, both sexes

2004.....	1.1	0.6	1.4	0.9	0.4	0.3	0.6	0.7	0.5	0.7
2003.....	1.2	0.7	1.4	0.7	0.4	0.4	0.6	0.6	0.6	0.8
2002.....	1.1	0.9	1.4	0.8	0.5	0.4	0.5	0.6	0.7	0.7
2001.....	1.2	1.2	1.5	0.8	0.4	0.4	0.7	0.6	0.6	0.9
2000.....	1.2	1.0	1.9	0.9	0.4	0.4	0.6	0.6	0.6	0.8
1999.....	1.2	1.1	2.0	0.8	0.5	0.4	0.6	0.7	0.7	0.9

All races 3/, male

2004.....	1.3	*	1.5	1.1	0.4	0.4	0.6	0.9	0.7	0.9
2003.....	1.4	*	1.6	0.7	0.4	0.4	0.7	0.7	0.8	1.1
2002.....	1.4	1.0	1.6	0.9	0.6	0.6	0.7	0.8	0.9	0.9
2001.....	1.4	1.5	1.8	1.0	0.4	0.4	0.9	0.8	0.8	1.2
2000.....	1.5	1.0	2.2	1.0	0.3	0.4	0.8	0.8	0.8	1.2
1999.....	1.5	1.1	2.2	0.9	0.5	0.6	0.8	0.9	1.0	1.2

All races 3/, female

2004.....	0.9	*	1.3	0.6	0.4	0.2	0.5	0.4	0.3	0.4
2003.....	1.0	1.1	1.2	0.7	0.4	0.3	0.5	0.5	0.5	0.5
2002.....	0.8	*	1.2	0.6	0.4	0.3	0.4	0.4	0.6	0.4
2001.....	0.9	1.0	1.1	0.6	0.4	0.3	0.4	0.4	0.4	0.5
2000.....	0.9	*	1.5	0.7	0.5	0.4	0.4	0.4	0.4	0.5
1999.....	0.9	1.1	1.7	0.7	0.4	0.3	0.4	0.5	0.4	0.5

White, both sexes

2004.....	1.0	*	1.1	0.6	0.3	0.3	0.5	0.6	0.5	0.7
2003.....	1.1	0.7	1.2	0.5	0.2	0.3	0.6	0.6	0.7	0.8
2002.....	1.0	*	1.0	0.5	0.4	0.4	0.5	0.5	0.7	0.6
2001.....	1.0	1.1	1.2	0.7	0.4	0.3	0.7	0.5	0.6	0.8
2000.....	1.0	0.8	1.3	0.7	0.3	0.4	0.5	0.5	0.5	0.7
1999.....	1.0	0.9	1.5	0.6	0.4	0.4	0.6	0.7	0.7	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental exposure to smoke, fire and flames (X00-X09)

White, male

2004.....	1.2	*	1.2	0.7	0.3	0.4	0.6	0.8	0.7	0.8
2003.....	1.3	*	1.4	0.5	*	0.4	0.8	0.7	0.8	1.1
2002.....	1.2	*	1.1	0.6	0.5	0.5	0.6	0.7	0.9	0.8
2001.....	1.3	*	1.6	0.8	0.4	0.4	0.9	0.7	0.8	1.0
2000.....	1.3	*	1.6	0.7	0.3	0.4	0.8	0.7	0.7	1.0
1999.....	1.3	*	1.7	0.6	0.4	0.6	0.7	0.9	1.0	1.0

White, female

2004.....	0.8	*	1.0	0.5	0.3	0.3	0.5	0.4	0.4	0.5
2003.....	0.9	*	1.0	0.5	0.3	0.3	0.5	0.5	0.5	0.5
2002.....	0.7	*	1.0	0.4	*	*	0.3	0.3	0.5	0.4
2001.....	0.8	*	0.8	0.5	0.3	0.3	0.4	*	0.4	0.5
2000.....	0.8	*	1.1	0.6	0.4	0.4	0.3	0.4	0.3	0.4
1999.....	0.8	*	1.3	0.6	0.3	0.3	0.4	0.4	0.3	0.5

Black, both sexes

2004.....	2.1	*	2.8	2.3	0.9	*	0.8	1.2	0.8	0.9
2003.....	2.0	*	3.0	1.9	1.0	0.6	*	0.9	0.8	1.0
2002.....	2.1	*	3.5	1.9	0.9	0.7	0.9	1.2	1.1	1.0
2001.....	2.3	*	3.3	1.8	0.8	*	1.0	1.4	1.0	1.5
2000.....	2.5	*	4.8	2.1	0.8	*	1.0	1.1	1.2	1.5
1999.....	2.5	*	4.5	2.1	0.9	*	0.9	1.1	0.9	1.9

Black, male

2004.....	2.6	*	2.9	3.1	*	*	*	1.6	*	*
2003.....	2.4	*	3.2	1.9	*	*	*	*	*	*
2002.....	2.6	*	4.1	2.2	*	*	*	*	*	1.5
2001.....	2.9	*	3.6	2.3	*	*	*	*	*	2.5
2000.....	3.2	*	5.6	2.6	*	*	*	*	1.5	2.2
1999.....	3.0	*	5.0	2.4	*	*	*	*	*	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ years	year 2/ years	years	years	years	years	years	years	years	years

Accidental exposure to smoke, fire and flames (X00-X09)

Black, female

2004.....	1.6	*	2.7	1.4	*	*	*	*	*	*
2003.....	1.7	*	2.7	1.8	*	*	*	*	*	*
2002.....	1.6	*	2.8	1.7	*	*	*	*	*	*
2001.....	1.7	*	2.9	1.2	*	*	*	1.5	*	*
2000.....	2.0	*	4.1	1.6	*	*	*	*	*	*
1999.....	2.0	*	3.9	1.8	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes

2004.....	7.1	*	*	*	0.2	3.1	7.7	8.9	9.3	12.8
2003.....	6.7	0.5	0.3	*	0.2	2.5	7.1	7.9	9.3	12.5
2002.....	6.1	0.6	0.2	*	0.1	2.4	5.9	6.9	8.6	11.9
2001.....	4.9	*	0.2	*	0.2	2.0	4.9	5.7	6.9	10.4
2000.....	4.5	*	0.2	*	0.1	1.7	4.3	5.1	6.8	9.5
1999.....	4.4	*	0.2	*	0.1	1.3	3.8	4.9	6.8	9.1

All races 3/, male

2004.....	9.6	*	*	*	0.2	4.6	11.8	13.3	13.0	16.8
2003.....	9.2	*	0.3	*	0.2	3.8	10.9	11.7	13.1	16.8
2002.....	8.5	1.0	*	*	*	3.5	9.1	10.4	12.5	16.5
2001.....	7.1	*	0.3	*	*	3.0	7.6	8.7	9.9	14.6
2000.....	6.6	*	0.3	*	*	2.6	6.7	7.9	10.1	13.7
1999.....	6.5	*	*	*	*	1.9	5.9	7.7	10.3	13.3

All races 3/, female

2004.....	4.7	*	*	*	0.2	1.5	3.4	4.4	5.4	8.7
2003.....	4.3	*	0.3	*	*	1.2	3.1	4.0	5.4	8.1
2002.....	3.7	*	*	*	*	1.2	2.6	3.3	4.7	7.2
2001.....	2.9	*	*	*	*	1.0	1.9	2.6	3.9	6.1
2000.....	2.5	*	*	*	*	0.8	1.7	2.1	3.5	5.3
1999.....	2.3	*	*	*	*	0.7	1.6	2.0	3.2	4.9

White, both sexes

2004.....	7.6	*	*	*	0.2	3.7	9.1	10.4	10.3	13.8
2003.....	7.1	*	0.3	*	0.1	3.0	8.5	9.1	10.2	13.3
2002.....	6.3	*	*	*	0.2	2.8	6.9	7.8	9.5	12.5
2001.....	5.0	*	0.2	*	0.2	2.3	5.6	6.4	7.4	10.8
2000.....	4.6	*	*	*	0.1	2.0	4.9	5.5	7.2	9.8
1999.....	4.3	*	0.2	*	0.1	1.5	4.2	5.2	6.9	9.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning and exposure to noxious substances (X40-X49)

White, male

2004.....	10.2	*	*	*	0.3	5.6	13.8	15.3	14.5	18.2
2003.....	9.7	*	*	*	*	4.6	12.9	13.4	14.4	17.7
2002.....	8.8	*	*	*	*	4.1	10.6	11.6	13.6	17.3
2001.....	7.2	*	*	*	*	3.5	8.7	9.5	10.7	15.2
2000.....	6.7	*	*	*	*	3.1	7.8	8.6	10.7	14.3
1999.....	6.4	*	*	*	*	2.1	6.5	8.3	10.6	13.7

White, female

2004.....	5.0	*	*	*	*	1.7	4.0	5.1	6.0	9.3
2003.....	4.5	*	*	*	*	1.4	3.8	4.5	5.8	8.7
2002.....	3.9	*	*	*	*	1.4	3.0	3.8	5.1	7.5
2001.....	2.9	*	*	*	*	1.0	2.3	3.0	3.9	6.2
2000.....	2.5	*	*	*	*	0.9	1.8	2.3	3.6	5.2
1999.....	2.3	*	*	*	*	0.8	1.7	2.1	3.1	4.8

Black, both sexes

2004.....	6.5	*	*	*	*	0.9	2.9	4.5	6.2	11.0
2003.....	6.3	*	*	*	*	*	2.3	4.0	7.3	11.8
2002.....	6.3	*	*	*	*	*	2.3	4.4	6.8	11.9
2001.....	5.8	*	*	*	*	0.9	2.3	4.0	6.3	11.1
2000.....	5.5	*	*	*	*	0.7	2.1	4.3	6.8	10.7
1999.....	5.7	*	*	*	*	*	2.8	4.2	8.0	10.3

Black, male

2004.....	8.9	*	*	*	*	*	4.4	6.5	8.7	13.5
2003.....	8.8	*	*	*	*	*	3.5	5.7	10.2	16.7
2002.....	9.1	*	*	*	*	*	3.5	7.0	9.9	17.0
2001.....	8.4	*	*	*	*	*	3.8	6.7	8.2	15.4
2000.....	8.2	*	*	*	*	*	2.7	7.1	9.6	14.9
1999.....	8.7	*	*	*	*	*	4.7	6.7	11.5	14.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Black, female

2004.....	4.3	*	*	*	*	*	1.4	2.5	4.0	8.8
2003.....	3.9	*	*	*	*	*	*	2.4	4.8	7.4
2002.....	3.7	*	*	*	*	*	*	2.0	4.0	7.4
2001.....	3.5	*	*	*	*	*	*	1.6	4.6	7.3
2000.....	3.1	*	*	*	*	*	1.5	1.7	4.3	7.1
1999.....	3.0	*	*	*	*	*	*	2.0	4.9	6.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

All races 3/, both sexes

2004.....	6.8	*	*	*	0.1	2.9	7.5	8.6	8.9	12.3
2003.....	6.3	*	0.2	*	*	2.3	6.8	7.6	8.9	11.9
2002.....	5.7	*	*	*	*	2.2	5.6	6.6	8.2	11.4
2001.....	4.6	*	*	*	*	1.7	4.5	5.4	6.6	9.8
2000.....	4.2	*	*	*	*	1.4	4.0	4.7	6.5	9.1
1999.....	4.0	*	*	*	*	1.0	3.4	4.6	6.4	8.6

All races 3/, male

2004.....	9.1	*	*	*	*	4.3	11.3	12.8	12.5	16.0
2003.....	8.6	*	*	*	*	3.4	10.4	11.2	12.5	16.0
2002.....	7.9	*	*	*	*	3.2	8.6	9.9	11.8	15.7
2001.....	6.5	*	*	*	*	2.6	7.1	8.2	9.5	13.8
2000.....	6.1	*	*	*	*	2.1	6.2	7.5	9.4	13.0
1999.....	5.9	*	*	*	*	1.5	5.4	7.2	9.8	12.5

All races 3/, female

2004.....	4.5	*	*	*	*	1.3	3.3	4.3	5.3	8.5
2003.....	4.1	*	*	*	*	1.2	3.0	3.8	5.3	7.9
2002.....	3.5	*	*	*	*	1.1	2.5	3.2	4.6	7.0
2001.....	2.7	*	*	*	*	0.8	1.8	2.5	3.7	5.8
2000.....	2.3	*	*	*	*	0.7	1.6	2.0	3.4	5.2
1999.....	2.1	*	*	*	*	0.5	1.5	2.0	3.0	4.7

White, both sexes

2004.....	7.2	*	*	*	0.1	3.4	8.8	10.0	10.0	13.2
2003.....	6.7	*	*	*	*	2.8	8.1	8.7	9.8	12.7
2002.....	5.9	*	*	*	*	2.6	6.6	7.4	9.1	12.0
2001.....	4.7	*	*	*	*	2.0	5.2	6.0	7.1	10.2
2000.....	4.2	*	*	*	*	1.7	4.6	5.2	6.8	9.3
1999.....	4.0	*	*	*	*	1.2	3.8	5.0	6.6	8.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

White, male

2004.....	9.6	*	*	*	*	5.2	13.3	14.7	13.9	17.3
2003.....	9.1	*	*	*	*	4.2	12.4	12.8	13.8	16.9
2002.....	8.2	*	*	*	*	3.7	10.0	11.0	13.0	16.5
2001.....	6.6	*	*	*	*	3.1	8.1	9.1	10.3	14.4
2000.....	6.1	*	*	*	*	2.5	7.3	8.1	10.0	13.5
1999.....	5.9	*	*	*	*	1.7	5.9	7.8	10.1	12.9

White, female

2004.....	4.8	*	*	*	*	1.6	3.9	5.0	5.8	9.0
2003.....	4.3	*	*	*	*	1.4	3.6	4.3	5.7	8.4
2002.....	3.7	*	*	*	*	1.3	2.9	3.7	5.0	7.3
2001.....	2.7	*	*	*	*	0.9	2.2	2.8	3.8	6.0
2000.....	2.3	*	*	*	*	0.8	1.7	2.2	3.5	5.1
1999.....	2.1	*	*	*	*	0.7	1.6	2.1	3.0	4.7

Black, both sexes

2004.....	6.1	*	*	*	*	0.7	2.7	4.3	6.0	10.7
2003.....	5.9	*	*	*	*	*	2.1	3.7	7.1	11.5
2002.....	5.8	*	*	*	*	*	2.2	4.2	6.4	11.2
2001.....	5.4	*	*	*	*	*	2.1	3.8	6.1	10.5
2000.....	5.2	*	*	*	*	*	2.0	3.9	6.6	10.3
1999.....	5.3	*	*	*	*	*	2.5	4.2	7.6	9.8

Black, male

2004.....	8.4	*	*	*	*	*	4.1	6.4	8.4	13.1
2003.....	8.2	*	*	*	*	*	3.2	5.1	9.7	16.3
2002.....	8.4	*	*	*	*	*	3.4	6.7	9.3	15.8
2001.....	7.8	*	*	*	*	*	3.5	6.3	8.0	14.5
2000.....	7.7	*	*	*	*	*	2.6	6.6	9.3	14.0
1999.....	8.1	*	*	*	*	*	4.2	6.7	10.9	13.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Black, female

2004.....	4.1	*	*	*	*	*	1.3	2.4	3.8	8.6
2003.....	3.8	*	*	*	*	*	*	2.4	4.7	7.3
2002.....	3.5	*	*	*	*	*	*	2.0	3.9	7.1
2001.....	3.2	*	*	*	*	*	*	1.4	4.4	6.9
2000.....	2.9	*	*	*	*	*	1.4	1.5	4.2	7.1
1999.....	2.8	*	*	*	*	*	*	2.0	4.6	6.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.3
2003.....	0.2	*	*	*	*	0.2	0.1	0.1	0.1	0.3
2002.....	0.2	*	*	*	*	0.1	*	0.2	0.2	0.2
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.1	0.3
2000.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.2
1999.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.3

All races 3/, male

2004.....	0.3	*	*	*	*	0.2	0.3	0.3	0.2	0.5
2003.....	0.3	*	*	*	*	0.3	*	0.2	0.3	0.5
2002.....	0.3	*	*	*	*	*	*	0.3	0.4	0.4
2001.....	0.2	*	*	*	*	*	*	*	0.2	0.5
2000.....	0.2	*	*	*	*	*	*	*	0.3	0.4
1999.....	0.3	*	*	*	*	*	0.2	0.2	0.2	0.4

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	*	*	*	*	0.1	0.2	0.2	0.2	0.4
2003.....	0.2	*	*	*	*	0.2	0.1	0.2	0.2	0.3
2002.....	0.2	*	*	*	*	0.1	*	0.2	0.2	0.2
2001.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3
2000.....	0.2	*	*	*	*	*	*	*	0.1	0.2
1999.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to other gases and vapors (X47)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	0.1	0.1	0.2	0.2	0.2
2003.....	0.2	*	0.1	*	0.1	*	0.2	0.2	0.2	0.2
2002.....	0.2	*	*	*	*	0.1	0.2	0.2	0.2	0.3
2001.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.2
2000.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.2
1999.....	0.2	*	*	*	*	0.1	0.2	*	0.2	0.2

All races 3/, male

2004.....	0.3	*	*	*	*	*	*	0.3	0.3	0.3
2003.....	0.3	*	*	*	*	*	0.3	0.3	0.3	0.4
2002.....	0.3	*	*	*	*	*	0.3	0.3	0.3	0.4
2001.....	0.3	*	*	*	*	0.2	0.4	0.3	0.2	0.4
2000.....	0.3	*	*	*	*	0.3	0.3	0.3	0.4	0.4
1999.....	0.3	*	*	*	*	0.2	0.3	*	0.3	0.4

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	*	*	*	*	0.1	*	0.2	0.2	0.2
2003.....	0.2	*	*	*	*	*	0.3	0.2	0.2	0.2
2002.....	0.2	*	*	*	*	0.1	0.2	0.2	0.2	0.2
2001.....	0.2	*	*	*	*	0.2	0.3	0.2	0.2	0.2
2000.....	0.2	*	*	*	*	0.2	0.2	0.2	0.3	0.2
1999.....	0.2	*	*	*	*	0.2	0.2	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

All races 3/, both sexes

2004.....	3.5	1.3	0.7	0.3	0.3	0.7	1.1	1.2	1.4	1.7
2003.....	3.7	1.1	0.7	0.2	0.3	0.7	1.3	1.1	1.2	1.5
2002.....	3.8	0.9	0.6	0.3	0.4	0.8	1.0	1.1	1.3	1.6
2001.....	4.0	1.3	0.8	0.3	0.3	0.8	1.2	1.1	1.3	1.8
2000.....	3.9	1.1	0.7	0.3	0.3	0.8	1.2	1.2	1.4	1.9
1999.....	4.3	1.2	0.8	0.4	0.4	0.9	1.2	1.3	1.4	1.7

All races 3/, male

2004.....	4.2	1.6	0.9	0.4	0.4	1.2	1.9	2.0	2.3	2.8
2003.....	4.3	1.1	0.8	0.3	0.5	1.2	2.0	1.8	2.1	2.4
2002.....	4.3	1.2	0.7	0.4	0.4	1.2	1.8	1.8	2.3	2.6
2001.....	4.5	1.6	0.9	0.4	0.4	1.2	2.1	2.0	2.3	3.0
2000.....	4.5	1.2	0.8	0.4	0.4	1.3	2.2	2.2	2.4	3.3
1999.....	4.8	1.4	1.0	0.5	0.5	1.4	2.0	2.3	2.3	2.9

All races 3/, female

2004.....	2.9	*	0.4	*	*	0.3	0.2	0.3	0.4	0.5
2003.....	3.1	1.1	0.5	*	*	0.2	0.4	0.4	0.4	0.6
2002.....	3.2	*	0.5	*	0.3	0.3	0.2	0.4	0.3	0.6
2001.....	3.5	1.1	0.7	0.2	*	0.4	0.3	0.3	0.4	0.6
2000.....	3.3	*	0.5	0.2	*	0.3	0.2	0.2	0.4	0.5
1999.....	3.7	*	0.5	0.3	0.2	0.4	0.3	0.4	0.4	0.6

White, both sexes

2004.....	3.8	1.2	0.7	0.3	0.2	0.8	1.1	1.3	1.4	1.6
2003.....	4.0	1.0	0.6	0.2	0.3	0.7	1.3	1.2	1.3	1.4
2002.....	4.1	0.8	0.5	0.3	0.3	0.8	1.1	1.2	1.3	1.5
2001.....	4.3	1.3	0.8	0.3	0.3	0.8	1.3	1.2	1.3	1.8
2000.....	4.2	1.0	0.6	0.3	0.3	0.8	1.3	1.3	1.4	1.9
1999.....	4.5	1.0	0.8	0.4	0.4	0.9	1.2	1.3	1.4	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

White, male										
2004.....	4.4	1.5	1.0	0.4	0.3	1.3	1.9	2.1	2.4	2.7
2003.....	4.5	*	0.8	0.3	0.5	1.2	2.1	2.0	2.1	2.3
2002.....	4.5	*	0.6	0.5	0.3	1.2	2.0	2.0	2.3	2.5
2001.....	4.8	1.5	0.9	0.3	0.4	1.3	2.2	2.0	2.3	3.1
2000.....	4.6	*	0.7	0.4	0.4	1.3	2.2	2.3	2.5	3.2
1999.....	5.0	*	1.0	0.5	0.5	1.4	2.1	2.3	2.3	2.9
White, female										
2004.....	3.3	*	0.4	*	*	0.3	*	0.4	0.3	0.5
2003.....	3.5	*	0.5	*	*	*	0.3	0.4	0.4	0.6
2002.....	3.6	*	0.4	*	0.3	0.3	*	0.4	0.3	0.5
2001.....	3.9	*	0.6	*	*	0.4	0.3	*	0.4	0.5
2000.....	3.7	*	0.5	*	*	0.3	*	0.3	0.4	0.5
1999.....	4.1	*	0.5	0.3	0.3	0.4	0.3	0.3	0.4	0.6
Black, both sexes										
2004.....	2.5	*	*	*	*	*	1.0	1.2	1.5	2.2
2003.....	2.8	*	0.9	*	*	*	1.5	1.0	1.2	1.9
2002.....	2.8	*	1.0	*	*	0.8	*	0.9	1.4	2.3
2001.....	2.9	*	0.9	*	*	*	0.8	1.3	1.3	1.8
2000.....	3.0	*	0.9	*	*	0.7	1.2	1.1	1.5	2.4
1999.....	3.6	*	0.8	*	*	0.8	1.1	1.5	1.6	1.7
Black, male										
2004.....	3.5	*	*	*	*	*	1.6	2.2	2.6	3.7
2003.....	3.7	*	*	*	*	*	1.9	1.6	2.0	3.1
2002.....	3.7	*	*	*	*	*	*	*	2.6	4.1
2001.....	3.7	*	*	*	*	*	*	2.0	2.4	2.7
2000.....	4.4	*	*	*	*	*	2.2	2.2	2.5	4.4
1999.....	4.9	*	*	*	*	1.4	2.1	2.4	2.8	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All races 3/, both sexes

2004.....	11.0	*	1.3	8.2	12.5	12.3	13.0	14.0
2003.....	10.8	*	1.2	7.3	12.1	11.9	13.5	13.8
2002.....	11.0	*	1.2	7.4	12.4	12.8	12.5	14.3
2001 4/.....	10.8	*	1.3	7.9	12.0	12.6	13.0	14.3
2000.....	10.4	*	1.5	8.0	12.5	12.1	12.0	13.7
1999.....	10.5	*	1.2	8.0	12.3	12.3	13.1	14.4

All races 3/, male

2004.....	17.7	*	1.7	12.6	20.8	20.2	20.6	21.7
2003.....	17.6	*	1.7	11.6	20.2	19.5	21.6	21.9
2002.....	17.9	*	1.8	12.2	20.8	20.8	20.2	22.7
2001 4/.....	17.6	*	1.9	12.9	20.5	21.0	21.0	22.7
2000.....	17.1	*	2.3	13.0	21.4	20.0	19.2	21.7
1999.....	17.1	*	1.9	13.0	20.8	20.3	21.1	22.7

All races 3/, female

2004.....	4.6	*	1.0	3.5	3.6	4.2	5.3	6.3
2003.....	4.3	*	0.5	2.7	3.4	4.0	5.2	5.7
2002.....	4.3	*	0.6	2.4	3.5	4.5	4.7	6.0
2001 4/.....	4.1	*	0.6	2.7	3.1	4.0	4.8	5.9
2000.....	4.0	*	0.6	2.7	3.2	4.0	4.6	5.8
1999.....	4.0	*	0.5	2.7	3.4	4.1	5.1	6.1

White, both sexes

2004.....	12.3	*	1.3	8.8	13.2	13.0	14.2	15.6
2003.....	12.1	*	1.1	7.9	12.5	12.6	14.7	15.3
2002.....	12.2	*	1.3	8.2	13.2	13.5	13.7	15.9
2001 4/.....	11.9	*	1.3	8.6	12.6	13.5	14.2	15.8
2000.....	11.5	*	1.5	8.5	13.1	12.7	13.0	15.1
1999.....	11.5	*	1.3	8.5	12.7	12.8	14.2	15.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

White, male

2004.....	19.6	*	1.7	13.6	22.0	21.2	22.3	24.0
2003.....	19.5	*	1.7	12.7	21.1	20.3	23.3	23.9
2002.....	19.9	*	1.9	13.4	22.0	21.4	21.7	24.7
2001 4/.....	19.5	*	2.0	14.0	21.3	22.2	22.6	24.7
2000.....	18.8	*	2.4	13.7	22.3	20.8	20.7	23.6
1999.....	18.7	*	2.0	13.7	21.5	21.0	22.5	24.3

White, female

2004.....	5.1	*	0.9	3.7	3.8	4.3	5.6	7.0
2003.....	4.7	*	0.5	2.9	3.3	4.3	5.7	6.5
2002.....	4.8	*	0.7	2.6	3.7	5.1	5.3	6.8
2001 4/.....	4.6	*	0.6	2.9	3.3	4.3	5.4	6.7
2000.....	4.4	*	0.7	2.9	3.3	4.1	5.1	6.5
1999.....	4.5	*	0.5	2.9	3.5	4.3	5.5	6.9

Black, both sexes

2004.....	5.2	*	1.4	4.7	9.8	10.1	8.7	7.0
2003.....	5.1	*	1.2	3.8	10.5	9.5	9.1	7.5
2002.....	5.1	*	1.0	4.0	9.1	10.3	8.2	8.1
2001 4/.....	5.3	*	1.1	4.3	10.1	9.6	8.6	7.8
2000.....	5.4	*	1.2	5.5	11.0	9.9	7.6	7.3
1999.....	5.4	*	0.9	5.8	10.7	9.6	8.7	8.4

Black, male

2004.....	9.0	*	1.6	7.4	17.2	18.0	14.4	11.7
2003.....	8.8	*	1.9	6.6	18.0	17.4	15.7	12.9
2002.....	9.1	*	1.6	6.9	16.1	19.5	15.2	14.5
2001 4/.....	9.2	*	1.7	7.3	19.4	17.7	15.4	13.8
2000.....	9.4	*	2.1	9.5	19.5	17.5	13.6	13.0
1999.....	9.6	*	1.3	9.9	19.5	17.9	15.7	15.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ year 1/	year 2/ year 1/	years	years	years	years	years	years	years	years

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Black, female

2004.....	1.8	*	*	1.9	2.4	2.8	3.6	2.7
2003.....	1.8	*	*	*	3.1	2.2	3.1	2.6
2002.....	1.5	*	*	*	2.3	2.0	2.0	2.5
2001 4/.....	1.7	*	*	1.3	*	2.2	2.5	2.5
2000.....	1.7	*	*	1.4	3.0	3.0	2.2	2.2
1999.....	1.6	*	*	1.6	2.3	2.1	2.5	2.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

All races 3/, both sexes

2004.....	1.4	*	*	0.5	0.8	1.2	1.6	2.3
2003.....	1.3	*	*	0.4	0.7	1.1	1.6	2.1
2002.....	1.3	*	*	0.4	0.7	1.1	1.6	2.2
2001.....	1.2	*	*	0.4	0.7	1.0	1.5	2.1
2000.....	1.2	*	*	0.4	0.7	1.0	1.3	1.8
1999.....	1.1	*	*	0.3	0.7	1.0	1.3	1.9

All races 3/, male

2004.....	1.4	*	*	0.4	1.0	1.3	1.7	2.4
2003.....	1.3	*	*	0.4	0.7	1.1	1.6	2.2
2002.....	1.3	*	*	0.4	0.7	1.3	1.7	2.2
2001.....	1.2	*	*	0.4	0.7	1.1	1.6	2.4
2000.....	1.1	*	*	0.4	0.7	0.9	1.5	1.9
1999.....	1.1	*	*	0.3	0.7	1.0	1.2	2.0

All races 3/, female

2004.....	1.5	*	*	0.5	0.7	1.1	1.6	2.2
2003.....	1.4	*	*	0.4	0.7	1.0	1.7	2.1
2002.....	1.4	*	*	0.4	0.8	1.0	1.4	2.1
2001.....	1.3	*	*	0.5	0.8	0.9	1.4	1.8
2000.....	1.2	*	*	0.4	0.6	1.1	1.2	1.8
1999.....	1.2	*	*	0.4	0.6	1.1	1.4	1.8

White, both sexes

2004.....	1.6	*	*	0.5	1.0	1.4	1.9	2.6
2003.....	1.5	*	*	0.4	0.8	1.2	1.9	2.4
2002.....	1.5	*	*	0.4	0.9	1.3	1.7	2.5
2001.....	1.4	*	*	0.5	0.8	1.1	1.7	2.4
2000.....	1.3	*	*	0.4	0.7	1.1	1.5	2.1
1999.....	1.3	*	*	0.4	0.7	1.1	1.4	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

White, male

2004.....	1.6	*	*	0.5	1.1	1.5	1.9	2.7
2003.....	1.5	*	*	0.5	0.8	1.3	1.8	2.5
2002.....	1.5	*	*	0.5	0.8	1.4	1.9	2.5
2001.....	1.4	*	*	0.5	0.8	1.2	1.8	2.7
2000.....	1.2	*	*	0.4	0.7	1.0	1.7	2.1
1999.....	1.3	*	*	0.3	0.8	1.1	1.3	2.2

White, female

2004.....	1.7	*	*	0.6	0.8	1.3	1.8	2.5
2003.....	1.6	*	*	0.4	0.7	1.1	2.0	2.4
2002.....	1.6	*	*	0.4	0.9	1.2	1.5	2.4
2001.....	1.5	*	*	0.5	0.9	1.0	1.6	2.1
2000.....	1.4	*	*	0.4	0.7	1.2	1.4	2.0
1999.....	1.3	*	*	0.5	0.7	1.2	1.5	2.1

Black, both sexes

2004.....	0.4	*	*	*	*	*	0.8	0.9
2003.....	0.4	*	*	*	*	*	*	0.9
2002.....	0.4	*	*	*	*	*	1.0	0.8
2001.....	0.4	*	*	*	*	*	*	1.0
2000.....	0.4	*	*	*	*	*	*	0.8
1999.....	0.4	*	*	*	*	*	0.8	0.7

Black, male

2004.....	0.4	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

Black, female

2004.....	0.5	*	*	*	*	*	*	*
2003.....	0.5	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	0.1	0.1	0.1
2003.....	0.1	*	*	*	*	*	0.1	0.1
2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.1	*	0.1
2000.....	0.1	*	*	*	*	*	*	0.1
1999.....	0.1	*	*	*	*	*	*	0.1

All races 3/, male

2004.....	0.1	*	*	*	*	*	0.2	*
2003.....	0.1	*	*	*	*	*	*	0.2
2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	0.2
1999.....	0.1	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	*	*	0.1	*
2003.....	0.1	*	*	*	*	*	*	0.1
2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	0.1
1999.....	0.1	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

White, male

2004.....	0.1	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	0.2
2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	0.2
1999.....	0.1	*	*	*	*	*	*	*

White, female

2004.....	0.1	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*

Black, both sexes

2004.....	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Black, male

2004.....	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

Black, female

2004.....	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

All races 3/, both sexes

2004.....	0.5	*	*	0.1	0.2	0.4	0.5	0.7
2003.....	0.5	*	*	*	0.2	0.3	0.7	0.9
2002.....	0.5	*	*	0.2	0.2	0.4	0.7	0.8
2001.....	0.5	*	*	0.1	0.3	0.4	0.7	0.8
2000.....	0.5	*	*	0.2	0.3	0.6	0.7	0.9
1999.....	0.6	*	*	0.2	0.5	0.5	0.9	0.9

All races 3/, male

2004.....	0.7	*	*	0.2	0.4	0.7	0.9	1.1
2003.....	0.7	*	*	*	0.4	0.6	1.2	1.4
2002.....	0.8	*	*	0.3	0.4	0.7	1.0	1.2
2001.....	0.8	*	*	0.2	0.6	0.7	1.1	1.3
2000.....	0.8	*	*	0.3	0.5	0.9	1.1	1.4
1999.....	0.9	*	*	0.3	0.9	0.9	1.3	1.4

All races 3/, female

2004.....	0.2	*	*	*	*	*	*	0.3
2003.....	0.2	*	*	*	*	*	0.2	0.3
2002.....	0.2	*	*	*	*	*	0.3	0.3
2001.....	0.2	*	*	*	*	*	0.2	0.3
2000.....	0.2	*	*	*	*	0.2	0.3	0.4
1999.....	0.2	*	*	*	*	*	0.4	0.3

White, both sexes

2004.....	0.5	*	*	0.2	0.3	0.5	0.6	0.9
2003.....	0.6	*	*	*	0.3	0.4	0.8	1.0
2002.....	0.6	*	*	0.2	0.3	0.5	0.8	0.9
2001.....	0.6	*	*	0.2	0.4	0.5	0.8	0.9
2000.....	0.6	*	*	0.2	0.4	0.6	0.8	1.0
1999.....	0.6	*	*	0.2	0.6	0.6	1.0	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

White, male										
2004.....	0.8	*	*	0.3	0.4	0.8	1.0	1.3
2003.....	0.9	*	*	*	0.4	0.7	1.4	1.6
2002.....	0.9	*	*	0.3	0.4	0.8	1.2	1.4
2001.....	0.9	*	*	0.2	0.7	0.8	1.3	1.5
2000.....	1.0	*	*	0.3	0.7	1.0	1.3	1.6
1999.....	1.0	*	*	0.3	1.1	0.9	1.6	1.6
White, female										
2004.....	0.2	*	*	*	*	*	*	0.4
2003.....	0.2	*	*	*	*	*	*	0.4
2002.....	0.3	*	*	*	*	*	0.4	0.4
2001.....	0.3	*	*	*	*	*	*	0.4
2000.....	0.3	*	*	*	*	*	0.3	0.4
1999.....	0.3	*	*	*	*	*	0.5	0.4
Black, both sexes										
2004.....	0.1	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*
Black, male										
2004.....	0.2	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ years	year 2/ years	years	years	years	years	years	years	years	years

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

Black, female

2004.....	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

All races 3/, both sexes

2004.....	2.5	*	1.0	3.3	4.0	3.9	4.1	3.9
2003.....	2.3	*	0.7	2.8	3.8	3.6	3.8	3.5
2002.....	2.2	*	0.7	2.7	3.6	3.6	3.5	3.4
2001.....	2.2	*	0.8	2.7	3.5	3.5	3.5	3.6
2000.....	2.0	*	0.8	2.5	3.3	3.3	3.0	3.0
1999.....	1.9	*	0.6	2.2	2.9	3.1	3.4	3.2

All races 3/, male

2004.....	4.1	*	1.2	4.7	6.6	6.6	6.8	6.5
2003.....	3.8	*	1.1	4.2	6.3	6.0	6.4	6.0
2002.....	3.8	*	1.1	4.3	6.2	6.0	6.0	5.9
2001.....	3.7	*	1.2	4.3	6.0	6.0	5.9	6.2
2000.....	3.4	*	1.3	4.0	5.7	5.6	5.2	5.2
1999.....	3.3	*	1.0	3.4	4.8	5.3	5.9	5.6

All races 3/, female

2004.....	0.9	*	0.7	1.7	1.2	1.1	1.2	1.2
2003.....	0.8	*	0.3	1.2	1.1	1.1	1.2	1.0
2002.....	0.8	*	0.3	1.0	1.0	1.2	0.8	1.0
2001.....	0.7	*	0.3	1.0	0.9	0.9	1.0	1.1
2000.....	0.7	*	0.3	1.0	0.8	0.9	0.8	0.8
1999.....	0.7	*	0.2	0.9	0.8	0.9	0.9	0.8

White, both sexes

2004.....	2.6	*	0.9	3.3	4.3	4.1	4.4	4.3
2003.....	2.4	*	0.7	3.0	3.9	3.8	4.2	3.7
2002.....	2.4	*	0.7	3.0	4.0	3.8	3.7	3.7
2001.....	2.3	*	0.8	2.9	3.8	3.6	3.6	4.0
2000.....	2.1	*	0.8	2.7	3.4	3.4	3.1	3.2
1999.....	2.0	*	0.6	2.2	3.0	3.3	3.6	3.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

White, male

2004.....	4.4	*	1.2	4.8	7.1	7.0	7.3	7.2
2003.....	4.1	*	1.0	4.6	6.6	6.3	7.0	6.3
2002.....	4.0	*	1.1	4.9	6.7	6.3	6.3	6.2
2001.....	4.0	*	1.2	4.8	6.6	6.2	6.1	6.6
2000.....	3.6	*	1.2	4.3	6.0	6.0	5.4	5.5
1999.....	3.4	*	1.0	3.5	5.1	5.6	6.2	5.9

White, female

2004.....	0.9	*	0.7	1.8	1.3	1.0	1.3	1.3
2003.....	0.8	*	0.2	1.3	1.1	1.2	1.3	1.0
2002.....	0.8	*	0.4	1.1	1.0	1.2	0.9	1.1
2001.....	0.7	*	0.3	1.0	0.9	0.9	1.0	1.2
2000.....	0.7	*	0.4	1.1	0.7	0.8	0.8	0.9
1999.....	0.7	*	*	0.9	0.8	0.9	0.9	0.9

Black, both sexes

2004.....	1.4	*	1.0	2.2	2.6	2.9	2.5	1.5
2003.....	1.3	*	1.0	1.3	2.7	2.3	2.0	2.0
2002.....	1.1	*	0.7	1.0	2.0	2.3	2.3	1.9
2001.....	1.1	*	0.6	1.1	1.8	2.5	2.6	1.8
2000.....	1.2	*	1.0	1.3	2.3	2.3	1.9	1.7
1999.....	1.0	*	0.7	1.3	1.8	2.0	2.2	1.9

Black, male

2004.....	2.4	*	1.2	3.4	4.4	5.0	4.4	2.8
2003.....	2.3	*	1.6	2.2	4.2	4.2	3.6	3.7
2002.....	2.0	*	1.1	1.6	3.1	4.0	4.3	3.6
2001.....	2.0	*	*	1.9	3.1	4.7	4.8	3.3
2000.....	2.1	*	1.8	2.1	4.1	3.9	3.7	3.0
1999.....	1.9	*	*	1.9	3.3	3.8	4.2	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Black, female

2004.....	0.4	*	*	*	*	*	*	*
2003.....	0.4	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes

2004.....	5.7	*	0.3	3.8	6.3	5.7	5.7	5.9
2003.....	5.8	*	0.3	3.6	6.5	5.8	6.1	6.0
2002.....	5.9	*	0.4	3.6	6.7	6.4	5.6	6.6
2001.....	5.9	*	0.4	4.1	6.6	6.7	6.3	6.4
2000.....	5.9	*	0.5	4.4	7.2	6.3	6.0	6.7
1999.....	5.9	*	0.5	4.9	7.2	6.6	6.5	7.1

All races 3/, male

2004.....	10.0	*	0.5	6.5	11.1	9.9	9.6	9.8
2003.....	10.4	*	0.6	6.3	11.4	10.1	10.6	10.2
2002.....	10.6	*	0.6	6.4	11.9	11.1	9.6	11.3
2001.....	10.6	*	0.6	7.1	11.8	11.7	10.8	10.8
2000.....	10.5	*	0.9	7.6	12.9	11.1	10.1	11.1
1999.....	10.6	*	0.8	8.4	12.8	11.5	10.9	11.8

All races 3/, female

2004.....	1.5	*	*	1.0	1.1	1.3	1.7	1.9
2003.....	1.4	*	*	0.8	1.2	1.3	1.5	1.8
2002.....	1.4	*	*	0.7	1.2	1.6	1.7	2.0
2001.....	1.5	*	0.2	1.0	1.0	1.5	1.7	2.1
2000.....	1.5	*	0.2	1.1	1.3	1.5	1.9	2.3
1999.....	1.5	*	0.2	1.1	1.4	1.6	1.9	2.3

White, both sexes

2004.....	6.5	*	0.3	4.2	6.6	6.0	6.3	6.6
2003.....	6.6	*	0.4	4.0	6.7	6.1	6.7	6.8
2002.....	6.7	*	0.4	3.9	7.0	6.8	6.2	7.4
2001.....	6.7	*	0.4	4.5	6.7	7.2	7.0	7.2
2000.....	6.6	*	0.6	4.6	7.6	6.6	6.6	7.5
1999.....	6.6	*	0.6	5.2	7.4	7.0	7.1	7.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

White, male

2004.....	11.3	*	0.5	7.1	11.7	10.2	10.4	11.0
2003.....	11.7	*	0.7	6.9	11.8	10.5	11.3	11.3
2002.....	11.9	*	0.7	6.9	12.3	11.3	10.4	12.4
2001.....	11.8	*	0.7	7.6	11.9	12.3	11.7	11.9
2000.....	11.6	*	1.0	7.9	13.5	11.4	11.0	12.2
1999.....	11.7	*	0.9	8.9	13.0	11.9	11.8	12.7

White, female

2004.....	1.7	*	*	1.1	1.2	1.5	2.0	2.1
2003.....	1.6	*	*	0.9	1.2	1.5	1.8	2.1
2002.....	1.6	*	*	0.8	1.3	2.0	1.9	2.3
2001.....	1.7	*	*	1.1	1.1	1.7	2.0	2.4
2000.....	1.7	*	*	1.1	1.3	1.6	2.0	2.7
1999.....	1.7	*	0.3	1.2	1.4	1.8	2.2	2.7

Black, both sexes

2004.....	2.6	*	*	2.2	5.9	5.4	4.2	2.9
2003.....	2.6	*	*	2.1	6.5	5.5	4.8	2.8
2002.....	2.8	*	*	2.5	5.8	6.1	4.0	4.0
2001.....	2.9	*	*	2.8	7.1	5.8	4.1	3.7
2000.....	3.0	*	*	3.8	6.8	6.0	4.0	3.4
1999.....	3.1	*	*	3.9	7.6	5.7	4.2	4.2

Black, male

2004.....	4.9	*	*	3.7	10.7	10.2	7.8	5.5
2003.....	5.0	*	*	3.9	11.9	11.0	9.1	5.5
2002.....	5.4	*	*	4.3	11.0	12.4	7.7	7.4
2001.....	5.5	*	*	5.0	13.9	10.9	8.0	7.1
2000.....	5.5	*	*	6.9	12.6	11.3	7.3	6.4
1999.....	5.9	*	*	7.0	14.2	11.2	8.1	8.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Black, female

2004.....	0.5	*	*	*	*	*	*	*
2003.....	0.5	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by jumping from a high place (X80)

All races 3/, both sexes

2004.....	0.2	*	*	0.2	0.3	0.3	0.3	0.3
2003.....	0.2	*	*	0.2	0.3	0.3	0.3	0.4
2002.....	0.3	*	*	0.2	0.4	0.4	0.4	0.3
2001.....	0.2	*	*	0.2	0.2	0.3	0.3	0.3
2000.....	0.2	*	*	0.2	0.4	0.3	0.3	0.3
1999.....	0.2	*	*	0.2	0.4	0.3	0.3	0.4

All races 3/, male

2004.....	0.3	*	*	0.3	0.5	0.5	0.5	0.4
2003.....	0.4	*	*	0.2	0.5	0.4	0.5	0.5
2002.....	0.4	*	*	0.2	0.6	0.6	0.6	0.5
2001.....	0.3	*	*	0.3	0.3	0.4	0.4	0.4
2000.....	0.3	*	*	0.3	0.5	0.4	0.3	0.5
1999.....	0.4	*	*	0.3	0.6	0.5	0.4	0.6

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	0.2
2003.....	0.1	*	*	*	*	*	*	0.2
2002.....	0.2	*	*	*	*	*	0.3	0.2
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	0.2	*	*	*
1999.....	0.1	*	*	*	*	*	*	0.2

White, both sexes

2004.....	0.2	*	*	0.2	0.3	0.3	0.3	0.3
2003.....	0.3	*	*	0.1	0.3	0.3	0.3	0.4
2002.....	0.3	*	*	0.1	0.4	0.4	0.4	0.4
2001.....	0.2	*	*	0.2	0.3	0.2	0.3	0.3
2000.....	0.2	*	*	0.2	0.4	0.2	0.2	0.3
1999.....	0.2	*	*	0.2	0.4	0.3	0.3	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by jumping from a high place (X80)

White, male

2004.....	0.3	*	*	0.3	0.5	0.4	0.4	0.4
2003.....	0.4	*	*	*	0.5	0.4	0.5	0.5
2002.....	0.4	*	*	*	0.6	0.5	0.5	0.5
2001.....	0.3	*	*	0.3	0.4	0.3	0.4	0.4
2000.....	0.3	*	*	0.3	0.5	0.4	0.3	0.5
1999.....	0.4	*	*	0.2	0.6	0.4	0.4	0.5

White, female

2004.....	0.1	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	0.3	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	0.2

Black, both sexes

2004.....	0.2	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*

Black, male

2004.....	0.3	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by jumping from a high place (X80)

Black, female

2004.....	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	-------------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)

All races 3/, both sexes

2004.....	0.6	*	*	0.4	0.7	0.6	0.6	0.9
2003.....	0.6	*	*	0.2	0.5	0.7	0.7	0.9
2002.....	0.6	*	*	0.3	0.6	0.7	0.7	0.9
2001 4/.....	0.6	*	*	0.3	0.6	0.7	0.7	0.9
2000.....	0.6	*	*	0.3	0.6	0.6	0.6	0.9
1999.....	0.6	*	*	0.3	0.6	0.6	0.7	0.8

All races 3/, male

2004.....	1.0	*	*	0.5	1.1	1.0	0.9	1.3
2003.....	0.9	*	*	0.3	0.8	1.1	1.1	1.4
2002.....	1.0	*	*	0.6	1.0	1.0	1.2	1.5
2001 4/.....	0.9	*	*	0.5	1.0	1.1	1.0	1.4
2000.....	0.9	*	*	0.4	1.0	0.9	0.9	1.4
1999.....	0.8	*	*	0.5	1.0	0.9	1.2	1.1

All races 3/, female

2004.....	0.3	*	*	*	0.3	0.2	0.4	0.5
2003.....	0.3	*	*	*	0.2	0.2	0.4	0.3
2002.....	0.3	*	*	*	0.3	0.4	0.3	0.4
2001 4/.....	0.3	*	*	*	0.2	0.2	0.3	0.4
2000.....	0.3	*	*	*	0.2	*	0.3	0.3
1999.....	0.3	*	*	*	0.2	*	0.3	0.5

White, both sexes

2004.....	0.7	*	*	0.4	0.7	0.7	0.7	0.9
2003.....	0.6	*	*	0.2	0.5	0.6	0.7	0.9
2002.....	0.7	*	*	0.4	0.6	0.7	0.8	1.0
2001 4/.....	0.6	*	*	0.4	0.6	0.7	0.7	0.9
2000.....	0.6	*	*	0.3	0.6	0.6	0.6	0.9
1999.....	0.6	*	*	0.3	0.6	0.5	0.7	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)

White, male										
2004.....	1.0	*	*	0.6	1.1	1.0	1.0	1.3
2003.....	1.0	*	*	0.3	0.8	1.1	1.1	1.4
2002.....	1.0	*	*	0.6	1.0	1.0	1.2	1.5
2001 4/.....	1.0	*	*	0.6	0.9	1.1	1.1	1.4
2000.....	0.9	*	*	0.4	0.9	1.0	0.9	1.4
1999.....	0.9	*	*	0.5	1.0	0.9	1.1	1.1
White, female										
2004.....	0.3	*	*	*	0.3	*	0.3	0.5
2003.....	0.3	*	*	*	*	*	0.3	0.4
2002.....	0.3	*	*	*	0.3	0.4	0.3	0.4
2001 4/.....	0.3	*	*	*	*	*	0.4	0.4
2000.....	0.3	*	*	*	*	*	0.3	0.3
1999.....	0.3	*	*	*	*	*	0.3	0.5
Black, both sexes										
2004.....	0.5	*	*	*	0.7	*	*	1.0
2003.....	0.5	*	*	*	*	0.8	1.0	0.9
2002.....	0.5	*	*	*	0.7	0.8	*	1.1
2001 4/.....	0.5	*	*	*	*	*	*	0.9
2000.....	0.5	*	*	*	1.0	*	*	0.9
1999.....	0.5	*	*	*	*	*	0.9	0.8
Black, male										
2004.....	0.7	*	*	*	*	*	*	1.5
2003.....	0.7	*	*	*	*	*	1.5	1.6
2002.....	0.8	*	*	*	*	*	*	1.8
2001 4/.....	0.7	*	*	*	*	*	*	1.5
2000.....	0.8	*	*	*	*	*	*	1.9
1999.....	0.7	*	*	*	*	*	1.5	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)

Black, female

2004.....	0.2	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*
2001 4/.....	0.2	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,X87.1)

All races 3/, both sexes

2004.....	5.9	8.0	2.4	0.6	1.0	9.3	15.0	13.3	9.3	7.3
2003.....	6.1	8.5	2.4	0.6	1.0	9.5	16.5	13.4	9.4	7.6
2002.....	6.1	7.5	2.7	0.7	1.0	9.3	16.5	13.3	9.4	7.9
2001 4/.....	7.1	8.2	2.7	0.7	0.9	9.4	17.3	14.9	11.5	10.4
2000.....	6.0	9.2	2.3	0.7	1.1	9.5	16.0	11.9	9.0	7.6
1999.....	6.1	8.7	2.5	0.9	1.2	10.4	15.6	12.1	9.1	7.7

All races 3/, male

2004.....	9.4	8.0	2.5	0.6	1.3	15.3	25.4	22.3	14.9	11.2
2003.....	9.7	10.0	2.5	0.7	1.3	15.9	27.6	22.4	14.8	11.4
2002.....	9.6	7.9	2.9	0.6	1.2	15.3	27.7	22.0	14.7	11.8
2001 4/.....	11.1	9.5	3.0	0.7	1.1	15.7	28.9	24.1	17.7	15.3
2000.....	9.3	10.4	2.5	0.7	1.4	15.5	26.7	19.6	14.0	10.9
1999.....	9.3	9.6	2.5	0.9	1.5	16.9	25.7	19.5	13.8	11.0

All races 3/, female

2004.....	2.5	7.9	2.2	0.6	0.7	3.0	4.0	3.9	3.6	3.3
2003.....	2.6	6.9	2.3	0.5	0.6	2.6	4.8	4.0	3.9	3.9
2002.....	2.7	7.1	2.5	0.8	0.8	2.9	4.6	4.2	4.1	4.1
2001 4/.....	3.3	6.9	2.4	0.6	0.7	2.7	5.1	5.5	5.2	5.5
2000.....	2.8	7.9	2.1	0.6	0.8	3.1	4.7	4.1	4.0	4.4
1999.....	2.9	7.8	2.4	0.9	0.9	3.5	5.1	4.5	4.3	4.3

White, both sexes

2004.....	3.6	6.5	1.6	0.5	0.7	5.1	7.9	7.0	5.2	4.6
2003.....	3.7	6.9	1.7	0.4	0.6	5.0	8.4	6.9	5.3	5.0
2002.....	3.7	5.4	2.0	0.5	0.7	5.2	8.2	6.8	5.1	5.0
2001 4/.....	4.9	6.2	1.9	0.7	0.7	5.1	9.4	8.9	7.8	7.8
2000.....	3.6	6.6	1.7	0.6	0.9	5.2	7.7	6.2	5.3	4.9
1999.....	3.8	6.7	1.7	0.7	1.0	5.6	8.0	6.5	5.5	5.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,X87.1)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
White, male										
2004.....	5.3	6.9	1.8	0.5	0.9	7.9	12.5	10.9	7.7	6.8
2003.....	5.4	8.1	1.5	0.4	0.9	8.0	13.3	10.7	7.8	6.9
2002.....	5.4	6.2	2.1	0.4	0.7	8.2	12.9	10.5	7.3	7.0
2001 4/.....	7.2	7.3	2.2	0.6	0.8	7.9	14.5	13.5	11.3	11.6
2000.....	5.2	8.2	1.9	0.7	1.1	8.0	12.0	9.2	7.6	6.6
1999.....	5.5	7.9	1.8	0.7	1.1	8.6	12.1	9.9	7.8	7.0
White, female										
2004.....	1.9	6.2	1.4	0.5	0.5	2.1	3.0	2.9	2.6	2.4
2003.....	2.0	5.7	1.8	0.4	0.3	1.8	3.2	2.9	2.6	3.0
2002.....	2.0	4.6	1.8	0.6	0.6	1.9	3.1	2.8	2.8	3.1
2001 4/.....	2.6	5.1	1.6	0.7	0.6	2.0	3.9	4.1	4.2	3.9
2000.....	2.1	5.0	1.5	0.5	0.7	2.1	3.2	3.0	2.9	3.2
1999.....	2.1	5.4	1.7	0.7	0.8	2.3	3.7	2.9	3.2	3.1
Black, both sexes										
2004.....	21.1	15.6	6.3	1.4	2.4	31.6	55.9	52.8	35.6	25.2
2003.....	22.0	16.1	6.1	1.7	2.7	33.2	62.4	54.5	36.3	25.4
2002.....	22.0	17.3	6.3	1.4	2.6	31.0	63.4	54.2	37.0	26.9
2001 4/.....	22.1	18.9	6.8	0.9	1.9	32.7	62.8	52.6	35.3	28.2
2000.....	21.5	22.8	5.8	1.2	2.5	33.1	64.1	48.2	33.5	25.9
1999.....	21.1	19.5	6.1	1.9	2.5	36.5	60.4	47.0	31.8	25.1
Black, male										
2004.....	37.1	14.2	6.3	1.4	3.3	54.8	101.8	98.4	64.7	43.1
2003.....	38.9	17.8	7.4	2.0	3.7	58.9	111.6	101.6	63.9	43.7
2002.....	38.4	16.3	7.1	1.4	3.5	53.3	115.3	99.9	65.4	45.2
2001 4/.....	38.3	21.0	8.0	1.2	2.7	58.2	116.3	96.0	62.3	42.8
2000.....	37.2	23.3	5.9	*	3.3	57.2	117.8	89.4	58.6	42.5
1999.....	36.1	19.5	5.9	2.0	3.3	62.4	110.9	84.0	54.7	40.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,X87.1)

Black, female

2004.....	6.4	17.0	6.3	1.4	1.5	7.8	9.8	10.2	9.3	9.2
2003.....	6.6	14.2	4.8	1.3	1.7	6.7	13.6	10.9	11.4	9.0
2002.....	7.0	18.5	5.5	1.3	1.6	8.0	12.7	12.3	11.5	10.6
2001 4/.....	7.4	16.7	5.6	*	*	6.4	11.4	13.1	11.1	15.3
2000.....	7.2	22.2	5.7	1.4	1.7	8.4	13.2	10.9	10.9	11.1
1999.....	7.6	19.5	6.3	1.8	1.6	10.0	12.5	13.6	11.2	11.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by hanging, strangulation and suffocation (X91)

All races 3/, both sexes

2004.....	0.2	1.0	0.2	*	*	0.1	0.4	0.3	0.3	0.3
2003.....	0.2	1.0	*	*	*	0.2	0.3	0.3	0.3	0.4
2002.....	0.2	0.8	0.2	*	*	0.2	0.3	0.3	0.3	0.4
2001.....	0.2	1.0	*	*	*	0.2	0.4	0.3	0.3	0.3
2000.....	0.2	0.7	0.2	*	*	0.2	0.4	0.3	0.4	0.3
1999.....	0.3	0.9	0.1	*	*	0.2	0.4	0.4	0.4	0.4

All races 3/, male

2004.....	0.1	*	*	*	*	*	0.2	*	0.2	*
2003.....	0.2	1.0	*	*	*	*	*	*	*	0.3
2002.....	0.2	*	*	*	*	*	0.2	0.2	*	0.2
2001.....	0.2	1.2	*	*	*	*	0.2	*	*	0.2
2000.....	0.2	*	*	*	*	*	*	*	0.3	*
1999.....	0.2	1.1	*	*	*	*	*	0.2	0.3	0.2

All races 3/, female

2004.....	0.3	1.3	*	*	*	0.2	0.5	0.6	0.4	0.4
2003.....	0.3	*	*	*	*	0.4	0.5	0.5	0.5	0.4
2002.....	0.3	*	*	*	*	0.3	0.5	0.5	0.4	0.6
2001.....	0.3	*	*	*	*	0.3	0.6	0.5	0.5	0.4
2000.....	0.3	*	*	*	*	0.3	0.6	0.5	0.4	0.4
1999.....	0.3	*	*	*	*	0.4	0.6	0.6	0.6	0.5

White, both sexes

2004.....	0.2	0.9	*	*	*	*	0.3	0.3	0.2	0.2
2003.....	0.2	0.9	*	*	*	0.2	0.3	0.3	0.3	0.3
2002.....	0.2	0.6	*	*	*	0.1	0.3	0.3	0.2	0.3
2001.....	0.2	0.7	*	*	*	0.2	0.3	0.3	0.3	0.3
2000.....	0.2	*	0.2	*	*	0.2	0.3	0.3	0.2	0.2
1999.....	0.2	0.7	*	*	*	0.2	0.3	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by hanging, strangulation and suffocation (X91)

Black, female

2004.....	0.7	*	*	*	*	*	*	*	*
2003.....	0.7	*	*	*	*	*	*	*	*
2002.....	0.7	*	*	*	*	*	*	*	1.6
2001.....	0.7	*	*	*	*	*	*	*	1.4
2000.....	0.8	*	*	*	*	*	1.4	1.4	1.4
1999.....	0.8	*	*	*	*	*	*	1.6	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All races 3/, both sexes

2004.....	4.0	*	0.2	0.2	0.7	7.6	12.2	10.5	7.0	4.9
2003.....	4.1	*	0.3	0.2	0.7	7.7	13.6	10.8	7.1	5.0
2002.....	4.1	*	0.3	0.3	0.7	7.7	13.6	10.6	6.9	5.2
2001.....	4.0	*	0.4	0.3	0.6	7.5	13.6	10.3	6.6	5.1
2000.....	3.8	*	0.2	0.2	0.7	7.7	12.7	9.2	6.3	4.7
1999.....	3.9	*	0.3	0.3	0.8	8.5	12.5	9.2	6.1	4.6

All races 3/, male

2004.....	6.9	*	*	0.2	1.0	13.1	21.5	18.8	11.9	8.3
2003.....	7.1	*	0.2	0.3	1.0	13.7	23.8	19.1	12.0	8.0
2002.....	7.0	*	0.4	0.3	0.9	13.2	23.8	18.6	11.6	8.4
2001.....	6.8	*	0.3	0.3	0.8	13.2	24.0	17.8	11.1	8.0
2000.....	6.5	*	*	0.3	1.0	13.3	22.4	16.1	10.4	7.4
1999.....	6.5	*	0.3	0.4	1.1	14.5	21.8	15.9	10.1	7.1

All races 3/, female

2004.....	1.1	*	*	0.2	0.3	1.8	2.2	2.0	2.0	1.4
2003.....	1.2	*	0.3	0.2	0.3	1.5	2.8	2.2	2.1	1.9
2002.....	1.3	*	*	0.3	0.5	1.8	2.9	2.4	2.1	2.0
2001.....	1.3	*	0.4	0.2	0.3	1.5	2.7	2.5	2.1	2.2
2000.....	1.3	*	*	0.2	0.3	1.7	2.7	2.2	2.1	1.9
1999.....	1.3	*	0.3	0.2	0.5	2.2	2.8	2.3	2.1	2.2

White, both sexes

2004.....	2.1	*	*	0.2	0.4	3.8	5.5	4.9	3.7	2.9
2003.....	2.2	*	0.2	0.1	0.4	3.8	6.3	4.9	3.6	3.0
2002.....	2.2	*	0.3	0.2	0.4	4.1	6.0	4.9	3.4	3.0
2001.....	2.2	*	0.3	0.3	0.4	3.7	6.3	5.0	3.5	3.2
2000.....	2.1	*	*	0.2	0.5	3.9	5.5	4.2	3.3	2.8
1999.....	2.2	*	0.2	0.2	0.6	4.2	5.6	4.3	3.4	3.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

White, male

2004.....	3.5	*	*	*	0.6	6.3	9.3	8.3	5.8	4.7
2003.....	3.5	*	*	*	0.6	6.4	10.6	8.2	5.7	4.4
2002.....	3.5	*	0.3	*	0.5	6.7	10.0	8.1	5.3	4.5
2001.....	3.5	*	*	0.3	0.6	6.2	10.5	8.2	5.4	4.9
2000.....	3.2	*	*	0.3	0.7	6.5	9.0	6.7	5.1	4.1
1999.....	3.4	*	*	0.2	0.8	6.9	9.1	7.1	5.2	4.3

White, female

2004.....	0.9	*	*	*	*	1.2	1.5	1.2	1.4	1.0
2003.....	0.9	*	*	*	*	1.0	1.7	1.4	1.4	1.6
2002.....	1.0	*	*	*	0.3	1.2	1.6	1.6	1.4	1.5
2001.....	1.0	*	*	*	0.3	1.1	1.9	1.6	1.6	1.6
2000.....	1.0	*	*	*	*	1.1	1.7	1.6	1.5	1.5
1999.....	0.9	*	*	*	0.4	1.2	1.9	1.4	1.5	1.6

Black, both sexes

2004.....	16.1	*	0.8	*	1.8	27.8	50.4	46.3	29.3	18.5
2003.....	16.8	*	*	0.7	2.0	29.0	55.3	47.9	30.1	18.1
2002.....	16.7	*	*	*	1.9	27.3	57.0	46.9	30.1	19.5
2001.....	15.8	*	*	*	1.3	28.7	55.7	43.6	26.9	18.4
2000.....	15.6	*	*	*	1.7	28.6	55.3	40.7	26.0	17.3
1999.....	15.4	*	*	0.7	1.8	32.0	53.4	39.3	23.7	15.9

Black, male

2004.....	30.3	*	*	*	2.9	49.8	93.9	88.9	55.2	34.6
2003.....	31.6	*	*	*	2.9	53.2	101.7	92.4	56.4	33.9
2002.....	31.0	*	*	*	2.7	48.4	105.9	90.2	56.3	35.7
2001.....	29.8	*	*	*	2.1	52.8	106.5	83.2	51.3	31.5
2000.....	29.2	*	*	*	2.8	51.7	105.2	78.7	48.1	31.7
1999.....	28.4	*	*	*	2.8	56.3	101.0	74.4	44.0	27.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, female

2004.....	3.0	*	*	*	*	5.1	6.6	6.4	5.9	4.0
2003.....	3.3	*	*	*	1.1	4.0	9.1	6.8	6.3	4.1
2002.....	3.6	*	*	*	1.2	5.4	9.2	7.3	6.5	5.0
2001.....	3.1	*	*	*	*	3.9	7.0	7.6	5.0	6.8
2000.....	3.2	*	*	*	*	4.9	8.0	6.2	6.0	4.5
1999.....	3.7	*	*	*	*	7.0	8.2	7.5	5.5	5.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by sharp object (X99)

All races 3/, both sexes

2004.....	0.7	*	*	*	0.1	0.8	1.5	1.3	1.1	1.0
2003.....	0.7	*	*	*	*	0.8	1.5	1.3	1.1	1.1
2002.....	0.7	*	*	0.1	*	0.7	1.4	1.3	1.2	1.1
2001.....	0.7	*	*	*	*	0.8	1.6	1.4	1.0	1.1
2000.....	0.6	*	*	*	0.1	0.8	1.3	1.1	1.0	1.1
1999.....	0.7	*	*	*	0.1	0.8	1.4	1.1	1.2	1.1

All races 3/, male

2004.....	1.0	*	*	*	*	1.3	2.3	1.9	1.5	1.4
2003.....	1.0	*	*	*	*	1.3	2.1	1.8	1.5	1.6
2002.....	1.0	*	*	*	*	1.1	2.2	1.9	1.5	1.6
2001.....	1.0	*	*	*	*	1.3	2.4	2.0	1.4	1.4
2000.....	0.9	*	*	*	*	1.1	2.1	1.6	1.5	1.4
1999.....	1.0	*	*	*	*	1.2	2.0	1.6	1.7	1.6

All races 3/, female

2004.....	0.4	*	*	*	*	0.4	0.6	0.8	0.6	0.7
2003.....	0.4	*	*	*	*	0.3	0.7	0.7	0.7	0.6
2002.....	0.4	*	*	*	*	0.3	0.5	0.8	0.9	0.7
2001.....	0.4	*	*	*	*	0.3	0.8	0.7	0.6	0.8
2000.....	0.4	*	*	*	*	0.4	0.5	0.5	0.6	0.7
1999.....	0.4	*	*	*	*	0.4	0.7	0.6	0.7	0.6

White, both sexes

2004.....	0.5	*	*	*	*	0.6	1.2	1.0	0.7	0.7
2003.....	0.5	*	*	*	*	0.5	0.9	0.9	0.8	0.7
2002.....	0.5	*	*	*	*	0.4	1.0	0.9	0.7	0.8
2001.....	0.5	*	*	*	*	0.5	1.2	1.0	0.7	0.7
2000.....	0.4	*	*	*	*	0.6	0.9	0.8	0.8	0.7
1999.....	0.5	*	*	*	*	0.5	1.0	0.9	0.8	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by sharp object (X99)

White, male

2004.....	0.7	*	*	*	*	0.9	1.8	1.3	0.8	1.0
2003.....	0.7	*	*	*	*	0.8	1.3	1.3	1.1	1.0
2002.....	0.7	*	*	*	*	0.7	1.6	1.3	0.9	1.2
2001.....	0.7	*	*	*	*	0.8	1.9	1.4	0.9	0.9
2000.....	0.6	*	*	*	*	0.9	1.5	1.2	1.0	1.0
1999.....	0.7	*	*	*	*	0.8	1.4	1.3	1.1	0.9

White, female

2004.....	0.3	*	*	*	*	*	0.5	0.7	0.5	0.5
2003.....	0.3	*	*	*	*	*	0.5	0.5	0.4	0.4
2002.....	0.3	*	*	*	*	*	0.4	0.5	0.6	0.4
2001.....	0.3	*	*	*	*	*	0.5	0.5	0.4	0.5
2000.....	0.3	*	*	*	*	0.3	0.3	0.4	0.5	0.5
1999.....	0.3	*	*	*	*	0.3	0.5	0.4	0.5	0.4

Black, both sexes

2004.....	2.1	*	*	*	*	2.1	3.3	3.4	3.3	3.0
2003.....	2.2	*	*	*	*	2.4	4.2	3.5	3.1	3.7
2002.....	2.2	*	*	*	*	2.0	3.3	4.2	3.4	3.2
2001.....	2.0	*	*	*	*	2.2	3.7	4.1	2.8	3.5
2000.....	1.9	*	*	*	*	1.6	3.5	2.9	2.7	3.4
1999.....	2.0	*	*	*	*	2.0	3.7	2.6	3.6	3.9

Black, male

2004.....	3.2	*	*	*	*	2.9	5.1	5.4	5.4	4.0
2003.....	3.4	*	*	*	*	3.7	6.5	5.6	4.5	5.4
2002.....	3.2	*	*	*	*	3.0	5.4	6.3	4.9	4.1
2001.....	3.0	*	*	*	*	3.3	5.4	6.3	4.2	4.7
2000.....	2.8	*	*	*	*	2.0	5.7	4.5	4.4	4.6
1999.....	3.1	*	*	*	*	3.1	5.3	3.6	5.5	6.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by sharp object (X99)

Black, female

2004.....	1.1	*	*	*	*	1.3	1.4	1.5	1.4	2.2
2003.....	1.1	*	*	*	*	*	2.0	1.6	1.9	2.1
2002.....	1.2	*	*	*	*	*	1.3	2.2	2.0	2.3
2001.....	1.1	*	*	*	*	*	2.0	2.1	1.6	2.5
2000.....	1.1	*	*	*	*	*	1.5	1.4	*	2.3
1999.....	1.0	*	*	*	*	*	2.2	1.7	1.9	1.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

All races 3/, both sexes

2004.....	0.1	2.5	0.4	*	*	*	*	*	*	*
2003.....	0.1	2.5	0.5	*	*	*	*	*	*	*
2002.....	0.1	2.4	0.6	*	*	*	*	*	*	*
2001.....	0.1	2.9	0.5	*	*	*	*	*	*	*
2000.....	0.1	3.4	0.5	*	*	*	*	*	*	*
1999.....	0.1	3.3	0.4	*	*	*	*	*	*	*

All races 3/, male

2004.....	0.1	2.9	0.5	*	*	*	*	*	*	*
2003.....	0.1	3.3	0.5	*	*	*	*	*	*	*
2002.....	0.1	2.5	0.6	*	*	*	*	*	*	*
2001.....	0.1	3.2	0.7	*	*	*	*	*	*	*
2000.....	0.1	4.0	0.5	*	*	*	*	*	*	*
1999.....	0.1	3.7	0.4	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.1	2.0	0.4	*	*	*	*	*	*	*
2003.....	0.1	1.7	0.6	*	*	*	*	*	*	*
2002.....	0.1	2.4	0.6	*	*	*	*	*	*	*
2001.....	0.1	2.5	0.4	*	*	*	*	*	*	*
2000.....	0.1	2.7	0.4	*	*	*	*	*	*	*
1999.....	0.1	2.9	0.4	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	2.1	0.3	*	*	*	*	*	*	*
2003.....	0.1	2.1	0.4	*	*	*	*	*	*	*
2002.....	0.1	1.9	0.5	*	*	*	*	*	*	*
2001.....	0.1	2.2	0.4	*	*	*	*	*	*	*
2000.....	0.1	2.6	0.4	*	*	*	*	*	*	*
1999.....	0.1	2.5	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

All races 3/, both sexes

2004.....	0.9	3.8	1.3	0.2	0.1	0.6	0.9	1.0	0.8	1.0
2003.....	0.9	4.4	1.3	0.2	0.1	0.5	1.0	0.9	0.8	1.1
2002.....	0.9	3.7	1.4	0.2	0.1	0.6	0.9	0.8	0.9	1.0
2001 4/.....	2.0	3.5	1.6	0.2	0.2	0.7	1.5	2.8	3.4	3.6
2000.....	1.0	4.3	1.3	0.3	0.2	0.7	1.3	1.1	1.2	1.4
1999.....	1.0	3.8	1.3	0.4	0.2	0.8	1.1	1.1	1.1	1.3

All races 3/, male

2004.....	1.2	3.8	1.3	0.2	*	0.7	1.3	1.4	1.2	1.3
2003.....	1.2	4.7	1.5	0.2	*	0.8	1.3	1.2	1.0	1.4
2002.....	1.2	4.0	1.6	*	*	0.7	1.2	1.1	1.2	1.3
2001 4/.....	2.9	4.0	1.7	0.3	*	0.9	2.0	3.8	4.9	5.5
2000.....	1.4	4.7	1.5	0.3	0.2	0.9	1.8	1.6	1.5	1.7
1999.....	1.3	4.0	1.4	0.3	0.2	1.0	1.4	1.4	1.4	1.8

All races 3/, female

2004.....	0.6	3.8	1.3	*	*	0.6	0.5	0.5	0.5	0.7
2003.....	0.6	4.0	1.2	*	*	0.3	0.7	0.6	0.6	0.8
2002.....	0.5	3.3	1.2	*	*	0.4	0.6	0.5	0.6	0.7
2001 4/.....	1.1	3.1	1.4	*	0.2	0.5	1.0	1.7	1.9	1.8
2000.....	0.7	3.9	1.1	0.3	0.3	0.6	0.8	0.7	0.8	1.1
1999.....	0.7	3.6	1.2	0.4	*	0.6	0.7	0.8	0.9	0.9

White, both sexes

2004.....	0.7	3.1	0.9	0.2	*	0.5	0.8	0.8	0.6	0.7
2003.....	0.7	3.5	0.9	*	*	0.5	0.8	0.7	0.6	0.9
2002.....	0.7	2.5	0.9	*	*	0.4	0.7	0.6	0.7	0.8
2001 4/.....	1.8	2.7	1.1	0.2	0.2	0.6	1.4	2.6	3.3	3.4
2000.....	0.7	3.0	0.8	0.2	0.2	0.5	0.9	0.8	0.8	1.0
1999.....	0.8	2.9	1.0	0.2	0.2	0.6	0.9	0.8	0.8	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

White, male

2004.....	0.9	3.5	1.0	*	*	0.6	1.1	1.0	0.8	0.9
2003.....	0.9	3.8	0.9	*	*	0.6	1.1	0.8	0.8	1.1
2002.....	0.9	2.9	1.1	*	*	0.5	0.9	0.9	0.9	1.0
2001 4/.....	2.6	3.1	1.2	*	*	0.7	1.8	3.5	4.7	5.5
2000.....	0.9	3.5	1.0	*	*	0.5	1.2	1.0	1.0	1.2
1999.....	1.0	3.1	1.1	*	*	0.7	1.2	1.0	1.0	1.3

White, female

2004.....	0.5	2.6	0.8	*	*	0.4	0.5	0.5	0.4	0.5
2003.....	0.5	3.1	0.9	*	*	0.3	0.6	0.5	0.4	0.6
2002.....	0.4	2.2	0.8	*	*	0.4	0.5	0.4	0.4	0.6
2001 4/.....	1.0	2.3	0.9	*	*	0.4	0.9	1.6	1.7	1.3
2000.....	0.5	2.4	0.7	*	*	0.5	0.6	0.6	0.6	0.7
1999.....	0.5	2.5	0.9	*	*	0.5	0.7	0.6	0.7	0.7

Black, both sexes

2004.....	2.0	7.7	3.7	*	*	1.3	1.5	2.2	2.2	2.5
2003.....	2.1	8.8	3.6	0.6	*	0.9	2.1	2.1	2.2	2.5
2002.....	2.1	9.2	3.8	*	*	1.0	2.0	1.9	2.3	2.6
2001 4/.....	3.1	8.0	4.1	*	*	1.3	2.2	3.8	4.5	5.1
2000.....	2.9	11.2	3.8	0.7	*	2.2	3.9	3.3	3.3	4.0
1999.....	2.5	8.9	3.2	0.9	*	1.7	2.0	3.1	3.0	3.6

Black, male

2004.....	2.8	5.9	3.4	*	*	1.4	2.3	3.5	3.6	3.7
2003.....	3.1	8.6	4.5	*	*	1.3	2.7	3.3	2.5	3.5
2002.....	3.1	9.3	4.3	*	*	1.5	2.9	2.5	3.2	3.7
2001 4/.....	4.3	8.4	4.8	*	*	1.6	3.4	5.7	5.9	6.1
2000.....	4.2	11.2	3.9	*	*	3.0	5.7	5.2	4.9	5.3
1999.....	3.4	9.1	3.6	*	*	2.5	3.3	4.2	3.9	4.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39
	ages 1/ year 2/	year 2/ year 2/	years	years	years	years	years	years	years	years

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

Black, female

2004.....	1.3	9.6	4.0	*	*	*	*	*	*	1.5
2003.....	1.2	9.0	2.6	*	*	*	1.4	*	1.8	1.7
2002.....	1.2	9.1	3.3	*	*	*	*	*	1.6	1.6
2001 4/.....	2.1	7.6	3.3	*	*	*	*	2.0	3.1	4.3
2000.....	1.7	11.2	3.7	*	*	1.3	2.2	1.7	1.9	2.8
1999.....	1.6	8.7	2.9	*	*	*	*	2.1	2.1	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2004.....	1.7	2.6	0.3	0.1	0.2	0.7	1.7	1.9	2.1	2.9
2003.....	1.7	1.9	0.2	*	0.2	0.8	1.9	1.8	2.3	3.2
2002.....	1.7	2.5	0.4	0.1	0.2	0.7	1.7	1.9	2.3	3.3
2001.....	1.5	2.1	0.3	*	0.2	0.7	1.4	1.6	2.0	3.0
2000.....	1.4	1.7	0.3	*	0.2	0.6	1.3	1.6	1.8	2.6
1999.....	1.4	1.4	0.3	*	0.2	0.8	1.5	1.6	2.1	2.8

All races 3/, male

2004.....	2.2	3.1	0.3	*	0.2	1.1	2.4	2.6	2.9	3.8
2003.....	2.3	2.4	*	*	0.3	1.2	2.7	2.5	3.2	4.2
2002.....	2.2	2.5	0.4	*	0.2	1.0	2.4	2.7	3.1	4.5
2001.....	2.0	2.4	*	*	0.3	1.0	2.1	2.4	2.7	3.9
2000.....	1.9	1.4	0.3	*	0.4	0.9	2.0	2.3	2.6	3.7
1999.....	2.0	1.7	0.3	*	0.3	1.3	2.2	2.3	2.8	3.9

All races 3/, female

2004.....	1.2	2.0	0.3	*	*	0.4	0.8	1.1	1.4	1.9
2003.....	1.2	1.3	*	*	*	0.5	1.1	1.1	1.5	2.2
2002.....	1.2	2.5	0.4	*	*	0.4	1.0	1.0	1.4	2.1
2001.....	0.9	1.7	0.3	*	*	0.3	0.7	0.9	1.3	2.1
2000.....	0.8	2.0	*	*	*	0.3	0.6	0.9	1.1	1.6
1999.....	0.9	1.2	0.3	*	*	0.3	0.6	0.9	1.3	1.7

White, both sexes

2004.....	1.7	2.1	0.2	*	0.2	0.8	1.9	2.1	2.2	3.0
2003.....	1.8	1.7	0.2	*	0.2	0.9	2.1	2.0	2.6	3.4
2002.....	1.7	1.8	0.4	*	0.2	0.7	1.9	1.9	2.3	3.4
2001.....	1.5	1.4	0.3	*	0.2	0.7	1.5	1.7	2.1	3.0
2000.....	1.3	1.4	0.2	*	0.2	0.6	1.3	1.6	1.8	2.7
1999.....	1.4	1.2	0.2	*	0.2	0.8	1.5	1.6	2.1	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)

White, male

2004.....	2.2	2.7	*	*	*	1.1	2.7	2.9	2.9	3.9
2003.....	2.3	1.9	*	*	0.3	1.3	2.9	2.7	3.3	4.4
2002.....	2.2	1.8	0.4	*	0.2	1.1	2.6	2.7	3.1	4.5
2001.....	2.0	1.7	*	*	0.3	1.1	2.2	2.5	2.9	3.8
2000.....	1.8	*	*	*	0.3	1.0	1.9	2.3	2.5	3.7
1999.....	1.9	1.4	*	*	0.3	1.3	2.3	2.3	2.8	3.7

White, female

2004.....	1.3	1.5	*	*	*	0.4	0.9	1.2	1.5	2.1
2003.....	1.3	1.4	*	*	*	0.5	1.2	1.2	1.7	2.3
2002.....	1.2	1.8	0.4	*	*	0.4	1.1	1.1	1.5	2.3
2001.....	1.0	*	*	*	*	0.3	0.8	0.9	1.4	2.3
2000.....	0.8	1.6	*	*	*	0.3	0.7	0.9	1.1	1.6
1999.....	0.9	*	*	*	*	*	0.6	1.0	1.4	1.8

Black, both sexes

2004.....	1.8	4.8	*	*	*	0.7	1.1	1.0	2.6	3.1
2003.....	1.8	3.2	*	*	*	*	1.1	1.4	2.1	3.2
2002.....	2.0	5.5	*	*	*	*	1.3	2.0	2.5	4.1
2001.....	1.7	5.5	*	*	*	*	1.1	1.5	1.9	3.8
2000.....	1.8	*	*	*	*	*	1.4	1.7	2.2	3.3
1999.....	1.9	*	*	*	*	0.8	1.7	1.9	2.5	3.9

Black, male

2004.....	2.6	*	*	*	*	*	1.5	1.5	4.0	4.6
2003.....	2.7	*	*	*	*	*	1.9	2.5	3.4	4.4
2002.....	2.9	*	*	*	*	*	1.9	3.2	3.7	6.5
2001.....	2.7	6.0	*	*	*	*	1.9	2.5	2.7	5.9
2000.....	2.8	*	*	*	*	*	2.5	2.7	3.4	4.8
1999.....	3.0	*	*	*	*	1.3	2.7	3.4	3.7	6.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Black, female

2004.....	1.1	*	*	*	*	*	*	*	1.4	1.8
2003.....	0.9	*	*	*	*	*	*	*	*	2.2
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.9
2001.....	0.9	*	*	*	*	*	*	*	*	2.0
2000.....	0.9	*	*	*	*	*	*	*	*	2.0
1999.....	0.9	*	*	*	*	*	*	*	1.4	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

All races 3/, both sexes

2004.....	1.1	*	*	*	*	0.4	1.1	1.3	1.6	2.3
2003.....	1.2	*	*	*	*	0.4	1.4	1.4	1.8	2.6
2002.....	1.1	*	*	*	*	0.4	1.0	1.4	1.7	2.5
2001.....	1.0	*	*	*	*	0.3	0.8	1.1	1.5	2.3
2000.....	0.9	*	*	*	*	0.3	0.7	1.0	1.2	2.1
1999.....	0.9	*	*	*	*	0.3	0.8	1.1	1.5	2.1

All races 3/, male

2004.....	1.4	*	*	*	*	0.6	1.5	1.8	2.0	2.9
2003.....	1.6	*	*	*	*	0.6	1.9	1.8	2.4	3.3
2002.....	1.4	*	*	*	*	0.5	1.4	1.9	2.1	3.3
2001.....	1.3	*	*	*	*	0.5	1.2	1.5	1.9	2.9
2000.....	1.2	*	*	*	*	0.4	1.0	1.4	1.7	2.8
1999.....	1.2	*	*	*	*	0.4	1.1	1.5	2.0	2.9

All races 3/, female

2004.....	0.9	*	*	*	*	*	0.6	0.9	1.1	1.6
2003.....	0.9	*	*	*	*	0.3	0.7	0.9	1.2	1.8
2002.....	0.8	*	*	*	*	0.2	0.6	0.8	1.2	1.7
2001.....	0.7	*	*	*	*	*	0.4	0.7	1.1	1.7
2000.....	0.5	*	*	*	*	*	0.4	0.6	0.7	1.3
1999.....	0.6	*	*	*	*	*	0.4	0.6	1.0	1.4

White, both sexes

2004.....	1.2	*	*	*	*	0.4	1.3	1.6	1.7	2.4
2003.....	1.3	*	*	*	*	0.5	1.6	1.6	2.0	2.8
2002.....	1.2	*	*	*	*	0.4	1.2	1.5	1.8	2.6
2001.....	1.0	*	*	*	*	0.4	1.0	1.2	1.6	2.4
2000.....	0.9	*	*	*	*	0.3	0.8	1.1	1.3	2.1
1999.....	0.9	*	*	*	*	0.3	0.9	1.1	1.7	2.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

White, male

2004.....	1.5	*	*	*	*	0.7	1.9	2.2	2.1	3.0
2003.....	1.6	*	*	*	*	0.7	2.3	2.1	2.6	3.5
2002.....	1.4	*	*	*	*	0.5	1.6	2.1	2.2	3.3
2001.....	1.3	*	*	*	*	0.6	1.4	1.6	2.1	2.9
2000.....	1.2	*	*	*	*	0.4	1.2	1.5	1.8	2.9
1999.....	1.1	*	*	*	*	0.5	1.2	1.5	2.2	2.8

White, female

2004.....	1.0	*	*	*	*	*	0.8	1.0	1.2	1.8
2003.....	1.0	*	*	*	*	0.3	0.9	1.1	1.4	2.0
2002.....	0.9	*	*	*	*	0.3	0.8	0.9	1.2	1.9
2001.....	0.7	*	*	*	*	*	0.6	0.8	1.1	1.8
2000.....	0.6	*	*	*	*	*	0.5	0.7	0.8	1.3
1999.....	0.6	*	*	*	*	*	0.5	0.7	1.1	1.4

Black, both sexes

2004.....	1.0	*	*	*	*	*	*	*	1.6	2.5
2003.....	1.1	*	*	*	*	*	*	*	1.4	2.4
2002.....	1.2	*	*	*	*	*	*	1.1	1.7	2.9
2001.....	1.1	*	*	*	*	*	*	0.8	1.6	2.8
2000.....	1.0	*	*	*	*	*	*	0.9	1.3	2.4
1999.....	1.2	*	*	*	*	*	*	1.2	1.5	2.9

Black, male

2004.....	1.4	*	*	*	*	*	*	*	2.2	3.6
2003.....	1.7	*	*	*	*	*	*	*	2.1	3.1
2002.....	1.6	*	*	*	*	*	*	1.6	2.2	4.4
2001.....	1.7	*	*	*	*	*	*	*	2.0	4.0
2000.....	1.5	*	*	*	*	*	*	*	2.0	3.3
1999.....	1.8	*	*	*	*	*	*	2.1	2.2	4.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Discharge of firearms, undetermined intent (Y22-Y24)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	0.1	0.2	0.1	0.1	*
2003.....	0.1	*	*	*	*	0.2	0.2	0.1	*	0.1
2002.....	0.1	*	*	*	*	0.2	0.2	0.1	*	0.1
2001.....	0.1	*	*	*	*	0.1	0.2	0.2	*	0.1
2000.....	0.1	*	*	*	*	0.1	0.2	0.2	0.1	*
1999.....	0.1	*	*	*	*	0.3	0.3	0.2	*	0.1

All races 3/, male

2004.....	0.1	*	*	*	*	*	0.3	*	*	*
2003.....	0.1	*	*	*	*	0.3	0.3	*	*	*
2002.....	0.1	*	*	*	*	0.3	0.4	0.2	*	0.2
2001.....	0.1	*	*	*	*	0.2	0.4	0.3	*	0.2
2000.....	0.1	*	*	*	*	0.2	0.4	0.3	*	*
1999.....	0.2	*	*	*	*	0.6	0.5	0.3	*	0.2

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	*	*	0.2	*	*	*
2003.....	0.1	*	*	*	*	0.1	0.1	*	*	*
2002.....	0.1	*	*	*	*	0.1	0.2	0.1	*	0.1
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	0.2
2000.....	0.1	*	*	*	*	0.1	0.2	0.1	*	*
1999.....	0.1	*	*	*	*	0.3	0.3	0.1	*	0.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2004.....	0.4	2.3	0.2	*	0.1	0.2	0.3	0.4	0.4	0.4
2003.....	0.4	1.7	0.1	*	0.1	0.2	0.3	0.3	0.3	0.4
2002.....	0.4	2.1	0.3	0.1	0.1	0.1	0.4	0.3	0.4	0.5
2001.....	0.3	1.7	0.2	*	0.1	0.1	0.3	0.3	0.3	0.5
2000.....	0.3	1.6	0.2	*	0.1	0.2	0.3	0.3	0.4	0.4
1999.....	0.3	1.2	0.2	*	0.1	0.2	0.3	0.3	0.4	0.5

All races 3/, male

2004.....	0.6	2.9	0.3	*	*	0.3	0.5	0.5	0.6	0.6
2003.....	0.5	2.2	*	*	*	0.3	0.4	0.5	0.5	0.6
2002.....	0.5	2.0	0.3	*	*	*	0.5	0.4	0.6	0.9
2001.....	0.5	2.0	*	*	*	0.2	0.5	0.5	0.4	0.7
2000.....	0.5	1.2	0.3	*	0.2	0.2	0.5	0.4	0.6	0.5
1999.....	0.5	1.4	*	*	*	0.3	0.5	0.5	0.5	0.7

All races 3/, female

2004.....	0.3	1.7	*	*	*	*	*	*	*	0.2
2003.....	0.2	1.2	*	*	*	*	*	*	0.2	0.3
2002.....	0.3	2.1	0.3	*	*	*	0.2	*	*	0.2
2001.....	0.2	1.5	0.3	*	*	*	*	*	*	0.2
2000.....	0.2	1.9	*	*	*	*	*	*	0.2	0.2
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.3

White, both sexes

2004.....	0.4	2.0	0.2	*	*	0.2	0.3	0.4	0.3	0.4
2003.....	0.3	1.5	*	*	0.1	0.2	0.2	0.2	0.3	0.4
2002.....	0.4	1.6	0.3	*	*	0.1	0.4	0.2	0.4	0.5
2001.....	0.3	1.1	0.2	*	*	0.2	0.3	0.3	0.3	0.4
2000.....	0.3	1.2	0.2	*	*	0.2	0.3	0.3	0.4	0.4
1999.....	0.3	1.1	0.2	*	*	0.2	0.3	0.3	0.3	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Legal intervention (Y35,Y89.0)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	0.1	0.3	0.3	0.3	0.3
2003.....	0.1	*	*	*	*	0.1	0.4	0.3	0.3	0.3
2002.....	0.1	*	*	*	*	0.1	0.3	0.3	0.2	0.3
2001.....	0.1	*	*	*	*	0.1	0.3	0.2	0.3	0.3
2000.....	0.1	*	*	*	*	0.1	0.2	0.3	0.2	0.3
1999.....	0.1	*	*	*	*	0.1	0.3	0.3	0.3	0.3

All races 3/, male

2004.....	0.3	*	*	*	*	0.2	0.5	0.5	0.5	0.5
2003.....	0.3	*	*	*	*	*	0.7	0.6	0.7	0.6
2002.....	0.3	*	*	*	*	0.2	0.6	0.5	0.5	0.5
2001.....	0.3	*	*	*	*	0.2	0.6	0.5	0.6	0.6
2000.....	0.3	*	*	*	*	0.3	0.5	0.6	0.5	0.5
1999.....	0.3	*	*	*	*	0.2	0.5	0.7	0.7	0.6

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	*	0.1	0.2	0.2	0.2	0.3
2003.....	0.1	*	*	*	*	*	0.3	0.2	0.3	0.3
2002.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.3
2001.....	0.1	*	*	*	*	*	0.3	0.2	0.3	0.3
2000.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
1999.....	0.1	*	*	*	*	*	0.1	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Legal intervention (Y35,Y89.0)

White, male

2004.....	0.2	*	*	*	*	0.3	0.4	0.3	0.5	0.5
2003.....	0.2	*	*	*	*	*	0.5	0.4	0.5	0.6
2002.....	0.2	*	*	*	*	*	0.4	0.4	0.5	0.5
2001.....	0.2	*	*	*	*	*	0.5	0.4	0.5	0.5
2000.....	0.2	*	*	*	*	*	0.4	0.4	0.4	0.4
1999.....	0.2	*	*	*	*	*	0.3	0.5	0.6	0.5

White, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, both sexes

2004.....	0.2	*	*	*	*	*	*	0.8	*	*
2003.....	0.3	*	*	*	*	*	0.9	1.1	0.9	*
2002.....	0.3	*	*	*	*	*	0.9	0.8	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	0.7
1999.....	0.3	*	*	*	*	*	0.8	0.9	*	0.7

Black, male

2004.....	0.5	*	*	*	*	*	*	1.6	*	*
2003.....	0.6	*	*	*	*	*	1.6	2.2	1.8	*
2002.....	0.6	*	*	*	*	*	1.7	1.6	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.6	*	*	*	*	*	*	*	*	1.4
1999.....	0.7	*	*	*	*	*	1.7	1.9	*	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	0.1	0.3	0.2	0.2	0.2
2003.....	0.1	*	*	*	*	0.1	0.4	0.3	0.3	0.3
2002.....	0.1	*	*	*	*	0.1	0.3	0.2	0.2	0.2
2001.....	0.1	*	*	*	*	0.1	0.3	0.2	0.3	0.2
2000.....	0.1	*	*	*	*	0.1	0.2	0.3	0.2	0.2
1999.....	0.1	*	*	*	*	0.1	0.2	0.3	0.3	0.2

All races 3/, male

2004.....	0.2	*	*	*	*	0.2	0.5	0.4	0.4	0.4
2003.....	0.2	*	*	*	*	*	0.7	0.6	0.5	0.5
2002.....	0.2	*	*	*	*	0.2	0.6	0.5	0.4	0.4
2001.....	0.2	*	*	*	*	0.2	0.6	0.4	0.5	0.4
2000.....	0.2	*	*	*	*	0.3	0.4	0.5	0.3	0.4
1999.....	0.2	*	*	*	*	0.2	0.5	0.6	0.6	0.4

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	*	0.1	0.2	0.2	0.2	0.2
2003.....	0.1	*	*	*	*	*	0.3	0.2	0.2	0.3
2002.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
2001.....	0.1	*	*	*	*	*	0.3	0.2	0.2	0.2
2000.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
1999.....	0.1	*	*	*	*	*	0.1	0.2	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)

White, male

2004.....	0.2	*	*	*	*	0.3	0.4	0.3	0.4	0.4
2003.....	0.2	*	*	*	*	*	0.5	0.4	0.4	0.5
2002.....	0.2	*	*	*	*	*	0.4	0.3	0.4	0.3
2001.....	0.2	*	*	*	*	*	0.5	0.4	0.5	0.4
2000.....	0.2	*	*	*	*	*	0.4	0.4	0.3	0.3
1999.....	0.2	*	*	*	*	*	0.3	0.4	0.5	0.3

White, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, both sexes

2004.....	0.2	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	0.9	1.0	*	*
2002.....	0.2	*	*	*	*	*	0.9	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	0.8	*	*	*

Black, male

2004.....	0.4	*	*	*	*	*	*	*	*	*
2003.....	0.5	*	*	*	*	*	1.6	2.0	*	*
2002.....	0.5	*	*	*	*	*	1.7	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	1.7	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes

2004.....	1.0	*	*	*	*	0.1	0.1	0.2	0.2	0.3
2003.....	1.0	*	0.2	*	*	0.1	0.1	0.1	0.2	0.4
2002.....	1.0	*	*	*	0.1	*	*	0.1	0.2	0.3
2001.....	1.1	0.6	0.2	*	*	*	0.1	0.1	0.3	0.3
2000.....	1.1	*	0.1	*	*	*	0.1	0.2	0.2	0.3
1999.....	1.0	1.4	0.2	*	*	*	*	0.1	0.2	0.3

All races 3/, male

2004.....	0.9	*	*	*	*	*	*	*	*	0.2
2003.....	0.9	*	*	*	*	*	*	*	0.2	0.3
2002.....	0.9	*	*	*	*	*	*	*	0.3	0.3
2001.....	1.0	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.0	*	*	*	*	*	*	0.2	0.2	0.3
1999.....	1.0	1.6	*	*	*	*	*	*	0.3	0.3

All races 3/, female

2004.....	1.0	*	*	*	*	*	*	*	0.3	0.3
2003.....	1.0	*	*	*	*	*	*	*	0.2	0.4
2002.....	1.1	*	*	*	*	*	*	*	0.2	0.3
2001.....	1.1	*	*	*	*	*	*	*	0.2	0.3
2000.....	1.1	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.1	1.2	*	*	*	*	*	0.2	*	0.3

White, both sexes

2004.....	1.0	*	*	*	*	*	*	0.2	0.2	0.2
2003.....	1.0	*	*	*	*	*	*	*	0.2	0.3
2002.....	1.0	*	*	*	*	*	*	0.1	0.2	0.3
2001.....	1.1	*	*	*	*	*	0.1	*	0.2	0.3
2000.....	1.1	*	*	*	*	*	*	0.1	0.2	0.3
1999.....	1.0	1.2	0.2	*	*	*	*	*	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

All races 3/, both sexes

2004.....	0.8	*	*	*	*	*	*	*	0.1	0.2
2003.....	0.8	*	0.1	*	*	*	*	*	0.1	0.2
2002.....	0.8	*	*	*	*	*	*	*	0.2	0.3
2001.....	0.9	*	*	*	*	*	*	*	0.2	0.2
2000.....	0.9	*	*	*	*	*	*	0.1	0.2	0.2
1999.....	0.9	1.2	0.1	*	*	*	*	*	0.2	0.2

All races 3/, male

2004.....	0.8	*	*	*	*	*	*	*	*	0.2
2003.....	0.8	*	*	*	*	*	*	*	*	0.2
2002.....	0.7	*	*	*	*	*	*	*	*	0.2
2001.....	0.8	*	*	*	*	*	*	*	*	0.2
2000.....	0.9	*	*	*	*	*	*	*	*	*
1999.....	0.8	1.4	*	*	*	*	*	*	0.2	0.2

All races 3/, female

2004.....	0.8	*	*	*	*	*	*	*	*	0.2
2003.....	0.9	*	*	*	*	*	*	*	*	0.3
2002.....	0.9	*	*	*	*	*	*	*	*	0.3
2001.....	1.0	*	*	*	*	*	*	*	*	0.2
2000.....	1.0	*	*	*	*	*	*	*	0.2	0.2
1999.....	0.9	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.8	*	*	*	*	*	*	*	0.1	0.2
2003.....	0.8	*	*	*	*	*	*	*	*	0.2
2002.....	0.9	*	*	*	*	*	*	*	0.2	0.2
2001.....	0.9	*	*	*	*	*	*	*	0.2	0.2
2000.....	0.9	*	*	*	*	*	*	*	0.2	0.2
1999.....	0.9	1.0	*	*	*	*	*	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
All races 3/, both sexes										
2004.....	231.7	352.3	511.7	734.8	1,140.1	1,727.4	2,676.4	4,193.2	6,717.2	13,823.5
2003.....	241.3	359.2	517.8	755.6	1,182.7	1,793.7	2,778.3	4,363.4	6,976.2	14,593.3
2002.....	242.6	357.1	512.8	770.1	1,187.7	1,832.7	2,845.9	4,449.1	7,103.8	14,828.3
2001.....	240.5	354.8	512.4	771.8	1,210.7	1,869.7	2,878.3	4,494.0	7,151.9	15,112.8
2000.....	239.1	352.5	509.0	792.6	1,241.0	1,906.3	2,929.4	4,553.8	7,335.0	15,524.4
1999.....	236.3	346.6	501.4	801.6	1,254.0	1,956.7	2,995.3	4,608.0	7,378.5	15,554.6
All races 3/, male										
2004.....	289.5	444.4	657.0	917.3	1,411.0	2,125.1	3,279.2	5,173.5	8,179.0	15,031.1
2003.....	304.1	457.9	660.8	942.3	1,462.2	2,216.6	3,428.7	5,400.3	8,513.7	15,794.0
2002.....	306.4	454.7	653.5	961.8	1,476.0	2,265.9	3,533.0	5,522.7	8,652.3	16,254.5
2001.....	304.8	455.4	646.8	958.4	1,496.8	2,315.6	3,587.4	5,603.2	8,781.9	16,744.8
2000.....	305.9	456.3	642.2	989.8	1,535.9	2,362.0	3,676.1	5,693.2	9,095.4	17,501.4
1999.....	304.0	448.0	632.4	1,003.8	1,560.8	2,444.1	3,785.1	5,806.4	9,216.8	17,597.2
All races 3/, female										
2004.....	174.5	262.5	372.7	562.7	893.6	1,381.2	2,191.0	3,488.7	5,816.6	13,280.3
2003.....	179.2	263.2	380.9	579.7	928.8	1,427.1	2,256.6	3,625.3	6,036.0	14,062.5
2002.....	179.7	262.4	378.2	590.0	926.0	1,458.7	2,296.7	3,690.9	6,162.2	14,209.6
2001.....	177.2	257.2	383.7	596.9	951.3	1,485.9	2,314.8	3,717.6	6,175.1	14,429.9
2000.....	173.4	252.0	381.3	608.1	973.7	1,515.7	2,341.2	3,760.2	6,296.6	14,719.2
1999.....	169.8	248.5	375.9	612.6	976.4	1,539.6	2,378.2	3,774.8	6,302.8	14,731.3
White, both sexes										
2004.....	215.0	325.0	471.5	688.5	1,090.5	1,675.5	2,634.1	4,160.5	6,736.0	14,010.1
2003.....	222.6	329.4	474.5	710.6	1,127.8	1,739.2	2,728.6	4,329.5	6,992.1	14,792.3
2002.....	224.0	327.4	471.3	723.5	1,133.0	1,777.6	2,796.3	4,407.8	7,111.3	15,015.9
2001.....	218.8	322.8	471.4	723.2	1,156.1	1,805.8	2,823.2	4,446.8	7,149.4	15,287.5
2000.....	215.9	317.9	467.4	743.9	1,184.9	1,844.5	2,866.8	4,496.6	7,333.2	15,707.0
1999.....	210.8	310.5	460.0	747.2	1,193.5	1,895.5	2,923.9	4,542.8	7,370.9	15,718.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
White, male										
2004.....	272.8	412.3	607.8	855.8	1,343.7	2,055.9	3,219.3	5,123.2	8,214.4	15,250.7
2003.....	284.0	423.5	606.9	884.2	1,390.6	2,146.2	3,359.5	5,349.4	8,522.6	16,037.9
2002.....	287.2	420.4	601.2	902.0	1,402.7	2,196.1	3,469.7	5,469.3	8,660.8	16,473.2
2001.....	281.4	417.1	595.5	896.0	1,427.9	2,237.6	3,513.9	5,543.5	8,771.1	16,935.4
2000.....	279.4	416.1	588.3	925.1	1,465.1	2,285.1	3,593.1	5,620.3	9,092.5	17,716.4
1999.....	274.0	402.1	579.3	934.6	1,484.0	2,367.8	3,698.3	5,727.2	9,211.2	17,752.9
White, female										
2004.....	156.6	238.0	338.1	527.6	855.6	1,338.1	2,153.2	3,456.3	5,817.7	13,450.9
2003.....	160.7	235.8	344.9	543.9	884.5	1,379.6	2,212.3	3,592.1	6,048.6	14,240.6
2002.....	160.5	235.1	344.3	552.6	883.7	1,409.1	2,247.4	3,648.0	6,161.6	14,382.8
2001.....	155.9	229.4	350.1	558.4	905.3	1,426.7	2,263.9	3,669.9	6,169.3	14,597.6
2000.....	152.2	220.7	349.3	571.5	926.5	1,459.3	2,284.0	3,706.2	6,286.6	14,890.7
1999.....	147.4	219.9	343.6	569.0	925.9	1,483.5	2,308.5	3,711.2	6,284.5	14,900.6
Black, both sexes										
2004.....	386.6	608.3	910.1	1,271.1	1,813.5	2,509.7	3,582.0	5,266.0	7,493.0	13,338.5
2003.....	410.4	632.9	940.3	1,295.1	1,889.9	2,595.8	3,754.9	5,421.2	7,745.3	13,976.7
2002.....	412.9	634.1	925.9	1,321.6	1,880.7	2,632.0	3,824.9	5,571.3	7,962.2	14,260.8
2001.....	428.4	644.1	928.5	1,328.9	1,891.7	2,738.8	3,904.2	5,660.2	8,028.9	14,550.8
2000.....	434.5	666.6	938.7	1,347.0	1,929.0	2,769.0	4,019.7	5,782.6	8,161.0	14,714.3
1999.....	448.6	668.3	929.9	1,399.3	1,969.2	2,813.7	4,148.0	5,878.3	8,169.4	14,757.9
Black, male										
2004.....	467.1	766.8	1,186.6	1,668.7	2,374.4	3,242.7	4,587.3	6,747.0	9,229.2	14,452.5
2003.....	507.1	800.8	1,227.5	1,688.7	2,454.5	3,334.8	4,843.9	6,978.2	9,827.8	14,903.4
2002.....	502.9	801.4	1,206.6	1,719.3	2,464.7	3,357.1	4,918.1	7,088.6	9,968.4	15,635.5
2001.....	527.8	830.2	1,201.4	1,718.2	2,424.6	3,497.3	5,053.2	7,255.7	10,256.3	16,439.9
2000.....	549.1	853.0	1,227.4	1,776.2	2,462.1	3,513.8	5,203.9	7,410.0	10,405.7	16,791.0
1999.....	572.7	885.6	1,220.1	1,822.6	2,549.7	3,599.1	5,336.3	7,538.1	10,433.8	17,304.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
Black, female										
2004.....	315.7	470.1	676.4	943.8	1,376.3	1,965.5	2,899.8	4,388.5	6,578.2	12,896.9
2003.....	325.1	486.7	697.2	971.3	1,449.4	2,047.3	3,020.5	4,495.8	6,654.1	13,616.7
2002.....	333.6	489.0	688.0	995.2	1,423.7	2,093.9	3,091.2	4,664.6	6,917.8	13,734.2
2001.....	340.6	482.8	696.2	1,009.5	1,472.5	2,178.4	3,138.8	4,707.8	6,894.3	13,849.9
2000.....	333.3	505.0	692.6	995.2	1,508.3	2,219.3	3,233.3	4,801.6	7,030.5	13,941.3
1999.....	339.3	480.9	683.9	1,053.5	1,514.0	2,231.6	3,359.6	4,878.6	7,047.2	13,805.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

I. Certain infectious and parasitic diseases (A00-B99)

All races 3/, both sexes

2004.....	18.2	22.8	27.2	26.5	29.9	40.2	60.3	91.6	142.7	249.5
2003.....	18.9	24.0	26.4	25.9	29.7	39.6	60.2	94.7	141.6	255.2
2002.....	19.6	24.7	25.1	24.0	28.8	41.5	60.8	94.0	140.5	252.8
2001.....	19.7	23.2	22.6	22.9	28.7	39.4	57.2	87.5	130.0	246.3
2000.....	19.5	23.5	22.2	21.9	27.5	37.5	54.7	85.1	131.5	256.0
1999.....	20.9	22.9	21.1	22.2	26.5	40.3	56.4	86.9	132.5	267.9

All races 3/, male

2004.....	24.9	32.1	38.0	35.8	35.8	46.6	68.8	103.7	163.7	273.1
2003.....	26.4	34.3	37.6	34.6	35.3	46.3	68.0	106.2	162.9	269.1
2002.....	27.5	35.6	36.6	31.2	35.3	47.9	69.6	109.1	164.0	270.9
2001.....	28.1	33.9	32.3	30.4	34.0	44.8	65.5	102.4	153.8	269.7
2000.....	27.8	34.9	32.3	28.8	33.4	43.8	62.7	99.8	154.8	273.0
1999.....	30.6	33.9	30.2	29.1	32.0	47.4	65.4	102.6	157.8	290.1

All races 3/, female

2004.....	11.5	13.6	16.8	17.7	24.6	34.7	53.5	82.9	129.7	238.8
2003.....	11.6	13.9	15.7	17.8	24.6	33.8	54.0	86.5	128.6	249.1
2002.....	11.8	14.0	14.1	17.2	22.9	36.0	53.8	83.4	126.2	245.0
2001.....	11.4	12.8	13.2	15.8	23.9	34.8	50.6	77.1	115.8	236.5
2000.....	11.3	12.5	12.6	15.4	22.1	32.1	48.4	74.9	117.8	249.1
1999.....	11.3	12.4	12.4	15.8	21.5	34.2	49.3	75.9	117.8	258.9

White, both sexes

2004.....	12.0	16.4	20.8	21.1	24.7	34.8	55.2	86.4	136.4	242.1
2003.....	12.9	17.6	19.7	20.4	24.4	33.9	55.1	88.6	136.1	245.0
2002.....	12.8	17.8	18.4	18.7	23.5	35.8	55.1	87.3	133.9	242.3
2001.....	12.8	16.3	16.3	17.9	23.2	34.0	51.5	81.4	123.5	235.4
2000.....	12.8	16.2	15.9	16.9	22.0	31.9	49.0	77.9	123.0	242.9
1999.....	13.2	15.3	15.3	16.6	21.6	34.7	49.5	78.9	124.1	251.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

I. Certain infectious and parasitic diseases (A00-B99)

White, male

2004.....	17.5	23.6	29.4	28.0	28.8	40.0	62.3	98.1	155.4	266.6
2003.....	18.8	26.0	28.4	26.7	28.5	39.0	61.5	98.9	156.5	259.4
2002.....	18.9	26.5	26.6	24.1	28.5	40.7	63.0	100.0	155.3	260.3
2001.....	19.5	24.4	23.5	23.3	27.2	38.3	58.3	95.3	146.1	256.7
2000.....	18.9	24.6	22.9	21.9	26.3	36.8	56.4	90.4	145.0	258.5
1999.....	20.3	22.6	21.8	21.2	25.5	40.2	57.2	94.2	146.7	270.4

White, female

2004.....	6.5	9.2	12.3	14.6	20.8	30.2	49.4	77.9	124.5	231.1
2003.....	6.9	9.2	11.3	14.5	20.6	29.4	49.9	81.1	123.6	238.6
2002.....	6.6	9.3	10.4	13.5	18.9	31.5	48.6	78.2	120.8	234.5
2001.....	6.1	8.4	9.2	12.6	19.5	30.3	46.0	71.6	109.8	226.4
2000.....	6.6	7.9	9.0	12.1	18.0	27.7	43.1	69.1	110.0	236.5
1999.....	6.2	8.1	8.9	12.3	18.1	29.8	43.4	68.2	110.7	244.5

Black, both sexes

2004.....	62.4	71.4	80.3	76.5	80.1	91.4	115.8	161.6	240.1	375.3
2003.....	64.0	73.9	81.9	76.6	80.0	92.2	115.9	171.4	230.0	416.6
2002.....	68.7	78.3	81.7	73.6	76.1	94.7	122.7	181.9	238.1	413.1
2001.....	69.7	77.3	75.6	69.7	79.2	89.6	119.4	166.2	226.7	404.7
2000.....	68.7	80.8	77.5	66.3	74.5	89.9	119.3	177.5	253.3	436.7
1999.....	76.9	83.1	71.9	72.0	69.0	93.0	131.6	181.7	249.4	469.0

Black, male

2004.....	82.4	102.3	115.2	114.8	108.6	115.1	147.6	192.4	299.2	408.3
2003.....	87.5	105.3	121.3	115.4	105.7	122.6	144.5	205.4	285.0	456.7
2002.....	93.7	112.7	126.8	104.1	101.6	123.4	151.2	232.9	297.3	448.1
2001.....	96.3	114.3	111.6	101.8	102.5	111.6	151.1	204.0	278.5	470.9
2000.....	97.4	121.5	120.5	96.7	100.2	116.0	144.1	229.2	311.1	486.7
1999.....	112.3	128.3	108.8	103.4	95.7	120.0	166.2	213.7	325.2	537.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

I. Certain infectious and parasitic diseases (A00-B99)

Black, female

2004.....	44.7	44.5	50.7	45.0	58.0	73.8	94.2	143.4	209.0	362.3
2003.....	43.4	46.6	48.5	44.7	60.0	69.6	96.6	151.1	201.2	401.0
2002.....	46.7	48.4	43.5	48.5	56.2	73.4	103.5	151.5	207.3	399.6
2001.....	46.1	45.2	44.9	43.4	60.8	73.4	98.2	143.7	200.3	380.2
2000.....	43.5	45.6	40.9	41.4	54.3	70.7	102.8	146.4	224.2	418.1
1999.....	45.7	44.0	40.7	46.4	48.1	73.1	108.7	162.5	211.7	443.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
 years years years years years years years years years and over

Intestinal infectious diseases (A00-A09)

All races 3/, both sexes

2004.....	0.1	0.2	0.3	0.6	1.3	2.2	4.7	9.4	17.5	34.4
2003.....	*	0.1	0.2	0.5	0.9	1.7	3.4	7.2	12.4	25.0
2002.....	*	0.1	0.2	0.4	0.7	1.4	2.7	5.1	9.8	21.0
2001.....	*	*	0.2	0.2	0.5	0.8	1.9	3.3	6.6	13.1
2000.....	*	*	0.1	0.3	0.4	0.9	1.6	2.8	5.4	11.9
1999.....	*	*	*	*	0.3	0.8	1.3	2.4	4.6	10.6

All races 3/, male

2004.....	*	0.2	0.3	0.6	1.4	2.2	4.3	9.4	17.9	32.2
2003.....	*	*	0.2	0.5	0.9	1.5	3.1	7.0	12.9	21.9
2002.....	*	*	*	0.4	0.8	1.6	3.1	5.1	10.3	20.4
2001.....	*	*	*	*	0.4	0.8	2.0	3.6	6.5	12.2
2000.....	*	*	*	0.4	0.4	1.0	1.7	3.0	5.9	11.5
1999.....	*	*	*	*	*	0.8	1.3	2.7	5.1	9.7

All races 3/, female

2004.....	*	*	0.3	0.6	1.2	2.2	4.9	9.3	17.3	35.4
2003.....	*	*	0.2	0.5	1.0	1.8	3.6	7.2	12.1	26.4
2002.....	*	*	*	0.5	0.6	1.3	2.3	5.1	9.5	21.3
2001.....	*	*	0.2	*	0.6	0.8	1.8	3.1	6.6	13.4
2000.....	*	*	*	*	*	0.8	1.6	2.7	5.1	12.0
1999.....	*	*	*	*	*	0.8	1.3	2.2	4.4	10.9

White, both sexes

2004.....	*	0.1	0.3	0.6	1.3	2.3	4.9	9.9	18.4	36.7
2003.....	*	0.1	0.2	0.5	0.9	1.8	3.6	7.3	13.2	26.3
2002.....	*	0.1	0.2	0.4	0.6	1.6	2.7	5.3	10.4	22.1
2001.....	*	*	0.2	0.2	0.5	0.8	2.0	3.5	6.7	13.8
2000.....	*	*	*	0.3	0.4	0.9	1.7	3.0	5.8	12.3
1999.....	*	*	*	*	0.3	0.8	1.3	2.5	4.8	11.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other intestinal infectious diseases (A01-A08)

All races 3/, both sexes

2004.....	0.1	0.2	0.3	0.6	1.2	2.2	4.5	9.1	17.1	33.6
2003.....	*	0.1	0.2	0.5	0.9	1.6	3.3	6.9	12.1	24.2
2002.....	*	0.1	0.2	0.4	0.7	1.4	2.6	5.0	9.4	20.2
2001.....	*	*	0.2	0.2	0.5	0.8	1.8	3.2	6.4	12.6
2000.....	*	*	0.1	0.3	0.3	0.8	1.5	2.7	5.3	11.3
1999.....	*	*	*	*	0.3	0.7	1.1	2.2	4.3	9.9

All races 3/, male

2004.....	*	0.2	0.3	0.6	1.3	2.2	4.3	9.2	17.5	31.7
2003.....	*	*	0.2	0.5	0.8	1.5	3.1	6.7	12.6	21.1
2002.....	*	*	*	0.4	0.7	1.5	3.0	5.0	9.9	19.7
2001.....	*	*	*	*	0.4	0.8	2.0	3.4	6.4	11.9
2000.....	*	*	*	0.4	*	1.0	1.6	2.9	5.8	10.6
1999.....	*	*	*	*	*	0.7	1.1	2.6	4.9	9.3

All races 3/, female

2004.....	*	*	0.3	0.6	1.2	2.1	4.7	9.0	16.9	34.5
2003.....	*	*	0.2	0.5	0.9	1.7	3.4	7.0	11.8	25.5
2002.....	*	*	*	0.4	0.6	1.3	2.2	5.0	9.1	20.4
2001.....	*	*	*	*	0.6	0.8	1.7	3.0	6.4	13.0
2000.....	*	*	*	*	*	0.7	1.5	2.6	5.0	11.5
1999.....	*	*	*	*	*	0.7	1.2	1.9	4.0	10.2

White, both sexes

2004.....	*	0.1	0.3	0.6	1.3	2.2	4.8	9.6	18.0	35.9
2003.....	*	0.1	0.2	0.5	0.9	1.7	3.5	7.0	12.9	25.4
2002.....	*	0.1	0.2	0.4	0.6	1.5	2.6	5.2	10.1	21.1
2001.....	*	*	0.2	0.2	0.5	0.7	1.9	3.3	6.5	13.3
2000.....	*	*	*	0.3	0.3	0.8	1.6	2.8	5.6	11.7
1999.....	*	*	*	*	0.3	0.7	1.2	2.2	4.5	10.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intestinal infections due to other specified organisms (A04,A07-A08)

All races 3/, both sexes

2004.....	0.1	0.2	0.3	0.6	1.2	2.1	4.5	9.1	17.0	33.5
2003.....	*	0.1	0.2	0.4	0.8	1.6	3.2	6.8	12.0	23.9
2002.....	*	0.1	0.2	0.4	0.7	1.3	2.5	4.9	9.3	20.2
2001.....	*	*	0.2	0.2	0.5	0.7	1.8	3.1	6.2	12.4
2000.....	*	*	*	0.2	0.3	0.8	1.5	2.6	5.1	11.1
1999.....	*	*	*	*	0.3	0.6	1.1	2.0	4.2	9.7

All races 3/, male

2004.....	*	*	0.3	0.5	1.3	2.1	4.2	9.1	17.4	31.4
2003.....	*	*	0.2	0.4	0.8	1.5	3.0	6.6	12.6	20.6
2002.....	*	*	*	0.4	0.7	1.4	2.9	5.0	9.7	19.6
2001.....	*	*	*	*	0.4	0.7	1.9	3.3	6.1	11.8
2000.....	*	*	*	0.3	*	0.9	1.5	2.8	5.7	10.5
1999.....	*	*	*	*	*	0.6	1.0	2.5	4.7	9.1

All races 3/, female

2004.....	*	*	0.3	0.6	1.2	2.1	4.7	9.0	16.8	34.4
2003.....	*	*	0.2	0.4	0.9	1.7	3.3	6.9	11.7	25.4
2002.....	*	*	*	0.4	0.6	1.2	2.2	4.9	9.1	20.4
2001.....	*	*	*	*	0.5	0.8	1.7	2.9	6.2	12.7
2000.....	*	*	*	*	*	0.7	1.5	2.5	4.8	11.4
1999.....	*	*	*	*	*	0.7	1.1	1.7	4.0	9.9

White, both sexes

2004.....	*	0.1	0.3	0.6	1.3	2.2	4.8	9.6	17.9	35.8
2003.....	*	*	0.2	0.4	0.8	1.7	3.4	7.0	12.8	25.2
2002.....	*	*	0.2	0.4	0.6	1.4	2.6	5.2	9.9	21.1
2001.....	*	*	0.2	0.2	0.4	0.7	1.9	3.2	6.3	13.1
2000.....	*	*	*	0.3	0.3	0.8	1.5	2.7	5.4	11.5
1999.....	*	*	*	*	0.3	0.7	1.1	2.1	4.4	10.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Tuberculosis (A16-A19)

All races 3/, both sexes

2004.....	0.1	0.1	0.2	0.2	0.3	0.6	1.0	1.1	1.5	2.4
2003.....	0.1	0.2	0.3	0.4	0.3	0.5	0.8	1.4	2.0	2.3
2002.....	0.1	0.2	0.2	0.3	0.5	0.9	0.9	1.5	2.3	2.6
2001.....	0.1	0.2	0.3	0.4	0.5	0.6	1.0	1.4	2.0	2.8
2000.....	0.1	0.2	0.3	0.2	0.5	0.7	1.0	1.8	2.4	2.7
1999.....	0.1	0.3	0.3	0.5	0.6	0.7	1.2	1.8	2.4	4.2

All races 3/, male

2004.....	0.2	0.2	0.3	0.3	0.6	0.8	1.4	1.2	2.2	3.8
2003.....	*	0.4	0.4	0.6	0.4	0.6	1.2	2.0	2.7	3.1
2002.....	0.2	0.2	0.4	0.5	0.7	1.2	1.2	1.8	3.2	4.0
2001.....	*	0.3	0.4	0.7	0.7	0.9	1.5	1.7	2.4	4.4
2000.....	*	0.4	0.5	0.4	0.7	1.0	1.1	2.3	3.8	4.6
1999.....	0.2	0.4	0.5	0.8	0.9	1.1	1.7	2.4	3.7	7.1

All races 3/, female

2004.....	*	*	*	*	*	*	0.6	1.0	1.1	1.7
2003.....	*	*	*	*	*	0.4	0.5	1.0	1.7	1.9
2002.....	*	*	*	*	0.4	0.5	0.6	1.3	1.7	2.1
2001.....	*	*	*	*	*	0.4	0.7	1.1	1.8	2.2
2000.....	*	*	*	*	*	0.5	0.9	1.4	1.6	2.0
1999.....	*	*	*	*	*	0.4	0.7	1.5	1.7	3.0

White, both sexes

2004.....	*	*	0.1	*	0.2	0.3	0.7	0.8	1.2	1.7
2003.....	*	0.1	0.1	0.3	0.2	0.3	0.7	1.0	1.7	1.8
2002.....	*	*	0.1	0.2	0.4	0.5	0.6	1.2	1.8	2.1
2001.....	*	0.1	0.1	0.3	0.3	0.5	0.7	1.1	1.4	2.2
2000.....	*	*	0.2	*	0.3	0.5	0.6	1.2	1.7	2.2
1999.....	*	0.2	0.2	0.3	0.4	0.6	0.8	1.3	2.0	3.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Respiratory tuberculosis (A16)

All races 3/, both sexes

2004.....	0.1	0.1	0.2	0.2	0.3	0.4	0.8	0.8	1.1	2.2
2003.....	0.1	0.2	0.2	0.3	0.2	0.4	0.6	1.2	1.6	1.8
2002.....	0.1	0.1	0.2	0.2	0.4	0.6	0.7	1.2	2.0	2.3
2001.....	*	0.1	0.2	0.4	0.4	0.5	0.8	1.1	1.6	2.4
2000.....	0.1	0.1	0.2	0.2	0.3	0.5	0.8	1.4	2.1	2.3
1999.....	0.1	0.2	0.2	0.4	0.5	0.6	0.9	1.5	2.1	3.7

All races 3/, male

2004.....	*	*	0.3	*	0.4	0.6	1.2	0.9	1.7	3.7
2003.....	*	0.3	0.3	0.5	*	0.5	1.0	1.7	2.2	2.5
2002.....	0.2	*	0.3	0.4	0.5	1.0	0.9	1.3	2.8	3.5
2001.....	*	0.2	0.3	0.6	0.7	0.7	1.2	1.5	2.0	3.8
2000.....	*	0.3	0.4	0.3	0.6	0.9	0.9	1.9	3.2	4.1
1999.....	*	0.3	0.4	0.6	0.9	0.9	1.4	2.0	3.3	6.4

All races 3/, female

2004.....	*	*	*	*	*	*	0.5	0.7	0.8	1.5
2003.....	*	*	*	*	*	*	*	0.8	1.3	1.5
2002.....	*	*	*	*	*	*	0.5	1.1	1.5	1.8
2001.....	*	*	*	*	*	*	0.5	0.8	1.4	1.8
2000.....	*	*	*	*	*	*	0.6	1.1	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	1.2	1.4	2.6

White, both sexes

2004.....	*	*	*	*	*	0.3	0.5	0.6	0.9	1.6
2003.....	*	*	*	0.2	*	*	0.5	0.9	1.4	1.4
2002.....	*	*	*	*	0.3	0.4	0.5	0.9	1.5	1.8
2001.....	*	*	*	0.2	0.3	0.4	0.5	0.9	1.2	1.8
2000.....	*	*	0.2	*	*	0.3	0.5	1.0	1.5	1.9
1999.....	*	0.1	0.1	0.2	0.4	0.5	0.6	1.1	1.7	3.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Zoonotic and other bacterial diseases (A20-A49)

All races 3/, both sexes

2004.....	2.5	4.4	7.3	11.0	17.2	26.0	43.1	68.3	108.7	194.8
2003.....	2.7	4.5	6.9	11.1	17.6	26.5	43.0	71.4	112.4	209.5
2002.....	2.5	4.5	6.7	10.6	16.8	28.1	45.1	73.3	113.9	210.5
2001.....	2.4	4.3	6.6	10.4	16.3	26.6	42.6	70.3	108.3	213.7
2000.....	2.6	4.2	6.5	9.9	15.9	25.2	40.0	66.9	109.2	222.5
1999.....	2.5	4.1	6.1	9.6	15.2	26.5	40.5	67.5	108.5	231.2

All races 3/, male

2004.....	2.5	4.8	8.0	12.2	18.0	28.5	49.0	77.9	126.8	214.7
2003.....	2.7	5.0	7.5	12.2	18.8	30.2	47.7	80.4	128.2	223.2
2002.....	2.7	5.1	7.4	11.3	18.6	30.2	50.7	84.9	132.3	222.5
2001.....	2.6	4.7	7.2	11.3	17.6	29.4	48.7	82.3	127.8	233.5
2000.....	2.7	4.8	7.6	10.7	17.4	28.2	45.0	77.3	127.0	233.1
1999.....	2.9	4.6	6.7	10.3	16.9	30.0	46.3	79.3	127.7	246.7

All races 3/, female

2004.....	2.5	3.9	6.7	9.9	16.4	23.9	38.3	61.4	97.6	185.8
2003.....	2.7	4.1	6.3	10.0	16.5	23.2	39.2	65.0	102.8	203.5
2002.....	2.3	4.0	6.0	10.0	15.2	26.3	40.7	65.1	102.7	205.3
2001.....	2.2	3.9	6.0	9.5	15.1	24.1	37.8	62.0	96.5	205.4
2000.....	2.5	3.6	5.5	9.1	14.6	22.7	36.1	59.6	98.7	218.2
1999.....	2.1	3.6	5.5	9.0	13.7	23.6	35.9	59.3	97.2	225.0

White, both sexes

2004.....	1.9	3.5	6.0	9.5	14.7	22.9	39.4	63.7	102.6	186.1
2003.....	2.2	3.7	5.5	9.4	15.2	22.9	39.0	66.6	107.3	198.2
2002.....	2.1	3.8	5.3	8.9	14.1	24.7	41.0	67.6	107.5	199.4
2001.....	1.9	3.4	5.3	8.5	14.0	23.4	38.4	65.0	102.8	202.9
2000.....	2.2	3.2	5.3	8.2	13.3	21.8	36.0	60.8	101.6	209.6
1999.....	2.0	3.1	4.9	7.8	13.0	23.1	35.6	60.9	100.8	216.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

White, male

2004.....	2.0	3.8	6.7	10.4	15.2	25.2	44.5	72.9	119.0	207.4
2003.....	2.2	4.2	6.0	10.1	16.2	25.9	43.3	74.9	122.5	212.7
2002.....	2.2	4.3	5.9	9.6	15.9	26.3	46.3	77.5	124.3	211.9
2001.....	2.2	3.7	5.8	9.1	15.3	25.7	43.5	76.2	121.5	221.0
2000.....	2.3	3.7	6.0	9.0	14.8	24.4	41.1	70.0	119.3	219.5
1999.....	2.3	3.5	5.4	8.2	13.9	25.9	40.6	72.4	117.9	230.1

White, female

2004.....	1.9	3.1	5.4	8.6	14.1	20.9	35.2	56.9	92.4	176.5
2003.....	2.2	3.3	5.0	8.7	14.2	20.1	35.5	60.7	97.9	191.9
2002.....	1.9	3.3	4.8	8.3	12.4	23.3	36.6	60.5	97.3	194.0
2001.....	1.7	3.2	4.8	8.0	12.7	21.4	34.4	57.1	91.5	195.4
2000.....	2.0	2.7	4.6	7.5	12.0	19.5	31.9	54.3	91.0	205.6
1999.....	1.7	2.8	4.5	7.4	12.1	20.6	31.7	52.8	90.6	210.5

Black, both sexes

2004.....	6.7	11.0	18.2	26.6	41.7	59.5	88.2	134.2	202.7	328.8
2003.....	6.6	11.2	18.6	26.9	43.1	62.8	90.6	137.8	197.7	379.6
2002.....	6.1	10.1	17.8	28.1	42.6	63.2	94.6	151.5	212.7	376.3
2001.....	5.9	11.0	17.9	28.6	40.5	59.8	92.3	143.8	198.0	373.1
2000.....	6.1	11.5	17.9	25.4	41.3	61.3	89.6	150.3	221.7	402.3
1999.....	6.6	11.7	16.6	26.7	37.7	61.3	98.2	151.0	219.6	428.2

Black, male

2004.....	7.2	12.6	20.0	31.8	46.7	67.8	108.1	160.6	254.0	352.7
2003.....	6.9	12.9	21.2	34.1	48.2	78.1	105.1	164.8	241.4	416.0
2002.....	6.8	10.9	21.3	30.2	47.1	73.1	110.5	191.1	265.2	402.0
2001.....	6.1	12.2	20.1	35.2	43.7	69.9	114.1	175.6	243.2	435.7
2000.....	6.1	13.1	23.0	29.0	46.2	71.3	101.0	188.1	256.3	441.7
1999.....	8.1	13.7	19.5	31.2	48.4	75.6	120.1	176.7	286.0	472.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Zoonotic and other bacterial diseases (A20-A49)

Black, female

2004.....	6.3	9.6	16.6	22.4	37.7	53.2	74.7	118.6	175.8	319.4
2003.....	6.4	9.8	16.3	21.0	39.1	51.5	80.9	121.8	174.9	365.4
2002.....	5.5	9.4	14.9	26.3	39.0	55.9	84.0	127.9	185.4	366.4
2001.....	5.7	9.9	16.0	23.2	37.9	52.4	77.8	124.8	175.0	349.8
2000.....	6.1	10.1	13.6	22.4	37.4	54.0	82.1	127.6	204.2	387.6
1999.....	5.3	9.9	14.1	23.0	29.3	50.8	83.7	135.6	186.7	411.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
All races 3/, both sexes										
2004.....	2.3	4.1	6.9	10.3	16.2	24.9	41.2	65.1	103.6	186.7
2003.....	2.6	4.2	6.5	10.3	16.7	25.2	40.9	68.4	107.8	202.5
2002.....	2.3	4.2	6.2	10.0	16.0	27.0	43.1	70.0	109.4	203.0
2001.....	2.3	4.0	6.2	9.8	15.5	25.3	40.9	67.3	103.9	205.9
2000.....	2.5	3.9	6.1	9.4	15.2	24.2	38.4	64.0	105.0	215.7
1999.....	2.2	3.7	5.6	9.0	14.4	24.7	38.3	63.8	102.9	220.7
All races 3/, male										
2004.....	2.4	4.5	7.6	11.4	17.1	26.9	46.6	74.5	120.9	203.9
2003.....	2.5	4.6	7.1	11.3	17.7	28.7	45.1	76.7	122.5	213.8
2002.....	2.5	4.7	6.8	10.6	17.8	29.1	48.2	80.6	127.0	213.9
2001.....	2.5	4.3	6.7	10.6	16.6	27.8	46.7	78.6	122.3	224.3
2000.....	2.6	4.4	7.0	10.1	16.7	26.9	43.2	73.9	121.7	223.0
1999.....	2.6	4.2	6.2	9.7	15.7	27.7	43.7	74.8	120.6	233.5
All races 3/, female										
2004.....	2.3	3.7	6.4	9.3	15.5	23.2	36.8	58.4	92.9	179.0
2003.....	2.7	3.8	6.0	9.4	15.7	22.2	37.5	62.5	98.8	197.5
2002.....	2.2	3.8	5.6	9.5	14.5	25.2	39.0	62.5	98.7	198.3
2001.....	2.1	3.8	5.7	9.1	14.4	23.1	36.2	59.5	92.9	198.3
2000.....	2.3	3.4	5.2	8.7	13.8	21.8	34.6	57.1	95.2	212.7
1999.....	1.9	3.3	5.1	8.4	13.1	22.1	34.1	56.2	92.5	215.5
White, both sexes										
2004.....	1.8	3.2	5.7	8.8	13.8	21.9	37.6	60.7	97.5	178.1
2003.....	2.1	3.4	5.2	8.8	14.3	21.7	36.9	63.7	102.5	191.1
2002.....	1.9	3.6	5.0	8.4	13.5	23.6	39.2	64.4	103.1	191.9
2001.....	1.8	3.2	4.9	8.1	13.2	22.2	36.8	62.1	98.5	195.3
2000.....	2.0	3.0	4.9	7.8	12.6	20.7	34.5	58.0	97.4	203.0
1999.....	1.7	2.8	4.5	7.2	12.1	21.3	33.6	57.6	95.5	205.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Septicemia (A40-A41)

White, male

2004.....	1.8	3.5	6.4	9.7	14.4	23.7	42.5	69.8	113.4	196.5
2003.....	2.0	3.8	5.7	9.3	15.3	24.6	40.8	71.3	116.7	203.3
2002.....	2.0	4.0	5.4	8.9	15.3	25.2	44.0	73.4	119.0	202.9
2001.....	2.1	3.3	5.3	8.5	14.4	24.4	41.6	72.5	116.0	211.8
2000.....	2.2	3.5	5.5	8.4	14.0	23.1	39.4	66.7	114.0	209.8
1999.....	2.0	3.1	4.9	7.6	12.8	23.7	38.2	68.3	111.6	217.4

White, female

2004.....	1.8	2.9	5.1	8.0	13.3	20.2	33.5	54.1	87.7	169.9
2003.....	2.2	3.0	4.7	8.2	13.4	19.2	33.7	58.2	93.7	185.8
2002.....	1.9	3.1	4.5	7.8	11.8	22.3	35.2	58.0	93.4	187.1
2001.....	1.6	3.0	4.5	7.7	12.2	20.4	32.9	54.8	87.9	188.3
2000.....	1.9	2.5	4.3	7.2	11.3	18.7	30.6	51.9	87.6	200.2
1999.....	1.4	2.4	4.1	6.9	11.5	19.2	29.9	50.1	86.1	201.4

Black, both sexes

2004.....	6.2	10.3	17.3	25.5	39.9	57.9	85.1	128.0	196.1	320.3
2003.....	6.3	10.5	17.5	25.3	41.5	61.0	88.0	133.4	193.4	372.0
2002.....	5.5	9.5	17.0	27.2	40.9	62.0	90.6	146.5	208.8	367.0
2001.....	5.6	10.6	16.9	27.1	38.4	57.4	89.6	139.1	193.1	364.8
2000.....	5.8	11.0	16.9	24.3	40.2	59.7	86.6	146.4	216.3	392.6
1999.....	6.2	11.0	15.5	25.4	36.3	57.9	94.6	142.8	209.5	412.3

Black, male

2004.....	6.7	12.0	19.0	30.4	44.5	65.2	102.3	153.8	244.6	341.7
2003.....	6.4	11.8	19.7	31.8	46.1	74.9	100.2	160.2	235.5	404.5
2002.....	6.1	10.3	20.0	29.0	45.1	71.6	105.5	183.7	259.8	393.5
2001.....	5.7	11.6	19.1	33.3	41.5	66.3	111.1	171.0	236.8	424.4
2000.....	5.8	12.4	21.3	27.8	45.1	68.9	97.0	182.9	250.4	427.8
1999.....	7.7	12.8	18.1	29.7	46.5	71.4	115.0	166.2	266.8	453.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
Black, female										
2004.....	5.7	8.9	15.8	21.5	36.4	52.5	73.4	112.7	170.5	311.9
2003.....	6.3	9.4	15.6	20.0	37.9	50.8	79.8	117.5	171.4	359.3
2002.....	5.0	8.9	14.4	25.7	37.6	54.9	80.7	124.3	182.2	356.9
2001.....	5.4	9.7	15.0	22.1	36.0	50.9	75.3	120.1	170.9	342.7
2000.....	5.8	9.7	13.2	21.5	36.3	52.8	79.7	124.5	199.1	379.5
1999.....	5.0	9.4	13.3	21.9	28.3	47.9	81.0	128.7	181.1	396.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

All races 3/, both sexes

2004.....	0.1	0.3	0.3	0.7	0.9	1.1	1.9	3.1	5.1	7.8
2003.....	0.1	0.3	0.3	0.7	0.9	1.2	2.1	3.0	4.6	7.0
2002.....	0.2	0.2	0.4	0.5	0.7	1.0	2.0	3.2	4.3	7.4
2001.....	0.1	0.2	0.4	0.5	0.8	1.2	1.6	2.9	4.3	7.5
2000.....	0.1	0.2	0.4	0.5	0.7	1.0	1.5	2.8	4.0	6.7
1999.....	0.2	0.3	0.4	0.6	0.8	1.8	2.2	3.5	5.4	10.3

All races 3/, male

2004.....	*	0.3	0.4	0.8	0.9	1.5	2.3	3.4	5.9	10.8
2003.....	*	0.4	0.4	0.9	1.0	1.5	2.6	3.7	5.6	9.3
2002.....	0.2	0.3	0.5	0.7	0.8	1.1	2.4	4.2	5.2	8.5
2001.....	*	0.3	0.5	0.6	1.0	1.5	1.9	3.6	5.4	8.9
2000.....	*	0.3	0.6	0.6	0.8	1.4	1.8	3.3	5.2	9.9
1999.....	0.2	0.4	0.4	0.6	1.1	2.2	2.6	4.4	7.0	13.1

All races 3/, female

2004.....	*	0.2	0.3	0.5	0.9	0.7	1.5	2.9	4.7	6.4
2003.....	*	0.2	0.3	0.5	0.8	0.9	1.6	2.5	4.0	5.9
2002.....	*	0.2	0.3	0.4	0.7	1.0	1.6	2.5	3.8	6.9
2001.....	*	*	0.3	0.3	0.7	1.0	1.5	2.4	3.6	6.9
2000.....	*	*	*	0.3	0.7	0.8	1.3	2.4	3.4	5.3
1999.....	0.2	0.3	0.4	0.6	0.6	1.4	1.8	2.9	4.5	9.2

White, both sexes

2004.....	*	0.2	0.3	0.6	0.9	1.0	1.8	2.9	5.1	7.7
2003.....	*	0.3	0.2	0.6	0.8	1.1	2.0	2.9	4.7	7.0
2002.....	0.1	0.2	0.3	0.5	0.6	1.0	1.8	3.1	4.3	7.3
2001.....	*	0.2	0.3	0.4	0.7	1.1	1.5	2.8	4.2	7.4
2000.....	*	0.2	0.3	0.4	0.7	1.0	1.4	2.7	4.0	6.4
1999.....	0.2	0.3	0.4	0.5	0.8	1.6	2.0	3.1	5.1	9.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

White, male

2004.....	*	0.3	0.3	0.7	0.8	1.4	2.0	3.1	5.6	10.9
2003.....	*	0.3	*	0.8	0.9	1.3	2.4	3.6	5.7	9.3
2002.....	*	0.3	0.4	0.6	0.6	1.1	2.2	4.1	5.1	8.7
2001.....	*	0.3	0.4	0.5	0.9	1.3	1.8	3.6	5.4	8.9
2000.....	*	0.2	0.5	0.6	0.7	1.2	1.6	3.1	5.2	9.5
1999.....	*	0.3	0.4	0.5	1.1	2.0	2.4	4.0	6.2	12.6

White, female

2004.....	*	*	*	0.5	0.9	0.7	1.6	2.7	4.7	6.3
2003.....	*	0.2	0.2	0.5	0.8	0.9	1.7	2.4	4.1	6.0
2002.....	*	*	0.3	0.4	0.6	1.0	1.4	2.4	3.7	6.7
2001.....	*	*	*	*	0.6	0.9	1.4	2.1	3.5	6.8
2000.....	*	*	*	*	0.6	0.7	1.2	2.4	3.2	5.1
1999.....	*	*	0.3	0.5	0.6	1.3	1.8	2.5	4.5	8.8

Black, both sexes

2004.....	*	*	*	*	1.7	*	3.0	6.3	6.7	8.2
2003.....	*	*	*	1.3	*	*	2.6	4.4	*	7.6
2002.....	*	*	*	*	*	*	4.0	5.0	*	9.2
2001.....	*	*	1.0	1.4	*	2.3	2.7	4.5	*	8.0
2000.....	*	*	*	*	*	*	2.9	3.9	*	9.7
1999.....	*	*	*	*	*	3.4	3.5	8.0	10.1	15.9

Black, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	10.0	19.2	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)										
Black, female										
2004.....	*	*	*	*	*	*	*	5.9	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	9.4
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	6.9	*	14.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Viral diseases (A80-B34)

All races 3/, both sexes

2004.....	14.5	16.1	16.4	11.7	8.4	7.5	6.5	6.5	6.6	7.8
2003.....	15.1	17.2	16.2	11.4	8.1	7.4	8.0	7.6	7.5	9.1
2002.....	15.9	18.4	15.9	10.6	8.3	7.9	7.8	7.8	7.7	10.1
2001.....	16.5	17.5	14.2	10.3	9.1	8.1	7.6	7.3	7.0	8.3
2000.....	16.1	18.1	14.2	10.1	8.7	7.9	8.0	8.1	8.1	9.6
1999.....	17.2	17.1	12.9	9.9	7.5	8.2	8.1	8.0	8.7	11.6

All races 3/, male

2004.....	20.9	24.1	25.1	18.6	12.5	10.5	8.1	7.7	6.7	9.0
2003.....	22.2	26.1	25.4	17.8	11.8	9.7	10.5	9.2	9.6	10.2
2002.....	23.2	28.2	25.5	16.3	12.1	10.7	9.2	10.0	9.1	11.7
2001.....	24.4	27.3	22.9	16.1	12.5	10.0	9.1	8.3	8.9	9.3
2000.....	24.0	28.4	22.7	15.7	12.4	10.2	9.5	10.7	9.9	10.8
1999.....	26.2	27.0	20.8	15.5	10.5	10.7	9.8	9.3	10.9	13.3

All races 3/, female

2004.....	8.2	8.3	8.0	5.3	4.7	4.9	5.2	5.5	6.5	7.2
2003.....	8.1	8.4	7.3	5.4	4.7	5.4	6.0	6.4	6.2	8.7
2002.....	8.8	9.0	6.7	5.2	4.8	5.4	6.7	6.3	6.9	9.3
2001.....	8.7	8.1	5.9	4.8	6.0	6.4	6.5	6.7	5.9	7.9
2000.....	8.3	8.1	6.0	4.8	5.3	6.0	6.9	6.3	7.0	9.1
1999.....	8.3	7.5	5.3	4.7	4.7	6.0	6.7	7.1	7.4	10.9

White, both sexes

2004.....	9.0	10.7	11.3	8.2	5.9	5.7	5.3	5.9	6.2	7.8
2003.....	9.6	11.8	11.2	7.9	5.6	5.5	6.9	6.9	7.0	9.3
2002.....	9.8	12.5	10.8	7.5	6.0	5.9	6.5	7.0	7.4	10.3
2001.....	10.2	11.7	9.5	7.4	6.4	6.3	6.5	6.8	6.6	8.4
2000.....	10.0	12.0	9.3	6.9	6.2	6.2	6.8	7.6	7.8	9.6
1999.....	10.4	10.9	8.7	6.6	5.5	6.4	6.8	7.4	8.6	11.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Viral diseases (A80-B34)

White, male

2004.....	14.2	16.7	17.9	13.0	8.6	7.9	6.5	7.2	6.2	8.8
2003.....	15.2	18.8	18.0	12.3	8.1	7.0	8.8	8.1	8.9	10.3
2002.....	15.4	19.9	17.4	11.6	8.4	7.8	7.5	8.9	8.6	11.7
2001.....	16.5	19.0	15.9	11.7	8.5	7.8	7.9	7.7	8.3	9.7
2000.....	15.8	19.6	15.3	10.8	8.6	7.6	7.6	9.9	9.3	10.9
1999.....	16.8	17.4	14.3	10.2	7.9	8.3	8.1	8.6	10.9	12.7

White, female

2004.....	3.8	4.7	4.8	3.5	3.4	3.8	4.4	5.0	6.2	7.3
2003.....	4.0	4.8	4.4	3.6	3.3	4.2	5.3	6.1	5.8	8.9
2002.....	4.1	5.1	4.3	3.6	3.8	4.1	5.8	5.7	6.7	9.7
2001.....	4.0	4.5	3.2	3.3	4.3	5.0	5.4	6.1	5.6	7.9
2000.....	4.3	4.5	3.5	3.2	4.0	5.0	6.1	5.9	6.8	9.1
1999.....	4.0	4.4	3.1	3.2	3.4	4.9	5.7	6.6	7.2	11.1

Black, both sexes

2004.....	53.7	57.6	58.0	44.1	31.7	22.5	15.2	10.7	9.9	7.3
2003.....	55.5	58.9	58.0	43.7	30.4	22.9	17.6	13.1	12.7	7.9
2002.....	60.5	65.2	59.0	38.4	27.5	24.2	18.4	16.3	9.4	7.4
2001.....	62.3	63.5	54.4	36.3	32.7	21.8	15.9	12.9	10.0	6.1
2000.....	60.6	66.4	55.8	36.7	27.8	21.0	19.3	13.7	11.3	8.8
1999.....	66.5	66.5	49.7	39.3	23.0	22.5	19.6	12.7	9.0	11.5

Black, male

2004.....	72.4	85.6	88.9	73.4	52.5	36.8	24.0	12.7	*	*
2003.....	78.3	87.4	92.8	72.9	49.4	36.4	28.6	18.7	16.2	*
2002.....	83.9	97.6	98.4	63.1	45.7	39.6	27.7	24.3	*	*
2001.....	87.8	97.7	86.4	60.2	51.5	31.5	21.9	17.2	*	*
2000.....	88.6	104.1	91.6	61.9	46.0	33.8	30.8	19.8	18.5	*
1999.....	100.2	108.7	82.0	66.3	35.9	33.2	28.8	15.7	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral diseases (A80-B34)										
Black, female										
2004.....	37.1	33.3	31.9	20.0	15.4	11.9	9.2	9.4	8.7	*
2003.....	35.5	34.1	28.6	19.7	15.6	12.8	10.2	9.8	10.8	*
2002.....	39.9	37.1	25.6	18.1	13.2	12.8	12.1	11.5	*	*
2001.....	39.7	33.9	27.1	16.7	17.9	14.6	11.9	10.3	8.2	*
2000.....	35.9	33.7	25.3	16.1	13.4	11.5	11.8	10.0	*	8.6
1999.....	36.8	30.1	22.2	17.3	13.0	14.6	13.5	10.9	8.6	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Herpesviral (herpes simplex) infections (B00)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	0.4	0.6	0.6
2003.....	*	*	*	*	*	*	0.2	*	0.4	0.7
2002.....	*	*	*	*	*	*	0.4	*	0.5	0.5
2001.....	*	*	*	*	*	*	0.3	0.3	0.4	0.5
2000.....	*	*	*	*	*	*	*	0.3	0.5	*
1999.....	*	*	*	*	*	*	0.2	0.3	*	0.5

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	0.7	*
2003.....	*	*	*	*	*	*	*	*	*	0.9
2002.....	*	*	*	*	*	*	0.5	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	*	*	*	*	*	*	*	0.4	0.6	0.7
2003.....	*	*	*	*	*	*	0.3	*	0.5	0.8
2002.....	*	*	*	*	*	*	0.4	*	0.5	0.6
2001.....	*	*	*	*	*	*	0.3	0.3	0.4	0.5
2000.....	*	*	*	*	*	*	*	0.4	0.5	*
1999.....	*	*	*	*	*	*	0.3	0.3	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Zoster (herpes zoster)(B02)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	*	0.4	1.9
2003.....	*	*	*	*	*	*	*	*	0.4	2.2
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	1.6
2000.....	*	*	*	*	*	*	*	*	*	1.7
1999.....	*	*	*	*	*	*	*	*	0.6	2.0

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	2.1
2003.....	*	*	*	*	*	*	*	*	*	1.8
2002.....	*	*	*	*	*	*	*	*	*	1.5
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	1.8
2003.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	1.8
2000.....	*	*	*	*	*	*	*	*	*	1.9
1999.....	*	*	*	*	*	*	*	*	*	2.4

White, both sexes

2004.....	*	*	*	*	*	*	*	*	0.4	2.0
2003.....	*	*	*	*	*	*	*	*	*	2.3
2002.....	*	*	*	*	*	*	*	*	*	1.8
2001.....	*	*	*	*	*	*	*	*	*	1.8
2000.....	*	*	*	*	*	*	*	*	*	1.8
1999.....	*	*	*	*	*	*	*	*	0.6	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Viral hepatitis (B15-B19)

All races 3/, both sexes

2004.....	2.0	4.3	6.5	4.9	3.5	3.3	3.6	3.6	3.6	2.7
2003.....	2.1	4.8	6.3	4.5	3.1	3.4	4.3	4.3	4.1	2.4
2002.....	2.3	5.6	6.1	4.3	3.7	3.9	4.5	4.7	4.4	3.9
2001.....	2.6	5.3	5.2	4.1	3.8	4.3	4.9	5.1	4.6	3.8
2000.....	2.5	5.2	4.8	3.7	3.8	4.3	4.9	5.4	5.2	4.0
1999.....	2.3	4.4	3.8	3.3	3.3	4.5	5.1	5.3	5.4	4.7

All races 3/, male

2004.....	2.8	6.4	9.9	7.3	4.5	4.1	3.7	3.8	3.3	3.0
2003.....	2.9	7.1	9.7	6.6	3.9	3.7	4.8	4.7	5.0	2.8
2002.....	3.2	8.5	9.5	6.2	4.6	4.3	4.7	5.7	4.8	5.0
2001.....	3.7	8.1	8.2	5.9	4.5	4.7	4.9	5.1	5.5	4.4
2000.....	3.5	8.0	7.4	5.2	4.7	4.6	5.0	6.5	6.2	5.7
1999.....	3.5	6.7	5.9	4.7	3.8	4.9	5.4	6.1	7.3	6.9

All races 3/, female

2004.....	1.3	2.3	3.3	2.5	2.6	2.6	3.5	3.4	3.7	2.5
2003.....	1.2	2.6	3.1	2.5	2.4	3.1	3.9	4.0	3.5	2.3
2002.....	1.4	2.8	2.8	2.5	2.9	3.4	4.4	4.1	4.2	3.3
2001.....	1.5	2.5	2.4	2.5	3.3	3.9	4.9	5.0	4.1	3.5
2000.....	1.5	2.5	2.4	2.2	3.1	4.1	4.9	4.6	4.7	3.3
1999.....	1.2	2.2	1.7	2.1	2.8	4.1	4.9	4.7	4.2	3.9

White, both sexes

2004.....	2.1	4.3	6.2	4.2	3.1	3.0	3.2	3.3	3.1	2.5
2003.....	2.1	4.9	6.0	3.8	2.7	2.9	3.8	4.0	3.8	2.3
2002.....	2.2	5.5	5.8	3.8	3.2	3.4	4.1	4.5	4.3	3.7
2001.....	2.6	5.2	4.7	3.7	3.5	3.8	4.4	4.9	4.4	3.6
2000.....	2.4	4.9	4.3	3.0	3.4	3.9	4.6	5.3	5.0	3.9
1999.....	2.2	4.2	3.3	2.9	2.9	4.0	4.6	4.9	5.3	4.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral hepatitis (B15-B19)

White, male

2004.....	2.9	6.4	9.4	6.4	4.0	3.7	3.4	3.6	2.8	2.7
2003.....	2.9	7.2	9.2	5.7	3.4	3.3	4.3	4.2	4.8	2.8
2002.....	3.1	8.3	9.1	5.5	3.9	4.0	4.3	5.5	4.8	4.9
2001.....	3.7	8.1	7.6	5.4	4.1	4.4	4.6	5.2	5.6	4.5
2000.....	3.4	7.6	6.6	4.4	4.2	4.0	4.6	6.4	5.9	5.7
1999.....	3.3	6.1	5.2	4.1	3.6	4.6	5.1	5.8	7.7	6.5

White, female

2004.....	1.3	2.3	3.0	2.2	2.4	2.4	3.0	3.1	3.3	2.4
2003.....	1.2	2.6	2.9	2.0	2.0	2.6	3.5	3.8	3.2	2.1
2002.....	1.3	2.6	2.7	2.2	2.6	2.8	3.9	3.8	4.0	3.2
2001.....	1.4	2.4	2.0	2.1	2.9	3.3	4.2	4.6	3.7	3.2
2000.....	1.4	2.3	2.0	1.8	2.7	3.7	4.5	4.5	4.4	3.2
1999.....	1.1	2.2	1.5	1.7	2.3	3.5	4.3	4.3	3.9	3.7

Black, both sexes

2004.....	2.0	5.2	10.3	10.2	6.5	4.2	5.1	3.9	6.7	*
2003.....	2.7	5.2	10.1	9.4	6.3	5.1	5.7	6.5	5.3	*
2002.....	2.8	7.2	9.0	7.7	6.3	6.1	6.2	6.0	*	*
2001.....	3.3	6.8	9.4	7.3	6.0	5.1	6.2	6.1	*	*
2000.....	3.1	7.3	8.8	6.9	4.6	5.2	6.3	5.7	7.1	*
1999.....	3.5	7.0	6.7	5.7	4.2	6.6	7.2	6.8	*	6.4

Black, male

2004.....	2.2	8.3	16.0	16.6	9.9	6.8	6.5	*	*	*
2003.....	3.6	7.6	16.2	14.9	7.5	5.4	6.9	*	*	*
2002.....	3.8	10.5	14.6	12.4	8.1	8.0	*	*	*	*
2001.....	4.7	10.1	13.9	10.2	7.3	*	*	*	*	*
2000.....	4.2	11.0	13.9	10.4	5.2	6.8	7.4	*	*	*
1999.....	5.1	11.9	11.2	7.6	4.5	5.5	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Viral hepatitis (B15-B19)

Black, female

2004.....	1.8	2.5	5.5	4.9	3.9	*	*	*	*	*
2003.....	1.9	3.2	5.0	4.8	5.3	4.9	4.9	5.7	*	*
2002.....	1.9	4.4	4.2	3.9	4.8	4.6	6.3	5.8	*	*
2001.....	2.0	3.9	5.5	4.9	5.0	5.4	6.7	7.2	*	*
2000.....	2.1	4.0	4.4	4.0	4.1	4.0	5.5	*	*	*
1999.....	2.0	2.8	2.8	4.2	4.0	7.5	7.7	6.6	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All races 3/, both sexes

2004.....	12.3	11.5	9.6	6.3	4.2	3.1	1.5	1.1	0.5	*
2003.....	12.8	12.1	9.5	6.4	4.1	2.8	1.9	1.0	0.4	*
2002.....	13.4	12.6	9.6	5.9	4.0	2.9	1.4	1.0	0.5	*
2001.....	13.7	12.1	8.7	5.8	4.5	2.7	1.4	0.9	0.4	*
2000.....	13.5	12.7	9.0	5.9	4.1	2.6	1.7	0.9	*	*
1999.....	14.6	12.4	8.7	5.9	3.5	2.7	1.5	0.7	*	*

All races 3/, male

2004.....	17.9	17.5	14.9	10.5	7.0	5.1	2.6	1.7	*	*
2003.....	19.1	18.8	15.3	10.7	6.9	4.6	3.3	1.8	1.0	*
2002.....	19.8	19.4	15.7	9.8	6.8	5.0	2.6	1.7	*	*
2001.....	20.5	19.0	14.4	9.8	7.1	4.2	2.7	1.8	*	*
2000.....	20.4	20.3	15.0	10.1	7.0	4.5	2.9	1.8	*	*
1999.....	22.6	20.0	14.5	10.2	5.9	4.8	2.7	1.3	*	*

All races 3/, female

2004.....	6.7	5.7	4.5	2.3	1.5	1.4	0.6	0.6	*	*
2003.....	6.6	5.6	3.9	2.5	1.6	1.3	0.7	*	*	*
2002.....	7.2	5.9	3.6	2.3	1.5	1.0	0.5	*	*	*
2001.....	7.0	5.4	3.3	2.0	2.0	1.4	0.4	*	*	*
2000.....	6.7	5.3	3.4	2.0	1.5	1.0	0.7	*	*	*
1999.....	6.9	5.0	3.1	1.9	1.2	1.0	0.6	*	*	*

White, both sexes

2004.....	6.8	6.1	4.8	3.3	2.0	1.5	0.8	0.6	*	*
2003.....	7.4	6.7	4.9	3.5	2.1	1.3	1.0	0.6	*	*
2002.....	7.3	6.8	4.6	3.3	2.2	1.3	0.6	0.5	*	*
2001.....	7.5	6.4	4.5	3.3	2.0	1.2	0.8	0.5	*	*
2000.....	7.5	6.9	4.7	3.4	2.1	1.3	0.7	0.5	*	*
1999.....	8.1	6.5	4.9	3.1	1.9	1.4	0.7	0.4	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

White, male

2004.....	11.2	10.1	8.2	5.9	3.7	2.7	1.5	1.2	*	*
2003.....	12.1	11.3	8.4	6.0	3.6	2.2	2.0	1.3	*	*
2002.....	12.0	11.4	8.0	5.8	3.8	2.4	1.2	0.8	*	*
2001.....	12.6	10.8	8.0	5.9	3.4	2.1	1.7	1.0	*	*
2000.....	12.3	11.7	8.3	6.0	3.7	2.5	1.3	1.0	*	*
1999.....	13.4	11.0	8.6	5.6	3.5	2.7	1.2	0.8	*	*

White, female

2004.....	2.4	2.1	1.5	0.8	0.4	0.5	*	*	*	*
2003.....	2.6	2.1	1.3	1.1	0.6	0.5	*	*	*	*
2002.....	2.6	2.2	1.3	0.9	0.7	*	*	*	*	*
2001.....	2.4	2.0	1.0	0.9	0.7	0.5	*	*	*	*
2000.....	2.7	2.1	1.2	0.9	0.7	*	*	*	*	*
1999.....	2.8	2.0	1.2	0.7	*	*	*	*	*	*

Black, both sexes

2004.....	51.4	52.1	47.4	33.4	24.6	17.6	8.5	6.3	*	*
2003.....	52.3	53.4	47.2	33.9	23.8	17.2	11.0	4.9	*	*
2002.....	57.3	57.4	49.7	30.2	20.7	17.5	10.5	7.0	*	*
2001.....	58.8	56.5	44.7	28.9	26.2	16.3	8.5	6.4	*	*
2000.....	57.3	58.7	46.6	29.3	22.0	15.1	11.6	6.2	*	*
1999.....	62.8	59.0	42.6	32.7	18.3	15.4	11.2	4.8	*	*

Black, male

2004.....	69.9	76.9	72.4	56.1	42.0	29.6	15.2	9.6	*	*
2003.....	74.0	79.5	75.7	57.6	41.3	30.0	20.0	*	*	*
2002.....	79.6	86.3	83.5	50.0	37.0	31.4	19.4	14.7	*	*
2001.....	82.8	87.5	72.2	49.9	43.5	26.4	15.5	13.0	*	*
2000.....	84.1	92.7	77.4	51.0	39.5	26.3	22.4	11.8	*	*
1999.....	94.9	96.1	70.6	57.5	30.6	27.3	21.4	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Black, female

2004.....	35.0	30.5	26.3	14.7	10.9	8.6	*	*	*	*
2003.....	33.1	30.7	23.1	14.3	10.2	7.7	4.9	*	*	*
2002.....	37.6	32.4	21.0	13.9	7.9	7.2	4.5	*	*	*
2001.....	37.5	29.7	21.2	11.7	12.6	8.8	*	*	*	*
2000.....	33.7	29.2	20.4	11.5	8.2	6.9	4.4	*	*	*
1999.....	34.6	26.9	18.8	12.5	8.7	6.5	4.5	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

All races 3/, both sexes

2004.....	5.0	4.6	3.7	2.4	1.3	1.2	0.5	0.4	*	*
2003.....	4.8	4.7	3.8	2.4	1.5	0.9	0.7	*	*	*
2002.....	5.3	5.1	3.9	2.3	1.4	1.0	0.4	0.4	*	*
2001.....	5.3	4.9	3.4	2.2	1.5	0.9	0.6	0.3	*	*
2000.....	5.4	5.2	3.5	2.0	1.4	0.9	0.6	0.4	*	*
1999.....	5.8	5.2	3.6	1.9	1.1	0.9	0.6	0.4	*	*

All races 3/, male

2004.....	7.3	6.9	5.5	3.9	2.1	2.0	0.9	0.7	*	*
2003.....	7.1	7.5	6.2	3.8	2.5	1.5	1.0	*	*	*
2002.....	7.9	7.8	6.5	3.7	2.2	1.7	0.9	0.7	*	*
2001.....	8.0	7.6	5.6	3.7	2.3	1.2	1.1	*	*	*
2000.....	7.9	8.2	5.8	3.3	2.4	1.7	1.1	*	*	*
1999.....	8.9	8.2	5.9	3.4	1.9	1.6	1.0	0.7	*	*

All races 3/, female

2004.....	2.7	2.4	1.9	0.9	0.5	0.6	*	*	*	*
2003.....	2.4	2.0	1.6	1.1	0.6	0.4	*	*	*	*
2002.....	2.7	2.4	1.4	0.9	0.7	0.4	*	*	*	*
2001.....	2.7	2.3	1.3	0.8	0.8	0.5	*	*	*	*
2000.....	2.9	2.2	1.2	0.9	0.5	*	*	*	*	*
1999.....	2.7	2.2	1.3	0.6	0.5	*	*	*	*	*

White, both sexes

2004.....	2.9	2.6	1.9	1.2	0.6	0.6	*	*	*	*
2003.....	3.0	2.8	1.9	1.3	0.8	0.5	0.3	*	*	*
2002.....	3.0	2.7	1.9	1.3	0.9	0.5	*	*	*	*
2001.....	3.1	2.7	1.7	1.3	0.7	0.4	0.3	*	*	*
2000.....	3.0	2.9	1.9	1.2	0.8	0.5	*	*	*	*
1999.....	3.2	2.6	2.1	1.1	0.7	0.5	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

White, male

2004.....	4.8	4.2	3.0	2.2	1.1	1.1	*	*	*	*
2003.....	4.9	4.7	3.3	2.1	1.3	0.8	0.6	*	*	*
2002.....	4.8	4.6	3.2	2.3	1.3	0.9	*	*	*	*
2001.....	5.0	4.6	3.0	2.3	1.2	0.6	0.6	*	*	*
2000.....	4.9	4.8	3.4	1.9	1.3	1.0	*	*	*	*
1999.....	5.2	4.4	3.6	1.9	1.2	0.9	*	*	*	*

White, female

2004.....	1.0	1.0	0.8	*	*	*	*	*	*	*
2003.....	1.1	0.9	0.6	0.6	*	*	*	*	*	*
2002.....	1.2	0.9	0.6	0.4	0.5	*	*	*	*	*
2001.....	1.1	0.9	0.5	0.4	*	*	*	*	*	*
2000.....	1.2	0.9	0.5	0.4	*	*	*	*	*	*
1999.....	1.2	0.9	0.6	*	*	*	*	*	*	*

Black, both sexes

2004.....	20.2	19.8	17.5	12.9	8.1	6.8	3.5	*	*	*
2003.....	17.8	19.0	19.2	12.5	8.5	5.0	4.1	*	*	*
2002.....	21.6	23.0	20.3	10.9	6.6	6.1	2.7	*	*	*
2001.....	21.9	22.2	17.5	10.7	8.5	5.2	3.6	*	*	*
2000.....	22.4	23.5	16.5	9.9	7.4	5.1	4.4	*	*	*
1999.....	24.5	25.1	16.8	10.1	5.8	4.6	4.3	*	*	*

Black, male

2004.....	27.6	28.5	26.1	20.8	13.5	11.1	*	*	*	*
2003.....	25.3	29.8	31.2	21.0	14.7	9.1	6.6	*	*	*
2002.....	31.3	34.8	35.1	17.7	11.1	9.5	*	*	*	*
2001.....	31.5	33.3	28.3	18.3	13.8	8.1	6.7	*	*	*
2000.....	31.4	36.9	28.0	16.3	13.3	9.4	9.0	*	*	*
1999.....	37.7	40.3	28.2	18.3	8.9	8.8	7.8	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

Black, female

2004.....	13.7	12.3	10.3	6.3	3.9	3.6	*	*	*	*
2003.....	11.2	9.5	9.1	5.4	3.8	*	*	*	*	*
2002.....	13.1	12.7	7.7	5.3	3.1	*	*	*	*	*
2001.....	13.5	12.6	8.3	4.5	4.3	*	*	*	*	*
2000.....	14.4	11.9	6.7	4.6	*	*	*	*	*	*
1999.....	12.8	12.0	7.2	3.3	3.3	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)

All races 3/, both sexes

2004.....	0.7	0.8	0.8	0.6	0.4	0.2	*	*	*	*
2003.....	0.9	0.8	0.6	0.6	0.3	0.3	*	*	*	*
2002.....	0.9	0.9	0.7	0.4	0.3	*	*	*	*	*
2001.....	1.0	0.8	0.6	0.6	0.4	0.2	*	*	*	*
2000.....	1.0	0.9	0.7	0.5	0.4	0.2	*	*	*	*
1999.....	1.1	0.9	0.7	0.6	0.3	*	*	*	*	*

All races 3/, male

2004.....	1.2	1.4	1.3	1.0	0.7	*	*	*	*	*
2003.....	1.5	1.2	1.0	0.9	0.5	0.4	*	*	*	*
2002.....	1.5	1.5	1.2	0.7	0.6	*	*	*	*	*
2001.....	1.7	1.2	1.1	0.9	0.8	*	*	*	*	*
2000.....	1.7	1.6	1.2	1.0	0.6	*	*	*	*	*
1999.....	1.8	1.5	1.2	1.1	0.5	*	*	*	*	*

All races 3/, female

2004.....	0.3	0.2	0.2	*	*	*	*	*	*	*
2003.....	0.3	0.4	0.3	*	*	*	*	*	*	*
2002.....	0.3	0.3	0.2	*	*	*	*	*	*	*
2001.....	0.3	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	0.2	0.2	*	*	*	*	*	*	*

White, both sexes

2004.....	0.5	0.6	0.4	0.4	0.3	*	*	*	*	*
2003.....	0.7	0.6	0.4	0.3	0.2	*	*	*	*	*
2002.....	0.6	0.6	0.4	0.3	0.2	*	*	*	*	*
2001.....	0.7	0.5	0.4	0.3	0.3	*	*	*	*	*
2000.....	0.7	0.7	0.5	0.3	0.2	*	*	*	*	*
1999.....	0.7	0.6	0.5	0.5	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)

White, male

2004.....	0.9	1.1	0.8	0.7	0.5	*	*	*	*	*
2003.....	1.2	0.9	0.7	0.6	*	*	*	*	*	*
2002.....	1.0	1.2	0.7	0.6	*	*	*	*	*	*
2001.....	1.3	0.9	0.8	0.6	0.5	*	*	*	*	*
2000.....	1.2	1.2	0.9	0.7	0.5	*	*	*	*	*
1999.....	1.3	1.1	0.9	0.9	*	*	*	*	*	*

White, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, both sexes

2004.....	2.6	2.4	3.3	2.3	*	*	*	*	*	*
2003.....	2.5	2.6	2.6	2.6	*	*	*	*	*	*
2002.....	3.1	3.1	3.0	1.3	*	*	*	*	*	*
2001.....	3.2	2.7	2.2	2.7	2.1	*	*	*	*	*
2000.....	3.5	3.1	2.7	1.9	*	*	*	*	*	*
1999.....	3.4	2.8	2.7	2.4	*	*	*	*	*	*

Black, male

2004.....	3.8	3.6	5.6	4.5	*	*	*	*	*	*
2003.....	3.8	3.1	4.0	4.2	*	*	*	*	*	*
2002.....	4.7	4.9	5.0	*	*	*	*	*	*	*
2001.....	5.3	4.0	3.8	4.5	4.3	*	*	*	*	*
2000.....	5.7	5.1	4.5	3.8	*	*	*	*	*	*
1999.....	6.0	5.1	4.4	3.6	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)

All races 3/, both sexes

2004.....	1.2	1.2	1.0	0.6	0.5	0.3	*	*	*	*
2003.....	1.3	1.2	0.8	0.6	0.5	0.3	*	*	*	*
2002.....	1.4	1.1	0.8	0.6	0.4	0.3	*	*	*	*
2001.....	1.2	1.1	0.8	0.6	0.5	0.4	*	*	*	*
2000.....	1.5	1.3	1.0	0.7	0.4	0.3	0.2	*	*	*
1999.....	1.6	1.2	0.9	0.7	0.4	0.3	*	*	*	*

All races 3/, male

2004.....	1.8	1.9	1.6	1.1	0.8	0.4	*	*	*	*
2003.....	2.1	1.9	1.5	1.1	0.8	0.5	*	*	*	*
2002.....	2.1	1.8	1.4	1.1	0.9	0.6	*	*	*	*
2001.....	1.9	1.8	1.4	1.1	0.8	0.6	*	*	*	*
2000.....	2.5	2.2	1.7	1.3	0.8	*	*	*	*	*
1999.....	2.6	2.1	1.6	1.2	0.7	0.5	*	*	*	*

All races 3/, female

2004.....	0.6	0.5	0.4	*	*	*	*	*	*	*
2003.....	0.5	0.5	0.2	*	*	*	*	*	*	*
2002.....	0.6	0.5	0.2	*	*	*	*	*	*	*
2001.....	0.5	0.4	0.3	*	*	*	*	*	*	*
2000.....	0.6	0.5	0.3	*	*	*	*	*	*	*
1999.....	0.6	0.4	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.7	0.8	0.6	0.3	0.3	*	*	*	*	*
2003.....	0.8	0.7	0.5	0.3	*	*	*	*	*	*
2002.....	0.9	0.8	0.5	0.4	0.3	*	*	*	*	*
2001.....	0.7	0.7	0.5	0.4	0.3	*	*	*	*	*
2000.....	1.0	0.7	0.6	0.5	*	*	*	*	*	*
1999.....	1.0	0.9	0.6	0.3	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
White, male										
2004.....	1.2	1.3	1.1	0.6	0.4	*	*	*	*	*
2003.....	1.3	1.3	1.0	0.6	*	*	*	*	*	*
2002.....	1.5	1.4	0.9	0.7	0.5	*	*	*	*	*
2001.....	1.2	1.2	0.9	0.8	0.4	*	*	*	*	*
2000.....	1.7	1.3	1.0	0.8	*	*	*	*	*	*
1999.....	1.7	1.6	1.1	0.6	*	*	*	*	*	*
White, female										
2004.....	0.2	0.3	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*
Black, both sexes										
2004.....	4.6	4.6	4.2	3.3	2.5	*	*	*	*	*
2003.....	5.0	4.7	3.5	3.1	3.0	*	*	*	*	*
2002.....	5.0	3.8	3.5	3.1	2.2	*	*	*	*	*
2001.....	5.0	4.4	3.5	2.8	3.0	2.2	*	*	*	*
2000.....	5.6	5.7	4.7	3.5	2.3	*	*	*	*	*
1999.....	6.5	4.2	3.7	4.0	2.0	*	*	*	*	*
Black, male										
2004.....	6.5	6.9	6.7	5.9	4.1	*	*	*	*	*
2003.....	8.2	7.2	5.8	5.8	5.6	*	*	*	*	*
2002.....	7.2	5.6	6.0	5.5	4.4	*	*	*	*	*
2001.....	7.6	7.0	5.5	4.7	4.3	*	*	*	*	*
2000.....	9.0	9.2	8.2	6.4	4.3	*	*	*	*	*
1999.....	10.1	7.0	6.2	7.1	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)										
All races 3/, both sexes										
2004.....	2.0	1.8	1.7	1.1	1.0	0.7	0.3	*	*	*
2003.....	3.0	3.0	2.3	1.6	1.1	0.8	0.5	0.3	*	*
2002.....	3.3	3.1	2.4	1.6	1.1	0.8	0.4	0.3	*	*
2001.....	3.3	3.0	2.2	1.4	1.3	0.7	0.4	0.3	*	*
2000.....	3.0	2.9	2.3	1.7	1.2	0.7	0.4	0.3	*	*
1999.....	3.3	2.9	2.2	1.7	1.1	0.9	0.4	*	*	*
All races 3/, male										
2004.....	2.8	2.8	2.6	1.7	1.6	1.1	*	*	*	*
2003.....	4.3	4.4	3.6	2.7	1.7	1.2	1.0	*	*	*
2002.....	4.4	4.6	3.8	2.6	2.0	1.5	0.5	*	*	*
2001.....	4.6	4.5	3.6	2.5	2.0	1.1	0.7	*	*	*
2000.....	4.4	4.4	3.6	2.8	1.8	1.1	0.7	0.7	*	*
1999.....	4.7	4.5	3.5	2.7	1.9	1.6	0.7	*	*	*
All races 3/, female										
2004.....	1.2	0.9	0.8	0.4	0.4	*	*	*	*	*
2003.....	1.8	1.7	1.0	0.6	0.5	0.4	*	*	*	*
2002.....	2.1	1.7	1.1	0.7	0.4	*	*	*	*	*
2001.....	2.1	1.6	0.9	0.5	0.6	*	*	*	*	*
2000.....	1.7	1.4	1.0	0.6	0.6	*	*	*	*	*
1999.....	1.9	1.3	0.9	0.7	0.4	*	*	*	*	*
White, both sexes										
2004.....	0.9	0.7	0.7	0.5	0.5	0.3	*	*	*	*
2003.....	1.5	1.3	1.1	0.8	0.5	0.3	0.3	*	*	*
2002.....	1.5	1.3	1.0	0.7	0.5	0.3	*	*	*	*
2001.....	1.6	1.3	0.9	0.7	0.5	0.3	*	*	*	*
2000.....	1.5	1.3	1.0	0.9	0.5	0.3	*	*	*	*
1999.....	1.5	1.2	1.0	0.7	0.6	0.4	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

White, male

2004.....	1.5	1.2	1.3	0.9	0.8	0.6	*	*	*	*
2003.....	2.3	2.2	1.8	1.4	0.8	0.5	*	*	*	*
2002.....	2.4	2.0	1.8	1.3	0.9	0.5	*	*	*	*
2001.....	2.5	2.1	1.7	1.3	1.0	0.5	*	*	*	*
2000.....	2.3	2.2	1.7	1.5	0.8	*	*	*	*	*
1999.....	2.4	2.0	1.7	1.2	1.1	0.8	*	*	*	*

White, female

2004.....	0.4	0.3	*	*	*	*	*	*	*	*
2003.....	0.6	0.5	0.3	*	*	*	*	*	*	*
2002.....	0.7	0.6	0.3	*	*	*	*	*	*	*
2001.....	0.7	0.6	0.3	*	*	*	*	*	*	*
2000.....	0.7	0.5	0.4	*	*	*	*	*	*	*
1999.....	0.7	0.5	*	*	*	*	*	*	*	*

Black, both sexes

2004.....	9.4	9.8	9.4	6.6	6.1	3.4	*	*	*	*
2003.....	14.3	15.7	12.5	9.0	6.8	5.5	3.6	*	*	*
2002.....	15.7	16.8	13.7	9.6	7.0	5.8	3.2	*	*	*
2001.....	15.7	15.5	12.9	7.8	8.0	4.1	*	*	*	*
2000.....	14.0	14.7	13.4	9.0	7.2	4.6	2.9	*	*	*
1999.....	16.2	15.5	13.1	10.4	5.7	5.6	3.2	*	*	*

Black, male

2004.....	12.6	15.5	14.6	10.6	9.9	6.1	*	*	*	*
2003.....	19.4	22.2	19.8	15.4	11.2	8.6	6.6	*	*	*
2002.....	20.1	25.2	21.6	15.5	12.5	11.2	*	*	*	*
2001.....	20.6	23.5	20.9	14.2	12.4	7.4	*	*	*	*
2000.....	20.1	22.9	21.4	15.1	11.6	8.1	*	*	*	*
1999.....	22.8	25.3	21.1	18.0	9.1	10.4	6.8	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

Black, female

2004.....	6.6	4.8	5.0	3.3	3.1	*	*	*	*	*
2003.....	9.7	10.1	6.4	3.7	3.3	*	*	*	*	*
2002.....	11.9	9.4	7.0	4.8	*	*	*	*	*	*
2001.....	11.4	8.6	6.1	2.6	4.5	*	*	*	*	*
2000.....	8.5	7.7	6.5	4.0	3.8	*	*	*	*	*
1999.....	10.4	7.2	6.3	4.3	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Unspecified human immunodeficiency virus (HIV) disease (B24)

All races 3/, both sexes

2004.....	3.4	3.1	2.5	1.6	1.1	0.7	0.3	*	*	*
2003.....	2.8	2.4	1.9	1.3	0.8	0.6	0.3	0.3	*	*
2002.....	2.6	2.4	1.8	1.0	0.7	0.5	0.3	*	*	*
2001.....	2.9	2.3	1.7	0.9	0.7	0.6	0.2	*	*	*
2000.....	2.6	2.4	1.6	1.0	0.7	0.5	0.3	*	*	*
1999.....	2.9	2.3	1.3	1.0	0.6	0.5	0.3	*	*	*

All races 3/, male

2004.....	4.8	4.5	3.8	2.7	1.9	1.1	0.6	*	*	*
2003.....	4.1	3.8	3.0	2.2	1.4	0.9	0.6	*	*	*
2002.....	3.9	3.7	2.9	1.7	1.2	0.8	0.6	*	*	*
2001.....	4.3	3.8	2.8	1.6	1.2	0.9	0.5	*	*	*
2000.....	4.0	3.9	2.6	1.7	1.4	0.9	0.5	*	*	*
1999.....	4.5	3.7	2.2	1.8	1.0	0.8	0.5	*	*	*

All races 3/, female

2004.....	1.9	1.7	1.2	0.7	0.3	*	*	*	*	*
2003.....	1.6	1.1	0.8	0.5	*	*	*	*	*	*
2002.....	1.4	1.2	0.7	0.4	*	*	*	*	*	*
2001.....	1.5	0.8	0.6	0.3	*	*	*	*	*	*
2000.....	1.3	1.0	0.7	*	*	*	*	*	*	*
1999.....	1.4	0.9	0.4	0.3	*	*	*	*	*	*

White, both sexes

2004.....	1.8	1.4	1.2	0.9	0.5	0.3	*	*	*	*
2003.....	1.5	1.2	0.9	0.7	0.4	*	*	*	*	*
2002.....	1.3	1.3	0.8	0.6	0.4	*	*	*	*	*
2001.....	1.5	1.1	0.9	0.5	0.2	*	*	*	*	*
2000.....	1.3	1.3	0.7	0.6	0.4	*	*	*	*	*
1999.....	1.6	1.1	0.8	0.5	0.3	0.3	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Unspecified human immunodeficiency virus (HIV) disease (B24)

White, male

2004.....	2.9	2.3	2.0	1.5	0.9	0.5	*	*	*	*
2003.....	2.4	2.1	1.7	1.3	0.8	*	*	*	*	*
2002.....	2.2	2.2	1.4	0.9	0.7	*	*	*	*	*
2001.....	2.6	1.9	1.6	0.9	*	*	*	*	*	*
2000.....	2.2	2.2	1.3	1.0	0.7	*	*	*	*	*
1999.....	2.7	1.9	1.4	0.9	0.6	0.5	*	*	*	*

White, female

2004.....	0.7	0.5	0.3	0.4	*	*	*	*	*	*
2003.....	0.5	0.4	*	*	*	*	*	*	*	*
2002.....	0.4	0.4	0.3	*	*	*	*	*	*	*
2001.....	0.4	0.3	*	*	*	*	*	*	*	*
2000.....	0.5	0.4	*	*	*	*	*	*	*	*
1999.....	0.5	0.3	*	*	*	*	*	*	*	*

Black, both sexes

2004.....	14.7	15.5	13.0	8.3	6.5	4.5	*	*	*	*
2003.....	12.7	11.4	9.3	6.7	4.1	3.9	*	*	*	*
2002.....	11.8	10.7	9.2	5.3	3.5	3.0	*	*	*	*
2001.....	12.8	11.7	8.6	4.8	4.7	3.7	*	*	*	*
2000.....	11.9	11.7	9.4	5.0	3.8	3.1	*	*	*	*
1999.....	12.2	11.3	6.2	5.9	3.5	2.2	*	*	*	*

Black, male

2004.....	19.5	22.4	19.5	14.3	11.8	7.5	*	*	*	*
2003.....	17.3	17.2	14.9	11.2	7.5	7.4	*	*	*	*
2002.....	16.3	15.8	15.8	9.1	6.5	5.2	*	*	*	*
2001.....	17.8	19.7	13.7	8.1	8.7	5.8	*	*	*	*
2000.....	17.8	18.5	15.2	9.4	7.9	5.2	*	*	*	*
1999.....	18.2	18.5	10.7	10.5	5.7	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified human immunodeficiency virus (HIV) disease (B24)										
Black, female										
2004.....	10.4	9.5	7.5	3.5	*	*	*	*	*	*
2003.....	8.6	6.4	4.6	3.0	*	*	*	*	*	*
2002.....	7.8	6.3	3.6	*	*	*	*	*	*	*
2001.....	8.4	4.8	4.1	*	*	*	*	*	*	*
2000.....	6.6	5.7	4.5	*	*	*	*	*	*	*
1999.....	6.9	5.1	2.4	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)

All races 3/, both sexes

2004.....	0.1	0.2	0.2	0.4	0.6	0.8	0.8	1.1	1.4	2.0
2003.....	0.2	0.2	0.2	0.4	0.6	0.8	1.2	1.2	1.4	2.8
2002.....	0.2	0.2	0.3	0.3	0.5	0.8	1.0	1.2	1.5	2.9
2001.....	0.1	0.1	0.2	0.3	0.6	0.9	0.9	0.9	1.2	2.2
2000.....	0.1	0.2	0.3	0.4	0.6	0.9	1.1	1.2	1.7	3.3
1999.....	0.2	0.2	0.4	0.6	0.6	0.8	1.1	1.5	2.1	4.0

All races 3/, male

2004.....	*	*	0.3	0.5	0.7	0.8	0.9	1.5	1.6	2.3
2003.....	0.2	0.2	0.3	0.4	0.8	0.9	1.4	1.3	1.7	3.0
2002.....	0.2	0.2	0.3	0.3	0.5	0.9	1.0	1.4	1.8	2.8
2001.....	0.2	*	0.2	0.3	0.7	0.9	1.0	1.0	1.5	2.4
2000.....	*	*	0.3	0.4	0.7	1.0	1.4	1.7	2.0	3.1
1999.....	*	0.2	0.4	0.5	0.7	0.8	1.4	1.4	2.0	3.9

All races 3/, female

2004.....	*	0.2	*	0.4	0.5	0.7	0.7	0.8	1.3	1.8
2003.....	0.2	*	*	0.4	0.4	0.7	0.9	1.2	1.3	2.7
2002.....	0.2	0.2	0.2	0.3	0.5	0.8	1.0	1.1	1.3	2.9
2001.....	*	*	0.2	0.3	0.5	0.9	0.8	0.8	1.0	2.1
2000.....	*	0.2	0.3	0.4	0.6	0.8	1.0	0.8	1.6	3.4
1999.....	*	*	0.4	0.6	0.6	0.8	0.9	1.5	2.1	4.0

White, both sexes

2004.....	*	0.2	0.2	0.4	0.6	0.8	0.9	1.2	1.5	2.0
2003.....	0.1	0.2	0.2	0.4	0.7	0.8	1.2	1.3	1.4	3.0
2002.....	0.2	0.2	0.3	0.3	0.5	0.9	1.1	1.2	1.6	3.1
2001.....	0.1	*	0.2	0.3	0.7	1.0	0.9	0.9	1.2	2.3
2000.....	0.1	0.2	0.3	0.5	0.6	1.0	1.2	1.2	1.8	3.4
1999.....	0.2	0.2	0.4	0.6	0.7	0.8	1.1	1.5	2.1	4.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

All races 3/, both sexes

2004.....	0.9	2.0	2.9	2.9	2.7	3.9	5.0	6.2	8.2	9.7
2003.....	0.9	1.9	2.8	2.6	2.8	3.5	5.0	7.0	7.1	9.0
2002.....	0.9	1.4	2.1	2.0	2.5	3.2	4.3	6.2	6.7	8.3
2001.....	0.6	1.1	1.2	1.6	2.3	3.3	4.0	5.1	5.9	8.2
2000.....	0.6	0.9	1.0	1.4	2.1	2.7	4.0	5.4	6.1	9.0
1999.....	1.0	1.4	1.7	2.0	2.8	4.1	5.3	7.0	8.2	10.0

All races 3/, male

2004.....	1.2	2.8	4.2	4.1	3.4	4.4	5.8	7.3	10.0	12.9
2003.....	1.2	2.7	3.9	3.5	3.4	4.1	5.4	7.6	9.2	10.4
2002.....	1.2	2.1	3.0	2.7	3.2	4.1	5.3	7.1	9.0	11.9
2001.....	0.9	1.6	1.6	2.1	2.6	3.6	4.1	6.5	8.0	10.1
2000.....	0.8	1.2	1.3	1.6	2.5	3.3	5.4	6.3	7.7	12.6
1999.....	1.3	1.7	2.1	2.2	3.2	4.8	6.2	8.9	10.3	13.1

All races 3/, female

2004.....	0.7	1.1	1.7	1.9	2.1	3.3	4.4	5.5	7.2	8.3
2003.....	0.6	1.2	1.7	1.7	2.2	3.1	4.8	6.6	5.8	8.3
2002.....	0.5	0.7	1.1	1.4	1.8	2.4	3.5	5.6	5.3	6.8
2001.....	0.4	0.6	0.9	1.1	2.0	3.1	3.9	4.2	4.7	7.4
2000.....	0.4	0.7	0.8	1.2	1.7	2.2	2.9	4.7	5.2	7.6
1999.....	0.8	1.1	1.4	1.8	2.4	3.5	4.6	5.6	6.9	8.8

White, both sexes

2004.....	0.9	2.0	3.0	2.7	2.5	3.6	4.8	6.0	7.9	9.6
2003.....	0.9	1.9	2.7	2.4	2.5	3.5	4.9	6.6	6.9	8.9
2002.....	0.8	1.3	1.9	1.7	2.3	3.1	4.2	6.2	6.6	8.3
2001.....	0.5	1.0	1.1	1.4	2.0	3.1	3.8	5.1	5.8	7.9
2000.....	0.5	0.9	0.9	1.3	1.8	2.5	3.8	5.3	6.0	8.9
1999.....	0.7	1.1	1.4	1.8	2.3	3.7	5.0	6.8	7.8	9.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

White, male

2004.....	1.1	2.9	4.2	3.7	3.2	4.3	5.7	7.0	9.8	13.2
2003.....	1.2	2.7	3.8	3.3	3.1	4.0	5.1	7.2	8.9	10.5
2002.....	1.1	2.0	2.8	2.3	3.0	4.0	5.2	6.9	8.9	11.8
2001.....	0.7	1.5	1.4	1.8	2.4	3.4	4.0	6.3	8.0	9.9
2000.....	0.7	1.1	1.2	1.6	2.1	3.1	5.2	5.9	7.5	12.0
1999.....	1.0	1.4	1.7	2.1	2.7	4.4	5.9	8.7	9.4	12.0

White, female

2004.....	0.7	1.1	1.7	1.8	2.0	3.0	4.1	5.3	6.7	7.9
2003.....	0.6	1.1	1.6	1.5	2.0	3.0	4.8	6.1	5.6	8.2
2002.....	0.4	0.7	1.0	1.2	1.7	2.3	3.4	5.6	5.2	6.7
2001.....	0.4	0.6	0.9	1.0	1.8	2.9	3.7	4.2	4.4	7.1
2000.....	0.3	0.6	0.7	1.1	1.5	2.1	2.8	4.9	5.0	7.6
1999.....	0.4	0.8	1.1	1.4	2.1	3.2	4.3	5.4	6.9	8.6

Black, both sexes

2004.....	1.5	2.0	2.8	4.2	4.4	5.3	5.6	8.0	11.1	13.3
2003.....	1.3	2.7	3.8	4.1	4.3	3.5	4.8	10.0	8.9	9.0
2002.....	1.5	2.2	3.8	4.8	3.3	3.5	4.0	7.0	5.5	9.5
2001.....	1.3	2.0	2.0	2.9	3.9	4.4	5.9	4.9	7.0	11.4
2000.....	1.3	1.9	2.2	2.5	3.7	3.9	5.3	5.9	9.0	10.9
1999.....	3.0	3.9	4.4	4.1	6.2	6.9	8.4	9.5	11.0	14.3

Black, male

2004.....	2.0	3.1	4.5	6.7	6.4	*	7.1	11.4	*	*
2003.....	1.5	3.3	5.0	5.3	5.8	*	6.6	9.6	*	*
2002.....	2.1	3.2	5.4	7.3	4.9	*	*	*	*	*
2001.....	2.1	3.0	2.9	3.6	4.7	5.6	*	*	*	*
2000.....	1.7	2.3	3.0	*	5.0	5.5	7.4	9.9	*	*
1999.....	3.0	4.2	5.1	*	7.6	7.8	9.8	10.9	18.3	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

Black, female

2004.....	*	*	*	*	2.9	5.9	4.6	5.9	10.6	13.0
2003.....	*	2.1	2.7	3.1	3.2	*	*	10.3	*	8.9
2002.....	*	*	2.5	2.7	*	*	*	6.6	*	*
2001.....	*	*	*	*	3.2	*	5.6	*	*	11.8
2000.....	*	*	*	*	*	*	*	*	*	9.0
1999.....	3.1	3.7	3.9	5.0	5.2	6.2	7.4	8.6	*	11.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mycoses (B35-B49)										
All races 3/, both sexes										
2004.....	0.2	0.3	0.3	0.6	0.8	1.0	1.3	1.7	2.4	2.5
2003.....	0.2	0.2	0.4	0.5	0.7	1.1	1.5	2.1	2.1	2.4
2002.....	0.2	0.3	0.5	0.6	0.8	1.1	1.7	2.3	2.7	2.5
2001.....	0.2	0.3	0.3	0.5	0.8	1.2	1.4	2.3	2.0	3.0
2000.....	0.2	0.3	0.5	0.7	1.0	1.2	1.7	2.3	2.1	3.2
1999.....	0.4	0.6	0.8	1.0	1.4	2.1	2.6	3.2	3.3	3.9
All races 3/, male										
2004.....	0.2	0.4	0.4	0.8	0.8	1.1	1.5	2.2	3.4	3.8
2003.....	0.3	0.3	0.4	0.6	0.9	1.1	1.7	2.7	3.2	3.3
2002.....	0.3	0.4	0.7	0.9	1.1	1.7	2.3	3.1	4.1	4.2
2001.....	0.3	0.5	0.4	0.7	1.0	1.4	1.5	3.2	2.9	3.8
2000.....	0.3	0.3	0.6	0.7	1.2	1.6	2.4	3.0	2.8	4.9
1999.....	0.5	0.6	0.8	1.2	1.7	2.4	3.4	4.4	4.9	6.6
All races 3/, female										
2004.....	*	*	0.3	0.4	0.7	0.8	1.2	1.3	1.8	1.9
2003.....	*	0.2	0.3	0.4	0.6	1.1	1.3	1.7	1.5	2.0
2002.....	*	*	0.3	0.4	0.6	0.7	1.2	1.6	1.9	1.7
2001.....	*	0.2	0.3	0.4	0.7	1.0	1.3	1.7	1.5	2.6
2000.....	0.2	0.3	0.4	0.6	0.8	1.0	1.2	1.8	1.7	2.6
1999.....	0.3	0.5	0.7	0.9	1.2	1.9	1.9	2.4	2.3	2.9
White, both sexes										
2004.....	0.1	0.2	0.2	0.5	0.7	0.9	1.2	1.5	2.2	2.4
2003.....	0.1	0.2	0.3	0.4	0.6	1.1	1.5	2.0	2.0	2.3
2002.....	0.2	0.2	0.3	0.5	0.7	1.0	1.6	2.1	2.7	2.4
2001.....	0.2	0.3	0.3	0.5	0.7	1.0	1.2	2.2	1.8	2.6
2000.....	0.2	0.3	0.4	0.6	0.9	1.2	1.6	2.2	2.1	3.0
1999.....	0.3	0.5	0.7	0.9	1.2	1.9	2.4	3.1	3.0	3.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Protozoal diseases (B50-B64)										
All races 3/, both sexes										
2004.....	0.1	*	0.1	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	0.1	0.1	*	*	*	*	*	*
2001.....	0.1	0.1	0.1	0.2	0.3	0.3	0.3	*	*	*
2000.....	0.1	0.1	*	0.1	0.2	*	0.3	*	*	*
1999.....	0.2	0.1	0.2	0.2	*	0.2	0.3	0.3	*	*
All races 3/, male										
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	0.6	*	*	*
1999.....	0.2	*	0.2	*	*	*	*	*	*	*
All races 3/, female										
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	0.4	0.3	*	*	*
2000.....	*	*	*	*	*	*	0.3	*	*	*
1999.....	0.1	*	*	*	*	*	0.3	0.3	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumocystosis (B59)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	0.1	*	*	*	*	*	*	*
2001.....	*	0.1	*	0.2	0.2	0.3	0.3	*	*	*
2000.....	*	0.1	*	*	0.2	*	0.3	*	*	*
1999.....	0.1	*	0.2	*	*	*	0.3	0.3	*	*

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	0.5	*	*	*
1999.....	*	*	0.2	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	0.3	0.3	*	*	*
2000.....	*	*	*	*	*	*	0.3	*	*	*
1999.....	0.1	*	*	*	*	*	0.3	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Sequelae of tuberculosis (B90)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	*	*	0.5
2003.....	*	*	*	*	*	*	*	*	0.4	0.5
2002.....	*	*	*	*	*	*	*	*	*	0.8
2001.....	*	*	*	*	*	*	*	*	0.5	0.6
2000.....	*	*	*	*	*	*	*	0.3	0.6	0.7
1999.....	*	*	*	*	*	*	0.3	0.4	1.0	0.8

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	0.9
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	1.0	0.8

White, both sexes

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	0.5
2002.....	*	*	*	*	*	*	*	*	*	0.9
2001.....	*	*	*	*	*	*	*	*	0.5	0.6
2000.....	*	*	*	*	*	*	*	*	0.5	0.7
1999.....	*	*	*	*	*	*	*	0.4	0.9	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Sequelae of poliomyelitis (B91)

All races 3/, both sexes

2004.....	*	*	*	*	0.2	0.4	0.6	0.7	1.0	1.4
2003.....	*	*	*	0.1	0.2	0.3	0.5	0.7	0.8	1.2
2002.....	*	*	*	0.1	0.2	0.4	0.6	0.9	0.9	1.3
2001.....	*	*	*	*	0.3	0.4	0.6	0.8	0.9	1.2
2000.....	*	*	*	0.2	0.2	0.3	0.5	0.8	0.8	1.2
1999.....	*	*	*	*	0.2	0.3	0.5	0.6	0.7	1.3

All races 3/, male

2004.....	*	*	*	*	*	*	0.6	0.8	1.4	1.9
2003.....	*	*	*	*	*	*	0.6	*	1.3	1.5
2002.....	*	*	*	*	*	0.5	0.5	0.7	1.1	1.9
2001.....	*	*	*	*	*	*	0.7	1.2	*	1.8
2000.....	*	*	*	*	*	*	0.6	0.8	*	*
1999.....	*	*	*	*	*	*	*	0.8	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	0.5	0.7	0.8	1.1
2003.....	*	*	*	*	*	*	0.5	0.8	*	1.1
2002.....	*	*	*	*	*	0.4	0.6	1.0	0.8	1.0
2001.....	*	*	*	*	*	0.4	0.6	0.6	0.9	1.0
2000.....	*	*	*	*	*	*	0.5	0.8	0.8	1.1
1999.....	*	*	*	*	*	*	0.5	0.5	0.7	1.2

White, both sexes

2004.....	*	*	*	*	0.2	0.4	0.6	0.8	1.1	1.5
2003.....	*	*	*	0.2	0.3	0.3	0.6	0.8	0.9	1.3
2002.....	*	*	*	0.2	0.2	0.5	0.6	1.0	1.0	1.4
2001.....	*	*	*	*	0.4	0.5	0.7	0.9	1.0	1.4
2000.....	*	*	*	0.2	0.2	0.4	0.6	0.9	0.9	1.4
1999.....	*	*	*	*	0.2	0.3	0.5	0.7	0.8	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

All races 3/, both sexes

2004.....	0.7	1.6	2.4	2.1	1.6	2.2	2.8	3.4	4.3	5.3
2003.....	0.7	1.5	2.3	1.8	1.6	1.9	2.7	3.9	3.7	4.8
2002.....	0.5	1.0	1.4	1.1	1.3	1.4	1.7	2.6	2.6	3.6
2001.....	0.3	0.6	0.7	0.7	0.8	1.3	1.4	1.5	2.3	3.2
2000.....	0.2	0.4	0.4	0.4	0.6	0.8	1.2	1.7	2.3	3.7
1999.....	0.5	0.7	0.7	0.7	1.0	1.4	1.6	2.4	3.0	3.9

All races 3/, male

2004.....	0.8	2.3	3.6	3.0	2.1	2.4	3.3	3.8	4.4	6.3
2003.....	0.9	2.2	3.4	2.6	1.9	2.4	2.8	4.1	4.3	4.8
2002.....	0.8	1.4	2.1	1.5	1.6	1.7	2.0	2.8	3.2	4.7
2001.....	0.5	0.9	1.0	1.0	0.9	1.3	1.3	1.5	3.2	3.4
2000.....	0.4	0.6	0.5	0.5	0.7	1.0	1.5	2.0	2.9	4.3
1999.....	0.6	0.9	0.9	0.7	1.0	1.7	1.6	2.9	3.7	3.8

All races 3/, female

2004.....	0.5	0.9	1.3	1.3	1.2	2.1	2.4	3.2	4.2	4.8
2003.....	0.5	0.9	1.2	1.0	1.4	1.5	2.7	3.8	3.3	4.8
2002.....	0.3	0.5	0.6	0.7	0.9	1.2	1.5	2.5	2.2	3.1
2001.....	0.2	0.3	0.4	0.4	0.7	1.2	1.5	1.5	1.8	3.2
2000.....	*	0.3	0.3	0.3	0.5	0.6	0.9	1.5	2.0	3.4
1999.....	0.3	0.5	0.5	0.7	0.9	1.1	1.6	2.1	2.6	3.9

White, both sexes

2004.....	0.7	1.7	2.6	2.1	1.6	2.1	2.7	3.4	4.1	5.1
2003.....	0.7	1.6	2.2	1.7	1.5	1.9	2.6	3.5	3.5	4.8
2002.....	0.5	1.0	1.4	0.9	1.2	1.4	1.8	2.6	2.6	3.5
2001.....	0.3	0.6	0.6	0.6	0.7	1.2	1.3	1.5	2.3	3.3
2000.....	0.2	0.5	0.4	0.3	0.5	0.7	1.2	1.8	2.2	3.6
1999.....	0.3	0.5	0.5	0.6	0.7	1.2	1.5	2.3	2.9	3.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

Black, female

2004.....	*	*	*	*	*	4.1	*	*	*	7.9
2003.....	*	*	1.8	*	*	*	*	5.7	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	1.4	2.1	2.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

II. Neoplasms (C00-D48)

All races 3/, both sexes

2004.....	45.1	88.1	157.6	265.0	433.4	647.1	909.1	1,199.4	1,472.0	1,731.0
2003.....	47.0	91.2	161.1	271.1	447.0	657.7	925.5	1,223.9	1,497.3	1,778.8
2002.....	48.5	92.1	163.6	280.5	452.5	673.5	949.9	1,235.8	1,504.6	1,802.6
2001.....	49.7	93.9	167.1	284.6	459.6	683.7	961.4	1,241.2	1,513.7	1,848.3
2000.....	49.8	95.0	168.5	292.8	470.0	701.2	969.0	1,251.3	1,555.9	1,904.3
1999.....	50.3	95.1	169.3	298.8	477.9	711.0	982.0	1,253.3	1,540.6	1,888.7

All races 3/, male

2004.....	40.2	87.5	170.5	293.9	499.6	768.6	1,112.1	1,538.1	1,955.7	2,448.4
2003.....	42.7	92.0	172.2	299.6	514.6	786.0	1,139.0	1,569.2	2,001.3	2,515.6
2002.....	43.9	91.4	173.9	313.2	526.0	809.1	1,177.8	1,591.4	2,010.6	2,592.2
2001.....	45.0	93.4	176.6	315.6	534.1	826.9	1,200.6	1,607.6	2,041.1	2,721.3
2000.....	45.3	95.1	176.6	325.4	543.2	848.0	1,211.6	1,622.8	2,115.8	2,819.6
1999.....	46.6	96.1	177.6	334.6	558.0	861.6	1,236.8	1,637.2	2,107.0	2,831.4

All races 3/, female

2004.....	49.9	88.8	145.2	237.8	373.1	541.3	745.6	956.0	1,174.0	1,408.2
2003.....	51.2	90.4	150.5	244.3	385.5	546.4	754.1	978.0	1,189.2	1,453.0
2002.....	53.1	92.8	153.8	249.8	385.7	556.4	767.8	984.7	1,196.9	1,460.0
2001.....	54.3	94.4	158.0	255.5	391.9	560.4	771.5	984.7	1,197.7	1,483.0
2000.....	54.1	95.0	160.7	262.4	403.7	575.3	777.8	992.7	1,225.7	1,531.6
1999.....	54.0	94.1	161.4	265.4	405.4	582.2	783.0	986.4	1,209.2	1,508.8

White, both sexes

2004.....	42.7	83.6	150.1	257.2	427.4	642.9	908.6	1,201.8	1,477.8	1,739.9
2003.....	44.8	86.1	153.3	264.3	440.7	654.6	925.3	1,224.7	1,499.4	1,784.4
2002.....	46.5	87.7	156.7	272.3	445.0	669.1	948.1	1,234.8	1,504.1	1,806.4
2001.....	47.3	89.3	160.7	276.2	453.3	676.3	956.5	1,237.9	1,507.6	1,849.1
2000.....	46.7	89.6	161.8	285.0	463.7	695.8	961.4	1,243.6	1,555.2	1,903.1
1999.....	47.7	89.8	162.1	289.3	468.4	703.7	974.3	1,242.7	1,537.1	1,884.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

II. Neoplasms (C00-D48)

White, male

2004.....	38.2	82.5	160.9	282.0	486.7	757.4	1,098.9	1,526.1	1,950.8	2,452.1
2003.....	40.8	86.4	162.0	288.9	501.8	775.0	1,129.8	1,556.3	1,993.4	2,517.5
2002.....	42.3	87.0	164.9	300.9	509.8	798.0	1,165.9	1,575.7	1,994.7	2,590.9
2001.....	42.9	88.3	167.6	302.3	521.3	810.7	1,183.4	1,588.2	2,014.4	2,707.4
2000.....	42.3	89.2	166.6	311.3	529.3	833.1	1,187.6	1,596.8	2,094.9	2,806.3
1999.....	44.2	89.4	167.0	320.0	540.8	844.6	1,216.4	1,608.8	2,087.0	2,810.7

White, female

2004.....	47.3	84.7	139.4	233.3	372.3	541.4	752.2	964.6	1,184.0	1,418.8
2003.....	49.0	85.8	144.8	240.7	384.1	548.3	757.9	984.9	1,194.9	1,459.6
2002.....	50.6	88.4	148.7	245.0	385.2	555.6	770.6	990.7	1,203.4	1,465.6
2001.....	51.8	90.3	154.0	251.2	390.6	558.3	772.7	989.8	1,201.4	1,489.8
2000.....	51.1	89.9	157.1	260.0	403.2	575.6	779.9	995.3	1,234.1	1,536.2
1999.....	51.2	90.3	157.3	260.1	401.8	580.9	781.9	985.6	1,212.5	1,511.8

Black, both sexes

2004.....	65.5	131.3	238.7	382.7	577.4	813.2	1,075.2	1,381.7	1,617.5	1,831.0
2003.....	67.5	140.6	245.9	386.7	595.7	813.9	1,095.5	1,408.1	1,680.6	1,910.9
2002.....	70.0	137.1	241.3	404.8	602.9	843.8	1,141.1	1,450.2	1,707.0	1,949.3
2001.....	71.6	139.2	244.2	409.2	597.0	869.3	1,175.3	1,457.5	1,759.5	2,005.1
2000.....	75.8	148.0	255.0	412.1	611.8	872.0	1,199.5	1,508.3	1,738.5	2,063.0
1999.....	74.6	146.6	258.7	434.4	641.8	892.8	1,209.9	1,531.6	1,742.9	2,070.3

Black, male

2004.....	57.5	135.6	278.2	462.5	738.9	1,047.9	1,461.9	1,960.9	2,387.5	2,810.4
2003.....	62.4	147.7	285.9	470.3	755.8	1,058.2	1,477.1	2,002.0	2,470.2	2,892.3
2002.....	61.7	138.8	274.3	491.8	784.4	1,092.0	1,552.3	2,077.8	2,567.7	3,055.5
2001.....	64.1	145.0	283.1	499.5	755.4	1,148.3	1,623.1	2,115.9	2,703.9	3,298.4
2000.....	71.7	154.0	297.3	517.2	779.4	1,152.9	1,678.6	2,200.5	2,722.0	3,340.9
1999.....	69.5	162.3	304.1	539.8	824.0	1,176.7	1,665.8	2,241.5	2,684.7	3,404.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

II. Neoplasms (C00-D48)

Black, female

2004.....	72.6	127.5	205.3	316.9	451.5	639.0	812.7	1,038.6	1,211.9	1,442.8
2003.....	72.0	134.4	212.1	318.0	470.8	632.5	838.1	1,055.1	1,266.9	1,529.6
2002.....	77.3	135.7	213.2	333.3	460.9	659.6	865.2	1,075.3	1,258.9	1,525.5
2001.....	78.3	134.3	211.1	335.1	472.5	663.3	877.1	1,064.5	1,278.4	1,525.3
2000.....	79.5	142.8	218.9	325.9	479.5	664.7	881.3	1,091.1	1,243.1	1,587.4
1999.....	79.1	133.2	220.2	348.4	498.8	682.4	907.4	1,103.9	1,276.1	1,571.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms (C00-C97)

All races 3/, both sexes

2004.....	44.3	86.7	155.6	261.5	427.6	637.8	892.4	1,172.1	1,424.8	1,653.3
2003.....	46.1	89.8	159.1	267.9	440.9	647.9	909.2	1,195.4	1,450.0	1,698.2
2002.....	47.5	90.6	161.5	276.8	447.0	663.8	933.6	1,207.0	1,458.3	1,723.9
2001.....	48.7	92.4	164.8	280.7	453.3	673.2	943.5	1,211.7	1,465.9	1,765.6
2000.....	48.8	93.6	166.2	289.0	463.4	690.6	951.6	1,222.5	1,505.3	1,819.4
1999.....	49.4	93.7	167.0	295.3	471.6	700.2	963.4	1,224.0	1,493.1	1,805.8

All races 3/, male

2004.....	39.2	86.1	168.2	289.9	492.5	756.9	1,091.6	1,503.3	1,894.3	2,349.5
2003.....	41.7	90.6	169.8	295.9	507.6	774.4	1,117.9	1,533.3	1,939.7	2,413.8
2002.....	42.8	89.8	171.7	308.6	519.7	796.8	1,157.8	1,554.1	1,951.5	2,491.1
2001.....	44.0	91.7	174.2	311.6	526.7	814.1	1,178.0	1,568.4	1,978.7	2,613.6
2000.....	44.4	93.5	173.9	321.3	535.7	834.7	1,190.4	1,586.0	2,050.4	2,710.7
1999.....	45.5	94.7	174.9	330.6	550.1	848.2	1,212.9	1,600.1	2,046.0	2,722.2

All races 3/, female

2004.....	49.2	87.4	143.5	234.8	368.5	534.2	732.0	934.2	1,135.5	1,340.1
2003.....	50.5	89.1	148.8	241.5	380.3	538.1	741.8	954.8	1,150.5	1,381.9
2002.....	52.2	91.4	151.8	246.8	381.1	548.9	754.4	961.9	1,158.4	1,391.1
2001.....	53.5	93.2	155.9	251.7	386.7	551.9	757.2	962.0	1,158.6	1,410.7
2000.....	53.2	93.7	158.8	258.9	398.0	567.2	763.5	969.3	1,183.8	1,456.4
1999.....	53.3	92.7	159.5	262.3	400.6	573.7	768.4	962.5	1,169.5	1,436.5

White, both sexes

2004.....	42.0	82.2	148.2	253.8	421.6	633.6	891.5	1,174.1	1,429.3	1,659.7
2003.....	44.0	84.8	151.3	261.2	434.5	644.8	908.5	1,195.1	1,450.7	1,701.4
2002.....	45.5	86.2	154.7	268.7	439.5	659.2	931.4	1,205.1	1,456.9	1,725.7
2001.....	46.4	87.8	158.4	272.2	447.1	665.6	938.4	1,207.5	1,458.8	1,765.2
2000.....	45.8	88.2	159.5	281.1	457.1	685.0	944.1	1,214.2	1,503.0	1,815.7
1999.....	46.8	88.5	159.9	285.8	462.2	693.1	955.3	1,213.0	1,488.4	1,798.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms (C00-C97)

White, male

2004.....	37.4	81.1	158.7	278.1	479.7	745.4	1,077.9	1,490.7	1,887.6	2,348.9
2003.....	39.8	85.1	159.7	285.4	494.7	763.2	1,108.1	1,518.9	1,930.1	2,412.1
2002.....	41.2	85.4	162.8	296.5	503.4	785.6	1,145.2	1,537.3	1,934.6	2,486.8
2001.....	41.9	86.5	165.1	298.2	513.9	797.6	1,160.4	1,547.8	1,950.7	2,597.6
2000.....	41.4	87.7	164.0	307.0	522.0	819.7	1,166.6	1,559.2	2,026.9	2,693.7
1999.....	43.1	88.0	164.3	316.0	532.7	831.3	1,191.9	1,571.1	2,024.1	2,696.3

White, female

2004.....	46.6	83.4	137.9	230.5	367.7	534.4	738.3	942.5	1,144.7	1,348.9
2003.....	48.3	84.6	143.1	237.8	378.9	540.3	745.2	961.0	1,155.2	1,386.5
2002.....	49.9	86.9	146.8	242.0	380.4	548.0	757.1	967.4	1,164.2	1,395.1
2001.....	51.0	89.0	152.0	247.4	385.4	549.7	758.6	966.4	1,161.4	1,416.7
2000.....	50.1	88.7	155.2	256.5	397.2	567.2	765.6	971.5	1,191.3	1,459.1
1999.....	50.5	89.1	155.5	257.1	397.2	572.5	767.3	961.5	1,172.1	1,438.2

Black, both sexes

2004.....	64.2	129.6	235.7	377.1	570.2	803.0	1,059.1	1,355.6	1,578.2	1,774.0
2003.....	66.5	138.3	243.2	382.6	589.1	802.7	1,080.1	1,385.5	1,640.8	1,853.1
2002.....	68.3	135.1	238.2	399.6	597.1	833.2	1,126.1	1,425.8	1,665.1	1,885.7
2001.....	70.4	137.8	241.5	404.7	589.4	858.4	1,156.8	1,433.0	1,716.9	1,931.7
2000.....	74.4	145.6	251.9	407.9	604.6	861.4	1,179.2	1,481.7	1,700.6	1,998.9
1999.....	73.3	144.5	255.6	429.5	633.9	879.1	1,191.7	1,502.1	1,704.2	2,008.4

Black, male

2004.....	56.2	134.0	274.7	455.9	729.5	1,035.5	1,442.5	1,926.3	2,340.8	2,748.8
2003.....	61.2	145.6	283.0	464.8	747.3	1,045.9	1,455.8	1,975.5	2,417.1	2,825.5
2002.....	60.4	137.1	271.5	484.4	777.8	1,079.3	1,536.2	2,047.9	2,514.3	2,976.1
2001.....	62.8	143.6	280.7	494.9	746.2	1,135.3	1,600.3	2,084.3	2,645.3	3,209.9
2000.....	70.3	151.8	293.7	512.8	769.2	1,137.3	1,654.9	2,166.0	2,683.2	3,264.8
1999.....	68.3	160.5	300.8	533.7	815.7	1,159.8	1,644.1	2,205.4	2,642.0	3,340.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms (C00-C97)

Black, female

2004.....	71.2	125.9	202.6	312.3	446.1	630.3	798.9	1,017.5	1,176.3	1,387.7
2003.....	71.1	132.0	209.5	314.9	465.7	622.1	826.8	1,034.8	1,234.0	1,475.3
2002.....	75.2	133.5	210.0	330.0	455.6	650.5	850.9	1,054.1	1,223.1	1,468.0
2001.....	77.1	132.8	208.1	330.6	466.1	653.9	861.4	1,044.2	1,244.0	1,457.5
2000.....	78.1	140.2	216.2	321.9	474.8	657.8	863.3	1,069.2	1,205.8	1,527.7
1999.....	77.7	130.6	217.2	344.4	491.4	671.1	891.6	1,078.5	1,239.4	1,510.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races 3/, both sexes

2004.....	0.9	2.0	3.8	5.1	7.5	9.4	11.0	12.7	14.9	18.0
2003.....	0.9	2.0	3.6	5.5	7.9	9.7	10.8	12.8	13.6	18.6
2002.....	0.8	2.0	3.6	5.8	8.1	8.8	12.1	13.1	14.1	18.7
2001.....	0.8	2.1	3.5	6.1	7.9	9.4	11.6	13.5	15.2	18.5
2000.....	0.9	2.0	3.8	5.7	8.1	9.9	11.5	12.6	14.0	19.4
1999.....	0.9	2.3	3.6	5.8	8.1	9.3	11.3	13.5	15.4	19.3

All races 3/, male

2004.....	1.2	3.2	6.2	8.6	11.9	14.9	16.4	19.3	21.7	24.9
2003.....	1.4	3.2	6.2	9.1	12.6	15.0	16.4	19.0	19.9	26.9
2002.....	1.2	3.2	5.8	9.4	12.9	13.9	18.1	19.2	20.3	26.9
2001.....	1.2	3.2	5.8	10.0	13.0	14.3	17.9	19.6	22.1	25.8
2000.....	1.4	3.1	6.0	9.5	12.7	15.2	16.9	18.7	20.2	27.1
1999.....	1.3	3.6	5.7	9.0	13.1	14.3	17.4	20.1	22.3	27.1

All races 3/, female

2004.....	0.5	0.9	1.5	1.7	3.6	4.7	6.7	8.0	10.8	14.9
2003.....	0.5	0.9	1.1	2.1	3.6	5.2	6.3	8.4	9.7	15.0
2002.....	0.4	0.9	1.4	2.5	3.7	4.4	7.3	8.8	10.3	15.1
2001.....	0.4	1.0	1.4	2.4	3.3	5.1	6.6	9.3	11.1	15.4
2000.....	0.4	1.0	1.6	2.2	3.9	5.4	7.2	8.3	10.4	16.3
1999.....	0.4	1.0	1.6	2.7	3.6	5.1	6.6	9.0	11.3	16.1

White, both sexes

2004.....	0.7	1.7	3.4	4.6	7.0	9.0	11.0	12.7	15.0	18.6
2003.....	0.8	1.8	3.2	5.2	7.3	9.3	10.4	12.6	13.4	19.0
2002.....	0.7	1.7	3.1	5.3	7.5	8.4	12.3	13.3	14.2	18.9
2001.....	0.7	1.7	3.2	5.5	7.3	8.7	11.4	13.6	15.5	19.0
2000.....	0.7	1.7	3.2	5.0	7.3	9.4	11.1	12.5	14.0	19.7
1999.....	0.7	1.8	3.0	5.0	7.5	8.9	11.0	13.2	15.5	19.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

White, male

2004.....	0.9	2.6	5.4	7.8	10.9	14.0	15.8	18.8	21.5	24.8
2003.....	1.3	2.7	5.5	8.5	11.3	14.1	15.2	18.2	19.2	27.2
2002.....	1.0	2.7	5.0	8.6	11.7	12.9	18.0	19.4	20.2	27.0
2001.....	1.0	2.6	5.2	8.9	11.8	13.2	17.1	19.2	22.3	26.4
2000.....	1.1	2.7	4.9	8.3	11.2	14.0	15.9	18.0	19.9	27.6
1999.....	1.1	2.7	4.7	7.6	12.0	13.5	16.8	19.3	22.0	27.3

White, female

2004.....	0.4	0.8	1.4	1.5	3.4	4.6	7.1	8.2	10.9	15.7
2003.....	0.3	0.9	0.9	2.1	3.7	5.1	6.4	8.5	9.8	15.4
2002.....	0.4	0.7	1.3	2.2	3.6	4.3	7.6	8.8	10.5	15.3
2001.....	0.3	0.8	1.2	2.2	3.2	4.8	6.7	9.6	11.3	15.9
2000.....	0.4	0.8	1.6	1.9	3.6	5.3	7.2	8.6	10.6	16.5
1999.....	0.4	0.8	1.3	2.5	3.3	4.9	6.4	9.0	11.7	16.7

Black, both sexes

2004.....	1.9	4.6	7.8	9.5	13.3	13.8	13.7	14.9	15.8	12.4
2003.....	1.7	3.9	6.5	8.8	14.4	15.5	15.3	14.8	16.0	15.5
2002.....	1.5	4.2	7.1	10.1	14.7	13.3	12.5	11.3	12.6	16.0
2001.....	1.7	4.3	6.2	12.2	13.1	15.3	15.7	14.2	13.3	12.9
2000.....	1.6	4.1	8.3	12.2	15.2	14.9	16.1	14.6	15.8	15.3
1999.....	1.5	5.9	8.8	12.6	14.6	14.4	16.2	17.7	15.6	14.3

Black, male

2004.....	2.9	7.6	13.8	17.0	23.7	26.2	27.2	25.9	27.3	26.8
2003.....	1.9	7.0	11.4	16.3	27.8	27.6	30.2	26.5	30.3	25.0
2002.....	2.4	6.7	12.9	18.2	26.9	24.2	21.4	16.5	21.4	26.8
2001.....	2.4	7.0	10.8	22.2	24.8	25.4	29.3	25.1	20.9	*
2000.....	2.7	6.3	15.9	21.9	27.3	27.6	29.1	28.8	28.7	*
1999.....	2.3	10.3	14.5	21.5	25.2	24.1	26.5	33.2	28.8	24.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
Black, female										
2004.....	*	1.9	2.7	3.4	5.2	4.7	4.6	8.4	9.8	*
2003.....	1.5	*	2.3	2.7	3.9	6.6	5.3	7.9	8.5	11.7
2002.....	*	1.9	2.2	3.4	5.1	5.2	6.5	8.2	8.0	11.9
2001.....	*	2.0	2.2	4.0	3.8	7.9	6.7	7.8	9.4	10.5
2000.....	*	2.1	*	4.2	5.6	5.6	7.5	6.0	9.3	14.2
1999.....	*	2.1	3.9	5.3	6.3	7.1	9.5	8.3	9.1	10.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of tongue (C01-C02)

All races 3/, both sexes

2004.....	0.2	0.5	1.0	1.3	1.7	2.5	2.5	2.5	3.4	4.1
2003.....	0.3	0.5	1.0	1.5	1.8	2.4	2.6	3.1	3.0	3.7
2002.....	0.2	0.5	0.9	1.4	2.1	2.1	2.8	3.2	3.4	4.0
2001.....	0.2	0.5	0.9	1.6	1.8	2.0	2.6	3.1	3.5	3.8
2000.....	0.3	0.5	0.9	1.4	2.1	2.3	2.4	2.8	3.2	4.0
1999.....	0.3	0.6	0.9	1.3	1.8	2.2	2.6	3.0	3.4	4.0

All races 3/, male

2004.....	0.3	0.7	1.6	2.3	2.8	3.8	3.4	3.7	4.5	4.8
2003.....	0.4	0.8	1.6	2.5	2.6	3.6	3.7	4.2	4.3	5.0
2002.....	0.3	0.7	1.4	2.3	3.1	3.5	3.8	4.5	4.3	5.2
2001.....	0.3	0.8	1.5	2.6	3.0	3.1	4.0	4.1	4.9	4.1
2000.....	0.3	0.8	1.5	2.2	3.4	3.5	3.3	3.6	3.8	5.1
1999.....	0.4	0.9	1.4	1.9	3.0	3.5	4.1	4.7	4.3	5.4

All races 3/, female

2004.....	0.2	0.3	0.4	0.4	0.8	1.3	1.7	1.7	2.8	3.8
2003.....	*	0.3	0.4	0.5	1.1	1.3	1.8	2.3	2.2	3.2
2002.....	*	0.2	0.4	0.6	1.2	0.9	2.0	2.3	2.8	3.5
2001.....	*	0.3	0.4	0.6	0.7	1.1	1.4	2.4	2.7	3.6
2000.....	0.2	0.3	0.4	0.6	1.0	1.2	1.7	2.3	2.8	3.6
1999.....	0.2	0.3	0.4	0.7	0.8	1.1	1.4	1.8	2.8	3.5

White, both sexes

2004.....	0.2	0.5	1.0	1.3	1.8	2.4	2.5	2.7	3.5	4.2
2003.....	0.2	0.5	0.9	1.4	1.8	2.3	2.6	3.1	3.0	3.8
2002.....	0.2	0.4	0.9	1.4	2.0	2.0	2.9	3.3	3.5	4.1
2001.....	0.2	0.5	0.9	1.6	1.8	1.9	2.6	3.2	3.5	3.9
2000.....	0.2	0.5	0.8	1.3	2.0	2.3	2.4	2.9	3.3	4.1
1999.....	0.3	0.5	0.8	1.1	1.8	2.2	2.6	3.0	3.5	4.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of tongue (C01-C02)										
White, male										
2004.....	0.2	0.6	1.5	2.2	2.8	3.7	3.4	3.9	4.4	4.8
2003.....	0.4	0.8	1.5	2.5	2.6	3.4	3.7	4.1	4.3	5.1
2002.....	0.3	0.6	1.4	2.2	3.0	3.2	3.9	4.6	4.4	5.2
2001.....	0.3	0.7	1.5	2.5	2.9	3.0	4.1	4.2	4.8	4.3
2000.....	0.3	0.7	1.3	2.2	3.3	3.5	3.3	3.5	3.8	5.1
1999.....	0.4	0.7	1.3	1.6	2.9	3.5	4.0	4.7	4.5	5.5
White, female										
2004.....	*	0.3	0.4	0.5	0.9	1.3	1.9	1.8	2.9	3.9
2003.....	*	0.3	0.3	0.5	1.1	1.3	1.8	2.4	2.2	3.3
2002.....	*	*	0.4	0.6	1.2	0.9	2.1	2.4	3.0	3.6
2001.....	*	0.3	0.4	0.7	0.7	1.0	1.4	2.5	2.8	3.8
2000.....	*	0.3	0.5	0.5	0.9	1.3	1.7	2.4	3.0	3.7
1999.....	*	0.3	0.4	0.7	0.8	1.0	1.4	1.9	3.0	3.5
Black, both sexes										
2004.....	*	0.9	1.5	1.6	*	3.0	*	*	*	*
2003.....	*	0.9	1.3	2.2	2.0	3.7	*	*	*	*
2002.....	*	1.0	1.2	1.9	3.4	3.8	*	*	*	*
2001.....	*	1.0	1.2	2.0	2.9	2.6	*	*	*	*
2000.....	*	*	2.0	2.1	3.1	2.5	3.1	*	*	*
1999.....	*	1.3	2.0	2.8	2.5	2.8	3.0	*	*	*
Black, male										
2004.....	*	*	2.7	3.0	*	6.1	*	*	*	*
2003.....	*	1.6	*	3.8	*	6.4	*	*	*	*
2002.....	*	*	*	3.1	6.1	7.2	*	*	*	*
2001.....	*	1.7	2.3	4.1	5.5	*	*	*	*	*
2000.....	*	*	4.0	3.6	5.4	5.2	*	*	*	*
1999.....	*	2.3	3.2	5.1	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of pharynx (C10-C13,C14.0)

All races 3/, both sexes

2004.....	0.4	1.0	1.6	2.2	3.6	4.2	4.9	5.6	5.6	5.1
2003.....	0.3	0.8	1.5	2.3	3.6	4.5	4.8	5.1	5.0	6.0
2002.....	0.3	0.8	1.6	2.5	3.5	3.8	5.2	5.2	4.9	5.4
2001.....	0.4	0.9	1.5	2.6	3.7	4.1	4.9	5.5	5.7	5.9
2000.....	0.4	0.8	1.6	2.7	3.7	4.3	5.1	5.2	5.3	5.6
1999.....	0.4	1.0	1.6	2.7	3.6	4.1	5.1	5.5	5.6	5.2

All races 3/, male

2004.....	0.6	1.6	2.7	3.8	5.8	7.0	7.9	9.2	8.9	8.7
2003.....	0.5	1.3	2.7	3.9	5.9	7.2	7.6	8.2	8.7	10.8
2002.....	0.5	1.4	2.6	4.0	5.9	6.0	8.2	8.2	8.2	9.8
2001.....	0.6	1.5	2.6	4.4	6.1	6.5	7.9	8.8	9.0	10.0
2000.....	0.6	1.3	2.6	4.6	5.9	6.8	8.2	8.6	8.6	9.9
1999.....	0.6	1.6	2.6	4.5	5.9	6.5	8.0	8.7	9.6	9.9

All races 3/, female

2004.....	0.2	0.4	0.6	0.8	1.6	1.9	2.5	3.1	3.5	3.5
2003.....	*	0.3	0.5	0.9	1.5	2.2	2.5	2.8	2.8	3.9
2002.....	*	0.3	0.6	1.2	1.4	1.8	2.9	3.1	2.8	3.5
2001.....	*	0.4	0.6	0.9	1.6	2.1	2.5	3.1	3.6	4.2
2000.....	*	0.3	0.6	1.0	1.6	2.2	2.8	2.9	3.3	3.9
1999.....	*	0.4	0.7	1.1	1.6	2.1	2.8	3.3	3.2	3.4

White, both sexes

2004.....	0.3	0.8	1.3	1.8	3.0	3.8	4.7	5.3	5.4	5.1
2003.....	0.3	0.6	1.2	2.1	3.1	4.2	4.4	4.8	4.6	5.8
2002.....	0.2	0.6	1.2	2.2	3.0	3.5	5.2	5.0	4.6	5.2
2001.....	0.2	0.7	1.2	2.2	3.3	3.7	4.6	5.2	5.5	6.0
2000.....	0.2	0.6	1.3	2.2	3.1	3.9	4.7	5.0	5.2	5.3
1999.....	0.2	0.6	1.2	2.3	3.2	3.8	4.7	5.1	5.3	5.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pharynx (C10-C13,C14.0)										
White, male										
2004.....	0.4	1.2	2.1	3.2	4.8	6.1	7.4	8.5	8.8	8.2
2003.....	0.4	0.9	2.2	3.4	4.8	6.4	6.7	7.7	7.8	10.6
2002.....	0.3	1.0	2.0	3.4	4.9	5.4	8.1	8.0	7.6	9.5
2001.....	0.4	1.1	2.0	3.6	5.3	5.6	7.4	8.3	8.8	9.9
2000.....	0.3	0.9	2.0	3.7	4.8	6.1	7.3	7.8	8.4	9.5
1999.....	0.3	1.0	1.9	3.7	5.1	5.8	7.4	7.9	8.9	9.9
White, female										
2004.....	*	0.3	0.5	0.6	1.4	1.7	2.5	3.0	3.4	3.7
2003.....	*	0.3	0.3	0.8	1.5	2.1	2.5	2.8	2.7	3.7
2002.....	*	0.2	0.5	1.0	1.3	1.8	2.8	2.9	2.7	3.4
2001.....	*	*	0.4	0.8	1.5	2.1	2.4	3.1	3.5	4.3
2000.....	*	0.2	0.6	0.8	1.5	1.9	2.7	2.9	3.3	3.7
1999.....	*	0.3	0.5	0.9	1.4	1.9	2.6	3.2	3.1	3.5
Black, both sexes										
2004.....	1.0	2.3	4.3	5.4	9.3	7.8	8.2	9.1	6.7	5.6
2003.....	0.7	2.0	3.3	4.2	8.4	8.6	8.5	7.7	8.1	7.3
2002.....	0.9	2.0	3.8	5.1	8.1	5.7	6.4	5.3	7.6	7.4
2001.....	0.9	2.2	3.7	6.5	6.4	7.4	7.8	7.3	7.6	*
2000.....	0.8	2.1	3.8	7.1	8.7	8.2	9.1	8.4	7.3	7.5
1999.....	0.8	3.0	4.6	6.5	7.5	7.5	8.9	10.2	9.0	*
Black, male										
2004.....	1.5	4.0	7.6	9.2	16.8	14.8	16.2	16.8	*	*
2003.....	*	3.7	6.0	8.0	16.4	16.5	17.1	14.6	17.7	*
2002.....	*	3.4	7.2	9.2	15.2	10.7	10.3	*	*	*
2001.....	*	3.4	6.5	11.7	11.6	14.5	14.5	13.5	*	*
2000.....	*	3.5	7.1	12.7	15.9	14.1	17.7	17.0	*	*
1999.....	*	5.4	8.1	11.2	12.9	13.0	14.6	19.9	17.4	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pharynx (C10-C13,C14.0)										
Black, female										
2004.....	*	*	*	2.2	3.4	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	3.8	*	*	*	*
1999.....	*	*	*	*	*	*	5.2	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)

All races 3/, both sexes

2004.....	0.2	0.5	1.2	1.5	2.1	2.7	3.6	4.4	5.6	8.4
2003.....	0.3	0.7	1.1	1.7	2.5	2.7	3.3	4.4	5.4	8.7
2002.....	0.2	0.7	1.1	1.9	2.4	2.9	4.0	4.5	5.7	8.7
2001.....	0.2	0.6	1.0	1.8	2.3	3.1	4.1	4.8	5.9	8.3
2000.....	0.3	0.7	1.2	1.6	2.2	3.3	3.8	4.4	5.2	9.5
1999.....	0.2	0.7	1.1	1.7	2.6	3.0	3.6	4.9	6.3	9.7

All races 3/, male

2004.....	0.4	0.9	1.9	2.5	3.2	4.0	5.0	6.1	7.8	10.8
2003.....	0.5	1.1	1.9	2.7	4.1	4.0	4.9	6.1	6.7	10.7
2002.....	0.3	1.1	1.8	3.1	3.8	4.4	5.9	6.2	7.5	11.1
2001.....	0.3	0.9	1.7	2.9	3.7	4.6	5.8	6.4	8.0	10.6
2000.....	0.4	1.1	1.9	2.6	3.3	4.8	5.2	6.3	7.4	11.4
1999.....	0.3	1.1	1.7	2.5	4.2	4.3	5.2	6.5	8.2	11.4

All races 3/, female

2004.....	*	0.2	0.5	0.5	1.2	1.5	2.5	3.2	4.3	7.3
2003.....	0.2	0.3	0.3	0.8	1.0	1.7	2.0	3.2	4.7	7.7
2002.....	*	0.3	0.4	0.7	1.1	1.6	2.4	3.4	4.6	7.7
2001.....	*	0.3	0.4	0.8	1.0	1.9	2.7	3.7	4.7	7.3
2000.....	*	0.4	0.5	0.6	1.3	1.9	2.8	3.1	4.0	8.8
1999.....	*	0.4	0.5	0.9	1.2	1.8	2.3	3.7	5.2	9.0

White, both sexes

2004.....	0.2	0.4	1.1	1.4	2.1	2.7	3.7	4.5	5.7	8.8
2003.....	0.3	0.7	1.0	1.7	2.4	2.8	3.2	4.4	5.6	9.0
2002.....	0.2	0.7	1.0	1.8	2.4	2.9	4.1	4.7	5.9	9.1
2001.....	0.2	0.5	1.0	1.7	2.2	3.0	4.0	5.0	6.3	8.5
2000.....	0.3	0.6	1.1	1.4	2.1	3.2	3.9	4.6	5.2	9.9
1999.....	0.2	0.6	1.0	1.6	2.4	2.9	3.6	4.9	6.5	10.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)

White, male										
2004.....	0.3	0.7	1.8	2.3	3.2	4.0	4.9	6.1	7.9	11.2
2003.....	0.5	1.0	1.8	2.6	3.9	4.0	4.7	5.9	6.8	11.1
2002.....	0.3	1.0	1.7	2.9	3.8	4.4	5.9	6.4	7.8	11.5
2001.....	0.3	0.8	1.6	2.6	3.5	4.5	5.5	6.5	8.4	11.0
2000.....	0.4	1.0	1.6	2.3	3.1	4.4	5.2	6.4	7.2	12.2
1999.....	0.3	0.9	1.6	2.3	3.9	4.0	5.2	6.5	8.3	11.5
White, female										
2004.....	*	*	0.5	0.5	1.1	1.6	2.7	3.4	4.4	7.7
2003.....	*	0.3	0.3	0.8	1.0	1.7	2.1	3.3	4.8	8.1
2002.....	*	0.3	0.4	0.7	1.1	1.6	2.6	3.5	4.7	8.0
2001.....	*	0.3	0.4	0.8	1.0	1.7	2.8	3.9	5.0	7.5
2000.....	*	0.3	0.5	0.6	1.3	2.1	2.8	3.2	4.1	8.9
1999.....	*	0.3	0.4	0.9	1.1	1.9	2.3	3.8	5.5	9.5
Black, both sexes										
2004.....	*	1.4	2.0	2.6	2.5	3.0	3.1	3.9	*	*
2003.....	*	1.0	1.9	2.4	3.9	3.3	4.4	4.8	5.1	*
2002.....	*	1.2	2.0	3.0	3.1	3.6	3.9	4.1	*	6.2
2001.....	*	1.2	1.2	3.7	3.7	5.3	5.5	4.5	*	*
2000.....	*	1.4	2.6	3.0	3.4	4.2	4.0	3.7	6.2	*
1999.....	*	1.6	2.2	3.3	4.7	4.1	4.3	4.3	*	6.4
Black, male										
2004.....	*	2.5	3.4	4.8	*	5.3	*	*	*	*
2003.....	*	1.6	3.5	4.5	7.7	*	8.9	*	*	*
2002.....	*	1.9	3.7	5.8	5.5	6.2	7.3	*	*	*
2001.....	*	1.8	*	6.4	7.5	7.1	10.1	*	*	*
2000.....	*	*	4.9	5.6	6.0	8.3	6.7	*	*	*
1999.....	*	2.6	3.2	5.3	8.3	7.3	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of digestive organs (C15-C26)

All races 3/, both sexes

2004.....	9.6	20.5	39.3	62.7	99.2	142.7	202.9	270.8	345.7	445.3
2003.....	9.3	21.2	40.1	62.7	100.4	148.1	209.2	277.0	355.1	461.2
2002.....	10.1	21.1	39.5	64.8	101.0	152.4	212.5	281.9	354.0	467.4
2001.....	9.8	21.6	39.7	63.4	102.1	154.5	215.5	282.2	362.2	487.9
2000.....	10.0	21.7	38.8	65.0	105.6	153.7	217.5	290.1	374.6	504.9
1999.....	10.2	21.8	38.1	65.5	106.6	157.7	218.4	288.9	367.2	507.4

All races 3/, male

2004.....	11.8	27.1	53.7	84.7	131.5	188.8	264.4	349.3	437.2	531.7
2003.....	11.4	28.0	53.7	84.2	133.1	196.2	271.8	356.3	455.5	546.7
2002.....	12.7	27.5	53.5	86.9	135.8	202.1	278.3	364.4	449.0	557.9
2001.....	11.6	28.1	52.7	84.4	136.1	203.6	278.0	366.3	458.9	593.2
2000.....	12.3	28.4	51.4	86.2	139.3	202.8	281.4	377.0	474.6	614.1
1999.....	12.5	29.0	50.4	86.9	138.8	205.8	285.0	374.7	472.3	620.8

All races 3/, female

2004.....	7.4	14.1	25.5	42.1	69.8	102.7	153.3	214.4	289.4	406.4
2003.....	7.2	14.7	27.0	42.4	70.7	106.4	159.0	220.5	293.6	423.4
2002.....	7.5	14.8	26.1	44.1	69.4	109.5	159.9	223.5	296.1	428.1
2001.....	7.9	15.3	27.2	43.8	71.3	112.3	165.8	223.4	304.3	443.9
2000.....	7.7	15.2	26.6	45.1	75.1	111.6	167.2	229.6	315.5	460.4
1999.....	8.0	14.9	26.2	45.5	77.5	116.7	166.3	229.3	305.7	461.7

White, both sexes

2004.....	8.7	18.6	36.2	58.5	93.9	136.5	195.2	264.7	340.6	440.0
2003.....	8.2	19.1	36.0	58.7	94.9	143.0	202.2	270.8	348.6	453.8
2002.....	9.1	19.2	36.3	60.5	95.8	145.8	205.6	274.7	346.3	460.4
2001.....	8.8	19.6	36.6	58.8	97.3	147.3	208.2	274.5	354.1	481.6
2000.....	8.8	19.6	35.6	60.3	100.2	147.0	210.3	281.8	365.9	497.0
1999.....	9.1	19.9	34.8	60.2	100.9	151.7	211.0	280.4	358.9	499.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of digestive organs (C15-C26)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
White, male										
2004.....	10.7	24.7	49.8	78.7	124.7	180.8	253.8	341.7	432.2	528.5
2003.....	10.2	25.1	48.3	79.4	125.7	189.7	263.8	348.6	450.2	539.4
2002.....	11.6	25.3	49.1	81.2	129.0	194.6	269.7	356.9	439.1	550.5
2001.....	10.5	25.5	48.6	78.4	130.0	194.4	269.8	358.3	449.9	588.1
2000.....	10.9	25.6	47.1	79.3	132.3	195.1	271.9	368.1	464.1	605.3
1999.....	11.3	26.3	45.8	79.5	131.0	198.2	277.1	363.3	463.6	614.4
White, female										
2004.....	6.7	12.5	23.0	39.0	65.3	97.3	147.1	208.4	283.7	400.0
2003.....	6.3	13.2	24.1	38.9	66.4	101.7	151.9	214.5	285.9	415.8
2002.....	6.7	13.1	23.9	40.6	65.2	102.9	153.3	215.9	289.5	421.3
2001.....	7.2	13.7	24.8	40.1	67.2	105.9	158.4	215.0	296.1	437.0
2000.....	6.8	13.6	24.4	42.3	70.6	105.0	160.8	221.1	307.4	453.1
1999.....	7.0	13.6	24.1	41.8	73.1	111.2	158.5	222.3	297.0	453.7
Black, both sexes										
2004.....	14.8	33.7	63.3	102.7	151.9	210.1	284.1	352.2	418.9	524.2
2003.....	15.5	35.7	71.8	102.0	156.0	201.4	287.0	359.2	450.1	567.1
2002.....	16.2	34.7	64.6	107.2	151.8	217.4	295.9	380.0	450.0	566.6
2001.....	15.7	36.7	65.8	107.4	150.7	224.7	297.8	380.2	475.1	585.6
2000.....	16.8	37.8	68.1	105.8	158.3	221.2	302.3	395.9	483.7	606.9
1999.....	17.0	36.1	66.2	112.9	160.5	222.0	306.6	395.8	484.5	613.1
Black, male										
2004.....	18.2	44.3	87.2	143.2	206.1	286.1	390.4	458.6	536.6	575.2
2003.....	19.6	49.0	101.0	138.7	214.4	268.9	386.3	467.4	547.8	654.8
2002.....	20.1	44.6	90.5	147.5	206.9	288.2	394.9	489.1	600.7	671.1
2001.....	19.1	48.4	89.0	146.3	199.9	305.9	385.4	489.7	602.7	676.3
2000.....	21.6	50.2	93.2	152.0	212.2	289.6	403.5	508.4	629.7	728.8
1999.....	21.1	50.3	92.5	159.0	216.8	292.9	397.5	543.8	615.6	715.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of digestive organs (C15-C26)

Black, female

2004.....	11.8	24.5	43.1	69.4	109.5	153.6	212.0	289.2	356.8	504.0
2003.....	11.9	24.1	47.1	71.8	110.4	151.3	220.0	294.9	398.8	533.1
2002.....	12.7	26.1	42.7	74.1	108.7	164.8	229.4	314.8	371.6	526.6
2001.....	12.7	26.5	46.1	75.6	112.0	164.6	239.4	314.8	410.0	551.9
2000.....	12.5	27.0	46.8	67.9	115.7	170.7	235.1	328.1	410.1	561.5
1999.....	13.4	23.9	44.0	75.2	116.3	169.5	246.2	306.7	419.6	574.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of esophagus (C15)

All races 3/, both sexes

2004.....	1.0	2.3	4.7	8.1	13.2	17.5	21.3	26.0	27.7	25.8
2003.....	0.9	2.4	4.6	8.8	12.8	17.6	22.2	25.4	27.6	25.0
2002.....	1.0	2.4	4.7	8.6	12.7	18.1	22.5	25.7	27.8	24.6
2001.....	1.1	2.4	5.0	7.9	13.0	18.5	22.9	26.1	26.5	25.8
2000.....	1.0	2.4	4.9	8.8	13.2	17.9	22.1	24.5	27.2	25.9
1999.....	1.0	2.4	4.7	8.7	13.0	18.2	22.0	24.3	24.6	26.8

All races 3/, male

2004.....	1.6	4.0	8.1	14.1	22.9	30.5	37.0	46.8	51.7	48.5
2003.....	1.5	4.1	8.1	15.3	22.7	30.6	38.3	45.7	51.0	45.3
2002.....	1.6	4.1	8.3	15.1	22.1	31.4	40.3	46.8	50.6	43.4
2001.....	1.8	4.1	8.7	13.7	22.6	32.6	39.9	46.5	48.8	48.1
2000.....	1.8	4.2	8.7	15.2	23.0	30.7	38.0	43.9	48.8	47.8
1999.....	1.7	4.2	8.3	15.3	21.9	31.7	39.3	44.1	44.8	48.0

All races 3/, female

2004.....	0.3	0.7	1.4	2.4	4.3	6.1	8.7	11.1	12.9	15.7
2003.....	0.3	0.7	1.3	2.6	3.8	6.3	9.2	11.0	13.3	16.1
2002.....	0.4	0.7	1.3	2.4	4.2	6.7	8.2	10.9	14.0	16.5
2001.....	0.3	0.7	1.4	2.4	4.2	6.4	9.4	11.9	13.2	16.5
2000.....	0.2	0.6	1.4	2.8	4.4	7.1	9.6	11.0	14.5	17.1
1999.....	0.3	0.6	1.3	2.5	5.0	6.7	8.5	10.5	12.8	18.3

White, both sexes

2004.....	1.0	2.2	4.6	7.9	12.8	17.4	21.2	26.5	28.0	26.2
2003.....	0.9	2.3	4.3	8.5	12.3	17.0	22.5	25.8	27.8	25.8
2002.....	0.9	2.3	4.6	8.1	12.2	17.5	22.4	25.7	28.2	25.1
2001.....	1.0	2.3	4.6	7.1	12.3	17.7	22.1	26.2	26.9	26.3
2000.....	0.9	2.3	4.5	8.1	12.4	17.4	21.8	24.0	27.2	26.6
1999.....	0.9	2.2	4.3	8.0	12.1	17.5	21.5	23.7	24.9	27.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of esophagus (C15)

White, male

2004.....	1.7	3.9	8.0	13.9	22.5	30.5	37.2	47.6	52.8	49.2
2003.....	1.6	3.9	7.6	15.0	22.0	29.5	39.0	46.5	52.3	46.7
2002.....	1.6	4.0	8.1	14.5	21.5	30.7	40.2	47.2	51.3	44.6
2001.....	1.8	3.9	8.1	12.5	21.6	31.2	39.0	46.9	49.5	48.8
2000.....	1.8	4.1	8.0	14.1	22.0	30.0	37.5	43.2	49.7	49.3
1999.....	1.6	4.0	7.6	14.1	20.5	30.8	38.4	43.0	45.6	48.0

White, female

2004.....	0.3	0.5	1.1	2.1	3.8	5.9	8.1	11.1	12.6	15.9
2003.....	0.3	0.6	1.0	2.3	3.4	6.0	9.0	10.9	12.7	16.5
2002.....	0.3	0.6	1.2	2.1	3.7	5.9	7.8	10.3	14.1	16.7
2001.....	0.3	0.6	1.2	1.9	3.7	5.8	8.5	11.6	13.2	16.8
2000.....	*	0.4	1.1	2.4	3.6	6.4	9.1	10.6	13.8	17.3
1999.....	*	0.5	1.0	2.1	4.4	5.8	8.0	10.1	12.6	18.9

Black, both sexes

2004.....	1.2	4.0	6.6	12.0	20.4	23.0	26.8	27.4	27.0	21.4
2003.....	1.3	4.1	8.4	14.1	20.7	26.1	24.9	26.9	29.4	21.3
2002.....	1.6	3.6	6.9	14.9	20.4	27.5	29.8	31.8	26.2	20.8
2001.....	1.6	3.7	8.6	15.9	21.9	29.6	35.8	30.7	27.6	23.6
2000.....	1.5	4.1	9.8	16.9	23.3	27.0	29.6	36.1	29.9	22.2
1999.....	1.7	4.3	9.4	17.1	23.5	28.7	31.3	35.7	23.7	23.9

Black, male

2004.....	1.7	6.3	10.8	19.9	33.4	40.0	43.1	49.6	45.2	38.7
2003.....	2.1	7.1	14.0	23.3	36.2	46.5	41.4	47.5	43.6	38.6
2002.....	2.5	5.7	11.5	26.8	34.4	45.4	54.4	54.2	48.1	37.5
2001.....	2.6	6.0	15.3	27.5	38.8	51.8	59.9	55.3	53.8	46.5
2000.....	2.4	6.3	17.4	30.5	38.5	44.8	49.8	65.7	45.5	38.1
1999.....	3.2	7.4	16.4	30.3	40.3	46.5	56.0	67.9	36.6	51.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of esophagus (C15)

Black, female

2004.....	*	1.9	3.1	5.5	10.2	10.4	15.8	14.3	17.4	14.6
2003.....	*	1.6	3.6	6.5	8.6	11.0	13.7	14.7	22.0	14.6
2002.....	*	1.8	3.0	5.1	9.4	14.2	13.2	18.4	14.7	14.4
2001.....	*	1.7	2.9	6.3	8.6	13.1	19.7	16.1	14.3	15.2
2000.....	*	2.1	3.4	5.8	11.3	13.8	16.2	18.2	22.1	16.3
1999.....	*	1.6	3.4	6.4	10.3	15.6	14.9	16.3	17.3	13.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of stomach (C16)

All races 3/, both sexes

2004.....	1.4	2.1	3.5	5.3	8.0	12.1	18.2	23.4	30.3	42.2
2003.....	1.3	2.3	3.8	5.3	8.6	12.4	18.4	24.2	32.2	43.6
2002.....	1.2	2.1	3.5	5.6	8.5	12.3	18.8	25.6	33.0	47.6
2001.....	1.3	2.2	3.7	5.3	9.2	13.7	19.5	27.0	34.7	45.4
2000.....	1.4	2.3	3.6	5.6	9.7	13.8	19.6	27.1	36.9	52.7
1999.....	1.4	2.3	3.8	6.2	9.8	14.2	20.2	28.1	36.8	50.1

All races 3/, male

2004.....	1.7	2.6	4.6	7.5	11.2	17.0	24.9	33.6	42.5	62.9
2003.....	1.5	2.8	5.0	7.5	12.3	17.9	26.6	34.2	44.7	60.0
2002.....	1.4	2.7	4.8	7.7	11.9	18.1	27.5	36.4	47.1	67.2
2001.....	1.4	2.8	4.9	7.5	13.7	19.5	27.5	39.1	49.4	62.6
2000.....	1.7	2.9	4.8	7.8	14.1	19.5	28.3	39.4	51.4	75.6
1999.....	1.7	3.1	5.4	8.8	13.7	20.3	28.6	41.5	54.9	71.3

All races 3/, female

2004.....	1.2	1.6	2.4	3.3	5.1	7.8	12.8	16.0	22.7	32.9
2003.....	1.1	1.7	2.6	3.1	5.3	7.7	11.9	17.1	24.5	36.4
2002.....	1.0	1.5	2.3	3.7	5.4	7.3	11.9	18.1	24.4	39.1
2001.....	1.1	1.6	2.5	3.2	5.1	8.8	13.1	18.5	25.9	38.3
2000.....	1.2	1.7	2.4	3.5	5.8	8.9	12.7	18.5	28.3	43.4
1999.....	1.2	1.5	2.4	3.7	6.4	9.1	13.7	18.8	26.2	41.6

White, both sexes

2004.....	1.2	1.7	2.9	4.5	6.9	10.2	15.9	20.9	27.3	38.8
2003.....	1.1	1.8	3.1	4.6	7.6	11.0	16.3	21.4	28.8	38.7
2002.....	1.0	1.8	2.9	4.9	7.3	10.5	16.5	22.6	29.3	42.2
2001.....	1.1	1.9	3.1	4.7	7.9	11.7	17.4	24.0	31.0	41.1
2000.....	1.1	1.9	3.1	4.9	8.5	11.9	17.2	23.6	32.7	47.3
1999.....	1.1	2.0	3.1	5.2	8.3	12.8	17.7	24.8	33.0	44.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of stomach (C16)

White, male

2004.....	1.4	2.1	3.9	6.5	9.5	14.4	21.3	29.9	38.0	58.4
2003.....	1.2	2.3	4.1	6.6	10.6	15.5	23.5	30.3	40.3	53.5
2002.....	1.1	2.4	3.9	6.8	10.3	15.7	24.3	32.8	42.4	60.5
2001.....	1.2	2.4	4.1	6.5	11.7	16.3	24.5	35.4	44.0	57.7
2000.....	1.2	2.3	4.2	6.7	12.5	17.1	25.1	34.6	46.1	68.9
1999.....	1.2	2.7	4.4	7.6	11.5	18.4	25.2	36.9	49.0	63.4

White, female

2004.....	1.0	1.3	2.0	2.7	4.4	6.5	11.5	14.2	20.7	29.9
2003.....	0.9	1.4	2.2	2.7	4.8	7.0	10.5	15.0	21.7	32.1
2002.....	0.8	1.2	1.8	3.0	4.6	5.9	10.1	15.4	21.3	34.2
2001.....	1.0	1.3	2.1	2.9	4.3	7.7	11.7	16.0	23.1	34.1
2000.....	1.0	1.5	2.0	3.1	4.9	7.4	10.9	15.8	24.7	38.6
1999.....	0.9	1.2	1.9	2.9	5.4	8.0	11.7	16.3	23.5	37.3

Black, both sexes

2004.....	2.7	3.9	6.9	10.6	15.8	27.3	34.0	43.8	59.9	71.3
2003.....	2.4	4.3	7.4	9.6	16.7	22.3	33.1	48.8	62.4	85.8
2002.....	2.2	3.6	7.5	10.4	16.1	24.6	38.9	52.7	63.1	100.4
2001.....	2.5	3.7	7.5	9.2	17.7	27.1	34.7	51.1	69.9	88.1
2000.....	2.9	4.7	7.0	11.1	17.3	26.8	38.2	56.9	75.1	101.7
1999.....	3.2	3.9	8.5	13.2	19.1	23.6	40.3	58.6	74.3	102.4

Black, male

2004.....	3.1	5.3	9.5	15.3	25.2	40.7	56.0	63.7	93.3	100.3
2003.....	3.3	6.2	10.9	14.9	26.0	36.4	54.2	74.9	87.1	106.3
2002.....	2.4	5.1	11.0	15.2	24.0	36.9	59.8	73.5	90.2	137.2
2001.....	2.8	5.0	11.5	13.8	27.5	43.7	54.2	75.7	106.7	104.4
2000.....	3.9	6.6	9.6	16.5	25.9	38.5	57.2	83.2	105.4	136.1
1999.....	4.0	5.7	12.2	19.0	28.2	35.1	60.4	89.3	121.2	147.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of stomach (C16)

Black, female

2004.....	2.3	2.7	4.7	6.7	8.4	17.3	19.1	32.1	42.3	59.9
2003.....	1.5	2.7	4.5	5.3	9.5	11.9	18.8	33.4	49.5	77.8
2002.....	2.1	2.3	4.6	6.5	9.9	15.5	24.9	40.3	48.9	86.3
2001.....	2.2	2.5	4.1	5.4	10.1	14.8	21.7	36.3	51.1	82.1
2000.....	1.9	3.0	4.7	6.6	10.5	18.2	25.5	41.0	59.9	88.9
1999.....	2.4	2.4	5.4	8.4	12.0	15.2	27.0	40.0	51.0	85.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of small intestine (C17)

All races 3/, both sexes

2004.....	0.1	0.2	0.4	0.5	0.9	1.4	1.6	2.1	2.7	3.1
2003.....	0.1	0.2	0.3	0.6	0.9	1.1	1.6	2.1	3.0	3.0
2002.....	0.1	0.2	0.4	0.6	0.7	1.2	1.6	2.1	2.6	2.8
2001.....	0.1	0.2	0.4	0.5	1.0	1.0	1.9	2.4	2.6	3.2
2000.....	0.1	0.3	0.4	0.6	1.0	1.2	1.7	2.2	2.8	3.0
1999.....	0.1	0.4	0.5	0.4	0.9	1.2	1.7	2.2	2.6	3.2

All races 3/, male

2004.....	*	0.2	0.4	0.7	1.1	1.8	2.1	2.4	3.9	4.3
2003.....	*	0.3	0.3	0.9	1.0	1.5	2.0	2.2	3.5	3.5
2002.....	0.2	0.3	0.4	0.6	1.0	1.4	1.9	2.7	2.9	3.7
2001.....	*	0.3	0.5	0.6	1.3	1.3	2.5	2.7	3.3	4.1
2000.....	*	0.3	0.4	0.6	1.2	1.4	2.2	2.5	3.7	4.2
1999.....	0.2	0.4	0.5	0.6	1.4	1.5	2.0	3.0	3.6	4.4

All races 3/, female

2004.....	*	0.2	0.4	0.4	0.8	1.1	1.2	1.9	2.0	2.5
2003.....	*	0.2	0.4	0.4	0.7	0.7	1.3	2.0	2.8	2.7
2002.....	*	*	0.3	0.5	0.5	1.1	1.3	1.7	2.5	2.4
2001.....	*	0.2	0.2	0.5	0.8	0.8	1.4	2.2	2.1	2.8
2000.....	*	0.2	0.3	0.5	0.8	1.1	1.4	1.9	2.3	2.5
1999.....	*	0.3	0.4	0.3	0.4	0.9	1.4	1.7	2.0	2.7

White, both sexes

2004.....	*	0.2	0.3	0.5	0.8	1.4	1.5	2.0	2.7	2.9
2003.....	*	0.2	0.3	0.6	0.8	1.1	1.6	2.0	3.0	2.9
2002.....	0.1	0.2	0.4	0.5	0.6	1.2	1.6	2.0	2.5	2.6
2001.....	0.1	0.2	0.3	0.5	1.0	1.0	1.8	2.4	2.5	3.0
2000.....	0.1	0.2	0.3	0.6	1.0	1.2	1.7	2.1	2.7	3.1
1999.....	0.1	0.3	0.4	0.4	0.8	1.2	1.6	2.1	2.4	3.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of colon, rectum and anus (C18-C21)

All races 3/, both sexes

2004.....	3.9	7.4	13.5	22.4	35.9	53.2	78.3	111.3	152.3	225.6
2003.....	3.8	7.9	14.3	23.0	38.6	59.3	83.3	117.0	159.0	239.7
2002.....	4.3	8.2	15.0	24.6	40.7	61.6	86.2	118.9	163.2	241.4
2001.....	4.1	8.4	15.2	25.4	39.3	62.9	88.8	121.1	165.8	255.7
2000.....	4.0	8.2	14.7	25.2	42.0	62.0	92.4	126.9	174.8	270.4
1999.....	4.0	8.3	14.4	25.3	43.6	63.9	91.9	129.0	173.2	271.7

All races 3/, male

2004.....	4.2	8.3	16.1	27.0	43.7	66.4	98.0	139.0	183.1	254.7
2003.....	4.2	9.0	16.6	27.6	47.1	75.0	103.4	143.0	196.5	273.3
2002.....	4.9	9.1	17.5	30.1	51.6	77.8	106.8	147.7	197.4	275.1
2001.....	4.3	9.5	17.1	30.8	48.0	78.9	110.4	149.4	201.3	298.3
2000.....	4.4	9.0	17.0	30.6	51.0	77.4	115.7	159.4	213.0	315.6
1999.....	4.1	9.1	16.6	30.6	52.9	78.0	114.3	161.3	216.2	324.0

All races 3/, female

2004.....	3.6	6.5	11.1	18.0	28.7	41.8	62.4	91.3	133.3	212.5
2003.....	3.4	6.9	12.1	18.6	30.9	45.6	67.3	98.5	136.1	224.8
2002.....	3.6	7.4	12.6	19.5	30.8	47.6	69.7	98.6	142.3	226.8
2001.....	3.8	7.4	13.4	20.4	31.5	49.2	71.5	101.2	144.6	237.8
2000.....	3.5	7.5	12.4	20.2	33.9	48.8	74.1	104.3	152.2	251.9
1999.....	3.9	7.5	12.3	20.4	35.1	51.8	74.3	106.6	148.0	250.6

White, both sexes

2004.....	3.6	6.8	12.6	21.3	34.2	51.2	76.2	109.5	150.9	225.1
2003.....	3.4	7.3	13.0	21.5	36.9	58.0	80.7	115.1	158.2	238.5
2002.....	3.9	7.5	14.2	23.3	38.9	59.9	84.6	116.9	160.3	240.7
2001.....	3.8	7.7	14.3	23.9	37.9	60.8	86.8	118.0	163.7	255.3
2000.....	3.6	7.5	13.8	23.7	40.5	60.3	90.6	125.0	172.1	269.2
1999.....	3.7	7.7	13.5	23.8	42.0	62.0	90.0	126.9	170.8	270.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of colon, rectum and anus (C18-C21)

White, male

2004.....	3.9	7.8	15.1	26.0	41.9	64.0	95.3	135.9	181.7	255.4
2003.....	3.8	8.3	15.2	26.2	45.0	73.6	100.8	140.7	196.0	272.9
2002.....	4.5	8.4	16.5	28.7	49.4	76.0	105.4	144.9	193.3	273.6
2001.....	4.0	8.7	16.4	29.3	46.7	76.7	108.7	146.7	198.9	297.3
2000.....	3.9	8.3	16.1	28.6	49.1	75.3	113.5	157.6	208.6	313.3
1999.....	3.9	8.5	15.8	28.8	51.1	76.1	113.3	158.1	214.4	324.5

White, female

2004.....	3.4	5.9	10.3	16.8	27.1	39.8	60.5	90.3	131.8	211.4
2003.....	2.9	6.3	10.8	17.0	29.5	44.1	64.2	96.6	134.8	223.2
2002.....	3.3	6.5	12.0	18.2	29.3	45.7	67.6	96.9	140.1	226.4
2001.....	3.6	6.7	12.2	18.8	29.8	46.9	69.1	97.7	142.5	237.7
2000.....	3.3	6.6	11.5	19.1	32.6	47.2	72.3	102.1	150.4	251.3
1999.....	3.5	6.9	11.2	19.2	33.6	49.7	71.5	104.9	145.1	249.3

Black, both sexes

2004.....	5.9	12.2	21.9	36.1	56.7	79.9	113.2	148.6	189.1	260.8
2003.....	6.6	13.2	25.5	39.4	59.7	81.2	124.6	154.7	197.7	281.5
2002.....	7.2	14.5	23.5	40.0	61.9	86.8	114.9	160.5	218.7	274.7
2001.....	6.5	14.5	24.6	43.1	58.3	90.9	120.3	172.6	211.0	286.5
2000.....	6.6	15.2	24.8	39.9	62.7	86.9	126.1	167.4	226.2	310.0
1999.....	6.6	13.3	23.4	42.0	65.7	90.5	125.4	172.3	226.8	306.1

Black, male

2004.....	6.6	13.2	26.3	42.0	69.7	100.5	147.0	198.8	239.6	278.2
2003.....	7.1	15.2	30.3	46.1	76.4	102.7	156.7	194.9	242.2	317.0
2002.....	8.4	15.7	27.9	47.5	79.4	109.2	138.2	209.0	285.1	328.1
2001.....	7.6	16.4	26.4	51.3	67.7	114.1	146.8	208.2	263.2	347.2
2000.....	8.0	16.4	29.2	51.4	76.4	109.3	160.1	208.9	298.4	377.1
1999.....	6.3	15.0	26.9	54.1	80.4	110.6	147.6	228.0	274.7	350.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of colon, rectum and anus (C18-C21)

Black, female

2004.....	5.3	11.3	18.2	31.3	46.6	64.6	90.3	118.9	162.5	254.0
2003.....	6.1	11.4	21.4	33.8	46.8	65.2	103.0	130.8	174.5	267.8
2002.....	6.1	13.5	19.7	33.9	48.2	70.1	99.2	131.5	184.2	254.2
2001.....	5.6	12.8	23.1	36.3	50.9	73.8	102.7	151.4	184.4	264.0
2000.....	5.3	14.2	21.1	30.5	52.0	70.5	103.4	142.4	189.8	285.0
1999.....	6.7	11.9	20.5	32.1	54.2	75.6	110.7	138.7	203.1	289.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Colon (C18)

All races 3/, both sexes

2004.....	2.8	5.5	10.5	17.6	28.4	43.7	65.7	94.0	130.9	195.3
2003.....	2.8	6.1	11.2	17.9	31.1	49.4	70.1	99.8	137.6	208.4
2002.....	3.3	6.3	11.9	19.8	32.5	50.5	72.7	101.5	142.3	211.3
2001.....	3.1	6.4	11.8	20.4	32.4	52.4	75.6	103.9	143.5	221.6
2000.....	3.1	6.3	11.8	20.1	34.4	51.4	78.2	108.3	153.7	236.6
1999.....	3.1	6.4	11.6	20.4	35.9	53.3	77.7	110.7	151.6	238.1

All races 3/, male

2004.....	2.9	6.2	12.4	20.5	34.2	53.7	80.4	115.7	155.2	219.9
2003.....	3.0	6.8	12.8	21.2	37.1	62.2	85.5	121.3	168.6	236.1
2002.....	3.7	6.8	13.5	23.8	40.2	63.2	89.8	124.7	169.8	240.2
2001.....	3.2	7.1	12.9	24.5	38.5	64.5	92.6	126.3	173.0	256.7
2000.....	3.4	6.7	13.6	24.0	40.9	62.7	96.6	135.2	185.0	275.8
1999.....	3.2	7.0	13.2	24.2	43.3	64.3	96.0	137.4	187.2	282.2

All races 3/, female

2004.....	2.7	4.8	8.7	14.9	23.2	34.9	53.8	78.4	116.0	184.2
2003.....	2.5	5.4	9.6	14.8	25.7	38.3	57.6	84.5	118.7	196.1
2002.....	2.9	5.8	10.3	16.1	25.4	39.5	59.1	85.1	125.6	198.7
2001.....	3.0	5.7	10.7	16.7	26.9	42.0	62.1	88.3	125.8	206.9
2000.....	2.8	5.9	10.1	16.6	28.5	41.7	63.7	89.6	135.2	220.6
1999.....	3.0	5.9	10.1	16.8	29.3	43.8	63.4	92.1	130.8	220.4

White, both sexes

2004.....	2.6	5.1	9.7	16.5	26.8	41.6	63.3	92.0	129.3	194.8
2003.....	2.4	5.5	10.0	16.6	29.5	48.1	67.5	98.1	136.6	206.9
2002.....	3.0	5.5	11.1	18.6	30.8	48.6	71.0	99.4	140.0	210.5
2001.....	2.8	5.8	10.9	19.1	30.9	50.1	73.7	100.7	141.6	221.2
2000.....	2.8	5.6	11.0	18.8	33.0	49.9	76.4	106.4	150.8	235.0
1999.....	2.8	5.9	10.8	19.0	34.5	51.6	75.8	108.5	149.3	237.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Colon (C18)

White, male

2004.....	2.8	5.8	11.4	19.4	32.4	51.5	77.4	112.3	153.5	220.7
2003.....	2.8	6.3	11.7	20.0	35.0	60.7	83.1	119.2	167.3	236.2
2002.....	3.4	6.1	12.6	22.5	38.1	61.2	88.2	121.7	166.9	238.7
2001.....	2.9	6.4	12.1	23.2	37.1	62.2	91.1	123.1	170.6	255.5
2000.....	3.0	6.1	12.7	22.3	39.0	61.2	94.6	133.3	180.4	272.5
1999.....	3.0	6.5	12.6	22.6	41.6	62.7	95.0	134.3	185.6	282.9

White, female

2004.....	2.4	4.3	7.9	13.7	21.7	32.8	51.7	77.2	114.2	183.1
2003.....	2.1	4.8	8.5	13.4	24.5	36.9	54.8	82.8	117.7	193.9
2002.....	2.6	4.9	9.7	14.8	24.0	37.6	57.0	83.4	123.5	198.2
2001.....	2.7	5.1	9.6	15.1	25.1	39.5	59.7	84.8	124.1	206.9
2000.....	2.6	5.1	9.3	15.5	27.3	40.1	61.8	87.5	133.2	219.8
1999.....	2.6	5.3	9.2	15.5	27.9	41.8	60.6	90.5	127.8	218.7

Black, both sexes

2004.....	4.2	9.2	18.0	30.6	47.7	68.9	100.8	131.9	168.3	227.3
2003.....	5.2	11.0	20.8	32.2	50.8	70.1	108.9	135.1	175.7	249.4
2002.....	5.7	12.1	19.7	33.6	52.1	74.8	100.3	142.8	190.2	244.1
2001.....	5.5	11.8	20.5	36.2	51.3	80.1	105.9	154.9	182.6	249.9
2000.....	5.4	12.5	21.0	33.5	53.6	73.0	109.7	148.6	206.7	279.7
1999.....	5.5	11.4	19.6	35.7	56.4	78.1	109.4	151.6	204.0	273.9

Black, male

2004.....	3.9	9.3	22.1	34.8	57.9	84.6	128.2	173.8	208.8	243.4
2003.....	5.3	11.8	24.3	36.7	64.4	89.1	133.7	168.0	220.7	270.0
2002.....	6.3	12.1	23.0	39.3	66.3	94.2	119.2	186.9	238.4	292.7
2001.....	5.9	13.5	22.0	41.6	57.2	97.8	124.9	184.9	231.1	307.5
2000.....	6.6	12.9	24.3	41.5	64.4	86.9	136.1	184.3	271.4	346.0
1999.....	5.1	12.3	21.6	44.4	67.7	94.8	126.5	200.9	238.9	304.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Colon (C18)

Black, female

2004.....	4.4	9.1	14.6	27.0	39.8	57.2	82.2	107.0	147.0	220.9
2003.....	5.2	10.2	17.8	28.4	40.2	55.9	92.2	115.6	152.0	241.4
2002.....	5.1	12.0	16.9	29.0	41.0	60.3	87.6	116.4	165.1	225.5
2001.....	5.1	10.3	19.2	31.8	46.7	67.0	93.3	137.0	157.8	228.6
2000.....	4.4	12.3	18.1	27.0	45.1	62.8	92.1	127.0	174.1	255.0
1999.....	5.9	10.7	18.0	28.5	47.6	65.8	98.1	121.9	186.7	262.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Rectosigmoid junction and rectum (C19-C20)

All races 3/, both sexes

2004.....	1.0	1.6	2.7	4.4	7.0	9.0	11.9	16.3	20.3	29.1
2003.....	0.9	1.6	2.8	4.7	7.0	9.3	12.6	16.3	20.4	29.9
2002.....	0.9	1.8	2.8	4.5	7.7	10.6	12.9	16.7	19.9	29.0
2001.....	0.8	1.8	3.2	4.7	6.6	10.0	12.5	16.3	21.4	32.6
2000.....	0.7	1.7	2.6	4.7	7.2	10.1	13.6	17.7	20.3	32.6
1999.....	0.8	1.6	2.5	4.6	7.3	10.1	13.5	17.6	20.6	32.1

All races 3/, male

2004.....	1.1	2.0	3.4	6.1	9.1	12.0	17.0	22.5	27.5	33.5
2003.....	1.0	2.0	3.6	6.1	9.7	12.2	17.4	21.2	27.5	36.4
2002.....	1.1	2.2	3.6	6.1	10.8	14.2	16.6	22.4	26.9	33.6
2001.....	1.0	2.2	4.0	6.0	9.2	13.9	17.2	22.3	27.7	40.7
2000.....	0.9	2.2	3.2	6.3	9.7	14.2	18.7	23.5	27.0	38.9
1999.....	0.9	1.9	3.2	6.2	9.4	13.3	17.8	23.3	28.2	40.7

All races 3/, female

2004.....	0.9	1.3	2.1	2.8	5.0	6.3	7.8	11.9	16.0	27.2
2003.....	0.7	1.2	2.1	3.3	4.6	6.8	8.8	12.8	16.0	27.0
2002.....	0.6	1.4	2.0	3.0	4.8	7.5	9.9	12.6	15.6	27.0
2001.....	0.7	1.5	2.4	3.3	4.2	6.6	8.8	12.0	17.6	29.2
2000.....	0.6	1.3	2.0	3.3	4.9	6.6	9.6	13.7	16.4	30.1
1999.....	0.8	1.4	1.9	3.1	5.3	7.4	10.1	13.6	16.2	28.7

White, both sexes

2004.....	0.9	1.5	2.7	4.4	6.9	9.0	12.2	16.6	20.5	29.1
2003.....	0.9	1.6	2.7	4.5	6.9	9.4	12.5	16.2	20.5	30.1
2002.....	0.8	1.7	2.8	4.4	7.6	10.7	13.0	16.8	19.3	29.0
2001.....	0.8	1.8	3.1	4.5	6.6	10.1	12.4	16.4	21.1	32.6
2000.....	0.7	1.7	2.5	4.6	7.1	9.8	13.5	17.8	20.5	33.0
1999.....	0.8	1.6	2.4	4.5	7.1	10.0	13.5	17.5	20.7	32.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Rectosigmoid junction and rectum (C19-C20)

White, male

2004.....	0.9	1.8	3.4	6.1	9.0	11.9	17.4	22.7	27.6	33.6
2003.....	1.0	1.9	3.4	5.9	9.6	12.3	17.2	20.9	28.3	36.0
2002.....	1.0	2.0	3.6	5.9	10.8	14.3	16.7	22.7	25.6	33.7
2001.....	0.9	2.1	4.0	5.8	9.2	14.0	16.9	22.8	27.8	41.0
2000.....	0.9	2.1	3.1	5.9	9.7	13.6	18.4	23.7	27.2	39.8
1999.....	0.8	1.8	3.0	5.9	9.3	13.0	17.8	23.2	28.0	40.6

White, female

2004.....	0.9	1.2	2.0	2.7	4.9	6.4	8.0	12.2	16.1	27.1
2003.....	0.7	1.2	1.9	3.1	4.5	6.7	8.6	12.7	15.8	27.5
2002.....	0.6	1.4	1.9	2.9	4.7	7.4	9.9	12.5	15.4	27.0
2001.....	0.7	1.4	2.2	3.3	4.2	6.8	8.7	11.9	17.1	29.1
2000.....	0.6	1.3	1.9	3.3	4.6	6.5	9.7	13.7	16.5	30.3
1999.....	0.8	1.4	1.8	3.2	5.2	7.4	10.1	13.5	16.4	28.9

Black, both sexes

2004.....	1.6	2.7	3.5	5.3	8.6	10.4	11.9	15.7	20.8	31.9
2003.....	1.2	2.1	4.3	6.8	8.5	10.5	14.7	18.2	21.3	31.5
2002.....	1.3	2.3	3.5	6.0	9.0	11.6	13.9	17.2	27.7	29.7
2001.....	0.9	2.6	3.7	6.5	6.9	10.3	14.1	17.2	26.8	35.6
2000.....	1.0	2.5	3.4	6.1	8.8	13.6	15.6	17.6	19.2	28.8
1999.....	0.9	1.7	3.3	6.1	9.1	11.9	14.7	20.4	21.4	31.9

Black, male

2004.....	2.4	3.6	3.8	7.0	11.2	15.0	18.1	24.6	30.8	31.8
2003.....	1.5	3.2	5.5	9.1	12.0	12.3	22.7	25.6	21.4	45.9
2002.....	1.7	3.5	4.4	7.9	12.5	14.5	18.7	21.1	45.9	34.3
2001.....	1.6	2.8	4.2	9.2	10.4	15.8	21.5	22.8	30.5	38.6
2000.....	*	3.5	4.2	9.6	11.4	21.9	23.4	22.7	27.0	30.0
1999.....	*	2.5	4.7	9.5	12.5	15.6	20.4	27.1	34.9	45.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Rectosigmoid junction and rectum (C19-C20)										
Black, female										
2004.....	*	1.9	3.3	3.9	6.6	7.0	7.7	10.5	15.5	31.9
2003.....	*	*	3.3	4.8	5.9	9.1	9.3	13.8	21.3	25.9
2002.....	*	*	2.7	4.5	6.3	9.4	10.8	14.8	18.3	27.9
2001.....	*	2.3	3.3	4.2	4.2	6.2	9.2	13.9	24.9	34.5
2000.....	*	1.7	2.8	3.2	6.7	7.5	10.4	14.5	15.3	28.3
1999.....	*	*	2.1	3.3	6.5	9.2	11.0	16.3	14.7	26.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Anus and anus canal (C21)

All races 3/, both sexes

2004.....	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.9	1.0	1.2
2003.....	0.1	0.2	0.3	0.4	0.5	0.5	0.7	0.9	1.0	1.4
2002.....	0.1	0.2	0.3	0.3	0.6	0.5	0.6	0.8	1.0	1.2
2001.....	0.1	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.0	1.4
2000.....	0.1	0.2	0.3	0.3	0.5	0.5	0.7	0.9	0.8	1.2
1999.....	0.1	0.2	0.2	0.3	0.4	0.5	0.7	0.7	0.9	1.4

All races 3/, male

2004.....	*	*	0.3	0.4	0.5	0.6	0.6	0.8	*	*
2003.....	*	*	*	0.3	*	0.6	*	*	*	*
2002.....	*	0.2	0.3	*	0.5	0.5	*	*	*	*
2001.....	*	*	0.2	0.3	*	0.5	0.6	0.8	*	*
2000.....	*	*	0.3	0.4	0.4	0.5	0.5	0.7	*	*
1999.....	*	0.2	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	0.3	0.4	0.3	0.5	0.6	0.8	1.0	1.4	1.1
2003.....	*	0.2	0.4	0.5	0.6	0.5	0.8	1.1	1.3	1.6
2002.....	*	0.2	0.3	0.4	0.6	0.6	0.7	0.9	1.1	1.2
2001.....	*	*	0.4	0.4	0.4	0.6	0.7	0.9	1.2	1.7
2000.....	*	0.3	0.3	0.3	0.5	0.5	0.8	1.0	0.7	1.3
1999.....	*	*	0.3	0.4	0.4	0.5	0.8	0.9	1.0	1.6

White, both sexes

2004.....	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.9	1.1	1.1
2003.....	0.1	0.2	0.3	0.4	0.5	0.5	0.6	0.9	1.0	1.5
2002.....	0.1	0.2	0.4	0.4	0.6	0.6	0.6	0.8	1.0	1.2
2001.....	0.1	0.2	0.3	0.4	0.4	0.5	0.7	0.9	1.0	1.5
2000.....	0.1	0.2	0.3	0.3	0.5	0.5	0.7	0.8	0.8	1.1
1999.....	*	0.2	0.2	0.4	0.4	0.5	0.6	0.8	0.9	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of liver and intrahepatic bile ducts (C22)

All races 3/, both sexes

2004.....	1.2	3.9	7.6	9.4	12.5	16.6	23.7	28.8	34.6	32.4
2003.....	1.2	3.7	7.4	8.5	11.7	16.5	23.5	29.6	32.1	33.4
2002.....	1.3	3.4	6.6	8.1	11.6	16.9	22.6	27.6	31.6	35.0
2001.....	1.2	3.8	5.8	7.5	11.6	16.3	21.5	27.0	32.0	34.1
2000.....	1.4	3.7	5.4	7.7	11.3	16.0	21.6	27.1	31.3	32.7
1999.....	1.4	3.4	4.9	7.1	10.9	16.3	21.8	26.1	30.9	30.6

All races 3/, male

2004.....	1.8	6.2	12.8	15.1	18.6	23.6	34.0	38.8	48.4	41.5
2003.....	1.8	6.0	12.2	13.0	17.3	23.3	33.4	43.7	46.1	47.5
2002.....	2.0	5.7	11.1	12.4	17.4	24.8	32.8	39.5	44.8	46.2
2001.....	1.8	6.0	9.7	11.4	17.2	23.5	30.6	39.2	43.9	50.3
2000.....	2.0	6.0	8.5	11.7	16.5	23.6	30.9	39.3	46.7	44.7
1999.....	2.1	5.5	7.6	10.6	15.6	23.7	31.9	36.6	42.9	40.1

All races 3/, female

2004.....	0.6	1.8	2.6	4.1	6.9	10.6	15.4	21.7	26.1	28.3
2003.....	0.7	1.6	2.8	4.3	6.6	10.7	15.6	19.6	23.6	27.2
2002.....	0.6	1.2	2.4	4.0	6.4	10.1	14.4	19.2	23.6	30.2
2001.....	0.5	1.7	2.1	3.9	6.5	10.2	14.2	18.4	24.8	27.2
2000.....	0.8	1.4	2.4	4.0	6.5	9.5	14.2	18.6	22.2	27.8
1999.....	0.8	1.3	2.3	3.8	6.6	10.1	13.8	18.7	23.9	26.8

White, both sexes

2004.....	1.0	3.3	6.6	8.0	11.2	15.1	21.9	27.3	33.7	31.2
2003.....	1.0	3.1	6.3	7.5	10.3	15.2	21.7	28.3	31.0	32.1
2002.....	1.1	3.0	5.5	6.8	10.6	15.8	21.1	26.7	30.7	33.8
2001.....	0.8	3.2	4.9	6.4	10.5	15.1	20.2	26.1	30.4	33.3
2000.....	1.1	3.1	4.4	6.6	10.0	14.6	20.2	25.8	30.7	32.0
1999.....	1.1	2.7	4.2	5.9	9.7	15.2	20.5	25.2	30.2	29.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of liver and intrahepatic bile ducts (C22)

White, male

2004.....	1.5	5.0	11.1	12.4	16.5	21.5	31.1	37.4	47.2	39.9
2003.....	1.4	4.9	10.2	11.4	15.2	21.4	31.4	41.8	44.8	45.6
2002.....	1.6	4.9	9.0	10.4	15.6	23.0	30.4	38.0	42.9	45.1
2001.....	1.3	5.0	8.1	9.5	15.7	22.1	29.1	37.7	42.6	49.7
2000.....	1.5	4.8	6.8	9.6	14.4	21.4	28.9	38.1	45.8	43.1
1999.....	1.7	4.3	6.3	8.8	13.9	21.9	30.2	35.3	42.4	40.6

White, female

2004.....	0.5	1.5	2.2	3.7	6.3	9.5	14.3	19.8	25.3	27.3
2003.....	0.6	1.3	2.5	3.8	5.7	9.7	13.9	18.6	22.5	26.2
2002.....	0.5	1.1	2.0	3.4	5.9	9.3	13.5	18.6	23.2	29.0
2001.....	0.4	1.4	1.9	3.4	5.7	8.9	13.0	17.8	23.1	26.5
2000.....	0.6	1.3	2.0	3.6	5.9	8.5	13.2	17.1	21.7	27.4
1999.....	0.6	1.2	2.1	3.1	5.8	9.3	12.8	18.1	23.0	25.6

Black, both sexes

2004.....	1.8	7.1	12.7	17.9	16.6	23.2	28.5	33.7	35.4	36.1
2003.....	1.9	6.2	14.2	14.7	18.4	18.9	27.9	30.3	32.7	41.7
2002.....	2.1	5.8	12.3	15.5	14.0	19.5	27.3	30.2	31.9	38.0
2001.....	2.3	7.1	10.7	15.2	14.3	20.0	24.0	26.9	40.9	35.0
2000.....	2.7	7.0	10.6	12.7	15.8	20.9	26.9	28.8	28.5	31.6
1999.....	2.2	6.6	8.1	12.0	14.7	21.1	28.0	26.6	33.8	34.4

Black, male

2004.....	2.9	11.7	22.9	31.9	27.1	36.8	45.0	39.6	50.9	46.7
2003.....	3.0	10.3	25.5	24.0	28.5	26.8	42.4	43.8	45.0	63.6
2002.....	3.3	9.8	21.7	25.3	22.8	32.2	45.1	44.6	51.2	41.8
2001.....	3.6	11.6	18.6	25.3	21.4	28.0	32.7	39.5	49.0	42.0
2000.....	3.8	12.6	17.7	22.1	25.3	34.1	41.5	36.4	45.5	48.4
1999.....	3.4	11.8	14.2	18.6	21.6	32.2	42.1	40.4	42.7	30.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of liver and intrahepatic bile ducts (C22)

Black, female

2004.....	*	3.0	4.1	6.4	8.4	13.1	17.4	30.2	27.2	31.9
2003.....	*	2.7	4.5	7.0	10.5	13.0	18.2	22.2	26.3	33.2
2002.....	*	2.3	4.3	7.5	7.1	10.2	15.5	21.7	21.9	36.6
2001.....	*	3.2	3.9	6.8	8.8	14.1	18.2	19.4	36.8	32.4
2000.....	1.8	2.2	4.7	5.0	8.3	11.1	17.3	24.2	20.0	25.3
1999.....	*	2.0	2.9	6.5	9.3	12.9	18.7	18.3	29.4	35.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Liver (C22.0,C22.2-C22.9)

All races 3/, both sexes

2004.....	0.9	3.3	6.6	7.8	9.4	12.7	18.2	21.3	25.1	23.2
2003.....	1.0	3.2	6.3	6.9	8.9	12.8	18.1	22.6	23.6	24.3
2002.....	1.0	3.0	5.7	6.4	9.1	13.5	18.1	21.2	23.6	24.7
2001.....	0.9	3.4	5.0	6.2	9.2	12.8	16.7	20.6	23.8	24.2
2000.....	1.2	3.2	4.5	6.4	8.2	12.7	17.1	20.7	24.1	23.4
1999.....	1.2	3.0	4.1	5.7	8.7	13.0	17.5	20.4	23.3	22.5

All races 3/, male

2004.....	1.6	5.5	11.6	13.2	14.9	19.5	27.6	30.6	36.8	30.6
2003.....	1.6	5.4	11.1	11.1	14.1	19.0	26.6	35.1	35.9	35.2
2002.....	1.7	5.1	10.0	10.6	14.5	20.9	27.6	31.4	34.8	34.6
2001.....	1.5	5.6	8.7	10.0	14.5	19.5	24.9	31.3	33.6	38.5
2000.....	1.8	5.4	7.6	10.2	14.0	19.7	25.5	31.4	37.6	33.9
1999.....	1.9	5.1	6.9	8.9	13.4	19.9	26.7	29.9	34.2	31.2

All races 3/, female

2004.....	0.3	1.2	1.8	2.7	4.4	6.8	10.6	14.6	17.9	19.8
2003.....	0.4	1.0	1.7	3.0	4.2	7.5	11.3	13.6	16.2	19.5
2002.....	0.4	0.9	1.6	2.5	4.2	7.1	10.5	14.0	16.8	20.4
2001.....	0.4	1.2	1.5	2.5	4.5	7.1	10.1	13.1	17.8	18.2
2000.....	0.6	1.1	1.5	2.7	4.4	6.6	10.4	13.3	16.1	19.1
1999.....	0.6	0.9	1.5	2.7	4.5	7.1	10.3	13.7	16.9	18.9

White, both sexes

2004.....	0.8	2.7	5.7	6.4	8.0	11.3	16.4	19.8	24.2	21.9
2003.....	0.8	2.6	5.3	5.9	7.6	11.6	16.4	21.2	22.4	23.0
2002.....	0.8	2.6	4.6	5.3	8.1	12.4	16.6	20.4	22.8	23.7
2001.....	0.6	2.8	4.2	5.1	8.2	11.6	15.5	19.6	22.5	23.5
2000.....	0.8	2.6	3.5	5.3	7.7	11.2	15.6	19.5	23.4	22.5
1999.....	0.9	2.3	3.4	4.6	7.6	11.8	16.1	19.4	22.5	21.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Liver (C22.0,C22.2-C22.9)

White, male

2004.....	1.2	4.4	10.0	10.6	12.8	17.5	24.8	29.2	35.2	28.8
2003.....	1.2	4.4	9.2	9.4	12.2	17.3	24.8	33.1	34.4	33.1
2002.....	1.4	4.3	8.0	8.7	12.8	19.3	25.1	30.2	33.2	33.5
2001.....	1.0	4.6	7.1	8.2	13.2	18.1	23.6	29.9	32.4	37.8
2000.....	1.3	4.2	5.9	8.4	12.0	17.5	23.4	30.2	36.6	31.8
1999.....	1.5	3.8	5.6	7.2	11.6	18.2	24.9	28.7	33.8	31.4

White, female

2004.....	0.3	1.0	1.5	2.3	3.7	5.8	9.5	12.9	17.4	18.8
2003.....	0.3	0.7	1.4	2.5	3.4	6.5	9.6	12.6	14.9	18.5
2002.....	0.3	0.8	1.3	2.0	3.8	6.2	9.7	13.3	16.5	19.4
2001.....	0.2	1.0	1.3	2.1	3.6	5.8	8.9	12.4	16.5	17.5
2000.....	0.4	0.9	1.2	2.3	3.8	5.7	9.3	11.9	15.5	18.7
1999.....	0.4	0.8	1.3	2.1	3.8	6.3	9.1	13.0	15.9	17.8

Black, both sexes

2004.....	1.4	6.4	11.8	16.6	14.4	19.0	23.9	27.1	28.7	29.0
2003.....	1.8	5.4	12.8	13.3	15.5	14.8	23.4	24.0	27.2	33.0
2002.....	1.9	5.3	11.5	13.8	12.0	16.5	23.9	24.2	24.9	28.8
2001.....	2.1	6.5	9.9	13.4	12.1	16.7	20.5	22.8	31.4	27.0
2000.....	2.5	6.5	9.8	11.3	13.1	18.1	23.1	23.8	24.3	24.7
1999.....	1.9	6.3	7.6	10.4	13.1	17.9	24.6	22.7	26.9	28.4

Black, male

2004.....	2.5	11.0	21.8	30.6	24.1	31.5	40.1	32.8	46.6	36.8
2003.....	2.9	9.6	23.9	22.0	25.5	21.7	35.8	37.9	36.2	51.1
2002.....	3.2	9.2	21.0	23.8	20.4	27.4	40.1	36.3	41.3	35.4
2001.....	3.4	11.2	17.8	23.2	17.9	23.9	27.9	32.1	40.9	34.0
2000.....	3.4	12.0	17.0	20.3	23.0	30.7	36.4	30.7	40.5	42.7
1999.....	3.1	11.5	13.8	17.5	20.4	28.6	37.3	34.7	34.9	24.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Liver (C22.0,C22.2-C22.9)

Black, female

2004.....	*	2.4	3.2	5.1	6.8	9.7	13.0	23.7	19.3	26.0
2003.....	*	1.6	3.4	6.1	7.7	9.7	15.1	15.7	22.4	25.9
2002.....	*	1.9	3.5	5.5	5.4	8.3	13.0	17.0	16.3	26.3
2001.....	*	2.3	3.1	5.4	7.5	11.5	15.5	17.2	26.6	24.4
2000.....	1.6	1.8	3.7	4.0	5.2	8.8	14.2	19.7	16.1	18.0
1999.....	*	1.9	2.3	4.7	7.3	10.0	16.2	15.4	22.9	29.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intrahepatic bile duct carcinoma (C22.1)

All races 3/, both sexes

2004.....	0.2	0.6	1.0	1.6	3.1	3.9	5.5	7.5	9.5	9.2
2003.....	0.2	0.6	1.1	1.6	2.8	3.7	5.4	7.1	8.5	9.1
2002.....	0.2	0.4	0.9	1.6	2.5	3.4	4.5	6.4	8.0	10.3
2001.....	0.2	0.4	0.8	1.3	2.3	3.5	4.8	6.4	8.2	9.9
2000.....	0.2	0.5	0.9	1.4	2.3	3.3	4.5	6.4	7.2	9.3
1999.....	0.2	0.4	0.7	1.4	2.2	3.3	4.3	5.7	7.7	8.1

All races 3/, male

2004.....	0.3	0.7	1.1	1.9	3.7	4.1	6.4	8.2	11.5	10.9
2003.....	0.2	0.6	1.1	1.9	3.2	4.3	6.7	8.6	10.2	12.4
2002.....	0.3	0.5	1.0	1.8	2.8	3.9	5.3	8.0	10.0	11.6
2001.....	0.3	0.4	1.0	1.3	2.7	4.0	5.7	7.9	10.3	11.9
2000.....	0.3	0.6	0.9	1.4	2.5	3.9	5.5	7.9	9.0	10.8
1999.....	0.2	0.4	0.7	1.7	2.3	3.8	5.2	6.7	8.8	8.9

All races 3/, female

2004.....	0.2	0.5	0.8	1.4	2.6	3.8	4.8	7.0	8.2	8.5
2003.....	0.3	0.6	1.1	1.3	2.4	3.2	4.3	6.0	7.5	7.6
2002.....	0.2	0.3	0.8	1.5	2.3	3.0	3.9	5.3	6.8	9.8
2001.....	0.2	0.4	0.6	1.3	2.0	3.0	4.1	5.3	7.0	9.0
2000.....	0.2	0.4	0.9	1.3	2.2	2.8	3.8	5.4	6.1	8.8
1999.....	0.2	0.4	0.8	1.1	2.1	3.0	3.6	5.0	7.0	7.8

White, both sexes

2004.....	0.2	0.6	0.9	1.6	3.2	3.8	5.5	7.5	9.5	9.3
2003.....	0.3	0.5	1.0	1.6	2.7	3.6	5.3	7.1	8.7	9.2
2002.....	0.2	0.4	0.8	1.5	2.5	3.4	4.5	6.4	7.9	10.2
2001.....	0.2	0.4	0.8	1.3	2.3	3.5	4.7	6.4	7.9	9.8
2000.....	0.2	0.5	0.9	1.3	2.2	3.4	4.6	6.3	7.3	9.5
1999.....	0.2	0.4	0.7	1.3	2.2	3.3	4.4	5.7	7.7	8.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of gallbladder and extrahepatic bile ducts (C23-C24)

All races 3/, both sexes

2004.....	0.2	0.2	0.6	1.3	2.3	3.6	5.9	8.0	9.7	13.3
2003.....	0.1	0.4	0.8	1.2	2.4	3.3	5.3	8.1	10.9	14.3
2002.....	0.2	0.4	0.8	1.4	2.0	3.7	5.7	8.3	9.9	14.5
2001.....	0.2	0.5	0.8	1.4	2.3	3.7	6.0	8.3	12.1	15.8
2000.....	0.2	0.4	0.9	1.5	2.6	4.2	6.4	9.1	11.4	16.0
1999.....	0.2	0.4	0.8	1.6	2.5	4.2	6.0	8.8	11.9	16.2

All races 3/, male

2004.....	*	0.2	0.5	1.2	1.7	3.2	5.2	8.1	9.4	12.9
2003.....	*	0.4	0.6	1.0	2.0	3.4	5.1	7.6	9.8	14.1
2002.....	*	0.2	0.6	1.2	1.8	3.3	5.5	8.2	9.1	15.0
2001.....	*	0.3	0.7	1.2	1.7	2.8	5.3	8.1	12.8	15.8
2000.....	*	0.3	0.6	1.1	2.2	3.9	6.2	8.6	11.0	15.8
1999.....	0.2	0.3	0.6	1.2	2.1	3.8	5.4	8.3	11.0	15.2

All races 3/, female

2004.....	0.2	0.3	0.8	1.5	2.9	4.0	6.5	7.9	10.0	13.5
2003.....	*	0.5	0.9	1.4	2.7	3.1	5.5	8.5	11.5	14.4
2002.....	0.3	0.6	1.0	1.6	2.2	4.1	5.8	8.3	10.3	14.3
2001.....	0.3	0.6	0.9	1.5	2.8	4.5	6.5	8.5	11.7	15.8
2000.....	0.2	0.5	1.1	1.8	2.9	4.5	6.6	9.4	11.7	16.0
1999.....	0.3	0.4	1.0	2.0	2.9	4.6	6.5	9.1	12.4	16.7

White, both sexes

2004.....	0.2	0.2	0.6	1.3	2.2	3.4	5.8	7.9	9.6	13.4
2003.....	*	0.4	0.7	1.2	2.4	3.3	5.3	8.0	10.7	14.4
2002.....	0.2	0.4	0.8	1.3	2.0	3.7	5.6	8.1	9.8	14.5
2001.....	0.2	0.5	0.8	1.4	2.2	3.5	5.9	8.3	12.2	15.8
2000.....	0.2	0.4	0.8	1.5	2.5	4.2	6.3	9.2	11.6	16.3
1999.....	0.2	0.3	0.8	1.6	2.5	4.2	6.0	8.7	11.5	16.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of gallbladder and extrahepatic bile ducts (C23-C24)

Black, female

2004.....	*	*	*	*	4.5	5.8	6.8	7.3	10.2	14.2
2003.....	*	*	*	*	*	*	5.5	7.9	12.8	14.2
2002.....	*	*	*	2.6	*	5.6	6.0	10.7	9.9	16.4
2001.....	*	*	*	*	4.2	6.4	6.5	7.5	10.2	16.0
2000.....	*	*	2.2	*	3.6	6.9	6.0	7.7	9.3	12.0
1999.....	*	*	*	2.9	*	4.0	6.5	8.3	13.8	14.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pancreas (C25)										
All races 3/, both sexes										
2004.....	1.7	4.2	8.8	15.2	25.8	37.3	52.4	68.9	85.4	97.7
2003.....	1.7	4.1	8.7	14.7	24.7	36.4	53.0	67.6	86.2	95.2
2002.....	1.9	4.2	8.2	15.5	24.0	37.2	53.2	70.2	81.8	94.5
2001.....	1.8	4.0	8.6	14.9	24.8	37.0	52.8	67.2	84.2	100.1
2000.....	1.8	4.2	8.7	15.1	24.8	37.2	51.4	70.4	85.7	96.4
1999.....	1.9	4.5	8.7	15.6	25.0	38.1	52.5	67.2	82.0	99.7
All races 3/, male										
2004.....	2.2	5.5	11.0	18.6	31.5	45.2	61.6	77.7	94.8	102.0
2003.....	2.0	5.2	10.7	18.2	29.8	42.5	60.9	76.1	99.5	95.6
2002.....	2.4	5.3	10.6	19.2	29.3	43.3	60.8	78.9	91.9	100.9
2001.....	2.0	4.9	10.7	18.5	30.3	43.6	59.0	77.2	94.9	105.1
2000.....	2.1	5.4	11.1	18.5	30.0	44.7	57.0	80.4	94.9	102.9
1999.....	2.4	5.9	11.0	19.1	29.9	44.9	60.9	75.7	92.0	107.5
All races 3/, female										
2004.....	1.3	2.9	6.7	12.0	20.6	30.5	45.1	62.6	79.6	95.8
2003.....	1.4	3.0	6.8	11.5	20.2	31.1	46.7	61.6	78.1	95.0
2002.....	1.4	3.1	6.0	12.0	19.2	31.9	47.1	64.1	75.6	91.7
2001.....	1.6	3.1	6.5	11.5	19.8	31.3	47.8	60.1	77.8	98.1
2000.....	1.5	3.0	6.4	11.8	20.0	30.8	47.1	63.5	80.2	93.8
1999.....	1.4	3.1	6.5	12.3	20.5	32.2	46.0	61.2	76.1	96.6
White, both sexes										
2004.....	1.6	4.0	8.4	14.6	25.1	36.8	51.3	68.3	85.1	97.1
2003.....	1.6	3.9	8.1	14.3	23.8	36.0	52.2	67.2	85.2	94.4
2002.....	1.9	4.1	7.8	15.0	23.3	36.1	52.0	69.2	81.4	94.3
2001.....	1.7	3.8	8.3	14.4	24.7	36.3	51.7	66.3	83.3	99.0
2000.....	1.7	4.1	8.4	14.5	24.3	36.2	50.4	69.2	84.5	94.7
1999.....	1.8	4.4	8.2	14.8	24.3	37.2	51.5	65.8	80.6	98.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pancreas (C25)										
White, male										
2004.....	2.0	5.3	10.5	17.7	30.8	44.5	60.0	77.7	95.8	102.9
2003.....	1.8	4.8	10.0	17.7	28.9	42.7	60.2	76.1	99.3	96.0
2002.....	2.4	5.1	10.1	18.6	28.8	42.6	59.7	78.9	92.2	101.6
2001.....	1.9	4.7	10.5	18.2	30.3	42.8	58.0	77.0	94.5	106.5
2000.....	2.0	5.3	10.7	17.8	29.6	44.3	55.9	79.9	94.0	102.4
1999.....	2.3	5.8	10.3	17.9	29.2	44.0	60.1	74.6	90.9	107.4
White, female										
2004.....	1.3	2.8	6.2	11.5	19.7	30.0	44.1	61.5	78.5	94.5
2003.....	1.4	2.9	6.3	11.0	19.1	30.0	45.8	60.7	76.6	93.8
2002.....	1.4	3.0	5.5	11.6	18.2	30.3	45.7	62.2	74.9	91.1
2001.....	1.6	2.9	6.3	10.8	19.6	30.6	46.5	58.7	76.5	95.9
2000.....	1.4	2.9	6.1	11.4	19.4	29.2	46.0	61.6	78.9	91.5
1999.....	1.3	3.1	6.3	11.9	19.9	31.3	44.6	59.5	74.5	95.0
Black, both sexes										
2004.....	2.6	6.0	13.3	22.6	36.3	48.0	70.5	84.3	93.3	113.6
2003.....	2.7	6.4	14.5	20.7	36.1	47.1	68.0	83.6	107.4	112.9
2002.....	2.5	5.6	12.3	22.4	34.4	50.4	73.7	88.6	91.8	106.6
2001.....	2.2	6.4	12.1	20.7	31.6	48.5	70.8	84.9	104.8	121.3
2000.....	2.6	5.8	13.2	22.8	33.3	50.3	69.8	93.8	104.2	119.8
1999.....	2.8	6.2	14.1	25.0	32.2	50.9	70.1	87.5	104.6	118.6
Black, male										
2004.....	3.4	7.6	16.1	30.0	44.8	60.6	88.1	88.7	91.1	100.3
2003.....	3.5	9.0	18.6	26.2	42.8	50.0	80.8	89.0	112.2	104.3
2002.....	2.9	7.2	16.5	29.0	40.8	56.6	84.1	90.5	105.5	100.8
2001.....	2.3	7.9	15.2	24.2	38.2	60.0	78.8	95.2	109.2	99.8
2000.....	3.0	7.3	17.5	29.3	40.4	55.7	78.9	101.6	116.3	109.6
1999.....	3.4	8.3	19.5	34.1	39.5	60.0	78.4	100.7	117.7	114.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of pancreas (C25)

Black, female

2004.....	1.9	4.6	11.0	16.6	29.7	38.7	58.6	81.7	94.5	118.9
2003.....	2.1	4.1	11.0	16.2	30.8	45.0	59.4	80.3	104.8	116.3
2002.....	2.1	4.3	8.8	17.0	29.4	45.9	66.8	87.5	84.7	108.8
2001.....	2.2	5.1	9.6	17.8	26.4	40.0	65.5	78.8	102.6	129.2
2000.....	2.3	4.4	9.4	17.4	27.8	46.3	63.7	89.1	98.1	123.6
1999.....	2.2	4.3	9.5	17.6	26.5	44.3	64.6	79.5	98.1	120.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other and ill-defined digestive organs (C26)

All races 3/, both sexes

2004.....	0.1	0.1	0.2	0.4	0.7	1.0	1.4	2.3	3.1	5.1
2003.....	*	0.2	0.2	0.5	0.7	1.5	1.9	2.9	4.0	7.0
2002.....	*	0.1	0.2	0.5	0.7	1.3	2.0	3.4	4.1	6.9
2001.....	*	0.2	0.3	0.5	0.9	1.3	2.2	3.2	4.3	7.8
2000.....	*	0.2	0.3	0.5	1.0	1.3	2.1	2.8	4.5	7.8
1999.....	0.1	0.3	0.3	0.6	1.0	1.6	2.3	3.3	5.3	9.1

All races 3/, male

2004.....	*	*	0.2	0.4	0.9	1.3	1.7	2.9	3.5	5.0
2003.....	*	*	0.3	0.7	0.9	2.0	2.2	3.8	4.5	7.3
2002.....	*	*	0.3	0.7	0.8	2.0	2.6	4.3	5.3	6.4
2001.....	*	0.2	0.4	0.7	1.3	1.4	2.7	4.1	4.5	8.9
2000.....	*	*	0.3	0.7	1.3	1.6	3.1	3.4	5.2	7.5
1999.....	0.2	0.3	0.5	0.7	1.3	1.9	2.6	4.2	6.9	10.4

All races 3/, female

2004.....	*	*	*	0.4	0.5	0.8	1.3	1.9	2.8	5.2
2003.....	*	*	*	0.4	0.5	1.0	1.6	2.3	3.7	6.8
2002.....	*	*	*	0.3	0.7	0.8	1.5	2.7	3.4	7.2
2001.....	*	*	*	0.3	0.7	1.2	1.8	2.6	4.2	7.3
2000.....	*	*	0.2	0.4	0.7	1.0	1.4	2.3	4.1	8.0
1999.....	*	*	*	0.4	0.7	1.4	2.1	2.7	4.4	8.5

White, both sexes

2004.....	*	0.1	0.2	0.4	0.7	0.9	1.4	2.3	3.2	5.3
2003.....	*	0.1	0.2	0.5	0.6	1.5	1.9	2.9	3.9	7.0
2002.....	*	*	0.2	0.5	0.7	1.3	1.9	3.5	4.0	7.0
2001.....	*	0.1	0.2	0.5	0.8	1.3	2.2	3.2	4.1	7.7
2000.....	*	0.2	0.2	0.6	1.0	1.2	2.0	2.8	4.3	7.9
1999.....	0.1	0.2	0.3	0.5	1.0	1.6	2.3	3.3	5.5	9.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)

All races 3/, both sexes

2004.....	8.8	21.6	42.2	81.0	149.6	232.7	318.6	381.1	383.2	306.3
2003.....	9.0	22.2	42.4	84.8	154.3	236.8	321.9	384.6	389.5	307.9
2002.....	9.6	21.5	43.4	89.5	157.8	239.7	328.9	386.5	387.5	305.1
2001.....	10.0	21.6	44.1	92.3	159.7	246.4	331.7	382.6	377.8	312.2
2000.....	9.8	22.0	45.6	95.9	165.1	252.9	333.8	377.8	385.1	310.8
1999.....	9.7	20.9	46.4	96.7	166.0	250.7	331.0	370.6	372.2	306.6

All races 3/, male

2004.....	9.6	25.1	52.6	100.8	187.3	296.4	413.5	523.7	545.3	485.7
2003.....	10.2	26.3	52.9	106.1	192.7	303.6	424.2	529.4	559.7	493.3
2002.....	10.1	25.3	54.2	112.3	201.0	307.8	436.0	532.5	563.8	498.8
2001.....	11.2	25.9	55.2	115.5	204.0	323.8	449.3	538.5	558.1	534.1
2000.....	11.2	27.2	56.2	121.6	211.2	333.9	455.0	533.6	578.3	541.7
1999.....	11.2	25.9	57.7	124.3	216.4	334.1	453.2	529.0	569.3	550.6

All races 3/, female

2004.....	8.1	18.3	32.3	62.3	115.3	177.3	242.2	278.7	283.2	225.6
2003.....	7.8	18.2	32.5	64.8	119.5	178.8	239.8	281.5	285.4	225.9
2002.....	9.0	17.8	32.9	68.0	118.6	180.9	243.4	283.4	280.3	221.1
2001.....	8.7	17.3	33.6	70.6	119.5	179.8	238.3	273.4	269.7	219.3
2000.....	8.4	17.0	35.5	71.8	123.3	183.5	238.3	269.3	271.1	216.8
1999.....	8.4	16.1	35.5	70.9	120.4	179.4	235.5	260.5	256.8	208.3

White, both sexes

2004.....	8.5	20.7	40.3	78.9	149.9	235.0	325.1	386.6	386.8	310.7
2003.....	8.7	20.7	40.6	83.5	154.9	239.9	327.1	389.3	393.6	311.2
2002.....	9.1	20.4	41.6	87.8	157.2	242.4	334.8	391.2	393.1	309.0
2001.....	9.5	20.2	42.6	91.1	160.7	248.3	335.4	385.9	380.2	314.8
2000.....	9.0	20.4	43.8	94.4	165.8	254.9	335.8	379.8	391.0	313.4
1999.....	9.2	19.3	44.3	95.3	165.4	251.2	333.1	371.7	374.7	308.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)

White, male										
2004.....	9.1	23.3	48.6	95.8	182.5	294.8	414.0	523.5	544.6	490.5
2003.....	9.5	24.2	49.3	101.6	188.7	301.6	424.3	527.2	559.1	496.2
2002.....	9.4	23.3	50.6	107.5	194.8	305.5	436.9	530.6	564.7	503.6
2001.....	10.4	23.6	51.5	110.8	201.1	319.8	446.3	534.5	554.0	535.5
2000.....	9.8	24.8	52.0	116.2	208.2	329.8	450.1	527.8	580.1	543.3
1999.....	10.2	23.2	53.4	119.7	210.9	328.7	448.4	524.2	565.4	551.9
White, female										
2004.....	7.9	18.1	32.1	62.5	119.7	182.0	252.0	286.5	288.7	229.6
2003.....	7.8	17.3	32.1	66.1	123.6	185.4	247.6	289.6	291.5	229.3
2002.....	8.9	17.5	32.7	68.9	122.4	186.8	251.5	291.5	287.9	224.5
2001.....	8.6	16.9	33.8	72.2	123.4	185.6	245.6	280.6	275.2	222.3
2000.....	8.2	16.0	35.7	73.7	126.7	189.5	244.1	275.7	278.5	220.1
1999.....	8.2	15.4	35.5	72.1	123.5	183.7	241.5	264.7	262.1	210.1
Black, both sexes										
2004.....	13.3	32.6	66.9	120.0	188.5	267.0	322.3	395.4	400.8	287.6
2003.....	13.2	37.4	65.8	121.3	193.3	268.4	343.1	401.4	391.2	293.2
2002.....	14.9	34.8	68.4	127.3	203.6	278.0	343.7	407.8	378.3	278.0
2001.....	15.4	35.0	67.8	127.4	191.9	284.8	363.2	410.1	381.9	290.2
2000.....	16.8	37.8	74.0	131.6	200.8	291.8	375.8	412.1	357.5	295.9
1999.....	15.7	37.5	75.3	133.2	209.9	294.7	364.5	414.0	375.3	300.4
Black, male										
2004.....	15.5	43.4	96.6	172.9	287.8	388.4	492.0	626.9	656.4	508.6
2003.....	16.7	48.1	93.7	179.7	286.1	401.1	522.9	650.0	662.2	530.7
2002.....	18.1	46.6	98.0	184.4	311.8	417.4	531.1	666.4	642.7	498.5
2001.....	19.6	48.9	100.0	189.0	285.8	440.1	582.0	691.4	685.4	571.9
2000.....	23.1	51.1	108.6	200.9	294.2	453.8	593.7	688.0	645.7	584.7
1999.....	20.7	53.3	111.5	201.4	321.8	451.3	581.4	680.1	694.9	599.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)

Black, female

2004.....	11.4	23.3	41.8	76.5	111.2	176.8	207.1	258.2	266.1	200.0
2003.....	10.1	28.1	42.3	73.3	120.8	170.0	221.8	253.7	249.1	200.9
2002.....	12.0	24.6	43.2	80.4	118.9	174.6	218.0	253.3	240.7	193.5
2001.....	11.7	23.0	40.3	77.0	118.1	170.1	217.5	242.1	227.3	185.7
2000.....	11.3	26.3	44.4	74.8	127.0	172.2	231.1	245.8	212.3	188.4
1999.....	11.3	23.9	44.7	77.5	122.1	178.7	220.5	253.7	216.9	188.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of nasal cavity, middle ear and accessory sinuses (C30-C31)

All races 3/, both sexes

2004.....	*	0.1	0.1	0.3	0.3	0.5	0.7	0.8	1.1	1.4
2003.....	0.1	0.1	0.2	0.2	0.4	0.5	0.6	1.0	0.9	1.2
2002.....	*	0.2	0.2	0.2	0.4	0.4	0.7	0.7	1.0	1.2
2001.....	0.1	0.1	0.2	0.3	0.4	0.4	0.6	0.8	1.4	1.7
2000.....	0.1	*	0.2	0.3	0.4	0.5	0.6	0.6	0.9	1.2
1999.....	0.1	0.1	0.3	0.3	0.3	0.5	0.6	0.8	1.0	1.5

All races 3/, male

2004.....	*	*	*	0.4	0.4	0.5	1.0	1.0	1.5	1.8
2003.....	*	*	0.3	0.3	0.7	0.7	0.9	1.4	1.3	*
2002.....	*	0.3	0.3	0.3	0.6	0.7	1.0	1.0	1.1	1.7
2001.....	*	0.2	0.3	0.4	0.5	0.5	0.8	0.9	1.9	1.8
2000.....	*	*	0.2	0.4	0.6	0.7	0.9	1.0	1.1	*
1999.....	*	*	0.4	0.5	0.4	0.7	0.6	1.2	1.1	1.8

All races 3/, female

2004.....	*	*	*	*	*	0.4	*	0.6	0.9	1.2
2003.....	*	*	*	*	*	*	*	0.7	0.6	1.1
2002.....	*	*	*	*	*	*	0.5	0.6	0.8	1.0
2001.....	*	*	*	*	*	*	0.4	0.8	1.1	1.6
2000.....	*	*	*	*	*	*	0.4	*	0.8	1.1
1999.....	*	*	*	*	*	*	0.6	0.6	0.9	1.4

White, both sexes

2004.....	*	*	0.1	0.3	0.3	0.4	0.7	0.8	1.1	1.4
2003.....	0.1	*	0.2	0.2	0.5	0.4	0.7	0.9	0.9	1.2
2002.....	*	0.2	0.2	0.3	0.4	0.4	0.7	0.8	1.0	1.3
2001.....	*	0.2	0.2	0.3	0.4	0.4	0.5	0.8	1.4	1.7
2000.....	*	*	0.2	0.3	0.4	0.5	0.6	0.7	0.9	1.3
1999.....	0.1	*	0.3	0.3	0.3	0.5	0.7	0.9	1.0	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of larynx (C32)										
All races 3/, both sexes										
2004.....	0.2	0.7	1.4	2.6	4.1	5.3	6.2	7.1	6.0	6.5
2003.....	0.3	0.7	1.4	2.6	4.6	5.8	6.6	6.6	7.4	6.5
2002.....	0.2	0.7	1.4	2.7	4.2	5.3	6.6	7.9	7.2	5.5
2001.....	0.3	0.7	1.5	2.9	4.5	5.6	7.3	7.4	6.9	6.7
2000.....	0.2	0.8	1.5	2.9	4.5	6.0	7.3	8.0	8.1	6.5
1999.....	0.3	0.8	1.5	3.1	4.6	6.1	7.4	7.2	7.4	7.0
All races 3/, male										
2004.....	0.4	1.2	2.4	4.4	6.8	9.4	11.0	13.3	12.4	15.6
2003.....	0.5	1.2	2.4	4.2	7.9	10.3	11.4	12.6	15.4	15.3
2002.....	0.4	1.2	2.4	4.7	7.2	8.9	11.3	14.8	14.9	13.2
2001.....	0.5	1.2	2.5	4.9	7.5	9.5	12.9	14.5	13.9	18.3
2000.....	0.4	1.2	2.5	4.9	7.9	10.0	12.9	14.8	17.1	17.4
1999.....	0.4	1.3	2.6	5.4	7.8	10.4	12.6	13.2	15.8	17.8
All races 3/, female										
2004.....	*	0.2	0.5	0.9	1.6	1.7	2.2	2.6	2.1	2.4
2003.....	*	0.3	0.4	1.0	1.6	2.0	2.8	2.3	2.5	2.7
2002.....	*	0.2	0.5	0.9	1.5	2.2	2.9	3.1	2.5	2.2
2001.....	*	0.2	0.5	1.0	1.7	2.2	2.8	2.5	2.8	1.9
2000.....	*	0.3	0.6	0.9	1.4	2.6	2.8	3.3	2.8	2.1
1999.....	*	0.2	0.4	1.0	1.7	2.5	3.3	3.0	2.6	2.6
White, both sexes										
2004.....	0.2	0.7	1.2	2.3	3.8	5.0	5.9	6.9	5.7	6.3
2003.....	0.3	0.6	1.1	2.3	4.2	5.5	6.2	6.3	7.1	6.2
2002.....	0.2	0.6	1.2	2.5	3.9	5.0	6.4	7.5	6.9	5.6
2001.....	0.3	0.6	1.2	2.5	4.1	5.2	6.7	7.0	6.8	6.6
2000.....	0.2	0.6	1.3	2.4	3.9	5.6	6.7	7.5	8.0	6.6
1999.....	0.2	0.6	1.3	2.7	4.2	5.7	7.1	7.1	7.3	6.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of larynx (C32)

White, male

2004.....	0.3	1.2	1.9	3.9	6.2	8.8	10.4	12.8	11.7	15.0
2003.....	0.5	1.1	2.0	3.7	7.2	9.3	10.4	11.9	14.4	14.5
2002.....	0.3	1.0	2.0	4.1	6.3	8.3	10.7	13.8	14.6	13.3
2001.....	0.4	0.9	2.0	4.2	6.9	8.8	11.9	13.3	13.2	18.3
2000.....	0.3	1.0	2.1	4.1	6.8	9.2	11.7	13.9	16.8	17.6
1999.....	0.3	1.1	2.2	4.6	7.0	9.6	12.0	12.9	15.1	16.8

White, female

2004.....	*	0.2	0.5	0.9	1.6	1.7	2.1	2.6	2.0	2.5
2003.....	*	*	0.3	0.9	1.4	2.1	2.7	2.3	2.5	2.5
2002.....	*	*	0.4	0.9	1.5	2.1	2.8	2.9	2.3	2.3
2001.....	*	*	0.4	0.9	1.5	2.1	2.6	2.5	2.9	1.7
2000.....	*	0.3	0.5	0.9	1.3	2.5	2.6	3.0	2.7	2.1
1999.....	*	*	0.3	1.0	1.5	2.3	3.3	3.0	2.7	2.7

Black, both sexes

2004.....	*	1.2	3.5	5.5	8.0	9.3	11.6	11.2	11.1	8.2
2003.....	*	1.7	3.6	5.8	9.7	10.6	12.7	11.1	12.7	11.1
2002.....	*	1.9	3.6	6.1	9.3	9.1	11.8	15.3	11.3	*
2001.....	*	1.9	3.8	7.1	8.9	9.5	13.9	13.4	9.2	8.3
2000.....	*	2.0	4.1	7.5	10.8	10.6	15.3	15.6	11.6	6.9
1999.....	0.9	2.1	3.9	7.5	10.0	11.4	11.8	10.2	9.8	8.3

Black, male

2004.....	*	2.1	6.7	11.0	14.9	18.7	22.7	24.1	25.8	23.8
2003.....	*	2.7	6.8	10.7	17.5	22.7	25.3	25.6	31.7	27.1
2002.....	*	3.2	6.0	11.5	18.8	17.0	22.4	33.1	22.9	*
2001.....	*	3.6	7.0	13.0	15.9	17.5	26.3	30.7	22.5	*
2000.....	*	3.2	7.6	14.3	20.7	19.8	30.1	29.3	27.0	*
1999.....	1.6	3.9	6.9	14.4	17.8	21.0	22.7	19.9	27.9	24.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of larynx (C32)										
Black, female										
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	3.6	*	*	*	*	*
2002.....	*	*	*	*	*	*	4.7	*	*	*
2001.....	*	*	*	*	3.4	*	5.6	*	*	*
2000.....	*	*	*	*	*	3.8	5.5	7.4	*	*
1999.....	*	*	*	*	3.8	4.2	4.5	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of trachea, bronchus and lung (C33-C34)

All races 3/, both sexes

2004.....	8.5	20.6	40.4	77.9	144.9	226.5	311.2	372.5	375.1	297.5
2003.....	8.5	21.2	40.7	81.8	149.0	230.0	314.0	376.1	380.1	298.9
2002.....	9.2	20.4	41.6	86.3	152.9	233.4	320.9	376.9	378.5	297.2
2001.....	9.5	20.5	42.3	88.8	154.5	239.7	323.2	373.4	368.6	302.7
2000.....	9.3	21.0	43.8	92.5	159.8	245.8	325.4	368.2	374.7	302.1
1999.....	9.3	19.9	44.4	93.0	160.7	243.6	322.3	361.7	362.6	297.0

All races 3/, male

2004.....	9.1	23.5	49.8	95.8	179.7	285.8	400.8	508.4	530.2	466.8
2003.....	9.5	24.8	50.0	101.3	183.7	292.1	411.0	514.2	541.5	475.1
2002.....	9.6	23.6	51.4	107.0	192.7	297.5	423.0	515.5	546.8	482.2
2001.....	10.5	24.3	52.2	109.6	195.7	312.9	434.7	522.3	541.0	512.4
2000.....	10.5	25.7	53.2	115.9	202.3	322.5	440.5	516.5	558.1	521.2
1999.....	10.5	24.3	54.4	118.1	207.7	322.4	439.2	513.4	551.3	529.2

All races 3/, female

2004.....	7.9	17.8	31.5	61.1	113.2	174.9	239.1	274.9	279.5	221.3
2003.....	7.5	17.8	31.9	63.5	117.4	176.2	236.1	277.9	281.4	221.0
2002.....	8.8	17.3	32.2	66.8	116.8	177.9	239.3	279.1	276.1	217.0
2001.....	8.4	16.8	32.8	69.2	117.2	176.7	234.5	269.1	265.2	215.0
2000.....	8.1	16.5	34.7	70.5	121.3	180.2	234.7	264.9	266.6	212.8
1999.....	8.1	15.7	34.9	69.5	118.1	176.1	230.9	256.2	252.2	203.5

White, both sexes

2004.....	8.1	19.7	38.8	76.1	145.5	229.2	318.0	378.3	378.9	302.0
2003.....	8.2	19.9	39.1	80.7	149.9	233.5	319.7	381.3	384.5	302.4
2002.....	8.8	19.4	40.0	84.9	152.6	236.4	327.0	382.1	384.1	300.9
2001.....	9.1	19.3	40.9	87.9	155.9	242.0	327.4	377.2	371.2	305.4
2000.....	8.6	19.5	42.2	91.5	161.2	248.2	328.0	370.7	380.7	304.5
1999.....	8.8	18.4	42.6	92.0	160.6	244.5	324.6	362.9	365.2	298.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of trachea, bronchus and lung (C33-C34)

White, male

2004.....	8.5	21.8	46.4	91.4	175.5	284.9	401.9	508.8	530.1	472.2
2003.....	8.9	22.8	46.9	97.3	180.4	291.1	412.1	513.0	541.9	478.5
2002.....	8.9	21.8	48.1	102.8	187.3	295.9	424.5	514.5	547.8	486.6
2001.....	9.7	22.2	49.0	105.6	193.4	309.8	432.8	519.4	537.4	513.9
2000.....	9.3	23.5	49.5	111.5	200.5	319.1	436.8	511.6	560.0	522.4
1999.....	9.7	21.9	50.5	114.3	203.1	317.9	435.0	508.8	548.0	531.5

White, female

2004.....	7.7	17.7	31.3	61.3	117.7	179.7	249.0	282.8	285.0	225.3
2003.....	7.5	17.0	31.5	64.8	121.6	182.6	244.0	286.0	287.4	224.4
2002.....	8.7	17.1	32.0	67.7	120.6	184.0	247.5	287.3	283.8	220.2
2001.....	8.4	16.4	33.1	71.0	121.3	182.5	242.1	276.4	270.7	218.1
2000.....	8.0	15.6	35.0	72.5	125.0	186.2	240.7	271.6	274.1	215.9
1999.....	7.9	15.0	35.0	70.8	121.6	180.5	236.9	260.4	257.3	205.3

Black, both sexes

2004.....	12.7	31.0	62.7	114.0	179.5	256.4	309.0	382.2	387.7	277.5
2003.....	12.4	35.4	62.0	114.8	182.9	256.6	328.9	387.1	377.2	280.9
2002.....	14.1	32.4	64.4	120.6	193.7	267.2	330.7	391.2	366.3	272.3
2001.....	14.5	32.6	63.5	119.3	182.2	273.6	347.6	395.1	370.3	280.7
2000.....	15.7	35.5	69.3	123.1	188.7	279.9	359.1	394.9	343.3	288.1
1999.....	14.5	35.1	70.8	125.2	198.4	282.1	351.1	402.9	363.5	290.5

Black, male

2004.....	14.5	40.8	89.1	161.1	271.7	367.8	467.4	600.5	627.7	481.8
2003.....	15.6	45.1	86.6	167.9	267.8	376.2	495.0	618.5	628.3	502.5
2002.....	17.0	42.6	91.6	172.1	292.0	398.4	506.4	630.6	619.0	484.6
2001.....	18.3	45.0	92.4	174.1	268.7	418.8	552.7	659.8	659.7	549.2
2000.....	21.4	47.4	100.4	185.2	271.7	432.2	561.6	656.8	617.1	562.8
1999.....	18.6	48.8	103.6	186.3	301.7	429.2	556.6	658.7	665.3	574.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of trachea, bronchus and lung (C33-C34)

Black, female

2004.....	11.1	22.5	40.4	75.2	107.7	173.8	201.4	252.8	261.2	196.5
2003.....	9.6	26.9	41.2	71.2	116.7	167.8	216.9	249.6	245.6	194.8
2002.....	11.5	23.6	41.5	78.3	116.8	169.8	212.8	248.1	234.7	191.0
2001.....	11.1	21.9	38.8	74.3	114.1	166.3	211.0	237.1	222.8	181.0
2000.....	10.7	25.1	42.7	72.2	123.2	167.6	224.7	237.0	205.5	185.9
1999.....	10.8	23.3	42.9	75.3	117.3	173.2	214.7	248.9	213.9	184.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of thymus, heart, mediastinum and pleura (C37-C38)

All races 3/, both sexes

2004.....	*	0.1	0.2	0.2	0.3	0.5	0.6	0.7	0.9	0.9
2003.....	0.1	0.1	*	0.2	0.3	0.4	0.6	0.8	0.9	1.3
2002.....	*	0.2	0.1	0.2	0.3	0.5	0.6	0.8	0.8	1.1
2001.....	0.1	0.2	0.2	0.3	0.3	0.6	0.6	0.8	0.9	1.0
2000.....	0.1	0.1	0.1	0.2	0.3	0.5	0.5	0.8	1.3	0.9
1999.....	*	0.1	0.2	0.3	0.3	0.4	0.7	0.8	1.0	1.0

All races 3/, male

2004.....	*	0.2	*	*	0.4	0.6	0.7	0.9	1.3	1.5
2003.....	*	*	*	0.3	0.4	0.5	0.7	1.0	1.2	1.7
2002.....	*	0.2	*	*	0.5	0.6	0.6	1.0	*	1.7
2001.....	*	0.2	0.2	0.5	*	0.7	0.8	0.8	1.2	1.5
2000.....	*	*	*	*	0.4	0.6	0.6	1.1	1.8	*
1999.....	*	*	0.3	0.3	0.4	*	0.7	1.0	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	0.5	0.5	0.6	0.7
2003.....	*	*	*	*	*	*	0.5	0.7	0.7	1.1
2002.....	*	*	*	*	*	0.4	0.6	0.6	0.7	0.9
2001.....	*	*	*	*	*	0.4	0.5	0.8	0.7	0.8
2000.....	*	*	*	*	*	0.4	*	0.6	0.9	0.7
1999.....	*	*	*	*	*	0.4	0.6	0.7	1.0	0.8

White, both sexes

2004.....	*	0.2	0.1	0.2	0.3	0.4	0.5	0.6	0.9	1.0
2003.....	*	0.1	*	0.2	0.3	0.4	0.6	0.7	1.0	1.4
2002.....	*	0.2	0.1	*	0.3	0.5	0.7	0.8	0.8	1.3
2001.....	0.1	0.2	0.2	0.3	0.3	0.5	0.6	0.7	0.9	1.0
2000.....	*	*	*	0.2	0.3	0.5	0.5	0.8	1.3	0.9
1999.....	*	*	0.2	0.3	0.3	0.4	0.6	0.8	1.1	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of bone and articular cartilage (C40-C41)										
All races 3/, both sexes										
2004.....	0.2	0.2	0.3	0.5	0.7	0.9	1.1	1.5	2.2	2.7
2003.....	0.3	0.3	0.4	0.4	0.5	0.6	1.1	1.5	2.5	2.8
2002.....	0.2	0.3	0.3	0.5	0.7	0.6	1.0	1.3	2.0	2.9
2001.....	0.2	0.3	0.3	0.5	0.6	0.9	1.3	1.7	2.2	3.5
2000.....	0.2	0.3	0.3	0.5	0.6	0.9	1.0	1.8	2.1	3.4
1999.....	0.2	0.3	0.4	0.3	0.6	0.8	1.1	1.6	2.1	3.0
All races 3/, male										
2004.....	0.2	0.4	0.5	0.6	0.8	1.1	1.1	1.6	2.7	3.2
2003.....	0.3	0.3	0.4	0.5	0.8	0.8	1.4	1.7	3.6	3.5
2002.....	0.3	0.4	0.4	0.6	0.7	0.7	1.1	1.8	2.4	3.7
2001.....	0.3	0.4	0.4	0.5	0.7	1.1	1.7	2.3	2.9	3.6
2000.....	0.2	0.4	0.5	0.7	0.7	1.2	1.2	2.2	2.7	4.2
1999.....	0.3	0.4	0.5	0.4	0.8	0.9	1.6	1.8	2.2	4.5
All races 3/, female										
2004.....	*	*	0.2	0.4	0.7	0.8	1.1	1.5	1.8	2.5
2003.....	0.2	*	0.3	0.4	*	0.5	1.0	1.3	1.8	2.5
2002.....	*	*	*	0.4	0.6	0.6	1.0	1.0	1.8	2.5
2001.....	*	*	0.2	0.5	0.5	0.7	1.0	1.3	1.8	3.4
2000.....	*	*	*	0.4	0.5	0.7	0.8	1.5	1.8	3.1
1999.....	*	0.3	0.3	*	0.4	0.8	0.8	1.5	2.0	2.4
White, both sexes										
2004.....	0.2	0.2	0.3	0.5	0.7	1.0	1.2	1.5	2.2	2.8
2003.....	0.3	0.3	0.4	0.4	0.5	0.6	1.2	1.4	2.5	2.9
2002.....	0.2	0.3	0.3	0.5	0.6	0.6	1.0	1.3	2.1	3.0
2001.....	0.2	0.2	0.3	0.5	0.6	0.8	1.3	1.7	2.3	3.6
2000.....	0.2	0.3	0.3	0.5	0.6	1.0	1.0	1.8	2.2	3.5
1999.....	0.2	0.3	0.4	0.3	0.5	0.9	1.2	1.7	2.2	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Melanoma and other malignant neoplasms of skin (C43-C44)

All races 3/, both sexes

2004.....	1.6	2.6	4.0	5.8	7.8	10.1	13.7	17.9	22.3	30.2
2003.....	1.8	2.8	4.0	6.0	7.9	9.5	13.5	18.3	21.6	30.8
2002.....	1.8	2.6	4.1	5.4	7.7	10.0	13.2	17.8	21.8	29.8
2001.....	1.9	2.9	4.4	5.9	8.0	10.2	14.2	17.1	21.0	31.2
2000.....	1.9	3.0	4.5	5.8	7.8	10.5	13.3	16.6	20.3	31.8
1999.....	1.9	3.1	4.2	5.7	8.0	10.6	13.1	17.2	21.3	29.2

All races 3/, male

2004.....	2.0	3.4	5.5	8.3	11.9	15.2	21.2	28.9	37.5	53.9
2003.....	2.2	3.6	5.6	8.6	11.6	14.0	20.4	29.8	35.4	53.5
2002.....	2.4	3.8	5.6	7.7	11.5	14.9	20.2	28.3	35.3	50.2
2001.....	2.4	3.8	5.9	8.3	11.9	15.6	22.2	27.1	34.5	54.0
2000.....	2.4	3.7	6.3	8.3	11.0	15.5	20.4	26.5	34.7	55.4
1999.....	2.5	4.2	5.7	8.4	11.6	15.5	20.5	26.9	35.2	50.6

All races 3/, female

2004.....	1.2	1.8	2.5	3.3	4.2	5.6	7.7	9.9	12.9	19.5
2003.....	1.4	2.1	2.5	3.5	4.7	5.5	7.9	10.2	13.2	20.7
2002.....	1.3	1.5	2.7	3.2	4.3	5.7	7.6	10.5	13.6	21.0
2001.....	1.5	2.1	2.9	3.6	4.5	5.5	7.8	10.1	12.9	21.7
2000.....	1.4	2.2	2.7	3.4	4.9	6.2	7.8	9.7	11.8	22.2
1999.....	1.3	2.0	2.8	3.2	4.7	6.3	7.2	10.4	13.1	20.6

White, both sexes

2004.....	1.8	3.1	4.6	6.5	8.8	11.3	15.4	19.7	24.1	32.7
2003.....	2.1	3.3	4.6	6.8	8.9	10.6	15.0	19.9	23.6	33.3
2002.....	2.2	3.1	4.7	6.0	8.6	11.1	14.5	19.6	23.8	32.2
2001.....	2.3	3.4	5.0	6.6	8.9	11.3	15.5	18.7	22.6	33.7
2000.....	2.2	3.4	5.1	6.4	8.7	11.6	14.5	17.9	21.8	34.3
1999.....	2.3	3.5	4.8	6.4	8.9	11.8	14.3	18.5	22.9	31.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Melanoma and other malignant neoplasms of skin (C43-C44)

White, male

2004.....	2.3	3.9	6.3	9.3	13.1	17.1	23.7	31.7	40.5	58.9
2003.....	2.6	4.2	6.4	9.7	12.9	15.6	22.7	32.4	38.6	58.2
2002.....	2.8	4.4	6.2	8.4	12.8	16.4	21.9	30.9	38.5	54.5
2001.....	2.8	4.3	6.7	9.3	13.1	17.1	24.1	29.6	37.4	58.7
2000.....	2.7	4.3	7.0	9.1	12.2	17.0	22.1	28.6	37.2	60.6
1999.....	3.0	4.7	6.3	9.3	12.8	17.2	22.3	28.7	37.9	55.0

White, female

2004.....	1.4	2.2	3.0	3.9	4.8	6.2	8.6	11.0	13.8	20.9
2003.....	1.7	2.4	2.8	3.9	5.2	6.3	8.8	10.9	14.3	22.2
2002.....	1.6	1.8	3.2	3.7	4.8	6.4	8.4	11.5	14.7	22.5
2001.....	1.8	2.5	3.4	4.1	5.0	6.2	8.5	11.0	13.7	23.2
2000.....	1.7	2.6	3.2	3.7	5.4	6.9	8.4	10.4	12.6	23.6
1999.....	1.5	2.3	3.3	3.7	5.3	7.0	7.9	11.3	14.1	22.1

Black, both sexes

2004.....	*	*	1.3	1.7	2.7	3.0	3.0	3.9	7.2	7.0
2003.....	*	0.9	1.1	1.5	2.5	3.1	2.9	5.6	*	6.7
2002.....	*	*	1.7	2.5	2.4	3.6	3.9	3.8	*	7.7
2001.....	*	*	1.2	1.6	3.0	4.0	5.1	3.8	6.0	8.0
2000.....	*	*	1.3	2.8	2.9	3.6	4.9	4.8	6.8	8.1
1999.....	*	1.0	1.5	1.7	2.9	2.8	3.7	5.4	*	6.7

Black, male

2004.....	*	*	2.0	2.8	4.9	*	*	*	*	*
2003.....	*	*	*	*	3.9	5.4	*	*	*	*
2002.....	*	*	2.8	4.5	*	6.2	*	*	*	*
2001.....	*	*	*	*	4.1	6.9	7.7	*	*	*
2000.....	*	*	*	4.1	4.3	5.2	*	*	*	*
1999.....	*	*	2.6	*	5.1	*	*	10.4	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant melanoma of skin (C43)

All races 3/, both sexes

2004.....	1.4	2.3	3.3	4.7	6.1	8.1	10.6	13.6	16.3	17.7
2003.....	1.6	2.5	3.4	4.9	6.4	7.3	10.3	13.9	15.5	17.6
2002.....	1.7	2.3	3.4	4.4	6.0	7.8	10.0	12.9	14.9	17.9
2001.....	1.8	2.6	3.6	4.6	6.1	7.8	10.7	12.4	14.7	17.8
2000.....	1.7	2.7	3.8	4.7	6.3	7.9	10.0	12.0	14.9	18.9
1999.....	1.7	2.6	3.6	4.7	6.1	8.3	9.7	12.9	14.2	16.4

All races 3/, male

2004.....	1.8	2.9	4.5	6.6	8.9	11.8	16.1	21.8	26.7	31.4
2003.....	1.9	3.1	4.6	6.9	8.9	10.4	15.1	22.3	24.9	30.2
2002.....	2.1	3.3	4.4	6.1	8.5	11.3	14.9	19.8	23.3	29.2
2001.....	2.2	3.2	4.7	6.3	8.8	11.9	16.3	19.6	24.0	29.9
2000.....	2.1	3.2	5.2	6.5	8.7	11.2	14.7	18.4	24.7	31.8
1999.....	2.2	3.5	4.6	6.7	8.4	12.0	14.6	19.8	22.7	27.5

All races 3/, female

2004.....	1.1	1.7	2.2	3.0	3.6	4.9	6.2	7.7	9.9	11.5
2003.....	1.3	1.9	2.2	3.1	4.0	4.6	6.4	7.8	9.8	12.1
2002.....	1.2	1.4	2.5	2.8	3.6	4.8	6.1	7.9	9.8	12.9
2001.....	1.4	1.9	2.6	3.0	3.7	4.3	6.4	7.3	9.2	12.8
2000.....	1.4	2.1	2.4	3.0	4.1	5.0	6.3	7.5	9.1	13.7
1999.....	1.3	1.8	2.6	2.8	4.0	5.2	5.8	8.1	9.2	11.9

White, both sexes

2004.....	1.7	2.8	3.9	5.5	7.0	9.2	12.0	15.1	17.6	19.2
2003.....	1.9	2.9	3.9	5.6	7.3	8.3	11.6	15.2	17.0	19.1
2002.....	2.0	2.8	4.0	5.0	6.8	8.9	11.0	14.2	16.3	19.2
2001.....	2.2	3.1	4.3	5.3	7.0	8.8	11.9	13.6	15.9	19.3
2000.....	2.0	3.1	4.4	5.3	7.2	8.9	11.0	13.1	16.0	20.3
1999.....	2.1	3.1	4.2	5.4	6.9	9.4	10.7	14.0	15.3	17.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other malignant neoplasms of skin (C44)										
All races 3/, both sexes										
2004.....	0.1	0.3	0.6	1.0	1.7	2.0	3.1	4.3	6.0	12.5
2003.....	0.2	0.3	0.7	1.0	1.6	2.2	3.2	4.4	6.1	13.1
2002.....	0.2	0.3	0.7	1.0	1.7	2.2	3.2	5.0	6.9	12.0
2001.....	0.1	0.3	0.7	1.3	1.9	2.4	3.5	4.8	6.3	13.4
2000.....	0.1	0.3	0.7	1.1	1.5	2.6	3.3	4.6	5.4	12.9
1999.....	0.2	0.4	0.7	1.0	1.9	2.3	3.4	4.3	7.0	12.8
All races 3/, male										
2004.....	0.2	0.4	1.0	1.7	3.0	3.5	5.1	7.2	10.9	22.5
2003.....	0.3	0.6	1.1	1.7	2.6	3.6	5.3	7.4	10.6	23.3
2002.....	0.2	0.5	1.2	1.6	3.0	3.7	5.3	8.5	12.0	21.0
2001.....	0.2	0.5	1.2	2.1	3.1	3.7	5.9	7.5	10.5	24.1
2000.....	0.2	0.5	1.1	1.9	2.4	4.2	5.7	8.1	9.9	23.6
1999.....	0.3	0.7	1.1	1.7	3.2	3.6	5.9	7.1	12.4	23.1
All races 3/, female										
2004.....	*	*	0.3	0.4	0.6	0.7	1.5	2.1	3.1	8.0
2003.....	*	*	0.3	0.4	0.6	0.9	1.5	2.3	3.3	8.6
2002.....	*	*	0.3	0.5	0.7	0.9	1.6	2.5	3.8	8.1
2001.....	*	*	0.2	0.5	0.8	1.2	1.5	2.8	3.7	8.9
2000.....	*	*	0.3	0.4	0.8	1.2	1.5	2.2	2.7	8.5
1999.....	*	*	0.3	0.4	0.7	1.1	1.4	2.4	3.9	8.6
White, both sexes										
2004.....	0.1	0.3	0.7	1.1	1.7	2.2	3.3	4.6	6.4	13.6
2003.....	0.2	0.3	0.7	1.1	1.7	2.3	3.4	4.8	6.6	14.2
2002.....	0.2	0.3	0.7	1.0	1.8	2.2	3.4	5.4	7.4	13.0
2001.....	0.1	0.3	0.7	1.3	1.9	2.5	3.6	5.1	6.7	14.3
2000.....	0.1	0.3	0.7	1.1	1.5	2.7	3.5	4.9	5.8	14.0
1999.....	0.2	0.4	0.6	1.0	2.0	2.4	3.6	4.5	7.6	13.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of mesothelial and soft tissue (C45-C49)

All races 3/, both sexes

2004.....	0.9	1.3	1.9	3.3	5.2	7.6	10.0	15.0	17.7	15.7
2003.....	0.8	1.3	2.1	3.1	5.5	7.6	10.1	14.8	16.9	14.9
2002.....	0.9	1.2	2.0	3.4	5.5	7.4	10.9	13.8	16.3	15.7
2001.....	0.9	1.4	2.3	3.4	5.3	7.7	10.6	14.7	15.7	14.9
2000.....	0.9	1.4	2.2	3.7	5.3	8.1	11.2	14.4	16.2	15.7
1999.....	1.0	1.4	2.3	3.5	5.6	7.7	11.0	14.0	15.9	14.8

All races 3/, male

2004.....	0.9	1.3	1.9	3.4	6.7	9.6	13.3	22.6	27.8	27.9
2003.....	0.9	1.5	2.2	3.3	6.3	9.7	13.9	22.5	27.1	25.1
2002.....	1.1	1.3	2.0	3.7	6.7	9.4	15.6	20.8	25.1	28.1
2001.....	1.0	1.5	2.6	3.6	6.3	10.0	15.3	21.3	24.9	25.9
2000.....	0.9	1.5	2.2	4.2	6.6	10.8	15.2	21.9	26.2	28.4
1999.....	1.3	1.4	2.5	4.3	6.9	10.4	15.6	20.8	25.7	24.6

All races 3/, female

2004.....	1.0	1.3	1.9	3.3	3.9	5.9	7.3	9.5	11.5	10.2
2003.....	0.8	1.2	2.0	3.0	4.7	5.8	7.1	9.3	10.6	10.3
2002.....	0.8	1.1	1.9	3.1	4.4	5.7	7.1	8.9	11.0	10.4
2001.....	0.9	1.2	2.0	3.2	4.3	5.7	6.8	10.1	10.2	10.3
2000.....	0.9	1.3	2.2	3.2	4.1	5.8	8.0	9.1	10.3	10.5
1999.....	0.7	1.5	2.0	2.7	4.4	5.3	7.4	9.3	10.1	10.8

White, both sexes

2004.....	0.9	1.3	1.9	3.4	5.6	8.1	10.6	15.8	18.5	16.5
2003.....	0.8	1.3	2.1	3.3	5.5	8.0	10.7	15.4	17.9	15.6
2002.....	0.9	1.2	2.0	3.4	5.6	7.7	11.4	14.6	17.0	16.5
2001.....	0.9	1.3	2.3	3.4	5.6	8.3	11.1	15.4	16.5	15.4
2000.....	0.9	1.4	2.2	3.8	5.5	8.3	11.7	15.1	17.1	16.2
1999.....	0.9	1.5	2.3	3.6	5.7	8.1	11.4	14.4	16.3	15.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of mesothelial and soft tissue (C45-C49)

White, male

2004.....	0.9	1.3	1.9	3.5	7.3	10.3	14.2	23.9	29.2	29.5
2003.....	0.8	1.4	2.2	3.5	6.4	10.2	14.8	23.6	29.2	26.1
2002.....	1.0	1.3	2.1	3.8	7.0	10.0	16.4	22.0	26.5	29.8
2001.....	0.9	1.4	2.7	3.6	6.8	10.8	16.3	22.4	26.2	27.0
2000.....	0.9	1.6	2.3	4.5	6.9	11.1	16.0	23.1	27.6	29.5
1999.....	1.2	1.4	2.6	4.6	7.1	11.0	16.2	21.6	26.9	26.2

White, female

2004.....	0.9	1.3	1.9	3.3	4.1	6.1	7.7	9.9	11.9	10.6
2003.....	0.8	1.2	2.0	3.1	4.7	6.1	7.4	9.5	10.9	10.9
2002.....	0.8	1.1	1.9	3.1	4.4	5.8	7.3	9.3	11.2	10.7
2001.....	0.8	1.2	1.9	3.2	4.5	6.0	6.9	10.4	10.7	10.5
2000.....	0.9	1.3	2.0	3.2	4.2	5.9	8.3	9.5	10.9	10.7
1999.....	0.6	1.5	2.0	2.7	4.4	5.5	7.7	9.4	10.1	10.9

Black, both sexes

2004.....	1.2	1.5	2.8	3.9	2.9	6.0	6.0	9.0	12.6	9.3
2003.....	1.2	2.0	2.5	2.5	5.8	5.9	6.9	11.2	8.4	9.9
2002.....	1.3	1.7	2.0	3.8	4.7	6.6	7.5	9.1	11.8	7.7
2001.....	1.3	1.7	2.9	3.9	3.4	4.5	7.5	8.7	8.1	11.1
2000.....	1.4	1.9	2.6	3.8	4.8	7.7	8.8	8.5	7.9	10.9
1999.....	1.5	1.5	2.4	2.9	6.1	5.1	8.4	11.4	11.6	10.5

Black, male

2004.....	*	*	2.7	3.8	*	6.5	7.8	12.7	18.7	*
2003.....	1.5	2.9	2.8	2.9	6.9	6.4	7.6	15.5	*	*
2002.....	1.7	2.3	*	4.0	4.8	6.7	9.7	11.9	*	*
2001.....	1.5	*	2.9	4.4	*	5.1	8.1	10.7	*	*
2000.....	*	1.8	*	4.0	6.2	9.4	11.0	12.3	*	*
1999.....	1.9	*	2.5	3.6	7.2	6.0	10.5	14.7	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of mesothelial and soft tissue (C45-C49)

Black, female

2004.....	1.4	1.5	2.8	3.9	*	5.6	4.8	6.7	9.4	*
2003.....	*	*	2.3	*	5.0	5.5	6.4	8.7	8.5	*
2002.....	*	*	2.1	3.5	4.6	6.5	6.0	7.4	10.3	*
2001.....	*	1.8	2.9	3.6	3.5	4.1	7.2	7.5	*	8.8
2000.....	1.3	1.9	3.7	3.6	3.8	6.5	7.3	6.3	*	8.6
1999.....	*	1.6	2.3	*	5.2	4.4	7.0	9.4	10.4	10.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Mesothelioma (C45)

All races 3/, both sexes

2004.....	0.1	0.2	0.4	0.8	2.0	2.9	4.2	7.6	8.4	5.7
2003.....	0.1	0.2	0.5	0.8	1.9	3.2	4.4	7.1	8.1	5.7
2002.....	0.1	0.1	0.4	1.0	1.9	3.0	5.1	7.0	7.4	5.9
2001.....	0.1	0.2	0.5	0.9	1.9	3.1	4.8	7.2	6.9	5.5
2000.....	0.1	0.2	0.5	1.1	1.9	3.3	5.2	6.6	7.2	5.7
1999.....	*	0.2	0.5	1.2	2.0	3.4	5.1	6.6	7.1	4.8

All races 3/, male

2004.....	*	0.2	0.6	1.2	3.5	5.1	7.6	15.5	17.4	14.4
2003.....	*	0.4	0.7	1.3	3.4	5.6	8.3	14.4	17.1	14.2
2002.....	*	*	0.6	1.6	3.4	5.6	9.8	14.0	15.6	15.9
2001.....	*	0.3	0.8	1.5	3.2	5.6	9.1	14.1	14.7	14.5
2000.....	*	0.2	0.7	1.9	3.4	5.9	9.8	13.4	15.5	15.5
1999.....	*	0.4	0.8	1.9	3.3	6.1	9.6	12.9	15.8	12.7

All races 3/, female

2004.....	*	*	0.2	0.4	0.6	1.0	1.5	1.9	2.9	1.8
2003.....	*	*	0.2	0.4	0.6	1.1	1.2	1.9	2.6	1.9
2002.....	*	*	*	0.4	0.5	0.7	1.3	2.0	2.3	1.6
2001.....	*	*	0.3	0.4	0.7	0.9	1.4	2.4	2.1	1.7
2000.....	*	*	0.2	0.4	0.6	1.1	1.5	1.9	2.3	1.8
1999.....	*	*	0.3	0.5	0.7	1.0	1.6	2.2	2.1	1.6

White, both sexes

2004.....	0.1	0.2	0.4	0.9	2.3	3.2	4.6	8.3	9.0	6.1
2003.....	0.1	0.2	0.5	1.0	2.1	3.5	4.8	7.6	8.7	6.0
2002.....	0.1	0.2	0.4	1.1	2.1	3.3	5.5	7.5	7.9	6.4
2001.....	*	0.2	0.5	1.0	2.2	3.5	5.3	7.9	7.3	5.9
2000.....	*	0.2	0.5	1.2	2.1	3.5	5.6	7.1	7.8	6.1
1999.....	*	0.3	0.5	1.3	2.1	3.7	5.5	7.0	7.5	5.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other mesothelial and soft tissue (C47-C49)

All races 3/, both sexes

2004.....	0.8	1.1	1.5	2.5	3.2	4.7	5.8	7.3	9.2	9.6
2003.....	0.7	1.1	1.7	2.3	3.5	4.4	5.7	7.6	8.7	8.8
2002.....	0.8	1.1	1.6	2.4	3.6	4.4	5.8	6.8	8.8	9.3
2001.....	0.8	1.1	1.8	2.4	3.4	4.6	5.8	7.4	8.7	9.1
2000.....	0.8	1.2	1.7	2.6	3.4	4.8	5.9	7.7	8.8	9.6
1999.....	0.9	1.2	1.8	2.3	3.6	4.3	5.9	7.4	8.6	9.6

All races 3/, male

2004.....	0.7	1.0	1.3	2.1	3.2	4.5	5.7	7.0	10.2	12.7
2003.....	0.7	1.1	1.5	2.0	2.9	4.0	5.5	7.9	9.9	10.7
2002.....	0.9	1.1	1.4	2.1	3.3	3.8	5.7	6.7	9.3	11.7
2001.....	0.9	1.1	1.8	2.1	3.1	4.4	6.1	7.1	10.0	10.7
2000.....	0.8	1.3	1.5	2.3	3.3	4.9	5.2	8.4	10.4	12.5
1999.....	1.1	1.0	1.7	2.4	3.6	4.4	5.9	7.8	9.6	11.3

All races 3/, female

2004.....	0.9	1.1	1.7	2.9	3.3	4.9	5.8	7.6	8.5	8.2
2003.....	0.7	1.0	1.8	2.6	4.1	4.7	5.9	7.3	8.0	8.0
2002.....	0.8	1.0	1.7	2.7	3.8	4.9	5.8	6.8	8.6	8.3
2001.....	0.8	1.1	1.8	2.8	3.6	4.8	5.4	7.6	8.0	8.4
2000.....	0.8	1.1	1.9	2.8	3.5	4.7	6.5	7.1	7.8	8.4
1999.....	0.7	1.3	1.8	2.2	3.7	4.3	5.8	7.1	8.0	8.9

White, both sexes

2004.....	0.8	1.1	1.5	2.5	3.4	4.9	6.0	7.6	9.5	9.9
2003.....	0.7	1.0	1.6	2.3	3.4	4.5	5.9	7.6	9.1	9.1
2002.....	0.8	1.0	1.6	2.4	3.6	4.4	5.9	7.1	9.0	9.6
2001.....	0.8	1.1	1.7	2.4	3.4	4.8	5.8	7.4	9.1	9.1
2000.....	0.8	1.2	1.7	2.6	3.4	4.8	6.1	7.9	9.1	9.7
1999.....	0.8	1.2	1.8	2.3	3.6	4.4	5.9	7.4	8.7	9.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other mesothelial and soft tissue (C47-C49)

White, male

2004.....	0.7	1.0	1.3	2.1	3.4	4.6	5.9	7.2	10.8	13.0
2003.....	0.6	1.0	1.5	2.0	2.8	4.0	5.7	8.1	10.5	10.9
2002.....	0.8	1.1	1.5	2.1	3.3	3.8	5.9	7.1	9.6	12.0
2001.....	0.8	1.0	1.8	2.0	3.2	4.5	6.4	7.1	10.4	10.6
2000.....	0.8	1.3	1.6	2.4	3.2	4.9	5.4	8.6	10.8	12.4
1999.....	1.1	1.0	1.8	2.4	3.6	4.3	5.9	7.9	9.8	12.0

White, female

2004.....	0.8	1.1	1.6	2.9	3.4	5.1	6.1	7.8	8.7	8.5
2003.....	0.7	1.0	1.8	2.6	4.1	4.9	6.0	7.3	8.2	8.3
2002.....	0.7	1.0	1.8	2.6	3.8	4.9	5.9	7.1	8.6	8.5
2001.....	0.7	1.1	1.7	2.8	3.7	5.1	5.4	7.7	8.3	8.4
2000.....	0.8	1.1	1.7	2.8	3.6	4.7	6.6	7.4	8.2	8.5
1999.....	0.5	1.3	1.7	2.1	3.6	4.4	6.0	7.1	8.0	8.9

Black, both sexes

2004.....	1.1	1.3	2.4	3.5	2.1	5.0	4.4	6.8	8.4	7.0
2003.....	1.2	1.7	2.1	2.3	4.4	4.6	5.6	8.2	5.6	7.0
2002.....	1.2	1.6	1.7	3.2	4.1	5.3	5.5	5.7	8.9	5.9
2001.....	1.2	1.4	2.5	3.5	3.0	3.8	6.1	7.3	6.0	9.8
2000.....	1.2	1.6	2.4	2.9	3.7	5.2	5.9	6.0	*	9.1
1999.....	1.3	1.3	1.9	2.4	4.8	4.0	6.6	8.0	7.5	8.3

Black, male

2004.....	*	*	2.1	3.2	*	5.1	*	*	*	*
2003.....	*	2.2	*	*	4.2	*	*	*	*	*
2002.....	1.5	2.1	*	*	*	*	*	*	*	*
2001.....	*	*	2.3	3.4	*	*	*	*	*	*
2000.....	*	*	*	*	4.1	*	*	*	*	*
1999.....	1.5	*	*	*	4.5	*	7.1	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other mesothelial and soft tissue (C47-C49)

Black, female

2004.....	1.3	1.4	2.7	3.7	*	4.9	4.4	6.7	9.4	*
2003.....	*	*	2.2	*	4.5	5.1	6.4	8.4	*	*
2002.....	*	*	2.0	3.4	4.3	6.3	5.4	6.3	9.5	*
2001.....	*	1.7	2.7	3.5	3.5	3.8	6.7	7.2	*	8.8
2000.....	*	1.7	3.7	3.4	3.4	5.6	6.7	*	*	*
1999.....	*	*	2.2	*	5.0	*	6.3	8.0	*	9.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of breast (C50)

All races 3/, both sexes

2004.....	7.3	11.7	18.7	26.1	33.1	40.8	49.6	64.7	82.4	124.2
2003.....	7.9	12.6	19.1	26.6	33.5	39.9	51.8	66.0	87.3	132.5
2002.....	7.9	12.9	19.7	26.2	33.7	41.3	52.4	67.1	89.8	134.6
2001.....	8.2	13.4	20.6	27.1	34.1	41.7	54.0	69.1	88.0	134.2
2000.....	8.3	13.6	20.8	28.1	35.0	43.4	55.2	70.9	90.9	147.4
1999.....	7.9	13.3	21.0	28.1	34.8	43.5	55.9	70.0	92.4	144.4

All races 3/, male

2004.....	*	*	0.2	0.4	0.6	0.9	1.3	2.1	2.1	3.3
2003.....	*	*	0.4	0.4	0.7	0.9	1.2	1.5	2.8	3.9
2002.....	*	*	0.3	0.4	0.7	0.8	1.3	1.9	2.7	3.4
2001.....	*	*	0.4	0.5	0.8	1.2	1.7	2.3	2.4	3.5
2000.....	*	*	0.3	0.7	0.8	1.1	1.7	2.1	2.4	4.2
1999.....	*	*	0.4	0.6	0.7	1.1	1.4	1.9	2.3	4.5

All races 3/, female

2004.....	14.5	23.0	36.3	50.4	62.6	75.6	88.5	109.6	131.8	178.6
2003.....	15.6	24.7	37.0	51.3	63.4	73.7	92.4	111.9	139.0	189.4
2002.....	15.6	25.3	38.3	50.4	63.7	76.2	93.2	113.2	142.7	191.5
2001.....	16.2	26.3	40.1	52.0	64.3	76.5	95.6	115.9	139.4	188.9
2000.....	16.4	26.6	40.4	53.8	66.0	79.6	97.3	118.8	143.1	205.7
1999.....	15.6	26.1	40.8	53.9	65.6	79.7	98.4	117.4	145.1	200.8

White, both sexes

2004.....	6.5	10.5	17.1	25.1	32.0	40.6	49.5	65.0	83.4	124.8
2003.....	7.1	11.4	17.8	25.3	32.6	39.2	51.7	66.0	87.5	132.5
2002.....	7.0	11.7	18.5	25.0	33.1	40.8	52.0	67.5	89.0	135.3
2001.....	7.3	12.4	19.4	25.8	33.3	40.9	53.6	69.4	88.2	134.8
2000.....	7.5	12.4	19.7	26.9	34.4	43.3	55.5	71.1	91.7	147.3
1999.....	7.1	12.1	19.8	26.8	33.8	42.8	55.5	69.7	92.8	146.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of breast (C50)

Black, female

2004.....	25.6	39.5	58.1	72.4	85.8	88.1	103.7	121.6	135.7	196.9
2003.....	26.3	42.0	57.3	78.3	84.7	94.1	107.4	127.2	160.5	222.4
2002.....	28.7	42.4	58.7	73.1	81.2	92.9	111.1	123.2	176.7	209.1
2001.....	27.5	41.9	60.5	76.7	83.7	99.5	115.0	130.9	157.8	205.9
2000.....	27.7	44.3	60.4	78.8	83.3	91.0	107.4	133.6	149.0	238.7
1999.....	27.5	44.9	63.3	79.6	85.2	99.5	114.1	139.3	154.7	199.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of female genital organs (C51-C58)

All races 3/, both sexes

2004.....	3.8	6.2	9.4	14.5	21.5	30.9	39.4	48.8	62.9	72.7
2003.....	3.7	5.9	9.9	15.2	22.9	30.9	38.2	49.9	63.8	75.2
2002.....	3.7	6.4	10.3	15.1	22.5	30.3	39.9	52.0	63.6	77.5
2001.....	3.8	6.4	10.6	14.7	22.8	30.1	39.6	52.1	66.9	79.9
2000.....	3.8	6.5	10.4	15.5	22.2	31.0	38.9	52.5	65.4	82.3
1999.....	3.9	6.6	10.3	15.1	22.7	30.8	39.4	51.7	64.6	79.1

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	7.5	12.1	18.3	28.3	41.0	57.8	71.1	83.8	101.6	105.4
2003.....	7.3	11.7	19.4	29.5	43.7	57.6	68.8	85.4	102.8	108.5
2002.....	7.3	12.6	20.2	29.2	42.9	56.4	71.8	88.8	102.3	111.2
2001.....	7.5	12.6	20.7	28.4	43.4	56.0	71.0	88.6	107.0	113.4
2000.....	7.5	12.7	20.3	30.1	42.4	57.5	69.6	89.1	103.9	115.8
1999.....	7.8	13.0	20.3	29.3	43.2	57.2	70.3	87.6	102.4	110.9

White, both sexes

2004.....	3.6	6.0	9.0	14.3	21.1	29.7	38.1	48.6	63.5	73.0
2003.....	3.6	5.8	9.7	15.2	22.2	30.0	37.4	49.2	63.6	75.2
2002.....	3.6	6.2	10.3	14.9	22.1	29.2	38.9	51.3	63.3	76.9
2001.....	3.7	6.3	10.4	14.5	21.9	29.0	38.9	51.6	66.9	79.7
2000.....	3.6	6.4	10.2	15.4	21.7	29.8	37.7	52.0	65.0	82.1
1999.....	3.8	6.7	10.3	14.9	22.2	29.7	38.3	50.7	64.5	78.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of female genital organs (C51-C58)

Black, female

2004.....	9.1	14.8	22.9	33.5	52.0	81.7	99.1	99.5	105.1	113.0
2003.....	8.5	14.5	22.6	32.7	57.2	76.2	88.6	103.9	116.4	124.0
2002.....	9.2	14.9	22.4	34.3	52.3	76.6	95.7	106.8	117.8	136.0
2001.....	9.8	13.1	23.0	32.6	56.8	77.5	91.7	107.1	117.7	130.1
2000.....	10.5	13.7	23.0	33.7	54.1	81.6	96.3	110.5	119.7	132.6
1999.....	9.6	12.0	21.9	36.4	55.9	80.0	101.3	113.3	110.6	125.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of vulva and vagina (C51-C52)

All races 3/, both sexes

2004.....	0.1	0.2	0.2	0.4	0.7	0.9	1.6	2.1	3.9	6.9
2003.....	*	0.2	0.2	0.3	0.6	1.0	1.4	2.4	3.6	7.2
2002.....	0.1	0.1	0.3	0.3	0.6	0.7	1.5	2.5	3.5	7.7
2001.....	0.1	0.2	0.3	0.3	0.5	0.9	1.4	2.5	3.9	7.3
2000.....	0.1	0.1	0.3	0.3	0.6	1.0	1.5	2.5	3.6	7.7
1999.....	0.1	0.1	0.3	0.3	0.6	0.9	1.5	2.5	3.9	8.6

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	0.3	0.4	0.5	0.8	1.3	1.7	2.9	3.7	6.3	10.1
2003.....	*	0.3	0.5	0.7	1.1	1.8	2.5	4.0	5.7	10.4
2002.....	0.2	0.3	0.5	0.7	1.2	1.3	2.7	4.3	5.7	11.0
2001.....	0.2	0.3	0.6	0.6	1.0	1.6	2.6	4.2	6.3	10.3
2000.....	0.2	0.3	0.6	0.6	1.1	1.9	2.8	4.3	5.7	10.9
1999.....	0.2	0.3	0.6	0.7	1.2	1.6	2.6	4.2	6.1	12.0

White, both sexes

2004.....	0.1	0.2	0.2	0.3	0.7	0.9	1.6	2.3	4.0	7.1
2003.....	*	0.2	0.2	0.3	0.6	1.0	1.4	2.5	3.7	7.5
2002.....	0.1	0.1	0.3	0.4	0.7	0.7	1.5	2.6	3.6	8.0
2001.....	*	0.1	0.3	0.3	0.4	0.9	1.4	2.5	4.1	7.4
2000.....	*	0.1	0.3	0.3	0.6	1.1	1.6	2.5	3.7	7.9
1999.....	*	0.1	0.3	0.3	0.6	0.9	1.5	2.5	3.9	9.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of cervix uteri (C53)

All races 3/, both sexes

2004.....	1.7	2.1	2.0	2.7	2.9	3.2	3.3	3.3	4.3	5.1
2003.....	1.7	2.0	2.4	2.8	2.9	3.2	3.0	3.7	5.1	5.7
2002.....	1.7	2.1	2.4	2.7	2.8	3.3	3.6	4.1	4.2	6.2
2001.....	1.8	2.3	2.6	2.6	3.2	2.9	3.5	4.0	5.5	7.2
2000.....	1.8	2.2	2.7	3.1	3.1	3.6	3.9	4.3	5.6	7.3
1999.....	1.8	2.4	2.7	3.0	3.5	3.7	3.9	4.7	4.6	7.9

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	3.4	4.1	4.0	5.2	5.6	6.0	6.0	5.6	7.0	7.3
2003.....	3.3	4.0	4.6	5.5	5.5	5.9	5.4	6.3	8.3	8.3
2002.....	3.4	4.1	4.7	5.2	5.3	6.1	6.5	7.0	6.7	8.9
2001.....	3.7	4.5	5.1	5.1	6.0	5.4	6.3	6.8	8.9	10.1
2000.....	3.5	4.4	5.2	6.0	5.9	6.7	6.9	7.3	9.0	10.3
1999.....	3.6	4.6	5.2	5.8	6.7	6.8	6.9	8.0	7.2	11.1

White, both sexes

2004.....	1.5	1.9	1.8	2.5	2.6	2.7	2.8	2.9	3.8	4.1
2003.....	1.6	1.9	2.1	2.6	2.6	2.8	2.4	2.8	4.6	4.9
2002.....	1.6	1.8	2.2	2.5	2.4	2.7	3.0	3.6	3.5	5.1
2001.....	1.7	2.2	2.3	2.3	2.9	2.4	3.0	3.4	4.7	6.2
2000.....	1.6	2.0	2.4	2.7	2.6	2.9	3.3	3.8	5.0	6.3
1999.....	1.7	2.3	2.5	2.7	3.0	3.1	3.1	4.0	4.1	6.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of cervix uteri (C53)

Black, female

2004.....	5.4	6.6	7.2	8.6	10.9	12.4	13.6	12.7	15.5	22.4
2003.....	4.5	6.5	7.1	9.3	10.8	12.4	12.8	17.9	17.4	20.3
2002.....	5.6	7.0	8.4	7.6	10.7	14.2	16.8	15.4	19.1	24.2
2001.....	5.6	6.1	8.2	8.3	10.6	13.7	12.8	15.5	22.9	26.1
2000.....	5.9	7.0	9.1	11.2	11.4	16.3	16.2	17.4	20.4	27.0
1999.....	5.7	5.5	7.5	11.1	14.8	15.2	21.8	19.7	14.3	29.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of corpus uteri and uterus, part unspecified (C54-C55)

All races 3/, both sexes

2004.....	0.5	0.9	1.8	3.4	5.8	8.8	11.5	13.8	17.7	22.3
2003.....	0.5	0.8	1.9	3.5	6.1	9.0	11.3	13.8	17.1	22.8
2002.....	0.4	0.9	2.0	3.5	5.8	8.4	11.8	14.2	17.0	24.6
2001.....	0.4	0.9	2.0	3.2	5.8	8.2	11.4	14.5	19.2	24.7
2000.....	0.4	0.9	1.8	3.5	6.0	8.4	11.1	14.0	18.6	24.7
1999.....	0.5	0.9	1.7	3.3	5.5	8.8	10.9	14.4	19.4	24.0

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	1.0	1.8	3.5	6.7	11.0	16.5	20.8	23.8	28.7	32.3
2003.....	1.0	1.6	3.7	6.8	11.7	16.8	20.4	23.5	27.6	33.0
2002.....	0.9	1.7	3.9	6.9	11.0	15.7	21.2	24.2	27.4	35.2
2001.....	0.8	1.8	4.0	6.2	11.1	15.3	20.4	24.6	30.7	35.0
2000.....	0.9	1.9	3.5	6.9	11.4	15.6	19.8	23.8	29.5	34.8
1999.....	0.9	1.8	3.3	6.3	10.5	16.2	19.5	24.5	30.7	33.7

White, both sexes

2004.....	0.5	0.9	1.6	3.2	5.2	7.7	10.2	13.0	17.4	22.4
2003.....	0.4	0.7	1.9	3.4	5.4	7.9	10.5	13.0	16.2	22.7
2002.....	0.4	0.8	1.9	3.2	5.2	7.2	10.8	13.1	16.3	23.8
2001.....	0.4	0.9	1.9	3.1	5.2	7.2	10.5	13.7	18.6	24.5
2000.....	0.4	0.9	1.8	3.4	5.4	7.3	9.9	13.0	17.5	24.4
1999.....	0.4	0.9	1.6	3.0	5.2	7.6	10.2	13.7	18.8	23.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of corpus uteri and uterus, part unspecified (C54-C55)

Black, female

2004.....	1.4	1.9	5.5	9.4	21.1	35.3	44.4	42.3	38.9	36.6
2003.....	1.6	2.1	4.7	9.6	24.8	34.7	37.4	40.7	47.6	41.7
2002.....	*	2.0	5.4	12.1	19.2	34.8	42.1	44.5	44.2	53.8
2001.....	*	2.6	5.4	8.3	20.8	32.7	40.4	42.4	47.0	44.2
2000.....	*	1.9	4.4	9.6	21.4	34.0	44.0	44.4	52.6	45.5
1999.....	1.6	*	4.1	10.5	18.8	36.4	36.2	41.2	44.5	48.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of ovary (C56)

All races 3/, both sexes

2004.....	1.4	2.9	5.2	7.9	11.9	17.8	22.5	28.9	36.4	37.7
2003.....	1.4	2.9	5.2	8.3	13.0	17.3	21.9	29.4	36.9	38.4
2002.....	1.4	3.2	5.5	8.3	12.9	17.3	22.3	30.3	37.8	38.1
2001.....	1.4	2.9	5.5	8.3	13.0	17.6	22.6	30.2	37.4	39.6
2000.....	1.5	3.1	5.5	8.3	12.2	17.5	21.8	30.9	36.5	40.9
1999.....	1.5	3.1	5.5	8.3	12.6	17.0	22.4	29.2	35.5	37.6

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	2.8	5.8	10.1	15.3	22.7	33.3	40.7	49.7	58.8	54.6
2003.....	2.8	5.7	10.2	16.2	24.8	32.3	39.5	50.4	59.5	55.4
2002.....	2.8	6.4	10.7	16.2	24.6	32.3	40.1	51.7	60.8	54.6
2001.....	2.8	5.8	10.8	16.0	24.7	32.8	40.5	51.4	59.8	56.1
2000.....	2.9	6.0	10.8	16.1	23.3	32.5	39.0	52.5	58.0	57.6
1999.....	3.0	6.2	10.8	16.0	24.0	31.6	39.9	49.5	56.2	52.8

White, both sexes

2004.....	1.5	3.0	5.3	8.0	12.4	18.2	23.1	29.8	37.8	38.6
2003.....	1.4	3.0	5.4	8.7	13.5	17.9	22.4	30.3	38.1	39.1
2002.....	1.4	3.3	5.8	8.6	13.5	18.0	22.9	31.1	38.8	39.0
2001.....	1.4	3.1	5.7	8.5	13.2	18.1	23.3	31.1	38.6	40.4
2000.....	1.5	3.2	5.6	8.7	12.7	18.0	22.4	32.0	37.6	42.0
1999.....	1.6	3.3	5.7	8.6	13.0	17.5	22.9	29.6	36.4	38.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of ovary (C56)

Black, female

2004.....	2.0	5.7	9.1	13.7	18.1	31.8	37.8	41.8	44.2	45.7
2003.....	2.3	5.3	9.4	12.5	19.6	26.7	34.6	41.8	44.9	51.0
2002.....	2.3	5.3	7.7	13.9	19.5	25.9	33.4	42.3	49.3	48.1
2001.....	2.7	3.6	8.1	14.4	22.6	28.3	34.5	41.9	42.1	50.1
2000.....	2.9	4.0	8.8	11.6	19.4	29.2	32.6	41.9	41.2	47.2
1999.....	2.1	4.5	9.3	13.2	20.3	26.2	37.8	46.3	42.3	39.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other and unspecified female genital organs (C57)

All races 3/, both sexes

2004.....	*	*	0.1	0.1	0.2	0.2	0.4	0.6	0.5	0.7
2003.....	*	*	0.1	0.2	0.3	0.4	0.6	0.6	1.1	1.1
2002.....	*	*	0.2	0.2	0.4	0.5	0.6	1.0	1.1	1.0
2001.....	*	*	0.1	0.3	0.3	0.4	0.7	0.9	0.9	1.2
2000.....	*	*	0.1	0.2	0.4	0.5	0.6	0.7	1.1	1.6
1999.....	*	*	0.2	0.3	0.4	0.5	0.7	0.8	1.3	0.9

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	*	*	0.2	0.3	0.4	0.4	0.7	1.0	0.8	1.1
2003.....	*	*	0.3	0.4	0.5	0.8	1.1	1.1	1.7	1.5
2002.....	*	*	0.3	0.4	0.8	1.0	1.2	1.6	1.7	1.5
2001.....	*	*	0.3	0.5	0.6	0.8	1.2	1.5	1.4	1.7
2000.....	*	*	0.2	0.5	0.8	0.8	1.1	1.3	1.7	2.2
1999.....	*	*	0.4	0.5	0.8	0.9	1.3	1.4	2.1	1.3

White, both sexes

2004.....	*	*	*	0.1	0.2	*	0.4	0.6	0.5	0.8
2003.....	*	*	0.1	0.2	0.2	0.5	0.6	0.6	1.1	1.0
2002.....	*	*	0.2	0.2	0.3	0.5	0.7	1.0	1.1	1.0
2001.....	*	*	0.1	0.3	0.3	0.4	0.7	0.9	0.9	1.3
2000.....	*	*	*	0.2	0.4	0.5	0.6	0.6	1.1	1.6
1999.....	*	*	0.2	0.3	0.4	0.5	0.7	0.8	1.2	0.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

Of placenta (C58)

All races 3/, both sexes

Table with 11 columns (years) and 6 rows (2004-1999) showing asterisks for all cells.

All races 3/, male

Table with 11 columns (years) and 6 rows (2004-1999) showing ellipses for all cells.

All races 3/, female

Table with 11 columns (years) and 6 rows (2004-1999) showing asterisks for all cells.

White, both sexes

Table with 11 columns (years) and 6 rows (2004-1999) showing asterisks for all cells.

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Of placenta (C58)

White, male

Table with 11 columns (years 1999-2004) and 11 rows (ages 40-44 to 85+). All cells contain '...'.

White, female

Table with 11 columns (years 1999-2004) and 11 rows (ages 40-44 to 85+). All cells contain '*'.

Black, both sexes

Table with 11 columns (years 1999-2004) and 11 rows (ages 40-44 to 85+). All cells contain '*'.

Black, male

Table with 11 columns (years 1999-2004) and 11 rows (ages 40-44 to 85+). All cells contain '...'.

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of male genital organs (C60-C63)

All races 3/, both sexes

2004.....	0.3	0.7	2.0	4.4	11.1	22.4	41.9	74.1	118.5	181.7
2003.....	0.3	0.7	1.9	4.4	12.0	23.9	44.0	76.8	122.9	185.9
2002.....	0.4	0.6	1.8	4.9	11.8	24.6	46.4	80.3	129.4	195.9
2001.....	0.4	0.7	1.9	4.8	11.6	25.3	47.9	83.0	133.8	201.9
2000.....	0.3	0.7	1.7	4.8	12.4	26.3	49.4	85.7	137.7	210.1
1999.....	0.4	0.8	1.8	5.1	12.6	28.9	52.3	89.5	142.3	208.7

All races 3/, male

2004.....	0.7	1.4	4.0	9.1	23.2	48.2	94.0	177.2	310.9	585.7
2003.....	0.6	1.4	3.9	9.1	25.2	51.5	98.9	184.7	323.8	606.5
2002.....	0.8	1.3	3.7	10.1	24.7	53.1	104.4	193.9	342.2	647.3
2001.....	0.7	1.5	3.9	9.9	24.4	54.8	108.2	201.5	357.2	684.2
2000.....	0.7	1.5	3.5	10.0	26.2	56.9	112.1	208.7	371.2	726.1
1999.....	0.9	1.5	3.7	10.6	26.4	62.6	119.3	218.2	385.6	726.5

All races 3/, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	0.4	0.6	1.7	3.9	9.7	19.9	37.7	67.7	111.3	173.1
2003.....	0.3	0.6	1.7	3.7	10.3	20.9	40.1	70.3	114.8	178.3
2002.....	0.4	0.5	1.5	4.2	10.0	21.6	41.3	73.2	121.0	187.6
2001.....	0.4	0.7	1.6	4.1	9.9	22.5	42.7	75.8	123.9	190.3
2000.....	0.3	0.6	1.5	4.1	10.8	22.8	43.6	78.1	127.0	199.5
1999.....	0.4	0.7	1.5	4.5	10.4	25.3	46.8	81.6	133.9	196.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of male genital organs (C60-C63)

White, male

2004.....	0.7	1.3	3.3	7.9	20.1	42.4	83.7	160.3	290.4	557.1
2003.....	0.7	1.2	3.3	7.5	21.5	44.6	89.0	167.6	301.1	581.0
2002.....	0.8	1.1	3.1	8.6	20.8	46.2	91.9	175.4	318.4	619.3
2001.....	0.8	1.4	3.2	8.5	20.7	48.1	95.5	182.8	328.9	645.0
2000.....	0.6	1.3	3.1	8.4	22.4	48.9	98.0	189.2	340.4	690.4
1999.....	0.9	1.4	3.1	9.2	21.7	54.4	105.7	197.7	360.7	685.1

White, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Black, both sexes

2004.....	*	1.5	4.8	10.6	26.6	51.5	93.3	162.2	228.5	310.8
2003.....	*	1.2	4.5	12.2	30.3	56.5	96.4	166.6	243.4	304.0
2002.....	*	1.5	4.4	11.9	29.8	58.2	111.3	177.3	262.7	328.7
2001.....	*	1.4	5.1	12.1	29.0	58.1	115.3	181.7	281.4	363.3
2000.....	*	1.7	3.7	12.9	29.7	65.1	120.7	191.2	294.2	357.9
1999.....	*	1.4	4.4	12.1	34.1	69.0	124.1	197.3	273.6	367.6

Black, male

2004.....	*	3.2	10.5	23.5	60.7	120.9	230.8	435.8	662.2	1,094.8
2003.....	*	2.6	9.9	26.9	69.2	132.7	239.5	446.9	708.0	1,086.4
2002.....	*	3.2	9.7	26.5	67.9	136.6	277.1	474.0	767.3	1,186.8
2001.....	*	3.1	11.0	26.7	65.9	136.7	288.5	486.0	833.9	1,342.3
2000.....	*	3.6	8.1	28.7	67.3	153.3	302.6	508.4	878.4	1,319.3
1999.....	*	3.0	9.6	27.0	77.6	162.0	311.0	524.8	825.7	1,351.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of prostate (C61)										
All races 3/, both sexes										
2004.....	0.1	0.5	1.7	4.2	10.8	22.1	41.3	73.5	117.8	180.9
2003.....	0.1	0.5	1.7	4.2	11.6	23.5	43.6	76.1	122.1	184.9
2002.....	0.1	0.4	1.6	4.6	11.4	24.2	45.8	79.7	128.7	195.0
2001.....	0.1	0.5	1.7	4.6	11.3	25.0	47.4	82.4	133.2	200.9
2000.....	*	0.5	1.5	4.6	12.1	25.9	49.0	85.1	136.6	209.3
1999.....	0.1	0.5	1.6	4.9	12.3	28.4	51.9	89.0	141.4	207.5
All races 3/, male										
2004.....	0.2	0.9	3.5	8.6	22.6	47.4	92.6	175.7	309.0	583.1
2003.....	0.2	0.9	3.4	8.7	24.4	50.6	97.9	183.0	321.8	603.3
2002.....	0.2	0.8	3.3	9.6	24.0	52.2	103.2	192.6	340.3	644.5
2001.....	0.2	1.0	3.5	9.4	23.8	53.9	107.1	200.1	355.4	681.2
2000.....	*	1.1	3.1	9.6	25.4	56.0	111.3	207.4	368.2	723.1
1999.....	0.3	1.0	3.3	10.1	25.8	61.6	118.3	216.9	383.0	722.4
All races 3/, female										
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
White, both sexes										
2004.....	*	0.4	1.4	3.6	9.4	19.6	37.1	67.1	110.5	172.3
2003.....	*	0.4	1.4	3.5	9.9	20.5	39.6	69.7	114.1	177.5
2002.....	*	0.3	1.3	3.9	9.6	21.2	40.7	72.6	120.3	186.7
2001.....	*	0.4	1.4	3.9	9.7	22.1	42.2	75.3	123.3	189.4
2000.....	*	0.4	1.3	3.8	10.4	22.4	43.2	77.6	125.8	198.6
1999.....	0.1	0.4	1.4	4.2	10.2	24.9	46.4	81.1	132.9	195.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of prostate (C61)

White, male

2004.....	*	0.7	2.8	7.3	19.5	41.7	82.3	158.8	288.4	554.4
2003.....	*	0.8	2.8	7.2	20.7	43.7	88.0	166.0	299.2	578.2
2002.....	*	0.5	2.7	8.1	20.1	45.2	90.7	174.1	316.4	616.3
2001.....	*	0.8	2.8	8.0	20.2	47.3	94.3	181.5	327.2	641.8
2000.....	*	0.8	2.6	7.9	21.6	48.0	97.0	188.0	337.4	687.5
1999.....	0.2	0.8	2.8	8.6	21.2	53.4	104.7	196.5	358.0	681.0

White, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Black, both sexes

2004.....	*	1.3	4.7	10.5	26.4	50.6	92.8	161.3	228.0	310.2
2003.....	*	1.1	4.5	11.8	29.9	56.1	96.2	165.4	242.2	301.1
2002.....	*	1.2	4.2	11.7	29.4	57.9	110.6	176.8	261.9	328.4
2001.....	*	1.1	4.8	11.8	28.7	57.4	114.7	180.8	280.3	362.9
2000.....	*	1.5	3.6	12.9	29.5	64.7	120.7	190.1	292.8	356.6
1999.....	*	1.4	4.2	12.0	33.4	68.7	123.4	196.6	273.0	366.4

Black, male

2004.....	*	2.8	10.3	23.2	60.2	118.7	229.5	433.6	660.7	1,092.8
2003.....	*	2.3	9.8	26.2	68.3	131.7	238.8	443.7	704.3	1,076.0
2002.....	*	2.6	9.1	25.9	66.9	135.9	275.4	472.6	765.0	1,185.7
2001.....	*	2.4	10.4	26.1	65.1	135.2	286.8	483.7	830.7	1,341.2
2000.....	*	3.2	7.8	28.7	66.9	152.5	302.6	505.6	874.2	1,314.7
1999.....	*	2.9	9.2	26.6	76.0	161.5	309.3	522.9	824.0	1,346.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of penis and other and unspecified male genital organs (C60,C63)

All races 3/, both sexes

2004.....	*	*	0.1	0.2	0.2	0.3	0.5	0.6	0.5	0.6
2003.....	*	*	0.1	*	0.3	0.3	0.3	0.5	0.6	0.8
2002.....	*	*	*	0.1	0.2	0.3	0.4	0.5	0.5	0.6
2001.....	*	*	*	*	0.2	0.3	0.4	0.4	0.6	0.8
2000.....	*	*	*	*	0.2	0.3	0.3	0.4	1.0	0.7
1999.....	*	*	*	*	0.2	0.3	0.3	0.5	0.7	1.1

All races 3/, male

2004.....	*	*	0.2	0.3	0.4	0.6	1.2	1.3	1.4	2.1
2003.....	*	*	0.2	*	0.6	0.7	0.7	1.3	1.7	2.6
2002.....	*	*	*	0.3	0.5	0.6	0.9	1.2	1.3	2.0
2001.....	*	*	*	*	0.4	0.6	0.8	1.0	1.5	2.8
2000.....	*	*	*	*	0.5	0.7	0.7	1.1	2.7	2.4
1999.....	*	*	*	*	0.4	0.7	0.7	1.2	1.8	3.7

All races 3/, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	*	*	*	0.2	*	0.2	0.5	0.6	0.6	0.6
2003.....	*	*	*	*	0.3	0.3	0.3	0.5	0.6	0.7
2002.....	*	*	*	*	*	0.3	0.4	0.5	0.5	0.7
2001.....	*	*	*	*	*	0.3	0.4	0.4	0.5	0.9
2000.....	*	*	*	*	0.3	0.4	0.3	0.4	1.0	0.7
1999.....	*	*	*	*	*	0.3	0.3	0.5	0.7	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of urinary tract (C64-C68)

All races 3/, both sexes

2004.....	1.2	3.1	5.9	10.6	18.3	26.6	39.0	57.3	77.5	102.7
2003.....	1.4	2.9	5.9	10.8	18.6	26.7	39.2	56.5	77.3	101.4
2002.....	1.4	3.3	6.1	11.2	17.6	28.7	41.0	58.2	74.7	102.0
2001.....	1.7	3.2	6.3	10.7	19.3	27.3	40.9	55.2	76.1	103.9
2000.....	1.5	3.1	6.2	11.4	17.9	27.5	40.1	55.2	77.4	106.0
1999.....	1.4	2.9	6.3	11.3	18.2	26.9	40.6	54.4	73.1	103.0

All races 3/, male

2004.....	1.6	4.3	8.9	15.7	27.9	40.1	59.9	89.6	128.3	182.8
2003.....	2.0	4.5	8.6	16.0	28.4	39.9	58.6	90.0	127.6	178.8
2002.....	1.9	4.7	9.0	16.8	26.8	42.7	61.3	92.6	119.6	183.6
2001.....	2.1	4.7	9.2	15.9	28.9	41.4	62.9	87.5	126.4	192.4
2000.....	2.0	4.3	9.2	17.1	26.7	41.5	61.0	87.4	129.7	198.9
1999.....	1.9	4.3	9.2	16.9	27.2	40.6	61.8	88.2	121.7	188.4

All races 3/, female

2004.....	0.8	1.9	3.1	5.7	9.7	14.9	22.2	34.1	46.1	66.7
2003.....	0.8	1.4	3.3	5.8	9.6	15.3	23.6	32.7	46.6	67.2
2002.....	0.9	2.0	3.4	6.0	9.4	16.5	24.8	33.9	47.4	66.6
2001.....	1.2	1.8	3.4	5.8	10.5	15.2	23.3	32.7	46.0	66.9
2000.....	1.0	1.9	3.3	6.2	10.0	15.4	23.7	32.7	46.6	68.1
1999.....	0.9	1.6	3.5	6.1	10.0	15.1	23.9	30.8	44.6	68.6

White, both sexes

2004.....	1.2	3.1	6.1	11.0	19.3	27.9	40.3	59.3	80.4	105.5
2003.....	1.3	2.9	6.1	11.0	19.2	27.6	40.9	58.3	79.3	104.1
2002.....	1.4	3.4	6.3	11.5	18.3	29.8	42.4	59.6	76.4	103.9
2001.....	1.6	3.3	6.3	10.9	19.7	28.6	42.2	56.6	77.4	106.4
2000.....	1.5	3.2	6.4	11.8	18.7	28.4	41.1	56.2	78.9	109.2
1999.....	1.4	2.8	6.4	11.5	18.8	27.6	41.4	55.5	74.7	104.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of urinary tract (C64-C68)

White, male

2004.....	1.6	4.3	9.2	16.4	29.2	42.2	62.1	92.9	134.5	190.9
2003.....	1.9	4.5	8.8	16.5	29.5	41.2	61.1	92.7	132.2	186.6
2002.....	1.9	4.9	9.3	17.4	27.8	44.6	63.7	95.5	122.7	190.7
2001.....	2.1	4.9	9.3	16.4	29.5	43.5	65.2	89.8	129.9	200.5
2000.....	2.1	4.4	9.4	17.5	28.0	42.9	62.8	89.8	132.5	206.8
1999.....	1.9	4.2	9.1	17.2	28.2	41.8	63.7	90.7	125.5	194.9

White, female

2004.....	0.7	1.8	3.1	5.8	10.1	15.3	22.3	34.7	46.7	67.0
2003.....	0.7	1.2	3.4	5.7	9.7	15.6	24.4	33.3	46.7	67.6
2002.....	0.9	1.9	3.3	5.9	9.5	16.8	25.0	33.9	48.1	66.2
2001.....	1.1	1.7	3.5	5.7	10.6	15.5	23.6	33.0	45.7	67.0
2000.....	1.0	1.9	3.5	6.4	10.1	15.8	23.7	32.5	47.0	69.6
1999.....	0.9	1.5	3.7	6.1	10.1	15.2	23.8	30.8	44.7	68.4

Black, both sexes

2004.....	1.8	3.6	6.4	10.1	16.0	21.8	36.5	50.1	59.4	88.8
2003.....	2.2	3.9	6.4	11.4	16.5	25.6	30.4	49.9	69.6	89.0
2002.....	1.3	3.5	6.4	11.7	16.5	25.7	36.1	54.1	64.9	93.2
2001.....	2.4	3.5	6.9	11.1	19.8	22.5	34.5	50.4	71.2	86.6
2000.....	1.5	3.4	6.0	10.9	15.9	25.1	37.5	53.5	74.0	80.4
1999.....	1.7	4.1	7.4	12.8	17.0	26.0	39.2	49.1	61.0	93.1

Black, male

2004.....	2.5	4.9	9.3	14.1	24.3	28.6	54.4	73.7	83.2	120.2
2003.....	3.4	5.4	9.3	16.0	23.5	38.4	43.7	78.5	101.9	126.2
2002.....	1.7	4.4	8.3	16.5	24.5	34.4	47.4	77.2	98.6	135.1
2001.....	2.7	4.4	10.7	14.9	30.3	31.0	50.2	76.2	101.9	127.1
2000.....	1.7	4.2	9.1	16.3	22.6	38.0	53.5	76.5	122.2	140.7
1999.....	2.3	5.8	11.9	18.6	24.0	36.1	52.2	73.1	89.8	138.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of urinary tract (C64-C68)

Black, female

2004.....	*	2.5	3.9	6.7	9.5	16.7	24.4	36.1	46.9	76.4
2003.....	*	2.6	3.9	7.6	11.0	16.1	21.5	32.8	52.6	74.5
2002.....	*	2.8	4.8	7.8	10.2	19.2	28.5	40.3	47.3	77.2
2001.....	2.2	2.7	3.7	8.1	11.5	16.1	24.0	34.9	55.6	71.6
2000.....	*	2.7	3.5	6.5	10.6	15.6	26.9	39.6	49.7	57.9
1999.....	*	2.5	3.6	8.0	11.5	18.5	30.6	34.6	46.7	76.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of kidney and renal pelvis (C64-C65)

All races 3/, both sexes

2004.....	0.8	2.1	4.0	6.8	11.3	14.6	19.1	24.5	29.8	32.7
2003.....	1.0	2.0	4.0	7.1	11.4	14.5	19.6	24.8	30.0	33.2
2002.....	0.9	2.4	4.3	7.2	10.9	16.0	19.6	24.6	28.7	32.5
2001.....	1.2	2.4	4.4	7.1	11.9	14.6	20.8	24.4	29.1	32.3
2000.....	1.1	2.2	4.5	7.6	10.8	15.0	19.4	24.5	29.6	33.8
1999.....	1.1	2.2	4.3	7.3	10.8	14.4	19.2	23.1	26.9	31.4

All races 3/, male

2004.....	1.2	2.9	5.9	9.8	16.6	20.7	27.8	34.2	44.0	47.0
2003.....	1.5	3.0	5.7	10.4	17.2	20.2	27.7	35.9	44.9	48.7
2002.....	1.3	3.4	6.4	10.6	16.1	23.0	27.6	36.6	39.9	50.1
2001.....	1.6	3.5	6.5	10.5	17.2	21.0	30.0	34.4	42.2	52.9
2000.....	1.6	3.1	6.6	11.2	15.6	21.4	27.4	34.6	44.2	55.1
1999.....	1.5	3.2	6.1	10.6	15.8	20.3	27.3	33.7	38.9	49.4

All races 3/, female

2004.....	0.5	1.3	2.2	3.8	6.4	9.2	12.1	17.6	21.0	26.3
2003.....	0.6	1.0	2.3	4.0	6.1	9.4	13.0	16.9	20.9	26.4
2002.....	0.5	1.5	2.3	3.9	6.2	10.0	13.2	16.2	21.9	24.9
2001.....	0.9	1.4	2.4	3.9	7.0	9.1	13.5	17.4	21.2	23.7
2000.....	0.7	1.4	2.4	4.3	6.5	9.5	13.0	17.5	21.0	25.1
1999.....	0.6	1.2	2.5	4.3	6.2	9.3	12.8	15.7	19.9	24.1

White, both sexes

2004.....	0.8	2.2	4.2	7.0	11.7	15.1	19.4	24.9	30.6	33.7
2003.....	1.0	2.0	4.2	7.2	11.6	15.0	20.2	25.4	30.8	33.8
2002.....	1.0	2.4	4.4	7.4	11.2	16.6	20.2	25.0	29.2	33.2
2001.....	1.2	2.5	4.4	7.2	12.1	15.3	21.3	24.7	29.5	32.8
2000.....	1.1	2.3	4.6	7.8	11.1	15.4	19.7	24.9	30.0	34.5
1999.....	1.1	2.1	4.3	7.5	11.2	14.8	19.4	23.4	27.3	31.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of kidney and renal pelvis (C64-C65)

White, male

2004.....	1.1	3.0	6.2	10.2	17.2	21.3	28.2	34.5	45.4	48.6
2003.....	1.5	3.1	5.9	10.7	17.5	20.8	28.2	36.5	46.2	50.0
2002.....	1.4	3.5	6.6	10.9	16.4	23.8	28.3	37.5	40.2	51.9
2001.....	1.6	3.6	6.5	10.8	17.2	21.9	30.6	34.4	42.8	54.4
2000.....	1.6	3.2	6.7	11.3	16.1	21.5	27.7	35.1	44.5	56.6
1999.....	1.5	3.1	6.0	10.7	16.4	20.6	27.6	34.1	40.1	49.8

White, female

2004.....	0.5	1.3	2.3	4.0	6.7	9.6	12.2	17.8	21.3	27.0
2003.....	0.6	0.9	2.5	3.8	6.2	9.7	13.7	17.4	21.3	26.6
2002.....	0.5	1.4	2.2	4.1	6.4	10.3	13.6	16.1	22.5	25.1
2001.....	0.8	1.3	2.4	3.8	7.4	9.5	13.8	17.8	21.4	23.8
2000.....	0.7	1.4	2.5	4.5	6.5	10.0	13.3	17.7	21.4	25.5
1999.....	0.6	1.1	2.6	4.4	6.3	9.7	13.0	15.8	19.8	24.2

Black, both sexes

2004.....	1.2	2.3	4.0	6.2	10.5	13.1	19.2	26.2	25.3	28.2
2003.....	1.5	2.5	4.1	8.2	11.2	13.4	16.1	22.3	26.1	33.8
2002.....	0.9	2.8	4.4	7.0	10.6	14.3	16.8	25.8	26.7	28.2
2001.....	1.8	2.6	5.0	7.4	12.2	11.7	19.2	24.0	28.2	29.8
2000.....	1.1	2.5	4.7	7.4	10.4	14.8	18.5	23.5	29.1	29.7
1999.....	1.2	2.9	4.9	7.8	9.7	13.5	19.3	23.0	24.3	31.2

Black, male

2004.....	1.6	3.2	6.1	9.2	16.1	19.6	29.8	38.2	35.9	38.7
2003.....	2.5	3.6	6.5	11.5	17.0	19.5	25.9	35.6	39.1	43.8
2002.....	*	3.5	6.4	11.8	16.8	19.9	23.7	34.4	39.0	39.7
2001.....	1.9	3.5	7.8	10.5	20.5	17.0	29.3	39.0	42.5	46.5
2000.....	*	3.2	7.1	11.4	15.3	25.0	28.1	32.1	48.9	49.6
1999.....	1.8	4.1	8.0	12.5	14.0	20.0	28.2	34.2	27.9	49.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of kidney and renal pelvis (C64-C65)

Black, female

2004.....	*	1.5	2.2	3.7	6.1	8.3	12.1	19.1	19.7	24.0
2003.....	*	1.5	2.1	5.5	6.6	9.0	9.5	14.4	19.3	30.0
2002.....	*	2.3	2.6	3.1	5.7	10.2	12.1	20.6	20.3	23.8
2001.....	1.7	1.8	2.7	4.9	5.6	7.7	12.6	15.0	20.8	23.6
2000.....	*	1.8	2.6	4.2	6.5	7.3	12.2	18.2	19.1	22.3
1999.....	*	1.9	2.3	3.9	6.3	8.7	13.5	16.3	22.5	24.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of bladder (C67)

All races 3/, both sexes

2004.....	0.4	0.9	1.8	3.6	6.8	11.5	18.9	31.1	45.7	67.9
2003.....	0.3	0.8	1.8	3.5	6.9	11.8	18.5	30.0	45.1	64.8
2002.....	0.4	0.9	1.7	3.8	6.4	12.0	20.2	31.9	44.1	66.6
2001.....	0.4	0.8	1.7	3.4	7.0	12.0	19.0	29.3	44.8	68.9
2000.....	0.3	0.8	1.6	3.6	6.8	11.7	19.6	29.1	45.5	69.6
1999.....	0.3	0.7	1.9	3.8	7.0	11.8	20.2	29.7	43.8	68.5

All races 3/, male

2004.....	0.5	1.3	2.8	5.7	10.8	18.5	30.6	53.1	81.6	133.0
2003.....	0.4	1.4	2.7	5.5	10.9	19.1	29.6	51.7	80.3	125.1
2002.....	0.5	1.2	2.4	5.9	10.1	19.0	32.2	54.0	76.6	128.9
2001.....	0.5	1.2	2.6	5.2	11.3	19.5	31.6	50.8	81.0	134.8
2000.....	0.4	1.1	2.5	5.7	10.6	19.0	32.2	50.6	82.4	139.4
1999.....	0.4	1.1	2.9	5.9	11.0	19.3	32.9	52.3	79.5	134.8

All races 3/, female

2004.....	0.3	0.5	0.8	1.7	3.1	5.3	9.5	15.3	23.5	38.6
2003.....	0.2	0.3	0.9	1.7	3.3	5.4	9.5	14.6	23.5	38.1
2002.....	0.3	0.5	1.0	1.9	2.9	6.0	10.6	16.3	24.2	39.6
2001.....	0.3	0.4	0.9	1.7	3.2	5.6	9.0	14.3	23.2	41.4
2000.....	0.3	0.5	0.8	1.7	3.4	5.5	9.7	14.0	23.8	41.2
1999.....	0.2	0.3	0.9	1.7	3.4	5.4	10.3	13.9	22.9	41.8

White, both sexes

2004.....	0.3	0.9	1.8	3.8	7.3	12.3	19.8	32.6	47.6	69.6
2003.....	0.3	0.8	1.8	3.7	7.3	12.2	19.6	31.1	46.2	66.6
2002.....	0.4	0.9	1.8	3.9	6.7	12.5	20.9	32.9	45.2	67.7
2001.....	0.4	0.8	1.8	3.5	7.3	12.6	19.8	30.3	45.7	70.8
2000.....	0.3	0.8	1.7	3.8	7.2	12.3	20.2	29.7	46.4	72.0
1999.....	0.3	0.7	2.0	3.8	7.2	12.1	20.8	30.5	45.0	69.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of bladder (C67)

White, male

2004.....	0.4	1.3	2.9	6.0	11.6	20.2	32.3	55.9	86.2	139.3
2003.....	0.4	1.4	2.8	5.7	11.7	19.8	31.6	53.8	83.5	131.0
2002.....	0.5	1.3	2.6	6.2	10.8	20.1	33.7	56.0	79.4	134.1
2001.....	0.5	1.2	2.7	5.4	12.0	20.7	33.1	53.1	83.7	141.1
2000.....	0.4	1.2	2.7	6.0	11.4	20.2	33.6	52.7	84.6	145.7
1999.....	0.4	1.1	3.0	6.2	11.4	20.1	34.4	54.3	82.1	140.7

White, female

2004.....	0.3	0.5	0.7	1.7	3.3	5.3	9.5	15.6	23.7	38.1
2003.....	*	0.2	0.8	1.7	3.3	5.5	9.8	14.6	23.3	38.1
2002.....	0.3	0.5	1.0	1.7	2.9	5.9	10.4	16.4	24.3	38.9
2001.....	0.3	0.3	0.9	1.7	3.0	5.5	9.0	14.2	22.7	41.3
2000.....	0.3	0.5	0.8	1.8	3.4	5.5	9.6	13.6	23.7	42.1
1999.....	0.2	0.3	1.0	1.5	3.4	5.1	10.0	13.8	23.2	41.4

Black, both sexes

2004.....	*	1.2	2.0	3.3	5.1	7.9	16.5	23.0	33.4	60.1
2003.....	*	1.2	2.2	3.2	5.0	11.4	12.7	26.5	40.9	53.4
2002.....	*	*	1.6	4.0	5.2	10.9	18.6	27.0	37.2	63.3
2001.....	*	0.9	1.7	3.4	6.9	10.0	14.1	24.8	41.2	55.0
2000.....	*	0.8	1.3	3.0	5.4	9.5	17.9	28.4	44.6	49.7
1999.....	*	1.0	2.2	4.5	6.9	11.8	18.9	25.4	35.3	60.3

Black, male

2004.....	*	1.7	3.0	4.4	8.0	8.2	23.0	35.0	46.6	81.5
2003.....	*	1.7	2.8	4.5	6.6	18.2	16.4	41.1	61.3	80.3
2002.....	*	*	*	4.5	7.1	14.2	23.7	41.8	57.3	91.1
2001.....	*	*	2.8	4.2	9.2	12.7	20.5	34.4	57.8	78.3
2000.....	*	*	*	4.6	7.2	12.5	24.1	41.1	73.3	88.8
1999.....	*	*	3.9	5.4	9.8	15.6	23.7	38.9	59.3	86.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of bladder (C67)

Black, female

2004.....	*	*	*	2.5	*	7.7	12.1	15.9	26.5	51.6
2003.....	*	*	1.7	*	3.8	6.4	10.2	17.9	30.2	42.9
2002.....	*	*	*	3.7	3.7	8.5	15.2	18.1	26.7	52.6
2001.....	*	*	*	2.7	5.1	8.1	9.9	19.1	32.7	46.3
2000.....	*	*	*	*	3.9	7.3	13.8	20.8	30.1	35.2
1999.....	*	*	*	3.7	4.7	9.0	15.8	17.2	23.3	50.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other and unspecified sites within the urinary tract (C66,C68)

All races 3/, both sexes

2004.....	*	*	0.1	0.2	0.3	0.6	1.0	1.6	2.0	2.1
2003.....	*	0.1	*	0.2	0.3	0.5	1.1	1.7	2.3	3.4
2002.....	*	*	0.1	0.2	0.4	0.7	1.2	1.6	1.9	2.9
2001.....	*	*	0.1	0.2	0.4	0.7	1.1	1.5	2.2	2.7
2000.....	*	*	*	0.2	0.3	0.7	1.2	1.6	2.3	2.6
1999.....	*	*	0.1	0.2	0.4	0.7	1.1	1.6	2.3	3.1

All races 3/, male

2004.....	*	*	*	*	0.4	0.8	1.5	2.3	2.7	2.8
2003.....	*	*	*	*	*	0.6	1.2	2.3	2.4	5.1
2002.....	*	*	*	0.3	0.6	0.7	1.5	2.0	3.1	4.7
2001.....	*	*	*	*	0.4	0.9	1.3	2.2	3.3	4.7
2000.....	*	*	*	*	0.4	1.1	1.5	2.1	3.1	4.4
1999.....	*	*	*	0.3	*	1.0	1.6	2.2	3.3	4.2

All races 3/, female

2004.....	*	*	*	*	*	0.4	0.6	1.2	1.6	1.8
2003.....	*	*	*	*	*	0.4	1.0	1.2	2.2	2.7
2002.....	*	*	*	*	*	0.6	1.0	1.4	1.2	2.2
2001.....	*	*	*	*	*	0.4	0.9	1.0	1.6	1.8
2000.....	*	*	*	*	*	*	0.9	1.2	1.8	1.8
1999.....	*	*	*	*	0.4	0.4	0.8	1.2	1.8	2.7

White, both sexes

2004.....	*	*	*	0.1	0.3	0.6	1.0	1.8	2.2	2.3
2003.....	*	*	*	0.2	0.3	0.5	1.1	1.8	2.3	3.6
2002.....	*	*	*	0.2	0.4	0.7	1.3	1.7	2.0	3.0
2001.....	*	*	0.1	*	0.3	0.7	1.1	1.6	2.3	2.8
2000.....	*	*	*	0.2	0.3	0.7	1.2	1.6	2.5	2.7
1999.....	*	*	*	0.2	0.4	0.7	1.2	1.7	2.4	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of eye and adnexa (C69)										
All races 3/, both sexes										
2004.....	*	*	*	*	0.2	0.3	0.3	0.4	*	0.5
2003.....	*	*	*	0.2	0.2	*	0.3	0.4	0.5	0.7
2002.....	*	*	*	0.2	0.2	0.2	0.3	0.4	0.5	0.7
2001.....	*	*	*	*	*	0.3	0.3	0.4	0.6	0.6
2000.....	*	*	*	0.2	0.2	0.3	0.3	0.4	0.5	0.8
1999.....	*	*	*	*	*	0.2	0.3	0.4	0.6	0.8
All races 3/, male										
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	0.5	*	*	*
All races 3/, female										
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	0.7
2001.....	*	*	*	*	*	*	*	*	*	0.6
2000.....	*	*	*	*	*	*	*	*	*	0.7
1999.....	*	*	*	*	*	*	*	*	0.8	0.8
White, both sexes										
2004.....	*	*	*	*	0.2	0.3	0.4	0.5	*	0.5
2003.....	*	*	*	0.2	0.2	*	0.3	0.4	0.6	0.7
2002.....	*	*	*	0.2	0.2	*	0.3	0.5	0.6	0.8
2001.....	*	*	*	*	*	0.3	0.3	0.4	0.7	0.7
2000.....	*	*	*	*	0.2	0.3	0.4	0.5	0.5	0.8
1999.....	*	*	*	*	*	*	0.4	0.4	0.7	0.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes

2004.....	2.4	3.8	5.5	8.4	10.9	14.0	17.7	19.0	20.4	16.6
2003.....	2.9	3.9	5.8	8.5	11.2	13.8	17.3	19.4	19.6	15.2
2002.....	2.8	4.2	6.2	8.5	11.1	14.7	17.8	19.6	18.5	16.4
2001.....	2.8	4.0	6.1	8.3	11.6	14.2	16.7	20.3	20.6	16.4
2000.....	2.9	4.1	6.3	8.2	11.4	14.8	18.3	19.4	21.1	16.9
1999.....	2.8	4.3	6.1	9.2	11.7	15.7	18.3	21.0	19.4	16.0

All races 3/, male

2004.....	2.9	4.7	7.1	10.4	13.7	16.0	21.3	23.1	24.5	19.4
2003.....	3.4	5.1	7.4	9.7	13.7	16.4	21.3	24.0	25.4	20.0
2002.....	3.5	5.3	7.2	10.6	13.1	18.4	22.2	24.4	22.9	20.1
2001.....	3.5	4.9	7.5	10.7	13.9	17.3	21.2	25.4	26.5	20.1
2000.....	3.7	4.9	7.7	10.1	14.0	17.9	22.8	23.2	26.3	22.4
1999.....	3.4	5.2	7.4	10.6	14.7	19.5	22.3	26.1	24.5	20.7

All races 3/, female

2004.....	1.9	2.8	4.0	6.4	8.4	12.2	14.7	16.0	17.9	15.4
2003.....	2.4	2.8	4.3	7.3	9.0	11.6	14.1	16.2	16.0	13.1
2002.....	2.1	3.1	5.1	6.5	9.2	11.4	14.2	16.2	15.8	14.8
2001.....	2.1	3.1	4.8	6.0	9.5	11.5	13.1	16.7	17.1	14.8
2000.....	2.0	3.3	4.9	6.5	9.0	12.1	14.8	16.7	18.0	14.7
1999.....	2.3	3.4	4.9	7.9	8.9	12.4	15.1	17.5	16.5	14.1

White, both sexes

2004.....	2.6	4.1	6.0	9.2	11.9	15.2	19.1	20.2	21.7	17.3
2003.....	3.3	4.3	6.3	9.1	12.2	14.9	18.5	20.6	20.7	15.7
2002.....	3.1	4.7	6.7	9.1	12.0	15.9	19.0	20.7	19.3	17.1
2001.....	3.0	4.4	6.6	8.8	12.6	15.2	17.8	21.6	21.6	17.0
2000.....	3.1	4.5	6.9	8.8	12.3	16.0	19.7	20.5	21.8	17.5
1999.....	3.0	4.7	6.6	10.0	12.5	16.9	19.6	22.2	20.3	16.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

White, male

2004.....	3.1	5.1	7.6	11.3	14.8	17.4	23.1	24.2	25.8	20.6
2003.....	3.8	5.5	8.0	10.3	14.8	17.4	22.6	25.3	26.6	20.7
2002.....	3.9	5.9	7.9	11.2	14.0	19.9	23.6	25.6	23.7	21.2
2001.....	3.8	5.4	8.1	11.4	14.9	18.6	22.6	26.7	27.8	20.5
2000.....	4.0	5.4	8.3	10.9	15.2	19.4	24.3	24.3	27.5	23.5
1999.....	3.6	5.7	7.9	11.4	15.7	20.9	23.9	27.4	25.8	21.8

White, female

2004.....	2.1	3.1	4.4	7.2	9.1	13.2	15.7	17.2	19.2	15.8
2003.....	2.7	3.0	4.8	7.9	9.8	12.7	15.2	17.3	17.0	13.4
2002.....	2.3	3.5	5.5	7.1	10.2	12.4	15.3	17.2	16.7	15.4
2001.....	2.3	3.4	5.2	6.4	10.5	12.2	13.9	18.0	17.9	15.6
2000.....	2.1	3.5	5.5	6.9	9.7	13.0	16.0	17.7	18.5	15.0
1999.....	2.5	3.8	5.4	8.7	9.7	13.4	16.2	18.5	17.1	14.5

Black, both sexes

2004.....	1.8	2.2	3.5	4.1	6.0	7.1	9.2	10.0	9.9	10.4
2003.....	1.6	2.7	3.6	5.5	5.6	8.3	9.7	10.6	10.4	10.5
2002.....	1.3	1.9	3.7	5.7	5.6	8.0	9.5	12.2	11.3	9.5
2001.....	1.6	2.6	3.4	5.4	6.4	9.0	9.4	9.4	10.8	11.7
2000.....	2.0	2.2	3.7	5.4	6.0	7.5	8.8	10.3	14.4	10.6
1999.....	1.8	2.6	4.0	5.3	6.7	8.5	7.8	12.0	11.6	9.9

Black, male

2004.....	2.8	2.9	5.0	5.9	7.8	7.8	8.1	14.1	14.3	*
2003.....	2.4	3.1	5.1	6.6	6.7	11.8	12.2	16.0	*	*
2002.....	*	2.1	3.3	8.2	7.9	9.5	11.7	15.6	17.6	*
2001.....	1.9	3.0	4.4	6.9	9.2	9.1	11.4	12.5	*	*
2000.....	2.7	2.2	5.7	6.6	7.7	7.8	12.4	11.8	*	*
1999.....	2.6	3.7	5.3	5.9	8.7	10.9	8.8	14.2	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Black, female

2004.....	*	1.6	2.2	2.6	4.5	6.7	9.9	7.5	7.6	11.4
2003.....	*	2.3	2.2	4.6	4.7	5.7	8.0	7.3	8.5	11.7
2002.....	*	1.8	4.0	3.7	3.7	6.8	8.1	10.2	8.0	10.3
2001.....	1.3	2.3	2.6	4.1	4.3	8.8	8.1	7.5	10.2	9.7
2000.....	1.5	2.2	2.1	4.5	4.7	7.3	6.4	9.4	14.0	10.7
1999.....	*	1.6	2.8	4.8	5.2	6.7	7.2	10.6	12.5	9.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of brain (C71)

All races 3/, both sexes

2004.....	2.4	3.7	5.4	8.2	10.7	13.9	17.4	18.7	20.1	16.1
2003.....	2.9	3.9	5.7	8.3	11.1	13.6	17.1	19.0	19.2	14.6
2002.....	2.7	4.2	6.0	8.4	10.9	14.4	17.3	19.2	18.0	16.0
2001.....	2.7	3.9	6.0	8.2	11.4	14.0	16.4	19.8	20.1	16.0
2000.....	2.8	4.0	6.2	8.1	11.2	14.5	18.0	19.1	20.6	16.3
1999.....	2.7	4.2	6.0	9.1	11.4	15.5	17.9	20.5	19.0	15.5

All races 3/, male

2004.....	2.9	4.7	7.0	10.2	13.4	15.9	21.1	22.8	24.1	19.1
2003.....	3.4	5.0	7.3	9.5	13.6	16.2	21.1	23.7	25.2	19.2
2002.....	3.4	5.3	7.1	10.4	12.9	18.0	21.6	23.9	21.9	19.7
2001.....	3.4	4.8	7.4	10.5	13.7	17.0	20.8	24.9	26.0	19.7
2000.....	3.6	4.8	7.6	10.0	13.9	17.7	22.5	22.8	25.8	21.8
1999.....	3.3	5.2	7.3	10.5	14.4	19.2	21.9	25.5	24.0	20.1

All races 3/, female

2004.....	1.9	2.8	3.9	6.4	8.2	12.1	14.5	15.8	17.6	14.7
2003.....	2.3	2.8	4.2	7.2	9.0	11.4	13.9	15.7	15.5	12.6
2002.....	2.0	3.1	5.0	6.4	9.1	11.2	13.9	15.9	15.5	14.4
2001.....	2.0	3.1	4.7	6.0	9.3	11.4	12.9	16.3	16.6	14.4
2000.....	2.0	3.2	4.8	6.3	8.8	11.9	14.5	16.5	17.6	14.1
1999.....	2.2	3.3	4.7	7.8	8.7	12.3	14.8	17.1	16.0	13.6

White, both sexes

2004.....	2.6	4.0	5.9	9.1	11.6	15.1	18.8	19.9	21.3	16.7
2003.....	3.2	4.2	6.2	9.0	12.1	14.6	18.3	20.2	20.3	15.0
2002.....	3.0	4.6	6.6	9.0	11.8	15.6	18.6	20.3	18.8	16.7
2001.....	3.0	4.3	6.6	8.7	12.4	14.9	17.4	21.1	21.2	16.6
2000.....	3.0	4.4	6.8	8.7	12.2	15.7	19.4	20.2	21.4	16.9
1999.....	3.0	4.6	6.5	9.9	12.3	16.6	19.2	21.7	19.8	16.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of brain (C71)										
White, male										
2004.....	3.0	5.0	7.5	11.1	14.5	17.3	22.9	23.9	25.3	20.4
2003.....	3.7	5.5	7.8	10.1	14.7	17.2	22.4	24.9	26.4	19.8
2002.....	3.8	5.8	7.9	11.0	13.9	19.4	23.0	25.1	22.7	20.8
2001.....	3.8	5.3	8.0	11.2	14.6	18.3	22.1	26.1	27.4	20.1
2000.....	3.9	5.3	8.1	10.7	15.0	19.1	24.0	24.1	27.0	23.0
1999.....	3.5	5.6	7.8	11.3	15.4	20.6	23.5	26.8	25.3	21.2
White, female										
2004.....	2.1	3.1	4.3	7.1	9.0	13.0	15.5	16.9	18.8	15.1
2003.....	2.7	3.0	4.7	7.8	9.8	12.4	15.0	16.9	16.5	12.9
2002.....	2.2	3.5	5.4	7.0	10.0	12.2	15.0	16.9	16.4	14.9
2001.....	2.3	3.3	5.2	6.3	10.3	12.0	13.6	17.5	17.4	15.1
2000.....	2.1	3.4	5.4	6.7	9.5	12.8	15.7	17.5	18.1	14.5
1999.....	2.5	3.7	5.2	8.5	9.5	13.2	15.9	18.1	16.6	14.1
Black, both sexes										
2004.....	1.7	2.2	3.3	3.9	5.7	7.0	8.9	9.8	9.7	10.4
2003.....	1.6	2.7	3.5	5.2	5.5	8.1	9.0	10.0	10.2	10.2
2002.....	1.3	1.8	3.6	5.3	5.3	7.5	9.0	11.9	10.5	9.2
2001.....	1.5	2.6	3.3	5.3	6.2	8.9	9.4	9.2	10.6	11.7
2000.....	1.9	2.2	3.6	5.3	5.8	7.2	8.8	9.8	14.1	10.0
1999.....	1.8	2.6	3.7	5.0	6.4	8.5	7.4	11.4	10.7	9.2
Black, male										
2004.....	2.8	2.9	4.9	5.5	7.7	7.8	7.4	13.6	*	*
2003.....	2.4	3.1	5.0	6.2	6.7	11.6	11.5	16.0	*	*
2002.....	*	2.0	3.1	7.7	7.5	9.0	10.7	15.2	16.0	*
2001.....	1.8	2.9	4.2	6.7	8.7	9.1	11.4	12.5	*	*
2000.....	2.6	2.2	5.6	6.4	7.5	7.5	12.4	10.9	*	*
1999.....	2.5	3.7	4.9	5.8	8.5	10.9	7.8	13.8	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of brain (C71)										
Black, female										
2004.....	*	1.6	1.9	2.6	4.2	6.5	9.9	7.5	7.6	11.4
2003.....	*	2.3	2.2	4.4	4.5	5.5	7.3	6.5	8.1	11.3
2002.....	*	1.7	4.0	3.3	3.6	6.5	7.8	9.9	*	10.3
2001.....	*	2.3	2.6	4.1	4.2	8.6	8.1	7.2	9.8	9.7
2000.....	*	2.1	2.0	4.3	4.6	6.9	6.4	9.1	14.0	10.3
1999.....	*	1.6	2.7	4.4	4.8	6.7	7.2	10.0	11.7	8.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other parts of central nervous system (C70,C72)

All races 3/, both sexes

2004.....	*	*	*	*	0.2	*	0.3	0.3	0.4	0.5
2003.....	*	*	*	0.1	*	0.2	0.2	0.4	0.4	0.6
2002.....	*	*	0.1	0.2	0.2	0.3	0.5	0.4	0.5	*
2001.....	*	*	*	*	0.2	0.2	0.4	0.5	0.4	*
2000.....	*	*	0.1	*	*	0.2	0.3	0.3	0.4	0.6
1999.....	*	*	0.2	*	0.2	0.2	0.4	0.5	0.5	*

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	0.6	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	0.7
2003.....	*	*	*	*	*	*	*	0.5	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	*	*	*	*	0.2	*	0.3	*	0.4	0.6
2003.....	*	*	*	*	*	0.3	*	0.4	*	0.7
2002.....	*	*	*	*	*	0.3	0.5	0.4	0.5	*
2001.....	*	*	*	*	*	0.3	0.4	0.5	0.5	*
2000.....	*	*	*	*	*	0.2	0.3	*	0.4	0.5
1999.....	*	*	0.2	*	*	0.2	0.4	0.5	0.5	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of thyroid and other endocrine glands (C73-C75)

All races 3/, both sexes

2004.....	0.2	0.4	0.5	0.9	1.4	2.1	2.8	3.4	4.7	6.0
2003.....	0.2	0.4	0.6	1.0	1.4	1.8	2.6	3.5	4.2	5.1
2002.....	0.2	0.5	0.6	1.0	1.4	1.9	2.7	3.9	4.7	5.3
2001.....	0.2	0.5	0.7	1.0	1.7	2.2	3.1	3.8	4.7	5.3
2000.....	0.2	0.4	0.5	1.0	1.6	2.3	2.9	3.9	4.5	6.2
1999.....	0.2	0.4	0.6	1.1	1.6	2.0	2.6	3.9	4.4	5.8

All races 3/, male

2004.....	0.2	0.3	0.5	1.0	1.4	2.2	3.3	3.3	4.2	5.5
2003.....	0.2	0.4	0.7	0.9	1.5	1.7	2.9	3.5	3.7	4.5
2002.....	0.2	0.5	0.7	1.0	1.6	1.9	2.6	4.1	4.6	4.2
2001.....	0.3	0.5	0.7	1.0	1.8	2.5	3.3	3.4	4.7	4.5
2000.....	0.2	0.5	0.5	1.0	1.8	2.5	3.0	3.9	5.0	5.3
1999.....	0.3	0.4	0.7	1.1	1.7	2.1	3.1	4.0	4.5	5.1

All races 3/, female

2004.....	*	0.5	0.5	0.9	1.4	2.1	2.4	3.5	4.9	6.3
2003.....	*	0.4	0.6	1.0	1.4	1.8	2.5	3.5	4.5	5.4
2002.....	0.2	0.4	0.6	0.9	1.3	1.9	2.8	3.8	4.8	5.8
2001.....	0.2	0.5	0.6	0.9	1.6	1.9	3.0	4.0	4.7	5.7
2000.....	0.3	0.3	0.6	1.1	1.5	2.1	2.8	3.8	4.3	6.5
1999.....	0.2	0.4	0.5	1.1	1.6	1.9	2.2	3.8	4.4	6.1

White, both sexes

2004.....	0.2	0.4	0.5	0.9	1.4	2.1	2.9	3.5	4.7	5.9
2003.....	0.2	0.4	0.7	1.0	1.5	1.9	2.7	3.5	4.1	5.2
2002.....	0.2	0.5	0.6	1.0	1.5	2.0	2.6	4.0	4.7	5.3
2001.....	0.3	0.5	0.7	1.0	1.7	2.3	3.3	3.7	4.5	5.3
2000.....	0.2	0.4	0.5	1.1	1.6	2.2	2.8	3.9	4.6	6.2
1999.....	0.2	0.5	0.6	1.1	1.6	2.0	2.6	3.8	4.5	5.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

All races 3/, both sexes

2004.....	2.9	6.0	10.4	18.0	27.9	42.6	60.1	82.9	109.6	141.2
2003.....	3.0	6.4	11.4	18.7	28.9	43.2	61.9	86.5	110.1	147.9
2002.....	3.2	6.5	11.3	18.8	31.0	44.0	64.4	85.4	112.6	154.5
2001.....	3.3	6.3	11.3	19.9	30.4	44.2	64.4	86.9	112.4	155.2
2000.....	3.4	6.5	11.8	19.6	32.0	46.9	66.7	90.0	119.6	159.0
1999.....	3.9	7.0	12.2	22.1	35.9	52.6	74.2	96.8	128.2	168.4

All races 3/, male

2004.....	2.9	6.7	12.7	22.0	33.6	53.0	72.7	103.5	137.1	170.3
2003.....	3.1	7.4	13.3	22.7	35.2	52.7	74.4	106.9	137.6	181.3
2002.....	3.3	7.2	14.0	21.9	38.3	54.3	79.5	105.3	138.8	196.9
2001.....	3.5	7.2	13.8	23.5	37.1	52.9	79.1	107.0	138.3	196.5
2000.....	3.6	7.5	13.9	23.3	38.6	56.6	81.7	110.4	148.1	200.1
1999.....	4.0	8.3	14.3	26.5	43.1	62.9	92.2	118.6	158.2	218.9

All races 3/, female

2004.....	2.9	5.3	8.3	14.2	22.6	33.6	49.9	68.2	92.7	128.1
2003.....	2.8	5.4	9.7	14.9	23.2	34.9	51.8	72.0	93.3	133.2
2002.....	3.2	5.8	8.6	15.9	24.4	35.1	52.3	71.4	96.6	136.0
2001.....	3.0	5.4	8.9	16.4	24.3	36.8	52.8	72.9	96.9	137.9
2000.....	3.1	5.5	9.7	16.2	26.0	38.6	54.8	75.8	102.8	142.3
1999.....	3.8	5.8	10.3	18.0	29.3	43.8	60.1	81.5	110.7	148.1

White, both sexes

2004.....	2.8	5.4	9.8	17.2	27.1	41.7	60.0	83.0	109.8	143.3
2003.....	2.7	5.9	10.7	17.7	28.3	42.5	61.5	87.0	111.1	149.3
2002.....	3.0	6.1	10.7	18.2	30.0	43.3	63.6	84.8	112.5	154.9
2001.....	3.0	5.8	10.6	18.8	29.3	43.2	62.9	86.8	111.6	157.1
2000.....	3.1	5.9	11.0	19.0	30.9	46.5	65.6	88.5	120.1	160.3
1999.....	3.6	6.5	11.5	20.9	34.6	52.1	72.9	96.0	128.0	168.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

White, male

2004.....	2.6	5.9	11.6	20.7	32.2	51.2	71.6	103.6	138.0	172.2
2003.....	2.8	6.7	12.3	21.3	34.2	51.4	73.5	107.2	138.3	184.1
2002.....	2.9	6.7	13.2	21.1	36.5	53.9	78.5	103.6	139.1	198.4
2001.....	3.2	6.6	12.5	21.8	35.2	51.0	76.9	106.2	136.9	199.0
2000.....	3.2	6.7	12.8	22.1	36.1	55.9	80.1	106.9	149.0	201.3
1999.....	3.7	7.3	13.3	24.8	41.4	61.7	90.2	117.7	158.1	218.7

White, female

2004.....	2.9	4.9	8.1	13.8	22.3	33.3	50.5	67.9	92.3	130.3
2003.....	2.7	5.2	9.2	14.2	22.8	34.6	51.7	72.4	94.3	133.9
2002.....	3.2	5.4	8.2	15.5	23.9	33.9	51.4	71.3	96.2	136.0
2001.....	2.9	5.0	8.7	16.0	23.9	36.4	51.5	73.0	96.3	139.6
2000.....	3.0	5.2	9.2	16.1	26.1	38.3	53.9	75.6	102.9	143.7
1999.....	3.5	5.6	9.8	17.1	28.3	43.8	59.2	80.8	110.2	147.7

Black, both sexes

2004.....	4.5	10.6	16.9	28.4	40.9	60.1	73.4	98.3	124.5	135.9
2003.....	4.7	10.4	19.4	30.8	42.0	58.1	76.9	97.5	118.5	150.0
2002.....	4.8	10.3	17.2	28.5	46.1	58.8	84.7	108.1	127.9	168.4
2001.....	4.8	11.1	18.8	31.3	45.0	61.9	88.7	101.9	134.1	152.9
2000.....	5.9	11.5	19.9	29.1	46.8	59.5	86.2	117.5	128.8	155.2
1999.....	6.3	12.4	20.1	36.2	53.4	65.5	95.9	117.3	143.9	184.3

Black, male

2004.....	5.4	13.2	24.5	38.4	55.9	81.6	101.6	126.9	153.5	178.8
2003.....	5.4	13.4	24.3	41.2	54.4	77.1	99.9	124.6	165.4	174.1
2002.....	5.9	11.7	22.7	35.6	63.4	71.6	108.2	145.6	162.0	212.3
2001.....	5.6	13.4	26.6	41.9	62.2	82.1	117.5	138.5	170.1	208.8
2000.....	7.5	15.1	26.0	38.7	67.5	76.0	110.7	169.6	160.2	204.1
1999.....	6.5	16.7	25.7	49.0	69.0	87.8	126.2	145.3	183.1	241.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

Black, female

2004.....	3.8	8.4	10.5	20.1	29.3	44.1	54.3	81.4	109.2	118.9
2003.....	4.1	7.7	15.2	22.2	32.3	44.0	61.4	81.4	94.0	140.6
2002.....	3.8	9.1	12.6	22.7	32.5	49.4	69.0	85.6	110.2	151.6
2001.....	4.1	9.2	12.1	22.6	31.5	46.9	69.5	80.2	115.7	132.2
2000.....	4.5	8.4	14.7	21.2	30.4	47.4	69.9	86.0	112.9	136.9
1999.....	6.1	8.7	15.2	25.8	41.3	49.1	75.8	100.4	124.5	163.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes

2004.....	4.1	6.7	11.6	20.2	33.1	54.5	84.2	122.7	162.6	189.4
2003.....	4.7	7.2	11.7	20.1	35.5	55.2	87.3	127.4	165.2	198.0
2002.....	4.6	7.5	12.6	21.5	36.9	59.2	90.1	125.6	168.8	197.5
2001.....	4.8	8.1	13.0	22.7	38.1	58.8	91.8	129.0	168.5	199.9
2000.....	4.8	8.3	13.4	23.4	38.2	62.2	91.6	131.4	176.0	204.6
1999.....	4.8	8.4	13.6	25.6	39.4	62.8	93.9	130.7	174.0	199.4

All races 3/, male

2004.....	5.0	8.1	14.4	24.9	41.7	70.0	108.7	158.6	214.5	254.9
2003.....	5.8	8.8	14.5	25.2	45.6	71.9	112.3	163.5	216.8	268.7
2002.....	5.5	9.1	15.1	26.8	45.7	76.4	116.8	164.3	224.2	269.1
2001.....	5.9	9.8	16.0	27.7	47.5	75.2	116.9	165.7	220.9	275.3
2000.....	5.7	10.3	16.1	28.4	45.8	78.3	117.7	169.8	230.6	281.8
1999.....	5.9	10.2	16.7	31.0	48.6	78.1	118.9	169.4	221.8	279.0

All races 3/, female

2004.....	3.2	5.3	9.0	15.7	25.2	40.9	64.5	96.8	130.6	160.0
2003.....	3.5	5.5	9.1	15.3	26.4	40.8	67.3	101.7	133.6	166.8
2002.....	3.7	6.0	10.1	16.4	28.9	44.4	68.8	98.2	135.1	166.5
2001.....	3.7	6.4	10.1	18.1	29.5	44.6	71.8	103.4	137.1	168.4
2000.....	3.9	6.5	10.8	18.8	31.3	48.4	71.0	104.7	143.8	173.1
1999.....	3.8	6.6	10.6	20.5	31.0	49.8	74.3	103.8	145.9	167.3

White, both sexes

2004.....	4.0	6.5	11.2	19.9	33.1	55.0	85.0	125.4	167.2	194.9
2003.....	4.6	7.0	11.3	20.1	35.8	56.1	88.7	130.3	169.5	204.5
2002.....	4.6	7.2	12.1	21.0	36.9	60.3	91.7	128.7	173.6	204.0
2001.....	4.6	7.9	12.8	22.3	38.0	59.0	93.7	131.9	172.8	205.9
2000.....	4.7	8.0	13.2	23.3	38.5	63.5	93.4	134.4	181.5	208.7
1999.....	4.7	8.3	13.4	25.3	39.2	63.8	95.6	133.1	178.4	204.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

White, male										
2004.....	5.0	7.8	13.9	24.8	41.9	70.4	109.7	162.3	221.1	264.0
2003.....	5.7	8.7	13.9	25.1	46.3	73.6	115.2	168.7	224.3	279.5
2002.....	5.5	8.8	14.8	26.5	45.7	77.9	119.2	169.0	231.0	279.4
2001.....	5.7	9.7	15.8	27.1	47.3	75.6	119.2	169.8	226.4	284.8
2000.....	5.8	9.9	15.6	28.3	46.1	80.4	119.0	174.1	238.0	290.2
1999.....	5.6	10.0	16.5	30.6	48.6	79.4	121.0	172.1	228.5	285.8
White, female										
2004.....	3.0	5.2	8.6	15.2	25.0	41.4	64.6	98.3	133.7	163.8
2003.....	3.4	5.4	8.8	15.3	26.1	40.7	67.0	102.6	135.8	171.3
2002.....	3.7	5.7	9.6	15.8	28.8	44.8	69.4	99.9	138.5	171.2
2001.....	3.5	6.1	9.8	17.7	29.3	44.4	73.0	105.0	140.4	172.8
2000.....	3.6	6.1	10.7	18.6	31.4	48.7	72.8	106.5	147.9	175.5
1999.....	3.8	6.6	10.3	20.2	30.5	50.3	75.5	105.7	148.8	171.3
Black, both sexes										
2004.....	5.2	9.1	16.8	26.3	40.8	60.7	93.3	115.8	134.7	154.0
2003.....	6.3	9.5	17.3	23.5	41.1	59.4	90.5	117.4	139.1	150.5
2002.....	5.7	10.5	17.3	29.8	43.9	62.3	93.2	111.8	138.2	152.3
2001.....	6.6	10.4	16.5	30.2	44.9	67.8	90.7	116.7	142.7	155.7
2000.....	6.3	12.5	17.5	29.1	44.0	62.2	90.5	114.3	131.6	175.5
1999.....	6.5	10.4	18.2	33.2	45.9	64.6	90.4	119.1	139.3	162.0
Black, male										
2004.....	5.9	11.3	21.4	31.5	51.2	80.4	122.7	139.2	168.6	193.7
2003.....	8.0	11.9	22.5	32.0	51.1	73.4	105.1	137.8	166.8	184.5
2002.....	7.2	13.1	19.7	36.0	55.6	78.8	123.2	137.8	177.3	202.6
2001.....	8.2	11.9	21.0	37.5	56.9	87.9	114.1	141.2	191.0	206.5
2000.....	6.6	14.9	21.8	36.7	54.2	73.1	125.4	147.9	167.8	224.9
1999.....	8.4	13.3	22.5	41.2	55.2	80.7	114.6	159.1	164.8	220.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Black, female

2004.....	4.6	7.2	12.9	21.9	32.8	46.1	73.4	101.9	116.8	138.2
2003.....	4.8	7.4	12.9	16.6	33.2	49.0	80.6	105.3	124.6	137.3
2002.....	4.3	8.3	15.2	24.7	34.7	50.1	73.0	96.3	117.8	133.1
2001.....	5.3	9.2	12.7	24.2	35.4	52.9	75.1	102.1	118.1	136.8
2000.....	6.0	10.4	13.9	22.9	36.0	54.1	67.3	94.0	113.4	157.1
1999.....	4.8	8.0	14.5	26.6	38.6	52.7	74.3	95.0	126.6	140.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hodgkin's disease (C81)

All races 3/, both sexes

2004.....	0.3	0.4	0.4	0.7	0.8	1.0	1.4	1.8	2.1	2.2
2003.....	0.3	0.3	0.4	0.6	0.8	0.9	1.5	2.1	2.3	2.4
2002.....	0.3	0.4	0.5	0.5	0.8	1.0	1.6	1.7	2.3	2.7
2001.....	0.4	0.4	0.5	0.4	0.7	1.1	1.2	2.0	2.5	2.2
2000.....	0.3	0.4	0.5	0.5	0.9	1.0	1.6	1.8	2.4	2.5
1999.....	0.3	0.4	0.5	0.7	0.9	1.3	1.8	2.0	2.3	2.4

All races 3/, male

2004.....	0.3	0.5	0.5	0.9	1.0	1.4	1.9	2.0	3.2	3.2
2003.....	0.5	0.4	0.5	0.7	1.1	1.1	2.0	2.4	2.9	2.6
2002.....	0.4	0.5	0.6	0.7	1.0	1.4	2.2	2.0	3.2	3.6
2001.....	0.5	0.5	0.6	0.6	0.9	1.5	1.6	2.5	2.9	2.6
2000.....	0.4	0.5	0.6	0.5	1.1	1.3	1.9	2.3	2.7	3.7
1999.....	0.4	0.5	0.6	0.9	1.2	1.6	2.3	2.3	2.8	3.8

All races 3/, female

2004.....	0.3	0.3	0.3	0.5	0.5	0.8	1.0	1.6	1.5	1.7
2003.....	0.2	0.2	0.4	0.5	0.6	0.6	1.2	1.8	1.9	2.3
2002.....	0.3	0.3	0.4	0.3	0.7	0.6	1.1	1.5	1.7	2.3
2001.....	0.3	0.4	0.4	0.3	0.6	0.7	0.9	1.7	2.2	2.0
2000.....	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.4	2.3	2.0
1999.....	0.3	0.3	0.4	0.5	0.5	0.9	1.5	1.8	2.0	1.8

White, both sexes

2004.....	0.3	0.4	0.4	0.7	0.8	1.1	1.5	1.9	2.2	2.3
2003.....	0.4	0.4	0.5	0.6	0.9	0.8	1.7	2.2	2.4	2.5
2002.....	0.4	0.4	0.5	0.5	0.8	1.0	1.7	1.8	2.4	2.9
2001.....	0.4	0.4	0.5	0.4	0.8	1.2	1.3	2.1	2.6	2.4
2000.....	0.3	0.4	0.5	0.5	0.9	1.0	1.6	2.0	2.5	2.6
1999.....	0.4	0.4	0.5	0.7	0.9	1.3	2.0	2.2	2.5	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

All races 3/, both sexes

2004.....	1.5	2.8	4.8	7.8	12.7	21.1	32.1	47.0	65.1	75.3
2003.....	1.7	3.0	4.7	8.1	14.3	21.6	32.9	49.4	65.5	79.0
2002.....	1.9	3.0	5.0	8.8	14.9	23.3	35.1	50.3	67.3	77.6
2001.....	2.0	3.4	5.6	9.6	15.6	23.1	36.2	51.0	70.1	80.1
2000.....	2.1	3.7	5.8	10.7	16.1	25.1	36.4	53.9	72.1	80.7
1999.....	2.1	3.6	6.2	10.9	16.3	25.7	38.5	54.8	72.8	78.0

All races 3/, male

2004.....	2.1	3.7	6.1	9.8	15.7	26.7	39.7	57.6	81.9	96.3
2003.....	2.2	3.9	6.0	10.1	18.7	27.9	41.2	61.0	81.8	104.0
2002.....	2.4	3.9	6.3	11.3	18.4	30.2	43.0	61.9	84.8	100.2
2001.....	2.7	4.4	7.4	11.9	19.1	29.3	45.0	61.4	87.8	107.1
2000.....	2.6	4.8	7.5	13.4	19.2	30.7	45.3	66.6	88.5	106.6
1999.....	2.8	4.8	7.9	13.5	19.6	30.9	46.8	68.4	85.9	104.7

All races 3/, female

2004.....	0.9	1.8	3.5	5.9	9.9	16.3	25.9	39.4	54.7	65.8
2003.....	1.2	2.1	3.5	6.1	10.4	16.2	26.2	41.3	55.5	68.0
2002.....	1.3	2.2	3.9	6.4	11.7	17.4	28.7	42.1	56.6	67.8
2001.....	1.3	2.5	3.8	7.5	12.4	17.8	29.2	43.7	59.5	68.8
2000.....	1.5	2.6	4.1	8.2	13.3	20.3	29.4	45.0	62.5	70.2
1999.....	1.4	2.4	4.6	8.3	13.3	21.2	31.9	45.3	65.2	67.3

White, both sexes

2004.....	1.5	2.7	4.7	8.0	13.1	22.2	33.6	49.4	68.9	79.0
2003.....	1.7	2.9	4.8	8.5	14.9	22.8	34.4	52.0	69.0	83.6
2002.....	1.8	3.0	5.0	9.0	15.6	24.5	36.6	53.0	70.6	82.4
2001.....	2.0	3.4	5.7	9.8	16.3	24.0	38.1	53.7	73.3	83.8
2000.....	2.1	3.6	5.9	11.2	17.0	26.4	38.3	56.7	76.2	84.8
1999.....	2.0	3.6	6.5	11.2	17.1	27.2	40.3	57.2	76.9	81.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

White, male

2004.....	2.1	3.6	6.1	10.0	16.2	27.7	41.4	60.3	86.6	101.7
2003.....	2.2	3.7	6.0	10.6	19.4	29.7	43.3	64.2	86.5	110.1
2002.....	2.3	4.0	6.2	11.6	19.1	31.7	44.8	65.2	88.9	106.1
2001.....	2.7	4.4	7.5	12.0	19.8	30.5	46.9	64.3	91.7	111.9
2000.....	2.7	4.7	7.6	13.8	20.1	32.4	46.9	70.1	93.1	112.5
1999.....	2.7	4.7	8.1	13.8	20.3	32.4	49.0	70.5	91.1	108.3

White, female

2004.....	0.9	1.8	3.4	6.1	10.2	17.4	27.2	41.4	57.9	68.8
2003.....	1.2	2.1	3.5	6.4	10.7	16.8	27.0	43.1	58.3	71.8
2002.....	1.3	2.1	3.7	6.5	12.3	18.2	29.9	44.3	59.5	72.1
2001.....	1.3	2.4	3.9	7.7	13.1	18.3	30.9	46.2	62.2	72.1
2000.....	1.4	2.6	4.3	8.8	14.1	21.2	31.3	47.2	66.1	73.5
1999.....	1.4	2.4	4.8	8.7	14.2	22.7	33.4	47.8	68.5	70.7

Black, both sexes

2004.....	1.7	3.9	5.9	7.7	11.0	15.9	21.7	29.1	27.0	39.2
2003.....	2.3	3.9	5.9	6.1	12.7	14.7	22.8	26.5	31.0	31.5
2002.....	2.7	3.5	6.1	9.3	12.8	17.7	25.7	27.1	35.6	31.2
2001.....	2.4	4.3	5.9	9.9	12.4	19.2	23.5	25.9	38.7	39.6
2000.....	2.3	4.6	5.5	8.9	12.6	16.6	22.7	26.3	28.5	36.9
1999.....	2.8	4.1	5.7	10.6	11.7	17.0	23.9	32.1	31.5	40.8

Black, male

2004.....	2.2	5.4	7.8	10.0	14.4	19.6	28.5	31.8	34.4	42.7
2003.....	3.0	5.8	7.5	7.7	16.0	17.2	24.6	28.8	34.7	34.4
2002.....	3.8	4.0	7.4	11.5	16.6	23.2	34.4	31.7	42.0	42.9
2001.....	3.4	5.1	8.7	12.4	15.5	24.9	31.6	31.1	48.2	55.6
2000.....	2.6	6.3	7.9	13.7	15.7	19.3	32.1	31.2	32.0	46.1
1999.....	3.9	5.6	8.0	13.9	15.7	23.4	28.5	47.5	28.8	59.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

Black, female

2004.....	1.3	2.5	4.3	5.9	8.3	13.1	17.1	27.5	23.1	37.8
2003.....	1.6	2.3	4.5	4.7	10.1	12.8	21.5	25.2	29.0	30.4
2002.....	1.7	3.1	5.0	7.6	9.8	13.7	19.9	24.4	32.2	26.7
2001.....	1.6	3.6	3.6	8.0	9.9	15.0	18.2	22.7	33.9	33.7
2000.....	2.1	3.2	3.5	5.0	10.1	14.6	16.4	23.4	26.7	33.5
1999.....	1.8	2.8	3.7	7.9	8.7	12.3	20.9	22.9	32.8	33.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Non-Hodgkin's lymphoma (C82-C85)

All races 3/, both sexes

2004.....	1.5	2.8	4.8	7.7	12.6	21.1	32.0	46.9	64.8	75.0
2003.....	1.7	3.0	4.7	8.1	14.3	21.6	32.8	49.3	65.4	78.8
2002.....	1.9	3.0	5.0	8.8	14.9	23.2	35.0	50.2	67.0	77.3
2001.....	2.0	3.4	5.5	9.6	15.6	23.0	36.0	50.9	69.8	79.6
2000.....	2.1	3.6	5.8	10.6	16.1	25.0	36.3	53.7	71.8	80.2
1999.....	2.1	3.5	6.2	10.8	16.3	25.6	38.4	54.6	72.6	77.7

All races 3/, male

2004.....	2.1	3.7	6.1	9.8	15.7	26.7	39.6	57.5	81.6	95.8
2003.....	2.2	3.9	6.0	10.1	18.7	27.8	41.0	60.7	81.8	103.6
2002.....	2.4	3.9	6.2	11.3	18.4	30.0	42.8	61.8	84.4	99.7
2001.....	2.7	4.4	7.4	11.9	19.1	29.2	44.7	61.3	87.3	106.6
2000.....	2.6	4.7	7.5	13.3	19.2	30.5	45.2	66.3	88.1	105.9
1999.....	2.8	4.7	7.9	13.5	19.6	30.8	46.8	68.1	85.7	104.4

All races 3/, female

2004.....	0.9	1.8	3.5	5.8	9.8	16.2	25.8	39.3	54.5	65.7
2003.....	1.2	2.1	3.5	6.1	10.4	16.2	26.1	41.2	55.4	67.8
2002.....	1.3	2.2	3.8	6.4	11.7	17.3	28.7	41.9	56.4	67.6
2001.....	1.3	2.5	3.8	7.5	12.4	17.7	29.1	43.6	59.3	68.3
2000.....	1.5	2.6	4.1	8.2	13.3	20.3	29.2	45.0	62.2	69.8
1999.....	1.4	2.4	4.5	8.3	13.3	21.1	31.8	45.2	65.0	66.9

White, both sexes

2004.....	1.5	2.7	4.7	7.9	13.1	22.2	33.5	49.2	68.6	78.8
2003.....	1.7	2.9	4.7	8.4	14.9	22.8	34.2	51.8	69.0	83.3
2002.....	1.8	3.0	4.9	9.0	15.6	24.5	36.5	52.9	70.4	82.1
2001.....	2.0	3.4	5.7	9.8	16.2	23.9	37.9	53.6	73.0	83.4
2000.....	2.1	3.6	5.9	11.2	16.9	26.3	38.1	56.5	75.8	84.3
1999.....	2.0	3.6	6.4	11.1	17.0	27.1	40.2	57.0	76.7	81.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Non-Hodgkin's lymphoma (C82-C85)

White, male

2004.....	2.1	3.6	6.1	9.9	16.2	27.7	41.3	60.1	86.2	101.2
2003.....	2.1	3.7	6.0	10.6	19.4	29.6	43.1	63.9	86.5	109.7
2002.....	2.3	4.0	6.2	11.6	19.1	31.6	44.5	65.0	88.6	105.6
2001.....	2.6	4.4	7.4	12.0	19.7	30.3	46.6	64.2	91.2	111.5
2000.....	2.7	4.7	7.6	13.7	20.0	32.2	46.8	69.8	92.6	111.8
1999.....	2.7	4.7	8.1	13.7	20.2	32.3	48.9	70.2	90.8	108.0

White, female

2004.....	0.9	1.8	3.4	6.0	10.2	17.3	27.1	41.3	57.7	68.7
2003.....	1.2	2.1	3.5	6.4	10.7	16.7	27.0	43.0	58.2	71.6
2002.....	1.3	2.1	3.7	6.5	12.3	18.2	29.9	44.2	59.3	71.8
2001.....	1.3	2.4	3.9	7.7	13.0	18.3	30.9	46.0	62.0	71.6
2000.....	1.4	2.6	4.3	8.8	14.1	21.2	31.1	47.2	65.8	73.1
1999.....	1.4	2.4	4.8	8.7	14.1	22.6	33.3	47.7	68.3	70.4

Black, both sexes

2004.....	1.7	3.9	5.9	7.6	11.0	15.9	21.7	28.9	26.5	38.6
2003.....	2.3	3.9	5.9	6.1	12.7	14.6	22.8	26.5	31.0	31.2
2002.....	2.7	3.5	6.1	9.3	12.8	17.6	25.7	26.8	35.3	30.6
2001.....	2.4	4.3	5.9	9.8	12.3	19.1	23.4	25.7	37.9	39.3
2000.....	2.3	4.5	5.5	8.8	12.6	16.6	22.7	25.9	28.2	36.3
1999.....	2.8	4.1	5.6	10.5	11.7	17.0	23.9	32.0	31.5	40.5

Black, male

2004.....	2.2	5.4	7.8	10.0	14.4	19.6	28.5	31.8	34.4	41.7
2003.....	3.0	5.7	7.5	7.7	16.0	17.2	24.6	28.8	34.7	34.4
2002.....	3.8	4.0	7.4	11.5	16.6	22.9	34.4	31.7	41.3	42.9
2001.....	3.4	5.1	8.7	12.2	15.5	24.6	31.6	30.7	47.4	54.5
2000.....	2.6	6.1	7.9	13.5	15.7	19.3	32.1	30.7	32.0	43.8
1999.....	3.9	5.6	7.9	13.7	15.7	23.4	28.5	47.0	28.8	59.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Non-Hodgkin's lymphoma (C82-C85)

Black, female

2004.....	1.3	2.5	4.2	5.6	8.3	13.1	17.1	27.2	22.3	37.4
2003.....	1.6	2.3	4.5	4.7	10.1	12.6	21.5	25.2	29.0	30.0
2002.....	1.7	3.1	5.0	7.6	9.8	13.7	19.9	23.9	32.2	25.9
2001.....	1.6	3.6	3.6	7.8	9.8	15.0	17.9	22.7	33.1	33.7
2000.....	2.1	3.2	3.5	5.0	10.1	14.6	16.4	23.1	26.3	33.5
1999.....	1.7	2.8	3.7	7.9	8.7	12.3	20.9	22.9	32.8	33.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant immunoproliferative diseases (C88)

All races 3/, both sexes

2004.....	*	*	*	*	*	0.3	0.6	0.9	1.1	1.5
2003.....	*	*	*	*	0.2	0.2	0.8	1.2	1.1	2.2
2002.....	*	*	*	*	0.2	0.3	0.6	1.2	1.7	2.0
2001.....	*	*	*	*	0.2	0.4	0.6	0.8	1.6	2.0
2000.....	*	*	*	*	*	0.3	0.6	0.9	1.7	2.1
1999.....	*	*	*	*	0.4	0.3	0.7	1.2	1.6	2.7

All races 3/, male

2004.....	*	*	*	*	*	*	0.8	1.2	1.6	2.5
2003.....	*	*	*	*	*	*	1.2	1.7	1.7	3.6
2002.....	*	*	*	*	0.4	0.5	0.9	1.7	2.9	2.8
2001.....	*	*	*	*	*	0.6	0.7	1.1	2.2	2.9
2000.....	*	*	*	*	*	*	0.8	1.3	2.4	3.4
1999.....	*	*	*	*	0.5	0.5	1.1	1.7	2.3	3.9

All races 3/, female

2004.....	*	*	*	*	*	*	*	0.7	0.8	1.1
2003.....	*	*	*	*	*	*	0.4	0.8	0.7	1.6
2002.....	*	*	*	*	*	*	*	0.8	1.0	1.6
2001.....	*	*	*	*	*	*	0.5	0.6	1.2	1.7
2000.....	*	*	*	*	*	*	0.4	0.5	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	0.9	1.2	2.2

White, both sexes

2004.....	*	*	*	*	*	0.3	0.6	1.0	1.2	1.7
2003.....	*	*	*	*	*	0.2	0.9	1.3	1.2	2.4
2002.....	*	*	*	*	0.3	0.3	0.6	1.3	1.8	2.1
2001.....	*	*	*	*	*	0.4	0.6	0.9	1.7	2.2
2000.....	*	*	*	*	*	0.3	0.6	0.9	1.8	2.2
1999.....	*	*	*	*	0.4	0.4	0.8	1.2	1.8	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Multiple myeloma and malignant plasma cell neoplasms (C90)

All races 3/, both sexes

2004.....	0.5	1.0	2.2	4.5	7.6	12.1	18.4	27.4	32.8	31.6
2003.....	0.5	1.1	2.4	4.3	7.8	12.9	19.6	27.4	33.4	34.1
2002.....	0.4	1.2	2.4	4.7	8.0	13.8	20.5	26.7	34.1	34.7
2001.....	0.5	1.2	2.3	5.0	8.3	13.6	20.5	27.4	33.1	34.2
2000.....	0.5	1.2	2.6	4.4	8.3	14.1	19.8	27.4	34.6	35.7
1999.....	0.5	1.2	2.4	5.4	8.7	13.8	19.8	26.3	34.5	33.9

All races 3/, male

2004.....	0.5	1.1	2.7	5.1	9.1	14.9	22.3	33.8	41.2	42.1
2003.....	0.7	1.3	3.0	5.3	9.8	15.7	23.1	32.2	41.6	46.8
2002.....	0.5	1.5	2.9	5.5	9.8	16.4	25.0	33.8	43.3	48.0
2001.....	0.6	1.5	2.8	5.8	9.9	16.2	24.6	34.5	41.2	45.1
2000.....	0.6	1.4	2.8	4.8	9.2	16.6	23.4	34.4	43.7	48.3
1999.....	0.6	1.3	2.9	6.4	10.5	16.0	23.1	32.4	43.2	47.4

All races 3/, female

2004.....	0.4	0.9	1.8	3.9	6.1	9.6	15.2	22.8	27.6	26.9
2003.....	0.4	0.9	1.9	3.4	5.9	10.5	16.8	23.9	28.4	28.5
2002.....	0.4	1.0	1.9	3.9	6.4	11.6	16.9	21.7	28.6	29.0
2001.....	0.4	0.8	1.9	4.2	7.0	11.3	17.2	22.5	28.3	29.7
2000.....	0.4	1.0	2.3	4.1	7.5	12.0	16.9	22.6	29.3	30.6
1999.....	0.4	1.1	2.0	4.6	7.1	11.8	17.3	22.1	29.4	28.4

White, both sexes

2004.....	0.4	0.9	1.9	3.9	6.9	11.0	16.8	26.0	31.3	30.3
2003.....	0.5	0.9	2.0	3.9	7.2	11.9	18.2	25.6	31.9	32.6
2002.....	0.4	1.1	2.1	4.1	7.2	12.7	19.2	25.6	33.1	33.6
2001.....	0.4	1.0	2.0	4.3	7.3	12.5	19.2	26.1	31.3	32.8
2000.....	0.4	1.0	2.2	3.7	7.6	13.1	18.5	25.6	33.4	33.0
1999.....	0.4	1.1	2.0	4.6	7.5	12.8	18.4	24.5	33.2	31.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Multiple myeloma and malignant plasma cell neoplasms (C90)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
White, male										
2004.....	0.5	1.0	2.3	4.6	8.5	13.8	20.8	32.5	39.6	40.7
2003.....	0.6	1.1	2.5	4.8	9.1	14.7	22.1	31.0	40.0	45.8
2002.....	0.4	1.3	2.6	4.9	9.0	15.4	23.9	32.9	42.3	47.3
2001.....	0.5	1.3	2.5	5.0	8.7	15.0	23.4	33.4	39.3	44.0
2000.....	0.5	1.2	2.4	4.1	8.6	15.6	21.9	32.5	42.9	46.7
1999.....	0.4	1.2	2.6	5.5	9.5	15.2	21.4	30.8	41.9	45.4
White, female										
2004.....	0.3	0.8	1.5	3.3	5.5	8.5	13.4	21.2	26.1	25.6
2003.....	0.3	0.7	1.6	3.0	5.3	9.4	15.1	21.7	26.9	26.7
2002.....	0.3	0.8	1.5	3.4	5.6	10.4	15.4	20.4	27.4	27.7
2001.....	*	0.7	1.6	3.6	6.0	10.2	15.9	21.0	26.5	28.1
2000.....	0.2	0.8	2.0	3.3	6.7	10.9	15.8	20.8	27.8	27.4
1999.....	0.3	0.9	1.5	3.8	5.7	10.7	16.0	20.1	28.1	26.5
Black, both sexes										
2004.....	1.0	2.0	5.2	10.2	15.3	24.3	38.0	48.2	58.9	53.9
2003.....	1.2	2.7	5.7	9.3	15.0	25.3	37.3	51.9	58.6	59.8
2002.....	1.0	2.9	5.6	10.5	16.2	26.3	37.7	44.1	54.2	54.3
2001.....	1.5	2.6	4.8	11.8	19.0	25.5	38.1	47.6	60.9	57.1
2000.....	1.1	3.1	6.1	11.6	16.5	27.2	37.6	52.8	57.0	70.1
1999.....	1.4	2.4	6.4	13.6	19.6	24.1	37.5	50.2	56.1	62.5
Black, male										
2004.....	*	2.1	6.1	11.2	18.1	30.5	44.0	56.9	72.5	71.5
2003.....	1.5	3.1	7.5	12.1	19.5	29.1	39.4	56.6	73.1	73.0
2002.....	*	3.5	5.9	12.2	20.0	30.7	44.7	51.4	69.5	70.8
2001.....	1.6	3.4	5.3	14.5	23.0	29.7	45.1	57.1	79.5	69.2
2000.....	*	3.3	7.0	12.2	18.4	31.0	48.1	66.2	67.4	76.1
1999.....	1.9	2.6	6.4	15.9	20.8	26.0	46.5	58.4	73.2	80.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and malignant plasma cell neoplasms (C90)										
Black, female										
2004.....	*	1.9	4.4	9.3	13.1	19.6	33.8	43.1	51.8	46.9
2003.....	*	2.4	4.3	7.0	11.4	22.5	35.9	49.1	51.1	54.7
2002.....	*	2.3	5.3	9.2	13.3	23.1	32.9	39.8	46.2	48.1
2001.....	1.5	1.9	4.3	9.5	15.8	22.3	33.4	41.9	51.5	52.6
2000.....	*	2.8	5.4	11.2	15.0	24.4	30.6	44.7	51.8	67.8
1999.....	*	2.3	6.4	11.8	18.6	22.7	31.5	45.2	47.5	55.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
All races 3/, both sexes										
2004.....	1.8	2.5	4.2	7.1	12.0	19.9	31.8	45.6	61.5	78.8
2003.....	2.1	2.7	4.0	7.0	12.4	19.6	32.5	47.4	62.9	80.3
2002.....	1.9	2.8	4.5	7.4	12.9	20.9	32.4	45.7	63.4	80.5
2001.....	1.9	3.0	4.6	7.6	13.1	20.6	33.3	47.7	61.3	81.4
2000.....	1.9	3.1	4.5	7.7	12.7	21.7	33.2	47.4	65.1	83.6
1999.....	1.9	3.2	4.4	8.5	13.1	21.8	33.1	46.3	62.7	82.4
All races 3/, male										
2004.....	2.1	2.8	5.0	9.0	15.7	26.7	43.9	64.1	86.6	110.7
2003.....	2.4	3.2	4.9	8.9	15.8	26.8	44.9	66.3	88.8	111.8
2002.....	2.2	3.2	5.3	9.2	16.1	28.0	45.8	65.0	90.0	114.5
2001.....	2.1	3.4	5.2	9.2	17.4	27.6	44.9	66.2	86.9	117.5
2000.....	2.1	3.6	5.1	9.7	16.0	29.2	46.2	65.1	93.4	119.8
1999.....	2.1	3.6	5.2	10.1	16.8	29.0	45.7	64.6	87.6	119.2
All races 3/, female										
2004.....	1.6	2.2	3.3	5.4	8.6	14.1	22.1	32.3	46.0	64.5
2003.....	1.7	2.2	3.2	5.2	9.4	13.3	22.7	33.9	47.0	66.4
2002.....	1.6	2.5	3.8	5.7	10.0	14.7	21.8	32.1	47.3	65.7
2001.....	1.7	2.7	4.0	6.0	9.3	14.6	24.1	34.8	45.9	66.3
2000.....	1.7	2.6	3.9	5.9	9.7	15.3	22.9	35.1	48.4	68.8
1999.....	1.7	2.8	3.6	7.0	9.8	15.6	23.2	33.6	48.1	67.6
White, both sexes										
2004.....	1.8	2.5	4.2	7.3	12.2	20.5	32.5	47.1	63.6	81.6
2003.....	2.1	2.8	4.0	7.1	12.7	20.3	33.6	49.3	65.0	83.4
2002.....	2.0	2.7	4.5	7.4	13.1	21.8	33.6	47.0	65.7	83.0
2001.....	1.9	3.1	4.5	7.6	13.5	20.9	34.5	49.1	63.8	84.7
2000.....	1.9	3.0	4.5	7.8	12.8	22.6	34.4	49.2	67.6	86.0
1999.....	1.9	3.2	4.3	8.7	13.3	22.2	34.2	48.0	64.1	85.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
White, male										
2004.....	2.1	2.8	4.9	9.2	16.0	27.2	44.4	66.2	89.8	115.3
2003.....	2.4	3.4	4.8	9.0	16.4	27.7	46.4	69.2	92.8	116.9
2002.....	2.3	3.0	5.3	9.3	16.3	29.0	47.2	67.0	93.3	119.1
2001.....	2.0	3.5	5.1	9.4	17.8	27.8	46.5	68.3	90.1	123.1
2000.....	2.2	3.5	5.0	9.8	16.1	30.5	47.4	67.5	96.7	123.6
1999.....	2.1	3.5	5.1	10.3	17.0	29.5	47.1	66.7	90.2	124.1
White, female										
2004.....	1.6	2.3	3.4	5.4	8.6	14.5	22.6	33.2	47.3	66.4
2003.....	1.7	2.3	3.2	5.4	9.3	13.8	23.1	34.9	47.9	68.6
2002.....	1.7	2.5	3.8	5.6	10.1	15.4	22.6	32.8	48.9	67.3
2001.....	1.7	2.7	3.9	6.0	9.5	14.8	24.7	35.4	47.9	68.6
2000.....	1.6	2.5	4.0	5.9	9.8	15.7	23.9	36.3	50.2	70.8
1999.....	1.8	2.9	3.6	7.2	9.8	15.8	23.9	34.9	48.7	69.9
Black, both sexes										
2004.....	2.2	2.9	5.3	7.7	13.8	19.4	32.5	37.1	47.0	60.1
2003.....	2.5	2.5	5.1	7.2	12.8	17.9	29.6	37.3	48.0	59.2
2002.....	1.7	3.6	5.1	9.1	13.8	17.3	28.6	39.2	46.6	64.7
2001.....	2.3	3.0	5.6	7.9	12.3	22.3	28.0	41.3	42.0	57.1
2000.....	2.4	4.5	5.2	7.9	14.1	17.9	28.4	34.7	43.8	66.0
1999.....	2.0	3.5	5.4	8.1	13.6	22.2	27.9	35.0	51.1	55.8
Black, male										
2004.....	2.3	3.1	7.1	9.4	17.6	28.3	48.6	48.2	58.8	79.5
2003.....	3.1	2.7	6.8	10.7	14.5	24.9	39.4	50.2	58.3	77.2
2002.....	1.7	4.8	5.7	11.0	17.2	23.4	42.4	52.4	64.2	86.8
2001.....	2.7	2.8	6.6	9.9	17.1	32.3	36.4	51.6	61.0	77.2
2000.....	2.5	4.9	6.2	10.0	18.6	22.4	43.5	49.6	64.1	98.0
1999.....	2.2	4.4	6.9	10.3	17.2	29.1	38.0	51.3	61.0	77.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Leukemia (C91-C95)

Black, female

2004.....	2.0	2.6	3.8	6.3	10.9	12.8	21.5	30.5	40.8	52.4
2003.....	2.1	2.4	3.8	4.3	11.4	12.8	23.0	29.6	42.6	52.3
2002.....	1.6	2.6	4.7	7.6	11.2	12.8	19.3	31.3	37.4	56.3
2001.....	2.0	3.3	4.7	6.3	8.5	15.0	22.4	35.2	32.3	49.7
2000.....	2.3	4.2	4.4	6.2	10.5	14.6	18.4	25.6	33.5	54.1
1999.....	1.7	2.6	4.2	6.2	10.8	17.1	21.2	25.2	46.2	47.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Lymphoid leukemia (C91)

All races 3/, both sexes

2004.....	0.4	0.5	1.0	1.9	3.0	5.0	8.4	12.0	17.0	28.9
2003.....	0.4	0.6	0.9	1.8	3.0	5.2	8.5	12.9	18.8	29.6
2002.....	0.5	0.7	1.1	1.8	3.3	5.3	8.2	12.2	18.6	30.1
2001.....	0.4	0.6	1.1	1.9	3.3	5.4	8.7	12.9	18.2	31.2
2000.....	0.4	0.6	1.0	1.8	3.2	5.5	8.9	13.1	19.5	32.0
1999.....	0.3	0.6	1.1	2.2	3.5	6.0	9.2	13.4	19.7	30.8

All races 3/, male

2004.....	0.5	0.7	1.3	2.7	4.6	6.8	12.0	17.0	24.7	41.7
2003.....	0.6	0.8	1.2	2.7	4.2	8.2	12.5	18.5	27.7	41.7
2002.....	0.6	0.8	1.5	2.4	4.4	7.8	12.5	17.7	26.9	43.0
2001.....	0.5	0.7	1.4	2.6	4.9	8.1	12.2	18.4	25.7	43.8
2000.....	0.5	0.8	1.2	2.7	4.6	8.2	12.8	18.9	28.4	43.8
1999.....	0.4	0.7	1.5	3.0	5.1	8.6	13.1	19.1	28.7	45.6

All races 3/, female

2004.....	0.3	0.4	0.8	1.1	1.6	3.3	5.4	8.3	12.3	23.1
2003.....	0.3	0.5	0.5	1.0	1.9	2.7	5.3	8.9	13.3	24.3
2002.....	0.3	0.5	0.7	1.3	2.3	3.2	4.8	8.3	13.6	24.5
2001.....	0.3	0.5	0.7	1.1	1.9	3.1	5.8	9.0	13.8	25.9
2000.....	0.2	0.5	0.7	1.0	1.9	3.2	5.8	9.1	14.2	27.3
1999.....	0.3	0.5	0.6	1.4	2.1	3.7	6.2	9.4	14.4	24.8

White, both sexes

2004.....	0.4	0.6	1.0	2.0	3.0	5.1	8.5	12.3	17.6	30.2
2003.....	0.4	0.6	0.9	1.9	3.1	5.4	8.7	13.4	19.7	31.0
2002.....	0.5	0.7	1.2	1.8	3.3	5.5	8.5	12.3	19.5	31.6
2001.....	0.4	0.6	1.0	1.9	3.4	5.3	9.0	13.2	19.1	32.8
2000.....	0.4	0.6	1.0	1.9	3.2	5.7	9.1	13.6	20.3	33.1
1999.....	0.4	0.6	1.1	2.1	3.5	6.1	9.6	13.9	20.3	32.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Lymphoid leukemia (C91)

White, male

2004.....	0.5	0.7	1.3	2.9	4.5	6.8	11.9	17.4	25.7	43.7
2003.....	0.6	0.8	1.1	2.7	4.4	8.5	13.0	19.3	29.2	43.5
2002.....	0.7	0.8	1.6	2.4	4.5	8.2	12.8	17.8	28.3	45.1
2001.....	0.5	0.7	1.4	2.7	4.9	8.0	12.7	19.0	26.6	46.2
2000.....	0.5	0.8	1.2	2.8	4.6	8.4	13.1	19.5	29.5	45.0
1999.....	0.4	0.7	1.6	3.0	5.2	8.7	13.5	19.8	29.8	47.7

White, female

2004.....	0.3	0.4	0.8	1.1	1.6	3.6	5.6	8.5	12.5	24.1
2003.....	0.3	0.5	0.6	1.0	1.9	2.7	5.2	9.1	13.9	25.4
2002.....	0.3	0.5	0.7	1.2	2.2	3.2	5.0	8.4	14.1	25.8
2001.....	0.3	0.4	0.6	1.1	2.0	2.9	5.9	9.0	14.5	27.3
2000.....	0.2	0.5	0.8	1.0	2.0	3.3	6.0	9.5	14.8	28.3
1999.....	0.3	0.5	0.6	1.4	2.1	3.8	6.5	9.8	14.7	25.7

Black, both sexes

2004.....	*	*	1.2	2.0	4.2	5.2	10.1	12.0	14.4	19.7
2003.....	*	*	1.1	2.0	2.9	5.6	9.5	11.2	12.2	19.8
2002.....	*	*	1.0	2.6	4.3	5.2	7.9	14.3	13.1	18.4
2001.....	*	0.9	1.7	2.4	4.0	7.4	7.9	13.0	13.5	19.0
2000.....	*	1.0	1.1	1.5	3.5	4.5	9.1	10.5	13.6	25.0
1999.....	*	*	1.2	2.8	4.0	7.0	8.0	10.0	16.2	22.0

Black, male

2004.....	*	*	*	2.6	6.7	9.0	17.2	17.7	20.1	30.8
2003.....	*	*	2.0	3.2	*	8.4	11.8	15.1	17.0	26.1
2002.....	*	*	*	3.3	5.7	7.0	11.7	21.6	18.3	31.1
2001.....	*	*	2.2	3.1	6.5	11.2	10.8	16.7	22.5	28.4
2000.....	*	*	*	*	6.0	7.3	13.7	16.5	21.1	40.4
1999.....	*	*	*	3.9	5.5	10.9	12.2	15.2	21.8	32.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lymphoid leukemia (C91)										
Black, female										
2004.....	*	*	*	*	*	*	5.3	8.6	11.3	15.4
2003.....	*	*	*	*	*	*	8.0	9.0	9.7	17.4
2002.....	*	*	*	*	3.1	3.9	5.4	9.9	10.3	13.6
2001.....	*	*	*	*	*	4.7	6.1	10.8	9.0	15.6
2000.....	*	*	*	*	*	*	6.0	6.8	9.8	19.3
1999.....	*	*	*	*	*	4.0	5.2	6.9	13.4	18.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Myeloid leukemia (C92)

All races 3/, both sexes

2004.....	1.1	1.6	2.4	3.7	6.6	10.7	15.5	21.7	26.9	25.6
2003.....	1.2	1.5	2.4	3.7	6.9	10.1	16.3	22.4	25.5	26.1
2002.....	1.1	1.6	2.5	4.0	7.1	10.9	16.1	21.6	26.9	25.7
2001.....	1.2	1.9	2.7	4.2	6.9	10.7	16.8	22.2	25.5	25.1
2000.....	1.2	1.8	2.7	4.3	6.8	11.1	16.0	22.0	26.7	25.3
1999.....	1.2	2.0	2.5	4.6	6.5	10.8	15.2	19.5	24.5	25.6

All races 3/, male

2004.....	1.2	1.6	2.8	4.4	7.9	14.5	20.7	29.9	36.8	36.1
2003.....	1.4	1.8	2.7	4.5	8.4	13.2	21.7	31.0	35.8	37.9
2002.....	1.2	1.8	2.8	5.0	8.6	14.1	22.1	30.5	38.6	37.6
2001.....	1.3	2.0	2.8	4.9	9.0	13.4	22.2	29.9	37.0	37.2
2000.....	1.2	2.0	3.1	5.0	8.2	14.1	21.9	29.7	39.1	39.1
1999.....	1.2	2.1	2.8	5.2	8.0	13.8	20.7	26.3	34.6	37.7

All races 3/, female

2004.....	1.0	1.5	2.0	3.1	5.3	7.4	11.3	15.8	20.8	20.8
2003.....	1.1	1.3	2.1	2.9	5.5	7.4	12.0	16.3	19.3	20.9
2002.....	1.0	1.5	2.3	3.1	5.7	8.2	11.3	15.3	19.9	20.5
2001.....	1.2	1.8	2.6	3.6	5.1	8.4	12.4	16.9	18.6	20.1
2000.....	1.2	1.6	2.3	3.7	5.6	8.5	11.3	16.6	19.4	19.7
1999.....	1.2	1.9	2.2	4.1	5.2	8.3	10.9	14.7	18.6	20.7

White, both sexes

2004.....	1.1	1.6	2.4	3.8	6.8	11.1	16.1	22.6	27.7	26.7
2003.....	1.2	1.6	2.4	3.8	7.1	10.6	17.0	23.5	26.4	26.9
2002.....	1.2	1.6	2.5	4.0	7.3	11.4	16.9	22.5	27.9	26.5
2001.....	1.2	1.9	2.7	4.3	7.3	11.0	17.4	23.0	26.5	26.0
2000.....	1.1	1.7	2.7	4.3	6.8	11.6	16.8	23.0	27.7	25.9
1999.....	1.2	2.0	2.5	4.8	6.6	11.1	15.7	20.2	25.2	26.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Myeloid leukemia (C92)

White, male

2004.....	1.2	1.6	2.7	4.6	8.3	15.0	21.6	30.9	38.0	38.3
2003.....	1.4	1.9	2.7	4.5	8.8	13.7	22.6	32.6	37.6	39.6
2002.....	1.2	1.7	2.8	5.0	8.8	14.5	23.1	31.8	39.9	38.7
2001.....	1.2	2.0	2.8	4.9	9.4	13.7	22.9	30.9	38.7	39.1
2000.....	1.2	2.0	3.1	5.0	8.2	14.8	22.6	31.4	40.5	41.0
1999.....	1.2	2.1	2.7	5.4	8.0	14.0	21.4	27.2	35.8	39.5

White, female

2004.....	1.0	1.5	2.1	3.1	5.4	7.7	11.7	16.5	21.4	21.5
2003.....	1.1	1.3	2.1	3.1	5.5	7.8	12.3	16.9	19.6	21.3
2002.....	1.1	1.5	2.2	3.0	5.9	8.6	11.8	15.9	20.6	21.2
2001.....	1.2	1.8	2.6	3.7	5.3	8.7	13.0	17.4	19.1	20.5
2000.....	1.1	1.5	2.4	3.7	5.5	8.8	12.1	17.2	20.1	19.8
1999.....	1.2	1.9	2.2	4.2	5.4	8.5	11.1	15.3	18.9	21.6

Black, both sexes

2004.....	1.1	1.7	3.1	3.7	6.8	9.4	11.1	14.7	20.3	17.8
2003.....	1.5	1.6	2.9	3.0	7.0	7.8	13.6	15.5	18.3	21.0
2002.....	0.8	1.9	2.4	4.8	6.6	8.4	12.5	15.1	18.1	19.3
2001.....	1.6	1.6	2.8	4.2	4.6	10.4	12.8	17.2	15.7	17.2
2000.....	1.7	2.5	3.2	4.5	7.9	8.6	11.2	13.3	16.1	20.3
1999.....	1.4	2.1	2.8	3.4	6.7	9.6	12.4	14.1	17.9	14.3

Black, male

2004.....	*	*	4.2	4.2	7.5	12.8	14.6	17.7	25.1	19.9
2003.....	1.5	*	3.2	4.8	7.1	11.6	17.7	22.4	19.9	31.3
2002.....	*	2.5	2.5	5.8	8.1	11.0	17.7	19.7	22.9	32.2
2001.....	1.7	*	3.1	5.3	6.3	14.0	17.8	20.9	19.3	*
2000.....	1.8	2.5	3.8	5.1	9.3	9.6	17.1	15.6	22.8	24.2
1999.....	*	2.5	3.7	4.4	8.7	12.5	16.6	19.9	20.9	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Myeloid leukemia (C92)

Black, female

2004.....	*	1.9	2.2	3.3	6.3	6.8	8.8	12.9	17.8	16.9
2003.....	1.4	1.8	2.6	*	6.9	4.9	10.9	11.4	17.4	17.0
2002.....	*	*	2.4	3.9	5.4	6.5	9.0	12.3	15.5	14.4
2001.....	1.4	1.8	2.5	3.3	3.2	7.7	9.4	15.0	13.9	16.4
2000.....	1.7	2.6	2.6	3.9	6.9	7.9	7.3	12.0	12.7	18.9
1999.....	1.4	1.6	*	*	5.2	7.5	9.7	10.6	16.4	12.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Monocytic leukemia (C93)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	0.4	0.5	0.6
2003.....	*	*	*	*	*	*	*	*	0.6	0.6
2002.....	*	*	*	*	*	*	*	0.4	0.4	0.6
2001.....	*	*	*	*	*	*	*	0.4	0.5	0.7
2000.....	*	*	*	*	*	*	0.2	0.3	0.4	0.7
1999.....	*	*	*	*	*	0.2	0.3	0.5	0.6	0.9

All races 3/, male

2004.....	*	*	*	*	*	*	*	0.7	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	0.7	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	0.7	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	0.7

White, both sexes

2004.....	*	*	*	*	*	*	*	0.4	0.5	0.6
2003.....	*	*	*	*	*	*	*	*	0.7	0.7
2002.....	*	*	*	*	*	*	*	0.4	0.4	0.6
2001.....	*	*	*	*	*	*	*	0.4	0.6	0.8
2000.....	*	*	*	*	*	*	*	0.4	*	0.7
1999.....	*	*	*	*	*	*	0.3	0.5	0.6	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified leukemia (C94-C95)

All races 3/, both sexes

2004.....	0.4	0.4	0.8	1.5	2.4	4.1	7.9	11.5	17.1	23.8
2003.....	0.4	0.5	0.7	1.4	2.4	4.2	7.5	11.8	17.9	24.0
2002.....	0.4	0.5	0.9	1.5	2.4	4.5	7.9	11.6	17.5	24.1
2001.....	0.3	0.5	0.8	1.4	2.8	4.4	7.7	12.2	17.0	24.4
2000.....	0.4	0.6	0.8	1.6	2.6	5.0	8.0	12.0	18.5	25.6
1999.....	0.4	0.6	0.8	1.7	3.0	4.8	8.3	13.0	17.8	25.2

All races 3/, male

2004.....	0.4	0.5	0.9	1.8	3.1	5.2	11.0	16.5	24.4	31.7
2003.....	0.5	0.7	0.9	1.7	3.0	5.3	10.4	16.4	24.6	31.1
2002.....	0.4	0.6	1.0	1.7	3.0	5.9	10.9	16.1	24.3	32.6
2001.....	0.4	0.7	0.9	1.7	3.4	6.0	10.2	17.3	23.6	35.3
2000.....	0.4	0.8	0.7	2.0	3.1	6.8	11.1	16.1	25.1	35.7
1999.....	0.4	0.7	0.8	1.8	3.7	6.3	11.5	18.4	23.4	34.6

All races 3/, female

2004.....	0.4	0.3	0.6	1.2	1.7	3.2	5.4	7.9	12.7	20.3
2003.....	0.3	0.4	0.5	1.2	1.9	3.2	5.2	8.6	13.9	20.8
2002.....	0.3	0.5	0.8	1.3	2.0	3.2	5.5	8.4	13.4	20.4
2001.....	0.2	0.4	0.7	1.2	2.2	3.0	5.6	8.6	13.1	19.8
2000.....	0.3	0.5	0.9	1.2	2.1	3.5	5.6	9.2	14.6	21.4
1999.....	0.3	0.5	0.8	1.5	2.4	3.4	5.9	9.2	14.6	21.3

White, both sexes

2004.....	0.4	0.4	0.7	1.5	2.4	4.1	7.7	11.9	17.7	24.1
2003.....	0.4	0.6	0.7	1.4	2.5	4.2	7.7	12.2	18.2	24.8
2002.....	0.4	0.5	0.8	1.5	2.4	4.7	8.1	11.8	18.0	24.2
2001.....	0.3	0.6	0.8	1.5	2.7	4.4	7.8	12.5	17.6	25.1
2000.....	0.4	0.6	0.8	1.6	2.7	5.1	8.2	12.2	19.2	26.3
1999.....	0.4	0.6	0.7	1.7	3.1	4.8	8.6	13.4	18.0	25.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified leukemia (C94-C95)

White, male

2004.....	0.4	0.5	0.9	1.8	3.2	5.2	10.8	17.2	25.4	32.1
2003.....	0.5	0.7	0.9	1.6	3.1	5.4	10.6	16.9	25.2	32.7
2002.....	0.4	0.5	0.9	1.8	3.0	6.0	10.9	16.6	24.9	34.0
2001.....	0.3	0.7	0.9	1.7	3.3	6.0	10.6	17.8	24.1	36.4
2000.....	0.4	0.7	0.7	1.9	3.2	7.0	11.3	16.1	26.0	36.4
1999.....	0.5	0.7	0.8	1.8	3.9	6.5	11.9	19.0	23.6	35.4

White, female

2004.....	0.3	0.3	0.6	1.2	1.7	3.2	5.2	8.0	13.0	20.5
2003.....	0.3	0.5	0.5	1.2	1.9	3.1	5.4	8.8	13.9	21.4
2002.....	0.3	0.4	0.8	1.3	1.9	3.5	5.7	8.4	13.7	20.0
2001.....	0.2	0.4	0.6	1.2	2.1	3.0	5.6	8.8	13.7	20.3
2000.....	0.3	0.5	0.9	1.2	2.2	3.5	5.7	9.4	15.1	22.2
1999.....	0.3	0.5	0.7	1.5	2.4	3.2	6.0	9.5	14.7	21.8

Black, both sexes

2004.....	0.7	*	1.0	2.0	2.9	4.7	11.3	10.2	12.4	22.3
2003.....	*	*	1.2	2.1	2.9	4.5	6.2	10.4	17.5	18.4
2002.....	*	1.0	1.6	1.7	3.0	3.7	8.0	9.4	15.4	26.1
2001.....	*	*	1.0	*	3.6	4.5	7.3	10.9	12.5	20.6
2000.....	*	0.9	*	2.0	2.6	4.6	8.0	10.8	13.8	20.0
1999.....	*	*	1.5	1.8	2.9	5.4	7.2	10.5	16.5	19.4

Black, male

2004.....	*	*	*	*	*	6.5	16.8	12.3	*	27.8
2003.....	*	*	*	*	*	*	9.5	12.8	21.4	*
2002.....	*	*	*	*	*	5.5	13.0	10.6	22.9	22.5
2001.....	*	*	*	*	4.3	7.1	7.7	13.9	18.5	29.5
2000.....	*	*	*	*	*	5.5	12.7	17.5	19.4	32.3
1999.....	*	*	*	*	*	5.7	8.8	15.2	18.3	24.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified leukemia (C94-C95)

Black, female

2004.....	*	*	*	*	*	*	7.5	8.9	11.7	20.1
2003.....	*	*	*	*	*	4.4	*	9.0	15.5	17.8
2002.....	*	*	*	*	*	*	4.7	8.8	11.5	27.5
2001.....	*	*	*	*	*	*	7.0	9.2	9.4	17.3
2000.....	*	*	*	*	*	4.0	4.9	6.8	11.0	15.5
1999.....	*	*	*	*	*	5.2	6.1	7.7	15.6	17.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Benign neoplasms (D10-D36)

All races 3/, both sexes

2004.....	0.1	0.3	0.3	0.5	0.8	0.9	1.7	3.0	4.5	7.9
2003.....	0.2	0.3	0.4	0.5	0.8	1.1	1.7	2.8	4.9	7.4
2002.....	0.2	0.4	0.4	0.5	0.8	1.4	1.8	3.1	4.3	7.8
2001.....	0.2	0.2	0.4	0.6	0.9	1.3	1.7	2.9	4.6	8.7
2000.....	0.2	0.2	0.4	0.4	0.7	1.2	1.9	3.2	5.5	9.0
1999.....	0.1	0.2	0.3	0.6	0.9	1.3	1.8	3.4	4.8	8.4

All races 3/, male

2004.....	*	0.2	0.3	0.5	0.6	0.8	1.5	2.9	4.2	7.4
2003.....	0.2	0.2	0.4	0.5	0.8	1.1	1.8	2.7	4.9	7.3
2002.....	*	0.3	0.3	0.5	0.8	1.4	1.9	3.1	3.8	6.3
2001.....	*	*	0.4	0.6	1.0	1.3	1.6	3.1	5.1	8.9
2000.....	*	0.3	0.4	0.4	0.5	1.3	1.7	3.2	4.9	8.7
1999.....	0.2	*	0.3	0.6	0.9	1.1	2.1	3.1	4.6	9.2

All races 3/, female

2004.....	0.2	0.4	0.4	0.6	0.9	1.0	1.8	3.0	4.7	8.1
2003.....	0.2	0.4	0.3	0.5	0.7	1.1	1.5	2.8	4.9	7.4
2002.....	0.3	0.4	0.4	0.5	0.7	1.5	1.8	3.2	4.6	8.4
2001.....	0.2	0.3	0.4	0.6	0.8	1.2	1.7	2.7	4.3	8.6
2000.....	0.3	0.2	0.4	0.5	0.9	1.1	2.0	3.3	5.9	9.1
1999.....	*	0.2	0.3	0.6	0.9	1.5	1.6	3.7	4.9	8.1

White, both sexes

2004.....	0.1	0.3	0.3	0.5	0.8	0.8	1.5	2.8	4.4	7.8
2003.....	0.2	0.3	0.4	0.5	0.8	1.0	1.5	2.8	4.8	7.4
2002.....	0.2	0.3	0.4	0.5	0.7	1.3	1.8	3.1	4.1	7.4
2001.....	0.2	0.2	0.4	0.5	0.7	1.1	1.6	2.8	4.3	8.2
2000.....	0.2	0.2	0.4	0.4	0.7	1.1	1.7	3.2	5.5	8.9
1999.....	0.1	0.2	0.3	0.5	0.8	1.2	1.7	3.2	4.7	8.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of eye, brain and other parts of central nervous system (D31-D33)

All races 3/, both sexes

2004.....	*	0.1	0.2	0.3	0.5	0.6	1.2	1.7	3.5	5.6
2003.....	0.1	0.1	0.2	0.3	0.5	0.8	1.1	1.8	3.4	5.4
2002.....	0.1	0.2	0.2	0.3	0.4	0.9	1.2	2.1	2.9	5.1
2001.....	*	0.1	0.2	0.3	0.5	0.8	1.2	1.8	3.0	5.9
2000.....	*	0.1	0.2	0.2	0.5	0.7	1.2	2.3	3.7	6.2
1999.....	*	*	0.2	0.4	0.6	0.9	1.1	2.2	3.5	5.5

All races 3/, male

2004.....	*	*	*	*	*	*	1.0	1.5	3.2	5.2
2003.....	*	*	*	0.3	0.4	0.7	1.1	1.6	3.4	5.2
2002.....	*	*	*	*	0.4	0.8	1.2	1.7	2.4	3.2
2001.....	*	*	*	0.3	0.4	0.8	1.0	1.8	2.7	5.4
2000.....	*	*	*	*	*	0.8	1.1	2.0	3.3	5.9
1999.....	*	*	*	0.3	0.4	0.8	1.2	1.9	3.2	5.5

All races 3/, female

2004.....	*	*	0.2	0.4	0.6	0.8	1.3	1.9	3.6	5.8
2003.....	*	*	*	0.3	0.5	0.8	1.1	2.0	3.3	5.5
2002.....	*	*	0.3	0.3	0.4	1.0	1.2	2.3	3.2	5.9
2001.....	*	*	0.2	0.3	0.5	0.8	1.3	1.9	3.1	6.2
2000.....	*	*	*	0.3	0.6	0.7	1.3	2.4	4.0	6.4
1999.....	*	*	*	0.4	0.7	1.1	1.1	2.5	3.7	5.5

White, both sexes

2004.....	*	0.1	0.2	0.2	0.4	0.4	1.0	1.6	3.3	5.7
2003.....	*	0.1	0.2	0.2	0.5	0.6	1.0	1.9	3.4	5.4
2002.....	*	0.2	0.2	0.3	0.4	0.8	1.2	2.0	2.8	4.9
2001.....	*	*	0.2	0.2	0.4	0.7	1.1	1.8	2.8	5.5
2000.....	*	*	0.2	0.2	0.5	0.7	1.1	2.2	3.7	6.3
1999.....	*	*	0.1	0.3	0.5	0.9	1.1	2.1	3.4	5.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of other and unspecified sites (D10-D24,D29-D30,D34-D36)										
All races 3/, both sexes										
2004.....	*	0.1	0.1	0.2	0.3	0.3	0.5	1.2	1.0	2.2
2003.....	0.1	0.1	0.2	0.2	0.3	0.4	0.5	0.9	1.4	1.9
2002.....	0.1	0.2	0.1	0.2	0.3	0.5	0.6	1.0	1.3	2.6
2001.....	0.1	0.1	0.2	0.3	0.4	0.5	0.5	1.0	1.6	2.5
2000.....	*	0.1	0.2	0.2	0.2	0.4	0.6	1.0	1.7	2.6
1999.....	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.3	2.8
All races 3/, male										
2004.....	*	*	*	0.3	0.4	0.5	*	1.5	1.0	2.3
2003.....	*	*	*	*	0.4	*	0.7	1.2	1.5	2.1
2002.....	*	*	*	0.3	0.4	0.6	0.7	1.4	1.4	3.1
2001.....	*	*	0.2	0.3	0.5	0.5	0.5	1.3	2.4	3.5
2000.....	*	*	0.2	*	*	0.5	0.6	1.1	1.6	2.9
1999.....	*	*	*	*	0.5	*	0.9	1.2	1.4	3.7
All races 3/, female										
2004.....	*	*	*	*	*	*	0.5	1.1	1.0	2.2
2003.....	*	*	*	*	*	*	*	0.7	1.4	1.8
2002.....	*	0.2	*	*	*	0.4	0.5	0.8	1.3	2.4
2001.....	*	*	*	*	*	0.4	*	0.8	1.1	2.1
2000.....	*	*	*	*	*	0.4	0.6	0.8	1.8	2.6
1999.....	*	*	*	*	*	0.4	0.5	1.2	1.2	2.4
White, both sexes										
2004.....	*	*	*	0.2	0.4	0.4	0.4	1.1	1.1	2.0
2003.....	*	0.1	0.2	0.2	0.3	0.4	0.5	0.9	1.4	1.9
2002.....	*	0.1	0.1	0.2	0.3	0.4	0.5	1.1	1.3	2.3
2001.....	*	*	0.2	0.3	0.4	0.4	0.5	1.0	1.4	2.5
2000.....	*	*	0.2	0.2	0.2	0.4	0.5	0.9	1.8	2.5
1999.....	*	*	*	0.2	0.3	0.3	0.6	1.1	1.3	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

All races 3/, both sexes

2004.....	0.6	1.0	1.5	2.7	4.7	7.9	14.3	23.1	40.9	65.6
2003.....	0.6	1.0	1.5	2.5	5.0	8.3	14.1	24.8	40.9	69.6
2002.....	0.7	1.0	1.5	3.0	4.4	7.9	14.0	24.7	40.0	67.0
2001.....	0.7	1.1	1.7	3.0	5.1	8.7	15.6	25.6	41.5	70.1
2000.....	0.7	1.1	1.8	3.2	5.5	9.0	14.9	24.6	43.2	72.0
1999.....	0.7	1.1	1.8	2.8	5.1	9.1	16.2	24.9	41.1	70.8

All races 3/, male

2004.....	0.8	1.1	1.8	3.2	6.1	10.5	18.1	30.8	55.3	87.9
2003.....	0.7	1.1	1.8	3.0	6.0	10.1	18.8	32.2	55.2	91.6
2002.....	0.8	1.2	1.7	3.9	5.3	10.5	17.5	33.2	53.7	91.9
2001.....	0.8	1.4	1.9	3.2	6.3	10.9	20.3	35.1	55.7	96.6
2000.....	0.8	1.2	2.1	3.6	6.7	11.6	18.9	32.6	58.8	97.6
1999.....	0.9	1.2	2.2	3.2	6.7	11.8	21.2	33.2	55.0	97.8

All races 3/, female

2004.....	0.5	0.9	1.1	2.2	3.4	5.6	11.2	17.6	32.0	55.5
2003.....	0.4	0.9	1.2	2.1	4.2	6.7	10.3	19.5	32.2	59.9
2002.....	0.6	0.9	1.3	2.3	3.5	5.6	11.1	18.7	31.7	56.2
2001.....	0.6	0.9	1.5	2.9	4.1	6.9	11.8	19.0	33.0	59.0
2000.....	0.6	1.0	1.4	2.8	4.4	6.8	11.7	19.0	33.9	61.5
1999.....	0.6	1.0	1.4	2.3	3.6	6.8	12.2	19.2	33.0	60.0

White, both sexes

2004.....	0.6	1.0	1.4	2.6	4.6	8.1	14.8	23.8	42.4	68.2
2003.....	0.6	1.0	1.5	2.5	5.1	8.5	14.6	25.8	42.2	72.1
2002.....	0.6	1.1	1.5	3.0	4.5	8.2	14.4	25.6	41.1	69.3
2001.....	0.7	1.2	1.7	3.2	5.3	9.2	15.9	26.5	42.8	71.6
2000.....	0.7	1.1	1.7	3.3	5.5	9.3	15.0	25.3	44.9	74.3
1999.....	0.7	1.1	1.8	2.8	5.1	9.2	16.7	25.6	42.5	73.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

White, male

2004.....	0.8	1.0	1.7	3.1	6.0	10.9	18.6	31.6	57.6	92.1
2003.....	0.8	1.1	1.8	2.9	6.0	10.5	19.5	33.7	56.9	95.6
2002.....	0.8	1.3	1.6	3.7	5.5	10.8	18.2	34.4	54.9	95.4
2001.....	0.8	1.4	2.0	3.4	6.4	11.4	21.0	36.1	57.3	99.2
2000.....	0.7	1.2	2.1	3.7	6.6	11.9	18.9	33.4	61.3	101.1
1999.....	0.9	1.2	2.2	3.3	6.9	11.8	21.9	34.2	57.1	102.7

White, female

2004.....	0.4	0.9	1.1	2.2	3.3	5.7	11.6	18.0	32.9	57.4
2003.....	0.5	0.9	1.2	2.2	4.3	6.7	10.7	20.0	33.1	61.7
2002.....	0.5	0.9	1.3	2.2	3.6	5.9	11.4	19.3	32.6	57.9
2001.....	0.6	0.9	1.5	2.9	4.3	7.2	11.8	19.7	34.1	60.1
2000.....	0.6	1.0	1.4	2.8	4.5	7.0	11.8	19.6	35.0	63.4
1999.....	0.6	1.0	1.4	2.3	3.5	6.9	12.5	19.6	33.9	61.6

Black, both sexes

2004.....	0.8	1.0	2.1	3.8	5.9	7.4	12.0	19.6	30.7	42.3
2003.....	*	1.5	2.1	3.2	5.2	7.8	11.9	19.1	32.5	46.7
2002.....	1.1	1.1	2.0	4.2	4.3	7.3	11.9	19.2	33.2	49.9
2001.....	1.0	0.9	1.9	2.8	5.1	6.9	14.8	19.8	33.3	55.6
2000.....	*	1.4	2.5	3.0	5.7	8.0	15.6	20.1	28.5	51.6
1999.....	1.0	1.4	2.1	2.9	5.6	10.5	14.2	21.6	30.3	48.8

Black, male

2004.....	*	*	2.6	5.0	8.0	9.7	16.5	25.9	35.9	48.7
2003.....	*	*	2.1	4.3	6.6	8.9	17.1	21.5	45.0	56.3
2002.....	*	*	2.4	6.3	4.2	9.5	13.4	24.8	45.1	66.5
2001.....	*	*	*	*	6.3	9.1	17.8	27.4	46.6	71.5
2000.....	*	*	2.9	3.5	7.9	11.4	19.4	27.9	32.9	64.6
1999.....	*	*	*	3.6	5.9	14.5	17.6	26.1	34.9	52.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

Black, female

2004.....	*	*	*	2.7	4.2	5.8	9.0	15.9	28.0	39.8
2003.....	*	1.4	2.0	*	4.2	6.9	8.4	17.6	25.9	42.9
2002.....	*	*	*	2.6	4.3	5.7	11.0	15.9	27.1	43.5
2001.....	*	*	2.1	3.0	4.2	5.3	12.8	15.3	26.6	49.7
2000.....	*	*	2.2	2.7	3.9	5.4	13.1	15.4	26.3	46.8
1999.....	*	*	*	*	5.3	7.5	11.9	18.9	28.1	47.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Neoplasms of uncertain or unknown behavior of unspecified sites (D48)

All races 3/, both sexes

2004.....	*	0.2	0.2	0.3	0.3	0.4	0.8	1.1	1.7	4.1
2003.....	*	*	0.1	0.2	0.3	0.4	0.5	0.9	1.5	3.4
2002.....	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.9	1.9	3.8
2001.....	*	0.1	0.2	0.2	0.3	0.5	0.6	0.9	1.6	3.8
2000.....	*	0.1	*	0.2	0.4	0.3	0.6	1.1	1.9	3.9
1999.....	*	*	0.2	*	0.3	0.3	0.6	0.9	1.5	3.5

All races 3/, male

2004.....	*	0.2	0.2	0.3	0.4	0.4	0.9	1.0	1.7	3.6
2003.....	*	*	*	*	*	*	0.5	1.0	1.5	2.6
2002.....	*	*	*	*	*	*	0.6	0.9	1.5	3.0
2001.....	*	*	*	*	*	0.5	0.6	0.9	1.4	1.8
2000.....	*	*	*	*	*	*	*	1.1	1.6	2.4
1999.....	*	*	*	*	*	0.5	0.6	0.9	1.2	2.1

All races 3/, female

2004.....	*	*	*	0.2	0.3	0.4	0.7	1.2	1.7	4.4
2003.....	*	*	*	*	*	0.4	0.5	0.8	1.5	3.8
2002.....	*	*	*	*	0.4	*	0.4	0.9	2.2	4.2
2001.....	*	*	*	0.3	*	0.4	0.7	0.9	1.7	4.6
2000.....	*	*	*	*	0.4	*	0.6	1.1	2.0	4.4
1999.....	*	*	*	*	*	*	0.6	0.9	1.7	4.1

White, both sexes

2004.....	*	0.2	0.2	0.3	0.3	0.4	0.8	1.2	1.7	4.2
2003.....	*	*	0.1	0.2	0.3	0.4	0.6	0.9	1.6	3.5
2002.....	*	0.1	*	0.2	0.3	0.4	0.5	1.0	2.0	3.9
2001.....	*	0.1	0.2	0.2	0.3	0.4	0.5	0.9	1.6	4.0
2000.....	*	0.1	*	0.2	0.4	0.3	0.5	1.0	1.9	4.1
1999.....	*	*	0.1	*	0.3	0.3	0.6	0.8	1.5	3.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

All races 3/, both sexes

2004.....	1.6	2.0	2.4	3.0	4.2	5.7	8.6	13.2	22.8	52.9
2003.....	1.6	2.0	2.6	3.0	4.1	5.8	9.1	13.6	23.0	56.7
2002.....	1.7	2.2	2.6	3.0	4.3	6.0	9.4	13.9	24.7	56.8
2001.....	1.8	2.2	2.4	3.0	4.3	6.1	8.7	15.0	24.5	58.5
2000.....	1.6	2.1	2.5	3.1	4.3	6.4	8.8	14.1	25.9	58.9
1999.....	1.6	2.1	2.1	2.9	3.9	5.4	9.3	14.5	23.7	60.3

All races 3/, male

2004.....	1.7	2.1	2.4	3.3	4.4	6.0	8.9	13.6	24.6	52.1
2003.....	1.8	2.2	2.5	2.8	4.2	6.1	10.1	14.8	24.9	55.1
2002.....	1.6	2.1	2.5	3.1	4.5	5.5	10.2	14.8	25.1	57.2
2001.....	1.6	2.1	2.4	2.7	4.1	5.9	8.6	15.3	27.3	58.1
2000.....	1.5	2.2	2.4	2.9	4.2	6.5	9.8	14.2	27.3	60.4
1999.....	1.6	2.2	2.1	2.7	3.6	5.4	9.9	15.7	25.4	63.2

All races 3/, female

2004.....	1.6	1.9	2.4	2.7	4.1	5.5	8.3	13.0	21.6	53.3
2003.....	1.5	1.9	2.6	3.2	4.0	5.6	8.4	12.8	21.9	57.5
2002.....	1.8	2.3	2.6	3.0	4.1	6.4	8.8	13.3	24.4	56.6
2001.....	2.0	2.2	2.5	3.4	4.4	6.2	8.8	14.8	22.8	58.7
2000.....	1.6	1.9	2.6	3.3	4.4	6.3	8.1	14.1	25.0	58.3
1999.....	1.7	2.0	2.2	3.0	4.1	5.4	8.8	13.6	22.8	59.2

White, both sexes

2004.....	0.9	1.2	1.5	2.1	3.4	5.1	8.0	13.0	22.7	52.6
2003.....	0.8	1.2	1.5	2.0	3.3	5.0	8.6	13.0	22.7	56.4
2002.....	0.8	1.3	1.6	2.3	3.5	5.3	8.8	13.1	24.3	56.2
2001.....	0.9	1.1	1.5	2.3	3.6	5.5	8.1	14.5	23.8	58.3
2000.....	0.8	1.1	1.6	2.4	3.5	5.7	8.3	13.7	25.5	58.8
1999.....	0.8	1.2	1.4	2.0	3.2	4.7	8.7	14.0	23.5	59.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

White, male

2004.....	1.0	1.4	1.7	2.3	3.8	5.4	8.5	13.3	24.4	52.5
2003.....	0.9	1.4	1.5	1.9	3.4	5.4	9.5	14.1	25.0	55.1
2002.....	0.8	1.4	1.6	2.4	3.8	5.0	9.5	14.2	24.5	56.3
2001.....	0.9	1.2	1.6	2.1	3.6	5.4	7.9	14.9	27.0	58.9
2000.....	0.9	1.3	1.6	2.4	3.5	6.2	9.2	13.8	26.8	59.7
1999.....	0.9	1.4	1.4	1.9	3.0	4.9	9.2	15.2	24.8	63.6

White, female

2004.....	0.8	1.0	1.3	1.8	3.1	4.8	7.6	12.7	21.7	52.6
2003.....	0.7	1.0	1.5	2.2	3.2	4.6	7.9	12.2	21.2	57.0
2002.....	0.8	1.3	1.6	2.2	3.2	5.6	8.2	12.3	24.2	56.2
2001.....	0.9	1.0	1.5	2.5	3.7	5.5	8.3	14.3	21.9	58.0
2000.....	0.7	1.0	1.6	2.3	3.6	5.3	7.6	13.6	24.6	58.4
1999.....	0.7	1.0	1.3	2.0	3.5	4.6	8.2	13.1	22.7	58.0

Black, both sexes

2004.....	6.8	7.8	9.7	11.4	12.6	13.0	15.3	19.8	25.5	65.7
2003.....	7.6	8.5	10.8	12.0	12.0	13.8	16.5	22.0	32.2	69.4
2002.....	7.6	9.0	10.7	10.5	12.1	13.4	16.9	25.6	31.9	71.9
2001.....	8.5	10.2	10.0	9.9	10.1	12.3	16.8	22.9	35.8	64.8
2000.....	7.0	9.2	10.5	10.3	11.8	13.0	14.7	21.0	35.9	66.6
1999.....	7.4	8.3	8.7	11.3	10.1	12.2	15.6	22.3	29.2	71.1

Black, male

2004.....	6.9	7.7	9.0	13.2	11.8	14.5	15.2	20.9	26.5	61.6
2003.....	8.1	9.3	10.9	12.1	12.5	12.3	20.0	26.5	30.3	66.7
2002.....	7.0	8.1	11.1	10.4	12.1	12.2	19.4	24.8	35.9	78.3
2001.....	7.6	9.6	9.7	8.6	9.6	11.2	18.2	24.6	36.1	52.2
2000.....	6.1	9.4	10.6	8.1	11.8	10.9	15.7	19.8	36.2	73.8
1999.....	6.8	8.2	8.3	10.2	10.4	11.7	16.6	23.3	34.9	63.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

Black, female

2004.....	6.8	7.8	10.3	9.9	13.3	11.9	15.4	19.1	24.9	67.3
2003.....	7.2	7.9	10.8	11.9	11.6	15.0	14.2	19.3	33.3	70.5
2002.....	8.0	9.8	10.4	10.5	12.1	14.2	15.2	26.1	29.8	69.4
2001.....	9.3	10.8	10.1	11.0	10.6	13.1	15.9	21.9	35.6	69.5
2000.....	7.7	9.0	10.5	12.1	11.8	14.6	14.0	21.6	35.7	64.0
1999.....	7.9	8.4	9.1	12.2	9.8	12.5	14.9	21.7	26.4	74.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Anemias (D50-D64)

All races 3/, both sexes

2004.....	0.5	0.5	0.6	0.8	1.3	1.8	3.4	5.9	12.2	38.1
2003.....	0.4	0.6	0.5	0.8	1.3	1.9	3.5	6.0	12.7	40.7
2002.....	0.4	0.6	0.6	0.8	1.2	1.9	3.7	6.4	12.9	40.3
2001.....	0.6	0.5	0.7	0.8	1.2	2.0	3.5	7.1	13.6	42.0
2000.....	0.5	0.5	0.7	0.9	1.2	2.2	3.4	6.5	13.9	42.4
1999.....	0.5	0.7	0.6	0.8	1.2	1.8	3.5	6.8	13.7	43.8

All races 3/, male

2004.....	0.5	0.5	0.6	1.0	1.4	1.9	3.6	6.3	12.8	36.5
2003.....	0.5	0.7	0.5	0.9	1.5	2.3	3.8	6.6	13.3	38.0
2002.....	0.4	0.5	0.6	0.8	1.3	1.7	4.3	6.6	12.4	38.3
2001.....	0.4	0.6	0.6	0.8	1.1	2.1	3.8	7.2	14.3	40.0
2000.....	0.4	0.5	0.7	1.0	1.2	2.2	3.6	6.5	14.6	40.1
1999.....	0.5	0.7	0.6	0.7	1.2	2.1	3.4	7.2	14.8	43.4

All races 3/, female

2004.....	0.5	0.6	0.6	0.7	1.3	1.7	3.3	5.7	11.8	38.8
2003.....	0.4	0.5	0.6	0.7	1.0	1.5	3.2	5.6	12.3	41.9
2002.....	0.5	0.7	0.7	0.8	1.1	2.1	3.2	6.2	13.2	41.2
2001.....	0.7	0.5	0.7	0.9	1.2	1.9	3.3	7.0	13.2	42.9
2000.....	0.5	0.6	0.7	0.9	1.2	2.2	3.3	6.5	13.6	43.3
1999.....	0.4	0.7	0.6	0.9	1.1	1.6	3.6	6.5	13.0	44.0

White, both sexes

2004.....	0.2	0.3	0.3	0.5	1.0	1.7	3.2	5.6	12.2	37.7
2003.....	0.1	0.3	0.3	0.4	0.9	1.6	3.2	5.7	12.4	40.6
2002.....	0.1	0.3	0.3	0.6	0.9	1.7	3.4	5.9	12.6	39.7
2001.....	0.2	0.2	0.4	0.6	1.0	1.8	3.2	6.7	13.2	41.7
2000.....	0.1	0.2	0.4	0.7	0.9	1.9	3.2	6.2	13.8	42.0
1999.....	0.1	0.3	0.3	0.4	0.8	1.6	3.2	6.5	13.6	43.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Anemias (D50-D64)

White, male

2004.....	0.2	0.3	0.3	0.5	1.2	1.7	3.5	5.7	12.7	36.7
2003.....	*	0.3	0.3	0.5	1.1	2.0	3.3	6.3	13.4	37.9
2002.....	*	0.3	0.3	0.6	1.0	1.6	3.7	6.3	11.9	37.2
2001.....	*	*	0.4	0.5	1.0	1.9	3.3	7.0	14.1	40.4
2000.....	*	*	0.4	0.8	0.9	2.1	3.4	6.1	14.2	39.1
1999.....	*	0.4	0.3	*	0.7	1.9	3.1	7.0	14.8	43.4

White, female

2004.....	*	0.2	0.3	0.5	0.9	1.6	3.0	5.6	11.9	38.2
2003.....	*	*	0.3	0.3	0.8	1.3	3.1	5.3	11.8	41.8
2002.....	*	0.3	0.3	0.5	0.8	1.8	3.1	5.6	13.0	40.8
2001.....	0.3	*	0.4	0.6	1.1	1.6	3.1	6.5	12.6	42.3
2000.....	*	0.3	0.4	0.7	0.9	1.8	3.0	6.3	13.5	43.2
1999.....	*	0.3	0.3	0.5	0.9	1.4	3.4	6.1	12.9	42.9

Black, both sexes

2004.....	2.4	2.5	3.0	3.9	4.2	3.7	6.2	10.7	13.4	48.2
2003.....	2.5	3.3	2.7	4.1	4.4	4.3	7.1	10.9	19.8	49.3
2002.....	2.4	3.1	3.1	3.2	4.1	4.4	7.9	13.7	19.4	52.3
2001.....	3.2	3.2	2.9	3.2	2.6	3.9	7.4	12.5	21.4	48.5
2000.....	2.7	2.9	3.2	3.0	3.9	4.8	6.1	10.3	18.1	51.0
1999.....	2.8	3.2	2.9	4.2	4.3	3.6	6.6	11.4	16.5	52.9

Black, male

2004.....	2.5	2.1	2.9	5.3	4.5	*	*	13.6	14.3	42.7
2003.....	2.9	3.9	2.6	4.9	6.2	5.7	10.8	12.8	18.5	49.0
2002.....	2.3	2.6	2.6	3.0	5.1	*	11.7	12.9	23.7	56.8
2001.....	2.7	3.9	3.2	3.3	*	*	10.4	12.1	20.1	42.0
2000.....	2.2	3.2	3.4	*	*	*	*	10.9	19.4	55.4
1999.....	2.9	3.3	3.2	4.2	6.2	*	7.8	10.9	*	44.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Anemias (D50-D64)

Black, female

2004.....	2.4	2.8	3.1	2.7	3.9	*	6.2	8.9	12.9	50.4
2003.....	2.1	2.8	2.7	3.3	3.0	*	4.7	9.8	20.5	49.4
2002.....	2.6	3.6	3.5	3.3	3.3	4.6	5.4	14.3	17.1	50.5
2001.....	3.7	2.6	2.6	3.2	*	3.9	5.4	12.8	22.1	50.9
2000.....	3.2	2.7	3.0	3.0	3.9	5.8	6.2	10.0	17.4	49.4
1999.....	2.6	3.2	2.7	4.2	*	*	5.9	11.7	16.4	56.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nutritional anemias (D50-D53)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	0.4	0.7	2.8
2003.....	*	*	*	*	*	*	0.3	0.4	0.7	3.5
2002.....	*	*	*	*	*	*	*	0.3	0.5	3.5
2001.....	*	*	*	*	*	*	*	0.5	0.8	3.6
2000.....	*	*	*	*	*	*	*	0.4	0.9	4.3
1999.....	*	*	*	*	*	*	0.2	0.4	1.1	4.0

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	2.7
2003.....	*	*	*	*	*	*	*	*	*	2.8
2002.....	*	*	*	*	*	*	*	*	*	2.8
2001.....	*	*	*	*	*	*	*	*	1.0	3.2
2000.....	*	*	*	*	*	*	*	*	*	5.1
1999.....	*	*	*	*	*	*	*	*	*	3.9

All races 3/, female

2004.....	*	*	*	*	*	*	*	0.5	0.7	2.8
2003.....	*	*	*	*	*	*	*	*	0.8	3.8
2002.....	*	*	*	*	*	*	*	*	0.7	3.8
2001.....	*	*	*	*	*	*	*	0.5	0.7	3.8
2000.....	*	*	*	*	*	*	*	*	1.2	4.0
1999.....	*	*	*	*	*	*	*	*	1.2	4.0

White, both sexes

2004.....	*	*	*	*	*	*	*	0.4	0.8	2.7
2003.....	*	*	*	*	*	*	*	0.4	0.7	3.6
2002.....	*	*	*	*	*	*	*	0.3	0.5	3.6
2001.....	*	*	*	*	*	*	*	0.5	0.8	3.7
2000.....	*	*	*	*	*	*	*	0.3	0.8	4.4
1999.....	*	*	*	*	*	*	*	0.3	1.1	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hemolytic anemias (D55-D59)

All races 3/, both sexes

2004.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.8	1.9
2003.....	0.3	0.3	0.3	0.3	0.2	*	0.3	0.6	1.1	1.9
2002.....	0.3	0.4	0.3	0.2	0.2	0.4	0.4	0.7	1.0	2.2
2001.....	0.4	0.3	0.3	0.3	0.2	0.3	0.4	0.7	1.2	2.2
2000.....	0.3	0.3	0.4	0.3	0.2	0.3	0.4	0.6	1.2	2.4
1999.....	0.3	0.4	0.2	0.3	0.2	0.3	0.4	0.7	1.2	2.2

All races 3/, male

2004.....	0.3	0.2	0.3	0.3	*	*	*	0.8	1.0	2.1
2003.....	0.3	0.4	0.2	0.3	*	*	*	0.7	1.1	1.5
2002.....	0.3	0.3	0.3	*	*	*	*	*	*	2.2
2001.....	0.3	0.4	0.3	*	*	*	*	0.8	1.4	2.2
2000.....	0.3	0.3	0.3	*	*	*	*	*	1.2	2.5
1999.....	0.4	0.4	0.2	*	*	*	*	0.8	1.4	1.8

All races 3/, female

2004.....	0.3	0.3	0.4	*	0.3	0.4	*	*	0.6	1.8
2003.....	0.3	0.3	0.3	*	*	*	*	0.5	1.1	2.1
2002.....	0.3	0.4	0.4	*	*	0.4	0.4	0.8	1.1	2.2
2001.....	0.5	0.3	0.3	0.3	*	*	*	0.7	1.1	2.2
2000.....	0.4	0.3	0.4	0.3	*	*	0.4	0.7	1.2	2.3
1999.....	0.3	0.4	*	0.4	*	*	0.5	0.7	1.0	2.4

White, both sexes

2004.....	*	*	*	*	*	*	*	0.5	0.9	1.9
2003.....	*	*	*	*	*	*	0.3	0.6	1.2	2.0
2002.....	*	*	*	*	*	0.3	0.3	0.6	1.1	2.3
2001.....	*	*	*	*	*	0.3	0.3	0.7	1.2	2.2
2000.....	*	*	*	*	*	*	0.3	0.7	1.3	2.5
1999.....	*	*	*	*	*	0.2	0.4	0.8	1.2	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hemolytic anemias (D55-D59)

Black, female

2004.....	2.0	2.1	2.6	*	*	*	*	*	*	*
2003.....	1.9	2.3	2.0	*	*	*	*	*	*	*
2002.....	2.3	3.0	2.6	*	*	*	*	*	*	*
2001.....	3.1	2.3	1.9	*	*	*	*	*	*	*
2000.....	2.6	2.1	2.5	*	*	*	*	*	*	*
1999.....	2.0	2.6	*	2.8	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Aplastic anemias (D60-D61)

All races 3/, both sexes

2004.....	0.1	0.1	0.1	0.3	0.5	0.6	1.2	1.5	2.6	5.6
2003.....	*	0.1	0.1	0.2	0.4	0.7	1.1	1.7	2.8	5.7
2002.....	*	0.1	0.1	0.2	0.4	0.6	1.1	1.8	3.2	5.3
2001.....	0.1	0.1	0.2	0.3	0.5	0.6	1.2	2.1	3.5	6.8
2000.....	*	0.1	0.2	0.3	0.5	0.8	1.1	1.8	3.6	7.4
1999.....	*	0.2	0.1	0.2	0.5	0.7	1.3	2.0	3.5	7.7

All races 3/, male

2004.....	*	*	*	*	0.5	0.7	1.4	1.7	3.2	6.2
2003.....	*	*	*	*	0.4	1.0	1.3	2.0	3.4	7.4
2002.....	*	*	*	*	0.5	*	1.3	2.3	3.7	7.0
2001.....	*	*	0.2	*	0.5	0.6	1.1	2.3	3.9	7.8
2000.....	*	*	*	0.3	0.5	1.0	1.2	1.8	4.4	7.6
1999.....	*	*	*	*	0.5	1.0	1.2	2.4	4.3	10.1

All races 3/, female

2004.....	*	*	*	0.3	0.5	0.6	1.0	1.4	2.2	5.4
2003.....	*	*	*	*	0.4	0.6	0.9	1.5	2.5	5.0
2002.....	*	*	*	0.3	*	0.8	0.9	1.5	2.8	4.5
2001.....	*	*	*	0.4	0.5	0.6	1.3	1.9	3.3	6.3
2000.....	*	*	*	*	0.5	0.7	1.0	1.8	3.1	7.3
1999.....	*	*	*	*	0.4	0.4	1.3	1.8	3.0	6.8

White, both sexes

2004.....	*	0.1	*	0.3	0.5	0.6	1.1	1.5	2.7	5.6
2003.....	*	0.1	*	0.2	0.4	0.7	1.0	1.8	2.8	5.8
2002.....	*	0.1	*	0.2	0.3	0.6	1.0	1.9	3.2	5.2
2001.....	*	0.1	0.2	0.3	0.5	0.6	1.2	2.2	3.6	7.1
2000.....	*	0.1	0.1	0.3	0.5	0.8	1.1	1.8	3.7	7.4
1999.....	*	0.1	0.1	*	0.4	0.7	1.2	2.0	3.6	7.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute posthemorrhagic and other anemias (D62,D64)

All races 3/, both sexes

2004.....	*	0.1	0.2	0.3	0.5	0.9	1.8	3.5	8.1	27.8
2003.....	*	0.1	0.2	0.3	0.6	0.9	1.9	3.3	8.1	29.6
2002.....	*	0.1	0.1	0.3	0.6	0.8	2.0	3.6	8.3	29.4
2001.....	*	*	0.1	0.2	0.4	0.9	1.7	3.8	8.1	29.5
2000.....	*	*	0.1	0.3	0.5	1.0	1.8	3.6	8.3	28.2
1999.....	*	0.1	0.2	0.2	0.4	0.8	1.6	3.6	7.9	29.9

All races 3/, male

2004.....	*	*	*	0.4	0.6	1.0	1.9	3.5	7.9	25.4
2003.....	*	*	*	0.4	0.8	1.1	1.9	3.2	8.2	26.2
2002.....	*	*	*	0.4	0.6	0.9	2.4	3.6	7.8	26.3
2001.....	*	*	*	*	*	1.0	2.1	3.7	8.0	26.9
2000.....	*	*	*	0.4	0.6	0.9	1.9	3.6	8.6	24.9
1999.....	*	*	*	*	0.4	0.8	1.7	3.5	8.0	27.7

All races 3/, female

2004.....	*	*	*	*	0.4	0.8	1.7	3.4	8.3	28.9
2003.....	*	*	*	*	0.4	0.7	1.8	3.4	8.0	31.1
2002.....	*	*	*	0.3	0.5	0.7	1.8	3.6	8.5	30.7
2001.....	*	*	*	*	0.4	0.9	1.4	3.9	8.1	30.5
2000.....	*	*	*	0.3	0.4	1.1	1.7	3.6	8.1	29.6
1999.....	*	*	*	*	0.5	0.8	1.6	3.7	7.8	30.8

White, both sexes

2004.....	*	*	*	0.2	0.4	0.8	1.7	3.2	8.0	27.4
2003.....	*	*	*	0.2	0.4	0.7	1.7	3.0	7.8	29.2
2002.....	*	*	*	0.3	0.4	0.7	1.9	3.0	7.8	28.7
2001.....	*	*	*	0.2	0.3	0.8	1.5	3.4	7.6	28.8
2000.....	*	*	*	0.3	0.3	0.8	1.6	3.4	8.0	27.8
1999.....	*	*	*	*	0.3	0.7	1.4	3.4	7.8	28.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute posthemorrhagic and other anemias (D62,D64)

Black, female

2004.....	*	*	*	*	*	*	*	6.7	11.3	39.4
2003.....	*	*	*	*	*	*	*	7.1	13.2	40.5
2002.....	*	*	*	*	*	*	*	11.3	13.9	40.3
2001.....	*	*	*	*	*	*	*	9.7	16.4	42.5
2000.....	*	*	*	*	*	*	*	7.7	12.7	38.6
1999.....	*	*	*	*	*	*	*	7.7	10.8	42.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)										
All races 3/, both sexes										
2004.....	0.4	0.6	0.8	1.0	1.5	2.0	3.0	4.9	7.5	10.1
2003.....	0.4	0.6	0.9	1.0	1.3	2.3	3.3	5.0	7.0	10.5
2002.....	0.4	0.7	0.7	0.9	1.4	2.2	3.5	4.8	7.7	11.3
2001.....	0.5	0.6	0.7	1.1	1.5	2.3	3.1	4.9	7.6	11.1
2000.....	0.4	0.5	0.8	1.1	1.6	2.6	3.3	4.8	7.9	11.6
1999.....	0.4	0.6	0.7	1.0	1.5	2.2	3.7	5.4	6.9	11.5
All races 3/, male										
2004.....	0.4	0.6	0.8	1.2	1.6	2.2	3.5	5.3	8.5	10.9
2003.....	0.4	0.6	0.9	1.1	1.3	2.3	3.8	5.5	7.9	11.9
2002.....	0.4	0.6	0.7	0.9	1.6	2.1	3.5	5.6	8.4	13.9
2001.....	0.4	0.6	0.7	1.0	1.6	2.3	3.3	4.8	9.6	12.3
2000.....	0.4	0.7	0.8	1.0	1.5	2.7	3.8	4.9	8.3	14.6
1999.....	0.4	0.6	0.8	1.0	1.3	2.2	4.1	5.8	6.8	14.3
All races 3/, female										
2004.....	0.4	0.5	0.8	0.8	1.3	1.9	2.6	4.7	6.8	9.7
2003.....	0.3	0.6	0.9	1.0	1.3	2.2	2.9	4.7	6.4	9.9
2002.....	0.4	0.8	0.7	0.9	1.2	2.3	3.6	4.3	7.2	10.1
2001.....	0.5	0.6	0.7	1.1	1.4	2.2	3.0	5.0	6.4	10.5
2000.....	0.3	0.4	0.8	1.3	1.7	2.5	2.9	4.7	7.7	10.4
1999.....	0.5	0.5	0.6	1.0	1.8	2.2	3.3	5.1	6.9	10.3
White, both sexes										
2004.....	0.3	0.5	0.7	0.8	1.4	1.9	2.9	4.9	7.5	10.0
2003.....	0.3	0.5	0.7	0.9	1.2	2.1	3.3	4.8	6.9	10.3
2002.....	0.4	0.6	0.6	0.8	1.3	2.0	3.5	4.6	7.8	11.3
2001.....	0.4	0.5	0.6	1.0	1.3	2.2	3.1	4.8	7.4	11.0
2000.....	0.3	0.5	0.6	1.0	1.5	2.5	3.1	4.7	7.9	11.8
1999.....	0.4	0.5	0.6	0.8	1.5	1.9	3.5	5.2	6.7	11.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)

White, male

2004.....	0.3	0.6	0.7	1.1	1.5	2.0	3.4	5.3	8.5	11.0
2003.....	0.4	0.6	0.7	0.9	1.2	2.1	3.8	5.3	8.0	11.8
2002.....	0.4	0.5	0.6	0.8	1.5	1.9	3.5	5.4	8.2	14.1
2001.....	0.4	0.4	0.6	0.9	1.4	2.2	3.1	4.6	9.6	12.5
2000.....	0.3	0.6	0.6	0.9	1.4	2.5	3.6	4.9	8.3	14.8
1999.....	0.4	0.5	0.7	0.8	1.2	1.9	3.9	5.4	6.3	14.6

White, female

2004.....	0.3	0.5	0.7	0.6	1.3	1.8	2.4	4.7	6.9	9.5
2003.....	0.3	0.4	0.8	0.9	1.2	2.2	2.9	4.4	6.3	9.7
2002.....	0.4	0.6	0.6	0.8	1.1	2.1	3.4	4.0	7.5	10.1
2001.....	0.4	0.5	0.6	1.1	1.2	2.2	3.0	5.0	6.1	10.4
2000.....	0.2	0.4	0.7	1.1	1.6	2.4	2.8	4.5	7.6	10.5
1999.....	0.3	0.4	0.5	0.7	1.8	2.0	3.1	5.0	7.0	10.2

Black, both sexes

2004.....	1.1	1.1	2.0	2.3	2.4	3.4	4.7	6.1	7.4	13.0
2003.....	0.8	1.5	2.1	2.2	2.3	3.9	4.8	7.5	7.9	13.7
2002.....	*	1.4	1.8	1.9	2.3	4.5	4.8	7.9	7.3	13.1
2001.....	1.0	1.9	1.8	1.6	2.9	3.0	4.2	6.4	10.3	12.3
2000.....	0.8	1.1	2.5	2.4	2.8	3.6	5.3	6.9	10.2	11.3
1999.....	1.1	1.1	1.8	2.8	2.3	4.6	5.4	8.2	8.7	13.7

Black, male

2004.....	*	*	2.0	*	*	*	*	*	*	*
2003.....	*	*	2.3	*	*	*	*	*	*	*
2002.....	*	*	2.3	*	*	*	*	*	*	*
2001.....	*	1.9	*	*	*	*	*	*	*	*
2000.....	*	*	2.6	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	5.5	*	10.0	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)

Black, female

2004.....	1.3	*	2.1	*	*	*	4.4	5.9	*	12.6
2003.....	*	1.7	1.9	*	*	*	*	7.1	8.1	13.8
2002.....	*	1.8	*	*	*	4.6	5.4	8.0	*	12.3
2001.....	*	1.8	*	*	*	*	*	5.8	9.8	13.9
2000.....	*	*	2.4	3.0	*	*	4.7	7.7	11.0	10.3
1999.....	*	*	*	2.9	*	4.0	5.4	7.2	*	13.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of blood and blood-forming organs (D70-D76)

All races 3/, both sexes

2004.....	0.1	0.1	0.2	0.3	0.4	0.6	1.1	1.3	1.9	3.0
2003.....	0.1	0.1	0.3	0.3	0.5	0.6	1.0	1.4	2.0	3.9
2002.....	0.1	0.1	0.3	0.4	0.5	0.7	1.1	1.6	2.6	3.4
2001.....	0.1	0.1	0.2	0.3	0.5	0.8	0.9	1.8	2.0	3.9
2000.....	0.1	0.2	0.3	0.3	0.5	0.8	1.0	1.6	2.2	3.4
1999.....	0.1	0.1	0.2	0.4	0.4	0.5	1.2	1.3	2.2	3.6

All races 3/, male

2004.....	*	*	0.3	0.3	0.5	0.7	1.2	1.5	2.3	3.2
2003.....	*	*	0.3	*	0.5	0.6	1.2	1.7	2.4	4.0
2002.....	*	*	0.3	0.4	0.6	0.8	1.4	1.8	2.7	3.5
2001.....	*	*	0.3	0.3	0.6	0.8	0.8	2.2	2.2	3.9
2000.....	*	0.2	*	*	0.5	1.0	1.2	1.8	2.7	4.1
1999.....	*	*	*	0.4	0.5	0.5	1.5	1.8	2.7	4.0

All races 3/, female

2004.....	*	*	*	*	0.3	0.5	1.0	1.1	1.7	2.9
2003.....	*	*	0.2	0.3	0.5	0.6	0.8	1.2	1.8	3.9
2002.....	*	*	0.2	0.3	0.5	0.6	0.8	1.4	2.5	3.3
2001.....	*	*	0.2	0.3	0.5	0.7	0.9	1.5	1.9	3.8
2000.....	*	*	0.3	*	0.6	0.7	0.8	1.4	1.9	3.1
1999.....	*	*	0.3	0.4	0.4	0.6	0.9	1.0	1.8	3.4

White, both sexes

2004.....	*	0.1	0.2	0.2	0.4	0.6	1.1	1.4	2.0	3.2
2003.....	*	0.1	0.2	0.2	0.6	0.6	1.1	1.4	2.1	4.1
2002.....	0.1	0.1	0.3	0.4	0.6	0.7	1.1	1.6	2.6	3.4
2001.....	0.1	0.1	0.2	0.3	0.5	0.7	0.9	1.9	2.0	3.9
2000.....	*	0.1	0.2	0.3	0.5	0.8	1.0	1.6	2.2	3.5
1999.....	0.1	*	0.2	0.4	0.4	0.5	1.2	1.4	2.2	3.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Certain disorders involving the immune mechanism (D80-D89)

All races 3/, both sexes

2004.....	0.6	0.7	0.8	0.9	1.0	1.3	1.1	1.1	1.2	1.7
2003.....	0.7	0.7	0.9	1.0	1.0	1.1	1.3	1.2	1.3	1.6
2002.....	0.7	0.8	1.0	1.0	1.1	1.1	1.0	1.1	1.5	1.8
2001.....	0.7	0.9	0.8	0.8	1.1	1.1	1.2	1.2	1.3	1.5
2000.....	0.6	0.8	0.8	0.8	0.9	0.8	1.1	1.3	1.8	1.6
1999.....	0.6	0.7	0.6	0.7	0.7	0.9	0.9	1.0	1.0	1.4

All races 3/, male

2004.....	0.7	0.8	0.8	0.8	0.8	1.2	0.7	0.6	1.0	1.5
2003.....	0.8	0.7	0.8	0.7	0.8	0.9	1.3	1.1	1.2	*
2002.....	0.7	0.9	0.9	1.0	1.0	0.9	1.0	0.8	1.5	1.5
2001.....	0.7	0.8	0.8	0.6	0.9	0.7	0.8	1.1	1.2	1.8
2000.....	0.6	0.8	0.7	0.7	0.9	0.7	1.1	1.0	1.7	1.6
1999.....	0.5	0.8	0.6	0.6	0.7	0.7	0.8	1.0	*	*

All races 3/, female

2004.....	0.5	0.6	0.8	1.0	1.2	1.3	1.5	1.5	1.2	1.9
2003.....	0.7	0.7	0.9	1.2	1.2	1.2	1.4	1.3	1.3	1.7
2002.....	0.8	0.7	1.0	1.0	1.2	1.4	1.1	1.3	1.5	1.9
2001.....	0.7	1.0	0.9	1.0	1.3	1.4	1.6	1.3	1.4	1.4
2000.....	0.6	0.8	0.8	0.9	0.9	0.9	1.0	1.5	1.9	1.5
1999.....	0.6	0.7	0.6	0.8	0.8	1.0	1.0	1.1	1.0	1.5

White, both sexes

2004.....	0.3	0.3	0.4	0.5	0.6	0.9	0.9	1.0	1.0	1.8
2003.....	0.2	0.3	0.3	0.5	0.6	0.7	1.1	1.1	1.2	1.5
2002.....	0.2	0.3	0.4	0.6	0.7	0.8	0.9	1.1	1.4	1.8
2001.....	0.2	0.3	0.4	0.4	0.8	0.8	0.9	1.1	1.2	1.6
2000.....	0.3	0.3	0.3	0.4	0.6	0.5	1.0	1.2	1.6	1.5
1999.....	0.2	0.3	0.3	0.4	0.5	0.7	0.7	1.0	0.9	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Certain disorders involving the immune mechanism (D80-D89)

White, male

2004.....	0.3	0.4	0.4	0.5	0.6	0.9	*	*	*	1.5
2003.....	0.3	0.4	0.3	0.3	0.6	0.8	1.2	0.9	1.1	*
2002.....	0.2	0.4	0.4	0.6	0.8	0.6	1.0	0.8	1.6	*
2001.....	0.2	0.4	0.4	*	0.7	0.6	0.6	0.9	1.1	2.0
2000.....	0.3	0.3	0.4	0.5	0.7	0.6	1.1	1.0	1.5	*
1999.....	0.3	0.4	0.3	0.4	0.6	0.6	0.6	1.0	*	*

White, female

2004.....	*	*	0.3	0.5	0.6	0.9	1.2	1.2	1.0	1.9
2003.....	*	0.3	0.3	0.6	0.7	0.6	1.0	1.3	1.2	1.7
2002.....	*	0.2	0.4	0.5	0.6	1.0	0.8	1.2	1.3	1.9
2001.....	*	0.3	0.3	0.5	0.9	1.0	1.1	1.2	1.3	1.5
2000.....	0.2	*	0.3	*	0.5	0.5	0.9	1.3	1.6	1.5
1999.....	*	*	*	0.4	0.5	0.7	0.8	1.0	0.9	1.4

Black, both sexes

2004.....	3.2	3.9	4.5	4.8	5.3	5.0	3.8	*	*	*
2003.....	4.2	3.5	5.8	5.4	4.8	4.9	4.1	*	*	*
2002.....	4.4	4.3	5.5	5.0	5.0	4.1	3.1	*	*	*
2001.....	4.0	5.0	4.8	4.8	4.0	4.0	4.6	*	*	*
2000.....	3.3	4.9	4.5	4.5	4.2	3.5	*	*	*	*
1999.....	3.4	3.7	3.6	3.3	2.7	3.1	*	*	*	*

Black, male

2004.....	3.6	3.9	4.1	4.6	3.9	4.8	*	*	*	*
2003.....	4.1	3.8	5.7	4.1	*	*	*	*	*	*
2002.....	4.0	4.3	5.8	4.8	4.0	*	*	*	*	*
2001.....	3.9	3.7	3.9	3.4	*	*	*	*	*	*
2000.....	3.1	4.7	4.3	*	*	*	*	*	*	*
1999.....	2.8	3.5	3.0	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Certain disorders involving the immune mechanism (D80-D89)

Black, female

2004.....	2.8	4.0	4.9	4.9	6.4	5.0	*	*	*	*
2003.....	4.3	3.3	5.9	6.5	5.9	7.1	5.1	*	*	*
2002.....	4.9	4.3	5.3	5.1	5.9	4.8	*	*	*	*
2001.....	4.2	6.2	5.6	5.9	4.6	5.3	5.8	*	*	*
2000.....	3.4	5.1	4.8	5.9	4.4	4.6	*	*	*	*
1999.....	4.0	3.9	4.1	4.0	3.5	4.2	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

IV. Endocrine, nutritional and metabolic diseases (E00-E88)

All races 3/, both sexes

2004.....	8.6	14.6	23.5	37.9	59.1	87.8	130.2	187.8	278.7	470.6
2003.....	8.8	14.8	24.3	39.0	60.7	91.7	132.9	194.0	284.5	492.8
2002.....	9.2	14.0	24.2	39.1	59.2	91.3	133.0	197.1	288.8	504.8
2001.....	8.0	14.0	23.8	37.3	59.1	90.6	132.7	194.8	286.9	519.2
2000.....	8.4	13.3	22.9	37.3	58.9	89.4	131.9	195.7	286.5	521.0
1999.....	8.1	13.3	21.7	36.5	58.8	90.3	132.3	191.6	285.9	529.8

All races 3/, male

2004.....	10.3	17.6	28.6	44.5	70.2	102.7	148.6	213.4	315.6	481.0
2003.....	10.8	17.6	28.7	45.5	71.0	106.9	151.4	220.3	318.9	490.8
2002.....	11.4	16.3	28.7	46.4	68.9	102.1	147.7	217.5	313.8	504.5
2001.....	9.5	16.8	27.9	42.9	67.4	102.8	148.3	215.1	310.2	507.7
2000.....	9.8	16.0	26.4	42.5	66.6	98.5	146.0	215.1	311.8	517.4
1999.....	9.4	16.1	25.2	41.9	65.3	98.8	146.0	213.4	312.5	521.0

All races 3/, female

2004.....	6.9	11.7	18.6	31.7	48.9	74.9	115.5	169.4	255.9	465.9
2003.....	6.9	12.1	20.1	32.7	51.3	78.6	118.1	175.2	263.5	493.7
2002.....	7.0	11.7	19.8	32.2	50.4	81.9	121.2	182.7	273.5	504.9
2001.....	6.5	11.3	19.9	31.9	51.6	80.0	120.3	180.6	272.9	524.0
2000.....	7.0	10.7	19.4	32.4	52.0	81.6	120.7	182.3	271.5	522.5
1999.....	6.8	10.6	18.3	31.4	52.9	83.0	121.6	176.4	270.3	533.3

White, both sexes

2004.....	7.6	13.2	20.7	33.8	52.8	79.4	118.7	174.8	264.7	453.5
2003.....	8.0	13.1	21.4	34.6	54.2	82.9	121.8	181.0	269.3	475.4
2002.....	8.3	12.3	21.2	34.5	52.5	81.0	121.2	183.8	274.3	487.1
2001.....	7.2	12.2	21.0	32.8	52.6	81.3	121.4	181.3	272.1	502.4
2000.....	7.7	11.6	19.9	32.9	52.9	80.1	120.6	181.8	270.5	503.5
1999.....	7.2	11.4	18.7	31.1	51.4	80.8	119.6	178.3	269.9	511.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

IV. Endocrine, nutritional and metabolic diseases (E00-E88)

White, male

2004.....	9.3	16.1	26.1	40.3	64.4	94.5	138.5	202.5	307.0	470.5
2003.....	9.9	15.9	26.0	41.6	64.9	98.9	141.8	210.2	308.1	484.1
2002.....	10.5	14.6	25.9	42.1	62.2	93.2	138.7	208.7	303.9	498.9
2001.....	8.9	14.8	25.1	38.7	61.2	95.1	139.9	204.3	297.8	500.5
2000.....	9.2	14.4	23.8	38.3	60.9	91.0	137.2	204.2	302.4	510.5
1999.....	8.5	14.2	22.5	36.8	58.9	90.4	136.6	203.3	301.4	504.7

White, female

2004.....	5.9	10.3	15.4	27.6	42.1	66.0	102.5	154.5	238.5	445.8
2003.....	6.0	10.2	16.9	27.9	44.4	68.9	105.4	159.9	245.4	471.5
2002.....	6.1	10.1	16.5	27.3	43.6	70.2	107.0	166.0	256.1	482.1
2001.....	5.4	9.6	17.0	27.1	44.6	69.2	106.5	165.0	256.6	503.2
2000.....	6.2	8.9	16.1	27.8	45.5	70.5	107.4	166.0	251.5	500.6
1999.....	5.8	8.7	15.0	25.6	44.5	72.4	106.1	160.7	251.2	514.0

Black, both sexes

2004.....	16.6	27.2	48.6	78.0	120.8	174.2	257.1	351.2	465.4	719.1
2003.....	16.5	29.2	50.1	83.1	123.9	179.3	255.5	354.1	484.1	736.9
2002.....	17.3	28.3	51.2	83.3	125.4	186.0	261.4	356.3	491.9	757.0
2001.....	15.1	28.7	50.2	81.1	121.3	180.4	263.0	363.7	482.9	763.1
2000.....	15.2	28.3	49.9	79.6	117.6	183.3	258.5	375.1	503.2	766.1
1999.....	16.2	28.7	49.8	84.7	127.7	183.9	268.1	349.9	497.3	763.6

Black, male

2004.....	19.2	31.5	54.5	88.4	134.5	199.6	277.1	370.3	476.3	683.5
2003.....	19.3	32.9	55.0	91.3	138.4	197.7	272.0	372.0	497.6	619.3
2002.....	20.6	32.0	56.7	92.7	143.3	192.0	264.0	352.7	492.9	610.0
2001.....	16.2	33.8	55.1	88.9	133.2	189.6	265.6	375.4	511.2	658.1
2000.....	16.5	31.6	53.2	87.3	126.7	180.1	265.1	376.6	480.5	655.0
1999.....	18.0	33.5	53.0	92.9	130.5	193.2	261.9	347.6	485.7	746.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

IV. Endocrine, nutritional and metabolic diseases (E00-E88)

Black, female

2004.....	14.3	23.5	43.6	69.5	110.1	155.2	243.6	339.9	459.6	733.2
2003.....	14.0	25.9	45.9	76.3	112.5	165.6	244.4	343.5	477.0	782.6
2002.....	14.4	25.0	46.5	75.5	111.4	181.5	259.7	358.4	491.4	813.3
2001.....	14.2	24.2	46.0	74.8	112.0	173.7	261.2	356.7	468.5	802.0
2000.....	14.0	25.4	47.1	73.3	110.5	185.7	254.2	374.2	514.6	807.4
1999.....	14.7	24.6	47.2	78.0	125.5	177.0	272.3	351.3	503.0	770.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Endocrine diseases (E00-E34)

All races 3/, both sexes

2004.....	6.0	10.4	17.6	29.9	48.2	72.6	107.7	153.4	220.9	330.4
2003.....	6.1	10.7	18.3	31.2	49.9	76.3	111.0	159.7	225.2	342.7
2002.....	6.3	10.2	18.5	30.7	48.4	76.9	111.3	161.9	226.5	347.6
2001.....	5.6	10.2	18.3	30.1	49.3	76.6	110.8	160.4	227.1	350.3
2000.....	5.8	9.8	17.7	30.3	48.9	75.2	110.7	160.9	222.4	345.2
1999.....	5.7	10.3	17.0	30.2	49.7	77.0	111.6	157.7	223.2	343.6

All races 3/, male

2004.....	7.3	12.7	21.5	34.9	57.5	84.1	122.8	175.5	252.8	349.1
2003.....	7.6	12.9	21.7	36.1	58.4	87.9	126.2	182.8	255.1	354.4
2002.....	7.9	12.1	21.9	36.6	56.4	85.4	122.9	178.9	248.9	365.2
2001.....	6.8	12.3	21.8	34.9	55.3	85.7	123.0	175.6	247.7	362.2
2000.....	7.0	11.6	20.1	34.0	54.5	81.7	121.4	177.0	247.4	360.9
1999.....	6.6	12.5	19.8	34.1	54.6	83.6	121.3	175.9	248.6	361.8

All races 3/, female

2004.....	4.6	8.0	13.9	25.2	39.6	62.6	95.5	137.6	201.2	322.0
2003.....	4.5	8.5	15.1	26.5	42.2	66.2	98.9	143.3	206.9	337.6
2002.....	4.7	8.3	15.3	25.2	41.2	69.6	102.0	149.9	212.9	339.9
2001.....	4.5	8.1	15.0	25.6	44.0	68.7	101.2	149.7	214.7	345.3
2000.....	4.7	8.1	15.4	26.9	43.8	69.5	102.3	149.7	207.7	338.8
1999.....	4.8	8.1	14.3	26.5	45.3	71.3	104.1	145.0	208.4	336.3

White, both sexes

2004.....	5.2	9.2	15.0	25.9	42.2	64.7	96.6	140.8	207.2	313.3
2003.....	5.3	9.1	15.6	27.0	43.6	67.7	100.2	147.6	211.2	326.4
2002.....	5.5	8.7	15.7	26.5	42.2	67.2	100.5	149.5	213.6	331.6
2001.....	5.0	8.8	15.7	25.8	43.1	67.9	100.1	147.7	214.1	335.2
2000.....	5.2	8.3	15.0	26.4	43.1	66.2	100.1	147.8	207.5	329.3
1999.....	4.9	8.6	14.3	25.0	42.6	68.1	99.9	145.2	208.9	325.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Endocrine diseases (E00-E34)										
White, male										
2004.....	6.5	11.3	19.0	30.8	51.7	76.2	113.0	165.2	244.2	338.1
2003.....	6.9	11.3	19.0	32.1	52.1	80.1	117.0	173.8	246.1	347.0
2002.....	7.0	10.5	19.1	32.4	50.2	76.9	114.5	170.9	240.8	361.5
2001.....	6.3	10.7	19.1	30.8	49.6	78.5	115.3	165.5	237.4	356.8
2000.....	6.4	10.0	17.6	30.1	49.3	74.6	113.2	167.2	238.9	353.0
1999.....	5.7	10.8	17.2	29.3	48.4	75.9	113.0	167.1	240.0	349.4
White, female										
2004.....	3.9	7.1	11.1	21.3	33.4	54.4	83.1	123.0	184.2	302.1
2003.....	3.8	6.9	12.2	22.2	35.7	56.8	86.4	128.7	189.6	317.3
2002.....	4.0	6.9	12.5	20.8	34.7	58.7	89.1	134.2	196.9	318.5
2001.....	3.6	6.9	12.5	21.0	37.2	58.5	87.8	135.1	200.0	326.1
2000.....	4.0	6.7	12.4	22.9	37.4	58.8	89.6	134.2	188.8	319.7
1999.....	4.1	6.5	11.4	21.0	37.4	61.3	89.4	129.9	190.5	316.4
Black, both sexes										
2004.....	11.9	20.1	39.8	66.1	104.0	150.5	224.4	304.5	394.4	555.3
2003.....	12.1	23.0	40.6	71.4	108.8	158.0	225.2	304.4	402.1	552.8
2002.....	12.6	22.0	42.1	69.3	107.6	163.3	226.6	305.3	400.6	559.8
2001.....	11.4	21.6	41.1	70.7	106.2	159.8	230.8	312.3	394.1	555.5
2000.....	10.8	22.4	41.4	67.6	103.3	162.5	228.0	325.0	415.6	550.6
1999.....	11.8	23.7	41.5	74.8	113.5	162.3	233.8	301.0	408.8	552.6
Black, male										
2004.....	14.3	25.1	45.0	75.5	118.1	172.0	242.8	315.7	399.6	525.5
2003.....	14.7	26.5	44.6	79.0	122.8	172.1	235.2	312.7	397.9	473.4
2002.....	15.4	25.9	47.9	77.6	121.2	167.3	228.0	299.5	391.3	427.8
2001.....	12.3	26.2	46.3	77.7	113.5	164.4	226.5	317.3	409.3	469.7
2000.....	12.1	25.8	44.2	73.8	108.9	156.9	228.7	317.5	389.5	474.0
1999.....	13.6	28.3	45.1	80.5	115.2	168.2	220.8	288.7	388.0	523.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Endocrine diseases (E00-E34)

Black, female

2004.....	9.8	15.7	35.5	58.3	93.1	134.6	212.0	297.9	391.6	567.0
2003.....	9.9	19.9	37.2	65.2	97.8	147.5	218.4	299.5	404.2	583.7
2002.....	10.1	18.7	37.2	62.4	97.0	160.4	225.6	308.7	405.4	610.4
2001.....	10.6	17.6	36.7	64.9	100.5	156.4	233.6	309.2	386.3	587.3
2000.....	9.7	19.4	39.0	62.5	98.9	166.7	227.5	329.5	428.8	579.1
1999.....	10.3	19.7	38.5	70.1	112.1	158.0	242.4	308.4	419.2	563.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Disorders of thyroid gland (E00-E07)

All races 3/, both sexes

2004.....	0.1	0.2	0.2	0.2	0.5	0.5	1.0	2.2	5.3	19.2
2003.....	0.1	0.2	0.2	0.3	0.4	0.7	1.2	2.9	6.1	21.1
2002.....	0.1	0.1	0.2	0.3	0.4	0.5	1.4	2.5	6.0	22.7
2001.....	0.1	0.1	0.2	0.3	0.4	0.6	1.1	3.0	5.8	23.2
2000.....	0.1	*	0.2	0.3	0.4	0.5	1.3	2.7	5.9	21.1
1999.....	0.1	0.2	0.2	0.3	0.3	0.6	1.3	2.5	6.2	22.4

All races 3/, male

2004.....	*	*	*	*	0.4	0.5	0.8	1.5	3.5	11.2
2003.....	*	*	*	*	*	0.6	0.6	1.9	2.9	10.5
2002.....	*	*	*	*	*	*	0.8	1.6	3.0	12.7
2001.....	*	*	*	*	*	*	0.7	2.3	2.8	14.1
2000.....	*	*	*	*	*	*	0.9	1.5	2.6	12.1
1999.....	*	*	*	*	*	0.5	0.8	1.5	4.4	12.9

All races 3/, female

2004.....	*	0.2	0.2	*	0.5	0.6	1.3	2.7	6.4	22.8
2003.....	*	0.2	0.3	0.4	0.6	0.8	1.7	3.6	8.0	25.8
2002.....	*	0.2	0.3	0.4	0.6	0.6	1.8	3.1	7.8	27.0
2001.....	*	*	0.3	0.4	0.5	0.8	1.4	3.6	7.6	27.1
2000.....	0.2	*	0.3	0.4	0.6	0.7	1.5	3.5	7.8	24.8
1999.....	*	0.3	*	0.4	0.4	0.8	1.7	3.2	7.3	26.3

White, both sexes

2004.....	*	0.1	0.2	0.2	0.4	0.5	0.9	2.0	5.3	19.8
2003.....	*	0.1	0.2	0.3	0.4	0.7	1.2	3.0	6.1	21.7
2002.....	*	*	0.1	0.2	0.4	0.6	1.3	2.5	6.2	23.5
2001.....	*	*	0.2	0.3	0.4	0.6	1.1	3.0	5.5	23.9
2000.....	*	*	0.2	0.3	0.5	0.4	1.2	2.6	5.8	21.6
1999.....	*	0.1	0.2	0.2	0.3	0.6	1.3	2.5	6.1	23.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diabetes mellitus (E10-E14)

All races 3/, both sexes

2004.....	5.7	10.0	17.2	29.4	47.2	71.3	105.7	149.8	213.1	307.0
2003.....	5.8	10.3	17.9	30.4	48.9	75.0	108.8	155.3	216.5	317.5
2002.....	6.0	9.9	18.0	30.1	47.5	75.7	108.8	157.7	217.9	320.6
2001.....	5.4	9.9	17.8	29.5	48.4	75.5	108.6	155.5	218.7	321.8
2000.....	5.5	9.5	17.3	29.6	48.1	74.0	108.6	156.5	213.9	319.7
1999.....	5.5	9.9	16.5	29.5	48.9	75.6	109.2	153.5	214.8	317.2

All races 3/, male

2004.....	7.1	12.4	21.2	34.5	56.6	82.7	121.2	172.2	246.9	333.9
2003.....	7.4	12.6	21.3	35.4	57.8	86.8	124.4	179.0	249.7	340.7
2002.....	7.7	11.9	21.5	36.0	55.5	84.3	121.1	175.9	243.5	348.5
2001.....	6.6	12.1	21.3	34.4	54.4	84.7	121.0	171.5	242.3	343.1
2000.....	6.8	11.4	19.7	33.4	53.9	80.8	119.7	173.7	242.5	345.1
1999.....	6.4	12.3	19.4	33.5	53.9	82.5	119.4	172.3	241.3	345.7

All races 3/, female

2004.....	4.3	7.7	13.4	24.6	38.6	61.4	93.2	133.6	192.2	294.9
2003.....	4.2	8.1	14.7	25.7	40.9	64.8	96.2	138.4	196.2	307.2
2002.....	4.4	7.9	14.7	24.5	40.2	68.2	99.0	144.9	202.3	308.4
2001.....	4.2	7.9	14.5	24.9	42.9	67.6	98.8	144.3	204.5	312.9
2000.....	4.3	7.8	14.9	26.1	42.8	68.1	99.9	144.6	197.0	309.4
1999.....	4.6	7.5	13.8	25.8	44.5	69.7	101.2	140.4	199.3	305.8

White, both sexes

2004.....	5.0	8.9	14.6	25.5	41.3	63.6	94.8	137.4	199.5	289.4
2003.....	5.1	8.8	15.2	26.4	42.7	66.5	98.1	143.1	202.7	300.8
2002.....	5.3	8.5	15.4	26.0	41.3	66.0	98.2	145.4	204.9	303.7
2001.....	4.8	8.6	15.3	25.3	42.3	66.8	98.0	143.0	205.9	306.0
2000.....	5.0	8.0	14.6	25.7	42.3	65.2	98.1	143.7	199.1	303.5
1999.....	4.7	8.3	13.8	24.5	41.9	66.9	97.6	141.2	200.8	298.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diabetes mellitus (E10-E14)

White, male

2004.....	6.3	11.1	18.7	30.3	51.0	75.0	111.6	162.1	238.5	323.0
2003.....	6.6	11.1	18.7	31.6	51.5	79.1	115.3	170.2	241.1	333.4
2002.....	6.9	10.3	18.8	32.0	49.5	75.9	112.9	167.9	235.5	344.3
2001.....	6.1	10.5	18.6	30.4	48.8	77.6	113.5	161.8	232.1	337.3
2000.....	6.3	9.7	17.3	29.5	48.7	73.8	111.6	164.0	234.1	337.4
1999.....	5.6	10.6	16.8	28.9	47.7	75.2	111.3	163.7	233.2	333.4

White, female

2004.....	3.6	6.7	10.6	20.8	32.4	53.4	81.1	119.4	175.3	274.3
2003.....	3.5	6.5	11.8	21.4	34.5	55.4	84.0	123.5	179.1	286.3
2002.....	3.8	6.6	12.0	20.2	33.7	57.3	86.3	129.3	186.1	286.1
2001.....	3.4	6.7	12.0	20.4	36.3	57.4	85.4	129.7	190.1	293.0
2000.....	3.7	6.4	12.0	22.1	36.3	57.6	87.3	129.3	178.3	289.7
1999.....	3.9	6.1	11.0	20.4	36.6	59.6	86.7	125.4	181.7	285.1

Black, both sexes

2004.....	11.5	19.2	38.8	64.8	102.0	147.7	219.3	297.6	384.2	532.4
2003.....	11.7	22.1	39.9	69.8	106.9	156.1	221.2	300.0	389.9	527.5
2002.....	12.0	21.4	40.7	67.8	105.5	161.6	221.6	300.3	392.7	537.8
2001.....	10.7	21.0	40.2	69.6	104.4	157.6	226.8	305.2	382.2	529.7
2000.....	10.1	21.9	40.7	66.3	102.4	159.2	223.7	316.5	404.9	524.9
1999.....	11.3	22.5	40.5	72.7	111.6	159.6	229.0	294.9	395.8	529.0

Black, male

2004.....	13.9	24.2	44.1	74.6	115.7	168.6	238.3	309.8	389.5	505.7
2003.....	14.4	25.4	44.0	77.3	121.7	170.6	232.6	306.3	387.6	453.5
2002.....	14.9	25.4	46.4	76.5	118.8	165.3	223.3	295.8	386.7	417.0
2001.....	11.7	25.5	45.6	76.7	112.5	162.1	221.5	308.1	400.5	450.5
2000.....	11.8	25.5	43.5	72.1	108.5	154.3	224.3	311.4	382.7	452.1
1999.....	13.2	27.5	44.1	78.2	113.3	163.8	216.1	282.6	374.1	506.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diabetes mellitus (E10-E14)

Black, female

2004.....	9.4	14.9	34.4	56.8	91.3	132.2	206.5	290.3	381.4	543.0
2003.....	9.3	19.2	36.4	63.6	95.4	145.3	213.6	296.3	391.1	556.2
2002.....	9.4	17.9	35.8	60.6	95.1	158.9	220.4	303.0	395.9	584.1
2001.....	9.8	17.0	35.6	63.9	98.1	154.3	230.3	303.4	372.8	559.1
2000.....	8.7	18.7	38.4	61.5	97.6	162.8	223.3	319.6	416.1	552.0
1999.....	9.7	18.1	37.4	68.3	110.3	156.4	237.6	302.4	406.6	537.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Disorders of pituitary gland (E22-E23)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	0.3	0.3	0.6	0.9
2003.....	*	*	*	*	*	*	*	0.3	0.6	0.9
2002.....	*	*	*	*	*	*	0.2	0.4	0.5	1.1
2001.....	*	*	*	*	*	*	*	0.4	0.6	1.0
2000.....	*	*	*	*	*	0.2	0.3	0.3	0.7	1.1
1999.....	*	*	*	*	*	0.2	0.2	0.5	0.5	1.1

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	0.7	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	0.7	1.0
2003.....	*	*	*	*	*	*	*	*	*	1.0
2002.....	*	*	*	*	*	*	*	0.5	*	1.1
2001.....	*	*	*	*	*	*	*	*	0.6	1.1
2000.....	*	*	*	*	*	*	*	*	1.0	1.1
1999.....	*	*	*	*	*	*	*	*	*	1.2

White, both sexes

2004.....	*	*	*	*	*	*	0.3	0.3	0.6	1.0
2003.....	*	*	*	*	*	*	*	0.3	0.5	0.9
2002.....	*	*	*	*	*	*	*	0.4	0.4	1.1
2001.....	*	*	*	*	*	*	*	0.4	0.7	1.0
2000.....	*	*	*	*	*	*	0.3	0.3	0.8	1.1
1999.....	*	*	*	*	*	*	*	0.4	0.5	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Disorders of adrenal glands (E24-E27)

All races 3/, both sexes

2004.....	*	*	*	*	0.2	*	0.3	0.4	0.7	1.2
2003.....	*	*	*	0.1	0.2	0.2	0.3	0.6	0.8	1.2
2002.....	*	*	*	*	0.2	0.2	0.3	0.5	0.9	1.1
2001.....	*	*	*	*	*	*	0.5	0.7	1.0	1.8
2000.....	*	*	*	0.2	*	0.2	0.2	0.5	1.0	1.0
1999.....	*	*	0.1	*	*	0.3	0.3	0.5	0.4	1.0

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	0.6	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	0.7	1.2	2.1
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	0.8	1.1
2003.....	*	*	*	*	*	*	*	0.5	0.9	1.3
2002.....	*	*	*	*	*	*	0.4	0.6	0.9	1.1
2001.....	*	*	*	*	*	*	0.5	0.7	0.8	1.7
2000.....	*	*	*	*	*	*	*	0.5	1.0	1.1
1999.....	*	*	*	*	*	*	0.4	0.6	*	1.0

White, both sexes

2004.....	*	*	*	*	*	*	*	0.4	0.8	1.2
2003.....	*	*	*	*	*	*	0.3	0.6	0.8	1.2
2002.....	*	*	*	*	*	*	0.3	0.5	0.9	1.1
2001.....	*	*	*	*	*	*	0.5	0.7	1.0	1.8
2000.....	*	*	*	0.2	*	*	*	0.5	1.0	1.1
1999.....	*	*	0.1	*	*	0.3	0.3	0.6	0.5	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other endocrine diseases (E15-E21,E28-E31,E34)

All races 3/, both sexes

2004.....	*	0.1	0.1	0.2	0.2	0.4	0.4	0.8	1.2	2.1
2003.....	*	*	*	0.2	0.2	0.3	0.5	0.6	1.2	2.1
2002.....	*	0.1	0.2	0.2	0.2	0.3	0.5	0.8	1.3	2.2
2001.....	*	*	0.1	0.1	0.3	0.2	0.5	0.7	1.0	2.5
2000.....	*	0.1	*	*	*	0.2	0.4	0.9	0.9	2.2
1999.....	*	*	0.2	0.2	0.2	0.2	0.6	0.7	1.2	1.9

All races 3/, male

2004.....	*	*	*	*	*	0.5	*	1.0	1.3	1.8
2003.....	*	*	*	*	*	*	0.6	0.7	1.2	1.7
2002.....	*	*	*	*	*	*	*	0.7	1.1	1.8
2001.....	*	*	*	*	*	*	0.6	0.8	*	2.2
2000.....	*	*	*	*	*	*	*	0.9	*	1.7
1999.....	*	*	*	*	*	*	0.7	0.8	1.4	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	0.7	1.1	2.2
2003.....	*	*	*	0.3	*	*	0.4	0.6	1.2	2.2
2002.....	*	*	*	*	*	*	0.6	0.8	1.3	2.3
2001.....	*	*	*	*	*	*	*	0.7	1.2	2.6
2000.....	*	*	*	*	*	*	0.4	0.9	0.9	2.4
1999.....	*	*	*	*	*	*	0.5	0.6	1.1	2.0

White, both sexes

2004.....	*	*	*	0.1	0.2	0.3	0.3	0.8	1.1	1.9
2003.....	*	*	*	*	0.2	*	0.4	0.6	1.0	2.0
2002.....	*	*	0.1	*	0.2	0.3	0.4	0.7	1.2	2.1
2001.....	*	*	*	*	0.2	*	0.3	0.6	0.9	2.4
2000.....	*	*	*	*	*	*	0.3	0.7	0.7	2.0
1999.....	*	*	*	*	*	*	0.5	0.6	1.1	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nutritional diseases (E40-E68)

All races 3/, both sexes

2004.....	1.7	2.5	3.4	3.7	4.3	4.8	5.8	7.7	12.8	32.3
2003.....	1.7	2.5	3.1	3.7	4.1	5.0	5.8	8.3	12.9	37.5
2002.....	1.9	2.2	3.0	3.8	3.9	4.7	5.0	8.7	15.0	42.9
2001.....	1.4	2.1	2.8	3.1	3.5	3.9	5.4	8.1	14.5	44.5
2000.....	1.6	2.0	2.7	2.9	3.6	4.5	5.4	9.1	17.9	51.7
1999.....	1.5	1.7	2.3	2.8	2.9	3.8	5.6	8.6	17.9	55.3

All races 3/, male

2004.....	1.8	2.5	3.5	3.8	4.0	4.7	5.0	7.6	12.3	28.8
2003.....	1.8	2.6	3.1	3.9	3.9	5.1	5.2	7.6	12.8	31.5
2002.....	2.2	2.2	3.1	3.7	3.6	4.3	4.9	8.2	13.5	37.8
2001.....	1.5	2.2	2.6	2.9	3.4	3.4	5.3	6.9	14.4	35.8
2000.....	1.7	2.2	2.8	3.3	3.2	4.0	5.3	8.5	15.8	44.6
1999.....	1.8	1.8	2.3	3.1	3.0	3.5	5.8	8.4	17.1	48.7

All races 3/, female

2004.....	1.6	2.6	3.2	3.6	4.6	4.9	6.4	7.8	13.1	33.9
2003.....	1.7	2.5	3.1	3.6	4.4	5.0	6.3	8.8	12.9	40.1
2002.....	1.6	2.2	2.9	3.9	4.2	5.1	5.2	9.0	15.9	45.1
2001.....	1.3	2.0	3.0	3.3	3.7	4.3	5.6	9.0	14.6	48.1
2000.....	1.6	1.8	2.7	2.6	3.9	4.9	5.5	9.6	19.2	54.6
1999.....	1.3	1.6	2.4	2.4	2.9	4.0	5.5	8.8	18.4	58.0

White, both sexes

2004.....	1.5	2.3	3.2	3.6	4.1	4.5	5.5	7.2	12.4	31.0
2003.....	1.6	2.4	3.0	3.7	4.0	5.1	5.5	8.1	12.3	36.1
2002.....	1.8	2.1	2.9	3.6	3.7	4.4	4.8	8.2	14.4	41.6
2001.....	1.2	1.9	2.6	3.0	3.4	3.6	5.0	7.6	13.6	43.8
2000.....	1.5	1.8	2.6	2.7	3.5	4.3	5.0	8.5	17.4	50.6
1999.....	1.4	1.5	2.2	2.6	2.8	3.6	4.9	8.0	16.8	54.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nutritional diseases (E40-E68)

White, male

2004.....	1.8	2.4	3.5	3.9	3.9	4.3	5.1	7.1	11.9	27.8
2003.....	1.7	2.5	3.1	4.0	3.9	5.3	4.8	7.1	12.1	30.5
2002.....	2.1	2.3	3.2	3.6	3.5	4.1	4.6	8.0	12.5	37.1
2001.....	1.4	2.0	2.5	2.9	3.2	3.1	4.9	6.6	13.5	34.5
2000.....	1.6	2.2	2.8	3.1	3.0	3.8	4.9	7.9	15.3	43.7
1999.....	1.8	1.7	2.3	2.9	2.8	3.3	5.0	7.6	15.8	46.9

White, female

2004.....	1.3	2.2	2.9	3.4	4.3	4.6	5.9	7.4	12.7	32.5
2003.....	1.5	2.3	3.0	3.4	4.1	5.0	6.1	8.8	12.3	38.6
2002.....	1.4	2.0	2.6	3.5	3.8	4.6	5.0	8.3	15.5	43.5
2001.....	1.1	1.7	2.7	3.1	3.6	4.1	5.1	8.3	13.8	47.7
2000.....	1.4	1.4	2.4	2.3	3.9	4.7	5.2	9.0	18.7	53.4
1999.....	1.1	1.4	2.2	2.2	2.7	3.8	4.8	8.3	17.4	57.2

Black, both sexes

2004.....	3.2	4.7	5.7	6.0	7.6	9.3	10.3	13.9	18.6	50.8
2003.....	3.0	4.1	4.8	5.6	6.8	5.6	9.9	13.4	22.8	58.1
2002.....	3.4	3.7	5.2	7.5	7.0	8.7	8.8	16.3	24.9	62.1
2001.....	2.9	4.3	5.5	5.4	5.7	6.8	10.6	15.5	26.5	56.2
2000.....	3.0	3.8	4.9	6.2	5.3	7.5	10.4	16.9	26.5	68.5
1999.....	2.6	3.2	4.0	5.3	5.5	6.6	13.4	16.2	34.4	72.4

Black, male

2004.....	2.8	3.3	4.8	5.0	6.0	10.4	*	15.5	17.9	45.7
2003.....	2.8	3.7	4.4	4.5	5.4	4.9	10.5	16.4	25.1	48.0
2002.....	3.7	2.6	3.8	6.7	5.7	7.0	8.7	12.4	29.8	48.2
2001.....	2.9	3.9	4.4	4.2	6.1	6.4	10.4	13.5	26.5	57.9
2000.....	2.8	2.8	4.0	6.4	5.6	6.5	12.0	18.0	25.3	65.7
1999.....	2.4	2.6	3.1	6.1	5.3	6.2	14.9	18.0	35.8	75.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nutritional diseases (E40-E68)

Black, female

2004.....	3.5	5.9	6.4	6.7	8.9	8.5	13.4	12.9	18.9	52.8
2003.....	3.2	4.5	5.1	6.5	8.0	6.0	9.5	11.7	21.7	62.0
2002.....	3.2	4.6	6.4	8.2	8.1	10.0	9.0	18.7	22.3	67.4
2001.....	2.8	4.5	6.4	6.3	5.4	7.1	10.8	16.6	26.6	55.6
2000.....	3.1	4.7	5.6	6.1	5.1	8.3	9.3	16.2	27.2	69.5
1999.....	2.9	3.7	4.8	4.6	5.7	6.9	12.4	15.2	33.7	71.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
 years years years years years years years years years and over

Malnutrition (E40-E46)

All races 3/, both sexes

2004.....	0.1	0.2	0.4	0.4	0.7	1.2	2.0	4.4	9.6	28.3
2003.....	0.1	0.2	0.3	0.5	0.7	1.2	2.4	5.5	10.0	33.4
2002.....	0.1	0.2	0.3	0.5	0.7	1.4	2.5	5.6	12.2	38.4
2001.....	0.1	0.2	0.3	0.4	0.8	1.3	2.6	5.5	12.0	40.2
2000.....	0.1	0.2	0.3	0.5	0.9	1.9	2.8	6.4	14.6	46.3
1999.....	0.2	0.2	0.3	0.5	0.8	1.5	3.0	6.0	14.7	50.0

All races 3/, male

2004.....	*	*	0.3	0.5	0.7	1.3	2.2	4.8	9.2	25.3
2003.....	*	*	0.3	0.5	0.7	1.3	2.5	5.4	10.4	28.2
2002.....	*	*	0.3	0.5	0.8	1.4	2.8	5.7	11.4	35.1
2001.....	*	0.2	0.3	0.5	0.8	1.3	2.6	5.1	12.6	33.6
2000.....	*	0.2	*	0.6	0.9	1.8	3.3	6.4	13.1	40.7
1999.....	*	0.2	0.3	0.6	1.1	1.5	3.6	6.2	14.8	44.4

All races 3/, female

2004.....	*	0.2	0.4	0.3	0.7	1.1	1.9	4.2	9.9	29.6
2003.....	*	*	0.3	0.5	0.8	1.1	2.2	5.7	9.7	35.6
2002.....	*	*	0.3	0.4	0.6	1.4	2.2	5.6	12.6	39.8
2001.....	*	0.2	0.3	0.4	0.7	1.3	2.5	5.7	11.6	43.0
2000.....	*	0.2	0.3	0.3	0.9	1.9	2.4	6.5	15.5	48.6
1999.....	0.2	0.2	0.3	0.4	0.6	1.4	2.5	5.8	14.7	52.3

White, both sexes

2004.....	*	0.1	0.3	0.4	0.6	1.0	1.9	4.1	9.3	26.9
2003.....	*	0.1	0.2	0.4	0.6	1.1	2.2	5.4	9.3	32.1
2002.....	0.1	0.2	0.3	0.4	0.6	1.3	2.4	5.4	11.7	37.0
2001.....	*	0.2	0.2	0.4	0.7	1.1	2.4	5.1	11.2	39.5
2000.....	*	0.2	0.2	0.4	0.8	1.8	2.4	6.1	14.2	45.2
1999.....	0.2	0.2	0.3	0.4	0.7	1.3	2.5	5.5	13.8	48.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malnutrition (E40-E46)

White, male

2004.....	*	*	0.3	0.4	0.6	1.0	2.0	4.3	8.7	24.1
2003.....	*	*	*	0.4	0.5	1.2	2.2	4.9	9.5	27.2
2002.....	*	*	*	0.4	0.6	1.3	2.7	5.5	10.5	34.3
2001.....	*	*	*	0.4	0.7	1.0	2.3	4.7	11.7	32.2
2000.....	*	*	*	0.5	0.7	1.6	2.8	5.9	12.5	39.7
1999.....	*	*	0.3	0.4	0.9	1.2	2.8	5.6	13.4	42.2

White, female

2004.....	*	*	0.3	0.4	0.6	1.1	1.7	3.9	9.7	28.2
2003.....	*	*	*	0.5	0.7	1.0	2.2	5.7	9.3	34.2
2002.....	*	*	0.3	0.4	0.6	1.2	2.1	5.3	12.4	38.2
2001.....	*	*	0.3	0.4	0.8	1.2	2.5	5.4	10.9	42.6
2000.....	*	*	0.3	*	0.9	1.9	2.1	6.2	15.1	47.4
1999.....	*	*	0.3	0.4	0.5	1.3	2.3	5.5	14.0	51.6

Black, both sexes

2004.....	*	*	0.9	*	*	2.5	4.2	8.1	13.6	46.0
2003.....	*	*	1.0	*	1.8	*	4.4	8.8	19.3	53.4
2002.....	*	*	*	1.5	1.9	3.0	4.4	10.1	19.9	57.6
2001.....	*	*	*	*	*	2.8	4.0	10.1	22.2	51.6
2000.....	*	*	*	*	1.9	3.1	6.5	11.2	20.9	63.8
1999.....	*	*	*	1.6	2.0	3.2	8.0	11.1	28.9	66.0

Black, male

2004.....	*	*	*	*	*	*	*	10.5	14.3	43.7
2003.....	*	*	*	*	*	*	6.9	13.2	23.6	45.9
2002.....	*	*	*	*	*	*	*	9.2	25.2	46.1
2001.....	*	*	*	*	*	*	*	11.2	24.1	55.6
2000.....	*	*	*	*	*	*	9.7	14.2	22.8	61.1
1999.....	*	*	*	*	*	*	12.2	13.3	33.1	73.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malnutrition (E40-E46)

Black, female

2004.....	*	*	*	*	*	*	*	6.7	13.2	46.9
2003.....	*	*	*	*	*	*	*	6.2	17.0	56.3
2002.....	*	*	*	*	*	*	*	10.7	17.1	62.0
2001.....	*	*	*	*	*	*	*	9.4	21.3	50.1
2000.....	*	*	*	*	*	*	4.4	9.4	20.0	64.8
1999.....	*	*	*	*	*	*	5.2	9.7	26.8	63.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nutritional marasmus (E41)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	*	0.5	1.3
2003.....	*	*	*	*	*	*	*	0.3	0.6	1.6
2002.....	*	*	*	*	*	*	*	*	0.6	2.4
2001.....	*	*	*	*	*	*	*	*	0.6	2.5
2000.....	*	*	*	*	*	*	*	*	0.8	2.7
1999.....	*	*	*	*	*	*	*	0.3	0.8	2.5

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	2.2
2000.....	*	*	*	*	*	*	*	*	*	2.9
1999.....	*	*	*	*	*	*	*	*	*	3.0

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	1.4
2003.....	*	*	*	*	*	*	*	*	0.6	1.7
2002.....	*	*	*	*	*	*	*	*	0.6	2.7
2001.....	*	*	*	*	*	*	*	*	*	2.6
2000.....	*	*	*	*	*	*	*	*	0.9	2.7
1999.....	*	*	*	*	*	*	*	*	0.6	2.2

White, both sexes

2004.....	*	*	*	*	*	*	*	*	0.5	1.2
2003.....	*	*	*	*	*	*	*	*	0.5	1.6
2002.....	*	*	*	*	*	*	*	*	0.6	2.4
2001.....	*	*	*	*	*	*	*	*	0.6	2.5
2000.....	*	*	*	*	*	*	*	*	0.9	2.7
1999.....	*	*	*	*	*	*	*	*	0.8	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified malnutrition (E40,E42-E46)

All races 3/, both sexes

2004.....	0.1	0.2	0.4	0.4	0.7	1.1	1.9	4.3	9.2	27.0
2003.....	0.1	0.2	0.3	0.5	0.7	1.1	2.3	5.3	9.4	31.8
2002.....	0.1	0.2	0.3	0.5	0.7	1.4	2.4	5.4	11.6	36.0
2001.....	0.1	0.2	0.3	0.4	0.7	1.2	2.4	5.2	11.4	37.7
2000.....	0.1	0.2	0.2	0.4	0.9	1.8	2.7	6.2	13.8	43.6
1999.....	0.2	0.2	0.3	0.5	0.8	1.4	2.8	5.7	14.0	47.5

All races 3/, male

2004.....	*	*	0.3	0.5	0.7	1.2	2.1	4.6	8.7	24.3
2003.....	*	*	0.3	0.5	0.6	1.2	2.5	5.1	9.9	27.1
2002.....	*	*	0.2	0.5	0.8	1.4	2.8	5.6	10.8	33.5
2001.....	*	0.2	0.3	0.5	0.7	1.2	2.5	4.8	11.9	31.3
2000.....	*	0.2	*	0.6	0.9	1.7	3.2	6.1	12.4	37.8
1999.....	*	0.2	0.3	0.6	1.0	1.5	3.3	6.0	13.8	41.4

All races 3/, female

2004.....	*	*	0.4	0.3	0.7	1.1	1.8	4.1	9.4	28.2
2003.....	*	*	0.3	0.5	0.7	1.1	2.1	5.4	9.1	33.9
2002.....	*	*	0.3	0.4	0.6	1.3	2.1	5.3	12.0	37.1
2001.....	*	0.2	0.3	0.4	0.7	1.2	2.4	5.5	11.1	40.4
2000.....	*	0.2	0.3	0.3	0.8	1.9	2.3	6.3	14.6	45.9
1999.....	0.2	*	0.3	0.4	0.6	1.4	2.4	5.5	14.1	50.0

White, both sexes

2004.....	*	0.1	0.3	0.4	0.6	1.0	1.8	3.9	8.8	25.7
2003.....	*	0.1	0.2	0.4	0.6	1.1	2.1	5.1	8.8	30.5
2002.....	0.1	0.2	0.2	0.4	0.6	1.2	2.3	5.2	11.1	34.7
2001.....	*	0.2	0.2	0.4	0.7	1.1	2.3	4.8	10.6	37.0
2000.....	*	0.2	0.2	0.4	0.7	1.7	2.3	5.9	13.3	42.5
1999.....	0.2	0.2	0.3	0.3	0.7	1.2	2.3	5.4	12.9	46.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified malnutrition (E40,E42-E46)

White, male

2004.....	*	*	0.3	0.4	0.5	0.9	1.9	4.2	8.2	23.2
2003.....	*	*	*	0.3	0.5	1.1	2.1	4.6	9.0	26.0
2002.....	*	*	*	0.4	0.6	1.3	2.6	5.4	10.0	32.5
2001.....	*	*	*	0.4	0.6	1.0	2.3	4.4	11.0	30.1
2000.....	*	*	*	0.4	0.7	1.6	2.6	5.7	11.7	37.0
1999.....	*	*	0.3	*	0.9	1.2	2.6	5.4	12.5	39.4

White, female

2004.....	*	*	0.3	0.3	0.6	1.1	1.7	3.8	9.2	26.9
2003.....	*	*	*	0.5	0.7	1.0	2.1	5.5	8.7	32.4
2002.....	*	*	0.3	0.4	0.6	1.2	2.0	5.0	11.8	35.6
2001.....	*	*	0.3	0.3	0.8	1.2	2.4	5.1	10.4	39.9
2000.....	*	*	*	*	0.8	1.8	2.0	6.0	14.2	44.7
1999.....	*	*	0.3	0.3	0.5	1.3	2.1	5.3	13.2	49.4

Black, both sexes

2004.....	*	*	0.9	*	*	2.4	3.9	8.1	13.6	44.8
2003.....	*	*	0.9	*	*	*	4.2	8.3	18.0	52.5
2002.....	*	*	*	1.4	1.8	2.9	4.3	9.3	18.8	55.8
2001.....	*	*	*	*	*	2.7	3.9	10.1	20.9	49.1
2000.....	*	*	*	*	1.9	3.0	6.3	10.7	20.9	59.7
1999.....	*	*	*	1.5	1.9	3.1	7.4	9.6	28.6	62.8

Black, male

2004.....	*	*	*	*	*	*	*	10.5	14.3	42.7
2003.....	*	*	*	*	*	*	6.9	12.3	22.9	44.8
2002.....	*	*	*	*	*	*	*	*	22.9	45.0
2001.....	*	*	*	*	*	*	*	11.2	23.3	53.3
2000.....	*	*	*	*	*	*	9.4	12.8	22.8	55.4
1999.....	*	*	*	*	*	*	11.5	12.8	32.3	69.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified malnutrition (E40,E42-E46)

Black, female

2004.....	*	*	*	*	*	*	*	6.7	13.2	45.7
2003.....	*	*	*	*	*	*	*	6.0	15.5	55.5
2002.....	*	*	*	*	*	*	*	9.6	16.7	60.0
2001.....	*	*	*	*	*	*	*	9.4	19.6	47.6
2000.....	*	*	*	*	*	*	*	9.4	20.0	61.4
1999.....	*	*	*	*	*	*	4.7	7.7	26.8	60.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other nutritional deficiencies (E50-E64)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	*	0.4	2.1
2003.....	*	*	*	*	*	*	*	*	0.5	2.1
2002.....	*	*	*	*	*	*	*	0.4	0.6	3.1
2001.....	*	*	*	*	*	*	*	0.4	0.8	2.7
2000.....	*	*	*	*	*	*	0.3	0.4	1.2	3.7
1999.....	*	*	*	*	*	*	*	0.4	1.1	3.9

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	2.1
2003.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	1.9
2001.....	*	*	*	*	*	*	*	*	*	1.6
2000.....	*	*	*	*	*	*	*	*	1.3	3.0
1999.....	*	*	*	*	*	*	*	*	1.3	3.9

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	2.1
2003.....	*	*	*	*	*	*	*	*	*	2.0
2002.....	*	*	*	*	*	*	*	*	0.6	3.6
2001.....	*	*	*	*	*	*	*	*	1.0	3.1
2000.....	*	*	*	*	*	*	*	*	1.1	4.0
1999.....	*	*	*	*	*	*	*	*	1.1	3.9

White, both sexes

2004.....	*	*	*	*	*	*	*	*	0.4	2.2
2003.....	*	*	*	*	*	*	*	*	0.5	2.2
2002.....	*	*	*	*	*	*	*	0.5	0.6	3.1
2001.....	*	*	*	*	*	*	*	0.4	0.8	2.7
2000.....	*	*	*	*	*	*	0.3	0.3	1.2	3.7
1999.....	*	*	*	*	*	*	*	0.4	1.1	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Obesity and other hyperalimentation (E65-E68)

All races 3/, both sexes

2004.....	1.6	2.3	3.0	3.3	3.6	3.6	3.7	3.1	2.7	2.0
2003.....	1.6	2.4	2.8	3.2	3.4	3.8	3.3	2.6	2.5	2.0
2002.....	1.8	2.0	2.7	3.3	3.1	3.1	2.4	2.6	2.2	1.4
2001.....	1.3	1.9	2.5	2.7	2.7	2.5	2.7	2.2	1.7	1.6
2000.....	1.5	1.7	2.4	2.4	2.6	2.5	2.4	2.3	2.2	1.7
1999.....	1.3	1.5	2.0	2.2	2.1	2.2	2.4	2.2	2.0	1.4

All races 3/, male

2004.....	1.8	2.3	3.2	3.3	3.2	3.4	2.8	2.7	2.7	1.4
2003.....	1.7	2.4	2.8	3.4	3.2	3.7	2.5	1.9	1.9	*
2002.....	2.1	2.0	2.8	3.2	2.7	2.7	1.9	1.9	1.5	*
2001.....	1.4	1.9	2.3	2.4	2.5	1.9	2.4	1.3	1.2	*
2000.....	1.6	1.9	2.5	2.6	2.2	2.0	1.9	1.7	1.4	*
1999.....	1.6	1.5	1.9	2.5	1.9	1.9	1.9	1.6	*	*

All races 3/, female

2004.....	1.4	2.3	2.8	3.3	3.9	3.8	4.5	3.5	2.8	2.2
2003.....	1.6	2.3	2.8	3.1	3.6	3.9	3.9	3.0	2.8	2.5
2002.....	1.5	2.1	2.6	3.4	3.5	3.5	2.8	3.1	2.6	1.6
2001.....	1.1	1.8	2.6	2.9	3.0	3.0	2.9	2.9	2.0	2.0
2000.....	1.4	1.6	2.3	2.2	3.0	2.9	2.8	2.7	2.6	2.0
1999.....	1.1	1.4	2.0	1.9	2.2	2.5	2.8	2.6	2.6	1.9

White, both sexes

2004.....	1.4	2.1	2.9	3.2	3.5	3.4	3.6	3.0	2.6	1.9
2003.....	1.5	2.3	2.8	3.2	3.3	3.9	3.2	2.5	2.4	1.9
2002.....	1.6	2.0	2.6	3.1	3.0	3.0	2.3	2.3	2.1	1.4
2001.....	1.1	1.7	2.3	2.6	2.6	2.4	2.4	2.1	1.7	1.6
2000.....	1.4	1.6	2.3	2.2	2.6	2.3	2.3	2.2	2.1	1.7
1999.....	1.2	1.4	1.9	2.2	2.0	2.2	2.2	2.0	2.0	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Obesity and other hyperalimentation (E65-E68)										
Black, female										
2004.....	3.3	5.5	5.5	6.4	7.4	7.2	9.2	6.2	*	*
2003.....	3.1	4.2	4.2	5.9	6.5	4.2	6.6	5.4	*	*
2002.....	3.0	4.5	6.0	7.3	7.1	6.5	*	7.7	*	*
2001.....	2.6	4.5	5.7	5.6	5.0	5.3	7.8	6.4	*	*
2000.....	2.8	4.4	4.9	5.7	3.8	5.6	4.9	6.0	*	*
1999.....	2.5	2.8	4.3	3.5	4.2	4.6	7.2	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Metabolic disorders (E70-E88)

All races 3/, both sexes

2004.....	0.9	1.7	2.5	4.3	6.6	10.4	16.8	26.7	45.1	107.9
2003.....	1.0	1.6	2.9	4.1	6.6	10.4	16.1	25.9	46.5	112.6
2002.....	1.0	1.5	2.6	4.6	6.9	9.7	16.6	26.5	47.3	114.3
2001.....	1.0	1.7	2.7	4.1	6.3	10.1	16.4	26.3	45.2	124.4
2000.....	1.0	1.5	2.5	4.0	6.5	9.8	15.7	25.7	46.1	124.1
1999.....	0.9	1.3	2.3	3.5	6.2	9.5	15.1	25.3	44.7	130.8

All races 3/, male

2004.....	1.1	2.4	3.6	5.8	8.7	13.9	20.7	30.3	50.5	103.1
2003.....	1.4	2.1	4.0	5.6	8.7	13.9	20.1	30.0	51.0	104.8
2002.....	1.3	2.0	3.7	6.1	8.9	12.5	20.0	30.4	51.5	101.5
2001.....	1.2	2.3	3.5	5.2	8.8	13.7	20.0	32.6	48.0	109.8
2000.....	1.2	2.2	3.6	5.3	8.9	12.8	19.2	29.6	48.6	112.0
1999.....	1.1	1.8	3.1	4.7	7.8	11.6	19.0	29.2	46.9	110.5

All races 3/, female

2004.....	0.7	1.1	1.5	2.9	4.7	7.4	13.6	24.1	41.7	110.1
2003.....	0.7	1.1	1.9	2.7	4.7	7.4	12.9	23.0	43.8	116.1
2002.....	0.8	1.1	1.6	3.2	5.0	7.2	13.9	23.8	44.7	119.9
2001.....	0.7	1.1	1.9	3.0	3.9	7.0	13.6	22.0	43.5	130.6
2000.....	0.8	0.8	1.4	2.8	4.3	7.2	12.9	22.9	44.7	129.1
1999.....	0.7	0.9	1.7	2.4	4.7	7.7	12.1	22.7	43.4	139.0

White, both sexes

2004.....	0.9	1.7	2.5	4.2	6.5	10.2	16.7	26.7	45.2	109.2
2003.....	1.0	1.6	2.8	3.9	6.6	10.1	16.1	25.3	45.9	112.9
2002.....	1.0	1.5	2.6	4.5	6.6	9.4	16.0	26.1	46.3	114.0
2001.....	1.0	1.5	2.7	4.0	6.1	9.9	16.3	26.0	44.4	123.4
2000.....	0.9	1.5	2.4	3.9	6.3	9.7	15.5	25.4	45.6	123.6
1999.....	0.8	1.3	2.2	3.5	6.0	9.1	14.8	25.0	44.2	131.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Metabolic disorders (E70-E88)

White, male

2004.....	1.0	2.4	3.5	5.7	8.8	13.9	20.5	30.2	50.9	104.6
2003.....	1.3	2.1	3.9	5.5	8.8	13.5	20.0	29.3	49.9	106.6
2002.....	1.3	1.8	3.7	6.0	8.4	12.1	19.6	29.8	50.6	100.2
2001.....	1.2	2.1	3.5	5.1	8.4	13.5	19.7	32.2	46.9	109.2
2000.....	1.1	2.2	3.5	5.2	8.6	12.6	19.1	29.1	48.1	113.9
1999.....	1.0	1.7	3.0	4.7	7.8	11.2	18.6	28.6	45.7	108.4

White, female

2004.....	0.7	1.0	1.5	2.8	4.4	7.0	13.5	24.1	41.6	111.2
2003.....	0.7	1.1	1.7	2.4	4.6	7.0	12.9	22.5	43.4	115.6
2002.....	0.7	1.1	1.5	3.0	5.0	6.9	13.0	23.4	43.7	120.0
2001.....	0.8	1.0	1.9	3.0	3.8	6.7	13.6	21.6	42.9	129.4
2000.....	0.7	0.8	1.2	2.7	4.2	7.1	12.6	22.8	44.1	127.6
1999.....	0.6	0.9	1.4	2.4	4.4	7.4	11.9	22.5	43.3	140.4

Black, both sexes

2004.....	1.5	2.4	3.1	6.0	9.2	14.3	22.4	32.8	52.5	113.1
2003.....	1.3	2.1	4.7	6.1	8.3	15.7	20.4	36.2	59.1	126.0
2002.....	1.3	2.5	3.8	6.5	10.8	13.9	26.0	34.7	66.5	135.1
2001.....	0.9	2.8	3.6	5.1	9.4	13.8	21.5	36.0	62.3	151.4
2000.....	1.4	2.1	3.6	5.8	9.0	13.3	20.1	33.2	61.0	147.0
1999.....	1.8	1.9	4.3	4.6	8.7	14.9	21.0	32.7	54.0	138.7

Black, male

2004.....	2.0	3.1	4.8	7.9	10.5	17.2	28.5	39.1	58.8	112.3
2003.....	1.8	2.8	6.0	7.9	10.2	20.7	26.3	42.9	74.6	98.0
2002.....	1.5	3.5	5.0	8.5	16.4	17.7	27.4	40.9	71.8	134.0
2001.....	*	3.7	4.3	6.9	13.6	18.8	28.6	44.6	75.4	130.5
2000.....	1.7	2.9	5.1	7.1	12.2	16.7	24.4	41.1	65.8	115.3
1999.....	2.0	2.5	4.8	6.3	10.0	18.7	26.1	40.8	61.9	147.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Metabolic disorders (E70-E88)

Black, female

2004.....	*	1.9	1.7	4.5	8.2	12.2	18.2	29.1	49.1	113.4
2003.....	*	1.5	3.6	4.6	6.8	12.1	16.4	32.3	51.1	136.9
2002.....	*	1.8	2.8	4.9	6.3	11.1	25.1	31.0	63.7	135.5
2001.....	*	2.1	2.9	3.6	6.1	10.1	16.8	30.8	55.6	159.1
2000.....	*	*	2.4	4.7	6.5	10.8	17.3	28.5	58.6	158.8
1999.....	1.6	*	3.9	3.3	7.7	12.1	17.6	27.7	50.1	135.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

All races 3/, both sexes

2004.....	0.2	0.5	0.5	0.9	1.3	2.1	4.5	9.1	19.7	70.7
2003.....	0.2	0.4	0.7	0.8	1.5	2.3	4.6	9.9	22.4	77.3
2002.....	0.3	0.5	0.6	1.0	1.6	2.5	5.2	9.8	24.7	83.2
2001.....	0.3	0.5	0.6	0.8	1.7	2.6	5.6	11.0	25.1	93.9
2000.....	0.4	0.4	0.6	1.0	1.7	3.0	5.2	11.3	26.6	96.8
1999.....	0.3	0.4	0.7	0.8	1.4	3.0	5.5	11.9	27.7	106.4

All races 3/, male

2004.....	0.3	0.6	0.7	1.0	1.3	2.4	4.8	9.0	19.6	59.8
2003.....	0.2	0.5	0.7	0.9	1.5	2.5	4.6	10.2	22.2	65.6
2002.....	0.3	0.5	0.6	1.0	1.7	2.6	5.2	9.6	24.1	67.9
2001.....	0.3	0.5	0.6	0.8	2.0	2.8	5.5	12.2	24.0	76.5
2000.....	0.4	0.5	0.7	1.2	1.9	3.4	5.8	12.0	27.2	84.3
1999.....	0.4	0.5	0.8	1.1	1.4	3.1	6.2	13.2	27.2	85.0

All races 3/, female

2004.....	0.2	0.4	0.4	0.7	1.2	1.9	4.2	9.2	19.8	75.7
2003.....	0.2	0.4	0.7	0.6	1.5	2.2	4.6	9.6	22.5	82.5
2002.....	0.2	0.4	0.5	1.0	1.5	2.3	5.2	10.0	25.0	89.8
2001.....	0.3	0.4	0.6	0.8	1.4	2.5	5.7	10.2	25.7	101.2
2000.....	0.3	0.3	0.5	0.9	1.5	2.6	4.7	10.8	26.2	101.9
1999.....	0.2	0.3	0.5	0.7	1.5	2.9	5.0	11.1	27.9	115.1

White, both sexes

2004.....	0.2	0.4	0.5	0.7	1.1	1.9	4.1	8.9	19.4	70.8
2003.....	0.2	0.4	0.6	0.7	1.4	2.0	4.4	9.2	21.7	76.6
2002.....	0.2	0.4	0.5	0.9	1.4	2.2	4.7	9.2	23.5	82.4
2001.....	0.3	0.3	0.5	0.7	1.4	2.4	5.1	10.4	24.0	93.0
2000.....	0.3	0.3	0.5	0.9	1.5	2.7	4.8	10.9	25.9	96.0
1999.....	0.2	0.4	0.5	0.7	1.2	2.5	5.2	11.4	27.2	106.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
White, male										
2004.....	*	0.5	0.7	0.8	1.0	2.3	4.5	8.6	19.3	59.0
2003.....	*	0.4	0.5	0.9	1.4	2.1	4.3	9.4	20.9	65.6
2002.....	0.2	0.4	0.5	0.8	1.5	2.2	4.8	8.9	22.7	66.1
2001.....	0.2	0.4	0.5	0.6	1.6	2.6	4.7	11.5	22.3	75.8
2000.....	0.3	0.4	0.6	1.0	1.6	3.0	5.3	11.2	26.3	84.9
1999.....	0.2	0.5	0.7	0.9	1.1	2.3	5.6	12.3	26.0	82.4
White, female										
2004.....	*	0.3	0.4	0.6	1.2	1.6	3.8	9.1	19.5	76.1
2003.....	*	0.3	0.6	0.5	1.4	1.9	4.4	9.0	22.2	81.5
2002.....	*	0.4	0.5	1.0	1.4	2.2	4.7	9.4	24.0	89.5
2001.....	0.3	0.3	0.5	0.7	1.3	2.1	5.4	9.7	25.0	100.2
2000.....	0.3	0.3	0.4	0.8	1.4	2.3	4.4	10.7	25.7	100.5
1999.....	*	0.3	0.4	0.6	1.2	2.6	4.8	10.8	27.9	116.4
Black, both sexes										
2004.....	0.7	1.0	*	2.4	3.0	4.5	9.3	13.9	28.0	84.3
2003.....	*	0.8	2.1	2.0	3.0	6.1	8.2	20.2	35.0	97.1
2002.....	0.8	1.2	1.3	2.0	3.1	5.4	11.4	18.9	42.9	104.2
2001.....	*	1.4	1.6	1.8	4.2	5.5	12.1	20.7	42.8	119.4
2000.....	0.8	1.3	1.6	2.2	3.4	6.6	10.1	18.5	40.9	119.8
1999.....	1.1	1.0	1.9	2.0	4.0	7.8	11.0	19.8	36.7	115.1
Black, male										
2004.....	*	*	*	3.6	4.1	5.3	10.4	16.4	31.6	86.4
2003.....	*	*	2.7	*	*	7.4	9.5	23.3	48.0	77.2
2002.....	*	1.8	*	*	4.4	7.0	11.3	20.7	46.6	99.7
2001.....	*	*	*	*	5.9	5.8	15.8	24.2	54.6	97.6
2000.....	*	*	*	*	4.6	8.1	12.0	26.0	47.2	98.0
1999.....	1.5	*	*	*	*	11.4	14.9	26.6	44.5	125.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

Black, female

2004.....	*	*	*	*	*	4.0	8.6	12.4	26.1	83.5
2003.....	*	*	*	*	3.0	5.1	7.3	18.4	28.2	104.9
2002.....	*	*	*	*	*	4.3	11.4	17.8	41.0	106.0
2001.....	*	*	*	*	*	5.3	9.6	18.6	36.8	127.6
2000.....	*	*	*	*	*	5.6	8.9	14.0	37.8	127.9
1999.....	*	*	*	*	4.0	5.2	8.3	15.7	32.8	111.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other metabolic disorders (E70-E83,E85,E88)

All races 3/, both sexes

2004.....	0.6	1.2	1.9	3.4	5.3	8.2	12.3	17.6	25.3	37.1
2003.....	0.6	1.1	2.2	3.3	5.1	8.1	11.5	16.0	24.1	35.2
2002.....	0.6	1.0	2.0	3.6	5.2	7.2	11.4	16.7	22.6	31.1
2001.....	0.5	1.1	2.0	3.2	4.6	7.4	10.8	15.3	20.0	30.5
2000.....	0.4	1.0	1.9	2.9	4.8	6.8	10.5	14.3	19.6	27.3
1999.....	0.5	0.9	1.6	2.6	4.7	6.5	9.5	13.4	17.0	24.3

All races 3/, male

2004.....	0.8	1.7	2.8	4.7	7.4	11.4	15.8	21.3	30.9	43.2
2003.....	0.9	1.5	3.2	4.6	7.2	11.4	15.5	19.7	28.8	39.2
2002.....	0.9	1.4	3.1	5.1	7.1	9.8	14.7	20.8	27.3	33.6
2001.....	0.7	1.6	2.8	4.4	6.8	10.9	14.5	20.3	23.8	33.3
2000.....	0.6	1.5	2.9	4.0	6.9	9.3	13.4	17.6	21.4	27.7
1999.....	0.6	1.3	2.2	3.6	6.4	8.6	12.7	15.9	19.6	25.5

All races 3/, female

2004.....	0.4	0.6	1.0	2.2	3.4	5.5	9.4	14.9	21.8	34.4
2003.....	0.3	0.7	1.2	2.0	3.2	5.2	8.4	13.4	21.2	33.4
2002.....	0.4	0.7	1.0	2.2	3.5	4.9	8.7	13.8	19.7	30.0
2001.....	0.4	0.6	1.2	2.1	2.6	4.5	7.8	11.7	17.8	29.3
2000.....	0.3	0.5	0.9	1.9	2.8	4.6	8.2	12.1	18.5	27.1
1999.....	0.4	0.5	1.0	1.7	3.2	4.8	7.1	11.6	15.5	23.9

White, both sexes

2004.....	0.6	1.2	1.9	3.5	5.4	8.3	12.5	17.8	25.7	38.4
2003.....	0.6	1.1	2.2	3.2	5.2	8.1	11.7	16.1	24.2	36.1
2002.....	0.6	1.0	2.0	3.6	5.2	7.2	11.2	16.9	22.7	31.6
2001.....	0.6	1.1	2.1	3.3	4.6	7.5	11.2	15.5	20.3	30.4
2000.....	0.4	1.1	1.9	3.0	4.8	7.0	10.7	14.5	19.7	27.5
1999.....	0.5	0.9	1.6	2.7	4.8	6.7	9.7	13.6	16.9	24.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other metabolic disorders (E70-E83,E85,E88)

White, male

2004.....	0.8	1.8	2.8	4.9	7.7	11.6	15.9	21.5	31.6	45.5
2003.....	0.9	1.5	3.3	4.6	7.4	11.4	15.7	19.8	29.0	40.9
2002.....	0.9	1.4	3.2	5.2	6.9	9.9	14.8	20.9	27.8	34.1
2001.....	0.8	1.6	2.9	4.4	6.8	10.9	15.0	20.7	24.4	33.3
2000.....	0.6	1.7	2.9	4.1	6.9	9.5	13.8	18.0	21.8	29.0
1999.....	0.6	1.2	2.2	3.7	6.6	8.9	13.0	16.2	19.6	25.9

White, female

2004.....	0.4	0.6	1.0	2.1	3.2	5.3	9.7	15.0	22.0	35.1
2003.....	0.3	0.7	1.1	1.9	3.2	5.2	8.5	13.4	21.2	34.0
2002.....	0.4	0.7	0.9	2.1	3.6	4.7	8.3	14.0	19.6	30.4
2001.....	0.4	0.6	1.3	2.2	2.6	4.5	8.1	11.8	17.9	29.1
2000.....	0.2	0.5	0.9	1.8	2.8	4.7	8.2	12.1	18.4	26.9
1999.....	0.3	0.5	0.9	1.7	3.2	4.7	7.0	11.8	15.4	24.0

Black, both sexes

2004.....	0.7	1.4	2.4	3.6	6.1	9.8	13.1	19.0	24.5	28.8
2003.....	0.7	1.3	2.5	4.1	5.3	9.7	12.2	15.8	24.1	28.9
2002.....	*	1.3	2.4	4.4	7.6	8.5	14.5	15.8	23.5	30.9
2001.....	*	1.5	1.9	3.2	5.2	8.3	9.4	15.3	19.5	31.9
2000.....	*	*	2.0	3.6	5.7	6.6	10.0	14.7	20.0	27.2
1999.....	0.7	0.9	2.3	2.7	4.7	7.1	10.0	12.9	17.3	23.6

Black, male

2004.....	*	1.9	3.6	4.4	6.4	11.9	18.1	22.7	27.3	25.8
2003.....	*	1.6	3.2	5.8	7.3	13.3	16.8	19.6	26.6	20.9
2002.....	*	1.7	3.3	5.8	12.1	10.7	15.7	20.2	25.2	34.3
2001.....	*	2.1	2.5	4.7	7.7	13.0	12.8	20.4	20.9	32.9
2000.....	*	*	3.1	4.0	7.7	8.6	12.4	15.1	18.5	*
1999.....	*	*	2.5	*	5.9	7.3	11.2	14.2	17.4	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other metabolic disorders (E70-E83,E85,E88)

Black, female

2004.....	*	*	*	3.0	6.0	8.3	9.7	16.7	23.1	29.9
2003.....	*	*	2.0	2.7	3.8	6.9	9.1	13.6	22.8	32.0
2002.....	*	*	*	3.3	4.0	6.8	13.7	13.2	22.7	29.6
2001.....	*	*	*	*	3.2	4.9	7.2	12.2	18.8	31.6
2000.....	*	*	*	3.2	4.1	5.2	8.4	14.5	20.8	30.9
1999.....	*	*	2.2	*	3.7	6.9	9.2	12.0	17.3	24.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

V. Mental and behavioral disorders (F01-F99)

All races 3/, both sexes

2004.....	6.0	8.6	9.5	9.2	10.5	13.8	28.6	69.6	180.4	671.4
2003.....	6.3	8.4	8.9	8.3	10.4	13.6	27.2	69.8	179.2	679.3
2002.....	6.5	7.9	8.3	8.3	9.5	13.0	25.6	64.2	168.2	646.7
2001.....	6.0	7.9	7.7	7.7	9.4	13.0	23.8	60.6	154.3	604.8
2000.....	5.8	7.5	7.1	7.8	8.6	11.7	22.5	54.1	143.7	561.6
1999.....	6.1	7.4	6.8	7.3	8.9	11.2	21.2	49.9	128.6	515.3

All races 3/, male

2004.....	8.8	12.5	14.9	14.5	15.8	18.7	34.0	73.2	174.5	529.7
2003.....	9.2	12.7	14.0	12.8	15.0	18.5	32.1	73.2	175.7	523.8
2002.....	9.3	12.0	13.4	12.9	14.4	18.4	30.2	68.2	166.1	506.0
2001.....	8.7	12.2	12.3	11.7	14.3	18.1	29.3	63.3	151.3	488.0
2000.....	8.5	11.6	11.4	11.9	13.0	16.5	28.0	59.4	140.5	451.0
1999.....	9.4	11.5	11.1	11.7	13.8	16.1	26.9	56.9	133.4	415.8

All races 3/, female

2004.....	3.2	4.9	4.3	4.3	5.6	9.6	24.2	67.0	184.1	735.1
2003.....	3.5	4.2	4.1	4.1	6.2	9.4	23.4	67.4	181.4	748.1
2002.....	3.6	4.0	3.3	4.0	5.1	8.4	21.9	61.3	169.4	707.8
2001.....	3.3	3.8	3.3	3.9	4.9	8.6	19.5	58.8	156.0	653.7
2000.....	3.2	3.5	3.1	4.0	4.6	7.7	18.3	50.4	145.6	606.6
1999.....	2.8	3.4	2.8	3.3	4.5	7.0	16.7	45.1	125.8	555.4

White, both sexes

2004.....	6.0	8.4	9.0	8.9	10.0	13.0	28.1	69.0	183.4	687.9
2003.....	6.1	8.1	8.3	7.8	9.8	13.5	27.1	69.8	180.9	697.4
2002.....	6.2	7.5	7.6	7.7	8.8	12.5	25.3	64.1	170.5	663.8
2001.....	5.5	7.1	6.8	6.9	8.9	12.5	23.5	61.1	157.4	621.6
2000.....	5.4	6.7	6.6	7.2	7.9	11.2	22.0	54.0	145.3	577.6
1999.....	5.5	6.5	6.1	6.5	8.3	10.7	20.7	50.0	130.3	534.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

V. Mental and behavioral disorders (F01-F99)

White, male

2004.....	8.9	12.3	13.8	13.7	14.6	17.0	33.0	70.7	175.8	538.9
2003.....	9.0	12.2	12.7	12.0	13.4	18.0	31.0	71.8	175.5	535.8
2002.....	9.2	11.3	12.3	11.6	13.0	16.9	29.0	66.5	167.4	514.9
2001.....	8.0	10.9	10.8	10.2	13.2	16.8	28.3	63.1	152.8	498.4
2000.....	7.9	10.3	10.4	10.8	11.6	15.2	26.7	58.9	140.3	459.4
1999.....	8.6	10.2	9.8	10.1	12.8	14.9	25.7	55.8	132.9	429.2

White, female

2004.....	3.1	4.5	4.2	4.3	5.7	9.5	24.0	67.8	188.1	755.1
2003.....	3.1	4.0	3.9	3.8	6.4	9.4	23.9	68.2	184.2	769.0
2002.....	3.3	3.7	3.1	4.0	4.9	8.7	22.3	62.4	172.4	728.4
2001.....	2.9	3.4	2.9	3.8	4.9	8.7	19.6	59.7	160.2	673.1
2000.....	2.9	3.1	2.8	3.8	4.4	7.7	18.3	50.6	148.3	625.6
1999.....	2.3	2.8	2.4	3.1	4.2	6.9	16.7	45.9	128.8	576.9

Black, both sexes

2004.....	6.7	12.6	15.5	14.7	16.6	24.0	41.0	95.5	188.4	589.4
2003.....	8.4	12.1	15.5	14.4	18.1	18.0	34.7	86.8	198.0	578.8
2002.....	9.5	12.6	15.0	15.9	16.8	19.8	34.5	78.3	173.2	540.8
2001.....	10.3	15.3	16.2	15.4	15.9	20.7	33.5	70.5	147.9	488.5
2000.....	9.4	15.2	13.7	14.2	16.2	18.7	33.4	68.4	145.4	442.6
1999.....	10.5	15.1	13.9	16.5	15.4	18.8	30.8	57.8	128.3	349.1

Black, male

2004.....	9.9	17.5	25.8	26.0	30.3	39.5	54.7	126.0	203.7	510.6
2003.....	11.5	18.9	27.0	23.1	33.0	27.3	51.6	106.4	216.3	482.7
2002.....	12.8	19.5	25.7	28.0	28.9	35.4	51.1	103.3	189.5	457.8
2001.....	14.9	24.5	27.3	27.4	29.1	34.6	47.8	84.6	171.8	423.2
2000.....	13.6	24.4	23.6	24.4	27.7	32.5	48.8	82.2	169.4	415.2
1999.....	15.4	23.1	23.2	29.5	26.5	31.4	46.5	79.8	168.3	309.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

V. Mental and behavioral disorders (F01-F99)

Black, female

2004.....	4.0	8.2	6.9	5.3	6.0	12.4	31.6	77.4	180.3	620.6
2003.....	5.7	6.2	5.8	7.2	6.5	11.1	23.3	75.2	188.4	616.1
2002.....	6.7	6.7	5.8	6.0	7.3	8.1	23.3	63.4	164.7	572.6
2001.....	6.2	7.4	6.7	5.5	5.6	10.5	24.0	62.1	135.7	512.8
2000.....	5.6	7.2	5.4	5.9	7.0	8.4	23.1	60.1	133.3	452.9
1999.....	6.2	8.1	6.1	6.0	6.7	9.4	20.5	44.6	108.5	364.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

Organic dementia (F01,F03)

All races 3/, both sexes

2004.....	*	0.1	0.2	0.7	2.2	5.7	19.7	58.5	163.4	628.2
2003.....	*	*	0.2	0.6	2.2	5.9	18.5	58.6	162.7	631.7
2002.....	*	0.1	0.2	0.5	1.8	5.4	17.0	53.8	150.9	598.3
2001.....	*	*	0.2	0.6	1.8	5.5	15.9	49.5	135.8	554.0
2000.....	*	0.1	0.1	0.6	1.4	4.5	14.4	43.0	125.3	505.6
1999.....	*	*	0.2	0.5	1.4	4.0	13.0	37.8	108.1	454.4

All races 3/, male

2004.....	*	*	0.3	0.8	2.5	6.2	20.5	58.4	154.1	487.8
2003.....	*	*	0.2	0.7	2.4	6.7	20.1	58.0	155.7	480.4
2002.....	*	*	*	0.5	1.9	6.2	18.7	55.1	147.0	465.7
2001.....	*	*	*	0.5	2.0	6.2	17.5	49.7	130.8	442.4
2000.....	*	*	*	0.5	1.7	5.2	16.3	45.4	120.5	398.9
1999.....	*	*	*	0.6	1.7	4.9	15.3	40.5	109.7	361.7

All races 3/, female

2004.....	*	*	*	0.6	2.0	5.3	19.0	58.6	169.2	691.3
2003.....	*	*	0.2	0.5	2.1	5.2	17.3	59.0	166.9	698.5
2002.....	*	*	*	0.6	1.6	4.6	15.7	52.9	153.3	655.9
2001.....	*	*	*	0.6	1.5	4.9	14.5	49.4	138.8	600.8
2000.....	*	*	*	0.7	1.2	4.0	12.8	41.4	128.2	549.1
1999.....	*	*	*	*	1.1	3.2	11.3	35.9	107.1	491.7

White, both sexes

2004.....	*	*	0.2	0.7	2.2	5.5	19.2	57.8	165.7	642.8
2003.....	*	*	0.2	0.6	2.2	5.8	18.4	58.3	163.9	647.9
2002.....	*	*	0.2	0.5	1.8	5.2	16.9	53.6	153.0	613.9
2001.....	*	*	0.2	0.6	1.8	5.4	15.6	49.9	138.3	569.1
2000.....	*	*	0.1	0.6	1.3	4.4	13.9	42.9	126.6	519.6
1999.....	*	*	0.2	0.5	1.4	3.8	12.9	37.9	109.6	471.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

Organic dementia (F01,F03)

White, male

2004.....	*	*	0.3	0.8	2.4	5.9	19.9	56.2	154.3	496.5
2003.....	*	*	*	0.7	2.3	6.6	19.4	56.6	155.3	491.1
2002.....	*	*	*	0.5	1.9	5.8	18.2	53.7	148.1	473.7
2001.....	*	*	*	0.5	2.0	5.9	16.9	49.8	132.0	452.2
2000.....	*	*	*	0.5	1.6	4.9	15.4	45.2	120.2	406.1
1999.....	*	*	*	0.6	1.7	4.8	14.9	40.2	109.4	373.7

White, female

2004.....	*	*	*	0.6	2.0	5.2	18.7	59.0	172.8	708.7
2003.....	*	*	*	0.5	2.2	5.1	17.6	59.5	169.2	717.4
2002.....	*	*	*	0.6	1.7	4.8	15.9	53.6	155.9	674.7
2001.....	*	*	*	0.6	1.6	5.0	14.5	50.0	142.2	618.0
2000.....	*	*	*	0.7	1.1	3.9	12.6	41.3	130.4	565.7
1999.....	*	*	*	*	1.1	3.0	11.2	36.3	109.7	510.4

Black, both sexes

2004.....	*	*	*	*	2.9	9.6	29.3	83.3	175.3	560.3
2003.....	*	*	*	*	3.2	8.3	24.3	75.0	182.0	545.3
2002.....	*	*	*	*	2.2	7.8	22.4	66.8	157.3	503.1
2001.....	*	*	*	*	2.2	8.2	23.0	57.5	132.7	452.3
2000.....	*	*	*	*	2.5	6.4	22.5	55.3	127.6	403.2
1999.....	*	*	*	*	*	6.0	17.7	42.8	107.5	309.3

Black, male

2004.....	*	*	*	*	3.9	12.6	32.7	104.2	192.3	468.9
2003.....	*	*	*	*	4.6	9.6	33.5	88.1	194.9	445.2
2002.....	*	*	*	*	*	11.0	28.7	84.0	169.7	425.6
2001.....	*	*	*	*	*	11.2	29.3	62.7	150.9	382.4
2000.....	*	*	*	*	*	9.1	29.4	61.4	145.8	372.5
1999.....	*	*	*	*	*	7.5	23.1	51.3	136.0	266.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Organic dementia (F01,F03)

Black, female

2004.....	*	*	*	*	*	7.4	27.0	70.9	166.3	596.6
2003.....	*	*	*	*	*	7.3	18.2	67.3	175.2	584.1
2002.....	*	*	*	*	*	5.4	18.1	56.5	150.8	532.7
2001.....	*	*	*	*	*	6.0	18.8	54.4	123.5	478.2
2000.....	*	*	*	*	*	4.4	18.0	51.6	118.5	414.7
1999.....	*	*	*	*	*	4.8	14.2	37.8	93.3	325.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified organic mental disorders (F04-F09)

All races 3/, both sexes

2004.....	*	*	*	*	0.2	0.4	0.8	2.2	6.0	22.3
2003.....	*	*	*	*	0.2	0.3	1.0	2.6	6.0	24.5
2002.....	*	*	*	0.1	0.2	0.5	1.2	2.4	6.5	26.7
2001.....	*	*	*	*	0.2	0.5	1.1	2.9	7.8	29.3
2000.....	*	*	*	0.2	0.3	0.5	1.2	3.2	7.5	33.7
1999.....	*	*	*	*	0.2	0.4	1.2	3.3	8.7	37.7

All races 3/, male

2004.....	*	*	*	*	*	0.5	1.0	2.7	6.4	19.4
2003.....	*	*	*	*	*	0.5	1.0	3.1	6.7	20.8
2002.....	*	*	*	*	*	0.7	1.3	2.4	6.9	20.1
2001.....	*	*	*	*	*	0.7	1.3	3.0	7.7	23.8
2000.....	*	*	*	*	*	0.6	1.4	3.4	6.4	28.4
1999.....	*	*	*	*	*	0.5	1.3	3.6	9.0	29.8

All races 3/, female

2004.....	*	*	*	*	*	*	0.6	1.9	5.7	23.6
2003.....	*	*	*	*	*	*	1.0	2.3	5.5	26.2
2002.....	*	*	*	*	*	*	1.1	2.4	6.4	29.6
2001.....	*	*	*	*	*	*	1.0	2.9	7.9	31.6
2000.....	*	*	*	*	*	*	1.1	3.1	8.2	35.8
1999.....	*	*	*	*	*	*	1.1	3.1	8.6	40.9

White, both sexes

2004.....	*	*	*	*	0.2	0.4	0.9	2.3	6.1	23.3
2003.....	*	*	*	*	0.2	0.3	1.0	2.7	6.1	25.1
2002.....	*	*	*	*	*	0.4	1.1	2.5	6.7	27.5
2001.....	*	*	*	*	*	0.5	1.1	2.9	8.0	30.2
2000.....	*	*	*	0.2	0.2	0.5	1.3	3.3	7.5	34.7
1999.....	*	*	*	*	0.2	0.4	1.1	3.2	8.8	38.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified organic mental disorders (F04-F09)

Black, female

2004.....	*	*	*	*	*	*	*	*	*	16.5
2003.....	*	*	*	*	*	*	*	*	*	20.7
2002.....	*	*	*	*	*	*	*	*	*	25.1
2001.....	*	*	*	*	*	*	*	*	*	24.8
2000.....	*	*	*	*	*	*	*	*	9.8	25.3
1999.....	*	*	*	*	*	*	*	*	8.6	28.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

All races 3/, both sexes

2004.....	5.6	8.0	8.6	7.7	7.2	6.7	6.1	5.5	5.3	4.6
2003.....	5.9	7.8	8.1	6.9	7.1	6.2	5.7	4.9	4.4	4.3
2002.....	6.0	7.3	7.6	6.9	6.7	6.1	5.3	4.7	3.8	3.7
2001.....	5.5	7.4	7.0	6.4	6.6	5.7	5.1	4.6	4.5	4.0
2000.....	5.5	7.0	6.5	6.4	6.0	5.8	5.1	4.7	4.5	4.1
1999.....	5.7	6.9	6.2	6.2	6.6	5.8	5.0	5.3	4.4	3.8

All races 3/, male

2004.....	8.4	11.7	13.8	12.8	12.2	10.8	10.1	9.0	8.8	8.2
2003.....	8.8	12.0	13.0	11.2	11.5	10.2	9.1	8.4	7.4	7.3
2002.....	8.8	11.2	12.6	11.5	11.2	10.4	8.1	7.8	6.2	7.0
2001.....	8.2	11.6	11.5	10.4	11.1	9.7	8.6	7.2	7.3	7.5
2000.....	8.1	11.1	10.7	10.4	10.1	9.7	8.3	8.0	7.9	7.3
1999.....	9.0	11.0	10.2	10.2	11.1	9.5	8.1	9.0	8.4	7.4

All races 3/, female

2004.....	2.8	4.3	3.6	2.9	2.6	3.1	2.9	2.9	3.1	3.0
2003.....	3.1	3.7	3.3	2.9	3.1	2.9	2.9	2.5	2.5	3.0
2002.....	3.2	3.4	2.8	2.6	2.5	2.4	3.1	2.5	2.4	2.2
2001.....	2.9	3.4	2.8	2.7	2.6	2.3	2.3	2.9	2.9	2.5
2000.....	2.9	3.0	2.5	2.7	2.3	2.4	2.6	2.5	2.6	2.8
1999.....	2.5	3.1	2.3	2.5	2.5	2.6	2.5	2.7	2.0	2.4

White, both sexes

2004.....	5.6	7.7	8.0	7.3	6.7	6.2	5.8	5.4	5.5	4.7
2003.....	5.7	7.5	7.4	6.4	6.4	6.2	5.6	5.0	4.5	4.5
2002.....	5.8	6.8	6.9	6.3	5.9	5.8	5.1	4.7	3.8	3.6
2001.....	5.0	6.7	6.2	5.7	6.1	5.3	5.0	4.6	4.6	4.0
2000.....	5.0	6.2	5.9	5.8	5.4	5.3	5.0	4.7	4.6	4.1
1999.....	5.1	6.0	5.4	5.4	6.0	5.4	4.7	5.1	4.3	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

White, male										
2004.....	8.4	11.5	12.7	12.0	11.1	9.6	9.3	8.7	9.3	8.2
2003.....	8.6	11.5	11.7	10.4	10.0	9.9	8.8	8.3	7.4	7.4
2002.....	8.7	10.6	11.5	10.2	9.8	9.4	7.4	7.7	5.9	6.8
2001.....	7.5	10.4	10.1	9.0	10.0	8.7	8.1	6.9	7.2	7.3
2000.....	7.6	9.8	9.6	9.3	9.0	8.7	7.9	7.6	7.8	7.2
1999.....	8.2	9.7	9.0	8.6	10.1	8.4	7.4	8.4	8.3	7.4
White, female										
2004.....	2.7	3.9	3.5	2.8	2.6	3.2	3.0	3.0	3.2	3.1
2003.....	2.8	3.6	3.2	2.6	3.1	2.9	2.9	2.6	2.7	3.2
2002.....	2.8	3.1	2.5	2.5	2.3	2.5	3.2	2.6	2.5	2.2
2001.....	2.6	3.0	2.3	2.6	2.5	2.4	2.4	3.0	2.9	2.6
2000.....	2.5	2.5	2.2	2.4	2.2	2.4	2.6	2.6	2.7	2.9
1999.....	2.0	2.5	1.8	2.3	2.3	2.7	2.5	2.7	2.0	2.5
Black, both sexes										
2004.....	6.0	11.8	14.6	12.7	12.5	11.7	9.9	5.9	*	*
2003.....	7.9	11.2	14.2	12.6	13.8	8.4	7.4	5.3	*	*
2002.....	8.6	11.9	14.2	14.3	13.5	9.6	8.3	5.8	*	*
2001.....	9.6	14.5	15.0	13.2	12.7	10.0	7.4	6.4	*	*
2000.....	9.0	14.6	13.1	12.9	12.4	11.0	7.5	6.9	*	*
1999.....	10.2	14.3	13.1	14.9	12.4	10.8	9.5	8.0	6.1	*
Black, male										
2004.....	9.1	16.7	24.6	23.8	24.9	23.7	20.1	13.6	*	*
2003.....	10.9	17.9	25.4	20.8	27.2	15.5	14.1	11.4	*	*
2002.....	11.5	18.5	24.8	26.5	25.3	20.7	16.7	11.5	*	*
2001.....	14.0	23.1	25.3	24.2	24.8	20.6	15.1	13.9	*	*
2000.....	12.7	23.6	22.9	23.1	22.8	21.6	15.0	15.6	*	*
1999.....	15.0	22.3	21.6	27.6	22.9	21.6	19.0	16.6	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

Black, female

2004.....	3.3	7.6	6.2	3.7	2.9	*	*	*	*	*
2003.....	5.2	5.3	4.7	5.9	3.3	*	*	*	*	*
2002.....	6.1	6.1	5.3	4.3	4.2	*	*	*	*	*
2001.....	5.7	7.0	6.3	4.1	3.2	*	*	*	*	*
2000.....	5.6	6.8	4.8	4.6	4.2	*	*	*	*	*
1999.....	5.9	7.5	5.8	4.6	4.2	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Mental and behavioral disorders due to use of alcohol (F10)

All races 3/, both sexes

2004.....	3.5	5.4	6.3	6.2	5.9	5.3	4.4	3.4	2.9	1.9
2003.....	3.9	5.6	6.1	5.7	6.0	4.9	4.1	3.1	2.6	2.0
2002.....	3.9	5.3	6.0	5.9	5.8	5.1	4.1	3.5	2.4	2.0
2001.....	3.7	5.4	5.6	5.5	5.9	5.1	4.1	3.3	3.0	2.1
2000.....	3.6	5.1	5.2	5.5	5.3	5.0	4.1	3.4	2.9	2.5
1999.....	3.6	5.1	5.0	5.3	5.7	4.9	3.8	3.8	2.9	2.3

All races 3/, male

2004.....	5.6	8.5	10.4	10.5	10.3	8.9	8.0	6.4	5.3	3.6
2003.....	6.1	8.7	10.1	9.4	10.0	8.2	7.1	5.6	5.4	3.5
2002.....	6.1	8.3	10.1	9.9	10.0	9.0	6.8	6.4	4.5	4.4
2001.....	5.7	8.6	9.2	9.0	10.0	8.8	7.3	5.8	5.4	4.7
2000.....	5.7	8.3	8.6	9.2	9.1	8.8	7.2	6.1	5.9	5.2
1999.....	5.9	8.1	8.4	8.9	9.7	8.4	6.7	7.2	6.2	4.9

All races 3/, female

2004.....	1.5	2.5	2.4	2.2	2.0	2.2	1.5	1.3	1.4	1.2
2003.....	1.8	2.5	2.4	2.2	2.5	2.0	1.7	1.3	0.9	1.3
2002.....	1.9	2.3	2.1	2.0	1.9	1.6	1.9	1.4	1.1	1.0
2001.....	1.8	2.3	2.1	2.2	2.1	1.9	1.5	1.6	1.6	1.1
2000.....	1.6	2.0	1.9	2.1	1.8	1.8	1.6	1.5	1.2	1.5
1999.....	1.5	2.1	1.7	2.0	2.1	2.0	1.6	1.5	1.0	1.3

White, both sexes

2004.....	3.7	5.6	6.3	6.1	5.6	4.8	4.0	3.4	3.0	2.0
2003.....	4.0	5.6	5.9	5.5	5.6	4.7	3.9	3.0	2.5	2.0
2002.....	4.0	5.3	5.8	5.4	5.1	4.7	3.8	3.4	2.3	2.0
2001.....	3.5	5.2	5.1	4.9	5.5	4.6	3.9	3.2	3.0	2.1
2000.....	3.5	4.7	4.9	5.1	4.7	4.5	3.9	3.2	2.9	2.5
1999.....	3.5	4.7	4.5	4.7	5.2	4.5	3.5	3.6	2.8	2.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Mental and behavioral disorders due to use of alcohol (F10)

White, male

2004.....	5.8	8.6	10.2	10.2	9.5	7.7	7.2	6.1	5.6	3.7
2003.....	6.2	8.6	9.4	9.1	8.9	7.9	6.7	5.3	5.2	3.4
2002.....	6.2	8.2	9.8	9.0	8.9	8.2	6.0	6.3	4.2	4.1
2001.....	5.3	8.2	8.4	7.9	9.1	7.7	6.8	5.5	5.4	4.4
2000.....	5.5	7.7	8.0	8.3	8.0	7.8	6.7	5.6	5.7	5.1
1999.....	5.6	7.6	7.7	7.7	8.7	7.3	5.9	6.7	5.9	4.7

White, female

2004.....	1.5	2.6	2.5	2.2	2.0	2.2	1.4	1.4	1.4	1.2
2003.....	1.8	2.6	2.5	2.1	2.5	2.0	1.6	1.4	0.9	1.4
2002.....	1.8	2.4	2.0	2.0	1.7	1.7	1.9	1.4	1.1	1.1
2001.....	1.7	2.2	1.9	2.1	2.0	1.9	1.5	1.6	1.6	1.1
2000.....	1.5	1.8	1.8	1.9	1.6	1.7	1.6	1.6	1.2	1.5
1999.....	1.3	1.9	1.4	1.9	1.9	2.0	1.6	1.4	0.9	1.3

Black, both sexes

2004.....	2.7	5.6	7.8	8.7	9.6	10.0	9.0	4.6	*	*
2003.....	3.6	6.0	8.4	8.4	11.1	7.0	6.2	4.8	*	*
2002.....	4.1	6.2	8.4	10.9	11.2	8.3	7.9	5.0	*	*
2001.....	5.2	7.9	10.3	11.0	11.0	9.6	6.3	5.2	*	*
2000.....	4.7	8.8	8.7	10.3	11.1	10.2	6.8	6.0	*	*
1999.....	4.9	7.9	8.9	11.6	11.3	9.6	8.4	7.1	*	*

Black, male

2004.....	4.4	9.4	14.0	16.8	19.2	20.4	17.8	10.5	*	*
2003.....	5.6	10.1	16.1	14.3	22.4	12.6	12.5	10.5	*	*
2002.....	5.9	10.4	14.6	20.7	21.0	18.0	16.4	9.6	*	*
2001.....	8.4	13.4	17.4	20.6	21.6	19.6	13.5	11.6	*	*
2000.....	7.3	14.5	15.7	18.9	21.3	20.3	14.4	14.6	*	*
1999.....	7.5	12.9	15.2	21.9	21.0	19.5	17.3	14.2	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Mental and behavioral disorders due to use of alcohol (F10)

Black, female

2004.....	*	2.4	2.6	*	*	*	*	*	*	*
2003.....	1.9	2.4	1.8	3.6	*	*	*	*	*	*
2002.....	2.4	2.5	3.2	2.8	3.6	*	*	*	*	*
2001.....	2.4	3.1	4.2	3.1	*	*	*	*	*	*
2000.....	2.3	3.8	2.8	3.2	*	*	*	*	*	*
1999.....	2.5	3.7	3.6	3.2	3.7	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

All races 3/, both sexes

2004.....	2.1	2.5	2.3	1.5	1.2	1.4	1.7	2.0	2.4	2.7
2003.....	2.0	2.2	1.9	1.2	1.0	1.4	1.6	1.8	1.8	2.3
2002.....	2.0	2.0	1.6	1.1	0.9	1.0	1.2	1.2	1.4	1.7
2001.....	1.8	2.1	1.4	0.9	0.7	0.7	1.0	1.3	1.5	1.8
2000.....	1.8	1.9	1.3	0.9	0.8	0.8	1.0	1.4	1.6	1.6
1999.....	2.1	1.9	1.2	0.9	0.9	0.9	1.2	1.4	1.5	1.5

All races 3/, male

2004.....	2.8	3.3	3.4	2.3	1.9	1.9	2.1	2.7	3.5	4.6
2003.....	2.7	3.3	3.0	1.8	1.5	2.0	2.0	2.8	2.0	3.8
2002.....	2.8	2.9	2.6	1.6	1.2	1.3	1.3	1.4	1.7	2.6
2001.....	2.5	3.0	2.3	1.3	1.0	0.9	1.3	1.4	1.9	2.8
2000.....	2.4	2.8	2.1	1.2	1.0	0.9	1.1	1.9	2.0	2.1
1999.....	3.1	2.8	1.8	1.3	1.4	1.1	1.5	1.8	2.3	2.5

All races 3/, female

2004.....	1.3	1.8	1.3	0.7	0.6	0.9	1.4	1.5	1.7	1.8
2003.....	1.3	1.2	1.0	0.7	0.6	0.8	1.2	1.1	1.6	1.7
2002.....	1.3	1.1	0.7	0.6	0.6	0.7	1.2	1.1	1.3	1.2
2001.....	1.2	1.1	0.6	0.5	0.5	0.4	0.9	1.3	1.3	1.4
2000.....	1.3	1.0	0.6	0.6	0.6	0.6	1.0	1.0	1.4	1.3
1999.....	1.0	1.0	0.6	0.5	0.4	0.7	0.9	1.2	1.0	1.1

White, both sexes

2004.....	1.9	2.1	1.7	1.2	1.1	1.4	1.8	2.1	2.5	2.7
2003.....	1.8	1.9	1.5	0.9	0.9	1.4	1.7	2.0	1.9	2.5
2002.....	1.8	1.6	1.1	0.9	0.8	1.0	1.3	1.3	1.5	1.6
2001.....	1.5	1.5	1.1	0.8	0.7	0.7	1.1	1.4	1.5	1.9
2000.....	1.6	1.4	1.0	0.7	0.8	0.8	1.1	1.4	1.7	1.6
1999.....	1.6	1.3	0.9	0.6	0.9	0.9	1.2	1.5	1.6	1.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

White, male

2004.....	2.7	2.9	2.5	1.8	1.6	1.8	2.2	2.6	3.7	4.5
2003.....	2.5	2.9	2.3	1.3	1.2	2.0	2.1	3.0	2.2	4.0
2002.....	2.5	2.4	1.7	1.2	0.9	1.2	1.4	1.4	1.8	2.7
2001.....	2.2	2.2	1.7	1.1	0.9	0.9	1.3	1.3	1.9	2.9
2000.....	2.1	2.1	1.6	0.9	1.0	0.9	1.1	2.0	2.1	2.1
1999.....	2.6	2.1	1.3	0.9	1.4	1.1	1.5	1.7	2.4	2.7

White, female

2004.....	1.2	1.3	1.0	0.6	0.6	1.0	1.6	1.7	1.8	2.0
2003.....	1.0	1.0	0.7	0.5	0.6	0.9	1.3	1.2	1.8	1.8
2002.....	1.0	0.7	0.5	0.5	0.6	0.8	1.3	1.2	1.4	1.1
2001.....	0.9	0.8	0.4	0.4	0.5	0.5	0.9	1.4	1.3	1.5
2000.....	1.0	0.8	0.4	0.5	0.5	0.7	1.0	1.0	1.5	1.4
1999.....	0.7	0.6	0.4	0.4	*	0.7	1.0	1.3	1.1	1.1

Black, both sexes

2004.....	3.4	6.2	6.8	4.1	2.9	*	*	*	*	*
2003.....	4.3	5.2	5.8	4.2	2.7	*	*	*	*	*
2002.....	4.6	5.7	5.8	3.4	2.3	*	*	*	*	*
2001.....	4.4	6.6	4.7	2.2	*	*	*	*	*	*
2000.....	4.3	5.8	4.4	2.6	*	*	*	*	*	*
1999.....	5.3	6.4	4.2	3.4	*	*	*	*	*	*

Black, male

2004.....	4.6	7.3	10.6	7.0	5.6	*	*	*	*	*
2003.....	5.3	7.7	9.3	6.5	4.8	*	*	*	*	*
2002.....	5.6	8.1	10.2	5.8	4.4	*	*	*	*	*
2001.....	5.6	9.7	7.9	3.6	*	*	*	*	*	*
2000.....	5.5	9.1	7.2	4.1	*	*	*	*	*	*
1999.....	7.5	9.3	6.4	5.8	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)										
Black, female										
2004.....	2.2	5.2	3.6	*	*	*	*	*	*	*
2003.....	3.3	2.9	2.9	2.3	*	*	*	*	*	*
2002.....	3.7	3.7	2.1	*	*	*	*	*	*	*
2001.....	3.3	3.9	2.0	*	*	*	*	*	*	*
2000.....	3.3	3.0	2.0	*	*	*	*	*	*	*
1999.....	3.4	3.8	2.3	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Schizophrenia, schizotypal and delusional disorders (F20-F29)

All races 3/, both sexes

2004.....	0.1	0.1	0.2	0.2	0.2	0.4	0.6	0.9	1.3	2.7
2003.....	*	0.1	0.2	*	0.2	0.3	0.8	0.9	1.5	2.7
2002.....	0.1	0.1	*	0.2	*	0.3	0.7	1.0	1.3	3.3
2001.....	*	0.1	*	0.1	0.2	0.3	0.4	1.1	1.3	2.7
2000.....	*	*	*	*	0.2	0.3	0.7	0.9	1.7	3.3
1999.....	*	*	*	*	0.2	0.3	0.6	1.0	2.1	4.9

All races 3/, male

2004.....	*	0.2	*	0.2	*	*	0.7	0.9	1.1	2.3
2003.....	*	*	0.2	*	*	*	0.6	0.6	1.1	1.8
2002.....	*	*	*	*	*	*	0.7	0.9	*	1.9
2001.....	*	*	*	*	*	0.5	0.5	0.9	1.1	1.5
2000.....	*	*	*	*	*	*	0.7	0.7	1.6	2.4
1999.....	*	*	*	*	*	*	0.7	1.0	1.4	4.6

All races 3/, female

2004.....	*	*	*	*	*	*	0.6	0.9	1.4	2.8
2003.....	*	*	*	*	*	*	0.9	1.0	1.7	3.0
2002.....	*	*	*	*	*	*	0.6	1.1	1.6	3.9
2001.....	*	*	*	*	*	*	*	1.2	1.4	3.1
2000.....	*	*	*	*	*	*	0.7	1.0	1.8	3.8
1999.....	*	*	*	*	*	*	0.5	1.0	2.5	5.0

White, both sexes

2004.....	*	0.1	0.1	0.2	0.2	0.3	0.6	0.9	1.3	2.6
2003.....	*	*	0.1	*	*	0.4	0.8	0.9	1.4	2.8
2002.....	*	*	*	0.2	*	0.3	0.6	1.0	1.3	3.4
2001.....	*	*	*	*	0.2	0.3	0.4	1.1	1.3	2.8
2000.....	*	*	*	*	*	0.4	0.7	0.9	1.8	3.4
1999.....	*	*	*	*	*	0.3	0.6	1.0	2.1	5.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Mood (affective) disorders (F30-F39)

All races 3/, both sexes

2004.....	*	*	*	0.1	*	0.2	0.7	1.4	2.6	8.4
2003.....	*	*	*	0.1	*	0.2	0.5	1.5	2.9	10.5
2002.....	*	*	*	*	0.2	0.2	0.7	1.3	3.5	9.1
2001.....	0.1	*	*	*	*	0.4	0.6	1.4	3.2	9.6
2000.....	*	*	*	*	*	0.3	0.5	1.2	2.9	9.8
1999.....	*	*	*	*	*	0.3	0.6	1.6	3.4	9.4

All races 3/, male

2004.....	*	*	*	*	*	*	0.7	1.0	2.1	7.2
2003.....	*	*	*	*	*	*	*	1.5	2.4	8.0
2002.....	*	*	*	*	*	*	0.5	0.9	2.6	6.6
2001.....	*	*	*	*	*	*	0.6	1.3	2.4	8.0
2000.....	*	*	*	*	*	*	0.5	0.9	2.3	9.1
1999.....	*	*	*	*	*	*	*	1.6	2.7	8.8

All races 3/, female

2004.....	*	*	*	*	*	*	0.6	1.6	3.0	9.0
2003.....	*	*	*	*	*	*	0.5	1.6	3.3	11.6
2002.....	*	*	*	*	*	*	0.7	1.6	4.0	10.1
2001.....	*	*	*	*	*	*	0.6	1.5	3.7	10.3
2000.....	*	*	*	*	*	*	0.5	1.5	3.2	10.1
1999.....	*	*	*	*	*	*	0.6	1.7	3.9	9.7

White, both sexes

2004.....	*	*	*	*	*	0.3	0.7	1.4	2.8	9.2
2003.....	*	*	*	*	*	0.2	0.5	1.6	3.1	11.3
2002.....	*	*	*	*	0.2	0.2	0.7	1.4	3.6	9.8
2001.....	*	*	*	*	*	0.4	0.6	1.6	3.4	10.1
2000.....	*	*	*	*	*	0.3	0.6	1.3	3.0	10.6
1999.....	*	*	*	*	*	0.3	0.6	1.8	3.7	10.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Mental retardation (F70-F79)

All races 3/, both sexes

2004.....	0.2	0.2	0.3	0.2	0.4	0.3	0.4	0.5	0.4	*
2003.....	0.1	0.2	0.3	0.4	0.4	0.4	0.3	0.4	0.4	0.4
2002.....	0.2	0.2	0.2	0.4	0.4	0.4	0.4	0.3	0.5	0.5
2001.....	0.1	0.2	0.2	0.3	0.3	0.3	0.4	0.4	0.5	*
2000.....	0.1	0.2	0.2	0.3	0.3	0.2	0.3	0.4	*	0.5
1999.....	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.5	*

All races 3/, male

2004.....	0.2	0.3	0.4	*	0.5	0.4	0.7	*	*	*
2003.....	0.2	0.3	0.3	0.5	0.4	*	*	*	*	*
2002.....	0.2	0.3	0.3	0.5	0.4	*	*	*	*	*
2001.....	*	0.2	0.2	*	0.5	*	*	*	*	*
2000.....	*	0.2	0.2	0.3	*	*	*	*	*	*
1999.....	0.2	0.2	0.3	0.3	*	0.5	*	*	*	*

All races 3/, female

2004.....	*	0.2	*	0.3	0.3	*	*	0.5	*	*
2003.....	*	*	0.2	0.3	0.3	0.4	*	*	*	*
2002.....	*	0.2	*	0.3	0.3	*	*	*	*	*
2001.....	*	*	0.2	0.3	*	*	*	*	*	*
2000.....	*	*	*	0.3	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	0.3	0.3	0.3	0.4	0.3	0.5	0.5	0.4	*
2003.....	0.2	0.2	0.3	0.4	0.4	0.4	0.3	0.4	0.4	0.5
2002.....	0.2	0.2	0.2	0.4	0.4	0.4	0.4	*	0.6	0.6
2001.....	0.1	0.2	0.2	0.3	0.3	0.3	0.4	0.3	0.6	*
2000.....	0.1	0.2	0.2	0.4	0.3	0.2	0.3	0.4	*	0.5
1999.....	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.6	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	0.5	1.0	3.9
2003.....	*	0.1	*	*	*	*	0.3	0.7	1.1	4.4
2002.....	*	*	*	*	*	*	0.3	0.4	1.2	4.4
2001.....	*	*	*	*	*	*	*	0.5	1.0	4.1
2000.....	*	*	*	*	*	*	*	0.4	1.2	3.9
1999.....	*	*	*	*	*	*	0.4	0.4	1.0	4.0

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	1.1	3.7
2003.....	*	*	*	*	*	*	*	0.7	1.3	4.2
2002.....	*	*	*	*	*	*	*	*	1.6	3.6
2001.....	*	*	*	*	*	*	*	*	*	3.5
2000.....	*	*	*	*	*	*	*	*	1.3	3.3
1999.....	*	*	*	*	*	*	*	*	*	2.5

All races 3/, female

2004.....	*	*	*	*	*	*	*	0.6	1.0	4.0
2003.....	*	*	*	*	*	*	*	0.7	1.0	4.5
2002.....	*	*	*	*	*	*	*	*	1.0	4.7
2001.....	*	*	*	*	*	*	*	0.5	1.0	4.4
2000.....	*	*	*	*	*	*	*	0.5	1.1	4.1
1999.....	*	*	*	*	*	*	*	*	1.1	4.7

White, both sexes

2004.....	*	*	*	*	*	*	*	0.5	1.1	4.0
2003.....	*	*	*	*	*	*	0.3	0.7	1.1	4.4
2002.....	*	*	*	*	*	*	*	0.4	1.3	4.4
2001.....	*	*	*	*	*	*	*	0.5	1.0	4.4
2000.....	*	*	*	*	*	*	*	0.4	1.2	4.0
1999.....	*	*	*	*	*	*	0.4	0.5	1.1	4.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

VI. Diseases of the nervous system (G00-G98)

All races 3/, both sexes

2004.....	4.9	7.6	11.0	16.3	25.0	42.3	83.4	190.8	405.4	1,014.7
2003.....	5.0	7.7	11.0	16.2	25.7	43.0	88.3	193.1	400.0	1,006.7
2002.....	5.1	7.8	10.5	15.7	24.9	42.7	83.2	185.2	388.8	951.0
2001.....	5.0	7.3	10.3	15.7	24.0	41.0	81.1	178.4	375.0	906.2
2000.....	5.0	7.3	11.0	15.6	23.8	40.2	79.5	172.9	361.9	864.9
1999.....	4.6	7.0	10.3	14.4	22.1	39.3	76.3	163.6	340.4	790.8

All races 3/, male

2004.....	5.6	8.4	12.0	17.5	26.9	47.0	95.4	211.4	441.6	910.7
2003.....	6.0	8.6	12.2	17.5	27.9	47.2	100.6	218.6	436.2	907.9
2002.....	5.7	8.5	11.2	16.5	27.4	48.2	95.7	210.0	422.2	867.4
2001.....	5.5	7.7	10.9	16.2	25.7	47.4	92.9	206.0	421.1	861.6
2000.....	5.7	8.3	11.8	16.5	25.9	46.5	92.8	198.3	407.5	826.2
1999.....	5.1	7.2	11.0	15.9	24.4	44.9	89.1	193.7	394.2	772.7

All races 3/, female

2004.....	4.2	6.8	10.1	15.2	23.3	38.3	73.8	176.0	383.1	1,061.5
2003.....	4.1	6.9	9.9	15.0	23.7	39.4	78.4	175.0	377.9	1,050.4
2002.....	4.4	7.0	9.9	14.9	22.6	37.8	73.2	167.7	368.5	987.3
2001.....	4.5	6.9	9.7	15.2	22.5	35.5	71.7	159.1	347.4	924.9
2000.....	4.4	6.4	10.3	14.8	21.8	34.9	69.0	155.1	335.0	880.7
1999.....	4.1	6.7	9.6	13.0	20.1	34.6	66.3	142.7	308.9	798.2

White, both sexes

2004.....	4.8	7.4	10.9	16.3	25.3	43.2	86.9	197.4	419.4	1,051.0
2003.....	4.9	7.4	10.7	16.3	25.9	44.0	91.4	200.1	416.1	1,047.3
2002.....	4.9	7.5	10.2	15.7	25.4	43.8	85.9	192.1	405.4	989.7
2001.....	4.8	7.0	10.2	15.5	24.3	42.0	84.1	184.4	389.0	942.3
2000.....	4.8	7.0	10.9	15.6	24.1	41.2	81.7	179.1	375.6	900.8
1999.....	4.4	6.6	10.1	14.3	22.2	40.1	78.1	169.0	353.7	825.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

VI. Diseases of the nervous system (G00-G98)

White, male

2004.....	5.6	8.2	11.8	17.4	27.3	48.0	99.2	218.0	459.2	945.7
2003.....	5.9	8.3	11.9	17.6	27.7	48.0	104.2	226.5	452.6	944.4
2002.....	5.5	8.2	10.7	16.4	28.0	49.0	98.4	217.7	438.9	904.3
2001.....	5.3	7.5	10.7	16.1	26.0	48.4	95.9	213.3	436.5	896.4
2000.....	5.3	8.1	11.9	16.3	26.0	47.8	95.3	205.0	422.6	859.9
1999.....	4.7	6.9	10.7	15.9	24.4	45.7	91.7	200.0	411.2	806.2

White, female

2004.....	4.0	6.5	10.0	15.2	23.5	39.0	76.7	182.3	394.7	1,098.4
2003.....	3.8	6.6	9.6	15.0	24.2	40.4	81.0	181.0	393.5	1,092.9
2002.....	4.3	6.9	9.7	15.1	22.9	39.2	75.8	173.7	384.8	1,026.8
2001.....	4.4	6.6	9.7	15.0	22.7	36.3	74.5	163.9	360.3	961.5
2000.....	4.3	5.9	10.0	14.9	22.3	35.4	70.7	160.9	347.7	917.4
1999.....	4.1	6.3	9.4	12.8	20.1	35.3	67.3	147.1	319.8	832.7

Black, both sexes

2004.....	6.9	11.1	14.8	21.4	30.2	45.2	72.9	166.6	327.0	776.9
2003.....	7.3	11.9	16.1	20.5	30.8	46.5	78.6	159.4	295.2	699.3
2002.....	7.7	11.7	16.0	18.8	28.2	43.4	76.3	150.1	269.2	655.4
2001.....	7.4	10.7	14.9	21.3	27.3	42.2	69.6	152.0	273.6	624.9
2000.....	7.9	12.1	15.4	20.0	26.2	41.6	73.1	136.1	259.8	561.2
1999.....	7.1	11.7	15.6	18.6	26.3	40.1	73.7	131.6	225.9	485.3

Black, male

2004.....	7.5	12.2	17.0	23.8	31.8	49.2	83.8	183.3	327.8	661.6
2003.....	8.1	13.0	17.9	21.5	37.0	53.9	89.7	180.3	324.8	620.4
2002.....	8.5	13.5	17.1	20.5	30.7	52.1	93.8	168.1	303.4	570.4
2001.....	8.5	11.5	16.5	22.1	29.5	49.3	81.8	170.5	302.6	593.4
2000.....	9.6	12.2	15.3	23.9	30.6	46.6	86.9	159.7	285.8	537.4
1999.....	8.9	12.3	18.1	19.3	29.7	46.2	81.1	152.4	231.1	481.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

VI. Diseases of the nervous system (G00-G98)

Black, female

2004.....	6.3	10.3	13.0	19.6	29.0	42.3	65.5	156.6	326.6	822.6
2003.....	6.6	10.9	14.7	19.7	25.9	40.9	71.1	147.0	279.7	730.0
2002.....	7.0	10.2	15.0	17.4	26.2	37.0	64.5	139.4	251.5	688.0
2001.....	6.4	10.0	13.6	20.7	25.6	37.0	61.4	140.9	258.8	636.5
2000.....	6.4	11.9	15.5	16.9	22.7	38.0	63.9	121.9	246.7	570.1
1999.....	5.6	11.2	13.5	18.0	23.6	35.6	68.9	119.0	223.4	486.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningitis (G00,G03)										
All races 3/, both sexes										
2004.....	0.2	0.3	0.3	0.3	0.3	0.5	0.5	0.6	0.7	0.9
2003.....	0.2	0.3	0.3	0.3	0.4	0.6	0.5	0.7	0.6	1.1
2002.....	0.2	0.3	0.3	0.3	0.3	0.5	0.7	0.6	0.7	1.2
2001.....	0.2	0.3	0.3	0.4	0.5	0.5	0.5	0.6	1.0	0.9
2000.....	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1	1.2
1999.....	0.2	0.3	0.3	0.3	0.5	0.5	0.7	0.9	1.2	1.5
All races 3/, male										
2004.....	0.3	0.3	0.4	0.3	0.3	*	*	*	*	*
2003.....	*	0.3	0.4	0.4	0.5	0.6	*	0.8	*	*
2002.....	0.2	0.4	0.3	0.4	*	0.6	0.9	0.7	1.0	*
2001.....	*	0.3	0.3	0.4	0.6	0.5	0.7	0.9	1.1	*
2000.....	0.2	0.3	0.4	0.4	0.5	0.6	0.9	1.3	1.1	*
1999.....	0.2	0.4	0.4	0.4	0.4	0.6	0.8	0.9	1.3	1.7
All races 3/, female										
2004.....	*	*	0.3	0.3	0.3	0.6	0.6	0.6	0.6	0.7
2003.....	*	0.2	0.2	0.3	0.3	0.6	0.4	0.7	0.6	1.0
2002.....	*	0.2	0.2	*	0.3	0.4	0.6	0.6	*	1.1
2001.....	*	0.2	0.2	0.4	0.4	0.4	0.4	0.5	0.9	0.8
2000.....	*	0.2	0.3	*	*	0.4	0.5	0.6	1.1	1.0
1999.....	*	0.2	*	*	0.5	0.4	0.7	0.9	1.1	1.4
White, both sexes										
2004.....	0.2	0.2	0.2	0.2	0.3	0.4	0.4	0.6	0.6	0.8
2003.....	0.1	0.2	0.2	0.3	0.3	0.5	0.4	0.7	0.6	1.0
2002.....	0.1	0.2	0.2	0.3	0.3	0.5	0.7	0.6	0.6	1.2
2001.....	0.1	0.2	0.2	0.3	0.5	0.4	0.6	0.6	0.9	0.8
2000.....	0.1	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1	1.0
1999.....	0.1	0.2	0.2	0.3	0.4	0.5	0.6	0.7	1.1	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other inflammatory diseases of central nervous system (G04-G09)

All races 3/, both sexes

2004.....	0.2	0.2	0.2	0.3	0.4	0.4	0.8	1.0	1.3	1.1
2003.....	0.2	0.2	0.2	0.3	0.4	0.7	0.7	1.0	1.5	1.1
2002.....	0.2	0.2	0.2	0.3	0.4	0.5	0.8	1.1	1.4	1.2
2001.....	0.1	0.2	0.2	0.3	0.4	0.5	0.8	0.9	1.1	1.0
2000.....	0.2	0.1	0.2	0.3	0.3	0.5	0.8	1.1	1.1	1.0
1999.....	0.1	0.2	0.2	0.3	0.3	0.5	0.7	0.7	1.0	1.1

All races 3/, male

2004.....	0.2	0.2	0.2	0.4	0.5	0.6	1.0	1.2	1.8	1.5
2003.....	0.2	0.3	0.3	0.4	0.5	0.8	0.9	1.3	1.7	*
2002.....	0.2	0.3	0.3	0.4	0.5	0.8	1.0	1.3	2.0	*
2001.....	*	0.2	0.3	0.3	0.4	0.7	0.8	1.1	1.3	1.8
2000.....	0.2	*	0.3	0.3	0.4	0.6	1.0	1.6	1.4	*
1999.....	0.2	*	*	0.5	*	0.6	0.6	0.9	1.6	*

All races 3/, female

2004.....	*	*	0.2	*	0.3	*	0.6	0.8	0.9	1.0
2003.....	*	*	*	0.3	0.3	0.6	0.5	0.8	1.4	0.9
2002.....	*	0.2	0.2	0.3	0.3	*	0.6	0.9	1.1	1.1
2001.....	*	0.2	*	0.3	0.4	0.4	0.8	0.8	0.9	0.7
2000.....	*	*	*	0.3	*	0.5	0.6	0.8	0.9	0.9
1999.....	*	*	*	*	0.4	0.4	0.8	0.6	0.7	0.9

White, both sexes

2004.....	0.1	0.2	0.2	0.3	0.4	0.4	0.8	1.0	1.2	1.1
2003.....	0.2	0.2	0.2	0.3	0.4	0.7	0.7	1.0	1.5	1.1
2002.....	0.1	0.2	0.2	0.3	0.4	0.5	0.7	1.0	1.4	1.3
2001.....	0.1	0.2	0.2	0.3	0.4	0.5	0.7	0.9	1.0	1.0
2000.....	0.2	*	0.2	0.3	0.4	0.5	0.7	1.1	1.0	1.0
1999.....	0.1	0.2	0.2	0.3	0.3	0.5	0.6	0.7	1.1	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

All races 3/, both sexes

2004.....	*	*	0.2	0.6	2.0	6.4	18.5	47.6	94.0	135.8
2003.....	*	*	0.2	0.7	2.0	6.4	19.9	49.3	93.3	138.2
2002.....	*	*	0.2	0.6	1.9	6.5	18.4	45.4	89.8	135.2
2001.....	*	*	0.2	0.7	1.8	5.8	18.1	46.4	90.9	134.2
2000.....	*	*	0.2	0.6	1.7	6.6	16.7	44.8	87.6	131.9
1999.....	*	*	0.2	0.6	1.6	5.6	16.7	41.8	82.8	124.4

All races 3/, male

2004.....	*	*	0.4	0.9	2.6	9.1	27.3	70.6	147.1	220.0
2003.....	*	*	0.2	0.8	2.7	8.9	29.2	74.2	140.7	222.4
2002.....	*	*	0.3	0.8	2.5	9.8	27.3	67.2	136.6	220.1
2001.....	*	*	0.3	1.0	2.7	8.2	26.5	68.8	141.7	224.4
2000.....	*	*	*	0.8	2.2	9.0	24.1	67.4	135.0	222.3
1999.....	*	*	0.3	0.7	2.3	7.7	24.8	64.4	131.3	205.6

All races 3/, female

2004.....	*	*	*	0.4	1.4	4.1	11.4	31.0	61.3	97.9
2003.....	*	*	0.2	0.6	1.4	4.3	12.4	31.5	64.3	101.0
2002.....	*	*	*	0.5	1.4	3.8	11.3	30.0	61.3	98.4
2001.....	*	*	*	0.4	1.0	3.7	11.5	30.7	60.4	96.4
2000.....	*	*	*	0.4	1.3	4.4	10.9	29.1	59.5	95.0
1999.....	*	*	*	0.4	0.9	3.9	10.4	26.0	54.4	91.7

White, both sexes

2004.....	*	*	0.2	0.7	2.1	6.8	20.2	50.6	99.9	143.1
2003.....	*	*	0.2	0.7	2.0	6.8	21.2	52.4	99.0	146.0
2002.....	*	*	0.2	0.6	2.1	6.9	19.6	48.3	95.7	143.3
2001.....	*	*	0.2	0.7	1.9	6.2	19.4	49.6	96.4	141.8
2000.....	*	*	0.2	0.6	1.9	7.1	17.8	47.5	92.5	139.5
1999.....	*	*	0.2	0.6	1.6	6.0	17.7	44.1	87.4	131.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Parkinson's disease (G20-G21)

White, male

2004.....	*	*	0.4	1.0	2.7	9.6	29.4	74.7	156.1	231.3
2003.....	*	*	*	0.8	2.7	9.4	30.9	78.4	148.9	233.3
2002.....	*	*	0.3	0.8	2.7	10.2	28.8	71.2	145.0	232.7
2001.....	*	*	0.3	1.0	3.0	8.8	28.1	73.4	149.6	237.6
2000.....	*	*	*	0.9	2.5	9.6	25.5	70.7	141.5	235.3
1999.....	*	*	0.3	0.7	2.4	8.1	26.4	67.7	138.6	216.7

White, female

2004.....	*	*	*	0.4	1.5	4.3	12.6	33.0	65.0	103.3
2003.....	*	*	*	0.6	1.4	4.5	13.3	33.6	68.2	107.4
2002.....	*	*	*	0.5	1.5	3.9	12.0	31.9	65.5	104.4
2001.....	*	*	*	0.5	0.9	4.0	12.3	32.7	64.2	101.7
2000.....	*	*	*	0.4	1.4	4.8	11.7	31.2	63.3	100.6
1999.....	*	*	*	0.4	0.9	4.2	10.8	27.6	57.3	96.8

Black, both sexes

2004.....	*	*	*	*	*	3.9	7.2	22.3	39.1	61.2
2003.....	*	*	*	*	1.9	4.1	10.7	22.8	38.3	52.8
2002.....	*	*	*	*	*	4.6	9.5	20.6	32.7	51.7
2001.....	*	*	*	*	*	2.7	8.9	18.9	34.9	59.0
2000.....	*	*	*	*	*	3.1	8.8	21.0	37.6	51.6
1999.....	*	*	*	*	*	3.6	8.7	19.5	32.4	52.0

Black, male

2004.....	*	*	*	*	*	5.3	12.6	34.1	58.1	106.3
2003.....	*	*	*	*	*	5.4	16.8	37.0	59.1	97.0
2002.....	*	*	*	*	*	7.5	16.0	30.3	55.0	89.0
2001.....	*	*	*	*	*	*	14.8	25.6	57.8	99.8
2000.....	*	*	*	*	*	*	13.7	37.8	65.8	86.5
1999.....	*	*	*	*	*	5.7	12.6	31.8	48.8	86.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Parkinson's disease (G20-G21)

Black, female

2004.....	*	*	*	*	*	*	*	15.4	29.1	43.3
2003.....	*	*	*	*	*	*	6.6	14.4	27.5	35.6
2002.....	*	*	*	*	*	*	5.2	14.8	21.1	37.4
2001.....	*	*	*	*	*	*	4.9	15.0	23.3	43.8
2000.....	*	*	*	*	*	*	5.5	10.8	23.4	38.6
1999.....	*	*	*	*	*	*	6.1	12.0	24.2	39.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alzheimer's disease (G30)

All races 3/, both sexes

2004.....	*	0.2	0.3	0.9	3.1	10.1	31.0	99.4	261.0	818.8
2003.....	*	0.1	0.3	1.0	3.3	10.0	33.3	98.6	255.0	802.4
2002.....	*	*	0.2	0.9	3.2	9.8	30.6	95.4	245.7	752.3
2001.....	*	*	0.3	1.2	3.3	9.0	29.2	88.5	232.5	710.3
2000.....	*	*	0.3	1.1	3.2	9.1	28.9	85.2	221.1	667.7
1999.....	*	0.1	0.2	1.0	3.0	8.7	26.7	78.6	206.2	601.3

All races 3/, male

2004.....	*	*	0.3	0.8	3.1	9.4	29.6	89.3	236.4	619.6
2003.....	*	*	0.3	1.0	2.9	10.1	32.1	92.1	231.2	606.5
2002.....	*	*	0.2	0.8	3.1	9.2	29.5	90.9	220.6	572.0
2001.....	*	*	0.3	0.9	3.2	9.0	27.7	85.6	216.6	556.1
2000.....	*	*	0.3	0.9	2.9	9.5	29.1	79.9	206.8	522.1
1999.....	*	*	*	1.1	3.0	8.8	26.9	77.4	198.2	485.7

All races 3/, female

2004.....	*	0.2	0.4	1.0	3.1	10.7	32.1	106.7	276.2	908.4
2003.....	*	*	0.3	0.9	3.7	10.0	34.2	103.2	269.5	889.0
2002.....	*	*	*	1.0	3.3	10.3	31.5	98.5	260.9	830.5
2001.....	*	*	0.3	1.4	3.4	9.0	30.4	90.5	242.1	774.9
2000.....	*	*	0.3	1.2	3.5	8.7	28.8	88.9	229.5	727.0
1999.....	*	*	0.3	0.8	3.0	8.6	26.6	79.4	210.9	648.0

White, both sexes

2004.....	*	0.2	0.4	1.0	3.3	10.4	32.0	102.3	268.5	846.8
2003.....	*	*	0.3	1.1	3.6	10.3	34.5	101.8	264.4	834.0
2002.....	*	*	0.2	1.0	3.4	10.3	31.9	99.1	255.3	781.8
2001.....	*	*	0.3	1.3	3.6	9.3	30.6	90.9	240.5	738.1
2000.....	*	*	0.3	1.1	3.4	9.6	29.9	88.1	229.1	695.3
1999.....	*	*	0.2	1.0	3.3	9.1	27.6	81.3	213.6	627.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alzheimer's disease (G30)

White, male

2004.....	*	*	0.4	0.9	3.3	9.7	30.2	91.6	243.9	640.9
2003.....	*	*	0.3	1.1	3.0	10.3	33.1	94.9	238.5	630.3
2002.....	*	*	*	0.8	3.3	9.4	30.5	94.3	227.6	594.8
2001.....	*	*	0.3	0.9	3.4	9.2	28.8	88.1	222.7	577.2
2000.....	*	*	0.3	1.0	3.1	10.1	29.9	82.6	214.4	542.1
1999.....	*	*	*	1.1	3.3	9.2	27.6	80.1	205.6	506.9

White, female

2004.....	*	0.3	0.4	1.1	3.3	11.1	33.5	110.2	283.8	939.6
2003.....	*	*	0.4	1.0	4.2	10.3	35.7	106.8	280.3	924.3
2002.....	*	*	*	1.2	3.5	11.0	33.1	102.6	272.2	863.0
2001.....	*	*	0.4	1.6	3.8	9.4	32.1	92.9	251.3	805.4
2000.....	*	*	0.3	1.3	3.7	9.2	29.9	92.0	237.7	757.6
1999.....	*	*	*	0.9	3.3	9.1	27.5	82.1	218.3	676.5

Black, both sexes

2004.....	*	*	*	*	2.2	10.3	30.6	96.0	236.7	657.9
2003.....	*	*	*	*	1.8	10.5	30.2	91.4	207.9	583.8
2002.....	*	*	*	*	2.5	8.5	27.2	81.3	186.8	543.7
2001.....	*	*	*	*	1.9	8.7	23.3	86.8	183.6	506.7
2000.....	*	*	*	*	2.1	7.4	25.9	71.1	171.7	445.1
1999.....	*	*	*	*	*	6.9	25.3	65.0	152.0	372.4

Black, male

2004.....	*	*	*	*	*	9.4	32.0	89.6	210.9	497.7
2003.....	*	*	*	*	*	11.1	32.5	93.1	206.0	459.8
2002.....	*	*	*	*	*	9.5	29.4	82.2	189.5	412.8
2001.....	*	*	*	*	*	9.9	24.2	84.6	185.4	414.1
2000.....	*	*	*	*	*	7.8	27.7	64.7	156.0	371.3
1999.....	*	*	*	*	*	8.0	27.5	62.7	139.5	317.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alzheimer's disease (G30)

Black, female

2004.....	*	*	*	*	*	11.0	29.7	99.7	250.2	721.4
2003.....	*	*	*	*	*	10.1	28.6	90.3	208.9	631.9
2002.....	*	*	*	*	*	7.8	25.8	80.7	185.4	593.9
2001.....	*	*	*	*	*	7.9	22.6	88.2	182.7	541.0
2000.....	*	*	*	*	*	7.1	24.6	74.9	179.6	472.6
1999.....	*	*	*	*	*	6.0	23.9	66.4	158.2	392.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Multiple sclerosis (G35)

All races 3/, both sexes

2004.....	0.7	1.4	2.3	3.2	3.5	3.8	3.7	3.9	3.2	2.6
2003.....	0.8	1.5	2.4	3.1	3.3	3.6	3.6	4.0	3.0	2.2
2002.....	0.8	1.7	2.3	3.0	3.5	3.4	3.5	3.4	3.3	2.2
2001.....	0.9	1.6	2.3	3.0	3.3	3.4	3.3	3.6	3.4	2.1
2000.....	0.9	1.4	2.5	3.0	3.1	3.1	3.1	3.0	3.1	1.7
1999.....	0.9	1.4	2.4	2.4	2.7	3.0	2.8	3.2	2.8	2.1

All races 3/, male

2004.....	0.5	1.0	1.5	2.4	2.4	2.9	2.9	2.8	2.4	1.7
2003.....	0.6	1.1	1.7	2.3	2.7	2.7	3.0	2.8	2.2	1.5
2002.....	0.5	1.1	1.6	2.2	3.1	2.4	2.6	2.7	2.4	1.9
2001.....	0.6	1.2	1.7	2.1	2.3	3.1	2.3	3.1	2.1	1.5
2000.....	0.7	1.1	1.8	2.2	2.3	2.3	2.4	2.4	2.2	1.8
1999.....	0.7	0.8	1.8	1.9	1.8	2.4	2.4	2.3	2.2	*

All races 3/, female

2004.....	0.9	1.8	3.0	4.0	4.6	4.6	4.4	4.6	3.7	3.1
2003.....	0.9	1.9	3.1	3.8	3.8	4.4	4.1	4.8	3.4	2.5
2002.....	1.1	2.3	3.0	3.8	3.8	4.2	4.2	3.9	3.8	2.4
2001.....	1.2	2.0	3.0	3.7	4.1	3.6	4.1	4.0	4.1	2.3
2000.....	1.1	1.8	3.1	3.8	3.8	3.7	3.6	3.5	3.7	1.7
1999.....	1.2	2.0	3.0	2.9	3.5	3.5	3.1	3.8	3.1	2.4

White, both sexes

2004.....	0.7	1.5	2.4	3.4	3.6	4.1	4.1	4.2	3.5	2.9
2003.....	0.8	1.5	2.5	3.3	3.4	3.8	3.8	4.2	3.2	2.3
2002.....	0.8	1.7	2.3	3.2	3.6	3.6	3.8	3.6	3.5	2.4
2001.....	0.9	1.6	2.4	3.2	3.5	3.6	3.6	3.9	3.6	2.2
2000.....	0.9	1.4	2.6	3.1	3.2	3.3	3.3	3.3	3.3	1.9
1999.....	0.9	1.5	2.5	2.7	2.9	3.1	2.9	3.5	3.0	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Multiple sclerosis (G35)

Black, female

2004.....	*	2.2	3.0	4.8	6.0	*	*	*	*	*
2003.....	1.4	2.9	3.1	3.0	3.3	4.0	*	*	*	*
2002.....	2.1	2.9	4.1	3.5	4.0	4.3	*	*	*	*
2001.....	1.6	2.5	3.3	3.5	4.0	*	*	*	*	*
2000.....	1.7	3.1	3.1	3.4	3.9	*	*	*	*	*
1999.....	1.6	1.8	2.8	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Epilepsy (G40-G41)

All races 3/, both sexes

2004.....	0.6	0.6	0.6	0.6	0.6	0.7	0.6	1.1	1.1	1.5
2003.....	0.5	0.7	0.6	0.7	0.6	0.7	0.8	1.0	1.5	1.5
2002.....	0.6	0.6	0.6	0.5	0.6	0.7	0.9	1.3	1.4	1.8
2001.....	0.6	0.5	0.6	0.6	0.6	0.7	0.8	1.1	1.2	1.6
2000.....	0.6	0.6	0.6	0.7	0.6	0.7	1.0	1.0	1.6	2.0
1999.....	0.6	0.5	0.6	0.5	0.6	0.8	1.0	1.1	1.5	1.8

All races 3/, male

2004.....	0.8	0.7	0.8	0.7	0.5	0.7	*	1.2	0.9	1.9
2003.....	0.7	0.8	0.8	0.7	0.6	0.6	0.9	1.0	1.8	1.5
2002.....	0.8	0.8	0.8	0.4	0.7	0.8	0.9	1.2	1.5	2.3
2001.....	0.8	0.6	0.7	0.7	0.7	0.8	0.8	1.0	*	2.0
2000.....	0.8	0.8	0.7	0.8	0.8	0.7	1.0	1.0	*	2.4
1999.....	0.7	0.7	0.6	0.6	0.7	0.9	1.0	0.9	1.3	2.3

All races 3/, female

2004.....	0.4	0.4	0.3	0.6	0.7	0.6	0.7	1.1	1.2	1.4
2003.....	0.4	0.5	0.5	0.6	0.6	0.8	0.8	1.0	1.3	1.6
2002.....	0.4	0.5	0.4	0.5	0.6	0.7	0.9	1.3	1.4	1.5
2001.....	0.5	0.4	0.5	0.5	0.6	0.6	0.7	1.2	1.5	1.4
2000.....	0.4	0.4	0.5	0.5	0.5	0.6	1.0	1.0	2.0	1.9
1999.....	0.4	0.4	0.5	0.5	0.5	0.7	0.9	1.2	1.6	1.7

White, both sexes

2004.....	0.5	0.5	0.5	0.5	0.5	0.6	0.5	0.9	1.1	1.4
2003.....	0.5	0.6	0.6	0.5	0.5	0.6	0.7	0.8	1.3	1.4
2002.....	0.6	0.6	0.5	0.4	0.6	0.6	0.7	1.1	1.3	1.6
2001.....	0.6	0.5	0.5	0.5	0.6	0.6	0.6	0.9	1.0	1.6
2000.....	0.6	0.5	0.6	0.6	0.5	0.6	0.8	0.8	1.5	1.7
1999.....	0.5	0.5	0.5	0.4	0.4	0.6	0.8	0.9	1.4	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Transient cerebral ischemic attacks and related syndromes (G45)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	0.3	0.6	2.1
2003.....	*	*	*	*	*	*	*	0.3	0.7	3.8
2002.....	*	*	*	*	*	*	*	0.4	0.8	2.9
2001.....	*	*	*	*	*	*	*	0.4	0.7	3.8
2000.....	*	*	*	*	*	*	*	0.5	1.0	3.9
1999.....	*	*	*	*	*	*	*	0.4	1.0	4.6

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	1.6
2003.....	*	*	*	*	*	*	*	*	*	2.8
2002.....	*	*	*	*	*	*	*	*	1.0	1.8
2001.....	*	*	*	*	*	*	*	*	*	4.2
2000.....	*	*	*	*	*	*	*	0.7	*	2.9
1999.....	*	*	*	*	*	*	*	0.7	1.2	3.8

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	0.7	2.4
2003.....	*	*	*	*	*	*	*	*	0.6	4.2
2002.....	*	*	*	*	*	*	*	*	0.6	3.4
2001.....	*	*	*	*	*	*	*	*	*	3.6
2000.....	*	*	*	*	*	*	*	*	1.1	4.2
1999.....	*	*	*	*	*	*	*	*	0.9	4.9

White, both sexes

2004.....	*	*	*	*	*	*	*	*	0.6	2.2
2003.....	*	*	*	*	*	*	*	0.3	0.6	4.0
2002.....	*	*	*	*	*	*	*	0.4	0.8	3.0
2001.....	*	*	*	*	*	*	*	0.5	0.8	3.9
2000.....	*	*	*	*	*	*	*	0.5	1.1	4.0
1999.....	*	*	*	*	*	*	*	0.4	1.1	4.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Infantile cerebral palsy (G80)										
All races 3/, both sexes										
2004.....	0.3	0.3	0.4	0.4	0.5	0.5	0.6	0.6	*	*
2003.....	0.2	0.3	0.4	0.4	0.5	0.3	0.5	0.5	0.6	*
2002.....	0.2	0.3	0.3	0.3	0.5	0.4	0.5	0.5	0.6	*
2001.....	0.2	0.2	0.3	0.4	0.4	0.3	0.6	0.6	0.4	*
2000.....	0.2	0.2	0.3	0.3	0.4	0.5	0.4	0.4	0.7	*
1999.....	0.2	0.2	0.2	0.2	0.3	0.3	0.4	0.4	0.5	0.5
All races 3/, male										
2004.....	0.2	0.3	0.5	0.4	0.7	0.7	0.7	0.9	*	*
2003.....	0.3	0.4	0.5	0.4	0.6	*	0.6	0.7	*	*
2002.....	0.2	*	0.4	0.4	0.6	0.5	0.5	*	*	*
2001.....	0.3	0.2	0.4	0.4	0.4	*	0.9	0.8	*	*
2000.....	0.2	0.3	0.3	0.3	*	0.7	0.7	*	*	*
1999.....	0.2	0.3	0.3	0.3	0.4	0.5	*	*	*	*
All races 3/, female										
2004.....	0.3	0.3	0.4	0.4	*	*	0.6	0.5	*	*
2003.....	*	0.3	0.3	0.4	0.4	*	*	*	*	*
2002.....	0.2	0.3	0.3	0.3	0.4	*	0.4	*	*	*
2001.....	*	0.2	0.3	0.4	*	*	*	*	*	*
2000.....	0.2	0.2	*	0.3	0.4	*	*	*	0.7	*
1999.....	0.2	0.2	*	*	*	*	*	*	*	*
White, both sexes										
2004.....	0.3	0.4	0.4	0.5	0.5	0.5	0.7	0.7	*	*
2003.....	0.2	0.3	0.4	0.4	0.5	0.4	0.5	0.5	0.6	*
2002.....	0.2	0.3	0.3	0.4	0.5	0.4	0.5	0.5	0.6	*
2001.....	0.2	0.2	0.3	0.4	0.4	0.3	0.6	0.6	0.4	*
2000.....	0.2	0.3	0.3	0.3	0.4	0.5	0.5	0.4	0.7	*
1999.....	0.2	0.3	0.2	0.3	0.3	0.4	0.4	0.4	0.5	0.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

All races 3/, both sexes

2004.....	3.0	4.6	6.6	9.9	14.6	19.9	27.6	36.3	43.1	51.4
2003.....	3.0	4.5	6.5	9.7	15.2	20.6	29.0	37.8	43.9	56.0
2002.....	3.1	4.5	6.4	9.7	14.4	20.7	27.7	37.2	45.2	53.8
2001.....	2.9	4.3	6.1	9.2	13.8	20.7	27.6	36.3	43.9	51.9
2000.....	2.9	4.5	6.7	9.3	13.9	19.2	27.7	35.9	44.6	55.4
1999.....	2.5	4.1	6.2	9.1	13.1	19.8	27.1	36.6	43.4	53.6

All races 3/, male

2004.....	3.6	5.5	7.8	11.6	16.7	23.2	32.9	44.4	51.3	62.4
2003.....	3.9	5.5	7.9	11.4	17.4	23.1	33.3	45.4	56.3	70.0
2002.....	3.7	5.6	7.3	11.1	16.5	24.2	32.9	44.8	56.2	66.1
2001.....	3.4	5.0	7.0	10.4	15.3	24.6	33.0	44.4	56.2	69.7
2000.....	3.4	5.5	7.7	10.7	16.3	22.9	33.5	43.6	58.4	72.0
1999.....	3.0	4.6	7.2	10.5	15.3	23.3	32.0	45.6	56.5	70.3

All races 3/, female

2004.....	2.3	3.7	5.4	8.3	12.7	17.1	23.3	30.5	38.1	46.5
2003.....	2.2	3.5	5.1	8.2	13.1	18.4	25.5	32.3	36.3	49.9
2002.....	2.4	3.4	5.5	8.3	12.4	17.7	23.6	31.8	38.4	48.5
2001.....	2.3	3.7	5.2	8.0	12.4	17.3	23.4	30.6	36.6	44.5
2000.....	2.4	3.6	5.7	8.0	11.8	16.0	23.2	30.6	36.5	48.7
1999.....	2.0	3.6	5.1	7.8	11.0	16.8	23.2	30.4	35.7	46.8

White, both sexes

2004.....	3.0	4.4	6.4	9.7	14.6	19.9	28.1	36.9	43.7	52.3
2003.....	2.9	4.4	6.2	9.6	15.0	20.9	29.4	38.3	44.8	56.9
2002.....	3.0	4.4	6.2	9.5	14.4	20.9	28.0	37.5	46.1	54.7
2001.....	2.8	4.1	5.9	8.8	13.6	20.8	27.9	36.6	44.4	52.5
2000.....	2.7	4.3	6.5	9.2	13.9	19.1	27.8	36.6	45.4	56.2
1999.....	2.5	3.7	6.0	8.8	12.9	19.8	27.2	36.9	44.5	54.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

White, male

2004.....	3.8	5.4	7.7	11.3	16.7	23.3	33.8	45.0	52.3	64.7
2003.....	3.9	5.4	7.8	11.4	17.2	23.3	34.2	46.3	57.5	72.0
2002.....	3.7	5.5	7.0	10.9	16.6	24.1	33.0	45.5	57.7	67.5
2001.....	3.3	4.8	6.7	10.1	15.2	24.8	33.3	44.7	57.4	70.1
2000.....	3.2	5.4	7.7	10.5	16.1	23.0	33.8	44.5	59.3	73.1
1999.....	3.0	4.5	7.1	10.4	15.2	23.4	32.3	46.3	58.5	71.5

White, female

2004.....	2.2	3.4	5.2	8.1	12.6	16.9	23.3	30.9	38.3	46.8
2003.....	2.0	3.4	4.7	7.9	12.9	18.8	25.4	32.6	37.0	50.3
2002.....	2.4	3.3	5.3	8.1	12.4	18.1	23.9	31.8	39.0	49.2
2001.....	2.3	3.5	5.1	7.5	12.1	17.3	23.5	30.8	36.5	45.1
2000.....	2.3	3.3	5.4	8.0	11.8	15.7	23.1	31.1	37.1	49.3
1999.....	2.0	3.0	4.9	7.3	10.8	16.7	23.1	30.4	36.2	47.3

Black, both sexes

2004.....	3.7	7.2	9.5	14.2	18.9	24.5	29.2	39.4	44.6	49.6
2003.....	4.4	6.5	9.9	13.0	19.9	24.3	30.7	37.1	39.9	52.5
2002.....	4.2	6.4	9.2	13.0	18.1	23.5	31.6	37.6	42.6	50.5
2001.....	4.0	6.5	9.7	14.6	17.7	24.3	30.7	38.4	45.8	48.8
2000.....	4.5	7.5	9.9	12.5	16.7	24.2	30.4	35.2	40.7	51.3
1999.....	3.5	7.6	9.4	13.1	16.5	22.5	31.4	38.0	34.7	50.7

Black, male

2004.....	3.8	8.2	11.8	17.6	21.3	27.1	32.7	46.9	50.2	50.7
2003.....	5.1	7.9	10.7	14.3	23.9	29.3	32.5	42.9	47.2	50.0
2002.....	5.1	7.9	10.5	15.6	20.4	29.2	40.1	42.3	49.7	57.9
2001.....	5.0	7.5	11.4	15.2	19.3	28.2	35.7	49.7	53.0	62.4
2000.....	5.6	8.3	10.2	16.0	20.1	27.6	37.4	44.9	56.5	63.4
1999.....	4.2	7.5	11.1	13.1	18.9	24.7	33.9	47.0	37.5	65.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

Black, female

2004.....	3.6	6.4	7.5	11.4	17.0	22.5	26.8	35.0	41.6	49.2
2003.....	3.8	5.3	9.3	12.0	16.7	20.7	29.5	33.6	36.0	53.5
2002.....	3.5	5.1	8.2	10.9	16.3	19.2	26.0	34.9	39.0	47.6
2001.....	3.1	5.5	8.2	14.1	16.5	21.4	27.4	31.6	42.1	43.8
2000.....	3.5	6.7	9.6	9.7	14.1	21.7	25.8	29.3	32.7	46.8
1999.....	2.9	7.6	7.9	13.2	14.6	21.0	29.7	32.6	33.3	45.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

IX. Diseases of the circulatory system (I00-I99)

All races 3/, both sexes

2004.....	48.7	87.2	140.6	213.0	345.4	540.6	889.4	1,521.5	2,741.8	6,639.9
2003.....	51.3	89.0	144.2	226.3	366.2	577.3	951.0	1,634.2	2,930.7	7,162.1
2002.....	50.4	89.0	145.7	235.5	376.5	605.9	997.5	1,707.6	3,047.6	7,414.0
2001.....	49.6	86.6	146.1	236.8	388.4	623.5	1,022.9	1,767.1	3,145.7	7,699.8
2000.....	50.5	89.8	148.2	250.1	410.1	651.3	1,071.3	1,832.2	3,265.6	8,055.8
1999.....	50.8	89.3	149.7	256.4	418.3	682.1	1,115.6	1,895.5	3,369.1	8,220.5

All races 3/, male

2004.....	65.6	121.7	200.3	298.4	474.9	718.1	1,139.5	1,908.7	3,295.5	6,837.7
2003.....	68.6	124.5	206.3	316.8	500.6	769.4	1,230.2	2,058.7	3,527.9	7,348.4
2002.....	68.7	125.2	207.7	329.0	515.2	805.6	1,291.7	2,147.8	3,646.7	7,659.3
2001.....	67.3	122.7	206.2	329.7	529.2	825.3	1,331.5	2,231.5	3,786.4	7,958.0
2000.....	69.1	128.3	210.8	352.5	559.6	861.1	1,409.4	2,313.7	3,969.1	8,470.3
1999.....	69.8	126.7	213.7	360.1	570.7	912.4	1,466.7	2,400.1	4,092.1	8,624.0

All races 3/, female

2004.....	32.0	53.7	83.4	132.5	227.5	386.2	688.0	1,243.2	2,400.8	6,551.0
2003.....	34.1	54.4	84.9	141.0	244.0	410.7	727.1	1,332.1	2,565.4	7,079.7
2002.....	32.4	53.9	86.3	147.7	250.7	433.5	762.4	1,396.8	2,683.2	7,307.6
2001.....	32.2	51.6	88.6	149.8	260.6	449.7	777.7	1,441.9	2,761.9	7,591.7
2000.....	32.1	52.4	88.3	154.3	274.6	471.5	804.9	1,496.7	2,850.6	7,887.0
1999.....	32.0	53.1	88.4	159.5	280.3	485.1	841.2	1,544.6	2,946.1	8,057.9

White, both sexes

2004.....	42.6	76.0	123.3	191.3	318.5	506.9	855.3	1,486.6	2,733.2	6,728.0
2003.....	44.2	77.3	126.2	204.9	335.4	540.6	911.7	1,600.0	2,921.5	7,256.7
2002.....	44.3	77.7	128.7	213.7	346.9	571.0	961.2	1,670.2	3,036.0	7,507.0
2001.....	42.3	75.2	128.8	214.1	357.9	583.9	985.3	1,728.3	3,134.6	7,796.7
2000.....	43.4	77.7	132.0	227.1	378.7	615.7	1,032.2	1,790.7	3,255.4	8,159.4
1999.....	43.1	77.5	133.3	231.7	387.3	647.6	1,072.5	1,854.7	3,357.7	8,322.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

IX. Diseases of the circulatory system (I00-I99)

White, male

2004.....	59.8	109.2	180.6	272.3	442.5	680.8	1,105.9	1,877.0	3,310.2	6,960.5
2003.....	61.4	111.5	185.2	292.0	466.1	730.5	1,188.2	2,029.2	3,531.1	7,485.4
2002.....	62.6	113.0	188.2	303.5	481.4	768.5	1,257.1	2,119.7	3,654.4	7,786.5
2001.....	59.8	110.2	187.0	303.7	496.2	785.5	1,291.8	2,200.7	3,789.8	8,082.3
2000.....	61.7	115.8	192.1	325.2	526.1	826.1	1,371.9	2,281.0	3,975.9	8,607.0
1999.....	61.6	113.9	194.5	331.6	535.7	877.3	1,424.9	2,368.9	4,101.9	8,738.2

White, female

2004.....	25.2	43.0	67.3	113.3	203.5	352.6	649.5	1,201.0	2,374.8	6,623.2
2003.....	26.8	43.2	68.4	121.2	214.4	372.9	685.4	1,289.6	2,545.7	7,155.4
2002.....	25.9	42.6	70.5	127.7	222.6	397.2	720.1	1,348.4	2,656.9	7,385.6
2001.....	24.6	40.5	72.0	128.6	230.3	406.9	737.1	1,393.7	2,738.6	7,677.1
2000.....	25.0	39.9	73.1	133.8	242.8	431.7	759.7	1,445.8	2,826.7	7,977.5
1999.....	24.5	41.5	73.6	136.8	250.6	447.2	792.4	1,493.6	2,918.3	8,155.3

Black, both sexes

2004.....	100.0	181.9	297.1	438.1	655.7	928.1	1,383.6	2,144.9	3,231.9	6,397.4
2003.....	106.5	186.5	308.3	455.3	700.9	989.8	1,498.2	2,263.5	3,423.0	6,895.6
2002.....	100.9	186.4	305.2	472.5	703.6	1,007.6	1,517.4	2,356.8	3,577.5	7,111.7
2001.....	107.5	184.1	309.6	475.0	719.8	1,064.4	1,557.8	2,436.9	3,638.1	7,346.8
2000.....	106.7	194.4	306.7	493.2	742.0	1,070.6	1,632.2	2,510.7	3,733.9	7,548.9
1999.....	111.7	190.9	313.3	511.4	747.2	1,100.0	1,711.0	2,564.7	3,805.3	7,610.9

Black, male

2004.....	121.0	236.6	398.3	597.9	896.8	1,216.2	1,737.7	2,639.2	3,693.8	6,222.9
2003.....	131.6	243.1	416.3	613.6	931.7	1,281.3	1,910.4	2,805.4	4,056.0	6,580.1
2002.....	124.8	240.2	409.4	638.6	940.4	1,300.5	1,916.4	2,815.9	4,121.4	6,961.1
2001.....	132.1	239.0	408.8	632.4	942.9	1,350.6	1,998.5	2,960.2	4,299.5	7,393.4
2000.....	133.7	250.3	410.7	675.6	960.7	1,337.4	2,049.3	3,012.7	4,411.3	7,717.4
1999.....	140.9	249.2	426.4	688.3	996.7	1,409.3	2,153.2	3,061.2	4,428.7	7,943.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

IX. Diseases of the circulatory system (I00-I99)

Black, female

2004.....	81.6	134.1	211.5	306.5	467.8	714.2	1,143.3	1,852.1	2,988.4	6,466.6
2003.....	84.4	137.1	216.9	325.0	520.9	773.4	1,220.2	1,941.4	3,091.2	7,018.2
2002.....	79.8	139.8	216.8	336.2	518.4	790.4	1,249.6	2,082.4	3,294.4	7,169.4
2001.....	85.8	136.5	225.1	345.8	544.3	853.0	1,264.3	2,124.5	3,301.2	7,329.5
2000.....	82.8	145.9	218.0	343.7	569.4	873.8	1,355.3	2,208.1	3,392.7	7,486.2
1999.....	86.0	140.6	217.4	367.0	551.6	870.8	1,417.5	2,265.6	3,496.4	7,486.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races 3/, both sexes

2004.....	0.3	0.4	0.6	0.9	1.4	2.6	4.2	7.3	10.9	17.1
2003.....	0.3	0.5	0.6	1.0	1.8	3.0	4.6	8.1	11.6	20.5
2002.....	0.2	0.5	0.7	1.3	2.1	2.8	5.0	8.4	11.4	17.8
2001.....	0.4	0.5	0.7	1.2	1.7	3.2	5.0	8.1	11.4	17.7
2000.....	0.4	0.6	0.8	1.4	2.2	3.4	5.6	8.1	11.6	16.8
1999.....	0.3	0.5	0.9	1.4	2.1	3.5	5.8	8.4	12.6	18.0

All races 3/, male

2004.....	0.2	0.4	0.5	0.7	0.9	2.1	2.8	5.7	9.5	11.5
2003.....	0.2	0.4	0.4	0.8	1.5	2.3	3.6	6.4	9.0	16.2
2002.....	0.2	0.4	0.6	1.0	1.5	1.9	4.2	5.8	8.2	11.8
2001.....	0.3	0.4	0.5	1.1	1.2	2.4	3.2	6.0	7.7	12.8
2000.....	0.3	0.5	0.6	1.1	1.8	2.2	3.6	5.8	7.8	12.2
1999.....	0.3	0.4	0.6	1.0	1.4	2.4	4.7	5.4	8.9	10.3

All races 3/, female

2004.....	0.3	0.4	0.7	1.1	1.9	3.0	5.3	8.3	11.8	19.5
2003.....	0.3	0.6	0.8	1.1	2.1	3.6	5.5	9.3	13.2	22.3
2002.....	0.3	0.5	0.9	1.5	2.6	3.7	5.6	10.2	13.4	20.4
2001.....	0.4	0.6	0.9	1.3	2.2	3.9	6.4	9.5	13.7	19.8
2000.....	0.5	0.7	0.9	1.6	2.5	4.5	7.1	9.7	13.9	18.7
1999.....	0.4	0.6	1.1	1.8	2.7	4.5	6.7	10.4	14.7	21.1

White, both sexes

2004.....	0.2	0.3	0.6	0.7	1.4	2.6	4.4	7.6	11.6	18.2
2003.....	0.2	0.4	0.5	0.8	1.6	3.1	4.7	8.6	12.4	21.8
2002.....	0.2	0.4	0.7	1.2	2.0	2.8	5.2	8.6	12.1	18.9
2001.....	0.3	0.4	0.6	1.0	1.7	3.2	5.0	8.4	12.2	18.7
2000.....	0.3	0.5	0.7	1.3	2.2	3.5	5.6	8.6	12.4	17.6
1999.....	0.3	0.5	0.8	1.4	2.0	3.6	6.0	8.8	13.1	18.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
Black, female										
2004.....	*	*	*	*	*	*	*	5.4	*	*
2003.....	*	1.6	1.7	*	3.3	*	6.2	*	*	8.1
2002.....	*	*	*	*	3.6	*	*	5.5	*	*
2001.....	*	*	2.2	2.7	*	3.8	7.2	6.4	*	*
2000.....	*	*	*	*	*	4.4	*	*	*	*
1999.....	*	*	*	2.8	*	*	4.7	*	*	11.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic rheumatic heart diseases (I05-I09)

All races 3/, both sexes

2004.....	0.3	0.4	0.6	0.9	1.4	2.5	4.1	7.2	10.8	17.0
2003.....	0.2	0.5	0.6	0.9	1.7	3.0	4.6	8.0	11.6	20.3
2002.....	0.2	0.4	0.7	1.3	2.1	2.8	4.9	8.2	11.3	17.5
2001.....	0.4	0.5	0.7	1.2	1.7	3.2	4.9	8.0	11.3	17.2
2000.....	0.4	0.6	0.7	1.4	2.1	3.4	5.5	8.0	11.6	16.7
1999.....	0.3	0.5	0.8	1.4	2.1	3.5	5.8	8.3	12.5	17.9

All races 3/, male

2004.....	0.2	0.3	0.5	0.7	0.9	2.0	2.8	5.7	9.4	11.4
2003.....	0.2	0.4	0.4	0.8	1.4	2.3	3.5	6.4	9.0	16.1
2002.....	0.2	0.4	0.6	1.0	1.5	1.9	4.1	5.7	8.1	11.4
2001.....	0.3	0.4	0.4	1.0	1.2	2.4	3.2	5.9	7.5	12.3
2000.....	0.3	0.5	0.6	1.1	1.8	2.2	3.5	5.7	7.8	12.1
1999.....	0.3	0.4	0.6	1.0	1.3	2.4	4.6	5.4	8.8	10.3

All races 3/, female

2004.....	0.3	0.4	0.7	1.1	1.8	3.0	5.2	8.3	11.7	19.5
2003.....	0.3	0.6	0.8	1.1	2.0	3.6	5.4	9.1	13.2	22.2
2002.....	0.3	0.5	0.9	1.5	2.6	3.6	5.5	10.0	13.3	20.2
2001.....	0.4	0.6	0.9	1.3	2.2	3.8	6.3	9.4	13.5	19.3
2000.....	0.5	0.7	0.8	1.6	2.5	4.4	7.1	9.7	13.8	18.5
1999.....	0.4	0.6	1.1	1.8	2.7	4.5	6.6	10.3	14.6	21.0

White, both sexes

2004.....	0.2	0.3	0.6	0.7	1.4	2.5	4.3	7.5	11.5	18.1
2003.....	0.2	0.4	0.5	0.8	1.6	3.1	4.6	8.5	12.4	21.7
2002.....	0.2	0.4	0.7	1.1	2.0	2.8	5.2	8.5	12.0	18.6
2001.....	0.3	0.4	0.6	1.0	1.7	3.1	4.9	8.3	12.0	18.3
2000.....	0.3	0.5	0.7	1.3	2.1	3.5	5.6	8.6	12.3	17.5
1999.....	0.3	0.5	0.7	1.4	2.0	3.6	6.0	8.7	13.1	18.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic rheumatic heart diseases (I05-I09)										
Black, female										
2004.....	*	*	*	*	*	*	*	5.4	*	*
2003.....	*	1.6	*	*	3.2	*	6.0	*	*	8.1
2002.....	*	*	*	*	3.6	*	*	*	*	*
2001.....	*	*	2.2	2.7	*	*	7.2	6.4	*	*
2000.....	*	*	*	*	*	4.2	*	*	*	*
1999.....	*	*	*	*	*	*	4.7	*	*	10.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Rheumatic mitral valve diseases (I05)

All races 3/, both sexes

2004.....	0.1	0.2	0.3	0.4	0.7	1.2	2.1	3.3	4.7	6.6
2003.....	0.1	0.3	0.4	0.4	1.0	1.7	2.3	4.0	5.6	7.6
2002.....	0.2	0.2	0.5	0.8	1.3	1.4	2.6	4.4	5.7	7.4
2001.....	0.2	0.3	0.5	0.7	1.0	1.8	2.8	4.5	6.0	8.5
2000.....	0.2	0.4	0.5	0.8	1.3	2.0	3.4	4.4	6.8	8.5
1999.....	0.2	0.3	0.5	0.8	1.2	2.1	3.2	4.9	6.8	9.0

All races 3/, male

2004.....	*	0.2	0.3	0.3	0.4	0.8	1.4	2.4	3.9	4.6
2003.....	*	0.2	0.2	0.4	1.0	1.4	1.6	3.1	4.2	5.9
2002.....	*	*	0.4	0.6	0.9	0.9	2.3	2.6	3.7	4.2
2001.....	0.2	0.3	0.3	0.6	0.5	1.4	1.8	2.9	3.7	6.2
2000.....	0.2	0.3	0.3	0.7	1.1	1.2	2.2	3.2	4.7	6.1
1999.....	*	0.3	0.4	0.6	0.7	1.5	2.4	3.2	4.4	4.9

All races 3/, female

2004.....	0.2	0.2	0.4	0.5	1.0	1.4	2.7	3.9	5.3	7.5
2003.....	0.2	0.4	0.5	0.5	1.0	1.9	2.9	4.7	6.5	8.4
2002.....	*	0.3	0.6	0.9	1.6	1.8	2.9	5.7	7.0	8.7
2001.....	0.2	0.4	0.6	0.8	1.4	2.1	3.6	5.5	7.4	9.4
2000.....	0.3	0.4	0.6	0.9	1.5	2.7	4.3	5.2	8.0	9.5
1999.....	0.2	0.4	0.7	1.1	1.6	2.7	3.8	6.1	8.3	10.7

White, both sexes

2004.....	*	0.2	0.3	0.4	0.7	1.2	2.2	3.4	4.9	7.1
2003.....	0.1	0.2	0.3	0.3	0.9	1.7	2.4	4.3	5.9	8.1
2002.....	0.1	0.2	0.5	0.7	1.2	1.3	2.8	4.6	6.2	7.9
2001.....	0.2	0.3	0.4	0.6	1.0	1.7	2.8	4.7	6.5	9.1
2000.....	0.2	0.4	0.5	0.8	1.3	2.2	3.4	4.6	7.3	9.1
1999.....	0.1	0.3	0.4	0.8	1.2	2.2	3.3	5.2	7.2	9.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of both mitral and aortic valves (I08.0)										
All races 3/, both sexes										
2004.....	*	*	*	0.1	0.2	0.5	0.7	1.5	2.6	5.2
2003.....	*	*	*	0.2	0.2	0.4	0.8	1.6	2.7	6.5
2002.....	*	*	*	*	*	0.3	0.7	1.3	1.8	3.5
2001.....	*	*	*	*	*	0.3	0.6	0.7	1.6	2.5
2000.....	*	*	*	*	0.2	0.2	0.5	0.8	1.0	2.0
1999.....	*	*	*	*	0.2	0.2	0.6	0.5	1.1	2.0
All races 3/, male										
2004.....	*	*	*	*	*	0.6	*	1.5	3.0	3.8
2003.....	*	*	*	*	*	*	0.9	1.6	2.4	6.0
2002.....	*	*	*	*	*	*	0.7	1.4	1.5	2.4
2001.....	*	*	*	*	*	*	0.6	*	1.6	2.3
2000.....	*	*	*	*	*	*	*	*	*	1.9
1999.....	*	*	*	*	*	*	0.6	*	*	*
All races 3/, female										
2004.....	*	*	*	*	*	0.4	0.9	1.5	2.3	5.9
2003.....	*	*	*	*	*	0.5	0.7	1.6	2.9	6.7
2002.....	*	*	*	*	*	*	0.6	1.2	2.1	4.0
2001.....	*	*	*	*	*	0.4	0.6	0.7	1.6	2.5
2000.....	*	*	*	*	*	*	0.5	0.9	1.2	2.1
1999.....	*	*	*	*	*	*	0.6	0.6	1.2	2.2
White, both sexes										
2004.....	*	*	*	*	0.2	0.5	0.8	1.5	2.8	5.6
2003.....	*	*	*	0.2	*	0.4	0.8	1.7	3.0	7.0
2002.....	*	*	*	*	*	0.4	0.7	1.4	1.9	3.8
2001.....	*	*	*	*	*	0.4	0.6	0.7	1.7	2.6
2000.....	*	*	*	*	*	*	0.5	0.8	1.1	2.0
1999.....	*	*	*	*	*	*	0.6	0.6	1.2	2.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)

All races 3/, both sexes

2004.....	0.1	0.1	0.2	0.2	0.4	0.8	1.2	2.2	3.1	4.4
2003.....	*	0.1	0.2	0.3	0.5	0.8	1.4	2.1	2.9	5.4
2002.....	*	0.1	0.2	0.4	0.6	1.0	1.5	2.2	3.2	5.6
2001.....	0.1	0.1	0.2	0.4	0.5	1.0	1.4	2.7	3.4	5.9
2000.....	0.1	0.2	0.2	0.4	0.7	1.1	1.6	2.9	3.7	5.8
1999.....	0.1	0.1	0.2	0.4	0.7	1.1	1.8	2.7	4.3	6.6

All races 3/, male

2004.....	*	*	*	*	*	0.6	0.7	1.6	2.1	2.5
2003.....	*	*	*	*	*	0.5	0.9	1.5	1.8	3.4
2002.....	*	*	*	0.4	0.4	0.6	1.0	1.5	2.4	3.8
2001.....	*	*	*	0.4	0.4	0.7	0.8	2.2	1.8	3.6
2000.....	*	*	*	*	0.5	0.7	0.9	1.9	2.2	4.0
1999.....	*	*	*	*	0.4	0.7	1.6	1.6	3.2	3.7

All races 3/, female

2004.....	*	*	0.2	0.3	0.5	1.1	1.6	2.7	3.7	5.3
2003.....	*	*	*	0.4	0.7	1.1	1.7	2.6	3.5	6.3
2002.....	*	*	*	0.4	0.8	1.3	1.9	2.7	3.8	6.3
2001.....	*	*	*	0.4	0.6	1.2	2.0	3.1	4.4	6.8
2000.....	*	0.2	0.3	0.5	0.8	1.4	2.2	3.5	4.6	6.6
1999.....	*	*	0.3	0.6	0.9	1.5	2.0	3.5	5.0	7.8

White, both sexes

2004.....	*	*	0.2	0.2	0.4	0.8	1.2	2.4	3.3	4.7
2003.....	*	0.1	0.1	0.3	0.5	0.9	1.3	2.3	3.1	5.7
2002.....	*	*	0.1	0.4	0.6	1.0	1.6	2.3	3.4	5.9
2001.....	*	*	*	0.3	0.5	1.0	1.5	2.8	3.6	6.3
2000.....	*	0.1	0.2	0.3	0.7	1.1	1.7	3.1	3.8	6.0
1999.....	*	*	0.2	0.4	0.6	1.2	1.9	2.8	4.6	6.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive diseases (I10-I13)

All races 3/, both sexes

2004.....	5.2	9.4	12.9	17.8	24.3	33.3	49.5	79.7	146.8	416.1
2003.....	5.2	8.9	13.3	17.6	24.0	34.0	49.3	83.1	146.3	409.5
2002.....	4.7	8.1	12.0	16.5	22.8	32.2	47.5	77.8	142.1	390.8
2001.....	4.5	7.2	11.1	14.9	21.2	30.3	45.3	77.2	140.5	384.7
2000.....	4.3	6.8	10.0	14.5	21.9	29.5	45.1	75.2	138.9	380.3
1999.....	4.0	6.6	9.4	14.6	20.6	30.4	43.8	75.7	136.9	366.0

All races 3/, male

2004.....	6.8	12.4	17.6	23.3	31.9	41.4	55.6	84.3	144.0	335.7
2003.....	6.8	12.0	17.8	23.0	29.3	41.8	54.8	85.9	139.5	322.4
2002.....	6.4	10.8	16.0	21.9	28.6	37.8	52.5	82.0	136.9	311.1
2001.....	6.2	9.8	14.6	18.8	25.8	36.4	50.0	80.0	134.2	309.4
2000.....	5.6	9.0	13.8	18.4	27.5	33.8	51.2	76.7	135.5	309.3
1999.....	5.4	8.6	12.2	18.7	24.7	35.1	47.9	79.8	131.9	297.3

All races 3/, female

2004.....	3.6	6.6	8.5	12.6	17.4	26.3	44.6	76.4	148.6	452.3
2003.....	3.7	5.8	9.0	12.4	19.1	27.3	44.8	81.1	150.4	448.0
2002.....	3.1	5.5	8.1	11.5	17.5	27.5	43.5	74.8	145.3	425.4
2001.....	2.9	4.7	7.7	11.3	17.0	25.2	41.5	75.2	144.3	416.3
2000.....	3.0	4.7	6.4	10.8	16.9	25.8	40.3	74.1	140.9	409.3
1999.....	2.7	4.7	6.7	10.8	16.9	26.3	40.6	72.9	139.8	393.7

White, both sexes

2004.....	3.5	6.3	8.8	12.5	17.8	25.9	39.8	67.5	132.1	394.8
2003.....	3.4	5.7	9.0	12.1	17.8	25.4	38.9	70.4	133.7	389.7
2002.....	3.1	5.4	8.1	11.5	16.5	24.3	38.0	65.5	126.8	368.7
2001.....	2.7	4.5	7.3	10.2	15.2	22.3	35.8	65.7	126.0	364.3
2000.....	2.8	4.3	6.4	9.6	15.3	21.9	35.2	63.1	123.9	357.1
1999.....	2.3	4.0	6.0	9.6	14.4	22.3	33.9	62.7	121.1	343.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hypertensive diseases (I10-I13)

White, male

2004.....	5.0	8.8	12.6	16.9	23.7	32.6	45.4	71.1	130.1	318.7
2003.....	4.9	8.2	12.8	16.7	22.4	32.0	43.9	73.7	127.4	301.5
2002.....	4.6	7.7	11.4	16.0	21.1	28.7	42.7	69.5	121.3	290.6
2001.....	4.0	6.3	10.0	13.5	19.2	27.5	40.3	68.4	119.3	288.6
2000.....	3.8	6.2	9.3	12.8	19.7	25.8	40.8	65.4	120.1	286.2
1999.....	3.5	5.6	8.1	12.9	17.9	26.2	37.7	66.4	116.4	273.5

White, female

2004.....	2.1	3.7	5.1	8.3	12.3	20.0	35.2	64.8	133.4	429.1
2003.....	1.9	3.1	5.3	7.7	13.6	19.5	34.8	68.0	137.5	428.7
2002.....	1.6	3.2	4.8	7.1	12.2	20.4	34.2	62.6	130.2	402.6
2001.....	1.4	2.7	4.6	7.1	11.6	17.8	32.2	63.7	130.1	396.0
2000.....	1.7	2.4	3.5	6.6	11.2	18.5	30.6	61.5	126.1	385.8
1999.....	1.2	2.5	3.9	6.5	11.2	19.0	30.9	60.1	123.9	371.5

Black, both sexes

2004.....	17.1	33.1	46.6	65.2	86.0	104.8	150.5	222.2	337.4	698.5
2003.....	18.2	33.2	48.7	67.7	82.8	113.4	157.0	231.9	315.3	686.5
2002.....	16.3	28.9	44.5	64.1	81.0	105.7	150.4	224.0	341.7	685.4
2001.....	17.2	28.3	43.3	58.1	76.4	104.6	145.9	213.3	325.6	664.2
2000.....	15.3	26.9	40.6	59.6	82.1	103.2	152.4	221.1	326.4	673.8
1999.....	16.3	26.7	39.8	60.1	75.9	105.6	151.7	230.5	335.2	639.9

Black, male

2004.....	20.7	40.8	61.3	86.2	117.2	136.4	173.2	258.9	343.6	572.2
2003.....	21.6	43.0	62.5	87.0	102.4	143.3	186.2	247.4	321.1	612.0
2002.....	20.3	36.6	57.8	82.5	104.1	126.9	170.9	247.6	354.6	602.5
2001.....	22.3	38.1	56.4	72.4	92.5	129.1	167.0	230.5	349.1	604.8
2000.....	19.5	33.2	54.3	75.6	103.7	119.4	173.5	224.0	349.8	606.6
1999.....	20.3	33.9	51.4	75.3	92.3	126.7	169.6	251.7	351.4	589.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive diseases (I10-I13)

Black, female

2004.....	13.9	26.5	34.1	47.9	61.6	81.3	135.1	200.5	334.1	748.6
2003.....	15.2	24.7	37.0	51.9	67.5	91.2	137.4	222.7	312.2	715.4
2002.....	12.8	22.2	33.2	49.0	62.9	89.9	136.7	209.9	335.0	717.1
2001.....	12.7	19.7	32.2	46.3	63.7	86.5	131.8	203.0	313.6	686.2
2000.....	11.5	21.5	28.9	46.6	65.0	91.2	138.3	219.3	314.6	698.8
1999.....	12.7	20.5	30.1	47.6	63.1	90.0	139.7	217.7	327.1	658.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Essential (primary) hypertension (I10)

All races 3/, both sexes

2004.....	0.5	1.2	2.0	2.8	4.6	6.9	11.6	20.8	38.9	127.4
2003.....	0.5	1.0	2.0	2.9	4.6	7.2	11.6	20.5	39.5	121.9
2002.....	0.5	1.0	1.6	2.5	4.2	6.6	10.9	18.5	37.1	112.0
2001.....	0.4	1.0	1.6	2.7	4.3	6.9	10.1	19.1	35.6	107.4
2000.....	0.5	1.1	1.7	2.6	4.3	6.1	10.4	18.2	33.1	101.1
1999.....	0.4	0.9	1.6	2.5	4.3	6.7	9.7	18.0	33.0	98.1

All races 3/, male

2004.....	0.6	1.5	2.7	3.2	5.9	8.4	11.9	20.7	35.3	86.5
2003.....	0.6	1.3	2.6	3.6	5.6	8.9	12.3	18.1	33.5	77.4
2002.....	0.7	1.3	2.1	3.3	5.0	7.3	11.9	17.5	32.0	74.2
2001.....	0.5	1.2	2.2	3.4	5.2	7.9	10.9	18.8	30.1	74.8
2000.....	0.6	1.5	2.3	3.3	5.4	6.5	11.7	17.4	29.5	70.3
1999.....	0.5	1.1	1.9	3.2	5.1	7.8	10.7	17.2	29.7	68.1

All races 3/, female

2004.....	0.3	1.0	1.3	2.4	3.5	5.6	11.3	20.8	41.1	145.8
2003.....	0.5	0.6	1.4	2.2	3.7	5.7	11.0	22.1	43.1	141.5
2002.....	0.4	0.7	1.2	1.8	3.5	6.0	10.2	19.1	40.2	128.3
2001.....	0.4	0.7	1.1	2.0	3.5	6.1	9.4	19.3	38.9	121.1
2000.....	0.4	0.6	1.2	1.9	3.4	5.7	9.4	18.7	35.2	113.6
1999.....	0.3	0.7	1.3	1.9	3.6	5.7	8.9	18.6	34.9	110.2

White, both sexes

2004.....	0.4	0.9	1.6	2.2	4.0	5.9	10.3	18.9	36.9	123.5
2003.....	0.4	0.7	1.6	2.2	3.8	5.9	9.6	18.3	36.6	119.9
2002.....	0.4	0.7	1.3	2.1	3.7	5.7	9.5	16.4	35.1	108.6
2001.....	0.3	0.7	1.2	2.1	3.6	5.7	8.6	17.6	33.1	104.6
2000.....	0.4	0.8	1.2	2.1	3.5	5.3	9.2	16.2	31.2	97.5
1999.....	0.3	0.7	1.1	2.0	3.6	5.6	8.7	16.0	30.9	94.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Essential (primary) hypertension (I10)

White, male

2004.....	0.5	1.2	2.3	2.5	5.2	7.4	10.6	18.4	33.5	84.0
2003.....	0.5	1.0	2.2	3.0	4.5	7.6	10.5	15.9	29.7	74.3
2002.....	0.5	1.0	1.7	2.8	4.4	6.3	10.4	15.6	29.1	71.0
2001.....	0.4	0.9	1.6	2.6	4.4	6.8	9.2	16.9	26.9	71.1
2000.....	0.5	1.1	1.7	2.7	4.3	5.8	10.3	15.6	26.8	66.4
1999.....	0.4	0.9	1.4	2.6	4.3	6.7	9.5	15.3	27.0	62.7

White, female

2004.....	0.2	0.5	0.9	1.8	2.8	4.7	10.1	19.2	39.0	141.3
2003.....	0.4	0.5	1.0	1.6	3.1	4.5	8.9	20.0	40.8	140.2
2002.....	0.3	0.4	0.9	1.4	3.0	5.1	8.7	17.1	38.7	124.9
2001.....	*	0.4	0.8	1.6	2.7	4.8	8.1	18.1	36.9	118.7
2000.....	0.3	0.4	0.8	1.5	2.8	4.8	8.4	16.7	33.8	110.2
1999.....	*	0.5	0.8	1.4	2.9	4.7	8.0	16.5	33.2	107.0

Black, both sexes

2004.....	1.3	3.8	5.4	8.5	11.5	16.5	25.0	44.5	64.9	183.6
2003.....	1.4	2.7	5.3	8.9	12.9	19.6	32.3	47.0	78.9	157.5
2002.....	1.5	3.4	4.7	6.9	9.1	15.4	27.5	43.6	67.0	158.6
2001.....	1.6	3.4	4.9	8.3	11.9	18.2	25.8	39.4	66.6	150.8
2000.....	1.4	3.2	5.9	7.9	11.9	14.7	23.6	41.4	57.0	149.2
1999.....	1.4	2.4	6.0	8.0	11.2	16.1	22.2	42.3	58.9	143.2

Black, male

2004.....	1.5	3.8	7.0	10.2	13.8	19.4	26.5	53.2	62.4	124.2
2003.....	1.5	3.7	6.8	10.5	17.2	24.4	32.8	48.4	87.1	119.9
2002.....	1.8	4.0	5.7	8.8	10.3	17.2	30.4	44.1	73.4	117.9
2001.....	1.7	3.8	6.9	11.3	13.6	20.3	31.3	46.0	72.2	132.8
2000.....	*	4.5	7.7	10.7	15.9	14.6	28.4	39.7	64.9	123.4
1999.....	1.7	2.6	6.3	9.5	13.6	19.0	26.5	41.3	66.3	124.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Essential (primary) hypertension (I10)										
Black, female										
2004.....	*	3.7	4.1	7.1	9.6	14.4	23.9	39.4	66.1	207.1
2003.....	1.4	1.8	4.0	7.6	9.6	16.1	31.9	46.1	74.7	172.2
2002.....	*	2.9	3.9	5.4	8.2	14.1	25.5	43.4	63.7	174.2
2001.....	1.5	3.0	3.1	5.9	10.6	16.7	22.2	35.5	63.8	157.5
2000.....	1.4	2.1	4.5	5.5	8.7	14.8	20.4	42.4	53.1	158.8
1999.....	*	2.1	5.7	6.8	9.3	14.0	19.4	42.9	55.3	150.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive heart disease (I11)

All races 3/, both sexes

2004.....	4.0	6.8	9.0	12.2	14.9	18.8	25.1	36.7	66.1	192.8
2003.....	3.9	6.6	9.4	11.7	14.9	19.4	25.1	39.5	66.8	195.0
2002.....	3.5	5.9	8.6	11.1	14.3	18.9	24.0	37.9	66.0	187.0
2001.....	3.2	5.1	7.7	9.7	12.5	16.3	23.5	36.6	66.0	189.5
2000.....	2.9	4.7	6.6	9.3	13.2	16.5	23.0	36.3	66.1	193.1
1999.....	2.8	4.5	6.2	9.6	11.9	16.3	22.1	36.4	64.7	184.9

All races 3/, male

2004.....	5.2	9.2	12.4	17.0	20.4	24.7	30.5	39.1	60.9	145.4
2003.....	5.2	9.0	12.8	16.0	18.8	25.2	29.4	42.8	63.3	142.2
2002.....	4.9	8.0	11.9	15.2	19.0	23.6	27.0	41.0	62.2	134.6
2001.....	4.5	7.0	10.3	12.8	16.5	20.5	27.1	38.9	63.5	134.4
2000.....	3.9	6.3	9.4	12.1	17.4	20.7	27.2	37.3	61.8	139.1
1999.....	3.9	6.1	8.1	12.7	15.0	19.5	24.6	39.4	59.2	133.6

All races 3/, female

2004.....	2.7	4.4	5.8	7.6	9.9	13.5	20.7	35.0	69.3	214.1
2003.....	2.6	4.4	6.1	7.7	11.2	14.4	21.6	37.2	69.0	218.3
2002.....	2.2	3.9	5.4	7.2	10.0	14.8	21.6	35.8	68.2	209.7
2001.....	2.0	3.2	5.1	6.8	8.9	12.7	20.6	35.1	67.5	212.6
2000.....	2.0	3.2	3.8	6.6	9.3	12.9	19.6	35.7	68.6	215.1
1999.....	1.7	3.0	4.3	6.8	9.2	13.5	20.2	34.3	67.9	205.5

White, both sexes

2004.....	2.8	4.8	6.4	9.0	11.2	15.1	20.4	31.0	59.1	183.7
2003.....	2.7	4.4	6.6	8.5	11.4	14.7	20.2	33.2	61.2	185.3
2002.....	2.4	4.3	6.0	8.1	10.5	14.4	19.3	31.9	58.1	177.3
2001.....	2.1	3.3	5.2	6.9	9.1	12.1	18.5	30.9	59.2	179.9
2000.....	2.0	3.1	4.3	6.2	9.4	12.5	17.6	30.2	58.4	181.8
1999.....	1.8	2.9	4.1	6.5	8.7	12.0	16.9	30.0	57.3	175.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive heart disease (I11)

White, male

2004.....	4.1	7.0	9.3	12.8	15.3	20.0	25.0	33.4	54.3	136.4
2003.....	4.0	6.4	9.7	12.2	15.0	19.4	23.6	36.1	57.9	131.9
2002.....	3.7	6.1	8.9	11.6	14.0	18.1	22.0	34.5	54.8	124.8
2001.....	3.1	4.8	7.5	9.6	12.5	15.9	21.9	32.9	56.1	123.7
2000.....	2.8	4.6	6.5	8.6	12.7	16.0	21.5	31.1	53.4	129.3
1999.....	2.7	4.2	5.8	8.9	11.3	14.4	19.1	32.4	51.8	123.7

White, female

2004.....	1.6	2.7	3.5	5.3	7.4	10.8	16.6	29.3	62.1	205.0
2003.....	1.3	2.4	3.5	5.0	8.1	10.7	17.4	31.1	63.3	209.0
2002.....	1.2	2.4	3.3	4.7	7.1	11.1	17.1	30.0	60.1	200.1
2001.....	1.0	1.8	3.0	4.4	6.0	8.8	15.8	29.5	61.1	203.5
2000.....	1.2	1.6	2.2	4.0	6.3	9.4	14.5	29.6	61.3	203.1
1999.....	0.8	1.6	2.5	4.2	6.3	10.0	15.1	28.2	60.6	196.0

Black, both sexes

2004.....	12.3	22.0	31.3	41.6	49.9	55.4	76.2	103.1	161.4	314.7
2003.....	12.8	23.9	32.8	41.9	48.3	63.0	77.1	113.3	145.7	332.3
2002.....	11.4	19.3	30.3	39.7	50.5	61.2	76.5	110.3	167.7	320.4
2001.....	11.5	18.6	28.4	35.7	43.6	55.8	77.8	105.1	154.7	326.1
2000.....	9.9	17.8	25.4	37.6	47.9	54.6	81.6	111.1	164.9	340.0
1999.....	10.2	17.9	24.8	38.0	41.6	57.6	79.0	113.7	161.2	305.1

Black, male

2004.....	14.5	27.8	40.8	59.1	72.3	78.3	97.8	114.6	158.5	267.2
2003.....	14.9	29.8	41.5	55.1	59.8	86.7	99.9	129.6	149.1	287.8
2002.....	14.6	24.4	39.9	53.1	71.1	79.0	89.8	125.4	167.4	267.0
2001.....	14.8	25.1	36.6	45.1	55.9	69.6	91.6	117.1	171.8	284.8
2000.....	12.7	21.4	35.8	47.9	63.8	70.0	95.9	117.2	178.7	273.3
1999.....	13.1	22.5	31.3	49.5	52.0	73.0	90.2	129.2	166.5	263.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive heart disease (I11)

Black, female

2004.....	10.3	17.0	23.2	27.2	32.5	38.5	61.5	96.2	162.9	333.5
2003.....	11.0	18.7	25.5	31.0	39.4	45.5	61.8	103.6	143.9	349.6
2002.....	8.6	14.9	22.3	28.7	34.4	47.9	67.7	101.3	167.9	340.9
2001.....	8.6	13.0	21.5	28.0	33.9	45.6	68.6	97.9	145.9	341.4
2000.....	7.5	14.6	16.5	29.2	35.3	43.2	72.1	107.4	157.9	364.9
1999.....	7.6	13.8	19.3	28.7	33.4	46.2	71.6	104.4	158.6	320.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive renal disease (I12)

All races 3/, both sexes

2004.....	0.5	1.0	1.4	2.2	3.5	6.0	10.3	17.8	32.4	71.1
2003.....	0.5	0.9	1.3	2.1	3.4	5.9	9.8	18.0	30.4	67.0
2002.....	0.5	0.8	1.2	2.0	3.1	5.2	9.7	16.5	29.6	68.4
2001.....	0.6	0.9	1.3	1.7	3.2	5.3	8.9	16.4	29.6	64.5
2000.....	0.5	0.7	1.3	1.8	3.5	5.4	8.7	15.7	29.9	61.8
1999.....	0.5	0.8	1.2	1.6	3.0	5.3	8.9	14.7	26.9	53.9

All races 3/, male

2004.....	0.6	1.2	1.7	2.3	3.9	6.5	10.4	19.4	37.4	79.5
2003.....	0.7	1.2	1.7	2.3	3.7	6.1	10.2	19.4	32.6	76.1
2002.....	0.6	0.9	1.4	2.2	3.2	5.4	10.3	18.4	32.7	79.2
2001.....	0.8	1.1	1.5	1.8	2.9	5.8	9.0	17.2	31.2	76.0
2000.....	0.7	0.8	1.6	2.1	3.7	5.0	8.9	17.0	33.4	72.0
1999.....	0.6	0.9	1.6	1.8	3.1	5.6	9.4	15.9	29.7	64.3

All races 3/, female

2004.....	0.4	0.8	1.0	2.0	3.1	5.7	10.2	16.6	29.3	67.4
2003.....	0.4	0.6	1.0	1.8	3.2	5.7	9.4	17.0	29.1	63.0
2002.....	0.4	0.6	1.1	1.9	2.9	5.0	9.3	15.1	27.7	63.7
2001.....	0.4	0.7	1.2	1.7	3.4	5.0	8.8	15.9	28.6	59.7
2000.....	0.4	0.6	1.0	1.6	3.3	5.7	8.5	14.7	27.8	57.7
1999.....	0.4	0.7	0.8	1.4	2.9	5.1	8.5	13.8	25.3	49.8

White, both sexes

2004.....	0.2	0.4	0.6	1.1	1.8	3.9	7.3	14.0	28.0	64.6
2003.....	0.3	0.4	0.7	1.0	2.0	3.8	7.0	14.7	27.5	61.1
2002.....	0.2	0.3	0.6	1.0	1.7	3.3	7.3	13.3	25.3	61.9
2001.....	0.2	0.4	0.6	0.8	1.8	3.4	6.7	13.3	25.5	58.7
2000.....	0.2	0.2	0.6	0.9	1.9	3.3	6.1	12.7	26.3	55.7
1999.....	0.3	0.3	0.5	0.7	1.4	3.5	6.5	11.5	22.4	47.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive renal disease (I12)

White, male

2004.....	0.3	0.5	0.7	1.3	2.0	4.2	7.6	15.3	33.1	75.4
2003.....	0.4	0.6	0.7	1.1	2.1	4.0	7.5	16.8	30.6	71.0
2002.....	0.3	0.4	0.7	1.1	1.9	3.5	7.9	15.2	28.6	74.2
2001.....	0.3	0.5	0.6	0.9	1.5	3.5	7.2	14.5	27.6	71.1
2000.....	0.3	0.3	0.8	1.1	2.2	3.0	6.5	14.3	30.6	66.1
1999.....	0.3	0.4	0.6	0.9	1.5	3.7	7.1	12.8	25.5	58.3

White, female

2004.....	*	0.4	0.5	0.9	1.6	3.7	7.0	13.1	24.9	59.8
2003.....	*	0.3	0.6	0.9	1.9	3.7	6.6	13.1	25.6	56.6
2002.....	*	0.3	0.5	0.9	1.5	3.2	6.9	12.0	23.3	56.5
2001.....	*	0.3	0.5	0.8	2.0	3.4	6.4	12.4	24.2	53.5
2000.....	*	*	0.3	0.7	1.7	3.5	5.8	11.5	23.6	51.4
1999.....	*	0.3	0.4	0.6	1.4	3.2	6.0	10.6	20.7	43.5

Black, both sexes

2004.....	2.2	4.6	7.0	11.3	18.2	25.3	39.7	59.4	85.2	152.0
2003.....	2.4	4.4	6.8	11.6	16.1	23.3	36.6	55.1	66.5	143.5
2002.....	2.3	4.0	6.1	11.5	15.2	21.1	34.1	52.7	84.0	150.3
2001.....	2.8	4.4	7.7	9.7	15.1	22.1	30.4	50.4	81.5	137.3
2000.....	2.8	4.0	6.8	10.1	17.1	25.6	35.2	51.0	73.4	134.2
1999.....	2.6	4.0	6.8	9.0	15.9	21.9	35.0	52.7	79.7	126.3

Black, male

2004.....	2.7	6.0	9.7	12.5	22.4	30.5	39.5	70.5	95.4	139.1
2003.....	3.0	6.1	9.6	13.7	19.1	25.6	41.7	53.9	60.5	154.3
2002.....	2.9	5.1	7.6	13.0	15.4	23.4	36.7	61.1	90.9	156.5
2001.....	3.8	6.2	9.4	10.5	16.7	28.5	30.6	49.3	85.9	143.0
2000.....	3.9	4.6	7.9	11.9	18.4	25.5	35.4	53.4	72.5	146.5
1999.....	2.9	5.2	10.3	10.0	17.4	24.1	35.6	57.5	88.9	141.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive renal disease (I12)

Black, female

2004.....	1.9	3.3	4.6	10.2	14.9	21.4	39.8	52.8	79.8	157.1
2003.....	1.9	2.9	4.4	9.8	13.7	21.6	33.2	55.9	69.6	139.4
2002.....	1.8	2.9	4.9	10.3	15.0	19.4	32.3	47.8	80.4	147.9
2001.....	1.9	2.8	6.2	9.1	13.9	17.5	30.3	51.0	79.3	135.1
2000.....	1.7	3.6	5.8	8.6	16.0	25.7	35.1	49.6	73.9	129.6
1999.....	2.2	2.9	3.8	8.2	14.6	20.2	34.7	49.8	75.2	120.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive heart and renal disease (I13)

All races 3/, both sexes

2004.....	0.3	0.5	0.5	0.7	1.3	1.6	2.6	4.5	9.4	24.8
2003.....	0.3	0.4	0.6	0.9	1.1	1.6	2.9	5.2	9.5	25.6
2002.....	0.2	0.4	0.5	0.9	1.2	1.6	2.8	4.9	9.5	23.5
2001.....	0.3	0.3	0.4	0.8	1.2	1.8	2.8	5.0	9.3	23.3
2000.....	0.3	0.3	0.4	0.8	1.0	1.6	3.0	5.1	9.8	24.3
1999.....	0.3	0.4	0.4	0.9	1.4	2.1	3.0	6.6	12.3	29.1

All races 3/, male

2004.....	0.3	0.5	0.7	0.7	1.7	1.7	2.8	5.1	10.4	24.3
2003.....	0.3	0.6	0.7	1.1	1.3	1.7	3.0	5.6	10.1	26.7
2002.....	0.2	0.5	0.6	1.1	1.4	1.5	3.4	5.1	9.9	23.1
2001.....	0.4	0.4	0.6	0.9	1.2	2.1	2.9	5.2	9.4	24.2
2000.....	0.3	0.4	0.5	0.9	1.0	1.7	3.4	5.0	10.8	27.8
1999.....	0.4	0.5	0.6	1.0	1.6	2.3	3.2	7.3	13.3	31.3

All races 3/, female

2004.....	*	0.4	0.4	0.7	0.9	1.5	2.3	4.0	8.8	25.0
2003.....	0.2	0.2	0.5	0.6	0.9	1.5	2.8	4.9	9.2	25.2
2002.....	0.2	0.2	0.4	0.7	1.0	1.7	2.4	4.7	9.2	23.7
2001.....	*	*	0.3	0.7	1.3	1.5	2.8	4.9	9.3	22.9
2000.....	0.2	0.2	0.4	0.7	1.0	1.5	2.7	5.1	9.3	22.9
1999.....	0.3	0.3	0.3	0.7	1.2	2.0	2.9	6.1	11.7	28.2

White, both sexes

2004.....	0.1	0.1	0.2	0.3	0.7	0.9	1.8	3.5	8.1	23.0
2003.....	*	0.1	0.2	0.3	0.6	0.9	2.1	4.2	8.3	23.4
2002.....	*	*	0.1	0.3	0.6	0.9	1.9	3.8	8.4	21.0
2001.....	*	*	0.2	0.4	0.7	1.1	2.0	3.9	8.3	21.0
2000.....	0.2	0.1	0.2	0.4	0.5	0.9	2.2	4.0	8.1	22.0
1999.....	*	0.1	0.2	0.4	0.7	1.2	1.9	5.2	10.4	26.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

Hypertensive heart and renal disease (I13)

White, male

2004.....	*	*	*	0.3	1.0	1.0	2.1	3.9	9.2	22.9
2003.....	*	*	*	0.5	0.8	1.1	2.2	4.9	9.3	24.3
2002.....	*	*	*	0.5	0.7	0.8	2.4	4.2	8.8	20.6
2001.....	*	*	0.3	0.4	0.7	1.3	2.0	4.1	8.7	22.7
2000.....	*	*	*	0.4	0.5	1.0	2.5	4.4	9.2	24.4
1999.....	*	*	*	0.5	0.8	1.4	2.0	5.8	12.1	28.8

White, female

2004.....	*	*	*	0.3	0.4	0.8	1.5	3.2	7.4	23.0
2003.....	*	*	*	*	0.5	0.7	1.9	3.7	7.8	22.9
2002.....	*	*	*	*	0.5	0.9	1.5	3.6	8.1	21.1
2001.....	*	*	*	0.3	0.8	0.8	2.0	3.7	8.0	20.3
2000.....	*	*	*	0.4	0.5	0.8	2.0	3.7	7.4	21.1
1999.....	*	*	*	0.4	0.6	1.1	1.8	4.7	9.5	25.0

Black, both sexes

2004.....	1.3	2.8	2.9	3.9	6.4	7.5	9.7	15.2	26.0	48.2
2003.....	1.5	2.3	3.8	5.3	5.5	7.4	11.0	16.5	24.1	53.1
2002.....	1.1	2.2	3.3	6.0	6.2	8.0	12.3	17.4	23.0	56.1
2001.....	1.4	1.9	2.3	4.4	5.7	8.4	11.8	18.4	22.8	50.1
2000.....	1.2	1.9	2.5	4.1	5.3	8.3	11.9	17.6	31.1	50.4
1999.....	2.1	2.5	2.3	5.0	7.3	10.1	15.4	21.8	35.3	65.4

Black, male

2004.....	2.0	3.3	3.8	4.4	8.6	8.2	9.4	20.5	27.3	41.7
2003.....	2.1	3.4	4.6	7.6	6.4	6.6	11.8	15.5	24.4	50.0
2002.....	*	3.0	4.7	7.6	7.3	7.2	14.0	17.0	22.9	61.1
2001.....	2.1	3.0	3.5	5.6	6.3	10.7	13.5	18.1	19.3	44.3
2000.....	1.5	2.7	2.9	5.1	5.6	9.4	13.7	13.7	33.7	63.4
1999.....	2.6	3.5	3.4	6.3	9.3	10.6	17.3	23.7	29.6	59.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive heart and renal disease (I13)

Black, female

2004.....	*	2.4	2.2	3.5	4.7	7.0	9.9	12.1	25.3	50.8
2003.....	*	*	3.1	3.5	4.8	8.0	10.4	17.1	24.0	54.3
2002.....	*	1.5	2.1	4.6	5.3	8.5	11.2	17.6	23.1	54.2
2001.....	*	*	*	3.3	5.3	6.8	10.8	18.6	24.5	52.2
2000.....	*	*	2.1	3.2	5.1	7.5	10.7	19.9	29.7	45.5
1999.....	1.6	1.6	*	4.0	5.7	9.6	14.2	20.6	38.0	67.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
All races 3/, both sexes										
2004.....	22.4	44.0	77.9	123.9	203.1	310.5	495.9	813.5	1,418.0	3,287.4
2003.....	23.5	45.9	80.7	132.4	218.7	335.0	538.1	886.4	1,538.2	3,610.2
2002.....	23.8	46.4	83.5	140.1	226.2	358.4	569.8	940.2	1,611.2	3,775.0
2001.....	23.4	45.6	84.8	142.8	236.6	372.4	591.8	984.3	1,685.2	3,958.2
2000.....	23.6	48.0	86.7	153.9	250.0	390.8	628.0	1,032.5	1,761.7	4,182.5
1999.....	24.2	48.5	90.1	157.2	260.0	420.1	661.9	1,081.1	1,841.7	4,318.1
All races 3/, male										
2004.....	34.0	68.2	121.1	187.5	297.8	440.2	682.0	1,106.0	1,855.0	3,680.3
2003.....	35.5	71.4	126.9	200.3	319.7	476.1	744.1	1,200.1	2,013.5	4,005.2
2002.....	36.5	72.3	130.3	211.4	331.4	508.5	791.5	1,274.8	2,094.2	4,205.3
2001.....	35.8	71.6	131.1	215.5	344.4	525.7	820.3	1,337.5	2,180.0	4,395.8
2000.....	37.0	76.9	134.2	232.8	362.9	550.5	879.6	1,394.4	2,305.0	4,701.7
1999.....	37.9	77.0	140.3	237.7	376.1	595.2	924.6	1,462.5	2,400.4	4,837.5
All races 3/, female										
2004.....	10.8	20.4	36.6	64.0	116.8	197.7	346.0	603.2	1,148.8	3,110.6
2003.....	11.6	21.1	36.5	68.4	126.9	212.6	373.0	663.1	1,247.5	3,435.5
2002.....	11.2	21.4	38.7	73.1	130.7	228.8	392.6	704.0	1,317.5	3,588.3
2001.....	11.1	20.3	40.3	74.8	138.8	240.5	410.2	737.0	1,388.7	3,775.1
2000.....	10.3	20.0	41.3	80.0	147.7	253.9	429.9	780.5	1,441.2	3,971.0
1999.....	10.7	20.8	42.1	82.1	155.0	270.2	456.7	815.9	1,514.7	4,108.8
White, both sexes										
2004.....	21.6	42.0	73.4	117.5	195.2	298.7	485.8	807.0	1,423.8	3,343.9
2003.....	22.4	43.5	76.1	126.4	208.2	322.6	526.3	877.2	1,542.2	3,671.7
2002.....	22.9	44.2	79.1	133.3	216.1	346.6	559.2	928.9	1,616.4	3,836.4
2001.....	22.1	43.1	80.3	135.4	225.8	357.9	580.5	972.9	1,689.9	4,022.7
2000.....	22.2	45.3	82.4	146.2	238.5	378.4	615.0	1,019.7	1,768.5	4,257.7
1999.....	22.6	46.0	85.4	149.1	249.5	408.5	647.3	1,070.6	1,849.5	4,389.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Ischemic heart diseases (I20-I25)

White, male

2004.....	33.7	66.3	116.1	180.0	289.4	428.5	673.9	1,105.8	1,880.8	3,768.8
2003.....	34.4	68.8	121.6	194.2	308.7	464.4	731.5	1,195.8	2,028.8	4,106.1
2002.....	35.9	70.3	125.6	203.9	320.5	498.1	784.6	1,273.1	2,115.1	4,299.7
2001.....	34.6	69.5	126.4	207.2	333.9	512.8	810.3	1,335.1	2,201.7	4,494.0
2000.....	35.9	74.5	130.0	223.7	352.1	541.2	870.1	1,390.0	2,328.3	4,813.9
1999.....	36.2	74.7	135.0	228.2	364.4	586.0	913.4	1,463.9	2,429.7	4,933.9

White, female

2004.....	9.4	17.9	31.5	57.3	107.8	183.6	331.3	588.5	1,140.0	3,152.3
2003.....	10.4	18.4	31.5	61.3	115.2	197.3	358.4	646.9	1,242.2	3,479.3
2002.....	9.8	18.2	33.6	65.7	119.6	213.2	375.5	682.5	1,310.6	3,635.2
2001.....	9.5	17.0	35.2	66.8	126.1	222.0	394.4	716.3	1,380.6	3,825.5
2000.....	8.5	16.4	36.0	72.5	133.8	236.0	410.4	759.2	1,435.5	4,031.8
1999.....	8.9	17.6	37.0	73.8	143.6	253.7	435.9	794.5	1,507.0	4,170.3

Black, both sexes

2004.....	33.2	67.4	130.6	207.3	320.6	471.5	691.7	1,030.8	1,554.7	3,014.3
2003.....	35.2	71.5	133.3	213.8	352.2	503.0	756.8	1,139.8	1,696.1	3,310.1
2002.....	35.1	72.8	136.5	230.9	362.0	522.5	779.6	1,196.0	1,768.0	3,462.9
2001.....	36.5	72.5	138.8	236.1	376.4	562.5	810.8	1,249.3	1,837.2	3,615.1
2000.....	37.1	78.2	140.3	250.9	392.9	566.1	866.5	1,296.9	1,882.8	3,684.7
1999.....	39.7	76.9	149.3	258.5	395.4	596.2	906.3	1,321.5	1,926.9	3,805.8

Black, male

2004.....	44.9	96.6	189.8	297.2	454.8	641.3	922.9	1,332.6	1,860.9	3,100.5
2003.....	50.2	103.2	195.6	302.7	490.5	677.1	1,027.8	1,501.3	2,138.8	3,285.3
2002.....	48.4	101.6	196.9	330.3	509.6	703.3	1,024.8	1,495.0	2,171.1	3,556.2
2001.....	50.8	101.9	199.9	333.7	515.8	743.8	1,093.0	1,582.5	2,244.8	3,775.0
2000.....	51.8	111.6	197.6	366.7	530.3	727.5	1,139.7	1,630.6	2,316.5	3,885.2
1999.....	56.8	110.4	217.6	372.5	553.5	788.8	1,188.9	1,630.3	2,312.4	4,080.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Ischemic heart diseases (I20-I25)

Black, female

2004.....	22.9	42.0	80.5	133.4	215.9	345.4	534.9	852.0	1,393.3	2,980.1
2003.....	21.9	43.9	80.5	140.7	244.3	373.7	574.0	924.9	1,464.2	3,319.7
2002.....	23.5	47.9	85.4	149.3	246.6	388.3	615.0	1,017.3	1,558.1	3,427.2
2001.....	23.9	47.1	86.8	156.0	266.7	428.6	622.8	1,050.3	1,629.6	3,555.7
2000.....	24.1	49.2	91.4	156.0	284.5	447.0	685.1	1,095.7	1,664.3	3,610.0
1999.....	24.7	48.0	91.4	165.5	271.4	453.5	718.8	1,135.4	1,735.9	3,703.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute myocardial infarction (I21-I22)

All races 3/, both sexes

2004.....	8.9	17.7	31.6	50.6	82.7	124.1	187.7	291.1	476.0	992.9
2003.....	9.6	18.8	33.2	55.5	90.8	134.5	207.4	325.6	532.5	1,118.8
2002.....	9.9	19.0	34.9	58.4	96.2	148.5	227.5	352.5	572.7	1,191.3
2001.....	9.9	19.7	36.5	61.3	100.1	154.1	235.6	374.0	611.5	1,265.4
2000.....	9.9	21.1	37.8	67.0	110.9	166.8	254.7	405.3	650.1	1,338.5
1999.....	10.4	21.5	39.0	69.7	114.9	181.7	271.6	422.9	689.4	1,379.3

All races 3/, male

2004.....	13.2	27.4	48.7	76.3	120.6	173.0	254.1	386.8	608.9	1,120.5
2003.....	14.4	29.2	52.1	83.4	131.6	188.5	281.4	430.0	682.4	1,263.5
2002.....	14.8	29.2	53.7	86.6	139.0	207.6	312.0	466.1	726.6	1,366.4
2001.....	15.0	30.8	55.6	91.3	143.5	214.2	322.0	495.6	776.1	1,461.3
2000.....	15.5	33.6	57.5	99.7	158.6	231.3	349.6	535.2	835.8	1,568.3
1999.....	16.0	33.7	60.9	105.3	164.9	253.5	373.8	564.2	885.1	1,622.1

All races 3/, female

2004.....	4.5	8.3	15.3	26.4	48.2	81.5	134.3	222.3	394.1	935.5
2003.....	5.0	8.7	15.1	29.3	53.8	87.7	148.0	251.3	440.9	1,054.8
2002.....	5.1	9.1	17.0	32.0	57.4	97.6	159.8	272.2	479.1	1,115.3
2001.....	4.9	9.0	18.2	33.2	60.7	102.4	167.0	288.8	512.9	1,183.5
2000.....	4.4	8.9	18.9	36.5	67.7	111.5	179.9	314.9	540.6	1,244.8
1999.....	4.9	9.7	18.0	36.5	69.6	120.2	191.7	324.7	574.9	1,281.5

White, both sexes

2004.....	8.7	17.4	30.6	49.4	81.8	121.7	185.3	291.1	478.6	1,004.1
2003.....	9.3	18.2	32.1	54.4	89.1	132.3	205.3	324.2	532.9	1,132.6
2002.....	9.7	18.4	34.2	57.3	94.3	146.4	225.0	349.5	572.7	1,201.2
2001.....	9.5	19.3	35.4	59.8	98.4	150.8	233.3	371.9	612.7	1,277.8
2000.....	9.5	20.6	36.7	65.6	108.9	164.1	251.8	401.7	652.6	1,353.5
1999.....	10.0	21.1	38.3	68.2	113.4	180.6	268.3	421.1	692.5	1,393.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute myocardial infarction (I21-I22)

White, male

2004.....	13.4	27.2	48.0	75.4	121.2	171.7	252.9	391.6	618.6	1,143.7
2003.....	14.1	28.7	51.2	83.0	130.8	188.3	280.0	430.3	685.7	1,289.3
2002.....	14.7	29.0	53.3	86.2	138.7	207.7	313.1	467.1	730.8	1,386.1
2001.....	14.9	30.9	55.1	90.6	143.7	213.3	321.9	497.6	784.0	1,481.1
2000.....	15.3	33.6	57.2	98.8	158.3	231.1	349.3	536.3	843.5	1,592.9
1999.....	15.8	33.8	60.8	104.0	165.1	254.3	374.0	568.8	896.6	1,644.0

White, female

2004.....	3.9	7.5	13.5	24.3	45.2	77.4	129.7	217.6	391.6	941.1
2003.....	4.5	7.7	13.4	26.8	50.5	82.8	144.1	247.5	438.7	1,063.2
2002.....	4.6	8.0	15.4	29.7	53.3	92.5	153.2	265.3	475.8	1,120.8
2001.....	4.2	7.7	16.1	30.3	56.7	96.0	161.5	282.8	509.1	1,192.7
2000.....	3.7	7.7	16.6	34.1	63.5	105.5	173.6	307.0	539.1	1,256.2
1999.....	4.3	8.4	16.2	34.2	65.8	116.3	184.3	317.3	572.1	1,292.4

Black, both sexes

2004.....	12.4	25.0	47.9	75.3	112.7	169.2	249.7	351.1	520.9	979.9
2003.....	14.4	27.5	49.5	79.4	124.3	180.3	270.2	401.4	597.0	1,094.9
2002.....	13.9	27.7	50.4	84.2	136.3	193.6	295.9	434.1	643.9	1,201.2
2001.....	14.3	27.8	53.2	89.2	135.9	211.1	304.3	452.3	671.2	1,239.9
2000.....	15.1	29.6	55.5	94.7	151.4	219.0	328.8	490.6	698.8	1,275.7
1999.....	15.4	30.0	54.9	96.9	150.2	224.0	347.5	495.1	729.6	1,318.8

Black, male

2004.....	15.4	36.3	67.6	104.8	150.1	223.1	322.7	419.5	591.1	1,009.3
2003.....	20.0	38.6	70.9	108.1	167.9	230.2	353.4	516.2	736.1	1,113.5
2002.....	19.0	37.1	70.4	114.5	177.6	247.8	370.2	531.4	776.4	1,280.1
2001.....	18.5	36.4	71.2	120.1	174.5	264.8	386.4	548.7	792.2	1,382.0
2000.....	20.6	41.0	72.7	131.4	196.7	270.9	419.2	592.5	856.5	1,418.5
1999.....	21.1	41.1	76.9	138.8	198.8	290.8	435.5	591.7	858.0	1,498.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute myocardial infarction (I21-I22)

Black, female

2004.....	9.7	15.2	31.2	51.0	83.6	129.2	200.1	310.5	483.8	968.3
2003.....	9.5	18.0	31.5	55.7	90.3	143.3	214.0	333.2	524.2	1,087.7
2002.....	9.3	19.5	33.4	59.4	104.1	153.4	246.0	376.0	574.9	1,171.0
2001.....	10.5	20.3	37.8	63.8	105.6	171.4	249.5	394.7	609.5	1,187.2
2000.....	10.1	19.8	40.8	64.6	115.7	180.7	268.8	429.2	619.4	1,222.5
1999.....	10.4	20.5	36.2	62.6	112.1	174.5	289.2	436.8	665.9	1,251.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other acute ischemic heart diseases (I24)

All races 3/, both sexes

2004.....	0.3	0.4	0.7	1.1	1.4	1.8	2.6	4.2	6.5	14.8
2003.....	0.3	0.5	0.9	1.3	1.7	2.7	3.6	5.2	8.3	20.0
2002.....	0.3	0.6	1.2	1.4	2.2	3.0	3.9	5.7	9.0	21.1
2001.....	0.3	0.6	1.0	1.4	2.4	2.9	4.1	5.3	9.2	21.5
2000.....	0.3	0.5	1.1	1.6	2.5	3.1	4.0	5.9	9.4	21.9
1999.....	0.3	0.5	0.9	1.6	2.2	3.1	4.3	5.5	9.2	21.9

All races 3/, male

2004.....	0.4	0.5	1.0	1.6	2.0	2.7	3.4	5.0	8.3	16.3
2003.....	0.5	0.7	1.3	1.8	2.4	3.8	4.9	7.1	9.4	19.5
2002.....	0.4	0.8	1.8	2.1	3.0	4.3	5.1	7.0	11.3	21.4
2001.....	0.4	0.9	1.6	2.0	3.5	4.1	5.5	6.7	11.3	20.7
2000.....	0.4	0.8	1.6	2.5	3.5	4.4	5.2	7.6	11.7	20.7
1999.....	0.4	0.7	1.3	2.5	3.1	4.3	5.6	7.6	10.8	22.3

All races 3/, female

2004.....	*	0.2	0.4	0.6	0.8	1.0	1.9	3.6	5.4	14.1
2003.....	*	0.3	0.5	0.8	1.0	1.8	2.6	3.9	7.6	20.2
2002.....	*	0.3	0.7	0.7	1.4	1.9	3.0	4.7	7.6	21.0
2001.....	*	0.3	0.5	0.9	1.3	2.0	3.1	4.4	8.0	21.8
2000.....	*	*	0.5	0.8	1.5	1.9	2.9	4.8	8.1	22.4
1999.....	*	0.2	0.6	0.7	1.5	2.0	3.2	4.1	8.3	21.7

White, both sexes

2004.....	0.3	0.3	0.6	0.9	1.3	1.6	2.4	3.9	6.3	15.0
2003.....	0.3	0.4	0.8	1.1	1.6	2.6	3.4	5.0	8.1	20.2
2002.....	0.2	0.5	1.1	1.3	2.0	2.9	3.8	5.6	8.9	21.4
2001.....	0.2	0.5	0.9	1.3	2.1	2.7	4.0	5.2	9.2	21.5
2000.....	0.2	0.4	1.0	1.4	2.2	2.9	3.8	5.6	9.3	22.1
1999.....	0.2	0.3	0.8	1.4	2.1	3.0	4.0	5.4	8.9	22.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other acute ischemic heart diseases (I24)

White, male

2004.....	0.4	0.4	0.8	1.4	1.8	2.3	3.1	4.6	8.3	16.2
2003.....	0.4	0.6	1.2	1.5	2.2	3.6	4.8	6.7	8.9	19.7
2002.....	0.4	0.7	1.6	2.0	2.8	4.1	4.8	7.0	11.1	21.9
2001.....	0.3	0.8	1.3	1.8	3.1	3.5	5.4	6.5	11.6	20.7
2000.....	0.4	0.7	1.5	2.2	3.2	4.0	4.9	7.2	11.6	21.0
1999.....	0.3	0.5	1.2	2.2	2.8	4.1	5.2	7.6	10.3	22.9

White, female

2004.....	*	*	0.3	0.5	0.8	1.0	1.9	3.4	5.1	14.4
2003.....	*	0.2	0.4	0.7	0.9	1.6	2.4	3.8	7.7	20.5
2002.....	*	0.3	0.6	0.7	1.2	1.8	2.9	4.6	7.5	21.1
2001.....	*	0.2	0.4	0.8	1.2	1.9	2.9	4.3	7.7	21.7
2000.....	*	*	0.5	0.6	1.3	1.9	2.8	4.5	8.0	22.6
1999.....	*	*	0.5	0.7	1.5	2.0	3.0	3.8	8.1	22.1

Black, both sexes

2004.....	*	1.0	1.9	2.9	3.0	3.9	4.3	8.1	9.9	15.2
2003.....	0.7	1.1	1.5	2.7	3.3	4.9	6.2	8.7	11.7	19.8
2002.....	*	1.1	2.4	2.1	4.1	5.0	7.1	7.7	11.8	21.1
2001.....	0.7	1.1	2.1	2.8	5.2	5.3	6.2	6.9	10.8	24.0
2000.....	*	1.0	2.1	3.8	4.7	5.2	6.1	10.0	12.1	21.9
1999.....	*	1.5	2.2	3.3	3.9	4.3	7.3	8.7	13.9	17.2

Black, male

2004.....	*	*	2.7	4.5	4.7	7.0	7.1	11.8	*	22.8
2003.....	*	1.7	2.2	4.6	5.0	7.4	7.9	14.2	19.2	24.0
2002.....	*	1.9	3.6	3.1	5.7	8.0	10.7	9.2	16.0	21.4
2001.....	*	*	3.6	3.4	8.1	9.1	7.7	11.2	*	23.8
2000.....	*	*	3.6	5.8	6.6	8.3	9.0	13.2	*	*
1999.....	*	2.5	2.8	5.8	6.6	6.8	10.2	10.0	17.4	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other acute ischemic heart diseases (I24)

Black, female

2004.....	*	*	*	*	*	*	*	5.9	9.1	12.2
2003.....	*	*	*	*	*	*	5.1	5.4	7.7	18.2
2002.....	*	*	*	*	*	*	4.7	6.9	9.5	20.9
2001.....	*	*	*	*	*	*	5.2	*	11.9	24.0
2000.....	*	*	*	*	3.3	*	*	8.0	10.6	22.3
1999.....	*	*	*	*	*	*	5.4	8.0	12.1	16.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other forms of chronic ischemic heart disease (I20,I25)

All races 3/, both sexes

2004.....	13.2	25.9	45.6	72.2	119.0	184.6	305.6	518.2	935.6	2,279.7
2003.....	13.5	26.6	46.6	75.6	126.1	197.8	327.1	555.6	997.3	2,471.4
2002.....	13.6	26.8	47.3	80.2	127.8	206.8	338.4	582.1	1,029.5	2,562.5
2001.....	13.2	25.2	47.3	80.1	134.1	215.4	352.0	605.0	1,064.5	2,671.3
2000.....	13.4	26.5	47.9	85.2	136.6	221.0	369.4	621.2	1,102.2	2,822.1
1999.....	13.5	26.5	50.2	85.9	142.9	235.3	386.1	652.6	1,143.1	2,917.0

All races 3/, male

2004.....	20.4	40.3	71.4	109.5	175.2	264.4	424.5	714.2	1,237.9	2,543.6
2003.....	20.7	41.5	73.5	115.1	185.6	283.8	457.8	763.1	1,321.6	2,722.2
2002.....	21.4	42.2	74.9	122.7	189.4	296.6	474.3	801.6	1,356.2	2,817.4
2001.....	20.4	40.0	74.0	122.2	197.4	307.4	492.9	835.3	1,392.6	2,913.8
2000.....	21.1	42.5	75.1	130.6	200.8	314.8	524.7	851.5	1,457.6	3,112.7
1999.....	21.5	42.6	78.1	129.9	208.1	337.3	545.2	890.7	1,504.6	3,193.1

All races 3/, female

2004.....	6.1	11.9	20.9	37.0	67.8	115.1	209.8	377.3	749.3	2,161.0
2003.....	6.5	12.1	20.9	38.4	72.0	123.2	222.4	408.0	799.1	2,360.5
2002.....	6.0	11.9	21.0	40.3	72.0	129.4	229.7	427.1	830.8	2,451.9
2001.....	6.1	11.0	21.7	40.6	76.8	136.2	240.1	443.8	867.9	2,569.8
2000.....	5.8	10.9	21.8	42.7	78.5	140.5	247.0	460.8	892.5	2,703.8
1999.....	5.6	10.9	23.4	44.9	83.9	148.1	261.7	487.1	931.6	2,805.7

White, both sexes

2004.....	12.6	24.4	42.2	67.2	112.1	175.4	298.1	512.0	939.0	2,324.8
2003.....	12.9	24.9	43.2	71.0	117.6	187.7	317.6	548.0	1,001.1	2,518.8
2002.....	13.0	25.2	43.9	74.7	119.8	197.3	330.5	573.8	1,034.7	2,613.9
2001.....	12.3	23.3	44.0	74.3	125.3	204.5	343.2	595.8	1,068.1	2,723.5
2000.....	12.5	24.3	44.8	79.2	127.3	211.4	359.4	612.4	1,106.6	2,882.1
1999.....	12.3	24.6	46.3	79.4	134.0	225.0	375.0	644.2	1,148.1	2,973.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other forms of chronic ischemic heart disease (I20,I25)

White, male

2004.....	19.9	38.7	67.2	103.3	166.4	254.5	417.9	709.7	1,254.0	2,608.9
2003.....	19.9	39.5	69.2	109.6	175.8	272.6	446.7	758.8	1,334.2	2,797.1
2002.....	20.8	40.6	70.7	115.8	179.1	286.2	466.8	799.0	1,373.2	2,891.6
2001.....	19.4	37.7	69.9	114.8	187.1	296.0	483.0	831.0	1,406.1	2,992.2
2000.....	20.3	40.1	71.3	122.7	190.6	306.1	515.9	846.5	1,473.2	3,200.0
1999.....	20.1	40.4	72.9	122.0	196.5	327.6	534.1	887.5	1,522.9	3,267.1

White, female

2004.....	5.3	10.2	17.7	32.5	61.8	105.2	199.7	367.4	743.3	2,196.8
2003.....	5.8	10.5	17.7	33.8	63.8	112.8	211.9	395.6	795.8	2,395.5
2002.....	5.1	10.0	17.6	35.3	65.1	118.9	219.3	412.6	827.2	2,493.2
2001.....	5.2	9.0	18.7	35.8	68.2	124.1	230.0	429.2	863.8	2,611.0
2000.....	4.7	8.6	18.8	37.8	69.0	128.7	233.9	447.7	888.4	2,753.0
1999.....	4.5	9.0	20.3	38.9	76.3	135.5	248.5	473.4	926.8	2,855.7

Black, both sexes

2004.....	20.3	41.4	80.8	129.1	204.8	298.4	437.7	671.6	1,023.9	2,019.1
2003.....	20.1	42.8	82.2	131.7	224.6	317.7	480.4	729.7	1,087.4	2,195.3
2002.....	20.7	44.0	83.7	144.6	221.6	323.9	476.6	754.2	1,112.3	2,240.6
2001.....	21.5	43.7	83.6	144.1	235.3	346.1	500.3	790.1	1,155.2	2,351.2
2000.....	21.5	47.6	82.7	152.4	236.7	341.9	531.5	796.3	1,171.8	2,387.1
1999.....	23.8	45.4	92.2	158.4	241.2	367.9	551.4	817.7	1,183.5	2,469.8

Black, male

2004.....	28.8	59.1	119.5	187.9	300.1	411.1	593.0	901.3	1,258.3	2,068.3
2003.....	29.4	62.9	122.5	189.9	317.6	439.5	666.5	970.9	1,383.5	2,147.8
2002.....	29.0	62.5	122.8	212.7	326.3	447.5	643.9	954.4	1,378.6	2,254.6
2001.....	31.2	64.0	125.1	210.1	333.1	469.8	698.8	1,022.7	1,443.9	2,369.1
2000.....	30.3	69.2	121.3	229.5	326.9	448.3	711.4	1,024.9	1,444.9	2,446.0
1999.....	34.9	66.9	138.0	227.8	348.1	491.2	743.2	1,028.6	1,437.0	2,561.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other forms of chronic ischemic heart disease (I20,I25)

Black, female

2004.....	12.8	26.0	48.2	80.7	130.5	214.6	332.3	535.6	900.4	1,999.6
2003.....	11.9	25.3	48.2	83.8	152.1	227.3	354.9	586.3	932.3	2,213.8
2002.....	13.3	28.0	50.5	88.6	139.7	232.2	364.3	634.5	973.6	2,235.2
2001.....	13.0	26.1	48.2	89.9	158.2	254.8	368.1	651.2	1,008.1	2,344.5
2000.....	13.7	28.8	49.7	89.2	165.6	263.4	412.0	658.5	1,034.3	2,365.2
1999.....	14.1	26.8	53.5	101.6	157.4	276.5	424.2	690.6	1,057.9	2,435.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All races 3/, both sexes

2004.....	4.8	9.5	18.2	26.6	39.1	49.8	69.0	100.8	166.6	400.4
2003.....	4.7	9.9	18.6	26.6	40.5	53.3	73.3	110.7	186.2	442.9
2002.....	4.8	9.8	17.5	27.9	39.7	53.9	76.5	116.6	187.6	474.5
2001.....	4.6	8.8	17.4	27.7	40.6	55.3	76.7	121.8	197.2	493.6
2000.....	4.4	9.4	17.5	28.5	40.7	55.7	81.9	125.0	207.8	534.6
1999.....	4.7	9.1	17.4	28.6	43.4	59.6	86.8	131.6	218.2	563.4

All races 3/, male

2004.....	7.7	15.3	29.6	42.0	58.4	73.1	96.4	133.8	210.5	414.0
2003.....	7.4	16.0	30.5	41.8	60.6	77.1	102.8	146.6	233.1	462.1
2002.....	7.8	15.6	28.4	44.8	59.7	78.8	106.7	154.5	232.4	491.8
2001.....	7.2	14.6	28.3	43.4	61.8	79.0	107.6	160.2	247.4	502.0
2000.....	7.1	15.5	28.4	45.8	61.3	80.8	115.6	167.1	252.8	550.8
1999.....	7.6	14.9	28.0	45.0	64.6	87.3	120.8	172.7	264.3	578.6

All races 3/, female

2004.....	2.0	3.9	7.4	12.0	21.5	29.5	47.0	77.0	139.6	394.3
2003.....	2.0	4.0	7.2	12.4	22.2	32.6	49.7	85.2	157.6	434.4
2002.....	1.8	4.1	7.1	12.1	21.4	32.4	52.5	89.7	160.4	467.1
2001.....	2.1	3.3	7.0	12.9	21.4	34.9	52.1	94.9	167.1	490.1
2000.....	1.8	3.5	7.2	12.4	22.1	34.1	55.4	95.7	181.2	528.0
1999.....	1.8	3.5	7.3	13.4	24.1	36.0	60.2	103.1	191.3	557.2

White, both sexes

2004.....	4.6	8.7	16.4	23.7	35.1	44.4	63.4	93.3	160.3	395.6
2003.....	4.4	9.0	16.7	24.2	35.4	47.5	66.6	103.0	179.6	439.3
2002.....	4.4	9.1	15.6	24.7	35.3	47.3	69.7	107.8	180.5	470.6
2001.....	4.2	7.9	15.6	23.9	35.0	47.9	69.4	112.2	190.6	489.7
2000.....	4.0	8.3	15.7	24.8	35.2	49.5	73.7	116.0	198.5	529.5
1999.....	4.2	8.0	15.2	24.6	37.5	52.3	77.9	121.5	209.8	557.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

White, male

2004.....	7.5	14.4	27.3	37.8	52.4	66.1	89.3	123.9	203.3	408.8
2003.....	6.9	14.8	27.7	38.3	53.4	69.2	92.5	136.9	224.4	458.7
2002.....	7.4	14.7	25.7	40.0	53.6	69.4	98.0	143.3	222.2	488.0
2001.....	6.6	13.3	25.7	38.3	54.0	69.3	97.6	148.7	239.9	498.6
2000.....	6.6	14.0	25.9	40.0	54.2	72.8	104.4	154.6	241.6	545.6
1999.....	7.0	13.2	24.8	39.4	56.2	77.6	109.0	159.8	253.9	571.2

White, female

2004.....	1.7	3.0	5.8	10.2	19.1	25.1	42.1	70.9	133.6	389.7
2003.....	1.8	3.2	5.9	10.6	18.7	28.3	45.4	78.5	152.0	430.7
2002.....	1.5	3.5	5.6	10.2	18.4	27.8	46.6	82.5	154.9	463.0
2001.....	1.8	2.6	5.7	10.1	17.6	29.1	46.7	86.4	160.8	486.0
2000.....	1.4	2.7	5.8	10.2	17.6	29.0	49.0	88.8	172.9	522.9
1999.....	1.3	2.8	5.8	10.6	20.3	30.3	53.1	94.6	183.8	552.0

Black, both sexes

2004.....	7.5	17.5	35.7	55.9	81.5	105.1	139.0	198.5	268.4	505.1
2003.....	7.3	17.4	35.9	53.2	91.8	111.4	149.2	214.1	282.8	543.2
2002.....	8.1	17.0	35.9	60.5	85.5	119.5	157.5	231.9	300.4	575.5
2001.....	8.2	16.8	35.7	64.2	94.4	128.0	162.6	244.0	301.5	594.5
2000.....	7.4	19.4	36.4	65.4	94.0	119.6	176.8	239.7	340.0	646.6
1999.....	9.1	19.1	38.4	67.8	98.2	133.9	188.9	261.7	338.9	668.6

Black, male

2004.....	10.7	25.1	54.0	88.3	127.8	153.1	197.1	278.0	350.8	531.5
2003.....	11.4	26.4	57.8	81.3	140.0	161.3	228.9	301.3	380.2	583.9
2002.....	11.8	25.5	55.0	97.1	130.5	180.0	220.3	321.0	417.3	599.3
2001.....	12.4	25.7	56.0	97.0	144.0	183.5	241.0	330.4	388.5	609.3
2000.....	11.0	30.3	56.8	106.4	137.1	168.6	257.1	338.3	425.7	701.2
1999.....	13.7	29.9	59.7	103.8	147.9	197.3	271.0	350.0	435.1	715.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Black, female

2004.....	4.7	10.9	20.2	29.3	45.3	69.4	99.5	151.5	224.9	494.6
2003.....	3.7	9.6	17.3	30.0	54.2	74.4	95.5	162.2	231.7	527.4
2002.....	4.8	9.7	19.7	30.4	50.3	74.6	115.4	178.7	239.5	566.4
2001.....	4.5	9.0	18.4	37.3	55.4	87.1	110.3	192.5	257.1	589.0
2000.....	4.2	10.0	19.0	31.8	60.0	83.5	123.4	180.3	296.8	626.3
1999.....	5.0	9.8	20.4	38.4	59.2	87.0	134.3	208.5	291.3	650.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races 3/, both sexes

2004.....	8.4	16.4	27.4	45.7	79.9	134.8	236.6	417.4	769.0	1,879.3
2003.....	8.9	16.7	28.0	49.0	85.6	144.5	253.8	444.9	811.1	2,028.5
2002.....	8.8	17.1	29.9	52.3	88.2	152.9	261.8	465.6	841.9	2,088.0
2001.....	8.6	16.4	29.8	52.4	93.5	160.1	275.3	483.2	867.3	2,177.7
2000.....	9.0	17.0	30.3	56.7	95.9	165.3	287.4	496.2	894.4	2,287.5
1999.....	8.8	17.4	32.8	57.3	99.5	175.7	299.3	521.0	924.9	2,353.6

All races 3/, male

2004.....	12.7	25.0	41.8	67.5	116.8	191.3	328.2	580.4	1,027.4	2,129.6
2003.....	13.3	25.5	43.0	73.3	125.0	206.7	355.0	616.5	1,088.5	2,260.1
2002.....	13.6	26.6	46.5	78.0	129.6	217.8	367.6	647.1	1,123.8	2,325.6
2001.....	13.2	25.4	45.7	78.8	135.6	228.4	385.2	675.1	1,145.1	2,411.8
2000.....	14.0	27.0	46.7	84.9	139.5	234.0	409.1	684.5	1,204.8	2,561.9
1999.....	13.9	27.7	50.2	84.9	143.5	250.0	424.4	718.0	1,240.3	2,614.4

All races 3/, female

2004.....	4.1	8.0	13.6	25.0	46.3	85.6	162.8	300.3	609.8	1,766.7
2003.....	4.5	8.2	13.7	26.0	49.8	90.5	172.7	322.7	641.5	1,926.2
2002.....	4.2	7.8	13.9	28.2	50.5	97.0	177.3	337.3	670.4	1,984.9
2001.....	4.0	7.6	14.7	27.7	55.4	101.2	188.0	348.9	700.7	2,079.8
2000.....	4.0	7.3	14.6	30.3	56.4	106.3	191.6	365.1	711.3	2,175.8
1999.....	3.9	7.4	16.1	31.5	59.7	112.1	201.5	384.0	740.3	2,248.4

White, both sexes

2004.....	8.1	15.7	25.8	43.5	77.0	131.0	234.7	418.8	778.7	1,929.2
2003.....	8.5	16.0	26.5	46.8	82.2	140.3	251.0	445.0	821.5	2,079.5
2002.....	8.6	16.1	28.3	49.9	84.5	150.0	260.7	466.0	854.2	2,143.3
2001.....	8.1	15.4	28.4	50.5	90.2	156.6	273.8	483.6	877.5	2,233.8
2000.....	8.5	16.0	29.1	54.4	92.2	162.0	285.8	496.4	908.0	2,352.7
1999.....	8.2	16.6	31.1	54.8	96.5	172.6	297.1	522.7	938.2	2,416.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
White, male										
2004.....	12.4	24.3	40.0	65.5	114.0	188.4	328.6	585.8	1,050.7	2,200.2
2003.....	12.9	24.7	41.5	71.3	122.4	203.4	354.1	622.0	1,109.8	2,338.4
2002.....	13.4	25.9	45.0	75.8	125.4	216.8	368.7	655.7	1,151.0	2,403.6
2001.....	12.8	24.4	44.2	76.5	133.1	226.8	385.4	682.4	1,166.2	2,493.6
2000.....	13.7	26.1	45.5	82.7	136.4	233.3	411.5	691.9	1,231.6	2,654.4
1999.....	13.1	27.2	48.1	82.6	140.3	249.9	425.1	727.7	1,269.0	2,695.9
White, female										
2004.....	3.7	7.2	11.9	22.3	42.8	80.1	157.6	296.5	609.7	1,807.0
2003.....	4.0	7.3	11.8	23.3	45.1	84.5	166.6	317.1	643.8	1,964.9
2002.....	3.6	6.5	12.0	25.2	46.7	91.1	172.7	330.1	672.3	2,030.2
2001.....	3.5	6.4	13.0	25.7	50.7	95.0	183.3	342.8	703.0	2,125.0
2000.....	3.3	5.9	13.0	27.5	51.4	99.6	184.9	358.9	715.6	2,230.1
1999.....	3.2	6.2	14.5	28.3	56.1	105.2	195.4	378.8	743.0	2,303.8
Black, both sexes										
2004.....	12.8	23.9	45.2	73.2	123.4	193.3	298.8	473.1	755.5	1,514.0
2003.....	12.8	25.4	46.4	78.5	132.8	206.3	331.2	515.6	804.7	1,652.1
2002.....	12.6	27.0	47.8	84.1	136.1	204.4	319.1	522.2	811.9	1,665.1
2001.....	13.3	26.9	47.9	79.9	140.9	218.1	337.8	546.0	853.7	1,756.7
2000.....	14.1	28.1	46.3	87.0	142.8	222.3	354.7	556.6	831.8	1,740.5
1999.....	14.7	26.3	53.8	90.6	143.1	233.9	362.6	555.9	844.6	1,801.2
Black, male										
2004.....	18.1	34.0	65.5	99.6	172.3	258.0	395.9	623.3	907.5	1,536.8
2003.....	18.0	36.5	64.7	108.6	177.6	278.2	437.5	669.6	1,003.3	1,563.9
2002.....	17.2	37.0	67.8	115.7	195.8	267.5	423.6	633.4	961.4	1,655.3
2001.....	18.8	38.3	69.2	113.1	189.2	286.4	457.8	692.3	1,055.4	1,759.8
2000.....	19.3	39.0	64.5	123.2	189.9	279.7	454.3	686.6	1,019.2	1,744.8
1999.....	21.2	37.0	78.2	124.1	200.3	293.9	472.2	678.6	1,001.9	1,845.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Black, female

2004.....	8.1	15.1	28.0	51.4	85.2	145.2	232.8	384.1	675.5	1,505.0
2003.....	8.3	15.7	30.9	53.8	97.9	153.0	259.4	424.0	700.6	1,686.4
2002.....	8.5	18.4	30.8	58.2	89.4	157.6	248.9	455.8	734.1	1,668.8
2001.....	8.5	17.0	29.8	52.6	102.9	167.6	257.8	458.7	751.0	1,755.5
2000.....	9.5	18.7	30.7	57.4	105.6	179.9	288.6	478.2	737.5	1,738.9
1999.....	9.1	17.1	33.1	63.3	98.2	189.5	289.9	482.0	766.6	1,784.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other heart diseases (I26-I51)

All races 3/, both sexes

2004.....	11.5	17.9	25.6	35.7	57.5	90.3	151.8	268.1	503.3	1,373.9
2003.....	12.6	17.9	25.3	39.1	59.7	97.3	160.2	279.1	525.5	1,427.2
2002.....	12.4	18.1	25.9	39.4	60.6	97.9	164.5	286.7	536.3	1,443.5
2001.....	11.9	17.9	25.9	39.7	62.0	100.7	164.7	287.2	537.0	1,475.5
2000.....	12.3	17.6	25.9	40.2	64.3	103.6	167.1	290.2	542.8	1,509.4
1999.....	12.5	17.8	25.9	41.8	66.5	105.2	175.0	295.4	553.5	1,512.9

All races 3/, male

2004.....	14.4	23.2	32.8	46.4	72.7	111.3	181.2	316.2	573.7	1,377.7
2003.....	15.6	22.7	32.4	49.9	74.9	119.4	195.4	334.0	599.9	1,431.3
2002.....	15.9	23.5	33.6	49.4	75.7	118.8	199.8	340.7	612.4	1,448.6
2001.....	14.9	23.1	32.7	49.4	77.7	121.5	200.9	344.7	622.1	1,473.4
2000.....	15.4	23.0	33.4	52.0	80.8	126.9	205.9	345.1	626.8	1,549.1
1999.....	15.6	23.0	33.3	54.1	84.3	133.4	217.0	350.9	641.4	1,548.9

All races 3/, female

2004.....	8.6	12.8	18.6	25.7	43.8	72.0	128.2	233.5	459.9	1,372.1
2003.....	9.5	13.2	18.6	28.8	45.9	78.1	131.9	240.0	480.0	1,425.4
2002.....	9.0	12.8	18.5	30.1	46.8	79.8	136.3	248.6	490.0	1,441.3
2001.....	8.9	12.8	19.5	30.6	47.8	82.8	135.9	247.0	486.0	1,476.4
2000.....	9.2	12.3	18.7	29.2	49.3	83.6	136.6	252.1	493.2	1,493.2
1999.....	9.4	12.8	18.9	30.2	50.5	81.0	142.2	256.8	502.0	1,498.5

White, both sexes

2004.....	9.7	15.0	21.8	31.3	52.7	84.7	146.5	262.9	507.6	1,408.8
2003.....	10.5	15.2	21.4	34.8	53.8	90.7	153.2	276.4	528.9	1,461.4
2002.....	10.7	15.0	22.1	35.0	55.3	92.4	158.5	282.4	539.1	1,475.5
2001.....	9.7	14.9	22.3	34.8	57.2	94.1	159.0	281.6	538.8	1,509.6
2000.....	10.2	14.6	22.1	35.9	59.0	97.3	160.6	284.5	545.7	1,541.6
1999.....	10.2	14.6	22.2	36.3	60.4	98.6	167.6	288.5	553.7	1,543.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other heart diseases (I26-I51)

White, male

2004.....	12.4	19.6	28.7	40.9	66.8	104.6	176.0	309.6	578.2	1,407.9
2003.....	13.5	19.7	27.5	44.5	68.3	111.8	188.4	332.0	604.3	1,467.6
2002.....	14.1	19.9	29.2	44.1	69.9	112.5	193.0	336.5	617.5	1,482.1
2001.....	12.7	19.7	28.9	43.5	72.1	114.9	193.7	338.2	621.9	1,503.4
2000.....	13.0	19.7	28.6	46.9	74.8	119.7	199.2	339.4	630.8	1,580.5
1999.....	13.3	19.4	28.7	47.5	76.9	125.6	209.0	344.0	641.4	1,571.5

White, female

2004.....	6.9	10.5	15.1	22.0	39.7	66.9	122.2	228.7	463.8	1,409.2
2003.....	7.5	10.8	15.4	25.5	40.4	72.1	124.3	236.1	482.4	1,458.6
2002.....	7.3	10.2	15.2	26.2	41.8	74.7	130.3	243.7	491.0	1,472.6
2001.....	6.6	10.1	15.8	26.5	43.5	75.9	130.9	241.5	488.5	1,512.2
2000.....	7.4	9.6	15.8	25.4	44.3	77.7	129.5	245.9	495.2	1,525.8
1999.....	7.2	9.8	15.8	25.7	45.1	75.0	134.8	249.5	501.9	1,531.9

Black, both sexes

2004.....	26.3	41.7	60.2	82.6	114.5	159.9	240.5	381.0	551.3	1,180.7
2003.....	28.9	40.8	61.1	85.3	124.6	176.1	265.5	364.8	574.9	1,238.1
2002.....	26.6	43.6	60.8	87.1	121.2	169.4	255.8	386.7	594.7	1,270.4
2001.....	29.2	43.1	60.9	91.4	118.8	179.7	254.5	402.9	601.8	1,277.7
2000.....	28.5	43.3	62.0	86.2	122.8	181.5	266.8	406.9	590.7	1,306.0
1999.....	30.2	44.9	62.2	96.8	133.5	185.5	284.2	417.4	621.2	1,302.8

Black, male

2004.....	31.6	54.3	75.0	109.0	149.1	202.5	284.5	469.5	653.5	1,224.9
2003.....	34.7	51.0	79.5	113.5	158.3	223.6	318.0	434.1	668.1	1,220.9
2002.....	32.0	56.7	77.1	110.7	150.7	208.7	317.5	454.2	675.6	1,238.3
2001.....	33.7	53.5	73.7	117.2	151.5	212.9	318.8	494.8	743.2	1,349.1
2000.....	34.8	54.3	82.1	112.1	153.6	224.8	318.9	480.5	696.3	1,402.3
1999.....	36.2	56.1	81.9	127.0	172.7	237.8	353.1	489.1	747.3	1,454.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other heart diseases (I26-I51)

Black, female

2004.....	21.6	30.6	47.7	60.9	87.6	128.3	210.7	328.6	497.4	1,163.2
2003.....	23.8	31.9	45.4	62.1	98.2	140.9	230.2	323.7	526.1	1,244.8
2002.....	21.8	32.3	47.0	67.7	98.2	140.2	214.4	346.3	552.6	1,282.7
2001.....	25.3	34.0	50.1	70.3	93.1	155.1	211.6	348.1	529.8	1,251.2
2000.....	22.9	33.9	44.8	64.9	98.4	149.6	232.2	362.6	537.5	1,270.2
1999.....	24.9	35.2	45.4	72.1	102.8	146.8	238.5	374.2	558.7	1,246.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)

All races 3/, both sexes

2004.....	1.9	2.9	3.7	4.7	7.1	11.0	16.3	23.3	33.1	54.8
2003.....	2.2	2.8	4.0	5.4	8.1	11.6	17.5	25.8	36.9	58.0
2002.....	1.9	2.9	3.8	5.9	8.1	12.7	17.5	25.7	35.5	56.3
2001.....	2.1	2.9	3.9	5.9	8.2	12.0	17.3	24.2	34.8	57.5
2000.....	2.1	2.8	4.2	5.2	8.5	12.3	16.8	24.1	34.7	57.4
1999.....	2.1	2.8	3.9	5.6	8.0	11.9	17.9	25.7	36.5	59.6

All races 3/, male

2004.....	1.7	2.7	3.6	5.1	6.8	10.3	15.2	21.5	30.6	48.3
2003.....	1.8	2.7	4.0	5.1	7.7	11.4	17.5	26.5	35.0	54.4
2002.....	1.6	2.8	3.5	5.3	8.0	12.3	16.3	23.8	33.8	52.1
2001.....	1.9	2.6	3.4	5.1	8.2	11.7	17.6	23.1	35.5	55.6
2000.....	1.7	2.7	4.0	4.9	8.3	11.5	16.6	25.1	33.5	54.8
1999.....	1.8	2.5	3.6	5.3	8.6	12.0	18.5	25.5	35.2	60.3

All races 3/, female

2004.....	2.0	3.1	3.7	4.5	7.3	11.7	17.2	24.5	34.7	57.7
2003.....	2.6	2.8	4.0	5.7	8.4	11.9	17.6	25.3	38.0	59.6
2002.....	2.2	3.1	4.1	6.4	8.3	13.0	18.4	27.1	36.6	58.1
2001.....	2.2	3.1	4.4	6.6	8.2	12.3	17.1	24.9	34.4	58.2
2000.....	2.5	3.0	4.4	5.5	8.8	12.9	17.1	23.5	35.3	58.5
1999.....	2.3	3.1	4.1	5.9	7.5	11.8	17.4	25.8	37.2	59.2

White, both sexes

2004.....	1.6	2.6	3.3	4.3	6.6	10.2	15.5	22.2	32.4	55.3
2003.....	1.8	2.5	3.6	4.9	7.4	10.7	16.1	25.4	36.5	57.8
2002.....	1.7	2.6	3.5	5.3	7.7	12.0	16.7	24.8	35.1	56.1
2001.....	1.8	2.5	3.6	5.3	7.7	11.2	16.7	23.2	34.6	56.8
2000.....	1.8	2.5	3.7	4.9	7.9	11.7	16.1	23.3	34.2	56.4
1999.....	1.7	2.4	3.5	4.9	7.2	11.0	16.7	24.7	35.3	59.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)

White, male

2004.....	1.6	2.5	3.3	4.5	6.5	9.9	14.6	20.5	28.6	48.0
2003.....	1.5	2.4	3.6	4.6	7.3	10.6	16.4	26.3	34.8	54.0
2002.....	1.5	2.5	3.5	4.8	7.9	11.8	15.9	23.2	33.1	51.9
2001.....	1.7	2.3	3.3	4.5	7.7	11.2	17.1	22.2	34.8	53.6
2000.....	1.4	2.5	3.6	4.7	7.8	11.3	16.2	24.6	33.0	52.9
1999.....	1.6	2.3	3.3	4.9	7.8	11.2	17.4	24.7	33.8	60.3

White, female

2004.....	1.6	2.6	3.3	4.0	6.8	10.6	16.2	23.4	34.8	58.6
2003.....	2.2	2.6	3.6	5.2	7.6	10.8	15.9	24.8	37.5	59.5
2002.....	1.9	2.7	3.5	5.8	7.4	12.1	17.4	25.9	36.3	58.0
2001.....	1.9	2.7	3.9	6.1	7.7	11.2	16.3	23.9	34.4	58.1
2000.....	2.1	2.6	3.8	5.2	8.1	12.1	16.0	22.5	35.0	57.8
1999.....	1.9	2.6	3.6	5.0	6.7	10.9	16.1	24.7	36.2	58.8

Black, both sexes

2004.....	4.2	5.9	7.6	10.2	13.3	21.0	29.1	42.8	46.8	62.0
2003.....	5.2	5.6	8.1	11.0	16.0	23.5	36.6	36.9	50.0	72.9
2002.....	3.9	6.4	7.3	12.8	15.0	22.6	30.3	43.6	49.5	68.6
2001.....	4.5	6.3	7.4	12.4	15.3	23.8	28.5	39.8	45.2	77.7
2000.....	4.9	5.8	9.3	8.9	16.3	20.9	29.2	39.3	46.9	76.6
1999.....	4.9	6.2	8.6	12.6	17.4	23.0	35.9	43.7	57.8	72.1

Black, male

2004.....	3.2	4.9	7.1	11.3	11.8	17.7	24.9	41.9	62.4	64.6
2003.....	4.7	5.9	8.2	11.5	14.5	23.1	34.5	37.0	50.2	76.1
2002.....	2.9	6.2	4.9	11.9	11.7	21.7	25.0	37.7	51.2	65.4
2001.....	3.8	5.5	5.2	11.9	16.3	21.1	28.6	38.1	54.6	95.3
2000.....	3.9	4.8	8.5	9.2	15.9	17.2	26.4	38.7	50.6	92.3
1999.....	3.9	5.0	7.9	10.3	18.9	23.9	38.3	43.2	62.8	71.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)

Black, female

2004.....	5.2	6.9	8.0	9.2	14.4	23.4	31.9	43.4	38.6	61.0
2003.....	5.7	5.3	8.0	10.5	17.1	23.8	38.1	36.9	49.9	71.7
2002.....	4.9	6.7	9.3	13.6	17.7	23.3	33.8	47.2	48.5	69.8
2001.....	5.2	7.0	9.3	12.8	14.6	25.7	28.5	40.8	40.5	71.1
2000.....	5.7	6.7	9.9	8.6	16.7	23.6	31.1	39.6	45.0	70.8
1999.....	5.7	7.3	9.2	14.4	16.3	22.3	34.2	44.1	55.3	72.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pulmonary embolism (I26)

All races 3/, both sexes

2004.....	1.2	2.0	2.4	3.0	4.5	6.8	9.7	14.4	20.7	33.4
2003.....	1.5	1.9	2.8	3.5	5.1	7.0	10.7	15.7	23.2	35.5
2002.....	1.3	2.0	2.5	3.9	5.3	8.1	11.0	15.8	22.2	35.4
2001.....	1.4	1.9	2.7	4.1	5.4	7.8	11.4	15.7	22.3	37.8
2000.....	1.4	1.9	2.8	3.5	5.8	7.9	10.8	15.7	22.5	38.4
1999.....	1.4	2.0	2.7	3.9	5.6	7.9	12.6	17.9	25.8	42.0

All races 3/, male

2004.....	1.2	1.9	2.4	3.5	4.7	7.2	9.9	14.3	20.5	30.6
2003.....	1.2	2.0	2.8	3.8	5.4	7.7	11.4	16.9	23.5	34.1
2002.....	1.1	2.0	2.4	3.8	5.5	8.2	10.8	15.0	21.3	34.1
2001.....	1.3	1.8	2.4	3.6	5.6	8.4	11.8	16.1	23.5	38.0
2000.....	1.1	2.0	3.0	3.5	6.0	8.0	11.2	17.4	22.4	37.5
1999.....	1.2	1.9	2.6	4.1	6.3	8.3	13.7	18.1	24.2	42.0

All races 3/, female

2004.....	1.3	2.2	2.4	2.4	4.4	6.6	9.6	14.5	20.8	34.7
2003.....	1.7	1.8	2.7	3.3	4.9	6.4	10.2	14.8	23.0	36.0
2002.....	1.4	1.9	2.6	4.0	5.1	8.0	11.1	16.4	22.8	36.0
2001.....	1.5	2.1	2.9	4.6	5.2	7.3	11.1	15.4	21.5	37.7
2000.....	1.6	1.9	2.6	3.5	5.6	7.8	10.6	14.5	22.5	38.8
1999.....	1.6	2.1	2.8	3.8	5.0	7.5	11.7	17.8	26.7	42.1

White, both sexes

2004.....	1.0	1.8	2.1	2.7	4.3	6.3	9.1	13.7	20.2	33.4
2003.....	1.2	1.7	2.5	3.3	4.8	6.3	9.8	15.5	22.8	34.8
2002.....	1.2	1.8	2.3	3.5	5.1	7.7	10.4	15.1	21.7	34.8
2001.....	1.2	1.7	2.5	3.8	5.1	7.4	10.9	15.0	22.1	37.0
2000.....	1.1	1.7	2.5	3.3	5.4	7.4	10.4	15.0	22.0	37.4
1999.....	1.2	1.7	2.4	3.5	5.1	7.3	11.6	17.0	25.1	41.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pulmonary embolism (I26)

White, male

2004.....	1.1	1.8	2.2	3.2	4.5	6.8	9.3	13.3	19.0	30.0
2003.....	1.1	1.7	2.6	3.4	5.0	7.1	10.7	16.6	23.0	33.2
2002.....	1.1	1.8	2.4	3.4	5.4	7.8	10.6	14.6	20.6	33.3
2001.....	1.1	1.6	2.4	3.3	5.2	8.1	11.5	15.4	22.9	36.1
2000.....	0.9	1.8	2.7	3.2	5.7	7.8	11.0	16.8	21.6	35.8
1999.....	1.1	1.7	2.3	3.7	5.7	7.6	12.8	17.2	23.0	41.3

White, female

2004.....	1.0	1.8	2.1	2.3	4.1	5.8	9.0	13.9	21.0	34.9
2003.....	1.4	1.7	2.4	3.1	4.5	5.6	9.1	14.7	22.6	35.6
2002.....	1.3	1.8	2.3	3.6	4.7	7.6	10.3	15.5	22.4	35.5
2001.....	1.3	1.8	2.5	4.4	5.0	6.8	10.5	14.7	21.6	37.4
2000.....	1.4	1.6	2.3	3.3	5.2	7.1	9.9	13.6	22.2	38.1
1999.....	1.3	1.7	2.5	3.4	4.4	7.1	10.6	16.8	26.4	41.8

Black, both sexes

2004.....	3.0	4.4	4.9	5.8	8.2	13.8	18.1	27.4	30.7	42.6
2003.....	3.3	3.6	5.8	7.3	9.6	15.3	23.6	23.0	34.8	51.1
2002.....	2.4	4.1	5.0	8.5	9.1	14.3	19.4	28.7	33.5	49.0
2001.....	3.1	4.4	5.0	7.9	9.9	14.3	19.2	27.3	30.3	55.9
2000.....	3.3	4.3	6.1	6.0	11.1	14.7	18.5	28.8	33.0	56.6
1999.....	3.2	4.4	6.0	8.8	12.1	15.7	26.4	33.4	40.7	53.2

Black, male

2004.....	2.4	3.6	4.6	7.9	7.8	12.8	17.8	31.8	45.2	45.7
2003.....	3.0	4.5	5.8	8.6	10.4	16.7	22.7	27.4	39.9	57.3
2002.....	1.8	4.6	3.8	8.6	8.5	14.5	16.0	24.8	35.9	51.5
2001.....	2.9	4.0	3.4	8.4	12.0	14.2	18.2	28.8	40.1	72.6
2000.....	3.0	3.9	7.0	7.4	12.0	12.7	17.1	30.2	38.8	71.5
1999.....	2.6	3.8	5.9	8.8	13.4	18.7	29.2	35.6	49.7	60.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pulmonary embolism (I26)

Black, female

2004.....	3.5	5.1	5.1	4.1	8.4	14.6	18.2	24.8	23.1	41.3
2003.....	3.6	2.8	5.8	6.2	9.0	14.3	24.1	20.3	32.1	48.6
2002.....	3.0	3.7	6.1	8.4	9.6	14.2	21.7	31.0	32.2	48.1
2001.....	3.2	4.8	6.4	7.4	8.3	14.3	20.0	26.3	25.3	49.7
2000.....	3.6	4.7	5.4	4.9	10.3	16.1	19.5	27.9	30.1	51.1
1999.....	3.7	5.0	6.1	8.7	11.1	13.5	24.5	32.0	36.3	50.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

All races 3/, both sexes

2004.....	0.6	0.9	1.3	1.8	2.6	4.2	6.6	8.9	12.4	21.4
2003.....	0.8	0.9	1.2	1.8	2.9	4.6	6.8	10.1	13.6	22.6
2002.....	0.6	1.0	1.3	2.0	2.9	4.6	6.5	9.9	13.3	20.9
2001.....	0.7	1.0	1.2	1.8	2.8	4.2	6.0	8.5	12.5	19.6
2000.....	0.7	0.9	1.4	1.7	2.7	4.4	6.0	8.4	12.2	19.0
1999.....	0.7	0.9	1.2	1.7	2.4	4.0	5.4	7.8	10.7	17.5

All races 3/, male

2004.....	0.5	0.8	1.2	1.5	2.2	3.1	5.3	7.2	10.0	17.7
2003.....	0.6	0.7	1.1	1.3	2.3	3.7	6.0	9.6	11.4	20.3
2002.....	0.4	0.7	1.1	1.5	2.5	4.1	5.5	8.8	12.5	18.0
2001.....	0.6	0.9	1.0	1.5	2.6	3.3	5.9	7.0	11.9	17.6
2000.....	0.5	0.7	1.0	1.4	2.3	3.5	5.4	7.7	11.1	17.3
1999.....	0.6	0.7	1.1	1.3	2.3	3.7	4.9	7.4	10.9	18.4

All races 3/, female

2004.....	0.7	0.9	1.4	2.0	3.0	5.1	7.6	10.0	13.9	23.1
2003.....	0.9	1.1	1.3	2.3	3.5	5.5	7.4	10.5	15.0	23.6
2002.....	0.8	1.2	1.5	2.4	3.2	4.9	7.3	10.7	13.8	22.1
2001.....	0.8	1.1	1.5	2.0	3.0	5.0	6.0	9.5	12.9	20.5
2000.....	0.9	1.1	1.8	2.0	3.2	5.2	6.5	8.9	12.8	19.7
1999.....	0.8	1.1	1.4	2.0	2.5	4.3	5.7	8.0	10.5	17.2

White, both sexes

2004.....	0.6	0.8	1.2	1.5	2.4	4.0	6.3	8.6	12.2	21.9
2003.....	0.6	0.8	1.1	1.7	2.7	4.4	6.3	9.9	13.7	23.0
2002.....	0.5	0.8	1.2	1.7	2.6	4.3	6.3	9.7	13.3	21.3
2001.....	0.6	0.8	1.1	1.5	2.6	3.8	5.8	8.2	12.5	19.7
2000.....	0.6	0.9	1.2	1.7	2.5	4.3	5.7	8.4	12.3	19.0
1999.....	0.5	0.7	1.1	1.4	2.1	3.7	5.1	7.7	10.2	17.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

White, male

2004.....	0.5	0.8	1.1	1.4	2.0	3.1	5.2	7.2	9.6	18.0
2003.....	0.5	0.6	1.0	1.2	2.3	3.5	5.7	9.7	11.8	20.8
2002.....	0.4	0.6	1.1	1.3	2.5	3.9	5.3	8.7	12.5	18.6
2001.....	0.5	0.8	0.9	1.3	2.5	3.1	5.6	6.8	11.9	17.4
2000.....	0.5	0.7	1.0	1.5	2.2	3.4	5.2	7.8	11.4	17.1
1999.....	0.4	0.6	1.0	1.3	2.0	3.5	4.6	7.5	10.8	19.0

White, female

2004.....	0.6	0.8	1.2	1.7	2.7	4.7	7.2	9.5	13.8	23.7
2003.....	0.8	0.9	1.2	2.1	3.0	5.1	6.8	10.1	14.9	23.9
2002.....	0.7	1.0	1.3	2.1	2.7	4.6	7.1	10.4	13.9	22.5
2001.....	0.6	0.9	1.3	1.7	2.7	4.4	5.9	9.2	12.8	20.7
2000.....	0.7	1.0	1.5	1.9	2.8	5.0	6.1	8.8	12.8	19.8
1999.....	0.6	0.8	1.2	1.6	2.2	3.8	5.4	7.8	9.8	17.0

Black, both sexes

2004.....	1.3	1.6	2.7	4.3	5.1	7.1	11.0	15.4	16.1	19.5
2003.....	1.9	2.0	2.3	3.7	6.3	8.2	13.1	14.0	15.2	21.9
2002.....	1.5	2.3	2.2	4.3	5.9	8.3	10.9	14.9	16.0	19.6
2001.....	1.5	1.9	2.4	4.5	5.4	9.5	9.3	12.5	14.9	21.8
2000.....	1.6	1.5	3.1	2.9	5.3	6.2	10.7	10.5	13.8	20.0
1999.....	1.7	1.8	2.6	3.8	5.3	7.3	9.5	10.4	17.0	18.8

Black, male

2004.....	*	*	2.5	3.4	3.9	4.8	7.1	10.0	17.2	*
2003.....	1.7	*	2.4	2.9	4.0	6.4	11.8	9.6	*	*
2002.....	*	*	*	3.3	*	7.2	9.0	12.9	15.3	*
2001.....	*	*	*	3.4	4.3	6.9	10.4	9.3	*	22.7
2000.....	*	*	*	*	*	*	9.4	*	*	*
1999.....	*	*	*	*	5.5	5.2	9.2	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

Black, female

2004.....	1.7	1.8	2.9	5.1	6.0	8.8	13.6	18.6	15.5	19.7
2003.....	2.1	2.5	2.2	4.3	8.1	9.5	14.0	16.5	17.8	23.1
2002.....	1.8	3.0	3.3	5.1	8.1	9.1	12.1	16.2	16.3	21.8
2001.....	2.0	2.3	2.9	5.4	6.2	11.5	8.5	14.4	15.1	21.5
2000.....	2.1	2.1	4.5	3.8	6.4	7.5	11.5	11.7	14.9	19.7
1999.....	1.9	2.4	3.1	5.7	5.2	8.9	9.7	12.0	19.0	21.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other non-pulmonary forms of heart disease (I30-I51)

All races 3/, both sexes

2004.....	9.6	15.0	21.9	31.0	50.4	79.2	135.6	244.8	470.2	1,319.1
2003.....	10.3	15.1	21.4	33.7	51.6	85.6	142.7	253.3	488.6	1,369.2
2002.....	10.5	15.1	22.1	33.6	52.4	85.2	147.1	261.0	500.8	1,387.2
2001.....	9.8	15.0	22.0	33.8	53.8	88.6	147.4	263.0	502.3	1,418.1
2000.....	10.2	14.8	21.7	35.0	55.7	91.3	150.3	266.1	508.1	1,452.0
1999.....	10.4	15.0	22.0	36.2	58.5	93.3	157.1	269.7	517.0	1,453.4

All races 3/, male

2004.....	12.7	20.5	29.2	41.3	65.8	101.0	166.0	294.7	543.2	1,329.4
2003.....	13.8	20.1	28.5	44.9	67.1	108.0	178.0	307.6	564.9	1,376.9
2002.....	14.3	20.7	30.0	44.1	67.8	106.5	183.6	316.9	578.7	1,396.5
2001.....	13.0	20.5	29.3	44.3	69.5	109.7	183.3	321.6	586.6	1,417.8
2000.....	13.7	20.3	29.4	47.1	72.5	115.4	189.3	320.0	593.3	1,494.4
1999.....	13.8	20.5	29.7	48.8	75.7	121.4	198.5	325.4	606.3	1,488.5

All races 3/, female

2004.....	6.6	9.7	14.9	21.2	36.4	60.3	111.0	209.0	425.2	1,314.4
2003.....	6.9	10.3	14.6	23.2	37.5	66.3	114.4	214.7	442.0	1,365.8
2002.....	6.8	9.7	14.4	23.6	38.5	66.8	117.9	221.5	453.4	1,383.1
2001.....	6.7	9.6	15.1	24.0	39.6	70.4	118.8	222.0	451.7	1,418.2
2000.....	6.7	9.3	14.3	23.7	40.6	70.7	119.5	228.6	457.9	1,434.8
1999.....	7.0	9.7	14.7	24.4	42.9	69.2	124.8	231.0	464.8	1,439.2

White, both sexes

2004.....	8.1	12.5	18.5	27.0	46.1	74.4	131.0	240.7	475.2	1,353.6
2003.....	8.7	12.8	17.8	29.9	46.4	80.0	137.1	250.9	492.4	1,403.6
2002.....	9.0	12.4	18.6	29.7	47.6	80.4	141.8	257.7	504.0	1,419.3
2001.....	7.9	12.3	18.7	29.5	49.5	83.0	142.3	258.4	504.2	1,452.9
2000.....	8.5	12.1	18.4	30.9	51.0	85.6	144.5	261.2	511.5	1,485.2
1999.....	8.5	12.1	18.7	31.4	53.2	87.6	151.0	263.8	518.4	1,484.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other non-pulmonary forms of heart disease (I30-I51)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
White, male										
2004.....	10.8	17.1	25.4	36.4	60.3	94.8	161.5	289.1	549.6	1,359.9
2003.....	12.0	17.3	23.9	39.9	61.0	101.1	172.1	305.7	569.5	1,413.6
2002.....	12.6	17.4	25.7	39.3	62.0	100.7	177.1	313.3	584.4	1,430.2
2001.....	11.0	17.3	25.6	38.9	64.5	103.7	176.6	316.0	587.2	1,449.9
2000.....	11.6	17.2	25.0	42.2	67.0	108.4	183.0	314.8	597.8	1,527.6
1999.....	11.7	17.1	25.5	42.6	69.2	114.4	191.6	319.3	607.6	1,511.1
White, female										
2004.....	5.3	7.9	11.8	18.0	32.9	56.4	106.0	205.3	429.0	1,350.7
2003.....	5.4	8.2	11.8	20.3	32.8	61.4	108.5	211.3	444.9	1,399.2
2002.....	5.4	7.5	11.7	20.5	34.3	62.5	112.9	217.8	454.8	1,414.6
2001.....	4.8	7.4	12.0	20.4	35.8	64.8	114.5	217.6	454.1	1,454.1
2000.....	5.3	7.0	11.9	20.2	36.3	65.6	113.5	223.4	460.2	1,468.0
1999.....	5.4	7.3	12.2	20.8	38.4	64.2	118.7	224.8	465.7	1,473.2
Black, both sexes										
2004.....	22.1	35.7	52.6	72.4	101.3	139.0	211.5	338.2	504.5	1,118.7
2003.....	23.7	35.2	53.0	74.3	108.6	152.6	228.9	327.9	524.9	1,165.2
2002.....	22.7	37.2	53.5	74.3	106.2	146.8	225.5	343.1	545.3	1,201.8
2001.....	24.7	36.7	53.5	79.0	103.5	155.9	225.9	363.2	556.6	1,200.0
2000.....	23.6	37.5	52.7	77.3	106.4	160.6	237.6	367.7	543.8	1,229.4
1999.....	25.3	38.6	53.6	84.2	116.1	162.6	248.4	373.7	563.4	1,230.8
Black, male										
2004.....	28.4	49.5	67.9	97.6	137.3	184.9	259.6	427.7	591.1	1,160.3
2003.....	30.0	45.2	71.4	102.0	143.8	200.4	283.5	397.1	617.9	1,144.8
2002.....	29.1	50.5	72.3	98.8	139.0	187.0	292.4	416.6	624.4	1,172.9
2001.....	29.9	48.0	68.5	105.3	135.3	191.8	290.2	456.7	688.6	1,253.8
2000.....	30.9	49.4	73.7	102.9	137.7	207.6	292.5	441.8	645.7	1,310.1
1999.....	32.3	51.1	74.0	116.6	153.8	213.9	314.8	445.9	684.5	1,382.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other non-pulmonary forms of heart disease (I30-I51)

Black, female

2004.....	16.4	23.8	39.7	51.7	73.1	104.9	178.8	285.2	458.9	1,102.2
2003.....	18.1	26.6	37.4	51.6	81.1	117.1	192.1	286.8	476.2	1,173.1
2002.....	17.0	25.6	37.7	54.1	80.5	116.9	180.6	299.1	504.1	1,212.9
2001.....	20.1	27.0	40.7	57.5	78.6	129.3	183.2	307.3	489.4	1,180.0
2000.....	17.2	27.1	34.9	56.3	81.7	126.0	201.1	323.0	492.5	1,199.3
1999.....	19.2	27.9	36.2	57.7	86.5	124.5	204.3	330.1	503.4	1,174.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute and subacute endocarditis (I33)

All races 3/, both sexes

2004.....	0.2	0.4	0.5	0.5	0.7	1.0	1.5	2.1	2.9	3.2
2003.....	0.3	0.4	0.5	0.6	0.9	1.0	1.5	2.3	2.7	3.0
2002.....	0.3	0.4	0.5	0.5	0.8	1.1	1.5	2.2	2.7	2.6
2001.....	0.3	0.4	0.5	0.5	0.8	1.1	1.7	2.2	3.0	2.5
2000.....	0.3	0.4	0.6	0.7	0.7	1.1	1.3	2.2	2.8	2.8
1999.....	0.3	0.3	0.4	0.7	0.7	1.1	1.5	2.1	2.7	2.9

All races 3/, male

2004.....	0.3	0.5	0.7	0.6	0.8	1.4	1.4	2.3	4.2	4.8
2003.....	0.4	0.6	0.7	0.8	1.1	1.3	1.6	2.9	3.7	4.1
2002.....	0.4	0.5	0.7	0.6	0.9	1.5	1.7	2.5	3.7	3.5
2001.....	0.3	0.7	0.7	0.6	1.1	1.1	2.1	3.1	4.3	3.8
2000.....	0.4	0.6	0.7	0.8	0.9	1.3	1.6	3.1	4.0	4.2
1999.....	0.4	0.5	0.7	1.0	0.8	1.2	1.7	2.8	3.9	4.7

All races 3/, female

2004.....	*	0.3	0.4	0.3	0.6	0.7	1.7	2.0	2.1	2.5
2003.....	0.2	0.3	0.3	0.4	0.8	0.9	1.4	1.9	2.1	2.5
2002.....	0.2	0.3	0.3	0.4	0.8	0.8	1.3	1.9	2.1	2.2
2001.....	0.3	0.2	0.3	0.5	0.5	1.0	1.3	1.6	2.2	2.0
2000.....	0.2	0.3	0.4	0.5	0.6	0.9	1.1	1.6	2.0	2.3
1999.....	*	*	0.2	0.5	0.7	1.0	1.4	1.7	2.0	2.1

White, both sexes

2004.....	0.2	0.3	0.4	0.4	0.6	1.0	1.5	2.0	2.9	3.2
2003.....	0.3	0.3	0.4	0.5	0.8	1.0	1.4	2.3	2.8	3.1
2002.....	0.2	0.3	0.4	0.4	0.8	1.0	1.4	2.2	2.9	2.7
2001.....	0.2	0.4	0.4	0.4	0.7	0.9	1.5	2.0	3.0	2.6
2000.....	0.3	0.3	0.4	0.5	0.7	1.1	1.3	2.2	2.7	2.7
1999.....	0.2	0.3	0.3	0.5	0.6	1.0	1.4	2.2	2.8	3.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races 3/, both sexes

2004.....	0.2	0.2	0.3	0.3	0.5	0.6	0.6	1.1	1.3	1.8
2003.....	0.2	0.3	0.3	0.4	0.4	0.8	0.8	0.9	1.5	1.6
2002.....	0.2	0.3	0.3	0.4	0.4	0.6	1.0	1.1	1.4	2.1
2001.....	0.2	0.3	0.3	0.4	0.5	0.5	0.8	1.2	1.3	1.9
2000.....	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.1	1.3	1.8
1999.....	0.3	0.3	0.3	0.5	0.6	0.7	0.9	1.2	1.5	2.0

All races 3/, male

2004.....	0.2	0.2	0.4	0.4	0.6	0.8	0.7	1.2	1.4	2.3
2003.....	0.3	0.4	0.3	0.4	0.4	1.0	1.0	1.1	1.5	1.5
2002.....	0.3	0.4	0.3	0.5	0.6	0.7	1.1	1.1	1.3	2.4
2001.....	*	0.3	0.3	0.4	0.6	0.6	0.8	1.3	1.1	1.8
2000.....	0.3	0.3	*	0.5	0.4	0.5	1.1	1.4	1.7	2.1
1999.....	0.3	0.3	0.3	0.6	0.6	0.8	0.9	1.3	1.5	2.5

All races 3/, female

2004.....	*	0.2	0.2	0.3	0.4	0.5	0.6	1.1	1.3	1.6
2003.....	0.2	0.2	*	0.3	0.3	0.7	0.7	0.8	1.5	1.7
2002.....	*	*	0.2	0.4	*	0.5	1.0	1.1	1.4	2.0
2001.....	0.3	0.3	0.3	0.4	*	*	0.9	1.1	1.4	2.0
2000.....	*	*	0.2	*	0.4	0.5	0.7	0.9	1.0	1.7
1999.....	0.3	0.3	0.2	0.3	0.6	0.7	0.8	1.2	1.5	1.9

White, both sexes

2004.....	0.1	0.2	0.3	0.3	0.4	0.6	0.6	1.0	1.3	1.8
2003.....	0.2	0.3	0.2	0.4	0.4	0.8	0.8	1.0	1.5	1.7
2002.....	0.2	0.2	0.2	0.4	0.4	0.6	1.0	1.0	1.3	2.2
2001.....	0.2	0.2	0.2	0.3	0.5	0.4	0.9	1.2	1.3	1.9
2000.....	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.0	1.4	1.8
1999.....	0.2	0.3	0.2	0.5	0.6	0.7	0.8	1.2	1.5	2.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of endocardium (I34-I38)

All races 3/, both sexes

2004.....	0.8	1.5	2.0	2.8	4.8	8.3	16.4	31.5	67.0	200.3
2003.....	0.9	1.2	1.9	3.1	4.5	9.1	15.9	32.4	68.7	201.9
2002.....	1.0	1.3	1.9	2.9	4.8	8.4	15.9	33.0	72.0	203.9
2001.....	0.8	1.2	2.1	2.8	4.7	9.0	17.3	33.9	70.3	204.2
2000.....	0.9	1.4	1.9	3.0	5.1	8.9	17.8	34.1	73.3	213.2
1999.....	1.1	1.3	2.0	2.9	5.1	9.2	17.6	35.2	76.6	209.8

All races 3/, male

2004.....	1.0	2.0	2.4	3.6	5.2	9.1	17.6	34.1	70.3	194.2
2003.....	1.1	1.4	2.1	3.9	4.9	9.6	17.4	35.9	73.9	195.3
2002.....	1.1	1.7	2.3	3.3	5.2	8.7	17.3	36.4	74.9	198.5
2001.....	0.9	1.6	2.7	3.3	5.3	9.7	18.9	38.6	75.4	200.8
2000.....	1.0	1.9	2.2	3.8	5.9	9.3	19.2	35.1	78.9	215.8
1999.....	1.2	1.5	2.4	3.5	5.8	10.4	19.6	38.5	86.3	206.4

All races 3/, female

2004.....	0.6	1.0	1.5	2.0	4.4	7.6	15.4	29.6	64.9	203.0
2003.....	0.7	1.0	1.7	2.4	4.1	8.6	14.7	29.8	65.6	204.8
2002.....	0.8	1.0	1.5	2.5	4.5	8.2	14.8	30.6	70.2	206.3
2001.....	0.6	0.9	1.5	2.4	4.2	8.4	16.1	30.6	67.2	205.6
2000.....	0.7	1.0	1.5	2.4	4.4	8.6	16.8	33.5	69.9	212.1
1999.....	0.9	1.2	1.7	2.4	4.4	8.2	16.0	32.9	70.9	211.2

White, both sexes

2004.....	0.7	1.3	1.7	2.6	4.7	8.2	17.1	32.8	70.3	212.5
2003.....	0.7	1.1	1.6	2.9	4.2	9.1	16.2	33.5	72.3	214.4
2002.....	0.9	1.1	1.7	2.7	4.8	8.2	16.4	34.7	75.8	216.2
2001.....	0.7	1.1	1.9	2.6	4.6	8.8	17.8	35.2	74.1	216.5
2000.....	0.8	1.3	1.7	2.7	5.0	8.9	18.1	35.1	77.0	225.8
1999.....	1.0	1.1	1.9	2.7	4.9	9.1	17.4	36.4	80.2	221.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of endocardium (I34-I38)

White, male

2004.....	0.9	1.7	2.1	3.4	5.1	9.2	18.5	35.2	73.7	205.2
2003.....	0.9	1.2	1.8	3.5	4.7	9.7	17.9	37.3	77.6	207.8
2002.....	1.1	1.4	2.1	3.1	5.1	8.7	17.8	38.2	78.9	211.7
2001.....	0.8	1.4	2.5	3.1	5.2	9.5	19.4	39.9	79.8	212.0
2000.....	1.0	1.7	2.0	3.5	5.9	9.4	19.5	36.2	82.7	229.0
1999.....	1.1	1.1	2.3	3.3	5.6	10.5	19.4	39.9	90.2	218.2

White, female

2004.....	0.6	0.8	1.3	1.9	4.3	7.2	15.9	31.0	68.1	215.8
2003.....	0.5	0.9	1.5	2.3	3.7	8.5	14.8	30.7	69.0	217.2
2002.....	0.6	0.7	1.3	2.3	4.4	7.8	15.2	32.1	73.9	218.2
2001.....	0.5	0.7	1.4	2.1	4.2	8.2	16.5	31.8	70.6	218.4
2000.....	0.6	0.8	1.4	2.0	4.2	8.4	17.0	34.3	73.6	224.5
1999.....	0.8	1.0	1.4	2.2	4.2	7.9	15.8	33.9	74.4	222.7

Black, both sexes

2004.....	1.6	3.1	4.7	4.6	6.6	11.6	13.9	22.7	40.6	88.3
2003.....	2.1	2.7	4.3	5.5	7.6	10.2	16.1	26.0	37.6	85.2
2002.....	1.8	3.6	3.8	4.8	6.7	12.1	13.8	22.2	37.2	86.4
2001.....	1.5	2.9	3.9	5.9	6.4	12.2	15.1	25.7	38.2	85.4
2000.....	1.9	3.0	3.6	5.9	7.2	10.9	18.9	29.9	39.5	92.9
1999.....	2.2	3.2	3.7	5.3	7.8	11.4	21.8	25.9	41.9	91.5

Black, male

2004.....	2.0	4.3	5.9	5.7	7.7	9.9	13.9	25.0	40.2	84.4
2003.....	2.6	3.3	5.5	8.4	7.5	10.6	15.1	26.9	39.9	70.9
2002.....	1.5	4.5	4.0	6.1	7.1	10.7	14.0	24.3	35.2	67.5
2001.....	1.9	3.8	5.1	6.7	7.9	13.0	15.8	29.7	32.9	86.2
2000.....	1.8	4.1	4.5	6.6	7.5	10.4	19.7	29.8	43.0	86.5
1999.....	2.3	3.8	4.1	5.4	9.3	11.2	25.1	27.1	47.1	77.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of endocardium (I34-I38)										
Black, female										
2004.....	1.3	2.1	3.6	3.8	5.8	12.8	13.8	21.3	40.8	89.8
2003.....	1.7	2.1	3.4	3.1	7.7	9.9	16.8	25.5	36.4	90.7
2002.....	2.0	2.8	3.6	3.8	6.3	13.1	13.7	20.9	38.2	93.6
2001.....	*	2.1	2.9	5.1	5.1	11.6	14.6	23.3	40.9	85.0
2000.....	1.9	2.0	2.8	5.3	7.0	11.3	18.4	29.9	37.8	95.3
1999.....	2.2	2.7	3.4	5.1	6.7	11.5	19.6	25.2	39.3	96.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nonrheumatic mitral valve disorders (I34)

All races 3/, both sexes

2004.....	0.2	0.3	0.3	0.4	0.7	1.3	2.5	4.7	8.4	20.1
2003.....	0.2	0.2	0.3	0.5	0.8	1.4	2.4	4.4	9.0	20.1
2002.....	0.2	0.3	0.3	0.5	1.0	1.5	2.8	5.0	9.8	22.1
2001.....	0.2	0.3	0.4	0.5	0.7	1.7	2.7	5.7	9.6	22.7
2000.....	0.3	0.3	0.3	0.5	1.0	1.6	3.0	5.8	10.6	24.3
1999.....	0.3	0.3	0.4	0.5	0.9	1.6	3.1	5.8	10.5	25.1

All races 3/, male

2004.....	0.2	0.3	0.3	0.4	0.6	1.1	2.2	4.4	7.8	16.6
2003.....	0.2	0.2	0.3	0.5	0.8	1.3	2.4	4.5	8.2	16.4
2002.....	0.2	0.2	0.3	0.4	0.9	1.3	2.4	4.2	8.2	21.0
2001.....	0.2	0.3	0.4	0.5	0.7	1.7	2.8	4.7	8.1	19.0
2000.....	0.2	0.4	0.3	0.5	0.8	1.6	2.6	5.6	9.8	22.9
1999.....	0.3	0.3	0.3	0.4	0.9	1.3	3.1	5.6	10.6	22.2

All races 3/, female

2004.....	0.3	0.3	0.3	0.4	0.8	1.4	2.7	5.0	8.7	21.7
2003.....	0.2	0.3	0.4	0.5	0.9	1.5	2.3	4.4	9.5	21.7
2002.....	0.2	0.3	0.3	0.5	1.1	1.8	3.1	5.6	10.9	22.6
2001.....	0.3	0.3	0.4	0.5	0.8	1.7	2.7	6.3	10.5	24.3
2000.....	0.3	0.2	0.4	0.6	1.1	1.5	3.2	5.9	11.1	24.9
1999.....	0.4	0.4	0.5	0.6	0.9	1.8	3.1	5.9	10.5	26.2

White, both sexes

2004.....	0.2	0.3	0.3	0.4	0.6	1.3	2.5	4.9	8.7	21.1
2003.....	0.2	0.2	0.3	0.5	0.8	1.4	2.5	4.6	9.3	21.4
2002.....	0.2	0.2	0.3	0.4	1.0	1.5	2.9	5.2	10.4	23.8
2001.....	0.2	0.3	0.4	0.5	0.7	1.6	2.8	6.0	10.1	24.2
2000.....	0.3	0.3	0.3	0.5	1.0	1.6	3.0	6.0	11.3	25.5
1999.....	0.3	0.3	0.3	0.5	0.9	1.5	3.1	5.9	11.1	26.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nonrheumatic aortic valve disorders (I35)

All races 3/, both sexes

2004.....	0.3	0.5	0.8	1.2	2.2	3.8	8.9	18.4	42.7	142.5
2003.....	0.3	0.4	0.6	1.3	1.9	4.5	8.5	18.4	44.1	144.9
2002.....	0.3	0.5	0.8	1.2	2.2	4.0	8.8	19.2	45.4	144.8
2001.....	0.2	0.5	0.9	1.2	2.3	4.5	9.2	19.9	44.6	144.1
2000.....	0.3	0.5	0.8	1.4	2.3	4.5	9.9	20.0	47.1	151.3
1999.....	0.3	0.5	0.8	1.3	2.3	4.8	9.3	20.7	49.9	147.8

All races 3/, male

2004.....	0.4	0.7	1.0	1.7	2.7	4.6	10.3	21.0	46.0	141.9
2003.....	0.4	0.6	0.8	1.8	2.4	5.5	9.9	21.9	50.4	144.7
2002.....	0.4	0.7	1.0	1.5	2.6	4.5	10.8	23.2	51.4	144.9
2001.....	0.3	0.8	1.3	1.5	2.9	5.3	11.0	24.7	50.7	145.9
2000.....	0.4	0.8	1.2	2.1	3.1	5.2	11.7	21.5	52.9	158.2
1999.....	0.4	0.7	1.2	1.7	3.0	5.9	11.7	24.3	58.9	150.3

All races 3/, female

2004.....	*	0.2	0.5	0.8	1.8	3.0	7.8	16.5	40.6	142.7
2003.....	*	0.2	0.5	0.8	1.4	3.6	7.4	15.9	40.2	145.0
2002.....	0.2	0.2	0.6	0.8	1.9	3.5	7.1	16.4	41.7	144.8
2001.....	*	0.3	0.5	0.9	1.8	3.8	7.8	16.6	40.9	143.4
2000.....	*	0.2	0.4	0.8	1.6	3.8	8.5	19.0	43.7	148.5
1999.....	0.2	0.4	0.5	0.9	1.7	3.8	7.4	18.1	44.7	146.8

White, both sexes

2004.....	0.3	0.5	0.7	1.3	2.3	3.9	9.5	19.1	45.0	151.6
2003.....	0.2	0.4	0.6	1.3	1.9	4.6	8.8	19.1	46.6	154.1
2002.....	0.3	0.4	0.7	1.2	2.3	3.9	9.1	20.3	47.9	153.2
2001.....	0.2	0.5	0.9	1.2	2.4	4.6	9.6	20.5	47.3	152.8
2000.....	0.3	0.5	0.8	1.4	2.3	4.4	10.2	20.9	49.5	160.7
1999.....	0.3	0.4	0.8	1.3	2.3	4.8	9.2	21.7	52.3	156.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nonrheumatic aortic valve disorders (I35)

White, male

2004.....	0.4	0.7	0.9	1.7	2.7	4.8	10.9	21.7	48.5	150.1
2003.....	0.3	0.5	0.7	1.7	2.5	5.8	10.3	22.7	53.4	154.4
2002.....	0.4	0.6	1.0	1.5	2.7	4.6	11.3	24.5	54.3	154.4
2001.....	0.3	0.7	1.2	1.5	3.0	5.4	11.4	25.2	53.8	153.5
2000.....	0.4	0.7	1.2	2.0	3.2	5.2	12.0	22.6	55.4	168.4
1999.....	0.4	0.6	1.1	1.7	3.0	6.1	11.5	25.6	61.2	159.0

White, female

2004.....	*	0.2	0.5	0.9	1.9	3.0	8.3	17.2	42.8	152.2
2003.....	*	0.2	0.5	0.9	1.4	3.7	7.6	16.5	42.3	154.0
2002.....	*	*	0.5	0.9	1.9	3.4	7.4	17.3	44.0	152.7
2001.....	*	0.3	0.5	0.8	2.0	3.9	8.2	17.2	43.3	152.5
2000.....	*	*	0.5	0.7	1.5	3.8	8.7	19.7	46.1	157.6
1999.....	0.2	0.3	0.4	0.8	1.8	3.7	7.4	19.0	47.0	155.0

Black, both sexes

2004.....	*	*	1.3	1.4	2.3	4.0	5.8	13.2	23.3	59.2
2003.....	*	*	0.9	1.7	1.8	3.3	7.8	14.1	21.1	58.6
2002.....	*	0.9	1.1	*	2.2	5.1	6.8	11.5	23.3	63.0
2001.....	*	*	1.3	1.7	2.0	4.4	6.9	16.3	21.4	59.6
2000.....	*	0.8	*	2.1	2.6	5.5	9.1	13.9	25.4	62.3
1999.....	*	1.0	1.4	1.7	2.8	5.0	10.7	12.0	26.9	64.4

Black, male

2004.....	*	*	*	*	4.1	*	6.5	14.1	23.7	58.6
2003.....	*	*	*	2.8	*	*	7.9	16.4	19.9	48.0
2002.....	*	*	*	*	*	5.2	7.7	13.3	22.2	52.5
2001.....	*	*	2.3	*	*	*	8.1	22.8	19.3	68.1
2000.....	*	*	*	*	*	6.2	10.7	13.2	30.3	57.7
1999.....	*	*	*	*	*	5.7	13.9	13.3	36.6	58.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nonrheumatic aortic valve disorders (I35)

Black, female

2004.....	*	*	*	*	*	4.1	5.3	12.7	23.1	59.5
2003.....	*	*	*	*	*	*	7.8	12.8	21.7	62.8
2002.....	*	*	*	*	*	5.0	6.3	10.4	23.9	66.9
2001.....	*	*	*	*	*	4.1	6.1	12.5	22.5	56.4
2000.....	*	*	*	*	*	5.0	8.0	14.2	22.9	64.0
1999.....	*	*	*	*	*	4.4	8.6	11.2	22.0	66.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases of endocardium (I36-I38)

All races 3/, both sexes

2004.....	0.3	0.7	0.9	1.1	1.9	3.3	5.0	8.4	15.9	37.7
2003.....	0.4	0.6	0.9	1.3	1.8	3.2	5.0	9.5	15.7	36.9
2002.....	0.4	0.6	0.8	1.2	1.6	2.9	4.4	8.7	16.7	37.0
2001.....	0.3	0.4	0.8	1.1	1.6	2.8	5.3	8.3	16.1	37.4
2000.....	0.3	0.6	0.8	1.1	1.8	2.9	5.0	8.3	15.6	37.6
1999.....	0.4	0.5	0.8	1.1	1.8	2.9	5.2	8.7	16.1	37.0

All races 3/, male

2004.....	0.4	1.0	1.1	1.4	2.0	3.4	5.1	8.7	16.5	35.7
2003.....	0.5	0.7	1.0	1.6	1.7	2.9	5.0	9.5	15.4	34.2
2002.....	0.5	0.8	1.0	1.3	1.7	2.9	4.1	9.1	15.3	32.6
2001.....	0.4	0.5	1.0	1.3	1.7	2.7	5.1	9.2	16.5	35.9
2000.....	0.4	0.7	0.8	1.2	2.0	2.5	4.9	8.0	16.2	34.7
1999.....	0.5	0.5	0.9	1.3	1.9	3.1	4.9	8.6	16.8	33.9

All races 3/, female

2004.....	0.2	0.5	0.7	0.8	1.8	3.2	5.0	8.1	15.6	38.5
2003.....	0.3	0.5	0.8	1.0	1.8	3.5	5.0	9.5	15.9	38.1
2002.....	0.4	0.5	0.6	1.1	1.5	3.0	4.6	8.5	17.6	38.9
2001.....	0.2	0.3	0.6	1.0	1.6	2.8	5.5	7.7	15.8	38.0
2000.....	0.2	0.5	0.7	0.9	1.7	3.2	5.0	8.6	15.2	38.7
1999.....	0.3	0.5	0.7	0.9	1.7	2.7	5.5	8.8	15.7	38.2

White, both sexes

2004.....	0.2	0.5	0.7	0.9	1.8	3.0	5.1	8.8	16.5	39.9
2003.....	0.3	0.4	0.7	1.1	1.5	3.0	4.9	9.8	16.4	38.9
2002.....	0.3	0.4	0.7	1.0	1.4	2.8	4.3	9.2	17.5	39.2
2001.....	0.3	0.3	0.7	0.9	1.5	2.6	5.4	8.6	16.7	39.5
2000.....	0.2	0.5	0.6	0.8	1.7	2.9	4.9	8.2	16.2	39.6
1999.....	0.3	0.3	0.7	0.9	1.6	2.8	5.1	8.8	16.8	38.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases of endocardium (I36-I38)

White, male

2004.....	0.3	0.7	0.9	1.3	1.9	3.3	5.3	9.0	17.1	37.6
2003.....	0.4	0.5	0.8	1.3	1.4	2.7	5.0	9.9	15.7	35.8
2002.....	0.4	0.5	0.9	1.1	1.5	2.9	4.1	9.4	15.9	34.8
2001.....	0.3	0.3	0.9	1.0	1.5	2.5	5.2	9.7	17.3	38.0
2000.....	0.4	0.6	0.6	1.0	1.8	2.5	4.8	7.8	16.9	36.4
1999.....	0.4	0.3	0.8	1.1	1.7	3.0	4.8	8.6	17.6	35.5

White, female

2004.....	*	0.4	0.5	0.6	1.6	2.8	4.9	8.6	16.1	40.9
2003.....	*	0.4	0.7	0.9	1.6	3.3	4.7	9.6	16.8	40.2
2002.....	0.2	0.3	0.4	1.0	1.3	2.7	4.6	9.0	18.5	41.1
2001.....	*	*	0.5	0.8	1.4	2.7	5.5	7.9	16.3	40.2
2000.....	*	0.4	0.6	0.7	1.6	3.2	4.9	8.4	15.7	40.9
1999.....	0.2	0.3	0.6	0.8	1.6	2.5	5.3	8.9	16.4	40.2

Black, both sexes

2004.....	0.9	2.1	2.9	2.6	2.9	5.9	5.9	5.8	12.1	19.2
2003.....	1.2	1.7	2.7	3.0	4.3	5.3	6.9	8.5	11.2	18.7
2002.....	1.3	2.2	2.1	2.8	3.6	4.9	4.8	6.9	9.7	16.6
2001.....	*	1.6	1.9	3.3	3.5	5.3	5.4	6.9	11.6	18.1
2000.....	1.0	1.5	2.1	2.9	3.6	3.8	7.2	11.0	10.4	18.5
1999.....	1.3	1.6	1.7	2.8	3.7	4.5	7.7	9.5	10.7	16.9

Black, male

2004.....	*	3.1	3.8	2.9	*	4.8	*	*	*	*
2003.....	*	2.0	3.6	4.6	4.2	5.4	*	*	15.5	*
2002.....	*	3.0	*	3.7	*	*	*	*	*	*
2001.....	*	1.9	2.6	4.4	4.3	5.8	*	*	*	*
2000.....	*	1.9	2.4	*	*	*	7.0	12.8	*	*
1999.....	*	*	*	*	4.9	*	*	10.0	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases of endocardium (I36-I38)

Black, female

2004.....	*	*	2.2	2.3	3.4	6.7	6.2	*	12.1	19.3
2003.....	*	1.4	2.0	*	4.4	5.3	7.8	9.2	8.9	19.0
2002.....	1.4	1.6	2.2	*	3.6	5.4	5.4	6.3	9.5	19.3
2001.....	*	*	*	*	*	4.9	5.6	7.8	11.9	18.9
2000.....	*	*	*	2.7	3.3	4.0	7.3	10.0	11.0	17.6
1999.....	*	*	*	*	*	4.6	8.6	9.2	11.2	18.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cardiomyopathy (I42)

All races 3/, both sexes

2004.....	3.5	5.3	6.9	9.0	13.0	18.3	28.0	43.8	68.9	127.6
2003.....	4.0	5.6	7.0	9.7	13.9	20.8	30.7	45.7	72.5	137.8
2002.....	4.1	5.9	7.2	10.5	13.8	21.1	33.6	49.5	77.3	142.0
2001.....	4.0	5.5	7.3	10.6	14.5	22.9	31.9	48.9	76.8	138.3
2000.....	4.1	5.6	8.0	10.8	15.8	24.1	33.7	51.5	76.2	142.3
1999.....	4.2	5.7	7.9	11.7	16.8	25.6	35.5	50.0	76.1	139.2

All races 3/, male

2004.....	5.0	7.7	9.7	12.5	18.5	25.4	39.2	62.4	96.3	174.6
2003.....	5.7	8.0	10.0	13.5	19.6	28.6	42.9	65.5	104.5	178.8
2002.....	5.9	8.6	10.4	14.7	19.4	28.9	47.9	71.8	110.8	189.7
2001.....	5.8	7.9	10.0	14.8	20.5	31.0	44.7	69.9	108.8	184.7
2000.....	5.9	7.9	11.5	15.5	21.6	33.1	47.9	74.4	109.5	195.3
1999.....	6.1	8.2	11.3	16.8	23.0	36.1	50.7	70.9	106.8	188.2

All races 3/, female

2004.....	2.0	2.9	4.3	5.7	8.0	12.2	19.0	30.5	52.1	106.4
2003.....	2.3	3.3	4.2	6.1	8.7	14.1	20.9	31.6	52.8	119.7
2002.....	2.4	3.2	4.1	6.5	8.7	14.3	22.1	33.7	56.9	121.3
2001.....	2.3	3.2	4.7	6.7	9.0	15.9	21.8	34.3	57.6	118.9
2000.....	2.2	3.4	4.7	6.4	10.5	16.4	22.5	35.6	56.5	120.7
1999.....	2.3	3.4	4.7	6.9	11.2	16.6	23.6	35.5	58.2	119.4

White, both sexes

2004.....	2.8	4.1	5.6	7.7	11.3	16.3	26.2	42.2	68.7	129.3
2003.....	3.3	4.6	5.6	8.1	11.9	18.8	28.8	44.3	71.3	139.8
2002.....	3.3	4.7	5.8	8.9	12.2	19.5	31.7	47.7	76.6	141.7
2001.....	3.0	4.2	5.9	8.8	12.7	21.0	30.3	46.9	75.3	139.2
2000.....	3.2	4.2	6.4	9.0	14.0	21.6	31.7	49.1	74.7	142.4
1999.....	3.1	4.4	6.3	9.6	14.2	23.4	33.1	48.0	74.3	138.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cardiomyopathy (I42)

White, male

2004.....	4.0	6.2	8.1	10.6	16.1	22.6	37.3	60.8	96.8	178.3
2003.....	4.8	6.7	8.0	11.4	16.9	25.8	40.6	64.2	103.5	182.6
2002.....	4.8	7.0	8.5	12.6	17.4	26.7	45.3	70.3	110.8	190.9
2001.....	4.5	6.2	8.3	12.4	18.2	28.5	42.7	67.7	107.1	187.0
2000.....	4.6	6.3	9.2	13.0	19.4	29.5	45.4	71.2	108.2	197.5
1999.....	4.7	6.4	9.1	13.8	19.4	33.2	47.2	68.7	105.1	188.0

White, female

2004.....	1.5	2.1	3.2	4.9	6.8	10.7	17.0	28.6	51.2	107.2
2003.....	1.7	2.5	3.2	4.9	7.2	12.7	19.2	30.0	51.4	120.8
2002.....	1.8	2.4	3.2	5.3	7.4	13.1	20.7	31.6	55.6	120.4
2001.....	1.5	2.2	3.5	5.4	7.6	14.3	20.2	32.2	56.1	119.2
2000.....	1.7	2.3	3.6	5.2	9.0	14.6	20.6	33.5	54.8	120.0
1999.....	1.5	2.4	3.6	5.6	9.4	14.9	21.9	33.4	56.1	118.8

Black, both sexes

2004.....	9.1	13.8	18.2	23.1	31.7	40.0	51.2	69.1	85.4	124.4
2003.....	9.8	14.1	19.0	25.7	34.1	42.5	55.0	68.6	99.5	135.7
2002.....	10.4	15.2	18.4	27.0	30.4	39.5	57.5	77.3	96.3	161.0
2001.....	11.8	15.5	19.8	27.8	33.3	44.7	54.5	80.9	104.5	141.2
2000.....	10.7	16.6	22.7	28.6	32.2	51.2	59.6	85.5	104.8	151.1
1999.....	12.1	16.6	22.7	30.9	40.5	49.4	64.3	79.1	103.7	153.0

Black, male

2004.....	13.1	20.6	24.3	33.8	46.5	58.1	69.3	96.0	112.6	158.0
2003.....	13.5	19.8	27.0	36.3	49.0	62.8	77.5	94.5	141.0	172.0
2002.....	14.8	22.4	27.4	38.6	42.0	55.8	85.5	103.8	130.7	204.8
2001.....	16.3	21.4	26.6	40.0	47.6	62.5	73.4	112.9	153.3	180.4
2000.....	16.2	22.0	33.1	42.8	44.5	73.6	82.2	123.3	149.2	193.7
1999.....	16.8	23.5	33.6	47.1	58.1	70.6	96.3	105.0	140.4	215.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cardiomyopathy (I42)

Black, female

2004.....	5.6	7.8	13.1	14.2	20.1	26.4	38.9	53.1	71.1	111.0
2003.....	6.5	9.2	12.1	17.0	22.6	27.4	39.9	53.2	77.8	121.5
2002.....	6.5	9.1	10.7	17.5	21.4	27.4	38.8	61.5	78.4	144.2
2001.....	7.8	10.3	14.0	17.8	22.1	31.5	41.9	61.8	79.7	126.7
2000.....	5.8	12.0	13.9	17.0	22.6	34.6	44.6	62.7	82.4	135.2
1999.....	7.9	10.7	13.4	17.7	26.6	33.7	43.0	63.5	85.6	129.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Conduction disorders and cardiac dysrhythmias (I44-I49)

All races 3/, both sexes

2004.....	2.4	3.9	6.2	9.1	15.6	22.4	36.3	62.1	114.4	295.3
2003.....	2.4	4.0	6.1	10.2	15.4	24.3	39.4	66.7	122.1	313.7
2002.....	2.6	3.6	6.4	10.1	15.8	24.7	39.0	70.1	126.0	323.2
2001.....	2.2	3.9	6.3	10.0	15.9	25.3	40.5	69.3	125.7	328.2
2000.....	2.5	3.8	5.9	10.5	16.2	26.1	41.0	68.8	127.3	340.2
1999.....	2.5	4.1	6.5	10.8	19.1	27.3	44.8	72.4	134.3	353.7

All races 3/, male

2004.....	3.1	5.0	8.1	12.4	21.2	28.9	45.1	71.8	126.1	278.5
2003.....	3.1	5.1	8.4	13.8	20.4	31.5	50.2	77.3	134.0	292.7
2002.....	3.4	4.8	8.7	13.5	21.2	31.1	48.1	81.3	138.2	310.5
2001.....	2.9	5.1	8.5	13.1	20.9	31.0	49.7	82.2	140.8	311.6
2000.....	3.0	5.1	8.2	14.3	21.7	34.5	52.3	80.6	137.4	333.2
1999.....	3.1	5.7	8.7	14.7	24.9	36.8	55.1	84.9	154.2	353.4

All races 3/, female

2004.....	1.8	2.8	4.3	6.1	10.5	16.7	29.3	55.2	107.3	302.9
2003.....	1.8	3.0	4.0	6.9	10.9	18.0	30.6	59.1	114.7	323.0
2002.....	1.8	2.4	4.3	6.9	10.9	19.1	31.8	62.2	118.6	328.8
2001.....	1.5	2.7	4.3	7.0	11.4	20.4	33.3	60.3	116.7	335.2
2000.....	1.9	2.5	3.7	7.0	11.2	18.8	32.2	60.6	121.4	343.0
1999.....	1.8	2.5	4.4	7.2	13.8	19.2	36.7	63.7	122.7	353.8

White, both sexes

2004.....	2.1	3.4	5.6	8.0	14.6	21.1	34.5	59.8	113.5	295.0
2003.....	2.2	3.6	5.4	9.4	14.3	22.2	37.4	65.0	120.9	314.5
2002.....	2.4	3.2	5.7	9.1	14.1	23.4	37.1	68.0	124.4	324.4
2001.....	1.9	3.5	5.6	8.9	14.7	23.7	38.6	66.9	123.8	329.7
2000.....	2.2	3.3	5.4	9.8	14.7	24.7	39.4	66.6	127.1	341.6
1999.....	2.3	3.6	5.9	9.8	17.9	26.0	43.0	70.0	133.1	355.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Conduction disorders and cardiac dysrhythmias (I44-I49)

White, male

2004.....	2.8	4.5	7.6	11.1	20.1	27.2	42.8	68.9	124.4	273.7
2003.....	2.9	4.7	7.5	12.7	19.0	29.1	47.7	74.8	132.6	291.3
2002.....	3.4	4.4	7.8	12.4	19.3	29.9	45.7	78.2	136.9	308.9
2001.....	2.7	4.7	7.7	11.6	19.4	29.7	46.9	78.8	137.5	310.2
2000.....	2.8	4.6	7.4	13.3	20.0	32.7	50.4	78.4	138.4	332.8
1999.....	3.0	5.3	8.0	13.3	23.6	35.1	53.3	81.3	152.3	350.9

White, female

2004.....	1.4	2.4	3.6	5.0	9.5	15.7	27.8	53.1	106.7	304.7
2003.....	1.5	2.5	3.3	6.1	9.9	16.1	28.9	58.0	113.7	324.7
2002.....	1.5	2.0	3.6	6.1	9.4	17.7	30.0	60.6	116.8	331.1
2001.....	1.2	2.2	3.6	6.2	10.4	18.4	31.8	58.5	115.5	337.9
2000.....	1.6	2.0	3.3	6.4	9.9	17.6	30.5	58.4	120.4	345.2
1999.....	1.6	2.0	3.9	6.5	12.6	18.1	34.9	62.1	121.8	357.9

Black, both sexes

2004.....	5.2	7.8	12.4	20.6	28.4	37.5	61.8	96.0	140.6	315.6
2003.....	4.8	8.2	13.3	19.6	28.8	46.5	66.2	95.0	145.2	327.3
2002.....	4.4	7.2	14.0	20.3	33.5	41.6	63.8	99.1	160.7	337.6
2001.....	4.0	8.0	13.6	21.1	29.8	44.7	65.7	105.2	162.5	340.2
2000.....	4.8	8.3	11.8	19.4	32.3	43.4	64.4	105.6	147.1	346.3
1999.....	4.2	8.9	12.8	21.7	33.6	44.6	70.6	109.4	170.2	359.0

Black, male

2004.....	5.4	9.9	15.5	26.8	38.9	49.9	81.3	117.8	166.4	332.8
2003.....	5.4	9.8	17.6	26.1	38.2	60.3	88.4	121.0	160.9	307.6
2002.....	5.1	9.3	18.4	26.5	43.9	51.4	82.5	120.3	173.5	336.6
2001.....	4.4	9.4	17.6	28.5	40.5	50.6	86.2	135.7	198.2	357.4
2000.....	5.1	10.8	17.1	26.2	42.9	58.8	79.6	121.4	152.6	361.0
1999.....	4.7	11.1	16.5	31.0	43.5	61.0	85.8	135.3	215.4	411.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Conduction disorders and cardiac dysrhythmias (I44-I49)

Black, female

2004.....	5.0	5.9	9.9	15.5	20.2	28.2	48.5	83.0	127.0	308.7
2003.....	4.2	6.9	9.7	14.2	21.5	36.2	51.2	79.5	136.9	335.0
2002.....	3.8	5.4	10.2	15.3	25.4	34.4	51.3	86.4	154.0	338.0
2001.....	3.7	6.7	10.2	15.0	21.4	40.4	52.0	87.1	144.3	333.8
2000.....	4.6	6.2	7.3	13.8	24.0	32.1	54.4	96.0	144.4	340.8
1999.....	3.7	6.9	9.6	14.1	25.8	32.5	60.5	93.8	147.8	339.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Heart failure (I50)

All races 3/, both sexes

2004.....	0.9	1.7	3.3	6.1	11.4	22.4	43.8	88.7	187.7	621.2
2003.....	1.0	1.6	3.3	6.4	12.1	23.6	44.7	89.9	192.7	641.1
2002.....	0.8	1.7	3.4	6.0	12.4	22.9	45.7	89.0	194.0	645.0
2001.....	0.8	1.7	3.3	6.5	13.0	23.8	45.6	92.6	199.2	674.6
2000.....	0.8	1.4	3.0	6.1	12.8	24.0	46.5	93.5	199.9	684.9
1999.....	0.8	1.5	2.9	6.5	11.7	23.5	48.3	94.3	201.6	681.9

All races 3/, male

2004.....	1.0	2.2	4.0	7.5	13.7	26.4	51.1	103.2	211.1	605.5
2003.....	1.3	2.1	3.9	7.9	14.7	28.3	53.1	105.3	213.9	629.8
2002.....	1.0	2.0	4.4	7.2	14.8	27.0	53.2	103.9	218.0	621.4
2001.....	0.9	2.2	4.0	8.1	15.0	28.0	54.2	107.9	226.6	646.4
2000.....	1.1	1.9	3.5	7.4	15.3	28.2	55.6	107.9	229.5	672.2
1999.....	1.0	1.9	3.6	8.0	14.2	28.0	58.9	108.4	224.9	667.0

All races 3/, female

2004.....	0.7	1.3	2.8	4.8	9.4	18.8	37.9	78.3	173.3	628.2
2003.....	0.7	1.2	2.7	4.9	9.7	19.6	37.8	78.9	179.8	646.1
2002.....	0.5	1.3	2.4	4.9	10.1	19.3	39.8	78.4	179.3	655.3
2001.....	0.7	1.2	2.6	5.1	11.3	20.2	38.7	81.9	182.7	686.4
2000.....	0.6	0.9	2.5	5.0	10.5	20.3	39.3	83.5	182.4	690.0
1999.....	0.6	1.1	2.1	5.1	9.5	19.6	39.9	84.5	188.0	687.9

White, both sexes

2004.....	0.7	1.4	2.5	5.1	10.3	21.5	42.9	87.9	191.0	641.5
2003.....	0.7	1.3	2.6	5.6	10.7	22.4	43.2	89.5	195.3	660.4
2002.....	0.6	1.2	2.7	5.3	11.4	21.7	44.1	88.1	195.6	664.5
2001.....	0.6	1.4	2.7	5.7	12.1	22.5	44.3	92.1	201.1	695.1
2000.....	0.6	1.1	2.5	5.4	11.8	22.6	44.9	92.8	201.8	703.7
1999.....	0.6	1.1	2.3	5.6	10.9	22.0	46.8	92.5	202.5	699.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
White, male										
2004.....	0.8	1.6	3.1	6.2	12.2	25.2	50.7	101.7	215.6	625.5
2003.....	0.9	1.7	3.0	7.0	13.1	27.0	52.1	105.9	217.1	650.8
2002.....	0.8	1.5	3.6	6.4	13.6	25.5	51.4	103.4	221.4	641.9
2001.....	0.7	1.9	3.4	7.2	14.2	26.5	52.8	107.6	228.2	666.5
2000.....	0.9	1.5	3.0	6.8	14.1	27.2	54.3	107.7	231.7	690.0
1999.....	0.8	1.4	2.9	7.1	13.3	26.2	58.1	107.2	226.3	680.0
White, female										
2004.....	0.6	1.1	2.0	4.0	8.6	18.1	36.5	77.8	175.7	648.7
2003.....	0.4	0.9	2.1	4.2	8.5	18.3	35.9	77.6	181.9	664.7
2002.....	0.4	0.9	1.9	4.2	9.3	18.4	38.2	77.1	179.8	674.3
2001.....	0.5	0.8	2.0	4.2	10.1	18.9	37.4	81.0	184.7	707.1
2000.....	0.4	0.6	2.0	4.1	9.6	18.6	37.3	82.4	183.9	709.3
1999.....	0.4	0.8	1.7	4.2	8.8	18.4	37.8	82.1	188.5	707.0
Black, both sexes										
2004.....	2.2	5.0	10.5	16.3	23.8	35.7	64.0	120.2	194.6	506.5
2003.....	3.3	4.7	9.4	15.3	27.7	41.1	70.9	114.7	203.6	531.8
2002.....	2.2	5.5	9.3	13.9	24.6	38.6	71.7	118.5	215.3	528.0
2001.....	2.7	4.5	8.7	15.7	24.6	41.2	71.1	121.1	213.7	543.8
2000.....	2.2	4.2	7.6	14.0	24.3	41.2	72.6	119.8	209.5	558.4
1999.....	2.4	4.8	8.4	16.4	22.0	42.0	74.0	130.5	216.7	552.9
Black, male										
2004.....	2.6	6.8	12.2	21.0	30.6	45.5	69.3	148.3	220.2	503.7
2003.....	3.9	5.5	11.6	19.4	35.9	51.0	79.5	124.6	227.4	513.0
2002.....	2.7	6.6	12.0	16.5	31.3	47.6	87.5	136.9	236.9	479.2
2001.....	2.6	5.0	10.4	18.9	26.4	49.6	86.2	140.8	256.8	541.2
2000.....	2.5	5.5	9.1	14.8	29.6	45.5	82.9	134.7	242.8	581.2
1999.....	2.7	6.4	11.0	19.3	27.4	51.4	83.1	146.7	238.9	603.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Heart failure (I50)

Black, female

2004.....	1.9	3.4	8.9	12.4	18.5	28.4	60.4	103.5	181.1	507.6
2003.....	2.8	4.0	7.6	11.9	21.4	33.8	65.1	108.8	191.1	539.2
2002.....	1.7	4.5	7.0	11.7	19.4	31.8	61.2	107.6	204.1	546.7
2001.....	2.8	4.1	7.2	13.1	23.2	35.1	61.0	109.3	191.7	544.8
2000.....	1.9	3.0	6.4	13.4	20.1	38.0	65.7	110.8	192.8	549.9
1999.....	2.1	3.5	6.3	14.0	17.8	35.0	68.0	120.7	205.7	534.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Congestive heart failure (I50.0)

All races 3/, both sexes

2004.....	0.7	1.5	2.8	5.1	9.9	19.8	40.0	82.1	175.8	588.9
2003.....	0.8	1.4	2.8	5.5	10.6	21.2	40.6	82.9	181.6	608.2
2002.....	0.6	1.3	2.8	5.1	10.7	20.3	41.4	82.3	181.6	609.0
2001.....	0.7	1.4	2.8	5.4	11.3	21.5	41.5	85.8	185.4	636.1
2000.....	0.7	1.1	2.5	5.4	11.1	20.9	41.8	86.1	186.9	644.4
1999.....	0.7	1.2	2.3	5.3	10.0	20.9	43.6	86.6	188.4	642.7

All races 3/, male

2004.....	0.8	1.8	3.3	6.2	12.0	23.0	46.5	95.6	196.2	571.8
2003.....	1.0	1.7	3.2	6.7	12.6	25.5	48.1	96.9	200.7	594.6
2002.....	0.8	1.6	3.5	6.1	12.5	23.8	48.2	95.6	203.2	584.6
2001.....	0.8	1.9	3.3	6.6	12.8	24.9	49.1	98.7	208.5	609.3
2000.....	0.8	1.6	2.9	6.4	13.0	24.7	49.6	99.1	214.1	632.2
1999.....	0.8	1.5	2.9	6.5	11.9	24.5	53.2	99.2	209.4	628.2

All races 3/, female

2004.....	0.6	1.2	2.4	4.1	8.0	16.9	34.8	72.4	163.3	596.6
2003.....	0.6	1.0	2.3	4.3	8.7	17.6	34.6	72.9	169.9	614.2
2002.....	0.5	1.0	2.1	4.2	9.0	17.2	36.0	72.8	168.4	619.6
2001.....	0.6	1.0	2.3	4.2	10.0	18.5	35.5	76.8	171.5	647.3
2000.....	0.5	0.7	2.1	4.4	9.4	17.7	35.7	77.0	170.9	649.4
1999.....	0.5	0.9	1.8	4.2	8.4	17.9	36.1	77.9	176.1	648.5

White, both sexes

2004.....	0.6	1.1	2.2	4.3	9.0	19.1	39.3	81.5	179.5	609.2
2003.....	0.5	1.0	2.2	4.8	9.3	20.3	39.5	82.7	184.5	628.4
2002.....	0.5	0.9	2.2	4.5	9.8	19.3	40.1	81.7	183.6	628.8
2001.....	0.5	1.1	2.3	4.7	10.5	20.2	40.5	85.7	187.6	656.5
2000.....	0.5	0.8	2.1	4.8	10.2	19.7	40.6	85.6	189.3	663.6
1999.....	0.5	0.8	1.8	4.5	9.4	19.6	42.3	85.3	189.4	660.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Congestive heart failure (I50.0)

White, male

2004.....	0.6	1.4	2.6	5.2	10.6	22.1	46.3	94.3	201.1	591.7
2003.....	0.7	1.4	2.6	5.9	11.2	24.5	47.2	97.8	204.3	616.2
2002.....	0.6	1.2	2.8	5.4	11.4	22.5	46.5	95.3	207.2	605.8
2001.....	0.6	1.6	2.8	6.0	12.1	23.5	47.9	99.1	210.8	629.0
2000.....	0.7	1.2	2.4	6.1	11.8	23.9	48.6	99.2	216.9	650.5
1999.....	0.7	1.1	2.3	5.7	11.0	22.9	52.5	98.4	210.7	641.7

White, female

2004.....	0.5	0.9	1.7	3.4	7.5	16.4	33.6	72.2	166.1	617.1
2003.....	0.3	0.7	1.9	3.7	7.6	16.6	33.1	71.8	172.3	633.9
2002.....	0.3	0.7	1.6	3.6	8.2	16.5	35.0	71.9	169.1	638.8
2001.....	0.4	0.7	1.9	3.5	9.1	17.2	34.4	76.1	173.6	667.9
2000.....	0.4	0.5	1.7	3.5	8.7	16.0	34.1	76.1	172.8	668.8
1999.....	0.4	0.6	1.3	3.4	7.8	16.8	34.2	76.2	176.9	667.4

Black, both sexes

2004.....	1.7	4.2	8.5	13.8	20.1	30.7	57.3	109.5	174.8	468.7
2003.....	2.9	4.1	7.6	12.9	24.3	35.8	63.5	102.9	185.8	484.3
2002.....	1.9	4.6	7.7	11.9	21.9	34.4	63.8	106.7	196.0	482.6
2001.....	2.2	4.0	7.3	13.3	21.0	37.6	63.1	108.5	194.2	502.4
2000.....	1.8	3.6	6.7	11.9	21.3	36.5	63.0	108.4	190.3	510.2
1999.....	1.8	4.2	7.3	13.6	18.7	37.2	66.6	116.4	200.2	509.5

Black, male

2004.....	1.9	5.5	9.6	17.9	26.7	38.5	61.2	135.6	194.4	466.9
2003.....	3.3	4.8	9.1	16.4	30.7	44.6	71.6	110.5	203.8	466.1
2002.....	2.2	5.6	9.6	14.3	27.3	42.9	81.5	122.6	209.4	428.8
2001.....	2.3	4.5	8.8	15.2	22.6	44.2	75.4	123.1	224.7	500.4
2000.....	2.0	4.7	8.0	11.7	26.1	38.2	71.2	120.0	216.6	528.2
1999.....	2.1	5.6	9.2	15.4	23.1	44.9	74.6	131.5	224.1	554.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Congestive heart failure (I50.0)

Black, female

2004.....	1.5	3.1	7.5	10.4	15.0	24.8	54.7	94.1	164.4	469.4
2003.....	2.5	3.5	6.3	9.9	19.4	29.2	58.0	98.5	176.4	491.4
2002.....	1.6	3.7	6.1	9.9	17.7	28.1	52.0	97.2	189.0	503.2
2001.....	2.2	3.6	6.0	11.7	19.7	32.7	54.9	99.8	178.7	503.1
2000.....	1.5	2.5	5.5	12.0	17.5	35.1	57.5	101.4	177.0	503.5
1999.....	1.6	3.0	5.7	12.0	15.3	31.6	61.2	107.3	188.4	492.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified heart failure (I50.1-I50.9)

All races 3/, both sexes

2004.....	0.2	0.3	0.5	1.0	1.6	2.6	3.8	6.6	11.9	32.3
2003.....	0.2	0.3	0.5	0.9	1.5	2.4	4.0	7.0	11.2	32.9
2002.....	0.1	0.4	0.6	0.9	1.7	2.6	4.3	6.7	12.4	36.1
2001.....	0.2	0.3	0.5	1.1	1.7	2.4	4.1	6.8	13.8	38.5
2000.....	0.2	0.3	0.5	0.8	1.7	3.0	4.7	7.4	12.9	40.5
1999.....	0.2	0.3	0.5	1.2	1.7	2.6	4.6	7.7	13.3	39.2

All races 3/, male

2004.....	0.2	0.4	0.7	1.2	1.7	3.4	4.7	7.6	15.0	33.7
2003.....	0.3	0.4	0.7	1.2	2.1	2.8	5.0	8.4	13.2	35.2
2002.....	0.2	0.4	1.0	1.1	2.3	3.2	5.0	8.3	14.8	36.8
2001.....	0.2	0.3	0.7	1.5	2.1	3.1	5.2	9.2	18.1	37.2
2000.....	0.3	0.4	0.7	0.9	2.3	3.6	6.1	8.8	15.4	40.0
1999.....	0.2	0.4	0.7	1.6	2.4	3.5	5.7	9.2	15.5	38.8

All races 3/, female

2004.....	*	0.2	0.4	0.7	1.4	1.9	3.1	5.8	10.0	31.6
2003.....	*	0.2	0.3	0.7	1.0	2.0	3.2	6.0	9.9	31.9
2002.....	*	0.3	0.3	0.7	1.1	2.1	3.8	5.6	10.9	35.7
2001.....	*	*	0.3	0.8	1.3	1.7	3.2	5.2	11.2	39.1
2000.....	*	*	0.4	0.6	1.1	2.5	3.6	6.5	11.5	40.6
1999.....	*	0.3	0.4	0.9	1.1	1.7	3.8	6.6	11.9	39.4

White, both sexes

2004.....	0.1	0.2	0.3	0.8	1.4	2.4	3.6	6.4	11.5	32.3
2003.....	0.2	0.3	0.3	0.8	1.3	2.1	3.7	6.7	10.9	32.0
2002.....	0.1	0.3	0.5	0.8	1.6	2.4	4.0	6.4	12.0	35.6
2001.....	0.1	0.2	0.4	1.0	1.5	2.3	3.8	6.4	13.5	38.7
2000.....	0.1	0.2	0.5	0.7	1.6	2.9	4.3	7.2	12.5	40.2
1999.....	0.1	0.3	0.5	1.1	1.6	2.4	4.5	7.2	13.1	39.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified heart failure (I50.1-I50.9)

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
White, male										
2004.....	*	0.3	0.5	1.1	1.6	3.1	4.4	7.4	14.5	33.8
2003.....	0.3	0.3	0.4	1.0	1.8	2.5	4.9	8.1	12.8	34.6
2002.....	*	0.3	0.8	1.0	2.2	3.0	4.9	8.0	14.2	36.0
2001.....	*	0.3	0.6	1.3	2.0	3.0	4.9	8.5	17.4	37.5
2000.....	0.2	0.3	0.6	0.7	2.2	3.2	5.7	8.4	14.8	39.5
1999.....	*	0.4	0.6	1.4	2.2	3.3	5.6	8.8	15.6	38.3
White, female										
2004.....	*	*	*	0.6	1.1	1.7	3.0	5.6	9.7	31.6
2003.....	*	*	*	0.5	0.9	1.7	2.8	5.7	9.7	30.8
2002.....	*	0.2	*	0.6	1.1	1.9	3.3	5.2	10.6	35.5
2001.....	*	*	*	0.8	1.1	1.7	3.0	4.9	11.1	39.2
2000.....	*	*	0.3	0.6	1.0	2.6	3.2	6.3	11.1	40.5
1999.....	*	*	0.3	0.8	1.0	1.6	3.6	6.0	11.6	39.5
Black, both sexes										
2004.....	*	0.8	2.0	2.4	3.7	5.1	6.7	10.7	19.8	37.8
2003.....	*	*	1.9	2.4	3.4	5.3	7.4	11.7	17.8	47.6
2002.....	*	0.9	1.6	2.0	2.7	4.1	7.9	11.9	19.4	45.4
2001.....	*	*	1.4	2.5	3.7	3.7	7.9	12.5	19.5	41.5
2000.....	*	*	*	2.2	3.0	4.8	9.6	11.4	19.2	48.2
1999.....	*	*	*	2.8	3.3	4.8	7.4	14.1	16.5	43.4
Black, male										
2004.....	*	*	2.6	3.0	3.9	7.0	8.1	12.7	25.8	36.8
2003.....	*	*	2.5	2.9	5.2	6.4	7.9	14.2	23.6	46.9
2002.....	*	*	2.5	*	4.0	*	*	14.2	27.5	50.4
2001.....	*	*	*	3.8	*	5.3	10.8	17.7	32.1	40.8
2000.....	*	*	*	*	*	7.3	11.7	14.6	26.1	53.0
1999.....	*	*	*	3.9	4.2	6.5	8.5	15.2	*	49.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified heart failure (I50.1-I50.9)

Black, female

2004.....	*	*	*	*	3.5	3.6	5.7	9.4	16.6	38.2
2003.....	*	*	*	*	*	4.6	7.1	10.3	14.7	47.8
2002.....	*	*	*	*	*	3.7	9.2	10.4	15.1	43.5
2001.....	*	*	*	*	3.5	*	6.1	9.4	13.1	41.7
2000.....	*	*	*	*	*	*	8.2	9.4	15.7	46.4
1999.....	*	*	*	*	*	*	6.8	13.4	17.3	41.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)

All races 3/, both sexes

2004.....	0.3	0.3	0.2	0.2	*	0.3	0.3	0.3	0.4	2.0
2003.....	0.2	0.2	0.2	0.2	0.2	*	*	0.3	0.5	2.2
2002.....	0.3	0.3	0.3	0.4	0.2	0.4	0.6	0.8	1.1	3.2
2001.....	0.3	0.3	0.3	0.2	0.3	0.4	0.4	0.5	0.6	2.0
2000.....	0.3	0.3	0.1	0.3	0.3	*	0.4	0.4	0.7	2.4
1999.....	0.3	0.3	0.3	0.2	0.2	0.3	0.3	0.4	0.7	2.4

All races 3/, male

2004.....	0.3	0.3	0.3	*	*	*	*	*	*	1.7
2003.....	*	0.2	0.2	0.3	*	*	*	*	*	2.6
2002.....	0.4	0.3	0.3	0.3	*	*	0.8	0.8	1.0	3.8
2001.....	0.3	0.3	0.3	*	*	0.5	0.5	0.8	*	2.2
2000.....	0.3	0.2	*	0.4	*	*	*	*	*	2.6
1999.....	0.3	0.3	0.3	*	*	*	*	*	*	2.1

All races 3/, female

2004.....	0.2	0.3	0.2	*	*	*	*	*	*	2.2
2003.....	0.3	0.3	0.2	*	*	*	*	*	*	2.0
2002.....	0.3	0.3	0.3	0.4	*	*	0.5	0.8	1.1	3.0
2001.....	0.3	0.3	*	*	*	*	*	*	*	1.9
2000.....	0.2	0.3	*	0.3	*	*	*	*	*	2.4
1999.....	0.3	0.3	0.2	*	*	*	*	*	*	2.5

White, both sexes

2004.....	0.2	0.2	0.2	0.2	*	0.3	*	0.4	*	2.1
2003.....	0.2	0.2	0.2	0.2	0.2	*	*	0.3	0.6	2.2
2002.....	0.3	0.3	0.2	0.4	0.2	0.3	0.6	0.8	1.1	3.2
2001.....	0.2	0.2	0.2	0.2	0.3	0.3	0.4	0.5	0.6	2.1
2000.....	0.2	0.2	*	0.4	0.2	*	0.4	0.4	0.7	2.6
1999.....	0.3	0.2	0.2	0.2	*	0.3	0.3	0.3	0.8	2.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

All races 3/, both sexes

2004.....	1.3	1.7	2.4	3.0	4.4	5.9	8.6	15.1	27.4	67.7
2003.....	1.3	1.7	2.1	3.1	4.2	5.8	9.6	15.2	27.9	67.8
2002.....	1.3	1.7	2.1	2.8	4.2	6.1	9.7	15.3	26.4	65.0
2001.....	1.2	1.7	2.0	2.8	4.1	5.7	9.0	14.4	25.4	66.3
2000.....	1.1	1.6	2.0	3.1	4.5	6.4	8.7	14.3	26.8	64.6
1999.....	1.0	1.4	1.8	2.8	4.3	5.6	8.3	14.1	23.4	61.5

All races 3/, male

2004.....	1.8	2.4	3.7	4.1	5.7	8.7	10.7	19.3	33.2	67.8
2003.....	1.9	2.3	2.9	4.3	5.8	7.5	11.5	19.3	32.5	72.1
2002.....	1.8	2.4	3.0	4.0	5.5	8.2	13.5	18.9	30.7	66.7
2001.....	1.7	2.4	2.8	3.9	5.8	7.8	12.3	17.8	28.8	66.4
2000.....	1.7	2.3	2.8	4.4	6.5	8.1	11.3	17.0	31.2	69.0
1999.....	1.5	2.1	2.5	4.0	6.2	7.8	11.2	18.3	27.7	64.3

All races 3/, female

2004.....	0.9	1.0	1.2	1.8	3.1	3.5	6.9	12.1	23.9	67.7
2003.....	0.7	1.1	1.3	2.0	2.7	4.3	8.1	12.3	25.0	66.0
2002.....	0.8	1.1	1.3	1.6	3.0	4.3	6.6	12.8	23.8	64.3
2001.....	0.7	0.9	1.2	1.9	2.6	3.9	6.5	12.0	23.3	66.3
2000.....	0.6	0.8	1.2	1.9	2.7	4.9	6.6	12.5	24.1	62.7
1999.....	0.6	0.8	1.1	1.7	2.7	3.7	6.0	11.2	20.8	60.4

White, both sexes

2004.....	1.2	1.5	2.3	2.7	4.1	5.6	8.0	14.6	27.1	68.1
2003.....	1.2	1.5	1.8	2.9	3.9	5.5	9.1	15.1	27.8	67.5
2002.....	1.1	1.5	1.8	2.6	3.7	5.7	9.4	15.2	26.3	64.5
2001.....	1.0	1.4	1.7	2.5	4.0	5.4	8.6	13.7	25.1	65.7
2000.....	1.0	1.5	1.7	2.8	4.3	6.1	7.9	13.9	26.1	64.7
1999.....	0.9	1.2	1.6	2.4	3.9	5.1	8.1	13.3	23.2	61.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

White, male

2004.....	1.6	2.3	3.4	3.9	5.4	8.3	9.9	18.7	33.0	68.2
2003.....	1.8	2.1	2.6	4.0	5.5	7.2	11.1	19.2	32.4	72.2
2002.....	1.6	2.1	2.6	3.7	4.9	7.6	13.3	18.7	30.2	66.9
2001.....	1.6	2.1	2.6	3.5	5.7	7.5	11.6	17.0	28.4	66.2
2000.....	1.5	2.2	2.4	3.9	6.3	7.7	10.3	16.4	30.0	69.2
1999.....	1.2	2.0	2.2	3.5	5.7	7.2	10.8	17.5	27.2	64.5

White, female

2004.....	0.8	0.8	1.1	1.6	2.8	3.2	6.4	11.7	23.5	68.1
2003.....	0.6	0.9	1.1	1.8	2.5	4.0	7.5	12.2	25.0	65.4
2002.....	0.7	0.8	1.0	1.5	2.7	4.0	6.3	12.8	23.9	63.4
2001.....	0.5	0.7	0.9	1.6	2.5	3.5	6.2	11.3	23.1	65.5
2000.....	0.5	0.7	0.9	1.7	2.4	4.6	6.0	12.1	23.8	62.8
1999.....	0.5	0.5	0.9	1.4	2.2	3.2	5.9	10.3	20.8	60.2

Black, both sexes

2004.....	2.6	3.6	4.0	5.8	8.2	10.8	17.0	24.5	37.9	76.4
2003.....	2.4	3.2	4.7	5.7	7.5	9.8	16.7	19.4	34.0	80.5
2002.....	2.3	3.6	5.4	5.1	8.7	11.2	14.5	19.9	32.2	80.5
2001.....	2.6	3.7	4.5	6.1	6.4	9.4	15.2	23.8	32.2	83.8
2000.....	2.4	2.9	4.8	6.5	8.0	11.0	18.7	22.2	37.8	73.8
1999.....	2.6	3.1	3.8	6.7	9.1	11.3	13.3	24.8	26.9	68.9

Black, male

2004.....	3.5	4.4	6.2	8.0	11.0	15.5	22.7	35.0	45.9	75.5
2003.....	3.4	3.8	6.5	7.7	11.0	13.0	18.7	25.6	44.3	80.3
2002.....	3.2	4.6	7.2	7.3	12.1	16.0	18.7	26.6	42.8	77.2
2001.....	3.0	5.1	5.3	8.4	8.3	10.9	21.9	28.8	39.3	84.0
2000.....	3.3	4.1	7.1	9.7	10.6	14.1	25.4	26.9	52.3	80.7
1999.....	3.6	3.8	5.3	9.7	11.7	16.1	19.0	29.9	36.6	72.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

Black, female

2004.....	1.8	2.9	2.2	3.9	6.0	7.4	13.2	18.3	33.6	76.8
2003.....	1.5	2.7	3.2	4.0	4.8	7.3	15.3	15.7	28.6	80.6
2002.....	1.5	2.6	4.0	3.3	6.0	7.6	11.6	15.9	26.7	81.7
2001.....	2.3	2.5	3.9	4.2	4.8	8.3	10.8	20.8	28.6	83.8
2000.....	1.7	1.8	2.8	3.9	6.0	8.8	14.2	19.4	30.6	71.3
1999.....	1.6	2.6	2.6	4.3	7.2	7.7	9.5	21.7	22.0	67.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cerebrovascular diseases (I60-I69)

All races 3/, both sexes

2004.....	7.1	11.9	18.3	26.5	44.5	78.4	142.2	276.7	532.1	1,245.9
2003.....	7.4	11.8	18.6	27.8	45.9	79.7	150.7	294.1	571.1	1,370.1
2002.....	7.1	12.3	18.3	28.6	48.3	83.8	160.7	308.8	601.7	1,445.9
2001.....	7.2	12.1	18.5	28.8	49.6	85.7	164.3	320.2	622.4	1,500.2
2000.....	7.7	13.1	19.3	30.8	53.8	91.0	169.1	332.8	654.0	1,589.2
1999.....	7.5	12.5	18.4	31.7	51.4	91.1	173.5	340.6	664.0	1,614.8

All races 3/, male

2004.....	7.4	12.8	21.1	29.9	52.2	90.2	158.8	301.2	551.6	1,118.1
2003.....	7.6	12.9	21.1	31.3	53.5	91.4	170.8	322.5	595.6	1,236.0
2002.....	6.9	13.4	20.4	32.7	55.8	97.0	178.7	334.8	615.6	1,317.9
2001.....	7.4	13.2	20.5	32.8	57.2	98.5	187.9	349.0	654.6	1,380.2
2000.....	7.8	14.2	21.3	35.2	62.3	102.6	192.9	367.6	695.1	1,484.3
1999.....	7.8	13.3	20.7	35.7	60.0	102.8	197.7	376.9	706.8	1,512.8

All races 3/, female

2004.....	6.8	11.0	15.5	23.2	37.5	68.1	128.8	259.1	520.2	1,303.4
2003.....	7.2	10.8	16.3	24.6	38.9	69.5	134.5	273.9	556.1	1,429.4
2002.....	7.2	11.2	16.3	24.8	41.4	72.4	146.2	290.4	593.3	1,501.5
2001.....	7.1	10.9	16.5	25.1	42.7	74.7	145.6	300.0	603.2	1,550.4
2000.....	7.5	12.0	17.3	26.6	46.0	81.0	150.3	308.5	629.7	1,632.0
1999.....	7.1	11.7	16.1	28.0	43.7	81.1	154.5	315.3	639.0	1,656.0

White, both sexes

2004.....	5.7	9.1	14.1	21.8	37.5	70.0	132.6	264.8	526.2	1,257.8
2003.....	5.6	9.0	14.2	22.6	38.3	70.8	140.3	283.2	565.9	1,382.2
2002.....	5.5	9.5	14.0	23.8	41.2	74.2	150.1	298.6	595.7	1,459.5
2001.....	5.6	9.3	13.8	23.9	41.7	75.2	153.2	309.7	617.1	1,511.2
2000.....	5.9	9.7	15.4	25.4	46.5	81.9	159.0	321.3	647.0	1,601.6
1999.....	5.7	9.4	14.4	26.2	43.5	82.9	161.3	329.3	658.4	1,631.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cerebrovascular diseases (I60-I69)

White, male

2004.....	6.1	9.7	16.3	24.1	43.3	80.6	147.0	288.0	546.9	1,129.3
2003.....	5.9	10.1	16.0	24.7	44.5	81.8	158.5	310.0	590.4	1,247.0
2002.....	5.5	10.6	15.5	26.9	47.2	85.1	167.3	323.4	610.1	1,327.4
2001.....	5.9	10.2	15.4	26.9	48.1	86.7	174.8	337.7	648.8	1,386.2
2000.....	6.1	10.8	16.9	28.6	53.8	91.7	180.5	354.6	686.1	1,490.7
1999.....	6.1	9.8	16.2	29.5	50.6	93.2	183.6	363.2	700.5	1,522.7

White, female

2004.....	5.2	8.5	11.9	19.5	32.2	60.5	120.7	247.9	513.3	1,315.7
2003.....	5.3	7.9	12.3	20.5	32.5	61.1	125.4	263.8	550.7	1,442.1
2002.....	5.5	8.4	12.6	20.9	35.6	64.5	136.0	280.8	586.8	1,516.9
2001.....	5.4	8.3	12.2	21.1	35.8	65.1	135.7	289.8	598.0	1,563.5
2000.....	5.7	8.7	13.9	22.4	39.8	73.3	141.9	297.9	623.8	1,646.7
1999.....	5.3	8.9	12.7	23.0	37.0	73.9	143.7	305.5	633.6	1,675.4

Black, both sexes

2004.....	17.2	31.5	48.0	66.3	108.1	154.9	246.5	418.6	647.9	1,211.2
2003.....	18.6	31.4	52.7	72.4	113.1	157.4	256.3	433.2	677.0	1,348.7
2002.....	17.4	32.1	51.2	71.1	109.9	169.6	272.5	447.2	712.5	1,383.9
2001.....	18.4	32.6	55.0	70.6	117.1	178.3	277.6	460.2	716.4	1,452.7
2000.....	19.7	37.2	51.2	77.4	114.0	174.8	281.1	475.2	769.4	1,530.0
1999.....	19.9	34.4	50.6	78.4	118.2	170.0	307.8	487.9	756.4	1,492.5

Black, male

2004.....	16.9	35.4	56.4	84.7	139.4	188.3	297.8	470.9	677.2	1,061.0
2003.....	18.8	34.2	62.6	90.6	141.7	184.9	307.4	503.0	727.9	1,180.3
2002.....	16.9	34.3	61.6	88.9	138.8	213.7	328.8	502.5	729.8	1,252.2
2001.....	17.8	36.5	63.9	87.6	143.4	218.5	336.3	524.6	768.1	1,349.1
2000.....	20.5	41.0	60.4	97.8	137.5	212.8	340.0	544.8	863.2	1,458.8
1999.....	20.8	38.8	61.3	92.7	149.4	204.3	366.7	559.0	818.8	1,448.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cerebrovascular diseases (I60-I69)

Black, female

2004.....	17.5	28.1	40.8	51.2	83.6	130.1	211.8	387.6	632.4	1,270.7
2003.....	18.5	29.0	44.4	57.3	90.7	137.1	221.8	391.8	650.3	1,414.2
2002.....	17.9	30.1	42.4	56.5	87.3	136.9	234.8	414.1	703.4	1,434.3
2001.....	19.0	29.1	47.4	56.7	96.5	148.5	238.6	421.9	690.1	1,491.2
2000.....	19.0	33.9	43.4	60.6	95.4	146.7	242.0	433.2	722.2	1,556.5
1999.....	19.0	30.5	41.6	66.6	93.7	144.5	268.7	445.1	725.6	1,509.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Subarachnoid hemorrhage (I60)

All races 3/, both sexes

2004.....	2.0	2.8	3.4	3.7	4.3	4.7	5.9	7.7	9.2	12.2
2003.....	2.1	2.6	3.6	3.8	4.3	5.0	5.8	7.7	10.1	11.5
2002.....	2.3	3.0	3.1	3.8	4.8	5.5	6.2	7.9	10.3	12.2
2001.....	2.2	3.0	3.4	4.2	4.3	5.3	6.3	8.0	9.8	13.2
2000.....	2.5	3.3	3.8	4.7	5.0	5.2	6.4	8.6	10.7	14.1
1999.....	2.6	3.3	3.7	4.4	5.1	5.7	6.7	8.4	10.9	13.1

All races 3/, male

2004.....	1.5	2.4	3.0	2.8	3.3	3.2	4.2	5.7	6.5	9.3
2003.....	1.6	2.1	2.9	2.8	3.4	4.0	4.0	5.4	7.4	9.1
2002.....	1.8	2.4	2.2	3.0	3.8	4.2	4.6	5.6	6.1	9.1
2001.....	1.9	2.4	2.8	3.2	3.1	3.5	5.2	5.7	7.5	9.5
2000.....	2.0	2.6	2.9	3.4	3.6	3.4	5.2	6.3	7.7	11.2
1999.....	2.2	2.7	2.9	3.3	3.6	4.2	4.4	6.1	7.7	9.0

All races 3/, female

2004.....	2.6	3.3	3.7	4.5	5.2	6.1	7.4	9.2	10.9	13.5
2003.....	2.5	3.2	4.3	4.7	5.1	5.8	7.2	9.3	11.8	12.5
2002.....	2.7	3.6	4.0	4.6	5.6	6.6	7.4	9.6	12.9	13.6
2001.....	2.6	3.6	3.9	5.2	5.3	6.8	7.2	9.7	11.2	14.8
2000.....	2.9	4.0	4.7	6.0	6.4	6.7	7.4	10.3	12.4	15.3
1999.....	3.0	3.9	4.4	5.5	6.4	6.9	8.5	10.0	12.8	14.8

White, both sexes

2004.....	1.8	2.6	3.0	3.5	4.0	4.6	5.8	7.4	9.1	12.4
2003.....	1.8	2.3	3.3	3.6	4.2	4.6	5.6	7.3	9.7	11.1
2002.....	2.0	2.8	2.8	3.5	4.6	5.3	5.7	7.7	10.1	12.0
2001.....	1.9	2.7	2.9	3.8	4.2	4.8	6.1	7.7	9.9	12.7
2000.....	2.1	2.9	3.5	4.6	4.8	4.8	6.0	8.4	10.4	13.6
1999.....	2.3	2.9	3.4	4.2	4.7	5.6	6.5	8.3	10.7	12.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Subarachnoid hemorrhage (I60)										
White, male										
2004.....	1.4	2.1	2.7	2.6	3.0	3.1	4.2	5.5	6.5	9.9
2003.....	1.4	1.9	2.7	2.6	3.2	3.7	4.2	5.1	7.2	8.6
2002.....	1.7	2.4	2.0	2.7	3.6	4.1	4.3	5.6	5.9	9.1
2001.....	1.7	2.1	2.5	2.8	3.1	3.3	5.0	5.5	7.9	9.2
2000.....	1.8	2.4	2.7	3.3	3.3	3.2	4.8	6.3	7.8	11.8
1999.....	2.1	2.5	2.7	3.2	3.4	4.0	4.4	6.3	7.6	9.0
White, female										
2004.....	2.2	3.0	3.4	4.4	4.9	5.9	7.1	8.7	10.8	13.5
2003.....	2.1	2.7	3.9	4.6	5.0	5.4	6.8	8.9	11.3	12.2
2002.....	2.4	3.1	3.6	4.3	5.5	6.4	6.8	9.3	12.7	13.3
2001.....	2.2	3.2	3.4	4.8	5.3	6.1	6.9	9.3	11.1	14.2
2000.....	2.5	3.4	4.3	5.9	6.2	6.1	7.0	10.0	12.0	14.3
1999.....	2.5	3.3	4.0	5.1	5.9	7.0	8.3	9.7	12.5	14.1
Black, both sexes										
2004.....	3.9	4.6	5.3	5.2	6.8	4.9	5.8	8.0	10.1	9.0
2003.....	3.8	4.4	5.7	5.3	4.8	6.6	5.4	8.7	11.7	14.6
2002.....	4.1	5.3	5.5	6.0	5.6	6.2	6.7	8.2	9.2	12.8
2001.....	4.4	5.1	6.5	7.2	4.8	8.2	6.3	9.7	6.8	13.8
2000.....	4.7	6.1	6.2	6.2	6.2	7.4	8.7	11.0	10.4	17.2
1999.....	5.0	6.0	6.3	6.7	6.7	5.5	8.7	7.5	12.4	14.7
Black, male										
2004.....	2.5	3.7	4.6	4.5	6.0	*	*	*	*	*
2003.....	2.8	2.8	3.9	5.0	5.8	5.7	*	*	*	*
2002.....	2.7	3.1	3.4	5.2	5.1	5.7	6.7	*	*	*
2001.....	3.4	3.9	5.3	5.8	4.1	*	*	*	*	*
2000.....	3.6	4.2	4.6	3.8	4.8	*	8.4	*	*	*
1999.....	3.5	4.5	5.3	5.1	4.9	5.7	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Subarachnoid hemorrhage (I60)

Black, female

2004.....	5.1	5.4	6.0	5.9	7.4	5.8	7.0	9.7	12.1	11.8
2003.....	4.8	5.8	7.2	5.5	4.1	7.3	7.8	10.3	12.8	15.0
2002.....	5.2	7.2	7.3	6.6	6.0	6.5	6.7	9.9	10.7	15.6
2001.....	5.3	6.1	7.6	8.3	5.4	10.9	7.2	10.8	8.6	16.4
2000.....	5.6	7.8	7.5	8.1	7.4	9.4	8.9	12.8	13.2	22.8
1999.....	6.4	7.3	7.2	8.0	8.2	5.4	10.8	10.3	13.0	18.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intracerebral and other intracranial hemorrhage (I61-I62)

All races 3/, both sexes

2004.....	2.9	5.1	7.5	9.8	13.9	21.2	34.1	57.5	89.6	124.3
2003.....	3.1	5.2	7.6	9.5	13.9	20.4	34.9	57.7	89.5	126.6
2002.....	2.8	5.0	7.2	9.7	14.3	20.9	36.0	59.1	90.6	127.2
2001.....	3.1	5.2	7.4	9.5	14.8	21.2	35.2	58.2	91.1	127.9
2000.....	3.2	5.6	7.6	10.6	15.9	23.5	35.5	60.1	95.1	137.9
1999.....	2.8	4.9	7.1	10.7	14.3	20.7	36.4	59.5	90.5	122.3

All races 3/, male

2004.....	3.5	6.0	9.1	11.7	16.8	24.9	37.7	63.2	98.2	134.4
2003.....	3.4	6.1	9.4	11.0	16.4	23.6	39.9	63.2	99.8	137.4
2002.....	3.2	6.0	8.8	11.0	17.3	24.7	41.0	63.4	98.4	135.0
2001.....	3.4	6.3	8.8	11.2	17.3	23.7	41.5	65.5	99.5	134.2
2000.....	3.6	6.8	8.9	13.4	19.0	27.0	40.0	65.9	100.6	149.6
1999.....	3.4	5.6	8.6	12.7	17.3	23.5	41.7	65.1	94.2	130.7

All races 3/, female

2004.....	2.3	4.2	6.1	8.0	11.3	18.0	31.2	53.4	84.3	119.8
2003.....	2.7	4.4	5.8	8.0	11.7	17.7	30.9	53.8	83.3	121.8
2002.....	2.4	3.9	5.7	8.4	11.7	17.6	32.1	56.1	85.9	123.8
2001.....	2.7	4.1	6.2	7.8	12.6	19.1	30.2	53.1	86.2	125.2
2000.....	2.8	4.4	6.4	7.9	13.0	20.4	32.0	56.1	91.8	133.1
1999.....	2.3	4.2	5.6	8.8	11.6	18.2	32.3	55.6	88.4	118.9

White, both sexes

2004.....	2.1	3.5	5.3	7.6	11.9	19.4	32.9	56.7	89.8	126.5
2003.....	2.1	3.6	5.0	7.3	11.3	18.5	33.6	57.1	89.5	127.7
2002.....	1.8	3.4	5.0	7.8	12.5	18.8	34.9	58.7	91.3	129.2
2001.....	2.2	3.5	5.1	7.8	12.5	19.1	33.7	57.8	91.7	128.9
2000.....	2.2	3.7	5.6	8.5	13.7	21.7	34.7	59.4	95.6	139.4
1999.....	1.9	3.3	5.1	8.5	12.2	19.4	35.4	58.9	90.7	123.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intracerebral and other intracranial hemorrhage (I61-I62)

White, male

2004.....	2.7	4.2	6.6	9.0	14.1	22.9	36.1	62.3	99.1	137.6
2003.....	2.5	4.4	6.3	8.3	13.2	21.7	38.0	62.7	100.0	138.9
2002.....	2.1	4.2	6.1	8.7	15.1	22.2	40.0	63.5	99.5	136.9
2001.....	2.5	4.5	6.0	9.2	14.9	21.2	39.7	65.1	100.4	134.9
2000.....	2.5	4.7	6.6	10.7	16.1	24.5	38.9	65.5	101.3	150.5
1999.....	2.3	3.7	6.2	10.1	14.8	22.2	40.7	64.0	93.7	131.7

White, female

2004.....	1.5	2.8	4.1	6.4	9.9	16.3	30.3	52.5	84.0	121.5
2003.....	1.7	2.8	3.8	6.2	9.6	15.7	30.1	53.0	83.0	122.8
2002.....	1.5	2.6	4.0	7.0	10.1	15.8	30.7	55.2	86.2	125.9
2001.....	1.9	2.6	4.1	6.5	10.2	17.3	28.8	52.7	86.4	126.3
2000.....	1.8	2.7	4.6	6.4	11.4	19.2	31.3	55.1	92.2	134.9
1999.....	1.5	2.8	4.1	7.1	9.7	16.9	31.2	55.3	89.0	119.7

Black, both sexes

2004.....	7.9	15.9	22.9	26.1	31.4	35.6	44.6	61.1	88.4	95.6
2003.....	8.9	16.4	25.8	27.3	34.1	33.8	43.4	64.2	86.1	107.4
2002.....	8.8	15.7	22.9	24.1	28.1	34.8	44.6	61.7	81.1	98.3
2001.....	8.7	16.6	25.5	22.6	32.0	37.0	43.7	59.2	83.2	106.6
2000.....	10.0	19.0	23.4	27.5	32.3	36.1	42.4	63.3	78.2	117.0
1999.....	9.4	16.1	22.3	26.6	29.6	30.5	44.4	62.5	77.4	96.9

Black, male

2004.....	8.6	19.0	27.1	35.1	42.0	42.6	53.1	64.1	96.1	100.3
2003.....	9.2	18.7	32.9	34.9	44.2	37.7	54.5	75.3	98.9	116.8
2002.....	10.4	19.3	30.0	30.8	36.0	43.4	52.4	61.5	81.0	101.8
2001.....	9.3	20.3	31.3	29.1	36.4	46.0	54.2	62.7	85.9	112.3
2000.....	11.6	23.4	28.5	38.4	43.5	46.3	47.1	69.9	78.4	130.3
1999.....	11.6	19.2	28.6	33.2	37.8	34.0	48.2	70.8	81.1	109.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intracerebral and other intracranial hemorrhage (I61-I62)

Black, female

2004.....	7.2	13.2	19.3	18.7	23.0	30.4	38.9	59.3	84.3	93.7
2003.....	8.6	14.3	19.7	21.1	26.2	30.9	35.9	57.5	79.3	103.7
2002.....	7.4	12.5	16.9	18.6	21.8	28.5	39.4	61.7	81.2	96.9
2001.....	8.1	13.4	20.6	17.2	28.5	30.4	36.8	57.1	81.8	104.4
2000.....	8.5	15.2	19.1	18.6	23.5	28.6	39.3	59.2	78.1	112.0
1999.....	7.4	13.3	16.9	21.2	23.1	27.9	41.9	57.5	75.6	92.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cerebral infarction (I63)

All races 3/, both sexes

2004.....	0.4	0.6	1.0	1.8	3.1	5.5	10.3	20.3	37.8	97.7
2003.....	0.4	0.8	1.1	1.8	3.0	6.2	10.9	21.6	44.5	112.2
2002.....	0.3	0.7	1.3	1.7	3.5	5.8	11.6	22.6	45.6	118.3
2001.....	0.4	0.8	1.1	1.9	3.1	6.1	13.0	24.3	47.6	124.8
2000.....	0.5	0.7	1.2	2.0	4.0	6.6	12.8	27.2	54.5	141.0
1999.....	0.5	0.7	1.2	2.4	4.0	8.1	14.9	29.7	58.5	153.1

All races 3/, male

2004.....	0.4	0.7	1.4	2.3	4.1	6.8	11.6	23.5	39.7	88.5
2003.....	0.5	1.0	1.3	2.2	4.0	7.6	13.2	23.4	47.7	100.3
2002.....	0.3	0.9	1.6	2.3	4.2	7.2	12.4	25.8	50.1	108.7
2001.....	0.4	0.9	1.2	2.6	3.8	7.7	14.9	28.0	53.3	111.2
2000.....	0.5	0.9	1.4	2.5	5.2	7.8	15.4	31.2	60.8	133.3
1999.....	0.5	0.9	1.5	2.9	4.8	10.0	17.4	34.0	63.3	144.6

All races 3/, female

2004.....	0.4	0.6	0.6	1.3	2.2	4.3	9.2	18.0	36.7	101.9
2003.....	0.3	0.5	0.9	1.5	2.1	5.1	9.0	20.4	42.6	117.5
2002.....	0.3	0.5	0.9	1.1	2.9	4.6	11.0	20.4	42.8	122.4
2001.....	0.3	0.7	0.9	1.3	2.5	4.6	11.4	21.8	44.2	130.4
2000.....	0.4	0.5	1.1	1.6	3.0	5.6	10.7	24.4	50.7	144.2
1999.....	0.4	0.5	0.9	1.8	3.3	6.4	12.9	26.7	55.7	156.6

White, both sexes

2004.....	0.3	0.5	0.8	1.5	2.6	5.0	9.6	19.5	37.5	97.6
2003.....	0.3	0.6	0.9	1.5	2.5	5.9	10.1	21.0	44.4	113.7
2002.....	0.3	0.6	1.0	1.5	3.2	5.3	10.9	22.1	44.8	119.1
2001.....	0.3	0.6	0.9	1.6	2.7	5.4	12.3	23.6	47.5	126.0
2000.....	0.3	0.6	1.0	1.6	3.7	6.0	12.0	26.3	54.0	141.7
1999.....	0.4	0.6	1.0	2.0	3.5	7.5	13.8	28.5	58.6	154.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cerebral infarction (I63)

White, male

2004.....	0.4	0.6	1.1	2.0	3.3	6.3	10.6	22.5	39.7	87.9
2003.....	0.4	0.9	1.2	1.8	3.4	7.2	12.2	22.4	48.5	101.5
2002.....	0.3	0.9	1.3	2.0	3.7	6.5	11.7	25.3	49.3	108.6
2001.....	0.4	0.7	1.1	2.2	3.4	7.1	14.3	26.8	52.9	110.5
2000.....	0.4	0.8	1.0	2.0	4.7	7.0	14.6	29.8	59.8	133.1
1999.....	0.4	0.7	1.3	2.6	4.2	9.5	15.9	32.3	63.5	144.0

White, female

2004.....	0.3	0.5	0.5	1.1	2.0	3.9	8.7	17.3	36.2	102.0
2003.....	0.2	0.3	0.6	1.3	1.7	4.7	8.5	19.9	41.8	119.1
2002.....	0.3	0.4	0.8	1.0	2.7	4.2	10.3	19.8	42.1	123.7
2001.....	0.2	0.5	0.8	1.1	2.1	4.0	10.7	21.3	44.2	132.5
2000.....	0.3	0.4	0.9	1.2	2.7	5.2	10.0	23.8	50.5	145.2
1999.....	0.3	0.5	0.8	1.5	2.9	5.7	12.1	25.9	55.7	158.5

Black, both sexes

2004.....	1.0	1.5	2.2	4.1	7.4	9.5	19.1	33.2	43.8	103.5
2003.....	0.9	1.6	2.6	4.7	7.7	10.6	18.8	32.3	51.8	107.7
2002.....	*	1.3	3.1	3.6	6.7	11.0	20.0	32.3	56.8	113.1
2001.....	0.9	2.0	2.2	4.4	7.3	12.3	21.7	34.9	53.4	121.3
2000.....	1.2	1.3	3.1	5.4	7.8	13.0	21.1	37.7	67.5	142.6
1999.....	1.1	1.8	2.7	5.0	8.7	14.4	27.7	47.0	62.7	150.8

Black, male

2004.....	*	2.0	3.1	5.1	11.2	13.6	22.7	41.9	45.9	97.4
2003.....	*	2.1	2.8	6.0	10.8	13.0	25.0	38.8	50.2	97.0
2002.....	*	*	4.4	4.9	8.9	14.7	22.7	35.8	63.4	114.7
2001.....	*	2.4	2.6	6.6	8.3	15.8	23.2	47.4	65.0	127.1
2000.....	*	*	4.5	7.1	10.1	17.7	24.7	46.8	83.5	137.2
1999.....	*	2.7	3.1	6.3	10.8	17.1	37.7	60.3	68.9	162.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cerebral infarction (I63)

Black, female

2004.....	*	*	*	3.1	4.4	6.5	16.7	28.0	42.7	105.9
2003.....	*	*	2.5	3.7	5.3	8.8	14.6	28.5	52.6	111.8
2002.....	*	*	2.0	2.4	5.0	8.3	18.1	30.2	53.3	112.5
2001.....	*	1.7	1.9	2.7	6.4	9.8	20.6	27.5	47.4	119.1
2000.....	1.3	*	*	4.0	5.9	9.6	18.6	32.2	59.4	144.7
1999.....	*	*	2.4	3.9	7.0	12.3	21.2	38.9	59.6	146.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Stroke, not specified as hemorrhage or infarction (I64)

All races 3/, both sexes

2004.....	1.3	2.6	4.9	8.6	17.9	36.0	70.1	141.8	288.1	713.4
2003.....	1.5	2.4	5.1	9.9	19.5	36.9	75.1	155.8	315.4	799.7
2002.....	1.2	2.8	5.3	10.7	20.5	40.3	82.7	165.7	338.0	859.4
2001.....	1.2	2.5	5.2	10.5	22.0	41.7	85.7	176.0	355.4	906.2
2000.....	1.2	2.6	5.2	10.7	23.0	44.6	89.8	181.1	372.2	954.7
1999.....	1.1	2.7	5.0	11.3	22.4	45.1	89.6	186.1	380.9	982.4

All races 3/, male

2004.....	1.5	3.0	6.1	10.0	21.4	41.6	80.0	152.1	292.2	615.5
2003.....	1.7	2.9	6.1	12.1	23.3	43.1	85.5	171.8	321.2	699.0
2002.....	1.2	3.2	6.1	13.2	24.3	47.5	93.1	179.9	334.5	762.8
2001.....	1.3	3.0	6.2	12.6	26.5	49.4	97.9	188.5	363.9	819.7
2000.....	1.3	3.1	6.5	12.8	27.8	51.0	102.3	198.1	390.9	863.2
1999.....	1.3	3.2	6.2	13.1	27.2	51.7	103.2	206.0	404.7	896.4

All races 3/, female

2004.....	1.1	2.2	3.8	7.3	14.8	31.1	62.2	134.3	285.5	757.4
2003.....	1.2	2.0	4.1	7.9	16.1	31.6	66.7	144.4	311.9	844.2
2002.....	1.2	2.4	4.4	8.4	16.9	34.1	74.3	155.8	340.2	901.3
2001.....	1.1	2.0	4.2	8.6	17.8	35.1	76.1	167.2	350.3	942.4
2000.....	1.1	2.2	3.9	8.6	18.7	39.2	80.0	169.2	361.2	992.0
1999.....	0.9	2.2	3.9	9.6	18.2	39.4	78.9	172.4	367.0	1,017.1

White, both sexes

2004.....	1.0	1.9	3.7	6.9	14.7	31.1	64.6	134.4	284.2	721.0
2003.....	1.0	1.8	3.9	8.0	16.0	32.2	69.0	148.7	312.5	807.1
2002.....	1.0	2.1	4.0	8.6	16.6	34.9	76.1	159.4	333.5	867.3
2001.....	0.9	1.9	3.9	8.5	17.8	35.8	78.8	169.6	350.8	913.6
2000.....	1.0	1.9	4.1	8.4	19.4	39.5	83.1	174.3	366.9	963.5
1999.....	0.8	1.9	3.8	9.0	18.4	39.9	82.3	178.9	377.1	993.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Stroke, not specified as hemorrhage or infarction (I64)

White, male

2004.....	1.1	2.3	4.7	7.9	17.6	36.1	73.3	144.3	288.8	622.4
2003.....	1.2	2.2	4.7	9.5	19.5	37.9	78.3	164.2	317.7	704.0
2002.....	1.1	2.5	4.7	10.6	19.6	40.6	85.5	172.6	329.7	766.7
2001.....	1.1	2.3	4.7	10.0	21.2	42.3	89.6	181.7	357.5	823.5
2000.....	1.1	2.2	5.3	10.1	23.6	45.2	94.2	189.8	384.8	865.2
1999.....	1.0	2.3	4.8	10.7	22.3	45.4	95.3	197.4	401.4	903.1

White, female

2004.....	0.8	1.6	2.7	6.0	12.1	26.7	57.5	127.1	281.3	765.4
2003.....	0.8	1.5	3.0	6.4	12.8	27.1	61.3	137.4	309.3	852.8
2002.....	0.9	1.6	3.3	6.7	13.8	30.0	68.4	149.9	335.9	910.9
2001.....	0.7	1.4	3.1	7.0	14.6	30.1	70.1	161.0	346.7	951.3
2000.....	0.8	1.5	2.9	6.8	15.5	34.6	74.2	163.4	356.3	1,003.5
1999.....	0.6	1.5	2.9	7.4	14.9	35.1	72.0	165.9	362.8	1,029.7

Black, both sexes

2004.....	3.7	7.4	14.4	24.1	48.5	82.1	132.2	236.6	367.9	706.1
2003.....	4.3	7.1	14.9	27.8	52.6	81.9	142.9	250.5	386.3	797.6
2002.....	3.2	8.0	15.7	30.9	55.6	91.8	156.7	259.7	425.2	840.7
2001.....	3.6	7.1	16.3	29.3	59.5	96.7	163.1	268.9	435.0	883.4
2000.....	2.9	8.6	14.7	31.0	55.2	97.3	167.8	275.6	473.0	919.4
1999.....	3.4	8.5	16.0	32.5	59.9	97.2	174.4	287.1	454.8	917.6

Black, male

2004.....	4.1	8.5	17.2	30.9	61.5	98.6	159.6	257.5	375.9	589.1
2003.....	5.3	8.1	18.3	36.2	63.0	95.5	170.8	285.3	410.5	681.9
2002.....	2.5	8.9	18.8	40.0	71.1	118.4	194.6	293.9	428.7	761.2
2001.....	3.6	8.2	19.8	37.9	77.9	122.7	199.6	302.9	470.3	816.9
2000.....	3.0	10.0	18.6	39.0	66.7	117.1	203.6	321.8	519.3	897.2
1999.....	3.8	10.1	19.8	37.8	77.2	120.2	202.2	328.6	489.2	877.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Stroke, not specified as hemorrhage or infarction (I64)

Black, female

2004.....	3.3	6.4	12.0	18.5	38.3	69.8	113.6	224.3	363.6	752.5
2003.....	3.5	6.2	12.0	20.9	44.4	71.8	124.1	229.8	373.7	842.6
2002.....	3.8	7.2	13.0	23.5	43.5	72.0	131.3	239.3	423.3	871.2
2001.....	3.7	6.1	13.3	22.2	45.0	77.5	138.8	248.5	417.0	908.1
2000.....	2.8	7.4	11.4	24.4	46.1	82.8	144.1	247.8	449.6	927.6
1999.....	3.1	7.2	12.7	28.1	46.3	80.2	156.0	262.0	437.8	932.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other cerebrovascular diseases and their sequelae (I67,I69)

All races 3/, both sexes

2004.....	0.5	0.7	1.4	2.6	5.2	11.0	21.7	49.4	107.4	298.3
2003.....	0.4	0.8	1.3	2.8	5.1	11.1	24.0	51.3	111.4	320.1
2002.....	0.5	0.8	1.5	2.7	5.2	11.3	24.2	53.4	117.2	328.9
2001.....	0.4	0.7	1.4	2.7	5.4	11.4	24.1	53.6	118.5	328.2
2000.....	0.4	0.8	1.4	2.8	5.8	11.0	24.6	55.7	121.5	341.4
1999.....	0.4	0.8	1.4	2.9	5.6	11.7	25.8	56.9	123.1	343.9

All races 3/, male

2004.....	0.5	0.7	1.6	3.1	6.5	13.7	25.4	56.8	115.0	270.4
2003.....	0.3	0.9	1.4	3.2	6.3	13.2	28.1	58.7	119.4	290.2
2002.....	0.4	0.8	1.7	3.2	6.2	13.3	27.6	60.2	126.5	302.3
2001.....	0.4	0.7	1.5	3.2	6.5	14.2	28.3	61.4	130.6	305.5
2000.....	0.4	0.8	1.6	3.1	6.8	13.3	30.1	66.1	135.0	326.9
1999.....	0.4	0.8	1.5	3.6	7.1	13.4	31.0	65.8	136.9	332.1

All races 3/, female

2004.....	0.4	0.7	1.3	2.2	4.1	8.7	18.8	44.1	102.8	310.8
2003.....	0.5	0.7	1.2	2.4	4.0	9.3	20.8	46.0	106.6	333.3
2002.....	0.6	0.8	1.2	2.4	4.3	9.5	21.5	48.6	111.5	340.4
2001.....	0.4	0.7	1.2	2.3	4.4	9.0	20.7	48.2	111.3	337.6
2000.....	0.4	0.9	1.3	2.4	5.0	9.0	20.3	48.5	113.6	347.4
1999.....	0.5	0.9	1.2	2.3	4.2	10.2	21.8	50.7	115.1	348.7

White, both sexes

2004.....	0.4	0.6	1.2	2.2	4.3	9.8	19.7	46.9	105.6	300.3
2003.....	0.3	0.6	1.0	2.3	4.2	9.6	22.0	49.2	109.8	322.5
2002.....	0.4	0.6	1.2	2.4	4.3	9.8	22.4	50.7	115.9	332.0
2001.....	0.3	0.6	1.0	2.2	4.5	10.0	22.3	51.0	117.3	329.9
2000.....	0.3	0.7	1.2	2.3	5.0	9.9	23.1	52.9	120.2	343.4
1999.....	0.4	0.7	1.1	2.4	4.7	10.5	23.3	54.7	121.3	348.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other cerebrovascular diseases and their sequelae (I67,I69)

White, male										
2004.....	0.5	0.5	1.2	2.6	5.3	12.2	22.9	53.4	112.8	271.5
2003.....	0.3	0.7	1.0	2.5	5.2	11.3	25.9	55.6	117.0	294.0
2002.....	0.3	0.7	1.4	2.7	5.1	11.7	25.8	56.5	125.7	306.3
2001.....	0.3	0.6	1.0	2.7	5.5	12.8	26.2	58.6	130.0	308.1
2000.....	0.3	0.7	1.3	2.5	6.2	11.8	27.9	63.3	132.5	330.0
1999.....	0.3	0.6	1.1	2.9	5.9	12.1	27.4	63.2	134.3	334.9
White, female										
2004.....	0.4	0.6	1.1	1.8	3.4	7.7	17.1	42.2	101.1	313.3
2003.....	0.4	0.5	1.0	2.0	3.3	8.2	18.8	44.5	105.3	335.2
2002.....	0.5	0.6	1.0	2.0	3.5	8.1	19.7	46.6	109.9	343.2
2001.....	0.3	0.6	0.9	1.8	3.6	7.6	19.2	45.6	109.6	339.1
2000.....	0.3	0.6	1.0	2.1	4.0	8.1	19.3	45.7	112.9	348.8
1999.....	0.4	0.7	1.1	1.9	3.6	9.2	20.2	48.7	113.6	353.4
Black, both sexes										
2004.....	0.7	2.0	3.1	6.9	14.1	22.8	44.9	79.7	137.6	296.9
2003.....	0.7	2.0	3.7	7.2	13.9	24.5	45.8	77.6	141.1	321.5
2002.....	0.8	1.8	4.0	6.5	13.9	25.8	44.5	85.2	140.2	318.9
2001.....	0.9	1.7	4.4	7.2	13.6	24.0	42.8	87.5	138.1	327.6
2000.....	0.9	2.2	3.8	7.3	12.5	20.9	41.1	87.6	140.3	333.8
1999.....	0.9	2.0	3.3	7.6	13.3	22.3	52.6	83.9	149.1	312.5
Black, male										
2004.....	*	2.1	4.3	9.1	18.7	29.8	58.6	102.4	152.8	272.2
2003.....	*	2.4	4.6	8.4	17.9	33.0	55.2	97.7	158.7	271.1
2002.....	*	*	5.0	7.9	17.6	31.4	52.4	105.6	150.5	269.1
2001.....	*	*	4.9	8.3	16.7	29.5	54.2	103.6	143.7	285.9
2000.....	*	*	4.2	9.6	12.4	27.1	56.2	98.3	177.0	291.8
1999.....	*	2.3	4.6	10.3	18.7	27.3	73.3	96.4	168.3	294.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other cerebrovascular diseases and their sequelae (I67,I69)

Black, female

2004.....	*	1.9	2.2	5.0	10.5	17.6	35.6	66.3	129.7	306.8
2003.....	*	1.6	3.0	6.1	10.8	18.3	39.4	65.7	131.9	341.1
2002.....	*	1.9	3.2	5.4	11.0	21.6	39.2	73.0	134.9	338.0
2001.....	*	1.8	4.1	6.3	11.2	19.9	35.2	77.9	135.3	343.1
2000.....	*	2.6	3.5	5.4	12.6	16.3	31.1	81.2	121.9	349.4
1999.....	*	1.7	2.3	5.4	9.2	18.7	38.9	76.4	139.6	319.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of arteries, arterioles and capillaries (I70-I78)

All races 3/, both sexes

2004.....	1.4	2.4	3.9	6.2	12.2	22.4	41.5	69.1	119.1	276.3
2003.....	1.4	2.7	4.2	6.8	13.8	25.0	43.2	76.4	126.2	300.6
2002.....	1.4	2.5	3.8	7.6	14.1	27.5	45.8	78.3	131.6	316.5
2001.....	1.4	2.3	3.9	7.4	14.6	27.3	46.5	82.5	137.2	337.3
2000.....	1.4	2.5	4.1	7.7	15.3	29.6	51.1	85.6	144.6	352.0
1999.....	1.5	2.4	4.0	8.0	15.4	28.6	51.3	87.6	147.8	367.2

All races 3/, male

2004.....	1.8	3.4	5.4	8.4	16.6	29.6	54.4	87.3	149.4	294.7
2003.....	1.9	3.7	5.9	9.3	19.1	34.8	56.6	102.8	159.5	318.3
2002.....	2.0	3.6	5.2	10.3	19.6	38.1	60.7	102.7	166.1	344.0
2001.....	2.0	3.3	5.2	10.0	20.0	37.0	63.3	106.0	175.7	364.5
2000.....	1.9	3.3	5.8	11.0	21.6	41.2	70.6	116.1	186.4	391.4
1999.....	1.8	3.4	5.5	11.3	22.0	39.8	70.0	117.2	189.4	396.5

All races 3/, female

2004.....	0.9	1.4	2.4	4.2	8.3	16.2	31.1	56.1	100.5	268.0
2003.....	0.9	1.8	2.5	4.4	8.9	16.4	32.4	57.7	105.8	292.8
2002.....	0.8	1.5	2.5	5.0	9.2	18.3	33.8	61.1	110.6	304.6
2001.....	0.9	1.4	2.7	5.0	9.7	18.9	33.1	66.0	114.2	325.9
2000.....	1.0	1.7	2.5	4.6	9.6	19.7	35.6	64.4	120.0	336.0
1999.....	1.2	1.5	2.7	5.0	9.5	19.1	36.7	67.0	123.5	355.4

White, both sexes

2004.....	1.2	2.1	3.5	5.8	11.6	22.1	42.3	69.9	120.5	281.3
2003.....	1.2	2.4	3.7	6.5	13.4	25.0	43.6	77.7	126.9	306.4
2002.....	1.2	2.1	3.3	7.1	13.7	27.7	46.1	79.0	133.2	323.7
2001.....	1.2	2.1	3.5	7.0	14.0	27.5	46.6	82.8	138.7	344.1
2000.....	1.2	2.1	3.5	7.2	14.8	29.7	51.8	86.1	146.1	358.3
1999.....	1.3	2.1	3.7	7.6	15.4	28.7	52.1	88.3	149.4	372.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of arteries, arterioles and capillaries (I70-I78)

White, male

2004.....	1.6	3.0	5.0	7.7	15.7	29.4	56.1	88.6	152.3	303.7
2003.....	1.7	3.3	5.3	9.1	18.6	35.0	57.5	104.4	160.3	327.4
2002.....	1.8	3.1	4.4	9.6	19.0	39.0	61.2	104.5	169.2	354.1
2001.....	1.6	2.9	4.6	9.6	19.3	37.6	63.8	107.1	178.0	375.1
2000.....	1.5	2.9	5.0	10.5	21.3	42.0	72.3	117.6	190.0	401.0
1999.....	1.6	3.1	5.0	10.8	22.3	40.5	71.7	118.9	191.8	404.8

White, female

2004.....	0.8	1.2	2.0	3.9	7.8	15.6	30.9	56.2	100.7	271.1
2003.....	0.8	1.4	2.1	4.0	8.6	16.2	32.2	58.5	106.4	297.1
2002.....	0.7	1.2	2.2	4.7	8.8	17.7	33.9	60.8	111.2	310.4
2001.....	0.8	1.3	2.4	4.5	9.0	18.6	32.7	65.6	115.0	331.0
2000.....	0.8	1.3	2.1	4.1	8.7	19.0	35.3	63.9	119.9	340.9
1999.....	1.0	1.2	2.4	4.5	9.2	18.4	36.5	66.8	124.4	360.0

Black, both sexes

2004.....	2.9	4.7	7.2	11.0	20.5	29.6	42.7	75.3	122.0	258.0
2003.....	2.7	5.7	8.6	10.4	19.9	30.5	48.6	76.1	137.6	271.0
2002.....	3.1	5.9	8.1	13.2	20.7	30.9	49.3	85.2	134.7	273.8
2001.....	3.3	4.4	7.5	12.7	23.0	29.2	54.9	92.2	138.9	300.3
2000.....	3.5	5.3	8.3	13.3	22.9	32.9	52.8	92.4	146.3	315.9
1999.....	3.0	4.8	7.4	13.4	18.2	33.0	50.5	91.8	143.3	333.2

Black, male

2004.....	3.8	6.6	9.7	15.4	29.2	39.5	49.5	91.4	139.2	236.4
2003.....	3.7	7.7	11.6	13.9	28.3	41.6	58.1	101.3	177.9	242.9
2002.....	4.4	7.4	12.0	19.1	29.3	36.9	64.1	100.6	159.7	276.6
2001.....	4.7	6.2	10.8	15.3	29.7	35.6	71.0	109.2	176.6	279.1
2000.....	4.6	6.5	12.0	17.5	28.6	41.4	65.9	116.2	171.1	325.2
1999.....	3.5	6.5	10.0	17.5	23.5	40.8	64.8	111.6	174.4	348.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of arteries, arterioles and capillaries (I70-I78)

Black, female

2004.....	2.1	3.0	5.0	7.4	13.8	22.3	38.0	65.8	113.0	266.6
2003.....	1.8	4.0	6.1	7.5	13.2	22.3	42.1	61.0	116.4	281.9
2002.....	1.9	4.5	4.8	8.3	13.9	26.5	39.4	76.0	121.7	272.7
2001.....	2.2	2.9	4.7	10.5	17.8	24.4	44.2	82.1	119.8	308.2
2000.....	2.5	4.4	5.2	9.9	18.5	26.7	44.2	78.0	133.7	312.5
1999.....	2.7	3.3	5.2	10.1	14.0	27.3	41.0	79.8	127.9	327.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerosis (I70)

All races 3/, both sexes

2004.....	0.1	0.3	0.5	1.0	2.3	4.1	8.5	17.5	38.0	137.0
2003.....	0.2	0.3	0.6	1.1	2.4	4.5	9.0	19.9	43.5	154.6
2002.....	0.1	0.3	0.6	1.4	2.4	5.2	9.3	20.5	47.8	168.1
2001.....	0.1	0.3	0.7	1.4	2.5	4.9	9.8	21.9	49.2	179.1
2000.....	0.1	0.4	0.6	1.3	2.6	5.2	10.5	22.7	52.4	188.8
1999.....	0.1	0.3	0.8	1.3	2.9	5.1	11.2	24.1	53.6	202.0

All races 3/, male

2004.....	*	0.4	0.7	1.2	3.0	4.9	10.3	20.2	43.5	128.2
2003.....	0.2	0.5	0.8	1.5	3.2	5.4	10.6	23.9	48.3	138.6
2002.....	0.2	0.4	0.8	1.7	3.2	6.4	11.5	24.4	52.5	156.6
2001.....	*	0.4	0.8	1.7	3.2	6.2	12.1	25.0	53.2	167.2
2000.....	0.2	0.5	0.8	1.7	3.6	6.6	13.1	27.5	59.8	176.5
1999.....	0.2	0.4	1.0	1.8	3.8	6.3	13.3	28.1	60.0	189.0

All races 3/, female

2004.....	*	*	0.3	0.8	1.7	3.4	7.1	15.5	34.6	141.0
2003.....	*	*	0.4	0.7	1.7	3.7	7.8	17.0	40.6	161.7
2002.....	*	*	0.5	1.0	1.7	4.1	7.5	17.8	45.0	173.1
2001.....	*	0.2	0.5	1.1	1.8	3.9	8.1	19.7	46.8	184.1
2000.....	*	0.2	0.4	0.9	1.7	4.0	8.4	19.4	48.1	193.8
1999.....	*	*	0.5	0.9	2.2	4.0	9.5	21.4	49.8	207.2

White, both sexes

2004.....	0.1	0.3	0.4	1.0	2.3	4.0	8.5	17.8	39.3	142.0
2003.....	0.2	0.3	0.6	1.0	2.4	4.4	8.9	20.2	44.1	159.5
2002.....	0.1	0.2	0.5	1.3	2.3	5.3	9.1	20.4	48.4	174.7
2001.....	0.1	0.3	0.6	1.4	2.3	4.9	9.8	22.2	50.3	185.9
2000.....	0.1	0.3	0.5	1.2	2.4	5.2	10.4	23.1	53.0	195.2
1999.....	*	0.3	0.7	1.2	2.9	5.0	11.3	24.2	54.7	207.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerosis (I70)

White, male

2004.....	*	0.4	0.7	1.2	2.9	4.9	10.4	20.5	44.8	134.4
2003.....	0.2	0.5	0.8	1.4	3.1	5.2	10.1	23.8	48.4	143.3
2002.....	*	0.4	0.6	1.7	3.0	6.5	11.3	24.0	53.1	163.2
2001.....	*	0.4	0.8	1.7	3.0	6.2	12.1	25.2	54.0	173.9
2000.....	*	0.4	0.7	1.7	3.5	6.5	13.2	28.2	60.3	181.4
1999.....	*	0.4	1.0	1.6	3.7	6.2	13.3	27.7	60.8	193.7

White, female

2004.....	*	*	0.2	0.8	1.7	3.2	7.0	15.9	35.9	145.4
2003.....	*	*	0.4	0.7	1.7	3.6	7.8	17.6	41.4	166.7
2002.....	*	*	0.4	1.0	1.6	4.2	7.3	17.7	45.5	179.7
2001.....	*	*	0.5	1.0	1.6	3.9	8.0	20.2	48.0	190.9
2000.....	*	*	0.3	0.8	1.5	4.1	8.2	19.5	48.7	200.8
1999.....	*	*	0.5	0.9	2.0	3.9	9.7	21.7	51.0	213.3

Black, both sexes

2004.....	*	*	*	1.7	3.2	5.4	9.9	18.4	31.9	102.9
2003.....	*	*	1.3	1.9	3.5	6.5	12.7	20.8	45.4	121.1
2002.....	*	*	1.5	1.9	4.2	5.2	11.5	26.8	51.3	117.0
2001.....	*	*	1.0	2.0	5.0	6.3	12.1	22.9	42.8	125.6
2000.....	*	*	1.5	2.1	4.7	5.6	12.9	23.5	53.9	138.3
1999.....	*	*	1.3	2.1	4.7	7.0	11.9	27.5	46.8	154.3

Black, male

2004.....	*	*	*	*	4.9	6.1	12.0	23.2	40.9	78.5
2003.....	*	*	*	*	4.8	8.1	18.1	28.8	59.1	104.3
2002.....	*	*	2.3	*	6.3	6.2	15.4	34.9	59.6	110.4
2001.....	*	*	*	*	5.9	7.6	15.1	30.2	49.0	113.5
2000.....	*	*	*	*	6.0	8.6	14.0	25.5	66.6	147.6
1999.....	*	*	*	3.4	5.7	8.8	15.6	37.0	56.7	164.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerosis (I70)

Black, female

2004.....	*	*	*	*	*	4.9	8.6	15.6	27.2	112.6
2003.....	*	*	*	*	*	5.3	9.1	16.0	38.3	127.6
2002.....	*	*	*	*	*	4.4	9.0	22.0	46.9	119.5
2001.....	*	*	*	*	4.3	5.3	10.1	18.6	39.7	130.1
2000.....	*	*	*	*	3.6	*	12.2	22.2	47.6	134.8
1999.....	*	*	*	*	3.8	5.6	9.5	21.7	41.9	150.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Aortic aneurysm and dissection (I71)

All races 3/, both sexes

2004.....	0.8	1.4	2.2	3.1	6.4	11.4	21.1	32.6	48.8	64.8
2003.....	0.8	1.7	2.4	3.7	7.3	13.5	22.7	36.6	50.7	70.0
2002.....	0.9	1.5	2.0	4.0	7.4	14.7	23.9	37.5	51.5	69.4
2001.....	0.9	1.4	2.0	3.7	7.8	15.0	23.8	39.0	55.4	77.5
2000.....	0.9	1.3	2.2	4.3	8.4	16.3	27.8	41.0	56.5	79.2
1999.....	0.9	1.4	2.1	4.4	8.1	16.5	27.3	42.2	58.2	80.3

All races 3/, male

2004.....	1.3	2.1	3.3	4.9	9.7	17.0	30.6	44.2	70.3	95.0
2003.....	1.3	2.5	3.7	5.8	11.4	20.2	33.0	53.8	74.2	105.5
2002.....	1.4	2.4	3.2	6.3	11.5	23.0	34.9	55.0	74.5	106.6
2001.....	1.4	2.1	3.2	5.8	12.0	22.2	36.0	56.3	82.5	114.8
2000.....	1.3	1.9	3.5	6.7	13.1	24.9	41.8	61.9	84.9	127.3
1999.....	1.2	2.2	3.3	6.8	13.0	25.3	41.0	63.3	88.6	125.3

All races 3/, female

2004.....	0.4	0.7	1.0	1.5	3.3	6.6	13.5	24.3	35.6	51.2
2003.....	0.4	0.9	1.1	1.7	3.6	7.8	14.5	24.3	36.3	54.2
2002.....	0.3	0.7	1.0	1.8	3.7	7.5	15.1	25.1	37.6	53.3
2001.....	0.3	0.6	0.9	1.8	3.9	8.8	14.0	26.9	39.2	61.9
2000.....	0.5	0.8	1.0	2.1	4.1	8.9	16.7	26.5	39.7	59.6
1999.....	0.5	0.7	1.0	2.1	3.7	9.0	16.6	27.5	40.4	62.2

White, both sexes

2004.....	0.7	1.2	1.9	2.8	6.1	11.7	22.1	33.7	49.8	66.8
2003.....	0.7	1.4	2.1	3.6	7.3	14.0	23.6	38.1	52.2	72.4
2002.....	0.7	1.2	1.7	3.8	7.5	15.3	24.8	39.0	53.1	71.2
2001.....	0.7	1.2	1.7	3.6	7.7	15.5	24.6	39.8	57.0	79.7
2000.....	0.7	1.2	2.0	4.1	8.4	16.7	29.0	42.0	58.1	80.8
1999.....	0.7	1.2	1.9	4.2	8.5	17.1	28.4	43.5	60.0	82.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Aortic aneurysm and dissection (I71)

White, male

2004.....	1.1	1.8	3.0	4.5	9.3	17.4	32.1	45.8	72.1	98.7
2003.....	1.2	2.1	3.4	5.7	11.4	21.1	34.6	56.2	76.5	110.0
2002.....	1.2	2.0	2.7	5.9	11.6	24.3	36.1	57.5	77.7	111.8
2001.....	1.1	1.9	2.8	5.6	11.9	23.2	37.3	57.7	85.5	120.3
2000.....	1.0	1.8	3.2	6.5	13.3	26.1	43.8	63.6	88.6	132.8
1999.....	1.0	1.9	2.9	6.7	13.6	26.5	43.0	66.3	91.5	129.8

White, female

2004.....	0.3	0.5	0.9	1.2	3.2	6.7	13.9	24.8	35.9	52.4
2003.....	0.3	0.7	0.8	1.5	3.5	7.9	14.6	25.0	37.2	55.7
2002.....	0.3	0.5	0.8	1.7	3.7	7.4	15.5	25.8	38.1	53.5
2001.....	0.3	0.5	0.7	1.6	3.9	8.8	14.3	27.1	39.8	62.6
2000.....	0.3	0.5	0.9	1.9	3.9	8.5	17.1	26.8	40.0	59.6
1999.....	0.4	0.5	0.9	1.8	3.7	8.9	16.7	27.5	41.3	63.1

Black, both sexes

2004.....	1.8	3.2	4.0	5.8	10.0	11.6	14.9	26.1	40.6	43.7
2003.....	1.8	3.8	4.7	5.7	9.0	11.7	17.9	25.9	36.0	46.4
2002.....	2.1	3.8	4.7	6.8	8.3	11.6	18.4	24.9	37.4	50.5
2001.....	2.1	2.7	4.4	6.2	9.3	11.6	18.4	34.4	42.5	55.3
2000.....	2.3	2.9	3.7	7.3	9.3	14.3	20.3	33.8	40.4	62.3
1999.....	2.0	3.5	4.0	6.5	6.5	14.1	19.8	29.6	40.5	60.3

Black, male

2004.....	2.8	5.1	6.0	8.8	15.9	17.9	18.5	29.6	48.1	56.6
2003.....	2.8	5.8	7.1	8.0	14.7	16.3	20.4	32.9	49.5	51.1
2002.....	3.5	5.7	7.7	11.3	13.1	14.7	24.7	31.2	41.3	49.3
2001.....	3.4	3.9	6.9	8.6	14.6	15.0	26.6	43.2	54.6	48.8
2000.....	3.5	3.6	6.0	10.4	12.8	17.7	27.7	47.7	45.5	69.2
1999.....	2.7	5.4	6.4	9.0	9.3	19.2	25.4	29.9	54.1	76.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Aortic aneurysm and dissection (I71)

Black, female

2004.....	*	1.6	2.4	3.3	5.4	6.8	12.5	24.0	36.7	38.6
2003.....	*	2.1	2.7	3.8	4.7	8.4	16.2	21.7	29.0	44.6
2002.....	*	2.1	2.2	3.2	4.5	9.3	14.1	21.1	35.4	50.9
2001.....	*	1.7	2.2	4.2	5.1	9.0	13.0	29.1	36.4	57.7
2000.....	*	2.3	*	4.7	6.5	11.7	15.3	25.3	37.8	59.7
1999.....	1.4	2.0	*	4.4	4.3	10.4	16.0	29.5	33.7	54.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races 3/, both sexes

2004.....	0.4	0.7	1.2	2.1	3.5	6.9	11.9	19.1	32.3	74.5
2003.....	0.4	0.7	1.2	2.0	4.0	7.0	11.4	20.0	32.0	76.0
2002.....	0.4	0.7	1.1	2.2	4.3	7.7	12.6	20.4	32.3	79.0
2001.....	0.5	0.6	1.2	2.2	4.3	7.3	12.9	21.5	32.7	80.7
2000.....	0.4	0.8	1.3	2.1	4.3	8.2	12.8	21.8	35.8	84.0
1999.....	0.5	0.7	1.2	2.3	4.4	7.0	12.9	21.3	36.1	84.9

All races 3/, male

2004.....	0.4	0.8	1.4	2.3	3.9	7.7	13.6	22.9	35.5	71.6
2003.....	0.3	0.7	1.3	2.0	4.4	9.3	13.0	25.1	37.1	74.2
2002.....	0.4	0.8	1.2	2.3	4.9	8.7	14.2	23.4	39.2	80.8
2001.....	0.4	0.7	1.2	2.4	4.8	8.5	15.2	24.7	40.0	82.5
2000.....	0.3	0.8	1.4	2.6	4.9	9.7	15.7	26.6	41.7	87.6
1999.....	0.4	0.8	1.2	2.6	5.2	8.1	15.7	25.9	40.8	82.2

All races 3/, female

2004.....	0.5	0.6	1.0	1.8	3.2	6.2	10.5	16.3	30.3	75.8
2003.....	0.5	0.7	1.0	1.9	3.6	5.0	10.1	16.3	28.9	76.9
2002.....	0.4	0.6	1.0	2.1	3.7	6.7	11.3	18.3	28.1	78.2
2001.....	0.5	0.6	1.3	2.0	3.9	6.3	11.1	19.3	28.3	79.9
2000.....	0.5	0.7	1.1	1.6	3.7	6.9	10.5	18.5	32.2	82.6
1999.....	0.5	0.6	1.2	2.1	3.6	6.0	10.6	18.2	33.3	86.0

White, both sexes

2004.....	0.4	0.7	1.1	1.9	3.2	6.4	11.7	18.4	31.4	72.5
2003.....	0.3	0.6	1.0	1.9	3.7	6.6	11.1	19.5	30.7	74.6
2002.....	0.4	0.6	1.0	2.0	3.9	7.1	12.2	19.7	31.7	77.8
2001.....	0.4	0.6	1.2	2.1	4.0	7.1	12.2	20.8	31.4	78.5
2000.....	0.3	0.7	1.0	1.9	3.9	7.8	12.4	21.0	34.9	82.3
1999.....	0.5	0.7	1.1	2.1	4.1	6.6	12.5	20.6	34.8	82.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

White, male

2004.....	0.4	0.8	1.3	2.1	3.6	7.1	13.6	22.3	35.4	70.7
2003.....	0.3	0.7	1.2	2.0	4.0	8.7	12.8	24.4	35.4	74.1
2002.....	0.4	0.8	1.1	2.0	4.4	8.2	13.7	23.0	38.4	79.1
2001.....	0.3	0.6	1.1	2.3	4.4	8.3	14.4	24.2	38.5	80.9
2000.....	0.3	0.7	1.1	2.4	4.5	9.4	15.3	25.9	41.1	86.9
1999.....	0.4	0.8	1.1	2.4	4.9	7.9	15.3	24.8	39.6	81.3

White, female

2004.....	0.4	0.5	0.9	1.8	2.8	5.7	10.1	15.6	28.9	73.3
2003.....	0.4	0.6	0.9	1.9	3.4	4.8	9.8	15.9	27.8	74.8
2002.....	0.3	0.5	0.9	2.0	3.4	6.1	11.0	17.3	27.5	77.2
2001.....	0.5	0.6	1.2	1.8	3.5	6.0	10.4	18.4	27.1	77.5
2000.....	0.4	0.6	0.9	1.4	3.3	6.5	10.0	17.6	31.2	80.4
1999.....	0.5	0.5	1.0	1.8	3.4	5.6	10.2	17.6	32.0	83.5

Black, both sexes

2004.....	0.8	1.1	2.3	3.6	7.4	12.7	17.8	30.8	49.5	111.4
2003.....	0.8	1.5	2.6	2.8	7.3	12.3	18.0	29.4	56.1	103.6
2002.....	0.7	1.5	2.0	4.4	8.3	14.1	19.4	33.5	46.1	106.3
2001.....	1.1	1.3	2.1	4.5	8.7	11.3	24.4	34.9	53.6	119.4
2000.....	0.9	1.7	3.1	3.9	9.0	13.0	19.6	35.2	52.0	115.4
1999.....	0.8	1.0	2.1	4.9	7.0	11.9	18.8	34.6	56.1	118.6

Black, male

2004.....	*	*	2.9	4.9	8.4	15.5	19.1	38.7	50.2	101.3
2003.....	*	*	2.8	3.2	8.9	17.2	19.7	39.7	69.4	87.6
2002.....	*	*	2.1	5.2	9.9	16.0	24.0	34.4	58.8	116.9
2001.....	*	*	2.5	4.5	9.2	13.0	29.3	35.8	73.0	116.9
2000.....	*	1.9	4.0	4.8	9.7	15.1	24.1	43.0	59.0	108.4
1999.....	*	*	*	5.1	8.5	12.7	23.7	44.6	63.7	107.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Black, female

2004.....	*	*	1.8	2.5	6.6	10.6	16.9	26.1	49.1	115.4
2003.....	*	1.8	2.4	2.4	6.0	8.6	16.8	23.3	49.1	109.8
2002.....	*	1.9	1.8	3.7	7.0	12.8	16.4	32.9	39.4	102.3
2001.....	*	*	1.8	4.5	8.3	10.1	21.1	34.4	43.7	120.4
2000.....	*	*	2.4	3.2	8.3	11.5	16.6	30.5	48.4	118.0
1999.....	*	*	*	4.7	5.8	11.4	15.5	28.6	52.3	122.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes

2004.....	0.9	1.3	1.4	1.9	2.5	3.1	4.3	7.1	11.5	23.3
2003.....	0.9	1.3	1.5	1.7	2.5	3.3	4.9	7.0	11.9	24.0
2002.....	0.8	1.2	1.5	2.0	2.5	3.4	4.3	7.4	13.2	24.4
2001.....	0.8	1.1	1.3	1.9	2.6	3.8	5.4	7.7	11.9	26.1
2000.....	0.9	1.2	1.5	1.7	2.6	3.4	5.3	7.7	11.9	25.6
1999.....	0.8	1.0	1.0	1.6	2.2	3.3	4.3	6.8	12.7	23.4

All races 3/, male

2004.....	1.0	1.4	1.8	2.2	3.0	3.3	4.6	7.9	12.2	19.6
2003.....	1.0	1.4	1.8	2.2	2.7	3.5	4.9	6.9	10.9	19.1
2002.....	0.8	1.3	1.6	2.3	2.6	3.6	4.2	7.0	13.3	20.5
2001.....	0.9	1.2	1.5	2.2	2.9	3.9	5.9	8.3	12.1	22.0
2000.....	1.1	1.5	1.7	1.9	2.7	3.9	5.6	8.0	12.4	22.2
1999.....	0.9	1.1	1.2	1.7	2.3	3.7	4.8	7.5	13.2	20.8

All races 3/, female

2004.....	0.8	1.1	1.1	1.6	2.0	2.9	4.0	6.5	11.0	25.0
2003.....	0.9	1.1	1.2	1.3	2.3	3.0	5.0	7.0	12.5	26.2
2002.....	0.8	1.1	1.4	1.7	2.5	3.2	4.4	7.7	13.1	26.2
2001.....	0.8	0.9	1.0	1.7	2.3	3.7	5.0	7.3	11.7	27.8
2000.....	0.7	1.0	1.3	1.5	2.6	3.0	5.0	7.5	11.6	26.9
1999.....	0.6	1.0	0.9	1.5	2.1	2.9	3.9	6.3	12.4	24.5

White, both sexes

2004.....	0.7	1.1	1.2	1.8	2.3	3.0	3.9	6.8	11.3	23.2
2003.....	0.8	1.1	1.4	1.6	2.2	3.0	4.7	6.5	11.6	23.6
2002.....	0.6	1.0	1.4	1.9	2.2	3.2	4.1	7.1	12.7	24.3
2001.....	0.7	0.9	1.1	1.8	2.3	3.6	5.1	7.2	11.8	26.0
2000.....	0.8	1.1	1.4	1.5	2.4	2.9	5.0	7.3	11.8	25.5
1999.....	0.6	0.9	0.9	1.6	2.1	2.9	4.1	6.5	12.4	23.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other disorders of circulatory system (I80-I99)

White, male

2004.....	0.8	1.4	1.5	2.1	2.7	3.1	4.3	7.6	11.9	19.7
2003.....	0.9	1.2	1.6	2.1	2.3	3.2	4.6	6.4	10.2	18.5
2002.....	0.6	1.1	1.5	2.1	2.3	3.3	4.0	6.7	12.4	19.9
2001.....	0.7	1.2	1.3	2.0	2.5	3.6	5.6	7.8	12.0	21.7
2000.....	1.0	1.3	1.6	1.7	2.4	3.4	5.5	7.8	12.3	22.0
1999.....	0.8	0.9	1.0	1.7	2.3	3.2	4.7	7.0	12.8	20.7

White, female

2004.....	0.6	0.9	0.9	1.4	1.8	2.8	3.6	6.2	11.0	24.8
2003.....	0.7	1.1	1.2	1.1	2.2	2.9	4.7	6.5	12.4	25.8
2002.....	0.7	1.0	1.2	1.6	2.1	3.0	4.2	7.4	12.9	26.3
2001.....	0.7	0.7	1.0	1.5	2.1	3.5	4.8	6.8	11.7	27.9
2000.....	0.5	0.9	1.1	1.4	2.4	2.5	4.7	7.0	11.5	26.9
1999.....	0.5	0.9	0.9	1.5	1.9	2.7	3.7	6.1	12.2	24.2

Black, both sexes

2004.....	2.5	2.4	3.6	3.6	5.0	4.9	8.9	12.9	14.6	28.8
2003.....	2.3	2.6	2.5	3.6	5.2	6.2	9.3	13.8	17.8	33.6
2002.....	1.8	2.4	2.9	4.2	5.9	6.4	7.2	12.9	22.2	30.0
2001.....	1.8	2.1	2.4	3.9	6.1	6.9	9.0	14.1	14.9	30.1
2000.....	1.8	2.5	3.2	3.9	5.1	8.7	9.1	14.6	14.1	30.0
1999.....	1.9	2.4	2.3	2.1	3.9	6.9	6.4	11.4	17.6	28.4

Black, male

2004.....	2.5	2.1	5.0	3.6	6.2	5.8	8.7	14.1	16.5	24.8
2003.....	1.8	3.4	3.4	3.6	7.3	7.6	10.2	15.5	20.7	32.3
2002.....	2.1	3.0	2.9	5.4	5.9	7.7	7.7	12.4	28.3	32.2
2001.....	2.2	1.7	3.1	4.7	8.5	7.9	10.4	16.3	*	29.5
2000.....	2.0	3.1	3.7	4.8	5.6	9.4	8.7	13.2	*	30.0
1999.....	2.6	2.7	3.0	*	*	9.3	6.8	14.7	20.9	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other disorders of circulatory system (I80-I99)

Black, female

2004.....	2.6	2.6	2.4	3.7	4.1	4.1	9.0	12.1	13.6	30.3
2003.....	2.7	1.9	1.8	3.6	3.6	5.1	8.6	12.8	16.2	34.0
2002.....	1.6	1.8	2.9	3.2	5.9	5.4	6.9	13.2	19.1	29.2
2001.....	1.5	2.4	*	3.2	4.2	6.2	8.1	12.8	14.7	30.3
2000.....	1.7	1.9	2.8	3.2	4.7	8.3	9.3	15.4	14.4	30.0
1999.....	*	2.1	*	*	4.2	5.0	6.1	9.4	16.0	30.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)

All races 3/, both sexes

2004.....	0.8	0.9	1.0	1.4	1.8	2.2	3.1	5.4	8.8	17.3
2003.....	0.8	1.0	1.1	1.3	1.9	2.6	3.7	5.2	9.0	18.1
2002.....	0.7	0.9	1.2	1.6	1.9	2.6	3.1	5.4	10.1	18.6
2001.....	0.6	0.8	0.9	1.4	2.0	2.8	4.1	5.9	8.9	19.6
2000.....	0.6	0.9	1.1	1.3	1.8	2.5	3.9	5.8	8.8	18.7
1999.....	0.6	0.7	0.7	1.1	1.4	2.2	2.8	4.3	8.0	14.6

All races 3/, male

2004.....	0.8	0.9	1.2	1.4	2.0	2.2	3.3	5.7	9.0	14.3
2003.....	0.7	1.0	1.2	1.5	2.0	2.7	3.5	4.8	7.8	13.8
2002.....	0.6	0.9	1.1	1.7	1.7	2.7	2.9	4.7	9.6	14.8
2001.....	0.6	0.8	1.0	1.5	2.1	2.7	4.4	5.9	8.3	15.2
2000.....	0.7	0.9	1.1	1.3	1.6	2.8	3.8	5.7	8.9	15.0
1999.....	0.6	0.6	0.6	1.0	1.2	2.3	3.1	4.4	7.9	11.2

All races 3/, female

2004.....	0.8	1.0	0.8	1.4	1.5	2.1	3.0	5.2	8.7	18.6
2003.....	0.8	1.0	1.0	1.1	1.7	2.5	3.9	5.4	9.8	20.1
2002.....	0.7	0.9	1.2	1.5	2.0	2.6	3.2	5.9	10.4	20.2
2001.....	0.7	0.8	0.8	1.3	1.9	2.9	3.8	5.9	9.2	21.4
2000.....	0.5	0.9	1.1	1.2	1.9	2.3	4.0	5.9	8.7	20.2
1999.....	0.6	0.7	0.7	1.1	1.5	2.0	2.5	4.2	8.1	15.9

White, both sexes

2004.....	0.6	0.8	0.8	1.2	1.5	2.1	2.8	5.1	8.7	17.1
2003.....	0.6	0.8	1.0	1.2	1.6	2.5	3.5	4.8	8.8	17.8
2002.....	0.5	0.8	1.0	1.5	1.6	2.5	2.9	5.2	9.6	18.4
2001.....	0.5	0.7	0.8	1.3	1.7	2.6	3.8	5.6	8.9	19.6
2000.....	0.6	0.8	1.0	1.1	1.6	2.1	3.7	5.5	8.8	18.8
1999.....	0.5	0.6	0.6	1.0	1.3	2.0	2.7	4.1	7.8	14.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)

White, male

2004.....	0.6	0.8	1.0	1.4	1.8	2.1	3.0	5.5	8.7	14.3
2003.....	0.7	0.8	1.1	1.5	1.7	2.5	3.3	4.3	7.5	13.3
2002.....	0.4	0.8	1.0	1.5	1.5	2.4	2.8	4.7	8.8	14.4
2001.....	0.5	0.7	0.9	1.4	1.7	2.6	4.1	5.7	8.3	15.0
2000.....	0.7	0.8	1.0	1.1	1.4	2.5	3.6	5.6	9.0	14.9
1999.....	0.5	0.6	0.5	1.0	1.2	2.0	3.0	4.2	7.7	11.4

White, female

2004.....	0.5	0.8	0.7	1.1	1.3	2.1	2.7	4.9	8.7	18.4
2003.....	0.6	0.9	1.0	0.8	1.6	2.4	3.7	5.1	9.7	19.8
2002.....	0.6	0.9	1.0	1.4	1.8	2.5	3.0	5.6	10.1	20.2
2001.....	0.6	0.6	0.8	1.2	1.7	2.6	3.5	5.6	9.2	21.5
2000.....	0.5	0.8	0.9	1.1	1.8	1.8	3.8	5.4	8.7	20.4
1999.....	0.5	0.6	0.7	1.1	1.4	1.9	2.4	4.1	7.8	15.5

Black, both sexes

2004.....	2.2	2.1	2.6	3.0	4.3	3.2	6.8	10.2	11.1	22.8
2003.....	2.1	2.2	1.9	2.9	4.5	4.8	7.4	10.6	13.7	26.3
2002.....	1.6	1.8	2.5	3.2	4.4	4.9	5.6	9.3	18.3	23.8
2001.....	1.5	1.8	1.7	2.9	5.3	5.3	7.8	10.4	11.6	23.3
2000.....	1.3	1.9	2.5	2.9	3.9	6.8	7.1	11.7	9.3	21.9
1999.....	1.5	1.6	1.5	1.6	2.4	4.8	4.5	7.1	12.7	18.8

Black, male

2004.....	2.0	1.7	3.3	2.6	5.0	*	7.4	10.0	*	19.9
2003.....	1.6	2.8	2.3	*	6.2	5.9	7.6	11.9	*	25.0
2002.....	1.7	2.1	2.4	4.2	4.4	5.7	*	*	22.2	22.5
2001.....	1.7	*	2.4	3.3	6.9	5.1	8.8	11.2	*	*
2000.....	*	2.2	2.4	*	*	6.5	7.0	*	*	*
1999.....	1.7	*	*	*	*	6.8	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)

Black, female

2004.....	2.4	2.4	1.9	3.3	3.8	*	6.4	10.2	10.6	24.0
2003.....	2.5	1.7	*	3.3	3.2	4.0	7.3	9.8	13.5	26.7
2002.....	1.6	1.6	2.6	*	4.5	4.3	5.8	11.0	16.3	24.2
2001.....	1.3	2.2	*	2.6	4.0	5.4	7.2	10.0	12.3	24.0
2000.....	*	1.7	2.6	2.7	3.9	7.1	7.1	13.4	10.6	22.3
1999.....	*	1.8	*	*	*	*	4.5	6.3	12.5	21.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified disorders of circulatory system (I83-I99)

All races 3/, both sexes

2004.....	0.1	0.3	0.4	0.5	0.7	0.9	1.2	1.7	2.7	6.0
2003.....	0.2	0.3	0.4	0.4	0.6	0.7	1.2	1.8	2.8	5.9
2002.....	0.1	0.3	0.3	0.4	0.7	0.7	1.2	2.1	3.1	5.9
2001.....	0.2	0.3	0.4	0.5	0.6	1.0	1.4	1.8	3.0	6.5
2000.....	0.2	0.3	0.4	0.4	0.9	0.9	1.4	1.9	3.1	6.8
1999.....	0.2	0.3	0.4	0.5	0.8	1.1	1.5	2.5	4.7	8.8

All races 3/, male

2004.....	0.2	0.5	0.6	0.8	0.9	1.1	1.3	2.3	3.3	5.4
2003.....	0.2	0.4	0.6	0.7	0.6	0.8	1.4	2.1	3.0	5.3
2002.....	0.2	0.4	0.5	0.6	0.8	0.9	1.3	2.3	3.7	5.7
2001.....	0.3	0.4	0.5	0.7	0.8	1.2	1.5	2.4	3.8	6.8
2000.....	0.3	0.5	0.6	0.6	1.1	1.0	1.8	2.4	3.5	7.3
1999.....	0.3	0.4	0.5	0.7	1.1	1.4	1.7	3.1	5.3	9.6

All races 3/, female

2004.....	*	*	0.2	0.3	0.5	0.8	1.0	1.3	2.3	6.4
2003.....	*	0.2	0.2	*	0.6	0.5	1.0	1.6	2.7	6.1
2002.....	*	*	0.2	0.3	0.5	0.6	1.2	1.9	2.7	5.9
2001.....	*	*	0.2	0.3	0.4	0.8	1.2	1.4	2.5	6.4
2000.....	*	*	0.2	*	0.7	0.8	1.0	1.6	2.9	6.7
1999.....	*	0.3	*	0.4	0.6	0.9	1.3	2.1	4.3	8.5

White, both sexes

2004.....	0.1	0.4	0.4	0.5	0.7	0.9	1.1	1.7	2.6	6.0
2003.....	0.1	0.3	0.4	0.4	0.6	0.6	1.2	1.7	2.7	5.8
2002.....	0.1	0.2	0.3	0.4	0.6	0.7	1.2	2.0	3.1	5.9
2001.....	0.2	0.3	0.3	0.5	0.6	0.9	1.3	1.6	3.0	6.5
2000.....	0.2	0.3	0.4	0.4	0.8	0.8	1.3	1.9	3.0	6.7
1999.....	0.2	0.3	0.3	0.5	0.8	1.0	1.5	2.4	4.6	8.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

X. Diseases of the respiratory system (J00-J98)

All races 3/, both sexes

2004.....	7.1	12.6	22.5	43.2	90.4	166.1	302.7	503.4	791.7	1,497.1
2003.....	7.3	13.6	23.5	44.5	94.9	176.0	311.6	524.2	827.7	1,625.4
2002.....	7.6	13.1	22.8	45.7	91.4	172.4	315.3	528.4	849.0	1,663.7
2001.....	7.4	12.7	23.1	46.9	93.4	178.6	319.0	526.3	833.1	1,671.8
2000.....	8.0	13.2	23.3	48.3	94.2	181.4	324.3	529.4	874.7	1,734.0
1999.....	7.4	12.9	22.4	49.4	97.1	185.1	330.2	537.8	873.1	1,733.0

All races 3/, male

2004.....	7.5	13.6	25.8	47.6	101.4	192.5	358.3	620.1	1,012.1	1,915.3
2003.....	7.5	15.2	26.2	49.5	106.5	202.6	371.4	646.6	1,052.6	2,057.2
2002.....	8.3	14.2	25.5	50.2	102.9	197.5	379.7	661.7	1,094.2	2,150.0
2001.....	8.0	14.3	25.7	51.9	104.0	204.9	385.2	662.7	1,080.0	2,187.2
2000.....	8.9	14.8	25.7	54.2	106.1	210.7	394.4	666.6	1,158.0	2,344.4
1999.....	7.9	14.6	24.5	54.1	111.1	220.6	411.8	691.1	1,170.2	2,356.4

All races 3/, female

2004.....	6.8	11.7	19.3	39.1	80.4	143.2	257.9	419.5	655.9	1,309.0
2003.....	7.2	12.1	20.8	39.8	84.3	153.0	263.6	437.0	690.2	1,434.5
2002.....	6.8	12.0	20.2	41.5	81.1	150.7	263.8	434.3	699.8	1,452.7
2001.....	6.8	11.0	20.6	42.2	83.9	155.9	266.4	430.8	685.2	1,456.2
2000.....	7.1	11.6	21.0	42.9	83.5	156.3	269.0	433.8	707.6	1,485.4
1999.....	7.0	11.4	20.5	45.0	84.3	154.8	266.4	431.1	699.3	1,481.7

White, both sexes

2004.....	6.5	11.5	21.4	42.9	92.2	172.9	316.6	521.6	817.9	1,541.7
2003.....	6.4	12.5	22.4	43.9	96.7	183.2	323.5	541.5	855.8	1,676.3
2002.....	6.7	12.1	21.6	45.3	93.0	179.4	326.6	545.8	873.3	1,713.3
2001.....	6.5	11.4	21.7	46.7	95.4	184.8	329.8	542.3	856.5	1,713.5
2000.....	7.0	11.6	21.8	47.7	95.6	185.4	333.7	544.9	898.9	1,783.5
1999.....	6.1	11.2	21.2	48.7	98.7	190.9	338.4	551.4	897.0	1,778.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

X. Diseases of the respiratory system (J00-J98)

White, male

2004.....	6.6	12.3	24.4	46.1	101.3	196.2	367.6	632.9	1,035.5	1,960.9
2003.....	6.4	14.0	24.9	48.0	106.0	207.1	377.9	655.2	1,072.5	2,106.7
2002.....	7.3	13.0	23.9	48.8	102.3	202.3	386.7	672.5	1,114.4	2,201.7
2001.....	7.1	13.1	23.6	51.2	103.7	208.2	391.6	672.0	1,098.6	2,230.5
2000.....	7.8	12.9	23.5	52.6	105.9	210.3	398.5	676.6	1,179.6	2,402.2
1999.....	6.3	12.6	22.5	52.3	110.8	223.6	415.4	699.7	1,188.8	2,409.4

White, female

2004.....	6.3	10.6	18.5	39.8	83.9	152.2	274.6	440.1	682.8	1,352.8
2003.....	6.3	11.0	19.9	40.0	88.1	162.1	279.0	459.2	722.2	1,485.6
2002.....	6.0	11.3	19.3	42.0	84.4	159.3	277.5	455.0	725.5	1,501.2
2001.....	5.9	9.6	19.8	42.3	87.8	164.3	279.7	450.5	710.2	1,497.0
2000.....	6.2	10.3	20.2	43.1	86.1	163.6	281.7	452.2	731.9	1,532.2
1999.....	5.9	9.9	19.9	45.3	87.5	162.3	277.1	447.2	724.8	1,525.0

Black, both sexes

2004.....	13.7	23.1	36.8	57.7	100.1	154.3	244.8	404.7	595.6	1,082.3
2003.....	15.1	24.7	37.8	60.3	102.7	160.3	264.1	427.8	606.9	1,135.5
2002.....	15.5	23.1	36.6	61.9	102.5	154.4	271.2	424.1	670.3	1,204.5
2001.....	14.9	24.9	39.3	60.7	99.0	162.5	275.3	430.7	647.1	1,258.0
2000.....	16.0	27.1	40.4	65.3	106.0	183.3	288.6	427.5	662.7	1,239.4
1999.....	17.6	28.0	38.2	68.0	105.5	173.1	302.0	456.5	655.6	1,298.0

Black, male

2004.....	15.2	26.0	44.3	74.2	131.2	215.1	344.8	585.5	873.1	1,501.1
2003.....	15.4	27.4	43.2	76.0	137.5	215.7	378.7	640.4	937.6	1,577.5
2002.....	17.9	26.6	43.7	77.9	135.8	204.2	378.9	634.7	986.6	1,685.3
2001.....	16.2	27.3	47.7	72.0	130.6	218.5	390.5	653.7	981.6	1,775.7
2000.....	18.5	32.3	48.7	82.0	134.0	254.7	419.2	634.6	980.4	1,808.3
1999.....	19.8	33.3	47.4	83.9	141.1	237.6	438.2	682.9	1,034.1	1,900.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

X. Diseases of the respiratory system (J00-J98)

Black, female

2004.....	12.3	20.6	30.5	44.1	75.9	109.2	177.0	297.6	449.4	916.3
2003.....	14.9	22.4	33.3	47.4	75.5	119.2	186.8	301.4	433.6	963.7
2002.....	13.4	20.1	30.5	48.8	76.5	117.5	198.9	298.3	505.7	1,020.3
2001.....	13.8	22.8	32.1	51.4	74.1	121.1	198.6	297.6	476.7	1,066.0
2000.....	13.7	22.6	33.4	51.7	84.0	130.6	201.8	302.8	502.7	1,027.6
1999.....	15.6	23.5	30.4	55.0	77.5	125.3	211.5	320.1	468.0	1,072.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute upper respiratory infections (J00-J06)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	*	*	0.8
2003.....	*	*	*	*	*	*	*	*	*	1.5
2002.....	*	*	*	*	*	*	*	*	*	1.1
2001.....	*	*	*	*	*	*	*	*	*	1.3
2000.....	*	*	*	*	*	*	*	*	0.4	1.7
1999.....	*	*	*	*	*	*	*	*	0.5	1.7

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	1.7
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	0.9
2003.....	*	*	*	*	*	*	*	*	*	1.4
2002.....	*	*	*	*	*	*	*	*	*	1.2
2001.....	*	*	*	*	*	*	*	*	*	1.6
2000.....	*	*	*	*	*	*	*	*	*	2.0
1999.....	*	*	*	*	*	*	*	*	*	2.0

White, both sexes

2004.....	*	*	*	*	*	*	*	*	*	0.8
2003.....	*	*	*	*	*	*	*	*	*	1.5
2002.....	*	*	*	*	*	*	*	*	*	1.1
2001.....	*	*	*	*	*	*	*	*	*	1.4
2000.....	*	*	*	*	*	*	*	*	*	1.8
1999.....	*	*	*	*	*	*	*	*	0.5	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	*	*	0.8
2003.....	*	*	*	*	*	*	*	*	*	1.4
2002.....	*	*	*	*	*	*	*	*	*	1.0
2001.....	*	*	*	*	*	*	*	*	*	1.1
2000.....	*	*	*	*	*	*	*	*	*	1.7
1999.....	*	*	*	*	*	*	*	*	0.5	1.6

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	1.6
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	0.9
2003.....	*	*	*	*	*	*	*	*	*	1.3
2002.....	*	*	*	*	*	*	*	*	*	1.2
2001.....	*	*	*	*	*	*	*	*	*	1.4
2000.....	*	*	*	*	*	*	*	*	*	1.9
1999.....	*	*	*	*	*	*	*	*	*	1.9

White, both sexes

2004.....	*	*	*	*	*	*	*	*	*	0.8
2003.....	*	*	*	*	*	*	*	*	*	1.4
2002.....	*	*	*	*	*	*	*	*	*	1.1
2001.....	*	*	*	*	*	*	*	*	*	1.2
2000.....	*	*	*	*	*	*	*	*	*	1.7
1999.....	*	*	*	*	*	*	*	*	0.5	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of the respiratory system (J10-J98)

All races 3/, both sexes

2004.....	7.1	12.6	22.5	43.2	90.3	166.1	302.6	503.2	791.4	1,496.3
2003.....	7.3	13.6	23.4	44.5	94.8	176.0	311.5	524.1	827.4	1,623.9
2002.....	7.5	13.1	22.7	45.6	91.4	172.3	315.1	528.3	848.8	1,662.6
2001.....	7.4	12.6	23.1	46.8	93.4	178.5	318.9	526.1	832.8	1,670.6
2000.....	8.0	13.2	23.2	48.3	94.2	181.3	324.1	529.1	874.3	1,732.2
1999.....	7.4	12.9	22.4	49.4	97.0	185.1	330.1	537.5	872.6	1,731.2

All races 3/, male

2004.....	7.4	13.5	25.8	47.5	101.4	192.5	358.2	620.0	1,011.8	1,914.6
2003.....	7.4	15.1	26.2	49.5	106.4	202.5	371.4	646.6	1,052.2	2,055.5
2002.....	8.3	14.2	25.4	50.1	102.9	197.4	379.5	661.6	1,094.0	2,149.3
2001.....	7.9	14.3	25.7	51.9	104.0	204.9	385.2	662.6	1,079.8	2,186.6
2000.....	8.9	14.8	25.6	54.2	106.1	210.6	394.3	666.5	1,157.7	2,343.3
1999.....	7.8	14.5	24.5	54.1	111.0	220.5	411.7	690.9	1,169.5	2,355.3

All races 3/, female

2004.....	6.8	11.7	19.3	39.1	80.3	143.1	257.9	419.3	655.6	1,308.1
2003.....	7.2	12.1	20.8	39.7	84.2	153.0	263.5	436.9	689.9	1,433.1
2002.....	6.8	11.9	20.1	41.4	81.0	150.6	263.7	434.1	699.7	1,451.5
2001.....	6.8	11.0	20.6	42.1	83.8	155.9	266.3	430.6	684.9	1,454.6
2000.....	7.1	11.6	20.9	42.8	83.5	156.2	268.8	433.5	707.2	1,483.4
1999.....	7.0	11.3	20.4	45.0	84.3	154.8	266.3	430.9	698.8	1,479.7

White, both sexes

2004.....	6.4	11.4	21.4	42.9	92.2	172.8	316.5	521.4	817.6	1,540.9
2003.....	6.3	12.5	22.4	43.9	96.7	183.2	323.5	541.4	855.4	1,674.8
2002.....	6.6	12.1	21.6	45.3	93.0	179.4	326.4	545.6	873.2	1,712.2
2001.....	6.5	11.3	21.7	46.6	95.4	184.8	329.7	542.2	856.2	1,712.1
2000.....	6.9	11.6	21.8	47.7	95.5	185.3	333.5	544.6	898.5	1,781.7
1999.....	6.1	11.2	21.2	48.7	98.6	190.8	338.3	551.1	896.5	1,776.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of the respiratory system (J10-J98)

White, male

2004.....	6.6	12.3	24.4	46.1	101.3	196.2	367.5	632.8	1,035.2	1,960.3
2003.....	6.4	14.0	24.9	48.0	105.9	207.0	377.9	655.2	1,072.2	2,105.2
2002.....	7.3	12.9	23.9	48.8	102.3	202.2	386.6	672.4	1,114.3	2,201.0
2001.....	7.0	13.1	23.6	51.2	103.6	208.1	391.6	671.9	1,098.3	2,229.8
2000.....	7.8	12.9	23.5	52.6	105.8	210.3	398.3	676.5	1,179.3	2,401.1
1999.....	6.2	12.5	22.5	52.3	110.8	223.5	415.3	699.4	1,188.2	2,408.6

White, female

2004.....	6.3	10.6	18.4	39.8	83.8	152.1	274.6	439.9	682.5	1,351.9
2003.....	6.3	11.0	19.9	40.0	88.1	162.1	279.0	459.1	721.8	1,484.2
2002.....	6.0	11.2	19.3	41.9	84.3	159.2	277.4	454.8	725.4	1,499.8
2001.....	5.9	9.6	19.8	42.3	87.8	164.2	279.6	450.3	709.9	1,495.4
2000.....	6.1	10.3	20.2	43.1	86.1	163.5	281.5	451.8	731.5	1,530.1
1999.....	5.9	9.9	19.8	45.3	87.4	162.3	277.0	447.0	724.3	1,522.9

Black, both sexes

2004.....	13.5	23.1	36.7	57.6	100.1	154.2	244.7	404.7	595.4	1,081.5
2003.....	15.0	24.6	37.7	60.3	102.4	160.1	264.1	427.8	606.9	1,134.0
2002.....	15.4	23.0	36.5	61.8	102.5	154.2	271.2	424.1	669.8	1,204.5
2001.....	14.9	24.8	39.2	60.6	98.8	162.5	275.2	430.5	647.1	1,257.1
2000.....	16.0	27.0	40.4	65.3	106.0	183.1	288.6	427.5	662.4	1,238.8
1999.....	17.6	28.0	38.1	67.9	105.3	173.1	301.6	456.5	655.3	1,296.4

Black, male

2004.....	15.0	26.0	44.2	74.0	131.2	215.1	344.4	585.5	872.3	1,500.1
2003.....	15.2	27.3	43.0	76.0	136.9	215.2	378.7	640.4	937.6	1,574.4
2002.....	17.8	26.6	43.5	77.9	135.8	203.9	378.9	634.7	985.8	1,685.3
2001.....	16.1	27.2	47.7	71.8	130.4	218.5	390.5	653.7	981.6	1,775.7
2000.....	18.5	32.3	48.7	82.0	134.0	254.2	419.2	634.6	979.5	1,806.0
1999.....	19.8	33.2	47.4	83.9	140.7	237.6	437.9	682.9	1,034.1	1,895.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of the respiratory system (J10-J98)

Black, female

2004.....	12.2	20.6	30.5	44.1	75.9	109.0	177.0	297.6	449.4	915.5
2003.....	14.8	22.3	33.1	47.4	75.5	119.2	186.8	301.4	433.6	962.9
2002.....	13.4	20.0	30.5	48.6	76.5	117.3	198.9	298.3	505.3	1,020.3
2001.....	13.8	22.8	32.0	51.4	73.9	121.1	198.4	297.3	476.7	1,064.7
2000.....	13.7	22.5	33.3	51.7	84.0	130.6	201.8	302.8	502.7	1,027.6
1999.....	15.6	23.5	30.3	54.8	77.5	125.3	211.1	320.1	467.6	1,072.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
 years years years years years years years years years and over

Influenza (J10-J11)

All races 3/, both sexes

2004.....	*	*	*	0.1	0.3	0.3	0.8	1.6	3.2	11.4
2003.....	*	0.1	0.2	0.3	0.3	0.7	1.3	2.7	4.7	17.0
2002.....	*	*	*	*	*	*	0.4	0.8	2.2	8.8
2001.....	*	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	0.1	0.1	0.3	0.5	0.8	1.4	2.5	6.1	20.2
1999.....	*	*	*	0.2	0.4	0.6	1.2	2.1	6.1	21.2

All races 3/, male

2004.....	*	*	*	*	*	*	1.1	1.8	3.3	11.1
2003.....	*	*	*	0.3	*	0.8	1.3	2.9	4.9	17.0
2002.....	*	*	*	*	*	*	*	0.9	2.6	8.9
2001.....	*	*	*	*	*	*	*	*	*	2.2
2000.....	*	*	*	0.3	0.5	1.0	1.6	2.7	6.3	20.7
1999.....	*	*	*	*	0.5	0.6	1.2	2.4	7.3	19.9

All races 3/, female

2004.....	*	*	*	*	*	*	0.6	1.4	3.1	11.5
2003.....	*	*	*	0.3	0.3	0.7	1.3	2.5	4.7	17.1
2002.....	*	*	*	*	*	*	*	0.7	2.0	8.7
2001.....	*	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	*	*	*	0.5	0.6	1.2	2.5	6.0	19.9
1999.....	*	*	*	*	*	0.6	1.1	1.9	5.4	21.7

White, both sexes

2004.....	*	*	*	0.2	0.3	0.3	0.9	1.5	3.4	12.3
2003.....	*	0.1	0.2	0.3	0.3	0.8	1.4	2.9	5.1	18.2
2002.....	*	*	*	*	*	*	0.4	0.9	2.4	9.4
2001.....	*	*	*	*	*	*	*	*	0.7	2.3
2000.....	*	*	0.1	0.3	0.5	0.8	1.4	2.7	6.5	21.4
1999.....	*	*	*	0.2	0.4	0.7	1.2	2.2	6.4	22.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonia (J12-J18)

All races 3/, both sexes

2004.....	2.5	3.6	5.6	8.2	13.9	23.5	46.4	93.1	195.6	571.2
2003.....	2.6	4.1	6.2	7.9	14.9	25.8	48.1	100.6	212.1	649.1
2002.....	2.6	4.1	5.4	8.4	14.6	25.1	50.5	105.7	225.1	687.8
2001.....	2.6	3.9	5.3	8.1	13.8	24.3	49.2	100.9	216.1	683.4
2000.....	2.9	3.8	5.5	8.6	15.1	25.8	51.2	104.1	234.7	723.9
1999.....	2.7	4.0	5.2	8.2	13.9	24.3	49.3	103.8	227.7	730.6

All races 3/, male

2004.....	2.7	4.2	7.0	9.7	17.0	29.4	57.6	116.5	249.1	666.1
2003.....	3.0	4.8	7.9	9.3	17.6	32.4	60.8	127.2	267.5	740.9
2002.....	3.0	4.9	6.7	10.1	17.8	30.9	63.4	133.0	285.0	796.9
2001.....	3.0	4.8	6.6	9.9	16.9	30.3	61.4	130.5	273.9	806.2
2000.....	3.7	4.6	6.8	10.6	18.4	31.9	64.1	131.5	302.6	871.3
1999.....	3.1	5.1	6.7	9.5	16.3	31.1	63.0	131.1	294.5	874.7

All races 3/, female

2004.....	2.3	2.9	4.1	6.7	11.1	18.4	37.3	76.4	162.6	528.5
2003.....	2.2	3.4	4.7	6.5	12.4	20.1	37.9	81.7	178.2	608.5
2002.....	2.2	3.4	4.3	6.8	11.8	20.0	40.2	86.5	188.6	640.5
2001.....	2.2	2.9	4.1	6.3	11.0	19.1	39.4	80.2	181.5	632.0
2000.....	2.1	3.0	4.2	6.7	12.0	20.5	41.0	85.1	194.6	663.9
1999.....	2.3	2.9	3.9	7.0	11.7	18.6	38.6	84.8	188.7	672.5

White, both sexes

2004.....	2.2	3.2	4.9	7.5	13.1	22.0	45.3	91.5	194.9	578.4
2003.....	2.3	3.6	5.6	7.1	13.7	24.4	46.5	99.5	213.3	661.1
2002.....	2.3	3.8	4.9	7.6	13.7	24.3	49.5	104.5	225.6	699.6
2001.....	2.1	3.3	4.8	7.4	12.9	23.2	48.0	98.3	215.8	692.7
2000.....	2.4	3.2	4.8	7.8	13.8	23.7	48.9	102.9	234.3	737.0
1999.....	2.1	3.3	4.6	7.4	12.6	23.4	47.6	103.3	229.2	741.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonia (J12-J18)

White, male

2004.....	2.3	3.8	6.2	9.0	15.7	26.7	55.6	112.9	247.3	673.4
2003.....	2.5	4.2	7.0	8.2	15.4	29.9	57.8	124.1	266.2	748.8
2002.....	2.7	4.3	6.0	9.0	16.3	29.6	61.9	130.4	283.7	806.2
2001.....	2.5	4.1	5.6	9.0	15.3	28.3	59.2	125.3	271.5	812.1
2000.....	3.1	3.8	5.7	9.2	16.7	28.6	60.6	129.1	299.7	884.4
1999.....	2.2	4.2	5.6	8.4	14.6	29.5	59.7	129.6	294.3	884.0

White, female

2004.....	2.1	2.6	3.7	6.1	10.6	17.8	36.9	75.9	162.3	535.6
2003.....	2.0	3.0	4.2	6.0	12.1	19.5	37.2	81.7	180.7	622.3
2002.....	1.9	3.2	3.9	6.2	11.3	19.7	39.4	86.0	190.0	653.3
2001.....	1.8	2.4	3.9	5.9	10.6	18.8	38.9	79.3	182.2	642.8
2000.....	1.8	2.7	3.9	6.4	11.1	19.4	39.6	84.5	195.4	677.1
1999.....	1.9	2.5	3.5	6.5	10.8	18.0	37.9	84.8	190.7	684.4

Black, both sexes

2004.....	4.9	6.9	11.5	15.5	23.5	39.5	61.9	115.6	214.1	502.5
2003.....	5.4	8.3	12.4	15.4	27.0	41.0	69.6	121.7	203.6	519.6
2002.....	5.2	7.5	10.1	17.2	26.1	35.0	68.6	123.2	230.8	572.8
2001.....	5.8	9.0	10.7	15.4	23.1	34.1	65.7	133.7	222.9	579.7
2000.....	6.0	8.1	12.1	17.7	26.9	45.4	76.0	123.5	243.1	591.5
1999.....	6.8	9.7	11.6	16.0	26.4	36.5	73.9	117.8	217.0	629.1

Black, male

2004.....	5.9	8.4	15.8	19.1	32.9	61.1	86.4	165.6	297.0	597.1
2003.....	7.0	10.0	16.5	20.8	39.5	60.8	100.2	176.6	283.5	638.1
2002.....	6.2	9.9	12.9	23.5	35.6	47.4	89.8	173.2	320.2	687.2
2001.....	7.0	11.4	15.7	20.6	33.2	49.6	94.9	203.5	309.0	712.6
2000.....	8.1	11.0	17.0	25.5	35.0	64.3	106.6	170.6	343.1	742.7
1999.....	8.9	13.3	17.1	21.7	33.9	51.4	108.2	159.6	307.8	800.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonia (J12-J18)

Black, female

2004.....	4.1	5.7	7.9	12.6	16.2	23.6	45.3	86.0	170.5	465.0
2003.....	4.1	6.7	8.9	11.0	17.1	26.3	49.0	89.0	161.7	473.5
2002.....	4.3	5.5	7.7	12.0	18.6	25.9	54.4	93.3	184.2	529.0
2001.....	4.7	6.9	6.5	11.2	15.2	22.7	46.2	92.1	179.1	530.5
2000.....	4.2	5.6	7.9	11.2	20.4	31.5	55.7	95.1	192.8	535.3
1999.....	5.0	6.6	7.0	11.4	20.5	25.4	51.1	92.7	172.0	565.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Viral pneumonia, not elsewhere classified (J12)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	0.3	0.6	1.3
2003.....	*	*	*	*	*	*	*	0.4	0.6	2.1
2002.....	*	*	*	*	*	*	0.3	0.4	0.7	1.7
2001.....	*	*	*	*	*	*	*	0.3	0.7	1.4
2000.....	*	*	*	*	*	0.2	0.3	0.6	1.0	2.9
1999.....	*	*	*	*	*	0.2	0.2	0.5	1.2	2.4

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	2.2
2002.....	*	*	*	*	*	*	*	*	1.0	1.9
2001.....	*	*	*	*	*	*	*	*	1.3	1.9
2000.....	*	*	*	*	*	*	*	0.7	*	2.4
1999.....	*	*	*	*	*	*	*	0.9	1.7	1.8

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	1.4
2003.....	*	*	*	*	*	*	*	*	*	2.0
2002.....	*	*	*	*	*	*	*	*	*	1.6
2001.....	*	*	*	*	*	*	*	*	*	1.2
2000.....	*	*	*	*	*	*	*	0.5	1.0	3.0
1999.....	*	*	*	*	*	*	*	*	0.9	2.6

White, both sexes

2004.....	*	*	*	*	*	*	*	0.3	0.7	1.3
2003.....	*	*	*	*	*	*	*	0.4	0.6	2.2
2002.....	*	*	*	*	*	*	0.3	0.4	0.8	1.8
2001.....	*	*	*	*	*	*	*	0.3	0.8	1.5
2000.....	*	*	*	*	*	0.3	0.3	0.6	1.0	3.0
1999.....	*	*	*	*	*	0.2	0.3	0.5	1.3	2.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Bacterial pneumonia (J13-J15)

All races 3/, both sexes

2004.....	0.3	0.5	0.8	1.2	1.7	2.6	4.3	7.9	13.9	30.1
2003.....	0.4	0.5	1.0	1.1	1.8	2.9	4.7	8.2	15.2	34.9
2002.....	0.4	0.6	0.7	1.2	1.6	2.6	4.7	8.3	15.5	35.7
2001.....	0.3	0.5	0.7	1.4	1.6	2.7	4.4	8.4	14.8	34.8
2000.....	0.5	0.6	0.8	1.3	1.9	2.7	4.9	8.1	15.3	35.7
1999.....	0.5	0.5	0.6	1.1	1.5	2.4	4.3	7.2	13.6	30.1

All races 3/, male

2004.....	0.3	0.6	1.0	1.2	1.9	3.3	5.3	10.2	19.9	41.1
2003.....	0.5	0.6	1.3	1.3	2.2	3.8	6.3	10.6	20.5	45.7
2002.....	0.4	0.6	0.8	1.4	1.8	3.0	6.1	11.3	20.9	46.8
2001.....	0.5	0.6	0.9	1.6	2.0	3.6	5.6	11.7	20.6	46.8
2000.....	0.6	0.6	0.9	1.5	2.3	3.3	6.4	11.9	21.7	52.3
1999.....	0.5	0.6	0.8	1.2	1.6	3.0	5.5	9.0	18.4	40.4

All races 3/, female

2004.....	0.4	0.4	0.6	1.1	1.6	2.0	3.4	6.2	10.3	25.2
2003.....	0.3	0.5	0.7	0.9	1.4	2.2	3.4	6.5	11.9	30.2
2002.....	0.4	0.5	0.6	0.9	1.4	2.3	3.6	6.2	12.2	31.0
2001.....	0.2	0.5	0.6	1.1	1.3	1.9	3.4	6.2	11.3	29.8
2000.....	0.4	0.5	0.6	1.1	1.6	2.1	3.7	5.4	11.6	29.0
1999.....	0.5	0.4	0.4	0.9	1.5	1.8	3.3	5.9	10.8	26.0

White, both sexes

2004.....	0.3	0.5	0.7	1.1	1.6	2.5	4.2	7.5	13.8	29.9
2003.....	0.4	0.5	0.9	1.0	1.6	2.9	4.6	8.0	15.3	34.9
2002.....	0.4	0.5	0.7	1.0	1.5	2.6	4.7	8.3	15.3	36.3
2001.....	0.3	0.5	0.7	1.4	1.5	2.6	4.2	8.1	14.8	35.0
2000.....	0.4	0.5	0.7	1.2	1.9	2.5	4.7	7.9	15.0	35.6
1999.....	0.4	0.5	0.5	1.0	1.4	2.3	4.1	7.1	13.6	30.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Bacterial pneumonia (J13-J15)

White, male

2004.....	0.3	0.6	0.9	1.0	1.7	2.9	5.0	9.7	19.7	41.7
2003.....	0.4	0.5	1.2	1.1	1.8	3.7	6.2	10.3	20.8	45.9
2002.....	0.4	0.5	0.7	1.2	1.7	2.9	5.9	11.1	20.1	47.9
2001.....	0.4	0.5	0.8	1.6	1.8	3.5	5.3	10.9	20.5	46.6
2000.....	0.6	0.6	0.9	1.4	2.1	3.0	6.1	11.4	21.7	51.1
1999.....	0.4	0.5	0.7	1.1	1.5	2.9	5.2	8.9	18.1	40.1

White, female

2004.....	0.3	0.4	0.6	1.1	1.6	2.0	3.5	5.9	10.2	24.7
2003.....	0.3	0.5	0.7	0.9	1.4	2.2	3.4	6.4	11.8	30.1
2002.....	0.3	0.5	0.6	0.8	1.4	2.3	3.7	6.3	12.3	31.2
2001.....	*	0.5	0.7	1.1	1.2	1.8	3.4	6.1	11.3	30.2
2000.....	0.3	0.4	0.6	1.0	1.6	2.0	3.6	5.5	11.0	29.2
1999.....	0.4	0.4	0.4	0.9	1.3	1.8	3.2	5.9	11.0	26.3

Black, both sexes

2004.....	*	0.9	1.7	2.1	2.8	4.3	5.4	11.8	15.6	35.0
2003.....	0.8	1.0	1.6	2.2	3.4	3.4	4.9	10.7	12.9	35.0
2002.....	0.8	1.1	1.0	2.3	2.2	2.9	5.4	8.4	16.0	28.5
2001.....	0.7	*	1.2	1.6	3.0	3.3	6.1	12.3	14.1	32.5
2000.....	*	1.0	1.3	2.0	2.8	4.6	7.1	8.9	17.5	34.7
1999.....	1.1	1.0	*	2.1	2.6	2.8	6.9	8.2	12.1	25.2

Black, male

2004.....	*	*	2.4	3.3	3.9	7.0	8.1	16.4	22.2	41.7
2003.....	*	*	2.2	3.2	5.4	*	7.6	16.0	*	41.7
2002.....	*	*	*	3.4	*	*	7.7	13.3	26.0	32.2
2001.....	*	*	*	*	*	5.1	10.1	21.4	18.5	49.9
2000.....	*	*	*	*	4.1	5.5	10.7	16.1	21.1	60.0
1999.....	*	*	*	*	*	*	9.8	10.9	20.9	38.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Bacterial pneumonia (J13-J15)

Black, female

2004.....	*	*	*	*	*	*	*	9.2	12.1	32.3
2003.....	*	*	*	*	*	*	*	7.6	12.4	32.4
2002.....	*	*	*	*	*	*	*	5.5	10.7	27.1
2001.....	*	*	*	*	*	*	*	6.9	11.9	26.1
2000.....	*	*	*	*	*	4.0	4.7	*	15.7	25.3
1999.....	*	*	*	*	*	*	5.0	6.6	*	20.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonia due to other or unspecified organisms (J16,J18)

All races 3/, both sexes

2004.....	2.1	3.0	4.7	7.0	12.1	20.8	42.0	85.0	181.0	539.8
2003.....	2.2	3.5	5.2	6.7	13.0	22.8	43.2	92.0	196.3	612.1
2002.....	2.2	3.5	4.7	7.2	13.0	22.4	45.5	97.0	208.8	650.3
2001.....	2.2	3.3	4.6	6.7	12.1	21.5	44.7	92.1	200.6	647.1
2000.....	2.4	3.2	4.7	7.2	13.0	22.9	46.0	95.5	218.3	685.3
1999.....	2.2	3.5	4.6	7.0	12.2	21.7	44.8	96.1	212.9	698.1

All races 3/, male

2004.....	2.3	3.6	6.0	8.5	15.1	26.0	52.2	106.1	228.3	623.9
2003.....	2.5	4.2	6.5	8.0	15.4	28.5	54.2	116.2	246.2	693.0
2002.....	2.6	4.2	5.8	8.6	15.9	27.8	56.9	121.3	263.1	748.2
2001.....	2.5	4.2	5.7	8.2	14.9	26.5	55.7	118.3	252.1	757.4
2000.....	3.1	3.9	5.8	9.0	16.0	28.3	57.5	118.9	279.9	816.5
1999.....	2.6	4.5	5.8	8.1	14.5	27.9	57.2	121.2	274.4	832.5

All races 3/, female

2004.....	1.9	2.5	3.5	5.6	9.5	16.2	33.8	69.8	151.8	502.0
2003.....	1.9	2.9	3.9	5.6	10.8	17.8	34.4	74.8	165.9	576.3
2002.....	1.7	2.8	3.6	5.8	10.3	17.7	36.4	79.8	175.8	607.9
2001.....	1.9	2.4	3.5	5.2	9.6	17.1	36.0	73.8	169.8	601.0
2000.....	1.7	2.5	3.6	5.6	10.3	18.2	37.0	79.2	182.0	631.9
1999.....	1.8	2.5	3.4	6.0	10.1	16.5	35.0	78.7	177.0	643.9

White, both sexes

2004.....	1.9	2.7	4.2	6.4	11.4	19.4	41.1	83.8	180.3	547.1
2003.....	1.9	3.1	4.6	6.1	12.0	21.4	41.6	91.1	197.4	624.0
2002.....	1.9	3.2	4.2	6.5	12.1	21.7	44.5	95.8	209.5	661.6
2001.....	1.8	2.7	4.0	6.0	11.3	20.5	43.6	89.9	200.3	656.2
2000.....	2.0	2.7	4.0	6.5	11.8	21.0	43.9	94.4	218.3	698.3
1999.....	1.7	2.9	4.0	6.4	11.1	20.8	43.2	95.8	214.3	708.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonia due to other or unspecified organisms (J16,J18)

White, male

2004.....	2.0	3.2	5.3	7.9	14.0	23.7	50.5	103.1	226.7	630.7
2003.....	2.1	3.7	5.8	7.1	13.6	26.1	51.4	113.4	244.6	700.5
2002.....	2.3	3.8	5.2	7.7	14.5	26.6	55.6	118.8	262.5	756.4
2001.....	2.1	3.5	4.8	7.3	13.5	24.6	53.8	113.8	249.6	763.5
2000.....	2.5	3.3	4.8	7.7	14.5	25.3	54.2	117.0	276.9	830.6
1999.....	1.9	3.7	4.9	7.2	12.9	26.4	54.2	119.9	274.5	841.9

White, female

2004.....	1.7	2.2	3.1	5.0	9.0	15.7	33.3	69.6	151.5	509.4
2003.....	1.6	2.5	3.5	5.1	10.5	17.3	33.6	75.0	168.3	590.1
2002.....	1.6	2.7	3.2	5.3	9.8	17.3	35.5	79.3	177.0	620.4
2001.....	1.6	2.0	3.2	4.7	9.3	16.9	35.4	73.0	170.5	611.3
2000.....	1.5	2.2	3.2	5.3	9.4	17.2	35.7	78.5	183.3	644.6
1999.....	1.5	2.1	3.1	5.6	9.4	15.9	34.5	78.8	178.7	655.4

Black, both sexes

2004.....	4.3	6.0	9.8	13.3	20.6	35.1	56.5	103.8	198.5	467.3
2003.....	4.6	7.2	10.7	13.2	23.4	37.7	64.7	110.9	190.1	484.0
2002.....	4.4	6.4	9.1	14.9	23.9	32.2	63.1	114.6	214.6	543.4
2001.....	5.1	8.2	9.5	13.8	20.1	30.6	59.5	121.4	208.6	546.9
2000.....	5.4	7.1	10.7	15.6	23.8	40.7	68.7	114.6	225.3	556.5
1999.....	5.7	8.6	10.6	13.9	23.8	33.5	67.0	109.1	204.3	602.6

Black, male

2004.....	5.3	7.5	13.4	15.7	29.0	54.0	78.3	149.2	274.8	554.3
2003.....	5.6	8.8	14.2	17.5	33.7	56.4	92.6	160.7	268.0	595.3
2002.....	5.5	8.6	11.7	20.1	32.7	44.4	82.1	159.4	293.5	652.9
2001.....	5.9	10.1	13.7	18.6	29.3	44.2	84.5	182.1	290.5	661.5
2000.....	7.5	10.0	15.3	22.6	30.6	58.8	95.6	154.5	322.0	682.7
1999.....	7.7	11.8	15.4	18.8	31.6	47.5	98.4	147.2	285.1	760.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonia due to other or unspecified organisms (J16,J18)

Black, female

2004.....	3.4	4.8	6.8	11.4	14.1	21.0	41.7	76.8	158.4	432.8
2003.....	3.7	5.8	7.7	9.6	15.3	23.8	45.9	81.4	149.3	440.7
2002.....	3.4	4.5	6.9	10.6	17.0	23.1	50.4	87.8	173.5	501.5
2001.....	4.3	6.4	5.9	9.9	12.8	20.5	42.8	85.1	166.8	504.4
2000.....	3.5	4.6	6.8	9.9	18.5	27.3	50.8	90.6	176.6	509.5
1999.....	4.0	5.9	6.6	9.8	17.6	23.1	46.1	86.1	164.2	543.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other acute lower respiratory infections (J20-J22)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	0.3	0.6	2.6
2003.....	*	*	*	*	*	*	*	0.5	0.8	3.5
2002.....	*	*	*	*	*	*	0.3	0.3	0.8	3.3
2001.....	*	*	*	*	*	*	*	0.3	0.7	3.3
2000.....	*	*	*	*	*	*	0.3	0.6	0.8	4.5
1999.....	*	*	*	*	*	*	0.2	0.6	1.3	6.0

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	2.5
2003.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	2.8
2001.....	*	*	*	*	*	*	*	*	*	2.3
2000.....	*	*	*	*	*	*	*	0.7	*	3.8
1999.....	*	*	*	*	*	*	*	0.7	1.6	4.0

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	0.7	2.7
2003.....	*	*	*	*	*	*	*	*	0.8	3.9
2002.....	*	*	*	*	*	*	*	*	0.9	3.5
2001.....	*	*	*	*	*	*	*	*	0.7	3.7
2000.....	*	*	*	*	*	*	*	0.5	0.9	4.8
1999.....	*	*	*	*	*	*	*	0.5	1.1	6.8

White, both sexes

2004.....	*	*	*	*	*	*	*	*	0.7	2.8
2003.....	*	*	*	*	*	*	*	0.5	0.9	3.7
2002.....	*	*	*	*	*	*	*	*	0.8	3.5
2001.....	*	*	*	*	*	*	*	0.3	0.7	3.5
2000.....	*	*	*	*	*	*	0.3	0.6	0.9	4.7
1999.....	*	*	*	*	*	*	*	0.6	1.3	6.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute bronchitis and bronchiolitis (J20-J21)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	*	*	1.4
2003.....	*	*	*	*	*	*	*	*	0.5	2.1
2002.....	*	*	*	*	*	*	*	*	0.5	2.0
2001.....	*	*	*	*	*	*	*	*	0.4	2.1
2000.....	*	*	*	*	*	*	*	0.4	0.5	2.6
1999.....	*	*	*	*	*	*	*	0.4	0.8	3.6

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	1.8
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	2.4
1999.....	*	*	*	*	*	*	*	*	*	2.6

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	1.3
2003.....	*	*	*	*	*	*	*	*	*	2.5
2002.....	*	*	*	*	*	*	*	*	*	2.2
2001.....	*	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	*	*	*	*	*	*	2.7
1999.....	*	*	*	*	*	*	*	*	0.7	4.0

White, both sexes

2004.....	*	*	*	*	*	*	*	*	*	1.5
2003.....	*	*	*	*	*	*	*	*	0.5	2.3
2002.....	*	*	*	*	*	*	*	*	0.5	2.1
2001.....	*	*	*	*	*	*	*	*	0.5	2.1
2000.....	*	*	*	*	*	*	*	0.4	0.5	2.7
1999.....	*	*	*	*	*	*	*	0.4	0.9	3.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)										
All races 3/, both sexes										
2004.....	*	*	*	*	*	*	*	*	*	1.2
2003.....	*	*	*	*	*	*	*	*	*	1.3
2002.....	*	*	*	*	*	*	*	*	0.4	1.3
2001.....	*	*	*	*	*	*	*	*	*	1.2
2000.....	*	*	*	*	*	*	*	*	*	1.9
1999.....	*	*	*	*	*	*	*	*	0.5	2.4
All races 3/, male										
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2004.....	*	*	*	*	*	*	*	*	*	1.4
2003.....	*	*	*	*	*	*	*	*	*	1.4
2002.....	*	*	*	*	*	*	*	*	*	1.3
2001.....	*	*	*	*	*	*	*	*	*	1.3
2000.....	*	*	*	*	*	*	*	*	*	2.1
1999.....	*	*	*	*	*	*	*	*	*	2.8
White, both sexes										
2004.....	*	*	*	*	*	*	*	*	*	1.3
2003.....	*	*	*	*	*	*	*	*	*	1.4
2002.....	*	*	*	*	*	*	*	*	*	1.4
2001.....	*	*	*	*	*	*	*	*	*	1.3
2000.....	*	*	*	*	*	*	*	*	*	2.0
1999.....	*	*	*	*	*	*	*	*	0.5	2.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of upper respiratory tract (J30-J39)

All races 3/, both sexes

2004.....	*	0.1	0.1	0.1	0.2	0.2	0.3	0.7	1.1	2.1
2003.....	*	*	*	0.2	0.3	0.3	0.4	0.7	1.0	2.3
2002.....	*	0.1	*	0.2	0.2	0.3	0.4	0.6	0.7	2.1
2001.....	*	0.1	0.1	*	0.2	0.3	0.4	0.6	1.3	2.4
2000.....	*	0.1	0.1	0.1	0.2	0.3	0.5	0.7	1.2	2.1
1999.....	*	*	*	*	0.3	0.4	0.4	0.9	1.1	2.4

All races 3/, male

2004.....	*	*	*	*	*	*	*	0.8	1.5	3.1
2003.....	*	*	*	*	*	0.4	*	0.9	1.5	3.0
2002.....	*	*	*	*	*	*	*	0.7	*	2.5
2001.....	*	*	*	*	*	*	*	0.7	1.9	3.4
2000.....	*	*	*	*	*	*	0.6	0.7	1.9	3.1
1999.....	*	*	*	*	*	0.5	0.5	1.1	1.3	3.5

All races 3/, female

2004.....	*	*	*	*	*	*	*	0.6	0.8	1.6
2003.....	*	*	*	*	*	*	*	0.5	0.7	2.0
2002.....	*	*	*	*	*	*	*	0.5	0.6	1.9
2001.....	*	*	*	*	*	*	*	0.6	1.0	1.9
2000.....	*	*	*	*	*	*	0.5	0.6	0.8	1.7
1999.....	*	*	*	*	*	*	*	0.7	1.0	2.0

White, both sexes

2004.....	*	0.1	*	*	*	*	*	0.6	1.1	2.1
2003.....	*	*	*	*	0.3	0.3	0.4	0.6	1.1	2.3
2002.....	*	*	*	0.2	*	0.2	0.4	0.6	0.6	2.2
2001.....	*	*	*	*	0.2	0.3	0.3	0.6	1.4	2.3
2000.....	*	*	0.1	*	0.2	0.3	0.5	0.6	1.3	2.2
1999.....	*	*	*	*	0.3	0.3	0.4	0.8	1.1	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic lower respiratory diseases (J40-J47)

All races 3/, both sexes

2004.....	2.6	5.6	11.7	25.5	60.0	113.7	200.6	315.7	434.7	601.7
2003.....	2.8	5.9	11.9	27.5	63.9	121.1	210.8	330.0	455.9	635.1
2002.....	2.9	5.8	12.0	28.2	60.8	119.0	211.5	330.1	465.7	637.6
2001.....	2.7	5.4	11.9	29.4	62.9	124.2	215.4	329.2	452.9	644.7
2000.....	2.8	5.8	11.8	29.6	62.5	124.1	218.2	327.8	473.4	648.6
1999.....	2.6	5.5	12.0	31.9	66.6	130.4	227.6	340.2	484.5	646.0

All races 3/, male

2004.....	2.6	5.6	12.5	26.9	64.6	126.1	228.2	378.8	543.4	811.1
2003.....	2.5	6.3	12.3	29.4	69.2	132.6	244.0	395.8	569.1	865.9
2002.....	3.0	5.9	12.8	29.5	65.9	131.5	246.0	405.6	595.9	894.8
2001.....	2.5	5.8	12.3	31.4	67.0	137.1	252.9	406.7	583.5	916.9
2000.....	2.6	6.2	12.1	31.8	68.0	139.7	257.8	404.2	628.5	967.9
1999.....	2.4	5.7	12.1	34.3	74.6	151.0	278.3	432.4	648.3	996.7

All races 3/, female

2004.....	2.7	5.5	10.9	24.2	55.8	102.9	178.4	270.4	367.7	507.4
2003.....	3.1	5.4	11.5	25.7	59.0	111.2	184.2	283.1	386.7	533.0
2002.....	2.8	5.6	11.2	26.9	56.1	108.3	183.8	276.8	386.5	526.0
2001.....	2.9	5.1	11.5	27.5	59.1	113.1	185.6	275.0	374.6	530.8
2000.....	2.9	5.5	11.5	27.5	57.5	110.7	187.0	274.6	381.9	518.5
1999.....	2.9	5.3	11.9	29.7	59.3	112.8	188.0	276.1	388.7	504.6

White, both sexes

2004.....	2.4	5.2	11.6	26.1	63.2	121.7	214.3	333.3	457.6	630.1
2003.....	2.5	5.6	11.7	28.0	67.3	129.5	223.2	347.2	478.9	664.5
2002.....	2.5	5.5	11.7	28.9	63.8	126.8	223.3	346.9	486.5	666.2
2001.....	2.4	5.1	11.7	30.2	66.2	131.3	227.1	346.1	473.5	671.8
2000.....	2.3	5.3	11.4	30.1	65.5	130.5	229.6	343.1	495.5	675.6
1999.....	2.2	4.9	11.8	32.6	69.6	137.2	237.9	353.4	504.9	672.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic lower respiratory diseases (J40-J47)

White, male

2004.....	2.4	5.3	12.3	26.6	66.5	132.4	238.5	392.8	563.4	837.7
2003.....	2.2	6.1	12.0	29.4	71.5	139.3	252.4	406.9	585.6	894.4
2002.....	2.5	5.6	12.4	29.6	67.5	137.1	254.3	417.4	612.9	923.4
2001.....	2.2	5.6	11.9	31.8	69.1	142.0	260.8	418.9	600.0	943.6
2000.....	2.2	5.7	11.5	31.8	69.8	143.1	265.8	415.5	648.5	997.4
1999.....	2.0	5.0	11.6	34.5	76.3	155.7	285.6	441.3	664.4	1,029.4

White, female

2004.....	2.5	5.1	11.0	25.6	60.1	112.2	194.4	289.7	391.9	536.5
2003.....	2.8	5.1	11.4	26.7	63.5	120.8	199.3	304.0	413.2	562.8
2002.....	2.5	5.4	11.0	28.3	60.4	117.6	198.0	296.4	408.9	554.5
2001.....	2.6	4.6	11.6	28.6	63.5	122.0	199.7	294.5	397.1	557.9
2000.....	2.4	4.9	11.3	28.4	61.5	119.4	200.5	292.1	404.4	544.8
1999.....	2.5	4.7	12.0	30.8	63.5	121.1	200.0	291.6	410.7	529.7

Black, both sexes

2004.....	4.8	9.6	16.3	27.9	52.1	81.7	128.1	204.6	249.3	333.9
2003.....	5.7	9.1	16.7	31.3	52.7	86.5	146.7	211.9	271.9	353.0
2002.....	6.5	9.1	17.4	30.9	53.4	86.1	148.1	215.9	298.5	368.2
2001.....	5.1	9.1	16.6	31.6	52.2	94.3	149.6	206.9	276.5	375.8
2000.....	6.1	11.0	18.2	33.0	57.4	97.3	151.2	211.6	271.3	373.8
1999.....	6.1	11.2	17.2	36.1	57.6	100.0	163.3	242.4	295.3	376.2

Black, male

2004.....	4.9	9.9	18.9	37.7	67.6	109.0	182.3	300.7	392.4	566.3
2003.....	5.1	9.8	17.7	38.0	67.7	112.0	212.5	326.8	471.8	601.6
2002.....	6.9	9.9	19.9	38.1	71.1	116.2	214.0	340.8	477.6	645.4
2001.....	5.1	8.9	18.8	37.1	66.3	127.1	222.2	329.9	468.7	671.7
2000.....	5.9	11.4	19.6	39.5	73.5	138.4	226.3	326.5	448.5	693.1
1999.....	6.2	11.9	19.7	43.2	79.3	139.4	243.5	387.0	521.4	701.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic lower respiratory diseases (J40-J47)

Black, female

2004.....	4.8	9.3	14.2	19.9	40.1	61.3	91.4	147.7	173.9	241.8
2003.....	6.2	8.6	15.8	25.7	41.1	67.6	102.4	143.5	167.1	256.4
2002.....	6.1	8.5	15.3	25.1	39.6	63.8	104.0	141.3	205.3	262.0
2001.....	5.1	9.2	14.7	27.1	41.1	70.0	101.3	133.4	178.7	266.1
2000.....	6.2	10.6	16.9	27.7	44.6	67.0	101.2	142.4	182.1	255.0
1999.....	6.1	10.7	15.0	30.2	40.6	70.8	110.0	155.3	183.2	254.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

All races 3/, both sexes

2004.....	2.6	5.5	11.6	25.4	59.7	113.3	199.5	313.2	430.7	595.3
2003.....	2.8	5.8	11.8	27.3	63.6	120.6	209.8	327.7	451.8	629.1
2002.....	2.9	5.7	11.9	28.0	60.4	118.4	210.3	327.6	461.0	631.8
2001.....	2.7	5.4	11.8	29.3	62.5	123.4	214.2	326.9	449.0	637.7
2000.....	2.7	5.8	11.8	29.3	62.2	123.4	216.9	325.4	468.8	642.4
1999.....	2.6	5.4	11.9	31.7	66.2	129.6	226.5	337.6	479.9	639.8

All races 3/, male

2004.....	2.6	5.6	12.4	26.8	64.3	125.8	227.5	376.9	541.1	806.6
2003.....	2.5	6.3	12.2	29.3	68.9	132.4	243.3	394.2	565.9	861.8
2002.....	2.9	5.9	12.7	29.4	65.6	131.1	245.1	404.0	592.8	890.1
2001.....	2.5	5.7	12.3	31.3	66.6	136.5	252.0	405.0	581.3	912.1
2000.....	2.6	6.2	12.1	31.6	67.7	139.3	256.8	402.6	625.4	963.6
1999.....	2.4	5.6	12.1	34.2	74.3	150.6	277.3	429.9	644.2	990.1

All races 3/, female

2004.....	2.7	5.5	10.8	24.0	55.5	102.3	176.9	267.5	362.8	500.3
2003.....	3.1	5.3	11.4	25.4	58.7	110.4	182.9	280.4	382.0	526.2
2002.....	2.8	5.6	11.1	26.7	55.6	107.4	182.4	273.7	380.9	519.7
2001.....	2.8	5.1	11.4	27.4	58.7	112.2	184.2	272.2	369.7	522.9
2000.....	2.8	5.4	11.4	27.2	57.2	109.9	185.6	271.7	376.5	511.6
1999.....	2.8	5.2	11.7	29.4	58.9	111.6	186.8	273.5	383.7	498.6

White, both sexes

2004.....	2.4	5.2	11.5	26.0	62.9	121.2	213.1	330.6	453.3	623.4
2003.....	2.5	5.6	11.7	27.9	67.0	129.0	222.1	344.7	474.6	658.1
2002.....	2.5	5.5	11.6	28.8	63.3	126.2	222.0	344.2	481.6	660.2
2001.....	2.4	5.1	11.6	30.1	65.8	130.6	225.8	343.6	469.4	664.3
2000.....	2.3	5.2	11.4	29.8	65.2	129.8	228.4	340.6	490.6	669.0
1999.....	2.2	4.8	11.7	32.3	69.2	136.4	236.8	350.8	500.1	666.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
White, male										
2004.....	2.4	5.2	12.1	26.5	66.2	132.0	237.7	390.9	560.9	832.8
2003.....	2.2	6.1	12.0	29.4	71.2	139.1	251.7	405.2	582.3	889.8
2002.....	2.5	5.5	12.3	29.5	67.1	136.8	253.5	415.7	609.7	918.7
2001.....	2.2	5.5	11.8	31.7	68.6	141.6	260.0	417.3	597.9	938.8
2000.....	2.2	5.6	11.5	31.6	69.6	142.8	264.8	414.0	645.1	992.8
1999.....	2.0	5.0	11.6	34.3	76.0	155.3	284.6	438.9	660.3	1,022.5
White, female										
2004.....	2.5	5.1	10.9	25.6	59.9	111.6	192.9	286.5	386.5	529.0
2003.....	2.7	5.1	11.4	26.5	63.2	120.0	197.9	301.0	408.2	555.5
2002.....	2.4	5.4	11.0	28.1	59.7	116.8	196.4	293.1	403.0	547.9
2001.....	2.6	4.6	11.5	28.5	63.2	121.0	198.2	291.4	391.8	549.4
2000.....	2.4	4.9	11.2	28.1	61.0	118.4	199.2	289.0	398.7	537.5
1999.....	2.5	4.7	11.8	30.4	63.0	119.9	198.8	288.9	405.5	523.1
Black, both sexes										
2004.....	4.8	9.5	16.3	27.8	52.0	81.5	127.5	204.0	248.5	332.5
2003.....	5.6	9.1	16.5	30.9	52.6	86.4	146.3	211.3	270.8	352.7
2002.....	6.5	9.0	17.2	30.9	53.3	85.6	147.6	215.4	297.0	367.0
2001.....	5.1	9.0	16.6	31.5	51.9	94.0	149.2	206.0	275.2	374.6
2000.....	6.1	10.9	18.1	32.6	57.1	96.6	150.4	211.1	270.5	373.2
1999.....	6.1	11.1	17.1	35.8	57.4	99.7	162.5	241.0	293.6	375.3
Black, male										
2004.....	4.9	9.8	18.9	37.5	67.5	109.0	181.9	300.7	392.4	564.3
2003.....	5.1	9.8	17.5	37.7	67.7	112.0	211.5	326.4	470.3	601.6
2002.....	6.9	9.8	19.7	38.0	70.9	115.7	212.6	339.9	476.1	645.4
2001.....	5.1	8.8	18.8	36.9	66.1	126.5	221.5	328.5	465.5	669.4
2000.....	5.9	11.3	19.6	39.5	73.1	137.1	225.0	325.1	448.5	693.1
1999.....	6.1	11.8	19.6	42.9	78.9	139.4	242.9	385.1	520.6	700.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
Black, female										
2004.....	4.8	9.2	14.2	19.8	39.9	61.0	90.5	146.6	172.7	240.6
2003.....	6.1	8.4	15.7	25.3	40.9	67.4	102.4	143.0	166.3	256.0
2002.....	6.1	8.3	15.1	25.1	39.5	63.3	104.0	141.1	203.7	260.4
2001.....	5.1	9.2	14.7	27.1	40.8	70.0	101.1	132.9	178.2	265.2
2000.....	6.2	10.6	16.8	27.0	44.5	66.6	100.8	142.4	180.9	254.1
1999.....	6.1	10.6	14.9	30.0	40.4	70.2	109.1	154.2	181.1	253.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Bronchitis, not specified as acute or chronic (J40)

All races 3/, both sexes

2004.....	*	*	*	*	*	0.2	0.3	0.5	1.0	3.8
2003.....	*	*	*	*	*	*	0.3	0.7	1.1	4.6
2002.....	*	*	*	*	*	*	0.4	0.7	1.4	5.2
2001.....	*	*	*	*	*	0.2	0.4	1.0	1.3	5.2
2000.....	*	*	*	*	0.2	0.3	0.4	1.1	2.0	6.8
1999.....	*	*	*	*	0.3	0.3	0.7	1.0	2.2	6.2

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	3.9
2003.....	*	*	*	*	*	*	*	0.8	1.1	3.9
2002.....	*	*	*	*	*	*	*	0.8	1.2	4.7
2001.....	*	*	*	*	*	*	*	1.0	1.2	5.1
2000.....	*	*	*	*	*	*	*	1.3	3.0	6.4
1999.....	*	*	*	*	*	*	0.7	1.2	2.3	5.9

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	1.0	3.7
2003.....	*	*	*	*	*	*	*	0.6	1.0	4.9
2002.....	*	*	*	*	*	*	0.5	0.7	1.5	5.4
2001.....	*	*	*	*	*	*	*	0.9	1.4	5.3
2000.....	*	*	*	*	*	*	0.5	1.0	1.4	7.0
1999.....	*	*	*	*	*	*	0.6	0.8	2.1	6.3

White, both sexes

2004.....	*	*	*	*	*	*	0.3	0.5	1.0	3.9
2003.....	*	*	*	*	*	*	0.3	0.8	1.1	4.7
2002.....	*	*	*	*	*	*	0.5	0.8	1.5	5.5
2001.....	*	*	*	*	*	*	0.4	1.0	1.4	5.5
2000.....	*	*	*	*	0.2	0.3	0.4	1.2	2.1	7.3
1999.....	*	*	*	*	0.3	0.3	0.7	0.9	2.2	6.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic bronchitis (J41-J42)

All races 3/, both sexes

2004.....	*	*	*	0.1	*	0.2	0.4	0.7	1.1	2.8
2003.....	*	*	*	*	*	0.3	0.4	0.7	1.5	2.7
2002.....	*	*	*	*	0.2	0.4	0.5	1.0	1.5	2.9
2001.....	*	*	*	*	0.2	0.3	0.4	0.8	1.2	3.9
2000.....	*	*	*	*	0.2	0.4	0.6	1.0	1.6	5.0
1999.....	*	*	*	0.2	0.3	0.3	0.6	0.9	1.9	4.5

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	1.3	3.1
2003.....	*	*	*	*	*	*	*	1.1	1.5	3.0
2002.....	*	*	*	*	*	0.5	0.7	1.5	1.6	4.0
2001.....	*	*	*	*	*	0.5	0.6	1.1	1.2	4.1
2000.....	*	*	*	*	*	0.6	0.6	1.4	1.9	6.4
1999.....	*	*	*	*	*	0.5	0.7	1.3	2.3	5.9

All races 3/, female

2004.....	*	*	*	*	*	*	0.4	0.7	0.9	2.6
2003.....	*	*	*	*	*	*	0.5	0.5	1.5	2.6
2002.....	*	*	*	*	*	*	*	0.7	1.4	2.4
2001.....	*	*	*	*	*	*	*	0.5	1.2	3.8
2000.....	*	*	*	*	*	*	0.6	0.7	1.4	4.4
1999.....	*	*	*	*	0.4	*	0.4	0.7	1.7	4.0

White, both sexes

2004.....	*	*	*	*	*	*	0.4	0.7	1.1	2.8
2003.....	*	*	*	*	*	0.3	0.4	0.7	1.5	2.8
2002.....	*	*	*	*	*	0.4	0.6	1.1	1.5	3.1
2001.....	*	*	*	*	0.2	0.4	0.4	0.8	1.3	4.1
2000.....	*	*	*	*	0.2	0.4	0.6	1.0	1.6	5.3
1999.....	*	*	*	0.2	0.3	0.4	0.6	1.0	1.9	4.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
All races 3/, both sexes										
2004.....	0.2	0.6	1.5	3.4	8.1	15.8	26.1	37.5	45.7	49.9
2003.....	0.3	0.7	1.5	4.0	9.3	16.6	28.5	41.9	51.6	54.6
2002.....	0.3	0.7	1.6	4.3	9.1	17.4	31.3	43.2	54.7	59.3
2001.....	0.2	0.7	1.7	5.0	10.1	19.8	33.2	46.9	55.5	62.1
2000.....	0.3	0.8	1.8	5.0	10.4	20.7	35.5	47.5	60.4	63.3
1999.....	0.4	0.7	2.0	5.5	12.1	23.5	38.4	51.3	65.3	62.1
All races 3/, male										
2004.....	0.3	0.8	1.6	3.8	9.5	18.4	31.6	46.5	60.5	74.4
2003.....	0.3	1.0	1.9	4.4	10.3	18.3	34.4	51.9	67.5	79.2
2002.....	0.5	0.9	1.9	4.6	10.7	19.7	37.8	56.1	74.9	90.0
2001.....	0.3	0.9	2.0	5.8	10.8	22.4	40.3	59.2	74.9	101.2
2000.....	0.3	1.0	2.1	5.6	11.2	24.5	42.6	60.4	83.6	104.2
1999.....	0.4	0.9	2.2	6.2	13.6	28.0	48.9	67.3	91.5	105.4
All races 3/, female										
2004.....	0.2	0.4	1.3	3.1	6.8	13.5	21.7	31.1	36.6	38.9
2003.....	0.3	0.5	1.2	3.6	8.4	15.2	23.7	34.8	41.9	43.7
2002.....	0.2	0.6	1.3	4.0	7.7	15.4	26.1	34.0	42.4	46.0
2001.....	*	0.5	1.4	4.2	9.5	17.6	27.5	38.3	43.8	45.8
2000.....	0.2	0.6	1.5	4.5	9.7	17.5	29.9	38.5	46.8	46.7
1999.....	0.3	0.5	1.7	4.9	10.8	19.8	30.2	40.2	49.9	44.7
White, both sexes										
2004.....	0.2	0.6	1.4	3.6	8.6	17.2	28.2	40.1	48.4	52.5
2003.....	0.3	0.7	1.6	4.2	9.9	18.0	30.5	44.3	54.2	57.0
2002.....	0.3	0.7	1.7	4.6	9.8	18.6	33.3	45.8	57.4	62.3
2001.....	0.2	0.7	1.7	5.1	10.9	21.5	35.2	49.8	58.7	65.1
2000.....	0.2	0.7	1.8	5.3	11.2	22.0	37.9	50.2	63.6	65.8
1999.....	0.3	0.6	2.0	5.8	13.0	25.4	40.9	53.8	68.8	64.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
White, male										
2004.....	0.3	0.8	1.6	3.8	9.8	19.4	33.4	48.5	63.2	77.1
2003.....	0.3	1.0	1.9	4.5	10.6	19.5	35.9	53.2	69.6	80.4
2002.....	0.4	0.9	1.9	4.8	11.1	20.2	39.5	58.2	77.4	93.0
2001.....	0.3	1.0	1.9	5.8	11.3	23.9	41.6	61.7	78.0	105.1
2000.....	0.2	0.9	2.0	5.8	11.8	25.0	44.6	63.1	87.0	105.5
1999.....	0.3	0.8	2.1	6.4	14.2	29.4	50.6	69.6	94.4	109.2
White, female										
2004.....	*	0.4	1.3	3.4	7.5	15.2	24.0	33.9	39.2	41.4
2003.....	0.3	0.5	1.4	3.9	9.2	16.7	26.0	37.8	44.7	46.6
2002.....	*	0.6	1.5	4.4	8.6	17.3	28.3	37.0	45.1	48.9
2001.....	*	0.5	1.4	4.5	10.6	19.4	30.0	41.3	47.0	48.4
2000.....	0.2	0.6	1.5	4.8	10.7	19.3	32.5	41.1	49.8	49.7
1999.....	0.3	0.5	1.9	5.3	11.9	21.9	33.1	42.6	53.7	46.9
Black, both sexes										
2004.....	*	*	2.1	3.1	6.5	8.4	13.7	20.7	24.3	27.6
2003.....	*	1.0	1.3	3.6	7.2	10.2	16.4	25.2	30.5	30.9
2002.....	0.8	1.0	1.5	3.3	5.9	11.9	20.9	23.5	34.8	32.1
2001.....	*	*	2.3	4.9	6.4	11.6	20.1	21.9	26.3	35.3
2000.....	*	1.6	2.5	4.7	7.7	14.9	19.5	24.5	30.8	37.2
1999.....	0.8	1.4	2.4	4.6	8.0	14.0	20.8	31.4	31.2	37.6
Black, male										
2004.....	*	*	3.0	4.9	9.7	13.8	21.4	34.6	39.5	56.6
2003.....	*	*	2.0	5.2	10.2	13.3	24.6	43.4	50.2	70.9
2002.....	*	*	2.4	4.5	9.3	20.4	30.7	41.3	61.1	64.3
2001.....	*	*	2.9	6.9	9.6	15.8	31.6	34.4	49.0	68.1
2000.....	*	2.4	3.5	5.9	10.8	25.2	30.1	35.0	53.1	87.6
1999.....	*	2.0	4.1	6.4	11.9	22.8	37.0	48.9	65.4	72.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Emphysema (J43)

Black, female

2004.....	*	*	*	*	3.9	4.3	8.6	12.4	16.3	16.1
2003.....	*	*	*	2.3	4.8	7.9	10.9	14.4	20.1	15.4
2002.....	*	*	*	*	3.3	5.6	14.3	12.9	21.1	19.7
2001.....	*	*	*	3.3	4.0	8.4	12.3	14.4	14.7	23.2
2000.....	*	*	*	3.6	5.2	7.3	12.4	18.2	19.5	18.5
1999.....	*	*	*	3.0	5.0	7.5	10.1	20.9	14.3	24.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic obstructive pulmonary disease (J44)

All races 3/, both sexes

2004.....	1.4	3.6	8.6	20.1	49.4	94.6	169.7	270.4	376.4	524.8
2003.....	1.3	3.5	8.5	21.3	51.9	100.9	177.4	279.4	390.4	552.9
2002.....	1.4	3.5	8.6	21.6	48.8	97.8	174.2	277.3	395.7	549.7
2001.....	1.3	3.3	8.3	22.3	49.4	100.0	176.2	273.1	382.9	550.3
2000.....	1.2	3.3	8.2	22.1	48.8	98.6	175.8	270.2	396.1	550.6
1999.....	1.1	3.1	8.0	23.3	51.2	101.7	182.1	278.2	400.8	549.6

All races 3/, male

2004.....	1.4	3.6	9.7	21.4	53.3	104.9	192.8	326.3	474.0	714.1
2003.....	1.2	4.0	9.1	23.2	56.7	111.9	205.7	336.6	490.1	764.9
2002.....	1.5	3.7	9.4	23.2	52.9	108.7	203.4	341.3	508.8	780.4
2001.....	1.3	3.7	8.9	24.1	53.5	111.0	207.7	339.9	498.3	789.6
2000.....	1.2	3.8	8.7	24.2	54.2	111.1	209.8	335.5	530.2	833.7
1999.....	1.2	3.6	8.5	25.7	58.4	119.2	222.9	355.4	540.9	860.0

All races 3/, female

2004.....	1.3	3.5	7.7	18.8	46.0	85.6	151.1	230.1	316.4	439.6
2003.....	1.4	3.1	8.0	19.5	47.5	91.3	154.7	238.7	329.5	459.2
2002.....	1.2	3.2	7.8	20.2	45.1	88.4	150.9	232.1	326.9	449.5
2001.....	1.4	2.8	7.7	20.6	45.7	90.5	151.2	226.3	313.7	450.2
2000.....	1.2	2.9	7.7	20.2	43.9	87.8	149.0	224.6	317.1	435.3
1999.....	1.1	2.5	7.6	21.1	44.7	86.7	150.2	224.6	318.9	424.5

White, both sexes

2004.....	1.4	3.7	8.9	20.9	52.6	101.4	181.4	285.7	396.6	550.5
2003.....	1.3	3.7	8.8	22.1	55.2	108.2	188.3	294.5	411.1	579.4
2002.....	1.4	3.6	8.8	22.7	51.4	104.8	184.5	291.4	413.7	574.7
2001.....	1.3	3.3	8.5	23.3	52.7	105.9	186.4	287.4	400.3	573.9
2000.....	1.1	3.3	8.3	23.0	51.5	104.3	185.5	283.0	414.8	574.7
1999.....	1.1	3.0	8.3	24.2	53.8	107.3	190.7	289.4	417.8	573.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic obstructive pulmonary disease (J44)

White, male

2004.....	1.5	3.7	9.8	21.5	55.3	110.4	201.6	339.0	491.5	738.3
2003.....	1.2	4.2	9.2	23.5	59.1	117.6	213.1	347.1	504.7	791.5
2002.....	1.5	3.8	9.4	23.6	54.4	114.4	210.6	351.3	523.6	805.8
2001.....	1.3	3.8	8.9	24.7	55.8	114.9	214.9	350.1	512.4	812.9
2000.....	1.2	3.7	8.5	24.7	56.0	114.7	216.3	344.3	546.6	862.1
1999.....	1.1	3.5	8.6	26.1	60.0	123.0	229.5	362.6	554.3	890.0

White, female

2004.....	1.4	3.6	8.1	20.4	50.1	93.5	164.8	246.7	337.6	465.8
2003.....	1.4	3.3	8.5	20.7	51.5	99.9	168.0	256.6	353.4	485.4
2002.....	1.3	3.5	8.1	21.9	48.7	96.3	163.2	248.5	346.4	474.3
2001.....	1.4	2.8	8.1	22.0	49.8	98.0	163.3	243.0	332.6	473.8
2000.....	1.1	3.0	8.1	21.4	47.3	95.2	160.8	239.8	336.4	457.9
1999.....	1.1	2.5	8.1	22.4	48.2	93.5	159.9	237.9	337.2	446.6

Black, both sexes

2004.....	1.5	4.0	9.8	19.3	38.8	67.3	107.6	173.5	214.1	284.5
2003.....	1.7	3.7	9.3	21.7	40.3	69.4	121.9	177.5	227.7	304.3
2002.....	1.7	3.8	10.0	20.3	41.6	67.2	116.4	183.8	250.9	318.9
2001.....	1.6	4.0	9.3	20.6	37.3	75.5	118.7	172.5	237.0	319.7
2000.....	1.8	3.8	10.1	21.3	41.9	72.0	119.9	174.1	227.0	310.0
1999.....	1.9	4.7	8.4	23.6	42.8	76.8	129.1	197.6	247.0	312.8

Black, male

2004.....	1.5	3.8	12.3	27.8	51.4	90.1	153.4	257.5	342.9	490.8
2003.....	1.8	4.5	10.8	28.5	52.4	93.8	180.3	272.5	409.0	520.3
2002.....	2.0	4.7	12.5	27.4	56.0	88.3	172.9	290.7	405.8	565.0
2001.....	1.6	4.3	11.3	26.4	50.2	105.2	178.7	283.9	405.3	580.9
2000.....	1.7	4.9	12.5	27.3	55.7	102.8	186.5	283.5	383.6	576.6
1999.....	2.4	5.9	10.8	30.9	60.9	109.6	191.6	323.9	440.3	605.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic obstructive pulmonary disease (J44)

Black, female

2004.....	1.4	4.2	7.6	12.3	29.0	50.4	76.4	123.7	146.3	202.8
2003.....	1.5	3.0	7.9	16.2	30.8	51.4	82.4	121.0	132.7	220.4
2002.....	1.4	2.9	7.8	14.4	30.4	51.6	78.4	119.9	170.3	224.7
2001.....	1.5	3.7	7.7	15.8	27.2	53.5	78.7	105.9	151.3	222.7
2000.....	1.9	2.8	7.9	16.3	31.1	49.3	75.7	108.2	148.2	210.8
1999.....	1.5	3.7	6.4	17.7	28.6	52.5	87.5	121.6	151.2	203.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Asthma (J45-J46)

All races 3/, both sexes

2004.....	0.9	1.3	1.4	1.7	1.9	2.5	3.0	4.2	6.6	14.1
2003.....	1.1	1.5	1.7	1.9	2.2	2.7	3.1	4.9	7.2	14.2
2002.....	1.2	1.4	1.6	1.9	2.1	2.7	3.8	5.4	7.7	14.7
2001.....	1.1	1.3	1.8	1.9	2.6	3.0	3.9	5.2	8.1	16.1
2000.....	1.2	1.6	1.7	2.0	2.5	3.4	4.6	5.7	8.7	16.7
1999.....	1.1	1.6	1.7	2.6	2.2	3.7	4.7	6.2	9.6	17.3

All races 3/, male

2004.....	0.8	1.1	1.0	1.4	1.3	2.0	2.3	2.8	4.4	11.1
2003.....	0.9	1.2	1.2	1.5	1.7	1.8	2.4	3.7	5.7	10.7
2002.....	0.9	1.1	1.3	1.5	1.7	2.0	2.8	4.3	6.2	10.9
2001.....	0.9	1.0	1.3	1.2	1.9	2.4	3.0	3.8	5.6	12.1
2000.....	1.0	1.3	1.3	1.6	1.8	2.7	3.4	3.9	6.7	12.9
1999.....	0.8	1.0	1.2	1.9	1.7	2.6	3.9	4.8	7.2	12.7

All races 3/, female

2004.....	1.1	1.5	1.8	2.0	2.5	2.9	3.6	5.1	7.9	15.4
2003.....	1.4	1.7	2.1	2.2	2.6	3.5	3.7	5.7	8.0	15.8
2002.....	1.4	1.8	1.9	2.3	2.5	3.3	4.5	6.2	8.7	16.3
2001.....	1.3	1.7	2.2	2.5	3.3	3.6	4.7	6.1	9.6	17.8
2000.....	1.4	1.9	2.1	2.4	3.1	4.0	5.5	6.9	9.8	18.2
1999.....	1.3	2.2	2.2	3.3	2.7	4.6	5.4	7.2	11.1	19.2

White, both sexes

2004.....	0.7	0.8	1.0	1.3	1.4	2.1	2.8	3.6	6.2	13.7
2003.....	0.8	1.1	1.1	1.5	1.8	2.3	2.6	4.4	6.7	14.3
2002.....	0.8	1.1	1.1	1.4	1.8	2.2	3.2	5.1	7.5	14.7
2001.....	0.8	1.0	1.4	1.4	1.9	2.6	3.4	4.7	7.8	15.7
2000.....	0.9	1.1	1.2	1.4	2.0	2.8	4.0	5.2	8.4	15.9
1999.....	0.8	1.2	1.3	2.0	1.8	3.1	4.0	5.8	9.4	16.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Asthma (J45-J46)

White, male

2004.....	0.6	0.6	0.7	0.9	0.8	1.6	1.9	2.2	3.8	10.3
2003.....	0.6	0.9	0.8	1.2	1.2	1.5	2.0	3.0	5.4	10.6
2002.....	0.6	0.8	0.9	1.0	1.4	1.5	2.3	3.8	5.8	10.7
2001.....	0.6	0.7	0.9	1.0	1.2	2.0	2.4	3.3	5.1	11.3
2000.....	0.7	1.0	1.0	1.0	1.4	2.0	3.0	3.7	6.3	11.4
1999.....	0.5	0.7	0.9	1.5	1.2	2.0	3.0	4.2	7.0	10.8

White, female

2004.....	0.8	1.0	1.4	1.6	2.1	2.6	3.5	4.7	7.7	15.2
2003.....	1.0	1.3	1.5	1.7	2.3	3.0	3.1	5.4	7.5	15.9
2002.....	1.0	1.3	1.3	1.7	2.2	2.8	3.9	6.1	8.6	16.4
2001.....	1.0	1.2	1.8	1.9	2.5	3.2	4.2	5.7	9.4	17.6
2000.....	1.0	1.2	1.5	1.8	2.6	3.4	4.8	6.3	9.7	17.7
1999.....	1.1	1.7	1.7	2.6	2.2	4.0	4.8	6.9	10.8	18.9

Black, both sexes

2004.....	2.9	4.8	4.1	5.1	6.3	5.7	5.6	9.0	9.4	15.5
2003.....	3.4	4.1	5.7	5.2	5.0	6.6	7.5	8.2	10.4	12.0
2002.....	4.0	4.1	5.7	7.1	5.6	6.1	10.1	7.4	8.9	11.9
2001.....	3.1	4.3	4.9	5.9	7.9	6.3	9.4	10.2	11.1	16.3
2000.....	3.6	5.4	5.3	6.6	7.2	8.8	9.7	10.7	9.9	22.2
1999.....	3.3	4.8	5.8	7.3	6.2	8.4	11.1	9.8	11.3	20.4

Black, male

2004.....	2.7	4.7	2.9	4.8	5.8	4.8	6.5	*	*	*
2003.....	2.7	3.5	4.4	3.8	4.8	4.9	*	9.1	*	*
2002.....	3.7	3.4	4.6	6.0	5.3	6.2	8.7	*	*	*
2001.....	2.9	3.8	4.5	3.6	5.9	*	9.8	*	*	*
2000.....	3.1	4.0	3.5	6.3	6.0	8.6	7.0	*	*	25.4
1999.....	2.9	3.6	4.2	5.1	5.7	6.8	12.2	9.5	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Asthma (J45-J46)

Black, female

2004.....	3.1	4.9	5.1	5.4	6.7	6.3	5.1	9.7	9.1	16.9
2003.....	4.1	4.7	6.7	6.3	5.1	7.9	8.6	7.6	11.6	13.8
2002.....	4.2	4.7	6.6	8.1	5.7	5.9	11.0	8.0	9.5	12.3
2001.....	3.3	4.8	5.3	7.8	9.4	7.5	9.2	11.4	11.9	16.8
2000.....	4.0	6.6	6.9	6.9	8.2	9.0	11.5	14.2	10.2	21.0
1999.....	3.6	5.9	7.1	9.1	6.5	9.6	10.4	10.0	13.8	21.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
 years years years years years years years years years and over

Bronchiectasis (J47)

All races 3/, both sexes

2004.....	*	*	0.1	0.1	0.2	0.5	1.2	2.5	4.0	6.3
2003.....	*	*	*	0.2	0.3	0.5	1.1	2.3	4.1	6.0
2002.....	*	*	*	0.1	0.4	0.6	1.2	2.5	4.6	5.8
2001.....	*	*	*	0.2	0.4	0.7	1.2	2.3	3.9	7.0
2000.....	*	*	*	0.3	0.3	0.7	1.2	2.4	4.6	6.1
1999.....	*	*	*	0.2	0.4	0.8	1.1	2.6	4.7	6.2

All races 3/, male

2004.....	*	*	*	*	*	*	0.8	1.9	2.3	4.5
2003.....	*	*	*	*	*	*	0.8	1.6	3.2	4.2
2002.....	*	*	*	*	*	*	0.9	1.6	3.1	4.7
2001.....	*	*	*	*	0.4	0.6	0.8	1.6	2.2	4.8
2000.....	*	*	*	*	*	0.5	1.0	1.6	3.2	4.3
1999.....	*	*	*	*	*	*	1.0	2.5	4.1	6.6

All races 3/, female

2004.....	*	*	*	*	*	0.6	1.5	2.9	5.0	7.2
2003.....	*	*	*	0.3	*	0.8	1.3	2.8	4.7	6.8
2002.....	*	*	*	*	0.5	0.9	1.4	3.1	5.5	6.3
2001.....	*	*	*	*	0.4	0.9	1.5	2.8	4.9	7.9
2000.....	*	*	*	0.3	0.4	0.9	1.4	3.0	5.5	6.9
1999.....	*	*	*	0.3	0.4	1.1	1.2	2.6	5.1	6.0

White, both sexes

2004.....	*	*	0.1	*	0.3	0.5	1.2	2.6	4.3	6.7
2003.....	*	*	*	0.1	0.3	0.5	1.1	2.4	4.3	6.4
2002.....	*	*	*	0.2	0.5	0.6	1.3	2.7	4.9	6.1
2001.....	*	*	*	*	0.4	0.8	1.2	2.4	4.1	7.4
2000.....	*	*	*	0.2	0.3	0.7	1.2	2.5	4.9	6.6
1999.....	*	*	*	0.2	0.4	0.8	1.1	2.6	4.8	6.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Lung diseases due to external agents (J60-J70)

All races 3/, both sexes

2004.....	0.5	0.8	1.3	2.1	3.8	7.0	15.4	31.6	66.5	171.1
2003.....	0.5	0.9	1.2	2.2	4.2	7.3	15.1	33.3	70.6	182.8
2002.....	0.6	0.7	1.2	2.0	3.9	7.5	15.2	34.9	71.7	192.9
2001.....	0.5	0.7	1.3	2.1	4.0	7.3	16.2	34.0	73.3	195.9
2000.....	0.5	0.8	1.2	2.1	3.5	7.7	16.5	34.0	70.6	195.3
1999.....	0.5	0.7	1.0	2.1	3.5	7.3	15.3	31.9	66.1	181.8

All races 3/, male

2004.....	0.6	1.0	1.6	2.6	5.3	10.0	22.2	46.0	98.0	256.1
2003.....	0.5	1.1	1.5	2.9	5.6	10.7	21.5	48.8	105.0	266.2
2002.....	0.7	0.9	1.6	2.6	5.1	10.5	22.0	51.6	105.9	290.1
2001.....	0.7	0.8	1.8	2.8	5.0	10.4	23.7	49.1	111.6	291.3
2000.....	0.7	1.1	1.6	2.7	4.6	11.2	25.2	50.6	107.8	301.5
1999.....	0.6	0.9	1.2	2.7	5.1	10.7	22.9	50.0	103.9	279.1

All races 3/, female

2004.....	0.3	0.6	0.9	1.5	2.5	4.3	10.0	21.2	47.1	132.9
2003.....	0.5	0.7	0.8	1.5	3.0	4.3	10.0	22.3	49.6	145.9
2002.....	0.4	0.5	0.8	1.5	2.9	5.0	9.9	23.2	50.9	150.7
2001.....	0.4	0.6	0.9	1.4	3.0	4.6	10.3	23.5	50.3	156.0
2000.....	0.4	0.6	0.9	1.6	2.5	4.8	9.7	22.4	48.6	152.1
1999.....	0.4	0.5	0.9	1.6	2.1	4.3	9.3	19.3	44.0	142.6

White, both sexes

2004.....	0.4	0.8	1.2	2.0	3.5	6.9	15.2	31.7	67.4	174.3
2003.....	0.4	0.9	1.2	2.2	3.8	7.0	15.1	33.1	70.4	185.7
2002.....	0.5	0.7	1.1	2.0	3.8	7.3	15.1	34.7	72.8	196.8
2001.....	0.5	0.7	1.2	2.1	3.8	7.2	15.8	34.1	73.7	197.6
2000.....	0.6	0.7	1.2	2.1	3.3	7.5	15.9	33.8	70.8	199.2
1999.....	0.4	0.6	0.9	2.0	3.3	7.0	14.7	31.5	66.1	184.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Lung diseases due to external agents (J60-J70)

White, male

2004.....	0.5	1.0	1.6	2.5	4.7	9.9	21.6	46.1	99.6	261.3
2003.....	0.5	1.1	1.6	2.9	5.1	10.5	21.2	48.2	105.0	272.3
2002.....	0.7	0.9	1.5	2.6	4.9	10.1	21.6	51.1	107.8	297.5
2001.....	0.7	0.8	1.6	2.8	4.7	10.4	23.2	49.4	112.8	297.5
2000.....	0.7	0.9	1.4	2.7	4.4	10.6	23.9	50.0	109.3	311.0
1999.....	0.5	0.8	1.0	2.5	4.7	10.2	21.9	49.2	103.7	285.0

White, female

2004.....	0.3	0.6	0.8	1.6	2.4	4.3	10.0	21.3	47.4	135.0
2003.....	0.4	0.7	0.8	1.5	2.7	3.9	10.0	22.1	49.1	147.3
2002.....	0.4	0.5	0.8	1.5	2.7	4.7	9.8	23.0	51.3	153.0
2001.....	0.4	0.5	0.9	1.4	3.0	4.3	9.8	23.3	50.0	155.8
2000.....	0.4	0.5	0.9	1.5	2.3	4.8	9.4	22.5	47.9	153.8
1999.....	0.4	0.4	0.9	1.5	2.1	4.2	9.0	19.0	43.9	144.0

Black, both sexes

2004.....	0.9	1.0	1.7	3.1	7.6	9.5	21.5	35.4	68.1	149.7
2003.....	0.9	1.4	1.3	2.8	8.6	11.6	17.9	42.2	78.9	166.0
2002.....	1.0	0.9	2.1	2.6	5.9	12.0	19.8	44.0	70.6	169.3
2001.....	0.8	1.1	2.5	2.6	6.4	9.8	23.4	39.9	79.6	189.2
2000.....	*	1.7	2.2	3.0	5.9	11.8	26.8	41.6	74.3	163.6
1999.....	1.0	1.2	2.0	4.0	5.9	10.9	24.8	42.7	73.4	172.2

Black, male

2004.....	*	*	2.3	4.5	11.8	15.3	33.7	53.2	101.9	220.5
2003.....	*	1.8	*	3.9	11.6	16.0	27.9	65.7	115.2	221.0
2002.....	*	*	3.2	3.7	7.3	17.5	30.7	67.5	99.3	247.7
2001.....	*	*	3.5	3.6	9.2	13.5	33.7	59.9	118.8	257.6
2000.....	*	2.4	3.3	3.3	7.7	18.7	45.1	69.0	103.7	223.7
1999.....	*	*	2.7	5.3	10.0	17.4	40.0	69.8	116.0	246.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Lung diseases due to external agents (J60-J70)

Black, female

2004.....	*	*	*	*	4.4	5.2	13.2	24.8	50.3	121.7
2003.....	*	*	*	*	6.3	8.4	11.1	28.2	60.0	144.6
2002.....	*	*	*	*	4.8	8.0	12.5	29.9	55.7	139.2
2001.....	*	*	*	*	4.3	7.1	16.6	28.0	59.7	163.8
2000.....	*	*	*	2.7	4.6	6.7	14.7	25.1	59.4	141.2
1999.....	*	*	*	2.9	*	6.2	14.6	26.3	52.3	144.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumoconioses and chemical effects (J60-J66,J68)

All races 3/, both sexes

2004.....	*	*	*	0.2	0.4	0.8	1.5	2.8	4.2	6.0
2003.....	*	*	*	0.2	0.4	0.9	1.7	3.0	4.9	6.4
2002.....	*	*	*	0.2	0.4	0.8	1.6	3.4	4.2	6.8
2001.....	*	*	*	0.2	0.3	1.0	2.0	3.4	5.1	6.2
2000.....	*	*	*	0.2	0.3	0.9	2.2	3.4	4.8	7.5
1999.....	*	*	*	*	0.4	1.1	1.9	3.4	5.2	6.1

All races 3/, male

2004.....	*	*	*	0.3	0.7	1.7	3.3	6.6	10.4	18.4
2003.....	*	*	*	0.3	0.7	2.0	3.5	6.9	12.5	19.8
2002.....	*	*	*	0.3	0.8	1.7	3.5	8.0	10.5	21.6
2001.....	*	*	*	0.4	0.7	2.0	4.4	7.8	13.2	20.2
2000.....	*	*	*	*	0.7	1.8	4.8	7.9	12.3	24.9
1999.....	*	*	*	*	0.7	2.3	4.4	8.1	13.5	20.4

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	*	*	*	0.2	0.4	0.9	1.6	3.2	4.4	6.2
2003.....	*	*	*	0.2	0.4	1.0	1.8	3.1	5.2	6.9
2002.....	*	*	*	0.2	0.4	0.9	1.8	3.6	4.4	7.2
2001.....	*	*	*	0.2	0.3	1.1	2.1	3.7	5.4	6.5
2000.....	*	*	*	0.2	0.4	1.0	2.4	3.5	5.0	7.9
1999.....	*	*	*	*	0.3	1.1	2.0	3.6	5.5	6.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonitis due to solids and liquids (J69)

All races 3/, both sexes

2004.....	0.5	0.8	1.2	1.9	3.4	6.0	13.7	28.4	62.2	164.9
2003.....	0.4	0.9	1.1	2.0	3.8	6.3	13.3	30.1	65.4	176.2
2002.....	0.5	0.7	1.1	1.8	3.4	6.5	13.4	31.3	67.3	186.0
2001.....	0.5	0.7	1.3	1.8	3.6	6.2	14.1	30.4	68.0	189.4
2000.....	0.5	0.8	1.2	1.9	3.2	6.7	14.2	30.4	65.7	187.6
1999.....	0.5	0.7	0.9	2.0	3.1	6.1	13.2	28.2	60.7	175.6

All races 3/, male

2004.....	0.6	1.0	1.5	2.3	4.5	8.1	18.7	39.0	87.4	237.2
2003.....	0.4	1.0	1.4	2.6	4.8	8.6	17.8	41.5	92.3	246.2
2002.....	0.7	0.9	1.5	2.2	4.2	8.6	18.2	43.4	95.1	268.5
2001.....	0.7	0.8	1.7	2.3	4.2	8.2	19.1	41.0	98.1	270.3
2000.....	0.6	1.0	1.5	2.4	3.9	9.2	20.2	42.4	95.2	276.0
1999.....	0.6	0.9	1.0	2.4	4.2	8.3	18.4	41.6	90.1	258.4

All races 3/, female

2004.....	0.3	0.5	0.8	1.5	2.5	4.2	9.8	20.9	46.6	132.4
2003.....	0.4	0.7	0.8	1.5	2.9	4.3	9.7	22.1	49.0	145.3
2002.....	0.4	0.5	0.8	1.4	2.8	4.8	9.6	22.8	50.4	150.2
2001.....	0.4	0.6	0.9	1.4	3.0	4.4	10.2	23.0	49.9	155.6
2000.....	0.4	0.6	0.9	1.5	2.5	4.6	9.4	22.1	48.2	151.6
1999.....	0.4	0.5	0.8	1.6	2.0	4.2	9.2	18.8	43.5	142.2

White, both sexes

2004.....	0.4	0.7	1.1	1.8	3.1	5.9	13.4	28.2	62.9	167.8
2003.....	0.4	0.8	1.1	2.0	3.3	5.9	13.1	29.7	65.0	178.7
2002.....	0.5	0.7	1.0	1.8	3.3	6.1	13.1	30.9	68.1	189.5
2001.....	0.5	0.7	1.2	1.8	3.5	6.0	13.6	30.2	68.1	190.8
2000.....	0.5	0.7	1.1	1.9	2.9	6.4	13.4	30.1	65.7	191.0
1999.....	0.4	0.6	0.8	1.8	2.9	5.8	12.5	27.6	60.3	178.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonitis due to solids and liquids (J69)

White, male										
2004.....	0.5	1.0	1.4	2.2	3.9	7.8	17.9	38.3	88.4	241.7
2003.....	0.4	1.0	1.4	2.5	4.2	8.2	17.2	40.6	91.7	250.8
2002.....	0.6	0.8	1.3	2.1	4.1	8.0	17.6	42.5	96.5	274.6
2001.....	0.7	0.8	1.5	2.3	4.0	8.0	18.4	40.6	98.7	275.4
2000.....	0.7	0.9	1.3	2.3	3.6	8.5	18.7	41.4	96.2	284.0
1999.....	0.5	0.8	0.9	2.2	3.9	7.7	17.1	40.4	88.9	263.6
White, female										
2004.....	0.3	0.5	0.8	1.5	2.3	4.2	9.7	20.8	47.0	134.5
2003.....	0.4	0.7	0.8	1.5	2.5	3.9	9.7	21.8	48.6	146.7
2002.....	0.4	0.5	0.7	1.5	2.6	4.5	9.5	22.6	50.7	152.5
2001.....	0.4	0.5	0.8	1.3	3.0	4.2	9.7	22.8	49.6	155.3
2000.....	0.4	0.5	0.9	1.4	2.3	4.6	9.1	22.2	47.5	153.2
1999.....	0.4	0.4	0.8	1.5	2.0	4.1	8.8	18.6	43.4	143.6
Black, both sexes										
2004.....	0.9	1.0	1.7	3.0	7.3	8.8	20.4	34.7	64.9	144.7
2003.....	0.9	1.3	1.2	2.7	8.5	11.0	17.2	39.8	76.2	163.4
2002.....	0.9	0.8	2.1	2.4	5.5	11.7	18.8	41.9	67.8	165.1
2001.....	0.7	1.0	2.4	2.4	6.0	9.3	22.3	38.6	76.1	185.5
2000.....	*	1.7	2.2	2.9	5.8	11.3	25.1	38.9	71.4	159.8
1999.....	1.0	1.2	1.8	3.9	5.0	10.1	23.7	40.5	71.1	167.1
Black, male										
2004.....	*	*	2.2	4.4	11.0	13.6	31.4	51.4	93.3	202.7
2003.....	*	1.7	*	3.8	11.6	14.5	26.3	59.3	107.8	212.7
2002.....	*	*	3.1	3.7	6.3	16.7	28.4	62.5	91.7	233.7
2001.....	*	*	3.4	3.1	8.1	12.5	31.0	56.2	108.3	243.9
2000.....	*	2.4	3.1	3.3	7.2	17.4	41.1	62.4	96.1	209.9
1999.....	*	*	*	5.1	8.1	15.6	37.3	65.1	109.9	228.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonitis due to solids and liquids (J69)

Black, female

2004.....	*	*	*	*	4.4	5.2	13.0	24.8	49.9	121.7
2003.....	*	*	*	*	6.0	8.4	11.1	28.2	59.6	144.2
2002.....	*	*	*	*	4.8	8.0	12.3	29.6	55.3	138.8
2001.....	*	*	*	*	4.3	6.9	16.6	28.0	59.7	163.8
2000.....	*	*	*	*	4.6	6.7	14.4	24.8	59.0	141.2
1999.....	*	*	*	2.9	*	6.0	14.6	25.7	51.9	144.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonitis due to food and vomit (J69.0)

All races 3/, both sexes

2004.....	0.5	0.8	1.2	1.9	3.4	6.0	13.7	28.4	62.2	164.9
2003.....	0.4	0.9	1.1	2.0	3.8	6.3	13.3	30.1	65.4	176.2
2002.....	0.5	0.7	1.1	1.8	3.4	6.5	13.4	31.3	67.3	186.0
2001.....	0.5	0.7	1.3	1.8	3.6	6.2	14.1	30.4	67.9	189.4
2000.....	0.5	0.8	1.2	1.9	3.1	6.7	14.2	30.4	65.7	187.6
1999.....	0.5	0.7	0.9	2.0	3.0	6.1	13.2	28.1	60.6	175.6

All races 3/, male

2004.....	0.6	1.0	1.5	2.3	4.5	8.1	18.7	38.9	87.3	237.2
2003.....	0.4	1.0	1.4	2.6	4.8	8.6	17.8	41.5	92.2	246.2
2002.....	0.7	0.9	1.5	2.2	4.2	8.5	18.2	43.4	95.1	268.5
2001.....	0.7	0.8	1.7	2.3	4.2	8.2	19.1	41.0	98.0	270.2
2000.....	0.6	1.0	1.5	2.3	3.9	9.2	20.2	42.4	95.2	276.0
1999.....	0.6	0.9	1.0	2.4	4.2	8.3	18.4	41.6	90.1	258.4

All races 3/, female

2004.....	0.3	0.5	0.8	1.5	2.5	4.2	9.8	20.9	46.6	132.4
2003.....	0.4	0.7	0.8	1.5	2.9	4.3	9.7	22.0	49.0	145.3
2002.....	0.4	0.5	0.8	1.4	2.8	4.8	9.6	22.8	50.3	150.2
2001.....	0.4	0.6	0.9	1.4	3.0	4.4	10.2	23.0	49.9	155.5
2000.....	0.4	0.6	0.9	1.5	2.5	4.6	9.4	22.1	48.2	151.6
1999.....	0.4	0.5	0.8	1.6	2.0	4.2	9.2	18.8	43.4	142.2

White, both sexes

2004.....	0.4	0.7	1.1	1.8	3.1	5.9	13.4	28.2	62.9	167.8
2003.....	0.4	0.8	1.1	2.0	3.3	5.9	13.1	29.7	65.0	178.7
2002.....	0.5	0.7	1.0	1.8	3.3	6.1	13.1	30.9	68.1	189.5
2001.....	0.5	0.7	1.2	1.8	3.5	5.9	13.6	30.2	68.1	190.7
2000.....	0.5	0.7	1.1	1.8	2.9	6.4	13.4	30.1	65.7	191.0
1999.....	0.4	0.6	0.8	1.8	2.9	5.7	12.5	27.6	60.2	178.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonitis due to food and vomit (J69.0)

White, male

2004.....	0.5	1.0	1.4	2.2	3.9	7.8	17.9	38.3	88.4	241.7
2003.....	0.4	1.0	1.4	2.5	4.2	8.2	17.2	40.6	91.6	250.8
2002.....	0.6	0.8	1.3	2.1	4.1	8.0	17.6	42.5	96.5	274.6
2001.....	0.7	0.8	1.5	2.3	4.0	8.0	18.4	40.6	98.6	275.3
2000.....	0.7	0.9	1.3	2.3	3.6	8.5	18.7	41.4	96.2	284.0
1999.....	0.5	0.8	0.9	2.2	3.9	7.7	17.1	40.4	88.9	263.6

White, female

2004.....	0.3	0.5	0.8	1.5	2.3	4.2	9.7	20.8	47.0	134.5
2003.....	0.4	0.7	0.8	1.5	2.5	3.9	9.7	21.8	48.5	146.7
2002.....	0.4	0.5	0.7	1.5	2.6	4.4	9.5	22.5	50.7	152.5
2001.....	0.4	0.5	0.8	1.3	3.0	4.2	9.7	22.8	49.6	155.3
2000.....	0.4	0.5	0.9	1.4	2.3	4.6	9.1	22.2	47.5	153.2
1999.....	0.4	0.4	0.8	1.5	2.0	4.1	8.8	18.6	43.3	143.6

Black, both sexes

2004.....	0.9	1.0	1.7	3.0	7.3	8.8	20.4	34.7	64.6	144.7
2003.....	0.9	1.3	1.2	2.7	8.5	11.0	17.2	39.8	76.2	163.1
2002.....	0.9	0.8	2.1	2.4	5.5	11.7	18.8	41.9	67.5	165.1
2001.....	0.7	1.0	2.4	2.4	6.0	9.3	22.3	38.6	76.1	185.5
2000.....	*	1.7	2.2	2.9	5.8	11.3	25.1	38.9	71.4	159.8
1999.....	1.0	1.2	1.8	3.9	5.0	9.9	23.7	40.3	71.1	167.1

Black, male

2004.....	*	*	2.2	4.4	11.0	13.6	31.4	51.4	92.5	202.7
2003.....	*	1.7	*	3.8	11.6	14.5	26.3	59.3	107.8	212.7
2002.....	*	*	3.1	3.7	6.3	16.7	28.4	62.5	91.7	233.7
2001.....	*	*	3.4	3.1	8.1	12.5	31.0	56.2	108.3	243.9
2000.....	*	2.4	3.1	3.3	7.2	17.4	41.1	62.4	96.1	209.9
1999.....	*	*	*	5.1	8.1	15.3	37.3	65.1	109.9	228.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to food and vomit (J69.0)										
Black, female										
2004.....	*	*	*	*	4.4	5.2	13.0	24.8	49.9	121.7
2003.....	*	*	*	*	6.0	8.4	11.1	28.2	59.6	143.8
2002.....	*	*	*	*	4.8	8.0	12.3	29.6	54.9	138.8
2001.....	*	*	*	*	4.3	6.9	16.6	28.0	59.7	163.8
2000.....	*	*	*	*	4.6	6.7	14.4	24.8	59.0	141.2
1999.....	*	*	*	2.9	*	6.0	14.6	25.5	51.9	144.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Suppurative and necrotic conditions of lower respiratory tract (J85-J86)

All races 3/, both sexes

2004.....	0.1	0.2	0.3	0.3	0.4	0.5	0.8	1.0	1.6	2.2
2003.....	0.1	0.3	0.3	0.4	0.5	0.6	0.9	1.1	1.7	2.1
2002.....	0.1	0.2	0.3	0.3	0.5	0.6	1.0	1.3	1.9	2.3
2001.....	0.1	0.2	0.3	0.4	0.5	0.7	0.9	1.5	1.8	3.0
2000.....	0.2	0.2	0.3	0.3	0.5	0.6	1.1	1.5	1.6	2.9
1999.....	0.2	0.2	0.3	0.4	0.5	1.0	1.0	1.3	1.7	2.8

All races 3/, male

2004.....	0.2	0.2	0.4	0.4	0.5	0.8	1.2	1.1	2.7	3.8
2003.....	0.2	0.3	0.4	0.5	0.6	0.9	1.3	1.2	2.2	3.3
2002.....	0.2	0.4	0.4	0.4	0.8	0.9	1.3	1.9	3.0	3.4
2001.....	0.2	0.2	0.5	0.5	0.7	0.9	1.3	2.2	2.5	4.7
2000.....	0.2	0.3	0.4	0.5	0.6	1.0	1.4	2.2	2.1	5.8
1999.....	0.3	0.3	0.5	0.5	0.7	1.6	1.5	1.7	2.8	4.0

All races 3/, female

2004.....	*	0.2	*	*	*	*	0.5	0.9	0.9	1.5
2003.....	*	0.2	*	0.2	0.4	*	0.7	1.0	1.3	1.5
2002.....	*	*	*	*	*	0.4	0.6	0.9	1.1	1.8
2001.....	*	*	*	0.3	*	0.5	0.7	1.1	1.4	2.3
2000.....	*	*	*	*	0.4	*	0.8	0.9	1.4	1.7
1999.....	*	*	*	*	*	0.4	0.6	1.1	1.0	2.3

White, both sexes

2004.....	0.1	0.2	0.2	0.3	0.4	0.5	0.8	0.9	1.5	2.2
2003.....	0.1	0.2	0.3	0.3	0.4	0.6	0.9	1.0	1.5	2.1
2002.....	0.1	0.2	0.2	0.3	0.4	0.6	0.9	1.1	1.8	2.2
2001.....	0.1	0.2	0.3	0.4	0.5	0.7	0.9	1.5	1.8	3.0
2000.....	0.1	0.1	0.3	0.3	0.4	0.6	1.0	1.5	1.6	2.7
1999.....	0.2	0.2	0.3	0.3	0.5	0.9	1.0	1.3	1.7	2.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pleural effusion and plaque (J90,J92)

All races 3/, both sexes

2004.....	*	*	*	0.2	0.2	0.5	0.9	1.7	3.6	8.8
2003.....	*	*	*	0.1	0.2	0.5	0.9	1.6	3.5	8.8
2002.....	*	*	*	*	0.2	0.5	0.9	1.6	3.7	9.9
2001.....	*	*	*	*	0.3	0.4	0.9	1.6	3.8	10.0
2000.....	*	*	*	*	0.2	0.5	0.8	1.6	3.6	8.9
1999.....	*	*	*	0.2	0.2	0.5	0.8	1.7	3.4	9.0

All races 3/, male

2004.....	*	*	*	0.3	*	0.7	0.9	2.2	4.3	11.1
2003.....	*	*	*	*	*	0.5	0.9	1.8	4.3	11.3
2002.....	*	*	*	*	*	*	0.9	1.8	4.7	11.0
2001.....	*	*	*	*	*	0.5	1.2	1.6	4.8	12.2
2000.....	*	*	*	*	*	0.5	1.0	2.0	4.2	10.8
1999.....	*	*	*	*	*	0.6	1.1	2.2	4.2	11.5

All races 3/, female

2004.....	*	*	*	*	0.3	*	0.9	1.3	3.1	7.8
2003.....	*	*	*	*	*	0.4	0.9	1.5	3.0	7.7
2002.....	*	*	*	*	*	0.6	0.8	1.4	3.1	9.4
2001.....	*	*	*	*	*	*	0.7	1.7	3.3	9.1
2000.....	*	*	*	*	*	0.4	0.6	1.3	3.2	8.1
1999.....	*	*	*	*	*	*	0.7	1.3	2.9	8.0

White, both sexes

2004.....	*	*	*	0.2	0.3	0.5	0.9	1.7	3.6	9.0
2003.....	*	*	*	*	0.2	0.5	0.9	1.5	3.6	9.2
2002.....	*	*	*	*	0.2	0.5	0.8	1.6	3.8	10.2
2001.....	*	*	*	*	0.3	0.4	0.9	1.6	3.9	10.0
2000.....	*	*	*	*	*	0.4	0.8	1.6	3.5	9.0
1999.....	*	*	*	*	0.3	0.5	0.8	1.7	3.5	9.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases of respiratory system (J80-J84,J93-J98)

All races 3/, both sexes

2004.....	1.3	2.2	3.5	6.6	11.5	20.3	37.3	57.6	84.7	125.2
2003.....	1.1	2.2	3.5	5.8	10.4	19.5	33.7	53.6	77.0	123.3
2002.....	1.2	2.0	3.5	6.3	10.9	19.0	35.0	53.0	77.0	118.1
2001.....	1.3	2.2	3.8	6.6	11.5	21.2	35.6	57.6	82.1	125.7
2000.....	1.4	2.2	4.0	7.1	11.7	21.3	34.2	56.4	82.3	125.8
1999.....	1.3	2.2	3.5	6.1	11.5	20.5	34.2	55.1	80.6	131.6

All races 3/, male

2004.....	1.2	2.2	3.9	7.1	13.2	24.7	46.6	72.7	109.1	149.7
2003.....	1.1	2.3	3.7	6.6	12.4	23.9	40.9	67.3	96.9	145.5
2002.....	1.3	2.0	3.6	7.0	12.5	22.7	44.7	65.9	95.2	138.9
2001.....	1.4	2.4	4.2	7.1	13.5	25.3	44.1	71.3	100.2	147.4
2000.....	1.4	2.3	4.3	8.0	13.5	24.6	42.3	71.9	103.5	158.3
1999.....	1.4	2.2	3.6	6.5	13.2	24.3	43.0	69.3	105.7	161.8

All races 3/, female

2004.....	1.3	2.2	3.0	6.2	9.9	16.5	29.8	46.8	69.7	114.2
2003.....	1.1	2.1	3.3	5.1	8.6	15.8	28.0	43.8	64.9	113.5
2002.....	1.2	2.1	3.5	5.7	9.5	15.8	27.3	43.9	65.9	109.0
2001.....	1.2	2.0	3.5	6.2	9.6	17.7	28.9	48.0	71.2	116.6
2000.....	1.4	2.1	3.7	6.3	10.0	18.4	27.8	45.6	69.8	112.6
1999.....	1.3	2.2	3.4	5.8	10.0	17.3	27.4	45.3	65.9	119.4

White, both sexes

2004.....	1.1	1.8	3.2	6.4	11.3	20.6	38.7	59.8	87.5	129.8
2003.....	0.9	1.9	3.2	5.6	10.5	20.0	34.9	55.1	80.6	128.0
2002.....	1.1	1.7	3.3	6.0	10.7	19.4	35.8	55.0	78.9	122.1
2001.....	1.1	1.9	3.4	6.3	11.3	21.4	36.5	59.3	84.7	129.0
2000.....	1.3	1.9	3.7	6.9	11.5	21.4	35.1	57.8	84.3	130.0
1999.....	1.1	1.9	3.2	5.9	11.5	20.8	34.5	56.2	82.4	134.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases of respiratory system (J80-J84,J93-J98)

White, male

2004.....	1.1	1.8	3.7	6.9	13.1	25.0	48.0	75.1	112.5	154.6
2003.....	0.9	2.1	3.5	6.3	12.4	24.4	42.3	68.6	101.3	150.4
2002.....	1.2	1.7	3.3	6.8	12.2	23.6	45.3	68.1	97.9	143.8
2001.....	1.4	2.1	3.8	7.0	13.1	25.8	45.5	73.7	103.5	151.5
2000.....	1.3	2.0	3.9	7.8	13.4	25.0	43.1	73.5	106.2	163.4
1999.....	1.2	1.9	3.3	6.1	13.3	24.9	43.5	71.0	108.0	166.3

White, female

2004.....	1.2	1.8	2.7	6.0	9.6	16.6	31.0	48.6	71.9	118.7
2003.....	0.9	1.7	3.0	4.9	8.7	16.2	28.8	45.3	67.8	118.0
2002.....	1.1	1.8	3.3	5.3	9.3	15.7	28.0	45.6	67.3	112.7
2001.....	0.9	1.7	3.0	5.7	9.6	17.6	29.2	49.1	73.3	119.6
2000.....	1.3	1.9	3.5	6.1	9.8	18.2	28.7	46.8	71.3	116.4
1999.....	1.0	1.9	3.1	5.6	9.9	17.1	27.3	45.9	67.4	122.2

Black, both sexes

2004.....	2.6	5.1	6.1	9.9	15.4	21.3	30.1	41.5	57.4	79.0
2003.....	2.6	4.7	6.2	9.2	12.3	18.5	26.5	44.9	43.7	77.3
2002.....	2.4	4.5	6.1	9.7	15.1	18.8	30.3	35.2	59.9	76.9
2001.....	2.6	5.0	7.9	10.5	15.2	22.2	32.0	44.6	59.0	90.0
2000.....	2.4	5.2	7.2	10.5	14.2	24.5	29.2	45.3	64.1	82.3
1999.....	3.0	5.0	6.4	9.1	13.3	22.0	34.8	48.2	58.9	94.1

Black, male

2004.....	2.7	5.7	6.2	11.1	17.0	26.9	38.5	55.5	71.0	100.3
2003.....	2.1	4.2	5.9	10.9	16.0	22.9	33.8	61.6	56.1	98.0
2002.....	2.8	4.2	6.3	10.4	18.6	20.2	40.4	45.0	73.4	86.8
2001.....	2.3	5.3	7.7	9.9	18.9	25.2	35.3	51.1	71.4	106.7
2000.....	2.5	5.9	8.0	11.7	15.7	27.1	35.4	61.4	74.2	104.9
1999.....	2.8	5.4	6.8	9.7	14.9	23.6	41.0	59.4	78.5	114.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases of respiratory system (J80-J84,J93-J98)

Black, female

2004.....	2.5	4.6	6.1	9.0	14.1	17.1	24.4	33.2	50.3	70.5
2003.....	3.1	5.3	6.5	7.7	9.3	15.2	21.5	35.0	37.1	69.3
2002.....	2.0	4.8	5.9	9.2	12.4	17.8	23.5	29.4	52.9	73.1
2001.....	3.0	4.6	8.0	11.0	12.3	20.1	29.8	40.8	52.7	83.8
2000.....	2.3	4.5	6.4	9.6	13.1	22.7	25.1	35.6	59.0	73.8
1999.....	3.3	4.6	6.1	8.7	12.1	20.8	30.6	41.5	49.3	86.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XI. Diseases of the digestive system (K00-K92)

All races 3/, both sexes

2004.....	14.4	24.6	35.1	40.1	50.3	65.7	93.3	134.1	200.3	387.5
2003.....	15.4	25.5	35.5	40.5	51.9	70.5	96.1	137.2	208.3	403.8
2002.....	15.9	26.3	33.8	40.2	50.5	68.6	96.0	139.2	212.9	415.3
2001.....	15.2	26.8	33.0	39.0	53.2	71.4	97.3	138.2	212.8	431.4
2000.....	15.9	25.8	31.7	39.8	52.9	68.9	96.3	140.7	214.9	440.7
1999.....	15.8	25.3	31.0	39.6	53.5	72.3	99.0	138.5	214.9	436.4

All races 3/, male

2004.....	18.8	33.4	50.3	54.7	63.8	79.2	106.7	151.3	212.4	366.9
2003.....	19.9	35.4	50.9	54.8	68.1	85.5	111.9	151.8	226.6	388.9
2002.....	20.7	37.0	48.7	54.2	65.3	83.9	111.8	155.6	227.7	404.4
2001.....	20.1	37.5	46.8	53.1	68.5	86.9	113.9	153.0	227.3	428.2
2000.....	21.5	37.2	44.7	53.3	68.7	84.7	112.4	160.6	229.3	441.9
1999.....	22.0	37.0	44.4	54.2	68.6	89.0	117.3	154.4	238.2	438.5

All races 3/, female

2004.....	10.1	16.1	20.5	26.2	38.0	53.9	82.5	121.7	192.8	396.8
2003.....	10.9	15.9	20.7	27.0	37.1	57.4	83.5	126.8	197.1	410.4
2002.....	11.2	16.0	19.5	27.1	37.0	55.5	83.4	127.6	203.9	420.0
2001.....	10.4	16.4	19.8	25.9	39.3	58.1	84.1	127.8	204.1	432.7
2000.....	10.4	14.9	19.3	27.1	38.7	55.5	83.7	126.8	206.5	440.2
1999.....	9.7	14.0	18.2	25.9	39.8	57.9	84.7	127.4	201.3	435.6

White, both sexes

2004.....	14.3	24.4	34.6	38.9	49.4	65.9	93.8	136.1	202.7	396.3
2003.....	15.3	25.2	34.2	39.3	50.6	70.4	97.0	139.0	211.9	412.6
2002.....	15.9	25.7	32.8	39.3	49.5	69.0	96.4	141.0	215.3	426.3
2001.....	14.7	26.0	31.4	37.0	52.3	71.1	97.7	138.3	216.0	441.6
2000.....	15.4	24.9	29.9	38.2	51.5	68.0	96.5	141.2	217.1	449.3
1999.....	15.0	23.8	29.7	37.8	52.0	71.2	98.7	138.4	217.3	445.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XI. Diseases of the digestive system (K00-K92)

White, male

2004.....	19.1	33.0	49.9	52.8	62.3	78.4	106.4	152.1	213.2	375.1
2003.....	20.0	35.4	49.1	52.8	66.4	84.9	111.5	152.8	228.0	395.9
2002.....	21.0	36.4	47.4	52.9	63.7	83.9	111.4	157.0	228.7	412.5
2001.....	19.4	36.7	44.6	50.1	67.0	85.5	113.6	151.9	230.6	434.7
2000.....	20.9	36.5	42.1	51.4	66.8	83.5	111.8	160.1	230.7	448.1
1999.....	21.1	35.0	42.5	51.8	67.1	87.8	117.1	154.3	240.5	443.1

White, female

2004.....	9.5	15.7	19.6	25.6	37.4	54.7	83.5	124.4	196.2	405.9
2003.....	10.6	15.1	19.7	26.5	36.0	57.5	85.1	129.1	202.0	420.0
2002.....	10.9	15.1	18.5	26.4	36.4	55.9	84.2	129.6	207.2	432.3
2001.....	10.0	15.5	18.5	24.5	38.7	58.4	84.9	128.7	207.2	444.5
2000.....	9.8	13.4	18.0	25.6	37.4	54.4	84.3	127.9	208.9	449.8
1999.....	8.9	12.7	17.2	24.6	38.1	56.7	84.1	127.3	203.5	446.0

Black, both sexes

2004.....	16.4	30.8	46.1	56.7	66.8	75.3	105.2	130.5	199.5	329.7
2003.....	17.3	30.9	51.6	58.1	70.0	82.9	102.7	137.5	187.3	346.0
2002.....	17.6	35.1	48.3	54.2	65.2	76.8	105.5	137.1	208.0	333.2
2001.....	20.1	36.6	51.1	62.2	68.6	87.7	108.1	151.8	201.0	351.0
2000.....	21.3	37.2	51.0	58.7	72.2	86.3	109.0	147.5	207.0	382.0
1999.....	22.6	40.1	46.6	59.1	74.3	90.3	112.7	150.0	207.5	364.4

Black, male

2004.....	19.4	42.6	65.0	83.4	91.8	102.2	134.3	162.9	229.6	322.9
2003.....	21.3	40.8	76.5	85.4	96.4	107.4	136.0	162.5	237.7	345.1
2002.....	20.9	47.7	69.4	76.8	89.3	99.5	133.2	158.0	241.5	355.9
2001.....	26.6	51.3	75.3	90.1	91.9	118.7	138.0	184.0	217.5	386.9
2000.....	28.0	49.6	73.8	81.0	97.3	109.0	138.7	181.4	241.1	392.1
1999.....	30.0	59.2	68.5	84.6	95.0	112.4	135.0	171.0	239.8	400.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XI. Diseases of the digestive system (K00-K92)

Black, female

2004.....	13.7	20.5	30.1	34.6	47.3	55.2	85.4	111.3	183.7	332.3
2003.....	13.8	22.3	30.5	35.5	49.3	64.7	80.2	122.6	160.9	346.4
2002.....	14.6	24.1	30.4	35.6	46.3	59.9	86.9	124.6	190.6	324.5
2001.....	14.3	23.9	30.5	39.4	50.2	64.8	88.1	132.6	192.6	337.6
2000.....	15.4	26.3	31.5	40.5	52.5	69.5	89.2	127.0	189.8	378.2
1999.....	16.1	23.6	28.0	38.2	58.1	73.9	97.9	137.3	191.4	350.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of other parts of digestive system (K20-K92)

All races 3/, both sexes

2004.....	14.4	24.6	35.1	40.0	50.2	65.5	93.1	133.8	199.6	385.7
2003.....	15.3	25.5	35.5	40.5	51.8	70.4	96.0	137.0	208.0	402.6
2002.....	15.9	26.3	33.8	40.2	50.4	68.5	95.9	139.0	212.5	413.8
2001.....	15.2	26.8	33.0	39.0	53.1	71.4	97.2	137.9	212.2	430.2
2000.....	15.9	25.8	31.7	39.8	52.9	68.9	96.2	140.6	214.5	439.5
1999.....	15.8	25.3	31.0	39.6	53.5	72.2	98.9	138.3	214.6	435.4

All races 3/, male

2004.....	18.8	33.4	50.3	54.7	63.7	79.1	106.6	150.9	211.7	365.5
2003.....	19.8	35.4	50.9	54.8	68.0	85.5	111.8	151.6	226.3	388.0
2002.....	20.7	37.0	48.7	54.1	65.2	83.8	111.6	155.3	227.4	403.1
2001.....	20.1	37.5	46.8	53.0	68.4	86.9	113.8	152.8	226.7	427.0
2000.....	21.4	37.1	44.7	53.3	68.6	84.6	112.2	160.4	229.0	440.8
1999.....	21.9	37.0	44.4	54.2	68.6	88.9	117.1	154.2	238.0	437.4

All races 3/, female

2004.....	10.1	16.0	20.5	26.2	37.9	53.8	82.3	121.5	192.2	394.7
2003.....	10.9	15.9	20.7	26.9	37.0	57.3	83.4	126.7	196.7	409.1
2002.....	11.2	16.0	19.5	27.0	37.0	55.4	83.3	127.5	203.4	418.4
2001.....	10.4	16.4	19.8	25.9	39.3	58.1	84.1	127.5	203.6	431.5
2000.....	10.4	14.9	19.2	27.1	38.7	55.4	83.6	126.7	206.0	439.0
1999.....	9.7	13.9	18.2	25.9	39.7	57.8	84.6	127.1	200.9	434.6

White, both sexes

2004.....	14.3	24.3	34.6	38.8	49.2	65.7	93.7	135.8	202.0	394.3
2003.....	15.3	25.2	34.2	39.3	50.6	70.3	96.9	138.8	211.5	411.3
2002.....	15.9	25.7	32.8	39.3	49.5	68.9	96.3	140.8	214.9	424.8
2001.....	14.7	26.0	31.4	37.0	52.2	71.0	97.7	138.1	215.4	440.4
2000.....	15.3	24.9	29.9	38.2	51.5	67.9	96.4	141.1	216.7	448.1
1999.....	15.0	23.7	29.7	37.8	52.0	71.1	98.6	138.2	217.0	444.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of other parts of digestive system (K20-K92)

White, male

2004.....	19.1	33.0	49.9	52.7	62.2	78.3	106.3	151.6	212.6	373.5
2003.....	20.0	35.4	49.1	52.7	66.4	84.9	111.5	152.6	227.7	394.9
2002.....	21.0	36.3	47.4	52.8	63.6	83.8	111.2	156.7	228.3	411.2
2001.....	19.4	36.6	44.6	50.0	67.0	85.5	113.6	151.7	230.0	433.4
2000.....	20.9	36.4	42.0	51.4	66.7	83.4	111.6	160.0	230.4	446.9
1999.....	21.1	34.9	42.5	51.7	67.1	87.7	116.9	154.1	240.4	441.9

White, female

2004.....	9.5	15.7	19.6	25.5	37.3	54.6	83.3	124.2	195.5	403.7
2003.....	10.5	15.1	19.7	26.5	36.0	57.4	85.0	128.9	201.6	418.6
2002.....	10.9	15.1	18.5	26.4	36.4	55.8	84.1	129.5	206.7	430.6
2001.....	10.0	15.5	18.5	24.5	38.7	58.3	84.8	128.5	206.6	443.3
2000.....	9.8	13.4	18.0	25.6	37.4	54.4	84.2	127.8	208.5	448.6
1999.....	8.8	12.7	17.2	24.6	38.0	56.6	84.1	127.1	203.2	445.0

Black, both sexes

2004.....	16.4	30.7	46.0	56.7	66.7	75.0	105.0	130.3	199.3	329.4
2003.....	17.3	30.9	51.6	58.0	69.8	82.9	102.4	137.5	187.3	346.0
2002.....	17.6	35.0	48.3	54.1	65.2	76.7	105.5	136.9	207.7	332.3
2001.....	20.1	36.6	51.1	62.2	68.6	87.7	108.1	150.9	200.2	350.1
2000.....	21.3	37.2	51.0	58.7	72.2	86.2	108.9	147.5	206.1	381.6
1999.....	22.5	40.0	46.6	59.1	74.2	90.3	112.6	149.6	206.9	363.8

Black, male

2004.....	19.4	42.5	64.8	83.4	91.8	102.0	134.0	162.4	229.6	322.9
2003.....	21.2	40.8	76.5	85.3	96.4	107.4	135.7	162.5	237.7	345.1
2002.....	20.9	47.5	69.3	76.8	89.3	99.2	133.2	158.0	240.7	355.9
2001.....	26.6	51.3	75.3	90.1	91.9	118.7	138.0	183.1	215.9	385.8
2000.....	28.0	49.6	73.8	81.0	97.3	109.0	138.4	181.4	241.1	392.1
1999.....	29.9	59.2	68.5	84.6	94.8	112.4	135.0	171.0	238.9	400.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of other parts of digestive system (K20-K92)

Black, female

2004.....	13.7	20.5	30.0	34.6	47.2	54.9	85.2	111.3	183.3	332.0
2003.....	13.8	22.3	30.5	35.5	49.0	64.7	80.0	122.6	160.9	346.4
2002.....	14.6	24.1	30.4	35.4	46.3	59.9	86.9	124.3	190.6	323.2
2001.....	14.3	23.9	30.5	39.4	50.2	64.8	88.1	131.7	192.1	336.8
2000.....	15.4	26.3	31.5	40.5	52.5	69.3	89.2	127.0	188.5	377.7
1999.....	16.1	23.5	28.0	38.2	58.1	73.9	97.7	136.7	191.0	350.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of esophagus (K20-K22)

All races 3/, both sexes

2004.....	0.1	0.2	0.2	0.3	0.6	1.0	1.5	2.8	5.8	17.9
2003.....	0.1	0.1	0.3	0.4	0.6	0.9	1.6	3.1	6.1	19.3
2002.....	0.2	0.3	0.3	0.3	0.5	0.9	1.7	3.5	6.8	20.7
2001.....	*	0.1	0.2	0.4	0.6	1.0	1.8	3.5	7.4	21.2
2000.....	0.1	0.2	0.3	0.4	0.6	1.1	1.8	3.6	7.7	23.3
1999.....	0.1	0.2	0.3	0.4	0.6	1.1	1.9	3.3	7.7	23.5

All races 3/, male

2004.....	0.2	0.2	0.4	0.4	0.8	1.1	1.6	3.8	6.7	19.1
2003.....	*	0.2	0.4	0.5	0.8	1.3	2.1	3.6	7.0	23.6
2002.....	0.2	0.4	0.4	0.3	0.6	1.4	1.9	4.2	8.3	25.4
2001.....	*	*	0.3	0.5	0.9	1.3	2.1	4.1	8.8	23.6
2000.....	0.2	0.3	0.4	0.5	1.0	1.5	2.1	4.6	9.3	28.6
1999.....	*	0.3	0.3	0.4	0.8	1.5	2.7	4.4	9.9	28.3

All races 3/, female

2004.....	*	*	*	*	0.4	0.8	1.4	2.1	5.3	17.3
2003.....	*	*	0.2	0.2	0.3	0.7	1.2	2.8	5.7	17.4
2002.....	*	*	*	0.3	0.4	0.6	1.6	3.0	5.9	18.7
2001.....	*	*	*	*	0.4	0.8	1.5	3.0	6.6	20.2
2000.....	*	*	*	*	*	0.7	1.6	2.8	6.8	21.1
1999.....	*	*	0.2	0.4	0.4	0.8	1.3	2.5	6.4	21.5

White, both sexes

2004.....	0.1	0.1	0.3	0.3	0.5	0.9	1.5	2.8	6.1	18.8
2003.....	0.1	0.2	0.3	0.4	0.6	1.0	1.7	3.2	6.4	20.2
2002.....	0.2	0.3	0.3	0.3	0.5	0.9	1.7	3.4	7.0	21.7
2001.....	*	0.1	0.2	0.4	0.6	1.0	1.7	3.6	7.6	22.2
2000.....	0.1	0.2	0.3	0.4	0.6	1.1	1.8	3.7	8.1	24.4
1999.....	*	0.2	0.2	0.4	0.6	1.1	1.9	3.4	7.9	24.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Peptic ulcer (K25-K28)

All races 3/, both sexes

2004.....	0.3	0.5	0.8	1.0	1.7	2.5	3.9	6.8	11.6	24.4
2003.....	0.3	0.6	0.8	1.2	1.7	2.3	4.2	7.3	13.6	26.4
2002.....	0.3	0.6	0.8	1.4	1.7	2.7	4.7	8.0	13.2	28.3
2001.....	0.4	0.7	0.8	1.3	2.1	3.1	4.8	8.3	16.0	33.4
2000.....	0.3	0.6	1.0	1.5	2.2	3.1	5.1	8.7	15.6	35.6
1999.....	0.4	0.6	0.8	1.4	2.2	3.3	5.9	9.1	17.1	34.6

All races 3/, male

2004.....	0.3	0.7	1.0	1.2	2.2	2.9	4.5	8.3	12.9	24.6
2003.....	0.4	0.8	1.2	1.4	2.3	3.1	5.4	8.8	16.2	28.9
2002.....	0.4	0.7	1.2	1.8	2.1	3.2	6.1	9.3	16.7	31.6
2001.....	0.5	0.9	1.1	1.8	2.8	4.1	5.6	10.3	17.3	34.9
2000.....	0.4	0.9	1.5	2.1	2.9	3.9	6.9	11.2	17.8	40.7
1999.....	0.4	0.7	1.2	1.8	3.0	4.5	7.9	11.0	21.1	37.5

All races 3/, female

2004.....	0.2	0.4	0.5	0.7	1.3	2.2	3.4	5.8	10.8	24.3
2003.....	0.2	0.4	0.5	1.0	1.1	1.7	3.2	6.2	11.9	25.3
2002.....	0.3	0.4	0.5	1.0	1.4	2.2	3.6	7.0	11.0	26.9
2001.....	0.2	0.4	0.6	0.8	1.5	2.2	4.2	6.9	15.2	32.7
2000.....	0.2	0.4	0.5	0.9	1.6	2.5	3.6	6.9	14.4	33.5
1999.....	0.3	0.5	0.5	1.1	1.5	2.3	4.3	7.7	14.7	33.5

White, both sexes

2004.....	0.2	0.4	0.7	0.9	1.7	2.5	3.8	6.8	11.7	25.1
2003.....	0.3	0.5	0.7	1.1	1.6	2.4	4.2	7.2	13.4	26.9
2002.....	0.2	0.5	0.8	1.3	1.6	2.6	4.8	7.9	13.6	29.1
2001.....	0.3	0.6	0.7	1.1	2.0	3.0	4.8	8.3	16.2	34.4
2000.....	0.3	0.6	0.9	1.4	2.2	3.0	5.1	8.5	16.0	36.3
1999.....	0.3	0.5	0.7	1.4	2.2	3.1	5.8	9.1	17.1	35.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Peptic ulcer (K25-K28)

White, male

2004.....	0.3	0.6	1.0	1.2	2.1	2.9	4.3	8.1	12.6	25.5
2003.....	0.3	0.7	1.0	1.3	2.2	3.1	5.3	8.6	15.6	29.0
2002.....	0.3	0.6	1.0	1.6	2.0	3.1	6.0	9.2	17.1	32.5
2001.....	0.5	0.8	0.9	1.5	2.7	3.8	5.3	10.1	17.3	35.7
2000.....	0.4	0.8	1.3	2.0	2.9	3.8	6.7	10.9	17.8	40.7
1999.....	0.3	0.6	1.1	1.6	2.9	4.1	7.8	11.1	21.1	38.4

White, female

2004.....	*	0.3	0.5	0.7	1.3	2.1	3.3	5.8	11.1	25.0
2003.....	0.2	0.3	0.5	0.9	1.0	1.7	3.3	6.2	12.1	26.0
2002.....	*	0.4	0.5	1.0	1.3	2.1	3.8	7.0	11.5	27.6
2001.....	*	0.4	0.6	0.8	1.4	2.2	4.4	7.0	15.6	33.9
2000.....	*	0.3	0.5	0.9	1.6	2.4	3.8	6.9	14.9	34.5
1999.....	0.3	0.4	0.4	1.1	1.5	2.2	4.2	7.7	14.8	34.1

Black, both sexes

2004.....	*	1.3	1.3	1.6	2.6	3.5	5.1	6.6	9.4	16.1
2003.....	*	1.2	1.8	2.1	2.4	2.1	4.5	8.7	14.5	16.9
2002.....	0.9	0.9	1.6	2.3	2.9	3.2	5.2	8.8	6.8	19.0
2001.....	*	1.1	1.6	2.5	3.0	3.6	5.5	7.8	11.1	20.9
2000.....	*	1.4	2.0	2.6	2.7	4.1	5.7	9.4	11.0	26.6
1999.....	1.0	1.2	1.8	2.4	3.2	5.0	6.5	7.7	16.2	25.5

Black, male

2004.....	*	2.1	*	*	3.9	*	*	10.0	*	*
2003.....	*	1.6	3.1	3.1	3.9	*	7.2	11.4	23.6	*
2002.....	*	*	2.8	3.6	4.2	*	8.0	11.9	*	*
2001.....	*	1.8	2.7	4.1	4.5	5.3	9.1	10.7	*	23.8
2000.....	*	*	3.3	3.5	*	*	10.0	12.8	*	31.1
1999.....	1.5	1.9	2.8	3.7	5.3	7.5	8.8	*	18.3	28.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Peptic ulcer (K25-K28)

Black, female

2004.....	*	*	*	*	*	3.6	4.4	*	7.6	16.5
2003.....	*	*	*	*	*	*	*	7.1	9.7	15.8
2002.....	*	*	*	*	*	*	*	6.9	*	18.5
2001.....	*	*	*	*	*	*	*	6.1	9.0	19.8
2000.....	*	*	*	*	*	*	*	7.4	8.5	24.9
1999.....	*	*	*	*	*	*	5.0	6.9	15.1	24.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Gastric ulcer (K25)

All races 3/, both sexes

2004.....	0.1	0.2	0.3	0.3	0.6	0.8	1.1	2.1	3.4	6.7
2003.....	0.1	0.2	0.3	0.5	0.6	0.7	1.1	2.1	3.8	7.0
2002.....	0.1	0.2	0.3	0.5	0.5	0.8	1.4	2.1	3.8	7.7
2001.....	0.1	0.3	0.3	0.4	0.6	1.0	1.5	2.4	4.6	8.0
2000.....	0.1	0.3	0.4	0.5	0.7	1.0	1.5	2.4	4.0	9.5
1999.....	0.2	0.2	0.3	0.5	0.6	0.9	1.8	2.7	4.8	8.8

All races 3/, male

2004.....	*	0.3	0.4	0.4	0.7	0.8	1.2	2.1	4.0	6.3
2003.....	*	0.3	0.5	0.5	0.7	0.9	1.1	2.1	3.9	6.4
2002.....	*	0.2	0.3	0.6	0.8	1.0	1.8	2.2	3.9	8.1
2001.....	0.2	0.3	0.3	0.5	0.7	1.2	1.8	2.7	4.6	6.0
2000.....	*	0.4	0.5	0.5	0.8	1.2	2.0	3.0	3.8	9.9
1999.....	0.2	0.3	0.5	0.6	0.7	1.0	2.2	2.7	5.4	8.7

All races 3/, female

2004.....	*	*	0.3	0.2	0.5	0.8	1.1	2.1	3.1	6.9
2003.....	*	*	*	0.4	0.4	0.6	1.1	2.1	3.7	7.3
2002.....	*	*	*	0.4	*	0.7	1.1	2.0	3.8	7.5
2001.....	*	*	0.2	*	0.6	0.9	1.4	2.2	4.6	8.8
2000.....	*	*	0.2	0.4	0.7	0.9	1.2	2.0	4.1	9.4
1999.....	*	*	*	0.4	0.5	0.8	1.5	2.7	4.4	8.8

White, both sexes

2004.....	*	0.2	0.3	0.3	0.5	0.7	1.1	2.1	3.4	6.8
2003.....	*	0.2	0.2	0.5	0.5	0.7	1.1	2.0	3.7	7.0
2002.....	*	0.2	0.2	0.5	0.5	0.8	1.5	2.0	3.8	7.8
2001.....	*	0.2	0.2	0.3	0.6	0.9	1.5	2.4	4.6	8.0
2000.....	*	0.2	0.3	0.4	0.7	1.0	1.5	2.3	4.0	9.4
1999.....	0.1	0.2	0.3	0.4	0.6	0.7	1.8	2.7	4.8	8.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Duodenal ulcer (K26)

All races 3/, both sexes

2004.....	0.1	0.2	0.2	0.4	0.6	0.9	1.5	2.5	3.9	7.7
2003.....	0.1	0.3	0.3	0.4	0.5	0.8	1.5	2.5	4.6	8.5
2002.....	0.1	0.2	0.3	0.5	0.7	0.9	1.7	3.1	4.3	9.0
2001.....	0.2	0.3	0.3	0.4	0.8	1.1	1.6	2.9	5.4	11.0
2000.....	0.1	0.2	0.4	0.6	0.9	1.0	1.7	3.0	5.6	10.1
1999.....	0.1	0.2	0.3	0.4	1.0	1.2	1.9	2.9	4.9	10.0

All races 3/, male

2004.....	*	0.3	0.3	0.5	0.9	1.1	2.0	3.3	4.2	9.2
2003.....	*	0.4	0.4	0.4	0.8	1.1	2.2	3.4	6.1	10.2
2002.....	*	0.3	0.4	0.8	0.9	1.2	2.2	3.6	6.0	10.1
2001.....	0.3	0.4	0.5	0.6	1.1	1.4	2.0	3.6	6.4	13.8
2000.....	0.2	0.3	0.7	0.8	1.1	1.4	2.3	3.8	6.4	12.3
1999.....	*	0.3	0.5	0.5	1.4	1.9	2.9	4.1	6.7	11.9

All races 3/, female

2004.....	*	*	*	0.3	0.4	0.7	1.1	1.9	3.7	7.1
2003.....	*	*	*	0.3	*	0.5	1.0	1.8	3.6	7.8
2002.....	*	0.2	*	0.3	0.5	0.6	1.3	2.8	3.3	8.5
2001.....	*	*	*	0.3	0.4	0.8	1.3	2.4	4.8	9.8
2000.....	*	*	*	0.3	0.7	0.7	1.3	2.5	5.1	9.2
1999.....	*	*	*	0.3	0.6	0.6	1.2	2.1	3.9	9.3

White, both sexes

2004.....	*	0.2	0.3	0.4	0.6	0.9	1.5	2.6	4.0	8.1
2003.....	*	0.2	0.3	0.3	0.5	0.8	1.6	2.4	4.6	8.8
2002.....	0.1	0.2	0.2	0.5	0.7	0.9	1.8	3.2	4.7	9.5
2001.....	0.2	0.2	0.3	0.4	0.8	1.1	1.6	2.9	5.6	11.6
2000.....	*	0.2	0.4	0.6	0.9	1.0	1.7	3.1	5.8	10.5
1999.....	0.1	0.2	0.3	0.4	1.0	1.2	1.9	3.0	5.1	10.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Peptic ulcer, site unspecified (K27)

All races 3/, both sexes

2004.....	*	0.1	0.2	0.3	0.5	0.8	1.2	2.1	4.1	9.7
2003.....	*	0.1	0.2	0.3	0.6	0.8	1.4	2.7	5.0	10.5
2002.....	*	0.1	0.3	0.3	0.5	0.8	1.5	2.7	4.9	11.3
2001.....	*	0.1	0.2	0.5	0.7	1.0	1.7	2.9	5.8	14.2
2000.....	*	0.2	0.2	0.5	0.6	1.0	1.7	3.1	5.9	15.5
1999.....	*	0.1	0.2	0.5	0.6	1.2	2.1	3.4	7.2	15.6

All races 3/, male

2004.....	*	*	0.3	0.4	0.6	0.9	1.3	2.8	4.5	9.0
2003.....	*	*	0.3	0.4	0.8	1.0	1.8	3.3	5.8	11.9
2002.....	*	*	0.4	0.4	0.5	1.0	2.0	3.4	6.5	12.8
2001.....	*	*	0.3	0.7	0.9	1.4	1.8	3.9	6.2	14.9
2000.....	*	0.2	0.3	0.7	1.0	1.3	2.5	4.3	7.2	18.2
1999.....	*	*	0.3	0.6	0.9	1.6	2.7	4.1	8.8	16.4

All races 3/, female

2004.....	*	*	*	*	0.4	0.7	1.2	1.7	3.8	10.0
2003.....	*	*	*	*	0.3	0.6	1.1	2.2	4.5	9.9
2002.....	*	*	*	0.3	0.5	0.7	1.1	2.2	3.8	10.7
2001.....	*	*	*	0.3	0.4	0.6	1.5	2.2	5.6	13.9
2000.....	*	*	*	*	*	0.8	1.2	2.3	5.1	14.5
1999.....	*	*	*	0.3	0.4	0.9	1.6	2.9	6.3	15.2

White, both sexes

2004.....	*	*	0.2	0.3	0.5	0.8	1.2	2.1	4.1	9.9
2003.....	*	*	0.2	0.3	0.6	0.8	1.4	2.7	4.9	10.7
2002.....	*	0.1	0.3	0.3	0.5	0.8	1.5	2.6	5.0	11.5
2001.....	*	0.2	0.2	0.4	0.6	1.0	1.7	2.9	5.8	14.6
2000.....	*	0.2	0.2	0.5	0.6	1.0	1.8	3.1	6.0	15.9
1999.....	*	*	0.2	0.5	0.6	1.2	2.0	3.4	7.2	16.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Gastrojejunal ulcer (K28)

All races 3/, both sexes

Table with 11 columns (years 2004-1999) and 11 rows of asterisks representing death rates for Gastrojejunal ulcer (K28) in all races 3/, both sexes.

All races 3/, male

Table with 11 columns (years 2004-1999) and 11 rows of asterisks representing death rates for Gastrojejunal ulcer (K28) in all races 3/, male.

All races 3/, female

Table with 11 columns (years 2004-1999) and 11 rows of asterisks representing death rates for Gastrojejunal ulcer (K28) in all races 3/, female.

White, both sexes

Table with 11 columns (years 2004-1999) and 11 rows of asterisks representing death rates for Gastrojejunal ulcer (K28) in White, both sexes.

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Gastritis and duodenitis (K29)

All races 3/, both sexes

2004.....	*	*	*	0.2	0.2	*	0.3	0.7	1.0	2.6
2003.....	*	*	0.1	*	0.2	*	0.4	0.5	1.0	2.9
2002.....	*	*	*	*	*	*	0.3	0.6	1.1	2.9
2001.....	*	*	*	*	*	0.2	0.3	0.6	1.0	3.8
2000.....	*	*	*	0.2	0.2	0.2	0.3	0.6	1.2	3.5
1999.....	*	*	*	*	*	0.2	0.3	0.6	1.5	3.8

All races 3/, male

2004.....	*	*	*	*	0.3	*	*	0.7	1.1	3.0
2003.....	*	*	0.2	*	*	*	*	0.7	1.2	2.8
2002.....	*	*	*	*	*	*	*	0.8	1.0	2.9
2001.....	*	*	*	*	*	*	*	*	*	4.8
2000.....	*	*	*	*	*	*	*	*	1.2	3.3
1999.....	*	*	*	*	*	*	*	0.8	1.7	3.7

All races 3/, female

2004.....	*	*	*	*	*	*	*	0.6	0.9	2.4
2003.....	*	*	*	*	*	*	*	*	0.9	3.0
2002.....	*	*	*	*	*	*	*	0.5	1.1	2.9
2001.....	*	*	*	*	*	*	*	0.5	1.0	3.3
2000.....	*	*	*	*	*	*	*	0.6	1.2	3.6
1999.....	*	*	*	*	*	*	*	*	1.4	3.9

White, both sexes

2004.....	*	*	*	0.2	*	*	0.3	0.6	0.8	2.6
2003.....	*	*	0.1	*	*	*	0.4	0.4	0.9	2.9
2002.....	*	*	*	*	*	*	0.3	0.7	1.0	2.9
2001.....	*	*	*	*	*	*	0.3	0.6	0.9	4.0
2000.....	*	*	*	0.2	*	*	0.3	0.6	1.3	3.5
1999.....	*	*	*	*	*	*	0.3	0.5	1.3	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Dyspepsia and other diseases of stomach and duodenum (K30-K31)

All races 3/, both sexes

2004.....	*	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.7	3.0
2003.....	0.1	*	0.1	0.2	0.2	0.4	0.7	1.0	1.9	3.5
2002.....	0.1	*	0.1	0.2	0.3	0.4	0.6	1.1	1.7	3.2
2001.....	*	*	0.2	*	0.2	0.4	0.6	1.0	1.7	3.6
2000.....	*	*	0.2	0.2	0.2	0.5	0.7	1.2	1.6	4.0
1999.....	*	0.1	*	*	0.3	0.3	0.6	1.0	1.9	4.2

All races 3/, male

2004.....	*	*	*	*	*	0.5	0.6	1.2	2.0	3.6
2003.....	*	*	*	*	*	*	0.9	1.1	2.0	2.7
2002.....	*	*	*	*	*	0.6	0.7	1.3	1.6	2.9
2001.....	*	*	*	*	*	0.5	0.6	0.9	2.2	3.3
2000.....	*	*	*	*	*	0.7	0.9	1.2	1.7	4.3
1999.....	*	*	*	*	*	*	0.6	1.2	1.6	4.8

All races 3/, female

2004.....	*	*	0.2	*	0.3	*	0.5	1.1	1.5	2.8
2003.....	*	*	*	*	*	0.5	0.6	1.0	1.8	3.9
2002.....	*	*	*	0.3	*	*	0.5	1.1	1.8	3.4
2001.....	*	*	*	*	*	*	0.6	1.0	1.5	3.7
2000.....	*	*	*	*	*	*	0.6	1.1	1.5	3.9
1999.....	*	*	*	*	0.4	*	0.6	0.9	2.1	4.0

White, both sexes

2004.....	*	*	0.2	0.2	0.3	0.4	0.5	1.2	1.8	3.0
2003.....	*	*	0.1	0.2	0.2	0.4	0.8	0.9	1.8	3.6
2002.....	*	*	0.1	0.2	0.3	0.4	0.6	1.2	1.6	3.3
2001.....	*	*	0.2	*	*	0.4	0.6	0.9	1.8	3.7
2000.....	*	*	0.2	0.2	0.2	0.4	0.7	1.1	1.6	4.1
1999.....	*	*	*	*	0.3	0.3	0.6	1.1	1.9	4.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
 years years years years years years years years years and over

Diseases of appendix (K35-K38)

All races 3/, both sexes

2004.....	*	*	0.2	0.1	0.2	0.3	0.6	0.8	1.4	1.7
2003.....	*	0.1	0.1	*	0.2	0.3	0.4	0.8	1.4	1.7
2002.....	*	0.1	0.2	0.2	0.3	0.3	0.7	0.8	1.1	2.0
2001.....	*	0.1	*	0.2	0.2	0.4	0.6	0.5	1.0	2.1
2000.....	*	*	0.1	0.2	0.3	0.3	0.5	0.7	1.4	1.6
1999.....	*	0.1	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.5

All races 3/, male

2004.....	*	*	0.2	*	*	*	0.7	1.1	1.7	2.5
2003.....	*	*	*	*	*	0.4	0.7	0.9	2.3	2.4
2002.....	*	*	0.2	*	0.5	0.5	0.8	1.1	1.6	2.2
2001.....	*	*	*	*	*	0.5	0.9	0.7	1.6	2.8
2000.....	*	*	*	0.3	0.4	*	0.6	1.0	1.6	2.0
1999.....	*	*	*	*	*	*	0.5	0.9	*	2.9

All races 3/, female

2004.....	*	*	*	*	*	*	0.5	0.6	1.2	1.4
2003.....	*	*	*	*	*	*	*	0.6	0.9	1.3
2002.....	*	*	*	*	*	*	0.5	0.6	0.8	2.0
2001.....	*	*	*	*	*	*	0.4	*	0.7	1.8
2000.....	*	*	*	*	*	*	0.4	0.5	1.2	1.5
1999.....	*	*	*	*	*	*	*	*	1.1	0.9

White, both sexes

2004.....	*	*	0.1	0.2	0.2	0.3	0.6	0.8	1.3	1.8
2003.....	*	*	*	*	*	0.3	0.4	0.7	1.5	1.7
2002.....	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.2	2.1
2001.....	*	*	*	*	*	0.3	0.7	0.5	0.9	2.1
2000.....	*	*	*	0.2	0.3	0.2	0.4	0.6	1.3	1.7
1999.....	*	*	0.2	*	*	0.3	0.4	0.6	1.0	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
 years years years years years years years years years and over

Hernia (K40-K46)

All races 3/, both sexes

2004.....	0.1	0.2	0.2	0.4	0.5	0.9	1.8	2.9	5.3	11.0
2003.....	0.1	0.2	0.3	0.5	0.7	1.0	1.6	2.7	4.8	11.9
2002.....	0.1	0.2	0.2	0.4	0.6	0.9	1.6	2.8	5.3	12.8
2001.....	0.1	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.6
2000.....	*	0.1	0.3	0.4	0.6	0.9	1.4	3.0	4.9	13.1
1999.....	0.1	*	0.2	0.3	0.7	0.8	1.6	2.5	5.4	12.7

All races 3/, male

2004.....	*	0.2	0.3	0.4	0.5	0.8	2.2	3.2	4.7	11.3
2003.....	*	0.2	0.4	0.4	0.7	1.1	1.6	3.0	5.2	12.3
2002.....	*	*	*	0.4	0.5	0.9	1.6	2.7	5.2	11.7
2001.....	*	*	0.3	0.3	0.8	0.9	1.4	2.3	5.3	12.0
2000.....	*	*	0.4	0.3	0.6	0.9	1.4	3.0	5.7	12.6
1999.....	*	*	0.2	0.3	0.8	0.8	1.5	2.5	6.5	13.2

All races 3/, female

2004.....	*	0.2	0.2	0.4	0.5	1.0	1.4	2.7	5.6	10.8
2003.....	*	0.2	0.3	0.6	0.7	1.0	1.5	2.5	4.6	11.8
2002.....	*	*	0.3	0.4	0.7	1.0	1.5	2.9	5.3	13.3
2001.....	*	*	0.3	0.5	0.6	0.8	1.6	2.6	4.8	12.8
2000.....	*	*	0.3	0.4	0.6	0.9	1.5	3.0	4.4	13.3
1999.....	*	*	*	*	0.6	0.8	1.6	2.6	4.8	12.6

White, both sexes

2004.....	*	0.2	0.2	0.4	0.5	0.9	1.8	3.1	5.4	11.2
2003.....	*	0.2	0.3	0.4	0.7	1.1	1.6	2.8	4.7	12.3
2002.....	*	0.1	0.2	0.4	0.6	0.9	1.6	3.0	5.4	13.4
2001.....	*	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.8
2000.....	*	0.1	0.3	0.4	0.6	0.8	1.5	3.0	5.0	13.3
1999.....	*	*	0.2	0.3	0.7	0.7	1.6	2.5	5.5	13.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Crohn's disease and ulcerative colitis (K50-K51)

All races 3/, both sexes

2004.....	0.1	0.2	0.3	0.4	0.6	0.7	1.1	1.8	2.4	3.3
2003.....	0.1	0.2	0.3	0.4	0.5	0.8	1.1	1.6	2.3	3.1
2002.....	0.2	0.2	0.3	0.3	0.4	0.7	1.0	1.7	2.3	3.1
2001.....	*	0.2	0.2	0.5	0.5	0.7	1.1	1.6	2.4	3.8
2000.....	*	0.1	0.2	0.4	0.4	0.6	1.1	1.6	2.2	3.3
1999.....	0.1	0.2	0.3	0.3	0.4	0.6	1.1	1.6	2.5	3.3

All races 3/, male

2004.....	*	0.2	0.3	0.4	0.7	0.7	1.1	1.8	2.2	2.7
2003.....	*	0.2	0.2	0.4	0.5	0.7	1.1	1.7	2.3	3.5
2002.....	0.2	0.2	*	*	*	0.8	1.0	1.7	2.3	2.7
2001.....	*	*	0.2	0.5	0.5	0.7	1.3	1.6	2.3	3.5
2000.....	*	*	*	0.3	*	0.6	1.1	1.6	1.6	2.9
1999.....	*	*	0.3	0.3	*	0.7	1.1	1.6	2.5	2.7

All races 3/, female

2004.....	*	0.2	0.3	0.4	0.5	0.7	1.1	1.8	2.6	3.6
2003.....	*	0.3	0.3	0.3	0.5	0.9	1.0	1.5	2.3	2.8
2002.....	0.2	0.2	0.3	0.3	0.5	0.7	1.0	1.7	2.3	3.2
2001.....	*	*	0.2	0.5	0.4	0.6	0.9	1.5	2.5	3.9
2000.....	*	*	0.3	0.4	0.4	0.7	1.2	1.6	2.6	3.4
1999.....	*	0.3	0.3	0.3	0.5	0.5	1.1	1.6	2.6	3.5

White, both sexes

2004.....	0.1	0.2	0.3	0.4	0.6	0.7	1.1	2.0	2.6	3.6
2003.....	0.1	0.2	0.3	0.4	0.5	0.8	1.1	1.8	2.4	3.3
2002.....	0.2	0.2	0.3	0.3	0.5	0.8	1.0	1.8	2.5	3.4
2001.....	*	0.2	0.2	0.5	0.5	0.7	1.2	1.7	2.5	3.9
2000.....	*	0.2	0.2	0.4	0.4	0.7	1.2	1.7	2.4	3.5
1999.....	0.2	0.2	0.3	0.3	0.4	0.7	1.2	1.6	2.8	3.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

All races 3/, both sexes

2004.....	0.6	0.8	1.5	2.6	4.8	8.7	16.1	29.5	49.9	106.9
2003.....	0.5	0.9	1.6	2.9	4.8	9.7	17.4	29.7	52.3	111.5
2002.....	0.5	0.9	1.4	2.8	4.8	9.0	17.0	29.6	53.0	113.3
2001.....	0.4	0.8	1.6	2.7	5.4	10.1	17.8	29.7	52.1	113.5
2000.....	0.5	0.9	1.7	2.8	5.3	9.5	17.5	31.5	52.9	116.2
1999.....	0.5	0.8	1.5	2.9	5.3	10.1	17.1	29.9	52.5	116.2

All races 3/, male

2004.....	0.6	0.8	1.5	2.4	4.5	7.9	14.6	28.0	45.1	88.8
2003.....	0.5	0.8	1.6	3.1	4.5	8.8	16.2	26.8	48.8	92.6
2002.....	0.6	0.8	1.5	2.5	4.7	8.1	15.8	26.8	49.7	96.7
2001.....	0.5	0.8	1.6	2.3	5.1	9.1	17.6	27.2	47.9	98.5
2000.....	0.4	0.9	1.6	2.7	5.2	8.9	16.6	30.0	48.2	98.5
1999.....	0.6	0.8	1.4	2.9	4.9	8.7	16.7	27.9	48.7	95.6

All races 3/, female

2004.....	0.5	0.9	1.4	2.7	5.1	9.5	17.3	30.5	53.0	115.1
2003.....	0.5	1.0	1.6	2.7	5.1	10.5	18.3	31.8	54.5	119.8
2002.....	0.4	0.9	1.4	3.0	4.9	9.8	18.0	31.6	55.0	120.5
2001.....	0.4	0.9	1.6	3.0	5.6	11.0	18.0	31.5	54.6	119.8
2000.....	0.5	0.9	1.8	2.8	5.4	10.0	18.2	32.5	55.7	123.4
1999.....	0.5	0.7	1.6	3.0	5.7	11.3	17.5	31.3	54.8	124.5

White, both sexes

2004.....	0.5	0.7	1.4	2.3	4.6	8.7	16.0	29.2	50.2	108.5
2003.....	0.4	0.8	1.4	2.6	4.5	9.5	17.2	29.4	52.8	113.7
2002.....	0.5	0.8	1.3	2.6	4.6	8.7	16.4	29.8	53.0	115.9
2001.....	0.4	0.8	1.3	2.4	5.0	9.7	17.7	29.1	52.9	116.2
2000.....	0.4	0.7	1.5	2.6	4.8	9.2	17.3	31.1	52.9	118.0
1999.....	0.4	0.6	1.4	2.9	5.3	9.7	16.5	29.2	52.9	118.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

White, male										
2004.....	0.6	0.7	1.4	2.2	4.2	7.6	13.8	27.0	44.1	88.6
2003.....	0.4	0.7	1.4	2.6	4.0	8.5	15.2	26.0	48.0	92.9
2002.....	0.5	0.7	1.3	2.3	4.2	7.8	14.9	26.7	48.7	97.5
2001.....	0.4	0.7	1.3	2.1	4.8	8.4	17.3	26.2	48.3	98.9
2000.....	0.3	0.6	1.4	2.4	4.5	8.2	16.2	29.0	47.0	98.9
1999.....	0.5	0.7	1.2	2.8	4.7	8.2	16.0	27.3	47.9	94.8
White, female										
2004.....	0.4	0.8	1.3	2.5	5.1	9.7	17.7	30.8	54.0	117.5
2003.....	0.5	1.0	1.4	2.6	4.9	10.5	18.9	31.9	55.8	122.9
2002.....	0.4	0.8	1.2	2.9	4.9	9.5	17.7	32.0	55.6	123.9
2001.....	0.4	0.8	1.4	2.7	5.2	10.9	18.0	31.1	55.7	123.4
2000.....	0.5	0.7	1.5	2.8	5.1	9.9	18.2	32.5	56.4	125.8
1999.....	0.4	0.6	1.5	2.9	5.8	11.0	16.8	30.6	55.9	127.4
Black, both sexes										
2004.....	1.1	1.6	2.5	5.0	7.4	10.5	20.8	37.7	56.7	106.0
2003.....	1.0	1.5	3.5	5.4	8.2	12.9	22.4	37.6	54.3	102.1
2002.....	1.0	1.8	3.0	4.6	7.2	13.3	25.3	32.3	63.3	99.8
2001.....	0.8	1.8	3.9	5.1	10.2	15.4	24.0	40.1	53.6	96.7
2000.....	0.9	2.4	4.0	5.4	10.4	14.3	21.7	40.7	59.0	112.0
1999.....	1.4	1.8	3.2	4.3	7.4	15.9	26.1	42.3	54.9	108.7
Black, male										
2004.....	*	1.7	2.4	5.0	9.3	12.4	24.9	44.6	66.0	107.3
2003.....	*	1.9	4.1	7.3	9.4	14.3	29.9	41.1	67.9	101.1
2002.....	*	1.9	3.5	4.8	9.5	12.5	27.4	30.3	71.8	105.1
2001.....	*	1.8	4.0	5.2	9.6	18.8	26.9	42.7	57.0	120.3
2000.....	*	3.0	4.1	6.9	13.0	16.4	23.4	45.8	70.8	110.7
1999.....	1.8	2.3	3.3	4.6	8.3	15.3	25.1	39.9	68.0	110.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

Black, female

2004.....	1.3	1.5	2.6	5.0	6.0	9.2	18.0	33.7	51.8	105.5
2003.....	*	*	3.0	3.8	7.2	11.9	17.3	35.5	47.2	102.5
2002.....	*	1.8	2.6	4.5	5.4	13.9	24.0	33.5	58.9	97.8
2001.....	*	1.8	3.8	5.1	10.7	13.0	22.0	38.6	51.9	88.0
2000.....	*	1.8	4.0	4.2	8.3	12.7	20.6	37.6	53.1	112.5
1999.....	*	*	3.0	4.2	6.7	16.4	26.8	43.8	48.4	108.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diverticular disease of intestine (K57)

All races 3/, both sexes

2004.....	0.1	*	0.3	0.5	1.1	1.8	3.8	6.5	13.0	27.0
2003.....	0.1	0.1	0.3	0.6	1.1	1.9	3.6	7.3	13.3	28.9
2002.....	*	0.2	0.3	0.5	0.9	1.9	3.5	7.3	13.4	27.7
2001.....	*	0.2	0.3	0.6	1.3	2.2	3.6	7.0	12.8	31.4
2000.....	0.1	0.2	0.3	0.4	1.1	2.2	3.7	6.5	13.3	30.9
1999.....	*	0.2	0.3	0.7	1.1	2.2	4.1	6.9	12.8	33.4

All races 3/, male

2004.....	*	*	0.4	0.5	1.1	1.6	3.3	5.8	10.7	18.8
2003.....	*	*	0.3	0.5	1.1	1.7	3.2	5.7	10.3	22.3
2002.....	*	0.3	0.4	0.6	0.9	1.7	3.1	6.5	10.4	19.9
2001.....	*	0.3	0.3	0.6	1.1	1.6	3.2	6.6	10.7	23.4
2000.....	*	0.2	0.3	0.6	1.0	2.1	3.4	5.2	10.8	22.3
1999.....	*	*	0.2	0.6	1.2	1.9	3.7	5.5	10.4	25.8

All races 3/, female

2004.....	*	*	0.2	0.5	1.1	2.0	4.3	7.0	14.4	30.6
2003.....	*	*	0.3	0.7	1.2	2.1	4.0	8.4	15.1	31.9
2002.....	*	*	*	0.5	1.0	2.0	3.8	7.9	15.2	31.0
2001.....	*	*	0.2	0.6	1.4	2.7	3.9	7.3	14.1	34.7
2000.....	*	*	0.3	0.3	1.2	2.4	4.0	7.4	14.8	34.4
1999.....	*	*	0.3	0.8	1.0	2.6	4.3	7.9	14.1	36.4

White, both sexes

2004.....	*	*	0.3	0.6	1.1	1.8	4.0	6.8	13.4	27.9
2003.....	*	0.1	0.3	0.5	1.1	2.0	3.7	7.5	13.7	30.3
2002.....	*	0.2	0.3	0.5	0.9	1.9	3.6	7.4	13.7	28.5
2001.....	*	0.2	0.3	0.6	1.3	2.0	3.7	7.2	13.5	32.4
2000.....	*	0.2	0.3	0.4	1.0	2.2	3.8	6.7	13.7	32.3
1999.....	*	0.2	0.3	0.7	1.1	2.2	4.2	7.2	13.1	34.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diverticular disease of intestine (K57)

Black, female

2004.....	*	*	*	*	*	*	4.4	*	10.2	23.2
2003.....	*	*	*	*	*	*	*	7.6	10.8	20.3
2002.....	*	*	*	*	*	*	*	6.0	13.1	24.6
2001.....	*	*	*	*	*	4.7	4.7	6.1	*	27.8
2000.....	*	*	*	*	*	*	*	6.0	12.3	24.5
1999.....	*	*	*	*	*	4.2	*	*	11.7	27.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of intestines and peritoneum (K52,K58-K63)

All races 3/, both sexes

2004.....	0.2	0.4	0.7	1.0	1.9	3.2	5.7	9.4	17.3	33.3
2003.....	0.2	0.4	0.7	1.2	2.1	3.3	5.5	9.8	16.5	32.8
2002.....	0.3	0.4	0.7	1.2	2.1	3.5	5.2	9.9	16.2	32.8
2001.....	0.3	0.4	0.7	1.1	1.9	3.5	5.6	10.1	16.8	34.1
2000.....	0.3	0.5	0.6	1.2	2.0	3.0	5.3	9.5	16.9	33.3
1999.....	0.3	0.4	0.7	1.2	1.9	3.4	5.6	9.7	15.7	32.0

All races 3/, male

2004.....	*	0.4	0.7	1.0	1.5	2.9	5.5	9.8	16.5	26.4
2003.....	*	0.4	0.8	1.2	2.2	3.0	5.5	10.1	15.7	27.1
2002.....	0.3	0.5	0.7	1.2	2.0	3.4	5.5	10.5	13.8	29.4
2001.....	0.2	0.4	0.7	1.0	1.9	3.2	5.4	9.9	15.6	29.9
2000.....	0.3	0.4	0.5	1.3	2.1	2.7	5.6	9.1	15.1	28.7
1999.....	0.3	0.6	0.7	1.2	1.8	3.1	5.5	9.8	15.5	28.9

All races 3/, female

2004.....	0.3	0.3	0.7	1.1	2.2	3.4	6.0	9.1	17.8	36.4
2003.....	0.3	0.3	0.7	1.1	2.0	3.6	5.4	9.6	17.0	35.2
2002.....	0.3	0.4	0.7	1.2	2.3	3.5	5.1	9.4	17.7	34.3
2001.....	0.3	0.5	0.8	1.1	2.0	3.7	5.7	10.2	17.5	35.8
2000.....	0.2	0.5	0.8	1.1	1.8	3.3	5.1	9.7	17.9	35.2
1999.....	0.3	0.3	0.6	1.2	2.1	3.6	5.7	9.7	15.8	33.2

White, both sexes

2004.....	0.2	0.3	0.7	1.0	1.8	3.1	5.8	9.6	17.8	34.4
2003.....	0.2	0.4	0.7	1.1	2.0	3.2	5.4	10.0	17.0	33.8
2002.....	0.3	0.4	0.7	1.2	2.1	3.6	5.1	10.1	16.7	34.1
2001.....	0.3	0.4	0.7	0.9	1.8	3.5	5.7	10.0	16.8	35.2
2000.....	0.2	0.5	0.6	1.2	1.8	3.0	5.4	9.6	17.2	33.7
1999.....	0.3	0.4	0.6	1.2	1.9	3.3	5.6	9.8	15.9	33.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of intestines and peritoneum (K52,K58-K63)

White, male

2004.....	*	0.4	0.7	0.9	1.5	2.6	5.5	9.8	16.8	26.9
2003.....	*	0.4	0.7	1.1	2.1	3.0	5.4	10.1	15.9	27.5
2002.....	0.3	0.5	0.7	1.2	1.9	3.4	5.1	10.7	13.7	30.3
2001.....	0.2	0.4	0.7	0.9	1.7	3.0	5.1	9.8	15.7	31.2
2000.....	0.3	0.4	0.4	1.2	1.8	2.7	5.7	9.1	15.2	28.1
1999.....	0.3	0.5	0.7	1.2	1.7	3.1	5.6	9.6	15.7	28.3

White, female

2004.....	0.2	0.3	0.7	1.1	2.1	3.5	6.1	9.4	18.4	37.8
2003.....	0.3	0.3	0.6	1.2	1.8	3.4	5.5	9.8	17.7	36.5
2002.....	0.3	0.4	0.7	1.2	2.2	3.8	5.1	9.7	18.5	35.8
2001.....	0.3	0.4	0.8	1.0	1.9	3.9	6.1	10.2	17.4	36.8
2000.....	*	0.5	0.7	1.1	1.8	3.2	5.1	10.0	18.4	36.1
1999.....	0.2	0.3	0.6	1.2	2.1	3.6	5.7	9.9	16.0	34.9

Black, both sexes

2004.....	*	*	*	1.5	3.1	4.3	6.5	9.1	14.9	24.0
2003.....	*	*	1.5	1.8	3.5	5.5	6.6	10.4	14.7	25.7
2002.....	*	*	1.0	1.7	3.3	3.3	7.8	8.9	13.9	23.2
2001.....	*	0.8	1.1	2.3	2.7	4.1	6.1	13.0	19.8	27.0
2000.....	*	*	1.1	1.8	3.5	3.4	6.1	8.0	15.0	29.7
1999.....	*	0.9	1.2	1.5	2.3	4.3	6.4	8.9	16.2	23.6

Black, male

2004.....	*	*	*	*	*	6.1	7.4	11.8	15.8	24.8
2003.....	*	*	*	*	*	4.9	7.9	11.9	16.2	25.0
2002.....	*	*	*	*	*	5.0	10.7	10.6	16.0	24.7
2001.....	*	*	*	*	*	5.8	8.8	15.8	16.9	*
2000.....	*	*	*	*	4.1	*	*	10.9	17.7	30.0
1999.....	*	*	*	*	*	*	*	10.4	17.4	39.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of intestines and peritoneum (K52,K58-K63)

Black, female

2004.....	*	*	*	*	3.5	*	5.9	7.5	14.4	23.6
2003.....	*	*	*	*	3.5	5.8	5.8	9.5	13.9	25.9
2002.....	*	*	*	*	3.4	*	5.8	8.0	12.7	22.6
2001.....	*	*	*	*	*	*	*	11.4	21.3	29.5
2000.....	*	*	*	*	*	4.0	6.4	6.3	13.6	29.6
1999.....	*	*	*	*	*	4.2	7.2	8.0	15.6	17.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of peritoneum (K65-K66)										
All races 3/, both sexes										
2004.....	0.2	0.4	0.4	0.6	0.9	1.4	2.1	3.4	5.1	8.6
2003.....	0.2	0.4	0.6	0.6	1.1	1.7	2.2	3.4	5.1	9.2
2002.....	0.2	0.4	0.6	0.7	1.0	1.3	2.3	3.7	6.2	10.3
2001.....	0.2	0.4	0.5	0.8	0.9	1.4	2.3	3.6	5.7	10.4
2000.....	0.3	0.4	0.5	0.9	1.2	1.4	2.7	3.9	5.3	10.6
1999.....	0.3	0.4	0.5	0.8	1.1	1.6	2.8	3.7	5.5	9.7
All races 3/, male										
2004.....	0.2	0.4	0.6	0.6	0.9	1.7	2.3	3.4	5.9	7.7
2003.....	0.3	0.3	0.6	0.7	1.2	1.7	2.3	3.7	5.7	8.7
2002.....	0.2	0.4	0.7	0.8	1.1	1.3	2.0	3.7	5.1	9.1
2001.....	0.3	0.5	0.6	0.9	1.0	1.7	2.2	3.6	6.2	10.1
2000.....	0.3	0.4	0.5	1.0	1.4	1.5	2.7	4.3	5.2	11.0
1999.....	0.3	0.5	0.7	0.8	1.3	1.7	3.1	4.1	6.3	10.1
All races 3/, female										
2004.....	0.2	0.3	0.3	0.6	0.8	1.2	1.9	3.4	4.7	9.0
2003.....	*	0.4	0.5	0.5	0.9	1.6	2.2	3.3	4.6	9.4
2002.....	0.2	0.3	0.5	0.6	1.0	1.3	2.5	3.7	6.8	10.8
2001.....	0.2	0.4	0.5	0.6	0.8	1.1	2.3	3.5	5.4	10.5
2000.....	0.3	0.3	0.5	0.9	1.0	1.4	2.6	3.6	5.3	10.5
1999.....	0.2	0.2	0.4	0.8	0.9	1.5	2.5	3.5	5.1	9.6
White, both sexes										
2004.....	0.2	0.3	0.4	0.6	0.8	1.4	2.0	3.5	5.0	8.8
2003.....	0.2	0.3	0.5	0.5	0.9	1.6	2.1	3.5	5.1	9.1
2002.....	0.2	0.3	0.5	0.7	0.9	1.2	2.2	3.8	6.1	10.5
2001.....	0.2	0.4	0.5	0.6	0.7	1.3	2.2	3.5	5.6	10.4
2000.....	0.2	0.3	0.5	0.8	1.1	1.3	2.6	3.8	5.2	10.6
1999.....	0.2	0.4	0.5	0.7	1.0	1.4	2.7	3.5	5.6	9.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of peritoneum (K65-K66)

Black, female

2004.....	*	*	*	*	*	*	*	*	7.6	7.9
2003.....	*	*	*	*	*	*	*	*	*	12.6
2002.....	*	*	*	*	*	*	*	*	8.0	11.1
2001.....	*	*	*	*	*	*	*	*	8.2	10.5
2000.....	*	*	*	*	*	*	*	*	*	10.7
1999.....	*	*	*	*	*	*	4.5	*	*	12.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of liver (K70-K76)

All races 3/, both sexes

2004.....	11.1	19.3	27.1	28.2	31.1	34.5	39.6	41.4	38.2	32.8
2003.....	11.9	20.2	27.1	28.5	32.1	37.3	40.4	41.5	40.2	33.6
2002.....	12.3	20.7	25.7	27.5	31.2	36.0	40.3	41.9	41.9	35.7
2001.....	11.9	21.1	25.0	26.4	31.7	36.6	39.7	42.2	40.2	36.1
2000.....	12.4	20.4	23.3	26.9	32.3	36.3	39.7	41.5	40.7	36.7
1999.....	12.1	19.9	23.0	26.6	32.9	38.0	40.9	42.4	41.9	37.2

All races 3/, male

2004.....	15.0	27.0	40.5	41.0	42.6	45.3	50.3	51.3	46.8	40.7
2003.....	15.8	29.2	40.8	40.8	45.7	49.6	51.2	50.0	49.1	44.4
2002.....	16.3	29.8	38.7	39.7	43.1	47.8	51.6	51.0	52.8	47.3
2001.....	16.0	30.6	37.2	38.3	44.4	49.4	51.0	51.2	50.4	49.4
2000.....	17.2	30.5	34.8	38.2	45.2	48.4	51.1	52.8	48.1	52.2
1999.....	17.4	30.2	34.9	39.4	45.5	52.4	52.6	51.1	53.5	52.3

All races 3/, female

2004.....	7.3	11.9	14.3	16.1	20.6	25.1	31.0	34.2	33.0	29.3
2003.....	8.1	11.4	14.1	16.8	19.8	26.6	31.7	35.5	34.8	28.9
2002.....	8.4	11.8	13.3	15.9	20.4	25.8	31.3	35.5	35.2	30.7
2001.....	7.9	11.8	13.3	15.3	20.3	25.5	30.7	35.9	34.1	30.5
2000.....	7.7	10.6	12.4	16.3	20.7	25.9	30.8	33.6	36.4	30.4
1999.....	6.9	10.0	11.6	14.6	21.5	25.8	31.9	36.3	35.0	31.1

White, both sexes

2004.....	11.3	19.6	27.2	27.7	31.1	35.3	41.0	43.2	39.3	33.4
2003.....	12.2	20.4	26.7	28.2	32.1	38.1	41.8	43.4	41.7	34.3
2002.....	12.6	20.5	25.3	27.2	31.4	37.0	41.9	43.6	42.9	36.6
2001.....	11.8	20.9	24.2	25.6	32.1	37.4	40.6	43.4	41.4	37.0
2000.....	12.3	20.2	22.3	26.2	32.3	36.8	40.7	42.9	41.6	37.5
1999.....	11.7	19.0	22.5	25.6	32.3	38.4	41.6	43.4	43.2	37.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of liver (K70-K76)

White, male

2004.....	15.3	27.2	40.8	40.1	42.5	46.0	51.7	53.2	48.3	42.6
2003.....	16.5	29.7	40.0	40.0	45.6	50.4	52.7	52.1	50.6	45.1
2002.....	16.9	30.0	38.2	39.3	43.2	48.8	53.3	53.0	53.9	48.5
2001.....	15.8	30.6	36.1	36.9	44.5	50.2	52.1	52.3	51.8	50.8
2000.....	17.2	30.7	33.3	37.4	45.3	49.2	52.0	54.3	49.8	53.3
1999.....	17.1	29.1	34.1	38.0	45.2	52.7	53.4	52.3	55.7	53.8

White, female

2004.....	7.2	12.0	13.8	15.8	20.5	25.8	32.2	35.9	33.7	29.3
2003.....	7.9	11.1	13.7	16.8	19.7	27.3	32.9	37.1	36.3	29.5
2002.....	8.3	11.1	12.8	15.6	20.4	26.6	32.5	36.9	36.1	31.4
2001.....	7.8	11.3	12.6	14.9	20.7	26.2	31.4	37.2	35.1	31.2
2000.....	7.3	9.8	11.6	15.6	20.3	26.0	31.5	34.8	36.8	31.1
1999.....	6.4	9.1	11.1	13.8	20.5	26.0	32.3	37.1	35.8	31.2

Black, both sexes

2004.....	10.7	21.0	31.6	35.8	35.0	32.9	31.7	23.9	27.2	24.3
2003.....	10.7	20.9	35.3	35.6	35.8	36.7	32.4	24.5	23.6	27.1
2002.....	11.2	24.4	33.4	34.0	33.9	32.6	30.4	30.1	28.8	27.6
2001.....	13.6	25.5	34.4	37.6	33.0	36.0	34.0	33.0	29.0	21.8
2000.....	14.2	25.2	34.8	34.5	36.0	36.7	35.6	29.1	28.2	26.0
1999.....	15.6	29.0	30.9	36.9	41.6	38.1	36.5	34.5	28.0	31.2

Black, male

2004.....	14.0	29.9	46.4	56.5	50.8	47.5	46.6	33.2	33.7	19.9
2003.....	12.4	29.1	55.5	56.0	54.6	53.7	44.7	31.0	34.7	40.7
2002.....	13.4	32.7	50.1	51.5	49.9	47.1	41.7	36.7	39.0	36.5
2001.....	18.6	36.9	53.4	56.8	50.6	53.4	48.1	44.1	37.7	27.2
2000.....	18.5	34.6	53.4	50.7	50.5	51.0	50.5	41.1	29.5	35.8
1999.....	21.2	43.5	47.7	55.8	56.4	55.8	50.5	42.3	33.1	37.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of liver (K70-K76)

Black, female

2004.....	7.9	13.3	19.0	18.8	22.6	22.1	21.5	18.3	23.8	26.0
2003.....	9.2	13.7	18.3	18.9	21.2	24.1	24.1	20.6	17.8	21.9
2002.....	9.4	17.1	19.3	19.6	21.4	21.8	22.9	26.1	23.5	24.2
2001.....	9.2	15.7	18.3	21.8	19.0	23.1	24.7	26.3	24.5	19.8
2000.....	10.4	17.0	18.8	21.2	24.5	26.1	25.8	21.9	27.6	22.3
1999.....	10.6	16.4	16.7	21.5	30.0	25.0	27.2	29.8	25.5	28.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alcoholic liver disease (K70)

All races 3/, both sexes

2004.....	5.8	9.1	12.2	11.8	11.5	10.8	9.0	6.5	4.1	2.1
2003.....	5.9	9.4	11.8	11.6	11.6	11.0	8.3	6.6	4.4	1.8
2002.....	5.9	9.4	11.4	11.4	11.8	10.7	9.5	6.7	4.4	2.1
2001.....	6.1	10.2	11.3	11.1	12.0	11.1	9.0	6.9	3.9	2.4
2000.....	6.6	10.0	10.7	11.7	12.9	11.4	9.1	7.2	4.4	2.2
1999.....	6.4	10.0	10.9	11.5	12.6	11.6	9.4	7.3	5.2	2.8

All races 3/, male

2004.....	8.0	12.9	18.6	18.2	18.3	17.8	15.0	11.5	8.0	4.6
2003.....	8.0	14.2	18.3	18.1	18.6	17.7	14.2	11.3	8.2	3.6
2002.....	8.0	13.8	17.7	17.6	19.2	17.4	15.4	10.9	8.0	4.7
2001.....	8.4	15.3	17.0	18.0	19.0	17.9	15.2	12.1	7.0	5.5
2000.....	9.4	15.3	16.9	18.0	20.3	18.7	15.7	13.8	7.7	5.3
1999.....	9.4	15.5	17.2	18.7	20.0	19.7	16.2	12.1	9.7	6.1

All races 3/, female

2004.....	3.6	5.5	6.0	5.7	5.4	4.8	4.1	2.9	1.7	1.0
2003.....	3.8	4.8	5.5	5.5	5.3	5.3	3.6	3.3	2.1	1.0
2002.....	3.9	5.1	5.4	5.6	5.1	4.9	4.8	3.7	2.2	1.0
2001.....	3.9	5.3	5.8	4.7	5.6	5.3	4.0	3.3	2.1	1.0
2000.....	3.9	4.8	4.8	5.8	6.2	5.1	3.9	2.6	2.4	1.0
1999.....	3.5	4.6	4.8	4.9	5.9	4.6	4.0	3.9	2.7	1.5

White, both sexes

2004.....	6.1	9.5	12.6	11.8	11.7	11.1	9.3	6.8	4.1	2.2
2003.....	6.2	9.7	11.8	11.8	11.6	11.2	8.6	7.0	4.5	1.9
2002.....	6.2	9.6	11.5	11.4	12.0	11.0	9.8	6.8	4.5	2.1
2001.....	6.2	10.5	11.0	11.0	12.2	11.2	9.2	7.1	4.0	2.3
2000.....	6.6	10.0	10.6	11.5	13.0	11.6	9.3	7.3	4.3	2.3
1999.....	6.2	9.8	10.8	11.2	12.4	11.5	9.4	7.2	5.3	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alcoholic liver disease (K70)

White, male

2004.....	8.5	13.3	19.4	18.1	18.4	18.1	15.3	12.0	8.0	4.8
2003.....	8.4	14.6	18.4	18.3	18.7	17.9	14.6	11.8	8.2	3.7
2002.....	8.3	14.3	17.9	17.9	19.4	17.7	15.8	11.1	8.2	4.5
2001.....	8.5	15.9	16.8	17.6	19.4	17.8	15.4	12.5	7.2	5.4
2000.....	9.4	15.6	16.8	18.0	20.6	19.0	15.9	14.1	7.8	5.4
1999.....	9.2	15.4	17.1	18.2	20.0	19.5	16.2	11.9	9.8	5.8

White, female

2004.....	3.7	5.7	5.9	5.7	5.4	4.9	4.4	3.0	1.7	1.0
2003.....	3.9	4.9	5.4	5.5	5.1	5.3	3.7	3.5	2.2	1.1
2002.....	4.0	5.0	5.2	5.3	5.1	5.0	4.9	3.7	2.2	1.1
2001.....	3.9	5.2	5.4	4.6	5.7	5.4	4.2	3.3	2.1	1.0
2000.....	3.8	4.4	4.5	5.3	6.0	5.1	4.0	2.6	2.3	1.0
1999.....	3.3	4.3	4.5	4.7	5.5	4.6	4.0	3.9	2.6	1.5

Black, both sexes

2004.....	4.1	7.5	10.8	12.9	12.0	10.8	6.5	*	*	*
2003.....	4.2	7.6	12.7	12.2	12.9	11.2	6.9	*	*	*
2002.....	4.2	8.2	12.1	12.5	12.0	9.6	7.4	6.2	*	*
2001.....	5.9	9.0	14.7	14.1	11.5	13.0	7.8	5.7	*	*
2000.....	6.9	11.0	13.3	13.9	12.6	11.8	8.4	6.2	*	*
1999.....	7.2	12.8	12.8	14.8	15.6	12.9	10.4	8.0	*	*

Black, male

2004.....	5.6	10.7	15.6	20.8	19.4	19.4	13.6	*	*	*
2003.....	5.4	12.0	20.6	19.5	20.8	18.7	12.5	*	*	*
2002.....	5.6	11.5	18.7	17.4	20.4	16.5	13.0	10.6	*	*
2001.....	8.2	12.9	22.3	23.5	18.7	22.9	15.1	*	*	*
2000.....	9.6	15.0	20.6	20.4	18.0	19.3	16.7	11.3	*	*
1999.....	10.1	19.2	20.2	24.9	23.3	24.1	18.7	14.2	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)										
Black, female										
2004.....	2.8	4.6	6.8	6.3	6.1	4.5	*	*	*	*
2003.....	3.0	3.7	6.0	6.1	6.8	5.7	*	*	*	*
2002.....	3.0	5.3	6.6	8.4	5.4	4.4	*	*	*	*
2001.....	3.9	5.5	8.2	6.4	5.8	5.6	*	*	*	*
2000.....	4.5	7.5	7.1	8.5	8.3	6.3	*	*	*	*
1999.....	4.7	7.2	6.6	6.5	9.5	4.6	5.0	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic hepatitis, not elsewhere classified (K73)

All races 3/, both sexes

2004.....	*	0.1	*	*	*	*	*	*	*	*
2003.....	*	*	0.1	*	*	0.2	*	*	*	*
2002.....	0.1	0.2	0.2	0.2	0.4	0.7	0.8	1.0	1.0	0.8
2001.....	*	0.2	0.2	0.3	0.3	0.6	0.8	0.8	0.8	0.7
2000.....	0.1	0.1	0.2	0.3	0.3	0.7	0.6	0.9	0.8	0.8
1999.....	*	0.2	0.2	0.3	0.4	0.4	0.7	0.8	0.9	0.6

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	0.2	*	0.4	0.6	1.0	1.1	1.3	1.3	0.9
2001.....	*	0.2	0.2	0.4	0.5	0.9	1.0	1.1	1.0	0.7
2000.....	*	*	*	0.4	0.6	1.1	0.9	1.2	1.0	0.9
1999.....	*	*	*	0.4	0.6	0.6	0.9	1.0	1.2	0.7

White, both sexes

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	0.2	0.2	0.2	0.4	0.7	0.9	1.1	0.9	0.8
2001.....	*	0.2	0.2	0.3	0.3	0.6	0.7	0.8	0.9	0.8
2000.....	*	*	0.2	0.3	0.4	0.7	0.6	0.9	0.8	0.9
1999.....	*	0.2	0.2	0.3	0.4	0.4	0.7	0.8	0.9	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Fibrosis and cirrhosis of liver (K74)

All races 3/, both sexes

2004.....	2.9	5.9	9.0	9.9	12.1	15.3	20.4	23.5	22.7	17.4
2003.....	3.3	6.4	9.2	10.3	12.6	16.8	22.7	24.3	23.9	18.0
2002.....	3.6	6.4	8.7	10.1	12.2	16.5	20.9	24.8	24.5	18.5
2001.....	3.6	6.7	8.6	9.7	12.4	17.2	21.4	23.5	23.9	19.1
2000.....	3.5	6.7	7.7	9.9	12.9	16.8	21.1	23.9	24.4	20.0
1999.....	3.4	6.3	7.5	9.6	13.5	17.7	21.6	24.6	24.6	19.8

All races 3/, male

2004.....	4.1	8.5	13.2	14.0	15.1	17.8	23.5	27.0	25.9	20.9
2003.....	4.7	9.1	13.8	14.3	16.9	20.5	25.6	26.7	26.7	23.7
2002.....	4.8	9.3	13.1	14.3	15.1	19.6	24.1	27.9	28.5	23.2
2001.....	4.7	9.8	12.9	13.3	16.2	21.5	25.2	25.2	27.8	25.7
2000.....	4.9	10.1	11.5	13.6	17.3	20.4	24.8	26.6	26.4	28.4
1999.....	5.1	9.6	11.3	13.6	17.0	21.9	24.9	27.5	28.5	26.6

All races 3/, female

2004.....	1.8	3.4	5.0	6.0	9.4	13.1	17.9	20.9	20.8	15.8
2003.....	2.0	3.8	4.8	6.6	8.8	13.5	20.4	22.6	22.2	15.5
2002.....	2.5	3.6	4.5	6.1	9.6	13.9	18.3	22.6	22.1	16.5
2001.....	2.4	3.7	4.5	6.3	9.1	13.4	18.5	22.4	21.6	16.4
2000.....	2.1	3.5	4.1	6.5	8.9	13.7	18.2	21.9	23.2	16.6
1999.....	1.8	3.0	3.8	5.8	10.2	14.1	19.0	22.5	22.4	17.0

White, both sexes

2004.....	2.9	6.0	8.9	9.8	12.3	15.9	21.6	24.8	23.8	17.9
2003.....	3.4	6.4	9.1	10.1	12.9	17.5	24.0	25.8	25.1	18.8
2002.....	3.8	6.3	8.5	10.1	12.4	17.3	22.0	26.2	25.5	19.4
2001.....	3.5	6.5	8.3	9.4	12.7	17.9	22.4	24.7	25.1	20.1
2000.....	3.5	6.8	7.4	9.9	13.1	17.2	21.9	25.2	25.4	20.8
1999.....	3.4	5.9	7.4	9.3	13.5	18.3	22.5	25.7	25.8	20.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Fibrosis and cirrhosis of liver (K74)

White, male

2004.....	4.1	8.5	13.0	13.8	15.2	18.2	24.7	28.4	27.2	21.9
2003.....	4.9	9.4	13.5	13.7	17.1	21.3	27.0	28.1	27.8	24.5
2002.....	5.1	9.3	12.8	14.2	15.1	20.2	25.3	29.8	29.5	24.4
2001.....	4.6	9.5	12.3	12.8	16.2	22.2	26.2	26.2	29.5	27.3
2000.....	5.0	10.2	10.9	13.4	17.5	21.0	25.7	27.8	27.7	29.4
1999.....	5.0	9.1	11.1	13.2	17.1	22.6	25.9	28.7	30.0	27.9

White, female

2004.....	1.7	3.5	4.9	5.9	9.5	13.8	19.0	22.2	21.7	16.1
2003.....	1.9	3.5	4.7	6.6	9.1	14.1	21.6	24.1	23.4	16.3
2002.....	2.5	3.3	4.4	6.2	9.9	14.7	19.3	23.7	23.0	17.2
2001.....	2.4	3.5	4.4	6.1	9.6	14.1	19.3	23.6	22.4	17.1
2000.....	2.0	3.4	4.0	6.5	9.0	13.8	18.9	23.3	24.1	17.3
1999.....	1.7	2.8	3.7	5.7	10.2	14.6	19.7	23.6	23.4	17.5

Black, both sexes

2004.....	3.6	6.9	12.1	12.7	12.8	11.8	13.6	11.2	11.4	11.3
2003.....	3.5	7.4	12.8	14.2	12.1	13.8	14.6	11.1	10.9	10.8
2002.....	3.4	8.8	12.4	12.0	13.0	12.6	12.3	13.9	11.0	9.2
2001.....	4.5	9.9	12.0	13.5	12.0	13.5	14.1	14.1	11.4	7.4
2000.....	3.9	8.0	12.2	11.8	13.2	14.8	15.9	11.9	12.7	11.6
1999.....	4.6	9.5	9.8	13.1	14.5	14.1	14.1	14.6	12.4	12.1

Black, male

2004.....	4.9	10.4	18.5	19.9	17.0	15.7	18.5	15.9	14.3	*
2003.....	4.2	9.4	20.9	22.6	18.1	17.7	16.1	14.6	*	*
2002.....	4.0	11.5	19.2	19.5	18.6	18.2	16.0	12.4	*	*
2001.....	6.2	15.1	19.8	20.3	19.5	19.8	18.5	18.1	*	*
2000.....	4.8	11.3	19.8	17.9	19.3	18.5	20.7	16.1	*	*
1999.....	7.0	15.0	16.1	19.2	19.1	19.2	18.0	17.6	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Fibrosis and cirrhosis of liver (K74)

Black, female

2004.....	2.4	3.8	6.7	6.8	9.5	8.8	10.3	8.4	9.8	11.0
2003.....	2.9	5.6	5.9	7.3	7.4	11.0	13.5	9.0	10.1	8.1
2002.....	2.8	6.4	6.7	5.9	8.7	8.5	9.9	14.8	9.5	9.0
2001.....	2.9	5.4	5.3	7.8	6.1	8.8	11.2	11.6	12.7	*
2000.....	3.1	5.1	5.8	6.7	8.3	12.1	12.7	9.4	13.2	10.3
1999.....	2.5	4.8	4.5	8.2	10.8	10.4	11.5	12.9	11.7	12.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of liver (K71-K72,K75-K76)

All races 3/, both sexes

2004.....	2.4	4.2	5.8	6.5	7.3	8.3	10.1	11.3	11.2	13.1
2003.....	2.6	4.3	6.1	6.4	7.7	9.3	9.2	10.5	11.7	13.5
2002.....	2.7	4.7	5.4	5.7	6.8	8.1	9.2	9.4	12.0	14.3
2001.....	2.1	3.9	4.9	5.3	7.0	7.7	8.6	10.9	11.5	13.8
2000.....	2.2	3.6	4.7	5.0	6.2	7.4	8.9	9.5	11.2	13.6
1999.....	2.2	3.5	4.5	5.1	6.4	8.4	9.3	9.8	11.0	14.0

All races 3/, male

2004.....	2.8	5.5	8.6	8.8	9.2	9.7	11.5	12.7	12.8	15.1
2003.....	3.0	5.8	8.5	8.4	10.1	11.2	11.2	11.9	14.1	16.9
2002.....	3.4	6.5	7.7	7.7	8.7	10.5	11.6	11.6	15.9	18.9
2001.....	2.8	5.4	7.1	6.8	9.1	9.8	10.2	13.5	15.0	17.4
2000.....	2.8	5.1	6.2	6.4	7.4	9.1	10.4	12.1	13.6	17.8
1999.....	2.9	4.9	6.1	6.9	8.2	10.6	11.0	11.0	14.9	19.2

All races 3/, female

2004.....	1.9	2.9	3.2	4.3	5.7	7.2	8.9	10.2	10.3	12.3
2003.....	2.2	2.8	3.8	4.6	5.6	7.6	7.6	9.5	10.3	12.1
2002.....	2.0	2.9	3.2	3.8	5.0	6.0	7.2	7.9	9.6	12.3
2001.....	1.5	2.6	2.8	3.9	5.1	6.0	7.2	9.0	9.4	12.3
2000.....	1.6	2.2	3.2	3.7	5.1	6.0	7.7	7.8	9.7	11.9
1999.....	1.6	2.1	2.9	3.5	4.7	6.4	7.9	8.9	8.8	11.9

White, both sexes

2004.....	2.3	4.0	5.5	6.1	7.0	8.2	9.9	11.4	11.2	13.1
2003.....	2.6	4.2	5.8	6.2	7.5	9.2	9.1	10.5	11.9	13.4
2002.....	2.6	4.4	5.1	5.4	6.6	8.1	9.2	9.4	11.9	14.3
2001.....	2.0	3.7	4.7	4.9	6.8	7.8	8.4	10.8	11.4	13.8
2000.....	2.1	3.4	4.2	4.6	5.8	7.4	8.8	9.5	11.1	13.6
1999.....	2.1	3.2	4.1	4.8	6.0	8.2	9.1	9.7	11.2	13.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of liver (K71-K72,K75-K76)

White, male

2004.....	2.7	5.2	8.3	8.2	8.7	9.6	11.4	12.6	13.0	15.9
2003.....	3.1	5.7	8.0	7.9	9.7	10.9	10.9	12.0	14.4	16.7
2002.....	3.3	6.2	7.3	7.1	8.5	10.6	11.7	11.3	15.8	18.9
2001.....	2.6	5.1	6.8	6.3	8.9	10.1	10.0	13.2	14.6	17.3
2000.....	2.8	4.8	5.5	5.9	7.0	8.9	10.3	12.0	13.9	17.8
1999.....	2.8	4.5	5.7	6.5	7.8	10.5	10.9	11.2	15.4	19.6

White, female

2004.....	1.8	2.7	2.9	4.2	5.5	7.0	8.7	10.4	10.2	11.9
2003.....	2.1	2.7	3.5	4.6	5.4	7.7	7.5	9.4	10.4	11.9
2002.....	1.8	2.6	3.0	3.7	4.9	5.9	7.2	8.0	9.6	12.3
2001.....	1.4	2.4	2.6	3.6	4.9	5.8	7.0	9.1	9.4	12.4
2000.....	1.4	2.0	2.9	3.3	4.7	6.0	7.6	7.6	9.4	11.9
1999.....	1.3	1.8	2.6	3.1	4.3	6.2	7.6	8.7	8.7	11.6

Black, both sexes

2004.....	3.0	6.6	8.6	10.1	10.2	10.3	11.4	10.0	12.1	12.1
2003.....	3.0	5.9	9.7	9.2	10.8	11.3	10.7	10.4	8.9	14.9
2002.....	3.3	7.0	8.5	9.3	8.4	9.7	10.5	10.0	12.8	15.4
2001.....	3.0	6.4	7.6	9.9	9.3	9.1	11.4	12.0	14.4	11.4
2000.....	3.3	6.0	8.9	8.6	10.0	9.2	10.9	10.8	10.7	12.2
1999.....	3.7	6.4	8.0	8.7	10.9	10.4	11.2	11.2	9.2	14.0

Black, male

2004.....	3.5	8.6	12.1	15.6	14.4	12.4	14.6	13.2	*	*
2003.....	2.6	7.7	14.0	13.7	15.6	17.0	15.8	11.4	*	*
2002.....	3.6	9.4	11.9	14.6	10.9	12.2	12.7	13.8	17.6	*
2001.....	4.1	8.7	11.1	13.0	12.2	10.2	14.5	17.2	23.3	*
2000.....	3.9	8.2	12.8	12.2	13.3	13.0	13.0	13.7	*	*
1999.....	4.1	9.0	11.1	11.7	14.0	12.5	13.2	10.4	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of liver (K71-K72,K75-K76)

Black, female

2004.....	2.6	4.9	5.5	5.6	7.0	8.8	9.2	8.1	12.1	14.6
2003.....	3.3	4.4	6.1	5.5	7.1	7.1	7.3	9.8	*	13.4
2002.....	3.0	4.9	5.5	5.0	6.5	7.8	9.0	7.7	10.3	13.6
2001.....	2.1	4.4	4.5	7.3	7.0	8.3	9.4	8.9	9.8	10.1
2000.....	2.7	4.0	5.5	5.7	7.5	6.3	9.5	9.1	11.0	10.7
1999.....	3.3	4.2	5.3	6.2	8.5	8.9	9.9	11.7	9.9	13.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races 3/, both sexes

2004.....	0.2	0.3	0.4	0.6	1.2	1.8	3.2	5.8	10.4	23.8
2003.....	0.1	0.2	0.4	0.5	0.9	2.0	3.4	5.9	11.3	21.8
2002.....	0.2	0.2	0.4	0.6	1.1	2.0	3.6	5.7	10.8	22.9
2001.....	0.1	0.3	0.4	0.7	1.3	2.0	3.8	5.8	10.0	23.7
2000.....	0.1	0.2	0.4	0.6	1.0	1.9	3.0	5.7	10.3	24.8
1999.....	0.2	0.3	0.4	0.7	1.0	1.8	3.4	5.7	10.9	23.9

All races 3/, male

2004.....	*	0.3	0.4	0.7	1.4	2.2	3.8	7.0	12.5	25.5
2003.....	*	0.2	0.5	0.6	1.1	2.2	4.3	7.5	13.9	22.7
2002.....	0.2	0.3	0.5	0.8	1.1	2.3	4.0	7.5	13.5	24.1
2001.....	*	0.4	0.4	0.9	1.5	2.2	4.4	7.6	12.1	26.9
2000.....	*	0.2	0.3	0.7	1.2	2.3	3.6	7.4	12.9	27.6
1999.....	*	0.3	0.4	0.9	1.3	2.0	4.2	6.9	13.3	27.7

All races 3/, female

2004.....	*	0.3	0.4	0.5	1.0	1.5	2.8	5.0	9.0	23.0
2003.....	*	0.2	0.4	0.4	0.7	1.8	2.8	4.8	9.8	21.4
2002.....	*	*	0.3	0.4	1.0	1.8	3.2	4.4	9.1	22.4
2001.....	*	0.3	0.3	0.5	1.1	1.9	3.3	4.6	8.7	22.4
2000.....	*	0.2	0.4	0.5	0.8	1.6	2.5	4.5	8.7	23.7
1999.....	*	0.3	0.5	0.6	0.7	1.6	2.8	4.8	9.5	22.4

White, both sexes

2004.....	0.2	0.2	0.3	0.5	1.1	1.8	3.0	5.7	10.3	24.2
2003.....	*	0.2	0.3	0.4	0.8	1.8	3.6	5.9	11.3	21.9
2002.....	0.2	0.2	0.3	0.6	1.0	2.0	3.5	5.5	10.6	23.5
2001.....	*	0.3	0.3	0.6	1.3	1.8	3.7	5.7	9.8	23.9
2000.....	0.1	0.2	0.3	0.6	0.9	1.8	2.7	5.6	10.1	25.6
1999.....	0.1	0.3	0.4	0.6	0.9	1.6	3.4	5.7	11.0	24.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Black, female

2004.....	*	*	*	*	*	*	5.5	6.7	12.9	21.7
2003.....	*	*	*	*	*	*	*	*	9.3	20.7
2002.....	*	*	*	*	*	*	5.4	7.7	11.1	16.0
2001.....	*	*	*	*	*	*	6.3	6.9	13.1	16.8
2000.....	*	*	*	*	*	*	4.9	6.3	11.5	19.3
1999.....	*	*	*	*	*	*	*	*	12.1	15.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Disorders of biliary tract and pancreas (K83-K86)

All races 3/, both sexes

2004.....	0.9	1.3	1.6	2.0	2.5	3.9	5.6	7.9	11.5	20.8
2003.....	1.0	1.3	1.5	1.9	2.9	3.9	5.9	8.3	12.3	21.9
2002.....	1.0	1.3	1.5	2.3	2.7	4.0	5.8	8.1	12.0	21.7
2001.....	1.0	1.4	1.7	2.1	3.1	4.2	6.1	7.6	11.9	22.9
2000.....	1.0	1.3	1.5	2.3	2.7	3.7	5.5	7.9	11.5	22.0
1999.....	1.0	1.3	1.6	2.0	2.8	3.6	5.4	7.2	10.3	22.2

All races 3/, male

2004.....	1.2	1.8	2.2	2.7	3.4	5.1	6.5	9.3	12.6	21.1
2003.....	1.2	1.7	2.1	2.6	3.7	5.1	7.3	10.1	14.4	20.7
2002.....	1.3	1.9	2.2	3.0	4.0	5.5	7.4	10.3	14.3	21.2
2001.....	1.3	1.9	2.2	2.8	3.9	5.1	8.0	8.8	12.5	22.4
2000.....	1.5	1.8	2.1	2.9	3.1	5.2	6.7	10.2	13.3	22.9
1999.....	1.4	1.8	2.1	2.8	3.6	4.6	6.7	9.1	11.7	23.1

All races 3/, female

2004.....	0.6	0.9	1.0	1.2	1.7	2.8	4.9	6.8	10.8	20.7
2003.....	0.7	0.8	0.9	1.1	2.1	2.8	4.8	7.1	10.9	22.4
2002.....	0.7	0.7	0.9	1.7	1.6	2.7	4.4	6.6	10.5	21.9
2001.....	0.7	1.0	1.2	1.4	2.4	3.3	4.5	6.8	11.6	23.2
2000.....	0.6	0.9	1.0	1.7	2.3	2.5	4.6	6.3	10.5	21.6
1999.....	0.7	0.8	1.1	1.4	2.2	2.8	4.4	5.9	9.5	21.8

White, both sexes

2004.....	0.8	1.2	1.4	1.7	2.3	3.9	5.4	7.8	11.5	21.0
2003.....	0.8	1.1	1.3	1.7	2.7	3.7	5.6	8.5	12.4	22.0
2002.....	0.9	1.1	1.3	2.2	2.5	3.9	5.7	8.2	12.1	22.0
2001.....	0.9	1.3	1.4	1.9	3.0	4.1	6.0	7.6	12.0	23.0
2000.....	0.9	1.1	1.4	2.0	2.5	3.4	5.4	7.8	11.5	21.9
1999.....	1.0	1.0	1.4	1.8	2.6	3.5	5.5	7.1	10.2	22.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Disorders of biliary tract and pancreas (K83-K86)

White, male

2004.....	1.2	1.6	1.9	2.4	3.1	5.1	6.3	9.2	12.8	21.4
2003.....	0.9	1.5	1.8	2.3	3.4	4.9	7.0	10.1	14.5	20.5
2002.....	1.2	1.7	1.9	2.8	3.6	5.5	7.2	10.5	14.5	21.2
2001.....	1.2	1.6	1.9	2.6	3.7	5.0	8.0	8.7	12.9	22.2
2000.....	1.3	1.5	1.8	2.7	2.9	4.7	6.6	10.2	13.2	22.0
1999.....	1.3	1.4	1.8	2.4	3.3	4.6	6.8	9.0	11.7	22.0

White, female

2004.....	0.5	0.8	0.9	1.1	1.6	2.8	4.7	6.8	10.6	20.8
2003.....	0.6	0.6	0.8	1.0	2.0	2.6	4.4	7.3	11.1	22.6
2002.....	0.5	0.6	0.8	1.7	1.4	2.6	4.5	6.6	10.6	22.3
2001.....	0.6	0.9	1.0	1.3	2.3	3.3	4.3	6.8	11.5	23.4
2000.....	0.6	0.8	1.0	1.5	2.2	2.2	4.5	6.2	10.4	21.9
1999.....	0.6	0.6	1.1	1.3	1.9	2.6	4.4	5.7	9.3	22.0

Black, both sexes

2004.....	1.8	2.4	3.5	4.5	5.3	4.3	8.8	10.0	11.4	20.3
2003.....	2.3	2.8	3.4	3.9	5.3	5.9	9.3	7.8	10.4	23.3
2002.....	1.8	2.9	3.2	3.8	5.3	5.0	7.1	6.7	11.5	21.4
2001.....	2.2	3.2	4.1	3.9	4.4	6.0	7.5	8.0	11.6	22.1
2000.....	2.0	3.1	2.8	4.5	4.6	6.7	7.7	8.7	12.1	23.5
1999.....	1.7	3.3	3.3	4.1	5.5	5.6	5.3	8.4	12.4	23.0

Black, male

2004.....	1.8	3.5	5.2	6.3	7.7	6.1	11.7	13.6	*	*
2003.....	3.6	3.5	4.8	6.2	7.7	7.1	11.8	10.0	14.8	26.1
2002.....	2.1	4.5	4.7	5.7	7.3	6.0	11.7	*	*	25.7
2001.....	2.7	4.7	5.5	5.9	5.9	8.4	9.1	11.2	*	*
2000.....	3.0	3.9	4.4	6.3	5.6	10.4	9.7	10.9	*	32.3
1999.....	2.5	5.0	5.4	6.6	6.4	6.8	*	10.4	*	32.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Disorders of biliary tract and pancreas (K83-K86)

Black, female

2004.....	1.7	1.5	2.1	2.9	3.5	*	6.8	7.8	12.5	20.9
2003.....	*	2.3	2.1	*	3.5	4.9	7.5	6.5	8.1	22.3
2002.....	1.5	1.5	1.9	*	3.7	4.3	*	5.5	11.5	19.7
2001.....	1.8	2.0	2.9	*	3.2	4.3	6.5	6.1	13.5	22.3
2000.....	*	2.4	*	3.1	3.8	4.0	6.4	7.4	11.0	20.2
1999.....	*	1.8	*	*	4.8	4.8	5.2	7.2	13.0	19.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pancreatitis (K85,K86.0-K86.1)

All races 3/, both sexes

2004.....	0.8	1.2	1.3	1.6	2.0	2.9	4.0	5.0	7.0	11.4
2003.....	0.9	1.1	1.3	1.5	2.2	2.8	4.1	5.7	7.9	11.8
2002.....	0.9	1.2	1.3	1.9	2.2	2.9	4.0	5.7	7.8	11.9
2001.....	0.9	1.2	1.4	1.8	2.5	3.2	4.5	5.4	7.7	13.2
2000.....	0.9	1.1	1.3	1.8	2.3	2.8	4.0	5.4	7.6	12.1
1999.....	0.9	1.1	1.4	1.6	2.2	2.9	4.0	5.2	6.8	12.8

All races 3/, male

2004.....	1.2	1.7	1.9	2.4	2.7	3.9	4.8	6.1	7.2	10.3
2003.....	1.1	1.5	1.8	2.1	3.1	4.0	5.1	6.7	9.1	10.6
2002.....	1.1	1.8	1.9	2.5	3.2	4.2	5.1	7.1	8.8	11.2
2001.....	1.2	1.7	1.8	2.5	3.1	3.9	6.1	6.1	7.8	12.9
2000.....	1.4	1.4	1.7	2.4	2.6	4.1	5.0	6.8	8.3	11.9
1999.....	1.3	1.6	1.8	2.2	2.9	3.8	5.1	6.4	7.5	12.7

All races 3/, female

2004.....	0.5	0.7	0.8	1.0	1.3	2.0	3.4	4.3	6.9	11.8
2003.....	0.6	0.7	0.8	0.8	1.5	1.9	3.4	5.0	7.2	12.3
2002.....	0.6	0.6	0.8	1.4	1.2	1.8	3.1	4.7	7.1	12.2
2001.....	0.6	0.8	1.0	1.1	1.9	2.7	3.3	4.9	7.7	13.4
2000.....	0.5	0.8	0.8	1.2	1.9	1.8	3.2	4.4	7.2	12.2
1999.....	0.6	0.7	0.9	1.0	1.6	2.2	3.2	4.3	6.5	12.8

White, both sexes

2004.....	0.8	1.1	1.1	1.4	1.8	2.9	3.9	5.0	6.9	11.4
2003.....	0.7	0.9	1.1	1.3	2.1	2.8	3.8	5.9	8.0	11.6
2002.....	0.8	1.0	1.1	1.9	1.9	2.9	4.0	5.8	7.9	12.1
2001.....	0.8	1.1	1.1	1.7	2.4	3.2	4.5	5.5	7.7	13.3
2000.....	0.8	0.9	1.2	1.6	2.1	2.6	3.9	5.3	7.7	12.2
1999.....	0.9	0.9	1.2	1.4	2.0	2.9	4.0	5.0	6.8	12.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pancreatitis (K85,K86.0-K86.1)

White, male

2004.....	1.1	1.5	1.6	2.0	2.4	3.8	4.6	6.1	7.1	10.4
2003.....	0.9	1.3	1.5	1.9	2.7	3.8	4.7	6.7	9.1	10.3
2002.....	1.1	1.5	1.6	2.4	2.8	4.3	5.1	7.2	8.9	11.0
2001.....	1.1	1.5	1.5	2.3	2.9	3.8	6.2	6.1	7.8	12.9
2000.....	1.2	1.2	1.5	2.1	2.4	3.7	4.9	6.7	8.3	11.5
1999.....	1.2	1.2	1.5	1.9	2.7	3.7	5.2	6.5	7.4	12.0

White, female

2004.....	0.4	0.6	0.6	0.8	1.2	2.1	3.2	4.3	6.9	11.8
2003.....	0.5	0.5	0.7	0.8	1.4	1.9	3.1	5.2	7.3	12.2
2002.....	0.5	0.5	0.6	1.4	1.1	1.8	3.2	4.8	7.2	12.6
2001.....	0.5	0.7	0.8	1.1	1.9	2.6	3.1	5.0	7.7	13.4
2000.....	0.5	0.7	0.8	1.0	1.8	1.6	3.2	4.4	7.3	12.5
1999.....	0.5	0.6	0.9	1.0	1.3	2.1	3.1	4.1	6.4	13.1

Black, both sexes

2004.....	1.6	2.3	3.1	4.0	4.1	3.3	6.4	6.4	8.2	13.3
2003.....	2.2	2.5	2.9	3.2	4.3	4.0	7.0	5.1	6.9	14.0
2002.....	1.4	2.5	2.8	3.0	4.7	3.5	3.9	4.3	6.5	11.3
2001.....	2.1	2.8	3.8	3.1	3.8	4.2	5.1	5.0	8.7	13.2
2000.....	1.9	2.7	2.4	3.9	3.8	5.2	5.6	5.3	7.6	11.9
1999.....	1.6	3.0	3.1	3.3	4.3	4.5	4.2	6.4	8.1	12.8

Black, male

2004.....	1.7	3.3	4.6	5.8	5.8	5.3	8.7	9.6	*	*
2003.....	3.3	3.3	4.2	5.5	6.7	5.9	9.2	*	*	*
2002.....	1.8	3.9	4.2	4.5	6.9	*	*	*	*	*
2001.....	2.6	4.2	4.9	5.0	5.5	5.6	*	*	*	*
2000.....	3.0	3.2	3.8	5.4	4.8	8.3	7.7	*	*	*
1999.....	2.4	4.5	5.1	5.4	4.9	6.0	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pancreatitis (K85,K86.0-K86.1)

Black, female

2004.....	1.4	1.4	1.9	2.5	*	*	4.8	*	8.7	13.4
2003.....	*	1.8	1.7	*	*	*	5.5	*	*	13.4
2002.....	*	*	*	*	*	*	*	*	*	9.4
2001.....	1.6	1.7	2.9	*	*	*	4.7	*	9.4	13.9
2000.....	*	2.2	*	2.7	*	*	*	*	*	9.9
1999.....	*	1.6	*	*	3.8	*	*	6.0	*	10.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)

All races 3/, both sexes

2004.....	0.1	0.1	0.2	0.3	0.6	1.0	1.6	2.8	4.4	9.4
2003.....	0.1	0.2	0.2	0.4	0.6	1.0	1.8	2.6	4.3	10.1
2002.....	0.1	0.1	0.2	0.4	0.6	1.1	1.8	2.5	4.2	9.8
2001.....	*	0.2	0.3	0.3	0.6	1.0	1.5	2.2	4.2	9.7
2000.....	0.1	0.2	0.3	0.5	0.4	0.9	1.5	2.5	3.9	9.9
1999.....	0.1	0.1	0.2	0.5	0.6	0.7	1.4	2.0	3.5	9.4

All races 3/, male

2004.....	*	*	0.2	0.4	0.8	1.2	1.7	3.2	5.4	10.7
2003.....	*	*	0.4	0.5	0.7	1.1	2.2	3.4	5.3	10.1
2002.....	*	*	0.3	0.5	0.8	1.3	2.3	3.3	5.5	10.0
2001.....	*	0.2	0.4	0.3	0.7	1.3	1.9	2.6	4.7	9.5
2000.....	*	0.3	0.4	0.6	0.5	1.1	1.6	3.5	5.0	11.0
1999.....	*	0.2	0.3	0.6	0.7	0.8	1.6	2.7	4.2	10.5

All races 3/, female

2004.....	*	*	0.2	0.3	0.4	0.8	1.5	2.6	3.9	8.9
2003.....	*	*	*	0.3	0.6	0.9	1.4	2.1	3.7	10.2
2002.....	*	*	*	0.3	0.3	0.9	1.4	1.9	3.4	9.8
2001.....	*	0.2	*	0.3	0.5	0.7	1.2	1.9	3.9	9.8
2000.....	*	*	*	0.4	0.4	0.7	1.4	1.9	3.2	9.4
1999.....	*	*	*	0.4	0.6	0.6	1.2	1.6	3.1	9.0

White, both sexes

2004.....	*	0.1	0.2	0.3	0.5	1.0	1.5	2.8	4.5	9.6
2003.....	*	0.1	0.2	0.3	0.6	0.9	1.8	2.7	4.4	10.3
2002.....	*	0.1	0.2	0.4	0.5	1.0	1.7	2.4	4.2	9.9
2001.....	*	0.2	0.3	0.2	0.6	0.9	1.5	2.1	4.3	9.7
2000.....	*	0.2	0.2	0.5	0.4	0.8	1.5	2.5	3.8	9.7
1999.....	*	0.1	0.3	0.4	0.6	0.7	1.4	2.0	3.4	9.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases of digestive system (K90,K92)

All races 3/, both sexes

2004.....	0.3	0.7	1.2	1.8	2.7	4.2	7.3	13.0	25.0	68.5
2003.....	0.4	0.6	1.1	1.7	2.9	4.7	7.7	14.0	26.0	74.2
2002.....	0.4	0.8	1.2	1.7	2.6	4.7	7.6	14.1	27.6	76.2
2001.....	0.4	0.7	1.0	1.8	3.1	4.8	7.7	14.1	28.2	77.8
2000.....	0.4	0.7	1.1	1.5	2.8	4.1	7.9	15.0	29.0	80.6
1999.....	0.4	0.7	1.0	1.7	2.8	4.7	7.7	14.1	27.8	77.2

All races 3/, male

2004.....	0.4	1.0	1.7	2.6	3.3	5.7	9.4	16.1	30.2	69.8
2003.....	0.6	0.9	1.5	2.3	3.6	6.1	9.7	17.8	32.3	73.3
2002.....	0.6	1.2	1.6	2.3	3.8	6.1	9.6	17.8	31.0	76.2
2001.....	0.6	1.0	1.6	2.5	3.9	6.3	9.7	17.5	33.0	81.4
2000.....	0.6	1.0	1.6	2.0	3.8	5.2	9.2	18.3	36.5	83.1
1999.....	0.5	1.0	1.4	2.3	3.6	6.1	10.0	17.3	34.3	80.8

All races 3/, female

2004.....	0.3	0.4	0.7	1.0	2.1	2.9	5.7	10.9	21.8	67.9
2003.....	0.2	0.4	0.6	1.1	2.2	3.5	6.1	11.3	22.1	74.6
2002.....	0.2	0.4	0.7	1.2	1.6	3.4	6.0	11.5	25.5	76.2
2001.....	0.2	0.4	0.5	1.1	2.4	3.6	6.1	11.8	25.3	76.3
2000.....	0.2	0.4	0.6	1.0	1.9	3.0	6.8	12.7	24.6	79.6
1999.....	0.2	0.4	0.6	1.2	2.1	3.6	6.0	11.8	24.0	75.8

White, both sexes

2004.....	0.3	0.7	1.1	1.6	2.5	3.9	7.0	12.8	24.8	69.8
2003.....	0.4	0.6	1.0	1.6	2.6	4.3	7.3	13.6	26.2	75.5
2002.....	0.3	0.7	1.1	1.5	2.4	4.5	7.3	13.6	27.6	77.8
2001.....	0.3	0.6	0.9	1.6	2.8	4.5	7.5	13.7	28.4	79.1
2000.....	0.3	0.6	0.9	1.3	2.4	3.7	7.6	14.4	28.8	81.7
1999.....	0.3	0.6	0.8	1.6	2.5	4.4	7.5	13.6	27.6	78.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other diseases of digestive system (K90,K92)

White, male

2004.....	0.4	1.0	1.6	2.3	2.9	5.3	8.9	15.9	29.9	71.6
2003.....	0.6	0.8	1.5	2.1	3.4	5.7	9.1	17.2	32.6	75.2
2002.....	0.5	1.0	1.6	2.1	3.5	5.8	9.1	17.3	31.2	77.3
2001.....	0.5	0.9	1.3	2.1	3.5	5.8	9.3	16.7	33.5	82.8
2000.....	0.5	0.8	1.4	1.7	3.4	4.9	8.8	17.5	36.2	84.9
1999.....	0.5	0.9	1.2	2.0	3.4	5.7	9.6	16.6	34.4	82.1

White, female

2004.....	0.2	0.3	0.6	1.0	2.1	2.6	5.5	10.6	21.7	69.0
2003.....	0.3	0.4	0.6	1.0	1.9	3.0	5.9	11.0	22.2	75.6
2002.....	*	0.4	0.6	1.0	1.4	3.3	5.8	11.0	25.4	78.1
2001.....	*	0.4	0.5	1.0	2.1	3.4	6.0	11.5	25.3	77.5
2000.....	*	0.3	0.5	0.8	1.6	2.7	6.6	12.2	24.4	80.5
1999.....	*	0.3	0.5	1.1	1.8	3.3	5.8	11.4	23.6	77.6

Black, both sexes

2004.....	*	1.0	2.4	3.5	4.8	7.7	12.3	19.0	32.4	62.0
2003.....	0.7	1.1	1.9	3.2	5.5	9.0	12.3	22.5	27.4	70.3
2002.....	0.8	1.6	2.0	3.6	5.3	7.8	12.2	21.6	33.5	65.3
2001.....	0.9	1.3	2.2	4.2	6.4	8.5	11.2	22.1	31.4	73.1
2000.....	0.9	1.6	2.5	3.9	6.3	7.2	12.9	22.6	36.7	77.9
1999.....	*	1.2	2.2	3.1	5.5	7.8	12.2	22.0	33.2	66.0

Black, male

2004.....	*	*	3.2	5.7	7.5	10.9	17.5	22.7	39.5	58.6
2003.....	*	*	2.6	4.8	6.7	10.6	18.1	29.7	34.0	59.4
2002.....	*	2.7	2.5	4.5	7.3	11.2	17.0	26.2	36.7	70.8
2001.....	*	2.2	3.7	6.3	7.9	11.4	16.5	30.2	36.1	71.5
2000.....	*	2.4	3.3	4.9	8.7	7.8	17.4	29.3	47.2	78.4
1999.....	*	*	3.3	5.3	6.2	9.9	16.6	29.0	38.4	71.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of digestive system (K90,K92)										
Black, female										
2004.....	*	*	1.7	*	*	5.4	8.8	16.7	28.7	63.4
2003.....	*	*	*	*	4.5	7.9	8.4	18.2	24.0	74.5
2002.....	*	*	*	2.8	3.7	5.2	9.0	18.9	31.8	63.3
2001.....	*	*	*	*	5.3	6.4	7.6	17.2	29.0	73.7
2000.....	*	*	*	3.0	4.4	6.7	10.0	18.5	31.4	77.7
1999.....	*	*	*	*	5.0	6.3	9.2	17.7	30.7	64.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XII. Diseases of the skin and subcutaneous tissue (L00-L98)

All races 3/, both sexes

2004.....	0.4	0.6	0.7	1.2	1.7	2.7	4.3	7.1	12.7	28.8
2003.....	0.4	0.5	0.8	1.2	1.7	2.4	4.3	6.7	12.5	30.5
2002.....	0.4	0.5	0.7	1.1	1.4	2.0	4.2	6.5	11.8	32.8
2001.....	0.3	0.5	0.7	1.0	1.4	2.2	3.5	6.5	12.0	32.2
2000.....	0.3	0.4	0.5	0.9	1.5	2.1	4.0	6.4	12.7	34.7
1999.....	0.3	0.5	0.6	0.8	1.3	2.2	3.5	6.6	12.7	35.0

All races 3/, male

2004.....	0.3	0.6	0.7	1.1	1.7	2.2	4.1	7.0	11.9	25.3
2003.....	0.4	0.5	0.8	1.1	1.8	2.3	4.4	6.3	11.7	26.6
2002.....	0.3	0.5	0.7	1.1	1.2	2.0	4.5	6.5	11.9	28.6
2001.....	0.2	0.6	0.7	1.0	1.4	2.5	3.6	6.7	10.8	24.9
2000.....	0.3	0.4	0.4	1.0	1.6	1.6	4.0	6.1	12.3	29.7
1999.....	0.2	0.5	0.6	0.8	1.1	2.3	3.5	6.1	12.2	31.7

All races 3/, female

2004.....	0.4	0.5	0.7	1.3	1.8	3.1	4.4	7.1	13.2	30.4
2003.....	0.4	0.5	0.8	1.2	1.6	2.5	4.3	7.1	13.0	32.2
2002.....	0.4	0.4	0.7	1.1	1.5	2.1	3.9	6.6	11.7	34.7
2001.....	0.3	0.5	0.6	1.0	1.5	1.9	3.4	6.4	12.7	35.2
2000.....	0.3	0.5	0.5	0.9	1.3	2.4	3.9	6.6	13.0	36.7
1999.....	0.4	0.5	0.6	0.8	1.4	2.2	3.6	7.0	12.9	36.3

White, both sexes

2004.....	0.3	0.5	0.7	1.0	1.5	2.3	3.4	6.4	11.3	25.7
2003.....	0.3	0.4	0.6	1.0	1.4	2.1	3.4	5.9	11.1	27.2
2002.....	0.3	0.4	0.5	0.9	1.0	1.7	3.3	5.6	10.0	28.3
2001.....	0.2	0.4	0.6	0.8	1.2	1.8	3.0	5.6	10.4	28.1
2000.....	0.2	0.3	0.4	0.8	1.2	1.7	3.1	5.2	11.0	29.7
1999.....	0.2	0.4	0.4	0.6	1.0	1.8	2.9	5.3	10.8	29.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XII. Diseases of the skin and subcutaneous tissue (L00-L98)

White, male

2004.....	0.3	0.5	0.7	0.9	1.5	1.9	3.3	6.5	10.8	23.1
2003.....	0.3	0.4	0.7	1.0	1.5	1.8	3.3	5.7	10.1	24.0
2002.....	0.3	0.5	0.5	1.0	0.9	1.6	3.6	5.7	10.7	26.1
2001.....	*	0.5	0.5	0.9	1.2	2.2	3.3	5.5	9.4	21.8
2000.....	*	0.3	0.4	0.8	1.4	1.2	3.1	4.9	10.4	25.8
1999.....	*	0.3	0.4	0.6	0.9	1.9	2.8	4.7	10.4	27.8

White, female

2004.....	0.3	0.5	0.6	1.0	1.5	2.6	3.6	6.3	11.7	27.0
2003.....	0.3	0.4	0.6	1.1	1.4	2.4	3.6	6.1	11.7	28.7
2002.....	0.3	0.3	0.5	0.9	1.1	1.8	3.1	5.5	9.6	29.3
2001.....	0.2	0.4	0.6	0.8	1.1	1.5	2.8	5.6	11.0	30.8
2000.....	0.2	0.2	0.4	0.7	1.1	2.1	3.2	5.5	11.4	31.2
1999.....	0.3	0.4	0.4	0.6	1.0	1.7	3.0	5.6	11.0	30.0

Black, both sexes

2004.....	0.7	1.3	1.7	3.3	3.9	6.6	13.3	16.8	33.2	70.8
2003.....	0.8	1.4	2.2	2.8	4.4	5.7	14.3	18.0	32.2	73.2
2002.....	1.1	1.0	2.0	3.2	4.4	5.1	12.5	19.1	35.3	92.9
2001.....	0.8	1.3	1.5	2.5	3.8	5.9	9.0	19.1	33.9	88.4
2000.....	0.8	1.4	1.1	2.5	3.7	6.0	12.5	19.9	36.1	97.3
1999.....	1.0	1.7	2.0	2.9	4.2	6.7	11.5	23.2	38.7	105.2

Black, male

2004.....	*	*	*	2.6	*	6.1	12.9	16.4	30.8	57.6
2003.....	*	*	*	*	5.6	6.6	17.7	16.4	34.7	65.7
2002.....	*	*	2.2	*	*	6.0	15.4	18.4	32.9	63.3
2001.....	*	*	*	*	*	6.6	8.1	24.6	34.5	71.5
2000.....	*	*	*	*	*	6.2	15.0	20.3	41.3	81.9
1999.....	*	2.2	*	*	*	6.8	12.2	24.2	39.2	85.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XII. Diseases of the skin and subcutaneous tissue (L00-L98)

Black, female

2004.....	*	*	1.7	3.9	4.4	7.0	13.6	17.0	34.4	76.0
2003.....	*	1.5	2.5	3.1	3.5	4.9	12.0	19.0	30.9	76.2
2002.....	1.4	*	1.8	3.7	5.1	4.4	10.5	19.5	36.6	104.3
2001.....	*	*	*	3.0	4.2	5.4	9.6	15.8	33.5	94.7
2000.....	*	2.0	*	2.7	3.6	5.8	10.9	19.7	33.5	103.0
1999.....	*	*	*	2.9	4.7	6.7	11.0	22.6	38.5	112.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Infections of skin and subcutaneous tissue (L00-L08)

All races 3/, both sexes

2004.....	0.2	0.3	0.4	0.6	0.8	1.2	1.7	3.0	4.9	10.4
2003.....	0.1	0.2	0.4	0.6	0.8	1.0	1.7	2.7	4.5	10.6
2002.....	0.1	0.2	0.3	0.5	0.6	0.8	1.3	2.2	3.6	9.3
2001.....	0.1	0.3	0.3	0.4	0.7	0.9	1.2	2.0	3.4	9.4
2000.....	0.1	0.1	0.2	0.4	0.6	0.7	1.4	2.0	3.4	9.3
1999.....	*	0.2	0.2	0.3	0.5	0.7	1.1	1.5	2.9	8.3

All races 3/, male

2004.....	0.2	0.4	0.5	0.6	0.8	1.1	1.8	3.3	5.0	10.3
2003.....	*	0.3	0.5	0.7	0.9	1.1	1.8	2.6	4.8	10.4
2002.....	0.2	0.3	0.4	0.6	0.6	0.9	1.5	2.2	4.1	9.1
2001.....	*	0.4	0.4	0.5	0.8	1.1	1.4	2.2	2.9	8.1
2000.....	*	*	*	0.6	0.7	0.7	1.6	2.1	4.0	8.9
1999.....	*	0.3	0.3	0.4	0.4	0.7	1.0	1.3	3.3	9.1

All races 3/, female

2004.....	*	0.2	0.3	0.5	0.9	1.3	1.6	2.8	4.9	10.4
2003.....	*	0.2	0.3	0.5	0.7	1.0	1.6	2.7	4.4	10.7
2002.....	*	*	0.2	0.4	0.5	0.8	1.2	2.2	3.2	9.4
2001.....	*	*	0.3	0.4	0.6	0.7	1.0	1.9	3.8	10.0
2000.....	*	*	*	0.3	0.5	0.7	1.2	1.9	3.1	9.5
1999.....	*	*	*	*	0.5	0.7	1.1	1.6	2.7	8.0

White, both sexes

2004.....	0.2	0.3	0.4	0.5	0.8	1.2	1.6	3.1	4.9	10.4
2003.....	0.1	0.2	0.4	0.6	0.8	1.0	1.6	2.7	4.5	10.8
2002.....	0.1	0.2	0.3	0.5	0.6	0.7	1.2	2.1	3.6	9.7
2001.....	*	0.2	0.3	0.4	0.6	0.9	1.2	2.0	3.4	9.6
2000.....	*	0.1	0.2	0.4	0.6	0.7	1.3	1.9	3.4	9.3
1999.....	*	0.2	0.2	0.3	0.4	0.7	1.1	1.5	3.0	8.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)

All races 3/, both sexes

2004.....	0.2	0.3	0.4	0.6	0.9	1.5	2.6	4.1	7.8	18.4
2003.....	0.2	0.3	0.4	0.6	0.9	1.4	2.7	4.1	8.0	19.9
2002.....	0.2	0.3	0.4	0.6	0.8	1.2	2.8	4.3	8.2	23.5
2001.....	0.2	0.3	0.3	0.6	0.8	1.3	2.3	4.5	8.5	22.7
2000.....	0.2	0.3	0.3	0.5	0.8	1.4	2.6	4.4	9.3	25.4
1999.....	0.2	0.3	0.4	0.5	0.8	1.5	2.5	5.1	9.7	26.6

All races 3/, male

2004.....	*	0.2	0.3	0.5	0.9	1.1	2.3	3.7	6.9	14.9
2003.....	0.2	0.2	0.3	0.4	1.0	1.2	2.6	3.6	6.9	16.2
2002.....	*	0.2	0.3	0.5	0.5	1.1	3.0	4.3	7.8	19.4
2001.....	*	0.2	0.3	0.5	0.6	1.4	2.2	4.5	7.9	16.8
2000.....	0.2	*	*	0.4	0.9	0.9	2.4	4.0	8.3	20.8
1999.....	*	0.2	0.3	0.4	0.7	1.6	2.4	4.8	8.9	22.6

All races 3/, female

2004.....	0.2	0.3	0.4	0.8	0.9	1.7	2.8	4.3	8.3	20.0
2003.....	0.3	0.4	0.5	0.7	0.9	1.6	2.7	4.4	8.6	21.5
2002.....	0.3	0.3	0.5	0.7	1.0	1.3	2.7	4.3	8.5	25.3
2001.....	0.2	0.3	0.4	0.6	0.9	1.2	2.4	4.5	8.9	25.2
2000.....	0.2	0.4	0.4	0.6	0.8	1.7	2.7	4.7	9.9	27.3
1999.....	0.3	0.3	0.4	0.6	0.8	1.5	2.5	5.4	10.2	28.3

White, both sexes

2004.....	0.1	0.2	0.3	0.5	0.7	1.1	1.8	3.3	6.4	15.3
2003.....	0.2	0.2	0.3	0.4	0.7	1.1	1.9	3.2	6.5	16.4
2002.....	0.2	0.2	0.2	0.4	0.5	1.0	2.1	3.5	6.4	18.7
2001.....	0.1	0.2	0.3	0.4	0.6	1.0	1.8	3.5	7.0	18.5
2000.....	0.1	0.2	0.2	0.3	0.6	1.0	1.8	3.3	7.6	20.4
1999.....	0.1	0.2	0.2	0.3	0.5	1.1	1.8	3.8	7.7	21.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)

White, male

2004.....	*	*	0.2	0.4	0.8	0.9	1.6	3.1	5.7	12.3
2003.....	*	*	*	*	0.7	0.9	1.6	3.1	5.3	13.5
2002.....	*	*	*	0.4	*	0.8	2.2	3.6	6.4	16.7
2001.....	*	*	*	0.4	0.4	1.1	1.8	3.3	6.4	13.3
2000.....	*	*	*	*	0.7	0.6	1.5	2.9	6.5	17.0
1999.....	*	*	*	*	0.5	1.1	1.7	3.6	7.0	18.9

White, female

2004.....	*	0.2	0.3	0.5	0.7	1.3	2.0	3.4	6.9	16.7
2003.....	*	0.2	0.3	0.5	0.7	1.3	2.2	3.3	7.3	17.8
2002.....	0.2	*	0.3	0.5	0.6	1.1	2.0	3.4	6.5	19.6
2001.....	*	0.2	0.3	0.5	0.7	0.9	1.8	3.6	7.4	20.6
2000.....	*	*	0.3	0.4	0.6	1.4	2.0	3.6	8.3	21.7
1999.....	0.2	0.3	0.3	0.4	0.6	1.1	1.9	3.9	8.1	21.9

Black, both sexes

2004.....	*	0.8	1.1	2.4	2.7	4.9	10.6	13.9	26.7	58.1
2003.....	*	0.9	1.5	2.2	3.1	4.1	11.1	14.8	27.4	63.6
2002.....	0.7	*	1.4	2.1	3.6	3.3	10.3	15.3	31.7	85.8
2001.....	*	*	1.0	1.7	2.6	4.5	8.2	16.8	29.3	79.2
2000.....	*	1.1	*	1.8	2.9	4.6	10.9	17.2	32.2	87.3
1999.....	0.8	0.9	1.5	1.9	3.1	5.9	10.1	22.0	36.4	96.3

Black, male

2004.....	*	*	*	*	*	*	10.7	12.7	25.1	48.7
2003.....	*	*	*	*	4.0	*	15.1	12.8	30.3	55.3
2002.....	*	*	*	*	*	*	12.7	15.2	29.8	56.8
2001.....	*	*	*	*	*	5.3	7.4	21.4	32.1	65.8
2000.....	*	*	*	*	*	*	12.7	18.0	36.2	72.7
1999.....	*	*	*	*	*	6.8	11.2	21.4	35.8	73.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)

Black, female

2004.....	*	*	*	3.0	2.9	5.6	10.5	14.6	27.6	61.8
2003.....	*	*	2.0	2.7	*	3.7	8.4	16.0	25.9	66.8
2002.....	*	*	*	2.7	4.0	*	8.7	15.4	32.6	96.9
2001.....	*	*	*	*	*	3.9	8.7	14.1	27.8	84.2
2000.....	*	1.7	*	*	*	4.8	9.8	16.8	30.1	92.7
1999.....	*	*	*	*	*	5.2	9.5	22.3	36.7	104.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

All races 3/, both sexes

2004.....	1.5	2.2	3.3	4.6	6.6	9.5	15.0	24.3	40.9	85.9
2003.....	1.6	2.2	3.3	4.4	7.0	9.8	15.9	25.3	40.5	90.4
2002.....	1.5	2.3	3.0	4.3	7.0	10.1	15.9	25.6	40.2	94.9
2001.....	1.7	2.1	3.1	4.2	6.6	10.1	15.8	25.4	42.0	93.4
2000.....	1.7	2.3	3.0	4.3	6.7	10.3	15.2	25.9	41.8	94.0
1999.....	1.6	2.4	3.1	4.3	6.7	10.1	15.5	25.5	39.3	91.2

All races 3/, male

2004.....	0.9	1.5	2.7	3.2	5.4	7.2	11.3	19.5	31.0	59.1
2003.....	1.2	1.5	2.4	3.3	5.1	8.4	12.6	20.8	30.1	59.9
2002.....	1.0	1.7	2.3	3.0	5.3	7.8	13.0	19.4	31.3	63.5
2001.....	1.1	1.5	2.4	3.2	4.7	7.1	12.3	20.1	32.4	63.2
2000.....	1.2	1.6	1.8	3.4	4.8	7.7	12.1	20.6	33.0	67.5
1999.....	1.1	1.5	2.1	3.1	4.7	6.9	12.5	20.2	30.9	63.7

All races 3/, female

2004.....	2.1	2.9	3.9	5.9	7.6	11.6	18.0	27.8	47.0	98.0
2003.....	2.1	3.0	4.2	5.5	8.8	11.0	18.5	28.5	46.8	103.8
2002.....	2.1	3.0	3.8	5.6	8.6	12.2	18.2	30.1	45.6	108.5
2001.....	2.2	2.8	3.7	5.2	8.3	12.7	18.5	29.1	47.8	106.0
2000.....	2.2	3.0	4.1	5.2	8.4	12.4	17.6	29.6	46.9	104.8
1999.....	2.1	3.3	4.1	5.5	8.5	12.8	18.0	29.3	44.3	102.3

White, both sexes

2004.....	1.1	1.6	2.7	4.0	6.2	9.2	15.0	24.4	41.8	88.8
2003.....	1.4	1.7	2.8	4.0	6.6	9.5	15.7	25.5	40.9	92.9
2002.....	1.1	1.8	2.7	3.9	6.6	9.9	15.8	25.9	40.7	98.2
2001.....	1.2	1.7	2.6	3.9	6.1	9.9	15.5	25.8	43.0	95.8
2000.....	1.2	1.7	2.5	3.8	6.3	10.2	15.0	26.0	42.5	96.9
1999.....	1.1	1.8	2.6	3.8	6.4	9.7	15.6	25.8	40.6	94.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

White, male

2004.....	0.7	1.1	2.2	2.9	5.2	6.9	10.9	19.3	31.4	59.8
2003.....	1.1	1.2	2.1	2.9	4.7	7.8	12.0	20.2	29.8	60.1
2002.....	0.8	1.4	2.1	2.7	5.0	7.3	12.7	19.4	31.0	64.3
2001.....	0.8	1.2	2.1	2.8	4.4	6.8	11.9	20.3	32.7	64.3
2000.....	0.9	1.3	1.6	3.2	4.5	7.6	11.3	20.2	32.9	67.3
1999.....	0.8	1.2	1.9	2.7	4.6	6.6	12.6	19.4	31.9	64.4

White, female

2004.....	1.5	2.2	3.1	5.2	7.1	11.2	18.5	28.2	48.2	101.8
2003.....	1.8	2.1	3.5	5.1	8.3	10.9	18.7	29.2	47.8	107.4
2002.....	1.4	2.3	3.2	5.0	8.1	12.2	18.4	30.6	46.6	112.9
2001.....	1.6	2.2	3.1	4.8	7.6	12.6	18.4	29.6	49.3	109.0
2000.....	1.5	2.0	3.4	4.4	7.9	12.6	17.9	30.1	48.2	108.9
1999.....	1.4	2.5	3.3	4.9	8.0	12.4	18.1	30.2	45.7	106.5

Black, both sexes

2004.....	4.0	6.6	8.2	9.4	11.2	14.5	17.7	27.1	36.4	64.9
2003.....	3.4	6.6	7.7	8.5	11.8	14.2	20.2	27.2	38.8	68.0
2002.....	4.5	6.0	6.5	8.9	11.4	13.5	18.8	25.6	39.0	67.4
2001.....	4.9	5.5	7.5	8.0	11.2	14.1	19.6	25.0	36.0	75.5
2000.....	5.4	7.2	6.9	8.9	11.1	12.4	18.7	26.8	37.6	68.2
1999.....	5.4	6.8	7.3	8.9	10.5	15.3	17.5	27.0	27.2	60.3

Black, male

2004.....	2.1	4.5	6.5	5.8	9.2	11.6	18.5	27.8	30.8	60.6
2003.....	2.0	3.5	4.9	6.7	10.0	15.8	21.0	30.6	38.4	58.4
2002.....	2.5	3.9	4.3	6.7	9.1	12.5	18.7	21.6	39.7	61.1
2001.....	3.4	4.2	5.7	6.6	9.2	11.7	17.2	22.8	36.9	59.0
2000.....	3.6	4.2	3.7	6.3	8.3	12.0	21.7	25.5	39.6	78.4
1999.....	3.4	4.0	4.2	6.4	6.8	12.2	14.2	33.7	20.9	63.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

Black, female

2004.....	5.8	8.3	9.7	12.4	12.8	16.5	17.1	26.7	39.3	66.5
2003.....	4.7	9.4	10.0	9.9	13.2	13.0	19.7	25.2	39.1	71.7
2002.....	6.3	7.8	8.4	10.6	13.2	14.2	18.8	28.0	38.6	69.8
2001.....	6.3	6.6	8.9	9.1	12.8	16.0	21.3	26.3	35.6	81.7
2000.....	7.1	9.8	9.6	11.1	13.2	12.7	16.6	27.6	36.5	64.4
1999.....	7.2	9.3	10.0	10.9	13.5	17.5	19.6	22.9	30.2	59.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)

All races 3/, both sexes

2004.....	*	0.2	0.3	0.7	1.1	2.3	4.1	6.8	9.6	11.9
2003.....	0.1	0.2	0.3	0.6	1.3	2.2	4.2	7.0	10.9	12.8
2002.....	*	0.2	0.3	0.7	1.4	2.5	4.2	8.0	10.6	14.2
2001.....	*	0.1	0.3	0.7	1.4	2.5	4.3	7.6	10.5	12.7
2000.....	*	0.2	0.3	0.6	1.4	2.7	4.4	7.5	11.6	12.9
1999.....	*	0.2	0.3	0.8	1.6	2.5	4.5	8.7	10.6	13.9

All races 3/, male

2004.....	*	*	*	0.4	0.9	1.6	2.2	4.1	4.6	6.5
2003.....	*	*	*	0.4	1.0	1.3	2.6	4.5	5.6	6.5
2002.....	*	*	*	0.5	1.0	1.7	2.6	4.5	6.6	7.6
2001.....	*	*	*	0.6	0.9	1.5	2.6	4.6	5.8	5.5
2000.....	*	*	*	0.4	0.9	1.7	3.1	4.4	6.6	6.9
1999.....	*	*	*	0.5	1.1	1.6	3.1	5.0	5.5	7.2

All races 3/, female

2004.....	*	0.2	0.4	0.9	1.4	2.9	5.6	8.7	12.7	14.3
2003.....	*	0.2	0.4	0.7	1.6	2.9	5.5	8.8	14.1	15.6
2002.....	*	0.2	0.4	0.9	1.8	3.2	5.4	10.4	13.1	17.0
2001.....	*	0.2	0.5	0.9	1.8	3.3	5.5	9.7	13.3	15.7
2000.....	*	0.3	0.5	0.8	1.9	3.5	5.4	9.7	14.6	15.3
1999.....	*	0.3	0.3	1.0	2.0	3.3	5.6	11.3	13.5	16.7

White, both sexes

2004.....	*	0.2	0.3	0.7	1.2	2.3	4.3	7.0	9.8	12.1
2003.....	*	0.2	0.3	0.6	1.4	2.2	4.5	7.3	11.1	13.2
2002.....	*	0.2	0.3	0.7	1.4	2.6	4.3	8.2	10.9	14.8
2001.....	*	0.1	0.3	0.7	1.4	2.7	4.4	8.0	10.9	12.8
2000.....	*	0.2	0.3	0.6	1.5	2.8	4.7	7.8	12.2	13.4
1999.....	*	0.2	0.3	0.8	1.7	2.6	4.8	9.0	11.0	14.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)

Black, female

2004.....	*	*	*	*	*	*	*	6.7	10.2	11.0
2003.....	*	*	*	*	*	*	*	5.4	11.6	11.3
2002.....	*	*	*	*	*	*	*	7.1	9.2	9.9
2001.....	*	*	*	*	*	*	*	*	*	14.7
2000.....	*	*	*	*	*	*	*	6.0	*	9.4
1999.....	*	*	*	*	*	*	*	6.3	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Systemic lupus erythematosus (M32)

All races 3/, both sexes

2004.....	0.5	0.5	0.8	0.8	0.8	0.7	1.0	1.2	1.5	1.1
2003.....	0.5	0.6	0.7	0.7	0.9	0.8	0.9	1.4	1.5	1.1
2002.....	0.5	0.6	0.6	0.7	1.0	0.9	1.0	1.3	1.4	1.3
2001.....	0.5	0.6	0.7	0.7	0.9	0.9	1.2	1.4	1.6	1.2
2000.....	0.6	0.6	0.7	0.8	1.0	1.0	0.9	1.4	1.5	1.2
1999.....	0.6	0.7	0.7	0.7	0.8	1.0	1.2	1.3	1.3	1.4

All races 3/, male

2004.....	*	*	0.3	*	*	*	*	*	*	*
2003.....	0.2	*	0.2	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	0.2	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.8	0.8	1.2	1.3	1.3	1.2	1.4	1.7	2.0	1.3
2003.....	0.8	0.9	1.2	1.3	1.5	1.2	1.4	2.2	2.1	1.3
2002.....	0.8	1.1	0.9	1.2	1.7	1.4	1.5	1.7	1.9	1.7
2001.....	0.8	1.1	1.1	1.1	1.5	1.5	1.8	2.2	2.0	1.3
2000.....	1.1	1.1	1.2	1.3	1.5	1.5	1.2	2.0	2.0	1.4
1999.....	1.0	1.2	1.2	1.2	1.4	1.7	1.8	1.8	1.7	1.6

White, both sexes

2004.....	0.3	0.3	0.5	0.6	0.6	0.6	0.9	1.1	1.4	1.2
2003.....	0.4	0.3	0.5	0.5	0.8	0.7	0.9	1.4	1.5	1.1
2002.....	0.3	0.4	0.4	0.5	0.7	0.7	1.0	1.3	1.4	1.4
2001.....	0.2	0.4	0.5	0.6	0.7	0.8	1.1	1.4	1.7	1.2
2000.....	0.4	0.3	0.5	0.6	0.8	0.9	0.8	1.4	1.6	1.3
1999.....	0.3	0.5	0.5	0.6	0.7	0.8	1.1	1.3	1.3	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Systemic lupus erythematosus (M32)										
Black, female										
2004.....	2.6	3.3	4.0	4.2	4.5	*	*	*	*	*
2003.....	2.3	3.8	3.8	3.6	*	*	*	*	*	*
2002.....	3.1	4.0	2.7	3.5	5.1	4.4	*	*	*	*
2001.....	3.4	3.3	3.2	2.7	3.5	*	*	*	*	*
2000.....	3.6	4.4	3.4	3.6	3.8	*	*	*	*	*
1999.....	3.9	3.9	3.3	3.2	*	4.8	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)

All races 3/, both sexes

2004.....	0.6	0.9	1.3	1.9	2.6	3.8	5.0	7.7	12.1	24.5
2003.....	0.7	0.9	1.3	1.8	2.8	3.8	5.6	8.3	10.9	25.9
2002.....	0.7	0.9	1.4	1.7	2.9	3.9	5.6	8.1	11.9	27.2
2001.....	0.7	0.9	1.3	1.7	2.7	4.1	5.9	8.2	13.0	27.7
2000.....	0.7	1.0	1.3	1.8	2.8	4.0	6.0	8.7	13.1	28.0
1999.....	0.7	1.1	1.5	1.8	2.9	4.2	6.0	7.9	11.9	27.6

All races 3/, male

2004.....	0.4	0.6	1.0	1.2	2.0	2.5	3.4	5.7	9.4	17.5
2003.....	0.5	0.6	0.9	1.3	1.8	2.9	4.2	7.2	8.4	19.4
2002.....	0.4	0.7	0.9	1.3	2.0	2.5	4.2	6.3	9.6	18.6
2001.....	0.5	0.8	1.0	1.1	2.0	2.3	4.6	6.5	10.6	20.5
2000.....	0.6	0.7	0.8	1.4	1.9	2.8	4.0	7.3	11.2	21.1
1999.....	0.5	0.7	1.1	1.3	1.9	2.8	4.4	6.5	10.0	22.5

All races 3/, female

2004.....	0.9	1.2	1.7	2.5	3.2	4.9	6.4	9.0	13.7	27.7
2003.....	0.9	1.2	1.7	2.3	3.8	4.5	6.7	9.0	12.4	28.8
2002.....	0.9	1.2	1.9	2.2	3.6	5.1	6.7	9.4	13.3	31.0
2001.....	1.0	1.1	1.5	2.2	3.3	5.6	7.0	9.4	14.4	30.7
2000.....	0.9	1.2	1.7	2.2	3.5	5.1	7.5	9.6	14.3	30.8
1999.....	0.8	1.4	1.9	2.3	3.8	5.4	7.2	8.8	13.0	29.7

White, both sexes

2004.....	0.5	0.6	1.1	1.6	2.5	3.6	5.1	7.7	12.5	25.5
2003.....	0.6	0.6	1.0	1.6	2.6	3.7	5.5	8.3	11.1	26.4
2002.....	0.5	0.7	1.2	1.6	2.8	4.0	5.5	8.3	12.1	28.0
2001.....	0.5	0.7	1.0	1.6	2.5	3.9	5.9	8.5	13.4	28.6
2000.....	0.5	0.7	1.0	1.6	2.6	4.1	5.9	8.8	13.4	28.8
1999.....	0.5	0.7	1.1	1.6	2.6	4.1	5.9	8.0	12.2	28.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)

White, male										
2004.....	0.3	0.3	0.9	1.0	1.9	2.4	3.4	5.8	9.8	18.0
2003.....	0.5	0.5	0.7	1.1	1.6	2.7	4.0	7.1	8.2	19.9
2002.....	0.3	0.5	0.8	1.1	1.9	2.5	4.0	6.5	9.7	19.1
2001.....	0.3	0.5	0.8	1.0	1.9	2.2	4.5	6.7	10.8	21.1
2000.....	0.4	0.6	0.7	1.3	1.7	2.7	3.8	7.2	11.1	20.6
1999.....	0.3	0.5	0.9	1.1	1.8	2.7	4.4	6.4	10.4	23.3
White, female										
2004.....	0.6	0.8	1.4	2.1	3.1	4.7	6.5	9.1	14.3	28.9
2003.....	0.7	0.8	1.4	2.1	3.5	4.5	6.7	9.3	12.8	29.2
2002.....	0.6	0.9	1.6	2.0	3.7	5.3	6.8	9.6	13.6	31.9
2001.....	0.8	0.9	1.2	2.1	3.1	5.5	7.0	9.8	14.9	31.7
2000.....	0.6	0.8	1.4	1.9	3.4	5.3	7.5	10.0	14.7	32.1
1999.....	0.6	1.0	1.4	2.1	3.4	5.3	7.2	9.2	13.3	30.4
Black, both sexes										
2004.....	1.7	3.1	3.3	3.8	4.2	5.3	5.4	6.9	8.9	16.6
2003.....	1.5	2.8	3.6	3.6	5.1	5.2	6.9	8.7	9.9	20.4
2002.....	2.1	2.5	3.0	3.6	4.1	4.4	7.0	7.4	10.7	21.7
2001.....	2.2	2.7	3.6	2.7	3.7	6.0	6.1	6.4	11.6	22.4
2000.....	2.3	3.0	3.3	3.6	4.4	4.0	7.9	7.5	12.1	21.9
1999.....	2.4	3.3	4.1	3.7	5.9	5.3	7.0	7.3	8.7	20.4
Black, male										
2004.....	*	2.3	2.4	*	*	*	*	*	*	*
2003.....	*	*	2.4	3.1	*	5.7	*	10.5	*	*
2002.....	*	2.0	2.2	3.6	4.2	*	*	*	*	*
2001.....	1.6	2.6	2.9	*	*	*	*	*	*	*
2000.....	1.8	*	*	*	4.1	*	*	*	*	27.7
1999.....	2.0	2.3	2.5	*	*	*	*	9.5	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)

Black, female

2004.....	2.5	3.7	4.1	4.9	4.7	6.7	5.9	8.1	9.8	17.7
2003.....	2.0	4.2	4.5	4.0	6.2	4.9	7.3	7.6	9.7	23.5
2002.....	2.8	3.0	3.8	3.5	4.0	4.8	7.4	8.8	11.9	24.6
2001.....	2.7	2.8	4.2	3.3	4.5	7.3	7.2	6.7	12.7	24.4
2000.....	2.8	4.2	4.5	4.3	4.6	4.0	8.9	7.1	10.6	19.7
1999.....	2.7	4.2	5.5	4.3	7.3	6.3	8.1	6.0	10.4	22.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Dorsopathies (M40-M54)										
All races 3/, both sexes										
2004.....	*	0.1	0.2	0.3	0.5	0.6	1.2	2.0	3.4	7.1
2003.....	*	0.1	0.2	0.3	0.5	0.7	1.1	2.2	3.4	5.7
2002.....	*	0.1	0.2	0.3	0.4	0.6	1.2	1.7	2.9	6.0
2001.....	*	*	0.2	0.3	0.3	0.5	1.0	1.7	3.3	7.4
2000.....	*	0.1	0.1	0.2	0.4	0.7	0.9	1.7	3.3	6.1
1999.....	*	0.1	0.2	0.2	0.3	0.6	0.9	1.5	2.5	5.4
All races 3/, male										
2004.....	*	*	0.2	0.3	0.6	0.6	1.3	2.3	3.3	6.6
2003.....	*	*	0.3	0.4	0.4	0.9	1.1	2.4	3.6	4.8
2002.....	*	*	0.3	0.3	0.4	0.5	1.3	2.0	3.1	5.5
2001.....	*	*	*	0.3	0.4	0.6	0.9	1.9	3.5	6.4
2000.....	*	*	*	*	0.4	0.8	1.2	1.5	3.8	6.5
1999.....	*	*	*	*	0.5	0.7	1.0	1.8	2.8	5.9
All races 3/, female										
2004.....	*	*	*	0.3	0.4	0.5	1.1	1.7	3.4	7.3
2003.....	*	*	*	0.3	0.5	0.5	1.2	2.0	3.3	6.1
2002.....	*	*	*	0.3	0.4	0.7	1.1	1.5	2.8	6.2
2001.....	*	*	*	*	*	0.5	1.1	1.6	3.2	7.8
2000.....	*	*	*	*	*	0.5	0.7	1.9	3.1	6.0
1999.....	*	*	0.2	*	*	0.5	0.9	1.3	2.4	5.2
White, both sexes										
2004.....	*	*	0.2	0.3	0.5	0.5	1.2	2.1	3.5	7.4
2003.....	*	0.1	0.2	0.3	0.4	0.7	1.0	2.1	3.5	5.9
2002.....	*	0.1	0.2	0.3	0.4	0.7	1.2	1.7	3.0	6.2
2001.....	*	*	0.2	0.2	0.3	0.5	1.0	1.7	3.3	7.8
2000.....	*	0.1	*	0.2	0.4	0.7	0.8	1.7	3.3	6.4
1999.....	*	*	0.2	0.2	0.3	0.5	0.9	1.5	2.7	5.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Soft tissue disorders (M60-M79)

All races 3/, both sexes

2004.....	0.2	0.4	0.5	0.6	0.7	0.9	1.5	2.1	3.4	5.1
2003.....	0.2	0.3	0.5	0.5	0.8	1.1	1.5	2.1	3.3	5.4
2002.....	0.2	0.3	0.4	0.5	0.6	0.9	1.3	1.9	3.1	5.3
2001.....	0.2	0.2	0.4	0.5	0.7	0.9	1.2	2.0	2.8	4.3
2000.....	0.2	0.4	0.4	0.5	0.6	0.8	1.2	2.0	2.3	4.2
1999.....	0.2	0.2	0.3	0.5	0.5	0.8	1.2	1.7	2.7	4.3

All races 3/, male

2004.....	0.2	0.4	0.6	0.7	0.8	0.9	1.6	2.6	4.5	7.8
2003.....	0.2	0.3	0.5	0.4	0.7	1.1	1.9	2.6	4.5	7.1
2002.....	0.3	0.4	0.4	0.4	0.8	1.3	1.6	2.0	4.3	7.8
2001.....	0.3	0.3	0.6	0.6	0.6	1.1	1.6	2.7	3.9	6.2
2000.....	0.2	0.4	0.3	0.6	0.5	0.9	1.6	2.6	3.5	6.7
1999.....	0.3	0.3	0.3	0.6	0.5	0.8	1.7	2.5	4.0	6.8

All races 3/, female

2004.....	*	0.4	0.3	0.5	0.6	0.9	1.4	1.7	2.7	3.9
2003.....	*	0.3	0.5	0.6	0.8	1.0	1.2	1.7	2.5	4.6
2002.....	*	*	0.3	0.6	0.5	0.6	1.1	1.8	2.4	4.2
2001.....	*	*	0.3	0.5	0.7	0.6	0.9	1.4	2.1	3.5
2000.....	*	0.3	0.4	0.4	0.6	0.8	0.9	1.6	1.7	3.2
1999.....	*	0.2	0.3	0.4	0.5	0.8	0.7	1.2	2.0	3.3

White, both sexes

2004.....	0.2	0.4	0.4	0.5	0.6	0.8	1.5	2.0	3.3	4.8
2003.....	0.2	0.2	0.5	0.5	0.6	1.0	1.5	1.9	3.2	5.4
2002.....	0.2	0.2	0.4	0.4	0.6	0.9	1.2	1.8	3.1	5.0
2001.....	0.2	0.2	0.4	0.5	0.5	0.8	1.1	1.8	2.7	4.2
2000.....	0.1	0.3	0.3	0.5	0.5	0.8	1.1	1.7	2.3	4.0
1999.....	0.2	0.2	0.3	0.4	0.4	0.7	1.1	1.7	2.6	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)

All races 3/, both sexes

2004.....	0.1	0.2	0.2	0.4	0.8	1.3	2.2	4.7	11.0	36.2
2003.....	0.1	0.2	0.3	0.5	0.7	1.3	2.5	4.3	10.5	39.5
2002.....	*	0.1	0.2	0.4	0.8	1.2	2.6	4.7	10.2	40.8
2001.....	*	0.2	0.2	0.3	0.7	1.3	2.2	4.4	10.8	40.1
2000.....	*	*	0.2	0.3	0.6	1.1	1.8	4.6	9.8	41.6
1999.....	*	0.1	0.2	0.4	0.5	1.0	1.8	4.4	10.3	38.6

All races 3/, male

2004.....	*	0.3	0.4	0.4	0.9	1.4	2.4	4.3	8.5	20.0
2003.....	*	0.2	0.3	0.7	0.8	1.8	2.5	3.7	7.5	21.5
2002.....	*	*	0.3	0.4	0.9	1.4	2.9	4.1	7.2	23.6
2001.....	*	*	0.2	0.5	0.7	1.3	2.1	4.0	7.7	23.6
2000.....	*	*	*	0.4	0.6	1.2	1.8	4.2	7.1	25.6
1999.....	*	*	*	0.3	0.5	0.8	1.8	3.7	7.8	20.4

All races 3/, female

2004.....	*	0.2	*	0.4	0.8	1.2	2.1	4.9	12.5	43.5
2003.....	*	0.2	0.2	0.3	0.7	0.9	2.6	4.8	12.4	47.5
2002.....	*	*	*	0.4	0.7	1.1	2.4	5.2	12.0	48.3
2001.....	*	*	*	*	0.7	1.3	2.2	4.8	12.7	47.0
2000.....	*	*	*	*	0.5	0.9	1.9	4.8	11.3	48.1
1999.....	*	*	*	0.4	0.6	1.2	1.7	4.8	11.8	45.9

White, both sexes

2004.....	0.1	0.2	0.2	0.4	0.8	1.3	2.1	4.5	11.1	37.7
2003.....	0.1	0.2	0.3	0.4	0.7	1.2	2.5	4.4	10.6	41.0
2002.....	*	0.1	0.2	0.4	0.8	1.1	2.7	4.6	10.1	42.8
2001.....	*	0.1	0.2	0.3	0.7	1.2	2.0	4.5	11.0	41.3
2000.....	*	*	0.2	0.3	0.5	1.0	1.8	4.6	9.8	43.0
1999.....	*	0.1	0.2	0.3	0.5	0.9	1.7	4.2	10.7	40.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)

Black, female

2004.....	*	*	*	*	*	*	*	5.7	11.3	26.0
2003.....	*	*	*	*	*	*	*	*	9.3	26.3
2002.....	*	*	*	*	*	*	*	6.3	10.3	23.4
2001.....	*	*	*	*	*	*	4.7	*	9.4	31.2
2000.....	*	*	*	*	*	*	*	*	10.2	25.8
1999.....	*	*	*	*	*	*	*	5.7	8.6	20.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Osteoporosis (M80-M81)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	0.6	1.7	5.3	24.4
2003.....	*	*	*	*	*	*	0.7	1.7	5.4	26.9
2002.....	*	*	*	*	*	0.2	0.6	1.9	5.4	29.0
2001.....	*	*	*	*	*	0.3	0.5	1.7	5.6	26.8
2000.....	*	*	*	*	*	0.3	0.4	2.1	5.5	28.5
1999.....	*	*	*	*	*	0.3	0.5	2.0	5.8	27.1

All races 3/, male

2004.....	*	*	*	*	*	*	*	0.7	1.9	7.2
2003.....	*	*	*	*	*	*	*	*	1.8	7.9
2002.....	*	*	*	*	*	*	*	0.7	1.7	9.9
2001.....	*	*	*	*	*	*	*	*	1.6	9.3
2000.....	*	*	*	*	*	*	*	0.9	1.9	9.9
1999.....	*	*	*	*	*	*	*	0.7	2.2	7.0

All races 3/, female

2004.....	*	*	*	*	*	*	0.8	2.4	7.4	32.2
2003.....	*	*	*	*	*	*	1.1	2.5	7.6	35.2
2002.....	*	*	*	*	*	*	1.0	2.9	7.6	37.2
2001.....	*	*	*	*	*	0.4	0.6	2.4	8.0	34.1
2000.....	*	*	*	*	*	0.4	0.6	2.9	7.7	36.0
1999.....	*	*	*	*	*	0.4	0.7	2.9	7.9	35.1

White, both sexes

2004.....	*	*	*	*	*	*	0.6	1.9	5.7	26.3
2003.....	*	*	*	*	*	*	0.8	1.8	5.8	29.1
2002.....	*	*	*	*	*	*	0.7	2.0	5.8	31.3
2001.....	*	*	*	*	*	0.3	0.5	1.8	6.0	28.8
2000.....	*	*	*	*	*	0.3	0.4	2.3	5.9	30.7
1999.....	*	*	*	*	*	0.3	0.5	2.2	6.2	29.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)

All races 3/, both sexes

2004.....	0.1	0.2	0.2	0.4	0.7	1.2	1.7	3.0	5.7	11.8
2003.....	0.1	0.2	0.2	0.5	0.7	1.2	1.8	2.7	5.1	12.6
2002.....	*	0.1	0.2	0.4	0.7	1.0	2.0	2.8	4.8	11.8
2001.....	*	0.2	0.2	0.3	0.6	1.0	1.7	2.8	5.3	13.3
2000.....	*	*	0.2	0.3	0.5	0.8	1.4	2.5	4.3	13.1
1999.....	*	0.1	0.2	0.3	0.4	0.7	1.3	2.4	4.6	11.5

All races 3/, male

2004.....	*	0.2	0.3	0.3	0.8	1.3	2.1	3.6	6.6	12.9
2003.....	*	0.2	0.3	0.6	0.7	1.7	2.1	3.3	5.6	13.6
2002.....	*	*	0.3	0.4	0.9	1.3	2.6	3.4	5.5	13.7
2001.....	*	*	0.2	0.4	0.7	1.2	1.8	3.3	6.1	14.2
2000.....	*	*	*	0.4	0.6	1.2	1.6	3.4	5.2	15.6
1999.....	*	*	*	0.3	0.5	0.7	1.7	3.0	5.7	13.3

All races 3/, female

2004.....	*	0.2	*	0.4	0.7	1.0	1.4	2.5	5.1	11.3
2003.....	*	0.2	0.2	0.3	0.6	0.7	1.5	2.3	4.8	12.2
2002.....	*	*	*	0.4	0.6	0.7	1.4	2.3	4.4	11.1
2001.....	*	*	*	*	0.5	0.8	1.7	2.4	4.8	12.9
2000.....	*	*	*	*	0.4	0.5	1.3	1.9	3.7	12.0
1999.....	*	*	*	0.3	0.4	0.7	1.0	1.9	3.9	10.8

White, both sexes

2004.....	*	0.2	0.2	0.3	0.7	1.1	1.5	2.7	5.4	11.4
2003.....	*	0.2	0.2	0.4	0.6	1.0	1.7	2.6	4.8	11.9
2002.....	*	0.1	0.2	0.3	0.7	0.8	2.0	2.5	4.4	11.5
2001.....	*	0.1	0.2	0.3	0.5	0.9	1.5	2.6	5.0	12.5
2000.....	*	*	0.1	0.3	0.4	0.7	1.3	2.3	3.8	12.4
1999.....	*	0.1	0.2	0.3	0.5	0.6	1.2	2.0	4.5	11.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XIV. Diseases of the genitourinary system (N00-N98)

All races 3/, both sexes

2004.....	2.7	4.6	7.9	13.0	21.2	37.3	63.2	115.4	204.1	458.5
2003.....	2.8	4.6	7.6	12.6	22.2	37.9	65.1	114.1	208.5	469.5
2002.....	2.6	4.5	7.1	12.2	20.5	36.5	64.2	115.2	206.9	469.4
2001.....	2.6	4.5	7.1	12.1	20.6	39.2	64.4	113.7	202.0	476.4
2000.....	2.6	4.3	6.8	11.5	20.7	36.4	60.5	111.8	202.9	471.4
1999.....	2.5	3.9	6.4	11.2	19.8	35.5	61.7	109.5	197.7	471.1

All races 3/, male

2004.....	3.0	5.3	9.0	14.5	23.2	40.8	69.7	132.9	246.3	533.9
2003.....	3.1	5.1	8.9	14.2	23.8	41.3	71.9	132.9	253.9	547.9
2002.....	2.7	5.1	8.1	12.9	22.3	40.4	73.7	134.0	246.8	553.0
2001.....	2.7	5.0	7.8	12.9	21.7	44.0	71.6	129.8	242.7	564.2
2000.....	2.9	5.1	7.6	12.6	22.1	39.6	67.3	131.0	243.1	567.5
1999.....	2.7	4.5	6.9	12.1	22.0	38.3	70.3	130.4	241.8	558.3

All races 3/, female

2004.....	2.4	4.0	6.8	11.6	19.4	34.2	58.0	102.9	178.1	424.5
2003.....	2.6	4.2	6.3	11.1	20.6	35.0	59.6	100.7	180.8	434.9
2002.....	2.4	4.0	6.2	11.6	18.9	33.1	56.7	102.0	182.6	433.2
2001.....	2.4	3.9	6.3	11.4	19.6	35.0	58.6	102.4	177.6	439.6
2000.....	2.2	3.4	6.0	10.4	19.4	33.7	55.2	98.5	179.1	432.3
1999.....	2.3	3.3	5.9	10.4	17.9	33.1	54.9	95.0	171.9	436.0

White, both sexes

2004.....	2.1	3.3	5.8	9.8	16.9	31.5	56.3	107.0	196.2	449.9
2003.....	2.0	3.3	5.4	9.7	17.7	31.9	57.1	106.2	199.8	461.3
2002.....	1.9	3.3	5.0	9.4	16.4	30.9	56.8	106.8	198.9	458.5
2001.....	1.9	3.1	5.1	9.2	16.2	32.5	57.2	105.3	192.3	465.1
2000.....	1.9	3.0	4.9	8.7	16.5	30.2	52.8	103.9	193.4	460.5
1999.....	1.7	2.7	4.6	8.8	15.8	29.3	54.0	101.0	187.8	456.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XIV. Diseases of the genitourinary system (N00-N98)

White, male

2004.....	2.4	3.7	6.4	11.2	18.6	34.3	63.1	124.6	240.8	527.3
2003.....	2.3	3.7	6.1	10.9	19.2	35.1	63.6	125.4	244.3	547.4
2002.....	2.0	3.6	5.6	9.8	17.5	34.7	65.9	125.5	240.2	543.2
2001.....	2.0	3.4	5.8	9.8	17.5	37.1	64.8	122.6	233.9	556.8
2000.....	2.1	3.5	5.4	9.5	17.9	33.7	60.4	123.0	234.8	561.2
1999.....	1.8	3.0	4.9	9.9	17.5	32.0	62.2	120.6	230.8	542.8

White, female

2004.....	1.7	2.9	5.1	8.4	15.4	29.0	50.7	94.1	168.5	415.0
2003.....	1.6	3.0	4.8	8.7	16.4	29.2	51.8	92.3	172.3	423.2
2002.....	1.8	2.9	4.4	9.0	15.4	27.5	49.3	93.5	173.5	421.7
2001.....	1.7	2.8	4.5	8.6	14.9	28.5	51.0	93.0	167.1	426.7
2000.....	1.6	2.6	4.4	7.9	15.3	27.0	46.7	90.5	168.8	419.6
1999.....	1.6	2.3	4.2	7.7	14.1	27.0	47.5	87.2	162.4	421.1

Black, both sexes

2004.....	7.7	14.6	25.0	41.6	63.0	92.4	140.9	222.4	331.0	634.5
2003.....	9.0	14.0	24.7	39.6	64.5	96.3	153.2	219.0	346.7	645.3
2002.....	7.6	13.6	24.1	37.9	58.7	91.9	150.1	225.4	332.3	662.5
2001.....	7.6	14.5	23.0	36.9	60.3	102.5	146.3	223.5	347.5	664.5
2000.....	7.5	14.0	22.6	36.6	58.1	96.6	145.9	219.5	343.1	637.5
1999.....	7.9	13.5	22.7	32.7	56.2	94.5	147.3	219.9	337.2	679.1

Black, male

2004.....	8.1	17.0	31.2	47.1	70.6	107.1	156.0	260.2	366.6	718.3
2003.....	9.2	15.5	33.3	47.7	73.7	108.3	181.0	249.7	437.1	681.9
2002.....	8.3	16.4	29.0	44.4	70.1	101.7	177.3	262.7	382.1	794.4
2001.....	7.8	18.3	25.9	42.9	63.3	118.4	158.5	241.6	404.5	755.7
2000.....	8.9	18.8	27.3	42.8	63.2	102.3	155.8	259.4	398.7	716.2
1999.....	9.2	17.3	26.4	32.9	65.8	102.0	172.3	270.7	433.4	802.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XIV. Diseases of the genitourinary system (N00-N98)

Black, female

2004.....	7.5	12.4	19.7	37.1	57.1	81.5	130.7	200.0	312.2	601.3
2003.....	8.7	12.7	17.3	32.9	57.3	87.4	134.5	200.8	299.4	631.1
2002.....	7.0	11.1	20.0	32.6	49.9	84.5	132.0	203.1	306.4	612.0
2001.....	7.4	11.3	20.5	32.1	57.9	90.7	138.1	212.7	318.5	630.6
2000.....	6.3	9.9	18.6	31.6	54.1	92.4	139.4	195.4	315.0	608.3
1999.....	6.7	10.2	19.5	32.5	48.8	88.9	130.7	189.4	289.5	633.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of urinary system (N00-N39)

All races 3/, both sexes

2004.....	2.7	4.5	7.7	12.7	20.9	36.7	62.2	113.8	200.2	448.9
2003.....	2.7	4.5	7.5	12.4	21.9	37.3	64.3	112.4	205.2	460.5
2002.....	2.5	4.4	7.0	12.1	20.3	36.1	63.5	113.9	203.4	460.8
2001.....	2.5	4.3	6.9	11.9	20.4	38.7	63.3	112.0	198.9	467.4
2000.....	2.5	4.2	6.6	11.3	20.4	35.9	59.7	110.2	199.3	462.5
1999.....	2.4	3.8	6.3	11.0	19.5	35.0	60.8	107.8	194.1	462.6

All races 3/, male

2004.....	3.0	5.1	8.8	14.2	22.7	40.0	68.3	130.1	238.8	513.6
2003.....	3.1	5.0	8.8	13.9	23.5	40.5	70.8	130.3	247.9	528.3
2002.....	2.7	5.0	8.0	12.8	21.9	39.8	72.4	132.0	239.9	533.8
2001.....	2.7	5.0	7.7	12.7	21.5	43.5	70.2	127.2	237.2	542.9
2000.....	2.9	5.1	7.5	12.4	21.7	39.0	66.0	128.5	236.8	546.9
1999.....	2.6	4.5	6.8	11.9	21.6	37.6	69.0	127.7	234.6	537.0

All races 3/, female

2004.....	2.3	3.9	6.7	11.4	19.2	33.8	57.3	102.1	176.4	419.9
2003.....	2.4	4.0	6.1	11.0	20.3	34.5	59.1	99.6	179.2	430.5
2002.....	2.3	3.8	6.1	11.4	18.8	32.9	56.3	101.2	181.1	429.1
2001.....	2.3	3.7	6.1	11.2	19.4	34.6	57.8	101.3	175.9	435.8
2000.....	2.1	3.3	5.8	10.3	19.3	33.3	54.7	97.5	177.2	428.1
1999.....	2.1	3.1	5.8	10.2	17.6	32.8	54.4	93.9	170.4	432.6

White, both sexes

2004.....	2.0	3.2	5.6	9.6	16.6	31.0	55.4	105.4	192.4	440.3
2003.....	1.9	3.2	5.4	9.6	17.5	31.4	56.3	104.4	196.4	451.9
2002.....	1.8	3.2	4.9	9.2	16.2	30.5	56.1	105.6	195.3	449.9
2001.....	1.8	3.0	5.0	9.0	16.0	32.1	56.1	103.6	189.1	456.0
2000.....	1.8	2.9	4.8	8.6	16.3	29.7	51.9	102.4	189.9	451.6
1999.....	1.6	2.6	4.5	8.6	15.5	28.9	53.3	99.4	184.2	447.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of urinary system (N00-N39)

White, male

2004.....	2.4	3.6	6.3	11.0	18.1	33.6	61.7	122.1	233.4	506.5
2003.....	2.3	3.7	6.1	10.7	18.9	34.4	62.4	122.9	238.2	527.0
2002.....	2.0	3.6	5.6	9.7	17.2	34.2	64.7	123.7	233.4	523.8
2001.....	2.0	3.3	5.7	9.7	17.4	36.5	63.4	119.9	228.5	535.0
2000.....	2.1	3.5	5.4	9.4	17.5	33.2	59.3	120.6	228.4	540.8
1999.....	1.8	3.0	4.8	9.8	17.2	31.4	61.1	117.9	223.8	521.3

White, female

2004.....	1.6	2.8	5.0	8.2	15.1	28.7	50.2	93.2	166.9	410.4
2003.....	1.5	2.8	4.7	8.5	16.2	28.7	51.4	91.1	170.6	418.6
2002.....	1.7	2.8	4.3	8.8	15.2	27.3	49.0	92.6	172.0	417.8
2001.....	1.6	2.7	4.4	8.4	14.8	28.2	50.2	92.0	165.3	423.0
2000.....	1.5	2.4	4.2	7.8	15.2	26.7	46.1	89.5	167.0	415.4
1999.....	1.5	2.2	4.1	7.5	13.9	26.7	47.1	86.3	160.9	417.7

Black, both sexes

2004.....	7.6	14.1	24.5	41.2	62.6	91.3	138.6	219.7	327.0	625.5
2003.....	8.7	13.8	24.2	39.2	63.5	95.1	152.3	216.8	344.7	638.9
2002.....	7.5	13.3	23.8	37.8	58.0	90.9	148.1	222.8	328.9	652.1
2001.....	7.3	14.2	22.5	36.7	59.1	101.9	144.9	220.6	343.2	655.3
2000.....	7.2	13.9	22.5	36.1	57.6	95.6	144.5	217.0	339.1	628.5
1999.....	7.7	13.1	22.3	32.2	55.4	93.3	145.0	216.7	332.3	669.3

Black, male

2004.....	8.0	16.9	30.7	46.4	69.9	105.6	153.4	253.9	359.4	701.4
2003.....	9.2	15.4	32.9	47.1	72.7	106.9	180.3	244.7	432.6	666.2
2002.....	8.3	16.4	28.8	44.4	68.9	100.5	173.6	256.7	373.7	776.2
2001.....	7.6	18.1	25.4	42.5	61.8	117.9	156.5	237.9	395.7	736.4
2000.....	8.8	18.8	27.0	42.0	62.1	101.2	153.5	254.2	392.0	695.4
1999.....	9.2	17.2	25.7	32.2	64.5	100.5	168.9	268.3	421.2	779.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of urinary system (N00-N39)

Black, female

2004.....	7.4	11.8	19.4	36.9	57.0	80.6	128.5	199.5	310.0	595.4
2003.....	8.3	12.4	16.9	32.8	56.4	86.4	133.4	200.2	298.6	628.3
2002.....	6.7	10.7	19.6	32.4	49.4	83.8	131.1	202.5	305.6	604.6
2001.....	7.0	10.8	19.9	31.8	57.0	90.1	137.2	210.2	316.4	625.2
2000.....	5.8	9.6	18.6	31.3	54.1	91.4	138.5	194.5	312.5	603.5
1999.....	6.4	9.7	19.3	32.3	48.3	87.9	129.2	185.7	288.2	628.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Glomerular and renal tubulo-interstitial diseases (N00-N15)

All races 3/, both sexes

2004.....	0.2	0.2	0.3	0.5	0.6	1.1	1.9	3.5	6.4	12.6
2003.....	0.2	0.3	0.4	0.5	0.6	1.3	1.9	3.2	6.9	13.3
2002.....	0.1	0.2	0.3	0.5	0.6	1.1	2.0	3.9	6.2	13.9
2001.....	0.1	0.2	0.3	0.4	0.6	1.2	2.0	3.8	5.8	13.7
2000.....	0.2	0.2	0.3	0.5	0.7	1.2	1.9	3.7	6.5	15.7
1999.....	0.2	0.2	0.5	0.6	0.8	1.3	2.2	4.2	7.2	16.4

All races 3/, male

2004.....	*	0.2	0.3	0.6	0.7	1.1	2.0	3.8	7.1	16.9
2003.....	*	0.3	0.4	0.4	0.6	1.1	1.8	3.5	7.7	17.8
2002.....	*	*	0.3	0.5	0.6	1.2	2.3	4.7	6.7	18.1
2001.....	*	*	0.3	0.4	0.5	1.3	2.4	3.9	7.6	17.2
2000.....	*	0.2	0.4	0.5	0.7	1.2	1.6	4.3	7.2	20.0
1999.....	0.2	0.2	0.4	0.6	0.8	1.2	2.0	4.8	9.8	22.6

All races 3/, female

2004.....	0.2	0.3	0.4	0.4	0.5	1.0	1.8	3.3	5.9	10.7
2003.....	0.2	0.2	0.3	0.5	0.7	1.4	1.9	3.0	6.4	11.3
2002.....	*	0.3	0.3	0.4	0.7	1.1	1.7	3.3	6.0	12.0
2001.....	*	0.2	0.4	0.4	0.7	1.2	1.6	3.7	4.7	12.3
2000.....	*	0.3	0.3	0.5	0.7	1.2	2.2	3.3	6.1	13.9
1999.....	*	0.3	0.5	0.6	0.7	1.3	2.5	3.8	5.6	13.9

White, both sexes

2004.....	0.2	0.2	0.3	0.4	0.5	1.0	1.9	3.4	6.4	12.4
2003.....	0.2	0.2	0.3	0.4	0.5	1.3	1.8	3.1	6.6	13.2
2002.....	0.2	0.2	0.3	0.4	0.6	1.2	2.1	3.8	6.2	13.9
2001.....	0.1	0.1	0.3	0.4	0.5	1.1	2.0	3.8	5.5	13.9
2000.....	0.1	0.2	0.2	0.5	0.6	1.1	1.8	3.5	6.3	15.2
1999.....	0.1	0.3	0.4	0.5	0.6	1.1	2.1	4.2	7.1	16.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Glomerular diseases (N00-N07)

All races 3/, both sexes

2004.....	*	*	0.2	0.2	0.3	0.5	0.7	1.3	2.0	3.3
2003.....	*	*	0.2	0.2	0.3	0.5	0.8	1.1	2.1	3.0
2002.....	*	0.1	0.1	0.2	0.2	0.4	0.7	1.3	2.2	3.3
2001.....	*	*	0.2	0.2	0.2	0.6	0.9	1.2	1.8	3.3
2000.....	0.1	0.1	0.2	0.2	0.3	0.6	0.8	1.2	2.4	3.6
1999.....	*	0.1	0.2	0.2	0.4	0.5	0.9	1.7	2.3	3.7

All races 3/, male

2004.....	*	*	*	*	0.4	0.5	0.8	1.5	2.5	4.2
2003.....	*	*	0.2	*	*	0.7	0.8	1.3	2.5	3.9
2002.....	*	*	*	0.3	*	0.5	0.9	1.6	2.4	4.3
2001.....	*	*	*	*	*	0.7	1.1	1.5	2.6	4.6
2000.....	*	*	0.3	*	0.4	0.7	0.8	1.7	2.6	5.6
1999.....	*	*	*	*	0.5	0.5	0.8	2.2	3.4	5.6

All races 3/, female

2004.....	*	*	*	*	*	0.5	0.6	1.1	1.6	2.9
2003.....	*	*	*	*	*	0.4	0.9	1.0	1.9	2.6
2002.....	*	*	*	*	*	*	0.6	1.1	2.0	2.8
2001.....	*	*	*	*	*	0.6	0.7	1.0	1.3	2.8
2000.....	*	*	*	*	*	0.5	0.8	0.8	2.3	2.8
1999.....	*	*	*	*	0.4	0.5	1.0	1.4	1.7	2.9

White, both sexes

2004.....	*	*	0.2	0.2	0.3	0.4	0.7	1.2	1.9	3.2
2003.....	*	*	0.1	0.2	0.3	0.5	0.7	1.0	2.0	2.8
2002.....	*	*	*	0.2	0.2	0.4	0.7	1.2	2.0	3.0
2001.....	*	*	*	*	*	0.5	0.9	1.2	1.7	3.3
2000.....	*	*	*	0.2	0.3	0.5	0.8	1.1	2.4	3.3
1999.....	*	*	0.1	0.2	0.3	0.4	0.9	1.7	2.1	3.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nephrotic syndrome (N04)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	*	0.4	0.6
2003.....	*	*	*	*	*	*	*	*	0.5	0.5
2002.....	*	*	*	*	*	*	*	*	0.6	0.7
2001.....	*	*	*	*	*	*	*	0.3	*	0.7
2000.....	*	*	*	*	*	*	*	*	0.6	1.0
1999.....	*	*	*	*	*	*	0.3	0.4	0.5	0.7

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	0.6
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	0.7
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	0.6	0.8
1999.....	*	*	*	*	*	*	*	0.5	*	*

White, both sexes

2004.....	*	*	*	*	*	*	*	*	*	0.6
2003.....	*	*	*	*	*	*	*	*	0.5	*
2002.....	*	*	*	*	*	*	*	*	0.5	0.6
2001.....	*	*	*	*	*	*	*	0.3	*	0.6
2000.....	*	*	*	*	*	*	*	*	0.6	0.9
1999.....	*	*	*	*	*	*	0.3	0.4	*	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified glomerular diseases (N02-N03,N05-N07)										
All races 3/, both sexes										
2004.....	*	*	0.1	0.2	0.3	0.4	0.5	0.9	1.5	2.5
2003.....	*	*	0.1	*	0.2	0.4	0.6	0.9	1.5	2.5
2002.....	*	0.1	0.1	0.2	*	0.3	0.6	1.1	1.6	2.4
2001.....	*	*	0.1	*	0.2	0.5	0.7	0.9	1.4	2.5
2000.....	*	*	0.1	0.2	0.2	0.4	0.6	1.0	1.7	2.5
1999.....	*	*	0.2	0.2	0.3	0.4	0.6	1.2	1.8	2.9
All races 3/, male										
2004.....	*	*	*	*	0.3	*	0.6	1.1	1.9	3.5
2003.....	*	*	0.2	*	*	0.5	0.5	1.0	1.9	3.6
2002.....	*	*	*	*	*	*	0.8	1.5	1.8	3.3
2001.....	*	*	*	*	*	0.5	1.0	1.2	2.2	3.5
2000.....	*	*	*	*	*	0.5	0.7	1.4	1.7	3.8
1999.....	*	*	*	*	*	*	0.5	1.7	2.4	4.4
All races 3/, female										
2004.....	*	*	*	*	*	*	0.5	0.8	1.3	2.1
2003.....	*	*	*	*	*	*	0.7	0.8	1.3	2.0
2002.....	*	*	*	*	*	*	0.4	0.8	1.4	1.9
2001.....	*	*	*	*	*	0.5	0.4	0.6	1.0	2.1
2000.....	*	*	*	*	*	*	0.6	0.7	1.7	1.9
1999.....	*	*	*	*	*	0.4	0.7	0.9	1.4	2.3
White, both sexes										
2004.....	*	*	*	0.1	0.2	0.3	0.5	0.9	1.5	2.4
2003.....	*	*	*	*	0.2	0.4	0.5	0.8	1.4	2.3
2002.....	*	*	*	0.2	*	0.3	0.6	1.0	1.5	2.2
2001.....	*	*	*	*	*	0.4	0.7	0.9	1.4	2.5
2000.....	*	*	*	0.2	*	0.4	0.6	0.9	1.6	2.1
1999.....	*	*	*	*	0.3	0.3	0.6	1.2	1.7	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Renal tubulo-interstitial diseases (N10-N15)

All races 3/, both sexes

2004.....	0.1	0.1	0.2	0.3	0.3	0.6	1.2	2.3	4.4	9.3
2003.....	0.1	0.2	0.2	0.3	0.4	0.7	1.0	2.1	4.8	10.3
2002.....	*	0.1	0.2	0.2	0.4	0.7	1.2	2.6	4.1	10.6
2001.....	*	0.1	0.2	0.2	0.4	0.6	1.1	2.5	4.0	10.4
2000.....	*	0.1	0.2	0.3	0.4	0.6	1.1	2.5	4.1	12.0
1999.....	0.1	0.1	0.3	0.3	0.4	0.8	1.3	2.5	4.9	12.7

All races 3/, male

2004.....	*	*	*	0.4	*	0.6	1.1	2.3	4.6	12.7
2003.....	*	*	*	*	0.3	0.5	1.0	2.3	5.2	13.8
2002.....	*	*	*	*	0.4	0.7	1.5	3.1	4.3	13.8
2001.....	*	*	*	*	*	0.6	1.3	2.3	5.0	12.5
2000.....	*	*	*	*	*	0.5	0.8	2.6	4.6	14.4
1999.....	*	*	*	0.3	0.4	0.8	1.1	2.6	6.4	17.0

All races 3/, female

2004.....	*	0.2	0.2	0.2	*	0.6	1.2	2.2	4.3	7.8
2003.....	*	0.2	0.2	0.4	0.4	1.0	1.0	2.0	4.5	8.7
2002.....	*	*	*	*	0.5	0.7	1.1	2.2	3.9	9.2
2001.....	*	*	0.3	0.3	0.5	0.5	0.9	2.7	3.4	9.5
2000.....	*	*	*	0.3	0.5	0.7	1.4	2.5	3.8	11.1
1999.....	*	*	0.3	0.4	*	0.8	1.5	2.4	4.0	11.0

White, both sexes

2004.....	0.1	0.1	0.1	0.3	0.3	0.6	1.2	2.2	4.5	9.3
2003.....	0.1	0.2	0.2	0.3	0.3	0.7	1.0	2.1	4.6	10.4
2002.....	*	*	0.2	0.2	0.4	0.8	1.3	2.5	4.2	10.9
2001.....	*	*	0.2	0.2	0.3	0.5	1.1	2.6	3.8	10.6
2000.....	*	0.1	*	0.3	0.4	0.6	1.0	2.4	3.9	12.0
1999.....	*	0.2	0.3	0.4	0.3	0.7	1.2	2.5	5.0	12.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Renal failure (N17-N19)

All races 3/, both sexes

2004.....	2.1	3.6	6.2	10.3	17.3	29.1	48.2	84.0	136.8	282.9
2003.....	2.2	3.7	5.9	10.2	17.6	30.2	49.7	83.2	141.7	289.7
2002.....	2.0	3.6	5.7	9.8	16.5	29.0	49.1	83.9	139.8	284.8
2001.....	2.0	3.5	5.7	9.7	16.6	31.1	48.4	80.9	133.8	284.0
2000.....	2.0	3.3	5.3	9.2	16.6	28.9	46.2	78.1	130.0	273.7
1999.....	1.9	3.0	4.9	8.7	15.3	27.7	45.7	75.7	124.8	264.8

All races 3/, male

2004.....	2.5	4.3	7.4	11.6	19.1	32.8	55.3	100.5	178.4	367.4
2003.....	2.6	4.3	7.4	12.0	19.3	33.9	57.2	101.3	185.2	383.1
2002.....	2.2	4.2	6.7	10.6	18.4	33.1	58.8	103.9	180.0	380.7
2001.....	2.2	4.2	6.5	10.7	17.9	35.9	55.8	98.9	173.9	388.7
2000.....	2.4	4.2	6.1	10.4	18.1	32.1	53.3	97.9	170.1	374.2
1999.....	2.2	3.6	5.6	9.7	17.3	30.5	53.9	96.7	165.1	364.6

All races 3/, female

2004.....	1.8	3.0	5.0	9.2	15.6	25.8	42.5	72.1	111.2	244.9
2003.....	1.8	3.2	4.6	8.5	16.0	27.0	43.6	70.2	115.0	248.4
2002.....	1.8	2.9	4.8	9.0	14.9	25.4	41.4	69.9	115.3	243.2
2001.....	1.8	2.8	4.8	8.8	15.5	27.0	42.5	68.3	109.8	240.1
2000.....	1.6	2.4	4.5	8.0	15.2	26.1	40.6	64.4	106.4	232.8
1999.....	1.6	2.3	4.3	7.9	13.5	25.3	39.3	61.1	101.2	224.5

White, both sexes

2004.....	1.5	2.5	4.3	7.6	13.3	23.8	42.1	76.8	130.1	274.7
2003.....	1.4	2.5	4.1	7.6	13.8	24.7	42.7	76.4	133.9	281.5
2002.....	1.4	2.5	3.8	7.1	12.9	23.9	42.3	76.8	133.2	275.3
2001.....	1.4	2.3	4.0	7.1	12.6	25.2	42.1	73.8	126.0	274.6
2000.....	1.4	2.1	3.7	6.6	12.8	23.3	39.6	71.7	122.6	265.0
1999.....	1.2	1.8	3.4	6.4	11.9	22.5	39.4	69.0	116.2	251.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Renal failure (N17-N19)

White, male

2004.....	1.9	2.9	5.1	8.8	14.9	27.1	49.5	94.0	173.7	362.3
2003.....	1.8	3.0	4.9	9.0	15.3	28.4	49.7	95.6	177.3	379.3
2002.....	1.5	2.9	4.5	7.7	14.0	27.9	52.0	97.3	175.3	372.7
2001.....	1.6	2.7	4.7	8.0	14.1	30.0	50.1	93.0	166.1	383.9
2000.....	1.7	2.7	4.2	7.6	14.5	27.2	47.7	91.6	163.8	370.4
1999.....	1.4	2.2	3.8	7.6	13.6	25.5	48.0	89.1	156.2	354.9

White, female

2004.....	1.1	2.1	3.5	6.4	11.8	21.0	36.0	64.2	103.1	235.2
2003.....	1.0	2.1	3.3	6.4	12.4	21.4	36.9	62.4	107.1	238.2
2002.....	1.2	2.1	3.2	6.6	11.8	20.3	34.4	62.1	107.4	232.9
2001.....	1.1	1.9	3.3	6.2	11.2	21.1	35.6	60.2	101.8	228.8
2000.....	1.1	1.6	3.1	5.6	11.3	19.9	33.2	57.8	98.1	222.1
1999.....	1.1	1.5	2.9	5.3	10.3	19.9	32.5	54.8	92.5	210.6

Black, both sexes

2004.....	6.6	12.2	21.3	35.2	56.1	78.0	114.8	173.3	239.6	426.4
2003.....	7.8	12.4	20.5	34.2	53.6	82.9	126.1	168.8	257.4	436.4
2002.....	6.6	11.2	21.1	33.3	50.1	78.8	125.7	175.4	238.1	434.2
2001.....	6.5	12.3	19.3	32.2	52.5	87.1	119.0	170.6	245.9	425.6
2000.....	6.1	12.3	18.9	32.0	50.1	82.1	118.6	163.1	238.9	400.4
1999.....	6.7	11.7	18.9	28.5	46.1	76.5	115.7	161.0	241.6	425.3

Black, male

2004.....	6.8	15.0	27.6	39.1	63.0	91.1	127.9	196.5	282.7	506.7
2003.....	8.6	14.4	28.7	43.3	61.9	92.6	153.7	188.1	328.5	513.0
2002.....	7.4	14.4	26.1	39.4	61.2	88.8	148.2	203.5	278.2	555.3
2001.....	6.8	16.1	22.1	37.2	54.5	99.1	129.9	187.7	307.4	508.3
2000.....	7.7	17.1	22.7	37.0	53.2	84.8	127.0	196.6	287.5	480.9
1999.....	8.0	15.3	22.7	28.7	53.5	80.5	128.9	204.7	320.9	513.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Renal failure (N17-N19)

Black, female

2004.....	6.3	9.8	16.0	31.9	50.7	68.4	105.9	159.6	217.0	394.6
2003.....	7.0	10.7	13.7	26.7	47.1	75.7	107.4	157.4	220.1	406.7
2002.....	5.9	8.5	16.8	28.2	41.5	71.4	110.7	158.6	217.2	387.7
2001.....	6.4	9.0	16.9	28.1	50.9	78.3	111.7	160.3	214.6	394.9
2000.....	4.7	8.2	15.7	27.8	47.6	80.1	113.0	143.0	214.4	370.4
1999.....	5.6	8.5	15.8	28.4	40.3	73.5	106.9	134.7	202.2	392.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Urolithiasis (N20-N23)

All races 3/, both sexes

2004.....	*	*	*	*	*	0.3	0.3	0.6	1.0	1.8
2003.....	*	*	*	*	*	0.2	0.3	0.6	0.9	2.2
2002.....	*	*	*	*	*	0.2	0.3	0.5	0.9	1.8
2001.....	*	*	*	*	*	*	0.3	0.6	0.7	1.8
2000.....	*	*	*	*	*	*	0.2	0.7	0.9	1.8
1999.....	*	*	*	*	0.2	*	0.3	0.3	0.9	1.9

All races 3/, male

2004.....	*	*	*	*	*	*	*	0.7	*	1.9
2003.....	*	*	*	*	*	*	*	*	*	2.6
2002.....	*	*	*	*	*	*	*	*	*	2.0
2001.....	*	*	*	*	*	*	*	*	*	2.0
2000.....	*	*	*	*	*	*	*	*	1.1	1.7
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	0.4	*	0.5	1.0	1.8
2003.....	*	*	*	*	*	*	*	0.6	0.8	2.1
2002.....	*	*	*	*	*	*	*	0.5	0.9	1.7
2001.....	*	*	*	*	*	*	*	0.7	0.6	1.7
2000.....	*	*	*	*	*	*	*	0.8	0.8	1.8
1999.....	*	*	*	*	*	*	*	*	0.8	2.0

White, both sexes

2004.....	*	*	*	*	*	0.3	0.3	0.6	1.0	1.8
2003.....	*	*	*	*	*	*	0.3	0.6	0.9	2.3
2002.....	*	*	*	*	*	0.2	0.3	0.5	0.9	1.9
2001.....	*	*	*	*	*	*	0.3	0.6	0.8	1.9
2000.....	*	*	*	*	*	*	*	0.7	1.0	1.8
1999.....	*	*	*	*	*	*	0.3	0.3	1.0	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Urinary tract infection, site not specified (N39.0)

All races 3/, both sexes

2004.....	0.2	0.4	0.9	1.4	2.2	5.3	10.4	23.4	51.9	143.5
2003.....	0.2	0.3	0.8	1.3	2.7	4.9	11.2	23.3	52.2	147.7
2002.....	0.2	0.4	0.7	1.4	2.4	4.8	10.8	23.7	52.2	152.2
2001.....	0.2	0.4	0.6	1.4	2.3	5.3	11.4	24.5	54.9	160.4
2000.....	0.2	0.4	0.7	1.2	2.5	4.9	9.9	25.6	57.7	163.4
1999.....	0.2	0.4	0.6	1.2	2.6	5.1	11.2	25.4	57.2	172.0

All races 3/, male

2004.....	0.2	0.4	0.7	1.3	2.0	4.9	9.0	22.2	46.7	117.7
2003.....	0.2	0.3	0.6	1.0	2.6	4.5	9.8	22.4	49.3	114.6
2002.....	0.2	0.4	0.5	1.2	2.2	4.3	9.5	20.5	47.1	123.5
2001.....	*	0.3	0.5	1.2	2.1	5.0	10.1	21.6	50.1	125.8
2000.....	*	0.4	0.6	1.0	2.1	4.5	9.2	23.4	52.4	139.8
1999.....	0.2	0.4	0.4	1.1	2.6	4.4	10.9	23.3	53.5	137.5

All races 3/, female

2004.....	0.2	0.4	1.0	1.4	2.4	5.7	11.5	24.3	55.2	155.1
2003.....	0.2	0.4	0.9	1.6	2.9	5.3	12.3	24.0	53.9	162.3
2002.....	0.2	0.4	0.8	1.6	2.6	5.3	11.9	26.0	55.3	164.7
2001.....	0.2	0.5	0.7	1.5	2.5	5.6	12.4	26.6	57.8	174.9
2000.....	*	0.4	0.7	1.3	2.8	5.1	10.5	27.1	60.8	173.0
1999.....	0.3	0.5	0.8	1.3	2.5	5.6	11.4	26.9	59.3	185.9

White, both sexes

2004.....	0.2	0.4	0.8	1.2	2.1	4.9	9.9	22.4	50.9	143.2
2003.....	0.2	0.3	0.7	1.1	2.4	4.6	10.4	22.2	51.2	147.2
2002.....	0.2	0.4	0.6	1.3	2.2	4.4	10.2	22.8	50.9	150.8
2001.....	0.2	0.4	0.5	1.2	2.1	4.8	10.6	23.3	53.2	158.2
2000.....	0.1	0.4	0.5	1.1	2.2	4.5	9.1	24.4	56.0	161.9
1999.....	0.2	0.4	0.5	1.2	2.2	4.4	10.3	24.0	56.3	169.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Urinary tract infection, site not specified (N39.0)

White, male

2004.....	*	0.4	0.6	1.1	1.7	4.3	8.5	20.9	45.7	115.9
2003.....	*	0.3	0.5	0.9	2.3	3.9	9.0	21.0	48.2	116.5
2002.....	*	0.4	0.5	1.1	2.0	3.9	8.6	19.3	45.6	121.1
2001.....	*	0.3	0.4	0.9	1.9	4.4	9.2	20.4	49.3	122.1
2000.....	*	0.3	0.5	0.9	1.8	4.0	8.3	22.1	50.2	137.8
1999.....	*	0.3	0.4	1.1	2.2	3.6	9.4	21.3	52.1	132.1

White, female

2004.....	0.2	0.4	0.9	1.2	2.4	5.5	11.1	23.5	54.1	155.5
2003.....	*	0.3	0.8	1.4	2.6	5.1	11.6	23.2	53.1	160.8
2002.....	*	0.4	0.7	1.6	2.3	4.7	11.5	25.3	54.2	163.8
2001.....	0.2	0.5	0.5	1.5	2.3	5.1	11.8	25.3	55.6	173.3
2000.....	*	0.4	0.6	1.3	2.5	4.9	9.7	26.0	59.4	171.7
1999.....	0.2	0.4	0.6	1.3	2.3	5.1	10.9	25.9	58.9	185.0

Black, both sexes

2004.....	*	1.0	1.8	3.7	4.2	9.4	17.5	37.2	73.3	172.9
2003.....	*	*	1.8	3.0	6.0	9.2	21.2	40.3	73.4	178.5
2002.....	*	*	1.4	2.8	5.0	9.0	18.6	38.7	76.9	191.5
2001.....	*	1.0	1.4	2.5	4.4	9.6	20.3	41.7	83.7	208.5
2000.....	*	*	1.7	2.2	5.6	9.1	19.3	43.4	85.8	196.8
1999.....	*	0.9	1.4	1.8	6.0	11.9	21.6	45.7	74.3	213.0

Black, male

2004.....	*	*	*	3.8	4.5	10.2	17.8	44.6	66.0	162.9
2003.....	*	*	*	*	6.2	10.6	21.0	45.2	79.0	125.1
2002.....	*	*	*	3.0	4.2	8.2	20.4	40.9	77.9	188.7
2001.....	*	*	*	3.1	4.5	10.7	20.2	41.4	72.2	204.2
2000.....	*	*	*	*	6.0	9.9	20.1	45.4	91.0	175.3
1999.....	*	*	*	*	7.0	13.0	29.8	51.8	80.2	215.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Urinary tract infection, site not specified (N39.0)

Black, female

2004.....	*	*	2.0	3.6	3.9	8.8	17.4	32.9	77.1	176.8
2003.....	*	*	2.0	3.7	5.9	8.2	21.3	37.4	70.4	199.3
2002.....	*	*	*	2.6	5.6	9.6	17.5	37.3	76.4	192.6
2001.....	*	*	*	*	4.3	8.8	20.4	41.9	89.5	210.1
2000.....	*	*	*	*	5.2	8.4	18.9	42.2	83.2	204.8
1999.....	*	*	*	*	5.2	11.2	16.2	42.1	71.3	212.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of urinary system (N28-N36,N39.1-N39.9)

All races 3/, both sexes

2004.....	*	0.2	0.2	0.4	0.6	0.9	1.3	2.1	3.8	7.8
2003.....	0.1	0.1	0.3	0.4	0.7	0.7	1.2	1.9	3.5	7.1
2002.....	0.1	0.2	0.3	0.3	0.6	0.9	1.2	1.7	3.9	7.5
2001.....	0.1	0.2	0.3	0.4	0.6	0.9	1.2	2.1	3.5	7.1
2000.....	0.2	0.2	0.3	0.4	0.6	0.8	1.2	2.0	3.9	7.5
1999.....	*	0.1	0.2	0.4	0.6	0.8	1.3	1.8	3.8	7.1

All races 3/, male

2004.....	*	0.2	0.3	0.5	0.8	1.0	1.7	2.7	5.4	9.2
2003.....	*	*	0.3	0.4	0.9	0.8	1.5	2.3	4.6	9.8
2002.....	*	0.2	0.4	0.4	0.7	1.0	1.4	2.1	4.9	9.0
2001.....	*	*	0.3	0.4	0.7	1.1	1.4	2.4	4.4	8.9
2000.....	0.2	*	0.3	0.5	0.6	1.1	1.5	2.3	5.6	10.4
1999.....	*	0.2	0.2	0.5	0.6	1.2	1.8	2.2	4.7	9.9

All races 3/, female

2004.....	*	*	*	0.2	0.5	0.8	1.1	1.7	2.9	7.1
2003.....	*	*	0.2	0.3	0.6	0.6	0.9	1.6	2.9	6.0
2002.....	*	*	*	*	0.4	0.8	1.0	1.4	3.4	6.9
2001.....	*	*	*	0.4	0.5	0.7	1.0	1.9	2.9	6.3
2000.....	*	*	0.2	0.4	0.5	0.6	1.0	1.8	2.9	6.3
1999.....	*	*	*	0.3	0.5	0.4	0.9	1.5	3.2	6.0

White, both sexes

2004.....	*	0.2	0.2	0.3	0.5	0.9	1.2	2.0	3.7	7.8
2003.....	*	0.1	0.2	0.3	0.5	0.7	1.1	1.9	3.6	7.2
2002.....	*	*	0.2	0.2	0.4	0.8	1.1	1.6	3.8	7.5
2001.....	*	0.1	0.2	0.3	0.5	0.8	1.0	2.1	3.4	7.2
2000.....	0.2	0.1	0.3	0.3	0.5	0.7	1.0	1.9	3.8	7.3
1999.....	*	*	0.2	0.3	0.5	0.7	1.2	1.7	3.4	7.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of male genital organs (N40-N50)										
All races 3/, both sexes										
2004.....	*	*	*	0.1	0.2	0.4	0.6	1.2	2.9	6.3
2003.....	*	*	*	*	*	0.3	0.5	1.1	2.2	6.0
2002.....	*	*	*	*	0.2	0.2	0.6	0.8	2.6	5.8
2001.....	*	*	*	*	*	0.2	0.6	1.1	2.1	6.3
2000.....	*	*	*	*	*	0.3	0.5	1.0	2.3	5.9
1999.....	*	*	*	*	0.2	0.3	0.6	1.1	2.7	6.1
All races 3/, male										
2004.....	*	*	*	0.3	0.5	0.8	1.4	2.8	7.5	20.3
2003.....	*	*	*	*	*	0.8	1.1	2.6	5.9	19.6
2002.....	*	*	*	*	0.4	0.5	1.3	2.1	6.9	19.2
2001.....	*	*	*	*	*	0.5	1.4	2.6	5.5	21.2
2000.....	*	*	*	*	*	0.5	1.2	2.5	6.3	20.5
1999.....	*	*	*	*	0.4	0.7	1.3	2.6	7.2	21.2
All races 3/, female										
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
White, both sexes										
2004.....	*	*	*	*	0.2	0.3	0.6	1.1	2.8	6.5
2003.....	*	*	*	*	*	0.3	0.5	1.0	2.3	6.2
2002.....	*	*	*	*	*	*	0.5	0.7	2.6	5.9
2001.....	*	*	*	*	*	0.2	0.6	1.1	2.1	6.4
2000.....	*	*	*	*	*	0.2	0.5	0.9	2.4	5.9
1999.....	*	*	*	*	*	0.3	0.5	1.1	2.6	6.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hyperplasia of prostate (N40)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	0.4	0.8	2.1	5.5
2003.....	*	*	*	*	*	*	0.4	0.8	2.0	5.3
2002.....	*	*	*	*	*	*	0.4	0.6	2.2	4.8
2001.....	*	*	*	*	*	*	0.4	0.8	1.6	5.5
2000.....	*	*	*	*	*	*	0.3	0.9	1.9	5.2
1999.....	*	*	*	*	*	*	0.4	0.8	2.1	5.0

All races 3/, male

2004.....	*	*	*	*	*	*	0.8	1.9	5.6	17.8
2003.....	*	*	*	*	*	*	0.9	1.9	5.2	17.2
2002.....	*	*	*	*	*	*	0.8	1.6	5.8	15.8
2001.....	*	*	*	*	*	*	0.8	2.0	4.2	18.6
2000.....	*	*	*	*	*	*	0.7	2.1	5.2	18.1
1999.....	*	*	*	*	*	*	0.8	2.0	5.6	17.5

All races 3/, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	*	*	*	*	*	*	0.4	0.8	2.1	5.7
2003.....	*	*	*	*	*	*	0.5	0.8	2.0	5.5
2002.....	*	*	*	*	*	*	0.3	0.6	2.2	4.9
2001.....	*	*	*	*	*	*	0.4	0.8	1.6	5.7
2000.....	*	*	*	*	*	*	0.3	0.9	2.0	5.2
1999.....	*	*	*	*	*	*	0.4	0.8	2.1	5.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of male genital organs (N41-N50)

All races 3/, both sexes

2004.....	*	*	*	*	*	0.2	0.3	0.4	0.7	0.8
2003.....	*	*	*	*	*	*	*	0.3	*	0.7
2002.....	*	*	*	*	*	*	*	*	0.4	1.0
2001.....	*	*	*	*	*	*	0.3	0.3	0.5	0.8
2000.....	*	*	*	*	*	*	0.2	*	0.4	0.7
1999.....	*	*	*	*	*	*	*	0.3	0.6	1.1

All races 3/, male

2004.....	*	*	*	*	*	0.5	0.6	0.9	1.9	2.5
2003.....	*	*	*	*	*	*	*	0.7	*	2.4
2002.....	*	*	*	*	*	*	*	*	1.1	3.4
2001.....	*	*	*	*	*	*	0.6	0.7	1.3	2.6
2000.....	*	*	*	*	*	*	0.5	*	1.1	2.4
1999.....	*	*	*	*	*	*	*	0.7	1.6	3.7

All races 3/, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	*	*	*	*	*	*	*	0.3	0.7	0.8
2003.....	*	*	*	*	*	*	*	*	*	0.7
2002.....	*	*	*	*	*	*	*	*	*	1.0
2001.....	*	*	*	*	*	*	*	*	0.5	0.7
2000.....	*	*	*	*	*	*	*	*	*	0.7
1999.....	*	*	*	*	*	*	*	*	0.6	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Disorders of female genital organs (N70-N98)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	0.4	0.4	0.9	2.5
2003.....	*	*	*	*	*	0.2	0.2	0.6	1.0	2.6
2002.....	*	0.1	*	*	*	*	*	0.4	0.9	2.4
2001.....	*	*	*	*	*	*	0.4	0.5	0.9	2.1
2000.....	*	*	*	*	*	0.2	0.3	0.5	1.0	2.5
1999.....	*	*	*	*	*	*	0.2	0.6	0.8	2.0

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	*	*	*	*	*	*	0.6	0.7	1.5	3.7
2003.....	*	*	*	*	*	0.4	0.4	1.1	1.5	3.7
2002.....	*	0.2	*	*	*	*	*	0.8	1.4	3.4
2001.....	*	*	*	*	*	*	0.6	0.9	1.5	3.0
2000.....	*	*	*	*	*	0.4	0.5	0.9	1.6	3.6
1999.....	*	*	*	*	*	*	0.4	1.0	1.3	2.9

White, both sexes

2004.....	*	*	*	*	*	*	0.3	0.5	0.9	2.5
2003.....	*	*	*	*	*	*	*	0.6	1.0	2.8
2002.....	*	*	*	*	*	*	*	0.5	0.9	2.3
2001.....	*	*	*	*	*	*	0.3	0.5	0.9	2.1
2000.....	*	*	*	*	*	*	0.3	0.5	1.0	2.6
1999.....	*	*	*	*	*	*	*	0.5	0.9	2.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Noninflammatory disorders of female genital tract (N80-N98)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	0.2	0.3	0.7	1.8
2003.....	*	*	*	*	*	*	*	0.4	0.7	1.9
2002.....	*	*	*	*	*	*	*	*	0.6	1.6
2001.....	*	*	*	*	*	*	0.2	0.3	0.6	1.2
2000.....	*	*	*	*	*	*	*	0.3	0.8	1.9
1999.....	*	*	*	*	*	*	*	0.4	0.5	1.6

All races 3/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2004.....	*	*	*	*	*	*	0.4	0.5	1.2	2.7
2003.....	*	*	*	*	*	*	*	0.7	1.2	2.8
2002.....	*	*	*	*	*	*	*	*	1.0	2.3
2001.....	*	*	*	*	*	*	0.4	0.6	0.9	1.8
2000.....	*	*	*	*	*	*	*	0.5	1.2	2.7
1999.....	*	*	*	*	*	*	*	0.6	0.7	2.3

White, both sexes

2004.....	*	*	*	*	*	*	*	0.3	0.8	1.8
2003.....	*	*	*	*	*	*	*	0.4	0.8	2.1
2002.....	*	*	*	*	*	*	*	*	0.7	1.5
2001.....	*	*	*	*	*	*	*	0.3	0.6	1.2
2000.....	*	*	*	*	*	*	*	0.3	0.7	2.0
1999.....	*	*	*	*	*	*	*	0.3	0.5	1.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Ectopic pregnancy (000)

White, male

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain '...'.

White, female

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain '*'.

Black, both sexes

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain '*'.

Black, male

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain '...'.

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Medical abortion (004)

All races 3/, both sexes

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain an asterisk (*).

All races 3/, male

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain three dots (...).

All races 3/, female

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain an asterisk (*).

White, both sexes

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain an asterisk (*).

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Obstetric embolism (088)

All races 3/, both sexes

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain an asterisk (*).

All races 3/, male

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain three dots (...).

All races 3/, female

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain an asterisk (*).

White, both sexes

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain an asterisk (*).

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Obstetric embolism (088)

White, male

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain '...'.

White, female

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain '*'.

Black, both sexes

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain '*'.

Black, male

Table with 11 columns (years 2004-1999) and 11 rows (ages 40-44 to 85+). All cells contain '...'.

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes

2004.....	1.2	1.5	1.8	2.2	2.2	2.3	2.3	3.1	4.3	6.1
2003.....	1.4	1.5	1.8	2.4	2.4	2.3	2.2	3.0	4.4	6.8
2002.....	1.3	1.6	1.9	2.3	2.4	2.3	2.5	3.1	4.8	6.6
2001.....	1.4	1.5	1.8	2.1	2.5	2.2	2.7	3.3	4.3	7.2
2000.....	1.3	1.4	1.8	2.2	2.5	2.6	2.4	3.3	5.0	7.5
1999.....	1.2	1.7	1.8	2.3	2.4	2.6	2.5	3.4	5.0	7.8

All races 3/, male

2004.....	1.3	1.7	2.0	2.3	2.2	2.3	2.7	3.7	3.8	6.8
2003.....	1.5	1.6	1.8	2.5	2.6	2.4	2.4	2.8	4.5	6.7
2002.....	1.4	1.9	2.0	2.3	2.7	2.4	2.7	3.4	4.8	7.0
2001.....	1.6	1.5	1.9	2.1	2.5	2.5	3.1	3.6	4.8	7.7
2000.....	1.5	1.6	2.2	2.0	2.6	2.7	2.6	3.3	5.3	10.6
1999.....	1.3	1.9	2.0	2.5	2.5	2.4	2.6	3.4	5.7	8.5

All races 3/, female

2004.....	1.2	1.3	1.7	2.0	2.1	2.3	2.0	2.7	4.6	5.8
2003.....	1.3	1.3	1.8	2.3	2.3	2.2	2.1	3.2	4.4	6.8
2002.....	1.2	1.4	1.8	2.3	2.2	2.3	2.3	2.9	4.9	6.5
2001.....	1.3	1.4	1.8	2.1	2.6	2.0	2.4	3.1	4.0	7.0
2000.....	1.1	1.3	1.5	2.4	2.5	2.5	2.3	3.3	4.8	6.3
1999.....	1.2	1.6	1.7	2.2	2.3	2.8	2.3	3.4	4.5	7.6

White, both sexes

2004.....	1.3	1.6	1.9	2.3	2.3	2.4	2.5	3.1	4.4	6.3
2003.....	1.5	1.5	1.9	2.5	2.5	2.3	2.3	3.1	4.5	6.9
2002.....	1.3	1.7	2.0	2.4	2.5	2.5	2.5	3.2	4.8	6.9
2001.....	1.5	1.4	1.9	2.2	2.7	2.3	2.7	3.4	4.4	7.4
2000.....	1.3	1.5	2.0	2.2	2.6	2.6	2.4	3.4	5.2	7.8
1999.....	1.3	1.8	1.8	2.4	2.4	2.7	2.4	3.3	5.1	8.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Congenital malformations of heart (Q20-Q24)

All races 3/, both sexes

2004.....	0.5	0.4	0.5	0.5	0.4	0.5	0.6	0.9	1.3	1.5
2003.....	0.6	0.5	0.5	0.5	0.5	0.7	0.6	0.8	1.5	2.1
2002.....	0.5	0.6	0.5	0.6	0.5	0.6	0.7	1.0	1.6	2.3
2001.....	0.6	0.5	0.6	0.5	0.5	0.6	0.7	1.1	1.4	2.6
2000.....	0.6	0.5	0.5	0.6	0.6	0.7	0.8	1.1	2.0	2.5
1999.....	0.6	0.6	0.6	0.6	0.7	0.9	0.9	1.2	1.7	3.3

All races 3/, male

2004.....	0.5	0.4	0.6	0.6	0.5	*	0.8	0.9	1.0	1.5
2003.....	0.6	0.5	0.4	0.6	0.5	0.8	*	*	1.3	*
2002.....	0.5	0.6	0.5	0.6	0.5	0.5	0.7	1.0	1.3	1.4
2001.....	0.6	0.6	0.6	0.5	0.5	0.7	1.0	1.0	1.6	1.7
2000.....	0.6	0.5	0.6	0.6	0.6	0.8	0.8	1.1	2.0	2.7
1999.....	0.6	0.7	0.7	0.6	0.7	0.7	0.9	1.0	1.8	3.1

All races 3/, female

2004.....	0.4	0.3	0.5	0.4	0.4	0.6	0.4	0.9	1.5	1.6
2003.....	0.6	0.5	0.5	0.5	0.5	0.7	0.7	1.0	1.6	2.4
2002.....	0.5	0.6	0.6	0.6	0.4	0.6	0.8	1.0	1.8	2.6
2001.....	0.6	0.5	0.6	0.4	0.5	0.6	0.5	1.2	1.3	3.0
2000.....	0.5	0.5	0.5	0.7	0.5	0.5	0.8	1.0	1.9	2.4
1999.....	0.6	0.6	0.5	0.7	0.6	1.0	0.8	1.3	1.7	3.4

White, both sexes

2004.....	0.5	0.4	0.6	0.5	0.5	0.5	0.6	0.9	1.4	1.6
2003.....	0.7	0.5	0.5	0.5	0.5	0.7	0.7	0.8	1.6	2.2
2002.....	0.5	0.6	0.5	0.6	0.4	0.6	0.8	1.0	1.6	2.4
2001.....	0.6	0.6	0.6	0.4	0.5	0.6	0.7	1.2	1.5	2.6
2000.....	0.6	0.5	0.6	0.6	0.6	0.7	0.7	1.1	2.1	2.7
1999.....	0.6	0.7	0.6	0.7	0.6	0.9	0.9	1.3	1.8	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other congenital malformations of circulatory system (Q25-Q28)

All races 3/, both sexes

2004.....	0.2	0.2	0.2	0.3	0.2	0.3	0.4	0.9	1.0	1.6
2003.....	0.2	0.2	0.3	0.2	0.4	0.4	0.5	0.7	0.9	1.5
2002.....	0.1	0.2	0.2	0.3	0.3	0.4	0.6	0.8	1.3	1.5
2001.....	0.2	0.2	0.2	0.3	0.3	0.4	0.8	0.6	0.9	1.7
2000.....	0.2	0.2	0.2	0.3	0.4	0.3	0.6	0.7	1.2	2.1
1999.....	0.1	0.3	0.3	0.3	0.3	0.6	0.5	0.7	1.2	1.7

All races 3/, male

2004.....	0.2	0.3	0.3	0.2	*	*	*	1.1	1.0	1.9
2003.....	0.2	*	0.3	*	0.5	*	0.6	0.8	1.3	1.7
2002.....	*	0.3	0.2	0.3	0.4	0.5	0.8	0.7	1.3	1.8
2001.....	0.2	0.3	*	0.4	*	*	0.8	0.9	1.2	2.1
2000.....	0.2	0.3	0.4	*	0.5	*	0.8	0.8	1.4	3.0
1999.....	*	0.3	0.4	0.4	*	0.6	*	0.9	1.4	2.2

All races 3/, female

2004.....	*	0.2	*	0.3	*	*	*	0.7	1.0	1.5
2003.....	0.2	*	0.2	*	*	*	0.5	0.5	0.7	1.4
2002.....	*	0.2	0.2	0.3	*	0.4	0.4	0.9	1.3	1.4
2001.....	*	*	*	*	*	0.4	0.7	*	0.7	1.6
2000.....	*	0.2	*	0.3	0.4	*	0.4	0.7	1.1	1.7
1999.....	*	0.3	0.2	*	0.4	0.6	0.5	0.5	1.1	1.5

White, both sexes

2004.....	0.2	0.3	0.2	0.2	0.2	0.3	0.5	0.8	1.1	1.7
2003.....	0.2	0.2	0.2	0.2	0.4	0.4	0.5	0.7	0.9	1.5
2002.....	*	0.2	0.2	0.2	0.3	0.5	0.6	0.8	1.3	1.6
2001.....	0.1	0.2	0.2	0.3	0.3	0.4	0.7	0.7	1.0	1.7
2000.....	0.2	0.2	0.2	0.2	0.4	0.3	0.6	0.8	1.2	2.2
1999.....	0.1	0.3	0.3	0.3	0.3	0.6	0.4	0.6	1.2	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other congenital malformations and deformations (Q10-Q18,Q35-Q89)

All races 3/, both sexes

2004.....	0.2	0.3	0.3	0.4	0.6	0.8	0.9	1.2	1.7	2.6
2003.....	0.2	0.3	0.3	0.5	0.5	0.6	0.8	1.3	1.7	2.8
2002.....	0.3	0.4	0.4	0.5	0.6	0.6	1.0	1.1	1.7	2.7
2001.....	0.3	0.3	0.3	0.4	0.6	0.7	0.9	1.3	1.8	2.5
2000.....	0.3	0.3	0.3	0.5	0.6	0.8	0.8	1.3	1.6	2.8
1999.....	0.2	0.3	0.3	0.4	0.6	0.7	0.9	1.3	1.8	2.6

All races 3/, male

2004.....	0.3	0.4	0.4	0.4	0.6	0.9	0.9	1.2	1.6	3.2
2003.....	0.2	0.4	0.4	0.5	0.4	0.7	0.9	1.3	1.6	3.3
2002.....	0.4	0.5	0.5	0.4	0.7	0.7	1.2	1.4	1.8	3.6
2001.....	0.4	0.3	0.3	0.4	0.6	0.8	1.0	1.4	1.8	3.7
2000.....	0.3	0.4	0.4	0.4	0.6	0.9	0.8	1.1	1.8	4.6
1999.....	0.3	0.4	0.4	0.5	0.5	0.7	1.1	1.4	2.1	3.0

All races 3/, female

2004.....	0.2	0.2	0.3	0.3	0.6	0.7	0.9	1.1	1.8	2.4
2003.....	0.2	0.3	0.2	0.5	0.5	0.6	0.7	1.4	1.8	2.6
2002.....	0.2	0.2	0.3	0.5	0.5	0.5	0.8	0.9	1.7	2.3
2001.....	0.2	0.3	0.2	0.4	0.6	0.6	0.8	1.2	1.8	2.0
2000.....	0.2	0.2	*	0.6	0.6	0.8	0.9	1.3	1.5	2.1
1999.....	0.2	0.3	0.2	0.4	0.6	0.8	0.7	1.2	1.7	2.5

White, both sexes

2004.....	0.3	0.3	0.3	0.3	0.6	0.8	1.0	1.1	1.7	2.7
2003.....	0.2	0.3	0.3	0.5	0.5	0.6	0.8	1.4	1.7	2.8
2002.....	0.3	0.4	0.4	0.5	0.6	0.6	1.0	1.1	1.7	2.7
2001.....	0.3	0.3	0.3	0.4	0.6	0.7	0.9	1.3	1.7	2.6
2000.....	0.3	0.3	0.3	0.5	0.6	0.8	0.9	1.3	1.7	2.8
1999.....	0.2	0.3	0.3	0.5	0.6	0.8	0.9	1.2	1.8	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Down's syndrome (Q90)

All races 3/, both sexes

2004.....	0.2	0.4	0.7	1.0	0.8	0.6	0.3	*	*	*
2003.....	0.3	0.4	0.6	1.1	0.9	0.4	*	*	*	*
2002.....	0.2	0.3	0.7	0.9	1.0	0.6	*	*	*	*
2001.....	0.3	0.3	0.6	0.9	1.0	0.4	*	*	*	*
2000.....	0.2	0.3	0.7	0.8	0.8	0.6	*	*	*	*
1999.....	0.2	0.4	0.5	0.8	0.7	0.3	*	*	*	*

All races 3/, male

2004.....	0.2	0.5	0.6	1.0	0.9	0.6	*	*	*	*
2003.....	0.3	0.4	0.6	1.1	1.1	0.4	*	*	*	*
2002.....	0.2	0.4	0.7	0.9	1.1	0.6	*	*	*	*
2001.....	0.3	0.3	0.6	0.8	1.0	0.5	*	*	*	*
2000.....	*	0.2	0.8	0.8	0.7	0.5	*	*	*	*
1999.....	*	0.4	0.4	0.8	0.8	*	*	*	*	*

All races 3/, female

2004.....	0.2	0.4	0.7	0.9	0.8	0.5	*	*	*	*
2003.....	0.2	0.4	0.7	1.1	0.8	0.4	*	*	*	*
2002.....	0.2	0.3	0.6	0.9	0.9	0.6	*	*	*	*
2001.....	0.3	0.3	0.7	1.0	1.0	*	*	*	*	*
2000.....	*	0.3	0.6	0.7	0.9	0.7	*	*	*	*
1999.....	*	0.3	0.6	0.8	0.6	0.4	*	*	*	*

White, both sexes

2004.....	0.2	0.5	0.7	1.1	0.9	0.6	0.3	*	*	*
2003.....	0.3	0.5	0.7	1.2	1.0	0.5	*	*	*	*
2002.....	0.2	0.3	0.7	1.0	1.1	0.6	*	*	*	*
2001.....	0.3	0.3	0.7	1.0	1.1	0.5	*	*	*	*
2000.....	0.2	0.3	0.8	0.8	0.9	0.7	*	*	*	*
1999.....	0.2	0.4	0.5	0.9	0.8	0.3	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2004.....	4.9	5.7	5.8	6.3	7.9	10.2	15.4	28.0	55.5	221.5
2003.....	5.9	7.3	6.8	6.8	8.2	10.6	17.3	26.7	55.7	224.1
2002.....	6.0	6.4	6.9	7.0	8.3	10.4	16.1	28.3	53.3	208.0
2001.....	7.7	9.1	8.8	8.4	9.9	12.7	17.8	29.8	54.3	206.4
2000.....	7.6	8.6	8.0	8.6	10.2	14.2	19.0	30.1	56.5	208.1
1999.....	5.1	5.9	5.8	6.9	8.9	11.9	17.6	26.5	50.5	187.7

All races 3/, male

2004.....	6.5	7.4	7.7	8.3	10.1	12.6	17.9	30.6	58.1	183.6
2003.....	7.4	9.4	8.9	8.9	10.6	13.7	21.3	31.2	58.0	177.9
2002.....	7.6	8.5	9.0	9.0	11.2	12.7	19.7	32.9	60.6	175.5
2001.....	9.9	11.9	11.8	11.0	13.2	15.5	20.9	33.2	60.6	175.0
2000.....	10.2	11.7	11.2	11.4	13.4	18.9	22.4	36.1	62.3	182.7
1999.....	6.7	8.2	7.9	9.1	11.9	15.1	21.8	31.5	54.4	166.3

All races 3/, female

2004.....	3.3	4.1	4.0	4.5	5.9	8.1	13.4	26.1	53.8	238.5
2003.....	4.4	5.1	4.8	4.9	6.0	7.9	14.2	23.5	54.3	244.6
2002.....	4.5	4.4	5.0	5.0	5.6	8.4	13.3	25.1	48.9	222.1
2001.....	5.5	6.3	5.9	5.9	6.9	10.2	15.4	27.4	50.4	219.5
2000.....	5.0	5.6	5.0	5.9	7.3	10.1	16.4	26.0	53.0	218.5
1999.....	3.4	3.7	3.9	4.9	6.2	9.2	14.2	23.0	48.3	196.4

White, both sexes

2004.....	4.7	5.6	5.4	5.8	7.4	9.8	14.7	27.6	55.1	224.4
2003.....	5.6	6.6	6.2	6.3	7.6	10.1	16.6	26.5	56.1	228.3
2002.....	5.7	5.8	6.1	6.4	7.9	9.8	15.9	27.9	53.3	211.6
2001.....	7.6	8.8	8.3	7.9	9.4	11.8	17.2	29.5	53.9	210.0
2000.....	7.4	8.2	7.3	7.9	9.5	13.0	18.5	29.5	56.0	212.2
1999.....	4.4	5.2	5.0	6.2	8.2	11.1	17.0	25.5	49.5	190.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, male

2004.....	6.3	7.2	6.9	7.5	9.3	11.9	17.4	30.1	57.8	186.0
2003.....	7.0	8.6	8.2	8.1	9.7	12.8	20.4	31.0	57.8	180.0
2002.....	7.2	7.6	7.7	8.1	10.7	11.9	19.2	31.6	60.0	178.2
2001.....	10.0	11.6	11.2	10.1	12.2	14.3	20.3	33.2	60.6	177.0
2000.....	10.1	11.2	10.1	10.2	12.4	17.1	21.5	35.4	62.0	186.8
1999.....	5.9	7.0	6.7	8.0	10.9	13.9	21.0	30.3	52.6	167.3

White, female

2004.....	3.1	4.0	3.8	4.2	5.6	7.9	12.5	25.8	53.5	241.7
2003.....	4.2	4.6	4.2	4.7	5.6	7.7	13.5	23.3	55.0	249.7
2002.....	4.2	4.0	4.4	4.7	5.3	7.9	13.1	25.3	49.2	226.1
2001.....	5.3	6.0	5.5	5.7	6.7	9.7	14.6	26.8	49.9	223.8
2000.....	4.6	5.2	4.6	5.6	6.7	9.4	16.0	25.4	52.4	222.5
1999.....	2.9	3.4	3.4	4.5	5.7	8.6	13.8	22.1	47.6	199.4

Black, both sexes

2004.....	7.6	8.6	11.0	12.5	14.6	16.1	25.9	39.9	72.0	220.8
2003.....	9.7	13.4	13.4	13.0	15.7	17.9	30.3	34.9	60.9	210.9
2002.....	9.5	12.1	15.7	13.8	13.6	17.8	23.1	39.5	64.9	196.0
2001.....	9.8	13.2	14.6	14.3	15.4	21.7	27.9	38.6	66.6	191.6
2000.....	11.2	13.8	15.9	16.0	19.0	28.3	29.7	44.2	73.4	186.4
1999.....	10.3	13.1	13.3	14.9	17.0	22.3	27.2	42.7	72.5	182.4

Black, male

2004.....	9.7	11.8	16.3	18.2	20.2	22.3	28.2	45.0	81.1	181.8
2003.....	12.5	17.5	17.8	18.5	22.8	25.9	38.8	41.5	70.1	193.9
2002.....	12.4	17.1	22.1	19.5	20.0	22.9	31.0	57.0	87.1	181.2
2001.....	11.7	17.3	18.9	21.7	25.4	29.5	31.6	40.9	70.6	184.9
2000.....	13.5	18.2	23.6	25.0	25.5	41.4	38.1	54.8	83.5	161.5
1999.....	13.6	19.7	19.1	22.4	24.0	30.1	35.6	54.1	88.1	184.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, female

2004.....	5.9	5.9	6.6	7.7	10.2	11.5	24.4	36.9	67.3	236.3
2003.....	7.3	9.9	9.7	8.4	10.2	12.1	24.6	30.9	56.1	217.5
2002.....	7.0	7.7	10.2	9.0	8.7	14.1	17.7	29.1	53.3	201.7
2001.....	8.1	9.7	11.0	8.2	7.5	16.0	25.3	37.2	64.6	194.1
2000.....	9.1	9.9	9.4	8.6	13.9	18.6	24.2	37.9	68.4	195.7
1999.....	7.3	7.3	8.5	8.9	11.5	16.5	21.6	35.8	64.8	181.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Senility (R54)										
All races 3/, both sexes										
2004.....	*	*	*	*	0.2	0.4	1.2	4.1	11.2	76.3
2003.....	*	*	*	*	0.2	0.4	1.4	3.5	11.2	76.7
2002.....	*	*	*	*	*	0.4	1.2	3.7	9.8	72.0
2001.....	*	*	*	*	*	0.4	1.3	3.3	10.8	73.0
2000.....	*	*	*	*	*	0.4	1.3	3.6	10.3	67.2
1999.....	*	*	*	*	*	0.3	1.1	2.9	9.0	56.5
All races 3/, male										
2004.....	*	*	*	*	*	0.6	1.3	4.3	11.6	57.2
2003.....	*	*	*	*	*	0.5	1.6	3.7	10.9	55.6
2002.....	*	*	*	*	*	0.5	1.2	3.6	10.3	56.5
2001.....	*	*	*	*	*	0.5	1.5	3.2	11.4	57.2
2000.....	*	*	*	*	*	*	1.3	3.8	10.5	51.7
1999.....	*	*	*	*	*	*	1.5	3.1	8.9	45.8
All races 3/, female										
2004.....	*	*	*	*	*	*	1.1	4.0	10.9	84.9
2003.....	*	*	*	*	*	*	1.3	3.4	11.4	86.0
2002.....	*	*	*	*	*	*	1.2	3.8	9.5	78.8
2001.....	*	*	*	*	*	*	1.2	3.3	10.4	79.6
2000.....	*	*	*	*	*	*	1.2	3.4	10.2	73.6
1999.....	*	*	*	*	*	*	0.8	2.8	9.0	60.8
White, both sexes										
2004.....	*	*	*	*	*	0.4	1.2	4.2	11.2	79.0
2003.....	*	*	*	*	*	0.4	1.5	3.7	11.8	79.3
2002.....	*	*	*	*	*	0.4	1.2	3.8	10.1	74.2
2001.....	*	*	*	*	*	0.3	1.4	3.3	11.0	75.1
2000.....	*	*	*	*	*	0.3	1.2	3.5	10.2	69.0
1999.....	*	*	*	*	*	0.4	1.1	2.8	8.8	58.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Senility (R54)

Black, female

2004.....	*	*	*	*	*	*	*	*	10.2	65.4
2003.....	*	*	*	*	*	*	*	*	*	62.8
2002.....	*	*	*	*	*	*	*	*	8.8	57.9
2001.....	*	*	*	*	*	*	*	*	10.6	56.4
2000.....	*	*	*	*	*	*	*	*	14.4	63.5
1999.....	*	*	*	*	*	*	*	*	11.7	41.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Ill-defined and unknown causes of mortality (R95-R99)

All races 3/, both sexes

2004.....	4.1	4.6	4.5	4.7	5.1	6.1	7.6	11.5	18.1	52.4
2003.....	4.9	5.9	5.5	5.3	5.6	6.7	9.1	11.8	19.7	59.5
2002.....	5.0	5.3	5.5	5.5	5.8	6.6	8.7	13.4	20.8	58.3
2001.....	6.6	8.0	7.6	6.8	7.6	8.8	10.4	14.9	22.9	62.1
2000.....	6.5	7.4	6.8	6.9	7.6	10.0	11.9	16.2	26.9	73.0
1999.....	4.2	4.9	4.6	5.2	6.3	7.9	10.5	13.5	22.9	66.7

All races 3/, male

2004.....	5.5	6.2	6.0	6.3	7.0	7.7	9.8	14.2	20.3	50.7
2003.....	6.3	7.8	7.3	7.1	7.6	9.4	12.2	15.0	23.4	53.6
2002.....	6.5	7.2	7.4	7.3	8.1	8.4	11.4	16.4	25.5	55.0
2001.....	8.6	10.5	10.2	9.1	10.2	11.2	13.2	18.1	27.6	58.1
2000.....	8.8	10.3	9.6	9.3	10.4	13.8	14.3	20.2	31.2	72.1
1999.....	5.7	6.9	6.3	6.9	8.8	10.5	13.8	17.0	26.4	64.4

All races 3/, female

2004.....	2.6	3.1	3.0	3.1	3.5	4.6	5.8	9.6	16.8	53.2
2003.....	3.5	4.1	3.8	3.6	3.7	4.3	6.5	9.4	17.4	62.1
2002.....	3.6	3.4	3.6	3.8	3.8	5.0	6.6	11.3	17.9	59.8
2001.....	4.7	5.5	5.0	4.8	5.2	6.8	8.1	12.6	20.0	63.8
2000.....	4.3	4.7	4.1	4.7	5.1	6.8	10.1	13.4	24.3	73.3
1999.....	2.8	3.1	3.0	3.6	4.0	5.7	7.9	11.2	20.9	67.7

White, both sexes

2004.....	4.0	4.5	4.2	4.4	5.0	5.9	7.2	11.6	18.1	53.1
2003.....	4.7	5.6	5.2	5.0	5.2	6.6	8.8	11.9	19.9	60.9
2002.....	4.9	4.8	4.8	5.1	5.6	6.3	8.7	13.3	20.8	59.3
2001.....	6.8	7.8	7.3	6.6	7.4	8.6	10.1	14.8	22.6	63.2
2000.....	6.4	7.2	6.3	6.5	7.3	9.3	11.9	16.0	27.1	74.5
1999.....	3.7	4.4	4.1	4.7	5.9	7.4	10.3	13.2	22.4	67.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Ill-defined and unknown causes of mortality (R95-R99)

White, male

2004.....	5.5	6.0	5.5	5.9	6.6	7.5	9.6	14.3	20.2	51.4
2003.....	6.1	7.4	6.9	6.5	7.0	9.0	11.7	15.0	23.6	54.7
2002.....	6.3	6.5	6.4	6.7	7.8	7.9	11.3	16.0	24.9	56.2
2001.....	8.9	10.3	9.9	8.5	9.9	10.8	13.0	18.1	27.2	59.3
2000.....	8.8	10.0	8.8	8.7	9.9	12.5	13.9	19.9	31.4	73.2
1999.....	5.1	6.0	5.4	6.1	8.2	9.7	13.4	16.5	25.3	63.1

White, female

2004.....	2.4	3.1	3.0	3.0	3.5	4.5	5.2	9.6	16.8	53.8
2003.....	3.4	3.8	3.5	3.6	3.6	4.5	6.4	9.6	17.6	63.7
2002.....	3.5	3.2	3.3	3.6	3.7	4.8	6.5	11.4	18.3	60.6
2001.....	4.6	5.3	4.7	4.7	5.1	6.7	7.9	12.4	19.8	64.9
2000.....	4.0	4.4	3.8	4.4	4.9	6.5	10.2	13.2	24.5	75.0
1999.....	2.4	2.8	2.7	3.3	3.7	5.4	7.8	10.9	20.7	68.5

Black, both sexes

2004.....	5.6	6.7	7.8	8.2	7.9	8.9	12.7	13.9	21.3	55.0
2003.....	7.8	10.0	9.6	9.2	10.1	9.2	15.0	13.4	21.3	54.6
2002.....	7.4	9.5	11.9	9.9	9.1	10.9	11.0	17.9	24.3	56.1
2001.....	7.4	10.9	11.2	10.4	10.2	11.9	14.0	18.6	28.2	55.6
2000.....	9.2	11.0	12.6	11.6	12.6	18.6	15.2	21.5	30.5	66.0
1999.....	8.2	10.2	10.1	11.0	11.5	13.7	13.8	19.3	32.4	71.4

Black, male

2004.....	6.8	9.4	11.9	12.0	12.1	11.4	14.6	16.4	26.5	52.7
2003.....	10.1	13.1	12.6	14.0	16.2	15.8	23.0	19.6	25.8	59.4
2002.....	9.9	13.8	17.3	14.6	14.1	15.5	16.0	25.7	40.5	53.6
2001.....	8.4	14.0	14.4	15.8	16.1	17.0	18.5	22.8	35.3	49.9
2000.....	11.2	14.8	18.7	17.0	18.2	28.6	21.4	29.3	38.8	70.3
1999.....	10.8	15.4	15.0	16.4	17.4	20.0	20.4	25.6	41.9	89.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Ill-defined and unknown causes of mortality (R95-R99)

Black, female

2004.....	4.5	4.3	4.3	5.1	4.7	7.0	11.4	12.4	18.5	55.9
2003.....	5.8	7.3	7.0	5.3	5.4	4.4	9.5	9.8	19.0	52.7
2002.....	5.1	5.7	7.2	6.1	5.3	7.6	7.6	13.2	15.9	57.1
2001.....	6.6	8.3	8.5	5.9	5.6	8.1	11.0	16.1	24.5	57.7
2000.....	7.4	7.8	7.4	7.2	8.2	11.1	11.1	16.8	26.3	64.4
1999.....	6.0	5.7	6.0	6.5	6.8	9.0	9.5	15.4	27.7	64.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other ill-defined and unknown causes of mortality (R96-R99)

All races 3/, both sexes

2004.....	4.1	4.6	4.5	4.7	5.1	6.1	7.6	11.5	18.1	52.4
2003.....	4.9	5.9	5.5	5.3	5.6	6.7	9.1	11.8	19.7	59.5
2002.....	5.0	5.3	5.5	5.5	5.8	6.6	8.7	13.4	20.8	58.3
2001.....	6.6	8.0	7.6	6.8	7.6	8.8	10.4	14.9	22.9	62.1
2000.....	6.5	7.4	6.8	6.9	7.6	10.0	11.9	16.2	26.9	73.0
1999.....	4.2	4.9	4.6	5.2	6.3	7.9	10.5	13.5	22.9	66.7

All races 3/, male

2004.....	5.5	6.2	6.0	6.3	7.0	7.7	9.8	14.2	20.3	50.7
2003.....	6.3	7.8	7.3	7.1	7.6	9.4	12.2	15.0	23.4	53.6
2002.....	6.5	7.2	7.4	7.3	8.1	8.4	11.4	16.4	25.5	55.0
2001.....	8.6	10.5	10.2	9.1	10.2	11.2	13.2	18.1	27.6	58.1
2000.....	8.8	10.3	9.6	9.3	10.4	13.8	14.3	20.2	31.2	72.1
1999.....	5.7	6.9	6.3	6.9	8.8	10.5	13.8	17.0	26.4	64.4

All races 3/, female

2004.....	2.6	3.1	3.0	3.1	3.5	4.6	5.8	9.6	16.8	53.2
2003.....	3.5	4.1	3.8	3.6	3.7	4.3	6.5	9.4	17.4	62.1
2002.....	3.6	3.4	3.6	3.8	3.8	5.0	6.6	11.3	17.9	59.8
2001.....	4.7	5.5	5.0	4.8	5.2	6.8	8.1	12.6	20.0	63.8
2000.....	4.3	4.7	4.1	4.7	5.1	6.8	10.1	13.4	24.3	73.3
1999.....	2.8	3.1	3.0	3.6	4.0	5.7	7.9	11.2	20.9	67.7

White, both sexes

2004.....	4.0	4.5	4.2	4.4	5.0	5.9	7.2	11.6	18.1	53.1
2003.....	4.7	5.6	5.2	5.0	5.2	6.6	8.8	11.9	19.9	60.9
2002.....	4.9	4.8	4.8	5.1	5.6	6.3	8.7	13.3	20.8	59.3
2001.....	6.8	7.8	7.3	6.6	7.4	8.6	10.1	14.8	22.6	63.2
2000.....	6.4	7.2	6.3	6.5	7.3	9.3	11.9	16.0	27.1	74.5
1999.....	3.7	4.4	4.1	4.7	5.9	7.4	10.3	13.2	22.4	67.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other ill-defined and unknown causes of mortality (R96-R99)

White, male

2004	5.5	6.0	5.5	5.9	6.6	7.5	9.6	14.3	20.2	51.4
2003	6.1	7.4	6.9	6.5	7.0	9.0	11.7	15.0	23.6	54.7
2002	6.3	6.5	6.4	6.7	7.8	7.9	11.3	16.0	24.9	56.2
2001	8.9	10.3	9.9	8.5	9.9	10.8	13.0	18.1	27.2	59.3
2000	8.8	10.0	8.8	8.7	9.9	12.5	13.9	19.9	31.4	73.2
1999	5.1	6.0	5.4	6.1	8.2	9.7	13.4	16.5	25.3	63.1

White, female

2004	2.4	3.1	3.0	3.0	3.5	4.5	5.2	9.6	16.8	53.8
2003	3.4	3.8	3.5	3.6	3.6	4.5	6.4	9.6	17.6	63.7
2002	3.5	3.2	3.3	3.6	3.7	4.8	6.5	11.4	18.3	60.6
2001	4.6	5.3	4.7	4.7	5.1	6.7	7.9	12.4	19.8	64.9
2000	4.0	4.4	3.8	4.4	4.9	6.5	10.2	13.2	24.5	75.0
1999	2.4	2.8	2.7	3.3	3.7	5.4	7.8	10.9	20.7	68.5

Black, both sexes

2004	5.6	6.7	7.8	8.2	7.9	8.9	12.7	13.9	21.3	55.0
2003	7.8	10.0	9.6	9.2	10.1	9.2	15.0	13.4	21.3	54.6
2002	7.4	9.5	11.9	9.9	9.1	10.9	11.0	17.9	24.3	56.1
2001	7.4	10.9	11.2	10.4	10.2	11.9	14.0	18.6	28.2	55.6
2000	9.2	11.0	12.6	11.6	12.6	18.6	15.2	21.5	30.5	66.0
1999	8.2	10.2	10.1	11.0	11.5	13.7	13.8	19.3	32.4	71.4

Black, male

2004	6.8	9.4	11.9	12.0	12.1	11.4	14.6	16.4	26.5	52.7
2003	10.1	13.1	12.6	14.0	16.2	15.8	23.0	19.6	25.8	59.4
2002	9.9	13.8	17.3	14.6	14.1	15.5	16.0	25.7	40.5	53.6
2001	8.4	14.0	14.4	15.8	16.1	17.0	18.5	22.8	35.3	49.9
2000	11.2	14.8	18.7	17.0	18.2	28.6	21.4	29.3	38.8	70.3
1999	10.8	15.4	15.0	16.4	17.4	20.0	20.4	25.6	41.9	89.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other ill-defined and unknown causes of mortality (R96-R99)

Black, female

2004.....	4.5	4.3	4.3	5.1	4.7	7.0	11.4	12.4	18.5	55.9
2003.....	5.8	7.3	7.0	5.3	5.4	4.4	9.5	9.8	19.0	52.7
2002.....	5.1	5.7	7.2	6.1	5.3	7.6	7.6	13.2	15.9	57.1
2001.....	6.6	8.3	8.5	5.9	5.6	8.1	11.0	16.1	24.5	57.7
2000.....	7.4	7.8	7.4	7.2	8.2	11.1	11.1	16.8	26.3	64.4
1999.....	6.0	5.7	6.0	6.5	6.8	9.0	9.5	15.4	27.7	64.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

All races 3/, both sexes

2004.....	0.9	1.1	1.3	1.7	2.6	3.7	6.6	12.4	26.2	92.7
2003.....	0.9	1.3	1.3	1.5	2.4	3.5	6.9	11.4	24.8	87.9
2002.....	1.0	1.1	1.5	1.4	2.3	3.4	6.2	11.2	22.8	77.7
2001.....	1.0	1.1	1.2	1.5	2.2	3.4	6.1	11.6	20.6	71.3
2000.....	1.1	1.1	1.3	1.6	2.5	3.8	5.9	10.4	19.3	67.9
1999.....	0.8	1.0	1.2	1.7	2.5	3.7	6.0	10.0	18.7	64.5

All races 3/, male

2004.....	1.0	1.2	1.7	1.9	2.9	4.4	6.8	12.2	26.2	75.7
2003.....	1.0	1.6	1.6	1.8	2.9	3.7	7.5	12.5	23.7	68.6
2002.....	1.1	1.3	1.6	1.7	3.0	3.8	7.1	12.9	24.8	64.0
2001.....	1.3	1.4	1.6	1.9	2.9	3.8	6.2	11.8	21.6	59.7
2000.....	1.4	1.4	1.7	2.1	2.9	4.7	6.8	12.1	20.7	58.9
1999.....	1.1	1.3	1.5	2.2	2.9	4.2	6.4	11.5	19.2	56.1

All races 3/, female

2004.....	0.7	1.0	1.0	1.4	2.3	3.1	6.5	12.6	26.1	100.4
2003.....	0.8	1.0	1.0	1.2	2.0	3.3	6.4	10.6	25.4	96.4
2002.....	0.9	1.0	1.3	1.2	1.7	3.1	5.5	10.1	21.5	83.6
2001.....	0.7	0.8	0.9	1.1	1.6	3.1	6.0	11.4	20.1	76.2
2000.....	0.7	0.9	0.9	1.2	2.1	3.0	5.1	9.2	18.5	71.6
1999.....	0.6	0.7	0.8	1.3	2.1	3.3	5.6	9.0	18.3	67.9

White, both sexes

2004.....	0.7	1.0	1.1	1.4	2.3	3.5	6.3	11.8	25.8	92.3
2003.....	0.8	1.0	1.0	1.3	2.2	3.1	6.4	11.0	24.4	88.1
2002.....	0.8	1.0	1.2	1.2	2.1	3.1	6.0	10.8	22.4	78.2
2001.....	0.9	1.0	1.0	1.3	1.9	3.0	5.6	11.4	20.3	71.6
2000.....	1.0	1.0	1.0	1.3	2.1	3.3	5.4	10.0	18.8	68.8
1999.....	0.7	0.8	1.0	1.5	2.2	3.3	5.6	9.4	18.2	64.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

White, male										
2004.....	0.8	1.1	1.4	1.6	2.6	3.9	6.4	11.4	26.1	74.7
2003.....	0.9	1.3	1.2	1.6	2.6	3.4	7.1	12.2	23.0	67.6
2002.....	0.9	1.1	1.3	1.3	2.7	3.5	6.8	12.0	24.3	64.0
2001.....	1.1	1.3	1.3	1.5	2.3	3.1	5.8	11.8	21.8	59.4
2000.....	1.3	1.1	1.3	1.5	2.5	4.2	6.2	11.8	20.1	59.8
1999.....	0.9	1.0	1.3	1.9	2.6	3.8	6.1	10.8	18.7	56.8
White, female										
2004.....	0.7	0.9	0.8	1.2	2.0	3.1	6.1	12.2	25.6	100.3
2003.....	0.8	0.8	0.8	1.1	1.8	2.9	5.8	10.1	25.3	97.1
2002.....	0.7	0.8	1.1	1.1	1.5	2.8	5.3	10.0	21.2	84.3
2001.....	0.6	0.7	0.7	1.0	1.6	2.8	5.5	11.1	19.4	76.7
2000.....	0.6	0.8	0.8	1.1	1.8	2.6	4.7	8.7	18.0	72.4
1999.....	0.5	0.5	0.7	1.2	1.9	2.9	5.2	8.4	17.9	67.6
Black, both sexes										
2004.....	2.0	2.0	3.2	4.2	6.1	6.5	11.9	21.5	37.9	109.7
2003.....	1.9	3.4	3.8	3.7	5.3	7.9	14.3	18.0	33.3	100.7
2002.....	2.2	2.5	3.8	3.8	4.5	6.4	10.3	18.0	33.0	84.3
2001.....	2.4	2.3	3.3	3.8	4.9	8.5	12.9	16.0	27.6	79.2
2000.....	2.0	2.8	3.3	4.5	6.2	8.8	12.3	17.8	28.8	64.8
1999.....	2.0	2.9	3.2	4.0	5.1	8.5	11.9	18.9	26.9	71.1
Black, male										
2004.....	2.8	2.4	4.3	6.1	7.3	9.9	12.0	23.7	36.6	96.4
2003.....	2.4	4.4	5.1	4.5	6.4	8.4	14.5	18.3	36.2	97.0
2002.....	2.5	3.2	4.8	4.9	5.9	7.0	12.7	26.6	41.3	78.3
2001.....	3.3	3.3	4.2	5.8	8.9	11.4	12.8	14.9	24.1	77.2
2000.....	2.3	3.5	4.9	8.1	7.0	11.7	15.4	19.4	31.2	56.5
1999.....	2.9	4.3	3.9	5.9	5.9	10.1	12.9	23.3	29.6	58.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

Black, female

2004.....	1.3	1.6	2.3	2.6	5.2	4.0	11.9	20.2	38.6	115.0
2003.....	1.5	2.6	2.6	3.0	4.5	7.5	14.2	17.9	31.7	102.1
2002.....	1.9	1.9	3.0	2.9	3.4	5.9	8.7	12.9	28.6	86.7
2001.....	1.6	*	2.5	*	*	6.4	13.0	16.6	29.4	80.0
2000.....	1.7	2.1	2.0	*	5.6	6.7	10.2	16.8	27.6	67.8
1999.....	*	1.6	2.5	*	4.5	7.3	11.3	16.3	25.5	75.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XX. External causes of mortality (*U01-*U03,V01-Y89)

All races 3/, both sexes

2004.....	66.1	69.2	62.6	53.2	52.3	55.9	70.5	103.6	163.9	307.5
2003.....	67.2	66.7	59.9	53.4	50.7	55.4	71.3	102.9	161.9	310.4
2002.....	66.2	64.5	56.6	51.0	50.3	56.5	72.1	104.7	161.6	308.3
2001 4/.....	64.3	62.5	54.8	50.2	49.5	56.0	69.9	106.1	161.2	310.1
2000.....	61.0	57.9	51.4	48.3	49.1	52.5	69.9	100.7	155.4	310.5
1999.....	59.9	56.7	49.0	48.3	48.0	57.0	72.3	104.8	164.4	318.2

All races 3/, male

2004.....	94.1	98.8	92.1	77.7	75.6	80.7	101.1	145.9	232.1	407.1
2003.....	97.8	97.2	87.4	78.4	74.8	80.0	101.3	146.8	228.3	417.7
2002.....	96.2	94.6	83.1	76.8	73.3	82.4	104.9	150.3	226.1	414.8
2001 4/.....	95.4	94.2	81.1	74.0	71.9	80.8	100.2	152.7	231.8	429.1
2000.....	91.6	87.5	76.8	71.2	70.9	74.9	100.4	145.6	225.1	438.2
1999.....	89.5	85.9	73.2	71.9	71.0	82.8	104.4	149.4	240.9	454.8

All races 3/, female

2004.....	38.3	40.3	34.4	30.1	31.2	34.3	45.9	73.2	121.9	262.7
2003.....	37.0	37.0	33.6	29.8	28.8	34.0	47.2	71.6	121.3	262.9
2002.....	36.6	35.3	31.1	26.9	29.5	34.1	45.9	72.5	122.4	262.0
2001 4/.....	33.7	31.8	29.6	27.8	29.3	34.6	45.8	73.4	118.9	260.3
2000.....	30.9	29.1	27.0	26.8	29.3	33.2	45.9	69.5	114.3	258.5
1999.....	30.7	28.5	25.9	26.4	27.3	35.0	47.3	73.8	119.6	263.2

White, both sexes

2004.....	67.8	70.3	62.7	53.0	52.4	56.2	70.9	105.2	168.8	321.4
2003.....	68.4	67.3	59.7	53.3	50.9	55.1	71.8	104.8	165.9	323.4
2002.....	67.2	64.6	56.2	51.0	50.4	56.0	73.1	106.1	166.2	319.7
2001 4/.....	64.3	61.7	54.4	49.9	48.9	55.9	69.6	107.6	164.7	319.5
2000.....	60.0	56.8	50.0	47.6	48.9	51.8	69.4	101.5	158.6	320.4
1999.....	59.0	55.2	47.8	47.4	46.6	56.5	71.4	104.5	166.6	326.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XX. External causes of mortality (*U01-*U03,V01-Y89)

White, male

2004.....	95.6	99.3	90.8	76.0	75.0	80.8	101.4	148.3	238.1	423.8
2003.....	98.6	96.8	86.3	77.4	74.7	79.5	102.4	149.2	233.1	434.4
2002.....	97.4	93.9	81.5	75.5	73.0	80.7	105.8	151.4	231.7	427.4
2001 4/.....	95.0	92.0	79.4	72.5	70.4	80.4	99.5	154.4	236.3	441.2
2000.....	89.7	84.9	73.6	69.1	69.8	72.8	99.3	146.2	228.6	451.7
1999.....	87.8	82.5	70.6	69.2	68.6	81.4	102.7	148.6	244.1	465.8

White, female

2004.....	39.8	41.5	35.2	30.9	31.5	34.4	45.9	73.7	125.8	275.2
2003.....	37.9	38.0	33.7	30.2	28.8	33.6	46.7	72.6	124.4	274.2
2002.....	36.9	35.5	31.5	27.5	29.5	34.2	46.4	73.7	126.0	272.8
2001 4/.....	33.4	31.7	29.9	28.4	29.2	34.3	45.4	74.5	121.4	268.6
2000.....	30.2	28.9	26.9	27.0	29.7	33.4	45.3	70.1	116.9	267.0
1999.....	30.1	28.2	25.6	26.6	26.3	34.7	46.6	73.6	120.7	270.9

Black, both sexes

2004.....	69.7	77.6	74.7	65.8	58.4	59.7	71.6	99.4	125.0	176.0
2003.....	75.2	76.8	72.8	62.2	57.1	62.4	72.6	89.0	124.9	184.4
2002.....	73.4	78.0	69.9	59.7	57.4	61.7	70.5	98.6	118.5	199.6
2001 4/.....	78.0	80.6	69.3	60.7	60.8	63.5	79.8	98.6	129.2	216.8
2000.....	79.5	76.2	70.5	60.6	56.3	64.2	82.0	97.2	127.1	212.4
1999.....	77.5	78.8	65.8	61.7	62.0	68.3	85.8	115.0	147.9	239.5

Black, male

2004.....	106.3	118.0	121.9	109.0	93.4	93.8	112.7	149.7	198.7	247.4
2003.....	116.4	121.6	114.5	102.0	91.6	98.2	100.8	136.9	183.1	256.5
2002.....	109.0	122.1	112.6	102.1	92.1	100.7	110.8	157.1	182.6	307.7
2001 4/.....	119.6	131.2	111.9	100.3	96.8	97.8	120.5	151.0	199.8	306.4
2000.....	124.9	123.6	116.0	101.8	91.3	106.9	123.7	150.3	199.8	318.3
1999.....	122.3	130.6	108.6	104.1	97.6	108.0	134.0	175.2	215.4	367.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XX. External causes of mortality (*U01-*U03,V01-Y89)										
Black, female										
2004.....	37.5	42.3	34.8	30.2	31.0	34.4	43.7	69.5	86.2	147.7
2003.....	39.0	37.8	37.5	29.4	30.2	35.8	53.6	60.5	94.4	156.4
2002.....	42.1	39.7	33.7	24.9	30.2	32.7	43.5	63.7	85.1	158.1
2001 4/.....	41.3	36.8	32.9	28.2	32.5	38.1	52.7	67.4	93.2	183.6
2000.....	39.4	35.0	31.8	26.9	28.6	32.6	54.4	65.2	90.4	173.0
1999.....	38.1	34.1	29.4	27.0	34.1	38.9	53.8	78.7	114.5	191.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes

2004.....	39.7	42.6	38.6	32.8	33.7	38.4	50.5	79.6	135.9	276.7
2003.....	40.4	40.7	36.6	32.8	32.9	38.1	50.9	78.8	133.5	278.9
2002.....	39.4	38.7	34.3	30.4	32.6	38.3	50.8	80.0	131.1	275.4
2001.....	36.7	36.0	31.9	29.3	31.5	36.8	49.3	80.4	130.4	276.4
2000.....	35.4	34.2	30.8	29.9	32.2	35.4	48.8	75.5	124.6	273.5
1999.....	35.4	34.1	29.2	30.1	31.2	38.7	50.9	78.6	133.3	282.4

All races 3/, male

2004.....	56.0	60.5	56.5	46.9	46.8	52.8	67.0	104.0	179.7	345.5
2003.....	58.0	58.9	52.3	47.2	47.0	52.3	66.8	104.9	175.2	352.5
2002.....	56.8	56.6	49.6	44.8	45.7	53.4	68.9	106.4	170.1	346.6
2001.....	54.1	54.0	46.9	41.8	44.5	49.9	65.6	108.3	173.2	358.5
2000.....	53.1	51.6	46.0	43.1	45.3	48.1	64.6	99.8	164.8	359.7
1999.....	53.1	51.7	43.6	43.6	44.5	53.1	68.2	102.8	181.8	375.4

All races 3/, female

2004.....	23.6	25.2	21.4	19.5	21.7	25.8	37.2	62.1	109.0	245.8
2003.....	23.0	23.0	21.6	19.3	20.1	25.8	38.1	60.3	108.1	246.4
2002.....	22.4	21.3	19.6	16.8	20.7	25.3	36.3	61.4	107.3	244.6
2001.....	19.6	18.5	17.5	17.6	19.6	25.5	36.3	60.9	104.8	242.0
2000.....	18.0	17.3	16.1	17.5	20.4	24.5	36.4	58.5	100.9	238.4
1999.....	18.1	17.0	15.5	17.6	19.2	26.4	37.3	61.8	105.0	244.8

White, both sexes

2004.....	41.3	43.2	37.9	32.0	33.3	38.2	50.5	80.6	140.2	289.9
2003.....	41.5	41.0	35.7	32.1	32.5	37.3	50.9	80.2	136.5	291.1
2002.....	40.5	38.5	33.6	29.6	32.1	37.3	51.0	81.1	134.6	286.1
2001.....	36.8	34.9	31.1	28.5	30.5	36.4	48.9	81.2	133.3	285.5
2000.....	35.2	33.2	29.2	28.9	31.8	34.5	48.1	75.7	127.3	282.8
1999.....	35.2	32.8	27.8	28.9	29.5	37.7	49.5	78.0	135.0	290.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

White, male

2004.....	57.9	61.0	54.7	44.8	45.4	52.4	66.5	104.6	183.9	359.4
2003.....	59.6	58.6	50.6	45.6	46.1	50.8	66.9	105.8	177.3	365.7
2002.....	58.2	56.4	47.8	42.8	44.5	50.9	68.6	106.5	173.0	356.6
2001.....	54.2	51.8	45.1	40.0	42.6	48.9	64.6	108.2	175.2	368.1
2000.....	52.5	49.8	43.2	40.8	43.8	45.9	63.3	98.7	166.4	369.8
1999.....	52.5	49.1	41.1	41.1	42.0	51.1	65.7	100.7	183.4	383.6

White, female

2004.....	24.5	25.5	21.5	19.7	22.0	25.6	37.3	63.0	113.0	258.6
2003.....	23.4	23.4	21.1	19.2	19.9	25.4	37.9	61.7	111.3	258.0
2002.....	22.8	20.7	19.7	16.9	20.7	25.3	36.7	62.9	111.2	255.4
2001.....	19.3	18.1	17.4	17.6	19.3	25.4	36.1	62.0	107.9	251.0
2000.....	17.8	16.8	15.5	17.5	20.7	24.6	35.8	59.6	104.0	247.4
1999.....	17.9	16.6	14.9	17.4	18.0	26.1	36.6	62.0	106.5	253.0

Black, both sexes

2004.....	38.4	49.0	50.7	45.2	41.7	43.8	53.8	77.9	100.5	152.3
2003.....	41.7	47.9	49.1	43.7	40.4	47.7	53.8	67.7	105.1	160.2
2002.....	40.0	48.3	45.1	42.0	40.6	47.3	51.9	74.9	96.8	173.4
2001.....	44.3	50.1	44.8	40.6	43.7	45.5	58.0	78.2	104.5	183.6
2000.....	43.8	47.0	47.0	42.0	39.4	46.1	60.2	75.3	100.0	181.1
1999.....	43.6	50.4	44.4	43.4	46.3	50.7	66.6	90.3	124.2	207.6

Black, male

2004.....	56.5	72.2	82.2	73.2	66.7	67.4	82.2	116.0	157.1	213.6
2003.....	60.8	73.9	75.2	70.1	64.8	74.4	74.2	105.0	152.8	225.2
2002.....	57.9	72.0	72.6	71.2	65.3	75.8	78.4	117.6	149.8	260.5
2001.....	66.6	81.7	71.8	65.9	69.8	69.1	82.5	119.4	164.5	255.3
2000.....	68.8	74.5	76.5	71.5	65.4	74.2	86.3	119.5	156.0	275.6
1999.....	69.0	82.7	73.2	71.5	70.6	77.9	101.8	135.3	177.0	316.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Black, female

2004.....	22.6	28.8	24.2	22.0	22.3	26.3	34.5	55.3	70.7	128.0
2003.....	24.8	25.3	27.0	22.0	21.4	28.0	40.1	45.6	80.1	134.9
2002.....	24.2	27.7	21.8	18.1	21.2	26.1	34.1	49.4	69.2	140.1
2001.....	24.5	22.7	21.8	19.9	23.2	28.0	41.7	53.5	74.0	157.0
2000.....	21.7	23.1	21.9	17.8	18.8	25.3	42.8	48.7	71.8	145.9
1999.....	21.3	22.5	20.1	20.4	27.3	30.6	43.2	63.2	98.1	166.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Transport accidents (V01-V99,Y85)

All races 3/, both sexes

2004.....	16.4	17.1	16.3	15.2	15.9	17.0	18.1	23.6	28.1	27.2
2003.....	17.5	16.8	16.3	15.9	16.0	16.2	19.0	23.9	29.4	29.7
2002.....	17.1	16.9	15.8	14.9	16.7	16.7	20.2	25.0	29.7	28.8
2001.....	17.3	16.3	15.6	15.2	15.4	16.4	19.5	26.2	30.4	28.9
2000.....	16.8	16.2	15.5	15.7	16.4	16.8	19.4	25.1	30.0	31.5
1999.....	16.6	15.9	14.9	16.3	15.7	18.1	20.9	25.9	32.1	32.0

All races 3/, male

2004.....	24.1	24.9	24.4	22.2	22.4	23.2	24.3	32.3	42.6	46.5
2003.....	25.5	24.9	23.6	23.2	22.7	22.5	25.0	32.9	42.1	52.0
2002.....	24.5	25.1	22.7	22.2	22.8	23.4	27.5	34.5	42.6	53.0
2001.....	25.4	24.0	22.5	21.4	22.0	22.1	26.8	36.2	45.5	52.7
2000.....	24.8	23.6	22.3	22.4	22.6	22.6	26.3	34.5	44.0	59.8
1999.....	24.0	23.1	21.5	23.0	21.8	24.8	27.5	36.2	48.3	63.5

All races 3/, female

2004.....	8.9	9.4	8.6	8.7	10.0	11.5	13.1	17.3	19.1	18.5
2003.....	9.5	9.0	9.2	9.0	9.9	10.7	14.2	17.5	21.7	19.9
2002.....	9.9	8.8	9.2	7.9	11.1	10.9	14.3	18.3	21.8	18.3
2001.....	9.4	8.8	8.9	9.4	9.3	11.6	13.8	19.2	21.3	18.9
2000.....	9.0	8.9	8.9	9.5	10.7	11.7	13.9	18.5	21.7	19.9
1999.....	9.4	8.9	8.6	10.0	10.2	12.4	15.7	18.8	22.6	19.3

White, both sexes

2004.....	16.8	17.2	16.0	15.2	15.9	17.1	18.2	23.6	29.0	28.4
2003.....	17.7	16.9	16.1	15.7	15.8	16.0	19.2	23.9	30.0	30.5
2002.....	17.5	16.9	15.9	14.8	16.7	16.5	20.5	25.5	30.3	29.6
2001.....	17.5	15.9	15.7	15.1	15.2	16.4	19.5	26.6	31.2	29.9
2000.....	16.8	16.0	15.1	15.6	16.4	16.6	19.2	25.3	30.7	32.2
1999.....	16.7	15.6	14.6	16.1	15.4	18.0	20.8	26.0	32.4	32.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Transport accidents (V01-V99,Y85)

White, male

2004.....	24.4	25.0	23.7	21.9	22.0	23.2	24.1	32.2	43.2	48.6
2003.....	25.8	24.9	23.3	22.8	22.6	22.2	25.4	32.7	42.6	52.8
2002.....	24.9	25.3	22.5	21.9	22.8	22.7	27.5	34.4	42.8	53.8
2001.....	25.4	23.2	22.6	21.0	21.5	21.7	26.6	36.2	46.6	54.1
2000.....	24.6	23.4	21.7	21.8	22.2	21.9	25.8	34.3	44.5	61.6
1999.....	24.0	22.5	21.0	22.5	21.1	24.6	27.1	35.9	48.4	63.7

White, female

2004.....	9.0	9.5	8.5	8.8	10.3	11.6	13.4	17.3	20.1	19.3
2003.....	9.6	9.1	9.1	8.8	9.6	10.5	14.0	17.6	22.2	20.6
2002.....	10.0	8.6	9.3	8.1	11.0	11.0	14.8	19.1	22.6	19.1
2001.....	9.5	8.7	8.9	9.5	9.3	11.6	13.8	19.8	21.8	19.7
2000.....	9.0	8.7	8.6	9.6	11.1	12.0	13.9	19.0	22.6	20.3
1999.....	9.4	8.8	8.4	9.9	10.1	12.3	15.8	19.1	23.0	19.7

Black, both sexes

2004.....	16.7	19.1	19.9	17.2	17.4	16.6	16.2	23.0	19.6	14.7
2003.....	18.7	18.1	18.2	18.9	17.8	18.5	17.9	22.8	21.8	21.0
2002.....	17.1	18.9	16.5	16.8	17.4	17.3	16.9	20.1	24.1	18.4
2001.....	19.4	20.5	16.8	17.8	18.6	18.6	19.0	22.1	19.8	17.5
2000.....	19.3	19.4	19.3	18.1	16.4	18.2	21.3	19.4	20.9	21.6
1999.....	18.0	19.6	18.3	18.7	17.7	18.0	21.0	24.6	28.3	27.7

Black, male

2004.....	25.8	29.6	32.6	27.1	28.6	27.1	26.2	36.9	40.9	28.8
2003.....	28.3	29.1	28.9	30.3	27.6	28.6	24.0	37.4	37.7	42.7
2002.....	25.2	29.1	26.3	29.0	27.3	28.7	28.4	38.1	44.3	46.1
2001.....	30.6	33.8	25.7	28.6	30.9	28.2	26.3	39.0	31.3	35.2
2000.....	31.0	29.3	30.0	31.0	27.5	31.2	33.4	34.5	39.6	45.0
1999.....	28.3	31.2	29.1	29.3	28.4	27.3	31.5	40.4	48.8	68.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
Black, female										
2004.....	8.6	10.0	9.1	9.1	8.7	8.8	9.4	14.8	8.3	9.1
2003.....	10.3	8.4	9.2	9.6	10.2	11.0	13.7	14.1	13.5	12.6
2002.....	10.0	10.0	8.2	6.8	9.6	8.9	9.2	9.3	13.5	*
2001.....	9.4	8.9	9.3	8.9	9.0	11.5	14.1	11.9	13.9	10.9
2000.....	9.0	10.8	10.2	7.6	7.7	8.6	13.3	10.3	11.5	12.9
1999.....	9.0	9.6	9.1	10.1	9.3	11.2	14.0	15.2	18.1	12.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Railway accidents (V05,V15,V80.6,V81.2-V81.9)

All races 3/, both sexes

2004.....	0.2	0.3	0.2	0.1	*	*	*	*	*	*
2003.....	0.2	0.3	0.2	0.2	*	*	*	*	*	*
2002.....	0.3	0.2	0.2	0.2	*	*	*	*	*	*
2001.....	0.3	0.3	0.2	0.1	*	*	*	*	*	*
2000.....	0.3	0.2	0.2	*	*	*	*	*	*	*
1999.....	0.3	0.2	0.2	*	*	*	*	*	*	*

All races 3/, male

2004.....	0.4	0.5	0.4	0.3	*	*	*	*	*	*
2003.....	0.4	0.5	0.3	0.3	*	*	*	*	*	*
2002.....	0.6	0.5	0.4	0.3	*	*	*	*	*	*
2001.....	0.5	0.4	0.4	*	*	*	*	*	*	*
2000.....	0.6	0.4	0.3	*	*	*	*	*	*	*
1999.....	0.5	0.4	0.3	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	0.2	0.2	*	*	*	*	*	*	*
2003.....	0.2	0.2	0.2	0.2	*	*	*	*	*	*
2002.....	0.3	0.2	0.2	*	*	*	*	*	*	*
2001.....	0.3	0.2	0.2	*	*	*	*	*	*	*
2000.....	0.3	0.2	0.1	*	*	*	*	*	*	*
1999.....	0.3	0.2	0.1	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes

2004.....	15.1	15.5	14.6	13.9	14.3	15.5	16.8	22.7	26.9	26.1
2003.....	16.0	15.1	14.6	14.1	14.3	14.8	17.7	22.5	28.2	28.8
2002.....	15.7	15.2	14.4	13.4	15.2	15.3	18.9	23.8	28.4	28.0
2001.....	15.5	14.5	13.9	13.7	13.7	14.8	18.0	24.6	28.9	27.8
2000.....	15.1	14.6	13.8	14.2	14.7	15.1	18.1	23.8	28.5	30.4
1999.....	14.8	14.0	13.1	14.4	13.9	16.4	19.3	24.1	30.1	30.4

All races 3/, male

2004.....	21.9	22.3	21.6	19.9	19.6	20.8	22.0	30.8	40.4	44.2
2003.....	23.1	22.0	20.8	20.3	19.6	20.1	22.8	30.5	40.0	50.0
2002.....	22.1	22.2	20.3	19.6	20.2	20.8	25.0	32.3	39.9	51.7
2001.....	22.5	21.2	19.7	18.9	19.2	19.2	24.2	33.6	43.1	51.1
2000.....	21.9	21.1	19.6	19.9	19.7	19.8	23.9	32.2	41.4	57.5
1999.....	21.0	20.0	18.6	19.8	18.7	22.0	24.7	33.6	45.2	59.8

All races 3/, female

2004.....	8.3	8.8	8.0	8.2	9.5	10.9	12.6	16.9	18.6	17.9
2003.....	9.0	8.3	8.7	8.3	9.4	10.3	13.7	16.9	21.0	19.5
2002.....	9.4	8.4	8.7	7.5	10.6	10.6	14.1	17.7	21.4	17.7
2001.....	8.7	8.0	8.3	8.8	8.8	11.0	13.1	18.2	20.4	18.0
2000.....	8.4	8.3	8.2	9.0	10.2	11.0	13.5	17.9	20.9	19.3
1999.....	8.6	8.1	8.0	9.4	9.5	11.6	15.0	17.5	21.3	18.5

White, both sexes

2004.....	15.4	15.6	14.3	13.9	14.3	15.6	16.8	22.8	27.7	27.3
2003.....	16.2	15.2	14.4	13.8	14.0	14.6	17.8	22.6	28.8	29.5
2002.....	16.0	15.2	14.3	13.3	15.1	15.1	19.2	24.3	29.0	28.8
2001.....	15.6	14.2	13.9	13.6	13.5	14.7	17.9	25.0	29.7	28.8
2000.....	15.0	14.5	13.4	14.1	14.7	14.9	17.9	24.0	29.2	31.1
1999.....	14.8	13.7	12.9	14.1	13.5	16.3	19.2	24.3	30.6	30.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, male

2004.....	22.2	22.3	20.9	19.6	19.2	20.7	21.7	30.8	40.8	46.3
2003.....	23.2	22.1	20.5	19.7	19.4	19.9	23.0	30.3	40.6	50.7
2002.....	22.5	22.3	20.0	19.3	20.0	20.1	25.0	32.4	40.1	52.4
2001.....	22.5	20.5	19.7	18.7	18.5	19.0	23.9	33.6	44.3	52.6
2000.....	21.6	20.9	18.9	19.3	19.2	19.1	23.4	32.0	41.7	59.2
1999.....	20.9	19.4	18.1	19.2	18.1	21.8	24.4	33.3	45.5	60.3

White, female

2004.....	8.4	8.9	7.9	8.3	9.8	11.1	12.8	16.9	19.6	18.7
2003.....	9.0	8.4	8.6	8.1	9.1	10.0	13.5	17.1	21.5	20.1
2002.....	9.5	8.2	8.8	7.6	10.5	10.7	14.5	18.5	22.2	18.5
2001.....	8.8	8.0	8.2	8.8	8.8	11.0	13.0	19.0	20.9	18.9
2000.....	8.4	8.1	7.9	9.1	10.6	11.3	13.5	18.4	21.8	19.7
1999.....	8.6	8.0	7.7	9.3	9.3	11.6	15.1	18.0	21.8	18.8

Black, both sexes

2004.....	15.4	17.3	18.3	15.9	15.5	15.6	15.3	20.8	19.3	14.7
2003.....	17.5	16.2	16.8	17.5	16.4	16.9	17.2	21.3	20.6	20.7
2002.....	15.9	16.9	15.1	15.0	16.2	15.9	15.8	18.4	22.5	18.1
2001.....	17.6	18.1	15.3	15.9	16.9	16.5	18.3	20.1	19.0	16.6
2000.....	17.9	17.6	18.3	16.7	15.4	16.6	20.1	18.1	20.0	20.3
1999.....	16.6	17.8	16.3	17.1	15.8	16.6	19.6	22.9	26.0	25.5

Black, male

2004.....	23.5	26.7	29.7	24.5	25.0	25.4	24.3	33.2	40.2	28.8
2003.....	26.4	25.8	26.0	27.6	24.5	25.1	23.0	35.1	34.7	41.7
2002.....	23.0	25.6	23.6	25.3	24.9	25.4	25.7	33.5	40.5	45.0
2001.....	27.5	29.9	23.2	24.7	28.5	23.9	24.6	36.2	30.5	35.2
2000.....	28.4	26.3	28.1	28.2	25.7	27.8	31.4	32.1	38.8	43.8
1999.....	25.6	28.2	25.5	26.3	24.6	24.9	28.5	38.9	45.3	60.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, female

2004.....	8.3	9.1	8.8	8.8	8.0	8.3	9.2	13.5	8.3	9.1
2003.....	9.7	7.9	9.1	9.2	10.1	10.8	13.3	13.0	13.2	12.6
2002.....	9.6	9.4	8.0	6.6	9.4	8.9	9.2	9.3	13.1	*
2001.....	8.9	8.0	8.7	8.7	7.8	11.1	14.1	10.5	13.1	9.7
2000.....	8.6	10.0	9.9	7.3	7.4	8.3	12.7	9.7	10.6	11.6
1999.....	8.7	8.9	8.5	9.6	8.8	10.4	13.7	13.2	16.4	12.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pedestrian involved in collision with motor vehicle (V02-V04)

All races 3/, both sexes

2004.....	1.2	1.4	1.4	1.4	1.5	1.3	1.8	2.2	2.8	2.7
2003.....	1.4	1.5	1.3	1.4	1.5	1.6	1.6	2.3	3.1	2.9
2002.....	1.4	1.4	1.3	1.4	1.4	1.9	2.1	2.4	2.8	3.3
2001.....	1.5	1.4	1.4	1.4	1.3	1.2	2.0	2.4	3.8	3.0
2000.....	1.4	1.4	1.3	1.5	1.5	1.5	1.7	2.5	3.3	3.6
1999.....	1.3	1.3	1.2	1.6	1.5	1.4	1.9	2.3	2.8	3.7

All races 3/, male

2004.....	1.9	2.0	2.3	2.0	2.2	1.8	2.1	2.9	4.2	4.9
2003.....	2.1	2.1	2.0	2.3	2.2	2.3	2.3	2.7	4.1	5.7
2002.....	2.0	2.2	2.0	2.0	1.9	2.6	2.8	3.0	3.9	5.8
2001.....	2.2	2.0	1.9	2.0	2.0	1.4	2.9	3.2	5.6	5.9
2000.....	2.0	2.2	1.9	2.3	2.1	2.1	2.3	3.4	4.5	7.0
1999.....	1.9	2.0	1.9	2.3	2.1	2.0	2.5	3.2	4.7	6.5

All races 3/, female

2004.....	0.6	0.8	0.5	0.8	0.9	0.9	1.5	1.7	2.0	1.7
2003.....	0.7	0.9	0.6	0.7	0.9	1.1	1.0	2.1	2.5	1.7
2002.....	0.9	0.7	0.7	0.8	1.0	1.3	1.5	2.0	2.1	2.2
2001.....	0.7	0.8	0.8	0.8	0.7	1.0	1.3	1.8	2.7	1.8
2000.....	0.8	0.6	0.8	0.8	0.9	1.0	1.3	1.9	2.5	2.3
1999.....	0.7	0.6	0.5	0.9	0.9	0.9	1.4	1.7	1.7	2.6

White, both sexes

2004.....	1.2	1.2	1.2	1.3	1.4	1.2	1.7	2.1	2.8	2.6
2003.....	1.2	1.4	1.1	1.4	1.5	1.4	1.5	2.1	2.9	2.9
2002.....	1.3	1.3	1.2	1.3	1.3	1.6	2.0	2.2	2.6	3.2
2001.....	1.2	1.3	1.2	1.3	1.2	1.1	1.8	2.2	3.5	2.9
2000.....	1.2	1.2	1.2	1.3	1.4	1.4	1.6	2.3	3.2	3.4
1999.....	1.0	1.2	1.1	1.4	1.4	1.3	1.7	2.1	2.7	3.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pedestrian involved in collision with motor vehicle (V02-V04)

White, male

2004.....	1.8	1.8	2.0	1.8	2.0	1.7	2.0	2.7	4.0	4.6
2003.....	1.9	2.0	1.8	2.2	2.1	2.0	2.1	2.4	3.9	5.4
2002.....	1.8	2.0	1.8	1.9	1.8	2.3	2.7	2.7	3.6	5.4
2001.....	1.9	1.9	1.7	1.9	1.7	1.3	2.6	2.9	5.5	5.5
2000.....	1.7	1.9	1.7	2.0	1.9	1.8	2.1	3.0	4.1	6.2
1999.....	1.5	1.7	1.7	2.0	2.0	1.9	2.1	2.8	4.4	6.0

White, female

2004.....	0.5	0.7	0.5	0.9	0.8	0.8	1.5	1.6	2.0	1.7
2003.....	0.6	0.8	0.5	0.6	0.8	1.0	0.9	2.0	2.4	1.8
2002.....	0.8	0.6	0.7	0.8	0.9	1.1	1.5	1.8	2.0	2.2
2001.....	0.6	0.7	0.7	0.8	0.7	0.9	1.1	1.7	2.4	1.9
2000.....	0.6	0.5	0.7	0.7	0.9	1.0	1.3	1.8	2.6	2.2
1999.....	0.6	0.6	0.5	0.8	0.8	0.7	1.4	1.6	1.7	2.6

Black, both sexes

2004.....	1.7	2.4	2.7	2.2	2.4	*	*	*	*	*
2003.....	2.5	2.4	2.2	2.0	1.7	2.5	*	*	*	*
2002.....	2.0	2.6	2.3	1.9	2.2	2.7	*	*	*	*
2001.....	2.7	2.5	2.6	2.0	2.5	*	*	*	*	*
2000.....	2.8	2.7	2.2	3.0	1.9	*	*	*	*	*
1999.....	3.1	2.4	2.1	3.4	*	*	3.1	*	*	*

Black, male

2004.....	2.5	3.5	5.2	4.0	4.3	*	*	*	*	*
2003.....	3.7	3.7	3.7	3.2	*	4.9	*	*	*	*
2002.....	2.9	4.4	3.7	3.4	*	*	*	*	*	*
2001.....	4.4	3.7	3.9	3.6	5.1	*	*	*	*	*
2000.....	4.0	4.4	3.3	5.3	*	*	*	*	*	*
1999.....	5.1	4.2	3.9	5.9	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)										
All races 3/, both sexes										
2004.....	0.3	0.3	0.3	0.2	0.2	0.3	0.3	0.3	*	*
2003.....	0.3	0.3	0.3	0.2	0.2	*	0.3	*	*	*
2002.....	0.3	0.4	0.2	0.2	0.2	*	*	*	*	*
2001.....	0.3	0.3	0.2	0.2	0.2	0.2	*	*	*	*
2000.....	0.3	0.3	0.2	0.3	*	*	*	*	*	*
1999.....	0.4	0.3	0.2	0.2	*	0.2	*	*	*	*
All races 3/, male										
2004.....	0.6	0.5	0.5	0.4	0.4	0.6	0.6	0.6	*	*
2003.....	0.6	0.4	0.5	0.4	0.3	*	0.5	*	*	*
2002.....	0.5	0.7	0.3	0.3	*	*	*	*	*	*
2001.....	0.6	0.6	0.5	0.5	0.4	0.5	*	*	*	*
2000.....	0.5	0.6	0.4	0.5	*	*	*	*	*	*
1999.....	0.6	0.5	0.4	*	*	0.5	*	*	*	*
All races 3/, female										
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2004.....	0.3	0.3	0.3	0.2	0.2	0.3	*	*	*	*
2003.....	0.3	0.2	0.3	0.2	*	*	0.3	*	*	*
2002.....	0.3	0.3	0.2	0.2	*	*	*	*	*	*
2001.....	0.3	0.3	0.2	0.2	0.2	*	*	*	*	*
2000.....	0.2	0.3	0.2	0.3	*	*	*	*	*	*
1999.....	0.4	0.2	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)

All races 3/, both sexes

2004.....	2.1	2.1	2.0	1.5	1.0	0.7	0.4	0.3	*	*
2003.....	2.1	2.0	1.8	1.5	0.8	0.6	0.5	*	*	*
2002.....	1.8	1.7	1.6	1.2	0.8	0.5	0.3	*	*	*
2001.....	1.7	1.5	1.3	0.8	0.6	0.4	0.3	*	*	*
2000.....	1.6	1.4	1.1	0.9	0.6	0.5	0.3	*	*	*
1999.....	1.4	1.1	0.9	0.8	0.6	0.3	*	*	*	*

All races 3/, male

2004.....	3.7	3.7	3.7	2.8	1.8	1.5	0.8	*	*	*
2003.....	3.6	3.4	3.2	2.7	1.5	1.2	1.0	*	*	*
2002.....	3.2	3.1	2.9	2.1	1.6	1.0	0.6	*	*	*
2001.....	3.0	2.7	2.3	1.6	1.2	0.8	0.6	*	*	*
2000.....	2.9	2.5	2.0	1.6	1.1	0.9	0.7	*	*	*
1999.....	2.3	2.0	1.6	1.5	1.1	0.7	*	*	*	*

All races 3/, female

2004.....	0.6	0.5	0.5	0.2	*	*	*	*	*	*
2003.....	0.6	0.5	0.4	0.3	*	*	*	*	*	*
2002.....	0.4	0.4	0.3	0.3	*	*	*	*	*	*
2001.....	0.4	0.3	0.3	*	*	*	*	*	*	*
2000.....	0.4	0.4	*	*	*	*	*	*	*	*
1999.....	0.4	0.3	*	*	*	*	*	*	*	*

White, both sexes

2004.....	2.4	2.4	2.3	1.6	1.1	0.8	0.4	0.4	*	*
2003.....	2.3	2.3	2.0	1.7	0.9	0.7	0.5	*	*	*
2002.....	2.0	1.9	1.9	1.3	0.9	0.5	0.3	*	*	*
2001.....	1.9	1.7	1.5	0.9	0.7	0.4	0.3	*	*	*
2000.....	1.9	1.7	1.2	1.0	0.7	0.5	0.3	*	*	*
1999.....	1.5	1.3	1.0	0.9	0.6	0.4	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

All races 3/, both sexes

2004.....	1.4	1.3	1.1	1.1	1.2	1.2	1.3	1.4	1.8	1.2
2003.....	1.6	1.5	1.4	1.1	1.1	1.1	1.6	1.5	1.7	1.9
2002.....	1.6	1.5	1.3	1.2	1.4	1.2	1.2	1.5	1.7	1.5
2001.....	1.4	1.3	1.1	0.9	1.0	1.1	1.1	1.7	1.6	1.7
2000.....	1.2	1.1	1.0	0.9	1.0	1.0	1.3	1.3	1.5	1.7
1999.....	1.2	1.1	1.0	1.0	1.0	1.3	1.1	1.4	1.6	1.2

All races 3/, male

2004.....	2.0	2.0	1.6	1.7	1.7	1.7	1.9	2.4	2.6	1.9
2003.....	2.4	2.2	1.9	1.7	1.6	1.6	2.2	2.2	2.5	3.7
2002.....	2.4	2.1	1.8	1.8	2.3	1.7	1.8	2.3	2.1	3.2
2001.....	2.2	2.0	1.6	1.6	1.5	1.6	1.5	2.5	2.6	3.1
2000.....	1.8	1.5	1.5	1.3	1.5	1.5	1.8	2.0	2.2	3.0
1999.....	1.8	1.6	1.6	1.4	1.5	1.9	1.6	2.1	2.8	2.5

All races 3/, female

2004.....	0.7	0.6	0.6	0.5	0.7	0.8	0.9	0.7	1.3	0.9
2003.....	0.8	0.8	0.8	0.5	0.6	0.7	1.1	1.1	1.1	1.1
2002.....	0.9	0.9	0.7	0.6	0.6	0.8	0.8	0.9	1.5	0.8
2001.....	0.6	0.6	0.6	0.3	0.5	0.7	0.8	1.1	0.9	1.2
2000.....	0.7	0.7	0.5	0.5	0.6	0.6	1.0	0.8	1.0	1.2
1999.....	0.6	0.7	0.4	0.7	0.6	0.8	0.7	1.0	0.9	0.7

White, both sexes

2004.....	1.4	1.3	1.1	1.0	1.1	1.2	1.3	1.5	1.9	1.2
2003.....	1.7	1.5	1.3	1.1	1.0	1.0	1.6	1.6	1.8	2.0
2002.....	1.8	1.5	1.2	1.1	1.4	1.1	1.2	1.6	1.8	1.6
2001.....	1.4	1.2	1.1	0.9	0.9	1.1	1.1	1.7	1.5	1.9
2000.....	1.2	1.0	0.9	0.8	1.0	1.0	1.3	1.2	1.5	1.8
1999.....	1.2	1.1	0.9	0.9	1.0	1.3	1.1	1.3	1.6	1.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

All races 3/, both sexes

2004.....	2.9	3.1	3.2	3.4	3.6	4.8	5.1	7.6	9.2	9.6
2003.....	3.3	3.1	3.4	3.3	3.7	4.1	5.8	7.7	10.0	10.2
2002.....	3.4	3.2	3.4	3.4	4.4	4.2	6.2	8.3	10.4	9.9
2001.....	3.3	3.1	3.1	3.5	3.6	4.6	5.7	7.8	9.1	10.1
2000.....	3.0	3.2	3.1	3.6	4.0	4.2	5.9	7.9	10.5	10.1
1999.....	3.3	3.2	3.2	3.8	3.7	4.5	6.1	7.7	10.4	9.2

All races 3/, male

2004.....	3.6	3.7	3.8	4.0	4.0	5.6	6.0	9.2	13.4	15.9
2003.....	4.2	3.9	4.2	4.0	4.4	4.7	6.3	9.7	13.4	16.6
2002.....	4.2	4.0	4.2	4.5	4.9	4.9	7.3	10.7	14.5	16.9
2001.....	4.1	3.9	3.7	4.1	4.2	5.5	6.8	9.6	13.0	18.1
2000.....	3.7	4.2	3.9	4.4	4.5	4.6	7.2	9.5	15.0	18.2
1999.....	4.1	4.0	3.9	4.2	4.4	5.2	7.3	9.7	14.9	17.8

All races 3/, female

2004.....	2.2	2.5	2.5	2.8	3.2	4.2	4.5	6.4	6.7	6.7
2003.....	2.4	2.3	2.6	2.7	3.1	3.6	5.3	6.3	7.9	7.4
2002.....	2.6	2.4	2.7	2.3	3.9	3.6	5.3	6.7	7.9	6.8
2001.....	2.5	2.4	2.5	2.9	3.1	3.8	4.8	6.5	6.9	6.8
2000.....	2.3	2.3	2.4	2.8	3.6	3.8	4.9	6.9	7.8	6.8
1999.....	2.4	2.4	2.5	3.4	3.1	4.0	5.2	6.4	7.8	5.7

White, both sexes

2004.....	3.0	3.1	3.1	3.4	3.7	5.0	5.3	7.7	9.6	10.1
2003.....	3.3	3.2	3.3	3.1	3.6	4.0	5.9	7.9	10.5	10.6
2002.....	3.5	3.1	3.4	3.4	4.4	4.3	6.4	8.7	10.9	10.4
2001.....	3.4	3.0	3.1	3.6	3.5	4.6	5.7	8.2	9.5	10.6
2000.....	3.0	3.2	3.0	3.6	4.0	4.3	6.0	8.2	10.9	10.6
1999.....	3.2	3.2	3.1	3.8	3.7	4.6	6.2	8.0	10.7	9.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

White, male										
2004.....	3.7	3.6	3.8	3.9	4.0	5.8	6.0	9.3	13.5	16.6
2003.....	4.1	4.0	4.1	3.7	4.3	4.6	6.4	9.8	13.8	17.3
2002.....	4.2	3.9	4.2	4.4	4.9	4.9	7.3	10.8	15.0	17.9
2001.....	4.0	3.6	3.6	4.3	4.0	5.4	6.9	9.9	13.4	18.9
2000.....	3.6	4.1	3.7	4.4	4.1	4.7	7.3	9.6	15.6	19.3
1999.....	4.0	4.0	3.7	4.1	4.3	5.0	7.3	9.8	15.2	18.1
White, female										
2004.....	2.2	2.6	2.5	2.8	3.4	4.3	4.8	6.6	7.1	7.2
2003.....	2.4	2.3	2.7	2.6	3.0	3.5	5.4	6.5	8.5	7.7
2002.....	2.7	2.3	2.7	2.3	4.0	3.7	5.7	7.1	8.4	7.1
2001.....	2.7	2.4	2.6	3.0	3.1	3.9	4.8	6.9	7.2	7.2
2000.....	2.3	2.3	2.3	2.9	3.8	4.0	5.0	7.2	8.2	7.1
1999.....	2.5	2.4	2.6	3.5	3.2	4.2	5.3	6.7	8.1	5.9
Black, both sexes										
2004.....	2.8	3.6	3.3	4.3	2.9	4.5	4.2	6.3	7.4	5.6
2003.....	3.8	3.0	3.9	4.9	4.4	5.1	4.9	6.5	6.6	6.4
2002.....	3.7	3.5	3.5	3.3	4.3	3.4	4.8	5.5	6.0	*
2001.....	3.7	4.1	3.6	3.1	4.7	5.4	6.5	4.3	6.2	*
2000.....	3.6	4.0	4.4	3.5	4.7	3.4	5.5	4.8	6.8	*
1999.....	3.7	3.5	3.7	4.1	3.9	4.5	5.7	5.2	7.8	7.7
Black, male										
2004.....	3.3	5.1	3.8	5.3	4.7	5.6	*	9.1	16.5	*
2003.....	5.4	3.7	5.5	6.7	5.6	6.6	*	9.6	*	*
2002.....	4.8	4.6	4.9	5.1	5.5	*	7.0	9.2	*	*
2001.....	5.6	6.6	4.9	3.8	6.1	7.6	*	*	*	*
2000.....	5.4	5.7	5.8	5.6	8.1	*	*	*	*	*
1999.....	5.5	4.8	5.1	5.4	5.3	6.8	8.8	9.5	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

Black, female

2004.....	2.3	2.4	2.8	3.6	*	3.8	*	*	*	*
2003.....	2.4	2.3	2.6	3.5	3.5	4.0	4.7	*	*	*
2002.....	2.8	2.6	2.4	*	3.3	*	*	*	*	*
2001.....	2.0	1.9	2.6	2.6	3.7	3.8	6.5	*	*	*
2000.....	2.1	2.5	3.3	*	*	*	4.9	*	*	*
1999.....	2.1	2.4	2.5	3.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

All races 3/, both sexes										
2004.....	1.5	1.3	1.2	1.1	1.1	1.1	0.9	0.9	0.8	1.0
2003.....	1.6	1.4	1.3	1.2	1.1	1.0	1.0	1.0	1.0	1.3
2002.....	1.5	1.4	1.2	1.1	1.3	0.9	1.0	1.0	1.0	0.9
2001.....	1.4	1.3	1.2	1.0	1.2	0.9	1.0	1.1	1.0	0.7
2000.....	1.3	1.2	1.1	0.9	1.0	0.8	0.8	0.9	0.9	1.0
1999.....	1.4	1.2	1.0	1.1	1.0	1.1	0.8	0.8	0.8	1.0
All races 3/, male										
2004.....	2.1	1.8	1.8	1.7	1.5	1.5	1.3	1.4	1.1	1.7
2003.....	2.2	2.0	1.8	1.6	1.5	1.5	1.5	1.5	1.7	2.0
2002.....	2.2	2.2	1.7	1.6	1.7	1.4	1.5	1.5	1.6	1.8
2001.....	2.0	2.0	1.8	1.5	1.7	1.3	1.5	1.5	1.4	*
2000.....	1.9	1.9	1.6	1.3	1.4	1.1	1.3	1.4	1.3	2.2
1999.....	2.0	1.7	1.5	1.6	1.5	1.6	0.9	1.3	*	2.0
All races 3/, female										
2004.....	0.9	0.8	0.6	0.5	0.7	0.6	0.5	0.6	*	0.7
2003.....	0.9	0.8	0.8	0.8	0.7	0.7	0.6	0.7	*	1.0
2002.....	0.8	0.7	0.7	0.6	0.9	0.6	0.6	0.7	0.6	*
2001.....	0.8	0.6	0.7	0.6	0.7	0.6	0.6	0.8	0.7	*
2000.....	0.7	0.6	0.5	0.6	0.6	0.6	*	0.5	0.7	*
1999.....	0.8	0.7	0.5	0.6	0.7	0.6	0.7	0.5	*	*
White, both sexes										
2004.....	1.5	1.3	1.2	1.1	1.1	1.1	0.8	0.9	0.8	1.1
2003.....	1.7	1.5	1.3	1.2	1.0	1.0	1.0	1.0	1.0	1.3
2002.....	1.5	1.5	1.2	1.1	1.3	1.0	1.0	1.0	1.0	0.8
2001.....	1.5	1.2	1.3	0.9	1.1	1.0	1.0	1.1	1.0	0.7
2000.....	1.3	1.3	1.0	0.8	0.9	0.9	0.8	0.9	0.9	1.0
1999.....	1.4	1.2	1.0	1.1	1.0	1.1	0.8	0.8	0.8	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Occupant of special-use motor vehicle involved in any accident (V83-V86)										
All races 3/, both sexes										
2004.....	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.4	*	*
2003.....	0.4	0.4	0.4	0.4	0.3	0.4	0.3	0.5	0.4	*
2002.....	0.3	0.3	0.2	0.3	0.3	0.3	0.5	0.4	0.5	*
2001.....	0.3	0.2	0.3	0.4	0.4	0.4	0.4	0.6	0.5	*
2000.....	0.3	0.3	0.3	0.4	0.5	0.5	0.4	0.5	0.4	*
1999.....	0.3	0.3	0.4	0.4	0.5	0.6	0.6	0.6	0.6	0.7
All races 3/, male										
2004.....	0.7	0.7	0.5	0.5	0.7	0.6	0.7	0.8	*	*
2003.....	0.7	0.7	0.6	0.7	0.6	0.8	0.6	1.1	*	*
2002.....	0.5	0.7	0.4	0.5	0.6	0.7	0.9	0.9	1.2	*
2001.....	0.6	0.4	0.5	0.7	0.8	0.9	0.8	1.5	1.3	*
2000.....	0.6	0.5	0.5	0.8	1.1	1.0	0.8	1.1	1.1	*
1999.....	0.6	0.6	0.7	0.7	0.9	1.3	1.2	1.4	1.3	2.2
All races 3/, female										
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2004.....	0.4	0.4	0.3	0.3	0.4	0.3	0.3	0.4	*	*
2003.....	0.5	0.5	0.4	0.4	0.3	0.5	0.3	0.5	*	*
2002.....	0.3	0.4	0.3	0.3	0.4	0.3	0.5	0.5	0.5	*
2001.....	0.4	0.2	0.3	0.4	0.4	0.5	0.4	0.7	0.6	*
2000.....	0.4	0.3	0.3	0.5	0.5	0.6	0.4	0.5	0.4	*
1999.....	0.4	0.3	0.4	0.4	0.5	0.7	0.6	0.6	0.6	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

All races 3/, both sexes

2004.....	5.1	5.4	4.9	4.7	5.2	5.7	6.6	9.5	11.6	10.8
2003.....	5.2	4.8	4.7	4.7	5.4	5.6	6.5	8.8	11.5	11.9
2002.....	5.3	5.1	4.9	4.6	5.2	5.9	7.3	9.6	11.4	11.4
2001.....	5.4	5.1	5.1	5.3	5.2	5.8	7.2	10.4	12.4	11.7
2000.....	5.7	5.4	5.5	5.6	5.8	6.2	7.3	10.2	11.5	13.1
1999.....	5.4	5.2	5.1	5.3	5.2	6.8	8.1	10.8	13.5	13.9

All races 3/, male

2004.....	7.2	7.5	6.8	6.5	6.9	7.4	8.5	12.6	17.3	17.9
2003.....	7.1	6.8	6.3	6.3	7.0	7.5	8.0	12.1	16.1	20.3
2002.....	6.9	7.0	6.5	6.5	6.5	8.0	9.5	13.0	15.3	21.7
2001.....	7.5	7.2	7.1	6.8	6.9	7.0	9.4	14.3	18.2	21.2
2000.....	8.0	7.4	7.5	7.4	7.6	7.9	9.1	13.6	16.6	24.7
1999.....	7.2	7.2	6.7	7.3	6.7	8.8	9.9	15.0	19.6	27.5

All races 3/, female

2004.....	3.1	3.4	3.1	3.1	3.7	4.3	5.1	7.2	8.1	7.7
2003.....	3.4	2.8	3.2	3.2	3.9	4.1	5.4	6.5	8.7	8.1
2002.....	3.7	3.3	3.4	2.8	3.9	4.1	5.5	7.2	9.0	6.9
2001.....	3.4	3.1	3.2	3.9	3.7	4.8	5.4	7.7	9.0	7.7
2000.....	3.4	3.5	3.6	3.9	4.2	4.8	5.8	7.8	8.6	8.4
1999.....	3.6	3.3	3.6	3.4	3.9	5.2	6.7	7.8	9.9	8.5

White, both sexes

2004.....	5.1	5.3	4.6	4.6	5.2	5.6	6.5	9.5	12.0	11.5
2003.....	5.1	4.5	4.5	4.5	5.2	5.7	6.6	8.9	11.6	12.1
2002.....	5.3	5.0	4.8	4.5	5.1	5.9	7.4	9.8	11.6	11.8
2001.....	5.4	5.0	5.1	5.1	5.1	5.7	7.2	10.6	13.0	12.2
2000.....	5.7	5.2	5.3	5.5	5.9	6.1	7.1	10.3	11.9	13.5
1999.....	5.4	5.0	4.9	5.2	5.0	6.8	8.2	10.8	13.6	14.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

White, male

2004.....	7.1	7.4	6.2	6.3	6.6	7.0	8.4	12.8	17.6	19.0
2003.....	7.0	6.4	6.0	5.9	6.8	7.5	8.1	12.2	16.3	20.4
2002.....	6.9	6.8	6.3	6.2	6.4	7.8	9.7	12.9	15.4	21.9
2001.....	7.4	6.9	7.1	6.4	6.7	6.9	9.3	14.3	18.8	22.3
2000.....	7.8	7.1	7.2	7.1	7.5	7.5	8.7	13.8	16.8	25.8
1999.....	7.2	6.8	6.5	7.2	6.3	8.7	10.1	14.7	19.6	28.3

White, female

2004.....	3.1	3.3	3.0	3.1	3.8	4.4	5.0	7.1	8.4	8.0
2003.....	3.2	2.7	3.1	3.0	3.7	4.0	5.3	6.6	8.7	8.4
2002.....	3.7	3.1	3.4	2.8	3.9	4.2	5.5	7.6	9.3	7.4
2001.....	3.4	3.2	3.2	3.9	3.7	4.7	5.6	8.0	9.4	7.9
2000.....	3.5	3.4	3.6	3.9	4.4	4.8	5.8	7.9	8.9	8.5
1999.....	3.6	3.2	3.5	3.4	3.8	5.1	6.7	8.1	10.1	8.7

Black, both sexes

2004.....	5.7	6.8	7.6	5.5	6.1	6.4	6.0	8.5	7.7	*
2003.....	6.7	6.5	6.2	6.1	7.0	6.0	6.7	6.8	9.9	10.8
2002.....	5.4	6.2	5.8	5.7	5.6	6.4	6.0	8.4	9.7	7.1
2001.....	6.4	6.7	5.6	7.2	6.1	6.9	6.9	10.4	7.0	7.1
2000.....	6.7	7.5	7.3	5.9	5.4	7.4	8.7	7.3	8.5	8.8
1999.....	5.4	7.9	6.7	5.9	6.9	7.2	8.3	11.2	13.3	10.2

Black, male

2004.....	8.5	10.3	12.0	8.0	9.5	11.1	9.7	11.8	15.1	*
2003.....	9.2	10.4	9.3	9.5	9.8	8.9	8.5	11.9	*	20.9
2002.....	7.5	8.5	8.2	9.5	8.1	10.2	9.0	16.1	*	21.4
2001.....	9.0	10.8	8.2	11.1	10.0	8.4	10.1	18.1	*	*
2000.....	10.9	10.7	11.2	10.0	8.3	11.7	14.4	11.3	16.9	*
1999.....	7.7	12.0	10.1	9.0	10.6	10.1	8.8	19.9	24.4	23.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

Black, female

2004.....	3.3	3.7	3.9	3.4	3.4	*	*	6.5	*	*
2003.....	4.4	3.1	3.5	3.3	4.8	3.8	5.5	*	8.5	*
2002.....	3.6	4.2	3.7	2.6	3.6	*	*	*	*	*
2001.....	4.1	3.0	3.4	4.0	*	5.8	4.7	5.8	*	*
2000.....	3.0	4.8	4.0	*	*	4.2	4.9	*	*	*
1999.....	3.4	4.3	3.9	3.5	4.0	5.0	7.9	6.0	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2004.....	0.3	0.3	0.3	0.3	0.3	0.3	0.4	*	0.5	*
2003.....	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.5	0.6	0.5
2002.....	0.2	0.3	0.3	0.2	0.3	0.2	0.4	0.4	0.5	*
2001.....	0.4	0.4	0.4	0.3	0.4	0.6	0.5	0.6	0.6	0.6
2000.....	0.5	0.4	0.4	0.4	0.4	0.5	0.5	0.6	0.7	0.7
1999.....	0.6	0.7	0.6	0.6	0.6	0.8	0.7	1.2	1.2	1.2

All races 3/, male

2004.....	0.4	0.4	0.5	0.5	0.5	0.5	0.6	*	0.9	*
2003.....	0.5	0.6	0.5	0.6	0.6	0.6	0.7	0.7	1.1	*
2002.....	0.3	0.6	0.5	0.4	0.5	0.5	0.6	0.7	1.0	*
2001.....	0.6	0.6	0.6	0.5	0.7	1.0	0.7	0.9	*	*
2000.....	0.7	0.5	0.5	0.5	0.7	0.7	0.8	0.9	1.1	*
1999.....	0.9	1.0	0.9	1.0	0.9	1.1	1.0	1.6	1.8	2.7

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	0.3	*	*	*	*	*	0.5	*	*
2000.....	0.2	0.3	0.3	*	*	*	*	*	*	*
1999.....	0.3	0.4	0.2	0.3	*	0.4	0.4	0.9	0.9	*

White, both sexes

2004.....	0.3	0.3	0.3	0.3	0.4	0.4	0.4	*	0.5	*
2003.....	0.4	0.3	0.3	0.4	0.4	0.3	0.4	0.5	0.6	0.5
2002.....	0.2	0.3	0.3	0.2	0.3	*	0.4	0.3	0.5	*
2001.....	0.3	0.4	0.3	0.3	0.4	0.5	0.5	0.6	0.6	0.5
2000.....	0.4	0.4	0.5	0.3	0.4	0.4	0.4	0.6	0.7	0.7
1999.....	0.6	0.7	0.6	0.7	0.6	0.7	0.6	1.1	1.2	1.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Water transport accidents (V90-V94)

All races 3/, both sexes

2004.....	0.2	0.3	0.3	0.2	0.2	0.3	*	*	*	*
2003.....	0.3	0.2	0.3	0.2	0.3	*	*	*	*	*
2002.....	0.3	0.2	0.3	0.3	0.2	0.4	0.3	*	*	*
2001.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	*	*	*
2000.....	0.3	0.2	0.3	0.3	0.2	0.3	0.2	*	*	*
1999.....	0.3	0.4	0.3	0.3	0.3	0.3	*	*	*	*

All races 3/, male

2004.....	0.4	0.5	0.5	0.4	0.4	0.5	*	*	*	*
2003.....	0.5	0.4	0.6	0.3	0.6	*	*	*	*	*
2002.....	0.5	0.4	0.5	0.5	0.5	0.7	0.6	*	*	*
2001.....	0.5	0.5	0.5	0.5	0.5	0.5	0.6	*	*	*
2000.....	0.6	0.5	0.6	0.5	0.4	0.6	*	*	*	*
1999.....	0.5	0.7	0.5	0.6	0.6	0.6	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	0.3	0.3	0.3	0.2	0.3	*	*	*	*
2003.....	0.3	0.2	0.3	0.2	0.3	*	*	*	*	*
2002.....	0.3	0.2	0.2	0.2	0.3	0.3	0.3	*	*	*
2001.....	0.3	0.3	0.3	0.2	0.3	*	0.3	*	*	*
2000.....	0.4	0.2	0.3	0.3	0.2	0.3	0.3	*	*	*
1999.....	0.3	0.4	0.3	0.3	0.3	0.3	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Air and space transport accidents (V95-V97)

All races 3/, both sexes

2004.....	0.3	0.3	0.5	0.3	0.6	0.4	0.3	*	*	*
2003.....	0.3	0.5	0.5	0.6	0.6	0.4	0.3	0.3	*	*
2002.....	0.3	0.4	0.4	0.5	0.5	0.4	0.4	*	*	*
2001.....	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.3	*	*
2000.....	0.3	0.4	0.5	0.5	0.6	0.5	0.2	*	*	*
1999.....	0.4	0.4	0.4	0.5	0.5	0.4	0.4	*	*	*

All races 3/, male

2004.....	0.4	0.5	0.9	0.5	1.1	0.8	0.6	*	*	*
2003.....	0.6	0.8	0.9	1.1	1.1	0.8	0.7	*	*	*
2002.....	0.5	0.8	0.7	0.9	0.9	0.8	0.8	*	*	*
2001.....	0.8	0.7	0.9	0.8	0.9	0.9	0.7	*	*	*
2000.....	0.6	0.7	0.9	1.0	0.9	0.9	*	*	*	*
1999.....	0.5	0.7	0.7	1.0	0.9	0.6	0.8	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	0.3	0.2	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.3	0.4	0.6	0.4	0.6	0.5	0.4	*	*	*
2003.....	0.4	0.5	0.6	0.7	0.6	0.4	0.4	0.3	*	*
2002.....	0.4	0.5	0.5	0.6	0.6	0.4	0.4	*	*	*
2001.....	0.6	0.6	0.6	0.6	0.6	0.6	0.4	0.3	*	*
2000.....	0.4	0.5	0.6	0.6	0.6	0.5	0.3	*	*	*
1999.....	0.4	0.5	0.5	0.6	0.5	0.4	0.4	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified transport accidents and their sequelae (V98-V99,Y85)

All races 3/, both sexes

2004.....	0.4	0.4	0.3	0.3	0.4	0.3	0.3	*	*	*
2003.....	0.3	0.4	0.3	0.4	0.4	0.3	0.4	0.3	*	*
2002.....	0.3	0.5	0.3	0.3	0.4	0.3	*	0.4	0.5	*
2001.....	0.4	0.3	0.3	0.2	0.3	0.2	0.3	0.5	0.4	*
2000.....	0.3	0.2	0.2	0.2	0.3	0.3	0.3	*	*	*
1999.....	0.2	0.2	0.3	0.3	0.3	*	0.3	*	*	*

All races 3/, male

2004.....	0.5	0.7	0.5	0.5	0.6	0.5	*	*	*	*
2003.....	0.5	0.5	0.5	0.7	0.6	0.5	0.6	*	*	*
2002.....	0.4	0.7	0.4	0.5	0.6	0.5	*	*	*	*
2001.....	0.6	0.6	0.5	0.4	0.5	*	*	0.7	*	*
2000.....	0.4	0.4	0.4	*	0.6	0.5	0.5	*	*	*
1999.....	0.4	0.3	0.5	0.5	0.4	*	*	*	*	*

All races 3/, female

2004.....	0.2	*	*	*	*	*	*	*	*	*
2003.....	0.2	0.2	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.4	0.4	0.3	0.3	0.4	0.3	0.3	*	*	*
2003.....	0.3	0.4	0.3	0.4	0.3	0.3	0.4	*	*	*
2002.....	0.3	0.5	0.3	0.3	0.3	0.2	*	0.3	0.5	*
2001.....	0.4	0.3	0.3	0.2	0.3	*	0.3	0.4	*	*
2000.....	0.4	0.2	0.2	0.2	0.3	0.4	*	*	*	*
1999.....	0.3	0.2	0.3	0.2	0.3	*	0.3	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Falls (W00-W19)

All races 3/, both sexes

2004.....	1.8	2.6	3.2	4.1	5.8	8.9	16.1	30.3	61.8	143.3
2003.....	1.7	2.3	2.8	3.7	5.3	8.1	14.7	28.7	57.5	135.9
2002.....	1.8	2.3	2.8	3.4	4.9	7.9	13.9	27.1	54.0	130.4
2001.....	1.6	2.5	2.7	3.3	4.9	7.8	12.4	25.2	49.9	121.5
2000.....	1.6	2.0	2.6	3.3	4.6	6.2	12.0	22.1	44.5	112.6
1999.....	1.7	2.3	2.2	3.0	4.6	6.9	11.3	22.3	45.5	110.2

All races 3/, male

2004.....	2.9	4.1	4.9	6.2	8.1	12.4	21.6	38.4	78.7	174.4
2003.....	2.7	3.6	4.4	5.3	7.3	10.7	18.9	36.9	73.8	165.1
2002.....	2.9	3.6	4.3	5.2	7.1	10.7	18.9	35.1	67.9	158.9
2001.....	2.8	4.2	4.4	5.0	7.1	10.6	16.0	34.0	64.1	155.7
2000.....	2.7	3.4	4.2	5.0	6.9	8.6	15.7	29.4	57.3	145.4
1999.....	2.8	3.8	3.6	4.7	7.0	9.8	15.3	27.9	61.6	142.5

All races 3/, female

2004.....	0.7	1.1	1.5	2.0	3.6	5.8	11.7	24.5	51.4	129.2
2003.....	0.6	1.1	1.3	2.1	3.5	5.8	11.3	22.8	47.5	122.9
2002.....	0.7	1.0	1.4	1.8	3.0	5.5	9.9	21.4	45.5	118.1
2001.....	0.6	0.9	1.2	1.6	2.9	5.4	9.5	19.0	41.3	107.2
2000.....	0.6	0.8	1.2	1.8	2.6	4.2	9.1	17.1	36.9	99.2
1999.....	0.6	0.8	0.9	1.5	2.4	4.5	8.1	18.4	36.0	97.2

White, both sexes

2004.....	1.9	2.6	3.2	4.0	5.8	9.2	16.7	32.0	65.3	152.6
2003.....	1.7	2.3	2.7	3.7	5.4	8.2	15.2	30.1	60.0	144.4
2002.....	1.8	2.3	2.8	3.4	5.1	8.1	14.4	28.3	57.1	138.0
2001.....	1.7	2.5	2.8	3.2	4.9	8.0	12.8	26.3	52.4	128.0
2000.....	1.7	2.1	2.5	3.2	4.6	6.2	12.3	23.1	46.6	119.1
1999.....	1.7	2.2	2.1	2.9	4.3	6.9	11.4	23.0	47.7	115.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Falls (W00-W19)

White, male

2004.....	3.1	4.2	4.8	6.1	7.9	13.0	22.0	40.4	82.7	183.7
2003.....	2.8	3.6	4.1	5.3	7.3	10.6	19.5	38.6	76.4	175.1
2002.....	3.0	3.7	4.2	5.1	7.2	10.8	19.3	36.4	71.2	166.4
2001.....	2.8	4.1	4.4	4.9	7.0	10.8	16.5	35.1	66.6	162.3
2000.....	2.7	3.5	3.9	4.8	6.9	8.4	16.2	30.3	59.7	152.8
1999.....	2.7	3.7	3.3	4.4	6.6	9.6	15.3	28.3	64.2	149.1

White, female

2004.....	0.7	1.1	1.6	2.1	3.8	5.8	12.3	25.9	54.6	138.6
2003.....	0.6	1.0	1.3	2.1	3.6	6.1	11.6	24.0	49.9	130.8
2002.....	0.7	1.0	1.5	1.8	3.1	5.8	10.3	22.6	48.5	125.6
2001.....	0.5	0.9	1.2	1.6	2.9	5.6	9.9	20.0	43.9	113.6
2000.....	0.6	0.7	1.2	1.8	2.6	4.3	9.3	18.1	38.8	105.4
1999.....	0.6	0.8	0.9	1.5	2.2	4.6	8.3	19.3	37.9	102.6

Black, both sexes

2004.....	1.7	2.6	3.5	5.1	6.3	7.9	12.8	16.8	25.5	46.5
2003.....	1.5	3.0	3.6	4.3	5.5	7.6	10.6	13.4	28.2	44.1
2002.....	1.6	2.5	3.2	3.8	4.7	6.1	10.9	14.8	18.1	49.3
2001.....	2.0	2.6	3.3	3.9	4.9	7.9	9.7	16.3	21.9	44.5
2000.....	1.6	2.0	3.6	4.5	5.1	7.1	9.7	11.6	20.3	38.8
1999.....	1.7	3.2	3.6	4.5	6.7	9.0	9.5	13.4	23.1	45.3

Black, male

2004.....	2.6	4.1	6.0	8.8	11.8	9.7	20.7	23.7	38.0	71.5
2003.....	2.6	4.8	5.8	6.5	8.1	12.8	12.2	18.3	35.4	56.3
2002.....	2.6	4.1	5.6	6.3	7.3	9.2	15.4	22.0	28.3	77.2
2001.....	3.3	4.9	6.0	6.4	7.7	11.9	13.5	24.2	36.1	68.1
2000.....	2.6	3.4	6.3	7.7	8.7	10.7	12.4	18.4	26.1	61.1
1999.....	3.3	5.6	6.0	7.6	10.2	14.8	14.6	20.4	31.4	58.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Falls (W00-W19)

Black, female

2004.....	*	*	*	*	*	6.7	7.5	12.7	18.9	36.6
2003.....	*	1.4	1.7	2.5	3.5	3.7	9.5	10.6	24.4	39.3
2002.....	*	*	*	*	*	3.7	7.8	10.4	12.7	38.6
2001.....	*	*	*	*	*	4.9	7.2	11.6	14.7	35.8
2000.....	*	*	*	*	*	4.4	8.0	7.4	17.4	30.5
1999.....	*	*	*	*	4.0	4.6	6.1	9.2	19.0	40.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Fall on same level (W00-W09,W18)

All races 3/, both sexes

2004.....	0.3	0.6	0.7	1.0	1.8	2.7	5.1	10.2	21.7	53.6
2003.....	0.4	0.4	0.8	1.0	1.5	2.3	4.4	9.5	20.5	50.3
2002.....	0.3	0.5	0.7	0.9	1.3	2.2	4.2	8.9	17.8	50.8
2001.....	0.2	0.5	0.6	0.8	1.2	1.9	3.3	6.8	14.4	37.3
2000.....	0.2	0.3	0.5	0.6	1.0	1.4	2.7	5.5	11.9	33.0
1999.....	0.1	0.3	0.3	0.4	0.5	1.1	1.9	4.2	8.2	21.4

All races 3/, male

2004.....	0.5	0.9	1.2	1.3	2.4	3.6	6.3	12.9	26.8	64.1
2003.....	0.5	0.6	1.1	1.4	1.9	2.7	5.5	12.2	26.8	60.1
2002.....	0.5	0.8	1.0	1.2	1.7	2.9	5.7	11.0	21.6	60.8
2001.....	0.3	0.8	0.8	1.0	1.7	2.4	3.7	9.0	18.5	45.3
2000.....	0.4	0.4	0.8	0.8	1.2	1.7	3.4	6.6	15.2	39.6
1999.....	0.2	0.5	0.5	0.5	0.6	1.5	2.5	5.3	10.5	26.6

All races 3/, female

2004.....	0.2	0.3	0.3	0.6	1.2	1.8	4.1	8.2	18.5	48.9
2003.....	0.2	0.3	0.4	0.7	1.1	2.0	3.6	7.6	16.7	45.9
2002.....	*	0.2	0.4	0.7	0.9	1.6	2.9	7.3	15.5	46.4
2001.....	*	0.3	0.4	0.5	0.8	1.4	2.9	5.2	11.9	33.9
2000.....	*	*	0.3	0.5	0.8	1.1	2.2	4.7	10.0	30.4
1999.....	*	*	*	*	0.4	0.8	1.4	3.5	6.8	19.4

White, both sexes

2004.....	0.3	0.6	0.8	0.9	1.7	2.8	5.2	10.5	22.9	57.5
2003.....	0.4	0.4	0.7	1.0	1.4	2.4	4.6	10.2	21.6	53.5
2002.....	0.3	0.5	0.7	0.9	1.4	2.3	4.3	9.2	18.8	53.7
2001.....	0.2	0.6	0.6	0.7	1.3	2.0	3.4	6.9	15.1	39.3
2000.....	0.3	0.3	0.5	0.6	1.0	1.4	2.8	5.8	12.5	35.0
1999.....	0.2	0.3	0.3	0.3	0.4	1.1	2.0	4.3	8.4	22.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Fall from one level to another (W10-W17)

All races 3/, both sexes

2004.....	0.8	1.0	1.1	1.4	1.8	2.2	3.4	4.7	7.3	10.6
2003.....	0.7	1.1	1.0	1.1	1.7	2.3	3.2	4.6	7.6	11.4
2002.....	0.8	1.0	0.9	1.3	1.7	2.3	3.4	5.1	7.5	10.9
2001.....	0.8	1.1	1.1	1.3	1.5	2.3	3.0	5.1	7.6	11.3
2000.....	0.9	1.0	1.0	1.2	1.5	1.9	3.0	4.2	6.5	8.8
1999.....	0.9	1.0	0.9	1.3	1.9	2.3	2.9	4.8	7.3	10.4

All races 3/, male

2004.....	1.5	1.7	2.0	2.4	3.0	3.6	5.5	6.7	11.2	16.4
2003.....	1.3	1.8	1.7	1.9	2.7	3.6	5.1	7.3	11.3	18.1
2002.....	1.5	1.7	1.5	2.2	2.8	3.5	5.3	8.2	12.0	17.4
2001.....	1.4	2.0	1.9	2.2	2.6	3.9	4.6	8.4	11.4	18.2
2000.....	1.5	1.7	1.8	2.1	2.4	2.9	4.9	7.0	10.8	16.4
1999.....	1.5	1.7	1.5	2.1	3.1	4.0	4.2	6.9	12.7	17.8

All races 3/, female

2004.....	0.2	0.3	0.3	0.5	0.7	1.0	1.7	3.2	5.0	7.9
2003.....	*	0.3	0.3	0.4	0.8	1.2	1.6	2.8	5.2	8.5
2002.....	0.2	0.2	0.4	0.4	0.7	1.2	1.9	3.0	4.8	8.1
2001.....	*	0.2	0.3	0.4	0.5	1.0	1.8	2.8	5.3	8.3
2000.....	0.2	0.3	0.3	0.5	0.6	1.0	1.5	2.2	3.9	5.7
1999.....	0.2	0.2	0.2	0.5	0.7	0.9	1.8	3.4	4.2	7.4

White, both sexes

2004.....	0.9	1.0	1.2	1.5	1.9	2.4	3.6	5.0	7.7	11.2
2003.....	0.7	1.0	1.0	1.2	1.8	2.3	3.3	4.9	7.9	12.1
2002.....	0.9	1.0	1.0	1.3	1.8	2.4	3.6	5.4	7.9	11.2
2001.....	0.8	1.1	1.1	1.3	1.5	2.4	3.2	5.3	8.1	11.7
2000.....	0.9	1.0	1.0	1.3	1.5	1.8	3.1	4.4	6.8	9.2
1999.....	0.9	1.0	0.8	1.3	2.0	2.4	3.0	5.0	7.7	10.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Unspecified fall (W19)

All races 3/, both sexes

2004.....	0.7	1.0	1.3	1.7	2.2	4.0	7.6	15.5	32.8	79.1
2003.....	0.6	0.8	1.1	1.5	2.1	3.5	7.1	14.5	29.4	74.2
2002.....	0.6	0.8	1.2	1.3	2.0	3.5	6.3	13.1	28.6	68.8
2001.....	0.6	0.9	1.1	1.3	2.1	3.6	6.1	13.4	27.9	73.0
2000.....	0.5	0.8	1.1	1.4	2.2	3.0	6.3	12.5	26.1	70.7
1999.....	0.7	1.0	1.0	1.4	2.2	3.5	6.5	13.2	30.0	78.4

All races 3/, male

2004.....	1.0	1.5	1.7	2.5	2.8	5.2	9.8	18.8	40.7	93.9
2003.....	0.9	1.2	1.6	2.0	2.7	4.5	8.3	17.5	35.6	86.9
2002.....	0.9	1.1	1.8	1.9	2.6	4.3	7.8	15.9	34.3	80.7
2001.....	1.0	1.3	1.7	1.8	2.8	4.3	7.6	16.6	34.2	92.1
2000.....	0.8	1.2	1.6	2.1	3.3	4.0	7.5	15.9	31.3	89.4
1999.....	1.0	1.6	1.5	2.1	3.2	4.3	8.5	15.7	38.4	98.1

All races 3/, female

2004.....	0.3	0.5	0.9	0.9	1.6	3.0	5.9	13.1	27.9	72.4
2003.....	0.3	0.4	0.6	1.0	1.6	2.6	6.1	12.3	25.6	68.5
2002.....	0.3	0.6	0.6	0.7	1.4	2.8	5.1	11.2	25.2	63.6
2001.....	0.3	0.4	0.5	0.7	1.5	3.0	4.8	11.1	24.2	64.9
2000.....	0.3	0.3	0.6	0.8	1.2	2.2	5.5	10.1	23.1	63.1
1999.....	0.3	0.4	0.5	0.8	1.3	2.7	4.9	11.5	25.0	70.4

White, both sexes

2004.....	0.7	1.0	1.3	1.6	2.2	4.0	7.9	16.5	34.7	83.9
2003.....	0.6	0.8	1.0	1.5	2.2	3.5	7.3	15.1	30.5	78.8
2002.....	0.6	0.8	1.1	1.2	1.9	3.4	6.4	13.7	30.4	73.0
2001.....	0.6	0.9	1.1	1.2	2.1	3.6	6.3	14.0	29.2	77.0
2000.....	0.5	0.8	1.0	1.4	2.1	3.0	6.4	12.9	27.3	74.9
1999.....	0.6	0.9	1.0	1.3	1.9	3.4	6.4	13.7	31.6	82.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Unspecified fall (W19)

White, male

2004.....	1.0	1.5	1.6	2.4	2.7	5.2	10.0	19.9	42.9	98.0
2003.....	0.9	1.2	1.5	2.0	2.7	4.5	8.5	18.3	36.4	92.2
2002.....	0.9	1.1	1.6	1.8	2.3	4.1	7.9	16.6	36.1	85.1
2001.....	1.0	1.3	1.7	1.7	2.8	4.2	7.9	17.3	35.3	95.7
2000.....	0.7	1.3	1.5	1.9	3.1	3.9	7.6	16.2	32.6	94.0
1999.....	1.0	1.5	1.4	1.8	2.8	4.1	8.4	15.9	40.3	101.9

White, female

2004.....	0.3	0.5	0.9	0.9	1.7	3.0	6.2	14.0	29.7	77.6
2003.....	0.3	0.5	0.5	1.0	1.7	2.6	6.3	12.7	26.9	72.8
2002.....	0.3	0.6	0.6	0.7	1.4	2.9	5.3	11.7	27.0	67.8
2001.....	0.2	0.4	0.5	0.7	1.5	3.0	5.0	11.7	25.5	69.2
2000.....	0.3	0.3	0.6	0.8	1.2	2.2	5.5	10.5	24.2	67.2
1999.....	0.3	0.3	0.5	0.7	1.1	2.8	4.8	12.1	26.4	74.4

Black, both sexes

2004.....	*	1.0	1.6	2.4	2.5	5.0	6.3	6.6	13.9	30.2
2003.....	*	1.2	1.7	2.0	2.5	3.9	6.5	8.3	17.5	28.6
2002.....	*	1.1	2.0	1.8	3.0	3.6	5.6	7.6	10.5	25.2
2001.....	0.8	1.0	1.2	1.8	2.5	4.8	4.7	8.9	13.8	26.4
2000.....	0.9	0.9	1.5	2.1	2.8	3.8	6.3	8.0	13.3	25.7
1999.....	0.8	1.9	1.6	2.6	4.5	4.9	6.4	8.6	15.0	33.5

Black, male

2004.....	*	*	2.5	4.4	4.7	6.5	10.0	9.6	20.8	48.7
2003.....	*	2.0	2.5	2.9	*	5.2	7.9	*	23.6	36.5
2002.....	*	1.8	3.7	3.0	5.1	5.2	8.0	*	*	38.6
2001.....	*	1.9	*	*	*	6.1	*	12.5	20.9	46.5
2000.....	*	*	2.7	3.6	5.2	5.2	7.4	11.8	18.5	38.1
1999.....	*	3.2	3.0	4.4	6.8	7.5	9.2	12.8	19.2	46.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Unspecified fall (W19)

Black, female

2004.....	*	*	*	*	*	3.8	*	*	10.2	22.8
2003.....	*	*	*	*	*	*	5.5	8.7	14.3	25.5
2002.....	*	*	*	*	*	*	*	7.1	8.4	20.1
2001.....	*	*	*	*	*	3.8	*	6.7	10.2	18.9
2000.....	*	*	*	*	*	*	5.5	5.7	10.6	21.0
1999.....	*	*	*	*	*	*	4.5	6.0	13.0	28.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental discharge of firearms (W32-W34)										
All races 3/, both sexes										
2004.....	0.2	0.2	0.3	0.2	*	0.2	0.2	*	*	*
2003.....	0.3	0.2	0.2	0.2	0.2	0.2	*	*	0.4	*
2002.....	0.3	0.2	0.2	0.2	0.2	*	*	*	*	*
2001.....	0.3	0.3	0.2	0.3	0.2	*	*	0.3	*	*
2000.....	0.3	0.2	0.2	0.3	*	0.3	*	*	*	*
1999.....	0.3	0.3	0.2	0.2	0.2	*	*	*	*	*
All races 3/, male										
2004.....	0.4	0.3	0.6	0.2	*	*	*	*	*	*
2003.....	0.5	0.4	0.4	0.5	0.4	0.4	*	*	*	*
2002.....	0.5	0.4	0.4	0.4	*	*	*	*	*	*
2001.....	0.6	0.5	0.4	0.5	0.4	*	*	0.7	*	*
2000.....	0.6	0.3	0.4	0.5	*	0.5	*	*	*	*
1999.....	0.5	0.5	0.4	0.3	*	*	*	*	*	*
All races 3/, female										
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2004.....	0.2	0.2	0.4	0.2	*	*	*	*	*	*
2003.....	0.3	0.2	0.3	0.3	0.2	0.3	*	*	*	*
2002.....	0.3	0.2	0.2	0.2	*	*	*	*	*	*
2001.....	0.3	0.3	0.3	0.3	0.3	*	*	0.3	*	*
2000.....	0.4	0.2	0.2	0.3	*	0.3	*	*	*	*
1999.....	0.3	0.3	0.3	0.2	0.2	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental drowning and submersion (W65-W74)

All races 3/, both sexes

2004.....	1.0	1.1	1.0	0.9	0.9	1.1	1.0	1.0	1.5	1.6
2003.....	1.1	1.0	1.0	1.0	0.9	1.0	1.2	1.3	1.3	1.8
2002.....	1.1	1.0	0.9	0.8	0.8	0.8	1.0	1.2	1.5	1.7
2001.....	1.1	1.0	0.8	0.8	0.9	0.8	0.9	1.1	1.4	1.7
2000.....	1.0	0.9	1.0	0.9	0.9	0.8	1.1	1.0	1.7	1.5
1999.....	1.0	1.0	1.0	0.9	0.9	0.9	1.1	1.2	1.4	1.6

All races 3/, male

2004.....	1.7	1.8	1.7	1.5	1.5	1.7	1.6	1.7	2.8	2.9
2003.....	1.8	1.6	1.6	1.6	1.5	1.6	1.9	1.8	2.3	3.3
2002.....	1.9	1.6	1.5	1.3	1.2	1.4	1.8	1.8	2.5	3.6
2001.....	1.7	1.8	1.3	1.2	1.4	1.4	1.6	1.8	2.4	3.5
2000.....	1.7	1.6	1.7	1.6	1.5	1.4	1.8	1.2	2.7	2.8
1999.....	1.7	1.7	1.8	1.5	1.5	1.5	1.7	2.1	2.7	2.3

All races 3/, female

2004.....	0.4	0.4	0.5	0.4	0.5	0.5	0.5	0.5	0.7	1.0
2003.....	0.3	0.3	0.4	0.3	0.4	0.6	0.6	0.8	0.6	1.1
2002.....	0.4	0.4	0.3	0.3	0.4	*	0.4	0.7	0.9	0.9
2001.....	0.4	0.3	0.3	0.4	0.4	*	*	0.7	0.8	1.0
2000.....	0.3	0.3	0.3	*	0.4	*	0.6	0.8	1.1	1.0
1999.....	0.3	0.3	0.3	0.4	0.4	0.4	0.7	0.6	0.7	1.4

White, both sexes

2004.....	1.0	1.1	1.0	0.9	0.9	1.1	0.9	1.0	1.4	1.5
2003.....	1.0	1.0	1.0	1.0	0.9	1.1	1.1	1.2	1.3	1.6
2002.....	1.2	0.9	0.8	0.8	0.8	0.8	0.9	1.1	1.4	1.6
2001.....	1.0	1.0	0.8	0.7	0.9	0.8	0.9	1.1	1.3	1.7
2000.....	1.0	0.9	0.9	0.9	0.9	0.8	1.1	0.9	1.8	1.5
1999.....	1.0	1.1	1.0	0.9	0.9	0.9	1.0	1.1	1.4	1.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

All races 3/, both sexes

2004.....	0.5	0.7	0.9	1.1	1.6	2.1	3.9	7.1	13.5	29.9
2003.....	0.5	0.7	0.9	1.2	1.7	2.4	3.9	6.6	13.0	27.4
2002.....	0.4	0.5	0.8	1.1	1.3	2.6	3.6	7.3	12.6	28.4
2001.....	0.5	0.8	1.0	1.2	1.5	2.3	3.6	7.2	12.8	30.0
2000.....	0.4	0.7	0.9	1.2	1.6	2.3	4.0	7.7	13.4	32.6
1999.....	0.5	0.5	0.8	1.0	1.4	2.2	4.0	6.6	12.1	30.3

All races 3/, male

2004.....	0.6	0.9	1.0	1.4	1.9	2.6	4.7	9.5	17.0	39.0
2003.....	0.6	0.8	0.9	1.4	2.3	2.8	4.6	9.0	17.9	37.5
2002.....	0.6	0.7	0.9	1.5	1.7	3.4	4.4	9.6	16.0	37.0
2001.....	0.6	1.0	1.2	1.6	1.6	2.9	4.3	9.2	17.5	39.8
2000.....	0.5	1.0	1.2	1.4	1.9	2.8	4.5	9.7	18.9	41.0
1999.....	0.6	0.6	1.2	1.1	1.6	2.5	5.2	7.9	15.9	39.2

All races 3/, female

2004.....	0.4	0.5	0.8	0.8	1.3	1.6	3.2	5.3	11.3	25.8
2003.....	0.4	0.6	0.8	0.9	1.1	2.0	3.3	5.0	10.1	23.0
2002.....	0.3	0.4	0.6	0.8	1.0	2.0	2.9	5.6	10.6	24.6
2001.....	0.3	0.6	0.8	0.8	1.3	1.8	3.1	5.8	9.9	26.0
2000.....	0.4	0.4	0.6	0.9	1.4	1.8	3.5	6.3	10.1	29.1
1999.....	0.3	0.5	0.4	0.8	1.2	2.0	3.0	5.6	9.9	26.7

White, both sexes

2004.....	0.5	0.7	0.9	1.1	1.5	1.9	3.6	6.9	13.4	30.1
2003.....	0.5	0.7	0.8	1.1	1.6	2.3	3.7	6.6	12.9	27.9
2002.....	0.4	0.5	0.7	1.1	1.2	2.3	3.4	7.1	12.5	28.3
2001.....	0.4	0.7	1.0	1.1	1.4	2.3	3.3	6.8	12.4	29.7
2000.....	0.5	0.7	0.8	1.1	1.5	2.1	3.7	7.6	13.3	32.5
1999.....	0.5	0.5	0.8	0.9	1.2	2.1	3.7	6.2	11.7	30.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

White, male										
2004.....	0.6	0.9	1.0	1.4	1.8	2.4	4.4	9.5	16.8	39.2
2003.....	0.5	0.8	0.9	1.3	2.1	2.8	4.4	8.9	17.5	38.2
2002.....	0.6	0.7	0.8	1.4	1.6	2.9	4.3	9.4	15.6	37.2
2001.....	0.6	0.9	1.1	1.3	1.5	2.8	3.9	8.6	16.4	39.3
2000.....	0.6	0.9	1.1	1.4	1.8	2.5	4.2	9.5	18.4	40.2
1999.....	0.6	0.6	1.1	1.0	1.3	2.4	4.9	7.3	15.4	39.7
White, female										
2004.....	0.5	0.6	0.9	0.8	1.2	1.5	3.0	5.0	11.3	26.0
2003.....	0.4	0.6	0.7	0.9	1.1	1.8	3.1	5.0	10.0	23.3
2002.....	0.3	0.4	0.6	0.8	0.9	1.8	2.7	5.6	10.6	24.5
2001.....	0.3	0.6	0.9	0.8	1.2	1.8	2.9	5.6	10.0	25.7
2000.....	0.4	0.5	0.6	0.9	1.3	1.7	3.3	6.3	10.2	29.4
1999.....	0.3	0.5	0.4	0.7	1.0	1.9	2.7	5.3	9.5	26.7
Black, both sexes										
2004.....	*	*	1.3	1.5	2.4	4.0	6.7	10.3	17.3	30.5
2003.....	0.8	*	1.4	1.5	2.9	3.7	6.5	6.6	16.5	24.5
2002.....	*	*	1.4	1.9	2.2	5.8	7.1	11.0	16.0	33.0
2001.....	*	1.3	1.0	2.3	2.8	3.6	7.5	11.5	18.7	36.8
2000.....	*	0.8	1.5	1.9	2.6	5.0	7.2	10.8	16.7	35.3
1999.....	*	1.0	1.2	2.1	3.0	3.6	7.7	12.3	18.2	33.2
Black, male										
2004.....	*	*	*	*	*	5.6	8.7	11.4	25.8	42.7
2003.....	*	*	*	*	4.4	*	7.9	10.5	25.8	30.2
2002.....	*	*	2.2	*	*	8.7	8.0	16.5	23.7	41.8
2001.....	*	*	*	3.6	*	*	9.8	15.8	36.1	51.1
2000.....	*	*	2.4	*	*	6.5	8.4	15.6	27.8	51.9
1999.....	*	*	*	*	*	*	10.5	17.6	25.3	43.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

Black, female

2004.....	*	*	*	*	*	*	5.3	9.7	12.9	25.6
2003.....	*	*	*	*	*	3.7	5.5	*	11.6	22.3
2002.....	*	*	*	*	*	3.7	6.5	7.7	11.9	29.6
2001.....	*	*	*	*	*	*	6.1	8.9	9.8	31.6
2000.....	*	*	*	*	*	3.8	6.4	8.0	11.0	29.2
1999.....	*	*	*	*	*	*	5.9	9.2	14.7	29.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
 race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

All races 3/, both sexes

2004.....	0.3	0.3	0.3	0.2	0.2	0.2	*	0.3	*	0.6
2003.....	0.3	0.3	0.3	0.2	0.2	*	0.3	*	0.4	0.8
2002.....	0.3	0.3	0.3	0.3	0.2	0.3	0.2	0.4	*	1.0
2001.....	0.4	0.4	0.2	0.2	0.2	0.2	0.4	0.4	0.4	0.9
2000.....	0.3	0.2	0.3	0.4	0.3	0.3	0.4	0.5	0.5	1.2
1999.....	0.4	0.4	0.3	0.3	0.3	0.5	0.8	1.4	1.8	4.7

All races 3/, male

2004.....	0.6	0.5	0.5	0.3	0.3	*	*	*	*	*
2003.....	0.5	0.4	0.5	0.4	0.3	*	*	*	*	*
2002.....	0.5	0.5	0.5	0.4	*	*	*	*	*	*
2001.....	0.6	0.6	0.4	0.3	*	*	*	*	*	*
2000.....	0.6	0.4	0.6	0.6	0.5	*	*	*	*	*
1999.....	0.6	0.6	0.6	0.6	*	0.7	1.1	1.9	1.8	5.4

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	0.6
2003.....	*	*	*	*	*	*	*	*	*	0.7
2002.....	*	*	*	*	*	*	*	*	*	1.1
2001.....	*	*	*	*	*	*	*	*	*	0.9
2000.....	*	*	*	*	*	*	*	*	*	1.1
1999.....	*	*	*	*	*	*	0.6	1.1	1.7	4.5

White, both sexes

2004.....	0.4	0.4	0.4	0.2	0.2	0.2	*	*	*	0.6
2003.....	0.4	0.3	0.3	0.2	0.2	*	0.3	*	0.4	0.8
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	*	0.4	*	1.0
2001.....	0.4	0.4	0.3	0.2	*	0.3	0.4	0.5	0.4	0.9
2000.....	0.4	0.3	0.4	0.4	0.3	0.3	0.3	0.5	0.6	1.2
1999.....	0.4	0.4	0.4	0.3	0.3	0.5	0.8	1.3	1.8	4.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to electric current (W85-W87)										
All races 3/, both sexes										
2004.....	0.2	0.2	0.2	0.1	*	*	*	*	*	*
2003.....	0.2	0.2	0.1	0.2	*	*	*	*	*	*
2002.....	0.3	0.2	0.2	*	*	*	*	*	*	*
2001.....	0.3	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	0.2	0.1	*	*	*	*	*	*	*
1999.....	0.3	0.2	*	0.2	*	*	*	*	*	*
All races 3/, male										
2004.....	0.3	0.4	0.3	*	*	*	*	*	*	*
2003.....	0.4	0.4	*	0.3	*	*	*	*	*	*
2002.....	0.5	0.4	0.3	*	*	*	*	*	*	*
2001.....	0.5	0.4	*	*	*	*	*	*	*	*
2000.....	0.4	0.5	0.2	*	*	*	*	*	*	*
1999.....	0.5	0.4	*	0.4	*	*	*	*	*	*
All races 3/, female										
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2004.....	0.2	0.2	0.2	*	*	*	*	*	*	*
2003.....	0.2	0.2	0.1	0.2	*	*	*	*	*	*
2002.....	0.3	0.2	0.2	*	*	*	*	*	*	*
2001.....	0.3	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	0.2	*	*	*	*	*	*	*	*
1999.....	0.3	0.2	*	0.2	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental exposure to smoke, fire and flames (X00-X09)

All races 3/, both sexes

2004.....	0.7	1.1	1.2	1.2	1.6	1.8	2.0	2.9	4.4	5.6
2003.....	1.0	1.2	1.3	1.3	1.6	2.0	2.6	3.0	4.3	5.4
2002.....	0.9	1.0	1.3	1.2	1.4	1.8	2.4	2.8	4.2	4.8
2001.....	1.1	1.0	1.1	1.5	1.5	2.0	2.5	3.0	3.8	5.7
2000.....	1.0	1.1	1.3	1.4	1.7	2.0	2.4	3.5	4.3	6.2
1999.....	1.0	1.0	1.0	1.2	1.5	2.0	2.5	3.5	4.7	5.9

All races 3/, male

2004.....	1.0	1.5	1.8	1.6	2.1	2.6	2.3	3.5	5.9	7.2
2003.....	1.2	1.7	1.6	1.8	2.0	2.5	3.4	4.0	5.1	7.4
2002.....	1.3	1.4	1.6	1.5	2.0	2.6	3.2	3.5	6.1	7.0
2001.....	1.6	1.5	1.5	2.1	1.8	2.5	3.0	3.5	5.1	8.3
2000.....	1.2	1.5	1.8	1.9	2.2	2.4	3.2	4.5	5.9	9.4
1999.....	1.5	1.5	1.3	1.6	1.9	2.5	3.4	4.3	6.6	9.3

All races 3/, female

2004.....	0.5	0.8	0.7	0.9	1.3	1.2	1.6	2.5	3.5	4.9
2003.....	0.7	0.8	1.1	0.8	1.1	1.6	1.9	2.3	3.9	4.5
2002.....	0.6	0.6	0.9	0.9	0.8	1.1	1.8	2.4	3.0	3.9
2001.....	0.7	0.6	0.7	1.0	1.2	1.5	2.1	2.7	3.1	4.6
2000.....	0.7	0.7	0.7	0.9	1.2	1.6	1.8	2.8	3.4	4.9
1999.....	0.5	0.5	0.7	0.9	1.2	1.6	1.8	3.0	3.6	4.5

White, both sexes

2004.....	0.7	1.0	1.1	1.1	1.4	1.6	1.6	2.5	3.9	4.9
2003.....	1.0	1.2	1.2	1.2	1.4	1.8	2.2	2.6	3.7	4.7
2002.....	0.9	0.9	1.1	1.1	1.1	1.6	2.1	2.5	3.7	4.0
2001.....	1.0	0.9	1.0	1.3	1.3	1.7	2.1	2.6	3.2	4.8
2000.....	0.9	0.9	1.2	1.1	1.4	1.7	2.0	2.8	3.5	5.1
1999.....	0.9	0.9	0.9	1.1	1.2	1.6	2.0	2.8	3.9	4.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental exposure to smoke, fire and flames (X00-X09)

White, male

2004.....	0.9	1.2	1.6	1.3	1.9	2.2	1.9	2.9	5.4	6.5
2003.....	1.3	1.5	1.5	1.6	1.9	2.2	3.0	3.2	4.2	6.7
2002.....	1.3	1.3	1.4	1.4	1.6	2.2	2.7	3.1	5.8	5.8
2001.....	1.4	1.2	1.2	1.8	1.5	2.1	2.4	2.8	4.4	7.0
2000.....	1.1	1.3	1.7	1.6	1.8	2.0	2.7	3.6	4.8	8.2
1999.....	1.4	1.3	1.2	1.4	1.7	2.0	2.6	3.3	5.8	8.0

White, female

2004.....	0.4	0.7	0.7	0.9	1.0	1.1	1.3	2.2	2.9	4.2
2003.....	0.7	0.8	0.9	0.8	1.0	1.5	1.6	2.2	3.4	3.8
2002.....	0.6	0.6	0.9	0.8	0.6	1.0	1.6	2.1	2.5	3.2
2001.....	0.6	0.5	0.7	0.8	1.1	1.3	1.9	2.4	2.5	3.8
2000.....	0.6	0.6	0.7	0.7	1.1	1.4	1.5	2.3	2.7	3.9
1999.....	0.5	0.5	0.6	0.9	0.8	1.3	1.5	2.5	2.8	3.3

Black, both sexes

2004.....	1.4	2.6	2.2	2.8	4.2	4.0	5.9	8.0	12.1	14.9
2003.....	1.4	2.3	2.9	2.8	2.9	4.4	6.4	8.7	12.4	14.9
2002.....	1.1	1.6	2.8	2.6	4.2	4.0	6.3	7.2	10.7	15.7
2001.....	2.1	2.3	2.5	3.7	3.3	5.2	7.3	8.5	11.6	17.8
2000.....	2.0	2.2	2.6	3.9	4.5	5.0	7.1	11.4	16.1	20.3
1999.....	1.7	2.0	2.3	2.6	4.3	5.7	8.3	12.0	15.3	21.7

Black, male

2004.....	2.0	3.9	3.7	4.2	4.7	7.3	7.8	12.3	15.1	*
2003.....	1.5	3.3	3.2	4.6	3.9	6.4	8.9	15.5	17.0	*
2002.....	1.8	2.1	4.0	3.4	6.3	6.7	9.3	9.2	*	25.7
2001.....	2.9	3.7	4.2	5.0	4.9	7.4	11.1	12.1	*	27.2
2000.....	2.7	3.2	4.0	5.8	7.0	7.5	10.4	16.5	21.9	27.7
1999.....	2.5	3.0	3.3	4.1	4.2	7.8	13.2	17.1	20.1	29.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental exposure to smoke, fire and flames (X00-X09)

Black, female

2004.....	*	1.5	*	*	3.8	*	4.6	5.4	10.6	13.4
2003.....	*	1.4	2.6	*	*	*	4.7	*	10.1	13.4
2002.....	*	*	*	*	*	*	*	6.0	9.2	11.9
2001.....	1.3	*	*	2.7	*	*	4.7	6.4	10.2	14.3
2000.....	1.3	*	*	*	*	*	4.9	8.3	13.2	17.6
1999.....	*	*	*	*	4.3	4.2	5.0	8.9	13.0	18.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes

2004.....	16.3	16.7	12.0	6.7	3.7	2.4	2.1	2.4	2.5	3.4
2003.....	15.5	15.4	10.9	6.1	3.4	2.3	2.0	2.0	2.5	3.5
2002.....	14.8	13.8	9.2	5.0	3.0	2.3	1.9	2.1	2.7	3.9
2001.....	12.0	10.9	7.0	3.7	2.4	1.7	1.5	2.1	2.4	3.5
2000.....	11.2	9.9	6.1	3.3	2.2	1.5	1.5	1.6	2.6	4.3
1999.....	11.1	9.8	5.4	3.4	2.0	1.7	1.6	1.9	2.6	3.4

All races 3/, male

2004.....	20.9	21.6	16.3	8.4	4.2	2.7	2.5	2.6	2.9	4.2
2003.....	20.7	20.8	14.7	7.6	4.5	3.0	2.2	2.3	3.5	4.0
2002.....	20.3	18.7	12.8	6.8	3.6	2.8	2.1	2.3	3.2	5.1
2001.....	16.8	15.7	9.9	5.0	3.0	2.0	2.1	2.7	2.8	3.8
2000.....	16.4	14.7	8.9	4.0	2.7	1.8	1.7	1.6	2.9	6.2
1999.....	16.6	15.1	7.9	4.7	3.0	2.2	1.9	2.1	2.9	4.4

All races 3/, female

2004.....	11.7	11.9	7.9	5.1	3.2	2.1	1.8	2.2	2.3	3.0
2003.....	10.3	10.2	7.3	4.6	2.5	1.8	1.8	1.7	2.0	3.3
2002.....	9.4	9.1	5.8	3.4	2.4	1.8	1.7	1.9	2.3	3.4
2001.....	7.3	6.2	4.1	2.6	1.9	1.4	1.0	1.6	2.2	3.3
2000.....	6.0	5.3	3.4	2.7	1.8	1.3	1.3	1.6	2.4	3.6
1999.....	5.7	4.7	3.1	2.2	1.2	1.2	1.4	1.7	2.4	3.0

White, both sexes

2004.....	17.3	17.1	11.7	6.2	3.5	2.3	2.1	2.3	2.4	3.4
2003.....	16.3	15.7	10.6	5.8	3.4	2.0	2.0	2.0	2.7	3.5
2002.....	15.4	13.7	8.7	4.6	2.7	2.2	1.9	2.1	2.6	3.9
2001.....	12.1	10.5	6.1	3.4	2.1	1.6	1.4	2.0	2.3	3.3
2000.....	11.0	9.4	5.3	3.0	2.1	1.5	1.4	1.7	2.6	4.1
1999.....	10.8	9.1	4.8	3.1	1.7	1.5	1.4	1.9	2.6	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

White, male

2004.....	22.2	22.0	15.4	7.3	3.6	2.3	2.4	2.4	2.7	4.3
2003.....	21.8	20.9	13.8	7.0	4.2	2.3	2.2	2.3	3.6	3.7
2002.....	21.0	18.3	11.7	5.8	2.9	2.6	2.0	2.3	3.0	5.2
2001.....	17.1	14.9	8.5	4.3	2.3	1.9	2.0	2.4	2.7	3.7
2000.....	16.2	13.7	7.6	3.3	2.4	1.6	1.6	1.7	2.9	5.9
1999.....	16.2	13.8	6.6	4.0	2.3	2.0	1.7	2.1	3.0	4.5

White, female

2004.....	12.5	12.2	8.2	5.2	3.4	2.2	1.8	2.3	2.2	3.0
2003.....	10.8	10.5	7.3	4.7	2.6	1.7	1.8	1.8	2.1	3.4
2002.....	9.8	9.0	5.8	3.4	2.5	1.9	1.8	1.9	2.5	3.4
2001.....	7.1	6.1	3.8	2.6	1.9	1.4	1.0	1.7	2.0	3.1
2000.....	5.8	5.1	3.1	2.7	1.8	1.4	1.3	1.7	2.5	3.4
1999.....	5.4	4.5	3.0	2.3	1.2	1.1	1.2	1.8	2.4	2.8

Black, both sexes

2004.....	14.6	19.6	17.9	12.6	6.0	4.1	3.1	*	*	*
2003.....	15.2	18.8	16.9	9.6	4.8	5.5	*	*	*	*
2002.....	15.7	19.4	16.1	10.3	6.0	3.0	*	*	*	*
2001.....	15.2	17.7	15.4	7.4	5.4	2.6	*	*	*	6.4
2000.....	15.3	16.5	14.2	6.8	3.9	2.7	*	*	*	8.1
1999.....	16.1	17.9	12.3	6.5	5.5	2.9	3.2	*	*	*

Black, male

2004.....	19.1	25.8	28.7	20.5	10.1	7.3	*	*	*	*
2003.....	20.5	26.7	25.7	15.8	8.7	10.3	*	*	*	*
2002.....	21.6	27.2	25.6	17.7	11.3	5.5	*	*	*	*
2001.....	20.3	28.0	24.8	12.7	9.6	*	*	*	*	*
2000.....	22.8	26.3	23.1	11.5	7.0	*	*	*	*	*
1999.....	24.2	29.7	20.9	11.9	10.0	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Black, female

2004.....	10.6	14.2	8.9	6.1	*	*	*	*	*	*
2003.....	10.5	11.9	9.5	4.5	*	*	*	*	*	*
2002.....	10.5	12.5	8.1	4.3	*	*	*	*	*	*
2001.....	10.7	8.9	7.5	3.1	*	*	*	*	*	*
2000.....	8.6	8.0	6.6	3.0	*	*	*	*	*	*
1999.....	9.1	7.8	4.9	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

All races 3/, both sexes

2004.....	15.7	16.0	11.4	6.3	3.3	2.0	1.6	1.8	1.8	2.5
2003.....	14.8	14.7	10.4	5.5	2.9	1.9	1.4	1.4	2.1	2.8
2002.....	14.2	13.2	8.7	4.4	2.5	1.8	1.3	1.5	1.9	3.1
2001.....	11.4	10.4	6.4	3.3	2.0	1.4	1.2	1.5	1.7	2.5
2000.....	10.6	9.3	5.6	2.8	1.7	1.2	1.0	1.2	1.7	3.3
1999.....	10.5	9.2	4.9	2.9	1.6	1.3	1.1	1.4	1.9	2.8

All races 3/, male

2004.....	20.0	20.6	15.3	7.8	3.7	2.1	1.7	1.9	1.7	2.6
2003.....	19.6	19.6	13.9	6.8	3.6	2.4	1.4	1.7	2.7	2.7
2002.....	19.2	17.7	12.0	5.8	2.9	2.0	1.4	1.6	2.1	3.7
2001.....	15.8	14.9	9.1	4.4	2.4	1.5	1.5	1.8	1.7	1.8
2000.....	15.6	13.9	8.2	3.2	2.0	1.5	1.2	1.1	1.5	4.4
1999.....	15.5	14.2	7.1	4.0	2.2	1.6	1.3	1.3	1.7	3.3

All races 3/, female

2004.....	11.5	11.6	7.6	4.8	3.0	1.9	1.4	1.8	1.8	2.5
2003.....	10.1	9.9	7.1	4.3	2.3	1.5	1.4	1.3	1.7	2.9
2002.....	9.2	8.9	5.6	3.2	2.1	1.5	1.3	1.4	1.8	2.8
2001.....	7.0	6.0	3.9	2.3	1.7	1.2	0.9	1.2	1.7	2.7
2000.....	5.8	4.9	3.2	2.4	1.4	1.0	0.9	1.3	1.9	2.8
1999.....	5.5	4.5	2.9	1.9	1.1	1.0	1.0	1.4	2.0	2.6

White, both sexes

2004.....	16.8	16.4	11.1	5.7	3.2	1.9	1.6	1.8	1.7	2.6
2003.....	15.6	15.0	10.0	5.3	2.9	1.7	1.4	1.5	2.1	2.8
2002.....	14.8	13.1	8.2	4.1	2.2	1.7	1.4	1.5	1.9	3.2
2001.....	11.5	9.9	5.7	3.0	1.8	1.3	1.1	1.4	1.7	2.3
2000.....	10.5	8.8	4.9	2.5	1.5	1.2	1.0	1.3	1.7	3.1
1999.....	10.2	8.6	4.3	2.7	1.3	1.2	1.0	1.4	1.9	2.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

White, male

2004.....	21.2	20.9	14.5	6.6	3.2	1.7	1.7	1.8	1.6	2.7
2003.....	20.6	19.8	13.0	6.2	3.4	1.9	1.4	1.7	2.7	2.4
2002.....	19.9	17.4	10.9	5.0	2.2	1.8	1.3	1.5	1.9	3.8
2001.....	16.1	14.0	7.8	3.7	1.8	1.4	1.4	1.6	1.8	1.7
2000.....	15.3	12.9	6.9	2.6	1.6	1.2	1.0	1.2	1.4	4.1
1999.....	15.1	12.9	5.9	3.4	1.6	1.4	1.2	1.4	1.7	3.2

White, female

2004.....	12.3	11.9	7.8	4.9	3.2	2.1	1.5	1.9	1.8	2.5
2003.....	10.5	10.2	7.1	4.5	2.4	1.5	1.4	1.4	1.8	3.0
2002.....	9.5	8.8	5.6	3.2	2.3	1.6	1.4	1.5	2.0	2.9
2001.....	6.8	5.9	3.6	2.3	1.7	1.2	0.9	1.3	1.7	2.6
2000.....	5.6	4.7	2.9	2.5	1.4	1.1	0.9	1.3	1.9	2.6
1999.....	5.2	4.3	2.8	2.0	1.1	1.0	0.8	1.4	2.0	2.3

Black, both sexes

2004.....	14.2	19.0	17.4	12.0	5.4	3.5	*	*	*	*
2003.....	14.6	18.1	16.3	9.0	4.2	4.5	*	*	*	*
2002.....	15.0	18.5	15.4	9.1	5.5	2.4	*	*	*	*
2001.....	14.6	17.2	14.3	7.0	4.7	2.2	*	*	*	*
2000.....	14.6	15.8	13.5	5.9	3.4	2.3	*	*	*	6.3
1999.....	15.7	17.1	11.4	5.6	4.7	2.4	*	*	*	*

Black, male

2004.....	18.4	24.8	27.6	19.7	9.0	6.5	*	*	*	*
2003.....	19.6	25.3	24.6	14.7	7.3	8.4	*	*	*	*
2002.....	20.5	25.8	24.4	15.3	10.5	*	*	*	*	*
2001.....	19.2	27.2	23.1	12.2	8.1	*	*	*	*	*
2000.....	21.8	25.2	22.3	9.9	6.2	*	*	*	*	*
1999.....	23.5	28.3	19.5	10.3	8.5	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Black, female

2004.....	10.5	14.0	8.7	5.6	*	*	*	*	*	*
2003.....	10.3	11.9	9.3	4.3	*	*	*	*	*	*
2002.....	10.1	12.1	7.7	3.9	*	*	*	*	*	*
2001.....	10.4	8.5	6.8	2.8	*	*	*	*	*	*
2000.....	8.3	7.8	6.0	*	*	*	*	*	*	*
1999.....	8.8	7.4	4.6	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)

All races 3/, both sexes

2004.....	0.3	0.4	0.4	0.2	*	*	*	*	*	*
2003.....	0.4	0.4	0.3	0.3	0.2	*	*	*	*	*
2002.....	0.4	0.3	0.3	0.3	0.2	*	*	*	*	*
2001.....	0.4	0.3	0.3	0.2	0.2	*	*	*	*	*
2000.....	0.3	0.3	0.2	0.3	0.2	*	*	*	*	*
1999.....	0.4	0.3	0.2	0.3	0.2	*	*	*	*	*

All races 3/, male

2004.....	0.5	0.6	0.5	0.3	*	*	*	*	*	*
2003.....	0.7	0.7	0.5	0.4	0.4	*	*	*	*	*
2002.....	0.6	0.5	0.5	0.5	0.4	*	*	*	*	*
2001.....	0.5	0.4	0.5	*	*	*	*	*	*	*
2000.....	0.5	0.5	0.4	0.4	*	*	*	*	*	*
1999.....	0.6	0.5	0.4	0.3	*	*	*	*	*	*

All races 3/, female

2004.....	*	0.2	0.2	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.3	0.4	0.4	0.2	*	*	*	*	*	*
2003.....	0.4	0.4	0.3	0.3	0.2	*	*	*	*	*
2002.....	0.4	0.3	0.3	0.2	*	*	*	*	*	*
2001.....	0.4	0.3	0.3	0.2	*	*	*	*	*	*
2000.....	0.3	0.3	0.2	0.3	0.2	*	*	*	*	*
1999.....	0.4	0.3	0.2	0.2	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning by and exposure to other gases and vapors (X47)

All races 3/, both sexes

2004.....	0.2	0.3	0.2	0.3	0.2	0.2	0.4	0.3	0.5	0.7
2003.....	0.2	0.3	0.2	0.3	0.3	0.3	0.5	0.4	*	0.6
2002.....	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.5	0.5	0.6
2001.....	0.2	0.3	0.2	0.2	0.2	*	0.2	0.5	0.6	0.9
2000.....	0.2	0.3	0.2	0.3	0.3	*	0.3	0.3	0.7	0.8
1999.....	0.3	0.2	0.2	0.3	0.2	0.2	0.3	0.3	0.5	0.5

All races 3/, male

2004.....	0.4	0.5	0.4	0.4	*	*	0.6	*	*	1.3
2003.....	0.4	0.5	0.4	0.4	0.5	0.4	0.7	*	*	*
2002.....	0.5	0.4	0.4	0.4	0.4	0.6	0.6	*	1.0	*
2001.....	0.4	0.4	0.4	0.3	*	*	*	0.8	*	1.7
2000.....	0.4	0.4	0.4	0.4	0.4	*	*	*	1.2	*
1999.....	0.5	0.4	0.4	0.4	0.4	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	0.3	0.2	0.3	0.2	*	0.4	0.3	0.4	0.7
2003.....	0.3	0.3	0.2	0.3	0.3	*	0.5	0.5	*	0.5
2002.....	0.3	0.3	0.3	0.2	0.3	0.4	0.4	0.5	0.5	0.7
2001.....	0.2	0.3	0.2	0.2	*	*	*	0.4	0.5	0.8
2000.....	0.2	0.3	0.2	0.2	0.3	*	0.3	*	0.7	0.7
1999.....	0.3	0.2	0.2	0.3	*	*	*	0.3	0.5	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

All races 3/, both sexes

2004.....	2.0	2.4	2.9	2.8	3.6	4.4	6.6	11.7	23.2	64.5
2003.....	2.2	2.5	2.6	3.1	3.4	5.3	6.7	12.7	24.3	73.5
2002.....	2.2	2.2	2.7	3.2	3.9	5.5	7.0	13.5	25.6	75.8
2001.....	2.0	2.5	2.9	2.9	4.3	4.9	8.0	14.4	28.4	83.4
2000.....	2.3	2.6	2.7	3.1	4.1	5.0	7.6	13.6	26.9	82.6
1999.....	2.4	2.5	2.9	3.2	4.3	5.9	8.1	15.3	32.3	92.8

All races 3/, male

2004.....	3.2	4.1	4.8	4.4	5.7	6.5	9.0	14.9	27.9	69.6
2003.....	3.8	4.1	4.2	4.9	5.6	7.9	9.5	16.9	28.7	80.4
2002.....	3.5	3.8	4.2	5.1	6.1	7.9	9.8	17.8	30.3	79.9
2001.....	3.3	4.2	4.7	4.4	6.6	7.1	10.6	19.0	33.9	92.4
2000.....	3.9	4.4	4.5	5.2	6.1	7.1	10.0	17.5	31.1	91.7
1999.....	3.9	4.1	4.9	5.2	6.6	8.2	11.1	19.5	40.3	107.0

All races 3/, female

2004.....	0.8	0.8	1.2	1.3	1.7	2.7	4.8	9.3	20.3	62.2
2003.....	0.7	0.9	1.1	1.4	1.4	3.1	4.5	9.7	21.6	70.5
2002.....	0.8	0.7	1.2	1.4	1.8	3.3	4.7	10.4	22.7	74.0
2001.....	0.7	0.9	1.3	1.4	2.2	3.1	5.8	11.2	25.1	79.6
2000.....	0.7	0.8	0.9	1.2	2.2	3.2	5.6	10.9	24.3	78.9
1999.....	0.9	0.9	1.0	1.4	2.2	3.9	5.8	12.4	27.6	87.1

White, both sexes

2004.....	2.1	2.5	2.9	2.7	3.6	4.4	6.7	11.6	23.8	67.6
2003.....	2.2	2.3	2.4	2.9	3.3	5.3	6.7	12.9	24.8	77.0
2002.....	2.2	2.2	2.6	3.1	3.8	5.1	7.1	13.4	25.9	79.1
2001.....	1.9	2.4	2.8	2.9	4.1	4.8	7.9	14.6	29.2	86.4
2000.....	2.2	2.4	2.6	3.0	4.0	4.8	7.5	13.4	27.5	86.2
1999.....	2.3	2.2	2.8	3.0	4.0	5.6	7.8	15.0	32.6	96.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

White, male										
2004.....	3.4	4.2	4.7	4.2	5.7	6.4	9.1	14.7	28.5	72.6
2003.....	3.8	3.9	3.9	4.6	5.4	7.8	9.2	17.1	28.8	83.8
2002.....	3.6	3.8	4.1	4.8	6.0	7.3	9.9	17.6	30.7	82.5
2001.....	3.1	4.1	4.4	4.4	6.3	6.9	10.5	19.3	34.4	95.8
2000.....	3.7	4.0	4.3	4.7	5.8	6.7	10.0	16.9	31.4	95.0
1999.....	3.7	3.6	4.6	4.8	6.3	7.7	10.5	19.0	40.5	108.9
White, female										
2004.....	0.9	0.8	1.1	1.3	1.6	2.6	4.7	9.4	20.9	65.4
2003.....	0.6	0.8	1.0	1.4	1.5	3.0	4.6	9.9	22.2	74.0
2002.....	0.7	0.6	1.2	1.5	1.8	3.1	4.8	10.4	23.0	77.6
2001.....	0.6	0.8	1.3	1.5	2.0	3.1	5.8	11.3	26.0	82.4
2000.....	0.6	0.7	0.9	1.3	2.3	3.2	5.4	11.0	25.3	82.6
1999.....	0.9	0.9	1.0	1.4	1.9	3.8	5.7	12.2	27.9	91.0
Black, both sexes										
2004.....	1.9	2.7	3.8	4.3	4.3	5.7	6.7	14.6	19.6	38.4
2003.....	2.5	3.8	4.3	4.6	4.6	7.0	8.1	11.9	22.3	45.8
2002.....	2.4	3.0	3.3	4.7	4.7	9.3	6.3	15.6	22.8	46.9
2001.....	2.8	3.5	4.3	2.8	6.8	6.4	9.6	15.1	23.0	58.3
2000.....	3.1	4.2	3.6	5.3	5.4	6.8	9.3	17.8	21.2	51.0
1999.....	3.4	4.7	4.6	5.7	7.1	9.4	13.1	20.7	32.6	67.0
Black, male										
2004.....	3.0	4.6	6.3	7.5	7.1	8.5	9.7	21.4	28.0	42.7
2003.....	4.0	6.3	6.9	7.7	8.5	10.8	14.1	16.9	29.5	54.2
2002.....	3.2	4.6	5.5	8.6	7.9	13.5	9.0	22.5	29.8	57.9
2001.....	4.7	5.4	7.3	5.0	10.0	10.4	13.1	18.6	32.9	62.4
2000.....	5.4	7.8	7.1	10.4	9.3	10.9	11.7	27.9	32.9	69.2
1999.....	5.9	8.4	7.8	9.8	10.8	15.1	20.4	28.0	41.0	106.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

Black, female

2004.....	*	*	1.7	*	*	3.6	4.6	10.5	15.1	36.6
2003.....	1.3	1.6	2.2	*	*	4.2	*	9.0	18.6	42.5
2002.....	1.8	1.6	*	*	*	6.3	4.5	11.5	19.1	42.7
2001.....	*	1.7	*	*	4.3	*	7.2	13.0	18.0	56.8
2000.....	*	*	*	*	*	3.8	7.8	11.7	15.3	44.2
1999.....	*	*	*	*	4.2	5.2	8.3	16.3	28.5	52.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All races 3/, both sexes

2004.....	16.0	16.9	16.2	14.2	13.2	11.4	13.5	16.0	16.7	16.4
2003.....	15.9	16.0	15.7	14.7	12.6	12.0	13.6	15.6	17.7	16.9
2002.....	16.1	16.3	15.1	14.6	12.3	12.5	14.6	16.6	19.3	18.0
2001 4/.....	15.2	15.7	14.6	14.0	12.0	12.7	13.9	16.4	18.9	17.5
2000.....	15.4	14.9	13.9	12.7	11.4	11.3	13.8	16.5	19.3	19.6
1999.....	14.3	14.3	13.4	12.8	11.4	12.4	14.5	17.4	19.2	19.3

All races 3/, male

2004.....	24.2	25.2	24.4	22.3	21.7	20.2	25.4	32.4	38.3	45.0
2003.....	24.5	24.4	24.4	23.1	21.1	21.2	26.0	31.8	40.1	47.8
2002.....	24.7	24.9	23.8	23.5	20.4	22.0	27.9	34.3	44.0	50.7
2001 4/.....	23.6	24.3	22.4	22.3	19.5	22.7	26.7	33.8	44.2	51.1
2000.....	23.9	23.3	21.4	20.1	18.7	19.8	26.1	34.9	44.8	57.5
1999.....	22.1	22.3	20.6	20.3	19.1	22.1	27.7	35.6	44.1	57.1

All races 3/, female

2004.....	7.9	8.9	8.4	6.6	5.5	3.7	3.9	4.3	3.5	3.6
2003.....	7.4	7.9	7.5	6.7	4.9	4.0	3.6	4.0	3.9	3.3
2002.....	7.7	8.0	6.7	6.2	5.0	4.3	3.9	4.1	4.4	3.8
2001 4/.....	6.9	7.4	7.0	6.2	5.2	4.1	3.7	4.3	3.8	3.4
2000.....	7.0	6.7	6.8	5.8	4.9	4.0	4.1	3.8	4.2	4.2
1999.....	6.7	6.7	6.5	5.8	4.4	4.2	4.1	4.8	4.6	4.1

White, both sexes

2004.....	18.1	19.0	18.3	15.7	14.4	12.4	14.5	17.3	17.7	17.5
2003.....	17.9	18.1	17.7	16.2	13.8	13.0	14.7	16.7	18.9	18.1
2002.....	18.2	18.4	16.8	16.1	13.6	13.7	15.7	17.6	20.9	19.1
2001 4/.....	16.9	17.6	16.2	15.3	13.1	13.8	14.9	17.6	20.0	18.6
2000.....	17.0	16.5	15.3	13.9	12.6	12.3	14.7	17.6	20.4	21.0
1999.....	16.0	15.9	14.8	14.0	12.3	13.3	15.6	18.6	20.4	20.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
White, male										
2004.....	27.0	27.8	27.3	24.1	23.6	21.9	27.1	34.8	40.4	48.4
2003.....	27.3	27.3	27.2	25.4	22.9	22.9	27.9	34.0	42.9	51.4
2002.....	27.6	27.6	26.4	25.7	22.5	23.8	30.1	36.3	47.2	53.9
2001 4/.....	26.1	27.1	24.7	24.3	21.0	24.5	28.3	36.0	46.8	55.0
2000.....	26.2	25.6	23.2	21.8	20.4	21.3	27.6	37.1	47.7	61.6
1999.....	24.5	24.4	22.3	21.9	20.5	23.6	29.5	37.9	46.8	61.5
White, female										
2004.....	9.1	10.2	9.5	7.6	5.8	4.1	4.1	4.5	3.6	3.6
2003.....	8.4	9.0	8.3	7.4	5.5	4.3	3.9	4.2	4.1	3.3
2002.....	8.7	9.3	7.4	6.9	5.3	4.8	3.9	4.2	4.7	4.0
2001 4/.....	7.8	8.1	7.9	6.8	5.8	4.3	4.0	4.6	3.8	3.4
2000.....	7.9	7.5	7.5	6.4	5.5	4.4	4.4	3.9	4.2	4.5
1999.....	7.5	7.4	7.4	6.4	4.8	4.4	4.5	5.0	4.8	4.3
Black, both sexes										
2004.....	6.4	6.4	5.1	6.4	4.8	3.5	5.5	5.6	6.7	*
2003.....	7.0	5.4	5.9	4.8	5.2	4.5	4.5	4.4	*	*
2002.....	7.1	5.9	5.9	4.6	5.0	3.5	6.6	6.7	*	5.9
2001 4/.....	6.8	6.5	5.9	5.8	4.4	4.6	6.5	5.2	6.5	*
2000.....	8.0	6.6	6.4	4.6	3.7	4.3	6.1	5.3	6.5	*
1999.....	6.8	6.4	4.7	5.2	4.7	6.4	5.1	6.2	*	*
Black, male										
2004.....	10.9	10.6	8.6	12.3	8.8	7.8	12.6	13.6	17.2	*
2003.....	11.4	8.9	9.4	8.0	9.6	7.9	8.9	11.0	*	*
2002.....	11.5	11.0	9.2	8.6	8.1	7.7	12.4	15.6	*	*
2001 4/.....	10.8	10.1	9.6	10.2	8.3	8.1	14.1	12.1	16.1	*
2000.....	13.4	11.8	10.5	8.4	6.4	9.4	13.4	11.8	*	*
1999.....	10.8	10.9	8.7	9.3	8.9	12.2	11.2	14.2	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

All races 3/, both sexes

2004.....	2.8	3.1	3.1	2.3	1.6	0.9	0.9	1.0	0.8	1.1
2003.....	2.8	3.0	2.8	2.1	1.4	1.1	0.8	0.9	0.9	0.9
2002.....	2.9	3.1	2.6	2.0	1.4	1.0	1.0	1.0	1.1	0.8
2001.....	2.7	2.9	2.4	1.7	1.2	1.0	0.9	0.9	0.6	0.9
2000.....	2.6	2.5	2.3	1.8	1.0	0.7	0.9	0.8	0.9	1.0
1999.....	2.5	2.4	2.2	1.6	1.1	1.0	0.9	1.1	1.1	0.8

All races 3/, male

2004.....	2.5	2.8	2.9	2.1	1.6	0.6	0.9	1.0	*	1.6
2003.....	2.9	2.7	2.6	1.9	1.2	0.9	0.8	0.7	*	1.7
2002.....	2.8	2.7	2.6	1.7	1.2	0.9	0.9	1.1	1.3	*
2001.....	2.8	2.8	2.0	1.4	0.8	1.2	0.8	0.8	*	*
2000.....	2.5	2.5	2.0	1.6	0.7	0.5	0.8	0.9	1.3	*
1999.....	2.5	2.5	2.0	1.4	1.0	1.1	0.9	0.7	1.2	*

All races 3/, female

2004.....	3.2	3.4	3.3	2.6	1.6	1.0	0.9	1.0	0.8	0.8
2003.....	2.7	3.3	2.9	2.3	1.6	1.3	0.9	1.1	1.0	*
2002.....	3.1	3.5	2.5	2.3	1.6	1.1	1.1	0.9	0.9	0.7
2001.....	2.6	3.1	2.7	1.9	1.5	0.9	1.0	0.9	0.6	0.6
2000.....	2.8	2.5	2.5	1.9	1.3	0.7	1.1	0.8	0.7	0.9
1999.....	2.5	2.4	2.4	1.8	1.2	0.9	0.9	1.3	1.0	*

White, both sexes

2004.....	3.3	3.6	3.5	2.6	1.8	0.9	1.0	1.1	0.9	1.1
2003.....	3.1	3.4	3.2	2.3	1.6	1.2	0.9	1.0	1.0	1.0
2002.....	3.4	3.6	2.9	2.3	1.5	1.1	1.0	1.1	1.1	0.9
2001.....	3.1	3.3	2.7	1.9	1.3	1.1	1.0	1.0	0.7	0.9
2000.....	3.0	2.8	2.5	2.0	1.2	0.7	1.1	0.9	1.0	1.1
1999.....	2.8	2.8	2.5	1.8	1.2	1.1	1.0	1.1	1.2	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

All races 3/, both sexes

2004.....	0.9	0.9	0.9	0.6	0.5	0.3	0.4	0.5	0.5	0.6
2003.....	0.9	0.8	0.8	0.8	0.5	0.3	0.5	0.4	0.6	0.7
2002.....	0.9	1.0	0.9	0.8	0.4	0.4	0.4	0.6	0.7	0.8
2001.....	0.9	1.0	0.8	0.9	0.5	0.5	0.4	0.6	0.7	0.8
2000.....	0.9	1.1	0.8	0.8	0.5	0.4	0.4	0.5	0.7	0.8
1999.....	1.0	1.0	0.9	0.7	0.6	0.3	0.4	0.8	0.9	1.2

All races 3/, male

2004.....	1.4	1.4	1.5	1.0	0.9	*	0.6	0.8	1.0	*
2003.....	1.4	1.3	1.2	1.2	0.9	0.6	0.9	0.6	1.0	1.6
2002.....	1.5	1.7	1.3	1.3	0.7	0.7	0.6	1.1	1.5	2.2
2001.....	1.4	1.6	1.2	1.3	0.7	0.8	0.7	1.0	1.5	1.9
2000.....	1.4	1.7	1.2	1.3	0.7	0.6	0.6	1.0	1.6	2.3
1999.....	1.5	1.5	1.2	1.0	0.8	0.6	0.6	1.3	1.7	3.9

All races 3/, female

2004.....	0.3	0.4	0.4	0.3	*	*	*	*	*	*
2003.....	0.4	0.3	0.4	0.3	*	*	*	*	*	*
2002.....	0.4	0.4	0.4	0.3	*	*	*	*	*	*
2001.....	0.4	0.4	0.4	0.6	*	*	*	*	*	*
2000.....	0.4	0.4	0.5	0.4	*	*	*	*	*	*
1999.....	0.4	0.5	0.5	0.4	0.4	*	*	*	*	*

White, both sexes

2004.....	1.0	1.1	1.1	0.7	0.6	0.3	0.5	0.6	0.5	0.7
2003.....	1.0	1.0	0.9	0.9	0.5	0.4	0.6	0.5	0.7	0.8
2002.....	1.1	1.2	1.0	0.9	0.4	0.5	0.5	0.7	0.8	0.9
2001.....	1.1	1.1	1.0	1.0	0.6	0.6	0.5	0.7	0.8	0.9
2000.....	1.1	1.2	0.9	0.9	0.6	0.4	0.4	0.5	0.8	0.9
1999.....	1.1	1.1	1.0	0.8	0.7	0.4	0.4	0.8	0.9	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

All races 3/, both sexes

2004.....	3.7	3.3	2.5	2.2	1.6	1.3	1.2	1.5	1.7	2.2
2003.....	3.4	2.9	2.4	2.1	1.4	1.2	1.2	1.5	1.9	2.1
2002.....	3.4	3.0	2.4	1.9	1.4	1.4	1.3	1.5	1.8	2.3
2001.....	3.2	2.7	2.1	1.7	1.4	1.3	1.3	1.4	2.1	2.1
2000.....	3.1	2.5	1.9	1.5	1.3	1.1	1.2	1.7	2.0	2.9
1999.....	2.8	2.1	1.9	1.4	1.3	1.4	1.3	1.8	2.0	2.8

All races 3/, male

2004.....	6.4	5.4	4.1	3.6	2.7	2.2	2.0	2.5	3.4	4.8
2003.....	5.8	4.9	4.0	3.5	2.2	2.1	2.3	2.6	3.7	5.0
2002.....	5.6	5.0	3.8	3.3	2.4	2.4	2.2	2.6	3.0	5.2
2001.....	5.5	4.7	3.6	2.9	2.1	2.2	2.4	2.3	4.3	5.2
2000.....	5.4	4.3	3.2	2.4	2.0	1.5	2.1	3.0	3.8	7.2
1999.....	4.7	3.6	3.0	2.5	2.1	2.1	2.2	2.9	3.6	6.6

All races 3/, female

2004.....	1.1	1.3	1.1	0.9	0.7	0.5	0.5	0.7	0.7	1.1
2003.....	1.1	0.9	0.9	0.7	0.6	0.5	*	0.6	0.8	0.9
2002.....	1.1	1.0	1.0	0.6	0.6	0.6	0.6	0.7	1.0	1.0
2001.....	0.9	0.8	0.7	0.6	0.7	0.6	*	0.7	0.8	0.8
2000.....	0.9	0.8	0.7	0.7	0.7	0.8	0.5	0.8	1.0	1.2
1999.....	0.8	0.7	0.7	0.4	0.6	0.8	0.5	1.0	1.1	1.3

White, both sexes

2004.....	4.1	3.6	2.7	2.4	1.6	1.3	1.1	1.4	1.8	2.3
2003.....	3.7	3.1	2.6	2.1	1.4	1.1	1.2	1.5	2.0	2.1
2002.....	3.7	3.2	2.6	2.0	1.6	1.3	1.3	1.5	1.8	2.3
2001.....	3.5	3.0	2.2	1.8	1.3	1.3	1.2	1.4	2.1	2.1
2000.....	3.3	2.7	2.0	1.6	1.4	1.1	1.2	1.6	2.0	2.9
1999.....	3.0	2.3	1.9	1.5	1.3	1.4	1.3	1.7	2.1	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes										
2004.....	7.1	8.1	8.0	7.8	8.3	8.0	9.8	12.2	12.3	11.1
2003.....	7.2	7.8	8.3	8.4	8.2	8.5	10.1	11.9	13.1	11.9
2002.....	7.4	7.8	7.9	8.6	8.1	8.8	10.7	12.4	14.3	12.0
2001.....	7.0	7.5	8.0	8.4	8.0	9.0	10.3	12.4	14.1	12.2
2000.....	7.4	7.4	7.8	7.4	7.7	8.3	10.3	12.4	14.0	13.2
1999.....	6.7	7.4	7.2	8.1	7.5	8.7	10.9	12.9	13.5	12.6
All races 3/, male										
2004.....	11.7	13.4	13.6	13.8	14.9	15.5	20.0	26.7	30.7	34.4
2003.....	12.1	13.2	14.5	14.6	15.3	16.5	20.7	26.3	32.4	36.6
2002.....	12.6	13.3	14.0	15.4	14.8	17.0	22.5	27.8	35.9	37.8
2001.....	12.0	12.8	13.8	14.9	14.2	17.3	21.4	27.7	35.5	38.9
2000.....	12.6	12.8	13.3	13.1	13.9	15.9	21.3	28.1	35.4	42.9
1999.....	11.2	12.6	12.4	14.0	14.0	16.9	22.4	29.1	34.6	41.6
All races 3/, female										
2004.....	2.6	2.8	2.7	2.1	2.3	1.4	1.7	1.7	1.0	*
2003.....	2.3	2.6	2.3	2.5	1.7	1.6	1.6	1.6	1.3	0.9
2002.....	2.3	2.4	2.1	2.3	2.0	1.8	1.3	1.5	1.2	0.8
2001.....	2.2	2.3	2.4	2.4	2.3	1.9	1.4	1.7	1.3	1.0
2000.....	2.4	2.2	2.5	2.1	2.1	1.8	1.6	1.4	1.4	1.1
1999.....	2.2	2.4	2.1	2.6	1.6	1.7	1.8	1.6	1.2	0.9
White, both sexes										
2004.....	8.1	9.2	9.2	8.7	9.3	8.8	10.8	13.3	13.3	12.1
2003.....	8.3	9.1	9.4	9.5	9.1	9.5	11.1	12.9	14.1	12.9
2002.....	8.5	8.9	8.9	9.7	9.0	9.9	11.7	13.3	15.6	12.9
2001.....	7.9	8.5	9.0	9.4	8.9	10.0	11.1	13.4	15.0	13.2
2000.....	8.4	8.3	8.7	8.2	8.5	9.2	11.1	13.4	15.2	14.3
1999.....	7.5	8.3	8.0	8.9	8.2	9.5	11.8	13.9	14.4	13.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

White, male										
2004.....	13.1	15.1	15.5	15.1	16.6	16.9	21.6	28.7	32.9	37.5
2003.....	13.8	15.1	16.2	16.4	16.8	18.3	22.5	28.3	34.8	39.9
2002.....	14.3	15.0	15.5	16.9	16.3	18.8	24.4	29.6	38.8	40.5
2001.....	13.3	14.5	15.4	16.3	15.7	19.0	22.9	29.7	37.8	42.1
2000.....	14.0	14.2	14.7	14.2	15.3	17.3	22.7	30.4	38.1	46.5
1999.....	12.5	14.0	13.7	15.1	15.2	18.3	24.0	31.2	36.8	44.9
White, female										
2004.....	3.0	3.3	3.1	2.5	2.6	1.7	1.9	2.0	1.1	*
2003.....	2.7	3.1	2.6	2.9	2.0	1.8	1.8	1.8	1.4	0.9
2002.....	2.7	2.8	2.4	2.7	2.3	2.1	1.4	1.6	1.3	0.9
2001.....	2.4	2.6	2.8	2.7	2.7	2.0	1.5	1.8	1.3	1.1
2000.....	2.7	2.5	2.8	2.4	2.3	2.1	1.8	1.5	1.5	1.2
1999.....	2.5	2.7	2.5	3.0	1.8	1.8	2.1	1.8	1.3	1.0
Black, both sexes										
2004.....	2.5	2.5	2.2	3.3	2.6	3.1	4.1	4.9	*	*
2003.....	2.4	1.9	3.2	2.5	3.0	3.4	3.2	3.9	*	*
2002.....	3.0	2.9	2.9	2.7	2.9	2.3	4.6	5.3	*	*
2001.....	3.2	2.6	3.0	3.0	2.4	2.9	4.7	4.5	6.2	*
2000.....	3.3	3.3	3.3	2.9	3.0	3.1	4.5	*	*	*
1999.....	2.7	3.4	2.4	3.5	3.4	4.0	3.9	5.0	*	*
Black, male										
2004.....	4.9	4.7	4.0	6.5	4.9	7.0	10.0	12.7	*	*
2003.....	4.2	3.7	6.0	4.2	6.4	6.4	6.9	9.6	*	*
2002.....	5.2	5.7	5.3	5.5	5.5	5.0	10.0	13.3	*	*
2001.....	5.4	4.4	5.4	6.1	5.1	5.6	10.4	10.7	*	*
2000.....	6.1	6.0	5.6	5.8	5.6	7.0	11.0	*	*	*
1999.....	5.1	6.0	4.9	7.1	6.8	7.8	9.8	11.9	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-harm (suicide) by jumping from a high place (X80)

All races 3/, both sexes

2004.....	0.3	0.2	0.3	0.3	0.3	0.2	0.3	*	0.4	0.5
2003.....	0.4	0.3	0.3	0.3	0.3	*	0.3	*	*	0.6
2002.....	0.4	0.3	0.3	0.3	0.2	*	0.3	*	0.5	0.6
2001.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.4	*	*
2000.....	0.2	0.3	0.3	0.2	0.2	*	0.2	*	0.5	0.6
1999.....	0.4	0.4	0.3	0.2	*	0.3	0.3	0.3	0.6	0.5

All races 3/, male

2004.....	0.5	0.3	0.4	0.4	0.4	*	*	*	*	*
2003.....	0.5	0.5	0.4	0.5	0.3	*	*	*	*	*
2002.....	0.5	0.4	0.4	0.5	0.4	*	*	*	*	*
2001.....	0.4	0.4	0.4	0.5	0.4	*	*	0.7	*	*
2000.....	0.3	0.5	0.4	*	*	*	*	*	*	*
1999.....	0.6	0.5	0.4	*	*	*	0.5	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	0.3	0.2	0.2	*	*	*	*	*	*	*
2002.....	0.2	0.2	0.2	*	*	*	*	*	*	*
2001.....	0.2	0.2	0.3	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.4	0.2	0.3	0.3	0.3	0.3	*	*	*	*
2003.....	0.4	0.3	0.3	0.4	0.3	*	*	*	*	0.5
2002.....	0.4	0.3	0.3	0.3	0.2	*	0.3	*	0.5	0.6
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	*	*
2000.....	0.2	0.4	0.3	0.2	0.2	*	*	*	0.5	0.5
1999.....	0.4	0.3	0.3	0.2	*	0.3	0.3	0.3	0.5	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)

All races 3/, both sexes

2004.....	1.0	1.1	1.1	0.8	0.8	0.7	0.8	0.5	0.8	0.9
2003.....	1.1	1.0	1.0	0.9	0.8	0.5	0.6	0.6	0.8	0.7
2002.....	1.0	1.0	1.0	0.9	0.7	0.6	0.8	0.8	0.9	1.4
2001 4/.....	0.8	1.1	0.9	0.8	0.6	0.6	0.7	0.8	1.0	1.0
2000.....	0.9	0.9	0.8	0.9	0.7	0.6	0.7	0.8	1.0	1.2
1999.....	0.9	0.9	0.9	0.6	0.6	0.7	0.7	0.6	1.1	1.3

All races 3/, male

2004.....	1.5	1.6	1.8	1.2	1.1	1.0	1.3	0.9	1.4	2.2
2003.....	1.6	1.6	1.5	1.3	1.1	0.7	1.0	1.1	1.5	1.7
2002.....	1.6	1.6	1.6	1.3	1.0	0.9	1.2	1.3	1.5	3.1
2001 4/.....	1.2	1.8	1.3	1.2	1.1	0.9	1.0	1.2	1.8	2.5
2000.....	1.4	1.4	1.1	1.4	0.9	1.0	1.0	1.4	1.8	2.4
1999.....	1.3	1.3	1.3	1.0	0.9	1.0	1.0	1.1	1.9	2.5

All races 3/, female

2004.....	0.5	0.7	0.6	0.5	0.4	0.4	0.4	*	*	*
2003.....	0.6	0.5	0.5	0.5	0.5	*	*	*	*	*
2002.....	0.5	0.4	0.4	0.5	0.5	*	0.5	*	*	0.7
2001 4/.....	0.5	0.5	0.5	0.5	*	*	0.4	0.5	*	*
2000.....	0.4	0.5	0.5	0.4	0.4	*	0.4	*	*	0.7
1999.....	0.6	0.4	0.5	0.3	0.4	0.5	0.5	*	*	0.8

White, both sexes

2004.....	1.0	1.1	1.2	0.8	0.8	0.7	0.8	0.5	0.7	0.9
2003.....	1.2	1.1	1.1	1.0	0.8	0.5	0.6	0.5	0.8	0.7
2002.....	1.1	1.1	1.0	0.9	0.8	0.6	0.8	0.8	0.9	1.5
2001 4/.....	0.9	1.2	0.9	0.9	0.7	0.5	0.7	0.8	1.0	1.1
2000.....	0.9	1.0	0.8	0.9	0.7	0.6	0.7	0.9	1.0	1.2
1999.....	1.0	0.9	0.9	0.6	0.6	0.7	0.7	0.7	1.1	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) (*U01-*U02,X85-Y09,X87.1)

All races 3/, both sexes

2004.....	6.3	5.3	4.2	3.2	2.8	2.6	2.2	2.2	2.2	2.1
2003.....	6.4	5.6	4.2	3.1	2.4	2.3	2.4	2.5	2.4	2.2
2002.....	6.5	5.4	4.0	3.5	2.7	2.4	2.2	2.3	2.4	2.1
2001 4/.....	8.6	7.0	5.5	4.4	3.5	3.3	2.5	2.5	2.5	2.4
2000.....	6.6	5.3	4.0	3.4	2.6	2.4	2.4	2.3	2.6	2.4
1999.....	6.6	5.1	4.0	3.2	2.8	2.7	2.4	2.5	2.4	2.4

All races 3/, male

2004.....	9.0	7.8	6.7	4.9	4.1	3.8	3.3	2.6	2.7	2.7
2003.....	9.5	8.3	6.4	4.6	3.7	3.4	3.4	3.5	3.0	3.2
2002.....	9.6	8.0	6.0	5.3	4.2	3.4	3.2	3.3	2.9	3.0
2001 4/.....	12.6	10.6	8.3	6.7	5.0	4.6	3.3	3.2	3.0	3.2
2000.....	9.7	7.8	5.9	5.2	3.9	3.3	3.4	2.8	3.8	3.3
1999.....	9.5	7.7	5.8	5.1	4.3	4.2	3.1	3.4	3.4	3.4

All races 3/, female

2004.....	3.6	3.0	1.9	1.6	1.6	1.6	1.3	1.9	2.0	1.9
2003.....	3.4	3.0	2.1	1.7	1.3	1.4	1.6	1.8	2.1	1.8
2002.....	3.5	2.9	2.2	1.8	1.4	1.5	1.4	1.6	2.1	1.7
2001 4/.....	4.7	3.6	2.7	2.3	2.2	2.1	1.9	1.9	2.2	2.0
2000.....	3.6	2.8	2.1	1.7	1.5	1.7	1.6	2.0	1.9	2.0
1999.....	3.6	2.5	2.2	1.4	1.5	1.5	1.9	1.9	1.9	2.0

White, both sexes

2004.....	4.3	3.8	3.0	2.4	2.3	2.2	1.8	1.8	2.0	1.9
2003.....	4.2	3.9	3.1	2.4	2.0	1.9	1.9	2.2	2.1	2.0
2002.....	4.5	3.8	2.9	2.9	2.2	1.9	1.9	1.8	2.1	1.8
2001 4/.....	6.8	5.6	4.4	3.8	3.0	2.7	2.2	2.1	2.2	2.2
2000.....	4.4	3.7	2.9	2.7	2.0	1.8	2.0	2.1	2.0	2.2
1999.....	4.5	3.6	3.0	2.6	2.3	2.3	2.0	2.0	2.1	2.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) (*U01-*U02,X85-Y09,X87.1)

White, male

2004.....	5.8	5.2	4.6	3.6	3.2	3.0	2.6	2.0	2.2	2.3
2003.....	5.8	5.4	4.5	3.3	2.9	2.7	2.6	2.9	2.5	3.0
2002.....	6.5	5.1	4.0	4.1	3.2	2.7	2.6	2.4	2.3	2.7
2001 4/.....	9.8	8.2	6.5	5.5	4.2	3.7	2.6	2.7	2.6	2.8
2000.....	6.4	5.1	4.0	4.0	2.8	2.2	2.6	2.4	2.5	2.8
1999.....	6.3	5.3	4.3	3.8	3.3	3.4	2.5	2.6	2.9	2.7

White, female

2004.....	2.9	2.3	1.4	1.3	1.4	1.6	1.1	1.6	1.9	1.8
2003.....	2.7	2.4	1.7	1.6	1.0	1.2	1.3	1.6	1.8	1.6
2002.....	2.5	2.5	1.8	1.7	1.3	1.3	1.4	1.4	1.9	1.5
2001 4/.....	3.8	3.0	2.4	2.1	1.9	1.9	1.8	1.7	1.9	2.0
2000.....	2.5	2.3	1.8	1.4	1.3	1.4	1.5	1.8	1.7	1.9
1999.....	2.6	1.9	1.7	1.3	1.3	1.4	1.7	1.6	1.6	1.9

Black, both sexes

2004.....	20.2	16.8	13.4	9.4	7.9	5.7	6.4	7.1	5.2	*
2003.....	21.4	17.7	12.7	9.1	6.8	5.7	6.9	6.6	6.6	*
2002.....	20.6	17.6	13.0	8.9	7.5	5.8	5.6	8.2	6.0	*
2001 4/.....	21.7	18.1	13.6	9.9	7.5	7.7	6.5	6.9	6.2	*
2000.....	21.8	16.8	12.4	9.1	7.4	8.2	6.8	5.3	9.6	*
1999.....	21.2	16.2	11.2	8.8	7.4	6.6	6.0	8.2	6.1	*

Black, male

2004.....	32.9	27.8	23.8	16.5	13.1	11.6	11.3	10.5	*	*
2003.....	36.8	30.0	21.8	17.3	11.4	10.8	11.5	12.3	*	*
2002.....	32.4	30.8	22.5	16.8	13.5	10.0	11.3	15.2	*	*
2001 4/.....	34.2	30.0	23.1	17.4	12.0	13.2	11.8	11.6	*	*
2000.....	34.1	28.7	22.4	16.0	13.0	15.1	12.4	*	19.4	*
1999.....	34.1	27.3	18.5	17.0	12.7	13.0	10.2	13.8	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) (*U01-*U02,X85-Y09,X87.1)

Black, female

2004.....	8.9	7.2	4.6	3.6	3.8	*	*	*	*	*
2003.....	7.9	7.0	5.0	2.3	3.2	*	*	*	*	*
2002.....	10.1	6.1	4.9	*	*	*	*	*	*	*
2001 4/.....	10.6	7.7	5.6	3.8	4.0	*	*	*	*	*
2000.....	10.9	6.5	3.8	3.5	*	*	*	*	*	*
1999.....	9.8	6.7	5.0	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by hanging, strangulation and suffocation (X91)

All races 3/, both sexes

2004.....	0.4	0.3	0.2	0.1	*	*	*	*	*	*
2003.....	0.3	0.2	0.2	0.1	*	*	*	*	*	*
2002.....	0.3	0.3	0.2	*	0.2	*	*	*	*	*
2001.....	0.3	0.3	0.2	0.1	0.2	0.2	0.3	*	*	*
2000.....	0.3	0.2	0.1	0.2	*	*	*	*	*	*
1999.....	0.3	0.2	0.2	*	*	0.3	*	*	*	*

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	*	0.3	*	*	*	*	*	*	*	*
2001.....	0.2	0.2	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	0.5	0.3	*	*	*	*	*	*	*	*
2003.....	0.4	0.3	0.3	*	*	*	*	*	*	*
2002.....	0.4	0.3	0.3	*	*	*	*	*	*	*
2001.....	0.5	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	0.3	*	*	*	*	*	*	*	*
1999.....	0.4	0.2	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.3	0.2	0.1	*	*	*	*	*	*	*
2003.....	0.3	0.1	0.2	*	*	*	*	*	*	*
2002.....	0.2	0.2	0.2	*	*	*	*	*	*	*
2001.....	0.2	0.2	0.2	*	*	*	0.3	*	*	*
2000.....	0.2	0.2	0.1	*	*	*	*	*	*	*
1999.....	0.2	0.1	0.1	*	*	0.3	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All races 3/, both sexes

2004.....	3.8	2.9	2.2	1.7	1.5	1.2	0.8	0.8	0.7	0.6
2003.....	3.8	3.1	2.2	1.6	1.2	0.9	0.9	0.7	0.6	0.5
2002.....	3.9	2.8	2.2	1.8	1.3	1.0	0.8	1.0	0.8	*
2001.....	3.7	2.7	2.1	1.5	1.4	1.2	0.9	0.8	0.7	*
2000.....	3.8	2.9	2.1	1.7	1.2	1.0	1.0	0.7	0.5	0.7
1999.....	3.7	2.8	2.0	1.6	1.6	1.1	1.0	0.8	0.7	0.6

All races 3/, male

2004.....	5.9	4.4	3.5	2.7	2.2	1.8	1.4	1.0	*	*
2003.....	5.9	5.0	3.5	2.4	1.8	1.4	1.4	0.9	*	*
2002.....	6.2	4.2	3.4	2.7	1.9	1.4	1.0	1.4	1.0	*
2001.....	5.6	4.2	3.2	2.3	2.0	1.8	1.1	1.2	*	*
2000.....	5.8	4.5	3.3	2.6	1.8	1.3	1.3	0.8	*	*
1999.....	5.5	4.4	3.0	2.6	2.5	1.5	1.4	1.1	*	*

All races 3/, female

2004.....	1.6	1.4	0.9	0.7	0.9	0.7	*	0.7	0.8	*
2003.....	1.7	1.4	1.0	0.8	0.6	0.4	0.5	0.6	0.7	*
2002.....	1.7	1.5	1.1	0.9	0.7	0.6	0.6	0.6	0.7	*
2001.....	1.7	1.1	1.0	0.7	0.8	0.7	0.7	0.5	0.7	*
2000.....	1.7	1.3	1.0	0.9	0.7	0.8	0.7	0.7	*	*
1999.....	1.8	1.3	1.1	0.6	0.7	0.7	0.7	0.6	0.6	*

White, both sexes

2004.....	2.5	2.0	1.5	1.3	1.2	1.1	0.7	0.7	0.7	0.5
2003.....	2.4	2.2	1.7	1.2	1.0	0.7	0.8	0.6	0.6	0.5
2002.....	2.6	1.9	1.6	1.5	0.9	0.8	0.7	0.8	0.8	*
2001.....	2.6	1.9	1.5	1.2	1.2	1.0	0.7	0.7	0.7	*
2000.....	2.5	2.0	1.5	1.4	1.0	0.8	0.8	0.7	*	0.6
1999.....	2.5	1.9	1.5	1.2	1.2	1.0	0.9	0.7	0.7	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
White, male										
2004.....	3.5	2.9	2.3	1.9	1.7	1.4	1.0	0.9	*	*
2003.....	3.4	3.1	2.5	1.8	1.4	1.1	1.1	0.7	*	*
2002.....	4.0	2.6	2.4	2.1	1.4	1.0	0.8	1.2	*	*
2001.....	3.7	2.8	2.3	1.6	1.7	1.3	0.8	1.0	*	*
2000.....	3.7	3.0	2.2	2.1	1.4	1.0	0.9	0.7	*	*
1999.....	3.6	2.8	2.3	1.9	1.8	1.3	1.1	0.7	*	*
White, female										
2004.....	1.5	1.1	0.8	0.7	0.8	0.8	*	0.6	0.9	*
2003.....	1.3	1.2	0.9	0.8	0.5	*	0.5	0.5	0.7	*
2002.....	1.3	1.3	0.9	0.9	0.5	0.5	0.6	0.6	0.7	*
2001.....	1.4	0.9	0.8	0.8	0.7	0.8	0.7	0.5	0.7	*
2000.....	1.3	1.1	0.9	0.8	0.7	0.7	0.7	0.7	*	*
1999.....	1.4	1.0	0.7	0.6	0.7	0.7	0.7	0.6	*	*
Black, both sexes										
2004.....	12.8	9.4	7.2	5.2	4.2	2.6	*	*	*	*
2003.....	13.7	10.2	6.8	4.6	3.1	2.4	*	*	*	*
2002.....	13.1	9.6	7.1	4.2	4.0	2.8	*	*	*	*
2001.....	11.6	8.4	6.5	4.3	3.0	2.9	*	*	*	*
2000.....	12.6	9.4	7.0	4.2	2.6	2.8	2.7	*	*	*
1999.....	12.2	9.7	5.9	4.5	4.2	2.2	*	*	*	*
Black, male										
2004.....	23.4	16.5	13.7	9.6	7.1	5.6	*	*	*	*
2003.....	24.3	19.1	11.9	9.1	5.2	*	*	*	*	*
2002.....	22.7	17.4	12.3	8.3	6.5	5.2	*	*	*	*
2001.....	20.3	15.3	11.8	8.6	5.1	6.4	*	*	*	*
2000.....	21.6	16.7	13.1	7.1	5.4	5.2	*	*	*	*
1999.....	20.5	17.3	9.5	9.3	8.3	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
Black, female										
2004.....	3.5	3.2	1.7	*	*	*	*	*	*	*
2003.....	4.4	2.5	2.5	*	*	*	*	*	*	*
2002.....	4.6	2.8	2.8	*	*	*	*	*	*	*
2001.....	3.9	2.4	1.9	*	*	*	*	*	*	*
2000.....	4.7	3.2	*	*	*	*	*	*	*	*
1999.....	4.8	3.2	2.8	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by sharp object (X99)

All races 3/, both sexes

2004.....	1.0	0.9	0.7	0.5	0.4	0.5	0.3	0.4	0.4	*
2003.....	1.0	0.9	0.7	0.5	0.4	0.4	0.5	0.5	0.4	*
2002.....	1.1	0.9	0.6	0.6	0.4	0.4	0.5	0.3	0.4	*
2001.....	0.9	0.8	0.6	0.5	0.3	0.6	0.3	0.5	*	*
2000.....	1.0	0.7	0.5	0.5	0.4	0.4	0.3	0.4	0.5	*
1999.....	1.0	0.8	0.6	0.5	0.4	0.3	0.3	0.4	*	*

All races 3/, male

2004.....	1.4	1.2	1.0	0.7	0.6	0.8	*	*	*	*
2003.....	1.5	1.3	1.0	0.6	0.5	*	0.7	0.7	*	*
2002.....	1.5	1.3	1.0	1.0	0.5	0.5	0.8	*	*	*
2001.....	1.3	1.2	0.9	0.8	*	0.7	*	*	*	*
2000.....	1.4	1.0	0.6	0.8	0.5	0.5	0.6	0.7	*	*
1999.....	1.6	1.1	0.9	0.7	0.6	0.5	*	*	*	*

All races 3/, female

2004.....	0.6	0.6	0.3	0.2	*	*	*	*	*	*
2003.....	0.6	0.5	0.3	0.3	*	*	*	*	*	*
2002.....	0.7	0.4	0.3	0.3	*	*	*	*	*	*
2001.....	0.6	0.4	0.3	0.3	0.3	0.5	*	0.5	*	*
2000.....	0.6	0.4	0.4	*	*	*	*	*	*	*
1999.....	0.4	0.4	0.3	*	*	*	*	*	*	*

White, both sexes

2004.....	0.6	0.6	0.4	0.3	0.3	0.4	*	*	*	*
2003.....	0.6	0.6	0.4	0.3	0.3	0.3	0.3	0.3	*	*
2002.....	0.7	0.5	0.4	0.4	0.3	0.3	0.4	*	*	*
2001.....	0.6	0.5	0.4	0.4	0.3	0.5	*	0.4	*	*
2000.....	0.6	0.4	0.4	0.3	0.3	*	*	0.3	*	*
1999.....	0.6	0.5	0.4	0.4	0.3	0.3	0.3	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by sharp object (X99)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
White, male										
2004.....	0.8	0.7	0.6	0.4	0.4	0.5	*	*	*	*
2003.....	0.8	0.8	0.6	0.4	*	*	*	*	*	*
2002.....	1.0	0.8	0.5	0.6	*	*	*	*	*	*
2001.....	0.8	0.7	0.6	0.6	*	0.5	*	*	*	*
2000.....	0.8	0.5	0.4	0.6	*	*	*	*	*	*
1999.....	1.0	0.7	0.6	0.4	*	*	*	*	*	*
White, female										
2004.....	0.4	0.4	*	*	*	*	*	*	*	*
2003.....	0.4	0.4	0.3	0.3	*	*	*	*	*	*
2002.....	0.5	0.3	*	*	*	*	*	*	*	*
2001.....	0.4	0.3	0.3	*	*	*	*	*	*	*
2000.....	0.3	0.3	0.3	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*
Black, both sexes										
2004.....	3.6	3.4	2.6	1.6	*	*	*	*	*	*
2003.....	4.2	3.1	2.3	1.6	*	*	*	*	*	*
2002.....	3.9	3.1	2.6	2.4	*	*	*	*	*	*
2001.....	3.5	3.0	1.7	1.7	*	*	*	*	*	*
2000.....	3.8	2.8	1.6	1.8	*	*	*	*	*	*
1999.....	3.4	2.8	1.9	1.6	*	*	*	*	*	*
Black, male										
2004.....	5.4	5.2	4.2	3.0	*	*	*	*	*	*
2003.....	6.8	5.1	4.1	2.8	*	*	*	*	*	*
2002.....	5.4	5.3	4.7	4.3	*	*	*	*	*	*
2001.....	5.3	5.4	3.2	*	*	*	*	*	*	*
2000.....	5.1	4.7	2.4	*	*	*	*	*	*	*
1999.....	6.0	4.1	3.1	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

All races 3/, both sexes

2004.....	1.0	1.2	1.1	0.8	0.7	0.7	0.8	0.7	0.7	0.9
2003.....	1.1	1.2	0.9	0.8	0.7	0.8	0.7	0.9	0.9	1.1
2002.....	1.1	1.2	0.8	0.8	0.8	0.8	0.7	0.8	0.7	0.9
2001 4/.....	3.4	3.1	2.5	2.1	1.4	1.1	0.9	0.8	1.0	1.1
2000.....	1.3	1.2	1.0	0.9	0.7	0.7	0.8	0.9	1.3	0.9
1999.....	1.3	1.1	1.0	0.8	0.7	0.8	0.7	0.9	1.0	1.1

All races 3/, male

2004.....	1.3	1.8	1.8	1.2	1.1	1.1	1.1	0.8	1.3	*
2003.....	1.7	1.7	1.5	1.3	1.2	1.3	1.0	1.5	1.5	2.0
2002.....	1.6	1.8	1.2	1.3	1.4	1.2	1.1	1.2	1.1	*
2001 4/.....	5.1	4.5	3.8	3.2	2.2	1.7	1.4	1.1	1.4	1.6
2000.....	1.9	1.8	1.6	1.3	1.3	1.2	1.3	1.1	2.0	*
1999.....	1.9	1.7	1.5	1.3	1.0	1.4	0.9	1.4	1.8	*

All races 3/, female

2004.....	0.7	0.6	0.5	0.4	0.3	*	0.5	0.6	*	0.8
2003.....	0.6	0.7	0.4	0.3	*	*	0.5	0.6	0.6	0.7
2002.....	0.6	0.6	0.4	0.4	*	0.4	*	0.5	*	0.8
2001 4/.....	1.7	1.6	1.2	1.1	0.8	0.6	0.4	0.6	0.8	0.9
2000.....	0.7	0.7	0.4	0.5	*	*	*	0.7	0.8	0.7
1999.....	0.8	0.6	0.5	0.4	*	*	0.5	0.6	*	0.9

White, both sexes

2004.....	0.8	0.9	0.9	0.7	0.5	0.6	0.7	0.6	0.7	0.9
2003.....	0.9	0.9	0.7	0.6	0.6	0.6	0.6	0.9	0.8	1.1
2002.....	0.9	0.9	0.6	0.7	0.7	0.6	0.6	0.6	0.6	0.7
2001 4/.....	3.2	2.8	2.2	2.0	1.3	0.9	0.7	0.7	0.9	1.0
2000.....	0.9	0.9	0.7	0.7	0.5	0.5	0.6	0.7	1.0	0.8
1999.....	0.9	0.9	0.7	0.7	0.6	0.7	0.6	0.8	0.9	0.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

White, male

2004.....	1.1	1.4	1.4	1.0	0.8	0.9	0.9	*	1.0	*
2003.....	1.3	1.2	1.1	1.0	1.0	1.0	0.8	1.3	1.3	1.9
2002.....	1.3	1.3	0.8	1.1	1.2	0.9	1.0	0.8	*	*
2001 4/.....	4.9	4.2	3.3	3.1	1.8	1.5	1.1	0.9	1.2	*
2000.....	1.4	1.1	1.1	1.0	0.9	0.8	1.0	0.9	1.3	*
1999.....	1.4	1.3	1.0	1.1	0.9	1.1	0.7	1.3	1.5	*

White, female

2004.....	0.6	0.5	0.4	0.3	*	*	*	*	*	0.8
2003.....	0.5	0.6	0.4	0.3	*	*	*	0.5	*	0.7
2002.....	0.4	0.5	0.4	0.4	*	*	*	*	*	0.7
2001 4/.....	1.5	1.4	1.0	1.0	0.7	*	*	0.5	0.7	0.9
2000.....	0.5	0.6	0.4	0.4	*	*	*	0.6	0.7	*
1999.....	0.5	0.4	0.5	*	*	*	0.5	*	*	0.9

Black, both sexes

2004.....	2.2	3.1	2.9	2.1	1.7	*	*	*	*	*
2003.....	2.8	3.2	2.5	2.0	2.2	*	*	*	*	*
2002.....	2.5	3.5	2.5	1.7	2.3	*	*	*	*	*
2001 4/.....	4.9	5.4	4.6	3.1	2.9	2.6	*	*	*	*
2000.....	3.6	3.7	3.1	2.2	2.7	2.2	*	*	*	*
1999.....	4.2	3.0	2.7	1.8	*	*	*	*	*	*

Black, male

2004.....	3.0	5.2	5.1	3.3	*	*	*	*	*	*
2003.....	4.7	4.9	4.5	3.9	4.2	*	*	*	*	*
2002.....	3.7	6.1	4.5	3.6	4.6	*	*	*	*	*
2001 4/.....	6.8	7.8	7.0	4.7	5.3	*	*	*	*	*
2000.....	5.5	6.5	6.2	4.3	5.4	*	*	*	*	*
1999.....	6.2	4.7	5.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2004.....	3.4	3.6	2.6	1.7	1.0	0.9	0.8	0.7	1.3	1.6
2003.....	3.9	3.6	2.6	1.6	1.0	0.8	0.7	0.6	0.9	1.1
2002.....	3.4	3.3	2.3	1.4	1.0	0.8	0.7	0.6	1.0	1.4
2001.....	3.2	3.0	2.0	1.1	0.7	0.6	0.5	0.7	1.1	1.0
2000.....	3.1	2.7	1.7	1.1	0.7	0.6	0.6	0.6	0.7	1.3
1999.....	2.9	2.7	1.6	1.0	0.8	0.8	0.5	0.7	0.9	1.4

All races 3/, male

2004.....	4.1	4.5	3.3	2.0	1.1	1.2	0.9	0.8	1.3	1.9
2003.....	4.9	4.6	3.4	2.0	1.2	0.8	0.7	0.8	1.1	1.5
2002.....	4.3	4.1	2.9	1.8	1.3	1.0	1.0	0.8	1.1	2.2
2001.....	4.4	4.2	2.5	1.5	0.8	0.9	0.6	0.9	1.5	1.5
2000.....	4.2	3.8	2.3	1.6	0.9	0.9	0.8	1.0	1.1	1.7
1999.....	4.0	3.7	2.1	1.5	1.1	1.1	0.6	1.0	1.3	2.1

All races 3/, female

2004.....	2.7	2.6	2.0	1.3	0.8	0.7	0.6	0.6	1.2	1.6
2003.....	2.9	2.6	1.8	1.1	0.9	0.7	0.6	*	0.8	0.9
2002.....	2.6	2.6	1.7	1.0	0.7	0.7	0.5	0.5	0.9	1.1
2001.....	2.0	1.8	1.5	0.6	0.5	*	*	0.6	0.8	0.8
2000.....	1.9	1.7	1.1	0.7	0.5	*	*	*	*	1.2
1999.....	1.9	1.7	1.0	0.6	0.6	0.5	*	0.5	0.7	1.1

White, both sexes

2004.....	3.5	3.7	2.6	1.7	1.0	0.8	0.8	0.6	1.3	1.6
2003.....	4.0	3.6	2.5	1.5	0.9	0.7	0.6	0.6	0.9	1.2
2002.....	3.5	3.3	2.2	1.4	0.9	0.8	0.7	0.6	0.9	1.4
2001.....	3.2	2.9	1.9	1.1	0.6	0.6	0.5	0.7	1.0	1.0
2000.....	2.9	2.6	1.6	1.0	0.7	0.5	0.5	0.6	0.8	1.3
1999.....	2.8	2.5	1.4	1.0	0.8	0.8	0.4	0.7	0.9	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)

White, male

2004.....	4.1	4.5	3.1	2.0	1.0	1.0	0.9	*	1.4	1.9
2003.....	5.0	4.5	3.2	1.8	1.0	0.7	0.7	0.9	1.1	1.5
2002.....	4.3	4.0	2.5	1.8	1.2	0.9	0.8	*	1.1	2.3
2001.....	4.2	4.0	2.3	1.4	0.7	0.8	0.6	0.9	1.5	*
2000.....	4.0	3.5	2.1	1.3	0.9	0.8	0.7	1.0	*	*
1999.....	3.8	3.2	1.8	1.3	1.0	1.1	*	1.0	1.3	2.1

White, female

2004.....	2.9	2.9	2.0	1.4	0.9	0.6	0.6	0.6	1.2	1.5
2003.....	3.1	2.8	1.9	1.2	0.8	0.7	0.5	*	0.8	1.0
2002.....	2.6	2.6	1.9	1.1	0.7	0.7	0.5	*	0.7	1.1
2001.....	2.1	1.9	1.6	0.7	0.6	*	*	0.6	0.7	0.8
2000.....	1.9	1.7	1.1	0.7	0.5	*	*	*	*	1.1
1999.....	1.7	1.8	1.0	0.7	0.7	0.5	*	*	0.8	1.1

Black, both sexes

2004.....	3.5	4.1	3.6	1.9	*	*	*	*	*	*
2003.....	4.3	4.4	3.5	2.0	1.9	*	*	*	*	*
2002.....	4.3	4.5	3.8	1.5	*	*	*	*	*	*
2001.....	4.0	4.1	3.3	1.5	*	*	*	*	*	*
2000.....	4.8	4.4	2.8	2.3	*	*	*	*	*	*
1999.....	4.6	4.8	3.3	1.5	*	*	*	*	*	*

Black, male

2004.....	4.6	6.2	5.4	3.2	*	*	*	*	*	*
2003.....	6.2	6.8	5.9	3.5	*	*	*	*	*	*
2002.....	5.5	6.2	6.5	*	*	*	*	*	*	*
2001.....	6.6	7.0	5.4	*	*	*	*	*	*	*
2000.....	7.2	7.0	4.4	4.0	*	*	*	*	*	*
1999.....	6.5	8.3	5.7	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

All races 3/, both sexes

2004.....	2.6	2.8	1.9	1.1	0.5	0.3	*	*	*	*
2003.....	3.2	2.8	1.9	1.0	0.5	0.3	*	*	*	*
2002.....	2.7	2.6	1.7	0.8	0.5	0.3	*	*	*	*
2001.....	2.5	2.2	1.4	0.6	0.3	*	*	*	*	*
2000.....	2.4	2.1	1.1	0.5	0.3	*	*	*	*	*
1999.....	2.3	2.0	1.0	0.6	0.3	0.3	*	*	*	*

All races 3/, male

2004.....	3.1	3.4	2.0	1.1	0.5	*	*	*	*	*
2003.....	3.9	3.5	2.2	1.2	0.5	*	*	*	*	*
2002.....	3.2	3.0	1.9	0.9	0.6	*	*	*	*	*
2001.....	3.3	3.2	1.7	0.7	*	*	*	*	*	*
2000.....	3.3	2.8	1.5	0.7	*	*	*	*	*	*
1999.....	3.1	2.6	1.3	0.7	*	*	*	*	*	*

All races 3/, female

2004.....	2.2	2.2	1.7	1.1	0.5	*	*	*	*	*
2003.....	2.5	2.1	1.5	0.9	0.5	*	*	*	*	*
2002.....	2.2	2.2	1.5	0.7	0.4	*	*	*	*	*
2001.....	1.7	1.3	1.1	0.5	0.3	*	*	*	*	*
2000.....	1.6	1.4	0.8	0.4	*	*	*	*	*	*
1999.....	1.6	1.4	0.7	0.4	*	*	*	*	*	*

White, both sexes

2004.....	2.8	2.9	1.9	1.1	0.5	0.3	*	*	*	*
2003.....	3.4	2.9	1.8	1.0	0.5	0.3	*	*	*	*
2002.....	2.7	2.6	1.6	0.8	0.5	0.3	*	*	*	*
2001.....	2.5	2.2	1.4	0.6	0.3	*	*	*	*	*
2000.....	2.3	2.0	1.1	0.5	0.3	*	*	*	*	*
1999.....	2.2	1.9	0.9	0.5	0.3	0.3	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

White, male

2004.....	3.1	3.4	2.0	1.0	0.4	*	*	*	*	*
2003.....	4.0	3.5	2.1	1.1	0.4	*	*	*	*	*
2002.....	3.3	3.0	1.7	0.8	0.6	*	*	*	*	*
2001.....	3.2	3.0	1.6	0.7	*	*	*	*	*	*
2000.....	3.1	2.7	1.4	0.6	*	*	*	*	*	*
1999.....	2.9	2.2	1.1	0.6	*	*	*	*	*	*

White, female

2004.....	2.4	2.4	1.8	1.2	0.6	*	*	*	*	*
2003.....	2.7	2.2	1.6	1.0	0.5	*	*	*	*	*
2002.....	2.2	2.2	1.6	0.7	0.5	*	*	*	*	*
2001.....	1.9	1.5	1.2	0.6	0.4	*	*	*	*	*
2000.....	1.5	1.3	0.8	0.4	*	*	*	*	*	*
1999.....	1.5	1.5	0.7	0.5	*	*	*	*	*	*

Black, both sexes

2004.....	2.6	3.0	2.2	1.2	*	*	*	*	*	*
2003.....	3.4	3.5	2.5	*	*	*	*	*	*	*
2002.....	3.4	3.4	2.9	*	*	*	*	*	*	*
2001.....	3.3	3.0	2.4	*	*	*	*	*	*	*
2000.....	3.8	3.3	1.9	*	*	*	*	*	*	*
1999.....	3.6	3.6	2.3	*	*	*	*	*	*	*

Black, male

2004.....	3.3	4.2	2.9	*	*	*	*	*	*	*
2003.....	4.8	5.5	4.1	*	*	*	*	*	*	*
2002.....	4.0	4.2	4.8	*	*	*	*	*	*	*
2001.....	5.6	5.2	3.7	*	*	*	*	*	*	*
2000.....	5.6	4.9	2.8	*	*	*	*	*	*	*
1999.....	4.8	6.0	3.7	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2004.....	0.5	0.6	0.6	0.4	0.3	0.4	0.4	0.5	1.0	1.3
2003.....	0.5	0.6	0.5	0.4	0.4	0.3	0.4	0.3	0.6	0.8
2002.....	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.6	1.1
2001.....	0.5	0.5	0.4	0.3	0.2	0.3	0.2	0.4	0.7	0.7
2000.....	0.4	0.5	0.4	0.5	0.3	0.3	0.4	0.3	0.5	0.9
1999.....	0.4	0.5	0.4	0.4	0.3	0.4	0.2	0.4	0.5	1.0

All races 3/, male

2004.....	0.8	0.9	0.9	0.7	0.4	0.6	*	*	1.0	1.4
2003.....	0.7	0.8	0.9	0.7	0.5	*	*	*	*	*
2002.....	0.7	0.8	0.7	0.7	0.6	0.7	0.5	*	*	1.7
2001.....	0.7	0.7	0.6	0.6	*	0.5	*	*	1.1	*
2000.....	0.6	0.8	0.6	0.6	0.5	0.5	0.5	*	*	*
1999.....	0.6	0.8	0.6	0.7	0.5	0.5	*	*	*	*

All races 3/, female

2004.....	0.3	0.3	0.3	*	*	*	*	*	0.9	1.3
2003.....	0.3	0.3	0.2	*	*	*	0.4	*	*	0.7
2002.....	0.4	0.3	*	*	*	*	*	*	*	0.8
2001.....	0.2	0.3	0.3	*	*	*	*	*	*	*
2000.....	0.3	0.2	0.3	*	*	*	*	*	*	0.9
1999.....	0.2	*	*	*	*	*	*	*	*	0.8

White, both sexes

2004.....	0.5	0.6	0.5	0.4	0.3	0.3	0.3	0.4	1.0	1.3
2003.....	0.5	0.6	0.5	0.3	0.3	0.3	0.4	0.3	0.6	0.9
2002.....	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.4	0.6	1.1
2001.....	0.4	0.5	0.4	0.3	*	0.3	*	0.4	0.7	0.7
2000.....	0.4	0.4	0.4	0.4	0.2	0.3	0.3	0.3	0.6	0.8
1999.....	0.3	0.4	0.3	0.4	0.3	0.3	*	0.4	0.5	0.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Legal intervention (Y35,Y89.0)

All races 3/, both sexes

2004.....	0.2	0.1	0.1	*	*	*	*	*	*	*
2003.....	0.2	0.1	0.1	*	*	*	*	*	*	*
2002.....	0.2	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	0.2	*	*	*	*	*	*	*

All races 3/, male

2004.....	0.4	0.3	0.3	*	*	*	*	*	*	*
2003.....	0.4	0.3	*	*	*	*	*	*	*	*
2002.....	0.4	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	0.4	*	*	*	*	*	*	*	*
1999.....	0.4	*	0.4	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	0.1	*	*	*	*	*	*	*	*
2003.....	0.2	0.1	*	*	*	*	*	*	*	*
2002.....	0.2	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes

2004.....	0.4	0.6	0.8	1.2	1.6	2.6	3.5	5.0	7.6	10.6
2003.....	0.5	0.7	0.7	1.1	1.7	2.2	3.7	5.3	7.3	11.2
2002.....	0.4	0.5	0.8	1.1	1.6	2.3	3.9	5.1	7.7	11.3
2001.....	0.4	0.7	0.8	1.3	1.9	2.6	3.6	5.9	8.3	12.8
2000.....	0.4	0.6	0.9	1.2	2.0	2.8	4.3	5.8	8.0	13.6
1999.....	0.4	0.5	0.7	1.1	1.7	2.4	4.0	5.5	8.4	12.7

All races 3/, male

2004.....	0.4	0.6	0.8	1.5	1.7	2.5	4.3	5.8	9.6	11.9
2003.....	0.5	0.7	0.7	1.3	1.6	2.3	4.2	5.6	8.8	12.9
2002.....	0.4	0.6	0.7	1.2	1.5	2.5	3.8	5.4	7.7	12.3
2001.....	0.4	0.7	0.8	1.3	2.0	2.6	3.8	6.3	9.9	14.5
2000.....	0.4	0.6	1.0	1.2	2.0	2.8	5.3	7.0	10.4	16.0
1999.....	0.4	0.4	0.7	1.2	1.7	2.3	4.5	6.4	10.1	16.5

All races 3/, female

2004.....	0.5	0.6	0.8	1.0	1.5	2.7	2.9	4.3	6.3	9.9
2003.....	0.4	0.6	0.6	0.9	1.7	2.1	3.3	5.1	6.4	10.5
2002.....	0.4	0.5	0.9	1.0	1.7	2.2	3.9	5.0	7.7	10.9
2001.....	0.5	0.6	0.8	1.2	1.7	2.6	3.5	5.7	7.3	12.1
2000.....	0.4	0.7	0.9	1.1	2.0	2.7	3.6	4.9	6.7	12.7
1999.....	0.4	0.5	0.7	0.9	1.6	2.5	3.6	4.8	7.4	11.2

White, both sexes

2004.....	0.4	0.6	0.8	1.1	1.5	2.4	3.4	4.8	7.3	10.4
2003.....	0.4	0.6	0.6	1.0	1.6	2.1	3.6	5.1	7.4	11.0
2002.....	0.4	0.5	0.7	1.0	1.5	2.2	3.8	5.0	7.6	11.2
2001.....	0.4	0.6	0.7	1.1	1.6	2.4	3.2	5.8	8.2	12.1
2000.....	0.3	0.6	0.9	1.0	1.8	2.6	4.1	5.4	8.0	13.1
1999.....	0.3	0.4	0.6	0.9	1.5	2.3	3.8	5.2	8.0	12.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Complications of medical and surgical care (Y40-Y84,Y88)

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
White, male										
2004.....	0.3	0.6	0.8	1.3	1.7	2.5	4.2	5.8	9.5	11.8
2003.....	0.6	0.6	0.5	1.2	1.6	2.3	4.1	5.5	9.1	12.8
2002.....	0.4	0.5	0.6	1.0	1.5	2.2	3.7	5.3	7.8	11.8
2001.....	0.3	0.6	0.6	1.1	1.7	2.5	3.2	6.2	10.0	13.6
2000.....	0.4	0.5	1.0	1.1	1.8	2.5	5.0	6.8	10.6	15.8
1999.....	0.3	0.3	0.6	1.0	1.6	2.3	4.2	6.2	9.7	15.6
White, female										
2004.....	0.4	0.5	0.7	0.9	1.4	2.4	2.7	4.0	6.0	9.8
2003.....	0.3	0.5	0.6	0.8	1.6	2.0	3.1	4.7	6.3	10.2
2002.....	0.3	0.4	0.7	0.9	1.5	2.1	3.9	4.8	7.5	10.9
2001.....	0.4	0.5	0.7	1.1	1.6	2.4	3.2	5.6	7.0	11.5
2000.....	0.3	0.6	0.9	0.9	1.8	2.7	3.3	4.5	6.4	12.0
1999.....	0.4	0.5	0.6	0.7	1.5	2.4	3.4	4.5	7.1	10.7
Black, both sexes										
2004.....	0.9	1.0	1.4	2.7	2.5	4.5	5.5	7.6	11.9	13.8
2003.....	*	1.3	1.6	2.5	3.0	3.6	5.8	9.5	7.6	15.2
2002.....	1.0	1.2	2.1	2.6	2.8	4.0	5.2	7.6	9.4	14.0
2001.....	1.0	1.6	1.5	2.7	4.2	4.6	8.5	8.0	11.1	23.3
2000.....	*	1.1	1.7	2.6	4.7	4.6	7.9	10.5	10.4	21.0
1999.....	0.8	0.9	1.8	2.6	2.6	3.5	7.3	9.5	12.4	20.1
Black, male										
2004.....	*	*	*	3.4	*	*	*	*	*	*
2003.....	*	*	2.1	2.8	*	*	*	*	*	*
2002.....	*	*	*	*	*	5.0	*	*	*	*
2001.....	*	1.7	*	3.4	5.1	5.3	11.1	*	*	29.5
2000.....	*	*	*	*	4.6	6.2	9.4	10.4	*	*
1999.....	*	*	*	*	*	*	9.2	10.4	*	23.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)										
Black, female										
2004.....	*	*	1.7	*	*	5.4	5.1	7.5	11.0	13.4
2003.....	*	*	*	*	3.0	3.8	6.4	10.6	7.7	15.4
2002.....	*	*	2.5	2.4	3.1	*	4.9	8.2	9.9	11.5
2001.....	*	*	*	*	3.5	4.1	6.7	8.0	11.0	21.0
2000.....	*	*	*	3.1	4.7	*	6.9	10.5	10.6	21.9
1999.....	*	*	*	*	*	*	6.1	8.9	10.8	18.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)

All races 3/, both sexes

2004.....	*	*	*	*	*	*	*	*	0.6	*
2003.....	*	*	*	*	*	*	*	*	*	0.6
2002.....	*	*	*	*	*	*	*	0.3	0.4	0.5
2001.....	*	*	*	*	*	0.2	*	0.4	0.4	0.5
2000.....	*	*	*	*	*	0.2	0.3	0.3	*	0.6
1999.....	*	*	*	*	*	*	0.3	0.3	0.5	*

All races 3/, male

2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	0.6	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	0.5	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	*	*	*	*	*	*	*	*	0.6	*
2003.....	*	*	*	*	*	*	*	*	*	0.6
2002.....	*	*	*	*	*	*	*	*	0.4	*
2001.....	*	*	*	*	*	*	*	0.3	0.5	0.5
2000.....	*	*	*	*	*	*	*	0.3	*	0.6
1999.....	*	*	*	*	*	*	*	0.3	0.5	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

All races 3/, both sexes

2004.....	0.3	0.4	0.7	1.0	1.2	2.2	3.1	4.1	6.2	9.3
2003.....	0.3	0.5	0.5	0.9	1.3	1.8	3.3	4.7	6.5	9.9
2002.....	0.3	0.4	0.7	0.8	1.4	2.0	3.3	4.6	6.8	10.4
2001.....	0.3	0.5	0.6	1.1	1.6	2.2	3.2	5.2	7.4	11.4
2000.....	0.3	0.5	0.8	0.9	1.8	2.4	3.8	5.0	7.0	12.2
1999.....	0.3	0.3	0.6	0.8	1.4	2.0	3.5	4.9	7.5	11.8

All races 3/, male

2004.....	0.3	0.4	0.7	1.2	1.4	2.2	3.8	4.7	8.4	10.7
2003.....	0.4	0.6	0.6	1.1	1.2	1.9	3.7	4.9	8.1	11.5
2002.....	0.3	0.4	0.6	0.9	1.3	2.1	3.3	4.9	7.0	11.7
2001.....	0.2	0.6	0.6	1.2	1.6	2.3	3.4	5.5	8.8	13.2
2000.....	0.3	0.5	0.8	1.0	1.8	2.5	4.7	6.0	8.9	14.3
1999.....	0.3	0.3	0.6	1.0	1.5	2.0	4.0	5.8	9.1	15.4

All races 3/, female

2004.....	0.3	0.4	0.6	0.8	1.0	2.2	2.5	3.7	4.9	8.7
2003.....	0.3	0.4	0.5	0.8	1.4	1.6	3.0	4.5	5.5	9.2
2002.....	0.4	0.4	0.7	0.8	1.5	1.9	3.3	4.4	6.7	9.9
2001.....	0.4	0.5	0.6	1.1	1.6	2.1	3.0	4.9	6.5	10.6
2000.....	0.3	0.5	0.7	0.9	1.8	2.3	3.1	4.4	5.9	11.4
1999.....	0.3	0.4	0.5	0.7	1.3	2.1	3.1	4.3	6.6	10.4

White, both sexes

2004.....	0.3	0.4	0.6	0.9	1.2	2.1	3.0	4.0	6.1	9.1
2003.....	0.3	0.4	0.4	0.8	1.3	1.7	3.2	4.4	6.6	9.6
2002.....	0.3	0.4	0.6	0.7	1.3	1.8	3.3	4.5	6.7	10.2
2001.....	0.3	0.5	0.5	1.0	1.4	2.0	2.8	5.1	7.2	10.7
2000.....	0.3	0.5	0.8	0.8	1.7	2.3	3.6	4.7	7.0	11.8
1999.....	0.3	0.3	0.5	0.7	1.3	2.0	3.4	4.7	7.2	11.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

White, male

2004.....	0.2	0.4	0.7	1.1	1.4	2.2	3.7	4.7	8.4	10.4
2003.....	0.4	0.5	0.4	1.0	1.3	2.0	3.6	4.8	8.4	11.5
2002.....	0.3	0.4	0.5	0.8	1.3	1.8	3.2	4.9	7.1	11.2
2001.....	*	0.6	0.5	1.0	1.4	2.2	2.9	5.5	8.9	12.3
2000.....	0.3	0.4	0.8	0.9	1.6	2.3	4.5	5.8	9.1	14.0
1999.....	0.2	*	0.5	0.7	1.5	1.9	3.8	5.6	8.7	14.5

White, female

2004.....	0.3	0.3	0.5	0.7	1.0	1.9	2.3	3.4	4.7	8.4
2003.....	0.2	0.4	0.4	0.6	1.3	1.5	2.8	4.2	5.5	8.8
2002.....	0.2	0.4	0.6	0.7	1.4	1.8	3.4	4.3	6.4	9.8
2001.....	0.4	0.4	0.6	1.0	1.5	1.9	2.8	4.9	6.2	10.0
2000.....	*	0.5	0.7	0.7	1.7	2.2	2.9	3.9	5.7	10.8
1999.....	0.3	0.4	0.5	0.6	1.2	2.0	3.0	4.0	6.2	9.9

Black, both sexes

2004.....	*	*	1.2	2.2	1.7	3.7	5.1	6.9	9.4	13.0
2003.....	*	1.0	1.3	2.1	2.1	2.9	5.2	8.7	6.6	14.0
2002.....	0.9	0.9	1.7	1.8	2.3	3.8	4.2	6.7	9.2	13.4
2001.....	0.8	1.0	1.3	2.5	3.6	3.8	7.3	6.4	10.3	22.1
2000.....	*	0.8	1.2	2.4	3.6	4.0	6.7	9.8	9.0	19.1
1999.....	*	*	1.5	2.2	2.5	2.9	6.0	8.4	11.0	19.1

Black, male

2004.....	*	*	*	2.6	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	3.3	4.7	*	9.4	*	*	28.4
2000.....	*	*	*	*	*	5.2	7.7	9.9	*	*
1999.....	*	*	*	*	*	*	8.1	9.5	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

Black, female

2004.....	*	*	*	*	*	4.5	4.8	7.0	8.7	12.6
2003.....	*	*	*	*	*	*	6.0	9.5	*	15.0
2002.....	*	*	1.8	*	*	*	*	7.1	9.5	11.1
2001.....	*	*	*	*	*	*	5.8	6.4	10.2	19.8
2000.....	*	*	*	2.8	3.4	*	6.0	9.7	8.9	19.3
1999.....	*	*	*	*	*	*	4.5	7.7	9.5	18.0

* Figure does not meet standards of reliability and precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Includes races other than white and black.

4/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.