

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
All origins, both sexes													
2003.....	2,448,288	28,025	4,965	6,954	33,568	41,300	89,461	176,781	262,519	413,497	703,024	687,852	342
2002.....	2,443,387	28,034	4,858	7,150	33,046	41,355	91,140	172,385	253,342	422,990	707,654	681,076	357
2001.....	2,416,425	27,568	5,107	7,095	32,252	41,683	91,674	168,065	244,139	430,960	701,929	665,531	422
2000.....	2,403,351	28,035	4,979	7,413	31,307	40,451	89,798	160,341	240,846	441,209	700,445	658,171	356
1999.....	2,391,399	27,937	5,249	7,595	30,656	41,066	89,256	152,974	238,979	452,600	698,590	646,141	356
All origins, male													
2003.....	1,201,964	15,902	2,826	4,150	24,670	28,602	56,435	110,682	156,461	231,421	342,332	228,212	271
2002.....	1,199,264	15,717	2,806	4,198	24,416	28,736	57,593	107,722	151,363	237,021	343,504	225,906	282
2001.....	1,183,421	15,477	2,899	4,168	23,963	28,757	58,164	104,848	144,958	241,581	340,742	217,533	331
2000.....	1,177,578	15,718	2,824	4,401	23,071	27,890	57,297	100,398	143,321	247,408	340,219	214,742	289
1999.....	1,175,460	15,646	2,975	4,492	22,414	28,276	57,118	95,659	142,724	254,920	340,970	209,989	277
All origins, female													
2003.....	1,246,324	12,123	2,139	2,804	8,898	12,698	33,026	66,099	106,058	182,076	360,692	459,640	71
2002.....	1,244,123	12,317	2,052	2,952	8,630	12,619	33,547	64,663	101,979	185,969	364,150	455,170	75
2001.....	1,233,004	12,091	2,208	2,927	8,289	12,926	33,510	63,217	99,181	189,379	361,187	447,998	91
2000.....	1,225,773	12,317	2,155	3,012	8,236	12,561	32,501	59,943	97,525	193,801	360,226	443,429	67
1999.....	1,215,939	12,291	2,274	3,103	8,242	12,790	32,138	57,315	96,255	197,680	357,620	436,152	79

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
Hispanic, both sexes													
2003.....	122,026	5,281	993	1,173	5,394	6,189	8,576	12,516	14,913	20,596	26,219	20,129	47
2002.....	117,135	4,943	933	1,124	5,255	5,816	8,467	11,745	14,215	20,476	24,838	19,265	58
2001.....	113,413	4,680	917	1,038	5,058	5,797	8,441	11,362	13,337	20,172	23,637	18,907	67
2000.....	107,254	4,598	872	1,065	4,793	5,509	8,085	10,356	12,733	19,420	21,951	17,818	54
1999.....	103,740	4,412	882	1,014	4,505	5,397	7,798	9,876	12,319	19,201	20,912	17,358	66
Hispanic, male													
2003.....	68,119	2,940	569	706	4,244	4,742	5,887	8,336	9,122	11,465	12,904	7,165	39
2002.....	65,703	2,746	546	647	4,182	4,474	5,828	7,719	8,762	11,476	12,291	6,977	55
2001.....	63,317	2,601	518	600	4,019	4,366	5,895	7,477	8,026	11,231	11,759	6,767	58
2000.....	60,172	2,514	475	622	3,837	4,200	5,597	6,808	7,760	10,842	10,973	6,493	51
1999.....	57,991	2,411	481	592	3,547	4,084	5,476	6,497	7,490	10,783	10,317	6,253	60
Hispanic, female													
2003.....	53,907	2,341	424	467	1,150	1,447	2,689	4,180	5,791	9,131	13,315	12,964	8
2002.....	51,432	2,197	387	477	1,073	1,342	2,639	4,026	5,453	9,000	12,547	12,288	3
2001.....	50,096	2,079	399	438	1,039	1,431	2,546	3,885	5,311	8,941	11,878	12,140	9
2000.....	47,082	2,084	397	443	956	1,309	2,488	3,548	4,973	8,578	10,978	11,325	3
1999.....	45,749	2,001	401	422	958	1,313	2,322	3,379	4,829	8,418	10,595	11,105	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
Non-Hispanic 1/, both sexes													
2003.....	2,319,476	22,475	3,956	5,761	28,044	34,955	80,503	163,533	246,665	391,778	675,309	666,350	147
2002.....	2,318,269	22,816	3,904	6,002	27,658	35,329	82,225	159,843	238,185	401,182	680,878	660,078	169
2001.....	2,295,244	22,629	4,171	6,033	27,059	35,656	82,763	155,916	229,812	409,500	676,512	645,020	173
2000.....	2,287,846	23,074	4,086	6,316	26,374	34,741	81,182	149,141	227,118	420,369	676,632	638,673	140
1999.....	2,279,325	23,126	4,343	6,560	26,043	35,471	80,949	142,302	225,659	431,952	675,730	627,035	155
Non-Hispanic 1/, male													
2003.....	1,129,927	12,799	2,247	3,432	20,328	23,740	50,284	101,828	146,696	219,240	328,640	220,575	118
2002.....	1,129,090	12,815	2,250	3,534	20,129	24,114	51,451	99,433	141,966	224,706	330,191	218,367	134
2001.....	1,115,683	12,739	2,371	3,554	19,836	24,225	51,918	96,815	136,214	229,539	328,079	210,256	137
2000.....	1,112,704	13,007	2,338	3,770	19,136	23,548	51,304	92,984	134,890	235,648	328,284	207,688	107
1999.....	1,112,718	12,998	2,479	3,885	18,797	24,047	51,272	88,567	134,540	243,210	329,682	203,128	113
Non-Hispanic 1/, female													
2003.....	1,189,549	9,676	1,709	2,329	7,716	11,215	30,219	61,705	99,969	172,538	346,669	445,775	29
2002.....	1,189,179	10,001	1,654	2,468	7,529	11,215	30,774	60,410	96,219	176,476	350,687	441,711	35
2001.....	1,179,561	9,890	1,800	2,479	7,223	11,431	30,845	59,101	93,598	179,961	348,433	434,764	36
2000.....	1,175,142	10,067	1,748	2,546	7,238	11,193	29,878	56,157	92,228	184,721	348,348	430,985	33
1999.....	1,166,607	10,128	1,864	2,675	7,246	11,424	29,677	53,735	91,119	188,742	346,048	423,907	42

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
Non-Hispanic white, both sexes													
2003.....	1,979,465	13,223	2,539	3,884	19,640	23,557	58,095	121,832	197,598	329,570	600,563	608,843	121
2002.....	1,981,973	13,463	2,502	4,031	19,458	23,758	59,529	119,812	190,843	338,961	606,431	603,063	122
2001.....	1,962,810	13,311	2,772	4,151	18,730	24,191	59,546	117,086	184,109	346,389	603,143	589,269	113
2000.....	1,959,919	13,513	2,641	4,397	18,261	23,191	58,231	111,964	182,348	356,869	604,658	583,757	89
1999.....	1,953,197	13,553	2,819	4,488	17,868	23,986	57,630	106,808	180,793	367,490	604,344	573,327	91
Non-Hispanic white, male													
2003.....	956,194	7,591	1,423	2,319	14,022	16,097	37,123	77,106	118,383	186,112	293,774	202,148	96
2002.....	957,645	7,706	1,431	2,365	13,957	16,339	38,049	75,724	114,535	191,844	295,763	199,834	98
2001.....	945,967	7,508	1,592	2,428	13,542	16,613	38,213	73,731	109,993	196,158	294,013	192,084	92
2000.....	944,781	7,660	1,541	2,652	13,066	15,934	37,622	70,788	109,031	202,038	294,842	189,537	70
1999.....	944,913	7,718	1,605	2,643	12,677	16,449	37,291	67,151	108,692	209,155	296,528	184,938	66
Non-Hispanic white, female													
2003.....	1,023,271	5,632	1,116	1,565	5,618	7,460	20,972	44,726	79,215	143,458	306,789	406,695	25
2002.....	1,024,328	5,757	1,071	1,666	5,501	7,419	21,480	44,088	76,308	147,117	310,668	403,229	24
2001.....	1,016,843	5,803	1,180	1,723	5,188	7,578	21,333	43,355	74,116	150,231	309,130	397,185	21
2000.....	1,015,138	5,853	1,100	1,745	5,195	7,257	20,609	41,176	73,317	154,831	309,816	394,220	19
1999.....	1,008,284	5,835	1,214	1,845	5,191	7,537	20,339	39,657	72,101	158,335	307,816	388,389	25

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
Non-Hispanic black, both sexes													
2003.....	287,968	8,159	1,169	1,530	7,021	9,795	19,561	36,584	42,350	52,583	61,770	47,426	20
2002.....	286,573	8,284	1,163	1,634	6,904	10,009	19,955	35,211	40,878	52,748	62,183	47,565	39
2001.....	284,343	8,277	1,156	1,557	7,024	9,859	20,436	34,098	39,523	53,803	61,592	46,968	50
2000.....	282,676	8,525	1,217	1,620	6,926	10,103	20,279	32,911	38,867	54,645	60,874	46,660	49
1999.....	281,979	8,589	1,283	1,771	6,980	9,992	20,770	31,377	39,063	55,567	60,634	45,895	58
Non-Hispanic black, male													
2003.....	146,136	4,601	681	911	5,343	6,592	11,405	21,708	24,468	27,957	28,297	14,156	17
2002.....	144,802	4,520	681	977	5,248	6,723	11,661	20,839	23,706	27,859	28,137	14,422	29
2001.....	143,971	4,651	655	941	5,377	6,548	11,971	20,307	22,600	28,446	28,083	14,352	40
2000.....	143,297	4,757	672	959	5,224	6,649	12,061	19,647	22,402	28,745	27,715	14,431	35
1999.....	143,883	4,765	734	1,081	5,286	6,633	12,384	19,026	22,493	29,259	27,552	14,628	42
Non-Hispanic black, female													
2003.....	141,832	3,558	488	619	1,678	3,203	8,156	14,876	17,882	24,626	33,473	33,270	3
2002.....	141,771	3,764	482	657	1,656	3,286	8,294	14,372	17,172	24,889	34,046	33,143	10
2001.....	140,372	3,626	501	616	1,647	3,311	8,465	13,791	16,923	25,357	33,509	32,616	10
2000.....	139,379	3,768	545	661	1,702	3,454	8,218	13,264	16,465	25,900	33,159	32,229	14
1999.....	138,096	3,824	549	690	1,694	3,359	8,386	12,351	16,570	26,308	33,082	31,267	16

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All causes													
Origin not stated 2/, both sexes													
2003.....	6,786	269	16	20	130	156	382	732	941	1,123	1,496	1,373	148
2002.....	7,983	275	21	24	133	210	448	797	942	1,332	1,938	1,733	130
2001.....	7,768	259	19	24	135	230	470	787	990	1,288	1,780	1,604	182
2000.....	8,251	363	21	32	140	201	531	844	995	1,420	1,862	1,680	162
1999.....	8,334	399	24	21	108	198	509	796	1,001	1,447	1,948	1,748	135
Origin not stated 2/, male													
2003.....	3,918	163	10	12	98	120	264	518	643	716	788	472	114
2002.....	4,471	156	10	17	105	148	314	570	635	839	1,022	562	93
2001.....	4,421	137	10	14	108	166	351	556	718	811	904	510	136
2000.....	4,702	197	11	9	98	142	396	606	671	918	962	561	131
1999.....	4,751	237	15	15	70	145	370	595	694	927	971	608	104
Origin not stated 2/, female													
2003.....	2,868	106	6	8	32	36	118	214	298	407	708	901	34
2002.....	3,512	119	11	7	28	62	134	227	307	493	916	1,171	37
2001.....	3,347	122	9	10	27	64	119	231	272	477	876	1,094	46
2000.....	3,549	166	10	23	42	59	135	238	324	502	900	1,119	31
1999.....	3,583	162	9	6	38	53	139	201	307	520	977	1,140	31

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
All origins, both sexes													
2003.....	43	3	-	1	-	1	3	2	6	8	10	9	-
2002.....	21	3	-	-	-	-	2	3	-	4	8	1	-
2001.....	41	2	-	-	-	2	1	3	3	7	14	9	-
2000.....	28	3	2	-	1	1	-	4	2	7	5	3	-
1999.....	38	3	-	-	2	-	2	2	3	7	12	7	-
All origins, male													
2003.....	23	1	-	-	-	-	2	2	4	4	4	6	-
2002.....	10	1	-	-	-	-	-	1	-	2	5	1	-
2001.....	21	-	-	-	-	1	1	3	2	6	7	1	-
2000.....	15	2	-	-	1	1	-	2	2	5	1	1	-
1999.....	21	2	-	-	2	-	1	2	2	5	5	2	-
All origins, female													
2003.....	20	2	-	1	-	1	1	-	2	4	6	3	-
2002.....	11	2	-	-	-	-	2	2	-	2	3	-	-
2001.....	20	2	-	-	-	1	-	-	1	1	7	8	-
2000.....	13	1	2	-	-	-	-	2	-	2	4	2	-
1999.....	17	1	-	-	-	-	1	-	1	2	7	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Salmonella infections (A01-A02)

Hispanic, both sexes													
2003.....	6	-	-	1	-	-	1	1	-	2	1	-	-
2002.....	3	-	-	-	-	-	-	1	-	2	-	-	-
2001.....	5	-	-	-	-	-	-	1	1	1	2	-	-
2000.....	2	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	4	-	-	-	-	-	-	-	1	1	2	-	-
Hispanic, male													
2003.....	3	-	-	-	-	-	1	1	-	1	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2001.....	5	-	-	-	-	-	-	1	1	1	2	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	-	-	-	1	1	1	-	-
Hispanic, female													
2003.....	3	-	-	1	-	-	-	-	-	1	1	-	-
2002.....	2	-	-	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
Non-Hispanic 1/, both sexes													
2003.....	37	3	-	-	-	1	2	1	6	6	9	9	-
2002.....	18	3	-	-	-	-	2	2	-	2	8	1	-
2001.....	36	2	-	-	-	2	1	2	2	6	12	9	-
2000.....	26	1	2	-	1	1	-	4	2	7	5	3	-
1999.....	34	3	-	-	2	-	2	2	2	6	10	7	-
Non-Hispanic 1/, male													
2003.....	20	1	-	-	-	-	1	1	4	3	4	6	-
2002.....	9	1	-	-	-	-	-	1	-	1	5	1	-
2001.....	16	-	-	-	-	1	1	2	1	5	5	1	-
2000.....	14	1	-	-	1	1	-	2	2	5	1	1	-
1999.....	18	2	-	-	2	-	1	2	1	4	4	2	-
Non-Hispanic 1/, female													
2003.....	17	2	-	-	-	1	1	-	2	3	5	3	-
2002.....	9	2	-	-	-	-	2	1	-	1	3	-	-
2001.....	20	2	-	-	-	1	-	-	1	1	7	8	-
2000.....	12	-	2	-	-	-	-	2	-	2	4	2	-
1999.....	16	1	-	-	-	-	1	-	1	2	6	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
Non-Hispanic white, both sexes													
2003.....	25	1	-	-	-	-	-	1	5	4	6	8	-
2002.....	15	2	-	-	-	-	2	2	-	1	7	1	-
2001.....	28	1	-	-	-	2	-	2	2	3	10	8	-
2000.....	23	-	2	-	-	1	-	4	2	6	5	3	-
1999.....	28	3	-	-	1	-	2	2	1	4	8	7	-
Non-Hispanic white, male													
2003.....	14	-	-	-	-	-	-	1	3	3	2	5	-
2002.....	8	1	-	-	-	-	-	1	-	-	5	1	-
2001.....	13	-	-	-	-	1	-	2	1	3	5	1	-
2000.....	12	-	-	-	-	1	-	2	2	5	1	1	-
1999.....	13	2	-	-	1	-	1	2	-	2	3	2	-
Non-Hispanic white, female													
2003.....	11	1	-	-	-	-	-	-	2	1	4	3	-
2002.....	7	1	-	-	-	-	2	1	-	1	2	-	-
2001.....	15	1	-	-	-	1	-	-	1	-	5	7	-
2000.....	11	-	2	-	-	-	-	2	-	1	4	2	-
1999.....	15	1	-	-	-	-	1	-	1	2	5	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Salmonella infections (A01-A02)													
Non-Hispanic black, both sexes													
2003.....	8	2	-	-	-	1	1	-	-	1	3	-	-
2002.....	3	1	-	-	-	-	-	-	-	1	1	-	-
2001.....	5	1	-	-	-	-	-	-	-	1	2	1	-
2000.....	2	1	-	-	1	-	-	-	-	-	-	-	-
1999.....	4	-	-	-	1	-	-	-	-	1	2	-	-
Non-Hispanic black, male													
2003.....	3	1	-	-	-	-	-	-	-	-	2	-	-
2002.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	2	1	-	-	1	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	1	-	-	-	-	1	1	-	-
Non-Hispanic black, female													
2003.....	5	1	-	-	-	1	1	-	-	1	1	-	-
2002.....	2	1	-	-	-	-	-	-	-	-	1	-	-
2001.....	4	1	-	-	-	-	-	-	-	-	2	1	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Shigellosis and amebiasis (A03,A06)													
All origins, both sexes													
2003.....	7	-	-	-	-	-	3	-	2	1	1	-	-
2002.....	8	-	1	-	1	-	1	2	1	-	1	1	-
2001.....	7	1	1	1	-	-	-	3	-	-	1	-	-
2000.....	12	-	-	-	-	-	1	1	1	1	7	1	-
1999.....	10	-	1	1	-	-	1	-	1	1	1	4	-
All origins, male													
2003.....	5	-	-	-	-	-	2	-	1	1	1	-	-
2002.....	4	-	-	-	1	-	1	2	-	-	-	-	-
2001.....	6	1	1	1	-	-	-	3	-	-	-	-	-
2000.....	5	-	-	-	-	-	1	-	1	-	3	-	-
1999.....	6	-	1	1	-	-	1	-	1	1	-	1	-
All origins, female													
2003.....	2	-	-	-	-	-	1	-	1	-	-	-	-
2002.....	4	-	1	-	-	-	-	-	1	-	1	1	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	7	-	-	-	-	-	-	1	-	1	4	1	-
1999.....	4	-	-	-	-	-	-	-	-	-	1	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Shigellosis and amebiasis (A03,A06)

Hispanic, both sexes													
2003.....	4	-	-	-	-	-	2	-	1	1	-	-	-
2002.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	3	-	1	1	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	4	-	1	1	-	-	1	-	-	-	1	-	-
Hispanic, male													
2003.....	2	-	-	-	-	-	1	-	-	1	-	-	-
2002.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	3	-	1	1	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	1	1	-	-	1	-	-	-	-	-	-
Hispanic, female													
2003.....	2	-	-	-	-	-	1	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)													
All origins, both sexes													
2003.....	3,090	10	4	7	4	8	14	64	176	444	1,192	1,167	-
2002.....	2,465	13	8	3	6	4	18	56	142	362	889	964	-
2001.....	1,537	10	18	4	-	4	12	42	81	235	566	565	-
2000.....	1,328	13	10	-	1	3	19	29	70	222	462	499	-
1999.....	1,108	12	8	3	5	6	7	23	49	176	390	429	-
All origins, male													
2003.....	1,109	6	4	2	2	4	5	32	82	185	478	309	-
2002.....	947	3	5	1	3	1	8	29	71	185	358	283	-
2001.....	572	7	11	2	-	2	8	15	35	108	227	157	-
2000.....	524	10	6	-	-	3	10	16	39	105	195	140	-
1999.....	420	8	4	3	3	5	4	11	26	74	169	113	-
All origins, female													
2003.....	1,981	4	-	5	2	4	9	32	94	259	714	858	-
2002.....	1,518	10	3	2	3	3	10	27	71	177	531	681	-
2001.....	965	3	7	2	-	2	4	27	46	127	339	408	-
2000.....	804	3	4	-	1	-	9	13	31	117	267	359	-
1999.....	688	4	4	-	2	1	3	12	23	102	221	316	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)													
Hispanic, both sexes													
2003.....	99	2	2	3	-	1	-	6	10	16	31	28	-
2002.....	78	5	5	-	1	-	1	3	6	10	31	16	-
2001.....	53	1	1	-	-	-	4	2	4	8	16	17	-
2000.....	46	3	2	-	-	-	2	-	8	9	11	11	-
1999.....	34	3	1	-	2	1	-	4	5	7	4	7	-
Hispanic, male													
2003.....	47	-	2	1	-	-	-	4	7	7	13	13	-
2002.....	31	1	3	-	-	-	-	2	4	5	12	4	-
2001.....	19	1	1	-	-	-	3	-	2	1	5	6	-
2000.....	23	2	1	-	-	-	1	-	5	6	4	4	-
1999.....	17	2	-	-	2	1	-	2	2	3	2	3	-
Hispanic, female													
2003.....	52	2	-	2	-	1	-	2	3	9	18	15	-
2002.....	47	4	2	-	1	-	1	1	2	5	19	12	-
2001.....	34	-	-	-	-	-	1	2	2	7	11	11	-
2000.....	23	1	1	-	-	-	1	-	3	3	7	7	-
1999.....	17	1	1	-	-	-	-	2	3	4	2	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)													
Non-Hispanic 1/, both sexes													
2003.....	2,986	8	2	4	4	7	14	57	164	428	1,160	1,138	-
2002.....	2,381	8	3	3	5	4	17	52	135	351	857	946	-
2001.....	1,474	9	16	4	-	4	8	40	77	225	546	545	-
2000.....	1,277	10	8	-	1	3	17	29	62	211	449	487	-
1999.....	1,071	9	7	3	3	5	7	19	44	169	386	419	-
Non-Hispanic 1/, male													
2003.....	1,061	6	2	1	2	4	5	28	75	178	465	295	-
2002.....	915	2	2	1	3	1	8	26	67	180	346	279	-
2001.....	550	6	10	2	-	2	5	15	33	105	222	150	-
2000.....	497	8	5	-	-	3	9	16	34	98	189	135	-
1999.....	403	6	4	3	1	4	4	9	24	71	167	110	-
Non-Hispanic 1/, female													
2003.....	1,925	2	-	3	2	3	9	29	89	250	695	843	-
2002.....	1,466	6	1	2	2	3	9	26	68	171	511	667	-
2001.....	924	3	6	2	-	2	3	25	44	120	324	395	-
2000.....	780	2	3	-	1	-	8	13	28	113	260	352	-
1999.....	668	3	3	-	2	1	3	10	20	98	219	309	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)													
Non-Hispanic white, both sexes													
2003.....	2,776	3	1	3	3	5	11	44	140	397	1,088	1,081	-
2002.....	2,205	2	1	3	5	2	14	45	104	319	814	896	-
2001.....	1,357	5	13	3	-	2	6	35	62	202	509	520	-
2000.....	1,175	5	5	-	1	2	9	22	54	191	430	456	-
1999.....	990	5	6	1	2	5	4	15	36	155	360	401	-
Non-Hispanic white, male													
2003.....	989	2	1	1	2	3	5	23	61	166	442	283	-
2002.....	840	-	1	1	3	1	6	24	52	159	328	265	-
2001.....	497	3	8	1	-	2	3	12	29	94	203	142	-
2000.....	453	4	4	-	-	2	6	12	29	89	180	127	-
1999.....	373	4	3	1	1	4	3	6	20	69	158	104	-
Non-Hispanic white, female													
2003.....	1,787	1	-	2	1	2	6	21	79	231	646	798	-
2002.....	1,365	2	-	2	2	1	8	21	52	160	486	631	-
2001.....	860	2	5	2	-	-	3	23	33	108	306	378	-
2000.....	722	1	1	-	1	-	3	10	25	102	250	329	-
1999.....	617	1	3	-	1	1	1	9	16	86	202	297	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)													
Non-Hispanic black, both sexes													
2003.....	183	5	1	1	1	1	2	10	21	25	64	52	-
2002.....	155	5	1	-	-	2	3	5	29	31	35	44	-
2001.....	98	3	3	1	-	2	2	5	13	22	28	19	-
2000.....	83	4	2	-	-	1	6	7	6	15	16	26	-
1999.....	74	4	-	2	1	-	3	3	6	14	25	16	-
Non-Hispanic black, male													
2003.....	65	4	1	-	-	1	-	4	12	9	23	11	-
2002.....	66	1	1	-	-	-	2	2	15	20	13	12	-
2001.....	42	2	2	1	-	-	2	3	4	10	12	6	-
2000.....	36	3	1	-	-	1	2	4	4	6	8	7	-
1999.....	25	2	-	2	-	-	1	2	2	2	9	5	-
Non-Hispanic black, female													
2003.....	118	1	-	1	1	-	2	6	9	16	41	41	-
2002.....	89	4	-	-	-	2	1	3	14	11	22	32	-
2001.....	56	1	1	-	-	2	-	2	9	12	16	13	-
2000.....	47	1	1	-	-	-	4	3	2	9	8	19	-
1999.....	49	2	-	-	1	-	2	1	4	12	16	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
All origins, both sexes													
2003.....	711	-	1	2	10	18	35	105	102	114	217	107	-
2002.....	784	1	-	-	9	18	52	81	107	161	234	121	-
2001.....	764	2	1	-	12	25	40	84	116	152	206	125	1
2000.....	776	1	6	1	11	20	42	91	81	155	252	115	1
1999.....	930	2	2	3	11	19	57	106	130	173	253	173	1
All origins, male													
2003.....	427	-	1	2	3	13	25	79	71	72	116	45	-
2002.....	462	-	-	-	5	11	38	55	75	102	121	55	-
2001.....	453	-	-	-	8	14	27	63	85	98	100	57	1
2000.....	468	-	1	-	7	14	28	76	59	87	139	56	1
1999.....	584	-	2	2	4	11	43	87	96	116	137	85	1
All origins, female													
2003.....	284	-	-	-	7	5	10	26	31	42	101	62	-
2002.....	322	1	-	-	4	7	14	26	32	59	113	66	-
2001.....	311	2	1	-	4	11	13	21	31	54	106	68	-
2000.....	308	1	5	1	4	6	14	15	22	68	113	59	-
1999.....	346	2	-	1	7	8	14	19	34	57	116	88	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
Hispanic, both sexes													
2003.....	89	-	1	-	5	6	8	13	13	15	20	8	-
2002.....	95	-	-	-	2	4	11	10	17	21	19	11	-
2001.....	97	-	-	-	6	9	9	9	18	18	20	8	-
2000.....	83	-	2	1	3	7	3	9	12	13	23	10	-
1999.....	125	1	-	1	4	3	13	19	20	24	27	13	-
Hispanic, male													
2003.....	60	-	1	-	3	4	7	11	9	11	11	3	-
2002.....	52	-	-	-	1	3	9	6	6	11	11	5	-
2001.....	67	-	-	-	6	7	5	9	13	14	10	3	-
2000.....	46	-	-	-	1	4	2	5	7	8	14	5	-
1999.....	78	-	-	1	3	3	9	16	11	16	13	6	-
Hispanic, female													
2003.....	29	-	-	-	2	2	1	2	4	4	9	5	-
2002.....	43	-	-	-	1	1	2	4	11	10	8	6	-
2001.....	30	-	-	-	-	2	4	-	5	4	10	5	-
2000.....	37	-	2	1	2	3	1	4	5	5	9	5	-
1999.....	47	1	-	-	1	-	4	3	9	8	14	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
Non-Hispanic 1/, both sexes													
2003.....	617	-	-	2	5	12	26	92	87	98	197	98	-
2002.....	682	1	-	-	6	14	39	70	88	139	215	110	-
2001.....	660	2	1	-	6	16	30	73	96	134	186	116	-
2000.....	683	1	4	-	8	13	39	80	68	140	226	104	-
1999.....	788	1	2	2	6	16	42	82	109	148	222	158	-
Non-Hispanic 1/, male													
2003.....	364	-	-	2	-	9	18	68	60	60	105	42	-
2002.....	404	-	-	-	4	8	27	48	67	90	110	50	-
2001.....	379	-	-	-	2	7	21	52	70	84	90	53	-
2000.....	414	-	1	-	6	10	26	69	51	78	122	51	-
1999.....	492	-	2	1	-	8	32	66	84	99	122	78	-
Non-Hispanic 1/, female													
2003.....	253	-	-	-	5	3	8	24	27	38	92	56	-
2002.....	278	1	-	-	2	6	12	22	21	49	105	60	-
2001.....	281	2	1	-	4	9	9	21	26	50	96	63	-
2000.....	269	1	3	-	2	3	13	11	17	62	104	53	-
1999.....	296	1	-	1	6	8	10	16	25	49	100	80	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
Non-Hispanic white, both sexes													
2003.....	347	-	-	2	1	2	3	32	47	60	130	70	-
2002.....	366	-	-	-	1	2	10	26	41	66	145	75	-
2001.....	368	1	1	-	2	2	12	32	48	72	120	78	-
2000.....	353	-	2	-	3	4	10	34	22	72	131	75	-
1999.....	430	-	-	-	-	2	11	30	53	84	142	108	-
Non-Hispanic white, male													
2003.....	195	-	-	2	-	1	2	26	34	34	66	30	-
2002.....	218	-	-	-	1	-	7	20	31	52	72	35	-
2001.....	200	-	-	-	-	2	10	20	37	39	59	33	-
2000.....	205	-	-	-	3	3	9	28	17	44	63	38	-
1999.....	263	-	-	-	-	1	9	27	41	57	77	51	-
Non-Hispanic white, female													
2003.....	152	-	-	-	1	1	1	6	13	26	64	40	-
2002.....	148	-	-	-	-	2	3	6	10	14	73	40	-
2001.....	168	1	1	-	2	-	2	12	11	33	61	45	-
2000.....	148	-	2	-	-	1	1	6	5	28	68	37	-
1999.....	167	-	-	-	-	1	2	3	12	27	65	57	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Tuberculosis (A16-A19)													
Non-Hispanic black, both sexes													
2003.....	168	-	-	-	4	6	15	47	29	20	33	14	-
2002.....	198	-	-	-	3	8	23	34	31	43	37	19	-
2001.....	200	-	-	-	1	9	16	36	37	44	35	22	-
2000.....	238	-	2	-	2	7	25	41	33	52	58	18	-
1999.....	239	1	2	1	4	9	24	41	39	42	50	26	-
Non-Hispanic black, male													
2003.....	111	-	-	-	-	5	11	35	19	15	20	6	-
2002.....	123	-	-	-	2	5	16	23	23	25	22	7	-
2001.....	124	-	-	-	-	3	10	29	26	30	14	12	-
2000.....	156	-	1	-	1	5	14	36	26	28	38	7	-
1999.....	154	-	2	-	-	4	18	33	31	28	25	13	-
Non-Hispanic black, female													
2003.....	57	-	-	-	4	1	4	12	10	5	13	8	-
2002.....	75	-	-	-	1	3	7	11	8	18	15	12	-
2001.....	76	-	-	-	1	6	6	7	11	14	21	10	-
2000.....	82	-	1	-	1	2	11	5	7	24	20	11	-
1999.....	85	1	-	1	4	5	6	8	8	14	25	13	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Tuberculosis (A16-A19)

Origin not stated 2/, both sexes

2003.....	5	-	-	-	-	-	1	-	2	1	-	1	-
2002.....	7	-	-	-	1	-	2	1	2	1	-	-	-
2001.....	7	-	-	-	-	-	1	2	2	-	-	1	1
2000.....	10	-	-	-	-	-	-	2	1	2	3	1	1
1999.....	17	-	-	-	1	-	2	5	1	1	4	2	1

Origin not stated 2/, male

2003.....	3	-	-	-	-	-	-	-	2	1	-	-	-
2002.....	6	-	-	-	-	-	2	1	2	1	-	-	-
2001.....	7	-	-	-	-	-	1	2	2	-	-	1	1
2000.....	8	-	-	-	-	-	-	2	1	1	3	-	1
1999.....	14	-	-	-	1	-	2	5	1	1	2	1	1

Origin not stated 2/, female

2003.....	2	-	-	-	-	-	1	-	-	-	-	1	-
2002.....	1	-	-	-	1	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	1	-	1	-
1999.....	3	-	-	-	-	-	-	-	-	-	2	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
All origins, both sexes													
2003.....	562	-	1	2	9	6	26	86	81	88	177	86	-
2002.....	626	-	-	-	6	14	40	64	84	120	192	106	-
2001.....	600	2	1	-	7	17	28	61	96	115	166	106	1
2000.....	615	-	3	-	7	11	35	72	60	119	209	98	1
1999.....	764	1	1	1	8	14	45	76	109	139	216	153	1
All origins, male													
2003.....	344	-	1	2	3	6	17	64	57	61	97	36	-
2002.....	372	-	-	-	4	9	31	43	60	80	96	49	-
2001.....	371	-	-	-	5	10	18	49	78	76	84	50	1
2000.....	393	-	1	-	4	8	26	64	50	73	116	50	1
1999.....	486	-	1	1	2	8	33	64	83	97	120	76	1
All origins, female													
2003.....	218	-	-	-	6	-	9	22	24	27	80	50	-
2002.....	254	-	-	-	2	5	9	21	24	40	96	57	-
2001.....	229	2	1	-	2	7	10	12	18	39	82	56	-
2000.....	222	-	2	-	3	3	9	8	10	46	93	48	-
1999.....	278	1	-	-	6	6	12	12	26	42	96	77	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
Hispanic, both sexes													
2003.....	77	-	1	-	5	4	7	10	11	14	19	6	-
2002.....	78	-	-	-	1	4	10	8	12	16	16	11	-
2001.....	77	-	-	-	3	8	5	7	15	17	17	5	-
2000.....	60	-	1	-	2	3	2	5	9	12	18	8	-
1999.....	93	-	-	-	2	1	10	10	16	18	24	12	-
Hispanic, male													
2003.....	55	-	1	-	3	4	6	9	7	11	11	3	-
2002.....	44	-	-	-	1	3	8	4	5	9	9	5	-
2001.....	56	-	-	-	3	6	3	7	12	13	10	2	-
2000.....	40	-	-	-	1	3	2	3	6	8	12	5	-
1999.....	55	-	-	-	1	1	6	9	9	13	11	5	-
Hispanic, female													
2003.....	22	-	-	-	2	-	1	1	4	3	8	3	-
2002.....	34	-	-	-	-	1	2	4	7	7	7	6	-
2001.....	21	-	-	-	-	2	2	-	3	4	7	3	-
2000.....	20	-	1	-	1	-	-	2	3	4	6	3	-
1999.....	38	-	-	-	1	-	4	1	7	5	13	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
Non-Hispanic 1/, both sexes													
2003.....	481	-	-	2	4	2	18	76	68	74	158	79	-
2002.....	543	-	-	-	5	10	28	56	70	103	176	95	-
2001.....	517	2	1	-	4	9	22	53	79	98	149	100	-
2000.....	546	-	2	-	5	8	33	65	50	105	188	90	-
1999.....	657	1	1	1	5	13	33	63	92	120	189	139	-
Non-Hispanic 1/, male													
2003.....	287	-	-	2	-	2	11	55	48	50	86	33	-
2002.....	323	-	-	-	3	6	21	39	53	70	87	44	-
2001.....	309	-	-	-	2	4	14	41	64	63	74	47	-
2000.....	345	-	1	-	3	5	24	59	43	64	101	45	-
1999.....	419	-	1	1	-	7	25	52	73	83	107	70	-
Non-Hispanic 1/, female													
2003.....	194	-	-	-	4	-	7	21	20	24	72	46	-
2002.....	220	-	-	-	2	4	7	17	17	33	89	51	-
2001.....	208	2	1	-	2	5	8	12	15	35	75	53	-
2000.....	201	-	1	-	2	3	9	6	7	41	87	45	-
1999.....	238	1	-	-	5	6	8	11	19	37	82	69	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
Non-Hispanic white, both sexes													
2003.....	270	-	-	2	1	-	2	25	37	43	106	54	-
2002.....	296	-	-	-	1	1	9	21	32	53	114	65	-
2001.....	288	1	1	-	1	-	8	23	41	52	97	64	-
2000.....	288	-	-	-	2	2	9	29	17	52	112	65	-
1999.....	371	-	-	-	-	2	9	27	45	69	119	100	-
Non-Hispanic white, male													
2003.....	154	-	-	2	-	-	1	20	28	28	53	22	-
2002.....	177	-	-	-	1	-	7	16	24	44	53	32	-
2001.....	162	-	-	-	-	-	6	16	35	29	49	27	-
2000.....	174	-	-	-	2	1	9	26	14	34	54	34	-
1999.....	234	-	-	-	-	1	8	24	36	48	67	50	-
Non-Hispanic white, female													
2003.....	116	-	-	-	1	-	1	5	9	15	53	32	-
2002.....	119	-	-	-	-	1	2	5	8	9	61	33	-
2001.....	126	1	1	-	1	-	2	7	6	23	48	37	-
2000.....	114	-	-	-	-	1	-	3	3	18	58	31	-
1999.....	137	-	-	-	-	1	1	3	9	21	52	50	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
Non-Hispanic black, both sexes													
2003.....	134	-	-	-	3	2	10	41	23	17	26	12	-
2002.....	160	-	-	-	2	8	14	29	26	32	34	15	-
2001.....	157	-	-	-	1	6	12	28	30	32	28	20	-
2000.....	187	-	2	-	1	6	20	31	22	43	46	16	-
1999.....	194	1	1	-	4	8	19	29	35	32	43	22	-
Non-Hispanic black, male													
2003.....	90	-	-	-	-	2	7	29	16	13	17	6	-
2002.....	101	-	-	-	1	5	10	20	20	19	21	5	-
2001.....	103	-	-	-	-	2	7	24	23	23	12	12	-
2000.....	129	-	1	-	-	4	12	28	22	26	30	6	-
1999.....	122	-	1	-	-	3	13	24	28	21	22	10	-
Non-Hispanic black, female													
2003.....	44	-	-	-	3	-	3	12	7	4	9	6	-
2002.....	59	-	-	-	1	3	4	9	6	13	13	10	-
2001.....	54	-	-	-	1	4	5	4	7	9	16	8	-
2000.....	58	-	1	-	1	2	8	3	-	17	16	10	-
1999.....	72	1	-	-	4	5	6	5	7	11	21	12	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Respiratory tuberculosis (A16)													
Origin not stated 2/, both sexes													
2003.....	4	-	-	-	-	-	1	-	2	-	-	1	-
2002.....	5	-	-	-	-	-	2	-	2	1	-	-	-
2001.....	6	-	-	-	-	-	1	1	2	-	-	1	1
2000.....	9	-	-	-	-	-	-	2	1	2	3	-	1
1999.....	14	-	-	-	1	-	2	3	1	1	3	2	1
Origin not stated 2/, male													
2003.....	2	-	-	-	-	-	-	-	2	-	-	-	-
2002.....	5	-	-	-	-	-	2	-	2	1	-	-	-
2001.....	6	-	-	-	-	-	1	1	2	-	-	1	1
2000.....	8	-	-	-	-	-	-	2	1	1	3	-	1
1999.....	12	-	-	-	1	-	2	3	1	1	2	1	1
Origin not stated 2/, female													
2003.....	2	-	-	-	-	-	1	-	-	-	-	1	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	1	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	-	1	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
All origins, both sexes													
2003.....	149	-	-	-	1	12	9	19	21	26	40	21	-
2002.....	158	1	-	-	3	4	12	17	23	41	42	15	-
2001.....	164	-	-	-	5	8	12	23	20	37	40	19	-
2000.....	161	1	3	1	4	9	7	19	21	36	43	17	-
1999.....	166	1	1	2	3	5	12	30	21	34	37	20	-
All origins, male													
2003.....	83	-	-	-	-	7	8	15	14	11	19	9	-
2002.....	90	-	-	-	1	2	7	12	15	22	25	6	-
2001.....	82	-	-	-	3	4	9	14	7	22	16	7	-
2000.....	75	-	-	-	3	6	2	12	9	14	23	6	-
1999.....	98	-	1	1	2	3	10	23	13	19	17	9	-
All origins, female													
2003.....	66	-	-	-	1	5	1	4	7	15	21	12	-
2002.....	68	1	-	-	2	2	5	5	8	19	17	9	-
2001.....	82	-	-	-	2	4	3	9	13	15	24	12	-
2000.....	86	1	3	1	1	3	5	7	12	22	20	11	-
1999.....	68	1	-	1	1	2	2	7	8	15	20	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
Hispanic, both sexes													
2003.....	12	-	-	-	-	2	1	3	2	1	1	2	-
2002.....	17	-	-	-	1	-	1	2	5	5	3	-	-
2001.....	20	-	-	-	3	1	4	2	3	1	3	3	-
2000.....	23	-	1	1	1	4	1	4	3	1	5	2	-
1999.....	32	1	-	1	2	2	3	9	4	6	3	1	-
Hispanic, male													
2003.....	5	-	-	-	-	-	1	2	2	-	-	-	-
2002.....	8	-	-	-	-	-	1	2	1	2	2	-	-
2001.....	11	-	-	-	3	1	2	2	1	1	-	1	-
2000.....	6	-	-	-	-	1	-	2	1	-	2	-	-
1999.....	23	-	-	1	2	2	3	7	2	3	2	1	-
Hispanic, female													
2003.....	7	-	-	-	-	2	-	1	-	1	1	2	-
2002.....	9	-	-	-	1	-	-	-	4	3	1	-	-
2001.....	9	-	-	-	-	-	2	-	2	-	3	2	-
2000.....	17	-	1	1	1	3	1	2	2	1	3	2	-
1999.....	9	1	-	-	-	-	-	2	2	3	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
Non-Hispanic 1/, both sexes													
2003.....	136	-	-	-	1	10	8	16	19	24	39	19	-
2002.....	139	1	-	-	1	4	11	14	18	36	39	15	-
2001.....	143	-	-	-	2	7	8	20	17	36	37	16	-
2000.....	137	1	2	-	3	5	6	15	18	35	38	14	-
1999.....	131	-	1	1	1	3	9	19	17	28	33	19	-
Non-Hispanic 1/, male													
2003.....	77	-	-	-	-	7	7	13	12	10	19	9	-
2002.....	81	-	-	-	1	2	6	9	14	20	23	6	-
2001.....	70	-	-	-	-	3	7	11	6	21	16	6	-
2000.....	69	-	-	-	3	5	2	10	8	14	21	6	-
1999.....	73	-	1	-	-	1	7	14	11	16	15	8	-
Non-Hispanic 1/, female													
2003.....	59	-	-	-	1	3	1	3	7	14	20	10	-
2002.....	58	1	-	-	-	2	5	5	4	16	16	9	-
2001.....	73	-	-	-	2	4	1	9	11	15	21	10	-
2000.....	68	1	2	-	-	-	4	5	10	21	17	8	-
1999.....	58	-	-	1	1	2	2	5	6	12	18	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
Non-Hispanic white, both sexes													
2003.....	77	-	-	-	-	2	1	7	10	17	24	16	-
2002.....	70	-	-	-	-	1	1	5	9	13	31	10	-
2001.....	80	-	-	-	1	2	4	9	7	20	23	14	-
2000.....	65	-	2	-	1	2	1	5	5	20	19	10	-
1999.....	59	-	-	-	-	-	2	3	8	15	23	8	-
Non-Hispanic white, male													
2003.....	41	-	-	-	-	1	1	6	6	6	13	8	-
2002.....	41	-	-	-	-	-	-	4	7	8	19	3	-
2001.....	38	-	-	-	-	2	4	4	2	10	10	6	-
2000.....	31	-	-	-	1	2	-	2	3	10	9	4	-
1999.....	29	-	-	-	-	-	1	3	5	9	10	1	-
Non-Hispanic white, female													
2003.....	36	-	-	-	-	1	-	1	4	11	11	8	-
2002.....	29	-	-	-	-	1	1	1	2	5	12	7	-
2001.....	42	-	-	-	1	-	-	5	5	10	13	8	-
2000.....	34	-	2	-	-	-	1	3	2	10	10	6	-
1999.....	30	-	-	-	-	-	1	-	3	6	13	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other tuberculosis (A17-A19)													
Non-Hispanic black, both sexes													
2003.....	34	-	-	-	1	4	5	6	6	3	7	2	-
2002.....	38	-	-	-	1	-	9	5	5	11	3	4	-
2001.....	43	-	-	-	-	3	4	8	7	12	7	2	-
2000.....	51	-	-	-	1	1	5	10	11	9	12	2	-
1999.....	45	-	1	1	-	1	5	12	4	10	7	4	-
Non-Hispanic black, male													
2003.....	21	-	-	-	-	3	4	6	3	2	3	-	-
2002.....	22	-	-	-	1	-	6	3	3	6	1	2	-
2001.....	21	-	-	-	-	1	3	5	3	7	2	-	-
2000.....	27	-	-	-	1	1	2	8	4	2	8	1	-
1999.....	32	-	1	-	-	1	5	9	3	7	3	3	-
Non-Hispanic black, female													
2003.....	13	-	-	-	1	1	1	-	3	1	4	2	-
2002.....	16	-	-	-	-	-	3	2	2	5	2	2	-
2001.....	22	-	-	-	-	2	1	3	4	5	5	2	-
2000.....	24	-	-	-	-	-	3	2	7	7	4	1	-
1999.....	13	-	-	1	-	-	-	3	1	3	4	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other tuberculosis (A17-A19)

Origin not stated 2/, both sexes

2003.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2002.....	2	-	-	-	1	-	-	1	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	1	-
1999.....	3	-	-	-	-	-	-	2	-	-	1	-	-

Origin not stated 2/, male

2003.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2002.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	2	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	1	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	1	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Whooping cough (A37)

Origin not stated 2/, both sexes

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing data for 'Origin not stated 2/, both sexes'.

Origin not stated 2/, male

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing data for 'Origin not stated 2/, male'.

Origin not stated 2/, female

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing data for 'Origin not stated 2/, female'.

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Scarlet fever and erysipelas (A38,A46)

All origins, both sexes													
2003.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2001.....	3	-	-	1	-	-	-	-	-	1	-	1	-
2000.....	2	-	-	-	-	-	-	1	-	-	1	-	-
1999.....	5	-	-	-	-	-	-	-	-	2	1	2	-
All origins, male													
2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	1	-	-	-	-	-	-	-	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	1	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-	-
All origins, female													
2003.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	1	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	5	-	-	-	-	-	-	-	-	2	1	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Scarlet fever and erysipelas (A38,A46)

Non-Hispanic 1/, both sexes

2003.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2001.....	3	-	-	1	-	-	-	-	-	1	-	1	-
2000.....	2	-	-	-	-	-	-	1	-	-	1	-	-
1999.....	5	-	-	-	-	-	-	-	-	2	1	2	-

Non-Hispanic 1/, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	1	-	-	-	-	-	-	-	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	1	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, female

2003.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	1	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	5	-	-	-	-	-	-	-	-	2	1	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Scarlet fever and erysipelas (A38,A46)

Non-Hispanic white, both sexes

2003.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2001.....	3	-	-	1	-	-	-	-	-	1	-	1	-
2000.....	2	-	-	-	-	-	-	1	-	-	1	-	-
1999.....	4	-	-	-	-	-	-	-	-	1	1	2	-

Non-Hispanic white, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	1	-	-	-	-	-	-	-	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	1	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic white, female

2003.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	1	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	4	-	-	-	-	-	-	-	-	1	1	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Scarlet fever and erysipelas (A38,A46)

Non-Hispanic black, both sexes

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-	-

Non-Hispanic black, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic black, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Scarlet fever and erysipelas (A38,A46)

Origin not stated 2/, both sexes

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing data for 'Origin not stated 2/, both sexes'. All cells contain dashes.

Origin not stated 2/, male

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing data for 'Origin not stated 2/, male'. All cells contain dashes.

Origin not stated 2/, female

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing data for 'Origin not stated 2/, female'. All cells contain dashes.

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningococcal infection (A39)													
All origins, both sexes													
2003.....	161	12	18	11	39	12	19	18	11	9	9	3	-
2002.....	161	17	12	20	33	16	9	19	13	7	10	5	-
2001.....	199	16	19	26	55	10	13	13	15	15	8	9	-
2000.....	211	22	19	18	56	15	13	29	9	8	15	7	-
1999.....	227	32	30	25	39	23	27	17	4	9	13	8	-
All origins, male													
2003.....	75	9	10	3	18	7	9	10	5	2	1	1	-
2002.....	83	8	9	13	17	10	5	10	4	2	3	2	-
2001.....	94	6	9	15	27	6	6	8	7	5	3	2	-
2000.....	106	14	12	7	35	6	9	11	2	2	7	1	-
1999.....	101	14	19	13	19	12	9	4	-	5	4	2	-
All origins, female													
2003.....	86	3	8	8	21	5	10	8	6	7	8	2	-
2002.....	78	9	3	7	16	6	4	9	9	5	7	3	-
2001.....	105	10	10	11	28	4	7	5	8	10	5	7	-
2000.....	105	8	7	11	21	9	4	18	7	6	8	6	-
1999.....	126	18	11	12	20	11	18	13	4	4	9	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningococcal infection (A39)													
Hispanic, both sexes													
2003.....	23	5	2	1	2	5	2	1	-	4	-	1	-
2002.....	19	7	-	1	5	3	1	1	1	-	-	-	-
2001.....	21	4	5	2	2	2	2	1	2	-	1	-	-
2000.....	34	9	3	2	9	5	-	3	1	-	2	-	-
1999.....	27	7	1	3	2	6	5	2	-	-	1	-	-
Hispanic, male													
2003.....	13	4	1	-	1	3	1	1	-	1	-	1	-
2002.....	9	3	-	1	2	2	-	-	1	-	-	-	-
2001.....	11	-	3	2	2	1	1	1	-	-	1	-	-
2000.....	16	4	2	-	6	2	-	1	-	-	1	-	-
1999.....	11	5	1	2	1	2	-	-	-	-	-	-	-
Hispanic, female													
2003.....	10	1	1	1	1	2	1	-	-	3	-	-	-
2002.....	10	4	-	-	3	1	1	1	-	-	-	-	-
2001.....	10	4	2	-	-	1	1	-	2	-	-	-	-
2000.....	18	5	1	2	3	3	-	2	1	-	1	-	-
1999.....	16	2	-	1	1	4	5	2	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningococcal infection (A39)													
Non-Hispanic 1/, both sexes													
2003.....	138	7	16	10	37	7	17	17	11	5	9	2	-
2002.....	142	10	12	19	28	13	8	18	12	7	10	5	-
2001.....	178	12	14	24	53	8	11	12	13	15	7	9	-
2000.....	177	13	16	16	47	10	13	26	8	8	13	7	-
1999.....	198	25	28	22	37	17	22	14	4	9	12	8	-
Non-Hispanic 1/, male													
2003.....	62	5	9	3	17	4	8	9	5	1	1	-	-
2002.....	74	5	9	12	15	8	5	10	3	2	3	2	-
2001.....	83	6	6	13	25	5	5	7	7	5	2	2	-
2000.....	90	10	10	7	29	4	9	10	2	2	6	1	-
1999.....	89	9	17	11	18	10	9	4	-	5	4	2	-
Non-Hispanic 1/, female													
2003.....	76	2	7	7	20	3	9	8	6	4	8	2	-
2002.....	68	5	3	7	13	5	3	8	9	5	7	3	-
2001.....	95	6	8	11	28	3	6	5	6	10	5	7	-
2000.....	87	3	6	9	18	6	4	16	6	6	7	6	-
1999.....	109	16	11	11	19	7	13	10	4	4	8	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningococcal infection (A39)													
Non-Hispanic white, both sexes													
2003.....	99	7	10	6	28	3	12	10	7	5	9	2	-
2002.....	106	7	8	9	22	10	4	15	10	6	10	5	-
2001.....	138	10	10	18	38	5	9	11	9	14	6	8	-
2000.....	129	10	11	8	37	6	7	16	7	7	13	7	-
1999.....	155	23	24	15	27	8	14	12	4	9	11	8	-
Non-Hispanic white, male													
2003.....	44	5	6	2	13	2	6	6	2	1	1	-	-
2002.....	58	5	6	8	12	7	3	8	3	1	3	2	-
2001.....	62	5	5	10	18	3	3	6	5	5	1	1	-
2000.....	60	7	6	4	22	1	5	4	2	2	6	1	-
1999.....	67	9	15	7	12	4	6	3	-	5	4	2	-
Non-Hispanic white, female													
2003.....	55	2	4	4	15	1	6	4	5	4	8	2	-
2002.....	48	2	2	1	10	3	1	7	7	5	7	3	-
2001.....	76	5	5	8	20	2	6	5	4	9	5	7	-
2000.....	69	3	5	4	15	5	2	12	5	5	7	6	-
1999.....	88	14	9	8	15	4	8	9	4	4	7	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Meningococcal infection (A39)

Non-Hispanic black, both sexes

2003.....	35	-	6	4	7	3	5	6	4	-	-	-	-
2002.....	29	3	3	8	5	2	4	3	1	-	-	-	-
2001.....	36	2	4	6	13	2	1	1	4	1	1	1	-
2000.....	44	3	5	8	7	4	6	9	1	1	-	-	-
1999.....	39	2	4	6	9	8	8	2	-	-	-	-	-

Non-Hispanic black, male

2003.....	16	-	3	1	3	1	2	3	3	-	-	-	-
2002.....	12	-	2	4	2	-	2	2	-	-	-	-	-
2001.....	18	1	1	3	6	1	1	1	2	-	1	1	-
2000.....	27	3	4	3	5	3	4	5	-	-	-	-	-
1999.....	20	-	2	3	6	5	3	1	-	-	-	-	-

Non-Hispanic black, female

2003.....	19	-	3	3	4	2	3	3	1	-	-	-	-
2002.....	17	3	1	4	3	2	2	1	1	-	-	-	-
2001.....	18	1	3	3	7	1	-	-	2	1	-	-	-
2000.....	17	-	1	5	2	1	2	4	1	1	-	-	-
1999.....	19	2	2	3	3	3	5	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Meningococcal infection (A39)

Origin not stated 2/, both sexes

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	1	-	-	-	-	1	-	-	-	-	-

Origin not stated 2/, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	1	-	-	-	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
All origins, both sexes													
2003.....	34,069	278	85	77	154	309	910	2,157	3,651	5,970	10,932	9,543	3
2002.....	33,865	296	79	95	118	314	856	2,074	3,360	6,336	11,010	9,324	3
2001.....	32,238	312	108	69	133	281	829	1,977	3,111	5,998	10,351	9,069	-
2000.....	31,224	274	99	63	100	280	877	1,845	2,899	5,704	9,938	9,144	1
1999.....	30,680	283	87	77	121	275	812	1,685	2,714	5,750	9,710	9,166	-
All origins, male													
2003.....	14,987	151	54	34	82	161	455	1,155	1,887	3,023	4,894	3,089	2
2002.....	14,947	158	41	52	59	166	441	1,124	1,748	3,154	5,028	2,973	3
2001.....	14,307	174	60	39	75	139	451	1,043	1,606	3,044	4,762	2,914	-
2000.....	13,537	139	53	38	60	158	451	1,039	1,513	2,867	4,482	2,736	1
1999.....	13,395	175	48	36	67	147	460	919	1,412	2,919	4,426	2,786	-
All origins, female													
2003.....	19,082	127	31	43	72	148	455	1,002	1,764	2,947	6,038	6,454	1
2002.....	18,918	138	38	43	59	148	415	950	1,612	3,182	5,982	6,351	-
2001.....	17,931	138	48	30	58	142	378	934	1,505	2,954	5,589	6,155	-
2000.....	17,687	135	46	25	40	122	426	806	1,386	2,837	5,456	6,408	-
1999.....	17,285	108	39	41	54	128	352	766	1,302	2,831	5,284	6,380	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
Hispanic, both sexes													
2003.....	1,505	50	10	7	17	35	86	148	190	270	397	293	2
2002.....	1,521	65	16	14	15	38	67	136	165	341	382	282	-
2001.....	1,386	50	18	18	13	23	72	133	151	288	346	274	-
2000.....	1,226	62	15	7	12	31	79	122	145	235	300	218	-
1999.....	1,215	43	14	8	9	26	74	92	131	265	296	257	-
Hispanic, male													
2003.....	721	29	6	4	8	20	48	94	95	137	175	104	1
2002.....	744	34	9	8	8	25	47	94	84	161	176	98	-
2001.....	684	29	5	14	9	15	43	80	86	149	170	84	-
2000.....	625	33	9	5	6	21	46	78	80	126	143	78	-
1999.....	605	27	8	4	5	14	51	56	78	134	144	84	-
Hispanic, female													
2003.....	784	21	4	3	9	15	38	54	95	133	222	189	1
2002.....	777	31	7	6	7	13	20	42	81	180	206	184	-
2001.....	702	21	13	4	4	8	29	53	65	139	176	190	-
2000.....	601	29	6	2	6	10	33	44	65	109	157	140	-
1999.....	610	16	6	4	4	12	23	36	53	131	152	173	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
Non-Hispanic 1/, both sexes													
2003.....	32,476	227	75	70	137	272	823	2,006	3,445	5,680	10,512	9,228	1
2002.....	32,225	227	63	81	102	275	787	1,931	3,182	5,969	10,593	9,012	3
2001.....	30,754	261	90	50	120	257	755	1,837	2,941	5,690	9,977	8,776	-
2000.....	29,864	208	83	56	88	248	793	1,713	2,739	5,442	9,601	8,892	1
1999.....	29,351	239	73	69	112	248	736	1,585	2,570	5,466	9,378	8,875	-
Non-Hispanic 1/, male													
2003.....	14,226	122	48	30	74	139	406	1,059	1,784	2,876	4,708	2,979	1
2002.....	14,136	120	32	44	50	141	392	1,025	1,657	2,975	4,830	2,867	3
2001.....	13,573	145	55	25	66	123	408	957	1,510	2,882	4,575	2,827	-
2000.....	12,842	104	44	33	54	137	401	957	1,424	2,724	4,316	2,647	1
1999.....	12,737	147	40	32	62	132	408	858	1,330	2,777	4,261	2,690	-
Non-Hispanic 1/, female													
2003.....	18,250	105	27	40	63	133	417	947	1,661	2,804	5,804	6,249	-
2002.....	18,089	107	31	37	52	134	395	906	1,525	2,994	5,763	6,145	-
2001.....	17,181	116	35	25	54	134	347	880	1,431	2,808	5,402	5,949	-
2000.....	17,022	104	39	23	34	111	392	756	1,315	2,718	5,285	6,245	-
1999.....	16,614	92	33	37	50	116	328	727	1,240	2,689	5,117	6,185	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
Non-Hispanic white, both sexes													
2003.....	25,664	96	43	47	86	155	513	1,295	2,471	4,315	8,803	7,840	-
2002.....	25,545	129	42	55	72	165	510	1,274	2,241	4,583	8,806	7,666	2
2001.....	24,389	122	63	35	69	145	458	1,181	2,081	4,385	8,352	7,498	-
2000.....	23,517	108	51	42	48	134	498	1,092	1,919	4,173	7,901	7,550	1
1999.....	23,023	121	47	47	71	155	439	1,015	1,792	4,140	7,717	7,479	-
Non-Hispanic white, male													
2003.....	11,181	46	29	22	48	74	255	698	1,280	2,207	3,972	2,550	-
2002.....	11,148	67	25	34	37	86	249	680	1,204	2,310	4,006	2,448	2
2001.....	10,668	71	41	18	37	69	266	615	1,071	2,223	3,851	2,406	-
2000.....	10,056	55	26	25	33	74	262	613	1,015	2,131	3,579	2,242	1
1999.....	9,863	79	25	24	44	87	243	551	904	2,111	3,538	2,257	-
Non-Hispanic white, female													
2003.....	14,483	50	14	25	38	81	258	597	1,191	2,108	4,831	5,290	-
2002.....	14,397	62	17	21	35	79	261	594	1,037	2,273	4,800	5,218	-
2001.....	13,721	51	22	17	32	76	192	566	1,010	2,162	4,501	5,092	-
2000.....	13,461	53	25	17	15	60	236	479	904	2,042	4,322	5,308	-
1999.....	13,160	42	22	23	27	68	196	464	888	2,029	4,179	5,222	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
Non-Hispanic black, both sexes													
2003.....	6,141	119	24	15	41	102	283	650	881	1,232	1,536	1,257	1
2002.....	6,074	85	17	20	24	102	254	593	868	1,251	1,633	1,226	1
2001.....	5,829	124	21	10	41	98	268	595	804	1,186	1,501	1,181	-
2000.....	5,863	96	25	11	33	107	269	572	756	1,173	1,576	1,245	-
1999.....	5,772	106	16	14	32	85	279	518	716	1,212	1,513	1,281	-
Non-Hispanic black, male													
2003.....	2,748	70	14	6	19	56	134	338	462	601	663	384	1
2002.....	2,668	46	5	7	10	47	130	312	418	598	729	365	1
2001.....	2,611	67	12	3	25	46	128	304	411	586	656	373	-
2000.....	2,559	48	13	6	18	59	125	319	381	545	677	368	-
1999.....	2,594	60	10	7	14	41	154	279	392	609	648	380	-
Non-Hispanic black, female													
2003.....	3,393	49	10	9	22	46	149	312	419	631	873	873	-
2002.....	3,406	39	12	13	14	55	124	281	450	653	904	861	-
2001.....	3,218	57	9	7	16	52	140	291	393	600	845	808	-
2000.....	3,304	48	12	5	15	48	144	253	375	628	899	877	-
1999.....	3,178	46	6	7	18	44	125	239	324	603	865	901	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Septicemia (A40-A41)													
Origin not stated 2/, both sexes													
2003.....	88	1	-	-	-	2	1	3	16	20	23	22	-
2002.....	119	4	-	-	1	1	2	7	13	26	35	30	-
2001.....	98	1	-	1	-	1	2	7	19	20	28	19	-
2000.....	134	4	1	-	-	1	5	10	15	27	37	34	-
1999.....	114	1	-	-	-	1	2	8	13	19	36	34	-
Origin not stated 2/, male													
2003.....	40	-	-	-	-	2	1	2	8	10	11	6	-
2002.....	67	4	-	-	1	-	2	5	7	18	22	8	-
2001.....	50	-	-	-	-	1	-	6	10	13	17	3	-
2000.....	70	2	-	-	-	-	4	4	9	17	23	11	-
1999.....	53	1	-	-	-	1	1	5	4	8	21	12	-
Origin not stated 2/, female													
2003.....	48	1	-	-	-	-	-	1	8	10	12	16	-
2002.....	52	-	-	-	-	1	-	2	6	8	13	22	-
2001.....	48	1	-	1	-	-	2	1	9	7	11	16	-
2000.....	64	2	1	-	-	1	1	6	6	10	14	23	-
1999.....	61	-	-	-	-	-	1	3	9	11	15	22	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Syphilis (A50-A53)

All origins, both sexes

2003.....	34	-	-	-	-	-	3	5	3	6	8	9	-
2002.....	41	1	1	-	-	-	2	5	5	6	12	9	-
2001.....	36	1	-	1	-	1	3	3	4	5	13	5	-
2000.....	41	1	-	-	-	1	-	6	3	7	13	10	-
1999.....	33	2	-	-	-	1	1	5	4	6	10	4	-

All origins, male

2003.....	22	-	-	-	-	-	3	5	2	5	4	3	-
2002.....	21	-	1	-	-	-	2	4	2	5	4	3	-
2001.....	19	1	-	1	-	-	2	3	3	3	5	1	-
2000.....	24	1	-	-	-	1	-	5	3	2	7	5	-
1999.....	13	-	-	-	-	1	-	3	3	5	-	1	-

All origins, female

2003.....	12	-	-	-	-	-	-	-	1	1	4	6	-
2002.....	20	1	-	-	-	-	-	1	3	1	8	6	-
2001.....	17	-	-	-	-	1	1	-	1	2	8	4	-
2000.....	17	-	-	-	-	-	-	1	-	5	6	5	-
1999.....	20	2	-	-	-	-	1	2	1	1	10	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Syphilis (A50-A53)

Hispanic, both sexes

2003.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2001.....	7	1	-	-	-	-	-	-	2	-	3	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	1	-	-
1999.....	5	2	-	-	-	-	-	2	-	-	1	-	-

Hispanic, male

2003.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2001.....	5	1	-	-	-	-	-	-	2	-	1	1	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	2	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	1	-	-
1999.....	4	2	-	-	-	-	-	1	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Syphilis (A50-A53)													
Non-Hispanic 1/, both sexes													
2003.....	32	-	-	-	-	-	3	4	3	5	8	9	-
2002.....	39	1	1	-	-	-	2	5	5	6	11	8	-
2001.....	28	-	-	1	-	1	3	3	1	5	10	4	-
2000.....	39	1	-	-	-	1	-	6	3	7	11	10	-
1999.....	27	-	-	-	-	1	1	2	4	6	9	4	-
Non-Hispanic 1/, male													
2003.....	21	-	-	-	-	-	3	4	2	5	4	3	-
2002.....	20	-	1	-	-	-	2	4	2	5	3	3	-
2001.....	14	-	-	1	-	-	2	3	1	3	4	-	-
2000.....	24	1	-	-	-	1	-	5	3	2	7	5	-
1999.....	11	-	-	-	-	1	-	1	3	5	-	1	-
Non-Hispanic 1/, female													
2003.....	11	-	-	-	-	-	-	-	1	-	4	6	-
2002.....	19	1	-	-	-	-	-	1	3	1	8	5	-
2001.....	14	-	-	-	-	1	1	-	-	2	6	4	-
2000.....	15	-	-	-	-	-	-	1	-	5	4	5	-
1999.....	16	-	-	-	-	-	1	1	1	1	9	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Syphilis (A50-A53)

Non-Hispanic white, both sexes

2003.....	16	-	-	-	-	-	1	-	2	2	5	6	-
2002.....	16	-	-	-	-	-	1	2	1	3	4	5	-
2001.....	14	-	-	-	-	-	1	1	1	2	7	2	-
2000.....	20	-	-	-	-	1	-	2	-	3	6	8	-
1999.....	14	-	-	-	-	1	1	2	3	3	3	1	-

Non-Hispanic white, male

2003.....	9	-	-	-	-	-	1	-	1	2	3	2	-
2002.....	8	-	-	-	-	-	1	2	-	3	-	2	-
2001.....	5	-	-	-	-	-	-	1	1	1	2	-	-
2000.....	7	-	-	-	-	1	-	1	-	-	2	3	-
1999.....	6	-	-	-	-	1	-	1	2	2	-	-	-

Non-Hispanic white, female

2003.....	7	-	-	-	-	-	-	-	1	-	2	4	-
2002.....	8	-	-	-	-	-	-	-	1	-	4	3	-
2001.....	9	-	-	-	-	-	1	-	-	1	5	2	-
2000.....	13	-	-	-	-	-	-	1	-	3	4	5	-
1999.....	8	-	-	-	-	-	1	1	1	1	3	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Syphilis (A50-A53)													
Non-Hispanic black, both sexes													
2003.....	15	-	-	-	-	-	2	4	1	3	3	2	-
2002.....	23	1	1	-	-	-	1	3	4	3	7	3	-
2001.....	14	-	-	1	-	1	2	2	-	3	3	2	-
2000.....	19	1	-	-	-	-	-	4	3	4	5	2	-
1999.....	12	-	-	-	-	-	-	-	1	3	5	3	-
Non-Hispanic black, male													
2003.....	11	-	-	-	-	-	2	4	1	3	1	-	-
2002.....	12	-	1	-	-	-	1	2	2	2	3	1	-
2001.....	9	-	-	1	-	-	2	2	-	2	2	-	-
2000.....	17	1	-	-	-	-	-	4	3	2	5	2	-
1999.....	5	-	-	-	-	-	-	-	1	3	-	1	-
Non-Hispanic black, female													
2003.....	4	-	-	-	-	-	-	-	-	-	2	2	-
2002.....	11	1	-	-	-	-	-	1	2	1	4	2	-
2001.....	5	-	-	-	-	1	-	-	-	1	1	2	-
2000.....	2	-	-	-	-	-	-	-	-	2	-	-	-
1999.....	7	-	-	-	-	-	-	-	-	-	5	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Acute poliomyelitis (A80)

Non-Hispanic 1/, both sexes

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing mortality data for Non-Hispanic 1/, both sexes. All cells contain dashes.

Non-Hispanic 1/, male

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing mortality data for Non-Hispanic 1/, male. All cells contain dashes.

Non-Hispanic 1/, female

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing mortality data for Non-Hispanic 1/, female. All cells contain dashes.

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Arthropod-borne viral encephalitis (A83-A84,A85.2)

Origin not stated 2/, both sexes

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing data for 'Origin not stated 2/, both sexes'.

Origin not stated 2/, male

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing data for 'Origin not stated 2/, male'.

Origin not stated 2/, female

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing data for 'Origin not stated 2/, female'.

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Measles (B05)

All origins, both sexes

2003.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	1	-	-	-	-	-	-	1	-	-	-	-

All origins, male

2003.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-	-

All origins, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	1	-	-	-	-	-	-	1	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Measles (B05)

Non-Hispanic 1/, both sexes

2003.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	1	-	-	-	-	-	-	1	-	-	-	-

Non-Hispanic 1/, male

2003.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	1	-	-	-	-	-	-	1	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Measles (B05)

Non-Hispanic white, both sexes

2003.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	1	-	-	-	-

Non-Hispanic white, male

2003.....	1	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic white, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	1	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Viral hepatitis (B15-B19)													
All origins, both sexes													
2003.....	5,431	1	-	-	14	64	652	2,259	1,085	698	543	115	-
2002.....	5,793	2	3	2	23	94	740	2,331	1,072	762	587	177	-
2001.....	5,585	1	-	3	20	74	797	2,060	1,014	836	613	166	1
2000.....	5,357	6	1	2	18	79	779	1,894	904	845	660	168	1
1999.....	4,853	2	3	3	14	88	722	1,506	786	881	651	197	-
All origins, male													
2003.....	3,546	-	-	-	7	42	461	1,666	729	352	249	40	-
2002.....	3,770	1	3	1	13	63	519	1,755	702	373	270	70	-
2001.....	3,539	1	-	2	11	42	553	1,570	643	396	263	57	1
2000.....	3,381	2	1	1	11	46	540	1,425	577	396	311	70	1
1999.....	3,043	2	2	2	10	53	520	1,138	491	426	317	82	-
All origins, female													
2003.....	1,885	1	-	-	7	22	191	593	356	346	294	75	-
2002.....	2,023	1	-	1	10	31	221	576	370	389	317	107	-
2001.....	2,046	-	-	1	9	32	244	490	371	440	350	109	-
2000.....	1,976	4	-	1	7	33	239	469	327	449	349	98	-
1999.....	1,810	-	1	1	4	35	202	368	295	455	334	115	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Viral hepatitis (B15-B19)													
Hispanic, both sexes													
2003.....	710	-	-	-	2	10	107	266	151	106	59	9	-
2002.....	758	-	-	-	5	17	101	291	164	102	63	15	-
2001.....	791	-	-	2	6	10	126	266	192	113	61	15	-
2000.....	708	3	-	-	3	14	126	218	147	120	66	11	-
1999.....	656	-	1	-	3	16	104	203	127	129	65	8	-
Hispanic, male													
2003.....	465	-	-	-	2	8	80	207	103	42	20	3	-
2002.....	491	-	-	-	5	14	75	218	108	40	25	6	-
2001.....	487	-	-	2	2	3	89	206	123	35	23	4	-
2000.....	456	1	-	-	2	11	100	165	94	52	25	6	-
1999.....	406	-	1	-	1	11	84	153	75	59	18	4	-
Hispanic, female													
2003.....	245	-	-	-	-	2	27	59	48	64	39	6	-
2002.....	267	-	-	-	-	3	26	73	56	62	38	9	-
2001.....	304	-	-	-	4	7	37	60	69	78	38	11	-
2000.....	252	2	-	-	1	3	26	53	53	68	41	5	-
1999.....	250	-	-	-	2	5	20	50	52	70	47	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Viral hepatitis (B15-B19)													
Non-Hispanic 1/, both sexes													
2003.....	4,704	1	-	-	12	54	545	1,984	931	587	484	106	-
2002.....	5,004	2	3	2	18	77	634	2,030	898	655	523	162	-
2001.....	4,776	1	-	1	14	64	669	1,785	817	721	552	151	1
2000.....	4,629	3	1	2	15	65	652	1,665	751	724	594	156	1
1999.....	4,176	2	2	3	11	71	613	1,294	656	750	585	189	-
Non-Hispanic 1/, male													
2003.....	3,068	-	-	-	5	34	381	1,451	625	306	229	37	-
2002.....	3,256	1	3	1	8	49	439	1,529	586	331	245	64	-
2001.....	3,038	1	-	-	9	39	462	1,357	515	361	240	53	1
2000.....	2,909	1	1	1	9	35	440	1,250	479	343	286	63	1
1999.....	2,620	2	1	2	9	41	432	978	414	365	298	78	-
Non-Hispanic 1/, female													
2003.....	1,636	1	-	-	7	20	164	533	306	281	255	69	-
2002.....	1,748	1	-	1	10	28	195	501	312	324	278	98	-
2001.....	1,738	-	-	1	5	25	207	428	302	360	312	98	-
2000.....	1,720	2	-	1	6	30	212	415	272	381	308	93	-
1999.....	1,556	-	1	1	2	30	181	316	242	385	287	111	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Viral hepatitis (B15-B19)													
Non-Hispanic white, both sexes													
2003.....	3,585	-	-	-	7	39	425	1,566	642	427	390	89	-
2002.....	3,857	1	-	2	15	49	496	1,590	635	489	440	140	-
2001.....	3,663	1	-	1	7	43	522	1,364	586	544	467	128	-
2000.....	3,513	1	-	2	5	37	501	1,243	520	559	506	138	1
1999.....	3,176	1	2	2	6	51	464	958	465	567	497	163	-
Non-Hispanic white, male													
2003.....	2,381	-	-	-	1	22	300	1,162	445	232	186	33	-
2002.....	2,551	1	-	1	6	33	351	1,212	415	263	214	55	-
2001.....	2,410	1	-	-	4	28	368	1,062	384	296	218	49	-
2000.....	2,234	-	-	1	4	19	342	958	339	265	248	57	1
1999.....	2,026	1	1	1	6	29	333	717	308	295	268	67	-
Non-Hispanic white, female													
2003.....	1,204	-	-	-	6	17	125	404	197	195	204	56	-
2002.....	1,306	-	-	1	9	16	145	378	220	226	226	85	-
2001.....	1,253	-	-	1	3	15	154	302	202	248	249	79	-
2000.....	1,279	1	-	1	1	18	159	285	181	294	258	81	-
1999.....	1,150	-	1	1	-	22	131	241	157	272	229	96	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Viral hepatitis (B15-B19)													
Non-Hispanic black, both sexes													
2003.....	842	-	-	-	4	11	97	352	221	89	57	11	-
2002.....	849	1	1	-	2	22	111	368	184	98	49	13	-
2001.....	819	-	-	-	6	15	121	349	167	91	53	16	1
2000.....	777	2	-	-	10	21	117	329	138	93	57	10	-
1999.....	718	1	-	1	4	17	119	277	118	109	53	19	-
Non-Hispanic black, male													
2003.....	536	-	-	-	3	9	63	250	145	41	21	4	-
2002.....	532	-	1	-	1	11	67	262	122	48	17	3	-
2001.....	479	-	-	-	4	7	79	240	99	34	13	2	1
2000.....	490	1	-	-	5	11	77	234	86	48	25	3	-
1999.....	433	1	-	1	2	9	79	215	65	38	15	8	-
Non-Hispanic black, female													
2003.....	306	-	-	-	1	2	34	102	76	48	36	7	-
2002.....	317	1	-	-	1	11	44	106	62	50	32	10	-
2001.....	340	-	-	-	2	8	42	109	68	57	40	14	-
2000.....	287	1	-	-	5	10	40	95	52	45	32	7	-
1999.....	285	-	-	-	2	8	40	62	53	71	38	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Viral hepatitis (B15-B19)

Origin not stated 2/, both sexes

2003.....	17	-	-	-	-	-	-	9	3	5	-	-	-
2002.....	31	-	-	-	-	-	5	10	10	5	1	-	-
2001.....	18	-	-	-	-	-	2	9	5	2	-	-	-
2000.....	20	-	-	-	-	-	1	11	6	1	-	1	-
1999.....	21	-	-	-	-	1	5	9	3	2	1	-	-

Origin not stated 2/, male

2003.....	13	-	-	-	-	-	-	8	1	4	-	-	-
2002.....	23	-	-	-	-	-	5	8	8	2	-	-	-
2001.....	14	-	-	-	-	-	2	7	5	-	-	-	-
2000.....	16	-	-	-	-	-	-	10	4	1	-	1	-
1999.....	17	-	-	-	-	1	4	7	2	2	1	-	-

Origin not stated 2/, female

2003.....	4	-	-	-	-	-	-	1	2	1	-	-	-
2002.....	8	-	-	-	-	-	-	2	2	3	1	-	-
2001.....	4	-	-	-	-	-	-	2	-	2	-	-	-
2000.....	4	-	-	-	-	-	1	1	2	-	-	-	-
1999.....	4	-	-	-	-	-	1	2	1	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
All origins, both sexes													
2003.....	13,658	5	2	36	178	1,588	5,340	4,442	1,517	438	95	14	3
2002.....	14,095	7	5	25	178	1,839	5,707	4,474	1,347	401	96	12	4
2001.....	14,175	10	13	38	225	2,101	5,867	4,120	1,317	384	88	10	2
2000.....	14,478	10	14	46	179	2,437	5,919	4,142	1,239	397	85	9	1
1999.....	14,802	14	24	66	198	2,729	6,232	3,907	1,149	398	70	14	1
All origins, male													
2003.....	10,119	2	2	19	95	1,038	3,884	3,438	1,215	336	75	12	3
2002.....	10,468	4	1	14	93	1,187	4,198	3,485	1,083	323	71	5	4
2001.....	10,590	7	8	23	105	1,421	4,360	3,241	1,051	292	73	7	2
2000.....	10,868	8	6	22	99	1,602	4,441	3,296	1,013	313	65	2	1
1999.....	11,234	9	16	36	95	1,914	4,706	3,134	948	317	50	8	1
All origins, female													
2003.....	3,539	3	-	17	83	550	1,456	1,004	302	102	20	2	-
2002.....	3,627	3	4	11	85	652	1,509	989	264	78	25	7	-
2001.....	3,585	3	5	15	120	680	1,507	879	266	92	15	3	-
2000.....	3,610	2	8	24	80	835	1,478	846	226	84	20	7	-
1999.....	3,568	5	8	30	103	815	1,526	773	201	81	20	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Hispanic, both sexes													
2003.....	1,890	-	1	2	31	244	727	606	204	64	10	1	-
2002.....	1,855	4	-	3	29	262	779	537	188	40	9	1	3
2001.....	1,853	1	-	3	32	300	759	526	169	51	10	2	-
2000.....	1,908	-	-	9	23	365	762	531	163	41	14	-	-
1999.....	1,958	3	5	9	33	407	855	455	137	49	4	1	-
Hispanic, male													
2003.....	1,465	-	1	-	20	184	550	499	153	50	7	1	-
2002.....	1,440	3	-	3	19	201	607	421	140	36	7	-	3
2001.....	1,437	-	-	2	21	222	589	420	129	43	9	2	-
2000.....	1,493	-	-	5	18	274	603	426	123	34	10	-	-
1999.....	1,525	1	4	3	20	314	666	354	118	41	4	-	-
Hispanic, female													
2003.....	425	-	-	2	11	60	177	107	51	14	3	-	-
2002.....	415	1	-	-	10	61	172	116	48	4	2	1	-
2001.....	416	1	-	1	11	78	170	106	40	8	1	-	-
2000.....	415	-	-	4	5	91	159	105	40	7	4	-	-
1999.....	433	2	1	6	13	93	189	101	19	8	-	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Non-Hispanic 1/, both sexes													
2003.....	11,629	5	1	34	147	1,333	4,561	3,785	1,298	366	84	13	2
2002.....	12,094	3	5	22	149	1,564	4,871	3,874	1,148	359	87	11	1
2001.....	12,141	9	13	35	193	1,779	5,044	3,533	1,123	325	77	8	2
2000.....	12,389	10	14	37	156	2,053	5,077	3,562	1,058	343	70	9	-
1999.....	12,638	11	19	56	165	2,301	5,305	3,377	987	339	65	13	-
Non-Hispanic 1/, male													
2003.....	8,550	2	1	19	75	846	3,299	2,900	1,050	278	67	11	2
2002.....	8,911	1	1	11	74	977	3,549	3,009	933	286	64	5	1
2001.....	9,011	7	8	21	84	1,187	3,725	2,766	902	241	63	5	2
2000.....	9,224	8	6	17	81	1,313	3,778	2,825	872	267	55	2	-
1999.....	9,538	8	12	33	75	1,584	3,984	2,715	806	268	45	8	-
Non-Hispanic 1/, female													
2003.....	3,079	3	-	15	72	487	1,262	885	248	88	17	2	-
2002.....	3,183	2	4	11	75	587	1,322	865	215	73	23	6	-
2001.....	3,130	2	5	14	109	592	1,319	767	221	84	14	3	-
2000.....	3,165	2	8	20	75	740	1,299	737	186	76	15	7	-
1999.....	3,100	3	7	23	90	717	1,321	662	181	71	20	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Non-Hispanic white, both sexes													
2003.....	4,126	1	-	3	28	375	1,703	1,360	485	125	38	7	1
2002.....	4,224	-	-	3	23	429	1,790	1,366	460	112	33	8	-
2001.....	4,286	2	3	8	30	519	1,911	1,248	423	110	27	5	-
2000.....	4,562	1	1	13	30	619	2,000	1,325	420	122	28	3	-
1999.....	4,744	2	2	15	34	794	2,082	1,286	369	121	32	7	-
Non-Hispanic white, male													
2003.....	3,487	1	-	2	17	294	1,429	1,172	428	103	35	5	1
2002.....	3,582	-	-	1	12	335	1,517	1,183	409	99	22	4	-
2001.....	3,669	2	-	5	15	410	1,639	1,106	370	95	24	3	-
2000.....	3,837	1	-	5	20	475	1,678	1,161	369	103	25	-	-
1999.....	4,043	2	1	10	22	643	1,768	1,136	331	104	24	2	-
Non-Hispanic white, female													
2003.....	639	-	-	1	11	81	274	188	57	22	3	2	-
2002.....	642	-	-	2	11	94	273	183	51	13	11	4	-
2001.....	617	-	3	3	15	109	272	142	53	15	3	2	-
2000.....	725	-	1	8	10	144	322	164	51	19	3	3	-
1999.....	701	-	1	5	12	151	314	150	38	17	8	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Non-Hispanic black, both sexes													
2003.....	7,351	3	1	31	117	938	2,786	2,382	802	239	45	6	1
2002.....	7,714	3	5	19	124	1,110	3,011	2,469	675	240	54	3	1
2001.....	7,710	7	10	26	161	1,234	3,072	2,249	690	208	49	2	2
2000.....	7,708	9	12	23	125	1,414	3,028	2,211	625	214	41	6	-
1999.....	7,751	9	17	41	130	1,479	3,154	2,057	613	213	33	5	-
Non-Hispanic black, male													
2003.....	4,947	1	1	17	57	537	1,814	1,693	614	175	31	6	1
2002.....	5,203	1	1	10	60	624	1,978	1,791	512	182	42	1	1
2001.....	5,227	5	8	15	68	753	2,040	1,629	524	144	38	1	2
2000.....	5,291	7	6	12	60	822	2,060	1,642	492	159	29	2	-
1999.....	5,377	6	11	23	52	919	2,157	1,552	471	160	21	5	-
Non-Hispanic black, female													
2003.....	2,404	2	-	14	60	401	972	689	188	64	14	-	-
2002.....	2,511	2	4	9	64	486	1,033	678	163	58	12	2	-
2001.....	2,483	2	2	11	93	481	1,032	620	166	64	11	1	-
2000.....	2,417	2	6	11	65	592	968	569	133	55	12	4	-
1999.....	2,374	3	6	18	78	560	997	505	142	53	12	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Origin not stated 2/, both sexes													
2003.....	139	-	-	-	-	11	52	51	15	8	1	-	1
2002.....	146	-	-	-	-	13	57	63	11	2	-	-	-
2001.....	181	-	-	-	-	22	64	61	25	8	1	-	-
2000.....	181	-	-	-	-	19	80	49	18	13	1	-	1
1999.....	206	-	-	1	-	21	72	75	25	10	1	-	1
Origin not stated 2/, male													
2003.....	104	-	-	-	-	8	35	39	12	8	1	-	1
2002.....	117	-	-	-	-	9	42	55	10	1	-	-	-
2001.....	142	-	-	-	-	12	46	55	20	8	1	-	-
2000.....	151	-	-	-	-	15	60	45	18	12	-	-	1
1999.....	171	-	-	-	-	16	56	65	24	8	1	-	1
Origin not stated 2/, female													
2003.....	35	-	-	-	-	3	17	12	3	-	-	-	-
2002.....	29	-	-	-	-	4	15	8	1	1	-	-	-
2001.....	39	-	-	-	-	10	18	6	5	-	-	-	-
2000.....	30	-	-	-	-	4	20	4	-	1	1	-	-
1999.....	35	-	-	1	-	5	16	10	1	2	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malaria (B50-B54)													
All origins, both sexes													
2003.....	4	-	-	-	-	-	1	-	2	1	-	-	-
2002.....	12	-	-	-	1	1	1	5	2	2	-	-	-
2001.....	9	-	-	1	-	-	2	2	2	1	1	-	-
2000.....	3	-	-	-	1	-	-	1	-	-	1	-	-
1999.....	7	1	-	-	-	2	1	-	1	2	-	-	-
All origins, male													
2003.....	4	-	-	-	-	-	1	-	2	1	-	-	-
2002.....	8	-	-	-	-	-	1	4	1	2	-	-	-
2001.....	5	-	-	1	-	-	-	1	1	1	1	-	-
2000.....	2	-	-	-	-	-	-	1	-	-	1	-	-
1999.....	4	1	-	-	-	1	-	-	1	1	-	-	-
All origins, female													
2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	-	-	-	1	1	-	1	1	-	-	-	-
2001.....	4	-	-	-	-	-	2	1	1	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	1	1	-	-	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malaria (B50-B54)													
Non-Hispanic 1/, both sexes													
2003.....	4	-	-	-	-	-	1	-	2	1	-	-	-
2002.....	12	-	-	-	1	1	1	5	2	2	-	-	-
2001.....	9	-	-	1	-	-	2	2	2	1	1	-	-
2000.....	3	-	-	-	1	-	-	1	-	-	1	-	-
1999.....	7	1	-	-	-	2	1	-	1	2	-	-	-
Non-Hispanic 1/, male													
2003.....	4	-	-	-	-	-	1	-	2	1	-	-	-
2002.....	8	-	-	-	-	-	1	4	1	2	-	-	-
2001.....	5	-	-	1	-	-	-	1	1	1	1	-	-
2000.....	2	-	-	-	-	-	-	1	-	-	1	-	-
1999.....	4	1	-	-	-	1	-	-	1	1	-	-	-
Non-Hispanic 1/, female													
2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	-	-	-	1	1	-	1	1	-	-	-	-
2001.....	4	-	-	-	-	-	2	1	1	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	1	1	-	-	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malaria (B50-B54)													
Non-Hispanic white, both sexes													
2003.....	4	-	-	-	-	-	1	-	2	1	-	-	-
2002.....	7	-	-	-	-	-	1	3	1	2	-	-	-
2001.....	5	-	-	-	-	-	2	1	-	1	1	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	4	-	-	-	-	1	1	-	1	1	-	-	-
Non-Hispanic white, male													
2003.....	4	-	-	-	-	-	1	-	2	1	-	-	-
2002.....	7	-	-	-	-	-	1	3	1	2	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	1	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	1	-	-	1	1	-	-	-
Non-Hispanic white, female													
2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	-	2	1	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	1	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malaria (B50-B54)

Non-Hispanic black, both sexes

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	-	-	-	1	1	-	2	-	-	-	-	-
2001.....	3	-	-	1	-	-	-	1	1	-	-	-	-
2000.....	1	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	3	1	-	-	-	1	-	-	-	1	-	-	-

Non-Hispanic black, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	2	-	-	1	-	-	-	1	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	1	1	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic black, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	-	1	1	-	1	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	1	-	-	-	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All origins, both sexes

2003.....	7,427	174	85	51	88	140	447	1,185	1,136	1,344	1,715	1,062	-
2002.....	6,707	224	61	84	97	169	453	916	889	1,223	1,594	997	-
2001.....	5,715	191	64	64	86	147	369	663	771	1,143	1,330	887	-
2000.....	5,532	167	71	67	96	141	350	585	707	1,074	1,375	899	-
1999.....	7,279	203	81	79	90	242	545	836	898	1,443	1,739	1,123	-

All origins, male

2003.....	3,964	86	43	25	52	83	284	792	689	711	816	383	-
2002.....	3,622	125	39	40	52	96	291	626	532	663	785	373	-
2001.....	2,931	101	37	29	51	83	227	426	449	565	656	307	-
2000.....	2,861	94	41	35	54	75	220	358	383	597	665	339	-
1999.....	3,655	119	51	44	49	131	333	478	482	760	827	381	-

All origins, female

2003.....	3,463	88	42	26	36	57	163	393	447	633	899	679	-
2002.....	3,085	99	22	44	45	73	162	290	357	560	809	624	-
2001.....	2,784	90	27	35	35	64	142	237	322	578	674	580	-
2000.....	2,671	73	30	32	42	66	130	227	324	477	710	560	-
1999.....	3,624	84	30	35	41	111	212	358	416	683	912	742	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

Hispanic, both sexes

2003.....	675	23	17	14	12	21	77	169	116	103	98	25	-
2002.....	593	41	11	17	19	51	67	127	85	75	68	32	-
2001.....	423	41	7	11	12	33	46	69	56	67	57	24	-
2000.....	412	23	18	13	19	26	50	61	54	62	57	29	-
1999.....	497	36	13	14	13	39	67	68	62	80	68	37	-

Hispanic, male

2003.....	416	9	8	7	8	14	57	130	76	53	43	11	-
2002.....	363	27	7	11	7	31	48	95	52	40	30	15	-
2001.....	250	21	4	5	9	22	31	47	38	38	27	8	-
2000.....	237	14	12	8	12	20	37	33	35	28	27	11	-
1999.....	260	22	10	7	6	22	49	40	29	36	29	10	-

Hispanic, female

2003.....	259	14	9	7	4	7	20	39	40	50	55	14	-
2002.....	230	14	4	6	12	20	19	32	33	35	38	17	-
2001.....	173	20	3	6	3	11	15	22	18	29	30	16	-
2000.....	175	9	6	5	7	6	13	28	19	34	30	18	-
1999.....	237	14	3	7	7	17	18	28	33	44	39	27	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic 1/, both sexes

2003.....	6,730	151	66	37	75	119	368	1,010	1,018	1,235	1,614	1,037	-
2002.....	6,097	181	49	67	77	117	384	787	801	1,148	1,522	964	-
2001.....	5,271	149	57	53	73	114	319	589	711	1,073	1,272	861	-
2000.....	5,100	144	53	54	75	115	297	521	649	1,012	1,312	868	-
1999.....	6,753	166	68	63	77	202	475	763	831	1,358	1,666	1,084	-

Non-Hispanic 1/, male

2003.....	3,536	77	33	18	43	69	227	659	613	654	771	372	-
2002.....	3,250	97	31	29	44	65	242	530	479	623	753	357	-
2001.....	2,666	79	33	24	42	61	193	374	408	525	628	299	-
2000.....	2,613	80	29	27	42	55	182	322	347	569	633	327	-
1999.....	3,379	96	41	36	43	108	282	435	449	722	797	370	-

Non-Hispanic 1/, female

2003.....	3,194	74	33	19	32	50	141	351	405	581	843	665	-
2002.....	2,847	84	18	38	33	52	142	257	322	525	769	607	-
2001.....	2,605	70	24	29	31	53	126	215	303	548	644	562	-
2000.....	2,487	64	24	27	33	60	115	199	302	443	679	541	-
1999.....	3,374	70	27	27	34	94	193	328	382	636	869	714	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

Non-Hispanic white, both sexes

2003.....	5,509	87	36	25	51	76	257	773	794	1,056	1,399	955	-
2002.....	4,939	113	34	49	47	66	243	582	606	973	1,344	882	-
2001.....	4,257	80	37	38	45	63	211	435	551	899	1,123	775	-
2000.....	4,131	79	33	40	51	58	189	383	505	851	1,165	777	-
1999.....	5,288	101	43	42	51	104	287	509	628	1,126	1,438	959	-

Non-Hispanic white, male

2003.....	2,871	42	17	12	31	43	163	512	478	554	675	344	-
2002.....	2,608	57	21	24	27	33	161	398	358	537	665	327	-
2001.....	2,148	40	21	17	24	39	129	281	320	443	562	272	-
2000.....	2,092	45	16	18	29	28	115	239	275	483	557	287	-
1999.....	2,644	64	27	24	28	55	190	303	345	608	681	319	-

Non-Hispanic white, female

2003.....	2,638	45	19	13	20	33	94	261	316	502	724	611	-
2002.....	2,331	56	13	25	20	33	82	184	248	436	679	555	-
2001.....	2,109	40	16	21	21	24	82	154	231	456	561	503	-
2000.....	2,039	34	17	22	22	30	74	144	230	368	608	490	-
1999.....	2,644	37	16	18	23	49	97	206	283	518	757	640	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic black, both sexes

2003.....	942	53	23	9	18	33	99	202	165	116	157	67	-
2002.....	927	58	13	10	25	46	121	179	151	122	130	72	-
2001.....	815	61	16	13	23	45	91	130	129	135	105	67	-
2000.....	815	60	14	10	21	50	95	124	116	129	120	76	-
1999.....	1,247	57	23	17	20	89	169	219	169	192	186	106	-

Non-Hispanic black, male

2003.....	515	30	12	5	9	20	57	124	97	73	66	22	-
2002.....	516	33	9	3	13	28	68	117	99	62	58	26	-
2001.....	419	36	10	6	15	19	56	80	70	66	42	19	-
2000.....	428	31	10	5	10	22	58	75	62	70	58	27	-
1999.....	617	29	14	10	12	48	81	114	82	94	94	39	-

Non-Hispanic black, female

2003.....	427	23	11	4	9	13	42	78	68	43	91	45	-
2002.....	411	25	4	7	12	18	53	62	52	60	72	46	-
2001.....	396	25	6	7	8	26	35	50	59	69	63	48	-
2000.....	387	29	4	5	11	28	37	49	54	59	62	49	-
1999.....	630	28	9	7	8	41	88	105	87	98	92	67	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Origin not stated 2/, both sexes

2003.....	22	-	2	-	1	-	2	6	2	6	3	-	-
2002.....	17	2	1	-	1	1	2	2	3	-	4	1	-
2001.....	21	1	-	-	1	-	4	5	4	3	1	2	-
2000.....	20	-	-	-	2	-	3	3	4	-	6	2	-
1999.....	29	1	-	2	-	1	3	5	5	5	5	2	-

Origin not stated 2/, male

2003.....	12	-	2	-	1	-	-	3	-	4	2	-	-
2002.....	9	1	1	-	1	-	1	1	1	-	2	1	-
2001.....	15	1	-	-	-	-	3	5	3	2	1	-	-
2000.....	11	-	-	-	-	-	1	3	1	-	5	1	-
1999.....	16	1	-	1	-	1	2	3	4	2	1	1	-

Origin not stated 2/, female

2003.....	10	-	-	-	-	-	2	3	2	2	1	-	-
2002.....	8	1	-	-	-	1	1	1	2	-	2	-	-
2001.....	6	-	-	-	1	-	1	-	1	1	-	2	-
2000.....	9	-	-	-	2	-	2	-	3	-	1	1	-
1999.....	13	-	-	1	-	-	1	2	1	3	4	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
All origins, both sexes													
2003.....	556,902	75	392	1,076	1,651	3,741	15,509	49,843	95,692	141,248	167,617	80,046	12
2002.....	557,271	74	402	1,072	1,730	3,872	16,085	49,637	93,391	144,757	167,062	79,182	7
2001.....	553,768	66	420	1,008	1,704	3,994	16,569	49,562	90,223	147,018	165,445	77,751	8
2000.....	553,091	92	420	1,014	1,713	3,916	16,520	48,034	89,005	150,131	165,099	77,136	11
1999.....	549,838	68	418	1,012	1,724	4,005	16,732	46,681	89,067	152,338	162,770	75,014	9
All origins, male													
2003.....	287,990	34	227	591	979	1,804	6,825	25,542	51,927	77,793	87,382	34,878	8
2002.....	288,768	42	216	600	1,016	1,868	7,056	25,192	51,115	80,085	86,952	34,621	5
2001.....	287,075	31	229	533	1,024	1,860	7,297	25,071	49,244	81,699	86,126	33,954	7
2000.....	286,082	50	233	569	1,022	1,854	7,344	24,215	48,427	83,191	85,907	33,260	10
1999.....	285,832	36	227	555	1,023	1,959	7,471	23,634	48,833	84,544	85,060	32,484	6
All origins, female													
2003.....	268,912	41	165	485	672	1,937	8,684	24,301	43,765	63,455	80,235	45,168	4
2002.....	268,503	32	186	472	714	2,004	9,029	24,445	42,276	64,672	80,110	44,561	2
2001.....	266,693	35	191	475	680	2,134	9,272	24,491	40,979	65,319	79,319	43,797	1
2000.....	267,009	42	187	445	691	2,062	9,176	23,819	40,578	66,940	79,192	43,876	1
1999.....	264,006	32	191	457	701	2,046	9,261	23,047	40,234	67,794	77,710	42,530	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
Hispanic, both sexes													
2003.....	24,070	13	98	217	288	522	1,355	3,081	4,482	5,929	5,701	2,383	1
2002.....	23,141	18	72	210	304	502	1,346	2,859	4,247	5,857	5,365	2,361	-
2001.....	22,371	14	84	177	289	497	1,396	2,815	4,126	5,724	5,068	2,181	-
2000.....	21,160	17	75	198	275	476	1,295	2,575	3,780	5,557	4,827	2,085	-
1999.....	20,233	16	72	176	259	466	1,204	2,516	3,686	5,394	4,449	1,995	-
Hispanic, male													
2003.....	12,671	4	60	123	186	274	566	1,550	2,442	3,347	3,035	1,084	-
2002.....	12,235	10	35	108	191	252	557	1,430	2,377	3,256	2,902	1,117	-
2001.....	11,825	4	44	90	199	233	615	1,372	2,202	3,219	2,814	1,033	-
2000.....	11,138	10	43	122	162	241	534	1,232	2,034	3,086	2,657	1,017	-
1999.....	10,668	11	34	92	158	222	504	1,237	2,060	3,045	2,403	902	-
Hispanic, female													
2003.....	11,399	9	38	94	102	248	789	1,531	2,040	2,582	2,666	1,299	1
2002.....	10,906	8	37	102	113	250	789	1,429	1,870	2,601	2,463	1,244	-
2001.....	10,546	10	40	87	90	264	781	1,443	1,924	2,505	2,254	1,148	-
2000.....	10,022	7	32	76	113	235	761	1,343	1,746	2,471	2,170	1,068	-
1999.....	9,565	5	38	84	101	244	700	1,279	1,626	2,349	2,046	1,093	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
Non-Hispanic 1/, both sexes													
2003.....	531,638	62	293	857	1,359	3,207	14,120	46,621	90,973	135,014	161,606	77,515	11
2002.....	532,712	55	329	861	1,418	3,350	14,693	46,628	88,886	138,545	161,303	76,639	5
2001.....	530,058	52	333	828	1,413	3,485	15,121	46,617	85,842	140,943	160,023	75,394	7
2000.....	530,531	75	341	812	1,432	3,431	15,170	45,305	84,977	144,200	159,906	74,871	11
1999.....	528,218	52	343	835	1,462	3,525	15,482	44,036	85,146	146,572	157,904	72,852	9
Non-Hispanic 1/, male													
2003.....	274,651	30	166	467	790	1,520	6,243	23,909	49,353	74,260	84,179	33,726	8
2002.....	275,770	31	181	492	821	1,610	6,475	23,682	48,587	76,621	83,839	33,427	4
2001.....	274,492	27	183	442	824	1,623	6,652	23,631	46,871	78,266	83,121	32,846	6
2000.....	274,185	40	186	446	858	1,607	6,787	22,902	46,253	79,878	83,047	32,171	10
1999.....	274,419	25	190	463	864	1,732	6,946	22,323	46,644	81,271	82,443	31,512	6
Non-Hispanic 1/, female													
2003.....	256,987	32	127	390	569	1,687	7,877	22,712	41,620	60,754	77,427	43,789	3
2002.....	256,942	24	148	369	597	1,740	8,218	22,946	40,299	61,924	77,464	43,212	1
2001.....	255,566	25	150	386	589	1,862	8,469	22,986	38,971	62,677	76,902	42,548	1
2000.....	256,346	35	155	366	574	1,824	8,383	22,403	38,724	64,322	76,859	42,700	1
1999.....	253,799	27	153	372	598	1,793	8,536	21,713	38,502	65,301	75,461	41,340	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
Non-Hispanic white, both sexes													
2003.....	457,127	51	221	637	1,039	2,354	10,723	36,168	75,733	116,198	144,012	69,981	10
2002.....	458,754	37	236	642	1,097	2,483	11,234	36,609	73,809	119,464	143,957	69,184	2
2001.....	456,709	42	268	607	1,076	2,594	11,551	36,836	71,389	121,529	142,760	68,053	4
2000.....	458,200	60	256	627	1,059	2,599	11,458	35,861	70,865	124,842	143,079	67,487	7
1999.....	456,315	37	281	608	1,070	2,682	11,848	34,944	70,788	127,119	141,295	65,636	7
Non-Hispanic white, male													
2003.....	236,150	25	123	348	607	1,159	4,862	18,601	41,026	64,019	74,968	30,404	8
2002.....	237,323	24	134	355	639	1,218	5,041	18,699	40,210	66,250	74,639	30,112	2
2001.....	236,004	21	144	310	632	1,260	5,160	18,679	38,884	67,554	73,885	29,471	4
2000.....	235,932	34	146	338	636	1,259	5,201	18,092	38,368	69,085	73,952	28,814	7
1999.....	236,384	20	160	345	629	1,341	5,425	17,588	38,658	70,518	73,535	28,161	4
Non-Hispanic white, female													
2003.....	220,977	26	98	289	432	1,195	5,861	17,567	34,707	52,179	69,044	39,577	2
2002.....	221,431	13	102	287	458	1,265	6,193	17,910	33,599	53,214	69,318	39,072	-
2001.....	220,705	21	124	297	444	1,334	6,391	18,157	32,505	53,975	68,875	38,582	-
2000.....	222,268	26	110	289	423	1,340	6,257	17,769	32,497	55,757	69,127	38,673	-
1999.....	219,931	17	121	263	441	1,341	6,423	17,356	32,130	56,601	67,760	37,475	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
Non-Hispanic black, both sexes													
2003.....	62,054	7	55	173	245	656	2,787	8,802	12,903	15,666	14,482	6,278	-
2002.....	61,996	12	71	168	242	663	2,870	8,376	12,693	16,084	14,518	6,296	3
2001.....	61,580	5	46	172	257	691	2,950	8,175	12,196	16,392	14,454	6,239	3
2000.....	61,390	11	66	147	297	641	3,072	8,068	12,019	16,485	14,244	6,337	3
1999.....	61,409	11	47	179	314	660	3,049	7,749	12,322	16,625	14,192	6,260	1
Non-Hispanic black, male													
2003.....	32,131	4	30	96	141	263	1,112	4,533	7,135	8,593	7,540	2,684	-
2002.....	32,287	4	33	103	137	287	1,158	4,203	7,106	8,835	7,669	2,750	2
2001.....	32,335	2	29	100	148	267	1,201	4,196	6,752	9,134	7,700	2,804	2
2000.....	32,493	4	31	86	178	259	1,292	4,152	6,763	9,235	7,689	2,802	2
1999.....	32,542	3	24	94	181	294	1,255	4,098	6,933	9,228	7,603	2,828	1
Non-Hispanic black, female													
2003.....	29,923	3	25	77	104	393	1,675	4,269	5,768	7,073	6,942	3,594	-
2002.....	29,709	8	38	65	105	376	1,712	4,173	5,587	7,249	6,849	3,546	1
2001.....	29,245	3	17	72	109	424	1,749	3,979	5,444	7,258	6,754	3,435	1
2000.....	28,897	7	35	61	119	382	1,780	3,916	5,256	7,250	6,555	3,535	1
1999.....	28,867	8	23	85	133	366	1,794	3,651	5,389	7,397	6,589	3,432	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms (C00-C97)													
Origin not stated 2/, both sexes													
2003.....	1,194	-	1	2	4	12	34	141	237	305	310	148	-
2002.....	1,418	1	1	1	8	20	46	150	258	355	394	182	2
2001.....	1,339	-	3	3	2	12	52	130	255	351	354	176	1
2000.....	1,400	-	4	4	6	9	55	154	248	374	366	180	-
1999.....	1,387	-	3	1	3	14	46	129	235	372	417	167	-
Origin not stated 2/, male													
2003.....	668	-	1	1	3	10	16	83	132	186	168	68	-
2002.....	763	1	-	-	4	6	24	80	151	208	211	77	1
2001.....	758	-	2	1	1	4	30	68	171	214	191	75	1
2000.....	759	-	4	1	2	6	23	81	140	227	203	72	-
1999.....	745	-	3	-	1	5	21	74	129	228	214	70	-
Origin not stated 2/, female													
2003.....	526	-	-	1	1	2	18	58	105	119	142	80	-
2002.....	655	-	1	1	4	14	22	70	107	147	183	105	1
2001.....	581	-	1	2	1	8	22	62	84	137	163	101	-
2000.....	641	-	-	3	4	3	32	73	108	147	163	108	-
1999.....	642	-	-	1	2	9	25	55	106	144	203	97	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
All origins, both sexes													
2003.....	7,778	-	4	8	30	59	274	1,131	1,826	1,874	1,692	879	1
2002.....	7,737	-	3	8	23	57	258	1,101	1,811	1,898	1,721	857	-
2001.....	7,702	-	-	4	29	56	283	1,076	1,740	1,912	1,787	814	1
2000.....	7,492	-	-	4	26	50	293	1,068	1,642	1,959	1,626	824	-
1999.....	7,486	-	2	6	27	43	291	1,061	1,616	1,896	1,744	800	-
All origins, male													
2003.....	5,298	-	2	4	21	39	195	921	1,423	1,304	999	389	1
2002.....	5,184	-	2	3	11	39	179	870	1,392	1,317	997	374	-
2001.....	5,156	-	-	1	12	35	202	846	1,372	1,327	1,025	335	1
2000.....	4,952	-	-	4	13	28	215	825	1,270	1,326	939	332	-
1999.....	4,944	-	1	3	20	27	204	820	1,231	1,308	1,007	323	-
All origins, female													
2003.....	2,480	-	2	4	9	20	79	210	403	570	693	490	-
2002.....	2,553	-	1	5	12	18	79	231	419	581	724	483	-
2001.....	2,546	-	-	3	17	21	81	230	368	585	762	479	-
2000.....	2,540	-	-	-	13	22	78	243	372	633	687	492	-
1999.....	2,542	-	1	3	7	16	87	241	385	588	737	477	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Hispanic, both sexes													
2003.....	321	-	1	-	4	5	14	50	75	80	63	29	-
2002.....	284	-	-	2	3	4	17	47	66	62	54	29	-
2001.....	311	-	-	-	4	1	14	56	69	78	58	31	-
2000.....	287	-	-	1	1	6	13	42	75	73	48	28	-
1999.....	282	-	-	1	-	1	18	41	66	72	63	20	-
Hispanic, male													
2003.....	233	-	1	-	2	4	8	41	64	56	43	14	-
2002.....	215	-	-	-	1	3	14	39	53	51	36	18	-
2001.....	219	-	-	-	2	1	11	41	55	58	39	12	-
2000.....	218	-	-	1	-	4	11	34	62	58	28	20	-
1999.....	204	-	-	1	-	-	9	33	55	62	35	9	-
Hispanic, female													
2003.....	88	-	-	-	2	1	6	9	11	24	20	15	-
2002.....	69	-	-	2	2	1	3	8	13	11	18	11	-
2001.....	92	-	-	-	2	-	3	15	14	20	19	19	-
2000.....	69	-	-	-	1	2	2	8	13	15	20	8	-
1999.....	78	-	-	-	-	1	9	8	11	10	28	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Non-Hispanic 1/, both sexes													
2003.....	7,437	-	3	8	26	53	260	1,079	1,743	1,791	1,627	846	1
2002.....	7,431	-	3	6	20	53	238	1,049	1,739	1,830	1,666	827	-
2001.....	7,370	-	-	4	25	55	268	1,018	1,666	1,829	1,724	781	-
2000.....	7,181	-	-	3	24	44	280	1,022	1,560	1,877	1,575	796	-
1999.....	7,181	-	2	5	27	42	272	1,018	1,542	1,818	1,678	777	-
Non-Hispanic 1/, male													
2003.....	5,050	-	1	4	19	34	187	878	1,351	1,246	955	374	1
2002.....	4,950	-	2	3	10	36	162	826	1,335	1,261	960	355	-
2001.....	4,920	-	-	1	10	34	190	804	1,313	1,264	982	322	-
2000.....	4,719	-	-	3	13	24	204	788	1,204	1,263	908	312	-
1999.....	4,725	-	1	2	20	27	194	785	1,172	1,241	970	313	-
Non-Hispanic 1/, female													
2003.....	2,387	-	2	4	7	19	73	201	392	545	672	472	-
2002.....	2,481	-	1	3	10	17	76	223	404	569	706	472	-
2001.....	2,450	-	-	3	15	21	78	214	353	565	742	459	-
2000.....	2,462	-	-	-	11	20	76	234	356	614	667	484	-
1999.....	2,456	-	1	3	7	15	78	233	370	577	708	464	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Non-Hispanic white, both sexes													
2003.....	6,079	-	2	5	15	36	182	774	1,385	1,483	1,419	777	1
2002.....	6,141	-	2	3	12	35	164	744	1,358	1,565	1,504	754	-
2001.....	6,052	-	-	2	18	36	165	736	1,288	1,524	1,560	723	-
2000.....	5,843	-	-	1	16	29	186	723	1,177	1,566	1,412	733	-
1999.....	5,826	-	2	3	14	29	176	677	1,180	1,527	1,497	721	-
Non-Hispanic white, male													
2003.....	4,043	-	-	2	11	21	145	634	1,055	1,014	819	341	1
2002.....	4,035	-	2	2	7	25	114	587	1,042	1,067	867	322	-
2001.....	3,961	-	-	1	6	22	120	586	1,004	1,045	880	297	-
2000.....	3,726	-	-	1	7	16	128	546	904	1,036	797	291	-
1999.....	3,746	-	1	1	9	17	125	523	892	1,043	850	285	-
Non-Hispanic white, female													
2003.....	2,036	-	2	3	4	15	37	140	330	469	600	436	-
2002.....	2,106	-	-	1	5	10	50	157	316	498	637	432	-
2001.....	2,091	-	-	1	12	14	45	150	284	479	680	426	-
2000.....	2,117	-	-	-	9	13	58	177	273	530	615	442	-
1999.....	2,080	-	1	2	5	12	51	154	288	484	647	436	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Non-Hispanic black, both sexes													
2003.....	1,100	-	1	3	9	10	63	242	309	261	150	52	-
2002.....	1,034	-	-	2	6	11	63	254	318	213	114	53	-
2001.....	1,056	-	-	1	4	6	71	229	318	256	129	42	-
2000.....	1,089	-	-	2	7	7	60	249	327	254	136	47	-
1999.....	1,131	-	-	2	8	8	67	287	316	248	150	45	-
Non-Hispanic black, male													
2003.....	831	-	1	2	6	9	32	197	260	201	99	24	-
2002.....	757	-	-	1	3	7	41	203	257	157	64	24	-
2001.....	773	-	-	-	2	5	45	179	263	184	78	17	-
2000.....	821	-	-	2	5	4	46	203	262	191	94	14	-
1999.....	819	-	-	1	6	6	47	224	243	169	102	21	-
Non-Hispanic black, female													
2003.....	269	-	-	1	3	1	31	45	49	60	51	28	-
2002.....	277	-	-	1	3	4	22	51	61	56	50	29	-
2001.....	283	-	-	1	2	1	26	50	55	72	51	25	-
2000.....	268	-	-	-	2	3	14	46	65	63	42	33	-
1999.....	312	-	-	1	2	2	20	63	73	79	48	24	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Origin not stated 2/, both sexes

2003.....	20	-	-	-	-	1	-	2	8	3	2	4	-
2002.....	22	-	-	-	-	-	3	5	6	6	1	1	-
2001.....	21	-	-	-	-	-	1	2	5	5	5	2	1
2000.....	24	-	-	-	1	-	-	4	7	9	3	-	-
1999.....	23	-	-	-	-	-	1	2	8	6	3	3	-

Origin not stated 2/, male

2003.....	15	-	-	-	-	1	-	2	8	2	1	1	-
2002.....	19	-	-	-	-	-	3	5	4	5	1	1	-
2001.....	17	-	-	-	-	-	1	1	4	5	4	1	1
2000.....	15	-	-	-	-	-	-	3	4	5	3	-	-
1999.....	15	-	-	-	-	-	1	2	4	5	2	1	-

Origin not stated 2/, female

2003.....	5	-	-	-	-	-	-	-	-	1	1	3	-
2002.....	3	-	-	-	-	-	-	-	2	1	-	-	-
2001.....	4	-	-	-	-	-	-	1	1	-	1	1	-
2000.....	9	-	-	-	1	-	-	1	3	4	-	-	-
1999.....	8	-	-	-	-	-	-	-	4	1	1	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
All origins, both sexes													
2003.....	12,860	-	-	-	5	29	288	1,407	2,940	3,619	3,392	1,180	-
2002.....	12,701	-	-	1	6	19	311	1,390	2,763	3,691	3,389	1,130	1
2001.....	12,530	-	-	-	6	29	309	1,408	2,554	3,779	3,307	1,137	1
2000.....	12,232	1	-	-	3	39	297	1,348	2,610	3,671	3,163	1,100	-
1999.....	11,918	-	-	-	-	37	293	1,262	2,531	3,694	2,986	1,114	1
All origins, male													
2003.....	9,921	-	-	-	4	23	240	1,202	2,483	2,851	2,464	654	-
2002.....	9,789	-	-	-	3	14	257	1,185	2,321	2,952	2,454	603	-
2001.....	9,607	-	-	-	5	26	255	1,204	2,137	2,989	2,365	625	1
2000.....	9,279	1	-	-	2	31	265	1,161	2,169	2,831	2,233	586	-
1999.....	9,094	-	-	-	-	29	255	1,094	2,081	2,925	2,136	573	1
All origins, female													
2003.....	2,939	-	-	-	1	6	48	205	457	768	928	526	-
2002.....	2,912	-	-	1	3	5	54	205	442	739	935	527	1
2001.....	2,923	-	-	-	1	3	54	204	417	790	942	512	-
2000.....	2,953	-	-	-	1	8	32	187	441	840	930	514	-
1999.....	2,824	-	-	-	-	8	38	168	450	769	850	541	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
Hispanic, both sexes													
2003.....	421	-	-	-	1	3	17	50	105	122	92	31	-
2002.....	411	-	-	-	1	1	23	53	80	120	91	42	-
2001.....	401	-	-	-	-	2	12	63	96	95	101	32	-
2000.....	384	-	-	-	-	2	10	38	86	120	98	30	-
1999.....	367	-	-	-	-	2	16	44	85	95	87	38	-
Hispanic, male													
2003.....	322	-	-	-	-	2	16	45	79	93	74	13	-
2002.....	329	-	-	-	1	1	19	49	75	96	66	22	-
2001.....	316	-	-	-	-	2	9	61	74	80	68	22	-
2000.....	292	-	-	-	-	2	8	34	64	92	72	20	-
1999.....	294	-	-	-	-	2	15	39	78	74	63	23	-
Hispanic, female													
2003.....	99	-	-	-	1	1	1	5	26	29	18	18	-
2002.....	82	-	-	-	-	-	4	4	5	24	25	20	-
2001.....	85	-	-	-	-	-	3	2	22	15	33	10	-
2000.....	92	-	-	-	-	-	2	4	22	28	26	10	-
1999.....	73	-	-	-	-	-	1	5	7	21	24	15	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
Non-Hispanic 1/, both sexes													
2003.....	12,415	-	-	-	4	26	271	1,354	2,826	3,491	3,297	1,146	-
2002.....	12,254	-	-	1	5	17	287	1,334	2,671	3,560	3,296	1,082	1
2001.....	12,085	-	-	-	6	27	296	1,339	2,447	3,664	3,201	1,104	1
2000.....	11,819	1	-	-	3	37	287	1,308	2,521	3,538	3,058	1,066	-
1999.....	11,520	-	-	-	-	35	276	1,215	2,439	3,587	2,893	1,074	1
Non-Hispanic 1/, male													
2003.....	9,580	-	-	-	4	21	224	1,154	2,396	2,753	2,389	639	-
2002.....	9,431	-	-	-	2	12	237	1,133	2,235	2,848	2,386	578	-
2001.....	9,254	-	-	-	5	24	245	1,138	2,052	2,894	2,292	603	1
2000.....	8,964	1	-	-	2	29	257	1,126	2,102	2,728	2,156	563	-
1999.....	8,775	-	-	-	-	27	239	1,053	1,996	2,841	2,068	550	1
Non-Hispanic 1/, female													
2003.....	2,835	-	-	-	-	5	47	200	430	738	908	507	-
2002.....	2,823	-	-	1	3	5	50	201	436	712	910	504	1
2001.....	2,831	-	-	-	1	3	51	201	395	770	909	501	-
2000.....	2,855	-	-	-	1	8	30	182	419	810	902	503	-
1999.....	2,745	-	-	-	-	8	37	162	443	746	825	524	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
Non-Hispanic white, both sexes													
2003.....	10,631	-	-	-	4	22	222	1,042	2,314	2,992	2,971	1,064	-
2002.....	10,484	-	-	1	5	15	227	1,076	2,173	3,027	2,961	999	-
2001.....	10,193	-	-	-	6	22	238	1,039	1,925	3,060	2,888	1,014	1
2000.....	10,005	-	-	-	3	36	239	1,013	1,997	3,017	2,714	986	-
1999.....	9,703	-	-	-	-	30	213	936	1,923	3,038	2,576	987	-
Non-Hispanic white, male													
2003.....	8,292	-	-	-	4	19	188	908	2,004	2,383	2,190	596	-
2002.....	8,169	-	-	-	2	11	196	938	1,849	2,465	2,171	537	-
2001.....	7,863	-	-	-	5	19	200	897	1,645	2,458	2,086	552	1
2000.....	7,680	-	-	-	2	29	221	895	1,702	2,366	1,941	524	-
1999.....	7,440	-	-	-	-	22	188	827	1,598	2,451	1,857	497	-
Non-Hispanic white, female													
2003.....	2,339	-	-	-	-	3	34	134	310	609	781	468	-
2002.....	2,315	-	-	1	3	4	31	138	324	562	790	462	-
2001.....	2,330	-	-	-	1	3	38	142	280	602	802	462	-
2000.....	2,325	-	-	-	1	7	18	118	295	651	773	462	-
1999.....	2,263	-	-	-	-	8	25	109	325	587	719	490	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
Non-Hispanic black, both sexes													
2003.....	1,578	-	-	-	-	3	47	285	463	435	273	72	-
2002.....	1,572	-	-	-	-	2	56	234	454	475	282	68	1
2001.....	1,676	-	-	-	-	3	54	264	466	534	278	77	-
2000.....	1,645	1	-	-	-	1	44	278	481	462	307	71	-
1999.....	1,629	-	-	-	-	3	53	261	472	486	280	73	1
Non-Hispanic black, male													
2003.....	1,122	-	-	-	-	2	36	222	351	313	162	36	-
2002.....	1,118	-	-	-	-	1	38	175	351	340	178	35	-
2001.....	1,222	-	-	-	-	3	41	210	363	379	185	41	-
2000.....	1,162	1	-	-	-	-	32	218	369	318	191	33	-
1999.....	1,187	-	-	-	-	3	44	209	365	340	183	42	1
Non-Hispanic black, female													
2003.....	456	-	-	-	-	1	11	63	112	122	111	36	-
2002.....	454	-	-	-	-	1	18	59	103	135	104	33	1
2001.....	454	-	-	-	-	-	13	54	103	155	93	36	-
2000.....	483	-	-	-	-	1	12	60	112	144	116	38	-
1999.....	442	-	-	-	-	-	9	52	107	146	97	31	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of esophagus (C15)													
Origin not stated 2/, both sexes													
2003.....	24	-	-	-	-	-	-	3	9	6	3	3	-
2002.....	36	-	-	-	-	1	1	3	12	11	2	6	-
2001.....	44	-	-	-	-	-	1	6	11	20	5	1	-
2000.....	29	-	-	-	-	-	-	2	3	13	7	4	-
1999.....	31	-	-	-	-	-	1	3	7	12	6	2	-
Origin not stated 2/, male													
2003.....	19	-	-	-	-	-	-	3	8	5	1	2	-
2002.....	29	-	-	-	-	1	1	3	11	8	2	3	-
2001.....	37	-	-	-	-	-	1	5	11	15	5	-	-
2000.....	23	-	-	-	-	-	-	1	3	11	5	3	-
1999.....	25	-	-	-	-	-	1	2	7	10	5	-	-
Origin not stated 2/, female													
2003.....	5	-	-	-	-	-	-	-	1	1	2	1	-
2002.....	7	-	-	-	-	-	-	-	1	3	-	3	-
2001.....	7	-	-	-	-	-	-	1	-	5	-	1	-
2000.....	6	-	-	-	-	-	-	1	-	2	2	1	-
1999.....	6	-	-	-	-	-	-	1	-	2	1	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
All origins, both sexes													
2003.....	12,110	-	-	1	19	127	480	1,210	1,874	2,793	3,549	2,057	-
2002.....	12,198	-	1	-	29	138	436	1,106	1,831	2,816	3,656	2,185	-
2001.....	12,319	-	-	-	20	122	464	1,127	1,771	3,021	3,793	2,001	-
2000.....	12,645	-	-	1	14	124	484	1,097	1,804	3,054	3,833	2,234	-
1999.....	12,711	-	-	1	13	136	505	1,101	1,860	3,152	3,860	2,083	-
All origins, male													
2003.....	7,055	-	-	1	7	63	262	767	1,282	1,826	1,980	867	-
2002.....	7,133	-	1	-	8	80	239	725	1,215	1,867	2,064	934	-
2001.....	7,191	-	-	-	9	71	253	731	1,238	1,928	2,148	813	-
2000.....	7,328	-	-	-	6	69	284	703	1,228	1,966	2,144	928	-
1999.....	7,459	-	-	1	6	76	283	746	1,252	2,005	2,239	851	-
All origins, female													
2003.....	5,055	-	-	-	12	64	218	443	592	967	1,569	1,190	-
2002.....	5,065	-	-	-	21	58	197	381	616	949	1,592	1,251	-
2001.....	5,128	-	-	-	11	51	211	396	533	1,093	1,645	1,188	-
2000.....	5,317	-	-	1	8	55	200	394	576	1,088	1,689	1,306	-
1999.....	5,252	-	-	-	7	60	222	355	608	1,147	1,621	1,232	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
Hispanic, both sexes													
2003.....	1,253	-	-	1	3	34	92	193	239	291	279	121	-
2002.....	1,185	-	-	-	6	38	97	154	218	259	283	130	-
2001.....	1,189	-	-	-	4	41	98	182	209	300	253	102	-
2000.....	1,111	-	-	1	4	24	105	151	167	306	254	99	-
1999.....	1,055	-	-	-	4	38	85	121	192	271	239	105	-
Hispanic, male													
2003.....	708	-	-	1	2	17	46	108	140	180	166	48	-
2002.....	671	-	-	-	2	26	44	85	128	172	150	64	-
2001.....	680	-	-	-	2	19	54	104	130	186	135	50	-
2000.....	655	-	-	-	1	16	52	78	110	205	144	49	-
1999.....	607	-	-	-	1	25	41	76	126	162	141	35	-
Hispanic, female													
2003.....	545	-	-	-	1	17	46	85	99	111	113	73	-
2002.....	514	-	-	-	4	12	53	69	90	87	133	66	-
2001.....	509	-	-	-	2	22	44	78	79	114	118	52	-
2000.....	456	-	-	1	3	8	53	73	57	101	110	50	-
1999.....	448	-	-	-	3	13	44	45	66	109	98	70	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
Non-Hispanic 1/, both sexes													
2003.....	10,827	-	-	-	16	91	388	1,014	1,630	2,496	3,260	1,932	-
2002.....	10,974	-	1	-	23	99	337	950	1,609	2,548	3,363	2,044	-
2001.....	11,099	-	-	-	16	81	365	943	1,558	2,712	3,529	1,895	-
2000.....	11,491	-	-	-	10	99	378	943	1,627	2,739	3,568	2,127	-
1999.....	11,615	-	-	1	9	98	420	970	1,662	2,867	3,615	1,973	-
Non-Hispanic 1/, male													
2003.....	6,326	-	-	-	5	45	216	656	1,138	1,641	1,808	817	-
2002.....	6,438	-	1	-	6	54	194	639	1,084	1,688	1,908	864	-
2001.....	6,492	-	-	-	7	52	199	625	1,105	1,737	2,007	760	-
2000.....	6,648	-	-	-	5	53	231	622	1,113	1,753	1,994	877	-
1999.....	6,827	-	-	1	5	51	242	662	1,122	1,834	2,096	814	-
Non-Hispanic 1/, female													
2003.....	4,501	-	-	-	11	46	172	358	492	855	1,452	1,115	-
2002.....	4,536	-	-	-	17	45	143	311	525	860	1,455	1,180	-
2001.....	4,607	-	-	-	9	29	166	318	453	975	1,522	1,135	-
2000.....	4,843	-	-	-	5	46	147	321	514	986	1,574	1,250	-
1999.....	4,788	-	-	-	4	47	178	308	540	1,033	1,519	1,159	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
Non-Hispanic white, both sexes													
2003.....	8,035	-	-	-	12	52	234	637	1,172	1,861	2,540	1,527	-
2002.....	8,124	-	1	-	15	55	205	616	1,137	1,861	2,613	1,621	-
2001.....	8,308	-	-	-	6	43	211	615	1,110	2,010	2,779	1,534	-
2000.....	8,600	-	-	-	5	53	213	625	1,180	2,031	2,781	1,712	-
1999.....	8,712	-	-	-	4	50	237	646	1,156	2,181	2,859	1,579	-
Non-Hispanic white, male													
2003.....	4,718	-	-	-	5	28	140	421	825	1,224	1,424	651	-
2002.....	4,864	-	1	-	3	31	123	431	791	1,265	1,523	696	-
2001.....	4,884	-	-	-	2	29	122	416	794	1,288	1,606	627	-
2000.....	5,033	-	-	-	2	29	129	424	825	1,330	1,580	714	-
1999.....	5,171	-	-	-	3	30	142	454	798	1,426	1,670	648	-
Non-Hispanic white, female													
2003.....	3,317	-	-	-	7	24	94	216	347	637	1,116	876	-
2002.....	3,260	-	-	-	12	24	82	185	346	596	1,090	925	-
2001.....	3,424	-	-	-	4	14	89	199	316	722	1,173	907	-
2000.....	3,567	-	-	-	3	24	84	201	355	701	1,201	998	-
1999.....	3,541	-	-	-	1	20	95	192	358	755	1,189	931	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
Non-Hispanic black, both sexes													
2003.....	2,017	-	-	-	1	22	101	271	347	458	528	289	-
2002.....	2,112	-	-	-	8	33	90	248	339	518	545	331	-
2001.....	2,051	-	-	-	5	25	121	237	324	505	548	286	-
2000.....	2,156	-	-	-	4	36	117	239	335	523	579	323	-
1999.....	2,183	-	-	1	3	36	127	237	373	507	580	319	-
Non-Hispanic black, male													
2003.....	1,175	-	-	-	-	10	53	183	239	310	279	101	-
2002.....	1,184	-	-	-	3	20	49	165	223	323	276	125	-
2001.....	1,177	-	-	-	2	14	63	161	220	332	293	92	-
2000.....	1,202	-	-	-	2	20	77	154	222	314	295	118	-
1999.....	1,252	-	-	1	1	19	72	158	241	309	325	126	-
Non-Hispanic black, female													
2003.....	842	-	-	-	1	12	48	88	108	148	249	188	-
2002.....	928	-	-	-	5	13	41	83	116	195	269	206	-
2001.....	874	-	-	-	3	11	58	76	104	173	255	194	-
2000.....	954	-	-	-	2	16	40	85	113	209	284	205	-
1999.....	931	-	-	-	2	17	55	79	132	198	255	193	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of stomach (C16)													
Origin not stated 2/, both sexes													
2003.....	30	-	-	-	-	2	-	3	5	6	10	4	-
2002.....	39	-	-	-	-	1	2	2	4	9	10	11	-
2001.....	31	-	-	-	-	-	1	2	4	9	11	4	-
2000.....	43	-	-	-	-	1	1	3	10	9	11	8	-
1999.....	41	-	-	-	-	-	-	10	6	14	6	5	-
Origin not stated 2/, male													
2003.....	21	-	-	-	-	1	-	3	4	5	6	2	-
2002.....	24	-	-	-	-	-	1	1	3	7	6	6	-
2001.....	19	-	-	-	-	-	-	2	3	5	6	3	-
2000.....	25	-	-	-	-	-	1	3	5	8	6	2	-
1999.....	25	-	-	-	-	-	-	8	4	9	2	2	-
Origin not stated 2/, female													
2003.....	9	-	-	-	-	1	-	-	1	1	4	2	-
2002.....	15	-	-	-	-	1	1	1	1	2	4	5	-
2001.....	12	-	-	-	-	-	1	-	1	4	5	1	-
2000.....	18	-	-	-	-	1	-	-	5	1	5	6	-
1999.....	16	-	-	-	-	-	-	2	2	5	4	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
All origins, both sexes													
2003.....	55,958	-	-	4	40	291	1,315	4,442	8,304	12,934	17,331	11,296	1
2002.....	56,741	1	-	1	50	290	1,425	4,573	8,421	13,392	17,498	11,089	1
2001.....	56,887	-	-	-	46	299	1,433	4,548	7,984	13,791	17,527	11,259	-
2000.....	57,477	-	-	3	40	272	1,369	4,237	7,940	14,101	18,053	11,462	-
1999.....	57,155	1	-	1	49	283	1,333	4,057	7,967	14,249	17,929	11,286	-
All origins, male													
2003.....	28,007	-	-	3	25	158	714	2,509	4,831	7,346	8,471	3,949	1
2002.....	28,501	1	-	1	34	161	797	2,564	5,035	7,580	8,504	3,823	1
2001.....	28,225	-	-	-	24	157	781	2,512	4,652	7,773	8,451	3,875	-
2000.....	28,462	-	-	2	24	153	759	2,359	4,608	7,924	8,760	3,873	-
1999.....	28,240	-	-	1	27	160	678	2,250	4,621	7,883	8,754	3,866	-
All origins, female													
2003.....	27,951	-	-	1	15	133	601	1,933	3,473	5,588	8,860	7,347	-
2002.....	28,240	-	-	-	16	129	628	2,009	3,386	5,812	8,994	7,266	-
2001.....	28,662	-	-	-	22	142	652	2,036	3,332	6,018	9,076	7,384	-
2000.....	29,015	-	-	1	16	119	610	1,878	3,332	6,177	9,293	7,589	-
1999.....	28,915	1	-	-	22	123	655	1,807	3,346	6,366	9,175	7,420	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Hispanic, both sexes													
2003.....	2,471	-	-	-	10	34	122	314	493	607	612	279	-
2002.....	2,364	-	-	1	2	27	121	282	454	605	573	299	-
2001.....	2,250	-	-	-	6	20	127	245	415	592	535	310	-
2000.....	2,117	-	-	-	8	33	118	263	395	530	497	273	-
1999.....	2,031	-	-	-	7	31	86	245	344	581	485	252	-
Hispanic, male													
2003.....	1,363	-	-	-	8	20	72	191	295	351	314	112	-
2002.....	1,287	-	-	1	-	13	62	170	291	351	295	104	-
2001.....	1,185	-	-	-	3	8	76	124	245	333	261	135	-
2000.....	1,124	-	-	-	3	17	52	140	227	301	267	117	-
1999.....	1,062	-	-	-	5	14	39	133	185	331	257	98	-
Hispanic, female													
2003.....	1,108	-	-	-	2	14	50	123	198	256	298	167	-
2002.....	1,077	-	-	-	2	14	59	112	163	254	278	195	-
2001.....	1,065	-	-	-	3	12	51	121	170	259	274	175	-
2000.....	993	-	-	-	5	16	66	123	168	229	230	156	-
1999.....	969	-	-	-	2	17	47	112	159	250	228	154	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Non-Hispanic 1/, both sexes													
2003.....	53,372	-	-	4	30	257	1,193	4,111	7,793	12,304	16,687	10,992	1
2002.....	54,244	1	-	-	48	259	1,299	4,281	7,948	12,758	16,882	10,767	1
2001.....	54,513	-	-	-	40	279	1,301	4,289	7,545	13,172	16,963	10,924	-
2000.....	55,244	-	-	3	32	238	1,247	3,963	7,523	13,544	17,527	11,167	-
1999.....	54,976	1	-	1	41	252	1,241	3,804	7,604	13,640	17,390	11,002	-
Non-Hispanic 1/, male													
2003.....	26,579	-	-	3	17	138	642	2,307	4,525	6,981	8,138	3,827	1
2002.....	27,140	1	-	-	34	146	730	2,388	4,731	7,208	8,188	3,713	1
2001.....	26,978	-	-	-	21	149	701	2,382	4,390	7,419	8,181	3,735	-
2000.....	27,287	-	-	2	21	135	705	2,215	4,373	7,609	8,476	3,751	-
1999.....	27,103	-	-	1	22	146	636	2,113	4,424	7,538	8,467	3,756	-
Non-Hispanic 1/, female													
2003.....	26,793	-	-	1	13	119	551	1,804	3,268	5,323	8,549	7,165	-
2002.....	27,104	-	-	-	14	113	569	1,893	3,217	5,550	8,694	7,054	-
2001.....	27,535	-	-	-	19	130	600	1,907	3,155	5,753	8,782	7,189	-
2000.....	27,957	-	-	1	11	103	542	1,748	3,150	5,935	9,051	7,416	-
1999.....	27,873	1	-	-	19	106	605	1,691	3,180	6,102	8,923	7,246	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Non-Hispanic white, both sexes													
2003.....	45,205	-	-	4	19	184	845	3,052	6,241	10,297	14,700	9,862	1
2002.....	46,147	1	-	-	36	206	947	3,271	6,401	10,782	14,821	9,681	1
2001.....	46,450	-	-	-	28	203	966	3,284	6,092	11,124	14,903	9,850	-
2000.....	47,317	-	-	3	23	167	898	3,034	6,085	11,543	15,519	10,045	-
1999.....	47,275	-	-	1	26	182	933	2,956	6,184	11,643	15,427	9,923	-
Non-Hispanic white, male													
2003.....	22,624	-	-	3	12	99	467	1,750	3,680	5,942	7,223	3,447	1
2002.....	23,119	1	-	-	24	114	534	1,853	3,859	6,189	7,211	3,333	1
2001.....	23,128	-	-	-	13	111	523	1,874	3,616	6,369	7,263	3,359	-
2000.....	23,369	-	-	2	13	101	515	1,732	3,561	6,553	7,539	3,353	-
1999.....	23,446	-	-	1	12	105	500	1,679	3,638	6,559	7,556	3,396	-
Non-Hispanic white, female													
2003.....	22,581	-	-	1	7	85	378	1,302	2,561	4,355	7,477	6,415	-
2002.....	23,028	-	-	-	12	92	413	1,418	2,542	4,593	7,610	6,348	-
2001.....	23,322	-	-	-	15	92	443	1,410	2,476	4,755	7,640	6,491	-
2000.....	23,948	-	-	1	10	66	383	1,302	2,524	4,990	7,980	6,692	-
1999.....	23,829	-	-	-	14	77	433	1,277	2,546	5,084	7,871	6,527	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Non-Hispanic black, both sexes													
2003.....	6,881	-	-	-	8	50	287	890	1,313	1,702	1,680	951	-
2002.....	6,817	-	-	-	10	38	283	857	1,297	1,656	1,756	920	-
2001.....	6,847	-	-	-	8	56	280	845	1,248	1,726	1,758	926	-
2000.....	6,820	-	-	-	6	54	290	819	1,215	1,720	1,732	984	-
1999.....	6,709	1	-	-	14	55	268	711	1,248	1,724	1,733	955	-
Non-Hispanic black, male													
2003.....	3,307	-	-	-	4	25	141	480	716	887	752	302	-
2002.....	3,322	-	-	-	8	23	156	452	717	840	822	304	-
2001.....	3,221	-	-	-	7	30	152	428	652	881	766	305	-
2000.....	3,307	-	-	-	5	26	160	430	676	893	792	325	-
1999.....	3,160	-	-	-	9	30	123	374	692	850	786	296	-
Non-Hispanic black, female													
2003.....	3,574	-	-	-	4	25	146	410	597	815	928	649	-
2002.....	3,495	-	-	-	2	15	127	405	580	816	934	616	-
2001.....	3,626	-	-	-	1	26	128	417	596	845	992	621	-
2000.....	3,513	-	-	-	1	28	130	389	539	827	940	659	-
1999.....	3,549	1	-	-	5	25	145	337	556	874	947	659	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Origin not stated 2/, both sexes													
2003.....	115	-	-	-	-	-	-	17	18	23	32	25	-
2002.....	133	-	-	-	-	4	5	10	19	29	43	23	-
2001.....	124	-	-	-	-	-	5	14	24	27	29	25	-
2000.....	116	-	-	-	-	1	4	11	22	27	29	22	-
1999.....	148	-	-	-	1	-	6	8	19	28	54	32	-
Origin not stated 2/, male													
2003.....	65	-	-	-	-	-	-	11	11	14	19	10	-
2002.....	74	-	-	-	-	2	5	6	13	21	21	6	-
2001.....	62	-	-	-	-	-	4	6	17	21	9	5	-
2000.....	51	-	-	-	-	1	2	4	8	14	17	5	-
1999.....	75	-	-	-	-	-	3	4	12	14	30	12	-
Origin not stated 2/, female													
2003.....	50	-	-	-	-	-	-	6	7	9	13	15	-
2002.....	59	-	-	-	-	2	-	4	6	8	22	17	-
2001.....	62	-	-	-	-	-	1	8	7	6	20	20	-
2000.....	65	-	-	-	-	-	2	7	14	13	12	17	-
1999.....	73	-	-	-	1	-	3	4	7	14	24	20	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
All origins, both sexes													
2003.....	14,706	6	23	14	36	89	395	2,223	2,765	3,630	3,949	1,576	-
2002.....	14,047	3	28	18	27	105	414	1,976	2,558	3,581	3,728	1,609	-
2001.....	13,351	5	18	15	29	75	401	1,868	2,351	3,440	3,649	1,500	-
2000.....	12,916	2	23	20	30	86	429	1,686	2,256	3,439	3,558	1,387	-
1999.....	12,382	8	29	11	43	78	444	1,492	2,091	3,491	3,425	1,270	-
All origins, male													
2003.....	9,482	2	18	9	20	53	288	1,779	1,995	2,330	2,301	687	-
2002.....	9,011	2	19	12	18	62	308	1,611	1,858	2,367	2,112	642	-
2001.....	8,483	3	7	7	15	43	302	1,490	1,691	2,224	2,047	654	-
2000.....	8,176	1	14	8	20	59	297	1,324	1,607	2,246	2,052	548	-
1999.....	7,689	5	20	9	24	48	318	1,161	1,465	2,284	1,877	478	-
All origins, female													
2003.....	5,224	4	5	5	16	36	107	444	770	1,300	1,648	889	-
2002.....	5,036	1	9	6	9	43	106	365	700	1,214	1,616	967	-
2001.....	4,868	2	11	8	14	32	99	378	660	1,216	1,602	846	-
2000.....	4,740	1	9	12	10	27	132	362	649	1,193	1,506	839	-
1999.....	4,693	3	9	2	19	30	126	331	626	1,207	1,548	792	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
Hispanic, both sexes													
2003.....	1,511	-	10	1	1	11	48	287	337	397	319	100	-
2002.....	1,363	-	3	-	1	8	48	217	310	403	265	108	-
2001.....	1,256	1	3	3	2	14	46	226	261	327	289	84	-
2000.....	1,176	1	3	7	4	7	59	176	234	335	262	88	-
1999.....	1,084	1	4	2	4	7	43	170	214	324	239	76	-
Hispanic, male													
2003.....	986	-	8	1	1	6	37	239	241	245	172	36	-
2002.....	891	-	2	-	1	4	33	179	225	256	146	45	-
2001.....	812	1	1	-	1	7	36	187	187	206	152	34	-
2000.....	762	1	2	4	3	5	45	141	165	208	152	36	-
1999.....	688	1	3	2	3	5	27	141	159	195	121	31	-
Hispanic, female													
2003.....	525	-	2	-	-	5	11	48	96	152	147	64	-
2002.....	472	-	1	-	-	4	15	38	85	147	119	63	-
2001.....	444	-	2	3	1	7	10	39	74	121	137	50	-
2000.....	414	-	1	3	1	2	14	35	69	127	110	52	-
1999.....	396	-	1	-	1	2	16	29	55	129	118	45	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
Non-Hispanic 1/, both sexes													
2003.....	13,157	6	13	13	35	77	346	1,927	2,418	3,222	3,626	1,474	-
2002.....	12,638	3	25	18	26	96	365	1,751	2,234	3,171	3,452	1,497	-
2001.....	12,053	4	15	12	27	61	352	1,637	2,079	3,103	3,352	1,411	-
2000.....	11,702	1	20	13	26	78	368	1,502	2,018	3,091	3,287	1,298	-
1999.....	11,265	7	23	9	39	71	398	1,320	1,869	3,158	3,180	1,191	-
Non-Hispanic 1/, male													
2003.....	8,469	2	10	8	19	46	250	1,531	1,747	2,078	2,128	650	-
2002.....	8,085	2	17	12	17	57	274	1,426	1,621	2,106	1,957	596	-
2001.....	7,640	2	6	7	14	36	263	1,299	1,495	2,009	1,889	620	-
2000.....	7,384	-	12	4	17	53	251	1,175	1,440	2,028	1,893	511	-
1999.....	6,976	4	15	7	21	43	290	1,019	1,300	2,081	1,752	444	-
Non-Hispanic 1/, female													
2003.....	4,688	4	3	5	16	31	96	396	671	1,144	1,498	824	-
2002.....	4,553	1	8	6	9	39	91	325	613	1,065	1,495	901	-
2001.....	4,413	2	9	5	13	25	89	338	584	1,094	1,463	791	-
2000.....	4,318	1	8	9	9	25	117	327	578	1,063	1,394	787	-
1999.....	4,289	3	8	2	18	28	108	301	569	1,077	1,428	747	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
Non-Hispanic white, both sexes													
2003.....	10,149	4	9	7	22	34	216	1,271	1,741	2,506	3,071	1,268	-
2002.....	9,878	2	19	13	18	49	227	1,167	1,613	2,504	2,970	1,296	-
2001.....	9,433	4	11	5	20	34	189	1,089	1,505	2,477	2,857	1,242	-
2000.....	9,121	-	14	8	16	37	201	987	1,447	2,448	2,824	1,139	-
1999.....	8,890	5	20	4	25	29	238	880	1,340	2,546	2,755	1,048	-
Non-Hispanic white, male													
2003.....	6,504	2	7	6	12	17	146	1,005	1,270	1,654	1,825	560	-
2002.....	6,215	2	12	7	9	23	168	951	1,167	1,672	1,681	523	-
2001.....	5,956	2	4	4	11	17	137	859	1,088	1,651	1,632	551	-
2000.....	5,679	-	9	4	10	22	134	759	1,020	1,624	1,655	442	-
1999.....	5,455	2	13	3	14	16	173	660	934	1,702	1,534	404	-
Non-Hispanic white, female													
2003.....	3,645	2	2	1	10	17	70	266	471	852	1,246	708	-
2002.....	3,663	-	7	6	9	26	59	216	446	832	1,289	773	-
2001.....	3,477	2	7	1	9	17	52	230	417	826	1,225	691	-
2000.....	3,442	-	5	4	6	15	67	228	427	824	1,169	697	-
1999.....	3,435	3	7	1	11	13	65	220	406	844	1,221	644	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
Non-Hispanic black, both sexes													
2003.....	1,851	2	1	3	7	25	77	462	444	385	304	141	-
2002.....	1,708	1	4	3	4	26	83	400	385	384	292	126	-
2001.....	1,649	-	3	5	4	15	95	385	373	357	299	113	-
2000.....	1,581	1	2	2	6	20	101	360	344	383	262	100	-
1999.....	1,498	1	3	4	7	21	91	291	312	395	265	108	-
Non-Hispanic black, male													
2003.....	1,215	-	-	2	4	15	58	371	316	234	155	60	-
2002.....	1,143	-	3	3	4	16	60	321	279	261	158	38	-
2001.....	1,036	-	2	2	1	10	70	302	266	204	142	37	-
2000.....	1,047	-	1	-	5	14	65	285	256	249	131	41	-
1999.....	939	1	2	3	2	14	65	238	210	245	133	26	-
Non-Hispanic black, female													
2003.....	636	2	1	1	3	10	19	91	128	151	149	81	-
2002.....	565	1	1	-	-	10	23	79	106	123	134	88	-
2001.....	613	-	1	3	3	5	25	83	107	153	157	76	-
2000.....	534	1	1	2	1	6	36	75	88	134	131	59	-
1999.....	559	-	1	1	5	7	26	53	102	150	132	82	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
Origin not stated 2/, both sexes													
2003.....	38	-	-	-	-	1	1	9	10	11	4	2	-
2002.....	46	-	-	-	-	1	1	8	14	7	11	4	-
2001.....	42	-	-	-	-	-	3	5	11	10	8	5	-
2000.....	38	-	-	-	-	1	2	8	4	13	9	1	-
1999.....	33	-	2	-	-	-	3	2	8	9	6	3	-
Origin not stated 2/, male													
2003.....	27	-	-	-	-	1	1	9	7	7	1	1	-
2002.....	35	-	-	-	-	1	1	6	12	5	9	1	-
2001.....	31	-	-	-	-	-	3	4	9	9	6	-	-
2000.....	30	-	-	-	-	1	1	8	2	10	7	1	-
1999.....	25	-	2	-	-	-	1	1	6	8	4	3	-
Origin not stated 2/, female													
2003.....	11	-	-	-	-	-	-	-	3	4	3	1	-
2002.....	11	-	-	-	-	-	-	2	2	2	2	3	-
2001.....	11	-	-	-	-	-	-	1	2	1	2	5	-
2000.....	8	-	-	-	-	-	1	-	2	3	2	-	-
1999.....	8	-	-	-	-	-	2	1	2	1	2	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
All origins, both sexes													
2003.....	30,777	-	-	1	10	60	548	2,540	5,320	8,104	9,708	4,486	-
2002.....	30,264	1	-	-	7	39	596	2,433	5,105	8,187	9,555	4,340	1
2001.....	29,803	-	1	2	2	55	574	2,410	4,868	8,158	9,322	4,410	1
2000.....	29,332	2	1	1	5	55	548	2,362	4,708	8,103	9,459	4,087	1
1999.....	29,082	1	-	-	3	55	594	2,353	4,707	8,293	8,933	4,142	1
All origins, male													
2003.....	15,060	-	-	1	6	35	329	1,548	3,106	4,251	4,402	1,382	-
2002.....	14,877	-	-	-	5	26	364	1,526	3,012	4,270	4,270	1,403	1
2001.....	14,467	-	1	-	2	30	321	1,467	2,874	4,215	4,191	1,365	1
2000.....	14,238	1	1	-	3	33	324	1,486	2,749	4,189	4,189	1,262	1
1999.....	14,176	-	-	-	1	33	369	1,482	2,721	4,347	3,940	1,283	-
All origins, female													
2003.....	15,717	-	-	-	4	25	219	992	2,214	3,853	5,306	3,104	-
2002.....	15,387	1	-	-	2	13	232	907	2,093	3,917	5,285	2,937	-
2001.....	15,336	-	-	2	-	25	253	943	1,994	3,943	5,131	3,045	-
2000.....	15,094	1	-	1	2	22	224	876	1,959	3,914	5,270	2,825	-
1999.....	14,906	1	-	-	2	22	225	871	1,986	3,946	4,993	2,859	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
Hispanic, both sexes													
2003.....	1,485	-	-	-	-	10	49	157	300	429	386	154	-
2002.....	1,363	-	-	-	1	4	51	148	273	410	333	143	-
2001.....	1,336	-	-	-	1	6	46	127	263	419	340	134	-
2000.....	1,243	1	-	-	2	6	37	128	234	397	313	125	-
1999.....	1,135	-	-	-	-	6	44	137	218	338	272	120	-
Hispanic, male													
2003.....	744	-	-	-	-	5	30	95	176	217	175	46	-
2002.....	667	-	-	-	1	2	35	96	147	214	136	36	-
2001.....	643	-	-	-	1	3	26	73	141	212	154	33	-
2000.....	620	1	-	-	1	2	21	80	133	214	128	40	-
1999.....	563	-	-	-	-	2	30	84	117	182	108	40	-
Hispanic, female													
2003.....	741	-	-	-	-	5	19	62	124	212	211	108	-
2002.....	696	-	-	-	-	2	16	52	126	196	197	107	-
2001.....	693	-	-	-	-	3	20	54	122	207	186	101	-
2000.....	623	-	-	-	1	4	16	48	101	183	185	85	-
1999.....	572	-	-	-	-	4	14	53	101	156	164	80	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
Non-Hispanic 1/, both sexes													
2003.....	29,233	-	-	1	10	50	498	2,379	5,004	7,665	9,302	4,324	-
2002.....	28,839	1	-	-	6	35	545	2,278	4,821	7,763	9,200	4,189	1
2001.....	28,407	-	1	2	1	48	526	2,279	4,598	7,719	8,963	4,269	1
2000.....	28,026	1	1	1	3	49	510	2,228	4,465	7,692	9,124	3,951	1
1999.....	27,869	1	-	-	3	49	547	2,207	4,480	7,928	8,638	4,015	1
Non-Hispanic 1/, male													
2003.....	14,285	-	-	1	6	30	299	1,452	2,921	4,028	4,216	1,332	-
2002.....	14,181	-	-	-	4	24	329	1,428	2,856	4,051	4,126	1,362	1
2001.....	13,792	-	1	-	1	27	294	1,393	2,728	3,989	4,031	1,327	1
2000.....	13,590	-	1	-	2	31	302	1,401	2,609	3,970	4,054	1,219	1
1999.....	13,572	-	-	-	1	31	338	1,391	2,598	4,150	3,822	1,241	-
Non-Hispanic 1/, female													
2003.....	14,948	-	-	-	4	20	199	927	2,083	3,637	5,086	2,992	-
2002.....	14,658	1	-	-	2	11	216	850	1,965	3,712	5,074	2,827	-
2001.....	14,615	-	-	2	-	21	232	886	1,870	3,730	4,932	2,942	-
2000.....	14,436	1	-	1	1	18	208	827	1,856	3,722	5,070	2,732	-
1999.....	14,297	1	-	-	2	18	209	816	1,882	3,778	4,816	2,774	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
Non-Hispanic white, both sexes													
2003.....	24,919	-	-	-	7	39	372	1,830	4,097	6,503	8,206	3,865	-
2002.....	24,647	-	-	-	6	25	424	1,789	3,971	6,533	8,134	3,765	-
2001.....	24,424	-	1	2	1	36	415	1,816	3,847	6,564	7,931	3,810	1
2000.....	24,034	1	1	1	2	39	391	1,797	3,690	6,545	8,066	3,500	1
1999.....	23,903	-	-	-	2	34	421	1,771	3,675	6,784	7,625	3,590	1
Non-Hispanic white, male													
2003.....	12,277	-	-	-	3	21	222	1,125	2,434	3,490	3,777	1,205	-
2002.....	12,242	-	-	-	4	17	257	1,141	2,392	3,480	3,715	1,236	-
2001.....	11,994	-	1	-	1	22	233	1,131	2,324	3,441	3,624	1,216	1
2000.....	11,764	-	1	-	2	25	236	1,141	2,183	3,444	3,638	1,093	1
1999.....	11,684	-	-	-	1	23	266	1,114	2,144	3,605	3,412	1,119	-
Non-Hispanic white, female													
2003.....	12,642	-	-	-	4	18	150	705	1,663	3,013	4,429	2,660	-
2002.....	12,405	-	-	-	2	8	167	648	1,579	3,053	4,419	2,529	-
2001.....	12,430	-	-	2	-	14	182	685	1,523	3,123	4,307	2,594	-
2000.....	12,270	1	-	1	-	14	155	656	1,507	3,101	4,428	2,407	-
1999.....	12,219	-	-	-	1	11	155	657	1,531	3,179	4,213	2,471	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
Non-Hispanic black, both sexes													
2003.....	3,579	-	-	1	3	10	103	477	743	958	903	381	-
2002.....	3,466	1	-	-	-	6	101	399	726	1,016	858	358	1
2001.....	3,352	-	-	-	-	9	89	399	638	961	865	391	-
2000.....	3,391	-	-	-	1	9	97	379	664	970	890	381	-
1999.....	3,368	1	-	-	-	12	112	388	669	972	846	368	-
Non-Hispanic black, male													
2003.....	1,662	-	-	1	3	8	64	292	403	448	343	100	-
2002.....	1,606	-	-	-	-	5	58	242	399	475	332	94	1
2001.....	1,501	-	-	-	-	4	47	229	339	460	335	87	-
2000.....	1,546	-	-	-	-	5	54	229	367	447	350	94	-
1999.....	1,601	-	-	-	-	8	63	246	386	457	344	97	-
Non-Hispanic black, female													
2003.....	1,917	-	-	-	-	2	39	185	340	510	560	281	-
2002.....	1,860	1	-	-	-	1	43	157	327	541	526	264	-
2001.....	1,851	-	-	-	-	5	42	170	299	501	530	304	-
2000.....	1,845	-	-	-	1	4	43	150	297	523	540	287	-
1999.....	1,767	1	-	-	-	4	49	142	283	515	502	271	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of pancreas (C25)													
Origin not stated 2/, both sexes													
2003.....	59	-	-	-	-	-	1	4	16	10	20	8	-
2002.....	62	-	-	-	-	-	-	7	11	14	22	8	-
2001.....	60	-	-	-	-	1	2	4	7	20	19	7	-
2000.....	63	-	-	-	-	-	1	6	9	14	22	11	-
1999.....	78	-	-	-	-	-	3	9	9	27	23	7	-
Origin not stated 2/, male													
2003.....	31	-	-	-	-	-	-	1	9	6	11	4	-
2002.....	29	-	-	-	-	-	-	2	9	5	8	5	-
2001.....	32	-	-	-	-	-	1	1	5	14	6	5	-
2000.....	28	-	-	-	-	-	1	5	7	5	7	3	-
1999.....	41	-	-	-	-	-	1	7	6	15	10	2	-
Origin not stated 2/, female													
2003.....	28	-	-	-	-	-	1	3	7	4	9	4	-
2002.....	33	-	-	-	-	-	-	5	2	9	14	3	-
2001.....	28	-	-	-	-	1	1	3	2	6	13	2	-
2000.....	35	-	-	-	-	-	-	1	2	9	15	8	-
1999.....	37	-	-	-	-	-	2	2	3	12	13	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
All origins, both sexes													
2003.....	3,792	-	-	-	-	1	79	420	956	1,136	891	308	1
2002.....	3,723	-	-	-	-	11	76	425	904	1,084	969	253	1
2001.....	3,797	-	-	-	-	3	86	419	913	1,170	910	296	-
2000.....	3,861	-	-	2	1	4	64	426	873	1,219	996	276	-
1999.....	3,815	-	-	-	-	6	78	405	904	1,239	892	291	-
All origins, male													
2003.....	3,020	-	-	-	-	1	60	354	777	899	708	221	-
2002.....	2,922	-	-	-	-	6	58	349	738	832	755	183	1
2001.....	3,018	-	-	-	-	2	66	346	737	919	710	238	-
2000.....	3,041	-	-	-	-	2	49	340	726	945	765	214	-
1999.....	2,992	-	-	-	-	5	64	346	735	948	681	213	-
All origins, female													
2003.....	772	-	-	-	-	-	19	66	179	237	183	87	1
2002.....	801	-	-	-	-	5	18	76	166	252	214	70	-
2001.....	779	-	-	-	-	1	20	73	176	251	200	58	-
2000.....	820	-	-	2	1	2	15	86	147	274	231	62	-
1999.....	823	-	-	-	-	1	14	59	169	291	211	78	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
Hispanic, both sexes													
2003.....	162	-	-	-	-	-	-	21	44	44	38	15	-
2002.....	165	-	-	-	-	1	8	19	43	48	33	13	-
2001.....	163	-	-	-	-	-	3	20	38	60	29	13	-
2000.....	143	-	-	1	-	1	4	17	29	45	35	11	-
1999.....	171	-	-	-	-	-	5	25	29	55	40	17	-
Hispanic, male													
2003.....	148	-	-	-	-	-	-	19	43	41	31	14	-
2002.....	139	-	-	-	-	-	7	16	40	42	27	7	-
2001.....	141	-	-	-	-	-	3	18	34	53	24	9	-
2000.....	124	-	-	-	-	-	3	16	28	37	30	10	-
1999.....	149	-	-	-	-	-	4	24	27	48	31	15	-
Hispanic, female													
2003.....	14	-	-	-	-	-	-	2	1	3	7	1	-
2002.....	26	-	-	-	-	1	1	3	3	6	6	6	-
2001.....	22	-	-	-	-	-	-	2	4	7	5	4	-
2000.....	19	-	-	1	-	1	1	1	1	8	5	1	-
1999.....	22	-	-	-	-	-	1	1	2	7	9	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
Non-Hispanic 1/, both sexes													
2003.....	3,610	-	-	-	-	1	78	396	908	1,086	847	293	1
2002.....	3,548	-	-	-	-	10	68	405	857	1,033	934	240	1
2001.....	3,614	-	-	-	-	3	82	398	865	1,106	877	283	-
2000.....	3,699	-	-	1	1	3	59	406	840	1,166	959	264	-
1999.....	3,627	-	-	-	-	6	73	377	871	1,180	847	273	-
Non-Hispanic 1/, male													
2003.....	2,853	-	-	-	-	1	59	332	730	853	671	207	-
2002.....	2,775	-	-	-	-	6	51	332	695	788	726	176	1
2001.....	2,858	-	-	-	-	2	62	327	694	862	682	229	-
2000.....	2,901	-	-	-	-	2	45	322	694	901	733	204	-
1999.....	2,832	-	-	-	-	5	60	320	706	897	646	198	-
Non-Hispanic 1/, female													
2003.....	757	-	-	-	-	-	19	64	178	233	176	86	1
2002.....	773	-	-	-	-	4	17	73	162	245	208	64	-
2001.....	756	-	-	-	-	1	20	71	171	244	195	54	-
2000.....	798	-	-	1	1	1	14	84	146	265	226	60	-
1999.....	795	-	-	-	-	1	13	57	165	283	201	75	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
Non-Hispanic white, both sexes													
2003.....	2,879	-	-	-	-	1	61	269	697	878	722	250	1
2002.....	2,838	-	-	-	-	6	40	276	653	850	793	220	-
2001.....	2,876	-	-	-	-	3	58	268	657	893	747	250	-
2000.....	2,940	-	-	1	1	3	44	281	612	936	822	240	-
1999.....	2,924	-	-	-	-	4	40	256	660	978	751	235	-
Non-Hispanic white, male													
2003.....	2,251	-	-	-	-	1	48	227	559	676	564	176	-
2002.....	2,205	-	-	-	-	3	30	230	518	645	619	160	-
2001.....	2,267	-	-	-	-	2	47	216	529	697	570	206	-
2000.....	2,291	-	-	-	-	2	31	220	500	721	632	185	-
1999.....	2,263	-	-	-	-	3	34	219	534	739	568	166	-
Non-Hispanic white, female													
2003.....	628	-	-	-	-	-	13	42	138	202	158	74	1
2002.....	633	-	-	-	-	3	10	46	135	205	174	60	-
2001.....	609	-	-	-	-	1	11	52	128	196	177	44	-
2000.....	649	-	-	1	1	1	13	61	112	215	190	55	-
1999.....	661	-	-	-	-	1	6	37	126	239	183	69	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
Non-Hispanic black, both sexes													
2003.....	687	-	-	-	-	-	16	120	206	193	114	38	-
2002.....	674	-	-	-	-	4	27	123	198	174	131	16	1
2001.....	668	-	-	-	-	-	23	124	195	188	111	27	-
2000.....	714	-	-	-	-	-	15	121	218	209	129	22	-
1999.....	657	-	-	-	-	2	32	117	203	188	89	26	-
Non-Hispanic black, male													
2003.....	566	-	-	-	-	-	10	99	167	166	98	26	-
2002.....	541	-	-	-	-	3	20	96	172	135	101	13	1
2001.....	531	-	-	-	-	-	14	105	156	144	94	18	-
2000.....	574	-	-	-	-	-	14	99	186	164	94	17	-
1999.....	532	-	-	-	-	2	25	97	168	147	72	21	-
Non-Hispanic black, female													
2003.....	121	-	-	-	-	-	6	21	39	27	16	12	-
2002.....	133	-	-	-	-	1	7	27	26	39	30	3	-
2001.....	137	-	-	-	-	-	9	19	39	44	17	9	-
2000.....	140	-	-	-	-	-	1	22	32	45	35	5	-
1999.....	125	-	-	-	-	-	7	20	35	41	17	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of larynx (C32)													
Origin not stated 2/, both sexes													
2003.....	20	-	-	-	-	-	1	3	4	6	6	-	-
2002.....	10	-	-	-	-	-	-	1	4	3	2	-	-
2001.....	20	-	-	-	-	-	1	1	10	4	4	-	-
2000.....	19	-	-	-	-	-	1	3	4	8	2	1	-
1999.....	17	-	-	-	-	-	-	3	4	4	5	1	-
Origin not stated 2/, male													
2003.....	19	-	-	-	-	-	1	3	4	5	6	-	-
2002.....	8	-	-	-	-	-	-	1	3	2	2	-	-
2001.....	19	-	-	-	-	-	1	1	9	4	4	-	-
2000.....	16	-	-	-	-	-	1	2	4	7	2	-	-
1999.....	11	-	-	-	-	-	-	2	2	3	4	-	-
Origin not stated 2/, female													
2003.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2002.....	2	-	-	-	-	-	-	-	1	1	-	-	-
2001.....	1	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	3	-	-	-	-	-	-	1	-	1	-	1	-
1999.....	6	-	-	-	-	-	-	1	2	1	1	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
All origins, both sexes													
2003.....	158,086	-	2	2	25	154	2,478	12,374	30,956	49,386	48,619	14,088	2
2002.....	157,713	-	1	4	21	165	2,698	12,150	30,684	50,256	48,082	13,652	-
2001.....	156,058	-	5	2	20	157	2,796	12,041	29,777	51,225	46,700	13,331	4
2000.....	155,521	2	2	1	22	197	2,756	11,915	29,721	52,258	45,836	12,806	5
1999.....	152,156	-	-	5	20	179	2,724	11,434	29,344	51,848	44,261	12,339	2
All origins, male													
2003.....	89,964	-	1	1	14	78	1,344	7,313	18,347	28,935	27,064	6,865	2
2002.....	90,171	-	1	3	16	90	1,386	7,194	18,415	29,543	26,821	6,702	-
2001.....	90,426	-	4	1	12	77	1,504	7,190	17,873	30,699	26,406	6,656	4
2000.....	90,469	2	1	-	14	102	1,542	7,123	17,937	31,382	25,966	6,395	5
1999.....	89,453	-	-	4	13	97	1,474	6,851	18,013	31,287	25,397	6,315	2
All origins, female													
2003.....	68,122	-	1	1	11	76	1,134	5,061	12,609	20,451	21,555	7,223	-
2002.....	67,542	-	-	1	5	75	1,312	4,956	12,269	20,713	21,261	6,950	-
2001.....	65,632	-	1	1	8	80	1,292	4,851	11,904	20,526	20,294	6,675	-
2000.....	65,052	-	1	1	8	95	1,214	4,792	11,784	20,876	19,870	6,411	-
1999.....	62,703	-	-	1	7	82	1,250	4,583	11,331	20,561	18,864	6,024	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Hispanic, both sexes													
2003.....	4,136	-	-	-	4	20	92	390	795	1,293	1,177	365	-
2002.....	4,038	-	-	2	4	27	90	362	808	1,303	1,101	341	-
2001.....	3,816	-	2	1	7	16	106	364	758	1,230	1,011	321	-
2000.....	3,643	-	-	-	3	17	90	342	672	1,191	1,006	322	-
1999.....	3,545	-	-	-	3	17	98	308	724	1,170	901	324	-
Hispanic, male													
2003.....	2,640	-	-	-	3	7	52	240	549	877	723	189	-
2002.....	2,617	-	-	1	2	14	45	229	567	846	729	184	-
2001.....	2,495	-	2	-	6	10	68	239	517	820	657	176	-
2000.....	2,411	-	-	-	3	10	56	229	471	793	666	183	-
1999.....	2,373	-	-	-	2	10	58	208	523	798	606	168	-
Hispanic, female													
2003.....	1,496	-	-	-	1	13	40	150	246	416	454	176	-
2002.....	1,421	-	-	1	2	13	45	133	241	457	372	157	-
2001.....	1,321	-	-	1	1	6	38	125	241	410	354	145	-
2000.....	1,232	-	-	-	-	7	34	113	201	398	340	139	-
1999.....	1,172	-	-	-	1	7	40	100	201	372	295	156	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Non-Hispanic 1/, both sexes													
2003.....	153,616	-	2	2	21	132	2,378	11,951	30,101	47,977	47,352	13,698	2
2002.....	153,292	-	1	2	17	135	2,601	11,752	29,802	48,836	46,862	13,284	-
2001.....	151,882	-	3	1	13	141	2,678	11,642	28,936	49,887	45,599	12,978	4
2000.....	151,483	2	2	1	19	180	2,653	11,525	28,965	50,940	44,742	12,449	5
1999.....	148,206	-	-	5	17	161	2,618	11,087	28,549	50,536	43,243	11,988	2
Non-Hispanic 1/, male													
2003.....	87,111	-	1	1	11	69	1,287	7,052	17,762	27,973	26,286	6,667	2
2002.....	87,335	-	1	2	14	76	1,337	6,942	17,806	28,628	26,026	6,503	-
2001.....	87,706	-	2	1	6	67	1,431	6,929	17,291	29,818	25,695	6,462	4
2000.....	87,816	2	1	-	11	92	1,479	6,866	17,409	30,511	25,245	6,195	5
1999.....	86,822	-	-	4	11	87	1,411	6,616	17,438	30,397	24,724	6,132	2
Non-Hispanic 1/, female													
2003.....	66,505	-	1	1	10	63	1,091	4,899	12,339	20,004	21,066	7,031	-
2002.....	65,957	-	-	-	3	59	1,264	4,810	11,996	20,208	20,836	6,781	-
2001.....	64,176	-	1	-	7	74	1,247	4,713	11,645	20,069	19,904	6,516	-
2000.....	63,667	-	1	1	8	88	1,174	4,659	11,556	20,429	19,497	6,254	-
1999.....	61,384	-	-	1	6	74	1,207	4,471	11,111	20,139	18,519	5,856	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Non-Hispanic white, both sexes													
2003.....	134,437	-	2	1	18	107	1,852	9,391	25,613	42,204	42,752	12,495	2
2002.....	134,435	-	1	1	13	97	2,026	9,335	25,218	43,104	42,506	12,134	-
2001.....	133,266	-	2	1	9	114	2,100	9,301	24,713	43,972	41,223	11,829	2
2000.....	133,061	1	2	1	11	139	1,996	9,176	24,777	44,952	40,666	11,337	3
1999.....	129,967	-	-	3	9	131	2,025	8,832	24,329	44,577	39,153	10,907	1
Non-Hispanic white, male													
2003.....	75,468	-	1	-	9	55	986	5,480	14,861	24,440	23,573	6,061	2
2002.....	75,754	-	1	1	11	58	1,016	5,415	14,818	25,102	23,408	5,924	-
2001.....	76,066	-	1	1	4	55	1,095	5,421	14,552	26,056	23,022	5,857	2
2000.....	76,218	1	1	-	7	73	1,076	5,355	14,682	26,704	22,718	5,598	3
1999.....	75,338	-	-	2	6	71	1,078	5,155	14,654	26,596	22,219	5,556	1
Non-Hispanic white, female													
2003.....	58,969	-	1	1	9	52	866	3,911	10,752	17,764	19,179	6,434	-
2002.....	58,681	-	-	-	2	39	1,010	3,920	10,400	18,002	19,098	6,210	-
2001.....	57,200	-	1	-	5	59	1,005	3,880	10,161	17,916	18,201	5,972	-
2000.....	56,843	-	1	1	4	66	920	3,821	10,095	18,248	17,948	5,739	-
1999.....	54,629	-	-	1	3	60	947	3,677	9,675	17,981	16,934	5,351	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Non-Hispanic black, both sexes													
2003.....	16,252	-	-	1	3	18	441	2,250	3,955	4,892	3,739	953	-
2002.....	16,170	-	-	1	4	23	503	2,165	3,994	4,930	3,641	909	-
2001.....	15,883	-	1	-	3	16	515	2,060	3,685	5,076	3,623	902	2
2000.....	15,924	1	-	-	5	31	578	2,113	3,688	5,178	3,414	915	1
1999.....	15,834	-	-	2	8	20	531	2,029	3,743	5,111	3,483	907	-
Non-Hispanic black, male													
2003.....	9,914	-	-	1	2	12	251	1,399	2,570	3,009	2,192	478	-
2002.....	9,983	-	-	1	3	14	287	1,378	2,606	3,083	2,163	448	-
2001.....	10,030	-	1	-	1	7	298	1,352	2,400	3,266	2,226	477	2
2000.....	10,056	1	-	-	2	14	359	1,372	2,410	3,309	2,105	483	1
1999.....	10,039	-	-	2	5	11	301	1,338	2,500	3,267	2,127	488	-
Non-Hispanic black, female													
2003.....	6,338	-	-	-	1	6	190	851	1,385	1,883	1,547	475	-
2002.....	6,187	-	-	-	1	9	216	787	1,388	1,847	1,478	461	-
2001.....	5,853	-	-	-	2	9	217	708	1,285	1,810	1,397	425	-
2000.....	5,868	-	-	-	3	17	219	741	1,278	1,869	1,309	432	-
1999.....	5,795	-	-	-	3	9	230	691	1,243	1,844	1,356	419	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Origin not stated 2/, both sexes													
2003.....	334	-	-	-	-	2	8	33	60	116	90	25	-
2002.....	383	-	-	-	-	3	7	36	74	117	119	27	-
2001.....	360	-	-	-	-	-	12	35	83	108	90	32	-
2000.....	395	-	-	-	-	-	13	48	84	127	88	35	-
1999.....	405	-	-	-	-	1	8	39	71	142	117	27	-
Origin not stated 2/, male													
2003.....	213	-	-	-	-	2	5	21	36	85	55	9	-
2002.....	219	-	-	-	-	-	4	23	42	69	66	15	-
2001.....	225	-	-	-	-	-	5	22	65	61	54	18	-
2000.....	242	-	-	-	-	-	7	28	57	78	55	17	-
1999.....	258	-	-	-	-	-	5	27	52	92	67	15	-
Origin not stated 2/, female													
2003.....	121	-	-	-	-	-	3	12	24	31	35	16	-
2002.....	164	-	-	-	-	3	3	13	32	48	53	12	-
2001.....	135	-	-	-	-	-	7	13	18	47	36	14	-
2000.....	153	-	-	-	-	-	6	20	27	49	33	18	-
1999.....	147	-	-	-	-	1	3	12	19	50	50	12	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
All origins, both sexes													
2003.....	7,818	-	1	2	38	191	561	1,179	1,547	1,593	1,875	831	-
2002.....	7,514	-	3	1	41	194	604	1,135	1,351	1,616	1,748	820	1
2001.....	7,542	-	-	-	41	186	622	1,208	1,333	1,689	1,678	785	-
2000.....	7,420	-	-	1	41	186	626	1,199	1,303	1,637	1,624	803	-
1999.....	7,215	1	-	1	39	194	625	1,120	1,263	1,652	1,639	681	-
All origins, male													
2003.....	4,963	-	-	-	24	119	336	753	1,042	1,050	1,203	436	-
2002.....	4,725	-	2	1	20	119	363	746	915	1,075	1,078	406	-
2001.....	4,763	-	-	-	24	103	380	756	893	1,156	1,063	388	-
2000.....	4,592	-	-	1	21	110	355	769	866	1,067	1,013	390	-
1999.....	4,529	1	-	-	23	110	397	717	851	1,096	1,006	328	-
All origins, female													
2003.....	2,855	-	1	2	14	72	225	426	505	543	672	395	-
2002.....	2,789	-	1	-	21	75	241	389	436	541	670	414	1
2001.....	2,779	-	-	-	17	83	242	452	440	533	615	397	-
2000.....	2,828	-	-	-	20	76	271	430	437	570	611	413	-
1999.....	2,686	-	-	1	16	84	228	403	412	556	633	353	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
Hispanic, both sexes													
2003.....	159	-	-	1	2	6	15	33	29	37	25	11	-
2002.....	147	-	-	-	2	7	21	27	24	19	29	18	-
2001.....	145	-	-	-	2	6	24	23	25	28	28	9	-
2000.....	144	-	-	-	2	5	26	28	27	26	19	11	-
1999.....	133	1	-	1	2	5	14	22	30	27	20	11	-
Hispanic, male													
2003.....	91	-	-	-	1	4	8	19	16	21	15	7	-
2002.....	77	-	-	-	1	6	13	13	16	11	13	4	-
2001.....	78	-	-	-	-	2	17	11	15	19	13	1	-
2000.....	92	-	-	-	1	4	17	19	19	14	13	5	-
1999.....	79	1	-	-	1	4	9	16	19	17	9	3	-
Hispanic, female													
2003.....	68	-	-	1	1	2	7	14	13	16	10	4	-
2002.....	70	-	-	-	1	1	8	14	8	8	16	14	-
2001.....	67	-	-	-	2	4	7	12	10	9	15	8	-
2000.....	52	-	-	-	1	1	9	9	8	12	6	6	-
1999.....	54	-	-	1	1	1	5	6	11	10	11	8	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
Non-Hispanic 1/, both sexes													
2003.....	7,640	-	1	1	36	183	544	1,144	1,513	1,555	1,846	817	-
2002.....	7,349	-	3	1	38	186	581	1,106	1,323	1,596	1,714	801	-
2001.....	7,387	-	-	-	39	179	597	1,184	1,307	1,660	1,646	775	-
2000.....	7,263	-	-	1	39	181	599	1,167	1,276	1,608	1,602	790	-
1999.....	7,062	-	-	-	37	189	611	1,094	1,231	1,620	1,612	668	-
Non-Hispanic 1/, male													
2003.....	4,859	-	-	-	23	113	326	732	1,025	1,028	1,185	427	-
2002.....	4,640	-	2	1	19	113	350	732	896	1,063	1,062	402	-
2001.....	4,678	-	-	-	24	100	363	744	877	1,136	1,047	387	-
2000.....	4,490	-	-	1	20	106	337	746	847	1,051	998	384	-
1999.....	4,438	-	-	-	22	106	388	698	831	1,076	994	323	-
Non-Hispanic 1/, female													
2003.....	2,781	-	1	1	13	70	218	412	488	527	661	390	-
2002.....	2,709	-	1	-	19	73	231	374	427	533	652	399	-
2001.....	2,709	-	-	-	15	79	234	440	430	524	599	388	-
2000.....	2,773	-	-	-	19	75	262	421	429	557	604	406	-
1999.....	2,624	-	-	-	15	83	223	396	400	544	618	345	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
Non-Hispanic white, both sexes													
2003.....	7,471	-	1	1	34	178	533	1,116	1,479	1,526	1,805	798	-
2002.....	7,190	-	3	1	37	184	568	1,085	1,291	1,554	1,687	780	-
2001.....	7,250	-	-	-	38	177	590	1,168	1,286	1,616	1,614	761	-
2000.....	7,090	-	-	1	38	175	579	1,147	1,247	1,570	1,566	767	-
1999.....	6,936	-	-	-	34	183	604	1,081	1,213	1,593	1,577	651	-
Non-Hispanic white, male													
2003.....	4,784	-	-	-	23	111	323	718	1,007	1,016	1,165	421	-
2002.....	4,555	-	2	1	18	111	344	719	876	1,039	1,050	395	-
2001.....	4,618	-	-	-	23	98	359	737	867	1,116	1,034	384	-
2000.....	4,414	-	-	1	20	104	327	737	833	1,032	981	379	-
1999.....	4,370	-	-	-	19	102	385	689	819	1,064	976	316	-
Non-Hispanic white, female													
2003.....	2,687	-	1	1	11	67	210	398	472	510	640	377	-
2002.....	2,635	-	1	-	19	73	224	366	415	515	637	385	-
2001.....	2,632	-	-	-	15	79	231	431	419	500	580	377	-
2000.....	2,676	-	-	-	18	71	252	410	414	538	585	388	-
1999.....	2,566	-	-	-	15	81	219	392	394	529	601	335	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
Non-Hispanic black, both sexes													
2003.....	120	-	-	-	1	3	10	16	23	26	28	13	-
2002.....	102	-	-	-	1	1	9	12	21	25	18	15	-
2001.....	102	-	-	-	1	1	7	9	13	36	24	11	-
2000.....	134	-	-	-	1	3	15	13	19	30	32	21	-
1999.....	94	-	-	-	1	4	4	9	14	23	25	14	-
Non-Hispanic black, male													
2003.....	49	-	-	-	-	2	3	8	12	12	10	2	-
2002.....	52	-	-	-	1	1	4	7	13	12	8	6	-
2001.....	36	-	-	-	1	1	4	3	4	16	6	1	-
2000.....	59	-	-	-	-	2	9	6	10	14	15	3	-
1999.....	47	-	-	-	1	2	1	6	10	9	12	6	-
Non-Hispanic black, female													
2003.....	71	-	-	-	1	1	7	8	11	14	18	11	-
2002.....	50	-	-	-	-	-	5	5	8	13	10	9	-
2001.....	66	-	-	-	-	-	3	6	9	20	18	10	-
2000.....	75	-	-	-	1	1	6	7	9	16	17	18	-
1999.....	47	-	-	-	-	2	3	3	4	14	13	8	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant melanoma of skin (C43)													
Origin not stated 2/, both sexes													
2003.....	19	-	-	-	-	2	2	2	5	1	4	3	-
2002.....	18	-	-	-	1	1	2	2	4	1	5	1	1
2001.....	10	-	-	-	-	1	1	1	1	1	4	1	-
2000.....	13	-	-	-	-	-	1	4	-	3	3	2	-
1999.....	20	-	-	-	-	-	-	4	2	5	7	2	-
Origin not stated 2/, male													
2003.....	13	-	-	-	-	2	2	2	1	1	3	2	-
2002.....	8	-	-	-	-	-	-	1	3	1	3	-	-
2001.....	7	-	-	-	-	1	-	1	1	1	3	-	-
2000.....	10	-	-	-	-	-	1	4	-	2	2	1	-
1999.....	12	-	-	-	-	-	-	3	1	3	3	2	-
Origin not stated 2/, female													
2003.....	6	-	-	-	-	-	-	-	4	-	1	1	-
2002.....	10	-	-	-	1	1	2	1	1	-	2	1	1
2001.....	3	-	-	-	-	-	1	-	-	-	1	1	-
2000.....	3	-	-	-	-	-	-	-	-	1	1	1	-
1999.....	8	-	-	-	-	-	-	1	1	2	4	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
All origins, both sexes													
2003.....	42,000	-	-	-	16	407	2,716	6,365	8,267	8,338	9,644	6,245	2
2002.....	41,883	-	-	-	10	425	2,722	6,454	7,838	8,505	9,748	6,181	-
2001.....	41,809	-	-	-	9	463	2,823	6,593	7,632	8,715	9,663	5,911	-
2000.....	42,300	-	-	-	11	467	2,841	6,385	7,573	9,024	9,750	6,249	-
1999.....	41,528	-	1	1	13	439	2,741	6,174	7,399	9,109	9,651	6,000	-
All origins, male													
2003.....	380	-	-	-	-	2	9	48	71	89	104	56	1
2002.....	369	-	-	-	-	3	12	44	67	84	112	47	-
2001.....	415	-	-	-	-	2	14	46	72	120	115	46	-
2000.....	428	-	-	-	-	3	19	46	85	115	109	51	-
1999.....	384	-	-	-	-	3	15	41	70	103	98	54	-
All origins, female													
2003.....	41,620	-	-	-	16	405	2,707	6,317	8,196	8,249	9,540	6,189	1
2002.....	41,514	-	-	-	10	422	2,710	6,410	7,771	8,421	9,636	6,134	-
2001.....	41,394	-	-	-	9	461	2,809	6,547	7,560	8,595	9,548	5,865	-
2000.....	41,872	-	-	-	11	464	2,822	6,339	7,488	8,909	9,641	6,198	-
1999.....	41,144	-	1	1	13	436	2,726	6,133	7,329	9,006	9,553	5,946	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
Hispanic, both sexes													
2003.....	1,893	-	-	-	2	55	256	402	428	331	269	150	-
2002.....	1,736	-	-	-	2	54	218	401	348	323	262	128	-
2001.....	1,760	-	-	-	1	53	244	408	407	318	219	110	-
2000.....	1,689	-	-	-	1	54	218	381	359	337	231	108	-
1999.....	1,551	-	-	-	2	47	197	362	321	290	209	123	-
Hispanic, male													
2003.....	9	-	-	-	-	-	-	2	2	2	-	3	-
2002.....	12	-	-	-	-	1	-	5	-	3	2	1	-
2001.....	14	-	-	-	-	-	2	1	1	5	5	-	-
2000.....	19	-	-	-	-	-	2	2	4	6	4	1	-
1999.....	13	-	-	-	-	-	-	3	3	5	2	-	-
Hispanic, female													
2003.....	1,884	-	-	-	2	55	256	400	426	329	269	147	-
2002.....	1,724	-	-	-	2	53	218	396	348	320	260	127	-
2001.....	1,746	-	-	-	1	53	242	407	406	313	214	110	-
2000.....	1,670	-	-	-	1	54	216	379	355	331	227	107	-
1999.....	1,538	-	-	-	2	47	197	359	318	285	207	123	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
Non-Hispanic 1/, both sexes													
2003.....	40,003	-	-	-	14	351	2,453	5,950	7,813	7,987	9,345	6,088	2
2002.....	40,037	-	-	-	8	368	2,495	6,033	7,470	8,161	9,464	6,038	-
2001.....	39,957	-	-	-	8	409	2,576	6,170	7,208	8,379	9,418	5,789	-
2000.....	40,506	-	-	-	10	413	2,615	5,989	7,192	8,670	9,494	6,123	-
1999.....	39,881	-	1	1	11	390	2,537	5,800	7,056	8,803	9,419	5,863	-
Non-Hispanic 1/, male													
2003.....	371	-	-	-	-	2	9	46	69	87	104	53	1
2002.....	355	-	-	-	-	2	12	38	66	81	110	46	-
2001.....	399	-	-	-	-	2	12	45	70	114	110	46	-
2000.....	409	-	-	-	-	3	17	44	81	109	105	50	-
1999.....	370	-	-	-	-	3	15	38	67	97	96	54	-
Non-Hispanic 1/, female													
2003.....	39,632	-	-	-	14	349	2,444	5,904	7,744	7,900	9,241	6,035	1
2002.....	39,682	-	-	-	8	366	2,483	5,995	7,404	8,080	9,354	5,992	-
2001.....	39,558	-	-	-	8	407	2,564	6,125	7,138	8,265	9,308	5,743	-
2000.....	40,097	-	-	-	10	410	2,598	5,945	7,111	8,561	9,389	6,073	-
1999.....	39,511	-	1	1	11	387	2,522	5,762	6,989	8,706	9,323	5,809	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
Non-Hispanic white, both sexes													
2003.....	33,451	-	-	-	9	216	1,728	4,475	6,354	6,824	8,356	5,488	1
2002.....	33,611	-	-	-	4	230	1,733	4,559	6,145	7,001	8,456	5,483	-
2001.....	33,629	-	-	-	3	255	1,828	4,737	5,895	7,185	8,472	5,254	-
2000.....	34,386	-	-	-	4	275	1,877	4,620	5,911	7,581	8,591	5,527	-
1999.....	33,790	-	-	1	6	257	1,802	4,470	5,779	7,623	8,488	5,364	-
Non-Hispanic white, male													
2003.....	317	-	-	-	-	1	4	37	57	74	95	48	1
2002.....	288	-	-	-	-	1	9	27	51	65	94	41	-
2001.....	329	-	-	-	-	1	6	33	59	95	96	39	-
2000.....	341	-	-	-	-	1	9	35	63	97	94	42	-
1999.....	299	-	-	-	-	2	8	28	51	79	84	47	-
Non-Hispanic white, female													
2003.....	33,134	-	-	-	9	215	1,724	4,438	6,297	6,750	8,261	5,440	-
2002.....	33,323	-	-	-	4	229	1,724	4,532	6,094	6,936	8,362	5,442	-
2001.....	33,300	-	-	-	3	254	1,822	4,704	5,836	7,090	8,376	5,215	-
2000.....	34,045	-	-	-	4	274	1,868	4,585	5,848	7,484	8,497	5,485	-
1999.....	33,491	-	-	1	6	255	1,794	4,442	5,728	7,544	8,404	5,317	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
Non-Hispanic black, both sexes													
2003.....	5,677	-	-	-	5	117	632	1,258	1,238	1,001	877	549	-
2002.....	5,579	-	-	-	3	123	670	1,247	1,122	1,005	900	509	-
2001.....	5,528	-	-	-	5	140	641	1,209	1,125	1,053	862	493	-
2000.....	5,384	-	-	-	5	120	641	1,167	1,103	961	825	562	-
1999.....	5,386	-	1	-	5	124	661	1,150	1,095	1,037	852	461	-
Non-Hispanic black, male													
2003.....	51	-	-	-	-	1	5	8	12	13	7	5	-
2002.....	62	-	-	-	-	1	3	10	14	16	13	5	-
2001.....	64	-	-	-	-	1	5	12	10	17	13	6	-
2000.....	63	-	-	-	-	2	8	8	16	11	10	8	-
1999.....	67	-	-	-	-	1	7	9	16	16	11	7	-
Non-Hispanic black, female													
2003.....	5,626	-	-	-	5	116	627	1,250	1,226	988	870	544	-
2002.....	5,517	-	-	-	3	122	667	1,237	1,108	989	887	504	-
2001.....	5,464	-	-	-	5	139	636	1,197	1,115	1,036	849	487	-
2000.....	5,321	-	-	-	5	118	633	1,159	1,087	950	815	554	-
1999.....	5,319	-	1	-	5	123	654	1,141	1,079	1,021	841	454	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of breast (C50)													
Origin not stated 2/, both sexes													
2003.....	104	-	-	-	-	1	7	13	26	20	30	7	-
2002.....	110	-	-	-	-	3	9	20	20	21	22	15	-
2001.....	92	-	-	-	-	1	3	15	17	18	26	12	-
2000.....	105	-	-	-	-	-	8	15	22	17	25	18	-
1999.....	96	-	-	-	-	2	7	12	22	16	23	14	-
Origin not stated 2/, male													
2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	1	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-	-
Origin not stated 2/, female													
2003.....	104	-	-	-	-	1	7	13	26	20	30	7	-
2002.....	108	-	-	-	-	3	9	19	19	21	22	15	-
2001.....	90	-	-	-	-	1	3	15	16	17	26	12	-
2000.....	105	-	-	-	-	-	8	15	22	17	25	18	-
1999.....	95	-	-	-	-	2	7	12	22	15	23	14	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)													
All origins, both sexes													
2003.....	3,919	-	-	-	16	201	633	890	793	564	552	270	-
2002.....	3,952	-	-	-	17	208	666	894	725	630	528	284	-
2001.....	4,092	-	-	1	12	209	711	953	723	585	583	315	-
2000.....	4,200	-	-	1	17	230	694	915	748	686	599	310	-
1999.....	4,205	-	-	-	12	217	699	914	769	696	568	329	1
All origins, male													
2003.....
2002.....
2001.....
2000.....
1999.....
All origins, female													
2003.....	3,919	-	-	-	16	201	633	890	793	564	552	270	-
2002.....	3,952	-	-	-	17	208	666	894	725	630	528	284	-
2001.....	4,092	-	-	1	12	209	711	953	723	585	583	315	-
2000.....	4,200	-	-	1	17	230	694	915	748	686	599	310	-
1999.....	4,205	-	-	-	12	217	699	914	769	696	568	329	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)													
Hispanic, both sexes													
2003.....	391	-	-	-	3	31	85	109	65	43	42	13	-
2002.....	388	-	-	-	2	30	86	87	71	55	40	17	-
2001.....	412	-	-	-	3	36	102	102	75	39	39	16	-
2000.....	434	-	-	-	4	33	108	87	78	60	43	21	-
1999.....	378	-	-	-	2	31	87	99	64	51	31	13	-
Hispanic, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Hispanic, female													
2003.....	391	-	-	-	3	31	85	109	65	43	42	13	-
2002.....	388	-	-	-	2	30	86	87	71	55	40	17	-
2001.....	412	-	-	-	3	36	102	102	75	39	39	16	-
2000.....	434	-	-	-	4	33	108	87	78	60	43	21	-
1999.....	378	-	-	-	2	31	87	99	64	51	31	13	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)													
Non-Hispanic 1/, both sexes													
2003.....	3,519	-	-	-	13	170	546	777	727	519	510	257	-
2002.....	3,551	-	-	-	15	177	577	805	650	573	487	267	-
2001.....	3,669	-	-	1	9	172	606	846	646	546	544	299	-
2000.....	3,755	-	-	1	13	197	583	825	668	625	554	289	-
1999.....	3,817	-	-	-	10	186	611	813	702	644	535	315	1
Non-Hispanic 1/, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic 1/, female													
2003.....	3,519	-	-	-	13	170	546	777	727	519	510	257	-
2002.....	3,551	-	-	-	15	177	577	805	650	573	487	267	-
2001.....	3,669	-	-	1	9	172	606	846	646	546	544	299	-
2000.....	3,755	-	-	1	13	197	583	825	668	625	554	289	-
1999.....	3,817	-	-	-	10	186	611	813	702	644	535	315	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)													
Non-Hispanic white, both sexes													
2003.....	2,564	-	-	-	10	110	392	565	549	371	370	197	-
2002.....	2,580	-	-	-	10	129	417	581	480	399	370	194	-
2001.....	2,725	-	-	-	8	125	448	632	478	392	409	233	-
2000.....	2,725	-	-	-	9	141	403	600	479	438	434	221	-
1999.....	2,854	-	-	-	7	147	451	630	508	454	414	242	1
Non-Hispanic white, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic white, female													
2003.....	2,564	-	-	-	10	110	392	565	549	371	370	197	-
2002.....	2,580	-	-	-	10	129	417	581	480	399	370	194	-
2001.....	2,725	-	-	-	8	125	448	632	478	392	409	233	-
2000.....	2,725	-	-	-	9	141	403	600	479	438	434	221	-
1999.....	2,854	-	-	-	7	147	451	630	508	454	414	242	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic black, both sexes

2003.....	788	-	-	-	2	49	126	173	152	125	111	50	-
2002.....	820	-	-	-	3	43	144	188	131	152	101	58	-
2001.....	778	-	-	1	1	42	137	165	131	129	111	61	-
2000.....	860	-	-	1	4	51	147	177	152	158	108	62	-
1999.....	825	-	-	-	3	34	140	140	167	173	101	67	-

Non-Hispanic black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic black, female

2003.....	788	-	-	-	2	49	126	173	152	125	111	50	-
2002.....	820	-	-	-	3	43	144	188	131	152	101	58	-
2001.....	778	-	-	1	1	42	137	165	131	129	111	61	-
2000.....	860	-	-	1	4	51	147	177	152	158	108	62	-
1999.....	825	-	-	-	3	34	140	140	167	173	101	67	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasm of cervix uteri (C53)

Origin not stated 2/, both sexes

2003.....	9	-	-	-	-	-	2	4	1	2	-	-	-
2002.....	13	-	-	-	-	1	3	2	4	2	1	-	-
2001.....	11	-	-	-	-	1	3	5	2	-	-	-	-
2000.....	11	-	-	-	-	-	3	3	2	1	2	-	-
1999.....	10	-	-	-	-	-	1	2	3	1	2	1	-

Origin not stated 2/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Origin not stated 2/, female

2003.....	9	-	-	-	-	-	2	4	1	2	-	-	-
2002.....	13	-	-	-	-	1	3	2	4	2	1	-	-
2001.....	11	-	-	-	-	1	3	5	2	-	-	-	-
2000.....	11	-	-	-	-	-	3	3	2	1	2	-	-
1999.....	10	-	-	-	-	-	1	2	3	1	2	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All origins, both sexes

2003.....	6,899	-	-	-	4	27	154	534	1,301	1,851	1,951	1,077	-
2002.....	6,853	-	-	-	-	23	146	565	1,199	1,835	1,957	1,128	-
2001.....	6,784	-	-	-	-	38	139	566	1,105	1,784	2,063	1,088	1
2000.....	6,586	-	-	-	3	27	138	507	1,122	1,781	1,958	1,049	1
1999.....	6,468	-	-	-	2	23	142	468	1,019	1,807	2,008	999	-

All origins, male

2003.....
2002.....
2001.....
2000.....
1999.....

All origins, female

2003.....	6,899	-	-	-	4	27	154	534	1,301	1,851	1,951	1,077	-
2002.....	6,853	-	-	-	-	23	146	565	1,199	1,835	1,957	1,128	-
2001.....	6,784	-	-	-	-	38	139	566	1,105	1,784	2,063	1,088	1
2000.....	6,586	-	-	-	3	27	138	507	1,122	1,781	1,958	1,049	1
1999.....	6,468	-	-	-	2	23	142	468	1,019	1,807	2,008	999	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Hispanic, both sexes

2003.....	321	-	-	-	1	3	24	44	61	85	63	40	-
2002.....	337	-	-	-	-	5	20	39	64	106	66	37	-
2001.....	319	-	-	-	-	7	13	48	71	85	65	30	-
2000.....	277	-	-	-	-	2	19	35	62	75	54	30	-
1999.....	272	-	-	-	1	3	15	33	61	86	51	22	-

Hispanic, male

2003.....
2002.....
2001.....
2000.....
1999.....

Hispanic, female

2003.....	321	-	-	-	1	3	24	44	61	85	63	40	-
2002.....	337	-	-	-	-	5	20	39	64	106	66	37	-
2001.....	319	-	-	-	-	7	13	48	71	85	65	30	-
2000.....	277	-	-	-	-	2	19	35	62	75	54	30	-
1999.....	272	-	-	-	1	3	15	33	61	86	51	22	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic 1/, both sexes

2003.....	6,561	-	-	-	3	24	130	488	1,237	1,761	1,881	1,037	-
2002.....	6,487	-	-	-	-	17	126	524	1,124	1,725	1,884	1,087	-
2001.....	6,449	-	-	-	-	31	126	517	1,029	1,697	1,996	1,052	1
2000.....	6,294	-	-	-	3	25	119	469	1,057	1,702	1,901	1,017	1
1999.....	6,188	-	-	-	1	20	127	435	956	1,719	1,954	976	-

Non-Hispanic 1/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 1/, female

2003.....	6,561	-	-	-	3	24	130	488	1,237	1,761	1,881	1,037	-
2002.....	6,487	-	-	-	-	17	126	524	1,124	1,725	1,884	1,087	-
2001.....	6,449	-	-	-	-	31	126	517	1,029	1,697	1,996	1,052	1
2000.....	6,294	-	-	-	3	25	119	469	1,057	1,702	1,901	1,017	1
1999.....	6,188	-	-	-	1	20	127	435	956	1,719	1,954	976	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)													
Non-Hispanic white, both sexes													
2003.....	5,321	-	-	-	2	13	86	386	955	1,366	1,584	929	-
2002.....	5,213	-	-	-	-	11	97	403	861	1,312	1,580	949	-
2001.....	5,283	-	-	-	-	19	94	398	802	1,319	1,712	939	-
2000.....	5,113	-	-	-	1	18	88	383	827	1,298	1,594	904	-
1999.....	5,114	-	-	-	1	15	96	353	753	1,351	1,687	858	-
Non-Hispanic white, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic white, female													
2003.....	5,321	-	-	-	2	13	86	386	955	1,366	1,584	929	-
2002.....	5,213	-	-	-	-	11	97	403	861	1,312	1,580	949	-
2001.....	5,283	-	-	-	-	19	94	398	802	1,319	1,712	939	-
2000.....	5,113	-	-	-	1	18	88	383	827	1,298	1,594	904	-
1999.....	5,114	-	-	-	1	15	96	353	753	1,351	1,687	858	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic black, both sexes

2003.....	1,092	-	-	-	1	8	30	80	245	357	269	102	-
2002.....	1,107	-	-	-	-	5	21	88	221	373	268	131	-
2001.....	1,037	-	-	-	-	9	22	93	190	352	265	105	1
2000.....	1,055	-	-	-	1	6	21	68	201	371	280	106	1
1999.....	980	-	-	-	-	5	27	57	189	346	245	111	-

Non-Hispanic black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic black, female

2003.....	1,092	-	-	-	1	8	30	80	245	357	269	102	-
2002.....	1,107	-	-	-	-	5	21	88	221	373	268	131	-
2001.....	1,037	-	-	-	-	9	22	93	190	352	265	105	1
2000.....	1,055	-	-	-	1	6	21	68	201	371	280	106	1
1999.....	980	-	-	-	-	5	27	57	189	346	245	111	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Origin not stated 2/, both sexes

2003.....	17	-	-	-	-	-	-	2	3	5	7	-	-
2002.....	29	-	-	-	-	1	-	2	11	4	7	4	-
2001.....	16	-	-	-	-	-	-	1	5	2	2	6	-
2000.....	15	-	-	-	-	-	-	3	3	4	3	2	-
1999.....	8	-	-	-	-	-	-	-	2	2	3	1	-

Origin not stated 2/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Origin not stated 2/, female

2003.....	17	-	-	-	-	-	-	2	3	5	7	-	-
2002.....	29	-	-	-	-	1	-	2	11	4	7	4	-
2001.....	16	-	-	-	-	-	-	1	5	2	2	6	-
2000.....	15	-	-	-	-	-	-	3	3	4	3	2	-
1999.....	8	-	-	-	-	-	-	-	2	2	3	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
All origins, both sexes													
2003.....	14,657	-	-	-	28	84	452	1,629	2,891	3,569	4,194	1,810	-
2002.....	14,682	1	1	2	37	97	469	1,717	2,748	3,601	4,259	1,750	-
2001.....	14,414	-	-	2	21	102	472	1,626	2,612	3,666	4,170	1,743	-
2000.....	14,060	-	-	2	23	101	472	1,583	2,444	3,601	4,098	1,736	-
1999.....	13,627	-	1	2	27	77	496	1,549	2,426	3,612	3,875	1,562	-
All origins, male													
2003.....
2002.....
2001.....
2000.....
1999.....
All origins, female													
2003.....	14,657	-	-	-	28	84	452	1,629	2,891	3,569	4,194	1,810	-
2002.....	14,682	1	1	2	37	97	469	1,717	2,748	3,601	4,259	1,750	-
2001.....	14,414	-	-	2	21	102	472	1,626	2,612	3,666	4,170	1,743	-
2000.....	14,060	-	-	2	23	101	472	1,583	2,444	3,601	4,098	1,736	-
1999.....	13,627	-	1	2	27	77	496	1,549	2,426	3,612	3,875	1,562	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
Hispanic, both sexes													
2003.....	645	-	-	-	3	5	48	104	134	155	141	55	-
2002.....	622	-	-	2	3	10	47	116	111	145	145	43	-
2001.....	603	-	-	1	-	13	43	91	131	164	111	49	-
2000.....	578	-	-	1	4	9	30	94	111	154	126	49	-
1999.....	478	-	1	-	2	6	29	80	100	126	100	34	-
Hispanic, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Hispanic, female													
2003.....	645	-	-	-	3	5	48	104	134	155	141	55	-
2002.....	622	-	-	2	3	10	47	116	111	145	145	43	-
2001.....	603	-	-	1	-	13	43	91	131	164	111	49	-
2000.....	578	-	-	1	4	9	30	94	111	154	126	49	-
1999.....	478	-	1	-	2	6	29	80	100	126	100	34	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
Non-Hispanic 1/, both sexes													
2003.....	13,985	-	-	-	25	79	403	1,519	2,749	3,406	4,050	1,754	-
2002.....	14,031	1	1	-	34	87	421	1,596	2,633	3,445	4,107	1,706	-
2001.....	13,783	-	-	1	21	89	428	1,533	2,475	3,495	4,052	1,689	-
2000.....	13,454	-	-	1	19	92	441	1,484	2,329	3,440	3,964	1,684	-
1999.....	13,103	-	-	2	25	71	466	1,466	2,316	3,474	3,756	1,527	-
Non-Hispanic 1/, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic 1/, female													
2003.....	13,985	-	-	-	25	79	403	1,519	2,749	3,406	4,050	1,754	-
2002.....	14,031	1	1	-	34	87	421	1,596	2,633	3,445	4,107	1,706	-
2001.....	13,783	-	-	1	21	89	428	1,533	2,475	3,495	4,052	1,689	-
2000.....	13,454	-	-	1	19	92	441	1,484	2,329	3,440	3,964	1,684	-
1999.....	13,103	-	-	2	25	71	466	1,466	2,316	3,474	3,756	1,527	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
Non-Hispanic white, both sexes													
2003.....	12,477	-	-	-	18	64	329	1,282	2,432	3,024	3,720	1,608	-
2002.....	12,560	1	1	-	27	67	343	1,369	2,329	3,082	3,766	1,575	-
2001.....	12,338	-	-	-	14	68	347	1,335	2,151	3,125	3,741	1,557	-
2000.....	12,138	-	-	-	13	75	367	1,284	2,072	3,087	3,680	1,560	-
1999.....	11,795	-	-	2	20	51	394	1,268	2,061	3,119	3,456	1,424	-
Non-Hispanic white, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic white, female													
2003.....	12,477	-	-	-	18	64	329	1,282	2,432	3,024	3,720	1,608	-
2002.....	12,560	1	1	-	27	67	343	1,369	2,329	3,082	3,766	1,575	-
2001.....	12,338	-	-	-	14	68	347	1,335	2,151	3,125	3,741	1,557	-
2000.....	12,138	-	-	-	13	75	367	1,284	2,072	3,087	3,680	1,560	-
1999.....	11,795	-	-	2	20	51	394	1,268	2,061	3,119	3,456	1,424	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
Non-Hispanic black, both sexes													
2003.....	1,180	-	-	-	6	12	56	181	236	297	267	125	-
2002.....	1,155	-	-	-	7	13	59	158	237	287	277	117	-
2001.....	1,140	-	-	1	4	15	65	134	251	301	252	117	-
2000.....	1,065	-	-	1	2	12	59	138	205	295	244	109	-
1999.....	1,065	-	-	-	4	13	47	143	211	301	256	90	-
Non-Hispanic black, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic black, female													
2003.....	1,180	-	-	-	6	12	56	181	236	297	267	125	-
2002.....	1,155	-	-	-	7	13	59	158	237	287	277	117	-
2001.....	1,140	-	-	1	4	15	65	134	251	301	252	117	-
2000.....	1,065	-	-	1	2	12	59	138	205	295	244	109	-
1999.....	1,065	-	-	-	4	13	47	143	211	301	256	90	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of ovary (C56)													
Origin not stated 2/, both sexes													
2003.....	27	-	-	-	-	-	1	6	8	8	3	1	-
2002.....	29	-	-	-	-	-	1	5	4	11	7	1	-
2001.....	28	-	-	-	-	-	1	2	6	7	7	5	-
2000.....	28	-	-	-	-	-	1	5	4	7	8	3	-
1999.....	46	-	-	-	-	-	1	3	10	12	19	1	-
Origin not stated 2/, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Origin not stated 2/, female													
2003.....	27	-	-	-	-	-	1	6	8	8	3	1	-
2002.....	29	-	-	-	-	-	1	5	4	11	7	1	-
2001.....	28	-	-	-	-	-	1	2	6	7	7	5	-
2000.....	28	-	-	-	-	-	1	5	4	7	8	3	-
1999.....	46	-	-	-	-	-	1	3	10	12	19	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
All origins, both sexes													
2003.....	12,286	1	14	39	24	56	321	1,201	2,498	3,090	3,475	1,567	-
2002.....	12,165	4	8	30	20	48	306	1,322	2,340	3,239	3,355	1,493	-
2001.....	12,078	1	22	35	20	52	358	1,311	2,328	3,221	3,306	1,424	-
2000.....	11,736	2	15	29	20	42	334	1,239	2,195	3,146	3,282	1,432	-
1999.....	11,116	2	19	33	17	54	342	1,147	2,111	3,076	3,010	1,305	-
All origins, male													
2003.....	7,667	1	4	19	16	36	231	859	1,787	1,975	2,036	703	-
2002.....	7,587	1	-	19	7	31	217	947	1,658	2,087	1,924	696	-
2001.....	7,524	1	10	9	16	27	240	941	1,634	2,093	1,866	687	-
2000.....	7,249	1	8	15	10	26	228	880	1,530	2,010	1,865	676	-
1999.....	6,867	1	11	16	11	37	236	816	1,474	1,958	1,717	590	-
All origins, female													
2003.....	4,619	-	10	20	8	20	90	342	711	1,115	1,439	864	-
2002.....	4,578	3	8	11	13	17	89	375	682	1,152	1,431	797	-
2001.....	4,554	-	12	26	4	25	118	370	694	1,128	1,440	737	-
2000.....	4,487	1	7	14	10	16	106	359	665	1,136	1,417	756	-
1999.....	4,249	1	8	17	6	17	106	331	637	1,118	1,293	715	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Hispanic, both sexes													
2003.....	698	-	2	4	6	3	30	82	170	198	151	52	-
2002.....	659	2	3	2	1	6	37	103	160	152	146	47	-
2001.....	640	-	6	6	-	8	38	112	127	163	129	51	-
2000.....	567	-	3	3	3	7	32	88	135	132	125	39	-
1999.....	547	-	1	4	1	4	31	84	129	144	108	41	-
Hispanic, male													
2003.....	426	-	-	2	5	3	23	54	109	121	84	25	-
2002.....	426	-	-	2	-	4	26	72	106	97	95	24	-
2001.....	426	-	1	4	-	5	34	80	86	107	79	30	-
2000.....	372	-	1	3	2	4	27	60	92	85	78	20	-
1999.....	353	-	1	2	-	2	23	63	91	96	57	18	-
Hispanic, female													
2003.....	272	-	2	2	1	-	7	28	61	77	67	27	-
2002.....	233	2	3	-	1	2	11	31	54	55	51	23	-
2001.....	214	-	5	2	-	3	4	32	41	56	50	21	-
2000.....	195	-	2	-	1	3	5	28	43	47	47	19	-
1999.....	194	-	-	2	1	2	8	21	38	48	51	23	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Non-Hispanic 1/, both sexes													
2003.....	11,566	1	12	35	17	53	291	1,117	2,325	2,884	3,320	1,511	-
2002.....	11,477	2	5	28	19	42	268	1,216	2,173	3,079	3,200	1,445	-
2001.....	11,415	1	16	29	20	44	319	1,196	2,198	3,051	3,172	1,369	-
2000.....	11,124	2	12	26	17	35	302	1,147	2,053	2,994	3,150	1,386	-
1999.....	10,542	2	18	29	16	49	309	1,060	1,979	2,927	2,892	1,261	-
Non-Hispanic 1/, male													
2003.....	7,224	1	4	17	10	33	208	804	1,675	1,848	1,948	676	-
2002.....	7,141	1	-	17	7	27	190	873	1,546	1,984	1,825	671	-
2001.....	7,084	1	9	5	16	22	205	859	1,546	1,981	1,784	656	-
2000.....	6,847	1	7	12	8	22	201	818	1,433	1,911	1,782	652	-
1999.....	6,501	1	10	14	11	35	212	751	1,381	1,858	1,657	571	-
Non-Hispanic 1/, female													
2003.....	4,342	-	8	18	7	20	83	313	650	1,036	1,372	835	-
2002.....	4,336	1	5	11	12	15	78	343	627	1,095	1,375	774	-
2001.....	4,331	-	7	24	4	22	114	337	652	1,070	1,388	713	-
2000.....	4,277	1	5	14	9	13	101	329	620	1,083	1,368	734	-
1999.....	4,041	1	8	15	5	14	97	309	598	1,069	1,235	690	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Non-Hispanic white, both sexes													
2003.....	10,225	1	12	26	10	32	219	936	2,004	2,570	3,031	1,384	-
2002.....	10,179	2	5	23	15	24	211	1,011	1,898	2,754	2,903	1,333	-
2001.....	10,073	-	11	17	8	23	235	998	1,904	2,741	2,880	1,256	-
2000.....	9,884	-	7	16	11	24	239	977	1,794	2,665	2,871	1,280	-
1999.....	9,319	2	11	21	9	28	238	877	1,740	2,613	2,636	1,144	-
Non-Hispanic white, male													
2003.....	6,408	1	4	12	6	17	159	682	1,463	1,650	1,787	627	-
2002.....	6,363	1	-	14	5	15	150	741	1,355	1,785	1,668	629	-
2001.....	6,263	-	6	1	5	11	157	726	1,340	1,790	1,616	611	-
2000.....	6,082	-	4	6	6	13	165	701	1,258	1,697	1,628	604	-
1999.....	5,754	1	7	9	7	18	165	618	1,220	1,662	1,528	519	-
Non-Hispanic white, female													
2003.....	3,817	-	8	14	4	15	60	254	541	920	1,244	757	-
2002.....	3,816	1	5	9	10	9	61	270	543	969	1,235	704	-
2001.....	3,810	-	5	16	3	12	78	272	564	951	1,264	645	-
2000.....	3,802	-	3	10	5	11	74	276	536	968	1,243	676	-
1999.....	3,565	1	4	12	2	10	73	259	520	951	1,108	625	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Non-Hispanic black, both sexes													
2003.....	1,105	-	-	9	7	19	63	153	261	246	232	115	-
2002.....	1,062	-	-	5	3	17	45	163	226	260	250	93	-
2001.....	1,105	-	3	12	10	19	72	163	240	249	241	96	-
2000.....	1,033	1	5	8	6	10	51	142	213	271	233	93	-
1999.....	1,028	-	4	6	7	20	61	154	202	263	213	98	-
Non-Hispanic black, male													
2003.....	667	-	-	5	4	14	43	107	169	154	129	42	-
2002.....	626	-	-	3	2	11	31	103	164	151	125	36	-
2001.....	673	-	1	4	10	10	40	111	168	153	136	40	-
2000.....	625	-	3	4	2	8	27	94	142	179	124	42	-
1999.....	618	-	2	3	4	16	42	109	137	159	104	42	-
Non-Hispanic black, female													
2003.....	438	-	-	4	3	5	20	46	92	92	103	73	-
2002.....	436	-	-	2	1	6	14	60	62	109	125	57	-
2001.....	432	-	2	8	-	9	32	52	72	96	105	56	-
2000.....	408	1	2	4	4	2	24	48	71	92	109	51	-
1999.....	410	-	2	3	3	4	19	45	65	104	109	56	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Origin not stated 2/, both sexes													
2003.....	22	-	-	-	1	-	-	2	3	8	4	4	-
2002.....	29	-	-	-	-	-	1	3	7	8	9	1	-
2001.....	23	-	-	-	-	-	1	3	3	7	5	4	-
2000.....	45	-	-	-	-	-	-	4	7	20	7	7	-
1999.....	27	-	-	-	-	1	2	3	3	5	10	3	-
Origin not stated 2/, male													
2003.....	17	-	-	-	1	-	-	1	3	6	4	2	-
2002.....	20	-	-	-	-	-	1	2	6	6	4	1	-
2001.....	14	-	-	-	-	-	1	2	2	5	3	1	-
2000.....	30	-	-	-	-	-	-	2	5	14	5	4	-
1999.....	13	-	-	-	-	-	1	2	2	4	3	1	-
Origin not stated 2/, female													
2003.....	5	-	-	-	-	-	-	1	-	2	-	2	-
2002.....	9	-	-	-	-	-	-	1	1	2	5	-	-
2001.....	9	-	-	-	-	-	-	1	1	2	2	3	-
2000.....	15	-	-	-	-	-	-	2	2	6	2	3	-
1999.....	14	-	-	-	-	1	1	1	1	1	7	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
All origins, both sexes													
2003.....	12,483	-	-	-	1	4	98	522	1,391	2,736	4,679	3,052	-
2002.....	12,628	-	-	3	1	14	123	503	1,310	2,907	4,707	3,059	1
2001.....	12,225	-	-	1	3	15	128	475	1,265	2,814	4,488	3,036	-
2000.....	12,002	-	-	1	3	10	111	440	1,223	2,855	4,407	2,952	-
1999.....	11,911	-	-	1	1	16	100	463	1,244	2,923	4,317	2,845	1
All origins, male													
2003.....	8,576	-	-	-	1	2	64	401	1,049	1,998	3,254	1,807	-
2002.....	8,504	-	-	1	-	9	78	352	984	2,089	3,199	1,791	1
2001.....	8,343	-	-	-	3	8	70	351	956	2,088	3,116	1,751	-
2000.....	8,163	-	-	-	3	2	62	322	916	2,094	3,054	1,710	-
1999.....	8,105	-	-	1	-	9	63	350	933	2,132	3,008	1,608	1
All origins, female													
2003.....	3,907	-	-	-	-	2	34	121	342	738	1,425	1,245	-
2002.....	4,124	-	-	2	1	5	45	151	326	818	1,508	1,268	-
2001.....	3,882	-	-	1	-	7	58	124	309	726	1,372	1,285	-
2000.....	3,839	-	-	1	-	8	49	118	307	761	1,353	1,242	-
1999.....	3,806	-	-	-	1	7	37	113	311	791	1,309	1,237	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
Hispanic, both sexes													
2003.....	394	-	-	-	-	1	4	25	54	96	133	81	-
2002.....	387	-	-	1	-	-	5	24	57	91	110	99	-
2001.....	336	-	-	-	2	-	4	21	41	90	104	74	-
2000.....	307	-	-	-	-	1	8	17	50	71	89	71	-
1999.....	301	-	-	-	-	2	4	22	51	88	79	55	-
Hispanic, male													
2003.....	265	-	-	-	-	-	2	15	40	71	87	50	-
2002.....	252	-	-	-	-	-	2	16	39	67	68	60	-
2001.....	223	-	-	-	2	-	1	11	37	59	67	46	-
2000.....	216	-	-	-	-	-	6	12	37	50	69	42	-
1999.....	222	-	-	-	-	2	4	16	43	69	52	36	-
Hispanic, female													
2003.....	129	-	-	-	-	1	2	10	14	25	46	31	-
2002.....	135	-	-	1	-	-	3	8	18	24	42	39	-
2001.....	113	-	-	-	-	-	3	10	4	31	37	28	-
2000.....	91	-	-	-	-	1	2	5	13	21	20	29	-
1999.....	79	-	-	-	-	-	-	6	8	19	27	19	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
Non-Hispanic 1/, both sexes													
2003.....	12,067	-	-	-	1	3	94	495	1,336	2,637	4,534	2,967	-
2002.....	12,211	-	-	2	1	14	118	476	1,248	2,809	4,591	2,952	-
2001.....	11,858	-	-	1	1	15	124	452	1,220	2,718	4,372	2,955	-
2000.....	11,665	-	-	1	3	9	102	423	1,169	2,780	4,304	2,874	-
1999.....	11,582	-	-	1	1	14	96	439	1,190	2,832	4,228	2,780	1
Non-Hispanic 1/, male													
2003.....	8,294	-	-	-	1	2	62	384	1,008	1,925	3,157	1,755	-
2002.....	8,235	-	-	1	-	9	76	335	941	2,019	3,128	1,726	-
2001.....	8,100	-	-	-	1	8	69	339	917	2,025	3,041	1,700	-
2000.....	7,925	-	-	-	3	2	56	310	876	2,040	2,974	1,664	-
1999.....	7,865	-	-	1	-	7	59	332	888	2,060	2,950	1,567	1
Non-Hispanic 1/, female													
2003.....	3,773	-	-	-	-	1	32	111	328	712	1,377	1,212	-
2002.....	3,976	-	-	1	1	5	42	141	307	790	1,463	1,226	-
2001.....	3,758	-	-	1	-	7	55	113	303	693	1,331	1,255	-
2000.....	3,740	-	-	1	-	7	46	113	293	740	1,330	1,210	-
1999.....	3,717	-	-	-	1	7	37	107	302	772	1,278	1,213	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
Non-Hispanic white, both sexes													
2003.....	11,003	-	-	-	1	3	72	407	1,198	2,399	4,168	2,755	-
2002.....	11,109	-	-	2	-	10	100	416	1,112	2,531	4,232	2,706	-
2001.....	10,823	-	-	1	1	14	100	386	1,076	2,479	4,026	2,740	-
2000.....	10,712	-	-	1	2	8	91	374	1,056	2,533	3,957	2,690	-
1999.....	10,548	-	-	1	-	8	77	375	1,042	2,556	3,921	2,567	1
Non-Hispanic white, male													
2003.....	7,692	-	-	-	1	2	48	328	921	1,777	2,952	1,663	-
2002.....	7,649	-	-	1	-	6	66	302	866	1,864	2,921	1,623	-
2001.....	7,541	-	-	-	1	8	56	296	830	1,884	2,856	1,610	-
2000.....	7,390	-	-	-	2	2	52	279	803	1,897	2,784	1,571	-
1999.....	7,290	-	-	1	-	3	46	281	800	1,907	2,771	1,480	1
Non-Hispanic white, female													
2003.....	3,311	-	-	-	-	1	24	79	277	622	1,216	1,092	-
2002.....	3,460	-	-	1	-	4	34	114	246	667	1,311	1,083	-
2001.....	3,282	-	-	1	-	6	44	90	246	595	1,170	1,130	-
2000.....	3,322	-	-	1	-	6	39	95	253	636	1,173	1,119	-
1999.....	3,258	-	-	-	-	5	31	94	242	649	1,150	1,087	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
Non-Hispanic black, both sexes													
2003.....	907	-	-	-	-	-	20	78	109	204	315	181	-
2002.....	931	-	-	-	-	4	13	49	116	242	295	212	-
2001.....	870	-	-	-	-	1	23	55	125	197	291	178	-
2000.....	842	-	-	-	1	-	10	44	99	218	314	156	-
1999.....	906	-	-	-	1	4	17	60	130	244	262	188	-
Non-Hispanic black, male													
2003.....	496	-	-	-	-	-	12	48	66	123	172	75	-
2002.....	475	-	-	-	-	3	7	26	63	128	164	84	-
2001.....	444	-	-	-	-	-	13	36	72	111	144	68	-
2000.....	460	-	-	-	1	-	4	27	63	119	171	75	-
1999.....	487	-	-	-	-	2	12	47	76	128	149	73	-
Non-Hispanic black, female													
2003.....	411	-	-	-	-	-	8	30	43	81	143	106	-
2002.....	456	-	-	-	-	1	6	23	53	114	131	128	-
2001.....	426	-	-	-	-	1	10	19	53	86	147	110	-
2000.....	382	-	-	-	-	-	6	17	36	99	143	81	-
1999.....	419	-	-	-	1	2	5	13	54	116	113	115	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasm of bladder (C67)													
Origin not stated 2/, both sexes													
2003.....	22	-	-	-	-	-	-	2	1	3	12	4	-
2002.....	30	-	-	-	-	-	-	3	5	7	6	8	1
2001.....	31	-	-	-	-	-	-	2	4	6	12	7	-
2000.....	30	-	-	-	-	-	1	-	4	4	14	7	-
1999.....	28	-	-	-	-	-	-	2	3	3	10	10	-
Origin not stated 2/, male													
2003.....	17	-	-	-	-	-	-	2	1	2	10	2	-
2002.....	17	-	-	-	-	-	-	1	4	3	3	5	1
2001.....	20	-	-	-	-	-	-	1	2	4	8	5	-
2000.....	22	-	-	-	-	-	-	-	3	4	11	4	-
1999.....	18	-	-	-	-	-	-	2	2	3	6	5	-
Origin not stated 2/, female													
2003.....	5	-	-	-	-	-	-	-	-	1	2	2	-
2002.....	13	-	-	-	-	-	-	2	1	4	3	3	-
2001.....	11	-	-	-	-	-	-	1	2	2	4	2	-
2000.....	8	-	-	-	-	-	1	-	1	-	3	3	-
1999.....	10	-	-	-	-	-	-	-	1	-	4	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
All origins, both sexes													
2003.....	12,901	18	104	336	236	405	1,079	1,962	2,698	2,836	2,510	717	-
2002.....	12,830	12	114	337	210	388	1,009	2,058	2,560	2,954	2,435	753	-
2001.....	12,609	16	100	300	224	416	1,028	1,951	2,464	2,823	2,566	721	-
2000.....	12,655	20	104	316	225	406	1,096	1,924	2,337	3,032	2,478	717	-
1999.....	12,765	10	113	308	218	444	1,065	1,880	2,453	3,119	2,492	663	-
All origins, male													
2003.....	7,122	9	56	175	136	232	639	1,233	1,530	1,556	1,267	289	-
2002.....	7,125	8	63	191	136	223	618	1,225	1,494	1,678	1,210	279	-
2001.....	7,015	8	52	145	129	262	638	1,173	1,470	1,589	1,288	261	-
2000.....	7,011	10	58	176	132	264	704	1,148	1,379	1,678	1,187	275	-
1999.....	7,001	8	56	154	131	276	638	1,120	1,417	1,728	1,226	247	-
All origins, female													
2003.....	5,779	9	48	161	100	173	440	729	1,168	1,280	1,243	428	-
2002.....	5,705	4	51	146	74	165	391	833	1,066	1,276	1,225	474	-
2001.....	5,594	8	48	155	95	154	390	778	994	1,234	1,278	460	-
2000.....	5,644	10	46	140	93	142	392	776	958	1,354	1,291	442	-
1999.....	5,764	2	57	154	87	168	427	760	1,036	1,391	1,266	416	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
Hispanic, both sexes													
2003.....	669	5	20	62	23	44	63	97	123	116	98	18	-
2002.....	649	2	21	52	24	33	55	123	115	118	82	24	-
2001.....	617	4	20	37	28	38	77	87	109	117	82	18	-
2000.....	631	4	16	63	28	45	76	92	106	105	72	24	-
1999.....	568	4	27	48	18	31	65	88	99	104	67	17	-
Hispanic, male													
2003.....	368	2	10	36	15	26	38	61	62	64	43	11	-
2002.....	359	2	12	27	16	16	30	80	62	65	39	10	-
2001.....	335	1	10	16	19	21	41	50	59	72	40	6	-
2000.....	346	1	10	41	16	27	40	41	62	59	38	11	-
1999.....	293	4	11	22	10	17	45	42	60	48	31	3	-
Hispanic, female													
2003.....	301	3	10	26	8	18	25	36	61	52	55	7	-
2002.....	290	-	9	25	8	17	25	43	53	53	43	14	-
2001.....	282	3	10	21	9	17	36	37	50	45	42	12	-
2000.....	285	3	6	22	12	18	36	51	44	46	34	13	-
1999.....	275	-	16	26	8	14	20	46	39	56	36	14	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
Non-Hispanic 1/, both sexes													
2003.....	12,213	13	84	273	213	361	1,012	1,863	2,573	2,715	2,408	698	-
2002.....	12,152	10	93	284	184	355	951	1,929	2,438	2,831	2,349	728	-
2001.....	11,967	12	79	261	196	377	950	1,862	2,346	2,703	2,479	702	-
2000.....	11,995	16	87	253	197	361	1,019	1,828	2,222	2,923	2,398	691	-
1999.....	12,173	6	86	260	199	413	1,000	1,789	2,348	3,010	2,418	644	-
Non-Hispanic 1/, male													
2003.....	6,746	7	46	139	121	206	598	1,171	1,468	1,489	1,223	278	-
2002.....	6,752	6	51	164	120	207	586	1,142	1,427	1,612	1,168	269	-
2001.....	6,667	7	41	128	110	241	596	1,122	1,406	1,516	1,245	255	-
2000.....	6,646	9	47	135	116	237	663	1,103	1,313	1,615	1,145	263	-
1999.....	6,696	4	45	132	121	259	593	1,076	1,353	1,678	1,191	244	-
Non-Hispanic 1/, female													
2003.....	5,467	6	38	134	92	155	414	692	1,105	1,226	1,185	420	-
2002.....	5,400	4	42	120	64	148	365	787	1,011	1,219	1,181	459	-
2001.....	5,300	5	38	133	86	136	354	740	940	1,187	1,234	447	-
2000.....	5,349	7	40	118	81	124	356	725	909	1,308	1,253	428	-
1999.....	5,477	2	41	128	78	154	407	713	995	1,332	1,227	400	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
Non-Hispanic white, both sexes													
2003.....	11,179	10	65	209	174	314	911	1,678	2,373	2,520	2,276	649	-
2002.....	11,139	5	59	214	150	305	865	1,756	2,243	2,646	2,209	687	-
2001.....	11,004	8	63	205	155	335	864	1,695	2,155	2,509	2,356	659	-
2000.....	11,026	12	65	194	156	315	896	1,678	2,047	2,756	2,260	647	-
1999.....	11,238	3	70	195	152	372	902	1,635	2,171	2,839	2,293	606	-
Non-Hispanic white, male													
2003.....	6,178	6	38	103	97	181	533	1,059	1,362	1,385	1,153	261	-
2002.....	6,195	3	34	116	101	175	540	1,051	1,306	1,515	1,097	257	-
2001.....	6,154	5	35	96	88	217	547	1,030	1,293	1,427	1,181	235	-
2000.....	6,128	8	35	101	93	200	591	1,015	1,219	1,531	1,086	249	-
1999.....	6,193	2	39	104	91	233	533	980	1,252	1,591	1,136	232	-
Non-Hispanic white, female													
2003.....	5,001	4	27	106	77	133	378	619	1,011	1,135	1,123	388	-
2002.....	4,944	2	25	98	49	130	325	705	937	1,131	1,112	430	-
2001.....	4,850	3	28	109	67	118	317	665	862	1,082	1,175	424	-
2000.....	4,898	4	30	93	63	115	305	663	828	1,225	1,174	398	-
1999.....	5,045	1	31	91	61	139	369	655	919	1,248	1,157	374	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
Non-Hispanic black, both sexes													
2003.....	799	2	17	48	34	33	82	145	151	149	102	36	-
2002.....	766	3	27	54	26	33	68	125	144	142	112	32	-
2001.....	736	1	10	44	29	28	63	133	145	153	92	38	-
2000.....	762	2	18	51	35	36	88	119	138	133	109	33	-
1999.....	754	3	11	54	37	36	76	128	142	132	104	31	-
Non-Hispanic black, male													
2003.....	440	1	6	27	21	18	55	88	81	82	54	7	-
2002.....	400	2	12	35	14	22	35	56	92	70	55	7	-
2001.....	378	-	3	25	16	14	33	75	87	70	40	15	-
2000.....	407	1	10	30	21	28	52	71	76	66	43	9	-
1999.....	399	2	4	24	24	23	47	81	76	68	40	10	-
Non-Hispanic black, female													
2003.....	359	1	11	21	13	15	27	57	70	67	48	29	-
2002.....	366	1	15	19	12	11	33	69	52	72	57	25	-
2001.....	358	1	7	19	13	14	30	58	58	83	52	23	-
2000.....	355	1	8	21	14	8	36	48	62	67	66	24	-
1999.....	355	1	7	30	13	13	29	47	66	64	64	21	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Origin not stated 2/, both sexes

2003.....	19	-	-	1	-	-	4	2	2	5	4	1	-
2002.....	29	-	-	1	2	-	3	6	7	5	4	1	-
2001.....	25	-	1	2	-	1	1	2	9	3	5	1	-
2000.....	29	-	1	-	-	-	1	4	9	4	8	2	-
1999.....	24	-	-	-	1	-	-	3	6	5	7	2	-

Origin not stated 2/, male

2003.....	8	-	-	-	-	-	3	1	-	3	1	-	-
2002.....	14	-	-	-	-	-	2	3	5	1	3	-	-
2001.....	13	-	1	1	-	-	1	1	5	1	3	-	-
2000.....	19	-	1	-	-	-	1	4	4	4	4	1	-
1999.....	12	-	-	-	-	-	-	2	4	2	4	-	-

Origin not stated 2/, female

2003.....	11	-	-	1	-	-	1	1	2	2	3	1	-
2002.....	15	-	-	1	2	-	1	3	2	4	1	1	-
2001.....	12	-	-	1	-	1	-	1	4	2	2	1	-
2000.....	10	-	-	-	-	-	-	-	5	-	4	1	-
1999.....	12	-	-	-	1	-	-	1	2	3	3	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
All origins, both sexes													
2003.....	55,679	31	138	394	606	865	1,728	3,785	7,471	12,885	18,442	9,334	-
2002.....	56,225	28	119	379	703	916	1,751	3,960	7,500	13,508	18,289	9,072	-
2001.....	56,341	27	157	367	705	988	1,857	4,064	7,455	13,659	18,257	8,805	-
2000.....	56,518	39	146	372	718	918	1,853	4,025	7,284	14,041	18,448	8,674	-
1999.....	56,318	30	137	351	720	960	1,913	3,950	7,554	14,332	18,089	8,282	-
All origins, male													
2003.....	30,076	10	80	225	375	535	1,043	2,294	4,557	7,550	9,524	3,883	-
2002.....	30,302	15	63	217	417	545	1,038	2,341	4,470	7,903	9,553	3,740	-
2001.....	30,060	12	82	217	426	574	1,133	2,442	4,412	7,859	9,327	3,576	-
2000.....	29,888	21	76	223	455	539	1,075	2,399	4,205	8,038	9,399	3,458	-
1999.....	29,721	12	75	206	437	557	1,144	2,368	4,426	8,067	9,100	3,329	-
All origins, female													
2003.....	25,603	21	58	169	231	330	685	1,491	2,914	5,335	8,918	5,451	-
2002.....	25,923	13	56	162	286	371	713	1,619	3,030	5,605	8,736	5,332	-
2001.....	26,281	15	75	150	279	414	724	1,622	3,043	5,800	8,930	5,229	-
2000.....	26,630	18	70	149	263	379	778	1,626	3,079	6,003	9,049	5,216	-
1999.....	26,597	18	62	145	283	403	769	1,582	3,128	6,265	8,989	4,953	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Hispanic, both sexes													
2003.....	2,836	6	44	97	131	147	202	323	425	576	635	250	-
2002.....	2,829	3	28	93	145	139	225	282	437	634	604	239	-
2001.....	2,747	7	38	87	144	154	211	279	442	597	573	215	-
2000.....	2,643	8	29	74	129	124	200	280	427	633	523	216	-
1999.....	2,538	8	28	75	128	135	190	299	396	606	485	188	-
Hispanic, male													
2003.....	1,528	1	26	56	89	96	130	182	241	311	299	97	-
2002.....	1,518	1	12	47	97	92	132	163	245	331	290	108	-
2001.....	1,577	2	21	48	99	99	142	164	251	338	318	95	-
2000.....	1,460	5	14	47	83	83	115	172	247	338	258	98	-
1999.....	1,319	3	16	42	73	75	108	175	226	312	226	63	-
Hispanic, female													
2003.....	1,308	5	18	41	42	51	72	141	184	265	336	153	-
2002.....	1,311	2	16	46	48	47	93	119	192	303	314	131	-
2001.....	1,170	5	17	39	45	55	69	115	191	259	255	120	-
2000.....	1,183	3	15	27	46	41	85	108	180	295	265	118	-
1999.....	1,219	5	12	33	55	60	82	124	170	294	259	125	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Non-Hispanic 1/, both sexes													
2003.....	52,736	25	94	296	474	717	1,525	3,450	7,029	12,284	17,777	9,065	-
2002.....	53,249	24	91	286	554	775	1,518	3,663	7,048	12,839	17,636	8,815	-
2001.....	53,458	20	117	279	559	828	1,640	3,772	6,994	13,027	17,645	8,577	-
2000.....	53,749	31	116	295	586	793	1,645	3,735	6,833	13,379	17,893	8,443	-
1999.....	53,669	22	108	276	592	823	1,719	3,642	7,143	13,704	17,561	8,079	-
Non-Hispanic 1/, male													
2003.....	28,485	9	54	168	285	438	912	2,103	4,304	7,229	9,208	3,775	-
2002.....	28,698	13	51	170	317	452	900	2,168	4,217	7,546	9,236	3,628	-
2001.....	28,410	10	60	169	326	473	987	2,269	4,151	7,505	8,985	3,475	-
2000.....	28,358	16	61	175	371	455	956	2,219	3,941	7,682	9,127	3,355	-
1999.....	28,347	9	58	164	364	481	1,032	2,190	4,195	7,740	8,853	3,261	-
Non-Hispanic 1/, female													
2003.....	24,251	16	40	128	189	279	613	1,347	2,725	5,055	8,569	5,290	-
2002.....	24,551	11	40	116	237	323	618	1,495	2,831	5,293	8,400	5,187	-
2001.....	25,048	10	57	110	233	355	653	1,503	2,843	5,522	8,660	5,102	-
2000.....	25,391	15	55	120	215	338	689	1,516	2,892	5,697	8,766	5,088	-
1999.....	25,322	13	50	112	228	342	687	1,452	2,948	5,964	8,708	4,818	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Non-Hispanic white, both sexes													
2003.....	46,553	23	72	218	355	524	1,166	2,727	5,996	10,779	16,253	8,440	-
2002.....	47,058	15	72	213	430	570	1,183	2,897	5,930	11,327	16,202	8,219	-
2001.....	47,190	17	97	198	422	595	1,255	3,057	5,878	11,492	16,199	7,980	-
2000.....	47,695	30	92	232	420	585	1,283	3,017	5,812	11,911	16,528	7,785	-
1999.....	47,608	16	91	199	437	618	1,352	2,980	6,042	12,208	16,172	7,493	-
Non-Hispanic white, male													
2003.....	25,390	7	42	127	214	324	708	1,682	3,719	6,471	8,552	3,544	-
2002.....	25,523	9	42	125	254	333	701	1,755	3,597	6,746	8,557	3,404	-
2001.....	25,162	8	47	111	252	352	748	1,873	3,519	6,693	8,307	3,252	-
2000.....	25,272	16	50	134	268	338	775	1,803	3,379	6,899	8,491	3,119	-
1999.....	25,219	9	48	120	277	357	816	1,812	3,591	6,967	8,202	3,020	-
Non-Hispanic white, female													
2003.....	21,163	16	30	91	141	200	458	1,045	2,277	4,308	7,701	4,896	-
2002.....	21,535	6	30	88	176	237	482	1,142	2,333	4,581	7,645	4,815	-
2001.....	22,028	9	50	87	170	243	507	1,184	2,359	4,799	7,892	4,728	-
2000.....	22,423	14	42	98	152	247	508	1,214	2,433	5,012	8,037	4,666	-
1999.....	22,389	7	43	79	160	261	536	1,168	2,451	5,241	7,970	4,473	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Non-Hispanic black, both sexes													
2003.....	5,059	1	14	57	88	155	299	620	853	1,237	1,223	512	-
2002.....	5,080	5	14	51	87	156	266	624	935	1,268	1,166	508	-
2001.....	5,162	3	13	62	107	181	322	584	915	1,288	1,183	504	-
2000.....	5,026	1	19	43	130	158	302	615	868	1,232	1,101	557	-
1999.....	5,038	4	14	61	133	157	307	554	917	1,247	1,142	502	-
Non-Hispanic black, male													
2003.....	2,517	1	8	31	52	87	170	366	491	613	521	177	-
2002.....	2,593	2	6	32	44	93	162	342	517	679	530	186	-
2001.....	2,642	2	9	43	57	98	199	326	512	679	538	179	-
2000.....	2,535	-	8	26	79	90	149	346	481	653	510	193	-
1999.....	2,566	-	8	38	73	97	180	318	500	645	521	186	-
Non-Hispanic black, female													
2003.....	2,542	-	6	26	36	68	129	254	362	624	702	335	-
2002.....	2,487	3	8	19	43	63	104	282	418	589	636	322	-
2001.....	2,520	1	4	19	50	83	123	258	403	609	645	325	-
2000.....	2,491	1	11	17	51	68	153	269	387	579	591	364	-
1999.....	2,472	4	6	23	60	60	127	236	417	602	621	316	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Origin not stated 2/, both sexes													
2003.....	107	-	-	1	1	1	1	12	17	25	30	19	-
2002.....	147	1	-	-	4	2	8	15	15	35	49	18	-
2001.....	136	-	2	1	2	6	6	13	19	35	39	13	-
2000.....	126	-	1	3	3	1	8	10	24	29	32	15	-
1999.....	111	-	1	-	-	2	4	9	15	22	43	15	-
Origin not stated 2/, male													
2003.....	63	-	-	1	1	1	1	9	12	10	17	11	-
2002.....	86	1	-	-	3	1	6	10	8	26	27	4	-
2001.....	73	-	1	-	1	2	4	9	10	16	24	6	-
2000.....	70	-	1	1	1	1	4	8	17	18	14	5	-
1999.....	55	-	1	-	-	1	4	3	5	15	21	5	-
Origin not stated 2/, female													
2003.....	44	-	-	-	-	-	-	3	5	15	13	8	-
2002.....	61	-	-	-	1	1	2	5	7	9	22	14	-
2001.....	63	-	1	1	1	4	2	4	9	19	15	7	-
2000.....	56	-	-	2	2	-	4	2	7	11	18	10	-
1999.....	56	-	-	-	-	1	-	6	10	7	22	10	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
All origins, both sexes													
2003.....	1,347	-	-	5	64	167	154	154	198	216	277	112	-
2002.....	1,352	-	-	6	86	152	159	177	168	232	247	125	-
2001.....	1,323	-	-	6	78	167	161	182	143	211	278	97	-
2000.....	1,287	-	-	8	73	145	151	161	158	232	255	104	-
1999.....	1,403	-	1	9	94	148	165	165	178	283	262	98	-
All origins, male													
2003.....	725	-	-	4	30	88	95	91	119	128	132	38	-
2002.....	751	-	-	5	41	81	92	105	106	146	125	50	-
2001.....	725	-	-	3	43	90	101	110	85	129	130	34	-
2000.....	686	-	-	4	35	79	82	99	87	134	121	45	-
1999.....	785	-	1	3	55	86	103	98	115	160	119	45	-
All origins, female													
2003.....	622	-	-	1	34	79	59	63	79	88	145	74	-
2002.....	601	-	-	1	45	71	67	72	62	86	122	75	-
2001.....	598	-	-	3	35	77	60	72	58	82	148	63	-
2000.....	601	-	-	4	38	66	69	62	71	98	134	59	-
1999.....	618	-	-	6	39	62	62	67	63	123	143	53	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
Hispanic, both sexes													
2003.....	91	-	-	1	8	16	6	15	8	19	12	6	-
2002.....	99	-	-	4	9	15	12	15	10	13	18	3	-
2001.....	99	-	-	1	7	13	13	14	14	21	11	5	-
2000.....	95	-	-	-	9	14	8	13	13	11	23	4	-
1999.....	113	-	-	3	8	7	12	17	17	26	18	5	-
Hispanic, male													
2003.....	52	-	-	1	5	8	5	8	5	11	8	1	-
2002.....	53	-	-	3	4	7	8	9	7	6	8	1	-
2001.....	66	-	-	1	4	8	8	11	11	15	5	3	-
2000.....	63	-	-	-	7	7	6	11	8	10	12	2	-
1999.....	65	-	-	2	6	3	8	11	9	13	11	2	-
Hispanic, female													
2003.....	39	-	-	-	3	8	1	7	3	8	4	5	-
2002.....	46	-	-	1	5	8	4	6	3	7	10	2	-
2001.....	33	-	-	-	3	5	5	3	3	6	6	2	-
2000.....	32	-	-	-	2	7	2	2	5	1	11	2	-
1999.....	48	-	-	1	2	4	4	6	8	13	7	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
Non-Hispanic 1/, both sexes													
2003.....	1,253	-	-	4	56	151	147	138	190	197	264	106	-
2002.....	1,248	-	-	2	76	136	146	161	158	219	228	122	-
2001.....	1,221	-	-	5	71	154	147	167	129	189	267	92	-
2000.....	1,189	-	-	8	64	131	143	148	143	221	231	100	-
1999.....	1,286	-	1	6	86	141	153	147	159	257	243	93	-
Non-Hispanic 1/, male													
2003.....	672	-	-	3	25	80	89	83	114	117	124	37	-
2002.....	693	-	-	2	36	73	83	95	99	140	116	49	-
2001.....	657	-	-	2	39	82	92	98	74	114	125	31	-
2000.....	621	-	-	4	28	72	76	88	78	124	108	43	-
1999.....	719	-	1	1	49	83	95	87	105	147	108	43	-
Non-Hispanic 1/, female													
2003.....	581	-	-	1	31	71	58	55	76	80	140	69	-
2002.....	555	-	-	-	40	63	63	66	59	79	112	73	-
2001.....	564	-	-	3	32	72	55	69	55	75	142	61	-
2000.....	568	-	-	4	36	59	67	60	65	97	123	57	-
1999.....	567	-	-	5	37	58	58	60	54	110	135	50	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
Non-Hispanic white, both sexes													
2003.....	1,099	-	-	4	43	118	125	123	162	177	246	101	-
2002.....	1,093	-	-	1	63	112	118	135	132	200	215	117	-
2001.....	1,069	-	-	4	55	113	124	146	110	178	249	90	-
2000.....	1,059	-	-	5	48	117	119	128	123	200	224	95	-
1999.....	1,149	-	1	5	67	116	134	129	144	234	235	84	-
Non-Hispanic white, male													
2003.....	585	-	-	3	19	66	78	73	96	101	115	34	-
2002.....	599	-	-	1	32	57	65	75	80	130	111	48	-
2001.....	577	-	-	2	34	62	76	85	63	109	117	29	-
2000.....	551	-	-	2	23	63	64	76	65	112	105	41	-
1999.....	643	-	1	1	39	69	84	73	97	135	105	39	-
Non-Hispanic white, female													
2003.....	514	-	-	1	24	52	47	50	66	76	131	67	-
2002.....	494	-	-	-	31	55	53	60	52	70	104	69	-
2001.....	492	-	-	2	21	51	48	61	47	69	132	61	-
2000.....	508	-	-	3	25	54	55	52	58	88	119	54	-
1999.....	506	-	-	4	28	47	50	56	47	99	130	45	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
Non-Hispanic black, both sexes													
2003.....	119	-	-	-	10	31	18	14	21	17	8	-	-
2002.....	126	-	-	1	10	23	25	20	20	15	9	3	-
2001.....	136	-	-	1	15	39	21	17	18	11	12	2	-
2000.....	109	-	-	2	14	12	24	17	17	15	5	3	-
1999.....	125	-	-	-	19	25	18	18	14	18	6	7	-
Non-Hispanic black, male													
2003.....	68	-	-	-	5	12	10	9	15	13	4	-	-
2002.....	80	-	-	1	4	15	17	15	15	8	4	1	-
2001.....	72	-	-	-	4	20	15	11	10	5	5	2	-
2000.....	57	-	-	1	5	8	12	9	11	7	3	1	-
1999.....	74	-	-	-	10	14	11	14	8	12	3	2	-
Non-Hispanic black, female													
2003.....	51	-	-	-	5	19	8	5	6	4	4	-	-
2002.....	46	-	-	-	6	8	8	5	5	7	5	2	-
2001.....	64	-	-	1	11	19	6	6	8	6	7	-	-
2000.....	52	-	-	1	9	4	12	8	6	8	2	2	-
1999.....	51	-	-	-	9	11	7	4	6	6	3	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hodgkin's disease (C81)													
Origin not stated 2/, both sexes													
2003.....	3	-	-	-	-	-	1	1	-	-	1	-	-
2002.....	5	-	-	-	1	1	1	1	-	-	1	-	-
2001.....	3	-	-	-	-	-	1	1	-	1	-	-	-
2000.....	3	-	-	-	-	-	-	-	2	-	1	-	-
1999.....	4	-	-	-	-	-	-	1	2	-	1	-	-
Origin not stated 2/, male													
2003.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	5	-	-	-	1	1	1	1	-	-	1	-	-
2001.....	2	-	-	-	-	-	1	1	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	1	-	1	-	-
1999.....	1	-	-	-	-	-	-	-	1	-	-	-	-
Origin not stated 2/, female													
2003.....	2	-	-	-	-	-	-	1	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	1	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	3	-	-	-	-	-	-	1	1	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
All origins, both sexes													
2003.....	21,475	-	5	45	140	252	630	1,548	3,007	4,919	7,217	3,712	-
2002.....	21,910	2	15	57	142	281	687	1,581	3,049	5,264	7,283	3,549	-
2001.....	22,305	1	1	38	148	298	756	1,732	3,098	5,358	7,369	3,506	-
2000.....	22,729	1	7	47	180	293	751	1,745	3,170	5,598	7,535	3,402	-
1999.....	22,802	2	16	37	144	289	789	1,741	3,154	5,848	7,554	3,228	-
All origins, male													
2003.....	11,399	-	5	26	88	165	406	972	1,848	2,829	3,563	1,497	-
2002.....	11,538	1	6	40	93	178	439	978	1,838	2,986	3,594	1,385	-
2001.....	11,710	1	1	29	94	179	500	1,113	1,825	3,025	3,558	1,385	-
2000.....	11,812	-	5	28	126	172	482	1,112	1,849	3,105	3,634	1,299	-
1999.....	11,794	-	9	30	91	178	510	1,111	1,847	3,176	3,596	1,246	-
All origins, female													
2003.....	10,076	-	-	19	52	87	224	576	1,159	2,090	3,654	2,215	-
2002.....	10,372	1	9	17	49	103	248	603	1,211	2,278	3,689	2,164	-
2001.....	10,595	-	-	9	54	119	256	619	1,273	2,333	3,811	2,121	-
2000.....	10,917	1	2	19	54	121	269	633	1,321	2,493	3,901	2,103	-
1999.....	11,008	2	7	7	53	111	279	630	1,307	2,672	3,958	1,982	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Hispanic, both sexes													
2003.....	1,057	-	3	10	27	46	82	116	176	233	259	105	-
2002.....	1,068	-	2	7	25	41	78	108	186	268	262	91	-
2001.....	1,008	-	-	6	22	37	77	122	201	208	240	95	-
2000.....	1,026	-	2	2	21	38	83	113	190	277	210	90	-
1999.....	958	-	1	3	19	37	68	131	153	261	205	80	-
Hispanic, male													
2003.....	569	-	3	5	19	33	56	65	104	129	118	37	-
2002.....	549	-	-	3	19	25	45	65	109	136	111	36	-
2001.....	579	-	-	6	15	24	56	74	115	122	131	36	-
2000.....	560	-	2	1	17	23	51	72	112	143	98	41	-
1999.....	493	-	-	2	10	18	43	84	91	125	92	28	-
Hispanic, female													
2003.....	488	-	-	5	8	13	26	51	72	104	141	68	-
2002.....	519	-	2	4	6	16	33	43	77	132	151	55	-
2001.....	429	-	-	-	7	13	21	48	86	86	109	59	-
2000.....	466	-	-	1	4	15	32	41	78	134	112	49	-
1999.....	465	-	1	1	9	19	25	47	62	136	113	52	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Non-Hispanic 1/, both sexes													
2003.....	20,379	-	2	35	112	206	548	1,427	2,823	4,679	6,947	3,600	-
2002.....	20,780	2	13	50	115	240	603	1,466	2,857	4,983	6,999	3,452	-
2001.....	21,246	1	1	32	125	258	678	1,605	2,892	5,134	7,113	3,407	-
2000.....	21,657	1	5	45	159	254	668	1,626	2,970	5,311	7,312	3,306	-
1999.....	21,792	2	15	34	125	252	718	1,606	2,996	5,575	7,327	3,142	-
Non-Hispanic 1/, male													
2003.....	10,810	-	2	21	68	132	350	904	1,739	2,697	3,440	1,457	-
2002.....	10,953	1	6	37	73	153	389	910	1,725	2,838	3,472	1,349	-
2001.....	11,105	1	1	23	79	153	443	1,036	1,708	2,897	3,416	1,348	-
2000.....	11,225	-	3	27	109	148	431	1,036	1,728	2,954	3,533	1,256	-
1999.....	11,271	-	9	28	81	160	464	1,025	1,754	3,044	3,490	1,216	-
Non-Hispanic 1/, female													
2003.....	9,569	-	-	14	44	74	198	523	1,084	1,982	3,507	2,143	-
2002.....	9,827	1	7	13	42	87	214	556	1,132	2,145	3,527	2,103	-
2001.....	10,141	-	-	9	46	105	235	569	1,184	2,237	3,697	2,059	-
2000.....	10,432	1	2	18	50	106	237	590	1,242	2,357	3,779	2,050	-
1999.....	10,521	2	6	6	44	92	254	581	1,242	2,531	3,837	1,926	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Non-Hispanic white, both sexes													
2003.....	18,526	-	2	22	85	132	417	1,158	2,500	4,242	6,533	3,435	-
2002.....	18,901	1	12	38	80	162	463	1,198	2,517	4,523	6,597	3,310	-
2001.....	19,300	1	1	26	87	182	523	1,337	2,535	4,683	6,702	3,223	-
2000.....	19,843	-	2	38	110	181	531	1,367	2,655	4,878	6,938	3,143	-
1999.....	19,956	-	12	23	91	177	559	1,372	2,667	5,145	6,935	2,975	-
Non-Hispanic white, male													
2003.....	9,871	-	2	12	53	88	271	738	1,556	2,490	3,267	1,394	-
2002.....	9,961	-	6	29	55	106	297	763	1,526	2,600	3,287	1,292	-
2001.....	10,052	1	1	17	54	115	339	873	1,497	2,655	3,226	1,274	-
2000.....	10,264	-	2	23	79	109	353	870	1,538	2,725	3,365	1,200	-
1999.....	10,259	-	8	21	61	112	362	879	1,554	2,822	3,302	1,138	-
Non-Hispanic white, female													
2003.....	8,655	-	-	10	32	44	146	420	944	1,752	3,266	2,041	-
2002.....	8,940	1	6	9	25	56	166	435	991	1,923	3,310	2,018	-
2001.....	9,248	-	-	9	33	67	184	464	1,038	2,028	3,476	1,949	-
2000.....	9,579	-	-	15	31	72	178	497	1,117	2,153	3,573	1,943	-
1999.....	9,697	-	4	2	30	65	197	493	1,113	2,323	3,633	1,837	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Non-Hispanic black, both sexes													
2003.....	1,359	-	-	10	23	58	108	229	244	307	275	105	-
2002.....	1,446	1	1	10	24	63	112	215	277	356	284	103	-
2001.....	1,478	-	-	5	32	62	129	221	269	348	284	128	-
2000.....	1,355	1	2	5	35	55	118	206	256	318	244	115	-
1999.....	1,427	2	2	9	32	58	135	189	259	330	286	125	-
Non-Hispanic black, male													
2003.....	682	-	-	8	12	34	63	143	138	143	108	33	-
2002.....	767	1	-	7	10	40	76	119	159	193	122	40	-
2001.....	795	-	-	5	22	32	88	135	151	190	124	48	-
2000.....	719	-	-	3	19	30	68	133	157	169	103	37	-
1999.....	778	-	1	6	18	39	86	120	153	174	131	50	-
Non-Hispanic black, female													
2003.....	677	-	-	2	11	24	45	86	106	164	167	72	-
2002.....	679	-	1	3	14	23	36	96	118	163	162	63	-
2001.....	683	-	-	-	10	30	41	86	118	158	160	80	-
2000.....	636	1	2	2	16	25	50	73	99	149	141	78	-
1999.....	649	2	1	3	14	19	49	69	106	156	155	75	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)													
Origin not stated 2/, both sexes													
2003.....	39	-	-	-	1	-	-	5	8	7	11	7	-
2002.....	62	-	-	-	2	-	6	7	6	13	22	6	-
2001.....	51	-	-	-	1	3	1	5	5	16	16	4	-
2000.....	46	-	-	-	-	1	-	6	10	10	13	6	-
1999.....	52	-	-	-	-	-	3	4	5	12	22	6	-
Origin not stated 2/, male													
2003.....	20	-	-	-	1	-	-	3	5	3	5	3	-
2002.....	36	-	-	-	1	-	5	3	4	12	11	-	-
2001.....	26	-	-	-	-	2	1	3	2	6	11	1	-
2000.....	27	-	-	-	-	1	-	4	9	8	3	2	-
1999.....	30	-	-	-	-	-	3	2	2	7	14	2	-
Origin not stated 2/, female													
2003.....	19	-	-	-	-	-	-	2	3	4	6	4	-
2002.....	26	-	-	-	1	-	1	4	2	1	11	6	-
2001.....	25	-	-	-	1	1	-	2	3	10	5	3	-
2000.....	19	-	-	-	-	-	-	2	1	2	10	4	-
1999.....	22	-	-	-	-	-	-	2	3	5	8	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
All origins, both sexes													
2003.....	21,535	30	133	342	402	433	796	1,365	2,605	4,706	6,937	3,786	-
2002.....	21,498	21	104	316	472	471	767	1,459	2,611	4,818	6,763	3,696	-
2001.....	21,451	25	155	323	479	513	779	1,475	2,535	4,886	6,696	3,585	-
2000.....	21,339	38	137	315	464	473	801	1,411	2,414	5,010	6,733	3,543	-
1999.....	21,014	27	119	305	482	500	799	1,370	2,516	5,012	6,460	3,424	-
All origins, male													
2003.....	12,073	9	75	195	257	274	444	801	1,593	2,930	3,880	1,615	-
2002.....	12,016	11	57	172	282	280	429	821	1,558	3,010	3,804	1,592	-
2001.....	11,850	10	81	185	289	300	432	813	1,552	2,962	3,699	1,527	-
2000.....	11,777	21	69	190	293	283	423	791	1,452	3,090	3,695	1,470	-
1999.....	11,519	11	65	173	291	276	435	777	1,489	3,054	3,525	1,423	-
All origins, female													
2003.....	9,462	21	58	147	145	159	352	564	1,012	1,776	3,057	2,171	-
2002.....	9,482	10	47	144	190	191	338	638	1,053	1,808	2,959	2,104	-
2001.....	9,601	15	74	138	190	213	347	662	983	1,924	2,997	2,058	-
2000.....	9,562	17	68	125	171	190	378	620	962	1,920	3,038	2,073	-
1999.....	9,495	16	54	132	191	224	364	593	1,027	1,958	2,935	2,001	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Hispanic, both sexes													
2003.....	1,156	6	41	86	96	84	100	139	132	170	211	91	-
2002.....	1,088	2	26	82	109	80	121	93	143	183	164	85	-
2001.....	1,114	7	38	80	115	101	103	100	120	197	185	68	-
2000.....	1,038	8	27	72	99	71	97	110	133	182	165	74	-
1999.....	994	8	26	69	101	86	98	100	115	179	149	63	-
Hispanic, male													
2003.....	649	1	23	50	65	54	58	78	75	99	106	40	-
2002.....	601	1	12	41	73	57	69	51	67	97	94	39	-
2001.....	640	2	21	41	80	66	65	56	73	98	102	36	-
2000.....	595	5	12	46	59	52	52	64	82	98	87	38	-
1999.....	534	3	16	38	57	49	48	52	66	107	75	23	-
Hispanic, female													
2003.....	507	5	18	36	31	30	42	61	57	71	105	51	-
2002.....	487	1	14	41	36	23	52	42	76	86	70	46	-
2001.....	474	5	17	39	35	35	38	44	47	99	83	32	-
2000.....	443	3	15	26	40	19	45	46	51	84	78	36	-
1999.....	460	5	10	31	44	37	50	48	49	72	74	40	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Non-Hispanic 1/, both sexes													
2003.....	20,340	24	92	255	306	348	696	1,221	2,467	4,526	6,718	3,687	-
2002.....	20,363	18	78	234	362	390	645	1,360	2,464	4,622	6,586	3,604	-
2001.....	20,285	18	115	242	363	409	673	1,371	2,406	4,677	6,501	3,510	-
2000.....	20,248	30	109	240	362	402	697	1,299	2,274	4,817	6,555	3,463	-
1999.....	19,981	19	92	236	381	412	700	1,266	2,395	4,826	6,298	3,356	-
Non-Hispanic 1/, male													
2003.....	11,394	8	52	144	192	219	386	718	1,513	2,825	3,768	1,569	-
2002.....	11,386	9	45	131	208	223	360	765	1,489	2,904	3,702	1,550	-
2001.....	11,180	8	59	144	208	234	365	754	1,474	2,857	3,590	1,487	-
2000.....	11,155	16	56	143	233	231	367	725	1,366	2,987	3,601	1,430	-
1999.....	10,968	8	48	135	234	226	386	724	1,421	2,940	3,446	1,400	-
Non-Hispanic 1/, female													
2003.....	8,946	16	40	111	114	129	310	503	954	1,701	2,950	2,118	-
2002.....	8,977	9	33	103	154	167	285	595	975	1,718	2,884	2,054	-
2001.....	9,105	10	56	98	155	175	308	617	932	1,820	2,911	2,023	-
2000.....	9,093	14	53	97	129	171	330	574	908	1,830	2,954	2,033	-
1999.....	9,013	11	44	101	147	186	314	542	974	1,886	2,852	1,956	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Non-Hispanic white, both sexes													
2003.....	18,190	22	70	190	227	265	534	1,004	2,151	4,054	6,221	3,452	-
2002.....	18,162	13	60	174	286	289	524	1,102	2,099	4,169	6,087	3,359	-
2001.....	18,105	15	95	168	280	298	524	1,123	2,084	4,185	6,031	3,302	-
2000.....	18,128	30	88	187	262	284	546	1,059	1,952	4,372	6,121	3,227	-
1999.....	17,885	15	78	171	279	315	564	1,040	2,075	4,348	5,848	3,152	-
Non-Hispanic white, male													
2003.....	10,278	6	40	112	142	164	296	601	1,335	2,560	3,540	1,482	-
2002.....	10,218	8	36	95	167	168	294	626	1,290	2,631	3,446	1,457	-
2001.....	10,018	6	46	92	164	173	277	632	1,289	2,579	3,349	1,411	-
2000.....	10,009	16	46	108	166	164	298	593	1,183	2,726	3,377	1,332	-
1999.....	9,853	8	39	98	177	170	313	593	1,246	2,673	3,220	1,316	-
Non-Hispanic white, female													
2003.....	7,912	16	30	78	85	101	238	403	816	1,494	2,681	1,970	-
2002.....	7,944	5	24	79	119	121	230	476	809	1,538	2,641	1,902	-
2001.....	8,087	9	49	76	116	125	247	491	795	1,606	2,682	1,891	-
2000.....	8,119	14	42	79	96	120	248	466	769	1,646	2,744	1,895	-
1999.....	8,032	7	39	73	102	145	251	447	829	1,675	2,628	1,836	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Non-Hispanic black, both sexes													
2003.....	1,745	1	14	47	55	64	130	176	262	389	404	203	-
2002.....	1,752	4	13	40	53	68	89	197	295	372	405	216	-
2001.....	1,741	3	13	56	60	75	118	187	248	408	390	183	-
2000.....	1,729	-	17	36	80	88	116	204	258	373	349	208	-
1999.....	1,694	2	12	52	82	67	104	174	252	405	371	173	-
Non-Hispanic black, male													
2003.....	908	1	8	23	35	40	73	99	150	218	187	74	-
2002.....	914	1	6	24	30	37	48	109	161	220	198	80	-
2001.....	915	2	9	38	31	44	67	93	147	232	187	65	-
2000.....	923	-	8	22	54	50	52	107	149	216	181	84	-
1999.....	887	-	7	32	45	38	54	103	142	222	178	66	-
Non-Hispanic black, female													
2003.....	837	-	6	24	20	24	57	77	112	171	217	129	-
2002.....	838	3	7	16	23	31	41	88	134	152	207	136	-
2001.....	826	1	4	18	29	31	51	94	101	176	203	118	-
2000.....	806	-	9	14	26	38	64	97	109	157	168	124	-
1999.....	807	2	5	20	37	29	50	71	110	183	193	107	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Leukemia (C91-C95)													
Origin not stated 2/, both sexes													
2003.....	39	-	-	1	-	1	-	5	6	10	8	8	-
2002.....	47	1	-	-	1	1	1	6	4	13	13	7	-
2001.....	52	-	2	1	1	3	3	4	9	12	10	7	-
2000.....	53	-	1	3	3	-	7	2	7	11	13	6	-
1999.....	39	-	1	-	-	2	1	4	6	7	13	5	-
Origin not stated 2/, male													
2003.....	30	-	-	1	-	1	-	5	5	6	6	6	-
2002.....	29	1	-	-	1	-	-	5	2	9	8	3	-
2001.....	30	-	1	-	1	-	2	3	5	7	7	4	-
2000.....	27	-	1	1	1	-	4	2	4	5	7	2	-
1999.....	17	-	1	-	-	1	1	1	2	7	4	-	-
Origin not stated 2/, female													
2003.....	9	-	-	-	-	-	-	-	1	4	2	2	-
2002.....	18	-	-	-	-	1	1	1	2	4	5	4	-
2001.....	22	-	1	1	-	3	1	1	4	5	3	3	-
2000.....	26	-	-	2	2	-	3	-	3	6	6	4	-
1999.....	22	-	-	-	-	1	-	3	4	-	9	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
All origins, both sexes													
2003.....	11,267	-	-	1	-	12	147	711	1,659	3,029	3,997	1,711	-
2002.....	11,392	2	-	-	2	12	138	736	1,669	3,176	3,971	1,686	-
2001.....	11,172	-	-	-	-	9	157	671	1,671	3,181	3,886	1,597	-
2000.....	11,063	-	-	1	-	7	146	700	1,530	3,179	3,896	1,604	-
1999.....	11,016	-	-	-	-	22	156	665	1,694	3,172	3,788	1,519	-
All origins, male													
2003.....	5,848	-	-	-	-	7	97	426	995	1,654	1,941	728	-
2002.....	5,953	-	-	-	-	6	78	433	966	1,746	2,018	706	-
2001.....	5,724	-	-	-	-	4	97	403	945	1,725	1,926	624	-
2000.....	5,553	-	-	-	-	5	85	391	808	1,697	1,932	635	-
1999.....	5,584	-	-	-	-	16	95	378	968	1,669	1,846	612	-
All origins, female													
2003.....	5,419	-	-	1	-	5	50	285	664	1,375	2,056	983	-
2002.....	5,439	2	-	-	2	6	60	303	703	1,430	1,953	980	-
2001.....	5,448	-	-	-	-	5	60	268	726	1,456	1,960	973	-
2000.....	5,510	-	-	1	-	2	61	309	722	1,482	1,964	969	-
1999.....	5,432	-	-	-	-	6	61	287	726	1,503	1,942	907	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Hispanic, both sexes													
2003.....	531	-	-	-	-	-	14	53	109	154	153	48	-
2002.....	572	1	-	-	1	3	14	66	98	169	160	60	-
2001.....	524	-	-	-	-	3	18	43	107	169	137	47	-
2000.....	478	-	-	-	-	1	12	43	90	161	124	47	-
1999.....	468	-	-	-	-	4	12	51	111	139	111	40	-
Hispanic, male													
2003.....	257	-	-	-	-	-	11	31	57	72	67	19	-
2002.....	314	-	-	-	-	3	10	38	62	92	77	32	-
2001.....	291	-	-	-	-	1	13	23	52	102	80	20	-
2000.....	239	-	-	-	-	1	6	24	44	87	61	16	-
1999.....	225	-	-	-	-	4	9	28	60	67	47	10	-
Hispanic, female													
2003.....	274	-	-	-	-	-	3	22	52	82	86	29	-
2002.....	258	1	-	-	1	-	4	28	36	77	83	28	-
2001.....	233	-	-	-	-	2	5	20	55	67	57	27	-
2000.....	239	-	-	-	-	-	6	19	46	74	63	31	-
1999.....	243	-	-	-	-	-	3	23	51	72	64	30	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Non-Hispanic 1/, both sexes													
2003.....	10,710	-	-	1	-	12	133	657	1,547	2,867	3,834	1,659	-
2002.....	10,787	1	-	-	1	9	124	669	1,566	2,998	3,798	1,621	-
2001.....	10,618	-	-	-	-	6	138	625	1,559	3,006	3,736	1,548	-
2000.....	10,561	-	-	1	-	6	133	655	1,435	3,010	3,767	1,554	-
1999.....	10,532	-	-	-	-	18	144	614	1,581	3,030	3,670	1,475	-
Non-Hispanic 1/, male													
2003.....	5,579	-	-	-	-	7	86	394	936	1,581	1,868	707	-
2002.....	5,623	-	-	-	-	3	68	394	902	1,649	1,934	673	-
2001.....	5,418	-	-	-	-	3	84	378	890	1,620	1,840	603	-
2000.....	5,300	-	-	-	-	4	79	365	761	1,605	1,868	618	-
1999.....	5,352	-	-	-	-	12	86	350	908	1,601	1,796	599	-
Non-Hispanic 1/, female													
2003.....	5,131	-	-	1	-	5	47	263	611	1,286	1,966	952	-
2002.....	5,164	1	-	-	1	6	56	275	664	1,349	1,864	948	-
2001.....	5,200	-	-	-	-	3	54	247	669	1,386	1,896	945	-
2000.....	5,261	-	-	1	-	2	54	290	674	1,405	1,899	936	-
1999.....	5,180	-	-	-	-	6	58	264	673	1,429	1,874	876	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Non-Hispanic white, both sexes													
2003.....	8,689	-	-	1	-	9	89	437	1,181	2,292	3,240	1,440	-
2002.....	8,843	-	-	-	1	7	78	456	1,179	2,419	3,283	1,420	-
2001.....	8,641	-	-	-	-	2	81	447	1,144	2,427	3,194	1,346	-
2000.....	8,582	-	-	1	-	3	84	458	1,072	2,442	3,220	1,302	-
1999.....	8,547	-	-	-	-	10	92	432	1,145	2,465	3,133	1,270	-
Non-Hispanic white, male													
2003.....	4,629	-	-	-	-	6	62	268	730	1,311	1,623	629	-
2002.....	4,708	-	-	-	-	2	45	287	699	1,371	1,703	601	-
2001.....	4,472	-	-	-	-	2	54	280	666	1,334	1,603	533	-
2000.....	4,400	-	-	-	-	2	58	261	586	1,325	1,628	540	-
1999.....	4,430	-	-	-	-	6	56	264	688	1,329	1,563	524	-
Non-Hispanic white, female													
2003.....	4,060	-	-	1	-	3	27	169	451	981	1,617	811	-
2002.....	4,135	-	-	-	1	5	33	169	480	1,048	1,580	819	-
2001.....	4,169	-	-	-	-	-	27	167	478	1,093	1,591	813	-
2000.....	4,182	-	-	1	-	1	26	197	486	1,117	1,592	762	-
1999.....	4,117	-	-	-	-	4	36	168	457	1,136	1,570	746	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Non-Hispanic black, both sexes													
2003.....	1,833	-	-	-	-	2	43	200	326	523	536	203	-
2002.....	1,750	-	-	-	-	2	40	192	343	524	465	184	-
2001.....	1,795	-	-	-	-	4	53	159	377	519	493	190	-
2000.....	1,824	-	-	-	-	3	44	186	336	526	500	229	-
1999.....	1,786	-	-	-	-	7	49	171	391	494	478	196	-
Non-Hispanic black, male													
2003.....	858	-	-	-	-	1	24	114	188	239	222	70	-
2002.....	830	-	-	-	-	1	21	99	182	257	205	65	-
2001.....	853	-	-	-	-	1	28	87	203	251	220	63	-
2000.....	829	-	-	-	-	2	17	95	163	261	222	69	-
1999.....	824	-	-	-	-	6	29	80	196	237	208	68	-
Non-Hispanic black, female													
2003.....	975	-	-	-	-	1	19	86	138	284	314	133	-
2002.....	920	-	-	-	-	1	19	93	161	267	260	119	-
2001.....	942	-	-	-	-	3	25	72	174	268	273	127	-
2000.....	995	-	-	-	-	1	27	91	173	265	278	160	-
1999.....	962	-	-	-	-	1	20	91	195	257	270	128	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Origin not stated 2/, both sexes													
2003.....	26	-	-	-	-	-	-	1	3	8	10	4	-
2002.....	33	-	-	-	-	-	-	1	5	9	13	5	-
2001.....	30	-	-	-	-	-	1	3	5	6	13	2	-
2000.....	24	-	-	-	-	-	1	2	5	8	5	3	-
1999.....	16	-	-	-	-	-	-	-	2	3	7	4	-
Origin not stated 2/, male													
2003.....	12	-	-	-	-	-	-	1	2	1	6	2	-
2002.....	16	-	-	-	-	-	-	1	2	5	7	1	-
2001.....	15	-	-	-	-	-	-	2	3	3	6	1	-
2000.....	14	-	-	-	-	-	-	2	3	5	3	1	-
1999.....	7	-	-	-	-	-	-	-	-	1	3	3	-
Origin not stated 2/, female													
2003.....	14	-	-	-	-	-	-	-	1	7	4	2	-
2002.....	17	-	-	-	-	-	-	-	3	4	6	4	-
2001.....	15	-	-	-	-	-	1	1	2	3	7	1	-
2000.....	10	-	-	-	-	-	1	-	2	3	2	2	-
1999.....	9	-	-	-	-	-	-	-	2	2	4	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
All origins, both sexes													
2003.....	55	1	-	1	-	1	1	7	2	15	14	13	-
2002.....	73	3	-	-	1	-	-	7	3	18	25	16	-
2001.....	90	1	1	-	-	1	4	4	8	23	28	20	-
2000.....	100	-	2	1	1	-	4	8	12	22	29	21	-
1999.....	83	1	1	-	-	1	4	9	12	17	25	13	-
All origins, male													
2003.....	31	1	-	-	-	1	1	4	2	9	8	5	-
2002.....	44	3	-	-	1	-	-	4	2	15	12	7	-
2001.....	51	1	-	-	-	1	3	3	5	18	14	6	-
2000.....	60	-	2	1	1	-	3	6	9	12	17	9	-
1999.....	39	1	-	-	-	1	1	4	7	8	14	3	-
All origins, female													
2003.....	24	-	-	1	-	-	-	3	-	6	6	8	-
2002.....	29	-	-	-	-	-	-	3	1	3	13	9	-
2001.....	39	-	1	-	-	-	1	1	3	5	14	14	-
2000.....	40	-	-	-	-	-	1	2	3	10	12	12	-
1999.....	44	-	1	-	-	-	3	5	5	9	11	10	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

Hispanic, both sexes

2003.....	1	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	2	-	-	-	1	-	-	-	-	1	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	2	-	-	-
2000.....	6	-	-	-	-	-	-	1	1	2	1	1	-
1999.....	5	-	1	-	-	1	-	-	-	1	2	-	-

Hispanic, male

2003.....	1	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	1	-	-	-	1	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	3	-	-	-	-	-	-	1	1	-	-	1	-
1999.....	2	-	-	-	-	1	-	-	-	-	1	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	3	-	-	-	-	-	-	-	-	2	1	-	-
1999.....	3	-	1	-	-	-	-	-	-	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
Non-Hispanic 1/, both sexes													
2003.....	54	1	-	1	-	-	1	7	2	15	14	13	-
2002.....	71	3	-	-	-	-	-	7	3	17	25	16	-
2001.....	88	1	1	-	-	1	4	4	8	21	28	20	-
2000.....	94	-	2	1	1	-	4	7	11	20	28	20	-
1999.....	78	1	-	-	-	-	4	9	12	16	23	13	-
Non-Hispanic 1/, male													
2003.....	30	1	-	-	-	-	1	4	2	9	8	5	-
2002.....	43	3	-	-	-	-	-	4	2	15	12	7	-
2001.....	50	1	-	-	-	1	3	3	5	17	14	6	-
2000.....	57	-	2	1	1	-	3	5	8	12	17	8	-
1999.....	37	1	-	-	-	-	1	4	7	8	13	3	-
Non-Hispanic 1/, female													
2003.....	24	-	-	1	-	-	-	3	-	6	6	8	-
2002.....	28	-	-	-	-	-	-	3	1	2	13	9	-
2001.....	38	-	1	-	-	-	1	1	3	4	14	14	-
2000.....	37	-	-	-	-	-	1	2	3	8	11	12	-
1999.....	41	-	-	-	-	-	3	5	5	8	10	10	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
Non-Hispanic white, both sexes													
2003.....	49	1	-	1	-	-	1	5	2	14	13	12	-
2002.....	59	1	-	-	-	-	-	6	3	16	20	13	-
2001.....	75	1	1	-	-	-	3	4	5	19	23	19	-
2000.....	83	-	2	1	-	-	3	5	10	19	25	18	-
1999.....	71	1	-	-	-	-	3	7	11	16	21	12	-
Non-Hispanic white, male													
2003.....	27	1	-	-	-	-	1	2	2	9	7	5	-
2002.....	37	1	-	-	-	-	-	4	2	14	10	6	-
2001.....	43	1	-	-	-	-	2	3	4	16	12	5	-
2000.....	48	-	2	1	-	-	2	3	7	11	16	6	-
1999.....	34	1	-	-	-	-	1	3	6	8	12	3	-
Non-Hispanic white, female													
2003.....	22	-	-	1	-	-	-	3	-	5	6	7	-
2002.....	22	-	-	-	-	-	-	2	1	2	10	7	-
2001.....	32	-	1	-	-	-	1	1	1	3	11	14	-
2000.....	35	-	-	-	-	-	1	2	3	8	9	12	-
1999.....	37	-	-	-	-	-	2	4	5	8	9	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

Non-Hispanic black, both sexes

2003.....	3	-	-	-	-	-	-	1	-	1	-	1	-
2002.....	6	-	-	-	-	-	-	-	-	1	3	2	-
2001.....	12	-	-	-	-	1	1	-	3	2	4	1	-
2000.....	9	-	-	-	1	-	-	2	1	-	3	2	-
1999.....	6	-	-	-	-	-	1	2	1	-	1	1	-

Non-Hispanic black, male

2003.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	1	1	1	-
2001.....	7	-	-	-	-	1	1	-	1	1	2	1	-
2000.....	7	-	-	-	1	-	-	2	1	-	1	2	-
1999.....	3	-	-	-	-	-	-	1	1	-	1	-	-

Non-Hispanic black, female

2003.....	2	-	-	-	-	-	-	-	-	1	-	1	-
2002.....	4	-	-	-	-	-	-	-	-	-	2	2	-
2001.....	5	-	-	-	-	-	-	-	2	1	2	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	2	-	-
1999.....	3	-	-	-	-	-	1	1	-	-	-	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over Age not stated

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

Origin not stated 2/, both sexes

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing mortality data for both sexes.

Origin not stated 2/, male

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing mortality data for males.

Origin not stated 2/, female

Table with 14 columns (years 2003-1999) and 14 rows (ages) showing mortality data for females.

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)													
All origins, both sexes													
2003.....	62,639	19	106	274	516	690	1,885	5,611	9,820	14,277	18,880	10,556	5
2002.....	62,969	24	124	288	527	734	2,043	5,493	9,723	14,757	18,685	10,570	1
2001.....	62,708	17	117	279	516	728	2,055	5,506	9,441	15,023	18,700	10,326	-
2000.....	63,060	24	129	260	511	701	2,090	5,307	9,293	15,716	18,860	10,165	4
1999.....	66,251	14	116	291	519	762	2,314	5,477	9,863	16,834	19,656	10,403	2
All origins, male													
2003.....	31,845	12	66	152	329	427	1,046	3,143	5,573	7,800	9,321	3,973	3
2002.....	32,122	15	65	152	340	459	1,110	3,131	5,521	8,141	9,146	4,041	1
2001.....	31,663	7	73	153	346	442	1,108	3,164	5,326	8,177	9,032	3,835	-
2000.....	31,728	13	75	140	319	432	1,141	2,959	5,223	8,572	9,161	3,689	4
1999.....	33,449	8	64	160	329	490	1,300	3,098	5,597	9,157	9,439	3,806	1
All origins, female													
2003.....	30,794	7	40	122	187	263	839	2,468	4,247	6,477	9,559	6,583	2
2002.....	30,847	9	59	136	187	275	933	2,362	4,202	6,616	9,539	6,529	-
2001.....	31,045	10	44	126	170	286	947	2,342	4,115	6,846	9,668	6,491	-
2000.....	31,332	11	54	120	192	269	949	2,348	4,070	7,144	9,699	6,476	-
1999.....	32,802	6	52	131	190	272	1,014	2,379	4,266	7,677	10,217	6,597	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Hispanic, both sexes

2003.....	3,085	2	21	51	94	110	193	375	494	717	701	326	1
2002.....	3,005	11	17	55	107	107	176	346	502	712	676	296	-
2001.....	2,844	2	15	42	85	82	188	334	481	690	615	310	-
2000.....	2,753	3	24	47	82	100	141	290	455	690	631	290	-
1999.....	2,747	2	11	45	84	99	175	314	471	671	587	288	-

Hispanic, male

2003.....	1,621	1	15	27	60	84	103	214	274	385	332	126	-
2002.....	1,566	7	9	30	69	69	94	189	277	362	338	122	-
2001.....	1,455	-	9	22	64	56	95	181	262	339	315	112	-
2000.....	1,394	2	16	26	49	67	78	148	235	349	309	115	-
1999.....	1,399	2	3	23	62	63	90	162	256	351	278	109	-

Hispanic, female

2003.....	1,464	1	6	24	34	26	90	161	220	332	369	200	1
2002.....	1,439	4	8	25	38	38	82	157	225	350	338	174	-
2001.....	1,389	2	6	20	21	26	93	153	219	351	300	198	-
2000.....	1,359	1	8	21	33	33	63	142	220	341	322	175	-
1999.....	1,348	-	8	22	22	36	85	152	215	320	309	179	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic 1/, both sexes

2003.....	59,402	17	84	223	420	578	1,686	5,216	9,289	13,529	18,150	10,206	4
2002.....	59,797	13	106	233	419	625	1,867	5,129	9,192	14,005	17,965	10,242	1
2001.....	59,685	15	102	237	431	645	1,858	5,155	8,935	14,278	18,042	9,987	-
2000.....	60,117	21	103	212	427	596	1,939	4,997	8,813	14,980	18,172	9,853	4
1999.....	63,328	12	105	245	434	655	2,130	5,149	9,361	16,121	19,020	10,094	2

Non-Hispanic 1/, male

2003.....	30,140	11	50	125	268	341	940	2,916	5,275	7,396	8,976	3,839	3
2002.....	30,463	8	56	122	270	389	1,016	2,929	5,225	7,755	8,782	3,910	1
2001.....	30,107	7	64	131	282	385	1,006	2,971	5,046	7,800	8,701	3,714	-
2000.....	30,237	11	57	114	269	362	1,059	2,803	4,972	8,195	8,823	3,568	4
1999.....	31,956	6	61	137	266	423	1,206	2,928	5,325	8,779	9,132	3,692	1

Non-Hispanic 1/, female

2003.....	29,262	6	34	98	152	237	746	2,300	4,014	6,133	9,174	6,367	1
2002.....	29,334	5	50	111	149	236	851	2,200	3,967	6,250	9,183	6,332	-
2001.....	29,578	8	38	106	149	260	852	2,184	3,889	6,478	9,341	6,273	-
2000.....	29,880	10	46	98	158	234	880	2,194	3,841	6,785	9,349	6,285	-
1999.....	31,372	6	44	108	168	232	924	2,221	4,036	7,342	9,888	6,402	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic white, both sexes

2003.....	51,534	13	58	165	328	424	1,287	4,066	7,743	11,712	16,368	9,366	4
2002.....	51,731	11	73	171	319	465	1,433	4,032	7,635	12,071	16,159	9,361	1
2001.....	51,581	13	83	176	338	492	1,430	4,024	7,342	12,292	16,233	9,158	-
2000.....	52,223	16	75	168	328	479	1,452	3,908	7,330	13,019	16,395	9,050	3
1999.....	54,977	10	87	178	324	513	1,626	4,077	7,722	14,076	17,158	9,204	2

Non-Hispanic white, male

2003.....	26,189	9	31	94	209	261	729	2,281	4,419	6,440	8,169	3,544	3
2002.....	26,457	8	40	88	201	295	769	2,332	4,362	6,790	7,966	3,605	1
2001.....	26,007	6	50	96	220	296	792	2,326	4,139	6,789	7,882	3,411	-
2000.....	26,258	9	46	89	204	303	797	2,213	4,111	7,208	7,989	3,286	3
1999.....	27,780	5	52	104	190	338	943	2,305	4,423	7,714	8,322	3,383	1

Non-Hispanic white, female

2003.....	25,345	4	27	71	119	163	558	1,785	3,324	5,272	8,199	5,822	1
2002.....	25,274	3	33	83	118	170	664	1,700	3,273	5,281	8,193	5,756	-
2001.....	25,574	7	33	80	118	196	638	1,698	3,203	5,503	8,351	5,747	-
2000.....	25,965	7	29	79	124	176	655	1,695	3,219	5,811	8,406	5,764	-
1999.....	27,197	5	35	74	134	175	683	1,772	3,299	6,362	8,836	5,821	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic black, both sexes

2003.....	6,537	2	22	51	70	122	326	978	1,316	1,498	1,458	694	-
2002.....	6,756	2	26	52	79	125	362	924	1,329	1,612	1,507	738	-
2001.....	6,725	1	16	46	76	124	339	963	1,333	1,659	1,468	700	-
2000.....	6,617	4	22	39	83	87	427	927	1,255	1,636	1,472	664	1
1999.....	7,009	1	14	49	83	106	421	928	1,407	1,704	1,539	757	-

Non-Hispanic black, male

2003.....	3,274	2	15	27	45	60	171	542	743	786	658	225	-
2002.....	3,340	-	12	28	54	67	200	509	739	813	675	243	-
2001.....	3,392	-	13	26	51	69	166	543	759	866	650	249	-
2000.....	3,337	1	9	24	56	46	227	510	733	827	687	216	1
1999.....	3,514	-	8	22	56	60	219	539	801	895	667	247	-

Non-Hispanic black, female

2003.....	3,263	-	7	24	25	62	155	436	573	712	800	469	-
2002.....	3,416	2	14	24	25	58	162	415	590	799	832	495	-
2001.....	3,333	1	3	20	25	55	173	420	574	793	818	451	-
2000.....	3,280	3	13	15	27	41	200	417	522	809	785	448	-
1999.....	3,495	1	6	27	27	46	202	389	606	809	872	510	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Origin not stated 2/, both sexes

2003.....	152	-	1	-	2	2	6	20	37	31	29	24	-
2002.....	167	-	1	-	1	2	-	18	29	40	44	32	-
2001.....	179	-	-	-	-	1	9	17	25	55	43	29	-
2000.....	190	-	2	1	2	5	10	20	25	46	57	22	-
1999.....	176	-	-	1	1	8	9	14	31	42	49	21	-

Origin not stated 2/, male

2003.....	84	-	1	-	1	2	3	13	24	19	13	8	-
2002.....	93	-	-	-	1	1	-	13	19	24	26	9	-
2001.....	101	-	-	-	-	1	7	12	18	38	16	9	-
2000.....	97	-	2	-	1	3	4	8	16	28	29	6	-
1999.....	94	-	-	-	1	4	4	8	16	27	29	5	-

Origin not stated 2/, female

2003.....	68	-	-	-	1	-	3	7	13	12	16	16	-
2002.....	74	-	1	-	-	1	-	5	10	16	18	23	-
2001.....	78	-	-	-	-	-	2	5	7	17	27	20	-
2000.....	93	-	-	1	1	2	6	12	9	18	28	16	-
1999.....	82	-	-	1	-	4	5	6	15	15	20	16	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All origins, both sexes													
2003.....	13,563	62	51	79	96	178	339	682	1,238	2,354	4,689	3,795	-
2002.....	13,299	69	60	89	87	163	366	714	1,193	2,348	4,597	3,613	-
2001.....	13,676	51	58	105	102	151	366	719	1,253	2,574	4,654	3,643	-
2000.....	13,546	60	53	99	95	165	368	690	1,224	2,546	4,645	3,600	1
1999.....	13,227	58	63	101	98	171	342	662	1,133	2,684	4,472	3,443	-
All origins, male													
2003.....	6,687	32	31	43	54	98	195	366	685	1,334	2,379	1,470	-
2002.....	6,536	30	30	49	50	84	196	365	680	1,315	2,332	1,405	-
2001.....	6,735	26	30	52	53	77	193	391	673	1,439	2,403	1,398	-
2000.....	6,559	32	33	49	62	82	188	385	657	1,415	2,319	1,336	1
1999.....	6,520	30	33	49	54	96	197	362	656	1,522	2,218	1,303	-
All origins, female													
2003.....	6,876	30	20	36	42	80	144	316	553	1,020	2,310	2,325	-
2002.....	6,763	39	30	40	37	79	170	349	513	1,033	2,265	2,208	-
2001.....	6,941	25	28	53	49	74	173	328	580	1,135	2,251	2,245	-
2000.....	6,987	28	20	50	33	83	180	305	567	1,131	2,326	2,264	-
1999.....	6,707	28	30	52	44	75	145	300	477	1,162	2,254	2,140	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Hispanic, both sexes													
2003.....	603	15	7	13	21	32	25	42	64	106	185	93	-
2002.....	563	17	7	26	8	16	34	54	60	105	148	88	-
2001.....	543	5	13	21	8	20	39	50	67	100	128	92	-
2000.....	521	6	7	13	10	25	37	43	53	98	123	106	-
1999.....	497	8	12	17	12	25	30	39	60	109	111	74	-
Hispanic, male													
2003.....	301	6	3	7	13	21	14	22	39	54	88	34	-
2002.....	282	9	4	15	5	8	15	26	35	54	77	34	-
2001.....	276	2	7	12	6	6	24	30	35	50	67	37	-
2000.....	237	4	2	8	8	15	14	18	25	46	54	43	-
1999.....	233	5	4	8	8	14	21	26	30	47	47	23	-
Hispanic, female													
2003.....	302	9	4	6	8	11	11	20	25	52	97	59	-
2002.....	281	8	3	11	3	8	19	28	25	51	71	54	-
2001.....	267	3	6	9	2	14	15	20	32	50	61	55	-
2000.....	284	2	5	5	2	10	23	25	28	52	69	63	-
1999.....	264	3	8	9	4	11	9	13	30	62	64	51	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic 1/, both sexes													
2003.....	12,938	46	44	66	75	146	313	640	1,170	2,241	4,499	3,698	-
2002.....	12,708	52	53	63	78	147	328	659	1,131	2,241	4,436	3,520	-
2001.....	13,098	46	45	84	94	131	327	667	1,182	2,468	4,510	3,544	-
2000.....	12,990	53	46	85	85	140	330	645	1,167	2,445	4,511	3,482	1
1999.....	12,699	50	51	83	85	146	311	620	1,070	2,572	4,349	3,362	-
Non-Hispanic 1/, male													
2003.....	6,372	26	28	36	41	77	180	344	645	1,275	2,286	1,434	-
2002.....	6,237	21	26	34	44	76	179	339	645	1,260	2,245	1,368	-
2001.....	6,442	24	23	40	47	71	169	360	636	1,385	2,328	1,359	-
2000.....	6,302	28	31	41	54	67	173	365	629	1,367	2,258	1,288	1
1999.....	6,276	25	29	41	46	82	176	334	623	1,474	2,168	1,278	-
Non-Hispanic 1/, female													
2003.....	6,566	20	16	30	34	69	133	296	525	966	2,213	2,264	-
2002.....	6,471	31	27	29	34	71	149	320	486	981	2,191	2,152	-
2001.....	6,656	22	22	44	47	60	158	307	546	1,083	2,182	2,185	-
2000.....	6,688	25	15	44	31	73	157	280	538	1,078	2,253	2,194	-
1999.....	6,423	25	22	42	39	64	135	286	447	1,098	2,181	2,084	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic white, both sexes													
2003.....	11,572	33	32	43	58	113	236	508	995	1,970	4,147	3,437	-
2002.....	11,332	42	36	41	59	105	251	530	962	1,985	4,063	3,258	-
2001.....	11,701	32	36	56	66	106	252	561	1,006	2,185	4,150	3,251	-
2000.....	11,636	41	37	63	63	105	255	516	1,004	2,144	4,171	3,237	-
1999.....	11,335	34	36	66	58	99	239	494	899	2,276	4,001	3,133	-
Non-Hispanic white, male													
2003.....	5,732	20	19	24	34	57	143	281	544	1,142	2,125	1,343	-
2002.....	5,607	18	17	22	32	62	149	283	547	1,135	2,069	1,273	-
2001.....	5,801	18	18	25	31	58	135	312	547	1,250	2,152	1,255	-
2000.....	5,653	21	23	31	40	55	132	301	539	1,202	2,108	1,201	-
1999.....	5,653	17	18	32	30	53	143	276	536	1,326	2,013	1,209	-
Non-Hispanic white, female													
2003.....	5,840	13	13	19	24	56	93	227	451	828	2,022	2,094	-
2002.....	5,725	24	19	19	27	43	102	247	415	850	1,994	1,985	-
2001.....	5,900	14	18	31	35	48	117	249	459	935	1,998	1,996	-
2000.....	5,983	20	14	32	23	50	123	215	465	942	2,063	2,036	-
1999.....	5,682	17	18	34	28	46	96	218	363	950	1,988	1,924	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic black, both sexes													
2003.....	1,111	11	9	17	15	28	67	117	143	222	284	198	-
2002.....	1,134	9	14	18	15	31	67	112	144	211	300	213	-
2001.....	1,143	11	5	26	21	18	61	89	147	236	296	233	-
2000.....	1,122	9	9	18	16	33	62	112	133	245	282	202	1
1999.....	1,161	14	13	17	25	30	65	104	148	256	295	194	-
Non-Hispanic black, male													
2003.....	516	5	8	8	6	17	31	54	82	114	127	64	-
2002.....	496	2	8	9	9	8	25	46	83	98	134	74	-
2001.....	522	5	3	13	13	11	30	38	73	119	141	76	-
2000.....	514	4	8	8	9	11	32	54	74	130	118	65	1
1999.....	516	7	10	9	15	20	30	44	75	128	124	54	-
Non-Hispanic black, female													
2003.....	595	6	1	9	9	11	36	63	61	108	157	134	-
2002.....	638	7	6	9	6	23	42	66	61	113	166	139	-
2001.....	621	6	2	13	8	7	31	51	74	117	155	157	-
2000.....	608	5	1	10	7	22	30	58	59	115	164	137	-
1999.....	645	7	3	8	10	10	35	60	73	128	171	140	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Origin not stated 2/, both sexes

2003.....	22	1	-	-	-	-	1	-	4	7	5	4	-
2002.....	28	-	-	-	1	-	4	1	2	2	13	5	-
2001.....	35	-	-	-	-	-	-	2	4	6	16	7	-
2000.....	35	1	-	1	-	-	1	2	4	3	11	12	-
1999.....	31	-	-	1	1	-	1	3	3	3	12	7	-

Origin not stated 2/, male

2003.....	14	-	-	-	-	-	1	-	1	5	5	2	-
2002.....	17	-	-	-	1	-	2	-	-	1	10	3	-
2001.....	17	-	-	-	-	-	-	1	2	4	8	2	-
2000.....	20	-	-	-	-	-	1	2	3	2	7	5	-
1999.....	11	-	-	-	-	-	-	2	3	1	3	2	-

Origin not stated 2/, female

2003.....	8	1	-	-	-	-	-	-	3	2	-	2	-
2002.....	11	-	-	-	-	-	2	1	2	1	3	2	-
2001.....	18	-	-	-	-	-	-	1	2	2	8	5	-
2000.....	15	1	-	1	-	-	-	-	1	1	4	7	-
1999.....	20	-	-	1	1	-	1	1	-	2	9	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
All origins, both sexes													
2003.....	4,594	22	24	47	105	157	183	241	276	484	1,136	1,919	-
2002.....	4,614	18	21	53	111	180	196	250	264	508	1,161	1,852	-
2001.....	4,627	16	22	55	109	160	215	232	249	492	1,226	1,851	-
2000.....	4,494	10	35	52	85	153	190	227	260	515	1,171	1,796	-
1999.....	4,503	12	30	53	121	160	194	232	231	486	1,164	1,820	-
All origins, male													
2003.....	1,832	11	14	23	52	83	91	125	156	251	477	549	-
2002.....	1,781	12	9	26	67	101	96	109	131	244	454	532	-
2001.....	1,774	11	12	27	59	95	91	115	111	237	496	520	-
2000.....	1,669	6	16	29	39	70	78	108	130	236	465	492	-
1999.....	1,746	8	17	23	67	81	101	116	105	226	484	518	-
All origins, female													
2003.....	2,762	11	10	24	53	74	92	116	120	233	659	1,370	-
2002.....	2,833	6	12	27	44	79	100	141	133	264	707	1,320	-
2001.....	2,853	5	10	28	50	65	124	117	138	255	730	1,331	-
2000.....	2,825	4	19	23	46	83	112	119	130	279	706	1,304	-
1999.....	2,757	4	13	30	54	79	93	116	126	260	680	1,302	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
Hispanic, both sexes													
2003.....	191	2	5	4	5	7	9	12	18	25	48	56	-
2002.....	208	2	3	4	8	11	11	12	14	37	44	62	-
2001.....	185	-	3	5	11	12	12	11	17	25	43	46	-
2000.....	144	1	1	7	5	13	8	10	13	24	36	26	-
1999.....	164	-	6	3	6	8	10	9	10	27	35	50	-
Hispanic, male													
2003.....	84	2	2	2	2	5	3	6	11	12	19	20	-
2002.....	106	1	1	3	5	5	9	7	8	17	18	32	-
2001.....	82	-	2	5	6	8	6	4	7	12	21	11	-
2000.....	55	1	-	3	1	5	1	4	4	12	15	9	-
1999.....	79	-	4	1	3	5	6	4	3	17	17	19	-
Hispanic, female													
2003.....	107	-	3	2	3	2	6	6	7	13	29	36	-
2002.....	102	1	2	1	3	6	2	5	6	20	26	30	-
2001.....	103	-	1	-	5	4	6	7	10	13	22	35	-
2000.....	89	-	1	4	4	8	7	6	9	12	21	17	-
1999.....	85	-	2	2	3	3	4	5	7	10	18	31	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
Non-Hispanic 1/, both sexes													
2003.....	4,395	20	19	43	99	150	173	228	258	457	1,087	1,861	-
2002.....	4,391	14	17	49	103	167	182	237	250	470	1,114	1,788	-
2001.....	4,424	15	19	50	97	146	202	219	231	466	1,177	1,802	-
2000.....	4,336	8	34	44	79	140	181	215	246	489	1,132	1,768	-
1999.....	4,321	12	24	50	115	152	183	221	220	455	1,125	1,764	-
Non-Hispanic 1/, male													
2003.....	1,745	9	12	21	49	78	87	118	145	239	458	529	-
2002.....	1,671	9	8	23	62	96	86	102	123	227	435	500	-
2001.....	1,687	10	10	22	53	86	84	110	104	225	474	509	-
2000.....	1,608	4	16	26	37	65	77	104	126	222	448	483	-
1999.....	1,659	8	13	22	64	76	95	111	101	207	465	497	-
Non-Hispanic 1/, female													
2003.....	2,650	11	7	22	50	72	86	110	113	218	629	1,332	-
2002.....	2,720	5	9	26	41	71	96	135	127	243	679	1,288	-
2001.....	2,737	5	9	28	44	60	118	109	127	241	703	1,293	-
2000.....	2,728	4	18	18	42	75	104	111	120	267	684	1,285	-
1999.....	2,662	4	11	28	51	76	88	110	119	248	660	1,267	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
Non-Hispanic white, both sexes													
2003.....	3,287	7	5	17	18	32	36	80	135	348	939	1,670	-
2002.....	3,253	5	10	19	23	33	34	90	147	361	944	1,587	-
2001.....	3,350	5	8	22	22	27	41	81	150	368	1,013	1,613	-
2000.....	3,299	3	14	14	16	16	36	82	155	384	994	1,585	-
1999.....	3,222	6	4	17	28	10	37	86	112	360	995	1,567	-
Non-Hispanic white, male													
2003.....	1,236	2	3	9	9	17	19	45	74	178	405	475	-
2002.....	1,164	2	4	8	13	20	18	45	75	172	371	436	-
2001.....	1,220	4	3	11	12	12	15	37	68	175	419	464	-
2000.....	1,178	1	8	9	7	10	20	39	83	185	392	424	-
1999.....	1,162	4	2	6	11	2	18	46	45	163	416	449	-
Non-Hispanic white, female													
2003.....	2,051	5	2	8	9	15	17	35	61	170	534	1,195	-
2002.....	2,089	3	6	11	10	13	16	45	72	189	573	1,151	-
2001.....	2,130	1	5	11	10	15	26	44	82	193	594	1,149	-
2000.....	2,121	2	6	5	9	6	16	43	72	199	602	1,161	-
1999.....	2,060	2	2	11	17	8	19	40	67	197	579	1,118	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
Non-Hispanic black, both sexes													
2003.....	1,029	6	12	25	78	114	133	143	116	94	139	169	-
2002.....	1,048	5	7	26	78	127	142	141	94	100	153	175	-
2001.....	993	6	11	26	72	117	154	135	75	90	149	158	-
2000.....	961	4	19	30	61	124	139	128	83	89	121	163	-
1999.....	1,010	3	20	33	84	139	142	125	100	81	119	164	-
Non-Hispanic black, male													
2003.....	483	4	7	12	39	58	68	73	67	55	53	47	-
2002.....	472	4	4	14	49	72	66	55	46	51	58	53	-
2001.....	443	4	7	10	41	73	66	73	35	46	51	37	-
2000.....	392	3	8	17	29	55	55	64	37	31	45	48	-
1999.....	453	2	11	16	50	72	73	61	53	37	41	37	-
Non-Hispanic black, female													
2003.....	546	2	5	13	39	56	65	70	49	39	86	122	-
2002.....	576	1	3	12	29	55	76	86	48	49	95	122	-
2001.....	550	2	4	16	31	44	88	62	40	44	98	121	-
2000.....	569	1	11	13	32	69	84	64	46	58	76	115	-
1999.....	557	1	9	17	34	67	69	64	47	44	78	127	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Anemias (D50-D64)													
Origin not stated 2/, both sexes													
2003.....	8	-	-	-	1	-	1	1	-	2	1	2	-
2002.....	15	2	1	-	-	2	3	1	-	1	3	2	-
2001.....	18	1	-	-	1	2	1	2	1	1	6	3	-
2000.....	14	1	-	1	1	-	1	2	1	2	3	2	-
1999.....	18	-	-	-	-	-	1	2	1	4	4	6	-
Origin not stated 2/, male													
2003.....	3	-	-	-	1	-	1	1	-	-	-	-	-
2002.....	4	2	-	-	-	-	1	-	-	-	1	-	-
2001.....	5	1	-	-	-	1	1	1	-	-	1	-	-
2000.....	6	1	-	-	1	-	-	-	-	2	2	-	-
1999.....	8	-	-	-	-	-	-	1	1	2	2	2	-
Origin not stated 2/, female													
2003.....	5	-	-	-	-	-	-	-	-	2	1	2	-
2002.....	11	-	1	-	-	2	2	1	-	1	2	2	-
2001.....	13	-	-	-	1	1	-	1	1	1	5	3	-
2000.....	8	-	-	1	-	-	1	2	1	-	1	2	-
1999.....	10	-	-	-	-	-	1	1	-	2	2	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
All origins, both sexes													
2003.....	74,219	3	8	31	160	657	2,049	5,658	10,731	16,656	23,299	14,964	3
2002.....	73,249	1	3	34	171	642	2,164	5,496	10,022	16,709	23,282	14,724	1
2001.....	71,372	2	6	39	151	595	1,958	5,343	9,570	16,731	22,805	14,171	1
2000.....	69,301	4	6	26	162	623	1,926	4,954	9,186	16,674	22,184	13,556	-
1999.....	68,399	2	8	32	155	582	1,942	4,735	9,097	16,908	21,757	13,178	3
All origins, male													
2003.....	35,438	-	6	16	96	379	1,264	3,341	6,044	8,687	10,679	4,923	3
2002.....	34,301	-	1	22	98	385	1,330	3,225	5,683	8,417	10,295	4,844	1
2001.....	32,841	1	2	23	85	341	1,181	3,154	5,241	8,437	9,918	4,457	1
2000.....	31,602	4	2	13	94	376	1,151	2,821	4,943	8,226	9,738	4,234	-
1999.....	31,150	-	4	16	80	319	1,139	2,791	4,856	8,279	9,539	4,125	2
All origins, female													
2003.....	38,781	3	2	15	64	278	785	2,317	4,687	7,969	12,620	10,041	-
2002.....	38,948	1	2	12	73	257	834	2,271	4,339	8,292	12,987	9,880	-
2001.....	38,531	1	4	16	66	254	777	2,189	4,329	8,294	12,887	9,714	-
2000.....	37,699	-	4	13	68	247	775	2,133	4,243	8,448	12,446	9,322	-
1999.....	37,249	2	4	16	75	263	803	1,944	4,241	8,629	12,218	9,053	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Hispanic, both sexes													
2003.....	6,179	-	1	3	19	64	197	593	1,121	1,611	1,743	826	1
2002.....	5,912	-	-	1	15	56	205	513	1,029	1,591	1,712	790	-
2001.....	5,663	1	1	2	19	49	167	507	982	1,604	1,548	783	-
2000.....	5,328	1	-	4	13	64	154	439	950	1,592	1,410	701	-
1999.....	5,182	-	-	4	14	43	157	413	927	1,551	1,369	704	-
Hispanic, male													
2003.....	2,976	-	1	1	11	33	121	345	618	793	775	277	1
2002.....	2,779	-	-	1	10	34	124	305	551	752	748	254	-
2001.....	2,590	-	1	2	11	28	109	311	510	758	638	222	-
2000.....	2,507	1	-	2	7	48	95	240	520	762	609	223	-
1999.....	2,335	-	-	2	8	21	84	252	480	709	562	217	-
Hispanic, female													
2003.....	3,203	-	-	2	8	31	76	248	503	818	968	549	-
2002.....	3,133	-	-	-	5	22	81	208	478	839	964	536	-
2001.....	3,073	1	-	-	8	21	58	196	472	846	910	561	-
2000.....	2,821	-	-	2	6	16	59	199	430	830	801	478	-
1999.....	2,847	-	-	2	6	22	73	161	447	842	807	487	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Non-Hispanic 1/, both sexes													
2003.....	67,862	3	7	27	138	592	1,847	5,046	9,583	15,014	21,498	14,105	2
2002.....	67,112	1	3	33	155	584	1,946	4,961	8,958	15,070	21,499	13,901	1
2001.....	65,498	1	5	37	131	542	1,787	4,810	8,562	15,066	21,198	13,358	1
2000.....	63,769	3	6	22	149	557	1,762	4,501	8,207	15,031	20,717	12,814	-
1999.....	63,017	2	8	28	140	534	1,781	4,305	8,138	15,318	20,325	12,436	2
Non-Hispanic 1/, male													
2003.....	32,368	-	5	15	82	345	1,139	2,986	5,408	7,876	9,875	4,635	2
2002.....	31,399	-	1	21	87	349	1,196	2,906	5,112	7,639	9,511	4,576	1
2001.....	30,144	1	1	21	74	311	1,069	2,828	4,715	7,643	9,255	4,225	1
2000.....	28,994	3	2	11	87	328	1,048	2,574	4,408	7,436	9,098	3,999	-
1999.....	28,719	-	4	14	72	295	1,053	2,525	4,358	7,548	8,955	3,894	1
Non-Hispanic 1/, female													
2003.....	35,494	3	2	12	56	247	708	2,060	4,175	7,138	11,623	9,470	-
2002.....	35,713	1	2	12	68	235	750	2,055	3,846	7,431	11,988	9,325	-
2001.....	35,354	-	4	16	57	231	718	1,982	3,847	7,423	11,943	9,133	-
2000.....	34,775	-	4	11	62	229	714	1,927	3,799	7,595	11,619	8,815	-
1999.....	34,298	2	4	14	68	239	728	1,780	3,780	7,770	11,370	8,542	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Non-Hispanic white, both sexes													
2003.....	52,894	2	5	19	90	366	1,240	3,412	6,873	11,316	17,601	11,968	2
2002.....	52,463	1	3	19	95	382	1,345	3,385	6,403	11,337	17,692	11,801	-
2001.....	51,482	-	5	19	66	335	1,233	3,332	6,118	11,464	17,507	11,402	1
2000.....	50,163	2	6	13	92	357	1,260	3,078	5,923	11,475	17,040	10,917	-
1999.....	49,347	2	6	19	94	363	1,231	2,940	5,674	11,664	16,821	10,533	-
Non-Hispanic white, male													
2003.....	25,940	-	4	12	54	212	794	2,122	4,041	6,206	8,417	4,076	2
2002.....	25,273	-	1	16	51	231	849	2,054	3,815	6,041	8,151	4,064	-
2001.....	24,302	-	1	13	42	184	767	2,014	3,524	6,142	7,874	3,740	1
2000.....	23,465	2	2	7	57	220	766	1,842	3,298	5,967	7,796	3,508	-
1999.....	23,177	-	3	9	52	203	747	1,801	3,200	6,066	7,722	3,374	-
Non-Hispanic white, female													
2003.....	26,954	2	1	7	36	154	446	1,290	2,832	5,110	9,184	7,892	-
2002.....	27,190	1	2	3	44	151	496	1,331	2,588	5,296	9,541	7,737	-
2001.....	27,180	-	4	6	24	151	466	1,318	2,594	5,322	9,633	7,662	-
2000.....	26,698	-	4	6	35	137	494	1,236	2,625	5,508	9,244	7,409	-
1999.....	26,170	2	3	10	42	160	484	1,139	2,474	5,598	9,099	7,159	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Non-Hispanic black, both sexes													
2003.....	12,787	1	2	7	45	208	547	1,431	2,347	3,145	3,265	1,789	-
2002.....	12,583	-	-	14	57	194	554	1,393	2,204	3,152	3,213	1,801	1
2001.....	12,167	1	-	15	56	198	506	1,306	2,133	3,110	3,132	1,710	-
2000.....	11,918	1	-	9	55	187	458	1,270	1,995	3,083	3,193	1,667	-
1999.....	11,830	-	2	9	43	160	502	1,219	2,136	3,120	2,995	1,642	2
Non-Hispanic black, male													
2003.....	5,380	-	1	3	26	122	309	741	1,169	1,392	1,188	429	-
2002.....	5,158	-	-	5	34	112	316	744	1,104	1,320	1,135	387	1
2001.....	4,983	1	-	6	27	122	275	701	1,031	1,278	1,150	392	-
2000.....	4,720	1	-	4	28	98	252	648	952	1,248	1,100	389	-
1999.....	4,718	-	1	5	19	85	276	641	986	1,262	1,016	426	1
Non-Hispanic black, female													
2003.....	7,407	1	1	4	19	86	238	690	1,178	1,753	2,077	1,360	-
2002.....	7,425	-	-	9	23	82	238	649	1,100	1,832	2,078	1,414	-
2001.....	7,184	-	-	9	29	76	231	605	1,102	1,832	1,982	1,318	-
2000.....	7,198	-	-	5	27	89	206	622	1,043	1,835	2,093	1,278	-
1999.....	7,112	-	1	4	24	75	226	578	1,150	1,858	1,979	1,216	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diabetes mellitus (E10-E14)													
Origin not stated 2/, both sexes													
2003.....	178	-	-	1	3	1	5	19	27	31	58	33	-
2002.....	225	-	-	-	1	2	13	22	35	48	71	33	-
2001.....	211	-	-	-	1	4	4	26	26	61	59	30	-
2000.....	204	-	-	-	-	2	10	14	29	51	57	41	-
1999.....	200	-	-	-	1	5	4	17	32	39	63	38	1
Origin not stated 2/, male													
2003.....	94	-	-	-	3	1	4	10	18	18	29	11	-
2002.....	123	-	-	-	1	2	10	14	20	26	36	14	-
2001.....	107	-	-	-	-	2	3	15	16	36	25	10	-
2000.....	101	-	-	-	-	-	8	7	15	28	31	12	-
1999.....	96	-	-	-	-	3	2	14	18	22	22	14	1
Origin not stated 2/, female													
2003.....	84	-	-	1	-	-	1	9	9	13	29	22	-
2002.....	102	-	-	-	-	-	3	8	15	22	35	19	-
2001.....	104	-	-	-	1	2	1	11	10	25	34	20	-
2000.....	103	-	-	-	-	2	2	7	14	23	26	29	-
1999.....	104	-	-	-	1	2	2	3	14	17	41	24	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
All origins, both sexes													
2003.....	3,338	10	2	2	7	10	35	100	168	340	992	1,672	-
2002.....	3,779	4	1	1	7	16	60	102	168	382	1,132	1,906	-
2001.....	3,704	4	4	2	10	14	44	110	153	374	1,100	1,889	-
2000.....	4,242	14	2	5	10	18	56	99	173	460	1,285	2,120	-
1999.....	4,289	9	2	1	8	12	64	106	170	429	1,248	2,240	-
All origins, male													
2003.....	1,176	7	2	2	4	7	19	48	79	166	400	442	-
2002.....	1,321	3	1	-	4	7	31	48	93	186	434	514	-
2001.....	1,233	4	1	2	7	4	21	56	81	174	426	457	-
2000.....	1,412	10	2	2	2	9	16	46	97	221	471	536	-
1999.....	1,486	4	2	1	5	4	28	52	98	221	494	577	-
All origins, female													
2003.....	2,162	3	-	-	3	3	16	52	89	174	592	1,230	-
2002.....	2,458	1	-	1	3	9	29	54	75	196	698	1,392	-
2001.....	2,471	-	3	-	3	10	23	54	72	200	674	1,432	-
2000.....	2,830	4	-	3	8	9	40	53	76	239	814	1,584	-
1999.....	2,803	5	-	-	3	8	36	54	72	208	754	1,663	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
Hispanic, both sexes													
2003.....	126	2	2	-	-	1	1	3	7	19	36	55	-
2002.....	141	2	-	-	3	1	3	5	4	21	49	53	-
2001.....	136	-	1	1	-	-	7	3	5	12	45	62	-
2000.....	147	-	-	-	1	3	4	5	7	24	43	60	-
1999.....	138	-	-	-	-	-	4	6	7	17	31	73	-
Hispanic, male													
2003.....	54	2	2	-	-	1	-	2	6	10	13	18	-
2002.....	61	1	-	-	1	1	3	3	2	13	19	18	-
2001.....	53	-	-	1	-	-	2	1	4	2	20	23	-
2000.....	57	-	-	-	-	2	1	3	3	13	17	18	-
1999.....	54	-	-	-	-	-	2	5	2	8	13	24	-
Hispanic, female													
2003.....	72	-	-	-	-	-	1	1	1	9	23	37	-
2002.....	80	1	-	-	2	-	-	2	2	8	30	35	-
2001.....	83	-	1	-	-	-	5	2	1	10	25	39	-
2000.....	90	-	-	-	1	1	3	2	4	11	26	42	-
1999.....	84	-	-	-	-	-	2	1	5	9	18	49	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
Non-Hispanic 1/, both sexes													
2003.....	3,204	8	-	2	7	9	34	96	161	320	952	1,615	-
2002.....	3,621	2	1	1	4	15	56	97	164	358	1,077	1,846	-
2001.....	3,563	4	3	1	9	14	37	107	148	362	1,054	1,824	-
2000.....	4,080	14	2	5	9	15	52	94	166	435	1,233	2,055	-
1999.....	4,142	9	2	1	8	12	60	100	162	411	1,215	2,162	-
Non-Hispanic 1/, male													
2003.....	1,118	5	-	2	4	6	19	45	73	155	385	424	-
2002.....	1,255	2	1	-	3	6	28	45	91	173	412	494	-
2001.....	1,178	4	1	1	6	4	19	55	77	172	405	434	-
2000.....	1,348	10	2	2	2	7	15	43	94	208	448	517	-
1999.....	1,428	4	2	1	5	4	26	47	95	213	479	552	-
Non-Hispanic 1/, female													
2003.....	2,086	3	-	-	3	3	15	51	88	165	567	1,191	-
2002.....	2,366	-	-	1	1	9	28	52	73	185	665	1,352	-
2001.....	2,385	-	2	-	3	10	18	52	71	190	649	1,390	-
2000.....	2,732	4	-	3	7	8	37	51	72	227	785	1,538	-
1999.....	2,714	5	-	-	3	8	34	53	67	198	736	1,610	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
Non-Hispanic white, both sexes													
2003.....	2,699	4	-	2	5	7	28	62	122	257	811	1,401	-
2002.....	3,064	1	1	-	3	12	37	76	117	280	922	1,615	-
2001.....	3,015	3	3	-	7	10	22	70	113	291	878	1,618	-
2000.....	3,469	8	1	2	7	6	36	68	123	345	1,062	1,811	-
1999.....	3,486	5	1	-	5	9	45	70	110	307	1,018	1,916	-
Non-Hispanic white, male													
2003.....	915	3	-	2	3	5	16	29	50	121	317	369	-
2002.....	1,039	1	1	-	2	5	19	31	57	137	348	438	-
2001.....	948	3	1	-	6	3	10	27	52	131	340	375	-
2000.....	1,112	6	1	2	2	2	10	30	62	159	382	456	-
1999.....	1,142	2	1	-	3	3	18	34	61	150	392	478	-
Non-Hispanic white, female													
2003.....	1,784	1	-	-	2	2	12	33	72	136	494	1,032	-
2002.....	2,025	-	-	-	1	7	18	45	60	143	574	1,177	-
2001.....	2,067	-	2	-	1	7	12	43	61	160	538	1,243	-
2000.....	2,357	2	-	-	5	4	26	38	61	186	680	1,355	-
1999.....	2,344	3	-	-	2	6	27	36	49	157	626	1,438	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
Non-Hispanic black, both sexes													
2003.....	448	3	-	-	1	2	4	32	37	54	127	188	-
2002.....	495	-	-	1	1	3	16	20	46	70	139	199	-
2001.....	472	1	-	1	2	4	15	33	29	61	149	177	-
2000.....	537	5	1	3	2	8	16	24	41	78	147	212	-
1999.....	585	4	1	1	3	3	15	27	47	90	175	219	-
Non-Hispanic black, male													
2003.....	181	1	-	-	-	1	2	16	22	31	62	46	-
2002.....	185	-	-	-	1	1	7	14	33	31	56	42	-
2001.....	202	1	-	1	-	1	9	25	21	38	55	51	-
2000.....	214	3	1	-	-	5	5	13	30	44	58	55	-
1999.....	250	2	1	1	2	1	8	12	30	55	74	64	-
Non-Hispanic black, female													
2003.....	267	2	-	-	1	1	2	16	15	23	65	142	-
2002.....	310	-	-	1	-	2	9	6	13	39	83	157	-
2001.....	270	-	-	-	2	3	6	8	8	23	94	126	-
2000.....	323	2	-	3	2	3	11	11	11	34	89	157	-
1999.....	335	2	-	-	1	2	7	15	17	35	101	155	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nutritional deficiencies (E40-E64)													
Origin not stated 2/, both sexes													
2003.....	8	-	-	-	-	-	-	1	-	1	4	2	-
2002.....	17	-	-	-	-	-	1	-	-	3	6	7	-
2001.....	5	-	-	-	1	-	-	-	-	-	1	3	-
2000.....	15	-	-	-	-	-	-	-	-	1	9	5	-
1999.....	9	-	-	-	-	-	-	-	1	1	2	5	-
Origin not stated 2/, male													
2003.....	4	-	-	-	-	-	-	1	-	1	2	-	-
2002.....	5	-	-	-	-	-	-	-	-	-	3	2	-
2001.....	2	-	-	-	1	-	-	-	-	-	1	-	-
2000.....	7	-	-	-	-	-	-	-	-	-	6	1	-
1999.....	4	-	-	-	-	-	-	-	1	-	2	1	-
Origin not stated 2/, female													
2003.....	4	-	-	-	-	-	-	-	-	-	2	2	-
2002.....	12	-	-	-	-	-	1	-	-	3	3	5	-
2001.....	3	-	-	-	-	-	-	-	-	-	-	3	-
2000.....	8	-	-	-	-	-	-	-	-	1	3	4	-
1999.....	5	-	-	-	-	-	-	-	-	1	-	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
All origins, both sexes													
2003.....	3,153	5	1	1	6	8	33	94	160	320	952	1,573	-
2002.....	3,510	2	1	1	6	13	55	91	156	354	1,067	1,764	-
2001.....	3,454	2	-	2	9	13	39	99	146	347	1,025	1,772	-
2000.....	3,926	11	1	3	8	16	53	90	158	423	1,199	1,964	-
1999.....	3,985	7	2	1	7	11	61	99	154	407	1,159	2,077	-
All origins, male													
2003.....	1,098	4	1	1	3	6	19	47	75	155	379	408	-
2002.....	1,233	1	1	-	4	5	28	42	84	174	406	488	-
2001.....	1,148	2	-	2	6	4	19	47	75	157	400	436	-
2000.....	1,302	8	1	1	2	8	15	41	86	207	434	499	-
1999.....	1,371	4	2	1	4	4	27	48	92	207	452	530	-
All origins, female													
2003.....	2,055	1	-	-	3	2	14	47	85	165	573	1,165	-
2002.....	2,277	1	-	1	2	8	27	49	72	180	661	1,276	-
2001.....	2,306	-	-	-	3	9	20	52	71	190	625	1,336	-
2000.....	2,624	3	-	2	6	8	38	49	72	216	765	1,465	-
1999.....	2,614	3	-	-	3	7	34	51	62	200	707	1,547	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
Hispanic, both sexes													
2003.....	112	1	1	-	-	1	-	3	7	19	30	50	-
2002.....	136	1	-	-	2	1	3	4	4	21	49	51	-
2001.....	132	-	-	1	-	-	6	3	5	12	44	61	-
2000.....	137	-	-	-	1	2	4	4	7	22	39	58	-
1999.....	132	-	-	-	-	-	4	6	7	17	30	68	-
Hispanic, male													
2003.....	47	1	1	-	-	1	-	2	6	10	10	16	-
2002.....	58	-	-	-	1	1	3	2	2	13	19	17	-
2001.....	52	-	-	1	-	-	2	1	4	2	20	22	-
2000.....	52	-	-	-	-	2	1	2	3	12	16	16	-
1999.....	50	-	-	-	-	-	2	5	2	8	12	21	-
Hispanic, female													
2003.....	65	-	-	-	-	-	-	1	1	9	20	34	-
2002.....	78	1	-	-	1	-	-	2	2	8	30	34	-
2001.....	80	-	-	-	-	-	4	2	1	10	24	39	-
2000.....	85	-	-	-	1	-	3	2	4	10	23	42	-
1999.....	82	-	-	-	-	-	2	1	5	9	18	47	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
Non-Hispanic 1/, both sexes													
2003.....	3,034	4	-	1	6	7	33	90	153	300	918	1,522	-
2002.....	3,358	1	1	1	4	12	51	87	152	330	1,013	1,706	-
2001.....	3,317	2	-	1	8	13	33	96	141	335	980	1,708	-
2000.....	3,774	11	1	3	7	14	49	86	151	400	1,151	1,901	-
1999.....	3,844	7	2	1	7	11	57	93	146	389	1,127	2,004	-
Non-Hispanic 1/, male													
2003.....	1,047	3	-	1	3	5	19	44	69	144	367	392	-
2002.....	1,171	1	1	-	3	4	25	40	82	161	385	469	-
2001.....	1,094	2	-	1	5	4	17	46	71	155	379	414	-
2000.....	1,243	8	1	1	2	6	14	39	83	195	412	482	-
1999.....	1,317	4	2	1	4	4	25	43	89	199	438	508	-
Non-Hispanic 1/, female													
2003.....	1,987	1	-	-	3	2	14	46	84	156	551	1,130	-
2002.....	2,187	-	-	1	1	8	26	47	70	169	628	1,237	-
2001.....	2,223	-	-	-	3	9	16	50	70	180	601	1,294	-
2000.....	2,531	3	-	2	5	8	35	47	68	205	739	1,419	-
1999.....	2,527	3	-	-	3	7	32	50	57	190	689	1,496	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
Non-Hispanic white, both sexes													
2003.....	2,545	1	-	1	5	5	28	57	115	239	779	1,315	-
2002.....	2,836	1	1	-	3	9	34	67	108	262	864	1,487	-
2001.....	2,806	1	-	-	6	9	19	64	109	270	816	1,512	-
2000.....	3,204	7	-	1	6	6	33	62	112	312	993	1,672	-
1999.....	3,232	4	1	-	5	8	43	67	99	287	945	1,773	-
Non-Hispanic white, male													
2003.....	852	1	-	1	3	4	16	28	47	111	301	340	-
2002.....	970	1	1	-	2	3	17	27	51	129	325	414	-
2001.....	881	1	-	-	5	3	8	23	49	119	316	357	-
2000.....	1,021	5	-	1	2	2	9	27	55	147	350	423	-
1999.....	1,047	2	1	-	3	3	17	31	58	138	359	435	-
Non-Hispanic white, female													
2003.....	1,693	-	-	-	2	1	12	29	68	128	478	975	-
2002.....	1,866	-	-	-	1	6	17	40	57	133	539	1,073	-
2001.....	1,925	-	-	-	1	6	11	41	60	151	500	1,155	-
2000.....	2,183	2	-	-	4	4	24	35	57	165	643	1,249	-
1999.....	2,185	2	-	-	2	5	26	36	41	149	586	1,338	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
Non-Hispanic black, both sexes													
2003.....	435	2	-	-	1	2	4	31	36	52	125	182	-
2002.....	465	-	-	1	1	3	14	19	44	60	134	189	-
2001.....	438	1	-	1	2	4	14	28	26	56	139	167	-
2000.....	505	3	1	2	1	7	16	23	37	77	136	202	-
1999.....	543	3	1	1	2	3	14	24	42	88	161	204	-
Non-Hispanic black, male													
2003.....	175	1	-	-	-	1	2	16	21	30	60	44	-
2002.....	172	-	-	-	1	1	6	13	31	27	52	41	-
2001.....	186	1	-	1	-	1	9	20	18	34	53	49	-
2000.....	202	2	1	-	-	4	5	12	26	43	56	53	-
1999.....	234	2	1	1	1	1	8	11	27	53	66	63	-
Non-Hispanic black, female													
2003.....	260	1	-	-	1	1	2	15	15	22	65	138	-
2002.....	293	-	-	1	-	2	8	6	13	33	82	148	-
2001.....	252	-	-	-	2	3	5	8	8	22	86	118	-
2000.....	303	1	-	2	1	3	11	11	11	34	80	149	-
1999.....	309	1	-	-	1	2	6	13	15	35	95	141	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Malnutrition (E40-E46)													
Origin not stated 2/, both sexes													
2003.....	7	-	-	-	-	-	-	1	-	1	4	1	-
2002.....	16	-	-	-	-	-	1	-	-	3	5	7	-
2001.....	5	-	-	-	1	-	-	-	-	-	1	3	-
2000.....	15	-	-	-	-	-	-	-	-	1	9	5	-
1999.....	9	-	-	-	-	-	-	-	1	1	2	5	-
Origin not stated 2/, male													
2003.....	4	-	-	-	-	-	-	1	-	1	2	-	-
2002.....	4	-	-	-	-	-	-	-	-	-	2	2	-
2001.....	2	-	-	-	1	-	-	-	-	-	1	-	-
2000.....	7	-	-	-	-	-	-	-	-	-	6	1	-
1999.....	4	-	-	-	-	-	-	-	1	-	2	1	-
Origin not stated 2/, female													
2003.....	3	-	-	-	-	-	-	-	-	-	2	1	-
2002.....	12	-	-	-	-	-	1	-	-	3	3	5	-
2001.....	3	-	-	-	-	-	-	-	-	-	-	3	-
2000.....	8	-	-	-	-	-	-	-	-	1	3	4	-
1999.....	5	-	-	-	-	-	-	-	-	1	-	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other nutritional deficiencies (E50-E64)													
All origins, both sexes													
2003.....	185	5	1	1	1	2	2	6	8	20	40	99	-
2002.....	269	2	-	-	1	3	5	11	12	28	65	142	-
2001.....	250	2	4	-	1	1	5	11	7	27	75	117	-
2000.....	316	3	1	2	2	2	3	9	15	37	86	156	-
1999.....	304	2	-	-	1	1	3	7	16	22	89	163	-
All origins, male													
2003.....	78	3	1	1	1	1	-	1	4	11	21	34	-
2002.....	88	2	-	-	-	2	3	6	9	12	28	26	-
2001.....	85	2	1	-	1	-	2	9	6	17	26	21	-
2000.....	110	2	1	1	-	1	1	5	11	14	37	37	-
1999.....	115	-	-	-	1	-	1	4	6	14	42	47	-
All origins, female													
2003.....	107	2	-	-	-	1	2	5	4	9	19	65	-
2002.....	181	-	-	-	1	1	2	5	3	16	37	116	-
2001.....	165	-	3	-	-	1	3	2	1	10	49	96	-
2000.....	206	1	-	1	2	1	2	4	4	23	49	119	-
1999.....	189	2	-	-	-	1	2	3	10	8	47	116	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other nutritional deficiencies (E50-E64)													
Non-Hispanic 1/, both sexes													
2003.....	170	4	-	1	1	2	1	6	8	20	34	93	-
2002.....	263	1	-	-	-	3	5	10	12	28	64	140	-
2001.....	246	2	3	-	1	1	4	11	7	27	74	116	-
2000.....	306	3	1	2	2	1	3	8	15	35	82	154	-
1999.....	298	2	-	-	1	1	3	7	16	22	88	158	-
Non-Hispanic 1/, male													
2003.....	71	2	-	1	1	1	-	1	4	11	18	32	-
2002.....	84	1	-	-	-	2	3	5	9	12	27	25	-
2001.....	84	2	1	-	1	-	2	9	6	17	26	20	-
2000.....	105	2	1	1	-	1	1	4	11	13	36	35	-
1999.....	111	-	-	-	1	-	1	4	6	14	41	44	-
Non-Hispanic 1/, female													
2003.....	99	2	-	-	-	1	1	5	4	9	16	61	-
2002.....	179	-	-	-	-	1	2	5	3	16	37	115	-
2001.....	162	-	2	-	-	1	2	2	1	10	48	96	-
2000.....	201	1	-	1	2	-	2	4	4	22	46	119	-
1999.....	187	2	-	-	-	1	2	3	10	8	47	114	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other nutritional deficiencies (E50-E64)													
Non-Hispanic white, both sexes													
2003.....	154	3	-	1	-	2	-	5	7	18	32	86	-
2002.....	228	-	-	-	-	3	3	9	9	18	58	128	-
2001.....	209	2	3	-	1	1	3	6	4	21	62	106	-
2000.....	265	1	1	1	1	-	3	6	11	33	69	139	-
1999.....	254	1	-	-	-	1	2	3	11	20	73	143	-
Non-Hispanic white, male													
2003.....	63	2	-	1	-	1	-	1	3	10	16	29	-
2002.....	69	-	-	-	-	2	2	4	6	8	23	24	-
2001.....	67	2	1	-	1	-	2	4	3	12	24	18	-
2000.....	91	1	1	1	-	-	1	3	7	12	32	33	-
1999.....	95	-	-	-	-	-	1	3	3	12	33	43	-
Non-Hispanic white, female													
2003.....	91	1	-	-	-	1	-	4	4	8	16	57	-
2002.....	159	-	-	-	-	1	1	5	3	10	35	104	-
2001.....	142	-	2	-	-	1	1	2	1	9	38	88	-
2000.....	174	-	-	-	1	-	2	3	4	21	37	106	-
1999.....	159	1	-	-	-	1	1	-	8	8	40	100	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other nutritional deficiencies (E50-E64)													
Non-Hispanic black, both sexes													
2003.....	13	1	-	-	-	-	-	1	1	2	2	6	-
2002.....	30	-	-	-	-	-	2	1	2	10	5	10	-
2001.....	34	-	-	-	-	-	1	5	3	5	10	10	-
2000.....	32	2	-	1	1	1	-	1	4	1	11	10	-
1999.....	42	1	-	-	1	-	1	3	5	2	14	15	-
Non-Hispanic black, male													
2003.....	6	-	-	-	-	-	-	-	1	1	2	2	-
2002.....	13	-	-	-	-	-	1	1	2	4	4	1	-
2001.....	16	-	-	-	-	-	-	5	3	4	2	2	-
2000.....	12	1	-	-	-	1	-	1	4	1	2	2	-
1999.....	16	-	-	-	1	-	-	1	3	2	8	1	-
Non-Hispanic black, female													
2003.....	7	1	-	-	-	-	-	1	-	1	-	4	-
2002.....	17	-	-	-	-	-	1	-	-	6	1	9	-
2001.....	18	-	-	-	-	-	1	-	-	1	8	8	-
2000.....	20	1	-	1	1	-	-	-	-	-	9	8	-
1999.....	26	1	-	-	-	-	1	2	2	-	6	14	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
All origins, both sexes													
2003.....	730	77	24	17	37	44	67	117	108	96	90	52	1
2002.....	700	74	19	31	26	36	61	116	88	109	84	55	1
2001.....	727	78	26	24	40	41	69	105	113	94	96	41	-
2000.....	758	91	36	30	15	43	72	100	90	108	123	50	-
1999.....	850	116	45	27	41	48	76	108	90	115	122	62	-
All origins, male													
2003.....	387	38	12	10	20	28	40	73	61	45	40	19	1
2002.....	389	41	13	16	15	19	43	71	48	62	41	19	1
2001.....	394	39	16	15	23	25	42	60	61	51	47	15	-
2000.....	405	48	20	16	4	20	46	56	54	60	62	19	-
1999.....	442	61	26	18	26	24	45	68	45	60	49	20	-
All origins, female													
2003.....	343	39	12	7	17	16	27	44	47	51	50	33	-
2002.....	311	33	6	15	11	17	18	45	40	47	43	36	-
2001.....	333	39	10	9	17	16	27	45	52	43	49	26	-
2000.....	353	43	16	14	11	23	26	44	36	48	61	31	-
1999.....	408	55	19	9	15	24	31	40	45	55	73	42	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
Hispanic, both sexes													
2003.....	82	18	4	3	7	11	7	11	11	5	2	3	-
2002.....	70	8	5	-	6	5	9	8	8	11	9	1	-
2001.....	58	8	4	3	3	4	7	6	7	6	8	2	-
2000.....	52	9	4	3	3	5	8	6	5	5	4	-	-
1999.....	76	17	7	1	5	6	4	12	7	9	4	4	-
Hispanic, male													
2003.....	46	12	2	3	5	6	4	7	4	2	-	1	-
2002.....	42	6	4	-	5	3	5	5	4	6	3	1	-
2001.....	33	5	3	2	1	1	6	3	3	5	4	-	-
2000.....	32	6	2	1	2	3	5	5	3	1	4	-	-
1999.....	41	13	5	1	1	3	3	6	3	5	-	1	-
Hispanic, female													
2003.....	36	6	2	-	2	5	3	4	7	3	2	2	-
2002.....	28	2	1	-	1	2	4	3	4	5	6	-	-
2001.....	25	3	1	1	2	3	1	3	4	1	4	2	-
2000.....	20	3	2	2	1	2	3	1	2	4	-	-	-
1999.....	35	4	2	-	4	3	1	6	4	4	4	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
Non-Hispanic 1/, both sexes													
2003.....	646	59	20	14	30	33	60	105	97	91	88	49	-
2002.....	625	66	14	31	20	31	51	107	79	98	73	54	1
2001.....	666	70	22	21	37	37	62	98	106	88	88	37	-
2000.....	705	82	32	27	12	37	64	94	85	103	119	50	-
1999.....	767	97	38	25	34	42	71	96	82	106	118	58	-
Non-Hispanic 1/, male													
2003.....	339	26	10	7	15	22	36	65	57	43	40	18	-
2002.....	344	35	9	16	10	16	37	66	43	56	37	18	1
2001.....	361	34	13	13	22	24	36	57	58	46	43	15	-
2000.....	372	42	18	15	2	16	41	51	51	59	58	19	-
1999.....	396	48	21	16	23	21	41	62	41	55	49	19	-
Non-Hispanic 1/, female													
2003.....	307	33	10	7	15	11	24	40	40	48	48	31	-
2002.....	281	31	5	15	10	15	14	41	36	42	36	36	-
2001.....	305	36	9	8	15	13	26	41	48	42	45	22	-
2000.....	333	40	14	12	10	21	23	43	34	44	61	31	-
1999.....	371	49	17	9	11	21	30	34	41	51	69	39	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
Non-Hispanic white, both sexes													
2003.....	444	35	14	10	18	15	36	68	61	71	77	39	-
2002.....	427	44	10	20	12	15	30	62	49	77	58	49	1
2001.....	469	39	15	15	24	25	33	66	77	73	74	28	-
2000.....	486	47	19	18	7	20	36	65	60	74	100	40	-
1999.....	524	51	26	18	15	28	45	60	63	80	92	46	-
Non-Hispanic white, male													
2003.....	247	21	7	6	11	11	22	43	39	35	37	15	-
2002.....	227	23	6	11	5	6	20	37	27	45	32	14	1
2001.....	243	17	10	8	12	17	18	38	42	37	35	9	-
2000.....	264	27	12	9	1	7	22	38	39	47	48	14	-
1999.....	265	33	15	11	10	10	24	35	33	41	38	15	-
Non-Hispanic white, female													
2003.....	197	14	7	4	7	4	14	25	22	36	40	24	-
2002.....	200	21	4	9	7	9	10	25	22	32	26	35	-
2001.....	226	22	5	7	12	8	15	28	35	36	39	19	-
2000.....	222	20	7	9	6	13	14	27	21	27	52	26	-
1999.....	259	18	11	7	5	18	21	25	30	39	54	31	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Meningitis (G00,G03)													
Non-Hispanic black, both sexes													
2003.....	183	22	5	3	12	18	22	31	35	18	8	9	-
2002.....	175	20	4	10	6	15	19	40	23	21	13	4	-
2001.....	172	29	5	4	10	11	26	30	25	13	11	8	-
2000.....	202	34	13	9	5	17	27	26	22	27	13	9	-
1999.....	217	40	11	7	17	11	21	35	18	21	24	12	-
Non-Hispanic black, male													
2003.....	82	5	2	1	4	11	12	17	18	8	2	2	-
2002.....	105	11	3	5	5	9	15	26	12	11	5	3	-
2001.....	104	16	2	3	8	6	17	18	14	8	6	6	-
2000.....	100	15	6	6	1	9	18	11	11	11	7	5	-
1999.....	119	13	6	5	11	9	14	27	8	12	10	4	-
Non-Hispanic black, female													
2003.....	101	17	3	2	8	7	10	14	17	10	6	7	-
2002.....	70	9	1	5	1	6	4	14	11	10	8	1	-
2001.....	68	13	3	1	2	5	9	12	11	5	5	2	-
2000.....	102	19	7	3	4	8	9	15	11	16	6	4	-
1999.....	98	27	5	2	6	2	7	8	10	9	14	8	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
All origins, both sexes													
2003.....	17,997	1	-	1	-	2	11	64	352	2,330	8,723	6,513	-
2002.....	16,959	-	-	-	-	-	6	58	318	2,226	8,139	6,212	-
2001.....	16,544	-	-	-	-	2	15	45	301	2,147	8,125	5,909	-
2000.....	15,682	-	1	-	-	4	7	49	270	2,108	7,652	5,590	1
1999.....	14,593	1	-	-	2	2	5	43	242	2,020	7,110	5,168	-
All origins, male													
2003.....	10,180	1	-	-	-	-	6	33	219	1,517	5,191	3,213	-
2002.....	9,593	-	-	-	-	-	3	40	193	1,487	4,811	3,059	-
2001.....	9,405	-	-	-	-	1	11	33	214	1,394	4,837	2,915	-
2000.....	8,804	-	-	-	-	3	5	29	170	1,338	4,530	2,728	1
1999.....	8,262	-	-	-	2	2	3	30	161	1,303	4,307	2,454	-
All origins, female													
2003.....	7,817	-	-	1	-	2	5	31	133	813	3,532	3,300	-
2002.....	7,366	-	-	-	-	-	3	18	125	739	3,328	3,153	-
2001.....	7,139	-	-	-	-	1	4	12	87	753	3,288	2,994	-
2000.....	6,878	-	1	-	-	1	2	20	100	770	3,122	2,862	-
1999.....	6,331	1	-	-	-	-	2	13	81	717	2,803	2,714	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
Hispanic, both sexes													
2003.....	521	-	-	-	-	-	1	1	15	96	258	150	-
2002.....	491	-	-	-	-	-	1	6	21	102	223	138	-
2001.....	483	-	-	-	-	1	-	3	13	98	230	138	-
2000.....	425	-	-	-	-	-	-	5	15	84	191	130	-
1999.....	389	-	-	-	-	-	1	2	8	84	184	110	-
Hispanic, male													
2003.....	313	-	-	-	-	-	1	-	9	67	152	84	-
2002.....	281	-	-	-	-	-	1	3	16	71	122	68	-
2001.....	274	-	-	-	-	1	-	2	7	56	141	67	-
2000.....	240	-	-	-	-	-	-	2	9	46	112	71	-
1999.....	222	-	-	-	-	-	1	2	5	52	113	49	-
Hispanic, female													
2003.....	208	-	-	-	-	-	-	1	6	29	106	66	-
2002.....	210	-	-	-	-	-	-	3	5	31	101	70	-
2001.....	209	-	-	-	-	-	-	1	6	42	89	71	-
2000.....	185	-	-	-	-	-	-	3	6	38	79	59	-
1999.....	167	-	-	-	-	-	-	-	3	32	71	61	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
Non-Hispanic 1/, both sexes													
2003.....	17,443	-	-	1	-	2	10	63	336	2,232	8,444	6,355	-
2002.....	16,435	-	-	-	-	-	5	52	296	2,120	7,899	6,063	-
2001.....	16,019	-	-	-	-	1	15	41	288	2,046	7,873	5,755	-
2000.....	15,223	-	1	-	-	4	7	43	255	2,020	7,442	5,450	1
1999.....	14,177	1	-	-	2	2	4	41	234	1,930	6,917	5,046	-
Non-Hispanic 1/, male													
2003.....	9,846	-	-	-	-	-	5	33	209	1,448	5,025	3,126	-
2002.....	9,290	-	-	-	-	-	2	37	176	1,413	4,677	2,985	-
2001.....	9,104	-	-	-	-	-	11	30	207	1,335	4,681	2,840	-
2000.....	8,544	-	-	-	-	3	5	26	161	1,290	4,407	2,651	1
1999.....	8,029	-	-	-	2	2	2	28	156	1,248	4,189	2,402	-
Non-Hispanic 1/, female													
2003.....	7,597	-	-	1	-	2	5	30	127	784	3,419	3,229	-
2002.....	7,145	-	-	-	-	-	3	15	120	707	3,222	3,078	-
2001.....	6,915	-	-	-	-	1	4	11	81	711	3,192	2,915	-
2000.....	6,679	-	1	-	-	1	2	17	94	730	3,035	2,799	-
1999.....	6,148	1	-	-	-	-	2	13	78	682	2,728	2,644	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
Non-Hispanic white, both sexes													
2003.....	16,499	-	-	-	-	2	9	54	292	2,061	8,020	6,061	-
2002.....	15,626	-	-	-	-	-	2	48	269	1,959	7,545	5,803	-
2001.....	15,239	-	-	-	-	1	11	36	257	1,911	7,521	5,502	-
2000.....	14,497	-	-	-	-	3	4	36	236	1,895	7,106	5,216	1
1999.....	13,475	1	-	-	2	1	4	37	207	1,807	6,594	4,822	-
Non-Hispanic white, male													
2003.....	9,303	-	-	-	-	-	5	29	180	1,343	4,784	2,962	-
2002.....	8,838	-	-	-	-	-	1	33	163	1,312	4,473	2,856	-
2001.....	8,685	-	-	-	-	-	8	27	190	1,252	4,486	2,722	-
2000.....	8,116	-	-	-	-	3	3	22	149	1,214	4,191	2,533	1
1999.....	7,648	-	-	-	2	1	2	27	142	1,182	4,005	2,287	-
Non-Hispanic white, female													
2003.....	7,196	-	-	-	-	2	4	25	112	718	3,236	3,099	-
2002.....	6,788	-	-	-	-	-	1	15	106	647	3,072	2,947	-
2001.....	6,554	-	-	-	-	1	3	9	67	659	3,035	2,780	-
2000.....	6,381	-	-	-	-	-	1	14	87	681	2,915	2,683	-
1999.....	5,827	1	-	-	-	-	2	10	65	625	2,589	2,535	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
Non-Hispanic black, both sexes													
2003.....	624	-	-	1	-	-	1	7	33	120	282	180	-
2002.....	547	-	-	-	-	-	1	3	18	111	243	171	-
2001.....	547	-	-	-	-	-	4	5	20	91	238	189	-
2000.....	528	-	-	-	-	1	3	6	14	93	246	165	-
1999.....	502	-	-	-	-	-	-	4	22	95	221	160	-
Non-Hispanic black, male													
2003.....	348	-	-	-	-	-	-	3	21	73	158	93	-
2002.....	306	-	-	-	-	-	-	3	9	76	138	80	-
2001.....	291	-	-	-	-	-	3	3	13	59	127	86	-
2000.....	304	-	-	-	-	-	2	4	9	60	154	75	-
1999.....	269	-	-	-	-	-	-	1	13	58	123	74	-
Non-Hispanic black, female													
2003.....	276	-	-	1	-	-	1	4	12	47	124	87	-
2002.....	241	-	-	-	-	-	1	-	9	35	105	91	-
2001.....	256	-	-	-	-	-	1	2	7	32	111	103	-
2000.....	224	-	-	-	-	1	1	2	5	33	92	90	-
1999.....	233	-	-	-	-	-	-	3	9	37	98	86	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Parkinson's disease (G20-G21)													
Origin not stated 2/, both sexes													
2003.....	33	1	-	-	-	-	-	-	1	2	21	8	-
2002.....	33	-	-	-	-	-	-	-	1	4	17	11	-
2001.....	42	-	-	-	-	-	-	1	-	3	22	16	-
2000.....	34	-	-	-	-	-	-	1	-	4	19	10	-
1999.....	27	-	-	-	-	-	-	-	-	6	9	12	-
Origin not stated 2/, male													
2003.....	21	1	-	-	-	-	-	-	1	2	14	3	-
2002.....	22	-	-	-	-	-	-	-	1	3	12	6	-
2001.....	27	-	-	-	-	-	-	1	-	3	15	8	-
2000.....	20	-	-	-	-	-	-	1	-	2	11	6	-
1999.....	11	-	-	-	-	-	-	-	-	3	5	3	-
Origin not stated 2/, female													
2003.....	12	-	-	-	-	-	-	-	-	-	7	5	-
2002.....	11	-	-	-	-	-	-	-	-	1	5	5	-
2001.....	15	-	-	-	-	-	-	-	-	-	7	8	-
2000.....	14	-	-	-	-	-	-	-	-	2	8	4	-
1999.....	16	-	-	-	-	-	-	-	-	3	4	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
All origins, both sexes													
2003.....	63,457	-	-	-	-	-	11	78	554	3,836	21,157	37,821	-
2002.....	58,866	-	-	1	-	2	12	52	510	3,602	20,135	34,552	-
2001.....	53,852	-	-	-	-	2	3	73	529	3,422	18,542	31,281	-
2000.....	49,558	-	-	-	-	2	6	64	491	3,431	17,253	28,309	2
1999.....	44,536	-	-	-	1	-	7	60	448	3,204	15,836	24,980	-
All origins, male													
2003.....	18,335	-	-	-	-	-	4	34	243	1,684	7,606	8,764	-
2002.....	16,989	-	-	-	-	2	8	27	228	1,547	7,227	7,950	-
2001.....	15,762	-	-	-	-	2	3	38	228	1,472	6,795	7,224	-
2000.....	14,438	-	-	-	-	1	4	34	210	1,554	6,228	6,406	1
1999.....	13,391	-	-	-	1	-	3	28	222	1,437	5,904	5,796	-
All origins, female													
2003.....	45,122	-	-	-	-	-	7	44	311	2,152	13,551	29,057	-
2002.....	41,877	-	-	1	-	-	4	25	282	2,055	12,908	26,602	-
2001.....	38,090	-	-	-	-	-	-	35	301	1,950	11,747	24,057	-
2000.....	35,120	-	-	-	-	1	2	30	281	1,877	11,025	21,903	1
1999.....	31,145	-	-	-	-	-	4	32	226	1,767	9,932	19,184	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
Hispanic, both sexes													
2003.....	1,835	-	-	-	-	-	1	12	27	179	664	952	-
2002.....	1,474	-	-	-	-	-	1	-	20	127	535	791	-
2001.....	1,239	-	-	-	-	-	1	3	19	112	465	639	-
2000.....	1,127	-	-	-	-	-	-	6	19	128	401	573	-
1999.....	981	-	-	-	1	-	-	3	14	100	362	501	-
Hispanic, male													
2003.....	633	-	-	-	-	-	-	6	11	75	264	277	-
2002.....	464	-	-	-	-	-	1	-	8	55	196	204	-
2001.....	401	-	-	-	-	-	1	3	6	44	186	161	-
2000.....	383	-	-	-	-	-	-	4	9	63	166	141	-
1999.....	314	-	-	-	1	-	-	-	5	51	128	129	-
Hispanic, female													
2003.....	1,202	-	-	-	-	-	1	6	16	104	400	675	-
2002.....	1,010	-	-	-	-	-	-	-	12	72	339	587	-
2001.....	838	-	-	-	-	-	-	-	13	68	279	478	-
2000.....	744	-	-	-	-	-	-	2	10	65	235	432	-
1999.....	667	-	-	-	-	-	-	3	9	49	234	372	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
Non-Hispanic 1/, both sexes													
2003.....	61,506	-	-	-	-	-	10	66	525	3,647	20,459	36,799	-
2002.....	57,246	-	-	1	-	2	11	52	488	3,463	19,552	33,677	-
2001.....	52,516	-	-	-	-	2	2	70	509	3,305	18,051	30,577	-
2000.....	48,331	-	-	-	-	2	6	58	469	3,296	16,829	27,670	1
1999.....	43,471	-	-	-	-	-	7	57	434	3,099	15,447	24,427	-
Non-Hispanic 1/, male													
2003.....	17,666	-	-	-	-	-	4	28	231	1,603	7,332	8,468	-
2002.....	16,491	-	-	-	-	2	7	27	220	1,487	7,015	7,733	-
2001.....	15,334	-	-	-	-	2	2	35	222	1,427	6,600	7,046	-
2000.....	14,034	-	-	-	-	1	4	30	200	1,489	6,058	6,252	-
1999.....	13,052	-	-	-	-	-	3	28	217	1,384	5,768	5,652	-
Non-Hispanic 1/, female													
2003.....	43,840	-	-	-	-	-	6	38	294	2,044	13,127	28,331	-
2002.....	40,755	-	-	1	-	-	4	25	268	1,976	12,537	25,944	-
2001.....	37,182	-	-	-	-	-	-	35	287	1,878	11,451	23,531	-
2000.....	34,297	-	-	-	-	1	2	28	269	1,807	10,771	21,418	1
1999.....	30,419	-	-	-	-	-	4	29	217	1,715	9,679	18,775	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
Non-Hispanic white, both sexes													
2003.....	57,268	-	-	-	-	-	10	62	489	3,269	18,937	34,501	-
2002.....	53,486	-	-	1	-	1	6	49	448	3,149	18,227	31,605	-
2001.....	49,030	-	-	-	-	2	2	64	480	3,020	16,742	28,720	-
2000.....	45,246	-	-	-	-	2	6	50	428	3,004	15,704	26,052	-
1999.....	40,835	-	-	-	-	-	5	51	399	2,821	14,453	23,106	-
Non-Hispanic white, male													
2003.....	16,427	-	-	-	-	-	4	26	212	1,443	6,796	7,946	-
2002.....	15,399	-	-	-	-	1	4	24	203	1,353	6,544	7,270	-
2001.....	14,295	-	-	-	-	2	2	31	205	1,305	6,137	6,613	-
2000.....	13,123	-	-	-	-	1	4	25	183	1,367	5,686	5,857	-
1999.....	12,253	-	-	-	-	-	2	26	200	1,263	5,432	5,330	-
Non-Hispanic white, female													
2003.....	40,841	-	-	-	-	-	6	36	277	1,826	12,141	26,555	-
2002.....	38,087	-	-	1	-	-	2	25	245	1,796	11,683	24,335	-
2001.....	34,735	-	-	-	-	-	-	33	275	1,715	10,605	22,107	-
2000.....	32,123	-	-	-	-	1	2	25	245	1,637	10,018	20,195	-
1999.....	28,582	-	-	-	-	-	3	25	199	1,558	9,021	17,776	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
Non-Hispanic black, both sexes													
2003.....	3,693	-	-	-	-	-	-	4	29	326	1,347	1,987	-
2002.....	3,303	-	-	-	-	1	4	3	31	277	1,173	1,814	-
2001.....	3,104	-	-	-	-	-	-	5	26	253	1,176	1,644	-
2000.....	2,719	-	-	-	-	-	-	5	30	261	1,004	1,418	1
1999.....	2,325	-	-	-	-	-	1	4	29	249	883	1,159	-
Non-Hispanic black, male													
2003.....	1,079	-	-	-	-	-	-	2	17	142	481	437	-
2002.....	940	-	-	-	-	1	3	3	14	122	418	379	-
2001.....	906	-	-	-	-	-	-	3	17	110	412	364	-
2000.....	774	-	-	-	-	-	-	3	16	113	320	322	-
1999.....	689	-	-	-	-	-	-	2	15	112	290	270	-
Non-Hispanic black, female													
2003.....	2,614	-	-	-	-	-	-	2	12	184	866	1,550	-
2002.....	2,363	-	-	-	-	-	1	-	17	155	755	1,435	-
2001.....	2,198	-	-	-	-	-	-	2	9	143	764	1,280	-
2000.....	1,945	-	-	-	-	-	-	2	14	148	684	1,096	1
1999.....	1,636	-	-	-	-	-	1	2	14	137	593	889	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alzheimer's disease (G30)													
Origin not stated 2/, both sexes													
2003.....	116	-	-	-	-	-	-	-	2	10	34	70	-
2002.....	146	-	-	-	-	-	-	-	2	12	48	84	-
2001.....	97	-	-	-	-	-	-	-	1	5	26	65	-
2000.....	100	-	-	-	-	-	-	-	3	7	23	66	1
1999.....	84	-	-	-	-	-	-	-	-	5	27	52	-
Origin not stated 2/, male													
2003.....	36	-	-	-	-	-	-	-	1	6	10	19	-
2002.....	34	-	-	-	-	-	-	-	-	5	16	13	-
2001.....	27	-	-	-	-	-	-	-	-	1	9	17	-
2000.....	21	-	-	-	-	-	-	-	1	2	4	13	1
1999.....	25	-	-	-	-	-	-	-	-	2	8	15	-
Origin not stated 2/, female													
2003.....	80	-	-	-	-	-	-	-	1	4	24	51	-
2002.....	112	-	-	-	-	-	-	-	2	7	32	71	-
2001.....	70	-	-	-	-	-	-	-	1	4	17	48	-
2000.....	79	-	-	-	-	-	-	-	2	5	19	53	-
1999.....	59	-	-	-	-	-	-	-	-	3	19	37	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Major cardiovascular diseases (I00-I78)													
All origins, both sexes													
2003.....	902,443	555	234	347	1,447	4,089	16,930	46,279	79,497	137,220	279,358	336,449	38
2002.....	918,628	640	223	358	1,283	3,986	16,962	45,800	78,430	144,071	287,415	339,409	51
2001.....	922,334	600	285	365	1,280	4,038	16,686	44,450	76,224	148,409	291,995	337,930	72
2000.....	936,923	636	234	362	1,309	3,792	16,624	43,583	77,482	156,187	296,202	340,450	62
1999.....	950,314	644	234	354	1,315	3,873	17,010	42,512	77,833	163,428	302,552	340,510	49
All origins, male													
2003.....	423,810	310	116	204	893	2,609	11,302	32,260	52,804	81,512	135,868	105,903	29
2002.....	429,682	343	122	194	845	2,568	11,356	31,928	52,037	85,319	138,770	106,164	36
2001.....	428,065	315	149	184	807	2,555	11,180	30,860	50,320	87,756	140,789	103,097	53
2000.....	435,912	341	128	206	783	2,378	11,110	30,535	51,423	92,510	142,797	103,658	43
1999.....	441,827	347	127	199	799	2,442	11,471	29,745	51,525	96,904	145,567	102,663	38
All origins, female													
2003.....	478,633	245	118	143	554	1,480	5,628	14,019	26,693	55,708	143,490	230,546	9
2002.....	488,946	297	101	164	438	1,418	5,606	13,872	26,393	58,752	148,645	233,245	15
2001.....	494,269	285	136	181	473	1,483	5,506	13,590	25,904	60,653	151,206	234,833	19
2000.....	501,011	295	106	156	526	1,414	5,514	13,048	26,059	63,677	153,405	236,792	19
1999.....	508,487	297	107	155	516	1,431	5,539	12,767	26,308	66,524	156,985	237,847	11

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Major cardiovascular diseases (I00-I78)													
Hispanic, both sexes													
2003.....	37,266	95	43	46	176	470	1,211	2,747	4,431	7,269	10,783	9,989	6
2002.....	36,543	109	36	51	176	445	1,184	2,690	4,460	7,342	10,362	9,687	1
2001.....	35,565	98	44	56	162	442	1,081	2,561	3,996	7,325	10,139	9,654	7
2000.....	34,059	106	37	47	169	419	1,094	2,351	4,041	7,114	9,380	9,296	5
1999.....	33,665	115	41	47	160	408	1,057	2,386	3,973	7,150	9,291	9,031	6
Hispanic, male													
2003.....	19,066	54	26	27	123	315	864	1,964	2,928	4,224	5,236	3,300	5
2002.....	18,824	52	16	21	122	322	829	1,876	2,958	4,304	5,039	3,284	1
2001.....	18,138	53	21	29	108	308	758	1,806	2,582	4,269	4,966	3,232	6
2000.....	17,442	54	21	30	110	283	739	1,671	2,657	4,123	4,622	3,128	4
1999.....	17,304	49	21	29	103	292	755	1,700	2,560	4,212	4,519	3,058	6
Hispanic, female													
2003.....	18,200	41	17	19	53	155	347	783	1,503	3,045	5,547	6,689	1
2002.....	17,719	57	20	30	54	123	355	814	1,502	3,038	5,323	6,403	-
2001.....	17,427	45	23	27	54	134	323	755	1,414	3,056	5,173	6,422	1
2000.....	16,617	52	16	17	59	136	355	680	1,384	2,991	4,758	6,168	1
1999.....	16,361	66	20	18	57	116	302	686	1,413	2,938	4,772	5,973	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Major cardiovascular diseases (I00-I78)													
Non-Hispanic 1/, both sexes													
2003.....	862,744	456	190	300	1,265	3,602	15,647	43,350	74,723	129,500	267,904	325,786	21
2002.....	879,170	528	187	307	1,105	3,522	15,707	42,913	73,650	136,175	276,232	328,806	38
2001.....	883,949	499	239	308	1,108	3,578	15,536	41,695	71,875	140,618	281,026	327,420	47
2000.....	899,801	526	196	313	1,137	3,357	15,433	41,021	73,064	148,517	285,960	330,234	43
1999.....	913,476	528	192	306	1,152	3,453	15,869	39,935	73,473	155,643	292,354	330,537	34
Non-Hispanic 1/, male													
2003.....	403,411	255	90	176	766	2,283	10,387	30,161	49,614	77,001	130,276	102,387	15
2002.....	409,343	289	106	173	721	2,234	10,477	29,901	48,851	80,666	133,303	102,597	25
2001.....	408,437	260	128	155	692	2,231	10,366	28,916	47,469	83,178	135,395	99,611	36
2000.....	416,837	284	107	176	672	2,088	10,294	28,702	48,495	88,005	137,754	100,232	28
1999.....	422,824	298	106	169	694	2,142	10,649	27,907	48,675	92,273	140,588	99,298	25
Non-Hispanic 1/, female													
2003.....	459,333	201	100	124	499	1,319	5,260	13,189	25,109	52,499	137,628	223,399	6
2002.....	469,827	239	81	134	384	1,288	5,230	13,012	24,799	55,509	142,929	226,209	13
2001.....	475,512	239	111	153	416	1,347	5,170	12,779	24,406	57,440	145,631	227,809	11
2000.....	482,964	242	89	137	465	1,269	5,139	12,319	24,569	60,512	148,206	230,002	15
1999.....	490,652	230	86	137	458	1,311	5,220	12,028	24,798	63,370	151,766	231,239	9

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Major cardiovascular diseases (I00-I78)													
Non-Hispanic white, both sexes													
2003.....	741,886	250	107	191	780	2,207	10,600	30,708	57,448	105,736	236,254	297,588	17
2002.....	758,738	314	125	195	695	2,121	10,838	30,798	56,865	112,288	244,074	300,399	26
2001.....	764,021	289	140	204	671	2,240	10,524	29,967	55,480	116,098	248,922	299,458	28
2000.....	780,445	330	126	195	690	2,084	10,556	29,780	56,666	123,708	254,089	302,192	29
1999.....	793,945	324	121	203	730	2,184	10,807	29,134	57,094	130,111	260,395	302,824	18
Non-Hispanic white, male													
2003.....	345,832	133	43	115	476	1,446	7,382	22,301	39,261	64,254	116,254	94,156	11
2002.....	352,520	177	77	111	462	1,387	7,559	22,435	38,736	68,016	119,331	94,211	18
2001.....	352,013	160	75	108	425	1,441	7,379	21,723	37,726	70,213	121,332	91,409	22
2000.....	360,757	180	75	118	418	1,361	7,422	21,785	38,681	75,026	123,834	91,837	20
1999.....	366,310	189	70	110	430	1,400	7,606	21,200	38,851	78,853	126,770	90,818	13
Non-Hispanic white, female													
2003.....	396,054	117	64	76	304	761	3,218	8,407	18,187	41,482	120,000	203,432	6
2002.....	406,218	137	48	84	233	734	3,279	8,363	18,129	44,272	124,743	206,188	8
2001.....	412,008	129	65	96	246	799	3,145	8,244	17,754	45,885	127,590	208,049	6
2000.....	419,688	150	51	77	272	723	3,134	7,995	17,985	48,682	130,255	210,355	9
1999.....	427,635	135	51	93	300	784	3,201	7,934	18,243	51,258	133,625	212,006	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Major cardiovascular diseases (I00-I78)													
Non-Hispanic black, both sexes													
2003.....	103,027	168	70	95	428	1,231	4,489	11,349	15,174	20,382	26,363	23,276	2
2002.....	103,054	183	50	93	356	1,252	4,363	10,923	14,834	20,461	26,932	23,596	11
2001.....	103,213	184	86	85	395	1,209	4,506	10,539	14,483	21,084	27,025	23,600	17
2000.....	103,239	166	60	95	396	1,160	4,396	10,109	14,502	21,561	26,954	23,826	14
1999.....	103,629	172	60	93	388	1,131	4,610	9,721	14,487	22,242	27,129	23,581	15
Non-Hispanic black, male													
2003.....	48,237	102	41	53	262	721	2,628	6,959	9,006	10,809	11,435	6,219	2
2002.....	47,656	96	24	50	217	739	2,551	6,635	8,862	10,770	11,323	6,383	6
2001.....	47,646	93	45	39	240	706	2,625	6,386	8,489	11,055	11,536	6,419	13
2000.....	47,408	88	27	47	218	652	2,544	6,143	8,568	11,074	11,434	6,605	8
1999.....	47,982	91	32	53	240	659	2,716	5,984	8,609	11,567	11,329	6,691	11
Non-Hispanic black, female													
2003.....	54,790	66	29	42	166	510	1,861	4,390	6,168	9,573	14,928	17,057	-
2002.....	55,398	87	26	43	139	513	1,812	4,288	5,972	9,691	15,609	17,213	5
2001.....	55,567	91	41	46	155	503	1,881	4,153	5,994	10,029	15,489	17,181	4
2000.....	55,831	78	33	48	178	508	1,852	3,966	5,934	10,487	15,520	17,221	6
1999.....	55,647	81	28	40	148	472	1,894	3,737	5,878	10,675	15,800	16,890	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Major cardiovascular diseases (I00-I78)													
Origin not stated 2/, both sexes													
2003.....	2,433	4	1	1	6	17	72	182	343	451	671	674	11
2002.....	2,915	3	-	-	2	19	71	197	320	554	821	916	12
2001.....	2,820	3	2	1	10	18	69	194	353	466	830	856	18
2000.....	3,063	4	1	2	3	16	97	211	377	556	862	920	14
1999.....	3,173	1	1	1	3	12	84	191	387	635	907	942	9
Origin not stated 2/, male													
2003.....	1,333	1	-	1	4	11	51	135	262	287	356	216	9
2002.....	1,515	2	-	-	2	12	50	151	228	349	428	283	10
2001.....	1,490	2	-	-	7	16	56	138	269	309	428	254	11
2000.....	1,633	3	-	-	1	7	77	162	271	382	421	298	11
1999.....	1,699	-	-	1	2	8	67	138	290	419	460	307	7
Origin not stated 2/, female													
2003.....	1,100	3	1	-	2	6	21	47	81	164	315	458	2
2002.....	1,400	1	-	-	-	7	21	46	92	205	393	633	2
2001.....	1,330	1	2	1	3	2	13	56	84	157	402	602	7
2000.....	1,430	1	1	2	2	9	20	49	106	174	441	622	3
1999.....	1,474	1	1	-	1	4	17	53	97	216	447	635	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
All origins, both sexes													
2003.....	685,089	439	186	264	1,133	3,250	13,600	37,732	65,060	107,263	207,331	248,796	35
2002.....	696,947	500	165	255	1,022	3,165	13,688	37,570	64,234	112,547	213,581	250,173	47
2001.....	700,142	479	225	272	999	3,160	13,326	36,399	62,486	116,299	216,992	249,439	66
2000.....	710,760	493	181	271	1,031	2,958	13,181	35,480	63,399	122,405	220,060	251,242	59
1999.....	725,192	524	183	277	1,069	3,066	13,600	34,994	64,167	129,253	226,152	251,860	47
All origins, male													
2003.....	336,095	247	89	155	718	2,129	9,463	27,304	44,522	65,568	104,646	81,227	27
2002.....	340,933	267	87	141	686	2,131	9,638	27,239	43,900	68,658	107,217	80,935	34
2001.....	339,095	243	121	138	647	2,068	9,346	26,280	42,519	70,637	108,576	78,473	47
2000.....	344,807	260	107	163	638	1,922	9,286	25,927	43,285	74,589	109,693	78,896	41
1999.....	351,617	270	104	157	664	1,988	9,640	25,489	43,662	78,836	112,470	78,300	37
All origins, female													
2003.....	348,994	192	97	109	415	1,121	4,137	10,428	20,538	41,695	102,685	167,569	8
2002.....	356,014	233	78	114	336	1,034	4,050	10,331	20,334	43,889	106,364	169,238	13
2001.....	361,047	236	104	134	352	1,092	3,980	10,119	19,967	45,662	108,416	170,966	19
2000.....	365,953	233	74	108	393	1,036	3,895	9,553	20,114	47,816	110,367	172,346	18
1999.....	373,575	254	79	120	405	1,078	3,960	9,505	20,505	50,417	113,682	173,560	10

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Hispanic, both sexes													
2003.....	28,298	75	35	31	135	347	879	2,035	3,454	5,635	8,077	7,589	6
2002.....	27,887	85	24	30	128	321	852	2,020	3,492	5,719	7,779	7,436	1
2001.....	27,090	79	33	38	112	314	750	1,875	3,152	5,730	7,632	7,369	6
2000.....	25,819	86	29	38	125	280	760	1,754	3,136	5,453	7,067	7,086	5
1999.....	25,866	96	33	36	116	286	761	1,806	3,155	5,557	7,046	6,968	6
Hispanic, male													
2003.....	14,867	42	19	19	101	244	654	1,520	2,334	3,387	4,025	2,517	5
2002.....	14,798	41	11	14	94	242	625	1,467	2,393	3,441	3,898	2,571	1
2001.....	14,195	43	16	19	76	226	565	1,397	2,104	3,424	3,808	2,512	5
2000.....	13,566	44	21	26	87	199	550	1,307	2,128	3,245	3,525	2,430	4
1999.....	13,554	40	20	23	73	219	572	1,348	2,091	3,355	3,483	2,324	6
Hispanic, female													
2003.....	13,431	33	16	12	34	103	225	515	1,120	2,248	4,052	5,072	1
2002.....	13,089	44	13	16	34	79	227	553	1,099	2,278	3,881	4,865	-
2001.....	12,895	36	17	19	36	88	185	478	1,048	2,306	3,824	4,857	1
2000.....	12,253	42	8	12	38	81	210	447	1,008	2,208	3,542	4,656	1
1999.....	12,312	56	13	13	43	67	189	458	1,064	2,202	3,563	4,644	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Non-Hispanic 1/, both sexes													
2003.....	654,828	362	150	232	993	2,891	12,662	35,545	61,305	101,261	198,713	240,694	20
2002.....	666,690	413	141	225	892	2,827	12,780	35,376	60,465	106,351	205,156	242,030	34
2001.....	670,787	397	191	234	877	2,834	12,515	34,359	59,038	110,181	208,704	241,415	42
2000.....	682,470	403	151	232	904	2,664	12,341	33,551	59,943	116,476	212,328	243,436	41
1999.....	696,719	427	149	240	950	2,770	12,767	33,029	60,684	123,161	218,359	244,151	32
Non-Hispanic 1/, male													
2003.....	320,095	205	70	135	614	1,878	8,765	25,665	41,957	61,943	100,321	78,528	14
2002.....	324,848	225	76	127	590	1,878	8,972	25,636	41,305	64,912	102,972	78,132	23
2001.....	323,673	198	105	119	564	1,832	8,733	24,759	40,188	66,949	104,429	75,766	31
2000.....	329,888	213	86	137	550	1,716	8,673	24,487	40,924	71,010	105,834	76,231	27
1999.....	336,615	230	84	133	589	1,762	9,009	24,019	41,321	75,118	108,605	75,721	24
Non-Hispanic 1/, female													
2003.....	334,733	157	80	97	379	1,013	3,897	9,880	19,348	39,318	98,392	162,166	6
2002.....	341,842	188	65	98	302	949	3,808	9,740	19,160	41,439	102,184	163,898	11
2001.....	347,114	199	86	115	313	1,002	3,782	9,600	18,850	43,232	104,275	165,649	11
2000.....	352,582	190	65	95	354	948	3,668	9,064	19,019	45,466	106,494	167,205	14
1999.....	360,104	197	65	107	361	1,008	3,758	9,010	19,363	48,043	109,754	168,430	8

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Non-Hispanic white, both sexes													
2003.....	565,808	200	83	146	605	1,788	8,894	25,982	48,091	83,479	176,035	220,489	16
2002.....	577,761	249	91	146	547	1,718	9,114	26,206	47,494	88,528	181,975	221,670	23
2001.....	582,349	234	108	152	522	1,775	8,734	25,544	46,433	91,855	185,536	221,430	26
2000.....	594,465	258	98	140	531	1,653	8,743	25,153	47,292	97,748	189,366	223,454	29
1999.....	607,704	259	92	156	585	1,755	8,936	24,823	47,950	103,750	195,187	224,193	18
Non-Hispanic white, male													
2003.....	276,241	110	34	88	368	1,212	6,429	19,504	33,871	52,148	89,985	72,482	10
2002.....	281,442	139	54	85	370	1,177	6,657	19,774	33,313	55,265	92,616	71,976	16
2001.....	280,713	127	59	80	336	1,184	6,395	19,178	32,504	57,059	93,996	69,775	20
2000.....	287,210	139	59	87	326	1,115	6,446	19,101	33,179	61,024	95,576	70,138	20
1999.....	293,206	145	55	84	357	1,152	6,590	18,747	33,466	64,699	98,422	69,476	13
Non-Hispanic white, female													
2003.....	289,567	90	49	58	237	576	2,465	6,478	14,220	31,331	86,050	148,007	6
2002.....	296,319	110	37	61	177	541	2,457	6,432	14,181	33,263	89,359	149,694	7
2001.....	301,636	107	49	72	186	591	2,339	6,366	13,929	34,796	91,540	151,655	6
2000.....	307,255	119	39	53	205	538	2,297	6,052	14,113	36,724	93,790	153,316	9
1999.....	314,498	114	37	72	228	603	2,346	6,076	14,484	39,051	96,765	154,717	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Non-Hispanic black, both sexes													
2003.....	76,452	129	55	76	346	974	3,392	8,625	11,664	15,409	19,046	16,734	2
2002.....	76,694	142	40	65	299	993	3,312	8,318	11,554	15,411	19,520	17,030	10
2001.....	76,794	140	73	70	322	961	3,424	7,956	11,209	15,940	19,681	17,004	14
2000.....	76,706	118	47	75	331	929	3,259	7,606	11,257	16,413	19,600	17,059	12
1999.....	77,713	141	49	77	336	913	3,506	7,448	11,346	17,019	19,808	17,056	14
Non-Hispanic black, male													
2003.....	36,962	79	31	42	225	572	2,066	5,443	7,040	8,377	8,488	4,597	2
2002.....	36,596	76	18	33	185	612	2,031	5,214	7,028	8,247	8,428	4,718	6
2001.....	36,541	65	40	34	205	578	2,071	4,955	6,713	8,491	8,621	4,758	10
2000.....	36,282	59	24	41	192	543	1,972	4,787	6,786	8,560	8,476	4,835	7
1999.....	37,077	71	27	45	212	546	2,165	4,716	6,907	9,016	8,416	4,945	11
Non-Hispanic black, female													
2003.....	39,490	50	24	34	121	402	1,326	3,182	4,624	7,032	10,558	12,137	-
2002.....	40,098	66	22	32	114	381	1,281	3,104	4,526	7,164	11,092	12,312	4
2001.....	40,253	75	33	36	117	383	1,353	3,001	4,496	7,449	11,060	12,246	4
2000.....	40,424	59	23	34	139	386	1,287	2,819	4,471	7,853	11,124	12,224	5
1999.....	40,636	70	22	32	124	367	1,341	2,732	4,439	8,003	11,392	12,111	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Origin not stated 2/, both sexes													
2003.....	1,963	2	1	1	5	12	59	152	301	367	541	513	9
2002.....	2,370	2	-	-	2	17	56	174	277	477	646	707	12
2001.....	2,265	3	1	-	10	12	61	165	296	388	656	655	18
2000.....	2,471	4	1	1	2	14	80	175	320	476	665	720	13
1999.....	2,607	1	1	1	3	10	72	159	328	535	747	741	9
Origin not stated 2/, male													
2003.....	1,133	-	-	1	3	7	44	119	231	238	300	182	8
2002.....	1,287	1	-	-	2	11	41	136	202	305	347	232	10
2001.....	1,227	2	-	-	7	10	48	124	227	264	339	195	11
2000.....	1,353	3	-	-	1	7	63	133	233	334	334	235	10
1999.....	1,448	-	-	1	2	7	59	122	250	363	382	255	7
Origin not stated 2/, female													
2003.....	830	2	1	-	2	5	15	33	70	129	241	331	1
2002.....	1,083	1	-	-	-	6	15	38	75	172	299	475	2
2001.....	1,038	1	1	-	3	2	13	41	69	124	317	460	7
2000.....	1,118	1	1	1	1	7	17	42	87	142	331	485	3
1999.....	1,159	1	1	-	1	3	13	37	78	172	365	486	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
All origins, both sexes													
2003.....	3,624	1	1	6	11	25	99	224	367	693	1,233	964	-
2002.....	3,579	4	1	6	18	36	95	237	433	703	1,227	819	-
2001.....	3,489	3	1	5	11	34	130	232	361	744	1,188	780	-
2000.....	3,582	6	2	4	17	33	141	252	420	819	1,176	712	-
1999.....	3,676	6	1	5	13	42	124	245	411	854	1,227	748	-
All origins, male													
2003.....	1,153	1	-	4	7	14	44	78	146	241	384	234	-
2002.....	1,078	3	1	2	8	18	45	95	156	245	341	164	-
2001.....	1,031	3	1	2	7	15	54	87	135	231	330	166	-
2000.....	1,072	5	2	3	11	12	61	104	166	237	321	150	-
1999.....	1,042	2	1	2	10	17	49	90	133	290	325	123	-
All origins, female													
2003.....	2,471	-	1	2	4	11	55	146	221	452	849	730	-
2002.....	2,501	1	-	4	10	18	50	142	277	458	886	655	-
2001.....	2,458	-	-	3	4	19	76	145	226	513	858	614	-
2000.....	2,510	1	-	1	6	21	80	148	254	582	855	562	-
1999.....	2,634	4	-	3	3	25	75	155	278	564	902	625	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Hispanic, both sexes													
2003.....	159	1	-	1	2	7	7	21	21	40	43	16	-
2002.....	198	-	-	-	2	11	11	24	35	54	42	19	-
2001.....	156	-	1	1	2	2	12	16	27	48	30	17	-
2000.....	154	1	-	-	1	6	13	19	30	47	27	10	-
1999.....	166	2	-	-	3	4	16	23	26	44	35	13	-
Hispanic, male													
2003.....	57	1	-	1	2	4	4	8	8	13	11	5	-
2002.....	55	-	-	-	1	5	4	9	13	15	4	4	-
2001.....	51	-	1	1	1	2	5	5	7	15	11	3	-
2000.....	49	1	-	-	-	1	4	9	12	13	6	3	-
1999.....	38	-	-	-	3	3	3	6	8	10	5	-	-
Hispanic, female													
2003.....	102	-	-	-	-	3	3	13	13	27	32	11	-
2002.....	143	-	-	-	1	6	7	15	22	39	38	15	-
2001.....	105	-	-	-	1	-	7	11	20	33	19	14	-
2000.....	105	-	-	-	1	5	9	10	18	34	21	7	-
1999.....	128	2	-	-	-	1	13	17	18	34	30	13	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Non-Hispanic 1/, both sexes													
2003.....	3,460	-	1	5	9	18	92	203	344	653	1,190	945	-
2002.....	3,375	4	1	6	16	25	84	212	397	647	1,183	800	-
2001.....	3,327	3	-	4	8	32	118	214	333	696	1,157	762	-
2000.....	3,424	5	2	4	16	27	128	233	388	772	1,149	700	-
1999.....	3,498	4	1	5	10	38	106	221	383	806	1,190	734	-
Non-Hispanic 1/, male													
2003.....	1,095	-	-	3	5	10	40	70	137	228	373	229	-
2002.....	1,021	3	1	2	7	13	41	85	142	230	337	160	-
2001.....	978	3	-	1	5	13	49	81	128	216	319	163	-
2000.....	1,022	4	2	3	11	11	57	95	153	224	315	147	-
1999.....	999	2	1	2	7	14	45	83	123	279	320	123	-
Non-Hispanic 1/, female													
2003.....	2,365	-	1	2	4	8	52	133	207	425	817	716	-
2002.....	2,354	1	-	4	9	12	43	127	255	417	846	640	-
2001.....	2,349	-	-	3	3	19	69	133	205	480	838	599	-
2000.....	2,402	1	-	1	5	16	71	138	235	548	834	553	-
1999.....	2,499	2	-	3	3	24	61	138	260	527	870	611	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Non-Hispanic white, both sexes													
2003.....	3,066	-	1	3	3	12	55	128	254	571	1,129	910	-
2002.....	3,007	4	1	3	8	14	57	156	313	575	1,109	767	-
2001.....	2,924	1	-	2	7	13	73	139	265	601	1,094	729	-
2000.....	3,073	4	2	4	9	19	83	178	320	687	1,102	665	-
1999.....	3,130	2	1	3	5	21	68	163	314	730	1,125	698	-
Non-Hispanic white, male													
2003.....	967	-	-	2	2	6	23	48	98	203	364	221	-
2002.....	897	3	1	2	3	7	26	63	113	205	319	155	-
2001.....	857	1	-	-	5	7	35	58	108	187	302	154	-
2000.....	917	3	2	3	6	9	42	78	136	197	303	138	-
1999.....	892	1	1	-	2	9	31	65	102	260	301	120	-
Non-Hispanic white, female													
2003.....	2,099	-	1	1	1	6	32	80	156	368	765	689	-
2002.....	2,110	1	-	1	5	7	31	93	200	370	790	612	-
2001.....	2,067	-	-	2	2	6	38	81	157	414	792	575	-
2000.....	2,156	1	-	1	3	10	41	100	184	490	799	527	-
1999.....	2,238	1	-	3	3	12	37	98	212	470	824	578	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Non-Hispanic black, both sexes													
2003.....	305	-	-	2	4	5	29	63	72	66	40	24	-
2002.....	249	-	-	2	5	9	20	44	62	48	42	17	-
2001.....	302	2	-	1	1	13	41	60	53	69	40	22	-
2000.....	263	1	-	-	7	8	35	43	48	58	36	27	-
1999.....	269	1	-	2	4	13	30	49	51	54	39	26	-
Non-Hispanic black, male													
2003.....	109	-	-	1	2	3	15	20	33	21	8	6	-
2002.....	91	-	-	-	3	4	12	17	21	20	11	3	-
2001.....	86	2	-	-	-	3	13	20	17	17	8	6	-
2000.....	77	1	-	-	5	2	12	13	12	16	8	8	-
1999.....	86	1	-	2	4	3	12	18	16	15	14	1	-
Non-Hispanic black, female													
2003.....	196	-	-	1	2	2	14	43	39	45	32	18	-
2002.....	158	-	-	2	2	5	8	27	41	28	31	14	-
2001.....	216	-	-	1	1	10	28	40	36	52	32	16	-
2000.....	186	-	-	-	2	6	23	30	36	42	28	19	-
1999.....	183	-	-	-	-	10	18	31	35	39	25	25	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Origin not stated 2/, both sexes													
2003.....	5	-	-	-	-	-	-	-	2	-	-	3	-
2002.....	6	-	-	-	-	-	-	1	1	2	2	-	-
2001.....	6	-	-	-	1	-	-	2	1	-	1	1	-
2000.....	4	-	-	-	-	-	-	-	2	-	-	2	-
1999.....	12	-	-	-	-	-	2	1	2	4	2	1	-
Origin not stated 2/, male													
2003.....	1	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	2	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	2	-	-	-	1	-	-	1	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	5	-	-	-	-	-	1	1	2	1	-	-	-
Origin not stated 2/, female													
2003.....	4	-	-	-	-	-	-	-	1	-	-	3	-
2002.....	4	-	-	-	-	-	-	-	-	2	2	-	-
2001.....	4	-	-	-	-	-	-	1	1	-	1	1	-
2000.....	3	-	-	-	-	-	-	-	1	-	-	2	-
1999.....	7	-	-	-	-	-	1	-	-	3	2	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
All origins, both sexes													
2003.....	28,345	-	1	-	53	305	1,306	3,230	3,651	4,045	6,565	9,189	-
2002.....	26,551	-	-	2	30	297	1,213	2,882	3,320	3,895	6,321	8,587	4
2001.....	24,689	-	-	1	24	246	1,091	2,464	2,769	3,619	6,119	8,345	11
2000.....	23,761	-	1	2	32	213	981	2,104	2,672	3,605	5,962	8,186	3
1999.....	22,702	1	-	-	33	210	964	1,930	2,537	3,517	5,830	7,679	1
All origins, male													
2003.....	12,536	-	1	-	40	208	877	2,157	2,308	2,264	2,627	2,054	-
2002.....	11,660	-	-	1	26	215	849	1,938	2,159	2,088	2,509	1,871	4
2001.....	10,451	-	-	-	17	169	740	1,646	1,749	1,961	2,414	1,746	9
2000.....	9,876	-	-	1	24	139	643	1,436	1,684	1,971	2,269	1,707	2
1999.....	9,316	-	-	-	25	149	660	1,259	1,560	1,816	2,252	1,594	1
All origins, female													
2003.....	15,809	-	-	-	13	97	429	1,073	1,343	1,781	3,938	7,135	-
2002.....	14,891	-	-	1	4	82	364	944	1,161	1,807	3,812	6,716	-
2001.....	14,238	-	-	1	7	77	351	818	1,020	1,658	3,705	6,599	2
2000.....	13,885	-	1	1	8	74	338	668	988	1,634	3,693	6,479	1
1999.....	13,386	1	-	-	8	61	304	671	977	1,701	3,578	6,085	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Hispanic, both sexes													
2003.....	1,586	-	-	-	12	27	109	218	247	280	347	346	-
2002.....	1,424	-	-	-	4	31	97	180	241	264	308	299	-
2001.....	1,211	-	-	-	5	22	93	165	184	239	260	243	-
2000.....	991	-	-	-	1	20	64	118	141	198	233	216	-
1999.....	933	-	-	-	3	18	61	99	146	174	215	217	-
Hispanic, male													
2003.....	876	-	-	-	10	22	91	158	168	167	166	94	-
2002.....	781	-	-	-	4	25	78	135	159	149	149	82	-
2001.....	648	-	-	-	4	15	75	118	120	132	108	76	-
2000.....	495	-	-	-	1	16	43	84	86	106	103	56	-
1999.....	455	-	-	-	2	16	45	69	96	95	82	50	-
Hispanic, female													
2003.....	710	-	-	-	2	5	18	60	79	113	181	252	-
2002.....	643	-	-	-	-	6	19	45	82	115	159	217	-
2001.....	563	-	-	-	1	7	18	47	64	107	152	167	-
2000.....	496	-	-	-	-	4	21	34	55	92	130	160	-
1999.....	478	-	-	-	1	2	16	30	50	79	133	167	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Non-Hispanic 1/, both sexes													
2003.....	26,632	-	1	-	41	277	1,189	2,990	3,378	3,741	6,199	8,816	-
2002.....	24,976	-	-	2	26	265	1,109	2,679	3,060	3,591	5,985	8,257	2
2001.....	23,343	-	-	1	18	222	991	2,270	2,554	3,364	5,837	8,077	9
2000.....	22,646	-	1	2	31	193	905	1,968	2,518	3,374	5,706	7,946	2
1999.....	21,639	1	-	-	30	192	898	1,815	2,363	3,315	5,591	7,433	1
Non-Hispanic 1/, male													
2003.....	11,588	-	1	-	30	186	780	1,981	2,121	2,082	2,452	1,955	-
2002.....	10,790	-	-	1	22	190	765	1,784	1,988	1,916	2,344	1,778	2
2001.....	9,725	-	-	-	12	152	660	1,509	1,608	1,815	2,299	1,663	7
2000.....	9,298	-	-	1	23	123	591	1,338	1,586	1,839	2,152	1,644	1
1999.....	8,790	-	-	-	23	133	610	1,180	1,445	1,702	2,157	1,539	1
Non-Hispanic 1/, female													
2003.....	15,044	-	-	-	11	91	409	1,009	1,257	1,659	3,747	6,861	-
2002.....	14,186	-	-	1	4	75	344	895	1,072	1,675	3,641	6,479	-
2001.....	13,618	-	-	1	6	70	331	761	946	1,549	3,538	6,414	2
2000.....	13,348	-	1	1	8	70	314	630	932	1,535	3,554	6,302	1
1999.....	12,849	1	-	-	7	59	288	635	918	1,613	3,434	5,894	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Non-Hispanic white, both sexes													
2003.....	19,286	-	-	-	15	133	623	1,606	2,087	2,465	4,823	7,534	-
2002.....	18,060	-	-	1	10	110	587	1,519	1,832	2,380	4,574	7,046	1
2001.....	16,991	-	-	-	5	102	493	1,213	1,520	2,198	4,540	6,915	5
2000.....	16,373	-	-	2	14	75	476	1,042	1,441	2,193	4,386	6,744	-
1999.....	15,654	1	-	-	9	91	428	954	1,371	2,147	4,294	6,359	-
Non-Hispanic white, male													
2003.....	8,056	-	-	-	11	101	463	1,172	1,383	1,389	1,908	1,629	-
2002.....	7,427	-	-	1	9	83	436	1,088	1,239	1,294	1,793	1,483	1
2001.....	6,728	-	-	-	3	80	367	866	1,016	1,237	1,781	1,374	4
2000.....	6,390	-	-	1	11	53	341	771	956	1,242	1,640	1,375	-
1999.....	6,013	-	-	-	7	75	323	676	882	1,122	1,647	1,281	-
Non-Hispanic white, female													
2003.....	11,230	-	-	-	4	32	160	434	704	1,076	2,915	5,905	-
2002.....	10,633	-	-	-	1	27	151	431	593	1,086	2,781	5,563	-
2001.....	10,263	-	-	-	2	22	126	347	504	961	2,759	5,541	1
2000.....	9,983	-	-	1	3	22	135	271	485	951	2,746	5,369	-
1999.....	9,641	1	-	-	2	16	105	278	489	1,025	2,647	5,078	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Non-Hispanic black, both sexes													
2003.....	6,759	-	1	-	25	134	539	1,321	1,210	1,170	1,233	1,126	-
2002.....	6,385	-	-	1	15	149	491	1,115	1,158	1,124	1,267	1,065	-
2001.....	5,880	-	-	1	13	114	474	1,007	972	1,078	1,164	1,053	4
2000.....	5,828	-	1	-	17	114	414	886	1,025	1,093	1,194	1,082	2
1999.....	5,556	-	-	-	20	98	447	838	932	1,084	1,182	954	1
Non-Hispanic black, male													
2003.....	3,248	-	1	-	19	78	299	762	687	646	482	274	-
2002.....	3,089	-	-	-	12	101	307	662	707	574	484	242	-
2001.....	2,760	-	-	-	9	67	275	609	551	538	458	250	3
2000.....	2,685	-	-	-	12	66	240	538	594	547	452	235	1
1999.....	2,571	-	-	-	15	55	267	487	528	536	458	224	1
Non-Hispanic black, female													
2003.....	3,511	-	-	-	6	56	240	559	523	524	751	852	-
2002.....	3,296	-	-	1	3	48	184	453	451	550	783	823	-
2001.....	3,120	-	-	1	4	47	199	398	421	540	706	803	1
2000.....	3,143	-	1	-	5	48	174	348	431	546	742	847	1
1999.....	2,985	-	-	-	5	43	180	351	404	548	724	730	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart disease (I11)													
Origin not stated 2/, both sexes													
2003.....	127	-	-	-	-	1	8	22	26	24	19	27	-
2002.....	151	-	-	-	-	1	7	23	19	40	28	31	2
2001.....	135	-	-	-	1	2	7	29	31	16	22	25	2
2000.....	124	-	-	-	-	-	12	18	13	33	23	24	1
1999.....	130	-	-	-	-	-	5	16	28	28	24	29	-
Origin not stated 2/, male													
2003.....	72	-	-	-	-	-	6	18	19	15	9	5	-
2002.....	89	-	-	-	-	-	6	19	12	23	16	11	2
2001.....	78	-	-	-	1	2	5	19	21	14	7	7	2
2000.....	83	-	-	-	-	-	9	14	12	26	14	7	1
1999.....	71	-	-	-	-	-	5	10	19	19	13	5	-
Origin not stated 2/, female													
2003.....	55	-	-	-	-	1	2	4	7	9	10	22	-
2002.....	62	-	-	-	-	1	1	4	7	17	12	20	-
2001.....	57	-	-	-	-	-	2	10	10	2	15	18	-
2000.....	41	-	-	-	-	-	3	4	1	7	9	17	-
1999.....	59	-	-	-	-	-	-	6	9	9	11	24	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
All origins, both sexes													
2003.....	3,108	-	1	-	5	28	93	201	267	401	903	1,209	-
2002.....	2,895	-	-	-	5	28	77	165	273	402	866	1,079	-
2001.....	2,826	1	-	-	7	28	99	146	247	417	855	1,026	-
2000.....	2,785	-	-	1	6	28	83	147	208	417	863	1,032	-
1999.....	3,327	-	-	2	6	21	114	156	262	470	1,087	1,209	-
All origins, male													
2003.....	1,326	-	1	-	4	13	62	130	159	189	382	386	-
2002.....	1,205	-	-	-	3	19	44	105	158	198	357	321	-
2001.....	1,169	1	-	-	4	17	63	99	126	206	339	314	-
2000.....	1,163	-	-	-	4	18	49	87	108	206	350	341	-
1999.....	1,390	-	-	1	3	16	67	100	146	223	460	374	-
All origins, female													
2003.....	1,782	-	-	-	1	15	31	71	108	212	521	823	-
2002.....	1,690	-	-	-	2	9	33	60	115	204	509	758	-
2001.....	1,657	-	-	-	3	11	36	47	121	211	516	712	-
2000.....	1,622	-	-	1	2	10	34	60	100	211	513	691	-
1999.....	1,937	-	-	1	3	5	47	56	116	247	627	835	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
Hispanic, both sexes													
2003.....	158	-	1	-	1	-	6	11	19	33	38	49	-
2002.....	172	-	-	-	2	3	4	11	26	34	51	41	-
2001.....	160	-	-	-	1	2	9	10	21	34	41	42	-
2000.....	138	-	-	-	1	3	8	14	15	27	32	38	-
1999.....	144	-	-	-	-	1	9	11	19	32	39	33	-
Hispanic, male													
2003.....	75	-	1	-	1	-	1	7	11	12	19	23	-
2002.....	80	-	-	-	1	1	2	5	18	18	20	15	-
2001.....	76	-	-	-	1	-	5	6	14	17	17	16	-
2000.....	67	-	-	-	1	2	4	7	5	19	18	11	-
1999.....	65	-	-	-	-	-	6	6	12	14	14	13	-
Hispanic, female													
2003.....	83	-	-	-	-	-	5	4	8	21	19	26	-
2002.....	92	-	-	-	1	2	2	6	8	16	31	26	-
2001.....	84	-	-	-	-	2	4	4	7	17	24	26	-
2000.....	71	-	-	-	-	1	4	7	10	8	14	27	-
1999.....	79	-	-	-	-	1	3	5	7	18	25	20	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
Non-Hispanic 1/, both sexes													
2003.....	2,944	-	-	-	4	28	86	190	247	368	863	1,158	-
2002.....	2,714	-	-	-	3	25	73	153	247	365	812	1,036	-
2001.....	2,650	1	-	-	6	26	87	135	225	380	810	980	-
2000.....	2,637	-	-	1	5	25	75	133	192	389	826	991	-
1999.....	3,171	-	-	2	6	20	102	142	242	436	1,047	1,174	-
Non-Hispanic 1/, male													
2003.....	1,248	-	-	-	3	13	60	123	147	177	362	363	-
2002.....	1,120	-	-	-	2	18	42	99	140	178	336	305	-
2001.....	1,086	1	-	-	3	17	56	92	111	188	321	297	-
2000.....	1,089	-	-	-	3	16	45	80	102	186	329	328	-
1999.....	1,318	-	-	1	3	16	59	91	134	208	446	360	-
Non-Hispanic 1/, female													
2003.....	1,696	-	-	-	1	15	26	67	100	191	501	795	-
2002.....	1,594	-	-	-	1	7	31	54	107	187	476	731	-
2001.....	1,564	-	-	-	3	9	31	43	114	192	489	683	-
2000.....	1,548	-	-	1	2	9	30	53	90	203	497	663	-
1999.....	1,853	-	-	1	3	4	43	51	108	228	601	814	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
Non-Hispanic white, both sexes													
2003.....	1,942	-	-	-	1	7	14	42	92	196	646	944	-
2002.....	1,752	-	-	-	-	8	17	29	82	184	604	828	-
2001.....	1,759	-	-	-	2	5	16	43	95	204	601	793	-
2000.....	1,769	-	-	-	-	5	30	40	75	220	593	806	-
1999.....	2,091	-	-	1	-	6	19	43	91	215	768	948	-
Non-Hispanic white, male													
2003.....	789	-	-	-	-	2	12	30	57	106	288	294	-
2002.....	679	-	-	-	-	5	13	19	47	96	256	243	-
2001.....	687	-	-	-	2	3	12	27	42	101	250	250	-
2000.....	710	-	-	-	-	4	18	24	41	109	256	258	-
1999.....	843	-	-	1	-	6	11	26	48	108	346	297	-
Non-Hispanic white, female													
2003.....	1,153	-	-	-	1	5	2	12	35	90	358	650	-
2002.....	1,073	-	-	-	-	3	4	10	35	88	348	585	-
2001.....	1,072	-	-	-	-	2	4	16	53	103	351	543	-
2000.....	1,059	-	-	-	-	1	12	16	34	111	337	548	-
1999.....	1,248	-	-	-	-	-	8	17	43	107	422	651	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
Non-Hispanic black, both sexes													
2003.....	910	-	-	-	3	20	71	142	148	154	191	181	-
2002.....	897	-	-	-	2	17	54	124	159	165	187	189	-
2001.....	821	1	-	-	4	21	65	92	125	164	187	162	-
2000.....	797	-	-	1	5	19	45	91	112	161	203	160	-
1999.....	989	-	-	1	6	13	82	92	143	205	243	204	-
Non-Hispanic black, male													
2003.....	411	-	-	-	3	10	47	88	86	63	66	48	-
2002.....	402	-	-	-	1	13	27	80	88	70	66	57	-
2001.....	370	1	-	-	1	14	41	65	66	81	62	39	-
2000.....	348	-	-	-	3	12	27	55	57	74	65	55	-
1999.....	426	-	-	-	3	9	47	60	81	92	83	51	-
Non-Hispanic black, female													
2003.....	499	-	-	-	-	10	24	54	62	91	125	133	-
2002.....	495	-	-	-	1	4	27	44	71	95	121	132	-
2001.....	451	-	-	-	3	7	24	27	59	83	125	123	-
2000.....	449	-	-	1	2	7	18	36	55	87	138	105	-
1999.....	563	-	-	1	3	4	35	32	62	113	160	153	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hypertensive heart and renal disease (I13)													
Origin not stated 2/, both sexes													
2003.....	6	-	-	-	-	-	1	-	1	-	2	2	-
2002.....	9	-	-	-	-	-	-	1	-	3	3	2	-
2001.....	16	-	-	-	-	-	3	1	1	3	4	4	-
2000.....	10	-	-	-	-	-	-	-	1	1	5	3	-
1999.....	12	-	-	-	-	-	3	3	1	2	1	2	-
Origin not stated 2/, male													
2003.....	3	-	-	-	-	-	1	-	1	-	1	-	-
2002.....	5	-	-	-	-	-	-	1	-	2	1	1	-
2001.....	7	-	-	-	-	-	2	1	1	1	1	1	-
2000.....	7	-	-	-	-	-	-	-	1	1	3	2	-
1999.....	7	-	-	-	-	-	2	3	-	1	-	1	-
Origin not stated 2/, female													
2003.....	3	-	-	-	-	-	-	-	-	-	1	2	-
2002.....	4	-	-	-	-	-	-	-	-	1	2	1	-
2001.....	9	-	-	-	-	-	1	-	-	2	3	3	-
2000.....	3	-	-	-	-	-	-	-	-	-	2	1	-
1999.....	5	-	-	-	-	-	1	-	1	1	1	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
All origins, both sexes													
2003.....	480,028	22	12	24	160	1,082	7,547	25,361	47,377	78,880	149,370	170,164	29
2002.....	494,382	23	7	20	133	996	7,683	25,575	47,262	83,867	155,391	173,388	37
2001.....	502,189	26	9	20	163	1,116	7,534	25,073	46,574	87,463	159,854	174,309	48
2000.....	515,204	30	10	16	123	935	7,519	24,890	47,737	92,885	163,692	177,319	48
1999.....	529,659	27	5	22	137	1,014	7,751	24,775	48,377	98,840	169,296	179,376	39
All origins, male													
2003.....	246,342	13	6	10	107	813	5,679	19,481	33,767	49,999	78,573	57,872	22
2002.....	252,760	15	4	13	92	741	5,840	19,551	33,656	53,136	81,239	58,446	27
2001.....	254,005	17	3	13	119	830	5,712	19,141	33,006	55,072	82,953	57,106	33
2000.....	260,574	17	7	9	81	704	5,787	19,149	33,795	58,554	84,747	57,690	34
1999.....	267,268	13	3	17	93	739	5,970	19,061	34,123	62,187	87,305	57,726	31
All origins, female													
2003.....	233,686	9	6	14	53	269	1,868	5,880	13,610	28,881	70,797	112,292	7
2002.....	241,622	8	3	7	41	255	1,843	6,024	13,606	30,731	74,152	114,942	10
2001.....	248,184	9	6	7	44	286	1,822	5,932	13,568	32,391	76,901	117,203	15
2000.....	254,630	13	3	7	42	231	1,732	5,741	13,942	34,331	78,945	119,629	14
1999.....	262,391	14	2	5	44	275	1,781	5,714	14,254	36,653	81,991	121,650	8

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Hispanic, both sexes													
2003.....	20,783	2	1	4	22	104	446	1,352	2,546	4,337	6,232	5,731	6
2002.....	20,941	4	1	3	18	88	442	1,389	2,683	4,510	6,060	5,742	1
2001.....	20,664	4	1	4	18	109	368	1,314	2,404	4,553	6,095	5,790	4
2000.....	19,744	6	1	1	12	69	392	1,240	2,457	4,364	5,626	5,572	4
1999.....	19,936	2	-	2	11	87	366	1,328	2,463	4,492	5,655	5,525	5
Hispanic, male													
2003.....	11,054	1	1	3	20	83	341	1,066	1,780	2,680	3,168	1,906	5
2002.....	11,295	2	-	2	12	76	346	1,058	1,907	2,775	3,099	2,017	1
2001.....	11,019	3	1	1	13	90	295	1,016	1,667	2,804	3,116	2,010	3
2000.....	10,562	4	1	1	9	56	322	973	1,726	2,661	2,888	1,918	3
1999.....	10,604	-	-	2	8	72	310	1,042	1,677	2,770	2,869	1,849	5
Hispanic, female													
2003.....	9,729	1	-	1	2	21	105	286	766	1,657	3,064	3,825	1
2002.....	9,646	2	1	1	6	12	96	331	776	1,735	2,961	3,725	-
2001.....	9,645	1	-	3	5	19	73	298	737	1,749	2,979	3,780	1
2000.....	9,182	2	-	-	3	13	70	267	731	1,703	2,738	3,654	1
1999.....	9,332	2	-	-	3	15	56	286	786	1,722	2,786	3,676	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Non-Hispanic 1/, both sexes													
2003.....	457,794	20	11	20	137	974	7,070	23,913	44,605	74,256	142,711	164,061	16
2002.....	471,655	19	6	17	114	900	7,214	24,069	44,359	78,987	148,834	167,107	29
2001.....	479,842	21	8	16	144	1,004	7,135	23,663	43,946	82,604	153,239	168,032	30
2000.....	493,541	23	9	15	111	861	7,082	23,523	45,022	88,137	157,523	171,202	33
1999.....	507,700	25	5	20	125	926	7,344	23,331	45,666	93,919	163,035	173,278	26
Non-Hispanic 1/, male													
2003.....	234,433	12	5	7	86	727	5,316	18,341	31,813	47,124	75,165	55,826	11
2002.....	240,462	13	4	11	79	658	5,475	18,400	31,581	50,116	77,866	56,240	19
2001.....	242,054	13	2	12	105	737	5,392	18,047	31,160	52,059	79,549	54,956	22
2000.....	248,932	12	6	8	72	645	5,426	18,074	31,880	55,623	81,578	55,585	23
1999.....	255,524	13	3	15	85	666	5,623	17,928	32,249	59,117	84,127	55,678	20
Non-Hispanic 1/, female													
2003.....	223,361	8	6	13	51	247	1,754	5,572	12,792	27,132	67,546	108,235	5
2002.....	231,193	6	2	6	35	242	1,739	5,669	12,778	28,871	70,968	110,867	10
2001.....	237,788	8	6	4	39	267	1,743	5,616	12,786	30,545	73,690	113,076	8
2000.....	244,609	11	3	7	39	216	1,656	5,449	13,142	32,514	75,945	115,617	10
1999.....	252,176	12	2	5	40	260	1,721	5,403	13,417	34,802	78,908	117,600	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Non-Hispanic white, both sexes													
2003.....	400,101	14	5	11	91	695	5,434	18,508	36,009	62,110	126,817	150,394	13
2002.....	413,230	11	4	12	76	650	5,565	18,799	35,799	66,553	132,548	153,193	20
2001.....	420,959	14	5	10	86	718	5,456	18,518	35,421	69,681	136,815	154,217	18
2000.....	434,505	16	9	12	80	606	5,437	18,531	36,367	74,867	141,111	157,445	24
1999.....	447,960	16	4	12	76	665	5,607	18,425	37,066	80,129	146,594	159,351	15
Non-Hispanic white, male													
2003.....	205,310	8	2	4	57	527	4,194	14,610	26,363	40,216	67,652	51,669	8
2002.....	211,391	8	2	8	54	493	4,346	14,856	26,108	43,205	70,370	51,928	13
2001.....	212,928	10	1	8	65	539	4,250	14,590	25,776	44,881	72,036	50,758	14
2000.....	219,732	9	6	6	52	473	4,319	14,723	26,398	48,374	74,042	51,313	17
1999.....	225,909	8	3	10	57	482	4,416	14,585	26,754	51,553	76,763	51,267	11
Non-Hispanic white, female													
2003.....	194,791	6	3	7	34	168	1,240	3,898	9,646	21,894	59,165	98,725	5
2002.....	201,839	3	2	4	22	157	1,219	3,943	9,691	23,348	62,178	101,265	7
2001.....	208,031	4	4	2	21	179	1,206	3,928	9,645	24,800	64,779	103,459	4
2000.....	214,773	7	3	6	28	133	1,118	3,808	9,969	26,493	67,069	106,132	7
1999.....	222,051	8	1	2	19	183	1,191	3,840	10,312	28,576	69,831	108,084	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Non-Hispanic black, both sexes													
2003.....	48,617	4	5	8	41	229	1,407	4,726	7,439	10,360	13,207	11,190	1
2002.....	49,522	7	2	5	31	210	1,441	4,665	7,504	10,598	13,541	11,509	9
2001.....	50,295	7	3	5	52	246	1,474	4,533	7,451	11,107	13,782	11,625	10
2000.....	50,659	4	-	2	24	229	1,451	4,418	7,587	11,485	13,809	11,641	9
1999.....	51,368	8	1	8	44	213	1,552	4,362	7,542	11,929	13,890	11,808	11
Non-Hispanic black, male													
2003.....	23,919	3	3	3	26	159	938	3,178	4,624	5,784	6,094	3,106	1
2002.....	23,976	4	2	3	18	131	947	3,053	4,722	5,811	6,013	3,266	6
2001.....	24,255	3	1	3	35	167	974	2,982	4,596	6,089	6,115	3,283	7
2000.....	24,289	2	-	2	14	150	946	2,886	4,713	6,121	6,118	3,331	6
1999.....	24,736	4	-	5	24	148	1,042	2,904	4,737	6,437	5,991	3,435	9
Non-Hispanic black, female													
2003.....	24,698	1	2	5	15	70	469	1,548	2,815	4,576	7,113	8,084	-
2002.....	25,546	3	-	2	13	79	494	1,612	2,782	4,787	7,528	8,243	3
2001.....	26,040	4	2	2	17	79	500	1,551	2,855	5,018	7,667	8,342	3
2000.....	26,370	2	-	-	10	79	505	1,532	2,874	5,364	7,691	8,310	3
1999.....	26,632	4	1	3	20	65	510	1,458	2,805	5,492	7,899	8,373	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Ischemic heart diseases (I20-I25)													
Origin not stated 2/, both sexes													
2003.....	1,451	-	-	-	1	4	31	96	226	287	427	372	7
2002.....	1,786	-	-	-	1	8	27	117	220	370	497	539	7
2001.....	1,683	1	-	-	1	3	31	96	224	306	520	487	14
2000.....	1,919	1	-	-	-	5	45	127	258	384	543	545	11
1999.....	2,023	-	-	-	1	1	41	116	248	429	606	573	8
Origin not stated 2/, male													
2003.....	855	-	-	-	1	3	22	74	174	195	240	140	6
2002.....	1,003	-	-	-	1	7	19	93	168	245	274	189	7
2001.....	932	1	-	-	1	3	25	78	179	209	288	140	8
2000.....	1,080	1	-	-	-	3	39	102	189	270	281	187	8
1999.....	1,140	-	-	-	-	1	37	91	197	300	309	199	6
Origin not stated 2/, female													
2003.....	596	-	-	-	-	1	9	22	52	92	187	232	1
2002.....	783	-	-	-	-	1	8	24	52	125	223	350	-
2001.....	751	-	-	-	-	-	6	18	45	97	232	347	6
2000.....	839	-	-	-	-	2	6	25	69	114	262	358	3
1999.....	883	-	-	-	1	-	4	25	51	129	297	374	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
All origins, both sexes													
2003.....	170,564	9	6	12	73	413	3,094	10,420	19,767	30,923	53,108	52,734	5
2002.....	179,514	11	5	7	62	400	3,161	10,619	19,933	34,003	56,590	54,718	5
2001.....	184,757	15	2	7	64	475	3,176	10,817	19,830	35,380	59,254	55,727	10
2000.....	192,898	20	1	6	52	375	3,132	10,877	21,013	38,459	62,210	56,745	8
1999.....	199,454	13	2	11	60	434	3,249	10,823	21,406	41,441	64,711	57,297	7
All origins, male													
2003.....	89,515	4	2	6	46	300	2,301	7,984	13,972	19,290	27,350	18,256	4
2002.....	93,830	7	2	2	41	292	2,347	7,995	13,966	21,264	28,919	18,991	4
2001.....	95,875	10	2	4	48	333	2,384	8,158	13,858	21,963	30,125	18,984	6
2000.....	100,306	10	1	4	30	269	2,387	8,265	14,638	23,823	31,631	19,243	5
1999.....	104,169	6	1	7	40	306	2,470	8,300	15,029	25,708	32,943	19,357	2
All origins, female													
2003.....	81,049	5	4	6	27	113	793	2,436	5,795	11,633	25,758	34,478	1
2002.....	85,684	4	3	5	21	108	814	2,624	5,967	12,739	27,671	35,727	1
2001.....	88,882	5	-	3	16	142	792	2,659	5,972	13,417	29,129	36,743	4
2000.....	92,592	10	-	2	22	106	745	2,612	6,375	14,636	30,579	37,502	3
1999.....	95,285	7	1	4	20	128	779	2,523	6,377	15,733	31,768	37,940	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Hispanic, both sexes													
2003.....	7,370	-	-	2	11	37	175	508	929	1,683	2,188	1,837	-
2002.....	7,631	3	1	1	7	34	177	518	1,089	1,726	2,184	1,891	-
2001.....	7,403	4	1	1	6	46	139	503	914	1,701	2,190	1,896	2
2000.....	7,106	5	-	-	5	31	154	486	941	1,661	2,049	1,774	-
1999.....	7,158	2	-	-	4	32	142	528	919	1,699	2,082	1,750	-
Hispanic, male													
2003.....	3,903	-	-	1	10	27	127	394	635	1,016	1,067	626	-
2002.....	4,125	2	-	-	5	29	126	382	755	1,043	1,121	662	-
2001.....	3,972	3	1	-	6	37	110	383	603	1,036	1,093	699	1
2000.....	3,825	3	-	-	4	24	125	363	643	984	1,047	632	-
1999.....	3,836	-	-	-	3	27	116	391	607	1,028	1,072	592	-
Hispanic, female													
2003.....	3,467	-	-	1	1	10	48	114	294	667	1,121	1,211	-
2002.....	3,506	1	1	1	2	5	51	136	334	683	1,063	1,229	-
2001.....	3,431	1	-	1	-	9	29	120	311	665	1,097	1,197	1
2000.....	3,281	2	-	-	1	7	29	123	298	677	1,002	1,142	-
1999.....	3,322	2	-	-	1	5	26	137	312	671	1,010	1,158	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Non-Hispanic 1/, both sexes													
2003.....	162,813	9	6	10	62	375	2,909	9,886	18,788	29,159	50,806	50,800	3
2002.....	171,441	8	4	6	55	365	2,975	10,076	18,801	32,172	54,285	52,689	5
2001.....	176,895	11	1	6	58	428	3,023	10,294	18,844	33,604	56,923	53,696	7
2000.....	185,283	14	1	6	47	344	2,963	10,363	20,000	36,695	60,020	54,823	7
1999.....	191,729	11	2	11	56	402	3,090	10,264	20,409	39,614	62,466	55,400	4
Non-Hispanic 1/, male													
2003.....	85,386	4	2	5	36	272	2,168	7,571	13,297	18,224	26,216	17,588	3
2002.....	89,480	5	2	2	36	262	2,215	7,597	13,183	20,161	27,736	18,277	4
2001.....	91,655	7	1	4	42	295	2,261	7,758	13,202	20,875	28,956	18,249	5
2000.....	96,218	6	1	4	26	245	2,250	7,886	13,950	22,772	30,516	18,558	4
1999.....	100,014	6	1	7	37	279	2,338	7,887	14,363	24,598	31,785	18,712	1
Non-Hispanic 1/, female													
2003.....	77,427	5	4	5	26	103	741	2,315	5,491	10,935	24,590	33,212	-
2002.....	81,961	3	2	4	19	103	760	2,479	5,618	12,011	26,549	34,412	1
2001.....	85,240	4	-	2	16	133	762	2,536	5,642	12,729	27,967	35,447	2
2000.....	89,065	8	-	2	21	99	713	2,477	6,050	13,923	29,504	36,265	3
1999.....	91,715	5	1	4	19	123	752	2,377	6,046	15,016	30,681	36,688	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Non-Hispanic white, both sexes													
2003.....	142,402	7	3	7	43	273	2,277	7,840	15,624	24,791	45,199	46,335	3
2002.....	150,051	6	4	4	36	267	2,345	8,094	15,624	27,467	48,317	47,883	4
2001.....	155,312	8	1	1	36	313	2,367	8,302	15,677	28,749	50,908	48,946	4
2000.....	163,203	10	1	4	34	243	2,336	8,410	16,678	31,568	53,840	50,073	6
1999.....	169,545	7	2	6	35	293	2,446	8,404	17,116	34,339	56,289	50,606	2
Non-Hispanic white, male													
2003.....	75,157	3	1	3	26	201	1,748	6,186	11,352	15,828	23,607	16,199	3
2002.....	79,037	4	2	2	25	200	1,793	6,306	11,266	17,673	25,036	16,727	3
2001.....	81,175	6	1	1	28	219	1,834	6,491	11,303	18,302	26,282	16,704	4
2000.....	85,412	5	1	2	20	183	1,839	6,635	11,919	20,062	27,749	16,994	3
1999.....	89,142	3	1	4	26	202	1,907	6,671	12,319	21,791	29,104	17,114	-
Non-Hispanic white, female													
2003.....	67,245	4	2	4	17	72	529	1,654	4,272	8,963	21,592	30,136	-
2002.....	71,014	2	2	2	11	67	552	1,788	4,358	9,794	23,281	31,156	1
2001.....	74,137	2	-	-	8	94	533	1,811	4,374	10,447	24,626	32,242	-
2000.....	77,791	5	-	2	14	60	497	1,775	4,759	11,506	26,091	33,079	3
1999.....	80,403	4	1	2	9	91	539	1,733	4,797	12,548	27,185	33,492	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Non-Hispanic black, both sexes													
2003.....	17,268	1	2	3	18	87	573	1,790	2,699	3,714	4,671	3,710	-
2002.....	18,162	2	-	2	15	88	556	1,756	2,800	3,993	4,939	4,010	1
2001.....	18,399	3	-	4	21	97	572	1,740	2,752	4,173	5,039	3,995	3
2000.....	18,934	2	-	2	11	93	560	1,720	2,910	4,415	5,189	4,031	1
1999.....	19,120	4	-	5	19	88	580	1,653	2,862	4,542	5,256	4,109	2
Non-Hispanic black, male													
2003.....	8,381	-	1	2	10	59	371	1,172	1,624	1,986	2,103	1,053	-
2002.....	8,607	1	-	-	7	54	359	1,109	1,651	2,086	2,154	1,185	1
2001.....	8,628	1	-	2	13	62	354	1,068	1,606	2,163	2,155	1,203	1
2000.....	8,959	1	-	2	5	55	355	1,068	1,731	2,279	2,247	1,215	1
1999.....	9,106	3	-	3	10	61	374	1,055	1,743	2,375	2,211	1,270	1
Non-Hispanic black, female													
2003.....	8,887	1	1	1	8	28	202	618	1,075	1,728	2,568	2,657	-
2002.....	9,555	1	-	2	8	34	197	647	1,149	1,907	2,785	2,825	-
2001.....	9,771	2	-	2	8	35	218	672	1,146	2,010	2,884	2,792	2
2000.....	9,975	1	-	-	6	38	205	652	1,179	2,136	2,942	2,816	-
1999.....	10,014	1	-	2	9	27	206	598	1,119	2,167	3,045	2,839	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute myocardial infarction (I21-I22)													
Origin not stated 2/, both sexes													
2003.....	381	-	-	-	-	1	10	26	50	81	114	97	2
2002.....	442	-	-	-	-	1	9	25	43	105	121	138	-
2001.....	459	-	-	-	-	1	14	20	72	75	141	135	1
2000.....	509	1	-	-	-	-	15	28	72	103	141	148	1
1999.....	567	-	-	-	-	-	17	31	78	128	163	147	3
Origin not stated 2/, male													
2003.....	226	-	-	-	-	1	6	19	40	50	67	42	1
2002.....	225	-	-	-	-	1	6	16	28	60	62	52	-
2001.....	248	-	-	-	-	1	13	17	53	52	76	36	-
2000.....	263	1	-	-	-	-	12	16	45	67	68	53	1
1999.....	319	-	-	-	-	-	16	22	59	82	86	53	1
Origin not stated 2/, female													
2003.....	155	-	-	-	-	-	4	7	10	31	47	55	1
2002.....	217	-	-	-	-	-	3	9	15	45	59	86	-
2001.....	211	-	-	-	-	-	1	3	19	23	65	99	1
2000.....	246	-	-	-	-	-	3	12	27	36	73	95	-
1999.....	248	-	-	-	-	-	1	9	19	46	77	94	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
All origins, both sexes													
2003.....	3,158	4	1	1	5	11	98	277	403	579	837	941	1
2002.....	3,407	2	-	1	4	18	77	345	460	631	898	971	-
2001.....	3,351	3	-	2	7	14	94	307	463	645	871	945	-
2000.....	3,363	1	1	-	8	16	92	282	484	644	906	928	1
1999.....	3,232	3	1	1	3	6	81	252	448	674	855	908	-
All origins, male													
2003.....	1,621	2	1	-	5	7	73	197	278	361	414	282	1
2002.....	1,767	1	-	1	1	12	50	250	320	390	444	298	-
2001.....	1,720	2	-	1	6	10	65	230	320	393	424	269	-
2000.....	1,749	1	1	-	6	13	71	217	341	398	446	254	1
1999.....	1,665	1	1	1	3	4	63	178	315	410	423	266	-
All origins, female													
2003.....	1,537	2	-	1	-	4	25	80	125	218	423	659	-
2002.....	1,640	1	-	-	3	6	27	95	140	241	454	673	-
2001.....	1,631	1	-	1	1	4	29	77	143	252	447	676	-
2000.....	1,614	-	-	-	2	3	21	65	143	246	460	674	-
1999.....	1,567	2	-	-	-	2	18	74	133	264	432	642	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
Hispanic, both sexes													
2003.....	47	-	-	-	1	1	1	5	6	10	12	10	1
2002.....	64	-	-	1	1	1	1	10	12	14	15	9	-
2001.....	63	-	-	-	-	-	1	4	12	19	14	13	-
2000.....	69	-	1	-	1	1	1	6	10	15	17	17	-
1999.....	53	-	-	1	-	-	1	7	13	8	12	11	-
Hispanic, male													
2003.....	21	-	-	-	1	1	1	2	3	5	4	3	1
2002.....	40	-	-	1	-	1	1	8	9	10	7	3	-
2001.....	42	-	-	-	-	-	1	2	9	14	7	9	-
2000.....	39	-	1	-	1	1	1	3	5	11	9	7	-
1999.....	26	-	-	1	-	-	1	4	6	4	8	2	-
Hispanic, female													
2003.....	26	-	-	-	-	-	-	3	3	5	8	7	-
2002.....	24	-	-	-	1	-	-	2	3	4	8	6	-
2001.....	21	-	-	-	-	-	-	2	3	5	7	4	-
2000.....	30	-	-	-	-	-	-	3	5	4	8	10	-
1999.....	27	-	-	-	-	-	-	3	7	4	4	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
Non-Hispanic 1/, both sexes													
2003.....	3,104	4	1	1	4	10	97	272	395	566	824	930	-
2002.....	3,339	2	-	-	2	17	75	335	447	617	883	961	-
2001.....	3,284	3	-	2	7	14	93	303	450	625	855	932	-
2000.....	3,282	1	-	-	7	15	90	273	473	626	887	909	1
1999.....	3,167	3	1	-	3	6	80	245	433	664	841	891	-
Non-Hispanic 1/, male													
2003.....	1,595	2	1	-	4	6	72	195	273	354	409	279	-
2002.....	1,725	1	-	-	-	11	49	242	311	380	437	294	-
2001.....	1,675	2	-	1	6	10	64	228	310	378	416	260	-
2000.....	1,705	1	-	-	5	12	69	212	336	386	436	247	1
1999.....	1,635	1	1	-	3	4	62	174	307	406	414	263	-
Non-Hispanic 1/, female													
2003.....	1,509	2	-	1	-	4	25	77	122	212	415	651	-
2002.....	1,614	1	-	-	2	6	26	93	136	237	446	667	-
2001.....	1,609	1	-	1	1	4	29	75	140	247	439	672	-
2000.....	1,577	-	-	-	2	3	21	61	137	240	451	662	-
1999.....	1,532	2	-	-	-	2	18	71	126	258	427	628	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
Non-Hispanic white, both sexes													
2003.....	2,613	2	-	1	3	5	68	200	304	463	716	851	-
2002.....	2,843	1	-	-	1	11	54	251	356	510	782	877	-
2001.....	2,756	3	-	2	4	11	62	222	342	511	757	842	-
2000.....	2,767	1	-	-	5	11	66	207	357	514	774	832	-
1999.....	2,689	1	1	-	3	5	52	169	344	553	736	825	-
Non-Hispanic white, male													
2003.....	1,314	1	-	-	3	4	55	146	209	296	346	254	-
2002.....	1,454	1	-	-	-	7	41	179	251	313	390	272	-
2001.....	1,386	2	-	1	3	9	45	170	240	307	374	235	-
2000.....	1,412	1	-	-	3	8	52	165	261	314	382	226	-
1999.....	1,361	1	1	-	3	4	40	123	240	339	366	244	-
Non-Hispanic white, female													
2003.....	1,299	1	-	1	-	1	13	54	95	167	370	597	-
2002.....	1,389	-	-	-	1	4	13	72	105	197	392	605	-
2001.....	1,370	1	-	1	1	2	17	52	102	204	383	607	-
2000.....	1,355	-	-	-	2	3	14	42	96	200	392	606	-
1999.....	1,328	-	-	-	-	1	12	46	104	214	370	581	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute ischemic heart diseases (I24)													
Non-Hispanic black, both sexes													
2003.....	432	1	1	-	1	3	27	62	81	91	97	68	-
2002.....	443	1	-	-	1	6	19	78	78	100	89	71	-
2001.....	453	-	-	-	3	2	29	69	98	95	79	78	-
2000.....	449	-	-	-	1	4	22	61	103	91	97	69	1
1999.....	429	2	-	-	-	1	26	73	83	93	97	54	-
Non-Hispanic black, male													
2003.....	254	1	1	-	1	1	16	43	58	53	57	23	-
2002.....	242	-	-	-	-	4	6	58	50	64	40	20	-
2001.....	249	-	-	-	3	-	19	50	62	59	35	21	-
2000.....	255	-	-	-	1	4	16	44	67	58	46	18	1
1999.....	246	-	-	-	-	-	20	49	63	56	41	17	-
Non-Hispanic black, female													
2003.....	178	-	-	-	-	2	11	19	23	38	40	45	-
2002.....	201	1	-	-	1	2	13	20	28	36	49	51	-
2001.....	204	-	-	-	-	2	10	19	36	36	44	57	-
2000.....	194	-	-	-	-	-	6	17	36	33	51	51	-
1999.....	183	2	-	-	-	1	6	24	20	37	56	37	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other acute ischemic heart diseases (I24)

Origin not stated 2/, both sexes

2003.....	7	-	-	-	-	-	-	-	2	3	1	1	-
2002.....	4	-	-	-	1	-	1	-	1	-	-	1	-
2001.....	4	-	-	-	-	-	-	-	1	1	2	-	-
2000.....	12	-	-	-	-	-	1	3	1	3	2	2	-
1999.....	12	-	-	-	-	-	-	-	2	2	2	6	-

Origin not stated 2/, male

2003.....	5	-	-	-	-	-	-	-	2	2	1	-	-
2002.....	2	-	-	-	1	-	-	-	-	-	-	1	-
2001.....	3	-	-	-	-	-	-	-	1	1	1	-	-
2000.....	5	-	-	-	-	-	1	2	-	1	1	-	-
1999.....	4	-	-	-	-	-	-	-	2	-	1	1	-

Origin not stated 2/, female

2003.....	2	-	-	-	-	-	-	-	-	1	-	1	-
2002.....	2	-	-	-	-	-	1	-	1	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	7	-	-	-	-	-	-	1	1	2	1	2	-
1999.....	8	-	-	-	-	-	-	-	-	2	1	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
All origins, both sexes													
2003.....	306,306	9	5	11	82	658	4,355	14,664	27,207	47,378	95,425	116,489	23
2002.....	311,461	10	2	12	67	578	4,445	14,611	26,869	49,233	97,903	117,699	32
2001.....	314,081	8	7	11	92	627	4,264	13,949	26,281	51,438	99,729	117,637	38
2000.....	318,943	9	8	10	63	544	4,295	13,731	26,240	53,782	100,576	119,646	39
1999.....	326,973	11	2	10	74	574	4,421	13,700	26,523	56,725	103,730	121,171	32
All origins, male													
2003.....	155,206	7	3	4	56	506	3,305	11,300	19,517	30,348	50,809	39,334	17
2002.....	157,163	7	2	10	50	437	3,443	11,306	19,370	31,482	51,876	39,157	23
2001.....	156,410	5	1	8	65	487	3,263	10,753	18,828	32,716	52,404	37,853	27
2000.....	158,519	6	5	5	45	422	3,329	10,667	18,816	34,333	52,670	38,193	28
1999.....	161,434	6	1	9	50	429	3,437	10,583	18,779	36,069	53,939	38,103	29
All origins, female													
2003.....	151,100	2	2	7	26	152	1,050	3,364	7,690	17,030	44,616	77,155	6
2002.....	154,298	3	-	2	17	141	1,002	3,305	7,499	17,751	46,027	78,542	9
2001.....	157,671	3	6	3	27	140	1,001	3,196	7,453	18,722	47,325	79,784	11
2000.....	160,424	3	3	5	18	122	966	3,064	7,424	19,449	47,906	81,453	11
1999.....	165,539	5	1	1	24	145	984	3,117	7,744	20,656	49,791	83,068	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Hispanic, both sexes													
2003.....	13,366	2	1	2	10	66	270	839	1,611	2,644	4,032	3,884	5
2002.....	13,246	1	-	1	10	53	264	861	1,582	2,770	3,861	3,842	1
2001.....	13,198	-	-	3	12	63	228	807	1,478	2,833	3,891	3,881	2
2000.....	12,569	1	-	1	6	37	237	748	1,506	2,688	3,560	3,781	4
1999.....	12,725	-	-	1	7	55	223	793	1,531	2,785	3,561	3,764	5
Hispanic, male													
2003.....	7,130	1	1	2	9	55	213	670	1,142	1,659	2,097	1,277	4
2002.....	7,130	-	-	1	7	46	219	668	1,143	1,722	1,971	1,352	1
2001.....	7,005	-	-	1	7	53	184	631	1,055	1,754	2,016	1,302	2
2000.....	6,698	1	-	1	4	31	196	607	1,078	1,666	1,832	1,279	3
1999.....	6,742	-	-	1	5	45	193	647	1,064	1,738	1,789	1,255	5
Hispanic, female													
2003.....	6,236	1	-	-	1	11	57	169	469	985	1,935	2,607	1
2002.....	6,116	1	-	-	3	7	45	193	439	1,048	1,890	2,490	-
2001.....	6,193	-	-	2	5	10	44	176	423	1,079	1,875	2,579	-
2000.....	5,871	-	-	-	2	6	41	141	428	1,022	1,728	2,502	1
1999.....	5,983	-	-	-	2	10	30	146	467	1,047	1,772	2,509	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Non-Hispanic 1/, both sexes													
2003.....	291,877	7	4	9	71	589	4,064	13,755	25,422	44,531	91,081	112,331	13
2002.....	296,875	9	2	11	57	518	4,164	13,658	25,111	46,198	93,666	113,457	24
2001.....	299,663	7	7	8	79	562	4,019	13,066	24,652	48,375	95,461	113,404	23
2000.....	304,976	8	8	9	57	502	4,029	12,887	24,549	50,816	96,616	115,470	25
1999.....	312,804	11	2	9	66	518	4,174	12,822	24,824	53,641	99,728	116,987	22
Non-Hispanic 1/, male													
2003.....	147,452	6	2	2	46	449	3,076	10,575	18,243	28,546	48,540	37,959	8
2002.....	149,257	7	2	9	43	385	3,211	10,561	18,087	29,575	49,693	37,669	15
2001.....	148,724	4	1	7	57	432	3,067	10,061	17,648	30,806	50,177	36,447	17
2000.....	151,009	5	5	4	41	388	3,107	9,976	17,594	32,465	50,626	36,780	18
1999.....	153,875	6	1	8	45	383	3,223	9,867	17,579	34,113	51,928	36,703	19
Non-Hispanic 1/, female													
2003.....	144,425	1	2	7	25	140	988	3,180	7,179	15,985	42,541	74,372	5
2002.....	147,618	2	-	2	14	133	953	3,097	7,024	16,623	43,973	75,788	9
2001.....	150,939	3	6	1	22	130	952	3,005	7,004	17,569	45,284	76,957	6
2000.....	153,967	3	3	5	16	114	922	2,911	6,955	18,351	45,990	78,690	7
1999.....	158,929	5	1	1	21	135	951	2,955	7,245	19,528	47,800	80,284	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Non-Hispanic white, both sexes													
2003.....	255,086	5	2	3	45	417	3,089	10,468	20,081	36,856	80,902	103,208	10
2002.....	260,336	4	-	8	39	372	3,166	10,454	19,819	38,576	83,449	104,433	16
2001.....	262,891	3	4	7	46	394	3,027	9,994	19,402	40,421	85,150	104,429	14
2000.....	268,535	5	8	8	41	352	3,035	9,914	19,332	42,785	86,497	106,540	18
1999.....	275,726	8	1	6	38	367	3,109	9,852	19,606	45,237	89,569	107,920	13
Non-Hispanic white, male													
2003.....	128,839	4	1	1	28	322	2,391	8,278	14,802	24,092	43,699	35,216	5
2002.....	130,900	3	-	6	29	286	2,512	8,371	14,591	25,219	44,944	34,929	10
2001.....	130,367	2	-	6	34	311	2,371	7,929	14,233	26,272	45,380	33,819	10
2000.....	132,908	3	5	4	29	282	2,428	7,923	14,218	27,998	45,911	34,093	14
1999.....	135,406	4	1	6	28	276	2,469	7,791	14,195	29,423	47,293	33,909	11
Non-Hispanic white, female													
2003.....	126,247	1	1	2	17	95	698	2,190	5,279	12,764	37,203	67,992	5
2002.....	129,436	1	-	2	10	86	654	2,083	5,228	13,357	38,505	69,504	6
2001.....	132,524	1	4	1	12	83	656	2,065	5,169	14,149	39,770	70,610	4
2000.....	135,627	2	3	4	12	70	607	1,991	5,114	14,787	40,586	72,447	4
1999.....	140,320	4	-	-	10	91	640	2,061	5,411	15,814	42,276	74,011	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Non-Hispanic black, both sexes													
2003.....	30,917	2	2	5	22	139	807	2,874	4,659	6,555	8,439	7,412	1
2002.....	30,917	4	2	3	15	116	866	2,831	4,626	6,505	8,513	7,428	8
2001.....	31,443	4	3	1	28	147	873	2,724	4,601	6,839	8,664	7,552	7
2000.....	31,276	2	-	-	12	132	869	2,637	4,574	6,979	8,523	7,541	7
1999.....	31,819	2	1	3	25	124	946	2,636	4,597	7,294	8,537	7,645	9
Non-Hispanic black, male													
2003.....	15,284	2	1	1	15	99	551	1,963	2,942	3,745	3,934	2,030	1
2002.....	15,127	3	2	3	11	73	582	1,886	3,021	3,661	3,819	2,061	5
2001.....	15,378	2	1	1	19	105	601	1,864	2,928	3,867	3,925	2,059	6
2000.....	15,075	1	-	-	8	91	575	1,774	2,915	3,784	3,825	2,098	4
1999.....	15,384	1	-	2	14	87	648	1,800	2,931	4,006	3,739	2,148	8
Non-Hispanic black, female													
2003.....	15,633	-	1	4	7	40	256	911	1,717	2,810	4,505	5,382	-
2002.....	15,790	1	-	-	4	43	284	945	1,605	2,844	4,694	5,367	3
2001.....	16,065	2	2	-	9	42	272	860	1,673	2,972	4,739	5,493	1
2000.....	16,201	1	-	-	4	41	294	863	1,659	3,195	4,698	5,443	3
1999.....	16,435	1	1	1	11	37	298	836	1,666	3,288	4,798	5,497	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Origin not stated 2/, both sexes													
2003.....	1,063	-	-	-	1	3	21	70	174	203	312	274	5
2002.....	1,340	-	-	-	-	7	17	92	176	265	376	400	7
2001.....	1,220	1	-	-	1	2	17	76	151	230	377	352	13
2000.....	1,398	-	-	-	-	5	29	96	185	278	400	395	10
1999.....	1,444	-	-	-	1	1	24	85	168	299	441	420	5
Origin not stated 2/, male													
2003.....	624	-	-	-	1	2	16	55	132	143	172	98	5
2002.....	776	-	-	-	-	6	13	77	140	185	212	136	7
2001.....	681	1	-	-	1	2	12	61	125	156	211	104	8
2000.....	812	-	-	-	-	3	26	84	144	202	212	134	7
1999.....	817	-	-	-	-	1	21	69	136	218	222	145	5
Origin not stated 2/, female													
2003.....	439	-	-	-	-	1	5	15	42	60	140	176	-
2002.....	564	-	-	-	-	1	4	15	36	80	164	264	-
2001.....	539	-	-	-	-	-	5	15	26	74	166	248	5
2000.....	586	-	-	-	-	2	3	12	41	76	188	261	3
1999.....	627	-	-	-	1	-	3	16	32	81	219	275	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
All origins, both sexes													
2003.....	67,232	2	-	-	16	211	1,488	5,684	9,110	11,494	18,339	20,874	14
2002.....	68,129	2	-	1	17	181	1,515	5,366	8,793	11,817	18,618	21,796	23
2001.....	68,103	-	-	-	20	179	1,457	5,047	8,438	12,005	19,196	21,736	25
2000.....	69,554	1	-	1	15	144	1,376	4,980	8,242	12,567	19,545	22,664	19
1999.....	71,926	-	1	1	13	160	1,488	4,737	8,383	13,395	20,321	23,403	24
All origins, male													
2003.....	36,042	2	-	-	12	181	1,157	4,552	6,694	7,422	9,333	6,677	12
2002.....	36,047	1	-	-	13	144	1,216	4,239	6,549	7,617	9,416	6,835	17
2001.....	35,440	-	-	-	16	133	1,130	4,037	6,245	7,668	9,671	6,521	19
2000.....	35,896	-	-	1	12	115	1,099	3,977	6,125	8,069	9,725	6,758	15
1999.....	36,633	-	-	1	10	119	1,192	3,752	6,123	8,546	9,963	6,905	22
All origins, female													
2003.....	31,190	-	-	-	4	30	331	1,132	2,416	4,072	9,006	14,197	2
2002.....	32,082	1	-	1	4	37	299	1,127	2,244	4,200	9,202	14,961	6
2001.....	32,663	-	-	-	4	46	327	1,010	2,193	4,337	9,525	15,215	6
2000.....	33,658	1	-	-	3	29	277	1,003	2,117	4,498	9,820	15,906	4
1999.....	35,293	-	1	-	3	41	296	985	2,260	4,849	10,358	16,498	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Hispanic, both sexes													
2003.....	3,295	-	-	-	3	21	93	374	614	700	798	689	3
2002.....	3,166	1	-	-	4	26	108	327	586	722	744	647	1
2001.....	3,150	-	-	-	1	25	81	343	548	748	735	667	2
2000.....	2,953	-	-	-	2	7	83	299	522	707	649	682	2
1999.....	3,063	-	-	-	1	14	73	308	540	747	708	669	3
Hispanic, male													
2003.....	2,094	-	-	-	3	20	80	327	482	480	442	258	2
2002.....	1,946	-	-	-	3	24	89	270	461	475	375	248	1
2001.....	1,928	-	-	-	-	21	63	288	416	502	405	231	2
2000.....	1,794	-	-	-	2	6	71	257	406	438	368	245	1
1999.....	1,847	-	-	-	1	12	63	260	384	527	372	225	3
Hispanic, female													
2003.....	1,201	-	-	-	-	1	13	47	132	220	356	431	1
2002.....	1,220	1	-	-	1	2	19	57	125	247	369	399	-
2001.....	1,222	-	-	-	1	4	18	55	132	246	330	436	-
2000.....	1,159	-	-	-	-	1	12	42	116	269	281	437	1
1999.....	1,216	-	-	-	-	2	10	48	156	220	336	444	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Non-Hispanic 1/, both sexes													
2003.....	63,550	2	-	-	13	189	1,385	5,269	8,403	10,715	17,436	20,132	6
2002.....	64,528	1	-	1	13	152	1,403	4,987	8,109	10,992	17,771	21,082	17
2001.....	64,534	-	-	-	19	154	1,366	4,664	7,816	11,159	18,346	20,999	11
2000.....	66,141	1	-	1	13	137	1,283	4,631	7,627	11,756	18,769	21,912	11
1999.....	68,373	-	1	1	12	146	1,404	4,387	7,754	12,524	19,465	22,661	18
Non-Hispanic 1/, male													
2003.....	33,677	2	-	-	9	160	1,067	4,193	6,136	6,883	8,828	6,394	5
2002.....	33,809	1	-	-	10	118	1,124	3,927	6,004	7,076	8,977	6,561	11
2001.....	33,247	-	-	-	16	112	1,059	3,717	5,770	7,093	9,200	6,270	10
2000.....	33,786	-	-	1	10	109	1,019	3,674	5,646	7,549	9,281	6,487	10
1999.....	34,451	-	-	1	9	107	1,118	3,457	5,663	7,922	9,504	6,654	16
Non-Hispanic 1/, female													
2003.....	29,873	-	-	-	4	29	318	1,076	2,267	3,832	8,608	13,738	1
2002.....	30,719	-	-	1	3	34	279	1,060	2,105	3,916	8,794	14,521	6
2001.....	31,287	-	-	-	3	42	307	947	2,046	4,066	9,146	14,729	1
2000.....	32,355	1	-	-	3	28	264	957	1,981	4,207	9,488	15,425	1
1999.....	33,922	-	1	-	3	39	286	930	2,091	4,602	9,961	16,007	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Non-Hispanic white, both sexes													
2003.....	52,268	1	-	-	9	136	1,047	3,870	6,273	8,237	14,706	17,985	4
2002.....	53,161	1	-	1	11	105	1,020	3,703	6,043	8,444	14,972	18,851	10
2001.....	53,031	-	-	-	11	108	999	3,421	5,649	8,504	15,521	18,811	7
2000.....	54,544	-	-	1	11	95	944	3,407	5,537	9,079	15,871	19,591	8
1999.....	56,375	-	-	1	5	96	1,017	3,155	5,626	9,665	16,517	20,283	10
Non-Hispanic white, male													
2003.....	27,329	1	-	-	6	114	814	3,161	4,680	5,338	7,490	5,722	3
2002.....	27,493	1	-	-	8	84	844	3,008	4,559	5,526	7,583	5,874	6
2001.....	26,960	-	-	-	8	82	794	2,809	4,291	5,496	7,849	5,625	6
2000.....	27,461	-	-	1	8	77	775	2,786	4,210	5,949	7,864	5,783	8
1999.....	27,943	-	-	1	5	73	838	2,558	4,208	6,227	8,097	5,928	8
Non-Hispanic white, female													
2003.....	24,939	-	-	-	3	22	233	709	1,593	2,899	7,216	12,263	1
2002.....	25,668	-	-	1	3	21	176	695	1,484	2,918	7,389	12,977	4
2001.....	26,071	-	-	-	3	26	205	612	1,358	3,008	7,672	13,186	1
2000.....	27,083	-	-	-	3	18	169	621	1,327	3,130	8,007	13,808	-
1999.....	28,432	-	-	-	-	23	179	597	1,418	3,438	8,420	14,355	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Non-Hispanic black, both sexes													
2003.....	9,764	1	-	-	3	46	281	1,213	1,880	2,157	2,344	1,838	1
2002.....	10,024	-	-	-	2	39	333	1,156	1,844	2,258	2,466	1,919	7
2001.....	10,156	-	-	-	7	38	324	1,115	1,927	2,361	2,472	1,909	3
2000.....	10,283	1	-	-	1	36	292	1,120	1,877	2,380	2,522	2,051	3
1999.....	10,643	-	1	-	7	41	358	1,105	1,904	2,553	2,593	2,073	8
Non-Hispanic black, male													
2003.....	5,431	1	-	-	2	39	203	881	1,267	1,327	1,158	552	1
2002.....	5,483	-	-	-	2	27	233	806	1,278	1,357	1,223	552	5
2001.....	5,476	-	-	-	7	24	230	799	1,299	1,412	1,174	528	3
2000.....	5,528	-	-	-	1	26	207	802	1,282	1,398	1,204	606	2
1999.....	5,692	-	-	-	4	27	254	792	1,278	1,518	1,210	601	8
Non-Hispanic black, female													
2003.....	4,333	-	-	-	1	7	78	332	613	830	1,186	1,286	-
2002.....	4,541	-	-	-	-	12	100	350	566	901	1,243	1,367	2
2001.....	4,680	-	-	-	-	14	94	316	628	949	1,298	1,381	-
2000.....	4,755	1	-	-	-	10	85	318	595	982	1,318	1,445	1
1999.....	4,951	-	1	-	3	14	104	313	626	1,035	1,383	1,472	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Origin not stated 2/, both sexes													
2003.....	387	-	-	-	-	1	10	41	93	79	105	53	5
2002.....	435	-	-	-	-	3	4	52	98	103	103	67	5
2001.....	419	-	-	-	-	-	10	40	74	98	115	70	12
2000.....	460	-	-	-	-	-	10	50	93	104	127	70	6
1999.....	490	-	-	-	-	-	11	42	89	124	148	73	3
Origin not stated 2/, male													
2003.....	271	-	-	-	-	1	10	32	76	59	63	25	5
2002.....	292	-	-	-	-	2	3	42	84	66	64	26	5
2001.....	265	-	-	-	-	-	8	32	59	73	66	20	7
2000.....	316	-	-	-	-	-	9	46	73	82	76	26	4
1999.....	335	-	-	-	-	-	11	35	76	97	87	26	3
Origin not stated 2/, female													
2003.....	116	-	-	-	-	-	-	9	17	20	42	28	-
2002.....	143	-	-	-	-	1	1	10	14	37	39	41	-
2001.....	154	-	-	-	-	-	2	8	15	25	49	50	5
2000.....	144	-	-	-	-	-	1	4	20	22	51	44	2
1999.....	155	-	-	-	-	-	-	7	13	27	61	47	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
All origins, both sexes													
2003.....	239,074	7	5	11	66	447	2,867	8,980	18,097	35,884	77,086	95,615	9
2002.....	243,332	8	2	11	50	397	2,930	9,245	18,076	37,416	79,285	95,903	9
2001.....	245,978	8	7	11	72	448	2,807	8,902	17,843	39,433	80,533	95,901	13
2000.....	249,389	8	8	9	48	400	2,919	8,751	17,998	41,215	81,031	96,982	20
1999.....	255,047	11	1	9	61	414	2,933	8,963	18,140	43,330	83,409	97,768	8
All origins, male													
2003.....	119,164	5	3	4	44	325	2,148	6,748	12,823	22,926	41,476	32,657	5
2002.....	121,116	6	2	10	37	293	2,227	7,067	12,821	23,865	42,460	32,322	6
2001.....	120,970	5	1	8	49	354	2,133	6,716	12,583	25,048	42,733	31,332	8
2000.....	122,623	6	5	4	33	307	2,230	6,690	12,691	26,264	42,945	31,435	13
1999.....	124,801	6	1	8	40	310	2,245	6,831	12,656	27,523	43,976	31,198	7
All origins, female													
2003.....	119,910	2	2	7	22	122	719	2,232	5,274	12,958	35,610	62,958	4
2002.....	122,216	2	-	1	13	104	703	2,178	5,255	13,551	36,825	63,581	3
2001.....	125,008	3	6	3	23	94	674	2,186	5,260	14,385	37,800	64,569	5
2000.....	126,766	2	3	5	15	93	689	2,061	5,307	14,951	38,086	65,547	7
1999.....	130,246	5	-	1	21	104	688	2,132	5,484	15,807	39,433	66,570	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
Hispanic, both sexes													
2003.....	10,071	2	1	2	7	45	177	465	997	1,944	3,234	3,195	2
2002.....	10,080	-	-	1	6	27	156	534	996	2,048	3,117	3,195	-
2001.....	10,048	-	-	3	11	38	147	464	930	2,085	3,156	3,214	-
2000.....	9,616	1	-	1	4	30	154	449	984	1,981	2,911	3,099	2
1999.....	9,662	-	-	1	6	41	150	485	991	2,038	2,853	3,095	2
Hispanic, male													
2003.....	5,036	1	1	2	6	35	133	343	660	1,179	1,655	1,019	2
2002.....	5,184	-	-	1	4	22	130	398	682	1,247	1,596	1,104	-
2001.....	5,077	-	-	1	7	32	121	343	639	1,252	1,611	1,071	-
2000.....	4,904	1	-	1	2	25	125	350	672	1,228	1,464	1,034	2
1999.....	4,895	-	-	1	4	33	130	387	680	1,211	1,417	1,030	2
Hispanic, female													
2003.....	5,035	1	-	-	1	10	44	122	337	765	1,579	2,176	-
2002.....	4,896	-	-	-	2	5	26	136	314	801	1,521	2,091	-
2001.....	4,971	-	-	2	4	6	26	121	291	833	1,545	2,143	-
2000.....	4,712	-	-	-	2	5	29	99	312	753	1,447	2,065	-
1999.....	4,767	-	-	-	2	8	20	98	311	827	1,436	2,065	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
Non-Hispanic 1/, both sexes													
2003.....	228,327	5	4	9	58	400	2,679	8,486	17,019	33,816	73,645	92,199	7
2002.....	232,347	8	2	10	44	366	2,761	8,671	17,002	35,206	75,895	92,375	7
2001.....	235,129	7	7	8	60	408	2,653	8,402	16,836	37,216	77,115	92,405	12
2000.....	238,835	7	8	8	44	365	2,746	8,256	16,922	39,060	77,847	93,558	14
1999.....	244,431	11	1	8	54	372	2,770	8,435	17,070	41,117	80,263	94,326	4
Non-Hispanic 1/, male													
2003.....	113,775	4	2	2	37	289	2,009	6,382	12,107	21,663	39,712	31,565	3
2002.....	115,448	6	2	9	33	267	2,087	6,634	12,083	22,499	40,716	31,108	4
2001.....	115,477	4	1	7	41	320	2,008	6,344	11,878	23,713	40,977	30,177	7
2000.....	117,223	5	5	3	31	279	2,088	6,302	11,948	24,916	41,345	30,293	8
1999.....	119,424	6	1	7	36	276	2,105	6,410	11,916	26,191	42,424	30,049	3
Non-Hispanic 1/, female													
2003.....	114,552	1	2	7	21	111	670	2,104	4,912	12,153	33,933	60,634	4
2002.....	116,899	2	-	1	11	99	674	2,037	4,919	12,707	35,179	61,267	3
2001.....	119,652	3	6	1	19	88	645	2,058	4,958	13,503	36,138	62,228	5
2000.....	121,612	2	3	5	13	86	658	1,954	4,974	14,144	36,502	63,265	6
1999.....	125,007	5	-	1	18	96	665	2,025	5,154	14,926	37,839	64,277	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
Non-Hispanic white, both sexes													
2003.....	202,818	4	2	3	36	281	2,042	6,598	13,808	28,619	66,196	85,223	6
2002.....	207,175	3	-	7	28	267	2,146	6,751	13,776	30,132	68,477	85,582	6
2001.....	209,860	3	4	7	35	286	2,028	6,573	13,753	31,917	69,629	85,618	7
2000.....	213,991	5	8	7	30	257	2,091	6,507	13,795	33,706	70,626	86,949	10
1999.....	219,351	8	1	5	33	271	2,092	6,697	13,980	35,572	73,052	87,637	3
Non-Hispanic white, male													
2003.....	101,510	3	1	1	22	208	1,577	5,117	10,122	18,754	36,209	29,494	2
2002.....	103,407	2	-	6	21	202	1,668	5,363	10,032	19,693	37,361	29,055	4
2001.....	103,407	2	-	6	26	229	1,577	5,120	9,942	20,776	37,531	28,194	4
2000.....	105,447	3	5	3	21	205	1,653	5,137	10,008	22,049	38,047	28,310	6
1999.....	107,463	4	1	5	23	203	1,631	5,233	9,987	23,196	39,196	27,981	3
Non-Hispanic white, female													
2003.....	101,308	1	1	2	14	73	465	1,481	3,686	9,865	29,987	55,729	4
2002.....	103,768	1	-	1	7	65	478	1,388	3,744	10,439	31,116	56,527	2
2001.....	106,453	1	4	1	9	57	451	1,453	3,811	11,141	32,098	57,424	3
2000.....	108,544	2	3	4	9	52	438	1,370	3,787	11,657	32,579	58,639	4
1999.....	111,888	4	-	-	10	68	461	1,464	3,993	12,376	33,856	59,656	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
Non-Hispanic black, both sexes													
2003.....	21,153	1	2	5	19	93	526	1,661	2,779	4,398	6,095	5,574	-
2002.....	20,893	4	2	3	13	77	533	1,675	2,782	4,247	6,047	5,509	1
2001.....	21,287	4	3	1	21	109	549	1,609	2,674	4,478	6,192	5,643	4
2000.....	20,993	1	-	-	11	96	577	1,517	2,697	4,599	6,001	5,490	4
1999.....	21,176	2	-	3	18	83	588	1,531	2,693	4,741	5,944	5,572	1
Non-Hispanic black, male													
2003.....	9,853	1	1	1	13	60	348	1,082	1,675	2,418	2,776	1,478	-
2002.....	9,644	3	2	3	9	46	349	1,080	1,743	2,304	2,596	1,509	-
2001.....	9,902	2	1	1	12	81	371	1,065	1,629	2,455	2,751	1,531	3
2000.....	9,547	1	-	-	7	65	368	972	1,633	2,386	2,621	1,492	2
1999.....	9,692	1	-	2	10	60	394	1,008	1,653	2,488	2,529	1,547	-
Non-Hispanic black, female													
2003.....	11,300	-	1	4	6	33	178	579	1,104	1,980	3,319	4,096	-
2002.....	11,249	1	-	-	4	31	184	595	1,039	1,943	3,451	4,000	1
2001.....	11,385	2	2	-	9	28	178	544	1,045	2,023	3,441	4,112	1
2000.....	11,446	-	-	-	4	31	209	545	1,064	2,213	3,380	3,998	2
1999.....	11,484	1	-	1	8	23	194	523	1,040	2,253	3,415	4,025	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)													
Origin not stated 2/, both sexes													
2003.....	676	-	-	-	1	2	11	29	81	124	207	221	-
2002.....	905	-	-	-	-	4	13	40	78	162	273	333	2
2001.....	801	1	-	-	1	2	7	36	77	132	262	282	1
2000.....	938	-	-	-	-	5	19	46	92	174	273	325	4
1999.....	954	-	-	-	1	1	13	43	79	175	293	347	2
Origin not stated 2/, male													
2003.....	353	-	-	-	1	1	6	23	56	84	109	73	-
2002.....	484	-	-	-	-	4	10	35	56	119	148	110	2
2001.....	416	1	-	-	1	2	4	29	66	83	145	84	1
2000.....	496	-	-	-	-	3	17	38	71	120	136	108	3
1999.....	482	-	-	-	-	1	10	34	60	121	135	119	2
Origin not stated 2/, female													
2003.....	323	-	-	-	-	1	5	6	25	40	98	148	-
2002.....	421	-	-	-	-	-	3	5	22	43	125	223	-
2001.....	385	-	-	-	-	-	3	7	11	49	117	198	-
2000.....	442	-	-	-	-	2	2	8	21	54	137	217	1
1999.....	472	-	-	-	1	-	3	9	19	54	158	228	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
All origins, both sexes													
2003.....	169,984	416	171	234	904	1,810	4,555	8,716	13,398	23,244	49,260	67,270	6
2002.....	169,540	473	157	227	836	1,808	4,620	8,711	12,946	23,680	49,776	66,300	6
2001.....	166,949	449	215	246	794	1,736	4,472	8,484	12,535	24,056	48,976	64,979	7
2000.....	165,428	457	168	248	853	1,749	4,457	8,087	12,362	24,679	48,367	63,993	8
1999.....	165,828	490	177	248	880	1,779	4,647	7,888	12,580	25,572	48,712	62,848	7
All origins, male													
2003.....	74,738	233	81	141	560	1,081	2,801	5,458	8,142	12,875	22,680	20,681	5
2002.....	74,230	249	82	125	557	1,138	2,860	5,550	7,771	12,991	22,771	20,133	3
2001.....	72,439	222	117	123	500	1,037	2,777	5,307	7,503	13,167	22,540	19,141	5
2000.....	72,122	238	98	150	518	1,049	2,746	5,151	7,532	13,621	22,006	19,008	5
1999.....	72,601	255	100	137	533	1,067	2,894	4,979	7,700	14,320	22,128	18,483	5
All origins, female													
2003.....	95,246	183	90	93	344	729	1,754	3,258	5,256	10,369	26,580	46,589	1
2002.....	95,310	224	75	102	279	670	1,760	3,161	5,175	10,689	27,005	46,167	3
2001.....	94,510	227	98	123	294	699	1,695	3,177	5,032	10,889	26,436	45,838	2
2000.....	93,306	219	70	98	335	700	1,711	2,936	4,830	11,058	26,361	44,985	3
1999.....	93,227	235	77	111	347	712	1,753	2,909	4,880	11,252	26,584	44,365	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
Hispanic, both sexes													
2003.....	5,612	72	33	26	98	209	311	433	621	945	1,417	1,447	-
2002.....	5,152	81	23	27	102	188	298	416	507	857	1,318	1,335	-
2001.....	4,899	75	31	33	86	179	268	370	516	856	1,206	1,277	2
2000.....	4,792	79	28	37	110	182	283	363	493	817	1,149	1,250	1
1999.....	4,687	92	33	34	99	176	309	345	501	815	1,102	1,180	1
Hispanic, male													
2003.....	2,805	40	17	15	68	135	217	281	367	515	661	489	-
2002.....	2,587	39	11	12	76	135	195	260	296	484	626	453	-
2001.....	2,401	40	14	17	57	119	185	252	296	456	556	407	2
2000.....	2,393	39	20	25	76	124	177	234	299	446	510	442	1
1999.....	2,392	40	20	21	60	128	208	225	298	466	513	412	1
Hispanic, female													
2003.....	2,807	32	16	11	30	74	94	152	254	430	756	958	-
2002.....	2,565	42	12	15	26	53	103	156	211	373	692	882	-
2001.....	2,498	35	17	16	29	60	83	118	220	400	650	870	-
2000.....	2,399	40	8	12	34	58	106	129	194	371	639	808	-
1999.....	2,295	52	13	13	39	48	101	120	203	349	589	768	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
Non-Hispanic 1/, both sexes													
2003.....	163,998	342	137	207	802	1,594	4,225	8,249	12,731	22,243	47,750	65,714	4
2002.....	163,970	390	134	200	733	1,612	4,300	8,263	12,402	22,761	48,342	64,830	3
2001.....	161,625	372	183	213	701	1,550	4,184	8,077	11,980	23,137	47,661	63,564	3
2000.....	160,222	375	139	210	741	1,558	4,151	7,694	11,823	23,804	47,124	62,597	6
1999.....	160,711	397	143	213	779	1,594	4,317	7,520	12,030	24,685	47,496	61,532	5
Non-Hispanic 1/, male													
2003.....	71,731	193	64	125	490	942	2,569	5,150	7,739	12,332	21,969	20,155	3
2002.....	71,455	209	71	113	480	999	2,649	5,268	7,454	12,472	22,089	19,649	2
2001.....	69,830	181	103	106	439	913	2,576	5,030	7,181	12,671	21,941	18,687	2
2000.....	69,547	197	78	125	441	921	2,554	4,900	7,203	13,138	21,460	18,527	3
1999.....	69,984	215	80	115	471	933	2,672	4,737	7,370	13,812	21,555	18,021	3
Non-Hispanic 1/, female													
2003.....	92,267	149	73	82	312	652	1,656	3,099	4,992	9,911	25,781	45,559	1
2002.....	92,515	181	63	87	253	613	1,651	2,995	4,948	10,289	26,253	45,181	1
2001.....	91,795	191	80	107	262	637	1,608	3,047	4,799	10,466	25,720	44,877	1
2000.....	90,675	178	61	85	300	637	1,597	2,794	4,620	10,666	25,664	44,070	3
1999.....	90,727	182	63	98	308	661	1,645	2,783	4,660	10,873	25,941	43,511	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
Non-Hispanic white, both sexes													
2003.....	141,413	186	77	132	495	941	2,768	5,698	9,649	18,137	42,620	60,707	3
2002.....	141,712	234	86	130	453	936	2,888	5,703	9,468	18,836	43,140	59,836	2
2001.....	139,716	219	103	140	422	937	2,696	5,631	9,132	19,171	42,486	58,776	3
2000.....	138,745	238	87	122	428	948	2,717	5,362	9,089	19,781	42,174	57,794	5
1999.....	138,869	240	87	140	495	972	2,814	5,238	9,108	20,529	42,406	56,837	3
Non-Hispanic white, male													
2003.....	61,119	102	32	82	298	576	1,737	3,644	5,970	10,234	19,773	18,669	2
2002.....	61,048	128	51	74	304	589	1,836	3,748	5,806	10,465	19,878	18,167	2
2001.....	59,513	116	58	72	261	555	1,731	3,637	5,562	10,653	19,627	17,239	2
2000.....	59,461	127	51	77	257	576	1,726	3,505	5,648	11,102	19,335	17,054	3
1999.....	59,549	136	51	73	291	580	1,809	3,395	5,680	11,656	19,365	16,511	2
Non-Hispanic white, female													
2003.....	80,294	84	45	50	197	365	1,031	2,054	3,679	7,903	22,847	42,038	1
2002.....	80,664	106	35	56	149	347	1,052	1,955	3,662	8,371	23,262	41,669	-
2001.....	80,203	103	45	68	161	382	965	1,994	3,570	8,518	22,859	41,537	1
2000.....	79,284	111	36	45	171	372	991	1,857	3,441	8,679	22,839	40,740	2
1999.....	79,320	104	36	67	204	392	1,005	1,843	3,428	8,873	23,041	40,326	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
Non-Hispanic black, both sexes													
2003.....	19,861	125	49	66	273	586	1,346	2,373	2,795	3,659	4,375	4,213	1
2002.....	19,641	135	38	57	246	608	1,306	2,370	2,671	3,476	4,483	4,250	1
2001.....	19,496	130	70	63	252	567	1,370	2,264	2,608	3,522	4,508	4,142	-
2000.....	19,159	113	46	72	278	559	1,314	2,168	2,485	3,616	4,358	4,149	1
1999.....	19,531	132	48	66	262	576	1,395	2,107	2,678	3,747	4,454	4,064	2
Non-Hispanic black, male													
2003.....	9,275	76	27	38	175	322	767	1,395	1,610	1,863	1,838	1,163	1
2002.....	9,038	72	16	30	151	363	738	1,402	1,490	1,772	1,854	1,150	-
2001.....	9,070	59	39	31	160	327	768	1,279	1,483	1,766	1,978	1,180	-
2000.....	8,883	56	24	39	158	313	747	1,295	1,410	1,802	1,833	1,206	-
1999.....	9,258	66	27	38	166	331	797	1,247	1,545	1,936	1,870	1,234	1
Non-Hispanic black, female													
2003.....	10,586	49	22	28	98	264	579	978	1,185	1,796	2,537	3,050	-
2002.....	10,603	63	22	27	95	245	568	968	1,181	1,704	2,629	3,100	1
2001.....	10,426	71	31	32	92	240	602	985	1,125	1,756	2,530	2,962	-
2000.....	10,276	57	22	33	120	246	567	873	1,075	1,814	2,525	2,943	1
1999.....	10,273	66	21	28	96	245	598	860	1,133	1,811	2,584	2,830	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other heart diseases (I26-I51)													
Origin not stated 2/, both sexes													
2003.....	374	2	1	1	4	7	19	34	46	56	93	109	2
2002.....	418	2	-	-	1	8	22	32	37	62	116	135	3
2001.....	425	2	1	-	7	7	20	37	39	63	109	138	2
2000.....	414	3	1	1	2	9	23	30	46	58	94	146	1
1999.....	430	1	1	1	2	9	21	23	49	72	114	136	1
Origin not stated 2/, male													
2003.....	202	-	-	1	2	4	15	27	36	28	50	37	2
2002.....	188	1	-	-	1	4	16	22	21	35	56	31	1
2001.....	208	1	-	-	4	5	16	25	26	40	43	47	1
2000.....	182	2	-	-	1	4	15	17	30	37	36	39	1
1999.....	225	-	-	1	2	6	14	17	32	42	60	50	1
Origin not stated 2/, female													
2003.....	172	2	1	-	2	3	4	7	10	28	43	72	-
2002.....	230	1	-	-	-	4	6	10	16	27	60	104	2
2001.....	217	1	1	-	3	2	4	12	13	23	66	91	1
2000.....	232	1	1	1	1	5	8	13	16	21	58	107	-
1999.....	205	1	1	-	-	3	7	6	17	30	54	86	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
All origins, both sexes													
2003.....	1,245	1	-	1	16	33	112	183	209	232	318	140	-
2002.....	1,154	2	2	2	7	44	90	174	168	238	306	121	-
2001.....	1,178	1	-	3	8	29	117	186	160	247	316	111	-
2000.....	1,172	3	1	3	13	40	116	186	165	224	302	119	-
1999.....	1,116	2	1	-	7	30	114	143	171	240	289	119	-
All origins, male													
2003.....	682	1	-	1	6	19	63	125	123	119	166	59	-
2002.....	643	1	1	1	6	31	60	117	91	133	153	49	-
2001.....	668	1	-	1	5	17	62	129	95	132	177	49	-
2000.....	665	2	-	1	7	24	73	120	96	122	169	51	-
1999.....	639	2	1	-	3	24	75	103	103	118	154	56	-
All origins, female													
2003.....	563	-	-	-	10	14	49	58	86	113	152	81	-
2002.....	511	1	1	1	1	13	30	57	77	105	153	72	-
2001.....	510	-	-	2	3	12	55	57	65	115	139	62	-
2000.....	507	1	1	2	6	16	43	66	69	102	133	68	-
1999.....	477	-	-	-	4	6	39	40	68	122	135	63	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
Hispanic, both sexes													
2003.....	80	1	-	1	3	4	10	17	12	16	13	3	-
2002.....	69	1	1	-	-	7	5	13	12	14	13	3	-
2001.....	56	-	-	1	1	6	8	10	9	9	10	2	-
2000.....	70	-	1	1	-	6	11	16	8	18	7	2	-
1999.....	63	-	1	-	2	6	10	7	10	13	12	2	-
Hispanic, male													
2003.....	42	1	-	1	1	3	7	9	6	9	3	2	-
2002.....	45	-	-	-	-	6	5	11	7	9	6	1	-
2001.....	34	-	-	-	1	5	6	8	4	5	4	1	-
2000.....	39	-	-	-	-	4	7	10	4	11	2	1	-
1999.....	42	-	1	-	-	5	6	6	5	9	8	2	-
Hispanic, female													
2003.....	38	-	-	-	2	1	3	8	6	7	10	1	-
2002.....	24	1	1	-	-	1	-	2	5	5	7	2	-
2001.....	22	-	-	1	-	1	2	2	5	4	6	1	-
2000.....	31	-	1	1	-	2	4	6	4	7	5	1	-
1999.....	21	-	-	-	2	1	4	1	5	4	4	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
Non-Hispanic 1/, both sexes													
2003.....	1,163	-	-	-	13	29	102	164	197	216	305	137	-
2002.....	1,082	1	1	2	7	37	85	159	155	224	293	118	-
2001.....	1,117	1	-	2	7	23	108	174	150	238	305	109	-
2000.....	1,096	3	-	2	13	34	102	170	157	205	293	117	-
1999.....	1,049	2	-	-	5	23	104	135	160	226	277	117	-
Non-Hispanic 1/, male													
2003.....	638	-	-	-	5	16	56	114	117	110	163	57	-
2002.....	596	1	1	1	6	25	55	105	83	124	147	48	-
2001.....	631	1	-	1	4	12	55	120	90	127	173	48	-
2000.....	625	2	-	1	7	20	65	110	92	111	167	50	-
1999.....	594	2	-	-	3	18	69	96	97	109	146	54	-
Non-Hispanic 1/, female													
2003.....	525	-	-	-	8	13	46	50	80	106	142	80	-
2002.....	486	-	-	1	1	12	30	54	72	100	146	70	-
2001.....	486	-	-	1	3	11	53	54	60	111	132	61	-
2000.....	471	1	-	1	6	14	37	60	65	94	126	67	-
1999.....	455	-	-	-	2	5	35	39	63	117	131	63	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
Non-Hispanic white, both sexes													
2003.....	912	-	-	-	11	22	68	100	137	172	271	131	-
2002.....	862	-	1	-	6	27	62	95	113	179	270	109	-
2001.....	859	1	-	2	5	15	65	114	107	182	265	103	-
2000.....	852	1	-	1	6	21	66	106	114	173	261	103	-
1999.....	814	1	-	-	4	17	59	82	107	182	252	110	-
Non-Hispanic white, male													
2003.....	500	-	-	-	5	11	39	67	85	89	148	56	-
2002.....	481	-	1	-	5	20	41	68	61	102	139	44	-
2001.....	482	1	-	1	3	7	34	79	65	93	154	45	-
2000.....	496	1	-	-	3	15	42	73	72	93	151	46	-
1999.....	466	1	-	-	2	12	39	59	68	93	139	53	-
Non-Hispanic white, female													
2003.....	412	-	-	-	6	11	29	33	52	83	123	75	-
2002.....	381	-	-	-	1	7	21	27	52	77	131	65	-
2001.....	377	-	-	1	2	8	31	35	42	89	111	58	-
2000.....	356	-	-	1	3	6	24	33	42	80	110	57	-
1999.....	348	-	-	-	2	5	20	23	39	89	113	57	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
Non-Hispanic black, both sexes													
2003.....	222	-	-	-	2	5	32	60	53	37	29	4	-
2002.....	197	-	-	1	1	10	22	57	41	38	19	8	-
2001.....	233	-	-	-	1	8	41	57	41	43	37	5	-
2000.....	213	2	-	-	6	11	31	57	39	30	26	11	-
1999.....	208	-	-	-	1	6	41	48	47	42	17	6	-
Non-Hispanic black, male													
2003.....	119	-	-	-	-	4	15	43	27	17	13	-	-
2002.....	104	-	-	-	1	5	14	33	22	19	6	4	-
2001.....	134	-	-	-	-	5	20	39	24	26	18	2	-
2000.....	111	1	-	-	4	3	21	35	18	16	11	2	-
1999.....	115	-	-	-	1	6	27	34	26	15	5	1	-
Non-Hispanic black, female													
2003.....	103	-	-	-	2	1	17	17	26	20	16	4	-
2002.....	93	-	-	1	-	5	8	24	19	19	13	4	-
2001.....	99	-	-	-	1	3	21	18	17	17	19	3	-
2000.....	102	1	-	-	2	8	10	22	21	14	15	9	-
1999.....	93	-	-	-	-	-	14	14	21	27	12	5	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and subacute endocarditis (I33)													
Origin not stated 2/, both sexes													
2003.....	2	-	-	-	-	-	-	2	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	2	1	-	-	-	-
2001.....	5	-	-	-	-	-	1	2	1	-	1	-	-
2000.....	6	-	-	-	-	-	3	-	-	1	2	-	-
1999.....	4	-	-	-	-	1	-	1	1	1	-	-	-
Origin not stated 2/, male													
2003.....	2	-	-	-	-	-	-	2	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	3	-	-	-	-	-	1	1	1	-	-	-	-
2000.....	1	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	3	-	-	-	-	1	-	1	1	-	-	-	-
Origin not stated 2/, female													
2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	1	-	-	1	-	-
2000.....	5	-	-	-	-	-	2	-	-	1	2	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
All origins, both sexes													
2003.....	868	33	22	28	55	48	86	111	104	151	153	77	-
2002.....	848	26	22	22	36	46	75	107	118	143	157	96	-
2001.....	839	32	26	21	40	50	93	111	106	120	154	85	1
2000.....	767	20	24	20	38	53	86	85	89	131	145	76	-
1999.....	890	27	25	21	32	50	114	105	125	146	160	85	-
All origins, male													
2003.....	449	15	8	12	34	29	52	71	57	84	65	22	-
2002.....	434	9	11	7	24	32	49	69	69	71	60	33	-
2001.....	401	18	15	7	22	33	41	59	61	59	61	24	1
2000.....	404	10	13	9	21	29	58	46	52	67	73	26	-
1999.....	442	13	14	7	20	34	64	56	70	69	65	30	-
All origins, female													
2003.....	419	18	14	16	21	19	34	40	47	67	88	55	-
2002.....	414	17	11	15	12	14	26	38	49	72	97	63	-
2001.....	438	14	11	14	18	17	52	52	45	61	93	61	-
2000.....	363	10	11	11	17	24	28	39	37	64	72	50	-
1999.....	448	14	11	14	12	16	50	49	55	77	95	55	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
Hispanic, both sexes													
2003.....	57	5	5	4	7	6	5	6	7	7	5	-	-
2002.....	64	4	2	3	5	3	6	10	14	7	5	5	-
2001.....	58	2	4	2	4	11	5	6	6	6	7	5	-
2000.....	61	5	4	1	5	9	10	4	9	5	7	2	-
1999.....	64	2	3	4	10	8	9	9	2	7	8	2	-
Hispanic, male													
2003.....	33	3	1	2	4	4	3	5	4	6	1	-	-
2002.....	35	3	-	-	3	3	3	7	9	3	1	3	-
2001.....	25	1	2	1	2	6	3	1	3	3	2	1	-
2000.....	29	2	4	-	1	4	6	2	3	-	6	1	-
1999.....	30	1	1	4	5	6	7	2	1	1	2	-	-
Hispanic, female													
2003.....	24	2	4	2	3	2	2	1	3	1	4	-	-
2002.....	29	1	2	3	2	-	3	3	5	4	4	2	-
2001.....	33	1	2	1	2	5	2	5	3	3	5	4	-
2000.....	32	3	-	1	4	5	4	2	6	5	1	1	-
1999.....	34	1	2	-	5	2	2	7	1	6	6	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
Non-Hispanic 1/, both sexes													
2003.....	806	28	17	24	48	41	81	102	97	144	148	76	-
2002.....	783	22	20	19	31	43	69	97	103	136	152	91	-
2001.....	779	30	22	19	36	38	87	105	100	114	147	80	1
2000.....	700	15	19	19	32	44	76	81	78	126	136	74	-
1999.....	824	25	22	17	22	41	104	96	123	139	152	83	-
Non-Hispanic 1/, male													
2003.....	414	12	7	10	30	25	49	64	53	78	64	22	-
2002.....	398	6	11	7	21	29	46	62	59	68	59	30	-
2001.....	375	17	13	6	20	26	38	58	58	56	59	23	1
2000.....	374	8	9	9	20	25	52	44	48	67	67	25	-
1999.....	410	12	13	3	15	27	56	54	69	68	63	30	-
Non-Hispanic 1/, female													
2003.....	392	16	10	14	18	16	32	38	44	66	84	54	-
2002.....	385	16	9	12	10	14	23	35	44	68	93	61	-
2001.....	404	13	9	13	16	12	49	47	42	58	88	57	-
2000.....	326	7	10	10	12	19	24	37	30	59	69	49	-
1999.....	414	13	9	14	7	14	48	42	54	71	89	53	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
Non-Hispanic white, both sexes													
2003.....	616	10	8	13	31	18	50	75	83	126	131	71	-
2002.....	601	9	11	9	21	30	49	69	76	116	126	85	-
2001.....	600	13	10	14	23	26	55	75	79	98	134	72	1
2000.....	528	6	11	9	20	25	49	55	62	103	122	66	-
1999.....	639	10	10	9	14	26	70	73	104	115	134	74	-
Non-Hispanic white, male													
2003.....	324	6	3	5	21	9	33	49	45	71	60	22	-
2002.....	308	3	7	3	14	21	33	43	45	60	51	28	-
2001.....	286	10	6	5	10	18	27	41	45	48	53	22	1
2000.....	280	4	7	5	12	14	34	30	41	52	59	22	-
1999.....	320	6	4	2	10	17	40	40	60	56	58	27	-
Non-Hispanic white, female													
2003.....	292	4	5	8	10	9	17	26	38	55	71	49	-
2002.....	293	6	4	6	7	9	16	26	31	56	75	57	-
2001.....	314	3	4	9	13	8	28	34	34	50	81	50	-
2000.....	248	2	4	4	8	11	15	25	21	51	63	44	-
1999.....	319	4	6	7	4	9	30	33	44	59	76	47	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
Non-Hispanic black, both sexes													
2003.....	168	14	9	9	15	22	30	23	12	16	13	5	-
2002.....	154	12	7	8	8	10	17	26	22	19	20	5	-
2001.....	157	14	12	4	13	10	24	30	19	13	11	7	-
2000.....	140	7	7	8	7	16	24	21	15	18	10	7	-
1999.....	158	10	11	8	6	14	30	20	17	20	16	6	-
Non-Hispanic black, male													
2003.....	81	4	4	5	9	15	16	13	7	5	3	-	-
2002.....	75	3	3	3	5	6	10	17	13	7	7	1	-
2001.....	78	5	7	1	10	7	7	17	11	8	5	-	-
2000.....	72	4	2	3	3	10	16	10	6	11	5	2	-
1999.....	76	4	9	1	3	10	13	12	9	11	4	-	-
Non-Hispanic black, female													
2003.....	87	10	5	4	6	7	14	10	5	11	10	5	-
2002.....	79	9	4	5	3	4	7	9	9	12	13	4	-
2001.....	79	9	5	3	3	3	17	13	8	5	6	7	-
2000.....	68	3	5	5	4	6	8	11	9	7	5	5	-
1999.....	82	6	2	7	3	4	17	8	8	9	12	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
All origins, both sexes													
2003.....	57,448	38	5	12	38	97	315	981	2,469	6,138	17,136	30,219	-
2002.....	56,494	23	11	10	27	101	282	993	2,342	6,168	16,910	29,627	-
2001.....	56,934	24	17	15	34	92	307	954	2,374	6,274	17,134	29,708	1
2000.....	55,704	31	9	21	34	63	262	811	2,211	6,403	16,820	29,035	4
1999.....	54,913	11	10	12	31	65	272	782	2,107	6,524	16,771	28,327	1
All origins, male													
2003.....	22,427	28	3	7	24	60	193	587	1,454	3,313	7,658	9,100	-
2002.....	21,698	14	6	5	15	60	173	617	1,345	3,254	7,573	8,636	-
2001.....	21,632	16	5	8	16	56	182	588	1,347	3,344	7,672	8,398	-
2000.....	21,175	22	7	9	20	36	159	496	1,264	3,414	7,498	8,248	2
1999.....	20,755	5	7	6	23	32	164	486	1,228	3,526	7,318	7,959	1
All origins, female													
2003.....	35,021	10	2	5	14	37	122	394	1,015	2,825	9,478	21,119	-
2002.....	34,796	9	5	5	12	41	109	376	997	2,914	9,337	20,991	-
2001.....	35,302	8	12	7	18	36	125	366	1,027	2,930	9,462	21,310	1
2000.....	34,529	9	2	12	14	27	103	315	947	2,989	9,322	20,787	2
1999.....	34,158	6	3	6	8	33	108	296	879	2,998	9,453	20,368	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Hispanic, both sexes													
2003.....	1,606	7	-	1	4	10	12	48	118	275	494	637	-
2002.....	1,412	4	-	2	-	3	13	43	91	235	451	570	-
2001.....	1,364	8	3	-	1	9	21	43	103	213	404	559	-
2000.....	1,270	7	1	2	-	3	10	22	100	203	365	557	-
1999.....	1,242	3	4	1	3	4	18	29	87	202	353	538	-
Hispanic, male													
2003.....	714	5	-	-	4	7	10	25	77	141	242	203	-
2002.....	593	2	-	2	-	2	6	20	54	121	209	177	-
2001.....	565	6	1	-	1	6	15	31	52	109	169	175	-
2000.....	535	5	1	1	-	2	6	9	62	109	151	189	-
1999.....	522	1	3	-	3	3	12	16	47	112	159	166	-
Hispanic, female													
2003.....	892	2	-	1	-	3	2	23	41	134	252	434	-
2002.....	819	2	-	-	-	1	7	23	37	114	242	393	-
2001.....	799	2	2	-	-	3	6	12	51	104	235	384	-
2000.....	735	2	-	1	-	1	4	13	38	94	214	368	-
1999.....	720	2	1	1	-	1	6	13	40	90	194	372	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Non-Hispanic 1/, both sexes													
2003.....	55,732	31	5	11	34	87	302	933	2,345	5,843	16,614	29,527	-
2002.....	54,932	19	11	8	27	97	268	943	2,245	5,914	16,416	28,984	-
2001.....	55,437	16	14	15	33	82	286	906	2,258	6,044	16,688	29,094	1
2000.....	54,307	24	8	19	34	60	252	789	2,105	6,185	16,417	28,411	3
1999.....	53,528	8	6	11	28	61	253	750	2,007	6,308	16,377	27,718	1
Non-Hispanic 1/, male													
2003.....	21,658	23	3	7	20	53	182	562	1,371	3,161	7,400	8,876	-
2002.....	21,050	12	6	3	15	58	166	592	1,288	3,124	7,344	8,442	-
2001.....	21,012	10	4	8	15	49	167	555	1,287	3,225	7,489	8,203	-
2000.....	20,599	17	6	8	20	34	153	487	1,197	3,296	7,335	8,045	1
1999.....	20,170	4	4	6	20	29	152	469	1,173	3,406	7,140	7,766	1
Non-Hispanic 1/, female													
2003.....	34,074	8	2	4	14	34	120	371	974	2,682	9,214	20,651	-
2002.....	33,882	7	5	5	12	39	102	351	957	2,790	9,072	20,542	-
2001.....	34,425	6	10	7	18	33	119	351	971	2,819	9,199	20,891	1
2000.....	33,708	7	2	11	14	26	99	302	908	2,889	9,082	20,366	2
1999.....	33,358	4	2	5	8	32	101	281	834	2,902	9,237	19,952	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Non-Hispanic white, both sexes													
2003.....	49,788	13	4	6	17	48	165	588	1,743	4,832	14,932	27,440	-
2002.....	49,162	11	8	5	16	49	160	593	1,719	4,914	14,725	26,962	-
2001.....	49,788	11	12	11	17	49	167	608	1,730	5,052	15,033	27,097	1
2000.....	48,782	13	4	14	17	35	154	536	1,611	5,176	14,827	26,392	3
1999.....	47,977	5	4	8	15	30	156	480	1,537	5,302	14,702	25,738	-
Non-Hispanic white, male													
2003.....	19,239	8	2	4	9	31	105	366	1,033	2,672	6,725	8,284	-
2002.....	18,705	8	5	2	7	28	105	387	1,000	2,620	6,663	7,880	-
2001.....	18,730	6	4	5	9	25	103	392	1,019	2,734	6,785	7,648	-
2000.....	18,386	10	4	6	10	20	103	342	942	2,828	6,671	7,449	1
1999.....	17,903	2	2	5	11	14	100	306	922	2,920	6,472	7,149	-
Non-Hispanic white, female													
2003.....	30,549	5	2	2	8	17	60	222	710	2,160	8,207	19,156	-
2002.....	30,457	3	3	3	9	21	55	206	719	2,294	8,062	19,082	-
2001.....	31,058	5	8	6	8	24	64	216	711	2,318	8,248	19,449	1
2000.....	30,396	3	-	8	7	15	51	194	669	2,348	8,156	18,943	2
1999.....	30,074	3	2	3	4	16	56	174	615	2,382	8,230	18,589	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Non-Hispanic black, both sexes													
2003.....	5,294	15	-	5	17	37	131	328	563	921	1,466	1,811	-
2002.....	5,143	8	3	3	9	45	104	331	487	891	1,495	1,767	-
2001.....	5,094	5	2	3	16	32	112	284	494	905	1,475	1,766	-
2000.....	4,936	11	4	4	13	24	94	241	453	909	1,407	1,776	-
1999.....	4,961	3	2	3	12	30	93	257	446	922	1,468	1,724	1
Non-Hispanic black, male													
2003.....	2,132	13	-	3	11	20	73	183	320	445	575	489	-
2002.....	2,055	4	1	1	6	27	59	192	266	450	602	447	-
2001.....	2,036	4	-	2	6	23	61	153	248	447	618	474	-
2000.....	1,942	7	2	2	8	13	47	138	231	420	572	502	-
1999.....	2,005	2	2	1	8	15	50	156	238	442	577	513	1
Non-Hispanic black, female													
2003.....	3,162	2	-	2	6	17	58	145	243	476	891	1,322	-
2002.....	3,088	4	2	2	3	18	45	139	221	441	893	1,320	-
2001.....	3,058	1	2	1	10	9	51	131	246	458	857	1,292	-
2000.....	2,994	4	2	2	5	11	47	103	222	489	835	1,274	-
1999.....	2,956	1	-	2	4	15	43	101	208	480	891	1,211	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Heart failure (I50)													
Origin not stated 2/, both sexes													
2003.....	110	-	-	-	-	-	1	-	6	20	28	55	-
2002.....	150	-	-	-	-	1	1	7	6	19	43	73	-
2001.....	133	-	-	-	-	1	-	5	13	17	42	55	-
2000.....	127	-	-	-	-	-	-	-	6	15	38	67	1
1999.....	143	-	-	-	-	-	1	3	13	14	41	71	-
Origin not stated 2/, male													
2003.....	55	-	-	-	-	-	1	-	6	11	16	21	-
2002.....	55	-	-	-	-	-	1	5	3	9	20	17	-
2001.....	55	-	-	-	-	1	-	2	8	10	14	20	-
2000.....	41	-	-	-	-	-	-	-	5	9	12	14	1
1999.....	63	-	-	-	-	-	-	1	8	8	19	27	-
Origin not stated 2/, female													
2003.....	55	-	-	-	-	-	-	-	-	9	12	34	-
2002.....	95	-	-	-	-	1	-	2	3	10	23	56	-
2001.....	78	-	-	-	-	-	-	3	5	7	28	35	-
2000.....	86	-	-	-	-	-	-	-	1	6	26	53	-
1999.....	80	-	-	-	-	-	1	2	5	6	22	44	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
All origins, both sexes													
2003.....	110,423	344	144	193	795	1,632	4,042	7,441	10,616	16,723	31,653	36,834	6
2002.....	111,044	422	122	193	766	1,617	4,173	7,437	10,318	17,131	32,403	36,456	6
2001.....	107,998	392	172	207	712	1,565	3,955	7,233	9,895	17,415	31,372	35,075	5
2000.....	107,785	403	134	204	768	1,593	3,993	7,005	9,897	17,921	31,100	34,763	4
1999.....	108,909	450	141	215	810	1,634	4,147	6,858	10,177	18,662	31,492	34,317	6
All origins, male													
2003.....	51,180	189	70	121	496	973	2,493	4,675	6,508	9,359	14,791	11,500	5
2002.....	51,455	225	64	112	512	1,015	2,578	4,747	6,266	9,533	14,985	11,415	3
2001.....	49,738	187	97	107	457	931	2,492	4,531	6,000	9,632	14,630	10,670	4
2000.....	49,878	204	78	131	470	960	2,456	4,489	6,120	10,018	14,266	10,683	3
1999.....	50,765	235	78	124	487	977	2,591	4,334	6,299	10,607	14,591	10,438	4
All origins, female													
2003.....	59,243	155	74	72	299	659	1,549	2,766	4,108	7,364	16,862	25,334	1
2002.....	59,589	197	58	81	254	602	1,595	2,690	4,052	7,598	17,418	25,041	3
2001.....	58,260	205	75	100	255	634	1,463	2,702	3,895	7,783	16,742	24,405	1
2000.....	57,907	199	56	73	298	633	1,537	2,516	3,777	7,903	16,834	24,080	1
1999.....	58,144	215	63	91	323	657	1,556	2,524	3,878	8,055	16,901	23,879	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Hispanic, both sexes													
2003.....	3,869	59	28	20	84	189	284	362	484	647	905	807	-
2002.....	3,607	72	20	22	97	175	274	350	390	601	849	757	-
2001.....	3,421	65	24	30	80	153	234	311	398	628	785	711	2
2000.....	3,391	67	22	33	105	164	252	321	376	591	770	689	1
1999.....	3,318	87	25	29	84	158	272	300	402	593	729	638	1
Hispanic, male													
2003.....	2,016	31	16	12	59	121	197	242	280	359	415	284	-
2002.....	1,914	34	11	10	73	124	181	222	226	351	410	272	-
2001.....	1,777	33	11	16	53	102	161	212	237	339	381	230	2
2000.....	1,790	32	15	24	75	114	158	213	230	326	351	251	1
1999.....	1,798	38	15	17	52	114	183	201	245	344	344	244	1
Hispanic, female													
2003.....	1,853	28	12	8	25	68	87	120	204	288	490	523	-
2002.....	1,693	38	9	12	24	51	93	128	164	250	439	485	-
2001.....	1,644	32	13	14	27	51	73	99	161	289	404	481	-
2000.....	1,601	35	7	9	30	50	94	108	146	265	419	438	-
1999.....	1,520	49	10	12	32	44	89	99	157	249	385	394	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Non-Hispanic 1/, both sexes													
2003.....	106,297	283	115	172	707	1,437	3,740	7,050	10,092	16,040	30,683	35,974	4
2002.....	107,173	348	102	171	668	1,435	3,878	7,064	9,899	16,487	31,481	35,637	3
2001.....	104,292	325	147	177	625	1,407	3,703	6,892	9,472	16,741	30,521	34,281	1
2000.....	104,119	333	112	170	662	1,420	3,721	6,654	9,483	17,288	30,278	33,995	3
1999.....	105,310	362	115	185	724	1,469	3,856	6,539	9,740	18,012	30,690	33,614	4
Non-Hispanic 1/, male													
2003.....	49,021	158	54	108	435	848	2,282	4,410	6,198	8,983	14,342	11,200	3
2002.....	49,411	190	53	102	438	887	2,382	4,509	6,024	9,156	14,539	11,129	2
2001.....	47,812	153	86	91	400	826	2,316	4,297	5,746	9,263	14,220	10,413	1
2000.....	47,949	170	63	107	394	842	2,284	4,259	5,866	9,664	13,891	10,407	2
1999.....	48,810	197	63	106	433	859	2,395	4,118	6,031	10,229	14,206	10,171	2
Non-Hispanic 1/, female													
2003.....	57,276	125	61	64	272	589	1,458	2,640	3,894	7,057	16,341	24,774	1
2002.....	57,762	158	49	69	230	548	1,496	2,555	3,875	7,331	16,942	24,508	1
2001.....	56,480	172	61	86	225	581	1,387	2,595	3,726	7,478	16,301	23,868	-
2000.....	56,170	163	49	63	268	578	1,437	2,395	3,617	7,624	16,387	23,588	1
1999.....	56,500	165	52	79	291	610	1,461	2,421	3,709	7,783	16,484	23,443	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Non-Hispanic white, both sexes													
2003.....	90,097	163	65	113	436	853	2,485	4,935	7,686	13,007	27,286	33,065	3
2002.....	91,087	214	66	116	410	830	2,617	4,946	7,560	13,627	28,019	32,680	2
2001.....	88,469	194	81	113	377	847	2,409	4,834	7,216	13,839	27,054	31,504	1
2000.....	88,583	218	72	98	385	867	2,448	4,665	7,302	14,329	26,964	31,233	2
1999.....	89,439	224	73	123	462	899	2,529	4,603	7,360	14,930	27,318	30,915	3
Non-Hispanic white, male													
2003.....	41,056	88	27	73	263	525	1,560	3,162	4,807	7,402	12,840	10,307	2
2002.....	41,554	117	38	69	278	520	1,657	3,250	4,700	7,683	13,025	10,215	2
2001.....	40,015	99	48	61	239	505	1,567	3,125	4,433	7,778	12,635	9,524	1
2000.....	40,299	112	40	66	232	527	1,547	3,060	4,593	8,129	12,454	9,537	2
1999.....	40,860	127	45	66	268	537	1,630	2,990	4,630	8,587	12,696	9,282	2
Non-Hispanic white, female													
2003.....	49,041	75	38	40	173	328	925	1,773	2,879	5,605	14,446	22,758	1
2002.....	49,533	97	28	47	132	310	960	1,696	2,860	5,944	14,994	22,465	-
2001.....	48,454	95	33	52	138	342	842	1,709	2,783	6,061	14,419	21,980	-
2000.....	48,284	106	32	32	153	340	901	1,605	2,709	6,200	14,510	21,696	-
1999.....	48,579	97	28	57	194	362	899	1,613	2,730	6,343	14,622	21,633	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Non-Hispanic black, both sexes													
2003.....	14,177	96	40	52	239	522	1,153	1,962	2,167	2,685	2,867	2,393	1
2002.....	14,147	115	28	45	228	543	1,163	1,956	2,121	2,528	2,949	2,470	1
2001.....	14,012	111	56	56	222	517	1,193	1,893	2,054	2,561	2,985	2,364	-
2000.....	13,870	93	35	60	252	508	1,165	1,849	1,978	2,659	2,915	2,355	1
1999.....	14,204	119	35	55	243	526	1,231	1,782	2,168	2,763	2,953	2,328	1
Non-Hispanic black, male													
2003.....	6,943	59	23	30	155	283	663	1,156	1,256	1,396	1,247	674	1
2002.....	6,804	65	12	26	139	325	655	1,160	1,189	1,296	1,239	698	-
2001.....	6,822	50	32	28	144	292	680	1,070	1,200	1,285	1,337	704	-
2000.....	6,758	44	20	34	143	287	663	1,112	1,155	1,355	1,245	700	-
1999.....	7,062	60	16	36	154	300	707	1,045	1,272	1,468	1,284	720	-
Non-Hispanic black, female													
2003.....	7,234	37	17	22	84	239	490	806	911	1,289	1,620	1,719	-
2002.....	7,343	50	16	19	89	218	508	796	932	1,232	1,710	1,772	1
2001.....	7,190	61	24	28	78	225	513	823	854	1,276	1,648	1,660	-
2000.....	7,112	49	15	26	109	221	502	737	823	1,304	1,670	1,655	1
1999.....	7,142	59	19	19	89	226	524	737	896	1,295	1,669	1,608	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Origin not stated 2/, both sexes													
2003.....	257	2	1	1	4	6	18	29	40	36	65	53	2
2002.....	264	2	-	-	1	7	21	23	29	43	73	62	3
2001.....	285	2	1	-	7	5	18	30	25	46	66	83	2
2000.....	275	3	-	1	1	9	20	30	38	42	52	79	-
1999.....	281	1	1	1	2	7	19	19	35	57	73	65	1
Origin not stated 2/, male													
2003.....	143	-	-	1	2	4	14	23	30	17	34	16	2
2002.....	130	1	-	-	1	4	15	16	16	26	36	14	1
2001.....	149	1	-	-	4	3	15	22	17	30	29	27	1
2000.....	139	2	-	-	1	4	14	17	24	28	24	25	-
1999.....	157	-	-	1	2	4	13	15	23	34	41	23	1
Origin not stated 2/, female													
2003.....	114	2	1	-	2	2	4	6	10	19	31	37	-
2002.....	134	1	-	-	-	3	6	7	13	17	37	48	2
2001.....	136	1	1	-	3	2	3	8	8	16	37	56	1
2000.....	136	1	-	1	-	5	6	13	14	14	28	54	-
1999.....	124	1	1	-	-	3	6	4	12	23	32	42	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All origins, both sexes

2003.....	21,940	1	2	2	22	98	372	1,031	1,755	3,104	6,649	8,904	-
2002.....	20,261	5	1	1	21	85	354	923	1,526	2,922	6,138	8,285	-
2001.....	19,250	-	1	1	23	104	331	931	1,462	2,832	5,993	7,572	-
2000.....	18,073	1	1	4	17	70	348	878	1,440	2,784	5,623	6,907	-
1999.....	16,968	3	3	1	13	78	313	799	1,318	2,799	5,324	6,317	-

All origins, male

2003.....	8,213	1	2	1	12	58	215	660	989	1,535	2,522	2,218	-
2002.....	7,647	2	1	1	11	47	219	560	850	1,419	2,405	2,132	-
2001.....	7,254	-	-	-	16	59	206	569	784	1,379	2,282	1,959	-
2000.....	6,895	1	-	3	4	39	216	555	818	1,308	2,204	1,747	-
1999.....	6,488	2	1	1	8	48	195	482	732	1,372	2,067	1,580	-

All origins, female

2003.....	13,727	-	-	1	10	40	157	371	766	1,569	4,127	6,686	-
2002.....	12,614	3	-	-	10	38	135	363	676	1,503	3,733	6,153	-
2001.....	11,996	-	1	1	7	45	125	362	678	1,453	3,711	5,613	-
2000.....	11,178	-	1	1	13	31	132	323	622	1,476	3,419	5,160	-
1999.....	10,480	1	2	-	5	30	118	317	586	1,427	3,257	4,737	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
Hispanic, both sexes													
2003.....	1,053	-	-	-	4	11	23	75	121	204	285	330	-
2002.....	942	2	-	-	5	9	30	39	123	183	283	268	-
2001.....	889	-	-	-	3	10	19	60	83	187	271	256	-
2000.....	821	1	1	-	4	7	27	55	109	162	232	223	-
1999.....	753	1	1	-	3	8	19	64	82	158	232	185	-
Hispanic, male													
2003.....	437	-	-	-	2	5	17	53	73	84	116	87	-
2002.....	385	1	-	-	1	5	18	22	66	80	113	79	-
2001.....	365	-	-	-	2	8	16	39	34	84	114	68	-
2000.....	386	1	-	-	2	5	20	44	65	72	107	70	-
1999.....	340	-	-	-	2	5	14	35	40	84	99	61	-
Hispanic, female													
2003.....	616	-	-	-	2	6	6	22	48	120	169	243	-
2002.....	557	1	-	-	4	4	12	17	57	103	170	189	-
2001.....	524	-	-	-	1	2	3	21	49	103	157	188	-
2000.....	435	-	1	-	2	2	7	11	44	90	125	153	-
1999.....	413	1	1	-	1	3	5	29	42	74	133	124	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
Non-Hispanic 1/, both sexes													
2003.....	20,831	1	2	2	18	85	349	953	1,628	2,883	6,350	8,560	-
2002.....	19,254	3	1	1	16	75	322	881	1,400	2,725	5,840	7,990	-
2001.....	18,306	-	1	1	20	94	310	868	1,370	2,636	5,706	7,300	-
2000.....	17,182	-	-	4	13	63	318	815	1,320	2,616	5,373	6,660	-
1999.....	16,158	2	2	1	10	69	292	734	1,232	2,633	5,073	6,110	-
Non-Hispanic 1/, male													
2003.....	7,751	1	2	1	10	51	198	605	911	1,441	2,401	2,130	-
2002.....	7,232	1	1	1	10	42	200	536	781	1,328	2,285	2,047	-
2001.....	6,856	-	-	-	14	51	188	528	743	1,288	2,160	1,884	-
2000.....	6,475	-	-	3	2	34	194	504	746	1,232	2,088	1,672	-
1999.....	6,123	2	1	1	6	42	179	447	689	1,285	1,958	1,513	-
Non-Hispanic 1/, female													
2003.....	13,080	-	-	1	8	34	151	348	717	1,442	3,949	6,430	-
2002.....	12,022	2	-	-	6	33	122	345	619	1,397	3,555	5,943	-
2001.....	11,450	-	1	1	6	43	122	340	627	1,348	3,546	5,416	-
2000.....	10,707	-	-	1	11	29	124	311	574	1,384	3,285	4,988	-
1999.....	10,035	-	1	-	4	27	113	287	543	1,348	3,115	4,597	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
Non-Hispanic white, both sexes													
2003.....	15,878	-	1	1	8	32	165	493	918	1,862	5,026	7,372	-
2002.....	14,617	2	1	1	9	26	126	449	805	1,838	4,570	6,790	-
2001.....	13,837	-	1	1	11	35	121	404	776	1,747	4,499	6,242	-
2000.....	12,917	-	-	3	3	25	132	381	736	1,744	4,252	5,641	-
1999.....	12,048	1	-	-	5	28	112	343	686	1,788	3,940	5,145	-
Non-Hispanic white, male													
2003.....	5,673	-	1	-	5	16	102	319	527	976	1,916	1,811	-
2002.....	5,325	1	1	1	7	16	82	290	477	920	1,784	1,746	-
2001.....	4,970	-	-	-	7	20	77	253	439	873	1,701	1,600	-
2000.....	4,721	-	-	3	1	17	89	255	434	852	1,671	1,399	-
1999.....	4,395	1	-	-	2	22	77	220	410	897	1,526	1,240	-
Non-Hispanic white, female													
2003.....	10,205	-	-	1	3	16	63	174	391	886	3,110	5,561	-
2002.....	9,292	1	-	-	2	10	44	159	328	918	2,786	5,044	-
2001.....	8,867	-	1	1	4	15	44	151	337	874	2,798	4,642	-
2000.....	8,196	-	-	-	2	8	43	126	302	892	2,581	4,242	-
1999.....	7,653	-	-	-	3	6	35	123	276	891	2,414	3,905	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
Non-Hispanic black, both sexes													
2003.....	4,445	1	1	-	10	51	174	441	664	918	1,162	1,023	-
2002.....	4,165	1	-	-	7	44	189	413	550	795	1,133	1,033	-
2001.....	4,012	-	-	-	9	55	177	439	553	789	1,056	934	-
2000.....	3,862	-	-	1	10	35	176	405	555	801	980	899	-
1999.....	3,708	1	1	1	5	37	175	370	506	764	1,006	842	-
Non-Hispanic black, male													
2003.....	1,875	1	1	-	5	33	92	278	359	424	419	263	-
2002.....	1,701	-	-	-	3	24	112	234	274	359	442	253	-
2001.....	1,698	-	-	-	7	29	102	263	284	374	399	240	-
2000.....	1,576	-	-	-	1	16	98	229	301	344	358	229	-
1999.....	1,550	1	1	1	4	17	97	218	256	348	381	226	-
Non-Hispanic black, female													
2003.....	2,570	-	-	-	5	18	82	163	305	494	743	760	-
2002.....	2,464	1	-	-	4	20	77	179	276	436	691	780	-
2001.....	2,314	-	-	-	2	26	75	176	269	415	657	694	-
2000.....	2,286	-	-	1	9	19	78	176	254	457	622	670	-
1999.....	2,158	-	-	-	1	20	78	152	250	416	625	616	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Origin not stated 2/, both sexes

2003.....	56	-	-	-	-	2	-	3	6	17	14	14	-
2002.....	65	-	-	-	-	1	2	3	3	14	15	27	-
2001.....	55	-	-	-	-	-	2	3	9	9	16	16	-
2000.....	70	-	-	-	-	-	3	8	11	6	18	24	-
1999.....	57	-	-	-	-	1	2	1	4	8	19	22	-

Origin not stated 2/, male

2003.....	25	-	-	-	-	2	-	2	5	10	5	1	-
2002.....	30	-	-	-	-	-	1	2	3	11	7	6	-
2001.....	33	-	-	-	-	-	2	2	7	7	8	7	-
2000.....	34	-	-	-	-	-	2	7	7	4	9	5	-
1999.....	25	-	-	-	-	1	2	-	3	3	10	6	-

Origin not stated 2/, female

2003.....	31	-	-	-	-	-	-	1	1	7	9	13	-
2002.....	35	-	-	-	-	1	1	1	-	3	8	21	-
2001.....	22	-	-	-	-	-	-	1	2	2	8	9	-
2000.....	36	-	-	-	-	-	1	1	4	2	9	19	-
1999.....	32	-	-	-	-	-	-	1	1	5	9	16	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
All origins, both sexes													
2003.....	157,689	101	46	69	221	583	2,460	6,127	9,946	20,708	52,847	64,579	2
2002.....	162,672	117	53	91	171	567	2,425	6,055	9,897	21,992	54,889	66,412	3
2001.....	163,538	108	54	80	196	601	2,491	5,910	9,608	22,598	55,822	66,066	4
2000.....	167,661	124	47	76	199	602	2,599	6,011	9,956	23,649	57,020	67,376	2
1999.....	167,366	103	46	65	182	580	2,574	5,563	9,652	24,092	57,427	67,080	2
All origins, male													
2003.....	61,426	57	25	41	116	314	1,283	3,350	5,478	10,668	22,234	17,859	1
2002.....	62,622	67	30	47	103	284	1,178	3,278	5,456	11,207	22,655	18,316	1
2001.....	63,177	63	25	42	103	311	1,267	3,205	5,277	11,649	23,301	17,930	4
2000.....	64,769	74	19	33	106	308	1,297	3,235	5,496	12,042	23,946	18,212	1
1999.....	64,485	65	22	34	95	299	1,308	2,996	5,305	12,220	24,088	18,052	1
All origins, female													
2003.....	96,263	44	21	28	105	269	1,177	2,777	4,468	10,040	30,613	46,720	1
2002.....	100,050	50	23	44	68	283	1,247	2,777	4,441	10,785	32,234	48,096	2
2001.....	100,361	45	29	38	93	290	1,224	2,705	4,331	10,949	32,521	48,136	-
2000.....	102,892	50	28	43	93	294	1,302	2,776	4,460	11,607	33,074	49,164	1
1999.....	102,881	38	24	31	87	281	1,266	2,567	4,347	11,872	33,339	49,028	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Hispanic, both sexes													
2003.....	6,658	17	8	14	28	89	267	569	736	1,202	2,029	1,699	-
2002.....	6,451	20	12	20	29	93	256	560	708	1,215	1,918	1,620	-
2001.....	6,416	17	10	16	40	99	275	539	654	1,206	1,892	1,668	-
2000.....	6,187	16	6	8	31	108	267	477	681	1,219	1,742	1,632	-
1999.....	5,907	16	7	9	32	102	241	464	617	1,211	1,672	1,536	-
Hispanic, male													
2003.....	3,070	11	7	8	15	51	162	345	431	617	879	544	-
2002.....	3,003	9	5	7	17	61	154	340	421	652	825	512	-
2001.....	2,982	9	4	10	26	60	151	315	378	637	864	528	-
2000.....	2,865	8	-	3	17	65	145	280	392	645	799	511	-
1999.....	2,809	8	1	6	21	62	146	285	354	641	758	527	-
Hispanic, female													
2003.....	3,588	6	1	6	13	38	105	224	305	585	1,150	1,155	-
2002.....	3,448	11	7	13	12	32	102	220	287	563	1,093	1,108	-
2001.....	3,434	8	6	6	14	39	124	224	276	569	1,028	1,140	-
2000.....	3,322	8	6	5	14	43	122	197	289	574	943	1,121	-
1999.....	3,098	8	6	3	11	40	95	179	263	570	914	1,009	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Non-Hispanic 1/, both sexes													
2003.....	150,719	82	38	55	192	491	2,182	5,536	9,179	19,458	50,742	62,763	1
2002.....	155,852	96	41	71	142	474	2,161	5,477	9,160	20,731	52,850	64,646	3
2001.....	156,741	91	43	63	156	498	2,211	5,350	8,916	21,346	53,810	64,253	4
2000.....	161,057	108	41	67	167	493	2,323	5,509	9,238	22,373	55,129	65,608	1
1999.....	161,057	87	39	56	150	477	2,323	5,071	8,991	22,814	55,643	65,404	2
Non-Hispanic 1/, male													
2003.....	58,226	45	18	33	100	261	1,115	2,993	5,026	10,023	21,322	17,289	1
2002.....	59,470	57	25	40	86	223	1,019	2,926	5,018	10,532	21,773	17,770	1
2001.....	60,030	54	21	32	77	247	1,111	2,882	4,872	10,989	22,377	17,364	4
2000.....	61,715	66	19	30	89	243	1,145	2,936	5,079	11,364	23,084	17,660	-
1999.....	61,499	57	21	28	74	237	1,156	2,697	4,922	11,542	23,275	17,489	1
Non-Hispanic 1/, female													
2003.....	92,493	37	20	22	92	230	1,067	2,543	4,153	9,435	29,420	45,474	-
2002.....	96,382	39	16	31	56	251	1,142	2,551	4,142	10,199	31,077	46,876	2
2001.....	96,711	37	22	31	79	251	1,100	2,468	4,044	10,357	31,433	46,889	-
2000.....	99,342	42	22	37	78	250	1,178	2,573	4,159	11,009	32,045	47,948	1
1999.....	99,558	30	18	28	76	240	1,167	2,374	4,069	11,272	32,368	47,915	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Non-Hispanic white, both sexes													
2003.....	127,969	46	23	37	124	298	1,239	3,294	6,293	15,259	44,277	57,078	1
2002.....	133,118	51	29	44	101	285	1,288	3,320	6,432	16,355	46,327	58,884	2
2001.....	133,879	46	28	42	98	320	1,333	3,200	6,208	16,838	47,304	58,461	1
2000.....	138,210	59	25	46	118	313	1,378	3,433	6,555	17,989	48,556	59,738	-
1999.....	138,716	55	27	40	105	301	1,406	3,143	6,311	18,332	49,178	59,818	-
Non-Hispanic white, male													
2003.....	48,549	22	8	23	72	159	637	1,813	3,427	7,947	18,699	15,741	1
2002.....	49,908	32	18	23	62	136	617	1,797	3,511	8,368	19,181	16,162	1
2001.....	50,382	27	14	24	55	161	684	1,745	3,400	8,753	19,762	15,756	1
2000.....	52,000	38	14	23	67	161	694	1,856	3,605	9,176	20,379	15,987	-
1999.....	51,968	36	15	21	51	151	712	1,662	3,479	9,337	20,618	15,886	-
Non-Hispanic white, female													
2003.....	79,420	24	15	14	52	139	602	1,481	2,866	7,312	25,578	41,337	-
2002.....	83,210	19	11	21	39	149	671	1,523	2,921	7,987	27,146	42,722	1
2001.....	83,497	19	14	18	43	159	649	1,455	2,808	8,085	27,542	42,705	-
2000.....	86,210	21	11	23	51	152	684	1,577	2,950	8,813	28,177	43,751	-
1999.....	86,748	19	12	19	54	150	694	1,481	2,832	8,995	28,560	43,932	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Non-Hispanic black, both sexes													
2003.....	18,649	31	14	16	58	167	786	1,949	2,449	3,411	5,176	4,592	-
2002.....	18,691	36	10	23	38	166	741	1,874	2,303	3,600	5,277	4,622	1
2001.....	18,844	42	12	14	51	153	758	1,886	2,289	3,684	5,257	4,695	3
2000.....	19,053	46	13	15	42	158	829	1,819	2,265	3,660	5,344	4,861	1
1999.....	18,731	27	10	15	40	149	808	1,662	2,269	3,792	5,314	4,644	1
Non-Hispanic black, male													
2003.....	7,798	19	9	9	25	86	385	1,031	1,363	1,673	2,071	1,127	-
2002.....	7,754	19	6	15	22	72	331	986	1,288	1,826	2,031	1,158	-
2001.....	7,828	26	5	5	20	75	355	998	1,252	1,844	2,068	1,177	3
2000.....	7,945	27	3	5	19	70	394	955	1,240	1,817	2,155	1,260	-
1999.....	7,823	17	4	7	19	72	387	900	1,235	1,857	2,100	1,225	-
Non-Hispanic black, female													
2003.....	10,851	12	5	7	33	81	401	918	1,086	1,738	3,105	3,465	-
2002.....	10,937	17	4	8	16	94	410	888	1,015	1,774	3,246	3,464	1
2001.....	11,016	16	7	9	31	78	403	888	1,037	1,840	3,189	3,518	-
2000.....	11,108	19	10	10	23	88	435	864	1,025	1,843	3,189	3,601	1
1999.....	10,908	10	6	8	21	77	421	762	1,034	1,935	3,214	3,419	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cerebrovascular diseases (I60-I69)													
Origin not stated 2/, both sexes													
2003.....	312	2	-	-	1	3	11	22	31	48	76	117	1
2002.....	369	1	-	-	-	-	8	18	29	46	121	146	-
2001.....	381	-	1	1	-	4	5	21	38	46	120	145	-
2000.....	417	-	-	1	1	1	9	25	37	57	149	136	1
1999.....	402	-	-	-	-	1	10	28	44	67	112	140	-
Origin not stated 2/, male													
2003.....	130	1	-	-	1	2	6	12	21	28	33	26	-
2002.....	149	1	-	-	-	-	5	12	17	23	57	34	-
2001.....	165	-	-	-	-	4	5	8	27	23	60	38	-
2000.....	189	-	-	-	-	-	7	19	25	33	63	41	1
1999.....	177	-	-	-	-	-	6	14	29	37	55	36	-
Origin not stated 2/, female													
2003.....	182	1	-	-	-	1	5	10	10	20	43	91	1
2002.....	220	-	-	-	-	-	3	6	12	23	64	112	-
2001.....	216	-	1	1	-	-	-	13	11	23	60	107	-
2000.....	228	-	-	1	1	1	2	6	12	24	86	95	-
1999.....	225	-	-	-	-	1	4	14	15	30	57	104	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
All origins, both sexes													
2003.....	13,053	3	-	-	3	7	47	190	466	1,211	3,837	7,288	1
2002.....	13,821	4	-	-	3	9	50	182	487	1,300	4,062	7,723	1
2001.....	14,086	1	1	1	1	3	40	186	473	1,335	4,156	7,888	1
2000.....	14,393	2	-	-	3	8	48	176	451	1,423	4,277	8,005	-
1999.....	14,979	2	-	-	2	8	45	185	486	1,476	4,385	8,390	-
All origins, male													
2003.....	4,859	2	-	-	2	5	35	131	298	649	1,734	2,002	1
2002.....	5,171	1	-	-	2	5	30	119	305	728	1,803	2,177	1
2001.....	5,143	1	1	1	1	2	26	118	286	743	1,791	2,172	1
2000.....	5,357	-	-	-	2	6	35	119	294	800	1,935	2,166	-
1999.....	5,443	1	-	-	-	5	32	121	307	795	1,927	2,255	-
All origins, female													
2003.....	8,194	1	-	-	1	2	12	59	168	562	2,103	5,286	-
2002.....	8,650	3	-	-	1	4	20	63	182	572	2,259	5,546	-
2001.....	8,943	-	-	-	-	1	14	68	187	592	2,365	5,716	-
2000.....	9,036	2	-	-	1	2	13	57	157	623	2,342	5,839	-
1999.....	9,536	1	-	-	2	3	13	64	179	681	2,458	6,135	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
Hispanic, both sexes													
2003.....	379	1	-	-	-	1	3	7	24	56	121	166	-
2002.....	416	1	-	-	2	-	2	9	20	50	136	196	-
2001.....	416	-	-	-	-	-	-	14	19	46	128	209	-
2000.....	414	-	-	-	-	-	2	9	16	63	124	200	-
1999.....	410	-	-	-	-	-	3	14	25	54	130	184	-
Hispanic, male													
2003.....	182	-	-	-	-	-	3	4	17	33	63	62	-
2002.....	175	1	-	-	2	-	1	6	12	27	65	61	-
2001.....	172	-	-	-	-	-	-	9	11	24	61	67	-
2000.....	173	-	-	-	-	-	2	7	8	39	65	52	-
1999.....	185	-	-	-	-	-	1	7	16	31	60	70	-
Hispanic, female													
2003.....	197	1	-	-	-	1	-	3	7	23	58	104	-
2002.....	241	-	-	-	-	-	1	3	8	23	71	135	-
2001.....	244	-	-	-	-	-	-	5	8	22	67	142	-
2000.....	241	-	-	-	-	-	-	2	8	24	59	148	-
1999.....	225	-	-	-	-	-	2	7	9	23	70	114	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
Non-Hispanic 1/, both sexes													
2003.....	12,627	2	-	-	3	6	44	182	440	1,152	3,696	7,102	-
2002.....	13,369	3	-	-	1	9	48	173	466	1,248	3,911	7,509	1
2001.....	13,626	1	1	1	1	3	40	172	452	1,285	4,011	7,658	1
2000.....	13,945	2	-	-	3	8	46	167	434	1,357	4,145	7,783	-
1999.....	14,519	2	-	-	2	8	42	169	457	1,416	4,242	8,181	-
Non-Hispanic 1/, male													
2003.....	4,660	2	-	-	2	5	32	126	279	615	1,663	1,936	-
2002.....	4,984	-	-	-	-	5	29	113	292	700	1,733	2,111	1
2001.....	4,952	1	1	1	1	2	26	109	274	718	1,721	2,097	1
2000.....	5,166	-	-	-	2	6	33	112	285	759	1,865	2,104	-
1999.....	5,236	1	-	-	-	5	31	112	287	761	1,861	2,178	-
Non-Hispanic 1/, female													
2003.....	7,967	-	-	-	1	1	12	56	161	537	2,033	5,166	-
2002.....	8,385	3	-	-	1	4	19	60	174	548	2,178	5,398	-
2001.....	8,674	-	-	-	-	1	14	63	178	567	2,290	5,561	-
2000.....	8,779	2	-	-	1	2	13	55	149	598	2,280	5,679	-
1999.....	9,283	1	-	-	2	3	11	57	170	655	2,381	6,003	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
Non-Hispanic white, both sexes													
2003.....	11,475	-	-	-	2	4	36	141	359	978	3,349	6,606	-
2002.....	12,191	2	-	-	1	6	38	121	376	1,082	3,518	7,046	1
2001.....	12,517	1	1	1	-	2	33	137	361	1,119	3,675	7,186	1
2000.....	12,740	2	-	-	3	5	34	114	349	1,181	3,782	7,270	-
1999.....	13,286	1	-	-	2	6	30	135	374	1,241	3,880	7,617	-
Non-Hispanic white, male													
2003.....	4,181	-	-	-	1	4	25	98	230	520	1,495	1,808	-
2002.....	4,508	-	-	-	-	4	23	79	237	613	1,561	1,990	1
2001.....	4,511	1	1	1	-	2	21	85	225	631	1,570	1,973	1
2000.....	4,688	-	-	-	2	5	26	79	238	673	1,716	1,949	-
1999.....	4,727	1	-	-	-	4	23	93	235	667	1,690	2,014	-
Non-Hispanic white, female													
2003.....	7,294	-	-	-	1	-	11	43	129	458	1,854	4,798	-
2002.....	7,683	2	-	-	1	2	15	42	139	469	1,957	5,056	-
2001.....	8,006	-	-	-	-	-	12	52	136	488	2,105	5,213	-
2000.....	8,052	2	-	-	1	-	8	35	111	508	2,066	5,321	-
1999.....	8,559	-	-	-	2	2	7	42	139	574	2,190	5,603	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
Non-Hispanic black, both sexes													
2003.....	981	2	-	-	1	2	5	38	72	152	296	413	-
2002.....	1,009	1	-	-	-	3	9	46	77	134	349	390	-
2001.....	957	-	-	-	1	1	6	31	84	145	285	404	-
2000.....	1,049	-	-	-	-	3	12	47	79	147	320	441	-
1999.....	1,061	1	-	-	-	2	9	29	76	150	312	482	-
Non-Hispanic black, male													
2003.....	400	2	-	-	1	1	4	26	44	83	140	99	-
2002.....	409	-	-	-	-	1	6	30	49	70	153	100	-
2001.....	365	-	-	-	1	-	4	21	43	74	124	98	-
2000.....	412	-	-	-	-	1	7	28	43	74	131	128	-
1999.....	430	-	-	-	-	1	6	15	46	80	142	140	-
Non-Hispanic black, female													
2003.....	581	-	-	-	-	1	1	12	28	69	156	314	-
2002.....	600	1	-	-	-	2	3	16	28	64	196	290	-
2001.....	592	-	-	-	-	1	2	10	41	71	161	306	-
2000.....	637	-	-	-	-	2	5	19	36	73	189	313	-
1999.....	631	1	-	-	-	1	3	14	30	70	170	342	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Atherosclerosis (I70)													
Origin not stated 2/, both sexes													
2003.....	47	-	-	-	-	-	-	1	2	3	20	20	1
2002.....	36	-	-	-	-	-	-	-	1	2	15	18	-
2001.....	44	-	-	-	-	-	-	-	2	4	17	21	-
2000.....	34	-	-	-	-	-	-	-	1	3	8	22	-
1999.....	50	-	-	-	-	-	-	2	4	6	13	25	-
Origin not stated 2/, male													
2003.....	17	-	-	-	-	-	-	1	2	1	8	4	1
2002.....	12	-	-	-	-	-	-	-	1	1	5	5	-
2001.....	19	-	-	-	-	-	-	-	1	1	9	8	-
2000.....	18	-	-	-	-	-	-	-	1	2	5	10	-
1999.....	22	-	-	-	-	-	-	2	4	3	6	7	-
Origin not stated 2/, female													
2003.....	30	-	-	-	-	-	-	-	-	2	12	16	-
2002.....	24	-	-	-	-	-	-	-	-	1	10	13	-
2001.....	25	-	-	-	-	-	-	-	1	3	8	13	-
2000.....	16	-	-	-	-	-	-	-	-	1	3	12	-
1999.....	28	-	-	-	-	-	-	-	-	3	7	18	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of circulatory system (I71-I78)													
All origins, both sexes													
2003.....	24,672	11	-	12	68	151	451	1,199	2,270	4,934	8,694	6,882	-
2002.....	24,927	14	4	11	66	160	445	1,070	2,286	5,310	8,745	6,816	-
2001.....	25,318	12	4	11	61	170	498	1,024	2,195	5,345	9,032	6,965	1
2000.....	26,036	16	5	11	59	154	448	1,038	2,236	5,926	9,222	6,920	1
1999.....	25,809	12	2	11	49	141	478	971	2,210	5,808	9,264	6,863	-
All origins, male													
2003.....	13,217	3	-	7	45	103	306	815	1,517	3,092	4,732	2,597	-
2002.....	13,309	6	4	5	43	101	291	732	1,526	3,307	4,690	2,604	-
2001.....	13,396	8	2	3	40	115	335	688	1,454	3,348	4,839	2,563	1
2000.....	14,084	6	2	7	33	103	276	699	1,530	3,771	5,019	2,637	1
1999.....	13,794	9	-	7	32	102	296	657	1,519	3,681	5,015	2,476	-
All origins, female													
2003.....	11,455	8	-	5	23	48	145	384	753	1,842	3,962	4,285	-
2002.....	11,618	8	-	6	23	59	154	338	760	2,003	4,055	4,212	-
2001.....	11,922	4	2	8	21	55	163	336	741	1,997	4,193	4,402	-
2000.....	11,952	10	3	4	26	51	172	339	706	2,155	4,203	4,283	-
1999.....	12,015	3	2	4	17	39	182	314	691	2,127	4,249	4,387	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of circulatory system (I71-I78)													
Hispanic, both sexes													
2003.....	878	2	-	1	9	22	39	61	96	172	271	205	-
2002.....	847	1	-	1	12	22	44	62	117	175	246	167	-
2001.....	754	2	1	2	7	19	37	73	88	156	216	152	1
2000.....	818	3	1	1	9	24	38	56	99	217	215	155	-
1999.....	729	2	-	2	9	12	33	38	94	170	211	158	-
Hispanic, male													
2003.....	510	1	-	-	5	15	28	42	73	103	153	90	-
2002.....	463	-	-	-	8	14	31	41	66	104	138	61	-
2001.....	424	1	1	-	4	14	26	46	55	100	119	57	1
2000.....	452	1	-	1	4	14	22	33	64	122	126	65	-
1999.....	416	1	-	-	7	6	22	25	59	101	119	76	-
Hispanic, female													
2003.....	368	1	-	1	4	7	11	19	23	69	118	115	-
2002.....	384	1	-	1	4	8	13	21	51	71	108	106	-
2001.....	330	1	-	2	3	5	11	27	33	56	97	95	-
2000.....	366	2	1	-	5	10	16	23	35	95	89	90	-
1999.....	313	1	-	2	2	6	11	13	35	69	92	82	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of circulatory system (I71-I78)													
Non-Hispanic 1/, both sexes													
2003.....	23,739	9	-	11	59	129	410	1,134	2,171	4,746	8,403	6,667	-
2002.....	24,005	13	4	10	54	137	396	1,006	2,159	5,120	8,475	6,631	-
2001.....	24,489	10	3	9	54	149	460	946	2,099	5,170	8,795	6,794	-
2000.....	25,147	13	4	10	50	129	405	979	2,129	5,695	8,985	6,747	1
1999.....	25,023	10	2	9	40	129	445	932	2,109	5,619	9,037	6,691	-
Non-Hispanic 1/, male													
2003.....	12,679	2	-	7	40	88	277	772	1,441	2,979	4,569	2,504	-
2002.....	12,809	6	4	5	35	86	257	690	1,455	3,194	4,540	2,537	-
2001.....	12,926	7	1	3	36	99	308	638	1,392	3,234	4,708	2,500	-
2000.....	13,593	5	2	6	29	89	249	663	1,461	3,640	4,883	2,565	1
1999.....	13,351	8	-	7	25	96	274	632	1,456	3,567	4,889	2,397	-
Non-Hispanic 1/, female													
2003.....	11,060	7	-	4	19	41	133	362	730	1,767	3,834	4,163	-
2002.....	11,196	7	-	5	19	51	139	316	704	1,926	3,935	4,094	-
2001.....	11,563	3	2	6	18	50	152	308	707	1,936	4,087	4,294	-
2000.....	11,554	8	2	4	21	40	156	316	668	2,055	4,102	4,182	-
1999.....	11,672	2	2	2	15	33	171	300	653	2,052	4,148	4,294	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of circulatory system (I71-I78)													
Non-Hispanic white, both sexes													
2003.....	20,756	4	-	7	41	85	266	798	1,787	4,158	7,567	6,043	-
2002.....	21,051	10	4	4	37	86	272	702	1,758	4,485	7,684	6,009	-
2001.....	21,439	8	2	8	40	108	303	682	1,702	4,539	7,908	6,139	-
2000.....	22,113	11	3	6	35	88	269	699	1,734	5,046	8,133	6,089	-
1999.....	22,191	8	2	7	33	94	323	690	1,773	5,000	8,210	6,051	-
Non-Hispanic white, male													
2003.....	11,188	1	-	4	30	55	189	567	1,206	2,663	4,159	2,314	-
2002.....	11,337	5	4	2	23	54	180	495	1,198	2,850	4,189	2,337	-
2001.....	11,437	5	1	3	27	74	202	462	1,158	2,897	4,303	2,305	-
2000.....	12,138	3	2	5	22	63	167	494	1,225	3,301	4,492	2,364	-
1999.....	12,014	6	-	5	20	71	204	478	1,261	3,253	4,514	2,202	-
Non-Hispanic white, female													
2003.....	9,568	3	-	3	11	30	77	231	581	1,495	3,408	3,729	-
2002.....	9,714	5	-	2	14	32	92	207	560	1,635	3,495	3,672	-
2001.....	10,002	3	1	5	13	34	101	220	544	1,642	3,605	3,834	-
2000.....	9,975	8	1	1	13	25	102	205	509	1,745	3,641	3,725	-
1999.....	10,177	2	2	2	13	23	119	212	512	1,747	3,696	3,849	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of circulatory system (I71-I78)													
Non-Hispanic black, both sexes													
2003.....	2,500	5	-	3	13	37	132	296	325	492	683	514	-
2002.....	2,495	3	-	5	12	46	112	272	350	521	653	521	-
2001.....	2,606	2	1	1	12	39	141	227	348	526	746	563	-
2000.....	2,569	2	-	4	13	35	120	232	346	540	710	566	1
1999.....	2,416	2	-	-	7	30	112	212	290	517	689	557	-
Non-Hispanic black, male													
2003.....	1,202	1	-	2	6	29	81	181	200	252	317	133	-
2002.....	1,196	1	-	2	7	30	71	171	223	268	269	154	-
2001.....	1,214	2	-	-	7	24	93	149	197	272	324	146	-
2000.....	1,193	2	-	1	6	22	73	144	198	279	314	153	1
1999.....	1,102	2	-	-	5	23	61	135	165	266	290	155	-
Non-Hispanic black, female													
2003.....	1,298	4	-	1	7	8	51	115	125	240	366	381	-
2002.....	1,299	2	-	3	5	16	41	101	127	253	384	367	-
2001.....	1,392	-	1	1	5	15	48	78	151	254	422	417	-
2000.....	1,376	-	-	3	7	13	47	88	148	261	396	413	-
1999.....	1,314	-	-	-	2	7	51	77	125	251	399	402	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of circulatory system (I71-I78)

Origin not stated 2/, both sexes

2003.....	55	-	-	-	-	-	2	4	3	16	20	10	-
2002.....	75	-	-	-	-	1	5	2	10	15	24	18	-
2001.....	75	-	-	-	-	2	1	5	8	19	21	19	-
2000.....	71	-	-	-	-	1	5	3	8	14	22	18	-
1999.....	57	-	-	-	-	-	-	1	7	19	16	14	-

Origin not stated 2/, male

2003.....	28	-	-	-	-	-	1	1	3	10	10	3	-
2002.....	37	-	-	-	-	1	3	1	5	9	12	6	-
2001.....	46	-	-	-	-	2	1	4	7	14	12	6	-
2000.....	39	-	-	-	-	-	5	3	5	9	10	7	-
1999.....	27	-	-	-	-	-	-	-	4	13	7	3	-

Origin not stated 2/, female

2003.....	27	-	-	-	-	-	1	3	-	6	10	7	-
2002.....	38	-	-	-	-	-	2	1	5	6	12	12	-
2001.....	29	-	-	-	-	-	-	1	1	5	9	13	-
2000.....	32	-	-	-	-	1	-	-	3	5	12	11	-
1999.....	30	-	-	-	-	-	-	1	3	6	9	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
All origins, both sexes													
2003.....	14,810	1	-	3	41	112	316	820	1,477	3,271	5,471	3,298	-
2002.....	14,818	2	-	5	39	110	305	710	1,460	3,482	5,518	3,187	-
2001.....	15,234	-	2	4	43	117	332	661	1,394	3,517	5,751	3,412	1
2000.....	15,810	-	-	5	36	107	303	662	1,493	4,012	5,835	3,357	-
1999.....	15,807	1	1	3	31	93	329	638	1,437	4,001	5,936	3,337	-
All origins, male													
2003.....	8,976	-	-	2	31	83	246	616	1,107	2,174	3,192	1,525	-
2002.....	9,013	-	-	3	32	82	229	540	1,092	2,369	3,185	1,481	-
2001.....	9,114	-	2	3	31	90	264	506	1,031	2,382	3,313	1,491	1
2000.....	9,661	-	-	3	23	78	219	492	1,111	2,730	3,443	1,562	-
1999.....	9,615	1	-	2	20	78	232	486	1,088	2,712	3,501	1,495	-
All origins, female													
2003.....	5,834	1	-	1	10	29	70	204	370	1,097	2,279	1,773	-
2002.....	5,805	2	-	2	7	28	76	170	368	1,113	2,333	1,706	-
2001.....	6,120	-	-	1	12	27	68	155	363	1,135	2,438	1,921	-
2000.....	6,149	-	-	2	13	29	84	170	382	1,282	2,392	1,795	-
1999.....	6,192	-	1	1	11	15	97	152	349	1,289	2,435	1,842	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Hispanic, both sexes													
2003.....	515	-	-	-	6	16	25	36	64	102	171	95	-
2002.....	431	-	-	-	8	16	24	36	75	89	127	56	-
2001.....	403	-	1	-	4	14	24	41	47	93	127	51	1
2000.....	465	-	-	1	8	15	21	41	58	121	137	63	-
1999.....	384	-	-	1	4	6	23	27	55	95	115	58	-
Hispanic, male													
2003.....	333	-	-	-	4	12	19	27	51	68	100	52	-
2002.....	284	-	-	-	5	12	20	29	51	63	78	26	-
2001.....	279	-	1	-	2	14	20	31	33	71	78	28	1
2000.....	294	-	-	1	4	11	14	28	42	76	86	32	-
1999.....	256	-	-	-	3	4	17	22	41	67	65	37	-
Hispanic, female													
2003.....	182	-	-	-	2	4	6	9	13	34	71	43	-
2002.....	147	-	-	-	3	4	4	7	24	26	49	30	-
2001.....	124	-	-	-	2	-	4	10	14	22	49	23	-
2000.....	171	-	-	-	4	4	7	13	16	45	51	31	-
1999.....	128	-	-	1	1	2	6	5	14	28	50	21	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Non-Hispanic 1/, both sexes													
2003.....	14,259	1	-	3	35	96	289	781	1,411	3,159	5,286	3,198	-
2002.....	14,341	2	-	5	31	93	277	672	1,379	3,384	5,373	3,125	-
2001.....	14,782	-	1	4	39	102	307	617	1,341	3,408	5,610	3,353	-
2000.....	15,296	-	-	4	28	91	277	618	1,431	3,881	5,682	3,284	-
1999.....	15,393	1	1	2	27	87	306	611	1,378	3,893	5,811	3,276	-
Non-Hispanic 1/, male													
2003.....	8,622	-	-	2	27	71	226	588	1,054	2,098	3,085	1,471	-
2002.....	8,701	-	-	3	27	69	207	510	1,036	2,299	3,099	1,451	-
2001.....	8,798	-	1	3	29	75	243	472	992	2,299	3,225	1,459	-
2000.....	9,338	-	-	2	19	67	200	461	1,066	2,648	3,349	1,526	-
1999.....	9,342	1	-	2	17	74	215	464	1,045	2,635	3,432	1,457	-
Non-Hispanic 1/, female													
2003.....	5,637	1	-	1	8	25	63	193	357	1,061	2,201	1,727	-
2002.....	5,640	2	-	2	4	24	70	162	343	1,085	2,274	1,674	-
2001.....	5,984	-	-	1	10	27	64	145	349	1,109	2,385	1,894	-
2000.....	5,958	-	-	2	9	24	77	157	365	1,233	2,333	1,758	-
1999.....	6,051	-	1	-	10	13	91	147	333	1,258	2,379	1,819	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Non-Hispanic white, both sexes													
2003.....	12,712	-	-	1	27	61	188	548	1,178	2,844	4,883	2,982	-
2002.....	12,831	2	-	2	23	57	192	457	1,156	3,057	4,988	2,897	-
2001.....	13,217	-	1	4	28	77	203	435	1,117	3,079	5,155	3,118	-
2000.....	13,722	-	-	1	18	61	182	454	1,201	3,526	5,250	3,029	-
1999.....	13,934	1	1	1	23	64	217	440	1,195	3,548	5,407	3,037	-
Non-Hispanic white, male													
2003.....	7,731	-	-	-	21	42	155	428	897	1,926	2,877	1,385	-
2002.....	7,834	-	-	1	19	43	142	355	875	2,105	2,917	1,377	-
2001.....	7,924	-	1	3	21	58	163	342	841	2,109	3,002	1,384	-
2000.....	8,488	-	-	1	14	45	134	355	914	2,453	3,136	1,436	-
1999.....	8,551	1	-	1	14	54	155	340	928	2,451	3,243	1,364	-
Non-Hispanic white, female													
2003.....	4,981	-	-	1	6	19	33	120	281	918	2,006	1,597	-
2002.....	4,997	2	-	1	4	14	50	102	281	952	2,071	1,520	-
2001.....	5,293	-	-	1	7	19	40	93	276	970	2,153	1,734	-
2000.....	5,234	-	-	-	4	16	48	99	287	1,073	2,114	1,593	-
1999.....	5,383	-	1	-	9	10	62	100	267	1,097	2,164	1,673	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Non-Hispanic black, both sexes													
2003.....	1,220	1	-	1	4	29	94	202	196	242	292	159	-
2002.....	1,200	-	-	3	5	33	79	192	192	244	285	167	-
2001.....	1,238	-	-	-	10	23	91	154	189	240	352	179	-
2000.....	1,262	-	-	3	9	25	86	135	196	279	331	198	-
1999.....	1,172	-	-	-	4	21	80	152	152	271	303	189	-
Non-Hispanic black, male													
2003.....	678	-	-	1	3	26	65	140	132	124	138	49	-
2002.....	664	-	-	2	5	24	61	138	138	132	119	45	-
2001.....	659	-	-	-	7	16	69	108	124	134	158	43	-
2000.....	661	-	-	1	5	19	60	89	123	151	153	60	-
1999.....	615	-	-	-	3	19	52	109	96	147	124	65	-
Non-Hispanic black, female													
2003.....	542	1	-	-	1	3	29	62	64	118	154	110	-
2002.....	536	-	-	1	-	9	18	54	54	112	166	122	-
2001.....	579	-	-	-	3	7	22	46	65	106	194	136	-
2000.....	601	-	-	2	4	6	26	46	73	128	178	138	-
1999.....	557	-	-	-	1	2	28	43	56	124	179	124	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Aortic aneurysm and dissection (I71)													
Origin not stated 2/, both sexes													
2003.....	36	-	-	-	-	-	2	3	2	10	14	5	-
2002.....	46	-	-	-	-	1	4	2	6	9	18	6	-
2001.....	49	-	-	-	-	1	1	3	6	16	14	8	-
2000.....	49	-	-	-	-	1	5	3	4	10	16	10	-
1999.....	30	-	-	-	-	-	-	-	4	13	10	3	-
Origin not stated 2/, male													
2003.....	21	-	-	-	-	-	1	1	2	8	7	2	-
2002.....	28	-	-	-	-	1	2	1	5	7	8	4	-
2001.....	37	-	-	-	-	1	1	3	6	12	10	4	-
2000.....	29	-	-	-	-	-	5	3	3	6	8	4	-
1999.....	17	-	-	-	-	-	-	-	2	10	4	1	-
Origin not stated 2/, female													
2003.....	15	-	-	-	-	-	1	2	-	2	7	3	-
2002.....	18	-	-	-	-	-	2	1	1	2	10	2	-
2001.....	12	-	-	-	-	-	-	-	-	4	4	4	-
2000.....	20	-	-	-	-	1	-	-	1	4	8	6	-
1999.....	13	-	-	-	-	-	-	-	2	3	6	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
All origins, both sexes													
2003.....	9,862	10	-	9	27	39	135	379	793	1,663	3,223	3,584	-
2002.....	10,109	12	4	6	27	50	140	360	826	1,828	3,227	3,629	-
2001.....	10,084	12	2	7	18	53	166	363	801	1,828	3,281	3,553	-
2000.....	10,226	16	5	6	23	47	145	376	743	1,914	3,387	3,563	1
1999.....	10,002	11	1	8	18	48	149	333	773	1,807	3,328	3,526	-
All origins, male													
2003.....	4,241	3	-	5	14	20	60	199	410	918	1,540	1,072	-
2002.....	4,296	6	4	2	11	19	62	192	434	938	1,505	1,123	-
2001.....	4,282	8	-	-	9	25	71	182	423	966	1,526	1,072	-
2000.....	4,423	6	2	4	10	25	57	207	419	1,041	1,576	1,075	1
1999.....	4,179	8	-	5	12	24	64	171	431	969	1,514	981	-
All origins, female													
2003.....	5,621	7	-	4	13	19	75	180	383	745	1,683	2,512	-
2002.....	5,813	6	-	4	16	31	78	168	392	890	1,722	2,506	-
2001.....	5,802	4	2	7	9	28	95	181	378	862	1,755	2,481	-
2000.....	5,803	10	3	2	13	22	88	169	324	873	1,811	2,488	-
1999.....	5,823	3	1	3	6	24	85	162	342	838	1,814	2,545	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
Hispanic, both sexes													
2003.....	363	2	-	1	3	6	14	25	32	70	100	110	-
2002.....	416	1	-	1	4	6	20	26	42	86	119	111	-
2001.....	351	2	-	2	3	5	13	32	41	63	89	101	-
2000.....	353	3	1	-	1	9	17	15	41	96	78	92	-
1999.....	345	2	-	1	5	6	10	11	39	75	96	100	-
Hispanic, male													
2003.....	177	1	-	-	1	3	9	15	22	35	53	38	-
2002.....	179	-	-	-	3	2	11	12	15	41	60	35	-
2001.....	145	1	-	-	2	-	6	15	22	29	41	29	-
2000.....	158	1	-	-	-	3	8	5	22	46	40	33	-
1999.....	160	1	-	-	4	2	5	3	18	34	54	39	-
Hispanic, female													
2003.....	186	1	-	1	2	3	5	10	10	35	47	72	-
2002.....	237	1	-	1	1	4	9	14	27	45	59	76	-
2001.....	206	1	-	2	1	5	7	17	19	34	48	72	-
2000.....	195	2	1	-	1	6	9	10	19	50	38	59	-
1999.....	185	1	-	1	1	4	5	8	21	41	42	61	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
Non-Hispanic 1/, both sexes													
2003.....	9,480	8	-	8	24	33	121	353	760	1,587	3,117	3,469	-
2002.....	9,664	11	4	5	23	44	119	334	780	1,736	3,102	3,506	-
2001.....	9,707	10	2	5	15	47	153	329	758	1,762	3,185	3,441	-
2000.....	9,851	13	4	6	22	38	128	361	698	1,814	3,303	3,463	1
1999.....	9,630	9	1	7	13	42	139	321	731	1,726	3,226	3,415	-
Non-Hispanic 1/, male													
2003.....	4,057	2	-	5	13	17	51	184	387	881	1,484	1,033	-
2002.....	4,108	6	4	2	8	17	50	180	419	895	1,441	1,086	-
2001.....	4,128	7	-	-	7	24	65	166	400	935	1,483	1,041	-
2000.....	4,255	5	2	4	10	22	49	202	395	992	1,534	1,039	1
1999.....	4,009	7	-	5	8	22	59	168	411	932	1,457	940	-
Non-Hispanic 1/, female													
2003.....	5,423	6	-	3	11	16	70	169	373	706	1,633	2,436	-
2002.....	5,556	5	-	3	15	27	69	154	361	841	1,661	2,420	-
2001.....	5,579	3	2	5	8	23	88	163	358	827	1,702	2,400	-
2000.....	5,596	8	2	2	12	16	79	159	303	822	1,769	2,424	-
1999.....	5,621	2	1	2	5	20	80	153	320	794	1,769	2,475	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
Non-Hispanic white, both sexes													
2003.....	8,044	4	-	6	14	24	78	250	609	1,314	2,684	3,061	-
2002.....	8,220	8	4	2	14	29	80	245	602	1,428	2,696	3,112	-
2001.....	8,222	8	1	4	12	31	100	247	585	1,460	2,753	3,021	-
2000.....	8,391	11	3	5	17	27	87	245	533	1,520	2,883	3,060	-
1999.....	8,257	7	1	6	10	30	106	250	578	1,452	2,803	3,014	-
Non-Hispanic white, male													
2003.....	3,457	1	-	4	9	13	34	139	309	737	1,282	929	-
2002.....	3,503	5	4	1	4	11	38	140	323	745	1,272	960	-
2001.....	3,513	5	-	-	6	16	39	120	317	788	1,301	921	-
2000.....	3,650	3	2	4	8	18	33	139	311	848	1,356	928	-
1999.....	3,463	5	-	4	6	17	49	138	333	802	1,271	838	-
Non-Hispanic white, female													
2003.....	4,587	3	-	2	5	11	44	111	300	577	1,402	2,132	-
2002.....	4,717	3	-	1	10	18	42	105	279	683	1,424	2,152	-
2001.....	4,709	3	1	4	6	15	61	127	268	672	1,452	2,100	-
2000.....	4,741	8	1	1	9	9	54	106	222	672	1,527	2,132	-
1999.....	4,794	2	1	2	4	13	57	112	245	650	1,532	2,176	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
Non-Hispanic black, both sexes													
2003.....	1,280	4	-	2	9	8	38	94	129	250	391	355	-
2002.....	1,295	3	-	2	7	13	33	80	158	277	368	354	-
2001.....	1,368	2	1	1	2	16	50	73	159	286	394	384	-
2000.....	1,307	2	-	1	4	10	34	97	150	261	379	368	1
1999.....	1,244	2	-	-	3	9	32	60	138	246	386	368	-
Non-Hispanic black, male													
2003.....	524	1	-	1	3	3	16	41	68	128	179	84	-
2002.....	532	1	-	-	2	6	10	33	85	136	150	109	-
2001.....	555	2	-	-	-	8	24	41	73	138	166	103	-
2000.....	532	2	-	-	1	3	13	55	75	128	161	93	1
1999.....	487	2	-	-	2	4	9	26	69	119	166	90	-
Non-Hispanic black, female													
2003.....	756	3	-	1	6	5	22	53	61	122	212	271	-
2002.....	763	2	-	2	5	7	23	47	73	141	218	245	-
2001.....	813	-	1	1	2	8	26	32	86	148	228	281	-
2000.....	775	-	-	1	3	7	21	42	75	133	218	275	-
1999.....	757	-	-	-	1	5	23	34	69	127	220	278	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Origin not stated 2/, both sexes

2003.....	19	-	-	-	-	-	-	1	1	6	6	5	-
2002.....	29	-	-	-	-	-	1	-	4	6	6	12	-
2001.....	26	-	-	-	-	1	-	2	2	3	7	11	-
2000.....	22	-	-	-	-	-	-	-	4	4	6	8	-
1999.....	27	-	-	-	-	-	-	1	3	6	6	11	-

Origin not stated 2/, male

2003.....	7	-	-	-	-	-	-	-	1	2	3	1	-
2002.....	9	-	-	-	-	-	1	-	-	2	4	2	-
2001.....	9	-	-	-	-	1	-	1	1	2	2	2	-
2000.....	10	-	-	-	-	-	-	-	2	3	2	3	-
1999.....	10	-	-	-	-	-	-	-	2	3	3	2	-

Origin not stated 2/, female

2003.....	12	-	-	-	-	-	-	1	-	4	3	4	-
2002.....	20	-	-	-	-	-	-	-	4	4	2	10	-
2001.....	17	-	-	-	-	-	-	1	1	1	5	9	-
2000.....	12	-	-	-	-	-	-	-	2	1	4	5	-
1999.....	17	-	-	-	-	-	-	1	1	3	3	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of circulatory system (I80-I99)													
All origins, both sexes													
2003.....	4,737	36	4	4	50	139	342	555	569	744	1,161	1,133	-
2002.....	4,711	27	4	2	35	139	305	531	595	697	1,253	1,123	-
2001.....	4,665	22	5	-	38	124	285	454	563	841	1,183	1,149	1
2000.....	4,603	27	4	12	37	133	319	515	516	795	1,161	1,084	-
1999.....	4,025	23	1	2	30	93	271	378	445	692	1,117	972	1
All origins, male													
2003.....	1,966	21	3	2	20	60	167	312	320	348	437	276	-
2002.....	1,944	15	1	-	21	69	154	284	309	323	483	285	-
2001.....	1,981	11	5	-	17	53	149	262	304	405	488	286	1
2000.....	1,981	16	2	6	16	57	190	295	264	390	472	273	-
1999.....	1,714	13	1	1	13	44	159	198	224	350	463	248	-
All origins, female													
2003.....	2,771	15	1	2	30	79	175	243	249	396	724	857	-
2002.....	2,767	12	3	2	14	70	151	247	286	374	770	838	-
2001.....	2,684	11	-	-	21	71	136	192	259	436	695	863	-
2000.....	2,622	11	2	6	21	76	129	220	252	405	689	811	-
1999.....	2,311	10	-	1	17	49	112	180	221	342	654	724	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of circulatory system (I80-I99)													
Hispanic, both sexes													
2003.....	239	11	2	1	7	15	26	41	29	32	41	34	-
2002.....	219	7	1	-	5	15	21	35	27	37	38	33	-
2001.....	219	5	-	-	3	7	15	32	16	48	41	52	-
2000.....	229	6	-	4	3	11	32	34	23	35	44	37	-
1999.....	186	2	-	1	4	6	12	34	24	42	26	35	-
Hispanic, male													
2003.....	120	5	1	1	2	5	19	27	17	18	16	9	-
2002.....	95	5	-	-	3	8	9	17	16	16	16	5	-
2001.....	104	4	-	-	2	4	7	22	9	20	20	16	-
2000.....	116	3	-	2	1	7	20	19	13	16	22	13	-
1999.....	83	-	-	1	2	4	8	15	11	17	13	12	-
Hispanic, female													
2003.....	119	6	1	-	5	10	7	14	12	14	25	25	-
2002.....	124	2	1	-	2	7	12	18	11	21	22	28	-
2001.....	115	1	-	-	1	3	8	10	7	28	21	36	-
2000.....	113	3	-	2	2	4	12	15	10	19	22	24	-
1999.....	103	2	-	-	2	2	4	19	13	25	13	23	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of circulatory system (I80-I99)													
Non-Hispanic 1/, both sexes													
2003.....	4,485	25	2	3	43	123	315	513	537	709	1,117	1,098	-
2002.....	4,479	20	3	2	30	124	281	494	566	659	1,212	1,088	-
2001.....	4,425	17	5	-	35	117	269	418	544	788	1,136	1,095	1
2000.....	4,359	20	4	8	34	120	283	479	491	758	1,117	1,045	-
1999.....	3,825	21	1	1	26	86	256	343	420	648	1,088	934	1
Non-Hispanic 1/, male													
2003.....	1,840	16	2	1	18	54	148	284	303	328	419	267	-
2002.....	1,842	10	1	-	18	61	143	265	292	306	466	280	-
2001.....	1,869	7	5	-	15	49	141	239	293	384	466	269	1
2000.....	1,856	12	2	4	15	49	167	274	250	373	450	260	-
1999.....	1,623	13	1	-	11	40	148	182	212	331	450	235	-
Non-Hispanic 1/, female													
2003.....	2,645	9	-	2	25	69	167	229	234	381	698	831	-
2002.....	2,637	10	2	2	12	63	138	229	274	353	746	808	-
2001.....	2,556	10	-	-	20	68	128	179	251	404	670	826	-
2000.....	2,503	8	2	4	19	71	116	205	241	385	667	785	-
1999.....	2,202	8	-	1	15	46	108	161	208	317	638	699	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of circulatory system (I80-I99)													
Non-Hispanic white, both sexes													
2003.....	3,607	18	2	2	23	72	198	381	413	575	952	971	-
2002.....	3,630	13	2	1	16	69	180	363	430	538	1,041	977	-
2001.....	3,617	12	3	-	18	73	175	306	414	642	987	987	-
2000.....	3,554	13	3	5	19	70	195	358	373	602	975	941	-
1999.....	3,149	17	-	1	12	57	168	243	342	524	949	835	1
Non-Hispanic white, male													
2003.....	1,449	11	2	-	9	31	110	202	236	266	349	233	-
2002.....	1,450	8	-	-	9	33	92	194	224	248	396	246	-
2001.....	1,496	4	3	-	8	30	92	182	220	313	405	239	-
2000.....	1,496	9	1	3	8	26	118	207	188	307	398	231	-
1999.....	1,310	9	-	-	7	26	95	127	180	268	388	210	-
Non-Hispanic white, female													
2003.....	2,158	7	-	2	14	41	88	179	177	309	603	738	-
2002.....	2,180	5	2	1	7	36	88	169	206	290	645	731	-
2001.....	2,121	8	-	-	10	43	83	124	194	329	582	748	-
2000.....	2,058	4	2	2	11	44	77	151	185	295	577	710	-
1999.....	1,839	8	-	1	5	31	73	116	162	256	561	625	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of circulatory system (I80-I99)													
Non-Hispanic black, both sexes													
2003.....	812	6	-	-	19	48	111	124	118	124	149	113	-
2002.....	789	7	1	1	13	54	94	120	128	113	159	99	-
2001.....	729	5	2	-	14	43	87	98	122	129	133	95	1
2000.....	751	5	1	3	14	48	86	116	109	145	130	94	-
1999.....	613	3	1	-	14	28	81	95	69	109	125	88	-
Non-Hispanic black, male													
2003.....	363	5	-	-	9	22	36	76	64	58	62	31	-
2002.....	359	2	1	-	9	27	48	63	64	53	63	29	-
2001.....	335	3	2	-	5	18	46	48	72	62	53	25	1
2000.....	333	2	1	1	7	22	48	66	56	61	43	26	-
1999.....	281	3	1	-	4	14	49	52	28	56	55	19	-
Non-Hispanic black, female													
2003.....	449	1	-	-	10	26	75	48	54	66	87	82	-
2002.....	430	5	-	1	4	27	46	57	64	60	96	70	-
2001.....	394	2	-	-	9	25	41	50	50	67	80	70	-
2000.....	418	3	-	2	7	26	38	50	53	84	87	68	-
1999.....	332	-	-	-	10	14	32	43	41	53	70	69	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other disorders of circulatory system (I80-I99)

Origin not stated 2/, both sexes													
2003.....	13	-	-	-	-	1	1	1	3	3	3	1	-
2002.....	13	-	-	-	-	-	3	2	2	1	3	2	-
2001.....	21	-	-	-	-	-	1	4	3	5	6	2	-
2000.....	15	1	-	-	-	2	4	2	2	2	-	2	-
1999.....	14	-	-	-	-	1	3	1	1	2	3	3	-
Origin not stated 2/, male													
2003.....	6	-	-	-	-	1	-	1	-	2	2	-	-
2002.....	7	-	-	-	-	-	2	2	1	1	1	-	-
2001.....	8	-	-	-	-	-	1	1	2	1	2	1	-
2000.....	9	1	-	-	-	1	3	2	1	1	-	-	-
1999.....	8	-	-	-	-	-	3	1	1	2	-	1	-
Origin not stated 2/, female													
2003.....	7	-	-	-	-	-	1	-	3	1	1	1	-
2002.....	6	-	-	-	-	-	1	-	1	-	2	2	-
2001.....	13	-	-	-	-	-	-	3	1	4	4	1	-
2000.....	6	-	-	-	-	1	1	-	1	1	-	2	-
1999.....	6	-	-	-	-	1	-	-	-	-	3	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J10-J18)													
All origins, both sexes													
2003.....	65,163	322	163	147	224	373	992	2,140	3,130	6,831	19,442	31,397	2
2002.....	65,681	263	110	91	167	345	971	1,918	2,987	6,847	19,984	31,995	3
2001.....	62,034	299	112	92	181	339	983	1,801	2,704	6,650	18,677	30,191	5
2000.....	65,313	289	103	87	189	364	1,068	1,774	2,879	7,189	19,821	31,547	3
1999.....	63,730	320	130	93	179	339	1,063	1,697	2,625	6,861	19,192	31,229	2
All origins, male													
2003.....	28,778	192	91	70	124	229	560	1,279	1,771	3,882	9,629	10,950	1
2002.....	28,918	162	63	38	97	190	546	1,138	1,738	3,850	9,895	11,199	2
2001.....	27,342	167	57	47	106	191	556	1,099	1,587	3,732	9,294	10,502	4
2000.....	28,658	163	54	48	102	204	630	1,066	1,680	4,013	9,751	10,945	2
1999.....	27,718	171	66	53	90	197	592	1,059	1,462	3,893	9,458	10,676	1
All origins, female													
2003.....	36,385	130	72	77	100	144	432	861	1,359	2,949	9,813	20,447	1
2002.....	36,763	101	47	53	70	155	425	780	1,249	2,997	10,089	20,796	1
2001.....	34,692	132	55	45	75	148	427	702	1,117	2,918	9,383	19,689	1
2000.....	36,655	126	49	39	87	160	438	708	1,199	3,176	10,070	20,602	1
1999.....	36,012	149	64	40	89	142	471	638	1,163	2,968	9,734	20,553	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J10-J18)													
Hispanic, both sexes													
2003.....	2,948	66	37	22	36	54	74	151	224	428	860	996	-
2002.....	2,824	48	17	14	19	43	77	124	207	399	855	1,021	-
2001.....	2,722	57	32	10	18	23	73	121	168	402	761	1,056	1
2000.....	2,625	60	17	21	21	48	82	120	191	405	690	970	-
1999.....	2,246	64	17	10	25	24	74	104	155	328	569	876	-
Hispanic, male													
2003.....	1,429	40	24	12	20	35	41	100	115	235	432	375	-
2002.....	1,398	33	11	6	11	26	52	86	130	229	422	392	-
2001.....	1,309	31	16	6	15	14	48	85	99	204	388	402	1
2000.....	1,303	30	5	10	10	27	56	82	119	218	367	379	-
1999.....	1,092	40	10	6	11	15	54	71	80	187	284	334	-
Hispanic, female													
2003.....	1,519	26	13	10	16	19	33	51	109	193	428	621	-
2002.....	1,426	15	6	8	8	17	25	38	77	170	433	629	-
2001.....	1,413	26	16	4	3	9	25	36	69	198	373	654	-
2000.....	1,322	30	12	11	11	21	26	38	72	187	323	591	-
1999.....	1,154	24	7	4	14	9	20	33	75	141	285	542	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J10-J18)													
Non-Hispanic 1/, both sexes													
2003.....	62,042	254	126	124	188	319	917	1,975	2,883	6,373	18,544	30,337	2
2002.....	62,657	213	92	76	147	300	889	1,783	2,773	6,424	19,059	30,898	3
2001.....	59,103	240	79	82	163	312	906	1,669	2,516	6,209	17,863	29,062	2
2000.....	62,440	228	84	66	167	314	974	1,639	2,674	6,750	19,051	30,492	1
1999.....	61,218	255	110	83	153	315	979	1,577	2,446	6,499	18,553	30,247	1
Non-Hispanic 1/, male													
2003.....	27,255	150	67	58	104	194	518	1,167	1,641	3,626	9,178	10,551	1
2002.....	27,417	128	52	32	85	162	491	1,043	1,603	3,600	9,437	10,782	2
2001.....	25,926	135	41	41	91	175	504	1,007	1,475	3,500	8,881	10,075	1
2000.....	27,218	132	47	38	91	175	566	972	1,549	3,772	9,339	10,536	1
1999.....	26,489	131	55	47	79	182	531	974	1,367	3,687	9,138	10,298	-
Non-Hispanic 1/, female													
2003.....	34,787	104	59	66	84	125	399	808	1,242	2,747	9,366	19,786	1
2002.....	35,240	85	40	44	62	138	398	740	1,170	2,824	9,622	20,116	1
2001.....	33,177	105	38	41	72	137	402	662	1,041	2,709	8,982	18,987	1
2000.....	35,222	96	37	28	76	139	408	667	1,125	2,978	9,712	19,956	-
1999.....	34,729	124	55	36	74	133	448	603	1,079	2,812	9,415	19,949	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J10-J18)													
Non-Hispanic white, both sexes													
2003.....	54,617	143	80	86	137	206	637	1,424	2,210	5,262	16,545	27,886	1
2002.....	55,419	114	46	55	103	207	617	1,329	2,152	5,419	17,001	28,373	3
2001.....	51,952	117	48	57	108	207	605	1,193	1,963	5,214	15,837	26,602	1
2000.....	55,135	114	51	49	116	213	678	1,165	2,061	5,546	17,068	28,073	1
1999.....	54,280	121	58	62	95	199	637	1,109	1,894	5,481	16,785	27,838	1
Non-Hispanic white, male													
2003.....	23,541	87	44	39	75	131	356	844	1,219	2,966	8,141	9,639	-
2002.....	23,934	73	24	20	62	111	340	768	1,228	3,050	8,402	9,854	2
2001.....	22,389	71	26	28	64	112	333	707	1,132	2,919	7,830	9,166	1
2000.....	23,609	68	24	29	68	111	394	667	1,176	3,084	8,346	9,641	1
1999.....	23,097	64	22	36	51	119	334	670	1,049	3,087	8,255	9,410	-
Non-Hispanic white, female													
2003.....	31,076	56	36	47	62	75	281	580	991	2,296	8,404	18,247	1
2002.....	31,485	41	22	35	41	96	277	561	924	2,369	8,599	18,519	1
2001.....	29,563	46	22	29	44	95	272	486	831	2,295	8,007	17,436	-
2000.....	31,526	46	27	20	48	102	284	498	885	2,462	8,722	18,432	-
1999.....	31,183	57	36	26	44	80	303	439	845	2,394	8,530	18,428	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J10-J18)													
Non-Hispanic black, both sexes													
2003.....	5,798	91	27	28	42	101	251	487	565	915	1,514	1,776	1
2002.....	5,803	86	40	18	38	80	250	399	554	837	1,584	1,917	-
2001.....	5,686	103	24	17	45	87	274	430	471	796	1,571	1,868	-
2000.....	5,920	101	29	11	44	92	254	423	531	984	1,550	1,901	-
1999.....	5,794	114	40	15	49	107	305	425	486	880	1,410	1,963	-
Non-Hispanic black, male													
2003.....	2,861	52	13	15	24	58	142	285	356	547	765	603	1
2002.....	2,734	47	22	10	20	43	136	239	338	454	789	636	-
2001.....	2,769	52	13	9	23	52	155	270	293	472	808	622	-
2000.....	2,880	55	20	7	21	58	151	271	322	562	763	650	-
1999.....	2,786	58	25	8	23	58	173	277	282	518	688	676	-
Non-Hispanic black, female													
2003.....	2,937	39	14	13	18	43	109	202	209	368	749	1,173	-
2002.....	3,069	39	18	8	18	37	114	160	216	383	795	1,281	-
2001.....	2,917	51	11	8	22	35	119	160	178	324	763	1,246	-
2000.....	3,040	46	9	4	23	34	103	152	209	422	787	1,251	-
1999.....	3,008	56	15	7	26	49	132	148	204	362	722	1,287	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza and pneumonia (J10-J18)													
Origin not stated 2/, both sexes													
2003.....	173	2	-	1	-	-	1	14	23	30	38	64	-
2002.....	200	2	1	1	1	2	5	11	7	24	70	76	-
2001.....	209	2	1	-	-	4	4	11	20	39	53	73	2
2000.....	248	1	2	-	1	2	12	15	14	34	80	85	2
1999.....	266	1	3	-	1	-	10	16	24	34	70	106	1
Origin not stated 2/, male													
2003.....	94	2	-	-	-	-	1	12	15	21	19	24	-
2002.....	103	1	-	-	1	2	3	9	5	21	36	25	-
2001.....	107	1	-	-	-	2	4	7	13	28	25	25	2
2000.....	137	1	2	-	1	2	8	12	12	23	45	30	1
1999.....	137	-	1	-	-	-	7	14	15	19	36	44	1
Origin not stated 2/, female													
2003.....	79	-	-	1	-	-	-	2	8	9	19	40	-
2002.....	97	1	1	1	-	-	2	2	2	3	34	51	-
2001.....	102	1	1	-	-	2	-	4	7	11	28	48	-
2000.....	111	-	-	-	-	-	4	3	2	11	35	55	1
1999.....	129	1	2	-	1	-	3	2	9	15	34	62	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J10-J11)													
All origins, both sexes													
2003.....	1,792	31	59	46	30	13	24	61	85	183	457	803	-
2002.....	727	7	5	12	7	4	8	17	30	56	177	404	-
2001.....	257	7	6	12	7	10	6	15	21	21	56	96	-
2000.....	1,765	9	10	11	13	13	22	48	93	199	492	855	-
1999.....	1,665	13	12	11	11	9	16	26	71	164	451	881	-
All origins, male													
2003.....	699	18	33	19	12	9	10	33	42	89	189	245	-
2002.....	289	5	3	2	4	2	4	12	20	32	81	124	-
2001.....	101	6	2	4	6	6	2	9	9	8	20	29	-
2000.....	664	5	4	5	6	5	14	25	44	106	196	254	-
1999.....	602	6	9	5	2	5	9	15	37	74	202	238	-
All origins, female													
2003.....	1,093	13	26	27	18	4	14	28	43	94	268	558	-
2002.....	438	2	2	10	3	2	4	5	10	24	96	280	-
2001.....	156	1	4	8	1	4	4	6	12	13	36	67	-
2000.....	1,101	4	6	6	7	8	8	23	49	93	296	601	-
1999.....	1,063	7	3	6	9	4	7	11	34	90	249	643	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J10-J11)													
Hispanic, both sexes													
2003.....	75	10	14	9	4	2	3	2	6	3	12	10	-
2002.....	19	2	-	4	-	1	1	1	-	2	3	5	-
2001.....	13	-	3	1	1	-	1	1	-	1	5	-	-
2000.....	47	2	1	3	1	4	2	2	2	4	10	16	-
1999.....	45	4	3	-	2	-	2	2	5	8	9	10	-
Hispanic, male													
2003.....	40	7	10	3	1	1	2	-	2	3	7	4	-
2002.....	10	2	-	2	-	1	1	1	-	1	1	1	-
2001.....	7	-	2	1	1	-	-	1	-	-	2	-	-
2000.....	21	-	1	1	1	1	1	1	2	2	5	6	-
1999.....	21	-	3	-	-	-	2	2	4	7	2	1	-
Hispanic, female													
2003.....	35	3	4	6	3	1	1	2	4	-	5	6	-
2002.....	9	-	-	2	-	-	-	-	-	1	2	4	-
2001.....	6	-	1	-	-	-	1	-	-	1	3	-	-
2000.....	26	2	-	2	-	3	1	1	-	2	5	10	-
1999.....	24	4	-	-	2	-	-	-	1	1	7	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J10-J11)													
Non-Hispanic 1/, both sexes													
2003.....	1,714	21	45	37	26	11	21	59	78	180	443	793	-
2002.....	704	5	5	8	7	3	7	16	30	54	173	396	-
2001.....	244	7	3	11	6	10	5	14	21	20	51	96	-
2000.....	1,716	7	9	8	12	9	20	46	91	195	481	838	-
1999.....	1,612	9	9	11	9	9	14	24	66	154	439	868	-
Non-Hispanic 1/, male													
2003.....	657	11	23	16	11	8	8	33	39	86	181	241	-
2002.....	277	3	3	-	4	1	3	11	20	31	79	122	-
2001.....	94	6	-	3	5	6	2	8	9	8	18	29	-
2000.....	642	5	3	4	5	4	13	24	42	104	190	248	-
1999.....	580	6	6	5	2	5	7	13	33	66	200	237	-
Non-Hispanic 1/, female													
2003.....	1,057	10	22	21	15	3	13	26	39	94	262	552	-
2002.....	427	2	2	8	3	2	4	5	10	23	94	274	-
2001.....	150	1	3	8	1	4	3	6	12	12	33	67	-
2000.....	1,074	2	6	4	7	5	7	22	49	91	291	590	-
1999.....	1,032	3	3	6	7	4	7	11	33	88	239	631	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J10-J11)													
Non-Hispanic white, both sexes													
2003.....	1,578	13	29	22	16	8	17	47	69	167	426	764	-
2002.....	663	3	2	6	5	3	7	15	25	45	168	384	-
2001.....	216	7	2	6	4	7	3	14	17	18	47	91	-
2000.....	1,613	5	8	6	11	7	16	38	85	176	458	803	-
1999.....	1,522	3	5	10	8	8	8	20	60	139	417	844	-
Non-Hispanic white, male													
2003.....	599	7	15	9	6	7	4	27	33	81	175	235	-
2002.....	258	1	1	-	2	1	3	11	16	27	76	120	-
2001.....	81	6	-	2	3	3	2	8	6	6	18	27	-
2000.....	595	5	3	3	4	2	9	18	39	97	181	234	-
1999.....	547	1	3	5	2	5	4	11	32	59	192	233	-
Non-Hispanic white, female													
2003.....	979	6	14	13	10	1	13	20	36	86	251	529	-
2002.....	405	2	1	6	3	2	4	4	9	18	92	264	-
2001.....	135	1	2	4	1	4	1	6	11	12	29	64	-
2000.....	1,018	-	5	3	7	5	7	20	46	79	277	569	-
1999.....	975	2	2	5	6	3	4	9	28	80	225	611	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J10-J11)													
Non-Hispanic black, both sexes													
2003.....	95	7	6	10	7	3	1	12	7	10	12	20	-
2002.....	30	1	3	2	2	-	-	-	4	6	4	8	-
2001.....	20	-	-	2	1	2	1	-	3	2	4	5	-
2000.....	81	2	1	1	1	1	3	7	6	16	13	30	-
1999.....	65	4	1	-	1	1	5	3	5	12	15	18	-
Non-Hispanic black, male													
2003.....	36	3	3	4	3	1	1	6	5	4	3	3	-
2002.....	15	1	2	-	2	-	-	-	3	3	2	2	-
2001.....	10	-	-	-	1	2	-	-	3	2	-	2	-
2000.....	35	-	-	1	1	1	3	5	3	5	5	11	-
1999.....	24	3	1	-	-	-	2	2	1	6	6	3	-
Non-Hispanic black, female													
2003.....	59	4	3	6	4	2	-	6	2	6	9	17	-
2002.....	15	-	1	2	-	-	-	-	1	3	2	6	-
2001.....	10	-	-	2	-	-	1	-	-	-	4	3	-
2000.....	46	2	1	-	-	-	-	2	3	11	8	19	-
1999.....	41	1	-	-	1	1	3	1	4	6	9	15	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Influenza (J10-J11)													
Origin not stated 2/, both sexes													
2003.....	3	-	-	-	-	-	-	-	1	-	2	-	-
2002.....	4	-	-	-	-	-	-	-	-	-	1	3	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	1	1	-
1999.....	8	-	-	-	-	-	-	-	-	2	3	3	-
Origin not stated 2/, male													
2003.....	2	-	-	-	-	-	-	-	1	-	1	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	1	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-	-
Origin not stated 2/, female													
2003.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	2	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	1	-
1999.....	7	-	-	-	-	-	-	-	-	1	3	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
All origins, both sexes													
2003.....	63,371	291	104	101	194	360	968	2,079	3,045	6,648	18,985	30,594	2
2002.....	64,954	256	105	79	160	341	963	1,901	2,957	6,791	19,807	31,591	3
2001.....	61,777	292	106	80	174	329	977	1,786	2,683	6,629	18,621	30,095	5
2000.....	63,548	280	93	76	176	351	1,046	1,726	2,786	6,990	19,329	30,692	3
1999.....	62,065	307	118	82	168	330	1,047	1,671	2,554	6,697	18,741	30,348	2
All origins, male													
2003.....	28,079	174	58	51	112	220	550	1,246	1,729	3,793	9,440	10,705	1
2002.....	28,629	157	60	36	93	188	542	1,126	1,718	3,818	9,814	11,075	2
2001.....	27,241	161	55	43	100	185	554	1,090	1,578	3,724	9,274	10,473	4
2000.....	27,994	158	50	43	96	199	616	1,041	1,636	3,907	9,555	10,691	2
1999.....	27,116	165	57	48	88	192	583	1,044	1,425	3,819	9,256	10,438	1
All origins, female													
2003.....	35,292	117	46	50	82	140	418	833	1,316	2,855	9,545	19,889	1
2002.....	36,325	99	45	43	67	153	421	775	1,239	2,973	9,993	20,516	1
2001.....	34,536	131	51	37	74	144	423	696	1,105	2,905	9,347	19,622	1
2000.....	35,554	122	43	33	80	152	430	685	1,150	3,083	9,774	20,001	1
1999.....	34,949	142	61	34	80	138	464	627	1,129	2,878	9,485	19,910	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Hispanic, both sexes													
2003.....	2,873	56	23	13	32	52	71	149	218	425	848	986	-
2002.....	2,805	46	17	10	19	42	76	123	207	397	852	1,016	-
2001.....	2,709	57	29	9	17	23	72	120	168	401	756	1,056	1
2000.....	2,578	58	16	18	20	44	80	118	189	401	680	954	-
1999.....	2,201	60	14	10	23	24	72	102	150	320	560	866	-
Hispanic, male													
2003.....	1,389	33	14	9	19	34	39	100	113	232	425	371	-
2002.....	1,388	31	11	4	11	25	51	85	130	228	421	391	-
2001.....	1,302	31	14	5	14	14	48	84	99	204	386	402	1
2000.....	1,282	30	4	9	9	26	55	81	117	216	362	373	-
1999.....	1,071	40	7	6	11	15	52	69	76	180	282	333	-
Hispanic, female													
2003.....	1,484	23	9	4	13	18	32	49	105	193	423	615	-
2002.....	1,417	15	6	6	8	17	25	38	77	169	431	625	-
2001.....	1,407	26	15	4	3	9	24	36	69	197	370	654	-
2000.....	1,296	28	12	9	11	18	25	37	72	185	318	581	-
1999.....	1,130	20	7	4	12	9	20	33	74	140	278	533	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Non-Hispanic 1/, both sexes													
2003.....	60,328	233	81	87	162	308	896	1,916	2,805	6,193	18,101	29,544	2
2002.....	61,953	208	87	68	140	297	882	1,767	2,743	6,370	18,886	30,502	3
2001.....	58,859	233	76	71	157	302	901	1,655	2,495	6,189	17,812	28,966	2
2000.....	60,724	221	75	58	155	305	954	1,593	2,583	6,555	18,570	29,654	1
1999.....	59,606	246	101	72	144	306	965	1,553	2,380	6,345	18,114	29,379	1
Non-Hispanic 1/, male													
2003.....	26,598	139	44	42	93	186	510	1,134	1,602	3,540	8,997	10,310	1
2002.....	27,140	125	49	32	81	161	488	1,032	1,583	3,569	9,358	10,660	2
2001.....	25,832	129	41	38	86	169	502	999	1,466	3,492	8,863	10,046	1
2000.....	26,576	127	44	34	86	171	553	948	1,507	3,668	9,149	10,288	1
1999.....	25,909	125	49	42	77	177	524	961	1,334	3,621	8,938	10,061	-
Non-Hispanic 1/, female													
2003.....	33,730	94	37	45	69	122	386	782	1,203	2,653	9,104	19,234	1
2002.....	34,813	83	38	36	59	136	394	735	1,160	2,801	9,528	19,842	1
2001.....	33,027	104	35	33	71	133	399	656	1,029	2,697	8,949	18,920	1
2000.....	34,148	94	31	24	69	134	401	645	1,076	2,887	9,421	19,366	-
1999.....	33,697	121	52	30	67	129	441	592	1,046	2,724	9,176	19,318	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Non-Hispanic white, both sexes													
2003.....	53,039	130	51	64	121	198	620	1,377	2,141	5,095	16,119	27,122	1
2002.....	54,756	111	44	49	98	204	610	1,314	2,127	5,374	16,833	27,989	3
2001.....	51,736	110	46	51	104	200	602	1,179	1,946	5,196	15,790	26,511	1
2000.....	53,522	109	43	43	105	206	662	1,127	1,976	5,370	16,610	27,270	1
1999.....	52,758	118	53	52	87	191	629	1,089	1,834	5,342	16,368	26,994	1
Non-Hispanic white, male													
2003.....	22,942	80	29	30	69	124	352	817	1,186	2,885	7,966	9,404	-
2002.....	23,676	72	23	20	60	110	337	757	1,212	3,023	8,326	9,734	2
2001.....	22,308	65	26	26	61	109	331	699	1,126	2,913	7,812	9,139	1
2000.....	23,014	63	21	26	64	109	385	649	1,137	2,987	8,165	9,407	1
1999.....	22,550	63	19	31	49	114	330	659	1,017	3,028	8,063	9,177	-
Non-Hispanic white, female													
2003.....	30,097	50	22	34	52	74	268	560	955	2,210	8,153	17,718	1
2002.....	31,080	39	21	29	38	94	273	557	915	2,351	8,507	18,255	1
2001.....	29,428	45	20	25	43	91	271	480	820	2,283	7,978	17,372	-
2000.....	30,508	46	22	17	41	97	277	478	839	2,383	8,445	17,863	-
1999.....	30,208	55	34	21	38	77	299	430	817	2,314	8,305	17,817	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Non-Hispanic black, both sexes													
2003.....	5,703	84	21	18	35	98	250	475	558	905	1,502	1,756	1
2002.....	5,773	85	37	16	36	80	250	399	550	831	1,580	1,909	-
2001.....	5,666	103	24	15	44	85	273	430	468	794	1,567	1,863	-
2000.....	5,839	99	28	10	43	91	251	416	525	968	1,537	1,871	-
1999.....	5,729	110	39	15	48	106	300	422	481	868	1,395	1,945	-
Non-Hispanic black, male													
2003.....	2,825	49	10	11	21	57	141	279	351	543	762	600	1
2002.....	2,719	46	20	10	18	43	136	239	335	451	787	634	-
2001.....	2,759	52	13	9	22	50	155	270	290	470	808	620	-
2000.....	2,845	55	20	6	20	57	148	266	319	557	758	639	-
1999.....	2,762	55	24	8	23	58	171	275	281	512	682	673	-
Non-Hispanic black, female													
2003.....	2,878	35	11	7	14	41	109	196	207	362	740	1,156	-
2002.....	3,054	39	17	6	18	37	114	160	215	380	793	1,275	-
2001.....	2,907	51	11	6	22	35	118	160	178	324	759	1,243	-
2000.....	2,994	44	8	4	23	34	103	150	206	411	779	1,232	-
1999.....	2,967	55	15	7	25	48	129	147	200	356	713	1,272	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonia (J12-J18)													
Origin not stated 2/, both sexes													
2003.....	170	2	-	1	-	-	1	14	22	30	36	64	-
2002.....	196	2	1	1	1	2	5	11	7	24	69	73	-
2001.....	209	2	1	-	-	4	4	11	20	39	53	73	2
2000.....	246	1	2	-	1	2	12	15	14	34	79	84	2
1999.....	258	1	3	-	1	-	10	16	24	32	67	103	1
Origin not stated 2/, male													
2003.....	92	2	-	-	-	-	1	12	14	21	18	24	-
2002.....	101	1	-	-	1	2	3	9	5	21	35	24	-
2001.....	107	1	-	-	-	2	4	7	13	28	25	25	2
2000.....	136	1	2	-	1	2	8	12	12	23	44	30	1
1999.....	136	-	1	-	-	-	7	14	15	18	36	44	1
Origin not stated 2/, female													
2003.....	78	-	-	1	-	-	-	2	8	9	18	40	-
2002.....	95	1	1	1	-	-	2	2	2	3	34	49	-
2001.....	102	1	1	-	-	2	-	4	7	11	28	48	-
2000.....	110	-	-	-	-	-	4	3	2	11	35	54	1
1999.....	122	1	2	-	1	-	3	2	9	14	31	59	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)													
All origins, both sexes													
2003.....	421	54	28	6	6	2	11	19	21	30	80	164	-
2002.....	386	52	23	8	3	5	8	18	18	33	66	152	-
2001.....	347	50	18	3	1	2	12	14	16	28	58	145	-
2000.....	425	43	11	7	5	8	12	12	10	43	82	192	-
1999.....	540	63	25	5	5	4	11	15	21	35	107	249	-
All origins, male													
2003.....	170	35	17	2	2	1	6	8	12	17	35	35	-
2002.....	154	39	9	7	-	1	3	10	6	13	27	39	-
2001.....	117	26	8	2	1	-	7	7	6	10	20	30	-
2000.....	160	25	4	3	4	6	7	4	3	22	35	47	-
1999.....	196	36	15	2	1	3	5	8	10	17	51	48	-
All origins, female													
2003.....	251	19	11	4	4	1	5	11	9	13	45	129	-
2002.....	232	13	14	1	3	4	5	8	12	20	39	113	-
2001.....	230	24	10	1	-	2	5	7	10	18	38	115	-
2000.....	265	18	7	4	1	2	5	8	7	21	47	145	-
1999.....	344	27	10	3	4	1	6	7	11	18	56	201	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other acute lower respiratory infections (J20-J22)

Hispanic, both sexes

2003.....	31	17	6	-	-	-	1	-	1	1	2	3	-
2002.....	21	10	2	2	-	1	-	-	2	-	-	4	-
2001.....	22	8	1	-	-	1	-	2	-	2	4	4	-
2000.....	24	9	2	-	-	1	-	1	-	1	2	8	-
1999.....	21	10	3	1	-	-	-	1	-	1	-	5	-

Hispanic, male

2003.....	20	12	4	-	-	-	-	-	1	-	1	2	-
2002.....	9	5	1	1	-	-	-	-	1	-	-	1	-
2001.....	9	6	-	-	-	-	-	1	-	1	1	-	-
2000.....	15	6	1	-	-	1	-	1	-	1	2	3	-
1999.....	10	5	2	1	-	-	-	-	-	1	-	1	-

Hispanic, female

2003.....	11	5	2	-	-	-	1	-	-	1	1	1	-
2002.....	12	5	1	1	-	1	-	-	1	-	-	3	-
2001.....	13	2	1	-	-	1	-	1	-	1	3	4	-
2000.....	9	3	1	-	-	-	-	-	-	-	-	5	-
1999.....	11	5	1	-	-	-	-	1	-	-	-	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)													
Non-Hispanic 1/, both sexes													
2003.....	388	37	22	6	6	2	10	19	19	29	77	161	-
2002.....	365	42	21	6	3	4	8	18	16	33	66	148	-
2001.....	325	42	17	3	1	1	12	12	16	26	54	141	-
2000.....	398	33	9	7	5	7	12	11	10	42	79	183	-
1999.....	518	53	22	4	5	4	10	14	21	34	107	244	-
Non-Hispanic 1/, male													
2003.....	148	23	13	2	2	1	6	8	10	17	33	33	-
2002.....	145	34	8	6	-	1	3	10	5	13	27	38	-
2001.....	108	20	8	2	1	-	7	6	6	9	19	30	-
2000.....	144	19	3	3	4	5	7	3	3	21	32	44	-
1999.....	185	31	13	1	1	3	4	8	10	16	51	47	-
Non-Hispanic 1/, female													
2003.....	240	14	9	4	4	1	4	11	9	12	44	128	-
2002.....	220	8	13	-	3	3	5	8	11	20	39	110	-
2001.....	217	22	9	1	-	1	5	6	10	17	35	111	-
2000.....	254	14	6	4	1	2	5	8	7	21	47	139	-
1999.....	333	22	9	3	4	1	6	6	11	18	56	197	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)													
Non-Hispanic white, both sexes													
2003.....	342	25	11	5	3	1	10	17	16	25	73	156	-
2002.....	314	25	18	6	2	3	7	14	12	28	59	140	-
2001.....	278	24	12	3	1	-	12	8	14	20	50	134	-
2000.....	345	15	6	3	5	4	8	8	10	40	74	172	-
1999.....	440	26	12	3	4	3	8	10	17	26	99	232	-
Non-Hispanic white, male													
2003.....	125	15	6	2	-	1	6	8	8	15	31	33	-
2002.....	122	22	6	6	-	-	3	9	3	12	25	36	-
2001.....	86	11	5	2	1	-	7	3	5	7	17	28	-
2000.....	118	6	3	2	4	3	6	3	3	19	30	39	-
1999.....	145	14	6	1	1	2	3	8	9	11	48	42	-
Non-Hispanic white, female													
2003.....	217	10	5	3	3	-	4	9	8	10	42	123	-
2002.....	192	3	12	-	2	3	4	5	9	16	34	104	-
2001.....	192	13	7	1	-	-	5	5	9	13	33	106	-
2000.....	227	9	3	1	1	1	2	5	7	21	44	133	-
1999.....	295	12	6	2	3	1	5	2	8	15	51	190	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)													
Non-Hispanic black, both sexes													
2003.....	40	12	7	1	3	1	-	2	2	4	3	5	-
2002.....	44	15	2	-	1	1	1	4	3	5	7	5	-
2001.....	33	14	2	-	-	1	-	4	1	4	-	7	-
2000.....	46	14	3	3	-	2	4	3	-	2	5	10	-
1999.....	64	24	6	-	1	1	2	4	2	8	6	10	-
Non-Hispanic black, male													
2003.....	21	8	6	-	2	-	-	-	1	2	2	-	-
2002.....	22	12	1	-	-	1	-	1	2	1	2	2	-
2001.....	15	8	1	-	-	-	-	3	-	1	-	2	-
2000.....	22	10	-	1	-	2	1	-	-	2	2	4	-
1999.....	30	15	3	-	-	1	1	-	1	5	1	3	-
Non-Hispanic black, female													
2003.....	19	4	1	1	1	1	-	2	1	2	1	5	-
2002.....	22	3	1	-	1	-	1	3	1	4	5	3	-
2001.....	18	6	1	-	-	1	-	1	1	3	-	5	-
2000.....	24	4	3	2	-	-	3	3	-	-	3	6	-
1999.....	34	9	3	-	1	-	1	4	1	3	5	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)													
All origins, both sexes													
2003.....	304	53	27	6	6	2	9	16	17	22	45	101	-
2002.....	279	49	23	8	2	5	6	17	13	23	39	94	-
2001.....	263	49	18	3	1	2	11	12	14	23	39	91	-
2000.....	290	42	11	6	4	6	10	9	7	31	53	111	-
1999.....	388	62	24	4	5	3	10	14	18	29	69	150	-
All origins, male													
2003.....	126	34	16	2	2	1	5	7	11	12	17	19	-
2002.....	122	37	9	7	-	1	3	9	5	10	18	23	-
2001.....	89	25	8	2	1	-	6	6	6	8	10	17	-
2000.....	119	24	4	2	3	4	6	3	3	14	27	29	-
1999.....	154	36	14	2	1	2	4	7	8	15	34	31	-
All origins, female													
2003.....	178	19	11	4	4	1	4	9	6	10	28	82	-
2002.....	157	12	14	1	2	4	3	8	8	13	21	71	-
2001.....	174	24	10	1	-	2	5	6	8	15	29	74	-
2000.....	171	18	7	4	1	2	4	6	4	17	26	82	-
1999.....	234	26	10	2	4	1	6	7	10	14	35	119	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)													
Hispanic, both sexes													
2003.....	31	17	6	-	-	-	1	-	1	1	2	3	-
2002.....	19	9	2	2	-	1	-	-	2	-	-	3	-
2001.....	17	8	1	-	-	1	-	2	-	1	2	2	-
2000.....	22	9	2	-	-	1	-	1	-	1	2	6	-
1999.....	20	10	3	1	-	-	-	1	-	1	-	4	-
Hispanic, male													
2003.....	20	12	4	-	-	-	-	-	1	-	1	2	-
2002.....	9	5	1	1	-	-	-	-	1	-	-	1	-
2001.....	7	6	-	-	-	-	-	1	-	-	-	-	-
2000.....	15	6	1	-	-	1	-	1	-	1	2	3	-
1999.....	10	5	2	1	-	-	-	-	-	1	-	1	-
Hispanic, female													
2003.....	11	5	2	-	-	-	1	-	-	1	1	1	-
2002.....	10	4	1	1	-	1	-	-	1	-	-	2	-
2001.....	10	2	1	-	-	1	-	1	-	1	2	2	-
2000.....	7	3	1	-	-	-	-	-	-	-	-	3	-
1999.....	10	5	1	-	-	-	-	1	-	-	-	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)													
Non-Hispanic 1/, both sexes													
2003.....	272	36	21	6	6	2	8	16	15	21	43	98	-
2002.....	260	40	21	6	2	4	6	17	11	23	39	91	-
2001.....	246	41	17	3	1	1	11	10	14	22	37	89	-
2000.....	267	32	9	6	4	5	10	8	7	30	51	105	-
1999.....	367	52	21	3	5	3	9	13	18	28	69	146	-
Non-Hispanic 1/, male													
2003.....	105	22	12	2	2	1	5	7	9	12	16	17	-
2002.....	113	32	8	6	-	1	3	9	4	10	18	22	-
2001.....	82	19	8	2	1	-	6	5	6	8	10	17	-
2000.....	104	18	3	2	3	3	6	2	3	13	25	26	-
1999.....	143	31	12	1	1	2	3	7	8	14	34	30	-
Non-Hispanic 1/, female													
2003.....	167	14	9	4	4	1	3	9	6	9	27	81	-
2002.....	147	8	13	-	2	3	3	8	7	13	21	69	-
2001.....	164	22	9	1	-	1	5	5	8	14	27	72	-
2000.....	163	14	6	4	1	2	4	6	4	17	26	79	-
1999.....	224	21	9	2	4	1	6	6	10	14	35	116	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Acute bronchitis and bronchiolitis (J20-J21)

Non-Hispanic white, both sexes

2003.....	231	24	10	5	3	1	8	15	12	17	41	95	-
2002.....	219	23	18	6	1	3	6	13	8	21	36	84	-
2001.....	202	23	12	3	1	-	11	6	12	17	34	83	-
2000.....	221	15	6	2	4	2	6	5	7	28	47	99	-
1999.....	299	25	12	2	4	3	7	9	14	20	63	140	-

Non-Hispanic white, male

2003.....	83	14	5	2	-	1	5	7	7	10	15	17	-
2002.....	92	20	6	6	-	-	3	8	3	9	17	20	-
2001.....	61	10	5	2	1	-	6	2	5	6	8	16	-
2000.....	83	6	3	1	3	1	5	2	3	11	24	24	-
1999.....	108	14	6	1	1	2	2	7	7	9	32	27	-

Non-Hispanic white, female

2003.....	148	10	5	3	3	-	3	8	5	7	26	78	-
2002.....	127	3	12	-	1	3	3	5	5	12	19	64	-
2001.....	141	13	7	1	-	-	5	4	7	11	26	67	-
2000.....	138	9	3	1	1	1	1	3	4	17	23	75	-
1999.....	191	11	6	1	3	1	5	2	7	11	31	113	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)													
Non-Hispanic black, both sexes													
2003.....	35	12	7	1	3	1	-	1	2	4	1	3	-
2002.....	35	15	2	-	1	1	-	4	2	2	3	5	-
2001.....	31	14	2	-	-	1	-	4	1	3	-	6	-
2000.....	40	13	3	3	-	2	4	3	-	2	4	6	-
1999.....	55	24	5	-	1	-	2	4	2	8	4	5	-
Non-Hispanic black, male													
2003.....	20	8	6	-	2	-	-	-	1	2	1	-	-
2002.....	20	12	1	-	-	1	-	1	1	1	1	2	-
2001.....	14	8	1	-	-	-	-	3	-	1	-	1	-
2000.....	18	9	-	1	-	2	1	-	-	2	1	2	-
1999.....	26	15	2	-	-	-	1	-	1	5	-	2	-
Non-Hispanic black, female													
2003.....	15	4	1	1	1	1	-	1	1	2	-	3	-
2002.....	15	3	1	-	1	-	-	3	1	1	2	3	-
2001.....	17	6	1	-	-	1	-	1	1	2	-	5	-
2000.....	22	4	3	2	-	-	3	3	-	-	3	4	-
1999.....	29	9	3	-	1	-	1	4	1	3	4	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)													
All origins, both sexes													
2003.....	117	1	1	-	-	-	2	3	4	8	35	63	-
2002.....	107	3	-	-	1	-	2	1	5	10	27	58	-
2001.....	84	1	-	-	-	-	1	2	2	5	19	54	-
2000.....	135	1	-	1	1	2	2	3	3	12	29	81	-
1999.....	152	1	1	1	-	1	1	1	3	6	38	99	-
All origins, male													
2003.....	44	1	1	-	-	-	1	1	1	5	18	16	-
2002.....	32	2	-	-	-	-	-	1	1	3	9	16	-
2001.....	28	1	-	-	-	-	1	1	-	2	10	13	-
2000.....	41	1	-	1	1	2	1	1	-	8	8	18	-
1999.....	42	-	1	-	-	1	1	1	2	2	17	17	-
All origins, female													
2003.....	73	-	-	-	-	-	1	2	3	3	17	47	-
2002.....	75	1	-	-	1	-	2	-	4	7	18	42	-
2001.....	56	-	-	-	-	-	-	1	2	3	9	41	-
2000.....	94	-	-	-	-	-	1	2	3	4	21	63	-
1999.....	110	1	-	1	-	-	-	-	1	4	21	82	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)													
Non-Hispanic 1/, both sexes													
2003.....	116	1	1	-	-	-	2	3	4	8	34	63	-
2002.....	105	2	-	-	1	-	2	1	5	10	27	57	-
2001.....	79	1	-	-	-	-	1	2	2	4	17	52	-
2000.....	131	1	-	1	1	2	2	3	3	12	28	78	-
1999.....	151	1	1	1	-	1	1	1	3	6	38	98	-
Non-Hispanic 1/, male													
2003.....	43	1	1	-	-	-	1	1	1	5	17	16	-
2002.....	32	2	-	-	-	-	-	1	1	3	9	16	-
2001.....	26	1	-	-	-	-	1	1	-	1	9	13	-
2000.....	40	1	-	1	1	2	1	1	-	8	7	18	-
1999.....	42	-	1	-	-	1	1	1	2	2	17	17	-
Non-Hispanic 1/, female													
2003.....	73	-	-	-	-	-	1	2	3	3	17	47	-
2002.....	73	-	-	-	1	-	2	-	4	7	18	41	-
2001.....	53	-	-	-	-	-	-	1	2	3	8	39	-
2000.....	91	-	-	-	-	-	1	2	3	4	21	60	-
1999.....	109	1	-	1	-	-	-	-	1	4	21	81	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)													
Non-Hispanic white, both sexes													
2003.....	111	1	1	-	-	-	2	2	4	8	32	61	-
2002.....	95	2	-	-	1	-	1	1	4	7	23	56	-
2001.....	76	1	-	-	-	-	1	2	2	3	16	51	-
2000.....	124	-	-	1	1	2	2	3	3	12	27	73	-
1999.....	141	1	-	1	-	-	1	1	3	6	36	92	-
Non-Hispanic white, male													
2003.....	42	1	1	-	-	-	1	1	1	5	16	16	-
2002.....	30	2	-	-	-	-	-	1	-	3	8	16	-
2001.....	25	1	-	-	-	-	1	1	-	1	9	12	-
2000.....	35	-	-	1	1	2	1	1	-	8	6	15	-
1999.....	37	-	-	-	-	-	1	1	2	2	16	15	-
Non-Hispanic white, female													
2003.....	69	-	-	-	-	-	1	1	3	3	16	45	-
2002.....	65	-	-	-	1	-	1	-	4	4	15	40	-
2001.....	51	-	-	-	-	-	-	1	2	2	7	39	-
2000.....	89	-	-	-	-	-	1	2	3	4	21	58	-
1999.....	104	1	-	1	-	-	-	-	1	4	20	77	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)													
Non-Hispanic black, both sexes													
2003.....	5	-	-	-	-	-	-	1	-	-	2	2	-
2002.....	9	-	-	-	-	-	1	-	1	3	4	-	-
2001.....	2	-	-	-	-	-	-	-	-	1	-	1	-
2000.....	6	1	-	-	-	-	-	-	-	-	1	4	-
1999.....	9	-	1	-	-	1	-	-	-	-	2	5	-
Non-Hispanic black, male													
2003.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	2	-	-	-	-	-	-	-	1	-	1	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2000.....	4	1	-	-	-	-	-	-	-	-	1	2	-
1999.....	4	-	1	-	-	1	-	-	-	-	1	1	-
Non-Hispanic black, female													
2003.....	4	-	-	-	-	-	-	1	-	-	1	2	-
2002.....	7	-	-	-	-	-	1	-	-	3	3	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	2	-
1999.....	5	-	-	-	-	-	-	-	-	-	1	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)													
All origins, both sexes													
2003.....	126,382	31	55	118	191	282	950	3,537	12,077	29,919	49,286	29,934	2
2002.....	124,816	39	65	136	192	301	1,008	3,475	11,280	29,788	49,241	29,284	7
2001.....	123,013	41	43	104	171	290	969	3,324	11,166	30,751	47,762	28,391	1
2000.....	122,009	36	51	139	190	296	930	3,251	10,739	31,157	47,722	27,496	2
1999.....	124,181	36	54	139	209	330	890	3,110	11,297	32,644	48,635	26,833	4
All origins, male													
2003.....	60,714	23	38	75	114	156	429	1,826	6,243	15,333	23,963	12,512	2
2002.....	60,713	23	44	74	120	165	497	1,795	5,783	15,339	24,432	12,436	5
2001.....	59,697	22	29	61	99	146	455	1,702	5,702	15,877	23,691	11,912	1
2000.....	60,004	23	28	80	120	165	437	1,661	5,564	16,210	23,838	11,876	2
1999.....	62,415	19	30	93	123	152	410	1,551	5,958	17,477	24,705	11,894	3
All origins, female													
2003.....	65,668	8	17	43	77	126	521	1,711	5,834	14,586	25,323	17,422	-
2002.....	64,103	16	21	62	72	136	511	1,680	5,497	14,449	24,809	16,848	2
2001.....	63,316	19	14	43	72	144	514	1,622	5,464	14,874	24,071	16,479	-
2000.....	62,005	13	23	59	70	131	493	1,590	5,175	14,947	23,884	15,620	-
1999.....	61,766	17	24	46	86	178	480	1,559	5,339	15,167	23,930	14,939	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)													
Hispanic, both sexes													
2003.....	3,174	7	11	15	20	29	58	134	266	651	1,095	888	-
2002.....	3,058	4	9	13	15	31	70	127	274	635	1,055	825	-
2001.....	2,832	7	8	9	10	22	52	124	258	634	940	768	-
2000.....	2,689	7	9	15	16	20	55	105	244	572	921	725	-
1999.....	2,859	6	8	18	15	31	79	115	239	649	935	764	-
Hispanic, male													
2003.....	1,697	5	8	10	12	15	30	61	164	369	603	420	-
2002.....	1,625	2	4	8	12	19	31	70	164	362	593	360	-
2001.....	1,482	4	4	6	6	13	21	54	140	378	501	355	-
2000.....	1,451	4	5	9	13	12	24	53	146	343	509	333	-
1999.....	1,547	4	3	11	9	13	36	52	135	373	523	388	-
Hispanic, female													
2003.....	1,477	2	3	5	8	14	28	73	102	282	492	468	-
2002.....	1,433	2	5	5	3	12	39	57	110	273	462	465	-
2001.....	1,350	3	4	3	4	9	31	70	118	256	439	413	-
2000.....	1,238	3	4	6	3	8	31	52	98	229	412	392	-
1999.....	1,312	2	5	7	6	18	43	63	104	276	412	376	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)													
Non-Hispanic 1/, both sexes													
2003.....	122,896	24	43	102	170	253	890	3,382	11,761	29,192	48,093	28,985	1
2002.....	121,424	34	56	121	176	267	930	3,335	10,975	29,069	48,058	28,397	6
2001.....	119,839	34	35	94	161	267	909	3,180	10,860	30,028	46,713	27,558	-
2000.....	118,963	29	42	123	173	272	872	3,122	10,447	30,486	46,687	26,709	1
1999.....	120,968	30	46	121	194	296	807	2,978	11,018	31,902	47,578	25,995	3
Non-Hispanic 1/, male													
2003.....	58,848	18	29	65	101	141	397	1,753	6,050	14,920	23,304	12,069	1
2002.....	58,896	20	40	64	107	143	460	1,719	5,604	14,924	23,761	12,050	4
2001.....	58,028	18	25	55	93	133	428	1,635	5,530	15,449	23,134	11,528	-
2000.....	58,349	19	23	71	107	150	411	1,593	5,388	15,805	23,266	11,515	1
1999.....	60,669	15	27	82	114	138	372	1,486	5,797	17,040	24,124	11,472	2
Non-Hispanic 1/, female													
2003.....	64,048	6	14	37	69	112	493	1,629	5,711	14,272	24,789	16,916	-
2002.....	62,528	14	16	57	69	124	470	1,616	5,371	14,145	24,297	16,347	2
2001.....	61,811	16	10	39	68	134	481	1,545	5,330	14,579	23,579	16,030	-
2000.....	60,614	10	19	52	66	122	461	1,529	5,059	14,681	23,421	15,194	-
1999.....	60,299	15	19	39	80	158	435	1,492	5,221	14,862	23,454	14,523	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)													
Non-Hispanic white, both sexes													
2003.....	113,541	17	15	39	82	140	617	2,725	10,465	26,902	45,177	27,361	1
2002.....	112,128	20	23	42	94	157	624	2,686	9,757	26,808	45,117	26,796	4
2001.....	110,753	16	13	39	75	134	613	2,570	9,678	27,693	43,972	25,950	-
2000.....	109,901	16	19	47	87	144	570	2,463	9,253	28,122	44,024	25,155	1
1999.....	111,620	10	23	55	110	146	533	2,372	9,786	29,400	44,662	24,521	2
Non-Hispanic white, male													
2003.....	53,593	16	7	21	48	64	275	1,426	5,327	13,604	21,553	11,251	1
2002.....	53,662	13	17	23	48	81	304	1,383	4,909	13,615	22,034	11,232	3
2001.....	52,952	10	9	22	39	62	293	1,331	4,885	14,089	21,496	10,716	-
2000.....	53,231	12	8	24	52	71	267	1,259	4,728	14,398	21,703	10,708	1
1999.....	55,314	6	11	36	59	66	239	1,174	5,091	15,559	22,377	10,695	1
Non-Hispanic white, female													
2003.....	59,948	1	8	18	34	76	342	1,299	5,138	13,298	23,624	16,110	-
2002.....	58,466	7	6	19	46	76	320	1,303	4,848	13,193	23,083	15,564	1
2001.....	57,801	6	4	17	36	72	320	1,239	4,793	13,604	22,476	15,234	-
2000.....	56,670	4	11	23	35	73	303	1,204	4,525	13,724	22,321	14,447	-
1999.....	56,306	4	12	19	51	80	294	1,198	4,695	13,841	22,285	13,826	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)													
Non-Hispanic black, both sexes													
2003.....	7,632	7	24	62	81	104	245	597	1,109	1,920	2,290	1,193	-
2002.....	7,730	14	29	74	78	102	292	589	1,067	1,892	2,366	1,226	1
2001.....	7,504	16	22	53	77	128	273	549	1,019	1,962	2,190	1,215	-
2000.....	7,528	11	21	70	80	114	272	593	1,063	1,993	2,128	1,183	-
1999.....	7,840	18	18	66	78	140	258	556	1,085	2,090	2,364	1,166	1
Non-Hispanic black, male													
2003.....	4,202	2	19	43	48	72	106	293	615	1,093	1,340	571	-
2002.....	4,282	7	20	39	58	56	150	304	607	1,093	1,350	597	1
2001.....	4,143	6	16	32	48	69	124	271	554	1,147	1,285	591	-
2000.....	4,184	6	14	43	51	70	129	289	587	1,195	1,207	593	-
1999.....	4,452	9	13	46	52	67	121	284	624	1,238	1,402	595	1
Non-Hispanic black, female													
2003.....	3,430	5	5	19	33	32	139	304	494	827	950	622	-
2002.....	3,448	7	9	35	20	46	142	285	460	799	1,016	629	-
2001.....	3,361	10	6	21	29	59	149	278	465	815	905	624	-
2000.....	3,344	5	7	27	29	44	143	304	476	798	921	590	-
1999.....	3,388	9	5	20	26	73	137	272	461	852	962	571	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)													
Origin not stated 2/, both sexes													
2003.....	312	-	1	1	1	-	2	21	50	76	98	61	1
2002.....	334	1	-	2	1	3	8	13	31	84	128	62	1
2001.....	342	-	-	1	-	1	8	20	48	89	109	65	1
2000.....	357	-	-	1	1	4	3	24	48	99	114	62	1
1999.....	354	-	-	-	-	3	4	17	40	93	122	74	1
Origin not stated 2/, male													
2003.....	169	-	1	-	1	-	2	12	29	44	56	23	1
2002.....	192	1	-	2	1	3	6	6	15	53	78	26	1
2001.....	187	-	-	-	-	-	6	13	32	50	56	29	1
2000.....	204	-	-	-	-	3	2	15	30	62	63	28	1
1999.....	199	-	-	-	-	1	2	13	26	64	58	34	1
Origin not stated 2/, female													
2003.....	143	-	-	1	-	-	-	9	21	32	42	38	-
2002.....	142	-	-	-	-	-	2	7	16	31	50	36	-
2001.....	155	-	-	1	-	1	2	7	16	39	53	36	-
2000.....	153	-	-	1	1	1	1	9	18	37	51	34	-
1999.....	155	-	-	-	-	2	2	4	14	29	64	40	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
All origins, both sexes													
2003.....	850	20	13	6	7	10	17	35	47	104	246	345	-
2002.....	955	24	13	5	6	6	18	36	54	135	286	372	-
2001.....	959	21	8	-	5	6	21	42	61	133	261	401	-
2000.....	1,167	18	14	5	5	8	20	38	67	158	333	501	-
1999.....	1,172	20	19	4	7	11	18	38	97	173	339	446	-
All origins, male													
2003.....	351	15	6	3	6	3	9	23	26	46	113	101	-
2002.....	420	15	8	2	3	2	12	27	28	74	127	122	-
2001.....	396	11	5	-	3	2	8	21	36	76	115	119	-
2000.....	503	9	7	3	4	7	14	16	32	79	174	158	-
1999.....	498	9	11	3	4	3	5	19	51	94	157	142	-
All origins, female													
2003.....	499	5	7	3	1	7	8	12	21	58	133	244	-
2002.....	535	9	5	3	3	4	6	9	26	61	159	250	-
2001.....	563	10	3	-	2	4	13	21	25	57	146	282	-
2000.....	664	9	7	2	1	1	6	22	35	79	159	343	-
1999.....	674	11	8	1	3	8	13	19	46	79	182	304	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
Hispanic, both sexes													
2003.....	40	5	5	-	1	1	-	1	1	3	9	14	-
2002.....	51	3	1	-	2	1	3	3	2	2	14	20	-
2001.....	37	4	1	-	1	1	2	1	2	3	9	13	-
2000.....	52	4	3	-	-	2	3	1	1	9	13	16	-
1999.....	52	3	3	1	-	2	7	-	2	4	13	17	-
Hispanic, male													
2003.....	20	3	3	-	1	-	-	-	-	2	5	6	-
2002.....	20	1	1	-	1	1	2	3	1	1	4	5	-
2001.....	15	2	-	-	-	1	1	-	1	1	6	3	-
2000.....	28	2	2	-	-	2	2	-	1	7	6	6	-
1999.....	17	1	2	1	-	-	2	-	-	3	3	5	-
Hispanic, female													
2003.....	20	2	2	-	-	1	-	1	1	1	4	8	-
2002.....	31	2	-	-	1	-	1	-	1	1	10	15	-
2001.....	22	2	1	-	1	-	1	1	1	2	3	10	-
2000.....	24	2	1	-	-	-	1	1	-	2	7	10	-
1999.....	35	2	1	-	-	2	5	-	2	1	10	12	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
Non-Hispanic 1/, both sexes													
2003.....	808	15	8	6	6	9	17	34	46	101	235	331	-
2002.....	900	20	12	5	4	5	15	33	52	132	271	351	-
2001.....	919	17	7	-	4	5	19	40	58	130	251	388	-
2000.....	1,112	14	11	5	5	6	17	37	66	149	320	482	-
1999.....	1,117	17	16	3	7	9	11	38	94	168	326	428	-
Non-Hispanic 1/, male													
2003.....	330	12	3	3	5	3	9	23	26	44	107	95	-
2002.....	398	13	7	2	2	1	10	24	27	73	122	117	-
2001.....	380	9	5	-	3	1	7	21	34	75	109	116	-
2000.....	474	7	5	3	4	5	12	16	31	72	168	151	-
1999.....	478	8	9	2	4	3	3	19	50	90	154	136	-
Non-Hispanic 1/, female													
2003.....	478	3	5	3	1	6	8	11	20	57	128	236	-
2002.....	502	7	5	3	2	4	5	9	25	59	149	234	-
2001.....	539	8	2	-	1	4	12	19	24	55	142	272	-
2000.....	638	7	6	2	1	1	5	21	35	77	152	331	-
1999.....	639	9	7	1	3	6	8	19	44	78	172	292	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
Non-Hispanic white, both sexes													
2003.....	712	11	4	3	5	5	13	23	37	92	214	305	-
2002.....	821	12	6	3	2	4	13	26	44	124	252	335	-
2001.....	839	7	5	-	3	4	16	35	54	113	232	370	-
2000.....	1,018	5	4	4	4	4	11	31	60	131	297	467	-
1999.....	1,002	8	12	3	6	6	9	24	86	150	293	405	-
Non-Hispanic white, male													
2003.....	290	10	2	1	4	2	7	17	23	39	96	89	-
2002.....	354	9	2	2	1	1	8	17	22	67	113	112	-
2001.....	338	5	5	-	2	-	6	17	32	66	97	108	-
2000.....	429	3	1	2	3	3	6	15	27	64	159	146	-
1999.....	421	5	6	2	3	2	3	11	45	81	135	128	-
Non-Hispanic white, female													
2003.....	422	1	2	2	1	3	6	6	14	53	118	216	-
2002.....	467	3	4	1	1	3	5	9	22	57	139	223	-
2001.....	501	2	-	-	1	4	10	18	22	47	135	262	-
2000.....	589	2	3	2	1	1	5	16	33	67	138	321	-
1999.....	581	3	6	1	3	4	6	13	41	69	158	277	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)													
Non-Hispanic black, both sexes													
2003.....	74	4	3	3	1	3	4	11	8	6	12	19	-
2002.....	66	8	6	2	2	1	2	7	5	6	13	14	-
2001.....	62	10	2	-	1	1	3	4	4	15	11	11	-
2000.....	80	7	6	1	1	1	6	6	4	16	20	12	-
1999.....	88	7	1	-	1	3	2	12	7	15	26	14	-
Non-Hispanic black, male													
2003.....	29	2	-	2	1	1	2	6	3	2	7	3	-
2002.....	35	4	5	-	1	-	2	7	2	4	5	5	-
2001.....	32	4	-	-	1	1	1	4	2	8	6	5	-
2000.....	35	3	3	1	1	1	6	1	3	6	7	3	-
1999.....	43	3	1	-	1	1	-	7	4	7	16	3	-
Non-Hispanic black, female													
2003.....	45	2	3	1	-	2	2	5	5	4	5	16	-
2002.....	31	4	1	2	1	1	-	-	3	2	8	9	-
2001.....	30	6	2	-	-	-	2	-	2	7	5	6	-
2000.....	45	4	3	-	-	-	-	5	1	10	13	9	-
1999.....	45	4	-	-	-	2	2	5	3	8	10	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
All origins, both sexes													
2003.....	14,861	-	1	-	4	5	87	450	1,755	4,063	5,920	2,574	2
2002.....	15,489	1	3	-	2	10	94	454	1,703	4,388	6,109	2,724	1
2001.....	16,242	3	-	1	2	7	75	448	1,829	4,804	6,337	2,736	-
2000.....	16,700	3	2	-	5	11	82	474	1,807	5,120	6,511	2,685	-
1999.....	17,787	1	-	1	1	9	97	468	2,019	5,656	6,955	2,580	-
All origins, male													
2003.....	7,549	-	1	-	2	3	49	279	932	2,141	2,995	1,145	2
2002.....	8,084	-	3	-	-	8	66	267	925	2,335	3,228	1,251	1
2001.....	8,415	1	-	1	1	4	52	266	966	2,554	3,255	1,315	-
2000.....	8,671	3	-	-	5	5	46	280	943	2,738	3,373	1,278	-
1999.....	9,480	-	-	-	1	6	49	268	1,085	3,136	3,677	1,258	-
All origins, female													
2003.....	7,312	-	-	-	2	2	38	171	823	1,922	2,925	1,429	-
2002.....	7,405	1	-	-	2	2	28	187	778	2,053	2,881	1,473	-
2001.....	7,827	2	-	-	1	3	23	182	863	2,250	3,082	1,421	-
2000.....	8,029	-	2	-	-	6	36	194	864	2,382	3,138	1,407	-
1999.....	8,307	1	-	1	-	3	48	200	934	2,520	3,278	1,322	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
Hispanic, both sexes													
2003.....	312	-	-	-	-	-	6	8	30	73	128	67	-
2002.....	295	-	-	-	-	2	1	5	29	79	113	66	-
2001.....	286	1	-	-	-	-	-	8	26	75	113	63	-
2000.....	266	-	-	-	1	-	3	11	26	65	113	47	-
1999.....	292	-	-	1	-	2	6	12	21	74	113	63	-
Hispanic, male													
2003.....	184	-	-	-	-	-	5	6	21	41	76	35	-
2002.....	168	-	-	-	-	1	1	4	17	48	65	32	-
2001.....	175	-	-	-	-	-	-	5	16	46	76	32	-
2000.....	167	-	-	-	1	-	3	8	17	43	66	29	-
1999.....	183	-	-	-	-	1	2	8	16	51	69	36	-
Hispanic, female													
2003.....	128	-	-	-	-	-	1	2	9	32	52	32	-
2002.....	127	-	-	-	-	1	-	1	12	31	48	34	-
2001.....	111	1	-	-	-	-	-	3	10	29	37	31	-
2000.....	99	-	-	-	-	-	-	3	9	22	47	18	-
1999.....	109	-	-	1	-	1	4	4	5	23	44	27	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
Non-Hispanic 1/, both sexes													
2003.....	14,514	-	1	-	4	5	81	438	1,716	3,982	5,781	2,505	1
2002.....	15,161	1	3	-	2	8	93	448	1,672	4,298	5,986	2,650	-
2001.....	15,914	2	-	1	2	7	74	439	1,794	4,718	6,210	2,667	-
2000.....	16,382	3	2	-	4	11	79	457	1,777	5,036	6,382	2,631	-
1999.....	17,441	1	-	-	1	7	88	450	1,992	5,571	6,825	2,506	-
Non-Hispanic 1/, male													
2003.....	7,342	-	1	-	2	3	44	270	908	2,094	2,910	1,109	1
2002.....	7,895	-	3	-	-	7	65	262	907	2,281	3,156	1,214	-
2001.....	8,213	1	-	1	1	4	51	260	943	2,500	3,172	1,280	-
2000.....	8,483	3	-	-	4	5	43	268	925	2,685	3,303	1,247	-
1999.....	9,260	-	-	-	1	5	45	255	1,063	3,076	3,598	1,217	-
Non-Hispanic 1/, female													
2003.....	7,172	-	-	-	2	2	37	168	808	1,888	2,871	1,396	-
2002.....	7,266	1	-	-	2	1	28	186	765	2,017	2,830	1,436	-
2001.....	7,701	1	-	-	1	3	23	179	851	2,218	3,038	1,387	-
2000.....	7,899	-	2	-	-	6	36	189	852	2,351	3,079	1,384	-
1999.....	8,181	1	-	-	-	2	43	195	929	2,495	3,227	1,289	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
Non-Hispanic white, both sexes													
2003.....	13,518	-	1	-	4	3	64	376	1,557	3,717	5,438	2,357	1
2002.....	14,157	1	1	-	1	6	66	385	1,529	3,992	5,663	2,513	-
2001.....	14,930	1	-	-	1	4	64	374	1,633	4,414	5,913	2,526	-
2000.....	15,301	3	-	-	1	10	60	367	1,616	4,704	6,068	2,472	-
1999.....	16,338	-	-	-	-	5	64	372	1,825	5,238	6,469	2,365	-
Non-Hispanic white, male													
2003.....	6,714	-	1	-	2	1	34	226	808	1,935	2,690	1,016	1
2002.....	7,241	-	1	-	-	5	47	219	810	2,081	2,946	1,132	-
2001.....	7,607	-	-	-	1	2	44	222	841	2,310	2,986	1,201	-
2000.....	7,805	3	-	-	1	4	31	206	830	2,470	3,121	1,139	-
1999.....	8,540	-	-	-	-	3	33	199	953	2,843	3,371	1,138	-
Non-Hispanic white, female													
2003.....	6,804	-	-	-	2	2	30	150	749	1,782	2,748	1,341	-
2002.....	6,916	1	-	-	1	1	19	166	719	1,911	2,717	1,381	-
2001.....	7,323	1	-	-	-	2	20	152	792	2,104	2,927	1,325	-
2000.....	7,496	-	-	-	-	6	29	161	786	2,234	2,947	1,333	-
1999.....	7,798	-	-	-	-	2	31	173	872	2,395	3,098	1,227	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
Non-Hispanic black, both sexes													
2003.....	805	-	-	-	-	2	15	55	142	220	266	105	-
2002.....	842	-	1	-	1	2	26	56	117	267	266	106	-
2001.....	800	1	-	1	1	3	9	59	138	254	219	115	-
2000.....	889	-	1	-	3	1	17	84	146	276	243	118	-
1999.....	923	1	-	-	-	-	22	75	146	279	283	117	-
Non-Hispanic black, male													
2003.....	495	-	-	-	-	2	8	37	90	129	162	67	-
2002.....	538	-	1	-	-	2	18	38	77	173	169	60	-
2001.....	480	1	-	1	-	2	7	33	87	155	134	60	-
2000.....	551	-	-	-	3	1	11	56	87	183	134	76	-
1999.....	594	-	-	-	-	-	11	54	94	195	178	62	-
Non-Hispanic black, female													
2003.....	310	-	-	-	-	-	7	18	52	91	104	38	-
2002.....	304	-	-	-	1	-	8	18	40	94	97	46	-
2001.....	320	-	-	-	1	1	2	26	51	99	85	55	-
2000.....	338	-	1	-	-	-	6	28	59	93	109	42	-
1999.....	329	1	-	-	-	-	11	21	52	84	105	55	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Emphysema (J43)													
Origin not stated 2/, both sexes													
2003.....	35	-	-	-	-	-	-	4	9	8	11	2	1
2002.....	33	-	-	-	-	-	-	1	2	11	10	8	1
2001.....	42	-	-	-	-	-	1	1	9	11	14	6	-
2000.....	52	-	-	-	-	-	-	6	4	19	16	7	-
1999.....	54	-	-	-	-	-	3	6	6	11	17	11	-
Origin not stated 2/, male													
2003.....	23	-	-	-	-	-	-	3	3	6	9	1	1
2002.....	21	-	-	-	-	-	-	1	1	6	7	5	1
2001.....	27	-	-	-	-	-	1	1	7	8	7	3	-
2000.....	21	-	-	-	-	-	-	4	1	10	4	2	-
1999.....	37	-	-	-	-	-	2	5	6	9	10	5	-
Origin not stated 2/, female													
2003.....	12	-	-	-	-	-	-	1	6	2	2	1	-
2002.....	12	-	-	-	-	-	-	-	1	5	3	3	-
2001.....	15	-	-	-	-	-	-	-	2	3	7	3	-
2000.....	31	-	-	-	-	-	-	2	3	9	12	5	-
1999.....	17	-	-	-	-	-	1	1	-	2	7	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
All origins, both sexes													
2003.....	4,099	7	37	110	158	227	411	632	562	532	752	671	-
2002.....	4,261	4	43	123	169	235	472	608	536	583	812	675	1
2001.....	4,269	10	31	99	140	233	455	603	553	634	802	709	-
2000.....	4,487	8	32	129	167	249	458	614	540	734	849	707	-
1999.....	4,657	5	27	126	182	262	447	607	583	773	925	720	-
All origins, male													
2003.....	1,493	4	29	71	91	127	160	236	214	173	234	154	-
2002.....	1,580	3	30	69	110	129	198	233	202	198	256	152	-
2001.....	1,479	6	22	57	80	116	195	219	182	220	225	157	-
2000.....	1,632	6	20	74	102	142	196	238	200	253	243	158	-
1999.....	1,620	3	16	85	108	128	184	197	207	267	273	152	-
All origins, female													
2003.....	2,606	3	8	39	67	100	251	396	348	359	518	517	-
2002.....	2,681	1	13	54	59	106	274	375	334	385	556	523	1
2001.....	2,790	4	9	42	60	117	260	384	371	414	577	552	-
2000.....	2,855	2	12	55	65	107	262	376	340	481	606	549	-
1999.....	3,037	2	11	41	74	134	263	410	376	506	652	568	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Hispanic, both sexes													
2003.....	299	-	6	15	14	23	27	61	45	41	39	28	-
2002.....	287	1	6	13	11	23	42	44	38	45	42	22	-
2001.....	274	2	5	8	8	17	31	49	45	43	44	22	-
2000.....	292	1	5	13	14	16	29	41	48	60	40	25	-
1999.....	320	-	3	14	14	21	46	48	39	60	50	25	-
Hispanic, male													
2003.....	131	-	5	10	8	12	13	27	22	13	15	6	-
2002.....	114	1	3	8	9	14	14	22	15	10	15	3	-
2001.....	103	2	3	5	6	10	13	14	17	18	13	2	-
2000.....	129	1	2	8	11	9	14	20	19	25	12	8	-
1999.....	119	-	-	9	8	11	22	20	12	19	12	6	-
Hispanic, female													
2003.....	168	-	1	5	6	11	14	34	23	28	24	22	-
2002.....	173	-	3	5	2	9	28	22	23	35	27	19	-
2001.....	171	-	2	3	2	7	18	35	28	25	31	20	-
2000.....	163	-	3	5	3	7	15	21	29	35	28	17	-
1999.....	201	-	3	5	6	10	24	28	27	41	38	19	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Non-Hispanic 1/, both sexes													
2003.....	3,786	7	30	94	143	204	382	568	516	491	713	638	-
2002.....	3,950	3	37	108	157	210	425	561	495	536	767	650	1
2001.....	3,976	8	26	90	132	215	421	550	507	587	756	684	-
2000.....	4,164	7	27	115	152	229	428	568	486	668	805	679	-
1999.....	4,324	5	24	112	168	238	400	557	543	711	874	692	-
Non-Hispanic 1/, male													
2003.....	1,356	4	23	61	82	115	145	208	192	160	219	147	-
2002.....	1,448	2	27	59	100	113	179	209	187	186	239	147	-
2001.....	1,365	4	19	52	74	106	180	202	165	200	211	152	-
2000.....	1,486	5	18	66	91	130	181	215	176	226	229	149	-
1999.....	1,495	3	16	76	100	116	162	176	195	246	260	145	-
Non-Hispanic 1/, female													
2003.....	2,430	3	7	33	61	89	237	360	324	331	494	491	-
2002.....	2,502	1	10	49	57	97	246	352	308	350	528	503	1
2001.....	2,611	4	7	38	58	109	241	348	342	387	545	532	-
2000.....	2,678	2	9	49	61	99	247	353	310	442	576	530	-
1999.....	2,829	2	8	36	68	122	238	381	348	465	614	547	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Non-Hispanic white, both sexes													
2003.....	2,589	4	8	34	64	108	215	318	334	348	581	575	-
2002.....	2,720	1	14	34	79	121	222	316	311	377	659	586	-
2001.....	2,717	4	7	37	59	97	219	326	306	432	628	602	-
2000.....	2,836	4	13	40	75	114	227	317	296	484	683	583	-
1999.....	3,011	1	8	47	92	107	222	327	351	511	747	598	-
Non-Hispanic white, male													
2003.....	831	4	4	19	36	48	78	113	114	114	169	132	-
2002.....	895	-	12	20	43	59	80	117	113	125	196	130	-
2001.....	834	3	4	21	29	42	91	113	95	141	167	128	-
2000.....	920	2	7	20	44	60	90	130	97	159	193	118	-
1999.....	930	1	5	31	52	48	80	97	124	164	218	110	-
Non-Hispanic white, female													
2003.....	1,758	-	4	15	28	60	137	205	220	234	412	443	-
2002.....	1,825	1	2	14	36	62	142	199	198	252	463	456	-
2001.....	1,883	1	3	16	30	55	128	213	211	291	461	474	-
2000.....	1,916	2	6	20	31	54	137	187	199	325	490	465	-
1999.....	2,081	-	3	16	40	59	142	230	227	347	529	488	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Non-Hispanic black, both sexes													
2003.....	1,019	3	20	59	74	90	155	229	140	120	89	40	-
2002.....	1,083	2	21	69	74	84	196	222	169	132	75	39	-
2001.....	1,092	3	19	51	67	115	186	203	170	126	99	53	-
2000.....	1,145	3	14	69	73	102	178	225	165	152	94	70	-
1999.....	1,134	4	14	65	71	126	166	211	162	157	94	64	-
Non-Hispanic black, male													
2003.....	439	-	18	41	42	63	61	86	52	38	31	7	-
2002.....	488	2	14	37	56	50	96	83	67	51	23	9	-
2001.....	450	-	15	30	41	62	79	83	52	46	29	13	-
2000.....	472	3	11	42	44	62	79	71	65	53	21	21	-
1999.....	474	2	10	45	46	66	72	72	57	61	27	16	-
Non-Hispanic black, female													
2003.....	580	3	2	18	32	27	94	143	88	82	58	33	-
2002.....	595	-	7	32	18	34	100	139	102	81	52	30	-
2001.....	642	3	4	21	26	53	107	120	118	80	70	40	-
2000.....	673	-	3	27	29	40	99	154	100	99	73	49	-
1999.....	660	2	4	20	25	60	94	139	105	96	67	48	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Asthma (J45-J46)													
Origin not stated 2/, both sexes													
2003.....	14	-	1	1	1	-	2	3	1	-	-	5	-
2002.....	24	-	-	2	1	2	5	3	3	2	3	3	-
2001.....	19	-	-	1	-	1	3	4	1	4	2	3	-
2000.....	31	-	-	1	1	4	1	5	6	6	4	3	-
1999.....	13	-	-	-	-	3	1	2	1	2	1	3	-
Origin not stated 2/, male													
2003.....	6	-	1	-	1	-	2	1	-	-	-	1	-
2002.....	18	-	-	2	1	2	5	2	-	2	2	2	-
2001.....	11	-	-	-	-	-	2	3	-	2	1	3	-
2000.....	17	-	-	-	-	3	1	3	5	2	2	1	-
1999.....	6	-	-	-	-	1	-	1	-	2	1	1	-
Origin not stated 2/, female													
2003.....	8	-	-	1	-	-	-	2	1	-	-	4	-
2002.....	6	-	-	-	-	-	-	1	3	-	1	1	-
2001.....	8	-	-	1	-	1	1	1	1	2	1	-	-
2000.....	14	-	-	1	1	1	-	2	1	4	2	2	-
1999.....	7	-	-	-	-	2	1	1	1	-	-	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
All origins, both sexes													
2003.....	106,572	4	4	2	22	40	435	2,420	9,713	25,220	42,368	26,344	-
2002.....	104,111	10	6	8	15	50	424	2,377	8,987	24,682	42,034	25,513	5
2001.....	101,543	7	4	4	24	44	418	2,231	8,723	25,180	40,362	24,545	1
2000.....	99,655	7	3	5	13	28	370	2,125	8,325	25,145	40,029	23,603	2
1999.....	100,565	10	8	8	19	48	328	1,997	8,598	26,042	40,416	23,087	4
All origins, male													
2003.....	51,321	4	2	1	15	23	211	1,288	5,071	12,973	20,621	11,112	-
2002.....	50,629	5	3	3	7	26	221	1,268	4,628	12,732	20,821	10,911	4
2001.....	49,407	4	2	3	15	24	200	1,196	4,518	13,027	20,096	10,321	1
2000.....	49,198	5	1	3	9	11	181	1,127	4,389	13,140	20,048	10,282	2
1999.....	50,817	7	3	5	10	15	172	1,067	4,615	13,980	20,598	10,342	3
All origins, female													
2003.....	55,251	-	2	1	7	17	224	1,132	4,642	12,247	21,747	15,232	-
2002.....	53,482	5	3	5	8	24	203	1,109	4,359	11,950	21,213	14,602	1
2001.....	52,136	3	2	1	9	20	218	1,035	4,205	12,153	20,266	14,224	-
2000.....	50,457	2	2	2	4	17	189	998	3,936	12,005	19,981	13,321	-
1999.....	49,748	3	5	3	9	33	156	930	3,983	12,062	19,818	12,745	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
Hispanic, both sexes													
2003.....	2,523	2	-	-	5	5	25	64	190	534	919	779	-
2002.....	2,425	-	2	-	2	5	24	75	205	509	886	717	-
2001.....	2,235	-	2	1	1	4	19	66	185	513	774	670	-
2000.....	2,079	2	1	2	1	2	20	52	169	438	755	637	-
1999.....	2,195	3	2	2	1	6	20	55	177	511	759	659	-
Hispanic, male													
2003.....	1,362	2	-	-	3	3	12	28	121	313	507	373	-
2002.....	1,323	-	-	-	2	3	14	41	131	303	509	320	-
2001.....	1,189	-	1	1	-	2	7	35	106	313	406	318	-
2000.....	1,127	1	1	1	1	1	5	25	109	268	425	290	-
1999.....	1,228	3	1	1	1	1	10	24	107	300	439	341	-
Hispanic, female													
2003.....	1,161	-	-	-	2	2	13	36	69	221	412	406	-
2002.....	1,102	-	2	-	-	2	10	34	74	206	377	397	-
2001.....	1,046	-	1	-	1	2	12	31	79	200	368	352	-
2000.....	952	1	-	1	-	1	15	27	60	170	330	347	-
1999.....	967	-	1	1	-	5	10	31	70	211	320	318	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
Non-Hispanic 1/, both sexes													
2003.....	103,788	2	4	2	17	35	410	2,342	9,483	24,618	41,364	25,511	-
2002.....	101,413	10	4	8	13	44	397	2,293	8,756	24,103	41,034	24,746	5
2001.....	99,030	7	2	3	23	40	395	2,151	8,501	24,593	39,496	23,819	-
2000.....	97,305	5	2	3	12	26	348	2,060	8,118	24,633	39,180	22,917	1
1999.....	98,086	7	6	6	18	42	308	1,933	8,389	25,452	39,553	22,369	3
Non-Hispanic 1/, male													
2003.....	49,820	2	2	1	12	20	199	1,252	4,924	12,622	20,068	10,718	-
2002.....	49,155	5	3	3	5	22	206	1,224	4,483	12,384	20,244	10,572	4
2001.....	48,070	4	1	2	15	22	190	1,152	4,388	12,674	19,642	9,980	-
2000.....	47,906	4	-	2	8	10	175	1,094	4,256	12,822	19,566	9,968	1
1999.....	49,436	4	2	4	9	14	162	1,036	4,489	13,628	20,112	9,974	2
Non-Hispanic 1/, female													
2003.....	53,968	-	2	1	5	15	211	1,090	4,559	11,996	21,296	14,793	-
2002.....	52,258	5	1	5	8	22	191	1,069	4,273	11,719	20,790	14,174	1
2001.....	50,960	3	1	1	8	18	205	999	4,113	11,919	19,854	13,839	-
2000.....	49,399	1	2	1	4	16	173	966	3,862	11,811	19,614	12,949	-
1999.....	48,650	3	4	2	9	28	146	897	3,900	11,824	19,441	12,395	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
Non-Hispanic white, both sexes													
2003.....	96,722	2	2	2	9	24	325	2,008	8,537	22,745	38,944	24,124	-
2002.....	94,430	6	2	5	12	26	323	1,959	7,873	22,315	38,543	23,362	4
2001.....	92,267	4	1	2	12	29	314	1,835	7,685	22,734	37,199	22,452	-
2000.....	90,746	4	2	3	7	16	272	1,748	7,281	22,803	36,976	21,633	1
1999.....	91,269	1	3	5	12	28	238	1,649	7,524	23,501	37,153	21,153	2
Non-Hispanic white, male													
2003.....	45,758	2	-	1	6	13	156	1,070	4,382	11,516	18,598	10,014	-
2002.....	45,172	4	2	1	4	16	169	1,030	3,964	11,342	18,779	9,858	3
2001.....	44,173	2	-	1	7	18	152	979	3,917	11,572	18,246	9,279	-
2000.....	44,077	4	-	2	4	4	140	908	3,774	11,705	18,230	9,305	1
1999.....	45,423	-	-	3	4	13	123	867	3,969	12,471	18,653	9,319	1
Non-Hispanic white, female													
2003.....	50,964	-	2	1	3	11	169	938	4,155	11,229	20,346	14,110	-
2002.....	49,258	2	-	4	8	10	154	929	3,909	10,973	19,764	13,504	1
2001.....	48,094	2	1	1	5	11	162	856	3,768	11,162	18,953	13,173	-
2000.....	46,669	-	2	1	3	12	132	840	3,507	11,098	18,746	12,328	-
1999.....	45,846	1	3	2	8	15	115	782	3,555	11,030	18,500	11,834	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
Non-Hispanic black, both sexes													
2003.....	5,734	-	1	-	6	9	71	302	819	1,574	1,923	1,029	-
2002.....	5,739	4	1	3	1	15	68	304	776	1,487	2,012	1,067	1
2001.....	5,550	2	1	1	8	9	75	283	707	1,567	1,861	1,036	-
2000.....	5,414	1	-	-	3	10	71	278	748	1,549	1,771	983	-
1999.....	5,695	6	3	1	6	11	68	258	770	1,639	1,961	971	1
Non-Hispanic black, male													
2003.....	3,239	-	1	-	5	6	35	164	470	924	1,140	494	-
2002.....	3,221	1	-	2	1	4	34	176	461	865	1,153	523	1
2001.....	3,181	1	1	1	6	4	37	151	413	938	1,116	513	-
2000.....	3,126	-	-	-	3	6	33	161	432	953	1,045	493	-
1999.....	3,341	4	2	1	5	-	38	151	469	975	1,181	514	1
Non-Hispanic black, female													
2003.....	2,495	-	-	-	1	3	36	138	349	650	783	535	-
2002.....	2,518	3	1	1	-	11	34	128	315	622	859	544	-
2001.....	2,369	1	-	-	2	5	38	132	294	629	745	523	-
2000.....	2,288	1	-	-	-	4	38	117	316	596	726	490	-
1999.....	2,354	2	1	-	1	11	30	107	301	664	780	457	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)													
Origin not stated 2/, both sexes													
2003.....	261	-	-	-	-	-	-	14	40	68	85	54	-
2002.....	273	-	-	-	-	1	3	9	26	70	114	50	-
2001.....	278	-	-	-	-	-	4	14	37	74	92	56	1
2000.....	271	-	-	-	-	-	2	13	38	74	94	49	1
1999.....	284	-	-	-	-	-	-	9	32	79	104	59	1
Origin not stated 2/, male													
2003.....	139	-	-	-	-	-	-	8	26	38	46	21	-
2002.....	151	-	-	-	-	1	1	3	14	45	68	19	-
2001.....	148	-	-	-	-	-	3	9	24	40	48	23	1
2000.....	165	-	-	-	-	-	1	8	24	50	57	24	1
1999.....	153	-	-	-	-	-	-	7	19	52	47	27	1
Origin not stated 2/, female													
2003.....	122	-	-	-	-	-	-	6	14	30	39	33	-
2002.....	122	-	-	-	-	-	2	6	12	25	46	31	-
2001.....	130	-	-	-	-	-	1	5	13	34	44	33	-
2000.....	106	-	-	-	-	-	1	5	14	24	37	25	-
1999.....	131	-	-	-	-	-	-	2	13	27	57	32	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
All origins, both sexes													
2003.....	1,114	-	-	-	1	1	7	14	70	233	487	301	-
2002.....	1,114	2	-	-	-	-	5	17	75	223	480	312	-
2001.....	1,136	-	1	-	-	1	5	11	64	266	516	272	-
2000.....	1,161	-	-	-	-	3	7	12	57	281	484	317	-
1999.....	1,113	-	3	-	1	-	7	15	57	274	503	253	-
All origins, male													
2003.....	1,065	-	-	-	1	1	6	13	67	222	469	286	-
2002.....	1,057	-	-	-	-	-	4	13	69	214	457	300	-
2001.....	1,095	-	1	-	-	1	5	9	62	261	494	262	-
2000.....	1,113	-	-	-	-	3	6	12	54	265	467	306	-
1999.....	1,079	-	3	-	1	-	5	15	57	269	486	243	-
All origins, female													
2003.....	49	-	-	-	-	-	1	1	3	11	18	15	-
2002.....	57	2	-	-	-	-	1	4	6	9	23	12	-
2001.....	41	-	-	-	-	-	-	2	2	5	22	10	-
2000.....	48	-	-	-	-	-	1	-	3	16	17	11	-
1999.....	34	-	-	-	-	-	2	-	-	5	17	10	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
Non-Hispanic 1/, both sexes													
2003.....	1,093	-	-	-	-	1	7	14	67	230	474	300	-
2002.....	1,078	2	-	-	-	-	4	15	70	211	473	303	-
2001.....	1,108	-	-	-	-	1	3	9	62	259	509	265	-
2000.....	1,133	-	-	-	-	2	6	11	56	276	470	312	-
1999.....	1,100	-	3	-	1	-	6	14	56	269	501	250	-
Non-Hispanic 1/, male													
2003.....	1,045	-	-	-	-	1	6	13	64	219	456	286	-
2002.....	1,023	-	-	-	-	-	3	12	64	203	450	291	-
2001.....	1,068	-	-	-	-	1	3	7	60	254	487	256	-
2000.....	1,086	-	-	-	-	2	5	11	53	260	453	302	-
1999.....	1,068	-	3	-	1	-	4	14	56	264	484	242	-
Non-Hispanic 1/, female													
2003.....	48	-	-	-	-	-	1	1	3	11	18	14	-
2002.....	55	2	-	-	-	-	1	3	6	8	23	12	-
2001.....	40	-	-	-	-	-	-	2	2	5	22	9	-
2000.....	47	-	-	-	-	-	1	-	3	16	17	10	-
1999.....	32	-	-	-	-	-	2	-	-	5	17	8	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
Non-Hispanic white, both sexes													
2003.....	1,037	-	-	-	-	-	5	12	64	218	447	291	-
2002.....	1,014	1	-	-	-	-	3	13	62	201	445	289	-
2001.....	1,046	-	-	-	-	1	3	5	53	245	487	252	-
2000.....	1,072	-	-	-	-	2	3	11	54	261	442	299	-
1999.....	1,024	-	-	-	-	-	3	10	46	250	482	233	-
Non-Hispanic white, male													
2003.....	992	-	-	-	-	-	5	11	61	207	430	278	-
2002.....	966	-	-	-	-	-	2	10	59	193	424	278	-
2001.....	1,008	-	-	-	-	1	3	5	51	240	465	243	-
2000.....	1,027	-	-	-	-	2	2	11	51	245	427	289	-
1999.....	995	-	-	-	-	-	2	10	46	246	465	226	-
Non-Hispanic white, female													
2003.....	45	-	-	-	-	-	-	1	3	11	17	13	-
2002.....	48	1	-	-	-	-	1	3	3	8	21	11	-
2001.....	38	-	-	-	-	-	-	-	2	5	22	9	-
2000.....	45	-	-	-	-	-	1	-	3	16	15	10	-
1999.....	29	-	-	-	-	-	1	-	-	4	17	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
Non-Hispanic black, both sexes													
2003.....	48	-	-	-	-	1	1	2	1	11	23	9	-
2002.....	56	1	-	-	-	-	1	2	6	10	22	14	-
2001.....	56	-	-	-	-	-	-	4	8	12	20	12	-
2000.....	54	-	-	-	-	-	3	-	2	15	23	11	-
1999.....	67	-	3	-	1	-	3	4	8	16	16	16	-
Non-Hispanic black, male													
2003.....	45	-	-	-	-	1	-	2	1	11	22	8	-
2002.....	50	-	-	-	-	-	1	2	4	10	20	13	-
2001.....	54	-	-	-	-	-	-	2	8	12	20	12	-
2000.....	52	-	-	-	-	-	3	-	2	15	21	11	-
1999.....	64	-	3	-	1	-	2	4	8	15	16	15	-
Non-Hispanic black, female													
2003.....	3	-	-	-	-	-	1	-	-	-	1	1	-
2002.....	6	1	-	-	-	-	-	-	2	-	2	1	-
2001.....	2	-	-	-	-	-	-	2	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	2	-	-
1999.....	3	-	-	-	-	-	1	-	-	1	-	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
All origins, both sexes													
2003.....	17,335	13	10	16	46	69	166	392	773	1,754	5,789	8,307	-
2002.....	17,593	2	7	16	44	74	192	351	671	1,793	5,900	8,543	-
2001.....	17,301	12	14	15	34	79	179	378	659	1,830	5,759	8,342	-
2000.....	16,636	10	11	13	37	62	181	366	598	1,899	5,505	7,954	-
1999.....	15,268	16	12	16	32	78	166	284	586	1,754	5,030	7,294	-
All origins, male													
2003.....	8,702	8	5	9	31	42	89	239	471	1,069	3,181	3,558	-
2002.....	8,876	-	3	9	27	42	119	226	393	1,083	3,243	3,731	-
2001.....	8,576	6	6	7	22	40	117	232	384	1,106	3,145	3,511	-
2000.....	8,390	4	5	9	23	35	115	225	354	1,195	3,038	3,387	-
1999.....	7,755	9	3	6	22	47	97	165	365	1,083	2,875	3,083	-
All origins, female													
2003.....	8,633	5	5	7	15	27	77	153	302	685	2,608	4,749	-
2002.....	8,717	2	4	7	17	32	73	125	278	710	2,657	4,812	-
2001.....	8,725	6	8	8	12	39	62	146	275	724	2,614	4,831	-
2000.....	8,246	6	6	4	14	27	66	141	244	704	2,467	4,567	-
1999.....	7,513	7	9	10	10	31	69	119	221	671	2,155	4,211	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
Hispanic, both sexes													
2003.....	524	2	3	2	6	5	9	18	35	80	146	218	-
2002.....	547	1	-	3	7	3	10	15	35	62	174	237	-
2001.....	555	-	3	5	3	4	10	18	33	69	152	258	-
2000.....	496	3	2	1	3	7	15	20	33	64	141	207	-
1999.....	413	2	2	5	2	4	3	8	26	57	118	186	-
Hispanic, male													
2003.....	263	2	2	1	5	3	6	13	20	47	84	80	-
2002.....	282	-	-	1	5	2	5	8	24	32	105	100	-
2001.....	283	-	1	2	1	3	5	10	22	42	81	116	-
2000.....	253	-	-	-	2	2	8	15	20	35	87	84	-
1999.....	195	1	1	1	-	3	1	4	14	31	65	74	-
Hispanic, female													
2003.....	261	-	1	1	1	2	3	5	15	33	62	138	-
2002.....	265	1	-	2	2	1	5	7	11	30	69	137	-
2001.....	272	-	2	3	2	1	5	8	11	27	71	142	-
2000.....	243	3	2	1	1	5	7	5	13	29	54	123	-
1999.....	218	1	1	4	2	1	2	4	12	26	53	112	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
Non-Hispanic 1/, both sexes													
2003.....	16,769	10	7	14	40	64	157	373	736	1,668	5,630	8,070	-
2002.....	17,006	1	7	13	36	71	182	336	634	1,723	5,717	8,286	-
2001.....	16,700	12	11	10	31	75	167	358	624	1,756	5,593	8,063	-
2000.....	16,094	7	9	12	34	55	166	343	562	1,833	5,346	7,727	-
1999.....	14,812	14	10	11	30	74	162	274	559	1,691	4,895	7,092	-
Non-Hispanic 1/, male													
2003.....	8,415	5	3	8	26	39	83	225	449	1,018	3,089	3,470	-
2002.....	8,565	-	3	8	21	40	114	218	367	1,045	3,130	3,619	-
2001.....	8,270	6	5	5	21	37	110	221	361	1,061	3,057	3,386	-
2000.....	8,107	4	5	9	21	33	107	207	333	1,158	2,937	3,293	-
1999.....	7,534	8	2	5	22	44	95	159	350	1,046	2,803	3,000	-
Non-Hispanic 1/, female													
2003.....	8,354	5	4	6	14	25	74	148	287	650	2,541	4,600	-
2002.....	8,441	1	4	5	15	31	68	118	267	678	2,587	4,667	-
2001.....	8,430	6	6	5	10	38	57	137	263	695	2,536	4,677	-
2000.....	7,987	3	4	3	13	22	59	136	229	675	2,409	4,434	-
1999.....	7,278	6	8	6	8	30	67	115	209	645	2,092	4,092	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
Non-Hispanic white, both sexes													
2003.....	14,845	4	6	11	23	51	115	307	579	1,397	4,978	7,374	-
2002.....	15,203	1	5	11	27	56	138	262	519	1,437	5,131	7,616	-
2001.....	14,843	9	9	6	22	58	131	280	518	1,469	5,012	7,329	-
2000.....	14,343	6	7	11	24	40	128	259	449	1,515	4,793	7,111	-
1999.....	13,139	11	6	7	20	53	118	209	446	1,400	4,361	6,508	-
Non-Hispanic white, male													
2003.....	7,454	1	3	6	16	30	65	187	354	857	2,741	3,194	-
2002.....	7,654	-	3	7	15	33	86	168	303	874	2,822	3,343	-
2001.....	7,411	6	3	3	17	32	87	179	298	902	2,765	3,119	-
2000.....	7,214	4	4	8	16	21	85	154	270	945	2,648	3,059	-
1999.....	6,642	6	1	4	14	30	70	122	276	860	2,492	2,767	-
Non-Hispanic white, female													
2003.....	7,391	3	3	5	7	21	50	120	225	540	2,237	4,180	-
2002.....	7,549	1	2	4	12	23	52	94	216	563	2,309	4,273	-
2001.....	7,432	3	6	3	5	26	44	101	220	567	2,247	4,210	-
2000.....	7,129	2	3	3	8	19	43	105	179	570	2,145	4,052	-
1999.....	6,497	5	5	3	6	23	48	87	170	540	1,869	3,741	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
Non-Hispanic black, both sexes													
2003.....	1,598	4	1	3	14	12	41	61	141	231	533	557	-
2002.....	1,528	-	2	2	8	14	38	64	98	250	499	553	-
2001.....	1,584	3	2	4	7	16	33	71	100	251	497	600	-
2000.....	1,507	1	2	1	9	14	37	79	102	289	468	505	-
1999.....	1,495	2	4	3	10	18	40	59	105	264	469	521	-
Non-Hispanic black, male													
2003.....	776	3	-	2	8	9	17	36	86	137	276	202	-
2002.....	755	-	-	1	5	6	23	42	56	152	253	217	-
2001.....	731	-	2	2	2	4	20	38	60	140	252	211	-
2000.....	757	-	1	1	4	11	21	52	55	190	243	179	-
1999.....	768	1	1	1	8	12	22	33	68	168	261	193	-
Non-Hispanic black, female													
2003.....	822	1	1	1	6	3	24	25	55	94	257	355	-
2002.....	773	-	2	1	3	8	15	22	42	98	246	336	-
2001.....	853	3	-	2	5	12	13	33	40	111	245	389	-
2000.....	750	1	1	-	5	3	16	27	47	99	225	326	-
1999.....	727	1	3	2	2	6	18	26	37	96	208	328	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)													
Origin not stated 2/, both sexes													
2003.....	42	1	-	-	-	-	-	1	2	6	13	19	-
2002.....	40	-	-	-	1	-	-	-	2	8	9	20	-
2001.....	46	-	-	-	-	-	2	2	2	5	14	21	-
2000.....	46	-	-	-	-	-	-	3	3	2	18	20	-
1999.....	43	-	-	-	-	-	1	2	1	6	17	16	-
Origin not stated 2/, male													
2003.....	24	1	-	-	-	-	-	1	2	4	8	8	-
2002.....	29	-	-	-	1	-	-	-	2	6	8	12	-
2001.....	23	-	-	-	-	-	2	1	1	3	7	9	-
2000.....	30	-	-	-	-	-	-	3	1	2	14	10	-
1999.....	26	-	-	-	-	-	1	2	1	6	7	9	-
Origin not stated 2/, female													
2003.....	18	-	-	-	-	-	-	-	-	2	5	11	-
2002.....	11	-	-	-	-	-	-	-	-	2	1	8	-
2001.....	23	-	-	-	-	-	-	1	1	2	7	12	-
2000.....	16	-	-	-	-	-	-	-	2	-	4	10	-
1999.....	17	-	-	-	-	-	-	-	-	-	10	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)													
All origins, both sexes													
2003.....	25,520	279	86	60	136	210	495	1,335	2,442	5,156	8,810	6,510	1
2002.....	25,039	278	85	84	126	185	528	1,288	2,440	5,237	8,658	6,130	-
2001.....	26,178	294	93	78	141	216	570	1,356	2,432	5,510	9,202	6,283	3
2000.....	25,535	273	87	74	131	253	615	1,332	2,411	5,448	8,900	6,008	3
1999.....	25,031	252	91	76	115	257	576	1,219	2,260	5,409	8,646	6,130	-
All origins, male													
2003.....	12,391	163	49	38	76	115	251	704	1,367	2,849	4,395	2,384	-
2002.....	12,057	164	56	39	68	88	281	656	1,339	2,929	4,261	2,176	-
2001.....	12,495	186	47	43	76	104	304	737	1,322	3,025	4,453	2,197	1
2000.....	12,391	154	45	38	73	127	335	712	1,324	2,961	4,415	2,205	2
1999.....	12,110	145	55	39	69	127	317	637	1,213	3,005	4,328	2,175	-
All origins, female													
2003.....	13,129	116	37	22	60	95	244	631	1,075	2,307	4,415	4,126	1
2002.....	12,982	114	29	45	58	97	247	632	1,101	2,308	4,397	3,954	-
2001.....	13,683	108	46	35	65	112	266	619	1,110	2,485	4,749	4,086	2
2000.....	13,144	119	42	36	58	126	280	620	1,087	2,487	4,485	3,803	1
1999.....	12,921	107	36	37	46	130	259	582	1,047	2,404	4,318	3,955	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Hispanic, both sexes													
2003.....	1,374	57	14	16	16	27	52	94	185	311	373	229	-
2002.....	1,273	43	18	14	11	27	39	99	164	308	348	202	-
2001.....	1,275	41	17	7	18	17	39	106	165	287	363	215	-
2000.....	1,225	43	17	10	15	34	56	81	167	273	325	204	-
1999.....	1,157	38	10	8	15	23	45	94	157	269	307	191	-
Hispanic, male													
2003.....	686	29	8	9	9	12	33	59	112	167	165	83	-
2002.....	635	21	14	8	9	18	23	48	92	181	156	65	-
2001.....	643	21	8	3	9	12	26	58	92	151	175	88	-
2000.....	608	23	8	4	8	19	34	46	96	139	165	66	-
1999.....	583	20	6	3	7	12	29	45	89	139	158	75	-
Hispanic, female													
2003.....	688	28	6	7	7	15	19	35	73	144	208	146	-
2002.....	638	22	4	6	2	9	16	51	72	127	192	137	-
2001.....	632	20	9	4	9	5	13	48	73	136	188	127	-
2000.....	617	20	9	6	7	15	22	35	71	134	160	138	-
1999.....	574	18	4	5	8	11	16	49	68	130	149	116	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)													
Non-Hispanic 1/, both sexes													
2003.....	24,094	221	72	43	120	180	443	1,237	2,251	4,833	8,425	6,268	1
2002.....	23,706	235	67	70	115	158	487	1,177	2,268	4,916	8,296	5,917	-
2001.....	24,835	251	76	71	123	198	528	1,245	2,265	5,203	8,815	6,057	3
2000.....	24,236	228	69	64	115	218	556	1,246	2,238	5,158	8,552	5,790	2
1999.....	23,802	211	81	68	100	233	529	1,120	2,092	5,127	8,315	5,926	-
Non-Hispanic 1/, male													
2003.....	11,675	133	41	29	67	101	218	642	1,250	2,674	4,222	2,298	-
2002.....	11,397	143	42	31	59	70	258	602	1,243	2,743	4,098	2,108	-
2001.....	11,818	165	39	40	67	92	276	677	1,228	2,860	4,265	2,108	1
2000.....	11,752	130	37	34	65	108	298	663	1,227	2,815	4,239	2,135	1
1999.....	11,483	123	49	36	62	115	286	589	1,117	2,857	4,153	2,096	-
Non-Hispanic 1/, female													
2003.....	12,419	88	31	14	53	79	225	595	1,001	2,159	4,203	3,970	1
2002.....	12,309	92	25	39	56	88	229	575	1,025	2,173	4,198	3,809	-
2001.....	13,017	86	37	31	56	106	252	568	1,037	2,343	4,550	3,949	2
2000.....	12,484	98	32	30	50	110	258	583	1,011	2,343	4,313	3,655	1
1999.....	12,319	88	32	32	38	118	243	531	975	2,270	4,162	3,830	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)													
Non-Hispanic white, both sexes													
2003.....	21,311	113	39	30	73	120	284	896	1,858	4,285	7,759	5,853	1
2002.....	20,879	109	39	43	70	105	339	859	1,850	4,329	7,619	5,517	-
2001.....	21,835	116	44	52	82	122	350	883	1,860	4,597	8,133	5,595	1
2000.....	21,397	122	46	43	71	148	385	928	1,847	4,557	7,866	5,384	-
1999.....	20,959	111	52	47	75	155	351	826	1,718	4,511	7,642	5,471	-
Non-Hispanic white, male													
2003.....	10,314	64	20	20	37	70	149	486	1,038	2,391	3,897	2,142	-
2002.....	10,043	67	26	19	39	45	178	446	1,023	2,448	3,782	1,970	-
2001.....	10,433	76	20	31	46	52	202	498	1,018	2,574	3,963	1,953	-
2000.....	10,349	77	22	24	38	74	205	493	1,021	2,508	3,913	1,974	-
1999.....	10,154	66	28	26	51	86	200	443	919	2,561	3,848	1,926	-
Non-Hispanic white, female													
2003.....	10,997	49	19	10	36	50	135	410	820	1,894	3,862	3,711	1
2002.....	10,836	42	13	24	31	60	161	413	827	1,881	3,837	3,547	-
2001.....	11,402	40	24	21	36	70	148	385	842	2,023	4,170	3,642	1
2000.....	11,048	45	24	19	33	74	180	435	826	2,049	3,953	3,410	-
1999.....	10,805	45	24	21	24	69	151	383	799	1,950	3,794	3,545	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic black, both sexes													
2003.....	2,207	97	27	7	41	54	139	304	329	410	493	306	-
2002.....	2,257	112	28	25	41	47	133	278	353	446	494	300	-
2001.....	2,462	116	26	17	38	67	163	326	343	482	527	355	2
2000.....	2,336	102	20	18	39	58	152	284	325	495	527	314	2
1999.....	2,339	90	26	19	21	73	160	258	317	503	523	349	-
Non-Hispanic black, male													
2003.....	1,083	60	18	6	27	27	59	138	183	217	239	109	-
2002.....	1,076	67	16	10	19	23	73	136	187	222	229	94	-
2001.....	1,116	76	14	8	19	34	68	157	173	222	230	114	1
2000.....	1,126	50	14	10	23	30	84	151	166	242	240	115	1
1999.....	1,081	52	19	9	9	27	74	132	164	237	235	123	-
Non-Hispanic black, female													
2003.....	1,124	37	9	1	14	27	80	166	146	193	254	197	-
2002.....	1,181	45	12	15	22	24	60	142	166	224	265	206	-
2001.....	1,346	40	12	9	19	33	95	169	170	260	297	241	1
2000.....	1,210	52	6	8	16	28	68	133	159	253	287	199	1
1999.....	1,258	38	7	10	12	46	86	126	153	266	288	226	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Origin not stated 2/, both sexes

2003.....	52	1	-	1	-	3	-	4	6	12	12	13	-
2002.....	60	-	-	-	-	-	2	12	8	13	14	11	-
2001.....	68	2	-	-	-	1	3	5	2	20	24	11	-
2000.....	74	2	1	-	1	1	3	5	6	17	23	14	1
1999.....	72	3	-	-	-	1	2	5	11	13	24	13	-

Origin not stated 2/, male

2003.....	30	1	-	-	-	2	-	3	5	8	8	3	-
2002.....	25	-	-	-	-	-	-	6	4	5	7	3	-
2001.....	34	-	-	-	-	-	2	2	2	14	13	1	-
2000.....	31	1	-	-	-	-	3	3	1	7	11	4	1
1999.....	44	2	-	-	-	-	2	3	7	9	17	4	-

Origin not stated 2/, female

2003.....	22	-	-	1	-	1	-	1	1	4	4	10	-
2002.....	35	-	-	-	-	-	2	6	4	8	7	8	-
2001.....	34	2	-	-	-	1	1	3	-	6	11	10	-
2000.....	43	1	1	-	1	1	-	2	5	10	12	10	-
1999.....	28	1	-	-	-	1	-	2	4	4	7	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
All origins, both sexes													
2003.....	3,913	2	1	3	7	25	101	281	383	587	1,279	1,244	-
2002.....	4,079	1	1	1	-	22	113	272	411	665	1,291	1,302	-
2001.....	4,491	-	1	5	6	32	118	284	418	718	1,438	1,470	1
2000.....	4,558	-	-	-	9	14	113	307	443	748	1,415	1,509	-
1999.....	4,604	2	2	-	2	21	123	255	422	836	1,500	1,439	2
All origins, male													
2003.....	1,880	-	1	1	6	13	60	192	237	346	606	418	-
2002.....	1,951	-	1	1	-	14	72	177	247	378	622	439	-
2001.....	2,074	-	1	2	3	21	87	186	273	397	649	454	1
2000.....	2,201	-	-	-	7	6	79	217	284	442	667	499	-
1999.....	2,189	-	-	-	2	12	76	169	265	504	712	448	1
All origins, female													
2003.....	2,033	2	-	2	1	12	41	89	146	241	673	826	-
2002.....	2,128	1	-	-	-	8	41	95	164	287	669	863	-
2001.....	2,417	-	-	3	3	11	31	98	145	321	789	1,016	-
2000.....	2,357	-	-	-	2	8	34	90	159	306	748	1,010	-
1999.....	2,415	2	2	-	-	9	47	86	157	332	788	991	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
Hispanic, both sexes													
2003.....	200	1	-	-	1	5	6	30	26	42	56	33	-
2002.....	187	-	-	-	-	1	5	20	32	39	53	37	-
2001.....	166	-	-	-	1	-	1	16	22	33	54	38	1
2000.....	186	-	-	-	-	-	9	17	28	36	50	46	-
1999.....	172	-	1	-	-	8	4	15	25	37	44	38	-
Hispanic, male													
2003.....	114	-	-	-	1	3	6	21	16	30	30	7	-
2002.....	115	-	-	-	-	1	3	17	23	28	27	16	-
2001.....	99	-	-	-	1	-	1	11	19	18	37	11	1
2000.....	97	-	-	-	-	-	6	14	23	21	17	16	-
1999.....	101	-	-	-	-	5	3	10	18	22	27	16	-
Hispanic, female													
2003.....	86	1	-	-	-	2	-	9	10	12	26	26	-
2002.....	72	-	-	-	-	-	2	3	9	11	26	21	-
2001.....	67	-	-	-	-	-	-	5	3	15	17	27	-
2000.....	89	-	-	-	-	-	3	3	5	15	33	30	-
1999.....	71	-	1	-	-	3	1	5	7	15	17	22	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
Non-Hispanic 1/, both sexes													
2003.....	3,692	1	1	3	6	20	94	246	354	544	1,215	1,208	-
2002.....	3,879	1	1	1	-	21	108	251	374	625	1,235	1,262	-
2001.....	4,308	-	1	5	5	32	117	267	393	681	1,380	1,427	-
2000.....	4,357	-	-	-	9	14	102	290	414	709	1,361	1,458	-
1999.....	4,415	2	1	-	2	13	118	239	393	796	1,451	1,398	2
Non-Hispanic 1/, male													
2003.....	1,753	-	1	1	5	10	53	166	219	315	574	409	-
2002.....	1,829	-	1	1	-	13	69	159	222	349	594	421	-
2001.....	1,966	-	1	2	2	21	86	175	252	375	609	443	-
2000.....	2,099	-	-	-	7	6	72	203	261	419	648	483	-
1999.....	2,079	-	-	-	2	7	72	158	244	481	682	432	1
Non-Hispanic 1/, female													
2003.....	1,939	1	-	2	1	10	41	80	135	229	641	799	-
2002.....	2,050	1	-	-	-	8	39	92	152	276	641	841	-
2001.....	2,342	-	-	3	3	11	31	92	141	306	771	984	-
2000.....	2,258	-	-	-	2	8	30	87	153	290	713	975	-
1999.....	2,336	2	1	-	-	6	46	81	149	315	769	966	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
Non-Hispanic white, both sexes													
2003.....	3,197	-	1	2	3	15	65	174	287	472	1,067	1,111	-
2002.....	3,394	1	1	-	-	11	72	192	288	537	1,123	1,169	-
2001.....	3,766	-	1	4	2	16	77	197	308	583	1,248	1,330	-
2000.....	3,825	-	-	-	9	9	70	210	343	609	1,233	1,342	-
1999.....	3,859	1	1	-	2	7	73	171	322	677	1,313	1,291	1
Non-Hispanic white, male													
2003.....	1,471	-	1	1	2	7	35	114	175	269	496	371	-
2002.....	1,569	-	1	-	-	9	47	116	169	296	539	392	-
2001.....	1,649	-	1	2	1	11	55	122	194	311	543	409	-
2000.....	1,799	-	-	-	7	4	49	150	217	357	580	435	-
1999.....	1,783	-	-	-	2	3	41	111	195	409	622	399	1
Non-Hispanic white, female													
2003.....	1,726	-	-	1	1	8	30	60	112	203	571	740	-
2002.....	1,825	1	-	-	-	2	25	76	119	241	584	777	-
2001.....	2,117	-	-	2	1	5	22	75	114	272	705	921	-
2000.....	2,026	-	-	-	2	5	21	60	126	252	653	907	-
1999.....	2,076	1	1	-	-	4	32	60	127	268	691	892	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
Non-Hispanic black, both sexes													
2003.....	376	1	-	1	1	5	23	68	59	53	107	58	-
2002.....	376	-	-	1	-	10	35	56	68	69	75	62	-
2001.....	403	-	-	1	2	15	35	59	68	73	83	67	-
2000.....	420	-	-	-	-	4	29	70	63	79	91	84	-
1999.....	443	1	-	-	-	4	43	60	65	92	98	79	1
Non-Hispanic black, male													
2003.....	219	-	-	-	1	3	13	49	41	36	57	19	-
2002.....	209	-	-	1	-	4	21	41	45	41	38	18	-
2001.....	238	-	-	-	1	10	27	46	47	47	39	21	-
2000.....	228	-	-	-	-	2	21	45	40	48	45	27	-
1999.....	240	-	-	-	-	3	30	43	46	55	39	24	-
Non-Hispanic black, female													
2003.....	157	1	-	1	-	2	10	19	18	17	50	39	-
2002.....	167	-	-	-	-	6	14	15	23	28	37	44	-
2001.....	165	-	-	1	1	5	8	13	21	26	44	46	-
2000.....	192	-	-	-	-	2	8	25	23	31	46	57	-
1999.....	203	1	-	-	-	1	13	17	19	37	59	55	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Peptic ulcer (K25-K28)													
Origin not stated 2/, both sexes													
2003.....	21	-	-	-	-	-	1	5	3	1	8	3	-
2002.....	13	-	-	-	-	-	-	1	5	1	3	3	-
2001.....	17	-	-	-	-	-	-	1	3	4	4	5	-
2000.....	15	-	-	-	-	-	2	-	1	3	4	5	-
1999.....	17	-	-	-	-	-	1	1	4	3	5	3	-
Origin not stated 2/, male													
2003.....	13	-	-	-	-	-	1	5	2	1	2	2	-
2002.....	7	-	-	-	-	-	-	1	2	1	1	2	-
2001.....	9	-	-	-	-	-	-	-	2	4	3	-	-
2000.....	5	-	-	-	-	-	1	-	-	2	2	-	-
1999.....	9	-	-	-	-	-	1	1	3	1	3	-	-
Origin not stated 2/, female													
2003.....	8	-	-	-	-	-	-	-	1	-	6	1	-
2002.....	6	-	-	-	-	-	-	-	3	-	2	1	-
2001.....	8	-	-	-	-	-	-	1	1	-	1	5	-
2000.....	10	-	-	-	-	-	1	-	1	1	2	5	-
1999.....	8	-	-	-	-	-	-	-	1	2	2	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of appendix (K35-K38)													
All origins, both sexes													
2003.....	439	4	5	11	11	16	25	50	43	64	132	78	-
2002.....	480	3	-	7	12	10	27	54	64	89	120	94	-
2001.....	428	1	8	6	15	15	26	40	45	89	90	93	-
2000.....	435	3	5	15	11	11	27	43	60	71	121	68	-
1999.....	390	3	2	26	5	6	24	49	48	69	96	62	-
All origins, male													
2003.....	258	2	2	5	9	9	16	30	30	46	75	34	-
2002.....	275	1	-	3	7	9	17	39	45	56	67	31	-
2001.....	252	-	7	5	8	11	17	28	30	57	52	37	-
2000.....	237	2	2	4	7	10	16	29	43	40	60	24	-
1999.....	236	1	2	15	5	5	17	37	34	38	47	35	-
All origins, female													
2003.....	181	2	3	6	2	7	9	20	13	18	57	44	-
2002.....	205	2	-	4	5	1	10	15	19	33	53	63	-
2001.....	176	1	1	1	7	4	9	12	15	32	38	56	-
2000.....	198	1	3	11	4	1	11	14	17	31	61	44	-
1999.....	154	2	-	11	-	1	7	12	14	31	49	27	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of appendix (K35-K38)													
Hispanic, both sexes													
2003.....	31	1	-	5	1	2	1	8	1	4	4	4	-
2002.....	28	1	-	2	3	1	5	1	6	3	4	2	-
2001.....	30	1	4	1	2	1	1	5	2	5	7	1	-
2000.....	25	1	1	4	5	2	1	1	5	2	3	-	-
1999.....	29	1	-	8	2	-	4	4	3	3	3	1	-
Hispanic, male													
2003.....	20	1	-	1	1	2	1	4	1	3	3	3	-
2002.....	16	-	-	2	1	1	2	1	5	2	2	-	-
2001.....	16	-	3	1	-	-	-	3	1	5	3	-	-
2000.....	16	-	1	3	3	2	-	1	4	-	2	-	-
1999.....	20	1	-	4	2	-	4	3	2	2	1	1	-
Hispanic, female													
2003.....	11	-	-	4	-	-	-	4	-	1	1	1	-
2002.....	12	1	-	-	2	-	3	-	1	1	2	2	-
2001.....	14	1	1	-	2	1	1	2	1	-	4	1	-
2000.....	9	1	-	1	2	-	1	-	1	2	1	-	-
1999.....	9	-	-	4	-	-	-	1	1	1	2	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of appendix (K35-K38)													
Non-Hispanic 1/, both sexes													
2003.....	407	3	5	6	10	14	24	41	42	60	128	74	-
2002.....	451	2	-	5	9	9	22	53	57	86	116	92	-
2001.....	398	-	4	5	13	14	25	35	43	84	83	92	-
2000.....	410	2	4	11	6	9	26	42	55	69	118	68	-
1999.....	358	2	2	17	3	6	20	44	45	65	93	61	-
Non-Hispanic 1/, male													
2003.....	237	1	2	4	8	7	15	25	29	43	72	31	-
2002.....	258	1	-	1	6	8	15	38	39	54	65	31	-
2001.....	236	-	4	4	8	11	17	25	29	52	49	37	-
2000.....	221	2	1	1	4	8	16	28	39	40	58	24	-
1999.....	213	-	2	10	3	5	13	33	32	35	46	34	-
Non-Hispanic 1/, female													
2003.....	170	2	3	2	2	7	9	16	13	17	56	43	-
2002.....	193	1	-	4	3	1	7	15	18	32	51	61	-
2001.....	162	-	-	1	5	3	8	10	14	32	34	55	-
2000.....	189	-	3	10	2	1	10	14	16	29	60	44	-
1999.....	145	2	-	7	-	1	7	11	13	30	47	27	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of appendix (K35-K38)													
Non-Hispanic white, both sexes													
2003.....	323	2	3	1	6	8	13	29	31	49	113	68	-
2002.....	377	1	-	2	4	7	13	39	50	69	108	84	-
2001.....	313	-	1	3	8	7	16	24	27	75	70	82	-
2000.....	323	2	3	9	3	2	17	28	45	52	97	65	-
1999.....	284	2	-	10	1	4	15	33	27	54	81	57	-
Non-Hispanic white, male													
2003.....	192	1	2	-	5	3	11	19	22	35	64	30	-
2002.....	210	-	-	-	2	6	11	28	35	43	60	25	-
2001.....	182	-	1	2	5	5	12	17	21	47	43	29	-
2000.....	167	2	1	1	2	2	9	18	30	31	49	22	-
1999.....	163	-	-	6	1	3	9	24	22	28	38	32	-
Non-Hispanic white, female													
2003.....	131	1	1	1	1	5	2	10	9	14	49	38	-
2002.....	167	1	-	2	2	1	2	11	15	26	48	59	-
2001.....	131	-	-	1	3	2	4	7	6	28	27	53	-
2000.....	156	-	2	8	1	-	8	10	15	21	48	43	-
1999.....	121	2	-	4	-	1	6	9	5	26	43	25	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Diseases of appendix (K35-K38)													
Non-Hispanic black, both sexes													
2003.....	70	-	2	5	4	5	11	8	10	9	11	5	-
2002.....	61	1	-	3	5	2	5	13	6	13	5	8	-
2001.....	72	-	2	2	5	6	8	9	16	7	11	6	-
2000.....	77	-	-	2	2	7	9	12	10	16	16	3	-
1999.....	67	-	2	7	1	2	5	10	17	9	11	3	-
Non-Hispanic black, male													
2003.....	37	-	-	4	3	3	4	4	6	7	5	1	-
2002.....	40	1	-	1	4	2	1	10	3	8	4	6	-
2001.....	47	-	2	2	3	6	5	7	8	4	5	5	-
2000.....	48	-	-	-	1	6	7	8	9	9	6	2	-
1999.....	47	-	2	4	1	2	4	9	9	6	8	2	-
Non-Hispanic black, female													
2003.....	33	-	2	1	1	2	7	4	4	2	6	4	-
2002.....	21	-	-	2	1	-	4	3	3	5	1	2	-
2001.....	25	-	-	-	2	-	3	2	8	3	6	1	-
2000.....	29	-	-	2	1	1	2	4	1	7	10	1	-
1999.....	20	-	-	3	-	-	1	1	8	3	3	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
All origins, both sexes													
2003.....	1,613	37	6	6	3	5	28	105	161	237	462	563	-
2002.....	1,595	29	3	7	5	11	35	77	126	225	489	588	-
2001.....	1,522	43	2	3	3	15	34	82	131	213	443	553	-
2000.....	1,522	46	4	3	5	5	31	84	113	210	466	555	-
1999.....	1,464	44	3	2	3	16	39	55	107	215	451	529	-
All origins, male													
2003.....	668	20	4	4	1	4	16	59	72	111	199	178	-
2002.....	599	18	3	5	4	7	17	35	56	103	189	162	-
2001.....	587	28	1	-	1	10	16	44	64	94	173	156	-
2000.....	586	28	3	3	3	1	13	45	49	92	195	154	-
1999.....	597	26	2	1	3	8	23	32	60	93	192	157	-
All origins, female													
2003.....	945	17	2	2	2	1	12	46	89	126	263	385	-
2002.....	996	11	-	2	1	4	18	42	70	122	300	426	-
2001.....	935	15	1	3	2	5	18	38	67	119	270	397	-
2000.....	936	18	1	-	2	4	18	39	64	118	271	401	-
1999.....	867	18	1	1	-	8	16	23	47	122	259	372	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
Hispanic, both sexes													
2003.....	88	3	2	-	-	-	2	14	15	15	23	14	-
2002.....	81	3	-	2	2	1	5	6	14	14	19	15	-
2001.....	92	9	-	-	2	1	7	8	9	12	24	20	-
2000.....	85	9	-	1	1	1	1	7	9	14	23	19	-
1999.....	66	3	1	-	-	-	3	4	5	16	17	17	-
Hispanic, male													
2003.....	38	2	2	-	-	-	1	8	5	7	7	6	-
2002.....	35	2	-	2	2	1	2	4	7	4	7	4	-
2001.....	32	7	-	-	1	-	3	3	3	4	6	5	-
2000.....	33	6	-	1	-	1	1	5	3	3	10	3	-
1999.....	27	2	1	-	-	-	2	3	4	6	6	3	-
Hispanic, female													
2003.....	50	1	-	-	-	-	1	6	10	8	16	8	-
2002.....	46	1	-	-	-	-	3	2	7	10	12	11	-
2001.....	60	2	-	-	1	1	4	5	6	8	18	15	-
2000.....	52	3	-	-	1	-	-	2	6	11	13	16	-
1999.....	39	1	-	-	-	-	1	1	1	10	11	14	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
Non-Hispanic 1/, both sexes													
2003.....	1,518	34	4	6	3	5	26	90	146	221	435	548	-
2002.....	1,504	26	3	5	3	10	29	71	111	209	465	572	-
2001.....	1,425	33	2	3	1	14	27	74	121	201	417	532	-
2000.....	1,431	37	4	2	4	4	30	76	103	194	442	535	-
1999.....	1,390	41	2	2	3	16	36	50	102	198	431	509	-
Non-Hispanic 1/, male													
2003.....	626	18	2	4	1	4	15	50	67	103	190	172	-
2002.....	558	16	3	3	2	6	14	31	49	98	178	158	-
2001.....	552	20	1	-	-	10	13	41	60	90	166	151	-
2000.....	550	22	3	2	3	-	12	40	45	88	184	151	-
1999.....	566	24	1	1	3	8	21	28	56	86	185	153	-
Non-Hispanic 1/, female													
2003.....	892	16	2	2	2	1	11	40	79	118	245	376	-
2002.....	946	10	-	2	1	4	15	40	62	111	287	414	-
2001.....	873	13	1	3	1	4	14	33	61	111	251	381	-
2000.....	881	15	1	-	1	4	18	36	58	106	258	384	-
1999.....	824	17	1	1	-	8	15	22	46	112	246	356	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
Non-Hispanic white, both sexes													
2003.....	1,327	26	1	3	2	3	18	70	116	192	389	507	-
2002.....	1,343	21	2	5	3	5	20	52	88	176	431	540	-
2001.....	1,248	22	2	1	-	12	18	59	97	173	374	490	-
2000.....	1,266	27	2	2	2	3	21	59	90	169	399	492	-
1999.....	1,227	31	2	2	2	10	27	36	89	167	390	471	-
Non-Hispanic white, male													
2003.....	544	14	1	2	1	2	10	40	57	88	172	157	-
2002.....	493	12	2	3	2	2	11	26	41	81	167	146	-
2001.....	469	11	1	-	-	8	10	29	46	81	149	134	-
2000.....	486	17	2	2	1	-	9	29	40	78	170	138	-
1999.....	505	18	1	1	2	5	18	22	53	79	171	135	-
Non-Hispanic white, female													
2003.....	783	12	-	1	1	1	8	30	59	104	217	350	-
2002.....	850	9	-	2	1	3	9	26	47	95	264	394	-
2001.....	779	11	1	1	-	4	8	30	51	92	225	356	-
2000.....	780	10	-	-	1	3	12	30	50	91	229	354	-
1999.....	722	13	1	1	-	5	9	14	36	88	219	336	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
Non-Hispanic black, both sexes													
2003.....	163	8	3	3	-	2	8	18	28	24	38	31	-
2002.....	137	4	1	-	-	3	8	17	21	27	30	26	-
2001.....	155	10	-	1	1	2	9	15	23	25	33	36	-
2000.....	142	7	2	-	2	1	8	13	12	21	36	40	-
1999.....	139	9	-	-	1	4	9	12	12	24	37	31	-
Non-Hispanic black, male													
2003.....	71	4	1	2	-	2	5	9	10	13	14	11	-
2002.....	56	3	1	-	-	2	3	5	7	15	11	9	-
2001.....	74	8	-	-	-	2	3	12	13	7	14	15	-
2000.....	52	3	1	-	2	-	2	9	4	9	11	11	-
1999.....	53	6	-	-	1	1	3	6	2	7	13	14	-
Non-Hispanic black, female													
2003.....	92	4	2	1	-	-	3	9	18	11	24	20	-
2002.....	81	1	-	-	-	1	5	12	14	12	19	17	-
2001.....	81	2	-	1	1	-	6	3	10	18	19	21	-
2000.....	90	4	1	-	-	1	6	4	8	12	25	29	-
1999.....	86	3	-	-	-	3	6	6	10	17	24	17	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Hernia (K40-K46)													
Origin not stated 2/, both sexes													
2003.....	7	-	-	-	-	-	-	1	-	1	4	1	-
2002.....	10	-	-	-	-	-	1	-	1	2	5	1	-
2001.....	5	1	-	-	-	-	-	-	1	-	2	1	-
2000.....	6	-	-	-	-	-	-	1	1	2	1	1	-
1999.....	8	-	-	-	-	-	-	1	-	1	3	3	-
Origin not stated 2/, male													
2003.....	4	-	-	-	-	-	-	1	-	1	2	-	-
2002.....	6	-	-	-	-	-	1	-	-	1	4	-	-
2001.....	3	1	-	-	-	-	-	-	1	-	1	-	-
2000.....	3	-	-	-	-	-	-	-	1	1	1	-	-
1999.....	4	-	-	-	-	-	-	1	-	1	1	1	-
Origin not stated 2/, female													
2003.....	3	-	-	-	-	-	-	-	-	-	2	1	-
2002.....	4	-	-	-	-	-	-	-	1	1	1	1	-
2001.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2000.....	3	-	-	-	-	-	-	1	-	1	-	1	-
1999.....	4	-	-	-	-	-	-	-	-	-	2	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
All origins, both sexes													
2003.....	27,503	2	-	-	16	358	3,020	7,466	6,428	5,406	3,858	946	3
2002.....	27,257	6	3	6	31	374	3,154	7,216	6,097	5,381	4,000	985	4
2001.....	27,035	3	-	2	26	387	3,336	7,259	5,750	5,486	3,801	979	6
2000.....	26,552	3	3	3	36	415	3,371	6,654	5,774	5,482	3,829	979	3
1999.....	26,259	5	2	2	26	407	3,302	6,368	5,637	5,642	3,901	965	2
All origins, male													
2003.....	17,912	2	-	-	8	252	2,033	5,514	4,537	3,265	1,901	397	3
2002.....	17,401	6	1	3	14	236	2,079	5,285	4,228	3,197	1,953	395	4
2001.....	17,393	1	-	2	12	259	2,206	5,295	4,028	3,335	1,833	416	6
2000.....	17,214	3	1	2	19	280	2,314	4,975	4,005	3,317	1,874	421	3
1999.....	17,115	4	-	1	11	257	2,314	4,829	3,943	3,454	1,905	395	2
All origins, female													
2003.....	9,591	-	-	-	8	106	987	1,952	1,891	2,141	1,957	549	-
2002.....	9,856	-	2	3	17	138	1,075	1,931	1,869	2,184	2,047	590	-
2001.....	9,642	2	-	-	14	128	1,130	1,964	1,722	2,151	1,968	563	-
2000.....	9,338	-	2	1	17	135	1,057	1,679	1,769	2,165	1,955	558	-
1999.....	9,144	1	2	1	15	150	988	1,539	1,694	2,188	1,996	570	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Hispanic, both sexes													
2003.....	3,382	1	-	-	2	100	469	927	802	624	397	60	-
2002.....	3,409	1	1	2	9	81	507	983	758	663	321	82	1
2001.....	3,301	-	-	-	5	81	530	951	721	618	335	59	1
2000.....	3,187	-	1	-	7	92	501	854	708	628	323	73	-
1999.....	2,972	-	-	-	8	80	518	787	638	583	285	73	-
Hispanic, male													
2003.....	2,387	1	-	-	2	79	380	768	616	332	186	23	-
2002.....	2,437	1	-	1	5	62	410	805	561	407	157	27	1
2001.....	2,410	-	-	-	2	67	425	782	555	386	170	22	1
2000.....	2,312	-	-	-	5	82	399	707	531	395	163	30	-
1999.....	2,143	-	-	-	5	62	421	647	481	360	136	31	-
Hispanic, female													
2003.....	995	-	-	-	-	21	89	159	186	292	211	37	-
2002.....	972	-	1	1	4	19	97	178	197	256	164	55	-
2001.....	891	-	-	-	3	14	105	169	166	232	165	37	-
2000.....	875	-	1	-	2	10	102	147	177	233	160	43	-
1999.....	829	-	-	-	3	18	97	140	157	223	149	42	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Non-Hispanic 1/, both sexes													
2003.....	24,030	1	-	-	14	257	2,541	6,508	5,596	4,771	3,454	886	2
2002.....	23,733	5	2	4	22	291	2,632	6,198	5,307	4,700	3,668	902	2
2001.....	23,629	3	-	1	21	305	2,796	6,280	5,002	4,846	3,455	917	3
2000.....	23,251	3	2	3	29	319	2,856	5,762	5,041	4,833	3,498	903	2
1999.....	23,185	5	2	2	18	326	2,767	5,547	4,978	5,044	3,606	888	2
Non-Hispanic 1/, male													
2003.....	15,452	1	-	-	6	172	1,646	4,719	3,894	2,925	1,713	374	2
2002.....	14,876	5	1	2	9	172	1,658	4,452	3,641	2,778	1,788	368	2
2001.....	14,914	1	-	1	10	192	1,777	4,493	3,450	2,934	1,661	392	3
2000.....	14,814	3	1	2	14	195	1,903	4,237	3,452	2,908	1,708	389	2
1999.....	14,896	4	-	1	6	194	1,881	4,155	3,444	3,081	1,765	363	2
Non-Hispanic 1/, female													
2003.....	8,578	-	-	-	8	85	895	1,789	1,702	1,846	1,741	512	-
2002.....	8,857	-	1	2	13	119	974	1,746	1,666	1,922	1,880	534	-
2001.....	8,715	2	-	-	11	113	1,019	1,787	1,552	1,912	1,794	525	-
2000.....	8,437	-	1	1	15	124	953	1,525	1,589	1,925	1,790	514	-
1999.....	8,289	1	2	1	12	132	886	1,392	1,534	1,963	1,841	525	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Non-Hispanic white, both sexes													
2003.....	20,602	-	-	-	7	186	2,068	5,310	4,706	4,256	3,239	828	2
2002.....	20,353	4	2	3	17	204	2,148	5,023	4,483	4,205	3,417	845	2
2001.....	20,065	3	-	1	5	227	2,212	5,035	4,189	4,307	3,222	862	2
2000.....	19,739	-	2	2	14	217	2,240	4,637	4,235	4,285	3,264	842	1
1999.....	19,602	4	-	1	7	243	2,148	4,430	4,117	4,482	3,347	821	2
Non-Hispanic white, male													
2003.....	13,254	-	-	-	3	126	1,366	3,894	3,285	2,627	1,603	348	2
2002.....	12,800	4	1	2	8	130	1,371	3,670	3,097	2,489	1,681	345	2
2001.....	12,609	1	-	1	4	145	1,403	3,645	2,875	2,606	1,557	370	2
2000.....	12,640	-	1	1	6	140	1,518	3,477	2,935	2,602	1,597	362	1
1999.....	12,613	4	-	-	3	151	1,478	3,365	2,885	2,746	1,641	338	2
Non-Hispanic white, female													
2003.....	7,348	-	-	-	4	60	702	1,416	1,421	1,629	1,636	480	-
2002.....	7,553	-	1	1	9	74	777	1,353	1,386	1,716	1,736	500	-
2001.....	7,456	2	-	-	1	82	809	1,390	1,314	1,701	1,665	492	-
2000.....	7,099	-	1	1	8	77	722	1,160	1,300	1,683	1,667	480	-
1999.....	6,989	-	-	1	4	92	670	1,065	1,232	1,736	1,706	483	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Non-Hispanic black, both sexes													
2003.....	2,582	1	-	-	3	41	323	936	700	401	135	42	-
2002.....	2,545	1	-	1	5	49	313	956	649	358	173	40	-
2001.....	2,733	-	-	-	10	45	420	993	649	412	169	34	1
2000.....	2,705	2	-	1	12	56	446	929	622	429	163	44	1
1999.....	2,793	-	2	1	10	54	453	905	691	432	193	52	-
Non-Hispanic black, male													
2003.....	1,696	1	-	-	2	25	188	664	492	233	70	21	-
2002.....	1,592	1	-	-	1	26	180	637	436	221	75	15	-
2001.....	1,819	-	-	-	2	25	273	702	462	268	75	11	1
2000.....	1,709	2	-	1	6	26	279	633	408	255	81	17	1
1999.....	1,812	-	-	1	3	27	293	649	453	273	96	17	-
Non-Hispanic black, female													
2003.....	886	-	-	-	1	16	135	272	208	168	65	21	-
2002.....	953	-	-	1	4	23	133	319	213	137	98	25	-
2001.....	914	-	-	-	8	20	147	291	187	144	94	23	-
2000.....	996	-	-	-	6	30	167	296	214	174	82	27	-
1999.....	981	-	2	-	7	27	160	256	238	159	97	35	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Origin not stated 2/, both sexes													
2003.....	91	-	-	-	-	1	10	31	30	11	7	-	1
2002.....	115	-	-	-	-	2	15	35	32	18	11	1	1
2001.....	105	-	-	1	-	1	10	28	27	22	11	3	2
2000.....	114	-	-	-	-	4	14	38	25	21	8	3	1
1999.....	102	-	-	-	-	1	17	34	21	15	10	4	-
Origin not stated 2/, male													
2003.....	73	-	-	-	-	1	7	27	27	8	2	-	1
2002.....	88	-	-	-	-	2	11	28	26	12	8	-	1
2001.....	69	-	-	1	-	-	4	20	23	15	2	2	2
2000.....	88	-	-	-	-	3	12	31	22	14	3	2	1
1999.....	76	-	-	-	-	1	12	27	18	13	4	1	-
Origin not stated 2/, female													
2003.....	18	-	-	-	-	-	3	4	3	3	5	-	-
2002.....	27	-	-	-	-	-	4	7	6	6	3	1	-
2001.....	36	-	-	-	-	1	6	8	4	7	9	1	-
2000.....	26	-	-	-	-	1	2	7	3	7	5	1	-
1999.....	26	-	-	-	-	-	5	7	3	2	6	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
All origins, both sexes													
2003.....	12,360	-	-	-	6	243	1,971	4,288	3,247	1,788	729	86	2
2002.....	12,121	-	1	-	8	242	1,971	4,137	3,083	1,849	730	98	2
2001.....	12,207	-	-	-	8	264	2,145	4,203	2,913	1,848	717	104	5
2000.....	12,109	-	-	-	8	279	2,226	3,888	2,969	1,891	751	95	2
1999.....	11,958	-	-	-	8	259	2,186	3,802	2,859	1,935	789	118	2
All origins, male													
2003.....	9,104	-	-	-	3	182	1,317	3,228	2,459	1,343	518	52	2
2002.....	8,814	-	1	-	6	162	1,305	3,074	2,337	1,367	495	65	2
2001.....	8,932	-	-	-	5	179	1,446	3,099	2,242	1,379	505	72	5
2000.....	8,985	-	-	-	5	202	1,533	2,968	2,216	1,433	561	65	2
1999.....	8,922	-	-	-	6	165	1,520	2,927	2,194	1,497	538	73	2
All origins, female													
2003.....	3,256	-	-	-	3	61	654	1,060	788	445	211	34	-
2002.....	3,307	-	-	-	2	80	666	1,063	746	482	235	33	-
2001.....	3,275	-	-	-	3	85	699	1,104	671	469	212	32	-
2000.....	3,124	-	-	-	3	77	693	920	753	458	190	30	-
1999.....	3,036	-	-	-	2	94	666	875	665	438	251	45	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
Hispanic, both sexes													
2003.....	1,712	-	-	-	1	67	332	551	447	216	90	8	-
2002.....	1,679	-	-	-	4	53	321	587	392	246	63	12	1
2001.....	1,729	-	-	-	-	65	341	582	396	240	94	10	1
2000.....	1,680	-	-	-	4	66	353	526	400	247	78	6	-
1999.....	1,547	-	-	-	2	50	342	499	356	213	69	16	-
Hispanic, male													
2003.....	1,425	-	-	-	1	53	273	487	380	157	69	5	-
2002.....	1,418	-	-	-	4	47	271	493	339	211	44	8	1
2001.....	1,457	-	-	-	-	55	282	494	338	210	68	9	1
2000.....	1,442	-	-	-	4	62	282	455	344	221	69	5	-
1999.....	1,294	-	-	-	2	41	276	426	307	180	52	10	-
Hispanic, female													
2003.....	287	-	-	-	-	14	59	64	67	59	21	3	-
2002.....	261	-	-	-	-	6	50	94	53	35	19	4	-
2001.....	272	-	-	-	-	10	59	88	58	30	26	1	-
2000.....	238	-	-	-	-	4	71	71	56	26	9	1	-
1999.....	253	-	-	-	-	9	66	73	49	33	17	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
Non-Hispanic 1/, both sexes													
2003.....	10,599	-	-	-	5	176	1,633	3,718	2,783	1,566	638	78	2
2002.....	10,377	-	1	-	4	188	1,639	3,527	2,671	1,597	663	86	1
2001.....	10,423	-	-	-	8	198	1,801	3,603	2,499	1,596	622	94	2
2000.....	10,370	-	-	-	4	211	1,865	3,337	2,554	1,639	671	88	1
1999.....	10,353	-	-	-	6	208	1,831	3,283	2,491	1,713	717	102	2
Non-Hispanic 1/, male													
2003.....	7,638	-	-	-	2	129	1,041	2,725	2,063	1,181	448	47	2
2002.....	7,342	-	1	-	2	114	1,027	2,562	1,981	1,150	447	57	1
2001.....	7,431	-	-	-	5	124	1,162	2,592	1,888	1,158	437	63	2
2000.....	7,492	-	-	-	1	139	1,243	2,492	1,859	1,208	490	59	1
1999.....	7,580	-	-	-	4	123	1,235	2,485	1,877	1,308	483	63	2
Non-Hispanic 1/, female													
2003.....	2,961	-	-	-	3	47	592	993	720	385	190	31	-
2002.....	3,035	-	-	-	2	74	612	965	690	447	216	29	-
2001.....	2,992	-	-	-	3	74	639	1,011	611	438	185	31	-
2000.....	2,878	-	-	-	3	72	622	845	695	431	181	29	-
1999.....	2,773	-	-	-	2	85	596	798	614	405	234	39	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
Non-Hispanic white, both sexes													
2003.....	8,902	-	-	-	3	131	1,325	3,068	2,342	1,368	592	71	2
2002.....	8,713	-	1	-	1	133	1,335	2,907	2,255	1,403	601	76	1
2001.....	8,635	-	-	-	2	146	1,431	2,935	2,092	1,381	566	80	2
2000.....	8,556	-	-	-	2	150	1,454	2,703	2,126	1,439	601	81	-
1999.....	8,470	-	-	-	3	155	1,414	2,649	2,045	1,480	636	86	2
Non-Hispanic white, male													
2003.....	6,467	-	-	-	1	96	858	2,262	1,753	1,039	413	43	2
2002.....	6,242	-	1	-	1	87	838	2,143	1,701	1,012	409	49	1
2001.....	6,209	-	-	-	2	96	922	2,152	1,583	997	401	54	2
2000.....	6,296	-	-	-	-	105	979	2,068	1,579	1,065	446	54	-
1999.....	6,288	-	-	-	2	95	967	2,041	1,566	1,131	431	53	2
Non-Hispanic white, female													
2003.....	2,435	-	-	-	2	35	467	806	589	329	179	28	-
2002.....	2,471	-	-	-	-	46	497	764	554	391	192	27	-
2001.....	2,426	-	-	-	-	50	509	783	509	384	165	26	-
2000.....	2,260	-	-	-	2	45	475	635	547	374	155	27	-
1999.....	2,182	-	-	-	1	60	447	608	479	349	205	33	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
Non-Hispanic black, both sexes													
2003.....	1,206	-	-	-	-	19	187	465	340	157	33	5	-
2002.....	1,173	-	-	-	3	26	169	461	315	143	48	8	-
2001.....	1,331	-	-	-	1	26	244	508	324	176	43	9	-
2000.....	1,370	-	-	-	-	34	285	505	321	169	50	5	1
1999.....	1,445	-	-	-	2	28	281	507	358	191	64	14	-
Non-Hispanic black, male													
2003.....	850	-	-	-	-	14	110	344	242	112	24	4	-
2002.....	783	-	-	-	1	14	103	314	212	103	30	6	-
2001.....	926	-	-	-	-	12	160	349	239	135	26	5	-
2000.....	910	-	-	-	-	17	185	339	208	123	33	4	1
1999.....	1,001	-	-	-	2	14	178	358	254	146	41	8	-
Non-Hispanic black, female													
2003.....	356	-	-	-	-	5	77	121	98	45	9	1	-
2002.....	390	-	-	-	2	12	66	147	103	40	18	2	-
2001.....	405	-	-	-	1	14	84	159	85	41	17	4	-
2000.....	460	-	-	-	-	17	100	166	113	46	17	1	-
1999.....	444	-	-	-	-	14	103	149	104	45	23	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Alcoholic liver disease (K70)													
Origin not stated 2/, both sexes													
2003.....	49	-	-	-	-	-	6	19	17	6	1	-	-
2002.....	65	-	-	-	-	1	11	23	20	6	4	-	-
2001.....	55	-	-	-	-	1	3	18	18	12	1	-	2
2000.....	59	-	-	-	-	2	8	25	15	5	2	1	1
1999.....	58	-	-	-	-	1	13	20	12	9	3	-	-
Origin not stated 2/, male													
2003.....	41	-	-	-	-	-	3	16	16	5	1	-	-
2002.....	54	-	-	-	-	1	7	19	17	6	4	-	-
2001.....	44	-	-	-	-	-	2	13	16	11	-	-	2
2000.....	51	-	-	-	-	1	8	21	13	4	2	1	1
1999.....	48	-	-	-	-	1	9	16	10	9	3	-	-
Origin not stated 2/, female													
2003.....	8	-	-	-	-	-	3	3	1	1	-	-	-
2002.....	11	-	-	-	-	-	4	4	3	-	-	-	-
2001.....	11	-	-	-	-	1	1	5	2	1	1	-	-
2000.....	8	-	-	-	-	1	-	4	2	1	-	-	-
1999.....	10	-	-	-	-	-	4	4	2	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
All origins, both sexes													
2003.....	15,143	2	-	-	10	115	1,049	3,178	3,181	3,618	3,129	860	1
2002.....	15,136	6	2	6	23	132	1,183	3,079	3,014	3,532	3,270	887	2
2001.....	14,828	3	-	2	18	123	1,191	3,056	2,837	3,638	3,084	875	1
2000.....	14,443	3	3	3	28	136	1,145	2,766	2,805	3,591	3,078	884	1
1999.....	14,301	5	2	2	18	148	1,116	2,566	2,778	3,707	3,112	847	-
All origins, male													
2003.....	8,808	2	-	-	5	70	716	2,286	2,078	1,922	1,383	345	1
2002.....	8,587	6	-	3	8	74	774	2,211	1,891	1,830	1,458	330	2
2001.....	8,461	1	-	2	7	80	760	2,196	1,786	1,956	1,328	344	1
2000.....	8,229	3	1	2	14	78	781	2,007	1,789	1,884	1,313	356	1
1999.....	8,193	4	-	1	5	92	794	1,902	1,749	1,957	1,367	322	-
All origins, female													
2003.....	6,335	-	-	-	5	45	333	892	1,103	1,696	1,746	515	-
2002.....	6,549	-	2	3	15	58	409	868	1,123	1,702	1,812	557	-
2001.....	6,367	2	-	-	11	43	431	860	1,051	1,682	1,756	531	-
2000.....	6,214	-	2	1	14	58	364	759	1,016	1,707	1,765	528	-
1999.....	6,108	1	2	1	13	56	322	664	1,029	1,750	1,745	525	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Hispanic, both sexes													
2003.....	1,670	1	-	-	1	33	137	376	355	408	307	52	-
2002.....	1,730	1	1	2	5	28	186	396	366	417	258	70	-
2001.....	1,572	-	-	-	5	16	189	369	325	378	241	49	-
2000.....	1,507	-	1	-	3	26	148	328	308	381	245	67	-
1999.....	1,425	-	-	-	6	30	176	288	282	370	216	57	-
Hispanic, male													
2003.....	962	1	-	-	1	26	107	281	236	175	117	18	-
2002.....	1,019	1	-	1	1	15	139	312	222	196	113	19	-
2001.....	953	-	-	-	2	12	143	288	217	176	102	13	-
2000.....	870	-	-	-	1	20	117	252	187	174	94	25	-
1999.....	849	-	-	-	3	21	145	221	174	180	84	21	-
Hispanic, female													
2003.....	708	-	-	-	-	7	30	95	119	233	190	34	-
2002.....	711	-	1	1	4	13	47	84	144	221	145	51	-
2001.....	619	-	-	-	3	4	46	81	108	202	139	36	-
2000.....	637	-	1	-	2	6	31	76	121	207	151	42	-
1999.....	576	-	-	-	3	9	31	67	108	190	132	36	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Non-Hispanic 1/, both sexes													
2003.....	13,431	1	-	-	9	81	908	2,790	2,813	3,205	2,816	808	-
2002.....	13,356	5	1	4	18	103	993	2,671	2,636	3,103	3,005	816	1
2001.....	13,206	3	-	1	13	107	995	2,677	2,503	3,250	2,833	823	1
2000.....	12,881	3	2	3	25	108	991	2,425	2,487	3,194	2,827	815	1
1999.....	12,832	5	2	2	12	118	936	2,264	2,487	3,331	2,889	786	-
Non-Hispanic 1/, male													
2003.....	7,814	1	-	-	4	43	605	1,994	1,831	1,744	1,265	327	-
2002.....	7,534	5	-	2	7	58	631	1,890	1,660	1,628	1,341	311	1
2001.....	7,483	1	-	1	5	68	615	1,901	1,562	1,776	1,224	329	1
2000.....	7,322	3	1	2	13	56	660	1,745	1,593	1,700	1,218	330	1
1999.....	7,316	4	-	1	2	71	646	1,670	1,567	1,773	1,282	300	-
Non-Hispanic 1/, female													
2003.....	5,617	-	-	-	5	38	303	796	982	1,461	1,551	481	-
2002.....	5,822	-	1	2	11	45	362	781	976	1,475	1,664	505	-
2001.....	5,723	2	-	-	8	39	380	776	941	1,474	1,609	494	-
2000.....	5,559	-	1	1	12	52	331	680	894	1,494	1,609	485	-
1999.....	5,516	1	2	1	10	47	290	594	920	1,558	1,607	486	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Non-Hispanic white, both sexes													
2003.....	11,700	-	-	-	4	55	743	2,242	2,364	2,888	2,647	757	-
2002.....	11,640	4	1	3	16	71	813	2,116	2,228	2,802	2,816	769	1
2001.....	11,430	3	-	1	3	81	781	2,100	2,097	2,926	2,656	782	-
2000.....	11,183	-	2	2	12	67	786	1,934	2,109	2,846	2,663	761	1
1999.....	11,132	4	-	1	4	88	734	1,781	2,072	3,002	2,711	735	-
Non-Hispanic white, male													
2003.....	6,787	-	-	-	2	30	508	1,632	1,532	1,588	1,190	305	-
2002.....	6,558	4	-	2	7	43	533	1,527	1,396	1,477	1,272	296	1
2001.....	6,400	1	-	1	2	49	481	1,493	1,292	1,609	1,156	316	-
2000.....	6,344	-	1	1	6	35	539	1,409	1,356	1,537	1,151	308	1
1999.....	6,325	4	-	-	1	56	511	1,324	1,319	1,615	1,210	285	-
Non-Hispanic white, female													
2003.....	4,913	-	-	-	2	25	235	610	832	1,300	1,457	452	-
2002.....	5,082	-	1	1	9	28	280	589	832	1,325	1,544	473	-
2001.....	5,030	2	-	-	1	32	300	607	805	1,317	1,500	466	-
2000.....	4,839	-	1	1	6	32	247	525	753	1,309	1,512	453	-
1999.....	4,807	-	-	1	3	32	223	457	753	1,387	1,501	450	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Non-Hispanic black, both sexes													
2003.....	1,376	1	-	-	3	22	136	471	360	244	102	37	-
2002.....	1,372	1	-	1	2	23	144	495	334	215	125	32	-
2001.....	1,402	-	-	-	9	19	176	485	325	236	126	25	1
2000.....	1,335	2	-	1	12	22	161	424	301	260	113	39	-
1999.....	1,348	-	2	1	8	26	172	398	333	241	129	38	-
Non-Hispanic black, male													
2003.....	846	1	-	-	2	11	78	320	250	121	46	17	-
2002.....	809	1	-	-	-	12	77	323	224	118	45	9	-
2001.....	893	-	-	-	2	13	113	353	223	133	49	6	1
2000.....	799	2	-	1	6	9	94	294	200	132	48	13	-
1999.....	811	-	-	1	1	13	115	291	199	127	55	9	-
Non-Hispanic black, female													
2003.....	530	-	-	-	1	11	58	151	110	123	56	20	-
2002.....	563	-	-	1	2	11	67	172	110	97	80	23	-
2001.....	509	-	-	-	7	6	63	132	102	103	77	19	-
2000.....	536	-	-	-	6	13	67	130	101	128	65	26	-
1999.....	537	-	2	-	7	13	57	107	134	114	74	29	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Origin not stated 2/, both sexes													
2003.....	42	-	-	-	-	1	4	12	13	5	6	-	1
2002.....	50	-	-	-	-	1	4	12	12	12	7	1	1
2001.....	50	-	-	1	-	-	7	10	9	10	10	3	-
2000.....	55	-	-	-	-	2	6	13	10	16	6	2	-
1999.....	44	-	-	-	-	-	4	14	9	6	7	4	-
Origin not stated 2/, male													
2003.....	32	-	-	-	-	1	4	11	11	3	1	-	1
2002.....	34	-	-	-	-	1	4	9	9	6	4	-	1
2001.....	25	-	-	1	-	-	2	7	7	4	2	2	-
2000.....	37	-	-	-	-	2	4	10	9	10	1	1	-
1999.....	28	-	-	-	-	-	3	11	8	4	1	1	-
Origin not stated 2/, female													
2003.....	10	-	-	-	-	-	-	1	2	2	5	-	-
2002.....	16	-	-	-	-	-	-	3	3	6	3	1	-
2001.....	25	-	-	-	-	-	5	3	2	6	8	1	-
2000.....	18	-	-	-	-	-	2	3	1	6	5	1	-
1999.....	16	-	-	-	-	-	1	3	1	2	6	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All origins, both sexes

2003.....	2,948	-	-	1	4	25	40	126	182	489	1,054	1,027	-
2002.....	2,979	-	-	-	9	21	54	129	216	504	992	1,054	-
2001.....	2,965	1	-	1	7	19	44	137	242	526	945	1,043	-
2000.....	2,810	-	-	2	9	19	53	109	189	450	926	1,053	-
1999.....	2,830	2	-	1	10	16	59	124	203	475	948	992	-

All origins, male

2003.....	1,318	-	-	-	4	11	21	64	106	265	519	328	-
2002.....	1,341	-	-	-	3	11	29	83	122	258	500	335	-
2001.....	1,341	1	-	-	3	9	25	79	142	269	464	349	-
2000.....	1,244	-	-	1	4	7	27	53	110	242	461	339	-
1999.....	1,248	1	-	1	5	4	30	58	120	252	447	330	-

All origins, female

2003.....	1,630	-	-	1	-	14	19	62	76	224	535	699	-
2002.....	1,638	-	-	-	6	10	25	46	94	246	492	719	-
2001.....	1,624	-	-	1	4	10	19	58	100	257	481	694	-
2000.....	1,566	-	-	1	5	12	26	56	79	208	465	714	-
1999.....	1,582	1	-	-	5	12	29	66	83	223	501	662	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)													
Hispanic, both sexes													
2003.....	178	-	-	-	2	2	6	13	14	44	61	36	-
2002.....	196	-	-	-	2	6	11	11	25	42	55	44	-
2001.....	197	-	-	-	2	3	8	13	23	51	47	50	-
2000.....	183	-	-	-	2	3	5	18	19	41	51	44	-
1999.....	185	-	-	-	-	3	11	14	20	34	57	46	-
Hispanic, male													
2003.....	94	-	-	-	2	-	3	11	7	24	36	11	-
2002.....	93	-	-	-	-	2	7	10	14	21	23	16	-
2001.....	92	-	-	-	-	1	3	5	16	25	25	17	-
2000.....	99	-	-	-	-	1	5	9	15	25	32	12	-
1999.....	84	-	-	-	-	1	6	6	12	15	29	15	-
Hispanic, female													
2003.....	84	-	-	-	-	2	3	2	7	20	25	25	-
2002.....	103	-	-	-	2	4	4	1	11	21	32	28	-
2001.....	105	-	-	-	2	2	5	8	7	26	22	33	-
2000.....	84	-	-	-	2	2	-	9	4	16	19	32	-
1999.....	101	-	-	-	-	2	5	8	8	19	28	31	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Non-Hispanic 1/, both sexes

2003.....	2,765	-	-	1	2	23	34	112	168	445	990	990	-
2002.....	2,776	-	-	-	7	15	43	118	190	460	935	1,008	-
2001.....	2,764	1	-	1	5	16	36	124	219	473	896	993	-
2000.....	2,619	-	-	2	7	16	48	91	170	405	873	1,007	-
1999.....	2,639	2	-	1	10	13	48	110	183	438	889	945	-

Non-Hispanic 1/, male

2003.....	1,223	-	-	-	2	11	18	52	99	241	483	317	-
2002.....	1,244	-	-	-	3	9	22	73	107	236	476	318	-
2001.....	1,248	1	-	-	3	8	22	74	126	243	439	332	-
2000.....	1,141	-	-	1	4	6	22	44	95	215	428	326	-
1999.....	1,163	1	-	1	5	3	24	52	108	236	418	315	-

Non-Hispanic 1/, female

2003.....	1,542	-	-	1	-	12	16	60	69	204	507	673	-
2002.....	1,532	-	-	-	4	6	21	45	83	224	459	690	-
2001.....	1,516	-	-	1	2	8	14	50	93	230	457	661	-
2000.....	1,478	-	-	1	3	10	26	47	75	190	445	681	-
1999.....	1,476	1	-	-	5	10	24	58	75	202	471	630	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Non-Hispanic white, both sexes

2003.....	2,391	-	-	1	1	16	22	72	128	377	878	896	-
2002.....	2,420	-	-	-	5	10	33	82	152	390	818	930	-
2001.....	2,368	1	-	1	3	12	25	86	168	381	787	904	-
2000.....	2,272	-	-	1	5	7	33	58	132	325	775	936	-
1999.....	2,299	-	-	1	6	10	26	85	134	361	805	871	-

Non-Hispanic white, male

2003.....	1,056	-	-	-	1	6	10	36	75	208	430	290	-
2002.....	1,087	-	-	-	3	6	17	49	87	207	424	294	-
2001.....	1,065	1	-	-	2	5	17	52	96	200	395	297	-
2000.....	1,012	-	-	1	3	4	17	31	77	181	391	307	-
1999.....	1,017	-	-	1	4	1	14	40	79	201	389	288	-

Non-Hispanic white, female

2003.....	1,335	-	-	1	-	10	12	36	53	169	448	606	-
2002.....	1,333	-	-	-	2	4	16	33	65	183	394	636	-
2001.....	1,303	-	-	1	1	7	8	34	72	181	392	607	-
2000.....	1,260	-	-	-	2	3	16	27	55	144	384	629	-
1999.....	1,282	-	-	-	2	9	12	45	55	160	416	583	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Non-Hispanic black, both sexes

2003.....	285	-	-	-	1	7	11	34	33	50	78	71	-
2002.....	266	-	-	-	1	5	8	29	28	54	91	50	-
2001.....	309	-	-	-	2	4	11	31	42	78	77	64	-
2000.....	274	-	-	1	2	8	14	27	29	61	75	57	-
1999.....	257	2	-	-	4	3	20	21	35	59	65	48	-

Non-Hispanic black, male

2003.....	126	-	-	-	1	5	8	14	18	23	37	20	-
2002.....	106	-	-	-	-	3	4	18	16	17	37	11	-
2001.....	130	-	-	-	1	3	5	18	24	34	21	24	-
2000.....	92	-	-	-	1	2	4	10	14	23	26	12	-
1999.....	102	1	-	-	1	2	9	9	19	25	23	13	-

Non-Hispanic black, female

2003.....	159	-	-	-	-	2	3	20	15	27	41	51	-
2002.....	160	-	-	-	1	2	4	11	12	37	54	39	-
2001.....	179	-	-	-	1	1	6	13	18	44	56	40	-
2000.....	182	-	-	1	1	6	10	17	15	38	49	45	-
1999.....	155	1	-	-	3	1	11	12	16	34	42	35	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Origin not stated 2/, both sexes

2003.....	5	-	-	-	-	-	-	1	-	-	3	1	-
2002.....	7	-	-	-	-	-	-	-	1	2	2	2	-
2001.....	4	-	-	-	-	-	-	-	-	2	2	-	-
2000.....	8	-	-	-	-	-	-	-	-	4	2	2	-
1999.....	6	-	-	-	-	-	-	-	-	3	2	1	-

Origin not stated 2/, male

2003.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	4	-	-	-	-	-	-	-	1	1	1	1	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	2	1	1	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-	-

Origin not stated 2/, female

2003.....	4	-	-	-	-	-	-	-	-	-	3	1	-
2002.....	3	-	-	-	-	-	-	-	-	1	1	1	-
2001.....	3	-	-	-	-	-	-	-	-	1	2	-	-
2000.....	4	-	-	-	-	-	-	-	-	2	1	1	-
1999.....	5	-	-	-	-	-	-	-	-	2	2	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
All origins, both sexes													
2003.....	42,453	181	14	22	95	282	796	2,001	3,806	7,345	14,093	13,816	2
2002.....	40,974	173	14	21	84	269	749	1,893	3,455	7,164	13,896	13,256	-
2001.....	39,480	133	16	18	92	225	768	1,822	3,284	7,356	13,097	12,668	1
2000.....	37,251	162	14	25	88	244	740	1,651	3,100	6,990	12,458	11,777	2
1999.....	35,525	166	15	31	72	252	707	1,480	2,864	6,841	11,927	11,170	-
All origins, male													
2003.....	20,481	106	7	9	59	147	459	1,193	2,065	3,785	7,049	5,600	2
2002.....	19,695	109	7	12	41	129	407	1,086	1,824	3,802	6,920	5,358	-
2001.....	18,852	76	7	6	45	112	414	1,053	1,714	3,826	6,485	5,114	-
2000.....	17,811	103	7	17	43	129	429	982	1,651	3,564	6,213	4,671	2
1999.....	17,016	90	9	18	33	125	402	846	1,530	3,504	6,031	4,428	-
All origins, female													
2003.....	21,972	75	7	13	36	135	337	808	1,741	3,560	7,044	8,216	-
2002.....	21,279	64	7	9	43	140	342	807	1,631	3,362	6,976	7,898	-
2001.....	20,628	57	9	12	47	113	354	769	1,570	3,530	6,612	7,554	1
2000.....	19,440	59	7	8	45	115	311	669	1,449	3,426	6,245	7,106	-
1999.....	18,509	76	6	13	39	127	305	634	1,334	3,337	5,896	6,742	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
Hispanic, both sexes													
2003.....	2,170	35	1	4	19	30	76	174	318	484	578	450	1
2002.....	1,898	35	5	5	9	33	65	157	267	427	532	363	-
2001.....	1,857	18	2	4	16	25	74	132	237	470	524	355	-
2000.....	1,646	25	4	3	14	26	53	109	232	404	429	346	1
1999.....	1,584	20	3	9	6	21	62	107	230	369	426	331	-
Hispanic, male													
2003.....	1,069	19	-	1	13	16	49	107	160	230	280	193	1
2002.....	937	21	1	3	5	12	38	88	141	214	273	141	-
2001.....	892	13	-	2	9	10	46	64	132	217	251	148	-
2000.....	805	16	3	2	7	11	36	69	115	194	206	145	1
1999.....	814	15	2	7	2	10	31	64	137	178	212	156	-
Hispanic, female													
2003.....	1,101	16	1	3	6	14	27	67	158	254	298	257	-
2002.....	961	14	4	2	4	21	27	69	126	213	259	222	-
2001.....	965	5	2	2	7	15	28	68	105	253	273	207	-
2000.....	841	9	1	1	7	15	17	40	117	210	223	201	-
1999.....	770	5	1	2	4	11	31	43	93	191	214	175	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
Non-Hispanic 1/, both sexes													
2003.....	40,172	143	13	18	76	252	717	1,821	3,472	6,848	13,481	13,330	1
2002.....	38,933	138	9	16	75	235	681	1,725	3,171	6,710	13,322	12,851	-
2001.....	37,511	113	14	14	76	198	691	1,677	3,036	6,866	12,538	12,288	-
2000.....	35,509	134	10	22	73	217	686	1,536	2,854	6,567	12,000	11,410	-
1999.....	33,830	145	12	22	66	231	643	1,368	2,621	6,451	11,474	10,797	-
Non-Hispanic 1/, male													
2003.....	19,358	87	7	8	46	131	409	1,082	1,895	3,546	6,753	5,393	1
2002.....	18,682	88	6	9	36	117	367	992	1,677	3,569	6,620	5,201	-
2001.....	17,908	61	7	4	36	100	366	982	1,577	3,598	6,220	4,957	-
2000.....	16,958	84	4	15	35	118	392	908	1,529	3,357	5,995	4,521	-
1999.....	16,143	74	7	11	31	115	370	777	1,384	3,312	5,808	4,254	-
Non-Hispanic 1/, female													
2003.....	20,814	56	6	10	30	121	308	739	1,577	3,302	6,728	7,937	-
2002.....	20,251	50	3	7	39	118	314	733	1,494	3,141	6,702	7,650	-
2001.....	19,603	52	7	10	40	98	325	695	1,459	3,268	6,318	7,331	-
2000.....	18,551	50	6	7	38	99	294	628	1,325	3,210	6,005	6,889	-
1999.....	17,687	71	5	11	35	116	273	591	1,237	3,139	5,666	6,543	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic white, both sexes

2003.....	31,497	80	9	11	38	128	340	977	2,197	4,891	11,179	11,647	-
2002.....	30,669	77	5	10	33	119	353	922	1,978	4,870	11,110	11,192	-
2001.....	29,449	58	11	6	45	100	359	904	1,861	4,984	10,398	10,723	-
2000.....	27,924	61	4	17	48	110	359	829	1,776	4,741	10,026	9,953	-
1999.....	26,418	81	10	13	36	116	319	717	1,636	4,718	9,504	9,268	-

Non-Hispanic white, male

2003.....	15,323	52	5	6	24	64	205	577	1,216	2,614	5,750	4,810	-
2002.....	14,883	49	3	6	17	64	191	533	1,045	2,694	5,687	4,594	-
2001.....	14,331	31	4	3	21	45	211	543	1,012	2,729	5,310	4,422	-
2000.....	13,567	41	2	12	25	62	206	491	990	2,574	5,141	4,023	-
1999.....	12,793	46	5	8	17	60	188	406	903	2,532	4,907	3,721	-

Non-Hispanic white, female

2003.....	16,174	28	4	5	14	64	135	400	981	2,277	5,429	6,837	-
2002.....	15,786	28	2	4	16	55	162	389	933	2,176	5,423	6,598	-
2001.....	15,118	27	7	3	24	55	148	361	849	2,255	5,088	6,301	-
2000.....	14,357	20	2	5	23	48	153	338	786	2,167	4,885	5,930	-
1999.....	13,625	35	5	5	19	56	131	311	733	2,186	4,597	5,547	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
Non-Hispanic black, both sexes													
2003.....	7,804	60	3	7	34	118	355	780	1,172	1,758	2,015	1,501	1
2002.....	7,410	55	4	5	36	108	306	745	1,072	1,670	1,941	1,468	-
2001.....	7,214	52	3	7	27	93	306	702	1,048	1,698	1,891	1,387	-
2000.....	6,853	69	5	4	24	99	305	664	980	1,637	1,776	1,290	-
1999.....	6,659	58	1	8	30	103	310	615	879	1,554	1,750	1,351	-
Non-Hispanic black, male													
2003.....	3,614	32	1	2	20	64	191	467	627	853	862	494	1
2002.....	3,389	35	3	2	15	49	161	428	573	794	809	520	-
2001.....	3,158	28	3	-	12	54	139	399	509	782	784	448	-
2000.....	3,045	40	2	2	10	50	175	394	484	706	759	423	-
1999.....	2,984	25	1	3	14	47	173	353	426	691	803	448	-
Non-Hispanic black, female													
2003.....	4,190	28	2	5	14	54	164	313	545	905	1,153	1,007	-
2002.....	4,021	20	1	3	21	59	145	317	499	876	1,132	948	-
2001.....	4,056	24	-	7	15	39	167	303	539	916	1,107	939	-
2000.....	3,808	29	3	2	14	49	130	270	496	931	1,017	867	-
1999.....	3,675	33	-	5	16	56	137	262	453	863	947	903	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Origin not stated 2/, both sexes													
2003.....	111	3	-	-	-	-	3	6	16	13	34	36	-
2002.....	143	-	-	-	-	1	3	11	17	27	42	42	-
2001.....	112	2	-	-	-	2	3	13	11	20	35	25	1
2000.....	96	3	-	-	1	1	1	6	14	19	29	21	1
1999.....	111	1	-	-	-	-	2	5	13	21	27	42	-
Origin not stated 2/, male													
2003.....	54	-	-	-	-	-	1	4	10	9	16	14	-
2002.....	76	-	-	-	-	-	2	6	6	19	27	16	-
2001.....	52	2	-	-	-	2	2	7	5	11	14	9	-
2000.....	48	3	-	-	1	-	1	5	7	13	12	5	1
1999.....	59	1	-	-	-	-	1	5	9	14	11	18	-
Origin not stated 2/, female													
2003.....	57	3	-	-	-	-	2	2	6	4	18	22	-
2002.....	67	-	-	-	-	1	1	5	11	8	15	26	-
2001.....	60	-	-	-	-	-	1	6	6	9	21	16	1
2000.....	48	-	-	-	-	1	-	1	7	6	17	16	-
1999.....	52	-	-	-	-	-	1	-	4	7	16	24	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
All origins, both sexes													
2003.....	148	8	2	1	4	2	4	7	14	30	50	26	-
2002.....	166	6	5	2	3	1	4	9	16	27	51	42	-
2001.....	156	8	4	1	-	3	8	6	13	27	48	38	-
2000.....	177	8	3	2	1	4	3	16	10	29	52	49	-
1999.....	182	3	2	5	2	5	5	11	16	37	62	34	-
All origins, male													
2003.....	65	7	1	-	4	1	2	3	7	15	20	5	-
2002.....	70	3	2	2	1	1	-	5	11	14	17	14	-
2001.....	73	4	3	-	-	2	6	5	9	10	19	15	-
2000.....	91	3	2	2	1	3	1	11	8	12	26	22	-
1999.....	90	3	2	2	2	4	2	6	6	17	32	14	-
All origins, female													
2003.....	83	1	1	1	-	1	2	4	7	15	30	21	-
2002.....	96	3	3	-	2	-	4	4	5	13	34	28	-
2001.....	83	4	1	1	-	1	2	1	4	17	29	23	-
2000.....	86	5	1	-	-	1	2	5	2	17	26	27	-
1999.....	92	-	-	3	-	1	3	5	10	20	30	20	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
Hispanic, both sexes													
2003.....	13	1	-	1	3	-	-	-	1	3	2	2	-
2002.....	10	-	2	-	-	-	2	-	2	2	2	-	-
2001.....	13	1	-	-	-	1	1	2	2	3	1	2	-
2000.....	9	2	1	-	-	-	-	2	-	2	1	1	-
1999.....	12	1	-	2	-	-	-	1	2	4	1	1	-
Hispanic, male													
2003.....	7	1	-	-	3	-	-	-	-	-	2	1	-
2002.....	4	-	-	-	-	-	-	-	2	1	1	-	-
2001.....	7	-	-	-	-	-	1	2	2	2	-	-	-
2000.....	4	-	1	-	-	-	-	1	-	-	1	1	-
1999.....	4	1	-	1	-	-	-	-	1	-	-	1	-
Hispanic, female													
2003.....	6	-	-	1	-	-	-	-	1	3	-	1	-
2002.....	6	-	2	-	-	-	2	-	-	1	1	-	-
2001.....	6	1	-	-	-	1	-	-	-	1	1	2	-
2000.....	5	2	-	-	-	-	-	1	-	2	-	-	-
1999.....	8	-	-	1	-	-	-	1	1	4	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
Non-Hispanic 1/, both sexes													
2003.....	135	7	2	-	1	2	4	7	13	27	48	24	-
2002.....	155	6	3	2	3	1	2	9	14	25	49	41	-
2001.....	142	7	4	1	-	2	7	4	11	24	46	36	-
2000.....	168	6	2	2	1	4	3	14	10	27	51	48	-
1999.....	170	2	2	3	2	5	5	10	14	33	61	33	-
Non-Hispanic 1/, male													
2003.....	58	6	1	-	1	1	2	3	7	15	18	4	-
2002.....	66	3	2	2	1	1	-	5	9	13	16	14	-
2001.....	65	4	3	-	-	2	5	3	7	8	18	15	-
2000.....	87	3	1	2	1	3	1	10	8	12	25	21	-
1999.....	86	2	2	1	2	4	2	6	5	17	32	13	-
Non-Hispanic 1/, female													
2003.....	77	1	1	-	-	1	2	4	6	12	30	20	-
2002.....	89	3	1	-	2	-	2	4	5	12	33	27	-
2001.....	77	3	1	1	-	-	2	1	4	16	28	21	-
2000.....	81	3	1	-	-	1	2	4	2	15	26	27	-
1999.....	84	-	-	2	-	1	3	4	9	16	29	20	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
Non-Hispanic white, both sexes													
2003.....	107	5	1	-	-	1	3	7	8	23	40	19	-
2002.....	119	5	2	2	1	-	1	5	12	19	40	32	-
2001.....	116	6	3	-	-	-	7	3	9	18	41	29	-
2000.....	132	4	-	1	1	3	1	4	10	20	45	43	-
1999.....	132	2	2	2	1	2	3	6	7	28	51	28	-
Non-Hispanic white, male													
2003.....	40	4	-	-	-	-	2	3	3	12	12	4	-
2002.....	52	2	1	2	1	-	-	4	8	10	13	11	-
2001.....	52	3	2	-	-	-	5	3	5	4	18	12	-
2000.....	68	3	-	1	1	2	1	4	8	9	22	17	-
1999.....	63	2	2	1	1	1	1	4	2	14	26	9	-
Non-Hispanic white, female													
2003.....	67	1	1	-	-	1	1	4	5	11	28	15	-
2002.....	67	3	1	-	-	-	1	1	4	9	27	21	-
2001.....	64	3	1	-	-	-	2	-	4	14	23	17	-
2000.....	64	1	-	-	-	1	-	-	2	11	23	26	-
1999.....	69	-	-	1	-	1	2	2	5	14	25	19	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
Non-Hispanic black, both sexes													
2003.....	18	1	-	-	-	1	1	-	3	4	6	2	-
2002.....	29	1	1	-	2	1	1	4	1	5	6	7	-
2001.....	21	-	1	1	-	2	-	1	1	6	5	4	-
2000.....	30	2	2	-	-	1	2	10	-	6	5	2	-
1999.....	28	-	-	1	1	2	2	4	5	3	6	4	-
Non-Hispanic black, male													
2003.....	12	1	-	-	-	1	-	-	2	3	5	-	-
2002.....	11	1	1	-	-	1	-	1	1	2	2	2	-
2001.....	9	-	1	-	-	2	-	-	1	4	-	1	-
2000.....	13	-	1	-	-	1	-	6	-	2	2	1	-
1999.....	16	-	-	-	1	2	1	2	2	2	3	3	-
Non-Hispanic black, female													
2003.....	6	-	-	-	-	-	1	-	1	1	1	2	-
2002.....	18	-	-	-	2	-	1	3	-	3	4	5	-
2001.....	12	-	-	1	-	-	-	1	-	2	5	3	-
2000.....	17	2	1	-	-	-	2	4	-	4	3	1	-
1999.....	12	-	-	1	-	-	1	2	3	1	3	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All origins, both sexes

2003.....	537	3	-	-	8	14	20	45	50	102	165	130	-
2002.....	553	2	-	3	4	23	26	53	52	85	178	127	-
2001.....	511	-	2	1	8	12	27	41	40	111	147	122	-
2000.....	556	1	-	4	9	17	34	39	59	103	172	118	-
1999.....	589	-	1	1	7	15	28	44	60	101	200	132	-

All origins, male

2003.....	269	1	-	-	5	5	14	35	22	48	80	59	-
2002.....	278	1	-	1	3	13	14	25	27	49	92	53	-
2001.....	264	-	1	-	5	1	13	24	25	63	84	48	-
2000.....	284	-	-	1	2	11	20	27	31	54	83	55	-
1999.....	304	-	-	1	6	10	18	25	32	45	107	60	-

All origins, female

2003.....	268	2	-	-	3	9	6	10	28	54	85	71	-
2002.....	275	1	-	2	1	10	12	28	25	36	86	74	-
2001.....	247	-	1	1	3	11	14	17	15	48	63	74	-
2000.....	272	1	-	3	7	6	14	12	28	49	89	63	-
1999.....	285	-	1	-	1	5	10	19	28	56	93	72	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Hispanic, both sexes

2003.....	35	1	-	-	1	2	2	5	7	5	9	3	-
2002.....	32	-	-	2	-	3	2	3	5	3	9	5	-
2001.....	34	-	1	-	1	1	6	4	4	6	6	5	-
2000.....	28	-	-	-	-	2	1	4	4	5	7	5	-
1999.....	33	-	1	-	1	3	2	2	6	8	5	5	-

Hispanic, male

2003.....	17	-	-	-	-	1	1	5	3	3	4	-	-
2002.....	15	-	-	-	-	3	-	-	4	-	6	2	-
2001.....	17	-	-	-	1	-	5	-	3	2	4	2	-
2000.....	17	-	-	-	-	1	-	3	3	2	3	5	-
1999.....	18	-	-	-	1	3	2	1	4	4	1	2	-

Hispanic, female

2003.....	18	1	-	-	1	1	1	-	4	2	5	3	-
2002.....	17	-	-	2	-	-	2	3	1	3	3	3	-
2001.....	17	-	1	-	-	1	1	4	1	4	2	3	-
2000.....	11	-	-	-	-	1	1	1	1	3	4	-	-
1999.....	15	-	1	-	-	-	-	1	2	4	4	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Non-Hispanic 1/, both sexes

2003.....	501	2	-	-	7	12	17	40	43	97	156	127	-
2002.....	520	2	-	1	4	19	24	50	47	82	169	122	-
2001.....	477	-	1	1	7	11	21	37	36	105	141	117	-
2000.....	528	1	-	4	9	15	33	35	55	98	165	113	-
1999.....	555	-	-	1	6	12	26	42	53	93	195	127	-

Non-Hispanic 1/, male

2003.....	251	1	-	-	5	4	12	30	19	45	76	59	-
2002.....	263	1	-	1	3	10	14	25	23	49	86	51	-
2001.....	247	-	1	-	4	1	8	24	22	61	80	46	-
2000.....	267	-	-	1	2	10	20	24	28	52	80	50	-
1999.....	285	-	-	1	5	7	16	24	27	41	106	58	-

Non-Hispanic 1/, female

2003.....	250	1	-	-	2	8	5	10	24	52	80	68	-
2002.....	257	1	-	-	1	9	10	25	24	33	83	71	-
2001.....	230	-	-	1	3	10	13	13	14	44	61	71	-
2000.....	261	1	-	3	7	5	13	11	27	46	85	63	-
1999.....	270	-	-	-	1	5	10	18	26	52	89	69	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Non-Hispanic white, both sexes

2003.....	394	2	-	-	2	6	13	26	35	74	128	108	-
2002.....	419	-	-	-	3	12	20	31	34	76	141	102	-
2001.....	384	-	1	-	6	8	9	17	24	87	123	109	-
2000.....	411	-	-	3	7	9	19	22	39	80	143	89	-
1999.....	436	-	-	1	3	8	13	30	36	72	169	104	-

Non-Hispanic white, male

2003.....	196	1	-	-	1	2	10	20	16	35	61	50	-
2002.....	218	-	-	-	2	7	11	16	19	45	72	46	-
2001.....	200	-	1	-	3	1	3	10	14	52	71	45	-
2000.....	209	-	-	1	1	6	12	15	20	43	72	39	-
1999.....	227	-	-	1	2	5	8	18	19	33	93	48	-

Non-Hispanic white, female

2003.....	198	1	-	-	1	4	3	6	19	39	67	58	-
2002.....	201	-	-	-	1	5	9	15	15	31	69	56	-
2001.....	184	-	-	-	3	7	6	7	10	35	52	64	-
2000.....	202	-	-	2	6	3	7	7	19	37	71	50	-
1999.....	209	-	-	-	1	3	5	12	17	39	76	56	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Non-Hispanic black, both sexes

2003.....	76	-	-	-	4	4	4	12	6	15	19	12	-
2002.....	80	2	-	-	1	6	3	18	8	5	20	17	-
2001.....	68	-	-	1	-	2	8	15	8	14	13	7	-
2000.....	90	1	-	1	2	4	10	12	12	14	19	15	-
1999.....	92	-	-	-	3	2	11	10	11	17	19	19	-

Non-Hispanic black, male

2003.....	39	-	-	-	3	1	2	9	2	7	10	5	-
2002.....	37	1	-	-	1	3	2	8	3	4	11	4	-
2001.....	36	-	-	-	-	-	5	12	6	7	5	1	-
2000.....	44	-	-	-	1	2	5	8	7	8	6	7	-
1999.....	46	-	-	-	3	1	6	6	6	6	10	8	-

Non-Hispanic black, female

2003.....	37	-	-	-	1	3	2	3	4	8	9	7	-
2002.....	43	1	-	-	-	3	1	10	5	1	9	13	-
2001.....	32	-	-	1	-	2	3	3	2	7	8	6	-
2000.....	46	1	-	1	1	2	5	4	5	6	13	8	-
1999.....	46	-	-	-	-	1	5	4	5	11	9	11	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
All origins, both sexes													
2003.....	41,737	170	11	21	82	266	771	1,944	3,741	7,206	13,870	13,653	2
2002.....	40,222	165	9	15	77	243	718	1,829	3,383	7,046	13,657	13,080	-
2001.....	38,784	125	9	16	82	209	732	1,770	3,226	7,211	12,898	12,505	1
2000.....	36,471	152	11	19	78	221	701	1,590	3,023	6,848	12,223	11,603	2
1999.....	34,719	163	12	25	63	230	673	1,422	2,784	6,697	11,652	10,998	-
All origins, male													
2003.....	20,135	98	5	9	49	141	443	1,151	2,036	3,719	6,947	5,535	2
2002.....	19,341	105	5	8	37	114	393	1,055	1,786	3,739	6,808	5,291	-
2001.....	18,509	72	3	6	40	109	394	1,022	1,680	3,752	6,381	5,050	-
2000.....	17,416	100	5	14	40	115	407	941	1,607	3,492	6,102	4,591	2
1999.....	16,611	87	7	15	25	110	382	814	1,492	3,440	5,888	4,351	-
All origins, female													
2003.....	21,602	72	6	12	33	125	328	793	1,705	3,487	6,923	8,118	-
2002.....	20,881	60	4	7	40	129	325	774	1,597	3,307	6,849	7,789	-
2001.....	20,275	53	6	10	42	100	338	748	1,546	3,459	6,517	7,455	1
2000.....	19,055	52	6	5	38	106	294	649	1,416	3,356	6,121	7,012	-
1999.....	18,108	76	5	10	38	120	291	608	1,292	3,257	5,764	6,647	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Hispanic, both sexes													
2003.....	2,121	33	1	3	15	28	74	168	310	476	567	445	1
2002.....	1,856	35	3	3	9	30	61	154	260	422	521	358	-
2001.....	1,806	17	1	4	15	23	66	125	230	460	517	348	-
2000.....	1,605	22	3	3	14	24	51	103	227	397	420	340	1
1999.....	1,535	19	2	7	5	18	60	104	222	356	417	325	-
Hispanic, male													
2003.....	1,044	18	-	1	10	15	48	101	157	227	274	192	1
2002.....	918	21	1	3	5	9	38	88	135	213	266	139	-
2001.....	867	13	-	2	8	10	39	62	127	213	247	146	-
2000.....	782	16	2	2	7	10	35	65	111	192	202	139	1
1999.....	790	14	2	6	1	7	29	63	132	174	209	153	-
Hispanic, female													
2003.....	1,077	15	1	2	5	13	26	67	153	249	293	253	-
2002.....	938	14	2	-	4	21	23	66	125	209	255	219	-
2001.....	939	4	1	2	7	13	27	63	103	247	270	202	-
2000.....	823	6	1	1	7	14	16	38	116	205	218	201	-
1999.....	745	5	-	1	4	11	31	41	90	182	208	172	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Non-Hispanic 1/, both sexes													
2003.....	39,506	134	10	18	67	238	695	1,770	3,415	6,717	13,269	13,172	1
2002.....	38,225	130	6	12	68	213	654	1,664	3,106	6,597	13,094	12,681	-
2001.....	36,867	106	8	12	67	184	663	1,632	2,985	6,731	12,347	12,132	-
2000.....	34,771	127	8	16	63	196	649	1,481	2,783	6,432	11,774	11,242	-
1999.....	33,074	143	10	18	58	212	611	1,313	2,550	6,320	11,208	10,631	-
Non-Hispanic 1/, male													
2003.....	19,038	80	5	8	39	126	395	1,046	1,869	3,483	6,657	5,329	1
2002.....	18,347	84	4	5	32	105	353	961	1,645	3,507	6,515	5,136	-
2001.....	17,591	57	3	4	32	97	353	953	1,548	3,528	6,121	4,895	-
2000.....	16,586	81	3	12	32	105	371	871	1,489	3,287	5,888	4,447	-
1999.....	15,763	72	5	9	24	103	352	746	1,352	3,252	5,668	4,180	-
Non-Hispanic 1/, female													
2003.....	20,468	54	5	10	28	112	300	724	1,546	3,234	6,612	7,843	-
2002.....	19,878	46	2	7	36	108	301	703	1,461	3,090	6,579	7,545	-
2001.....	19,276	49	5	8	35	87	310	679	1,437	3,203	6,226	7,237	-
2000.....	18,185	46	5	4	31	91	278	610	1,294	3,145	5,886	6,795	-
1999.....	17,311	71	5	9	34	109	259	567	1,198	3,068	5,540	6,451	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Non-Hispanic white, both sexes													
2003.....	30,973	73	7	11	36	121	323	941	2,153	4,791	11,004	11,513	-
2002.....	30,104	72	3	8	29	105	331	885	1,928	4,769	10,922	11,052	-
2001.....	28,932	52	6	6	38	91	343	881	1,825	4,873	10,233	10,584	-
2000.....	27,350	57	4	13	40	97	339	800	1,722	4,632	9,830	9,816	-
1999.....	25,831	79	8	10	32	104	302	679	1,592	4,614	9,279	9,132	-
Non-Hispanic white, male													
2003.....	15,080	47	4	6	23	62	193	552	1,197	2,566	5,675	4,755	-
2002.....	14,610	47	2	4	14	56	180	513	1,018	2,639	5,600	4,537	-
2001.....	14,076	28	1	3	18	44	203	528	993	2,672	5,221	4,365	-
2000.....	13,275	38	2	10	23	54	193	471	959	2,516	5,045	3,964	-
1999.....	12,496	44	3	6	14	53	179	383	882	2,484	4,787	3,661	-
Non-Hispanic white, female													
2003.....	15,893	26	3	5	13	59	130	389	956	2,225	5,329	6,758	-
2002.....	15,494	25	1	4	15	49	151	372	910	2,130	5,322	6,515	-
2001.....	14,856	24	5	3	20	47	140	353	832	2,201	5,012	6,219	-
2000.....	14,075	19	2	3	17	43	146	329	763	2,116	4,785	5,852	-
1999.....	13,335	35	5	4	18	51	123	296	710	2,130	4,492	5,471	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Non-Hispanic black, both sexes													
2003.....	7,704	59	3	7	29	113	350	767	1,163	1,736	1,989	1,487	1
2002.....	7,297	52	3	4	33	101	302	722	1,063	1,660	1,914	1,443	-
2001.....	7,119	52	2	5	26	89	298	686	1,038	1,678	1,871	1,374	-
2000.....	6,724	66	3	3	22	93	292	639	968	1,616	1,751	1,271	-
1999.....	6,527	58	1	7	26	99	297	600	860	1,533	1,720	1,326	-
Non-Hispanic black, male													
2003.....	3,559	31	1	2	16	62	189	457	623	841	847	489	1
2002.....	3,339	33	2	1	14	45	159	418	569	788	796	514	-
2001.....	3,111	28	2	-	12	52	134	387	502	771	778	445	-
2000.....	2,986	40	1	2	9	47	170	378	477	696	751	415	-
1999.....	2,920	25	1	3	10	44	166	345	418	682	789	437	-
Non-Hispanic black, female													
2003.....	4,145	28	2	5	13	51	161	310	540	895	1,142	998	-
2002.....	3,958	19	1	3	19	56	143	304	494	872	1,118	929	-
2001.....	4,008	24	-	5	14	37	164	299	536	907	1,093	929	-
2000.....	3,738	26	2	1	13	46	122	261	491	920	1,000	856	-
1999.....	3,607	33	-	4	16	55	131	255	442	851	931	889	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Renal failure (N17-N19)													
Origin not stated 2/, both sexes													
2003.....	110	3	-	-	-	-	2	6	16	13	34	36	-
2002.....	141	-	-	-	-	-	3	11	17	27	42	41	-
2001.....	111	2	-	-	-	2	3	13	11	20	34	25	1
2000.....	95	3	-	-	1	1	1	6	13	19	29	21	1
1999.....	110	1	-	-	-	-	2	5	12	21	27	42	-
Origin not stated 2/, male													
2003.....	53	-	-	-	-	-	-	4	10	9	16	14	-
2002.....	76	-	-	-	-	-	2	6	6	19	27	16	-
2001.....	51	2	-	-	-	2	2	7	5	11	13	9	-
2000.....	48	3	-	-	1	-	1	5	7	13	12	5	1
1999.....	58	1	-	-	-	-	1	5	8	14	11	18	-
Origin not stated 2/, female													
2003.....	57	3	-	-	-	-	2	2	6	4	18	22	-
2002.....	65	-	-	-	-	-	1	5	11	8	15	25	-
2001.....	60	-	-	-	-	-	1	6	6	9	21	16	1
2000.....	47	-	-	-	-	1	-	1	6	6	17	16	-
1999.....	52	-	-	-	-	-	1	-	4	7	16	24	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
All origins, both sexes													
2003.....	31	-	1	-	1	-	1	5	1	7	8	7	-
2002.....	33	-	-	1	-	2	1	2	4	6	10	7	-
2001.....	29	-	1	-	2	1	1	5	5	7	4	3	-
2000.....	47	1	-	-	-	2	2	6	8	10	11	7	-
1999.....	35	-	-	-	-	2	1	3	4	6	13	6	-
All origins, male													
2003.....	12	-	1	-	1	-	-	4	-	3	2	1	-
2002.....	6	-	-	1	-	1	-	1	-	-	3	-	-
2001.....	6	-	-	-	-	-	1	2	-	1	1	1	-
2000.....	20	-	-	-	-	-	1	3	5	6	2	3	-
1999.....	11	-	-	-	-	1	-	1	-	2	4	3	-
All origins, female													
2003.....	19	-	-	-	-	-	1	1	1	4	6	6	-
2002.....	27	-	-	-	-	1	1	1	4	6	7	7	-
2001.....	23	-	1	-	2	1	-	3	5	6	3	2	-
2000.....	27	1	-	-	-	2	1	3	3	4	9	4	-
1999.....	24	-	-	-	-	1	1	2	4	4	9	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
Hispanic, both sexes													
2003.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	-	-	1	1	1	1	-	-	-
2000.....	4	1	-	-	-	-	1	-	1	-	1	-	-
1999.....	4	-	-	-	-	-	-	-	-	1	3	-	-
Hispanic, male													
2003.....	1	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	1	-	1	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	-	2	-	-
Hispanic, female													
2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	-	-	1	1	1	-	-	-
2000.....	2	1	-	-	-	-	-	-	-	-	1	-	-
1999.....	2	-	-	-	-	-	-	-	-	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
Non-Hispanic 1/, both sexes													
2003.....	30	-	1	-	1	-	1	4	1	7	8	7	-
2002.....	33	-	-	1	-	2	1	2	4	6	10	7	-
2001.....	25	-	1	-	2	1	-	4	4	6	4	3	-
2000.....	42	-	-	-	-	2	1	6	6	10	10	7	-
1999.....	31	-	-	-	-	2	1	3	4	5	10	6	-
Non-Hispanic 1/, male													
2003.....	11	-	1	-	1	-	-	3	-	3	2	1	-
2002.....	6	-	-	1	-	1	-	1	-	-	3	-	-
2001.....	5	-	-	-	-	-	-	2	-	1	1	1	-
2000.....	18	-	-	-	-	-	-	3	4	6	2	3	-
1999.....	9	-	-	-	-	1	-	1	-	2	2	3	-
Non-Hispanic 1/, female													
2003.....	19	-	-	-	-	-	1	1	1	4	6	6	-
2002.....	27	-	-	-	-	1	1	1	4	6	7	7	-
2001.....	20	-	1	-	2	1	-	2	4	5	3	2	-
2000.....	24	-	-	-	-	2	1	3	2	4	8	4	-
1999.....	22	-	-	-	-	1	1	2	4	3	8	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
Non-Hispanic white, both sexes													
2003.....	23	-	1	-	-	-	1	3	1	3	7	7	-
2002.....	27	-	-	-	-	2	1	1	4	6	7	6	-
2001.....	17	-	1	-	1	1	-	3	3	6	1	1	-
2000.....	31	-	-	-	-	1	-	3	5	9	8	5	-
1999.....	19	-	-	-	-	2	1	2	1	4	5	4	-
Non-Hispanic white, male													
2003.....	7	-	1	-	-	-	-	2	-	1	2	1	-
2002.....	3	-	-	-	-	1	-	-	-	2	-	-	-
2001.....	3	-	-	-	-	-	-	2	-	1	-	-	-
2000.....	15	-	-	-	-	-	-	1	3	6	2	3	-
1999.....	7	-	-	-	-	1	-	1	-	1	1	3	-
Non-Hispanic white, female													
2003.....	16	-	-	-	-	-	1	1	1	2	5	6	-
2002.....	24	-	-	-	-	1	1	1	4	6	5	6	-
2001.....	14	-	1	-	1	1	-	1	3	5	1	1	-
2000.....	16	-	-	-	-	1	-	2	2	3	6	2	-
1999.....	12	-	-	-	-	1	1	1	1	3	4	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other disorders of kidney (N25,N27)													
Non-Hispanic black, both sexes													
2003.....	6	-	-	-	1	-	-	1	-	3	1	-	-
2002.....	4	-	-	1	-	-	-	1	-	-	1	1	-
2001.....	6	-	-	-	1	-	-	-	1	-	2	2	-
2000.....	9	-	-	-	-	1	1	3	-	1	1	2	-
1999.....	12	-	-	-	-	-	-	1	3	1	5	2	-
Non-Hispanic black, male													
2003.....	4	-	-	-	1	-	-	1	-	2	-	-	-
2002.....	2	-	-	1	-	-	-	1	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2000.....	2	-	-	-	-	-	-	2	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	1	1	-	-
Non-Hispanic black, female													
2003.....	2	-	-	-	-	-	-	-	-	1	1	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	1	-
2001.....	4	-	-	-	1	-	-	-	1	-	1	1	-
2000.....	7	-	-	-	-	1	1	1	-	1	1	2	-
1999.....	10	-	-	-	-	-	-	1	3	-	4	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
All origins, both sexes													
2003.....	823	4	2	1	10	16	36	62	62	105	247	278	-
2002.....	788	2	-	3	5	13	28	40	59	110	238	290	-
2001.....	748	2	4	-	5	11	33	46	46	87	249	265	-
2000.....	813	5	3	-	2	9	21	42	59	104	248	320	-
1999.....	846	3	1	1	4	9	28	55	59	133	242	311	-
All origins, male													
2003.....	246	3	-	1	2	1	10	25	21	31	74	78	-
2002.....	251	2	-	1	1	6	12	15	21	47	72	74	-
2001.....	213	1	2	-	2	3	10	11	12	40	76	56	-
2000.....	214	3	-	-	1	3	8	15	15	26	74	69	-
1999.....	263	1	-	-	1	-	7	19	27	44	83	81	-
All origins, female													
2003.....	577	1	2	-	8	15	26	37	41	74	173	200	-
2002.....	537	-	-	2	4	7	16	25	38	63	166	216	-
2001.....	535	1	2	-	3	8	23	35	34	47	173	209	-
2000.....	599	2	3	-	1	6	13	27	44	78	174	251	-
1999.....	583	2	1	1	3	9	21	36	32	89	159	230	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

Hispanic, both sexes													
2003.....	46	-	-	-	1	-	2	2	8	11	13	9	-
2002.....	47	1	-	2	-	3	2	3	2	12	14	8	-
2001.....	33	-	-	-	-	1	3	5	1	8	9	6	-
2000.....	39	1	-	-	-	-	2	4	4	7	10	11	-
1999.....	55	-	-	-	-	1	7	6	9	8	15	9	-
Hispanic, male													
2003.....	11	-	-	-	-	-	-	1	1	1	4	4	-
2002.....	19	1	-	-	-	2	1	2	1	4	5	3	-
2001.....	10	-	-	-	-	-	-	3	-	3	1	3	-
2000.....	11	1	-	-	-	-	-	3	1	3	2	1	-
1999.....	21	-	-	-	-	-	2	2	5	4	6	2	-
Hispanic, female													
2003.....	35	-	-	-	1	-	2	1	7	10	9	5	-
2002.....	28	-	-	2	-	1	1	1	1	8	9	5	-
2001.....	23	-	-	-	-	1	3	2	1	5	8	3	-
2000.....	28	-	-	-	-	-	2	1	3	4	8	10	-
1999.....	34	-	-	-	-	1	5	4	4	4	9	7	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
Non-Hispanic 1/, both sexes													
2003.....	774	4	2	1	9	16	33	59	54	94	233	269	-
2002.....	740	1	-	1	5	10	26	37	57	98	224	281	-
2001.....	714	2	4	-	5	10	30	41	45	79	239	259	-
2000.....	771	4	3	-	2	9	19	38	55	96	236	309	-
1999.....	786	3	1	1	4	8	20	49	49	125	225	301	-
Non-Hispanic 1/, male													
2003.....	233	3	-	1	2	1	9	23	20	30	70	74	-
2002.....	232	1	-	1	1	4	11	13	20	43	67	71	-
2001.....	203	1	2	-	2	3	10	8	12	37	75	53	-
2000.....	201	2	-	-	1	3	8	12	14	22	71	68	-
1999.....	240	1	-	-	1	-	4	17	22	40	76	79	-
Non-Hispanic 1/, female													
2003.....	541	1	2	-	7	15	24	36	34	64	163	195	-
2002.....	508	-	-	-	4	6	15	24	37	55	157	210	-
2001.....	511	1	2	-	3	7	20	33	33	42	164	206	-
2000.....	570	2	3	-	1	6	11	26	41	74	165	241	-
1999.....	546	2	1	1	3	8	16	32	27	85	149	222	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
Non-Hispanic white, both sexes													
2003.....	656	1	1	1	7	11	25	44	38	82	201	245	-
2002.....	645	1	-	-	1	7	20	23	44	89	203	257	-
2001.....	614	1	2	-	2	7	23	29	36	65	210	239	-
2000.....	647	4	3	-	1	7	10	26	44	74	198	280	-
1999.....	689	1	1	1	2	7	12	38	38	102	206	281	-
Non-Hispanic white, male													
2003.....	202	-	-	1	2	1	8	19	14	27	58	72	-
2002.....	200	1	-	-	1	4	9	10	16	36	58	65	-
2001.....	166	-	1	-	1	2	9	6	9	27	64	47	-
2000.....	169	2	-	-	-	3	4	9	12	13	62	64	-
1999.....	209	1	-	-	1	-	2	12	17	30	70	76	-
Non-Hispanic white, female													
2003.....	454	1	1	-	5	10	17	25	24	55	143	173	-
2002.....	445	-	-	-	-	3	11	13	28	53	145	192	-
2001.....	448	1	1	-	1	5	14	23	27	38	146	192	-
2000.....	478	2	3	-	1	4	6	17	32	61	136	216	-
1999.....	480	-	1	1	1	7	10	26	21	72	136	205	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
Non-Hispanic black, both sexes													
2003.....	95	3	1	-	2	2	7	12	14	10	24	20	-
2002.....	73	-	-	-	3	2	6	8	10	8	15	21	-
2001.....	80	1	-	-	1	3	7	10	7	13	22	16	-
2000.....	101	-	-	-	1	2	7	11	8	19	28	25	-
1999.....	85	1	-	-	2	1	6	10	11	20	16	18	-
Non-Hispanic black, male													
2003.....	29	3	-	-	-	-	1	4	5	3	11	2	-
2002.....	22	-	-	-	-	-	2	1	3	6	5	5	-
2001.....	28	1	-	-	1	1	1	2	3	9	6	4	-
2000.....	27	-	-	-	1	-	2	3	2	8	7	4	-
1999.....	29	-	-	-	-	-	2	5	5	10	4	3	-
Non-Hispanic black, female													
2003.....	66	-	1	-	2	2	6	8	9	7	13	18	-
2002.....	51	-	-	-	3	2	4	7	7	2	10	16	-
2001.....	52	-	-	-	-	2	6	8	4	4	16	12	-
2000.....	74	-	-	-	-	2	5	8	6	11	21	21	-
1999.....	56	1	-	-	2	1	4	5	6	10	12	15	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

Origin not stated 2/, both sexes													
2003.....	3	-	-	-	-	-	1	1	-	-	1	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	3	-	-	-	-	-	-	-	-	1	2	-	-
1999.....	5	-	-	-	-	-	1	-	1	-	2	1	-
Origin not stated 2/, male													
2003.....	2	-	-	-	-	-	1	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	1	1	-	-
1999.....	2	-	-	-	-	-	1	-	-	-	1	-	-
Origin not stated 2/, female													
2003.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	1	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	1	-	-
1999.....	3	-	-	-	-	-	-	-	1	-	1	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

All origins, both sexes													
2003.....	125	1	-	1	6	3	14	14	10	16	29	31	-
2002.....	114	1	-	-	2	4	17	11	6	8	29	36	-
2001.....	127	1	-	-	2	3	10	12	12	18	31	38	-
2000.....	120	2	-	1	2	6	15	9	8	21	29	27	-
1999.....	108	2	1	-	3	6	12	6	13	15	32	18	-
All origins, male													
2003.....
2002.....
2001.....
2000.....
1999.....
All origins, female													
2003.....	125	1	-	1	6	3	14	14	10	16	29	31	-
2002.....	114	1	-	-	2	4	17	11	6	8	29	36	-
2001.....	127	1	-	-	2	3	10	12	12	18	31	38	-
2000.....	120	2	-	1	2	6	15	9	8	21	29	27	-
1999.....	108	2	1	-	3	6	12	6	13	15	32	18	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

Hispanic, both sexes													
2003.....	9	-	-	-	1	1	1	1	-	3	1	1	-
2002.....	3	-	-	-	-	1	2	-	-	-	-	-	-
2001.....	5	-	-	-	-	-	-	2	-	2	-	1	-
2000.....	7	-	-	-	-	1	1	-	2	2	-	1	-
1999.....	7	-	-	-	-	-	1	2	1	-	1	2	-
Hispanic, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Hispanic, female													
2003.....	9	-	-	-	1	1	1	1	-	3	1	1	-
2002.....	3	-	-	-	-	1	2	-	-	-	-	-	-
2001.....	5	-	-	-	-	-	-	2	-	2	-	1	-
2000.....	7	-	-	-	-	1	1	-	2	2	-	1	-
1999.....	7	-	-	-	-	-	1	2	1	-	1	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

Non-Hispanic 1/, both sexes

2003.....	115	1	-	1	5	2	13	12	10	13	28	30	-
2002.....	110	1	-	-	2	3	15	11	6	8	28	36	-
2001.....	122	1	-	-	2	3	10	10	12	16	31	37	-
2000.....	111	2	-	1	2	5	14	7	6	19	29	26	-
1999.....	101	2	1	-	3	6	11	4	12	15	31	16	-

Non-Hispanic 1/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 1/, female

2003.....	115	1	-	1	5	2	13	12	10	13	28	30	-
2002.....	110	1	-	-	2	3	15	11	6	8	28	36	-
2001.....	122	1	-	-	2	3	10	10	12	16	31	37	-
2000.....	111	2	-	1	2	5	14	7	6	19	29	26	-
1999.....	101	2	1	-	3	6	11	4	12	15	31	16	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

Non-Hispanic white, both sexes													
2003.....	93	-	-	1	3	1	9	8	8	10	24	29	-
2002.....	88	1	-	-	-	2	7	7	5	7	27	32	-
2001.....	98	1	-	-	2	3	6	6	8	12	27	33	-
2000.....	87	1	-	-	2	3	6	6	5	15	26	23	-
1999.....	73	1	1	-	2	3	3	2	11	10	25	15	-
Non-Hispanic white, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic white, female													
2003.....	93	-	-	1	3	1	9	8	8	10	24	29	-
2002.....	88	1	-	-	-	2	7	7	5	7	27	32	-
2001.....	98	1	-	-	2	3	6	6	8	12	27	33	-
2000.....	87	1	-	-	2	3	6	6	5	15	26	23	-
1999.....	73	1	1	-	2	3	3	2	11	10	25	15	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

Non-Hispanic black, both sexes

2003.....	19	1	-	-	2	1	4	4	2	2	2	1	-
2002.....	20	-	-	-	2	1	8	3	1	1	1	3	-
2001.....	23	-	-	-	-	-	4	4	4	4	4	3	-
2000.....	21	1	-	1	-	1	7	1	1	4	3	2	-
1999.....	26	1	-	-	1	2	7	2	1	5	6	1	-

Non-Hispanic black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic black, female

2003.....	19	1	-	-	2	1	4	4	2	2	2	1	-
2002.....	20	-	-	-	2	1	8	3	1	1	1	3	-
2001.....	23	-	-	-	-	-	4	4	4	4	4	3	-
2000.....	21	1	-	1	-	1	7	1	1	4	3	2	-
1999.....	26	1	-	-	1	2	7	2	1	5	6	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

All origins, both sexes													
2003.....	545	1	116	229	166	30	2	1	-	-	-
2002.....	379	1	89	175	105	8	-	-	-	1	-
2001.....	416	3	110	190	106	6	-	-	-	1	-
2000.....	404	-	95	182	121	6	-	-	-	-	-
1999.....	406	-	97	181	121	7	-	-	-	-	-
All origins, male													
2003.....
2002.....
2001.....
2000.....
1999.....
All origins, female													
2003.....	545	1	116	229	166	30	2	1	-	-	-
2002.....	379	1	89	175	105	8	-	-	-	1	-
2001.....	416	3	110	190	106	6	-	-	-	1	-
2000.....	404	-	95	182	121	6	-	-	-	-	-
1999.....	406	-	97	181	121	7	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

Hispanic, both sexes													
2003.....	104	-	24	46	28	4	1	1	-	-	-
2002.....	70	-	19	28	21	2	-	-	-	-	-
2001.....	83	1	26	38	17	1	-	-	-	-	-
2000.....	84	-	24	36	23	1	-	-	-	-	-
1999.....	70	-	19	26	25	-	-	-	-	-	-
Hispanic, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Hispanic, female													
2003.....	104	-	24	46	28	4	1	1	-	-	-
2002.....	70	-	19	28	21	2	-	-	-	-	-
2001.....	83	1	26	38	17	1	-	-	-	-	-
2000.....	84	-	24	36	23	1	-	-	-	-	-
1999.....	70	-	19	26	25	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic 1/, both sexes													
2003.....	437	1	89	182	138	26	1	-	-	-	-
2002.....	305	1	69	145	83	6	-	-	-	1	-
2001.....	332	2	84	151	89	5	-	-	-	1	-
2000.....	319	-	71	146	97	5	-	-	-	-	-
1999.....	333	-	76	154	96	7	-	-	-	-	-
Non-Hispanic 1/, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic 1/, female													
2003.....	437	1	89	182	138	26	1	-	-	-	-
2002.....	305	1	69	145	83	6	-	-	-	1	-
2001.....	332	2	84	151	89	5	-	-	-	1	-
2000.....	319	-	71	146	97	5	-	-	-	-	-
1999.....	333	-	76	154	96	7	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic white, both sexes													
2003.....	213	-	41	87	66	19	-	-	-	-	-
2002.....	133	-	21	69	39	3	-	-	-	1	-
2001.....	156	1	38	77	37	2	-	-	-	1	-
2000.....	161	-	33	72	54	2	-	-	-	-	-
1999.....	152	-	26	74	48	4	-	-	-	-	-
Non-Hispanic white, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic white, female													
2003.....	213	-	41	87	66	19	-	-	-	-	-
2002.....	133	-	21	69	39	3	-	-	-	1	-
2001.....	156	1	38	77	37	2	-	-	-	1	-
2000.....	161	-	33	72	54	2	-	-	-	-	-
1999.....	152	-	26	74	48	4	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic black, both sexes

2003.....	190	1	42	82	58	7	-	-	-	-	-
2002.....	153	1	45	64	41	2	-	-	-	-	-
2001.....	154	1	45	65	40	3	-	-	-	-	-
2000.....	139	-	34	65	37	3	-	-	-	-	-
1999.....	158	-	46	70	40	2	-	-	-	-	-

Non-Hispanic black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic black, female

2003.....	190	1	42	82	58	7	-	-	-	-	-
2002.....	153	1	45	64	41	2	-	-	-	-	-
2001.....	154	1	45	65	40	3	-	-	-	-	-
2000.....	139	-	34	65	37	3	-	-	-	-	-
1999.....	158	-	46	70	40	2	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

Origin not stated 2/, both sexes

2003.....	4	-	3	1	-	-	-	-	-	-	-
2002.....	4	-	1	2	1	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-
1999.....	3	-	2	1	-	-	-	-	-	-	-

Origin not stated 2/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Origin not stated 2/, female

2003.....	4	-	3	1	-	-	-	-	-	-	-
2002.....	4	-	1	2	1	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-
1999.....	3	-	2	1	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy with abortive outcome (000-007)													
All origins, both sexes													
2003.....	40	-	11	19	9	1	-	-	-	-	-
2002.....	22	1	4	9	7	1	-	-	-	-	-
2001.....	38	2	13	14	9	-	-	-	-	-	-
2000.....	37	-	6	17	14	-	-	-	-	-	-
1999.....	31	-	8	15	8	-	-	-	-	-	-
All origins, male													
2003.....
2002.....
2001.....
2000.....
1999.....
All origins, female													
2003.....	40	-	11	19	9	1	-	-	-	-	-
2002.....	22	1	4	9	7	1	-	-	-	-	-
2001.....	38	2	13	14	9	-	-	-	-	-	-
2000.....	37	-	6	17	14	-	-	-	-	-	-
1999.....	31	-	8	15	8	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pregnancy with abortive outcome (000-007)

Hispanic, both sexes

2003.....	8	-	1	6	1	-	-	-	-	-	-
2002.....	1	-	-	1	-	-	-	-	-	-	-
2001.....	1	1	-	-	-	-	-	-	-	-	-
2000.....	6	-	2	3	1	-	-	-	-	-	-
1999.....	4	-	2	-	2	-	-	-	-	-	-

Hispanic, male

2003.....
2002.....
2001.....
2000.....
1999.....

Hispanic, female

2003.....	8	-	1	6	1	-	-	-	-	-	-
2002.....	1	-	-	1	-	-	-	-	-	-	-
2001.....	1	1	-	-	-	-	-	-	-	-	-
2000.....	6	-	2	3	1	-	-	-	-	-	-
1999.....	4	-	2	-	2	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Pregnancy with abortive outcome (000-007)

Non-Hispanic 1/, both sexes

2003.....	32	-	10	13	8	1	-	-	-	-	-
2002.....	20	1	4	8	6	1	-	-	-	-	-
2001.....	37	1	13	14	9	-	-	-	-	-	-
2000.....	30	-	4	14	12	-	-	-	-	-	-
1999.....	26	-	5	15	6	-	-	-	-	-	-

Non-Hispanic 1/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 1/, female

2003.....	32	-	10	13	8	1	-	-	-	-	-
2002.....	20	1	4	8	6	1	-	-	-	-	-
2001.....	37	1	13	14	9	-	-	-	-	-	-
2000.....	30	-	4	14	12	-	-	-	-	-	-
1999.....	26	-	5	15	6	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy with abortive outcome (000-007)													
Non-Hispanic white, both sexes													
2003.....	11	-	2	6	3	-	-	-	-	-	-
2002.....	6	-	1	4	1	-	-	-	-	-	-
2001.....	19	-	7	8	4	-	-	-	-	-	-
2000.....	12	-	-	5	7	-	-	-	-	-	-
1999.....	11	-	2	6	3	-	-	-	-	-	-
Non-Hispanic white, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic white, female													
2003.....	11	-	2	6	3	-	-	-	-	-	-
2002.....	6	-	1	4	1	-	-	-	-	-	-
2001.....	19	-	7	8	4	-	-	-	-	-	-
2000.....	12	-	-	5	7	-	-	-	-	-	-
1999.....	11	-	2	6	3	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Pregnancy with abortive outcome (000-007)													
Non-Hispanic black, both sexes													
2003.....	18	-	8	5	4	1	-	-	-	-	-
2002.....	11	1	2	4	4	-	-	-	-	-	-
2001.....	16	1	6	5	4	-	-	-	-	-	-
2000.....	18	-	4	9	5	-	-	-	-	-	-
1999.....	14	-	3	8	3	-	-	-	-	-	-
Non-Hispanic black, male													
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic black, female													
2003.....	18	-	8	5	4	1	-	-	-	-	-
2002.....	11	1	2	4	4	-	-	-	-	-	-
2001.....	16	1	6	5	4	-	-	-	-	-	-
2000.....	18	-	4	9	5	-	-	-	-	-	-
1999.....	14	-	3	8	3	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

All origins, both sexes

2003.....	505	1	105	210	157	29	2	1	-	-	-
2002.....	357	-	85	166	98	7	-	-	-	1	-
2001.....	378	1	97	176	97	6	-	-	-	1	-
2000.....	367	-	89	165	107	6	-	-	-	-	-
1999.....	375	-	89	166	113	7	-	-	-	-	-

All origins, male

2003.....
2002.....
2001.....
2000.....
1999.....

All origins, female

2003.....	505	1	105	210	157	29	2	1	-	-	-
2002.....	357	-	85	166	98	7	-	-	-	1	-
2001.....	378	1	97	176	97	6	-	-	-	1	-
2000.....	367	-	89	165	107	6	-	-	-	-	-
1999.....	375	-	89	166	113	7	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Hispanic, both sexes

2003.....	96	-	23	40	27	4	1	1	-	-	-
2002.....	69	-	19	27	21	2	-	-	-	-	-
2001.....	82	-	26	38	17	1	-	-	-	-	-
2000.....	78	-	22	33	22	1	-	-	-	-	-
1999.....	66	-	17	26	23	-	-	-	-	-	-

Hispanic, male

2003.....
2002.....
2001.....
2000.....
1999.....

Hispanic, female

2003.....	96	-	23	40	27	4	1	1	-	-	-
2002.....	69	-	19	27	21	2	-	-	-	-	-
2001.....	82	-	26	38	17	1	-	-	-	-	-
2000.....	78	-	22	33	22	1	-	-	-	-	-
1999.....	66	-	17	26	23	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Non-Hispanic 1/, both sexes

2003.....	405	1	79	169	130	25	1	-	-	-	-
2002.....	285	-	65	137	77	5	-	-	-	1	-
2001.....	295	1	71	137	80	5	-	-	-	1	-
2000.....	289	-	67	132	85	5	-	-	-	-	-
1999.....	307	-	71	139	90	7	-	-	-	-	-

Non-Hispanic 1/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 1/, female

2003.....	405	1	79	169	130	25	1	-	-	-	-
2002.....	285	-	65	137	77	5	-	-	-	1	-
2001.....	295	1	71	137	80	5	-	-	-	1	-
2000.....	289	-	67	132	85	5	-	-	-	-	-
1999.....	307	-	71	139	90	7	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Non-Hispanic white, both sexes

2003.....	202	-	39	81	63	19	-	-	-	-	-
2002.....	127	-	20	65	38	3	-	-	-	1	-
2001.....	137	1	31	69	33	2	-	-	-	1	-
2000.....	149	-	33	67	47	2	-	-	-	-	-
1999.....	141	-	24	68	45	4	-	-	-	-	-

Non-Hispanic white, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic white, female

2003.....	202	-	39	81	63	19	-	-	-	-	-
2002.....	127	-	20	65	38	3	-	-	-	1	-
2001.....	137	1	31	69	33	2	-	-	-	1	-
2000.....	149	-	33	67	47	2	-	-	-	-	-
1999.....	141	-	24	68	45	4	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Non-Hispanic black, both sexes

2003.....	172	1	34	77	54	6	-	-	-	-	-
2002.....	142	-	43	60	37	2	-	-	-	-	-
2001.....	138	-	39	60	36	3	-	-	-	-	-
2000.....	121	-	30	56	32	3	-	-	-	-	-
1999.....	144	-	43	62	37	2	-	-	-	-	-

Non-Hispanic black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic black, female

2003.....	172	1	34	77	54	6	-	-	-	-	-
2002.....	142	-	43	60	37	2	-	-	-	-	-
2001.....	138	-	39	60	36	3	-	-	-	-	-
2000.....	121	-	30	56	32	3	-	-	-	-	-
1999.....	144	-	43	62	37	2	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Origin not stated 2/, both sexes

2003.....	4	-	3	1	-	-	-	-	-	-	-
2002.....	3	-	1	2	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	1	1	-	-	-	-	-	-	-

Origin not stated 2/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Origin not stated 2/, female

2003.....	4	-	3	1	-	-	-	-	-	-	-
2002.....	3	-	1	2	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	1	1	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Certain conditions originating in the perinatal period (P00-P96)

All origins, both sexes

2003.....	14,378	14,254	79	19	16	3	3	2	-	2	-	-	-
2002.....	14,254	14,106	65	31	21	10	10	2	1	3	2	1	2
2001.....	13,887	13,734	72	36	17	6	9	5	2	3	1	1	1
2000.....	14,069	13,918	79	41	15	7	4	2	2	-	-	-	1
1999.....	14,259	14,084	92	43	16	7	3	2	3	2	1	1	5

All origins, male

2003.....	8,197	8,129	43	11	8	2	1	1	-	2	-	-	-
2002.....	8,047	7,960	40	19	12	4	5	2	1	1	1	-	2
2001.....	7,868	7,775	44	24	7	2	6	5	1	2	1	-	1
2000.....	7,968	7,872	52	23	11	6	2	2	-	-	-	-	-
1999.....	8,044	7,941	56	23	10	2	2	2	1	1	1	1	4

All origins, female

2003.....	6,181	6,125	36	8	8	1	2	1	-	-	-	-	-
2002.....	6,207	6,146	25	12	9	6	5	-	-	2	1	1	-
2001.....	6,019	5,959	28	12	10	4	3	-	1	1	-	1	-
2000.....	6,101	6,046	27	18	4	1	2	-	2	-	-	-	1
1999.....	6,215	6,143	36	20	6	5	1	-	2	1	-	-	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Certain conditions originating in the perinatal period (P00-P96)

Hispanic, both sexes													
2003.....	2,628	2,599	15	8	3	1	1	-	-	1	-	-	-
2002.....	2,402	2,375	14	6	3	1	2	-	-	-	-	1	-
2001.....	2,227	2,203	14	3	2	4	-	1	-	-	-	-	-
2000.....	2,145	2,120	12	10	2	-	1	-	-	-	-	-	-
1999.....	2,153	2,127	18	2	4	1	-	-	-	-	-	-	1
Hispanic, male													
2003.....	1,459	1,442	8	4	2	1	1	-	-	1	-	-	-
2002.....	1,352	1,335	9	5	2	1	-	-	-	-	-	-	-
2001.....	1,268	1,258	6	2	-	1	-	1	-	-	-	-	-
2000.....	1,194	1,180	7	6	1	-	-	-	-	-	-	-	-
1999.....	1,192	1,181	5	2	3	-	-	-	-	-	-	-	1
Hispanic, female													
2003.....	1,169	1,157	7	4	1	-	-	-	-	-	-	-	-
2002.....	1,050	1,040	5	1	1	-	2	-	-	-	-	1	-
2001.....	959	945	8	1	2	3	-	-	-	-	-	-	-
2000.....	951	940	5	4	1	-	1	-	-	-	-	-	-
1999.....	961	946	13	-	1	1	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Certain conditions originating in the perinatal period (P00-P96)

Non-Hispanic 1/, both sexes

2003.....	11,559	11,464	64	11	13	2	2	2	-	1	-	-	-
2002.....	11,673	11,553	50	25	18	9	8	2	1	3	2	-	2
2001.....	11,500	11,371	58	33	15	2	9	4	2	3	1	1	1
2000.....	11,672	11,546	67	31	13	7	3	2	2	-	-	-	1
1999.....	11,815	11,668	73	41	12	6	3	2	3	2	1	1	3

Non-Hispanic 1/, male

2003.....	6,622	6,571	35	7	6	1	-	1	-	1	-	-	-
2002.....	6,596	6,527	30	14	10	3	5	2	1	1	1	-	2
2001.....	6,511	6,428	38	22	7	1	6	4	1	2	1	-	1
2000.....	6,637	6,555	45	17	10	6	2	2	-	-	-	-	-
1999.....	6,680	6,589	50	21	7	2	2	2	1	1	1	1	3

Non-Hispanic 1/, female

2003.....	4,937	4,893	29	4	7	1	2	1	-	-	-	-	-
2002.....	5,077	5,026	20	11	8	6	3	-	-	2	1	-	-
2001.....	4,989	4,943	20	11	8	1	3	-	1	1	-	1	-
2000.....	5,035	4,991	22	14	3	1	1	-	2	-	-	-	1
1999.....	5,135	5,079	23	20	5	4	1	-	2	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Certain conditions originating in the perinatal period (P00-P96)

Non-Hispanic white, both sexes

2003.....	6,359	6,303	31	8	12	2	2	1	-	-	-	-	-
2002.....	6,384	6,313	22	15	14	6	6	2	1	2	2	-	1
2001.....	6,323	6,239	34	18	11	2	8	4	1	3	1	1	1
2000.....	6,318	6,242	34	20	11	3	3	2	2	-	-	-	1
1999.....	6,384	6,286	37	31	11	6	3	2	3	2	1	1	1

Non-Hispanic white, male

2003.....	3,673	3,643	17	6	6	1	-	-	-	-	-	-	-
2002.....	3,663	3,628	10	7	7	3	3	2	1	-	1	-	1
2001.....	3,574	3,518	22	14	6	1	5	4	-	2	1	-	1
2000.....	3,605	3,553	23	14	9	2	2	2	-	-	-	-	-
1999.....	3,627	3,570	24	15	7	2	2	2	1	1	1	1	1

Non-Hispanic white, female

2003.....	2,686	2,660	14	2	6	1	2	1	-	-	-	-	-
2002.....	2,721	2,685	12	8	7	3	3	-	-	2	1	-	-
2001.....	2,749	2,721	12	4	5	1	3	-	1	1	-	1	-
2000.....	2,713	2,689	11	6	2	1	1	-	2	-	-	-	1
1999.....	2,757	2,716	13	16	4	4	1	-	2	1	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
All origins, both sexes													
2003.....	10,518	5,621	541	386	451	426	564	667	666	413	464	319	-
2002.....	10,687	5,623	530	417	492	475	572	709	634	440	489	305	1
2001.....	10,442	5,513	557	376	505	458	586	636	581	448	465	317	-
2000.....	10,578	5,743	495	399	441	477	569	612	569	464	488	320	1
1999.....	10,393	5,473	549	428	434	465	552	647	557	469	491	326	2
All origins, male													
2003.....	5,460	3,020	275	198	272	233	304	349	336	198	178	97	-
2002.....	5,515	2,942	282	215	282	284	296	381	321	213	201	97	1
2001.....	5,448	2,954	282	210	306	249	317	323	276	228	203	100	-
2000.....	5,517	3,028	259	207	268	279	316	347	264	222	197	130	-
1999.....	5,346	2,875	310	215	262	242	296	347	283	209	204	102	1
All origins, female													
2003.....	5,058	2,601	266	188	179	193	260	318	330	215	286	222	-
2002.....	5,172	2,681	248	202	210	191	276	328	313	227	288	208	-
2001.....	4,994	2,559	275	166	199	209	269	313	305	220	262	217	-
2000.....	5,061	2,715	236	192	173	198	253	265	305	242	291	190	1
1999.....	5,047	2,598	239	213	172	223	256	300	274	260	287	224	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Hispanic, both sexes													
2003.....	1,787	1,324	122	65	57	58	44	38	35	17	13	14	-
2002.....	1,715	1,257	111	70	67	53	40	42	25	18	23	9	-
2001.....	1,632	1,195	103	58	73	45	42	36	24	22	24	10	-
2000.....	1,612	1,182	111	52	54	46	43	36	31	21	24	12	-
1999.....	1,498	1,072	101	57	56	54	34	42	27	30	16	9	-
Hispanic, male													
2003.....	937	693	62	28	30	37	25	22	19	10	6	5	-
2002.....	903	653	60	35	39	36	19	23	15	6	13	4	-
2001.....	892	639	54	36	39	26	27	25	14	14	13	5	-
2000.....	857	605	60	28	39	26	27	22	17	11	15	7	-
1999.....	756	534	63	29	30	25	15	24	9	17	7	3	-
Hispanic, female													
2003.....	850	631	60	37	27	21	19	16	16	7	7	9	-
2002.....	812	604	51	35	28	17	21	19	10	12	10	5	-
2001.....	740	556	49	22	34	19	15	11	10	8	11	5	-
2000.....	755	577	51	24	15	20	16	14	14	10	9	5	-
1999.....	742	538	38	28	26	29	19	18	18	13	9	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Non-Hispanic 1/, both sexes													
2003.....	8,687	4,269	418	317	393	367	516	628	629	395	451	304	-
2002.....	8,904	4,316	415	347	422	422	530	662	608	421	465	295	1
2001.....	8,748	4,271	452	317	432	412	542	596	557	422	441	306	-
2000.....	8,900	4,514	384	345	386	427	525	574	532	443	462	307	1
1999.....	8,825	4,349	447	370	378	410	516	601	529	435	473	315	2
Non-Hispanic 1/, male													
2003.....	4,500	2,312	213	167	241	196	276	327	317	187	172	92	-
2002.....	4,577	2,262	221	180	241	248	276	355	306	206	188	93	1
2001.....	4,527	2,297	227	173	267	222	289	294	262	211	190	95	-
2000.....	4,623	2,400	199	177	229	250	288	323	243	211	181	122	-
1999.....	4,550	2,311	246	185	232	217	280	320	274	191	196	97	1
Non-Hispanic 1/, female													
2003.....	4,187	1,957	205	150	152	171	240	301	312	208	279	212	-
2002.....	4,327	2,054	194	167	181	174	254	307	302	215	277	202	-
2001.....	4,221	1,974	225	144	165	190	253	302	295	211	251	211	-
2000.....	4,277	2,114	185	168	157	177	237	251	289	232	281	185	1
1999.....	4,275	2,038	201	185	146	193	236	281	255	244	277	218	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Non-Hispanic white, both sexes													
2003.....	6,594	2,977	272	222	285	285	430	530	560	342	411	280	-
2002.....	6,830	3,068	281	253	315	331	423	562	527	374	420	275	1
2001.....	6,709	3,053	287	226	302	330	439	503	503	378	405	283	-
2000.....	6,792	3,199	240	240	279	344	419	501	460	390	431	288	1
1999.....	6,782	3,127	284	244	277	334	426	514	473	383	427	291	2
Non-Hispanic white, male													
2003.....	3,353	1,596	136	117	175	152	233	271	279	158	151	85	-
2002.....	3,510	1,643	151	125	178	195	223	295	263	183	165	88	1
2001.....	3,487	1,664	144	135	183	186	236	251	235	196	174	83	-
2000.....	3,519	1,720	124	119	163	203	233	276	217	177	172	115	-
1999.....	3,498	1,693	154	119	175	178	230	271	248	169	171	89	1
Non-Hispanic white, female													
2003.....	3,241	1,381	136	105	110	133	197	259	281	184	260	195	-
2002.....	3,320	1,425	130	128	137	136	200	267	264	191	255	187	-
2001.....	3,222	1,389	143	91	119	144	203	252	268	182	231	200	-
2000.....	3,273	1,479	116	121	116	141	186	225	243	213	259	173	1
1999.....	3,284	1,434	130	125	102	156	196	243	225	214	256	202	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Non-Hispanic black, both sexes													
2003.....	1,671	1,015	118	73	92	75	69	79	60	40	30	20	-
2002.....	1,716	1,015	113	81	98	75	92	83	69	40	32	18	-
2001.....	1,674	999	127	73	113	67	89	76	48	35	28	19	-
2000.....	1,739	1,062	123	89	91	74	91	67	58	43	25	16	-
1999.....	1,680	994	137	105	82	60	76	73	46	46	38	23	-
Non-Hispanic black, male													
2003.....	905	558	62	38	55	38	32	42	33	25	16	6	-
2002.....	887	506	58	47	59	43	45	47	40	21	16	5	-
2001.....	861	525	65	33	72	30	46	34	23	12	12	9	-
2000.....	911	545	67	50	58	43	49	41	17	26	9	6	-
1999.....	880	518	79	53	50	32	41	42	22	16	19	8	-
Non-Hispanic black, female													
2003.....	766	457	56	35	37	37	37	37	27	15	14	14	-
2002.....	829	509	55	34	39	32	47	36	29	19	16	13	-
2001.....	813	474	62	40	41	37	43	42	25	23	16	10	-
2000.....	828	517	56	39	33	31	42	26	41	17	16	10	-
1999.....	800	476	58	52	32	28	35	31	24	30	19	15	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Origin not stated 2/, both sexes													
2003.....	44	28	1	4	1	1	4	1	2	1	-	1	-
2002.....	68	50	4	-	3	-	2	5	1	1	1	1	-
2001.....	62	47	2	1	-	1	2	4	-	4	-	1	-
2000.....	66	47	-	2	1	4	1	2	6	-	2	1	-
1999.....	70	52	1	1	-	1	2	4	1	4	2	2	-
Origin not stated 2/, male													
2003.....	23	15	-	3	1	-	3	-	-	1	-	-	-
2002.....	35	27	1	-	2	-	1	3	-	1	-	-	-
2001.....	29	18	1	1	-	1	1	4	-	3	-	-	-
2000.....	37	23	-	2	-	3	1	2	4	-	1	1	-
1999.....	40	30	1	1	-	-	1	3	-	1	1	2	-
Origin not stated 2/, female													
2003.....	21	13	1	1	-	1	1	1	2	-	-	1	-
2002.....	33	23	3	-	1	-	1	2	1	-	1	1	-
2001.....	33	29	1	-	-	-	1	-	-	1	-	1	-
2000.....	29	24	-	-	1	1	-	-	2	-	1	-	-
1999.....	30	22	-	-	-	1	1	1	1	3	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins, both sexes

2003.....	31,444	3,318	256	178	837	1,327	2,382	2,874	2,067	2,522	5,006	10,565	112
2002.....	29,975	3,456	248	139	740	1,319	2,402	2,672	2,002	2,398	4,936	9,554	109
2001.....	32,326	3,491	272	181	949	1,530	3,126	3,504	2,292	2,774	5,003	9,089	115
2000.....	31,876	3,639	248	177	868	1,554	2,987	3,142	2,253	3,040	5,029	8,824	115
1999.....	26,672	3,538	236	137	609	1,095	2,062	2,151	1,860	2,698	4,413	7,798	75

All origins, male

2003.....	14,583	1,974	143	92	563	907	1,526	1,840	1,294	1,431	2,160	2,570	83
2002.....	14,269	2,032	145	82	501	905	1,540	1,723	1,273	1,324	2,228	2,439	77
2001.....	15,779	2,041	162	99	654	1,040	2,007	2,286	1,457	1,497	2,184	2,274	78
2000.....	15,863	2,156	137	99	581	1,060	1,994	2,123	1,429	1,708	2,244	2,242	90
1999.....	12,877	2,072	136	77	422	721	1,339	1,446	1,181	1,513	1,930	1,984	56

All origins, female

2003.....	16,861	1,344	113	86	274	420	856	1,034	773	1,091	2,846	7,995	29
2002.....	15,706	1,424	103	57	239	414	862	949	729	1,074	2,708	7,115	32
2001.....	16,547	1,450	110	82	295	490	1,119	1,218	835	1,277	2,819	6,815	37
2000.....	16,013	1,483	111	78	287	494	993	1,019	824	1,332	2,785	6,582	25
1999.....	13,795	1,466	100	60	187	374	723	705	679	1,185	2,483	5,814	19

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Hispanic, both sexes													
2003.....	1,623	492	30	20	98	170	181	141	73	75	118	214	11
2002.....	1,548	472	40	11	91	149	177	144	95	83	112	165	9
2001.....	2,190	496	47	36	184	286	327	283	127	112	107	173	12
2000.....	2,071	506	51	31	140	260	342	251	114	107	104	154	11
1999.....	1,281	415	30	16	83	138	148	93	63	73	93	124	5
Hispanic, male													
2003.....	998	309	16	12	65	135	135	108	55	44	49	62	8
2002.....	926	283	26	7	62	116	120	85	59	50	57	53	8
2001.....	1,410	280	28	15	139	221	239	212	92	71	56	46	11
2000.....	1,369	310	31	16	102	201	262	192	72	68	52	53	10
1999.....	768	237	14	10	64	99	110	68	42	38	46	36	4
Hispanic, female													
2003.....	625	183	14	8	33	35	46	33	18	31	69	152	3
2002.....	622	189	14	4	29	33	57	59	36	33	55	112	1
2001.....	780	216	19	21	45	65	88	71	35	41	51	127	1
2000.....	702	196	20	15	38	59	80	59	42	39	52	101	1
1999.....	513	178	16	6	19	39	38	25	21	35	47	88	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic 1/, both sexes

2003.....	29,617	2,805	226	157	731	1,151	2,181	2,708	1,975	2,439	4,876	10,324	44
2002.....	28,211	2,968	207	126	643	1,162	2,206	2,504	1,890	2,307	4,804	9,351	43
2001.....	29,931	2,978	225	144	757	1,227	2,782	3,193	2,154	2,653	4,886	8,896	36
2000.....	29,514	3,108	196	144	720	1,279	2,604	2,849	2,109	2,908	4,908	8,654	35
1999.....	25,181	3,097	204	121	521	951	1,885	2,023	1,780	2,601	4,310	7,658	30

Non-Hispanic 1/, male

2003.....	13,448	1,650	127	80	495	767	1,375	1,711	1,225	1,379	2,102	2,499	38
2002.....	13,206	1,739	119	74	434	783	1,408	1,619	1,198	1,267	2,156	2,376	33
2001.....	14,246	1,750	134	84	510	807	1,754	2,058	1,355	1,422	2,125	2,221	26
2000.....	14,288	1,834	106	82	474	848	1,700	1,901	1,331	1,621	2,178	2,183	30
1999.....	11,962	1,817	121	67	355	619	1,209	1,347	1,125	1,463	1,876	1,944	19

Non-Hispanic 1/, female

2003.....	16,169	1,155	99	77	236	384	806	997	750	1,060	2,774	7,825	6
2002.....	15,005	1,229	88	52	209	379	798	885	692	1,040	2,648	6,975	10
2001.....	15,685	1,228	91	60	247	420	1,028	1,135	799	1,231	2,761	6,675	10
2000.....	15,226	1,274	90	62	246	431	904	948	778	1,287	2,730	6,471	5
1999.....	13,219	1,280	83	54	166	332	676	676	655	1,138	2,434	5,714	11

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)													
Non-Hispanic white, both sexes													
2003.....	24,452	1,665	136	99	562	843	1,661	2,022	1,562	2,013	4,367	9,487	35
2002.....	23,099	1,777	124	71	468	849	1,664	1,822	1,493	1,934	4,275	8,591	31
2001.....	24,596	1,752	139	89	563	907	2,175	2,510	1,713	2,191	4,350	8,178	29
2000.....	24,064	1,805	134	88	518	918	2,027	2,173	1,652	2,394	4,358	7,975	22
1999.....	20,200	1,812	129	81	348	660	1,341	1,459	1,374	2,166	3,775	7,033	22
Non-Hispanic white, male													
2003.....	10,752	1,025	75	51	389	570	1,050	1,297	958	1,143	1,881	2,282	31
2002.....	10,432	1,056	66	44	326	581	1,071	1,177	945	1,065	1,897	2,180	24
2001.....	11,404	1,046	79	52	376	614	1,387	1,634	1,058	1,184	1,920	2,032	22
2000.....	11,351	1,067	79	52	356	625	1,366	1,465	1,036	1,328	1,945	2,013	19
1999.....	9,249	1,083	82	45	244	431	884	963	870	1,222	1,638	1,771	16
Non-Hispanic white, female													
2003.....	13,700	640	61	48	173	273	611	725	604	870	2,486	7,205	4
2002.....	12,667	721	58	27	142	268	593	645	548	869	2,378	6,411	7
2001.....	13,192	706	60	37	187	293	788	876	655	1,007	2,430	6,146	7
2000.....	12,713	738	55	36	162	293	661	708	616	1,066	2,413	5,962	3
1999.....	10,951	729	47	36	104	229	457	496	504	944	2,137	5,262	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic black, both sexes

2003.....	4,628	1,022	80	49	150	267	476	632	385	400	439	721	7
2002.....	4,556	1,070	66	50	152	275	490	626	356	335	471	655	10
2001.....	4,606	1,081	69	42	158	266	518	605	373	401	466	622	5
2000.....	4,892	1,186	54	46	179	308	514	609	411	472	505	595	13
1999.....	4,508	1,149	67	33	152	260	504	520	371	395	488	563	6

Non-Hispanic black, male

2003.....	2,404	554	44	25	92	173	290	383	247	223	183	185	5
2002.....	2,482	615	44	28	92	174	305	407	229	180	234	167	7
2001.....	2,414	614	44	26	109	150	300	364	262	207	174	161	3
2000.....	2,614	694	26	28	105	191	283	390	266	267	214	139	11
1999.....	2,460	658	35	20	100	172	299	352	238	218	214	152	2

Non-Hispanic black, female

2003.....	2,224	468	36	24	58	94	186	249	138	177	256	536	2
2002.....	2,074	455	22	22	60	101	185	219	127	155	237	488	3
2001.....	2,192	467	25	16	49	116	218	241	111	194	292	461	2
2000.....	2,278	492	28	18	74	117	231	219	145	205	291	456	2
1999.....	2,048	491	32	13	52	88	205	168	133	177	274	411	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Origin not stated 2/, both sexes													
2003.....	204	21	-	1	8	6	20	25	19	8	12	27	57
2002.....	216	16	1	2	6	8	19	24	17	8	20	38	57
2001.....	205	17	-	1	8	17	17	28	11	9	10	20	67
2000.....	291	25	1	2	8	15	41	42	30	25	17	16	69
1999.....	210	26	2	-	5	6	29	35	17	24	10	16	40
Origin not stated 2/, male													
2003.....	137	15	-	-	3	5	16	21	14	8	9	9	37
2002.....	137	10	-	1	5	6	12	19	16	7	15	10	36
2001.....	123	11	-	-	5	12	14	16	10	4	3	7	41
2000.....	206	12	-	1	5	11	32	30	26	19	14	6	50
1999.....	147	18	1	-	3	3	20	31	14	12	8	4	33
Origin not stated 2/, female													
2003.....	67	6	-	1	5	1	4	4	5	-	3	18	20
2002.....	79	6	1	1	1	2	7	5	1	1	5	28	21
2001.....	82	6	-	1	3	5	3	12	1	5	7	13	26
2000.....	85	13	1	1	3	4	9	12	4	6	3	10	19
1999.....	63	8	1	-	2	3	9	4	3	12	2	12	7

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All origins, both sexes

2003.....	201,676	1,190	635	927	2,023	3,330	8,407	15,883	18,734	26,475	52,955	71,100	17
2002.....	194,591	1,144	661	868	1,853	3,136	8,506	15,151	17,378	25,689	51,618	68,574	13
2001.....	186,406	1,131	646	860	1,799	3,072	8,084	14,203	16,048	25,803	49,711	65,032	17
2000.....	178,378	1,105	609	879	1,764	3,097	8,113	13,326	15,156	25,007	48,019	61,292	11
1999.....	171,683	1,148	647	921	1,771	3,062	7,989	12,383	14,390	25,165	45,598	58,599	10

All origins, male

2003.....	81,907	711	342	541	1,189	1,885	4,932	9,456	10,274	12,993	21,393	18,178	13
2002.....	79,007	690	363	486	1,072	1,837	5,019	9,115	9,528	12,667	20,667	17,552	11
2001.....	75,162	649	358	454	1,064	1,729	4,774	8,397	8,652	12,732	19,817	16,520	16
2000.....	72,227	629	338	474	1,057	1,811	4,801	7,911	8,236	12,321	19,102	15,536	11
1999.....	69,784	680	354	483	1,045	1,790	4,835	7,272	7,870	12,332	18,495	14,623	5

All origins, female

2003.....	119,769	479	293	386	834	1,445	3,475	6,427	8,460	13,482	31,562	52,922	4
2002.....	115,584	454	298	382	781	1,299	3,487	6,036	7,850	13,022	30,951	51,022	2
2001.....	111,244	482	288	406	735	1,343	3,310	5,806	7,396	13,071	29,894	48,512	1
2000.....	106,151	476	271	405	707	1,286	3,312	5,415	6,920	12,686	28,917	45,756	-
1999.....	101,899	468	293	438	726	1,272	3,154	5,111	6,520	12,833	27,103	43,976	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Hispanic, both sexes

2003.....	9,218	235	140	159	287	452	829	1,156	1,117	1,306	1,822	1,712	3
2002.....	8,377	209	120	151	250	409	757	1,006	958	1,313	1,632	1,571	1
2001.....	7,847	187	112	122	246	354	671	892	918	1,218	1,539	1,585	3
2000.....	7,325	196	109	113	214	381	706	862	813	1,113	1,438	1,376	4
1999.....	7,291	200	106	107	218	409	690	837	842	1,133	1,265	1,482	2

Hispanic, male

2003.....	4,571	144	75	87	165	295	519	734	640	631	787	492	2
2002.....	4,221	116	66	89	148	261	517	647	527	659	707	483	1
2001.....	3,849	106	66	65	147	222	451	584	513	587	612	493	3
2000.....	3,663	100	54	63	140	246	483	552	464	568	557	432	4
1999.....	3,655	124	50	54	128	275	479	541	481	564	530	428	1

Hispanic, female

2003.....	4,647	91	65	72	122	157	310	422	477	675	1,035	1,220	1
2002.....	4,156	93	54	62	102	148	240	359	431	654	925	1,088	-
2001.....	3,998	81	46	57	99	132	220	308	405	631	927	1,092	-
2000.....	3,662	96	55	50	74	135	223	310	349	545	881	944	-
1999.....	3,636	76	56	53	90	134	211	296	361	569	735	1,054	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic 1/, both sexes

2003.....	191,961	946	493	765	1,727	2,865	7,544	14,662	17,558	25,109	51,044	69,241	7
2002.....	185,608	927	538	715	1,600	2,714	7,711	14,060	16,347	24,307	49,835	66,846	8
2001.....	177,985	934	532	736	1,545	2,709	7,370	13,230	15,063	24,506	48,059	63,290	11
2000.....	170,462	898	497	762	1,541	2,697	7,366	12,376	14,283	23,799	46,458	59,780	5
1999.....	163,832	935	537	814	1,547	2,645	7,247	11,467	13,479	23,956	44,232	56,968	5

Non-Hispanic 1/, male

2003.....	77,081	561	266	452	1,016	1,580	4,393	8,680	9,595	12,326	20,567	17,640	5
2002.....	74,500	572	294	396	923	1,568	4,475	8,410	8,953	11,964	19,905	17,033	7
2001.....	71,014	539	291	388	912	1,499	4,293	7,751	8,087	12,104	19,155	15,985	10
2000.....	68,263	522	282	411	911	1,556	4,287	7,294	7,734	11,700	18,493	15,068	5
1999.....	65,835	546	302	429	913	1,509	4,316	6,676	7,341	11,723	17,924	14,153	3

Non-Hispanic 1/, female

2003.....	114,880	385	227	313	711	1,285	3,151	5,982	7,963	12,783	30,477	51,601	2
2002.....	111,108	355	244	319	677	1,146	3,236	5,650	7,394	12,343	29,930	49,813	1
2001.....	106,971	395	241	348	633	1,210	3,077	5,479	6,976	12,402	28,904	47,305	1
2000.....	102,199	376	215	351	630	1,141	3,079	5,082	6,549	12,099	27,965	44,712	-
1999.....	97,997	389	235	385	634	1,136	2,931	4,791	6,138	12,233	26,308	42,815	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic white, both sexes

2003.....	165,261	565	324	531	1,245	1,910	5,404	10,965	14,071	21,224	45,510	63,507	5
2002.....	159,945	578	352	505	1,102	1,831	5,545	10,567	13,116	20,530	44,489	61,325	5
2001.....	153,156	556	362	541	1,089	1,808	5,201	9,814	11,982	20,807	42,990	58,000	6
2000.....	146,518	544	316	537	1,113	1,807	5,202	9,186	11,332	20,188	41,562	54,729	2
1999.....	141,002	554	375	569	1,100	1,845	5,036	8,437	10,749	20,267	39,663	52,406	1

Non-Hispanic white, male

2003.....	65,306	357	168	319	788	1,119	3,280	6,649	7,742	10,416	18,274	16,191	3
2002.....	63,109	376	191	279	677	1,113	3,386	6,464	7,230	10,109	17,748	15,531	5
2001.....	60,022	311	209	286	677	1,040	3,130	5,873	6,476	10,312	17,140	14,563	5
2000.....	57,565	310	183	307	683	1,105	3,128	5,569	6,149	9,933	16,519	13,677	2
1999.....	55,517	321	214	291	681	1,110	3,144	5,029	5,906	9,977	16,013	12,830	1

Non-Hispanic white, female

2003.....	99,955	208	156	212	457	791	2,124	4,316	6,329	10,808	27,236	47,316	2
2002.....	96,836	202	161	226	425	718	2,159	4,103	5,886	10,421	26,741	45,794	-
2001.....	93,134	245	153	255	412	768	2,071	3,941	5,506	10,495	25,850	43,437	1
2000.....	88,953	234	133	230	430	702	2,074	3,617	5,183	10,255	25,043	41,052	-
1999.....	85,485	233	161	278	419	735	1,892	3,408	4,843	10,290	23,650	39,576	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic black, both sexes

2003.....	22,997	333	133	182	401	813	1,880	3,308	3,048	3,335	4,682	4,880	2
2002.....	22,297	298	146	172	416	775	1,935	3,147	2,827	3,247	4,559	4,772	3
2001.....	21,752	329	131	165	386	758	1,944	3,106	2,691	3,236	4,415	4,586	5
2000.....	21,081	296	144	189	373	810	1,948	2,897	2,607	3,153	4,248	4,413	3
1999.....	20,137	332	129	201	385	704	1,997	2,728	2,437	3,251	3,963	4,006	4

Non-Hispanic black, male

2003.....	10,019	183	78	99	182	386	952	1,802	1,610	1,643	1,915	1,167	2
2002.....	9,722	167	80	95	205	388	948	1,733	1,501	1,591	1,795	1,217	2
2001.....	9,561	192	62	83	196	381	1,030	1,713	1,411	1,570	1,767	1,151	5
2000.....	9,351	180	77	87	197	400	1,033	1,562	1,411	1,548	1,711	1,142	3
1999.....	8,967	192	67	118	202	344	1,040	1,487	1,270	1,518	1,630	1,097	2

Non-Hispanic black, female

2003.....	12,978	150	55	83	219	427	928	1,506	1,438	1,692	2,767	3,713	-
2002.....	12,575	131	66	77	211	387	987	1,414	1,326	1,656	2,764	3,555	1
2001.....	12,191	137	69	82	190	377	914	1,393	1,280	1,666	2,648	3,435	-
2000.....	11,730	116	67	102	176	410	915	1,335	1,196	1,605	2,537	3,271	-
1999.....	11,170	140	62	83	183	360	957	1,241	1,167	1,733	2,333	2,909	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Origin not stated 2/, both sexes

2003.....	497	9	2	3	9	13	34	65	59	60	89	147	7
2002.....	606	8	3	2	3	13	38	85	73	69	151	157	4
2001.....	574	10	2	2	8	9	43	81	67	79	113	157	3
2000.....	591	11	3	4	9	19	41	88	60	95	123	136	2
1999.....	560	13	4	-	6	8	52	79	69	76	101	149	3

Origin not stated 2/, male

2003.....	255	6	1	2	8	10	20	42	39	36	39	46	6
2002.....	286	2	3	1	1	8	27	58	48	44	55	36	3
2001.....	299	4	1	1	5	8	30	62	52	41	50	42	3
2000.....	301	7	2	-	6	9	31	65	38	53	52	36	2
1999.....	294	10	2	-	4	6	40	55	48	45	41	42	1

Origin not stated 2/, female

2003.....	242	3	1	1	1	3	14	23	20	24	50	101	1
2002.....	320	6	-	1	2	5	11	27	25	25	96	121	1
2001.....	275	6	1	1	3	1	13	19	15	38	63	115	-
2000.....	290	4	1	4	3	10	10	23	22	42	71	100	-
1999.....	266	3	2	-	2	2	12	24	21	31	60	107	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
All origins, both sexes													
2003.....	109,277	945	1,717	2,618	15,272	12,541	16,766	15,837	9,170	8,081	13,108	13,146	76
2002.....	106,742	946	1,641	2,718	15,412	12,569	16,710	14,675	8,345	8,086	12,904	12,651	85
2001.....	101,537	976	1,714	2,836	14,411	11,839	15,945	13,344	7,658	7,835	12,688	12,171	120
2000.....	97,900	881	1,826	2,979	14,113	11,769	15,413	12,278	7,505	7,698	11,758	11,595	85
1999.....	97,860	845	1,898	3,091	13,656	11,890	15,231	11,639	7,285	8,208	12,282	11,729	106
All origins, male													
2003.....	70,532	539	1,048	1,662	11,291	9,534	12,003	11,196	6,324	4,921	6,851	5,093	70
2002.....	69,257	553	1,024	1,713	11,438	9,635	12,012	10,492	5,781	5,031	6,687	4,817	74
2001.....	66,060	541	1,045	1,803	10,832	9,092	11,612	9,743	5,228	4,751	6,652	4,657	104
2000.....	63,817	501	1,098	1,861	10,460	9,059	11,444	9,063	5,133	4,639	6,063	4,413	83
1999.....	63,535	494	1,127	1,974	10,040	9,097	11,248	8,602	5,015	4,992	6,372	4,480	94
All origins, female													
2003.....	38,745	406	669	956	3,981	3,007	4,763	4,641	2,846	3,160	6,257	8,053	6
2002.....	37,485	393	617	1,005	3,974	2,934	4,698	4,183	2,564	3,055	6,217	7,834	11
2001.....	35,477	435	669	1,033	3,579	2,747	4,333	3,601	2,430	3,084	6,036	7,514	16
2000.....	34,083	380	728	1,118	3,653	2,710	3,969	3,215	2,372	3,059	5,695	7,182	2
1999.....	34,325	351	771	1,117	3,616	2,793	3,983	3,037	2,270	3,216	5,910	7,249	12

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Hispanic, both sexes													
2003.....	10,418	131	325	431	2,366	2,167	1,877	1,254	639	466	454	293	15
2002.....	10,106	127	341	389	2,413	2,027	1,841	1,168	566	466	464	276	28
2001.....	9,523	154	310	379	2,182	1,938	1,760	1,078	564	454	409	266	29
2000.....	8,830	118	306	383	2,095	1,776	1,618	953	492	418	365	278	28
1999.....	8,650	134	341	362	1,946	1,790	1,607	926	466	440	334	260	44
Hispanic, male													
2003.....	7,899	77	200	289	1,905	1,827	1,488	966	449	289	267	127	15
2002.....	7,698	83	215	244	1,958	1,709	1,468	888	407	307	258	134	27
2001.....	7,157	75	195	224	1,773	1,619	1,418	822	391	275	235	104	26
2000.....	6,696	67	174	210	1,717	1,499	1,317	750	347	271	191	125	28
1999.....	6,578	80	197	227	1,575	1,494	1,307	734	334	288	182	120	40
Hispanic, female													
2003.....	2,519	54	125	142	461	340	389	288	190	177	187	166	-
2002.....	2,408	44	126	145	455	318	373	280	159	159	206	142	1
2001.....	2,366	79	115	155	409	319	342	256	173	179	174	162	3
2000.....	2,134	51	132	173	378	277	301	203	145	147	174	153	-
1999.....	2,072	54	144	135	371	296	300	192	132	152	152	140	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Non-Hispanic 1/, both sexes													
2003.....	98,406	812	1,386	2,183	12,866	10,326	14,813	14,488	8,485	7,580	12,620	12,824	23
2002.....	96,175	816	1,296	2,318	12,956	10,489	14,792	13,422	7,728	7,584	12,394	12,347	33
2001.....	91,522	817	1,399	2,445	12,187	9,857	14,089	12,184	7,042	7,347	12,245	11,874	36
2000.....	88,558	759	1,512	2,585	11,968	9,935	13,706	11,219	6,966	7,247	11,361	11,288	12
1999.....	88,669	710	1,550	2,721	11,658	10,034	13,529	10,622	6,753	7,738	11,896	11,432	26
Non-Hispanic 1/, male													
2003.....	62,290	460	844	1,369	9,355	7,670	10,454	10,158	5,839	4,604	6,564	4,953	20
2002.....	61,201	468	806	1,461	9,448	7,882	10,482	9,537	5,325	4,693	6,398	4,671	30
2001.....	58,528	462	845	1,570	9,024	7,438	10,117	8,852	4,795	4,455	6,397	4,544	29
2000.....	56,736	431	921	1,648	8,707	7,512	10,054	8,231	4,747	4,344	5,855	4,275	11
1999.....	56,551	414	926	1,739	8,428	7,550	9,871	7,791	4,627	4,681	6,158	4,343	23
Non-Hispanic 1/, female													
2003.....	36,116	352	542	814	3,511	2,656	4,359	4,330	2,646	2,976	6,056	7,871	3
2002.....	34,974	348	490	857	3,508	2,607	4,310	3,885	2,403	2,891	5,996	7,676	3
2001.....	32,994	355	554	875	3,163	2,419	3,972	3,332	2,247	2,892	5,848	7,330	7
2000.....	31,822	328	591	937	3,261	2,423	3,652	2,988	2,219	2,903	5,506	7,013	1
1999.....	32,118	296	624	982	3,230	2,484	3,658	2,831	2,126	3,057	5,738	7,089	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Non-Hispanic white, both sexes													
2003.....	82,815	496	985	1,535	10,638	8,238	12,106	11,721	7,043	6,466	11,487	12,078	22
2002.....	80,605	462	865	1,581	10,664	8,344	12,035	10,859	6,393	6,481	11,316	11,579	26
2001.....	76,262	500	1,006	1,688	9,867	7,798	11,340	9,704	5,776	6,309	11,157	11,092	25
2000.....	73,536	455	1,033	1,843	9,735	7,730	11,008	8,882	5,753	6,181	10,356	10,552	8
1999.....	73,326	435	1,042	1,908	9,410	7,855	10,844	8,392	5,455	6,568	10,757	10,647	13
Non-Hispanic white, male													
2003.....	51,888	283	610	954	7,698	6,138	8,574	8,219	4,841	3,952	5,957	4,643	19
2002.....	50,652	279	527	997	7,746	6,273	8,550	7,745	4,374	3,991	5,805	4,342	23
2001.....	48,180	281	624	1,066	7,297	5,906	8,188	7,034	3,905	3,837	5,800	4,220	22
2000.....	46,470	266	638	1,176	7,065	5,861	8,113	6,521	3,871	3,692	5,292	3,968	7
1999.....	46,113	258	617	1,205	6,749	5,932	7,949	6,142	3,730	3,962	5,551	4,006	12
Non-Hispanic white, female													
2003.....	30,927	213	375	581	2,940	2,100	3,532	3,502	2,202	2,514	5,530	7,435	3
2002.....	29,953	183	338	584	2,918	2,071	3,485	3,114	2,019	2,490	5,511	7,237	3
2001.....	28,082	219	382	622	2,570	1,892	3,152	2,670	1,871	2,472	5,357	6,872	3
2000.....	27,066	189	395	667	2,670	1,869	2,895	2,361	1,882	2,489	5,064	6,584	1
1999.....	27,213	177	425	703	2,661	1,923	2,895	2,250	1,725	2,606	5,206	6,641	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Non-Hispanic black, both sexes													
2003.....	12,154	283	333	517	1,590	1,602	2,195	2,289	1,152	850	798	544	1
2002.....	12,285	324	350	607	1,680	1,676	2,227	2,145	1,075	818	799	580	4
2001.....	12,239	272	331	630	1,713	1,600	2,282	2,103	1,045	836	830	589	8
2000.....	12,087	268	396	619	1,662	1,740	2,253	1,961	983	857	768	576	4
1999.....	12,533	251	428	701	1,694	1,738	2,245	1,903	1,047	943	925	646	12
Non-Hispanic black, male													
2003.....	8,237	158	197	330	1,205	1,197	1,539	1,625	824	518	429	214	1
2002.....	8,428	172	231	391	1,260	1,279	1,562	1,523	791	527	448	240	4
2001.....	8,381	158	188	432	1,285	1,192	1,608	1,574	750	505	461	222	6
2000.....	8,380	145	232	406	1,233	1,316	1,649	1,454	736	537	432	236	4
1999.....	8,629	140	260	467	1,281	1,313	1,628	1,441	742	596	483	268	10
Non-Hispanic black, female													
2003.....	3,917	125	136	187	385	405	656	664	328	332	369	330	-
2002.....	3,857	152	119	216	420	397	665	622	284	291	351	340	-
2001.....	3,858	114	143	198	428	408	674	529	295	331	369	367	2
2000.....	3,707	123	164	213	429	424	604	507	247	320	336	340	-
1999.....	3,904	111	168	234	413	425	617	462	305	347	442	378	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Origin not stated 2/, both sexes													
2003.....	453	2	6	4	40	48	76	95	46	35	34	29	38
2002.....	461	3	4	11	43	53	77	85	51	36	46	28	24
2001.....	492	5	5	12	42	44	96	82	52	34	34	31	55
2000.....	512	4	8	11	50	58	89	106	47	33	32	29	45
1999.....	541	1	7	8	52	66	95	91	66	30	52	37	36
Origin not stated 2/, male													
2003.....	343	2	4	4	31	37	61	72	36	28	20	13	35
2002.....	358	2	3	8	32	44	62	67	49	31	31	12	17
2001.....	375	4	5	9	35	35	77	69	42	21	20	9	49
2000.....	385	3	3	3	36	48	73	82	39	24	17	13	44
1999.....	406	-	4	8	37	53	70	77	54	23	32	17	31
Origin not stated 2/, female													
2003.....	110	-	2	-	9	11	15	23	10	7	14	16	3
2002.....	103	1	1	3	11	9	15	18	2	5	15	16	7
2001.....	117	1	-	3	7	9	19	13	10	13	14	22	6
2000.....	127	1	5	8	14	10	16	24	8	9	15	16	1
1999.....	135	1	3	-	15	13	25	14	12	7	20	20	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
All origins, both sexes													
2003.....	48,071	148	649	1,765	11,432	7,280	7,594	6,754	4,441	3,210	3,377	1,401	20
2002.....	48,366	127	640	1,738	11,816	7,484	7,688	6,563	4,165	3,351	3,433	1,322	39
2001.....	47,288	151	678	1,823	11,139	7,454	7,827	6,251	3,861	3,282	3,507	1,272	43
2000.....	46,749	174	691	1,922	11,028	7,400	7,661	5,965	3,883	3,315	3,342	1,335	33
1999.....	46,423	189	712	1,975	10,689	7,434	7,553	5,645	3,812	3,579	3,468	1,330	37
All origins, male													
2003.....	33,291	81	366	1,084	8,171	5,511	5,497	4,865	3,087	1,975	1,885	751	18
2002.....	33,533	72	381	1,039	8,471	5,676	5,499	4,727	2,877	2,101	1,917	737	36
2001.....	32,794	68	392	1,115	8,111	5,661	5,613	4,484	2,631	2,015	1,983	684	37
2000.....	32,166	94	391	1,138	7,890	5,570	5,560	4,260	2,617	2,021	1,858	734	33
1999.....	31,642	101	400	1,206	7,558	5,511	5,376	4,016	2,561	2,162	1,959	758	34
All origins, female													
2003.....	14,780	67	283	681	3,261	1,769	2,097	1,889	1,354	1,235	1,492	650	2
2002.....	14,833	55	259	699	3,345	1,808	2,189	1,836	1,288	1,250	1,516	585	3
2001.....	14,494	83	286	708	3,028	1,793	2,214	1,767	1,230	1,267	1,524	588	6
2000.....	14,583	80	300	784	3,138	1,830	2,101	1,705	1,266	1,294	1,484	601	-
1999.....	14,781	88	312	769	3,131	1,923	2,177	1,629	1,251	1,417	1,509	572	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
Hispanic, both sexes													
2003.....	6,045	39	139	308	1,834	1,412	973	577	329	227	151	48	8
2002.....	5,921	26	182	277	1,875	1,340	930	554	300	217	157	46	17
2001.....	5,774	43	159	265	1,740	1,293	942	557	331	229	167	38	10
2000.....	5,217	42	154	279	1,616	1,126	831	508	268	220	121	38	14
1999.....	4,880	40	157	250	1,457	1,125	758	445	253	228	120	36	11
Hispanic, male													
2003.....	4,574	25	77	209	1,437	1,175	740	427	212	142	92	30	8
2002.....	4,481	16	113	173	1,479	1,094	722	394	207	149	91	27	16
2001.....	4,334	16	92	155	1,383	1,067	731	390	214	141	114	23	8
2000.....	3,921	24	85	155	1,285	921	639	373	189	140	74	22	14
1999.....	3,630	22	85	151	1,136	904	590	330	161	144	77	19	11
Hispanic, female													
2003.....	1,471	14	62	99	397	237	233	150	117	85	59	18	-
2002.....	1,440	10	69	104	396	246	208	160	93	68	66	19	1
2001.....	1,440	27	67	110	357	226	211	167	117	88	53	15	2
2000.....	1,296	18	69	124	331	205	192	135	79	80	47	16	-
1999.....	1,250	18	72	99	321	221	168	115	92	84	43	17	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
Non-Hispanic 1/, both sexes													
2003.....	41,856	109	505	1,454	9,570	5,846	6,597	6,150	4,090	2,966	3,217	1,348	4
2002.....	42,268	101	457	1,456	9,912	6,120	6,730	5,975	3,844	3,120	3,264	1,274	15
2001.....	41,339	108	517	1,554	9,371	6,141	6,848	5,668	3,513	3,044	3,330	1,230	15
2000.....	41,341	132	533	1,637	9,375	6,244	6,801	5,430	3,595	3,082	3,214	1,295	3
1999.....	41,330	149	553	1,719	9,192	6,272	6,766	5,175	3,535	3,338	3,329	1,291	11
Non-Hispanic 1/, male													
2003.....	28,591	56	286	872	6,713	4,320	4,737	4,418	2,858	1,820	1,789	718	4
2002.....	28,906	56	267	862	6,970	4,562	4,755	4,305	2,651	1,940	1,816	708	14
2001.....	28,327	52	298	957	6,706	4,578	4,855	4,071	2,405	1,869	1,865	659	12
2000.....	28,096	70	304	982	6,580	4,624	4,896	3,865	2,411	1,871	1,779	711	3
1999.....	27,850	79	313	1,049	6,396	4,578	4,764	3,666	2,379	2,009	1,869	738	10
Non-Hispanic 1/, female													
2003.....	13,265	53	219	582	2,857	1,526	1,860	1,732	1,232	1,146	1,428	630	-
2002.....	13,362	45	190	594	2,942	1,558	1,975	1,670	1,193	1,180	1,448	566	1
2001.....	13,012	56	219	597	2,665	1,563	1,993	1,597	1,108	1,175	1,465	571	3
2000.....	13,245	62	229	655	2,795	1,620	1,905	1,565	1,184	1,211	1,435	584	-
1999.....	13,480	70	240	670	2,796	1,694	2,002	1,509	1,156	1,329	1,460	553	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
Non-Hispanic white, both sexes													
2003.....	34,220	73	352	1,095	7,826	4,480	5,276	5,019	3,414	2,543	2,892	1,246	4
2002.....	34,689	59	300	1,076	8,146	4,703	5,389	4,909	3,246	2,695	2,974	1,182	10
2001.....	33,753	74	353	1,122	7,563	4,771	5,494	4,611	2,937	2,628	3,044	1,148	8
2000.....	33,808	79	357	1,246	7,669	4,759	5,471	4,402	3,039	2,649	2,937	1,197	3
1999.....	33,803	98	349	1,291	7,493	4,889	5,425	4,211	2,946	2,897	3,020	1,180	4
Non-Hispanic white, male													
2003.....	23,332	39	208	648	5,439	3,319	3,796	3,620	2,411	1,584	1,606	658	4
2002.....	23,628	35	168	643	5,707	3,501	3,816	3,570	2,240	1,661	1,630	648	9
2001.....	23,047	36	208	686	5,401	3,561	3,887	3,326	2,009	1,617	1,698	611	7
2000.....	22,855	44	205	748	5,361	3,518	3,935	3,160	2,014	1,597	1,609	661	3
1999.....	22,682	58	192	782	5,181	3,561	3,827	2,992	1,986	1,745	1,686	668	4
Non-Hispanic white, female													
2003.....	10,888	34	144	447	2,387	1,161	1,480	1,399	1,003	959	1,286	588	-
2002.....	11,061	24	132	433	2,439	1,202	1,573	1,339	1,006	1,034	1,344	534	1
2001.....	10,706	38	145	436	2,162	1,210	1,607	1,285	928	1,011	1,346	537	1
2000.....	10,953	35	152	498	2,308	1,241	1,536	1,242	1,025	1,052	1,328	536	-
1999.....	11,121	40	157	509	2,312	1,328	1,598	1,219	960	1,152	1,334	512	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
Non-Hispanic black, both sexes													
2003.....	5,684	28	125	276	1,221	1,039	1,040	862	501	303	217	72	-
2002.....	5,665	36	121	302	1,275	1,089	1,033	810	444	284	206	62	3
2001.....	5,803	18	127	354	1,322	1,058	1,071	829	455	307	198	57	7
2000.....	5,757	40	139	309	1,262	1,152	1,059	809	417	322	179	69	-
1999.....	5,810	44	168	360	1,262	1,075	1,074	758	430	314	232	86	7
Non-Hispanic black, male													
2003.....	4,053	12	63	173	917	786	755	637	354	183	132	41	-
2002.....	4,051	18	79	177	917	837	733	582	327	196	139	43	3
2001.....	4,143	7	69	234	966	797	779	615	332	185	123	31	5
2000.....	4,127	21	77	191	907	875	796	571	314	218	118	39	-
1999.....	4,097	18	96	229	909	817	769	557	303	196	139	58	6
Non-Hispanic black, female													
2003.....	1,631	16	62	103	304	253	285	225	147	120	85	31	-
2002.....	1,614	18	42	125	358	252	300	228	117	88	67	19	-
2001.....	1,660	11	58	120	356	261	292	214	123	122	75	26	2
2000.....	1,630	19	62	118	355	277	263	238	103	104	61	30	-
1999.....	1,713	26	72	131	353	258	305	201	127	118	93	28	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Transport accidents (V01-V99,Y85)													
Origin not stated 2/, both sexes													
2003.....	170	-	5	3	28	22	24	27	22	17	9	5	8
2002.....	177	-	1	5	29	24	28	34	21	14	12	2	7
2001.....	175	-	2	4	28	20	37	26	17	9	10	4	18
2000.....	191	-	4	6	37	30	29	27	20	13	7	2	16
1999.....	213	-	2	6	40	37	29	25	24	13	19	3	15
Origin not stated 2/, male													
2003.....	126	-	3	3	21	16	20	20	17	13	4	3	6
2002.....	146	-	1	4	22	20	22	28	19	12	10	2	6
2001.....	133	-	2	3	22	16	27	23	12	5	4	2	17
2000.....	149	-	2	1	25	25	25	22	17	10	5	1	16
1999.....	162	-	2	6	26	29	22	20	21	9	13	1	13
Origin not stated 2/, female													
2003.....	44	-	2	-	7	6	4	7	5	4	5	2	2
2002.....	31	-	-	1	7	4	6	6	2	2	2	-	1
2001.....	42	-	-	1	6	4	10	3	5	4	6	2	1
2000.....	42	-	2	5	12	5	4	5	3	3	2	1	-
1999.....	51	-	-	-	14	8	7	5	3	4	6	2	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)													
All origins, both sexes													
2003.....	44,757	145	621	1,642	10,972	6,833	6,961	6,074	3,958	2,967	3,210	1,358	16
2002.....	45,380	123	610	1,614	11,459	7,092	7,077	5,937	3,764	3,113	3,271	1,285	35
2001.....	43,788	144	626	1,686	10,725	6,937	7,083	5,564	3,465	2,990	3,312	1,223	33
2000.....	43,354	168	651	1,772	10,560	6,884	6,927	5,361	3,506	3,038	3,173	1,288	26
1999.....	42,401	184	650	1,771	10,128	6,778	6,738	4,972	3,370	3,276	3,243	1,261	30
All origins, male													
2003.....	30,655	80	353	1,009	7,806	5,156	4,984	4,303	2,683	1,778	1,767	722	14
2002.....	31,064	68	361	963	8,178	5,349	4,984	4,189	2,540	1,888	1,793	718	33
2001.....	30,064	66	356	1,029	7,778	5,236	5,016	3,938	2,309	1,786	1,859	664	27
2000.....	29,451	90	368	1,043	7,516	5,135	4,940	3,780	2,306	1,802	1,739	706	26
1999.....	28,552	97	366	1,062	7,127	4,993	4,735	3,473	2,200	1,931	1,827	714	27
All origins, female													
2003.....	14,102	65	268	633	3,166	1,677	1,977	1,771	1,275	1,189	1,443	636	2
2002.....	14,316	55	249	651	3,281	1,743	2,093	1,748	1,224	1,225	1,478	567	2
2001.....	13,724	78	270	657	2,947	1,701	2,067	1,626	1,156	1,204	1,453	559	6
2000.....	13,903	78	283	729	3,044	1,749	1,987	1,581	1,200	1,236	1,434	582	-
1999.....	13,849	87	284	709	3,001	1,785	2,003	1,499	1,170	1,345	1,416	547	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)													
Hispanic, both sexes													
2003.....	5,753	39	134	295	1,782	1,332	911	541	304	213	148	47	7
2002.....	5,659	26	176	259	1,810	1,294	885	509	283	211	146	45	15
2001.....	5,272	40	142	246	1,677	1,197	834	470	279	201	146	32	8
2000.....	4,900	40	144	260	1,557	1,066	756	472	246	198	113	34	14
1999.....	4,525	40	148	231	1,384	1,035	684	405	238	204	113	33	10
Hispanic, male													
2003.....	4,340	25	73	198	1,397	1,109	685	397	196	134	89	30	7
2002.....	4,262	16	108	163	1,421	1,052	683	358	192	144	84	26	15
2001.....	3,987	15	82	144	1,330	992	653	340	184	123	99	19	6
2000.....	3,658	24	78	142	1,236	869	574	341	170	123	68	19	14
1999.....	3,337	22	82	136	1,077	828	530	294	146	124	71	17	10
Hispanic, female													
2003.....	1,413	14	61	97	385	223	226	144	108	79	59	17	-
2002.....	1,397	10	68	96	389	242	202	151	91	67	62	19	-
2001.....	1,285	25	60	102	347	205	181	130	95	78	47	13	2
2000.....	1,242	16	66	118	321	197	182	131	76	75	45	15	-
1999.....	1,188	18	66	95	307	207	154	111	92	80	42	16	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)													
Non-Hispanic 1/, both sexes													
2003.....	38,849	106	482	1,344	9,165	5,480	6,031	5,507	3,636	2,738	3,053	1,306	1
2002.....	39,567	97	433	1,350	9,621	5,777	6,169	5,400	3,464	2,890	3,114	1,238	14
2001.....	38,362	104	483	1,436	9,021	5,722	6,217	5,072	3,173	2,780	3,156	1,187	11
2000.....	38,295	128	505	1,507	8,969	5,792	6,147	4,866	3,243	2,829	3,054	1,252	3
1999.....	37,684	144	500	1,534	8,707	5,712	6,026	4,546	3,110	3,059	3,112	1,225	9
Non-Hispanic 1/, male													
2003.....	26,202	55	277	808	6,391	4,032	4,283	3,887	2,473	1,632	1,674	689	1
2002.....	26,677	52	252	796	6,735	4,280	4,284	3,808	2,333	1,734	1,700	690	13
2001.....	25,962	51	273	882	6,426	4,230	4,340	3,579	2,116	1,658	1,756	643	8
2000.....	25,670	66	288	900	6,257	4,243	4,346	3,421	2,122	1,671	1,667	686	3
1999.....	25,072	75	282	920	6,027	4,141	4,184	3,163	2,034	1,798	1,744	696	8
Non-Hispanic 1/, female													
2003.....	12,647	51	205	536	2,774	1,448	1,748	1,620	1,163	1,106	1,379	617	-
2002.....	12,890	45	181	554	2,886	1,497	1,885	1,592	1,131	1,156	1,414	548	1
2001.....	12,400	53	210	554	2,595	1,492	1,877	1,493	1,057	1,122	1,400	544	3
2000.....	12,625	62	217	607	2,712	1,549	1,801	1,445	1,121	1,158	1,387	566	-
1999.....	12,612	69	218	614	2,680	1,571	1,842	1,383	1,076	1,261	1,368	529	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Non-Hispanic white, both sexes

2003.....	31,662	70	333	1,005	7,495	4,182	4,792	4,479	3,008	2,343	2,748	1,206	1
2002.....	32,405	56	280	1,000	7,897	4,412	4,923	4,421	2,918	2,496	2,844	1,149	9
2001.....	31,283	71	323	1,036	7,266	4,424	4,980	4,121	2,652	2,398	2,891	1,113	8
2000.....	31,219	75	340	1,136	7,330	4,379	4,923	3,925	2,728	2,431	2,791	1,158	3
1999.....	30,743	94	315	1,144	7,099	4,425	4,797	3,678	2,579	2,655	2,832	1,122	3

Non-Hispanic white, male

2003.....	21,309	38	200	602	5,177	3,085	3,408	3,176	2,067	1,419	1,505	631	1
2002.....	21,743	32	155	594	5,505	3,262	3,423	3,148	1,968	1,484	1,532	632	8
2001.....	21,073	35	185	631	5,164	3,270	3,468	2,915	1,764	1,436	1,600	598	7
2000.....	20,789	40	197	680	5,094	3,196	3,473	2,781	1,759	1,424	1,505	637	3
1999.....	20,362	55	175	683	4,884	3,201	3,338	2,565	1,689	1,561	1,575	633	3

Non-Hispanic white, female

2003.....	10,353	32	133	403	2,318	1,097	1,384	1,303	941	924	1,243	575	-
2002.....	10,662	24	125	406	2,392	1,150	1,500	1,273	950	1,012	1,312	517	1
2001.....	10,210	36	138	405	2,102	1,154	1,512	1,206	888	962	1,291	515	1
2000.....	10,430	35	143	456	2,236	1,183	1,450	1,144	969	1,007	1,286	521	-
1999.....	10,381	39	140	461	2,215	1,224	1,459	1,113	890	1,094	1,257	489	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Non-Hispanic black, both sexes

2003.....	5,349	28	121	261	1,168	991	974	784	463	285	203	71	-
2002.....	5,338	35	118	278	1,243	1,049	959	734	404	263	191	61	3
2001.....	5,423	18	123	324	1,285	1,007	985	747	410	283	184	54	3
2000.....	5,424	40	132	294	1,216	1,095	977	750	388	298	169	65	-
1999.....	5,388	43	152	331	1,209	1,006	985	685	387	291	214	79	6

Non-Hispanic black, male

2003.....	3,771	12	62	159	871	747	702	569	319	167	123	40	-
2002.....	3,774	17	77	163	891	802	673	516	290	175	125	42	3
2001.....	3,842	7	67	214	935	757	706	550	295	163	116	31	1
2000.....	3,857	21	72	179	868	828	727	524	289	199	112	38	-
1999.....	3,763	17	84	203	869	761	696	499	267	178	132	52	5

Non-Hispanic black, female

2003.....	1,578	16	59	102	297	244	272	215	144	118	80	31	-
2002.....	1,564	18	41	115	352	247	286	218	114	88	66	19	-
2001.....	1,581	11	56	110	350	250	279	197	115	120	68	23	2
2000.....	1,567	19	60	115	348	267	250	226	99	99	57	27	-
1999.....	1,625	26	68	128	340	245	289	186	120	113	82	27	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Origin not stated 2/, both sexes

2003.....	155	-	5	3	25	21	19	26	18	16	9	5	8
2002.....	154	-	1	5	28	21	23	28	17	12	11	2	6
2001.....	154	-	1	4	27	18	32	22	13	9	10	4	14
2000.....	159	-	2	5	34	26	24	23	17	11	6	2	9
1999.....	192	-	2	6	37	31	28	21	22	13	18	3	11

Origin not stated 2/, male

2003.....	113	-	3	3	18	15	16	19	14	12	4	3	6
2002.....	125	-	1	4	22	17	17	23	15	10	9	2	5
2001.....	115	-	1	3	22	14	23	19	9	5	4	2	13
2000.....	123	-	2	1	23	23	20	18	14	8	4	1	9
1999.....	143	-	2	6	23	24	21	16	20	9	12	1	9

Origin not stated 2/, female

2003.....	42	-	2	-	7	6	3	7	4	4	5	2	2
2002.....	29	-	-	1	6	4	6	5	2	2	2	-	1
2001.....	39	-	-	1	5	4	9	3	4	4	6	2	1
2000.....	36	-	-	4	11	3	4	5	3	3	2	1	-
1999.....	49	-	-	-	14	7	7	5	2	4	6	2	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All origins, both sexes

2003.....	1,357	2	17	60	264	204	239	234	147	81	80	25	4
2002.....	1,086	2	17	70	159	138	223	211	109	70	66	17	4
2001.....	1,405	3	28	80	198	177	285	247	133	117	98	29	10
2000.....	1,492	2	31	98	243	212	317	234	122	100	91	35	7
1999.....	2,118	4	48	128	362	317	408	304	182	145	159	54	7

All origins, male

2003.....	1,044	-	9	38	205	166	186	192	113	63	52	16	4
2002.....	882	2	10	47	134	108	187	185	92	56	50	8	3
2001.....	1,067	-	20	49	158	152	235	189	104	84	56	10	10
2000.....	1,125	2	17	61	194	171	258	166	98	75	57	19	7
1999.....	1,559	4	26	91	279	244	319	234	135	97	88	35	7

All origins, female

2003.....	313	2	8	22	59	38	53	42	34	18	28	9	-
2002.....	204	-	7	23	25	30	36	26	17	14	16	9	1
2001.....	338	3	8	31	40	25	50	58	29	33	42	19	-
2000.....	367	-	14	37	49	41	59	68	24	25	34	16	-
1999.....	559	-	22	37	83	73	89	70	47	48	71	19	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Hispanic, both sexes

2003.....	167	-	3	10	32	48	34	17	13	7	1	1	1
2002.....	144	-	2	10	42	21	28	22	7	2	8	-	2
2001.....	200	-	8	7	39	38	42	29	14	5	13	3	2
2000.....	201	-	10	15	34	36	44	25	14	14	5	4	-
1999.....	264	-	8	14	55	60	57	31	8	21	7	2	1

Hispanic, male

2003.....	139	-	2	8	24	43	32	15	9	4	1	-	1
2002.....	119	-	1	6	38	18	25	17	6	2	5	-	1
2001.....	173	-	6	4	37	35	39	22	12	4	10	2	2
2000.....	166	-	7	11	31	32	36	22	11	10	3	3	-
1999.....	220	-	3	12	45	53	47	27	8	17	6	1	1

Hispanic, female

2003.....	28	-	1	2	8	5	2	2	4	3	-	1	-
2002.....	25	-	1	4	4	3	3	5	1	-	3	-	1
2001.....	27	-	2	3	2	3	3	7	2	1	3	1	-
2000.....	35	-	3	4	3	4	8	3	3	4	2	1	-
1999.....	44	-	5	2	10	7	10	4	-	4	1	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Non-Hispanic 1/, both sexes

2003.....	1,185	2	14	50	232	156	203	216	133	73	79	24	3
2002.....	929	2	15	60	116	117	192	184	100	68	57	17	1
2001.....	1,187	3	19	73	158	137	241	214	115	112	85	26	4
2000.....	1,262	2	19	82	206	172	268	207	105	85	85	31	-
1999.....	1,836	4	40	114	304	252	350	271	172	124	151	52	2

Non-Hispanic 1/, male

2003.....	901	-	7	30	181	123	153	176	103	58	51	16	3
2002.....	752	2	9	41	96	90	159	164	84	54	44	8	1
2001.....	878	-	13	45	121	115	194	163	89	80	46	8	4
2000.....	936	2	10	50	161	137	217	142	84	64	53	16	-
1999.....	1,323	4	23	79	231	187	271	205	126	80	81	34	2

Non-Hispanic 1/, female

2003.....	284	2	7	20	51	33	50	40	30	15	28	8	-
2002.....	177	-	6	19	20	27	33	20	16	14	13	9	-
2001.....	309	3	6	28	37	22	47	51	26	32	39	18	-
2000.....	326	-	9	32	45	35	51	65	21	21	32	15	-
1999.....	513	-	17	35	73	65	79	66	46	44	70	18	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Non-Hispanic white, both sexes

2003.....	952	2	11	39	195	117	159	162	112	61	69	22	3
2002.....	719	1	11	44	95	94	140	141	74	55	48	15	1
2001.....	901	2	16	49	126	103	165	165	89	92	73	21	-
2000.....	997	2	9	67	168	132	201	161	84	70	75	28	-
1999.....	1,480	3	25	86	239	195	282	219	151	102	131	46	1

Non-Hispanic white, male

2003.....	715	-	6	21	153	91	122	128	85	48	44	14	3
2002.....	582	1	7	32	81	71	116	124	63	43	36	7	1
2001.....	673	-	12	30	99	89	133	129	73	63	40	5	-
2000.....	741	2	3	41	130	106	165	110	68	51	49	16	-
1999.....	1,051	3	12	57	177	143	218	163	110	67	71	29	1

Non-Hispanic white, female

2003.....	237	2	5	18	42	26	37	34	27	13	25	8	-
2002.....	137	-	4	12	14	23	24	17	11	12	12	8	-
2001.....	228	2	4	19	27	14	32	36	16	29	33	16	-
2000.....	256	-	6	26	38	26	36	51	16	19	26	12	-
1999.....	429	-	13	29	62	52	64	56	41	35	60	17	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Non-Hispanic black, both sexes

2003.....	188	-	3	8	27	30	39	46	19	8	7	1	-
2002.....	168	1	3	13	17	19	40	41	18	9	7	-	-
2001.....	216	-	3	23	21	24	59	38	22	15	6	1	4
2000.....	203	-	7	11	30	32	54	34	15	10	7	3	-
1999.....	248	1	12	20	33	42	52	40	17	14	11	5	1

Non-Hispanic black, male

2003.....	154	-	1	7	21	25	28	42	17	8	4	1	-
2002.....	143	1	2	7	12	17	34	39	16	9	6	-	-
2001.....	159	-	1	15	15	18	48	28	14	13	3	-	4
2000.....	155	-	5	8	24	25	43	26	12	9	3	-	-
1999.....	192	1	9	19	27	34	41	32	12	9	3	4	1

Non-Hispanic black, female

2003.....	34	-	2	1	6	5	11	4	2	-	3	-	-
2002.....	25	-	1	6	5	2	6	2	2	-	1	-	-
2001.....	57	-	2	8	6	6	11	10	8	2	3	1	-
2000.....	48	-	2	3	6	7	11	8	3	1	4	3	-
1999.....	56	-	3	1	6	8	11	8	5	5	8	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Origin not stated 2/, both sexes

2003.....	5	-	-	-	-	-	2	1	1	1	-	-	-
2002.....	13	-	-	-	1	-	3	5	2	-	1	-	1
2001.....	18	-	1	-	1	2	2	4	4	-	-	-	4
2000.....	29	-	2	1	3	4	5	2	3	1	1	-	7
1999.....	18	-	-	-	3	5	1	2	2	-	1	-	4

Origin not stated 2/, male

2003.....	4	-	-	-	-	-	1	1	1	1	-	-	-
2002.....	11	-	-	-	-	-	3	4	2	-	1	-	1
2001.....	16	-	1	-	-	2	2	4	3	-	-	-	4
2000.....	23	-	-	-	2	2	5	2	3	1	1	-	7
1999.....	16	-	-	-	3	4	1	2	1	-	1	-	4

Origin not stated 2/, female

2003.....	1	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	2	-	-	-	1	-	-	1	-	-	-	-	-
2001.....	2	-	-	-	1	-	-	-	1	-	-	-	-
2000.....	6	-	2	1	1	2	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	1	-	-	1	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All origins, both sexes

2003.....	1,957	1	11	63	196	243	394	446	336	162	87	18	-
2002.....	1,900	2	13	54	198	254	388	415	292	168	96	20	-
2001.....	2,095	4	24	57	216	340	459	440	263	175	97	20	-
2000.....	1,903	4	9	52	225	304	417	370	255	177	78	12	-
1999.....	1,904	1	14	76	199	339	407	369	260	158	66	15	-

All origins, male

2003.....	1,592	1	4	37	160	189	327	370	291	134	66	13	-
2002.....	1,587	2	10	29	159	219	328	353	245	157	74	11	-
2001.....	1,663	2	16	37	175	273	362	357	218	145	68	10	-
2000.....	1,590	2	6	34	180	264	362	314	213	144	62	9	-
1999.....	1,531	-	8	53	152	274	322	309	226	134	44	9	-

All origins, female

2003.....	365	-	7	26	36	54	67	76	45	28	21	5	-
2002.....	313	-	3	25	39	35	60	62	47	11	22	9	-
2001.....	432	2	8	20	41	67	97	83	45	30	29	10	-
2000.....	313	2	3	18	45	40	55	56	42	33	16	3	-
1999.....	373	1	6	23	47	65	85	60	34	24	22	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
Hispanic, both sexes													
2003.....	125	-	2	3	20	32	28	19	12	7	2	-	-
2002.....	118	-	4	8	23	25	17	23	10	4	3	1	-
2001.....	302	3	9	12	24	58	66	58	38	23	8	3	-
2000.....	116	2	-	4	25	24	31	11	8	8	3	-	-
1999.....	91	-	1	5	18	30	17	9	7	3	-	1	-
Hispanic, male													
2003.....	95	-	2	3	16	23	23	15	7	4	2	-	-
2002.....	100	-	4	4	20	24	14	19	9	3	2	1	-
2001.....	174	1	4	7	16	40	39	28	18	14	5	2	-
2000.....	97	-	-	2	18	20	29	10	8	7	3	-	-
1999.....	73	-	-	3	14	23	13	9	7	3	-	1	-
Hispanic, female													
2003.....	30	-	-	-	4	9	5	4	5	3	-	-	-
2002.....	18	-	-	4	3	1	3	4	1	1	1	-	-
2001.....	128	2	5	5	8	18	27	30	20	9	3	1	-
2000.....	19	2	-	2	7	4	2	1	-	1	-	-	-
1999.....	18	-	1	2	4	7	4	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
Non-Hispanic 1/, both sexes													
2003.....	1,822	1	9	60	173	210	363	427	321	155	85	18	-
2002.....	1,772	2	9	46	175	226	369	391	280	162	93	19	-
2001.....	1,790	1	15	45	192	282	390	382	225	152	89	17	-
2000.....	1,784	2	9	48	200	280	386	357	247	168	75	12	-
1999.....	1,810	1	13	71	181	308	390	358	253	155	66	14	-
Non-Hispanic 1/, male													
2003.....	1,488	1	2	34	141	165	301	355	282	130	64	13	-
2002.....	1,477	2	6	25	139	192	312	333	234	152	72	10	-
2001.....	1,487	1	12	30	159	233	321	329	200	131	63	8	-
2000.....	1,490	2	6	32	162	244	333	302	205	136	59	9	-
1999.....	1,455	-	8	50	138	250	309	298	219	131	44	8	-
Non-Hispanic 1/, female													
2003.....	334	-	7	26	32	45	62	72	39	25	21	5	-
2002.....	295	-	3	21	36	34	57	58	46	10	21	9	-
2001.....	303	-	3	15	33	49	69	53	25	21	26	9	-
2000.....	294	-	3	16	38	36	53	55	42	32	16	3	-
1999.....	355	1	5	21	43	58	81	60	34	24	22	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Non-Hispanic white, both sexes

2003.....	1,606	1	8	51	136	181	325	378	294	139	75	18	-
2002.....	1,565	2	9	32	154	197	326	347	254	144	82	18	-
2001.....	1,569	1	14	37	171	244	349	325	196	138	80	14	-
2000.....	1,592	2	8	43	171	248	347	316	227	148	71	11	-
1999.....	1,580	1	9	61	155	269	346	314	216	140	57	12	-

Non-Hispanic white, male

2003.....	1,308	1	2	25	109	143	266	316	259	117	57	13	-
2002.....	1,303	2	6	17	121	168	277	298	209	134	62	9	-
2001.....	1,301	1	11	25	138	202	286	282	172	118	58	8	-
2000.....	1,325	2	5	27	137	216	297	269	187	122	55	8	-
1999.....	1,269	-	5	42	120	217	271	264	187	117	40	6	-

Non-Hispanic white, female

2003.....	298	-	6	26	27	38	59	62	35	22	18	5	-
2002.....	262	-	3	15	33	29	49	49	45	10	20	9	-
2001.....	268	-	3	12	33	42	63	43	24	20	22	6	-
2000.....	267	-	3	16	34	32	50	47	40	26	16	3	-
1999.....	311	1	4	19	35	52	75	50	29	23	17	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Non-Hispanic black, both sexes

2003.....	147	-	1	7	26	18	27	32	19	10	7	-	-
2002.....	159	-	-	11	15	21	34	35	22	12	8	1	-
2001.....	164	-	1	7	16	27	27	44	23	9	8	2	-
2000.....	130	-	-	4	16	25	28	25	14	14	3	1	-
1999.....	174	-	4	9	20	27	37	33	26	9	7	2	-

Non-Hispanic black, male

2003.....	128	-	-	7	25	14	25	26	18	8	5	-	-
2002.....	134	-	-	7	14	18	26	27	21	12	8	1	-
2001.....	142	-	1	5	16	22	25	37	23	9	4	-	-
2000.....	115	-	-	4	15	22	26	21	13	10	3	1	-
1999.....	142	-	3	7	13	22	32	26	24	9	4	2	-

Non-Hispanic black, female

2003.....	19	-	1	-	1	4	2	6	1	2	2	-	-
2002.....	25	-	-	4	1	3	8	8	1	-	-	-	-
2001.....	22	-	-	2	-	5	2	7	-	-	4	2	-
2000.....	15	-	-	-	1	3	2	4	1	4	-	-	-
1999.....	32	-	1	2	7	5	5	7	2	-	3	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
All origins, both sexes													
2003.....	61,206	797	1,068	853	3,840	5,261	9,172	9,083	4,729	4,871	9,731	11,745	56
2002.....	58,376	819	1,001	980	3,596	5,085	9,022	8,112	4,180	4,735	9,471	11,329	46
2001.....	54,249	825	1,036	1,013	3,272	4,385	8,118	7,093	3,797	4,553	9,181	10,899	77
2000.....	51,151	707	1,135	1,057	3,085	4,369	7,752	6,313	3,622	4,383	8,416	10,260	52
1999.....	51,437	656	1,186	1,116	2,967	4,456	7,678	5,994	3,473	4,629	8,814	10,399	69
All origins, male													
2003.....	37,241	458	682	578	3,120	4,023	6,506	6,331	3,237	2,946	4,966	4,342	52
2002.....	35,724	481	643	674	2,967	3,959	6,513	5,765	2,904	2,930	4,770	4,080	38
2001.....	33,266	473	653	688	2,721	3,431	5,999	5,259	2,597	2,736	4,669	3,973	67
2000.....	31,651	407	707	723	2,570	3,489	5,884	4,803	2,516	2,618	4,205	3,679	50
1999.....	31,893	393	727	768	2,482	3,586	5,872	4,586	2,454	2,830	4,413	3,722	60
All origins, female													
2003.....	23,965	339	386	275	720	1,238	2,666	2,752	1,492	1,925	4,765	7,403	4
2002.....	22,652	338	358	306	629	1,126	2,509	2,347	1,276	1,805	4,701	7,249	8
2001.....	20,983	352	383	325	551	954	2,119	1,834	1,200	1,817	4,512	6,926	10
2000.....	19,500	300	428	334	515	880	1,868	1,510	1,106	1,765	4,211	6,581	2
1999.....	19,544	263	459	348	485	870	1,806	1,408	1,019	1,799	4,401	6,677	9

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Hispanic, both sexes													
2003.....	4,373	92	186	123	532	755	904	677	310	239	303	245	7
2002.....	4,185	101	159	112	538	687	911	614	266	249	307	230	11
2001.....	3,749	111	151	114	442	645	818	521	233	225	242	228	19
2000.....	3,613	76	152	104	479	650	787	445	224	198	244	240	14
1999.....	3,770	94	184	112	489	665	849	481	213	212	214	224	33
Hispanic, male													
2003.....	3,325	52	123	80	468	652	748	539	237	147	175	97	7
2002.....	3,217	67	102	71	479	615	746	494	200	158	167	107	11
2001.....	2,823	59	103	69	390	552	687	432	177	134	121	81	18
2000.....	2,775	43	89	55	432	578	678	377	158	131	117	103	14
1999.....	2,948	58	112	76	439	590	717	404	173	144	105	101	29
Hispanic, female													
2003.....	1,048	40	63	43	64	103	156	138	73	92	128	148	-
2002.....	968	34	57	41	59	72	165	120	66	91	140	123	-
2001.....	926	52	48	45	52	93	131	89	56	91	121	147	1
2000.....	838	33	63	49	47	72	109	68	66	67	127	137	-
1999.....	822	36	72	36	50	75	132	77	40	68	109	123	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Non-Hispanic 1/, both sexes													
2003.....	56,550	703	881	729	3,296	4,480	8,216	8,338	4,395	4,614	9,403	11,476	19
2002.....	53,907	715	839	862	3,044	4,369	8,062	7,447	3,884	4,464	9,130	11,073	18
2001.....	50,183	709	882	891	2,816	3,716	7,241	6,516	3,529	4,303	8,915	10,644	21
2000.....	47,217	627	979	948	2,593	3,691	6,905	5,789	3,371	4,165	8,147	9,993	9
1999.....	47,339	561	997	1,002	2,466	3,762	6,763	5,447	3,218	4,400	8,567	10,141	15
Non-Hispanic 1/, male													
2003.....	33,699	404	558	497	2,642	3,350	5,717	5,740	2,981	2,784	4,775	4,235	16
2002.....	32,295	412	539	599	2,478	3,320	5,727	5,232	2,674	2,753	4,582	3,963	16
2001.....	30,201	410	547	613	2,318	2,860	5,262	4,781	2,390	2,586	4,532	3,885	17
2000.....	28,640	361	617	666	2,127	2,888	5,158	4,366	2,336	2,473	4,076	3,564	8
1999.....	28,701	335	613	690	2,032	2,972	5,107	4,125	2,248	2,672	4,289	3,605	13
Non-Hispanic 1/, female													
2003.....	22,851	299	323	232	654	1,130	2,499	2,598	1,414	1,830	4,628	7,241	3
2002.....	21,612	303	300	263	566	1,049	2,335	2,215	1,210	1,711	4,548	7,110	2
2001.....	19,982	299	335	278	498	856	1,979	1,735	1,139	1,717	4,383	6,759	4
2000.....	18,577	266	362	282	466	803	1,747	1,423	1,035	1,692	4,071	6,429	1
1999.....	18,638	226	384	312	434	790	1,656	1,322	970	1,728	4,278	6,536	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Non-Hispanic white, both sexes													
2003.....	48,595	423	633	440	2,812	3,758	6,830	6,702	3,629	3,923	8,595	10,832	18
2002.....	45,916	403	565	505	2,518	3,641	6,646	5,950	3,147	3,786	8,342	10,397	16
2001.....	42,509	426	653	566	2,304	3,027	5,846	5,093	2,839	3,681	8,113	9,944	17
2000.....	39,728	376	676	597	2,066	2,971	5,537	4,480	2,714	3,532	7,419	9,355	5
1999.....	39,523	337	693	617	1,917	2,966	5,419	4,181	2,509	3,671	7,737	9,467	9
Non-Hispanic white, male													
2003.....	28,556	244	402	306	2,259	2,819	4,778	4,599	2,430	2,368	4,351	3,985	15
2002.....	27,024	244	359	354	2,039	2,772	4,734	4,175	2,134	2,330	4,175	3,694	14
2001.....	25,133	245	416	380	1,896	2,345	4,301	3,708	1,896	2,220	4,102	3,609	15
2000.....	23,615	222	433	428	1,704	2,343	4,178	3,361	1,857	2,095	3,683	3,307	4
1999.....	23,431	200	425	423	1,568	2,371	4,122	3,150	1,744	2,217	3,865	3,338	8
Non-Hispanic white, female													
2003.....	20,039	179	231	134	553	939	2,052	2,103	1,199	1,555	4,244	6,847	3
2002.....	18,892	159	206	151	479	869	1,912	1,775	1,013	1,456	4,167	6,703	2
2001.....	17,376	181	237	186	408	682	1,545	1,385	943	1,461	4,011	6,335	2
2000.....	16,113	154	243	169	362	628	1,359	1,119	857	1,437	3,736	6,048	1
1999.....	16,092	137	268	194	349	595	1,297	1,031	765	1,454	3,872	6,129	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Non-Hispanic black, both sexes													
2003.....	6,470	255	208	241	369	563	1,155	1,427	651	547	581	472	1
2002.....	6,620	288	229	305	405	587	1,194	1,335	631	534	593	518	1
2001.....	6,436	254	204	276	391	542	1,211	1,274	590	529	632	532	1
2000.....	6,330	228	257	310	400	588	1,194	1,152	566	535	589	507	4
1999.....	6,723	207	260	341	432	663	1,171	1,145	617	629	693	560	5
Non-Hispanic black, male													
2003.....	4,184	146	134	157	288	411	784	988	470	335	297	173	1
2002.....	4,377	154	152	214	343	442	829	941	464	331	309	197	1
2001.....	4,238	151	119	198	319	395	829	959	418	320	338	191	1
2000.....	4,253	124	155	215	326	441	853	883	422	319	314	197	4
1999.....	4,532	122	164	238	372	496	859	884	439	400	344	210	4
Non-Hispanic black, female													
2003.....	2,286	109	74	84	81	152	371	439	181	212	284	299	-
2002.....	2,243	134	77	91	62	145	365	394	167	203	284	321	-
2001.....	2,198	103	85	78	72	147	382	315	172	209	294	341	-
2000.....	2,077	104	102	95	74	147	341	269	144	216	275	310	-
1999.....	2,191	85	96	103	60	167	312	261	178	229	349	350	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)													
Origin not stated 2/, both sexes													
2003.....	283	2	1	1	12	26	52	68	24	18	25	24	30
2002.....	284	3	3	6	14	29	49	51	30	22	34	26	17
2001.....	317	5	3	8	14	24	59	56	35	25	24	27	37
2000.....	321	4	4	5	13	28	60	79	27	20	25	27	29
1999.....	328	1	5	2	12	29	66	66	42	17	33	34	21
Origin not stated 2/, male													
2003.....	217	2	1	1	10	21	41	52	19	15	16	10	29
2002.....	212	2	2	4	10	24	40	39	30	19	21	10	11
2001.....	242	4	3	6	13	19	50	46	30	16	16	7	32
2000.....	236	3	1	2	11	23	48	60	22	14	12	12	28
1999.....	244	-	2	2	11	24	48	57	33	14	19	16	18
Origin not stated 2/, female													
2003.....	66	-	-	-	2	5	11	16	5	3	9	14	1
2002.....	72	1	1	2	4	5	9	12	-	3	13	16	6
2001.....	75	1	-	2	1	5	9	10	5	9	8	20	5
2000.....	85	1	3	3	2	5	12	19	5	6	13	15	1
1999.....	84	1	3	-	1	5	18	9	9	3	14	18	3

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
All origins, both sexes													
2003.....	17,229	14	54	45	230	285	636	1,043	1,220	2,048	5,249	6,404	1
2002.....	16,257	16	37	42	247	307	664	1,017	1,089	1,967	4,880	5,990	1
2001.....	15,019	23	32	66	256	340	647	1,024	1,004	1,833	4,440	5,350	4
2000.....	13,322	8	36	37	237	303	608	871	949	1,660	3,841	4,772	-
1999.....	13,162	12	55	53	242	343	645	824	887	1,663	3,856	4,578	4
All origins, male													
2003.....	8,910	9	39	30	186	237	525	800	827	1,209	2,661	2,386	1
2002.....	8,463	9	28	27	211	269	522	772	771	1,202	2,444	2,208	-
2001.....	8,089	15	25	46	232	295	544	821	718	1,090	2,277	2,023	3
2000.....	7,122	6	23	30	203	272	496	690	680	991	1,947	1,784	-
1999.....	7,109	8	37	40	209	294	528	661	650	1,027	1,950	1,701	4
All origins, female													
2003.....	8,319	5	15	15	44	48	111	243	393	839	2,588	4,018	-
2002.....	7,794	7	9	15	36	38	142	245	318	765	2,436	3,782	1
2001.....	6,930	8	7	20	24	45	103	203	286	743	2,163	3,327	1
2000.....	6,200	2	13	7	34	31	112	181	269	669	1,894	2,988	-
1999.....	6,053	4	18	13	33	49	117	163	237	636	1,906	2,877	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
Hispanic, both sexes													
2003.....	840	1	11	10	38	60	83	100	105	113	163	156	-
2002.....	832	4	9	6	51	69	94	90	87	118	175	129	-
2001.....	757	7	6	14	44	73	100	88	78	103	131	112	1
2000.....	693	3	9	3	46	73	66	77	74	86	134	122	-
1999.....	652	4	9	3	45	68	70	65	63	84	116	123	2
Hispanic, male													
2003.....	557	1	8	5	33	56	80	83	74	64	94	59	-
2002.....	563	3	7	3	49	65	81	70	65	72	90	58	-
2001.....	525	4	5	13	44	71	93	75	62	57	62	38	1
2000.....	485	2	8	3	44	70	62	69	53	60	62	52	-
1999.....	464	4	6	3	43	65	62	56	55	52	60	56	2
Hispanic, female													
2003.....	283	-	3	5	5	4	3	17	31	49	69	97	-
2002.....	269	1	2	3	2	4	13	20	22	46	85	71	-
2001.....	232	3	1	1	-	2	7	13	16	46	69	74	-
2000.....	208	1	1	-	2	3	4	8	21	26	72	70	-
1999.....	188	-	3	-	2	3	8	9	8	32	56	67	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
Non-Hispanic 1/, both sexes													
2003.....	16,337	13	42	35	192	223	548	937	1,109	1,930	5,073	6,235	-
2002.....	15,365	12	28	36	196	238	568	916	995	1,842	4,685	5,849	-
2001.....	14,204	15	26	51	212	266	544	929	921	1,718	4,299	5,223	-
2000.....	12,585	5	27	34	191	229	537	784	869	1,569	3,700	4,640	-
1999.....	12,447	8	46	50	197	273	573	753	813	1,572	3,721	4,440	1
Non-Hispanic 1/, male													
2003.....	8,320	8	30	25	153	179	440	713	750	1,140	2,560	2,322	-
2002.....	7,856	6	21	24	162	204	439	691	699	1,123	2,342	2,145	-
2001.....	7,527	10	20	33	188	223	448	739	651	1,024	2,209	1,982	-
2000.....	6,603	4	15	27	159	201	429	612	622	928	1,881	1,725	-
1999.....	6,602	4	31	37	166	227	464	599	587	969	1,878	1,639	1
Non-Hispanic 1/, female													
2003.....	8,017	5	12	10	39	44	108	224	359	790	2,513	3,913	-
2002.....	7,509	6	7	12	34	34	129	225	296	719	2,343	3,704	-
2001.....	6,677	5	6	18	24	43	96	190	270	694	2,090	3,241	-
2000.....	5,982	1	12	7	32	28	108	172	247	641	1,819	2,915	-
1999.....	5,845	4	15	13	31	46	109	154	226	603	1,843	2,801	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
Non-Hispanic white, both sexes													
2003.....	14,995	9	28	29	167	190	446	744	943	1,711	4,747	5,981	-
2002.....	14,157	8	15	28	168	203	466	762	853	1,644	4,419	5,591	-
2001.....	13,037	8	23	42	193	226	435	768	780	1,540	4,044	4,978	-
2000.....	11,537	2	15	27	172	192	438	646	725	1,393	3,487	4,440	-
1999.....	11,295	4	34	39	168	216	477	593	653	1,379	3,500	4,231	1
Non-Hispanic white, male													
2003.....	7,531	5	22	20	136	154	362	566	638	1,008	2,395	2,225	-
2002.....	7,107	3	11	19	137	175	359	574	596	1,002	2,199	2,032	-
2001.....	6,785	5	17	27	173	191	360	603	551	921	2,067	1,870	-
2000.....	5,949	1	6	22	144	169	354	505	519	829	1,764	1,636	-
1999.....	5,860	3	23	29	144	181	380	474	471	846	1,759	1,549	1
Non-Hispanic white, female													
2003.....	7,464	4	6	9	31	36	84	178	305	703	2,352	3,756	-
2002.....	7,050	5	4	9	31	28	107	188	257	642	2,220	3,559	-
2001.....	6,252	3	6	15	20	35	75	165	229	619	1,977	3,108	-
2000.....	5,588	1	9	5	28	23	84	141	206	564	1,723	2,804	-
1999.....	5,435	1	11	10	24	35	97	119	182	533	1,741	2,682	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
Non-Hispanic black, both sexes													
2003.....	898	2	11	3	17	25	74	150	131	150	185	150	-
2002.....	821	3	7	6	17	23	75	128	110	135	154	163	-
2001.....	837	6	3	6	13	26	90	128	108	143	173	141	-
2000.....	745	2	12	7	12	22	71	111	114	134	137	123	-
1999.....	865	2	11	10	21	42	73	136	127	150	153	140	-
Non-Hispanic black, male													
2003.....	519	1	7	3	10	18	57	112	86	87	85	53	-
2002.....	518	2	6	5	15	20	58	98	78	81	85	70	-
2001.....	537	4	3	4	9	19	72	109	76	85	97	59	-
2000.....	471	2	9	5	10	18	54	88	87	76	70	52	-
1999.....	554	1	8	7	16	34	65	107	90	100	77	49	-
Non-Hispanic black, female													
2003.....	379	1	4	-	7	7	17	38	45	63	100	97	-
2002.....	303	1	1	1	2	3	17	30	32	54	69	93	-
2001.....	300	2	-	2	4	7	18	19	32	58	76	82	-
2000.....	274	-	3	2	2	4	17	23	27	58	67	71	-
1999.....	311	1	3	3	5	8	8	29	37	50	76	91	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Falls (W00-W19)													
Origin not stated 2/, both sexes													
2003.....	52	-	1	-	-	2	5	6	6	5	13	13	1
2002.....	60	-	-	-	-	-	2	11	7	7	20	12	1
2001.....	58	1	-	1	-	1	3	7	5	12	10	15	3
2000.....	44	-	-	-	-	1	5	10	6	5	7	10	-
1999.....	63	-	-	-	-	2	2	6	11	7	19	15	1
Origin not stated 2/, male													
2003.....	33	-	1	-	-	2	5	4	3	5	7	5	1
2002.....	44	-	-	-	-	-	2	11	7	7	12	5	-
2001.....	37	1	-	-	-	1	3	7	5	9	6	3	2
2000.....	34	-	-	-	-	1	5	9	5	3	4	7	-
1999.....	43	-	-	-	-	2	2	6	8	6	12	6	1
Origin not stated 2/, female													
2003.....	19	-	-	-	-	-	-	2	3	-	6	8	-
2002.....	16	-	-	-	-	-	-	-	-	-	8	7	1
2001.....	21	-	-	1	-	-	-	-	-	3	4	12	1
2000.....	10	-	-	-	-	-	-	1	1	2	3	3	-
1999.....	20	-	-	-	-	-	-	-	3	1	7	9	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
All origins, both sexes													
2003.....	730	-	7	49	200	120	118	92	62	39	37	6	-
2002.....	762	1	11	48	210	143	123	95	56	36	32	7	-
2001.....	802	-	15	57	206	122	146	103	69	33	41	8	2
2000.....	776	1	18	67	202	131	153	78	50	41	24	11	-
1999.....	824	-	12	76	251	143	136	92	48	33	23	10	-
All origins, male													
2003.....	656	-	7	46	182	110	105	78	59	30	33	6	-
2002.....	667	1	7	39	195	129	98	84	50	32	26	6	-
2001.....	690	-	12	49	193	103	121	87	55	26	36	7	1
2000.....	671	-	11	62	183	109	125	69	44	34	23	11	-
1999.....	707	-	7	60	229	122	112	78	41	31	20	7	-
All origins, female													
2003.....	74	-	-	3	18	10	13	14	3	9	4	-	-
2002.....	95	-	4	9	15	14	25	11	6	4	6	1	-
2001.....	112	-	3	8	13	19	25	16	14	7	5	1	1
2000.....	105	1	7	5	19	22	28	9	6	7	1	-	-
1999.....	117	-	5	16	22	21	24	14	7	2	3	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
Hispanic, both sexes													
2003.....	71	-	-	8	33	14	8	3	3	1	1	-	-
2002.....	60	-	3	5	24	18	5	3	1	-	1	-	-
2001.....	86	-	1	4	36	21	12	6	1	1	2	1	1
2000.....	59	-	2	6	25	12	11	2	1	-	-	-	-
1999.....	66	-	-	3	31	17	8	4	1	2	-	-	-
Hispanic, male													
2003.....	64	-	-	6	30	14	7	2	3	1	1	-	-
2002.....	51	-	1	3	21	18	3	3	1	-	1	-	-
2001.....	76	-	1	3	33	16	12	5	1	1	2	1	1
2000.....	50	-	1	4	24	10	9	2	-	-	-	-	-
1999.....	61	-	-	3	29	16	7	3	1	2	-	-	-
Hispanic, female													
2003.....	7	-	-	2	3	-	1	1	-	-	-	-	-
2002.....	9	-	2	2	3	-	2	-	-	-	-	-	-
2001.....	10	-	-	1	3	5	-	1	-	-	-	-	-
2000.....	9	-	1	2	1	2	2	-	1	-	-	-	-
1999.....	5	-	-	-	2	1	1	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
Non-Hispanic 1/, both sexes													
2003.....	658	-	7	41	167	106	109	89	59	38	36	6	-
2002.....	700	1	8	43	185	125	118	92	55	36	30	7	-
2001.....	715	-	14	53	169	101	134	97	68	32	39	7	1
2000.....	717	1	16	61	177	119	142	76	49	41	24	11	-
1999.....	758	-	12	73	220	126	128	88	47	31	23	10	-
Non-Hispanic 1/, male													
2003.....	591	-	7	40	152	96	97	76	56	29	32	6	-
2002.....	614	1	6	36	173	111	95	81	49	32	24	6	-
2001.....	613	-	11	46	159	87	109	82	54	25	34	6	-
2000.....	621	-	10	58	159	99	116	67	44	34	23	11	-
1999.....	646	-	7	57	200	106	105	75	40	29	20	7	-
Non-Hispanic 1/, female													
2003.....	67	-	-	1	15	10	12	13	3	9	4	-	-
2002.....	86	-	2	7	12	14	23	11	6	4	6	1	-
2001.....	102	-	3	7	10	14	25	15	14	7	5	1	1
2000.....	96	1	6	3	18	20	26	9	5	7	1	-	-
1999.....	112	-	5	16	20	20	23	13	7	2	3	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
Non-Hispanic white, both sexes													
2003.....	515	-	4	30	115	69	95	77	56	33	32	4	-
2002.....	531	1	6	31	118	82	97	78	49	35	27	7	-
2001.....	591	-	9	41	117	82	116	92	61	29	36	7	1
2000.....	555	1	9	46	117	83	118	65	46	37	22	11	-
1999.....	585	-	5	54	145	98	108	77	40	28	20	10	-
Non-Hispanic white, male													
2003.....	461	-	4	29	103	61	87	65	54	26	28	4	-
2002.....	459	1	4	25	110	70	77	69	45	31	21	6	-
2001.....	506	-	7	38	109	71	96	78	48	22	31	6	-
2000.....	483	-	5	44	106	66	99	59	41	31	21	11	-
1999.....	495	-	3	43	131	80	87	65	36	26	17	7	-
Non-Hispanic white, female													
2003.....	54	-	-	1	12	8	8	12	2	7	4	-	-
2002.....	72	-	2	6	8	12	20	9	4	4	6	1	-
2001.....	85	-	2	3	8	11	20	14	13	7	5	1	1
2000.....	72	1	4	2	11	17	19	6	5	6	1	-	-
1999.....	90	-	2	11	14	18	21	12	4	2	3	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental discharge of firearms (W32-W34)													
Non-Hispanic black, both sexes													
2003.....	127	-	3	9	49	34	12	7	3	5	3	2	-
2002.....	153	-	2	9	60	40	20	13	6	-	3	-	-
2001.....	110	-	5	8	48	17	15	5	6	3	3	-	-
2000.....	137	-	5	12	52	32	19	10	3	3	1	-	-
1999.....	156	-	7	17	67	23	18	11	7	3	3	-	-
Non-Hispanic black, male													
2003.....	115	-	3	9	46	33	8	6	2	3	3	2	-
2002.....	140	-	2	8	57	38	17	11	4	-	3	-	-
2001.....	97	-	4	6	46	14	12	4	5	3	3	-	-
2000.....	117	-	3	12	46	29	14	7	3	2	1	-	-
1999.....	135	-	4	12	61	22	16	10	4	3	3	-	-
Non-Hispanic black, female													
2003.....	12	-	-	-	3	1	4	1	1	2	-	-	-
2002.....	13	-	-	1	3	2	3	2	2	-	-	-	-
2001.....	13	-	1	2	2	3	3	1	1	-	-	-	-
2000.....	20	-	2	-	6	3	5	3	-	1	-	-	-
1999.....	21	-	3	5	6	1	2	1	3	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental drowning and submersion (W65-W74)													
All origins, both sexes													
2003.....	3,306	58	456	268	567	356	462	397	265	202	162	83	30
2002.....	3,447	63	454	321	629	433	526	374	212	169	167	79	20
2001.....	3,281	68	458	333	596	374	462	359	206	159	157	75	34
2000.....	3,482	75	493	375	646	419	480	354	217	179	156	64	24
1999.....	3,529	68	490	369	647	446	475	370	219	185	158	68	34
All origins, male													
2003.....	2,632	34	301	199	525	323	392	320	211	145	104	48	30
2002.....	2,761	40	312	244	578	380	439	301	161	132	106	50	18
2001.....	2,560	30	291	237	544	328	376	299	155	124	102	45	29
2000.....	2,735	43	316	267	583	364	406	299	179	133	88	34	23
1999.....	2,794	39	310	275	592	397	402	311	170	129	111	28	30
All origins, female													
2003.....	674	24	155	69	42	33	70	77	54	57	58	35	-
2002.....	686	23	142	77	51	53	87	73	51	37	61	29	2
2001.....	721	38	167	96	52	46	86	60	51	35	55	30	5
2000.....	747	32	177	108	63	55	74	55	38	46	68	30	1
1999.....	735	29	180	94	55	49	73	59	49	56	47	40	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental drowning and submersion (W65-W74)													
Hispanic, both sexes													
2003.....	458	12	82	45	112	85	51	27	15	14	6	2	7
2002.....	464	18	65	29	143	80	55	30	15	12	10	2	5
2001.....	421	16	78	40	101	74	57	25	7	5	4	2	12
2000.....	433	9	75	43	147	62	41	20	11	12	4	1	8
1999.....	480	16	78	46	145	79	45	25	8	7	6	-	25
Hispanic, male													
2003.....	399	7	67	33	109	79	47	25	14	7	3	1	7
2002.....	409	13	48	23	137	78	48	28	10	10	7	2	5
2001.....	348	9	53	28	99	65	48	21	6	3	3	2	11
2000.....	350	6	44	22	135	60	36	18	8	10	3	-	8
1999.....	405	12	43	35	140	75	38	23	7	7	4	-	21
Hispanic, female													
2003.....	59	5	15	12	3	6	4	2	1	7	3	1	-
2002.....	55	5	17	6	6	2	7	2	5	2	3	-	-
2001.....	73	7	25	12	2	9	9	4	1	2	1	-	1
2000.....	83	3	31	21	12	2	5	2	3	2	1	1	-
1999.....	75	4	35	11	5	4	7	2	1	-	2	-	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental drowning and submersion (W65-W74)													
Non-Hispanic 1/, both sexes													
2003.....	2,816	46	374	222	452	271	407	364	248	187	156	81	8
2002.....	2,952	45	387	289	485	347	467	342	195	156	155	77	7
2001.....	2,832	51	380	291	493	300	403	329	198	153	153	73	8
2000.....	3,018	65	417	331	496	355	437	332	205	166	152	61	1
1999.....	3,029	52	409	323	500	363	429	343	210	178	152	67	3
Non-Hispanic 1/, male													
2003.....	2,202	27	234	165	413	244	342	289	195	137	101	47	8
2002.....	2,326	27	262	219	440	298	387	271	149	121	97	48	7
2001.....	2,188	21	238	207	443	263	326	275	148	120	99	43	5
2000.....	2,358	37	272	244	445	302	368	279	170	122	85	33	1
1999.....	2,370	27	265	240	450	318	363	286	162	122	107	27	3
Non-Hispanic 1/, female													
2003.....	614	19	140	57	39	27	65	75	53	50	55	34	-
2002.....	626	18	125	70	45	49	80	71	46	35	58	29	-
2001.....	644	30	142	84	50	37	77	54	50	33	54	30	3
2000.....	660	28	145	87	51	53	69	53	35	44	67	28	-
1999.....	659	25	144	83	50	45	66	57	48	56	45	40	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental drowning and submersion (W65-W74)													
Non-Hispanic white, both sexes													
2003.....	2,152	32	306	118	306	187	320	297	208	163	139	68	8
2002.....	2,143	32	303	134	316	245	360	260	165	126	131	64	7
2001.....	2,082	36	321	135	322	202	306	265	158	133	134	66	4
2000.....	2,235	38	342	176	316	258	344	255	175	144	134	53	-
1999.....	2,230	30	337	142	317	250	349	291	174	147	132	59	2
Non-Hispanic white, male													
2003.....	1,656	17	187	86	279	166	267	230	160	121	93	42	8
2002.....	1,669	18	203	96	284	209	294	205	125	99	85	44	7
2001.....	1,577	16	205	87	284	171	245	220	114	105	88	39	3
2000.....	1,737	22	226	129	278	222	287	212	150	104	78	29	-
1999.....	1,708	14	217	99	279	216	290	241	133	99	93	25	2
Non-Hispanic white, female													
2003.....	496	15	119	32	27	21	53	67	48	42	46	26	-
2002.....	474	14	100	38	32	36	66	55	40	27	46	20	-
2001.....	505	20	116	48	38	31	61	45	44	28	46	27	1
2000.....	498	16	116	47	38	36	57	43	25	40	56	24	-
1999.....	522	16	120	43	38	34	59	50	41	48	39	34	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental drowning and submersion (W65-W74)													
Non-Hispanic black, both sexes													
2003.....	466	13	52	87	100	59	48	45	33	16	8	5	-
2002.....	604	10	62	134	126	74	74	57	21	18	17	11	-
2001.....	570	11	47	133	133	62	70	51	32	13	13	4	1
2000.....	580	25	54	134	128	58	77	52	17	15	12	7	1
1999.....	624	21	59	158	137	86	58	36	28	21	15	5	-
Non-Hispanic black, male													
2003.....	394	9	37	66	94	54	45	42	29	10	5	3	-
2002.....	502	6	43	107	122	65	66	47	18	15	9	4	-
2001.....	471	3	28	102	126	61	58	42	26	12	9	3	1
2000.....	470	14	33	101	123	52	67	45	13	13	4	4	1
1999.....	521	12	40	124	128	78	53	34	22	17	11	2	-
Non-Hispanic black, female													
2003.....	72	4	15	21	6	5	3	3	4	6	3	2	-
2002.....	102	4	19	27	4	9	8	10	3	3	8	7	-
2001.....	99	8	19	31	7	1	12	9	6	1	4	1	-
2000.....	110	11	21	33	5	6	10	7	4	2	8	3	-
1999.....	103	9	19	34	9	8	5	2	6	4	4	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
All origins, both sexes													
2003.....	3,369	30	224	215	199	252	395	526	392	419	460	253	4
2002.....	3,159	36	221	253	193	266	361	448	340	385	432	221	3
2001.....	3,309	50	225	252	209	247	441	419	378	411	423	251	3
2000.....	3,377	37	290	266	192	241	402	439	369	401	472	263	5
1999.....	3,348	41	302	262	201	282	421	368	319	416	487	245	4
All origins, male													
2003.....	1,948	8	131	112	121	154	257	329	254	244	228	107	3
2002.....	1,935	21	126	151	132	170	251	292	219	242	231	97	3
2001.....	1,998	30	142	152	137	161	308	288	236	229	204	108	3
2000.....	2,025	20	174	145	121	157	268	303	241	232	244	115	5
1999.....	2,025	21	173	147	133	197	300	254	194	244	247	111	4
All origins, female													
2003.....	1,421	22	93	103	78	98	138	197	138	175	232	146	1
2002.....	1,224	15	95	102	61	96	110	156	121	143	201	124	-
2001.....	1,311	20	83	100	72	86	133	131	142	182	219	143	-
2000.....	1,352	17	116	121	71	84	134	136	128	169	228	148	-
1999.....	1,323	20	129	115	68	85	121	114	125	172	240	134	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Hispanic, both sexes													
2003.....	203	4	27	26	15	25	24	32	14	12	15	9	-
2002.....	223	5	36	27	25	17	27	22	17	20	16	10	1
2001.....	176	9	18	23	19	21	28	17	13	9	9	10	-
2000.....	178	4	21	19	21	19	25	11	17	18	17	6	-
1999.....	192	5	41	23	24	17	21	16	14	11	15	5	-
Hispanic, male													
2003.....	128	2	11	12	9	18	17	26	10	9	9	5	-
2002.....	144	3	18	17	19	9	21	18	12	14	8	4	1
2001.....	109	5	13	11	10	16	19	11	9	6	5	4	-
2000.....	105	2	11	10	11	14	17	7	9	10	10	4	-
1999.....	127	3	23	12	16	12	16	12	11	8	9	5	-
Hispanic, female													
2003.....	75	2	16	14	6	7	7	6	4	3	6	4	-
2002.....	79	2	18	10	6	8	6	4	5	6	8	6	-
2001.....	67	4	5	12	9	5	9	6	4	3	4	6	-
2000.....	73	2	10	9	10	5	8	4	8	8	7	2	-
1999.....	65	2	18	11	8	5	5	4	3	3	6	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Non-Hispanic 1/, both sexes													
2003.....	3,147	26	197	189	183	227	367	491	375	404	443	243	2
2002.....	2,918	31	184	225	168	247	334	424	317	362	414	211	1
2001.....	3,106	41	205	226	190	223	410	398	361	398	412	240	2
2000.....	3,178	31	268	246	169	222	375	426	349	382	451	256	3
1999.....	3,133	36	260	239	173	264	395	347	302	404	470	240	3
Non-Hispanic 1/, male													
2003.....	1,809	6	120	100	111	136	240	302	241	233	218	101	1
2002.....	1,780	18	108	133	113	159	230	274	201	227	223	93	1
2001.....	1,873	25	127	139	127	145	286	274	224	222	199	103	2
2000.....	1,907	16	163	135	108	143	250	295	230	222	232	110	3
1999.....	1,879	18	150	135	113	184	279	238	181	235	237	106	3
Non-Hispanic 1/, female													
2003.....	1,338	20	77	89	72	91	127	189	134	171	225	142	1
2002.....	1,138	13	76	92	55	88	104	150	116	135	191	118	-
2001.....	1,233	16	78	87	63	78	124	124	137	176	213	137	-
2000.....	1,271	15	105	111	61	79	125	131	119	160	219	146	-
1999.....	1,254	18	110	104	60	80	116	109	121	169	233	134	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Non-Hispanic white, both sexes													
2003.....	2,313	18	118	89	141	170	292	362	286	305	340	190	2
2002.....	2,055	16	90	112	113	171	258	317	227	268	325	157	1
2001.....	2,204	26	124	136	139	153	297	284	261	291	311	180	2
2000.....	2,194	17	139	139	120	151	267	320	242	278	328	190	3
1999.....	2,172	21	137	136	125	199	286	253	218	281	343	171	2
Non-Hispanic white, male													
2003.....	1,344	5	74	46	91	102	197	227	183	178	159	81	1
2002.....	1,271	12	52	73	78	115	173	204	146	166	182	69	1
2001.....	1,322	14	82	82	96	105	207	190	160	157	149	78	2
2000.....	1,330	7	88	76	79	99	179	222	160	159	172	86	3
1999.....	1,325	11	80	76	83	145	204	173	136	160	174	81	2
Non-Hispanic white, female													
2003.....	969	13	44	43	50	68	95	135	103	127	181	109	1
2002.....	784	4	38	39	35	56	85	113	81	102	143	88	-
2001.....	882	12	42	54	43	48	90	94	101	134	162	102	-
2000.....	864	10	51	63	41	52	88	98	82	119	156	104	-
1999.....	847	10	57	60	42	54	82	80	82	121	169	90	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Non-Hispanic black, both sexes													
2003.....	767	7	75	94	38	47	66	122	79	90	99	50	-
2002.....	765	13	88	95	48	61	60	98	84	83	82	53	-
2001.....	829	14	80	85	42	63	103	105	89	99	92	57	-
2000.....	915	14	120	101	42	64	97	97	100	98	117	65	-
1999.....	887	15	110	102	40	52	100	86	80	113	120	68	1
Non-Hispanic black, male													
2003.....	430	1	42	50	18	28	39	72	53	53	56	18	-
2002.....	454	6	53	53	31	35	45	64	52	53	38	24	-
2001.....	501	10	44	53	24	34	73	79	56	61	44	23	-
2000.....	539	9	70	57	25	39	63	67	67	60	58	24	-
1999.....	507	7	63	58	24	29	69	59	44	69	59	25	1
Non-Hispanic black, female													
2003.....	337	6	33	44	20	19	27	50	26	37	43	32	-
2002.....	311	7	35	42	17	26	15	34	32	30	44	29	-
2001.....	328	4	36	32	18	29	30	26	33	38	48	34	-
2000.....	376	5	50	44	17	25	34	30	33	38	59	41	-
1999.....	380	8	47	44	16	23	31	27	36	44	61	43	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Origin not stated 2/, both sexes													
2003.....	19	-	-	-	1	-	4	3	3	3	2	1	2
2002.....	18	-	1	1	-	2	-	2	6	3	2	-	1
2001.....	27	-	2	3	-	3	3	4	4	4	2	1	1
2000.....	21	2	1	1	2	-	2	2	3	1	4	1	2
1999.....	23	-	1	-	4	1	5	5	3	1	2	-	1
Origin not stated 2/, male													
2003.....	11	-	-	-	1	-	-	1	3	2	1	1	2
2002.....	11	-	-	1	-	2	-	-	6	1	-	-	1
2001.....	16	-	2	2	-	-	3	3	3	1	-	1	1
2000.....	13	2	-	-	2	-	1	1	2	-	2	1	2
1999.....	19	-	-	-	4	1	5	4	2	1	1	-	1
Origin not stated 2/, female													
2003.....	8	-	-	-	-	-	4	2	-	1	1	-	-
2002.....	7	-	1	-	-	-	-	2	-	2	2	-	-
2001.....	11	-	-	1	-	3	-	1	1	3	2	-	-
2000.....	8	-	1	1	-	-	1	1	1	1	2	-	-
1999.....	4	-	1	-	-	-	-	1	1	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
All origins, both sexes													
2003.....	19,457	20	49	61	1,999	3,435	6,230	5,434	1,370	403	284	166	6
2002.....	17,550	26	31	43	1,679	3,116	6,007	4,682	1,098	383	295	181	9
2001.....	14,078	15	31	50	1,362	2,507	5,036	3,547	798	291	279	152	10
2000.....	12,757	14	32	45	1,160	2,380	4,663	3,061	688	278	245	184	7
1999.....	12,186	12	34	40	964	2,355	4,549	2,844	668	300	265	143	12
All origins, male													
2003.....	13,176	9	27	31	1,562	2,514	4,166	3,598	841	223	142	58	5
2002.....	12,059	21	14	28	1,306	2,319	4,122	3,136	690	209	136	71	7
2001.....	9,885	8	21	23	1,080	1,862	3,522	2,503	500	170	137	49	10
2000.....	9,138	11	21	27	920	1,820	3,384	2,223	401	147	102	76	6
1999.....	8,887	11	17	25	754	1,828	3,349	2,106	448	172	116	53	8
All origins, female													
2003.....	6,281	11	22	30	437	921	2,064	1,836	529	180	142	108	1
2002.....	5,491	5	17	15	373	797	1,885	1,546	408	174	159	110	2
2001.....	4,193	7	10	27	282	645	1,514	1,044	298	121	142	103	-
2000.....	3,619	3	11	18	240	560	1,279	838	287	131	143	108	1
1999.....	3,299	1	17	15	210	527	1,200	738	220	128	149	90	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Hispanic, both sexes													
2003.....	1,802	4	13	5	219	421	610	401	91	22	10	6	-
2002.....	1,652	6	4	9	195	364	586	368	79	26	13	2	-
2001.....	1,324	1	5	10	144	334	475	263	64	19	7	1	1
2000.....	1,308	2	4	4	132	344	493	251	53	15	5	5	-
1999.....	1,502	4	3	6	146	383	585	289	58	18	5	2	3
Hispanic, male													
2003.....	1,428	2	5	3	179	347	483	313	73	17	3	3	-
2002.....	1,324	5	-	4	161	312	466	288	60	18	9	1	-
2001.....	1,061	1	3	2	111	274	385	218	50	13	3	-	1
2000.....	1,090	2	2	2	113	295	415	211	35	10	2	3	-
1999.....	1,271	4	3	4	123	332	493	240	51	15	2	1	3
Hispanic, female													
2003.....	374	2	8	2	40	74	127	88	18	5	7	3	-
2002.....	328	1	4	5	34	52	120	80	19	8	4	1	-
2001.....	263	-	2	8	33	60	90	45	14	6	4	1	-
2000.....	218	-	2	2	19	49	78	40	18	5	3	2	-
1999.....	231	-	-	2	23	51	92	49	7	3	3	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Non-Hispanic 1/, both sexes													
2003.....	17,537	16	36	56	1,776	2,994	5,590	4,986	1,269	377	273	160	4
2002.....	15,789	20	27	34	1,476	2,735	5,382	4,290	1,011	351	280	179	4
2001.....	12,631	14	26	40	1,210	2,157	4,519	3,250	719	270	271	151	4
2000.....	11,311	12	27	38	1,022	2,018	4,131	2,758	624	259	240	179	3
1999.....	10,542	8	31	33	814	1,955	3,907	2,516	596	280	258	139	5
Non-Hispanic 1/, male													
2003.....	11,655	7	22	28	1,381	2,151	3,659	3,250	759	202	138	55	3
2002.....	10,649	16	14	24	1,140	1,993	3,626	2,831	622	185	125	70	3
2001.....	8,720	7	18	21	962	1,574	3,103	2,257	437	155	133	49	4
2000.....	7,944	9	18	24	802	1,512	2,938	1,975	357	134	100	73	2
1999.....	7,510	7	14	20	627	1,482	2,817	1,833	387	155	114	51	3
Non-Hispanic 1/, female													
2003.....	5,882	9	14	28	395	843	1,931	1,736	510	175	135	105	1
2002.....	5,140	4	13	10	336	742	1,756	1,459	389	166	155	109	1
2001.....	3,911	7	8	19	248	583	1,416	993	282	115	138	102	-
2000.....	3,367	3	9	14	220	506	1,193	783	267	125	140	106	1
1999.....	3,032	1	17	13	187	473	1,090	683	209	125	144	88	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Non-Hispanic white, both sexes													
2003.....	14,829	8	19	31	1,659	2,611	4,708	4,058	1,044	292	253	143	3
2002.....	13,165	7	15	28	1,357	2,375	4,496	3,402	772	299	250	162	2
2001.....	10,269	8	16	30	1,078	1,836	3,698	2,468	546	226	229	130	4
2000.....	9,124	8	15	24	916	1,683	3,331	2,071	482	216	224	152	2
1999.....	8,311	6	17	21	694	1,581	3,096	1,867	444	224	238	120	3
Non-Hispanic white, male													
2003.....	9,855	6	11	17	1,292	1,892	3,098	2,625	596	144	127	45	2
2002.....	8,849	7	6	21	1,048	1,737	3,045	2,225	436	150	108	65	1
2001.....	7,111	4	12	17	864	1,358	2,579	1,688	304	129	109	43	4
2000.....	6,413	6	9	16	736	1,272	2,399	1,464	251	104	93	62	1
1999.....	5,921	5	6	13	536	1,217	2,265	1,337	263	124	105	48	2
Non-Hispanic white, female													
2003.....	4,974	2	8	14	367	719	1,610	1,433	448	148	126	98	1
2002.....	4,316	-	9	7	309	638	1,451	1,177	336	149	142	97	1
2001.....	3,158	4	4	13	214	478	1,119	780	242	97	120	87	-
2000.....	2,711	2	6	8	180	411	932	607	231	112	131	90	1
1999.....	2,390	1	11	8	158	364	831	530	181	100	133	72	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Non-Hispanic black, both sexes													
2003.....	2,354	7	15	17	86	303	770	848	204	68	19	16	1
2002.....	2,326	12	11	6	86	299	786	825	221	41	26	12	1
2001.....	2,122	6	8	6	93	273	749	736	157	39	34	21	-
2000.....	1,985	4	11	8	76	294	732	645	132	41	15	26	1
1999.....	2,008	1	12	7	93	328	729	613	139	49	18	17	2
Non-Hispanic black, male													
2003.....	1,579	1	9	7	64	204	497	570	154	52	10	10	1
2002.....	1,601	8	8	3	66	213	517	564	174	31	15	1	1
2001.....	1,452	3	5	2	67	186	477	540	121	25	20	6	-
2000.....	1,400	3	8	6	49	209	496	480	101	29	7	11	1
1999.....	1,450	1	7	5	76	231	503	471	114	29	9	3	1
Non-Hispanic black, female													
2003.....	775	6	6	10	22	99	273	278	50	16	9	6	-
2002.....	725	4	3	3	20	86	269	261	47	10	11	11	-
2001.....	670	3	3	4	26	87	272	196	36	14	14	15	-
2000.....	585	1	3	2	27	85	236	165	31	12	8	15	-
1999.....	558	-	5	2	17	97	226	142	25	20	9	14	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Origin not stated 2/, both sexes													
2003.....	118	-	-	-	4	20	30	47	10	4	1	-	2
2002.....	109	-	-	-	8	17	39	24	8	6	2	-	5
2001.....	123	-	-	-	8	16	42	34	15	2	1	-	5
2000.....	138	-	1	3	6	18	39	52	11	4	-	-	4
1999.....	142	-	-	1	4	17	57	39	14	2	2	2	4
Origin not stated 2/, male													
2003.....	93	-	-	-	2	16	24	35	9	4	1	-	2
2002.....	86	-	-	-	5	14	30	17	8	6	2	-	4
2001.....	104	-	-	-	7	14	34	28	13	2	1	-	5
2000.....	104	-	1	1	5	13	31	37	9	3	-	-	4
1999.....	106	-	-	1	4	14	39	33	10	2	-	1	2
Origin not stated 2/, female													
2003.....	25	-	-	-	2	4	6	12	1	-	-	-	-
2002.....	23	-	-	-	3	3	9	7	-	-	-	-	1
2001.....	19	-	-	-	1	2	8	6	2	-	-	-	-
2000.....	34	-	-	2	1	5	8	15	2	1	-	-	-
1999.....	36	-	-	-	-	3	18	6	4	-	2	1	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
All origins, both sexes													
2003.....	17,115	675	278	215	645	813	1,331	1,591	1,420	1,760	3,539	4,833	15
2002.....	17,201	677	247	273	638	820	1,341	1,496	1,385	1,795	3,665	4,851	13
2001.....	17,760	669	275	255	643	795	1,386	1,641	1,342	1,826	3,841	5,063	24
2000.....	17,437	572	266	267	648	895	1,446	1,510	1,349	1,824	3,678	4,966	16
1999.....	18,388	523	293	316	662	887	1,452	1,496	1,332	2,032	4,025	5,355	15
All origins, male													
2003.....	9,919	398	177	160	544	685	1,061	1,206	1,045	1,095	1,798	1,737	13
2002.....	9,839	389	156	185	545	692	1,081	1,180	1,013	1,113	1,827	1,648	10
2001.....	10,044	390	162	181	535	682	1,128	1,261	933	1,097	1,913	1,741	21
2000.....	9,960	327	162	192	560	767	1,205	1,219	971	1,081	1,801	1,659	16
1999.....	10,371	314	183	221	565	748	1,181	1,176	951	1,227	1,969	1,822	14
All origins, female													
2003.....	7,196	277	101	55	101	128	270	385	375	665	1,741	3,096	2
2002.....	7,362	288	91	88	93	128	260	316	372	682	1,838	3,203	3
2001.....	7,716	279	113	74	108	113	258	380	409	729	1,928	3,322	3
2000.....	7,477	245	104	75	88	128	241	291	378	743	1,877	3,307	-
1999.....	8,017	209	110	95	97	139	271	320	381	805	2,056	3,533	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Hispanic, both sexes

2003.....	999	71	53	29	115	150	128	114	82	77	108	72	-
2002.....	954	68	42	36	100	139	144	101	67	73	92	87	5
2001.....	985	78	43	23	98	122	146	122	70	88	89	102	4
2000.....	942	58	41	29	108	140	151	84	68	67	84	106	6
1999.....	878	65	53	31	98	101	120	82	69	90	72	94	3

Hispanic, male

2003.....	749	40	32	21	108	138	114	90	63	49	65	29	-
2002.....	726	43	28	21	92	133	127	87	52	44	52	42	5
2001.....	704	40	28	12	93	110	130	102	49	54	46	36	4
2000.....	695	31	23	14	105	129	139	70	53	41	40	44	6
1999.....	620	35	37	19	88	90	101	70	48	60	30	39	3

Hispanic, female

2003.....	250	31	21	8	7	12	14	24	19	28	43	43	-
2002.....	228	25	14	15	8	6	17	14	15	29	40	45	-
2001.....	281	38	15	11	5	12	16	20	21	34	43	66	-
2000.....	247	27	18	15	3	11	12	14	15	26	44	62	-
1999.....	258	30	16	12	10	11	19	12	21	30	42	55	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
Non-Hispanic 1/, both sexes													
2003.....	16,055	602	225	186	526	659	1,195	1,471	1,335	1,678	3,422	4,751	5
2002.....	16,183	606	205	235	534	677	1,193	1,383	1,311	1,717	3,566	4,750	6
2001.....	16,695	588	231	230	542	669	1,231	1,513	1,262	1,732	3,741	4,950	6
2000.....	16,408	513	224	238	538	748	1,283	1,413	1,275	1,748	3,580	4,846	2
1999.....	17,430	457	239	284	562	781	1,331	1,400	1,250	1,935	3,943	5,245	3
Non-Hispanic 1/, male													
2003.....	9,122	356	145	139	432	544	939	1,110	980	1,043	1,726	1,704	4
2002.....	9,070	344	128	163	450	555	950	1,084	954	1,065	1,771	1,601	5
2001.....	9,280	347	133	167	439	568	990	1,154	876	1,040	1,858	1,702	6
2000.....	9,207	295	139	178	454	631	1,057	1,138	913	1,033	1,755	1,612	2
1999.....	9,694	279	146	201	476	655	1,079	1,094	891	1,162	1,933	1,775	3
Non-Hispanic 1/, female													
2003.....	6,933	246	80	47	94	115	256	361	355	635	1,696	3,047	1
2002.....	7,113	262	77	72	84	122	243	299	357	652	1,795	3,149	1
2001.....	7,415	241	98	63	103	101	241	359	386	692	1,883	3,248	-
2000.....	7,201	218	85	60	84	117	226	275	362	715	1,825	3,234	-
1999.....	7,736	178	93	83	86	126	252	306	359	773	2,010	3,470	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
Non-Hispanic white, both sexes													
2003.....	13,791	356	158	143	424	531	969	1,164	1,092	1,419	3,084	4,446	5
2002.....	13,865	339	136	172	446	565	969	1,131	1,081	1,414	3,190	4,416	6
2001.....	14,326	348	160	182	455	528	994	1,216	1,033	1,462	3,359	4,583	6
2000.....	14,083	310	156	185	425	604	1,039	1,123	1,044	1,464	3,224	4,509	-
1999.....	14,930	276	163	225	468	622	1,103	1,100	980	1,612	3,504	4,876	1
Non-Hispanic white, male													
2003.....	7,709	211	104	108	358	444	767	886	799	891	1,549	1,588	4
2002.....	7,669	203	83	120	382	466	786	898	786	882	1,580	1,478	5
2001.....	7,832	206	93	129	370	449	814	929	719	886	1,658	1,573	6
2000.....	7,703	186	99	141	361	515	860	899	736	868	1,555	1,483	-
1999.....	8,122	167	96	163	395	532	896	860	705	962	1,717	1,628	1
Non-Hispanic white, female													
2003.....	6,082	145	54	35	66	87	202	278	293	528	1,535	2,858	1
2002.....	6,196	136	53	52	64	99	183	233	295	532	1,610	2,938	1
2001.....	6,494	142	67	53	85	79	180	287	314	576	1,701	3,010	-
2000.....	6,380	124	57	44	64	89	179	224	308	596	1,669	3,026	-
1999.....	6,808	109	67	62	73	90	207	240	275	650	1,787	3,248	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic black, both sexes

2003.....	1,858	226	52	31	79	95	185	255	201	218	267	249	-
2002.....	1,951	250	59	55	68	90	179	214	189	257	311	279	-
2001.....	1,968	217	61	38	62	101	184	249	198	232	317	309	-
2000.....	1,968	183	55	48	90	118	198	237	200	244	307	286	2
1999.....	2,183	168	61	47	74	132	193	263	236	293	384	330	2

Non-Hispanic black, male

2003.....	1,147	134	36	22	56	74	138	186	146	130	138	87	-
2002.....	1,162	132	40	38	52	71	126	157	138	151	159	98	-
2001.....	1,180	131	35	31	47	81	137	185	134	134	165	100	-
2000.....	1,256	96	32	34	73	94	159	196	151	139	174	106	2
1999.....	1,365	101	42	32	67	102	153	203	165	182	185	131	2

Non-Hispanic black, female

2003.....	711	92	16	9	23	21	47	69	55	88	129	162	-
2002.....	789	118	19	17	16	19	53	57	51	106	152	181	-
2001.....	788	86	26	7	15	20	47	64	64	98	152	209	-
2000.....	712	87	23	14	17	24	39	41	49	105	133	180	-
1999.....	818	67	19	15	7	30	40	60	71	111	199	199	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)													
Origin not stated 2/, both sexes													
2003.....	61	2	-	-	4	4	8	6	3	5	9	10	10
2002.....	64	3	-	2	4	4	4	12	7	5	7	14	2
2001.....	80	3	1	2	3	4	9	6	10	6	11	11	14
2000.....	87	1	1	-	2	7	12	13	6	9	14	14	8
1999.....	80	1	1	1	2	5	1	14	13	7	10	16	9
Origin not stated 2/, male													
2003.....	48	2	-	-	4	3	8	6	2	3	7	4	9
2002.....	43	2	-	1	3	4	4	9	7	4	4	5	-
2001.....	60	3	1	2	3	4	8	5	8	3	9	3	11
2000.....	58	1	-	-	1	7	9	11	5	7	6	3	8
1999.....	57	-	-	1	1	3	1	12	12	5	6	8	8
Origin not stated 2/, female													
2003.....	13	-	-	-	-	1	-	-	1	2	2	6	1
2002.....	21	1	-	1	1	-	-	3	-	1	3	9	2
2001.....	20	-	-	-	-	-	1	1	2	3	2	8	3
2000.....	29	-	1	-	1	-	3	2	1	2	8	11	-
1999.....	23	1	1	-	1	2	-	2	1	2	4	8	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
All origins, both sexes													
2003.....	31,484	250	3,988	5,065	6,602	6,481	3,843	2,335	2,115	798	7
2002.....	31,655	264	4,010	5,046	6,851	6,308	3,618	2,463	2,259	826	10
2001 3/.....	30,622	279	3,971	5,070	6,635	5,942	3,317	2,432	2,192	769	15
2000.....	29,350	307	3,994	4,792	6,562	5,437	2,945	2,292	2,181	833	7
1999.....	29,199	244	3,901	5,106	6,466	5,081	2,896	2,471	2,217	801	16
All origins, male													
2003.....	25,203	194	3,381	4,156	5,138	4,887	2,987	1,954	1,809	690	7
2002.....	25,409	199	3,432	4,135	5,305	4,796	2,837	2,053	1,938	704	10
2001 3/.....	24,672	214	3,409	4,199	5,180	4,504	2,563	2,040	1,885	664	14
2000.....	23,618	244	3,424	3,938	5,114	4,149	2,265	1,888	1,884	705	7
1999.....	23,458	194	3,326	4,194	5,019	3,854	2,255	2,051	1,868	681	16
All origins, female													
2003.....	6,281	56	607	909	1,464	1,594	856	381	306	108	-
2002.....	6,246	65	578	911	1,546	1,512	781	410	321	122	-
2001 3/.....	5,950	65	562	871	1,455	1,438	754	392	307	105	1
2000.....	5,732	63	570	854	1,448	1,288	680	404	297	128	-
1999.....	5,741	50	575	912	1,447	1,227	641	420	349	120	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
Hispanic, both sexes													
2003.....	2,007	43	491	499	402	275	145	80	51	21	-
2002.....	1,954	37	455	484	407	260	142	89	55	23	2
2001 3/.....	1,850	37	414	490	397	239	114	86	49	22	2
2000.....	1,787	29	447	420	385	244	102	84	59	16	1
1999.....	1,695	30	391	423	369	210	112	73	63	22	2
Hispanic, male													
2003.....	1,711	31	421	434	341	220	129	71	47	17	-
2002.....	1,651	21	388	426	340	206	123	77	47	21	2
2001 3/.....	1,576	26	343	419	350	192	100	77	45	22	2
2000.....	1,525	22	387	370	320	194	89	74	53	15	1
1999.....	1,430	21	338	364	307	169	94	63	54	18	2
Hispanic, female													
2003.....	296	12	70	65	61	55	16	9	4	4	-
2002.....	303	16	67	58	67	54	19	12	8	2	-
2001 3/.....	274	11	71	71	47	47	14	9	4	-	-
2000.....	262	7	60	50	65	50	13	10	6	1	-
1999.....	265	9	53	59	62	41	18	10	9	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
Non-Hispanic 1/, both sexes													
2003.....	29,367	206	3,483	4,554	6,171	6,184	3,682	2,247	2,063	775	2
2002.....	29,543	226	3,536	4,538	6,413	6,015	3,455	2,360	2,199	797	4
2001 3/.....	28,647	242	3,545	4,560	6,213	5,682	3,184	2,334	2,138	744	5
2000.....	27,426	276	3,529	4,350	6,145	5,165	2,827	2,198	2,119	813	4
1999.....	27,379	212	3,500	4,665	6,068	4,842	2,773	2,388	2,146	776	9
Non-Hispanic 1/, male													
2003.....	23,406	162	2,950	3,712	4,775	4,651	2,844	1,878	1,761	671	2
2002.....	23,623	177	3,026	3,689	4,941	4,563	2,696	1,963	1,887	677	4
2001 3/.....	22,983	188	3,055	3,762	4,809	4,292	2,446	1,952	1,835	639	5
2000.....	21,981	221	3,023	3,553	4,767	3,931	2,161	1,805	1,828	688	4
1999.....	21,929	171	2,983	3,816	4,689	3,663	2,152	1,978	1,807	661	9
Non-Hispanic 1/, female													
2003.....	5,961	44	533	842	1,396	1,533	838	369	302	104	-
2002.....	5,920	49	510	849	1,472	1,452	759	397	312	120	-
2001 3/.....	5,664	54	490	798	1,404	1,390	738	382	303	105	-
2000.....	5,445	55	506	797	1,378	1,234	666	393	291	125	-
1999.....	5,450	41	517	849	1,379	1,179	621	410	339	115	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
Non-Hispanic white, both sexes													
2003.....	26,434	151	2,832	3,811	5,563	5,774	3,462	2,110	1,980	749	2
2002.....	26,691	177	2,951	3,809	5,795	5,603	3,254	2,228	2,105	765	4
2001 3/.....	25,813	183	2,926	3,871	5,604	5,292	2,979	2,187	2,049	718	4
2000.....	24,616	225	2,868	3,658	5,540	4,774	2,659	2,075	2,028	786	3
1999.....	24,534	174	2,830	3,927	5,469	4,487	2,580	2,249	2,058	754	6
Non-Hispanic white, male													
2003.....	21,085	122	2,412	3,099	4,296	4,357	2,678	1,769	1,696	654	2
2002.....	21,323	138	2,527	3,065	4,462	4,252	2,546	1,865	1,812	652	4
2001 3/.....	20,710	144	2,522	3,190	4,329	4,016	2,284	1,836	1,763	622	4
2000.....	19,711	179	2,467	2,989	4,287	3,627	2,031	1,701	1,761	666	3
1999.....	19,620	141	2,416	3,205	4,210	3,388	2,001	1,869	1,740	644	6
Non-Hispanic white, female													
2003.....	5,349	29	420	712	1,267	1,417	784	341	284	95	-
2002.....	5,368	39	424	744	1,333	1,351	708	363	293	113	-
2001 3/.....	5,103	39	404	681	1,275	1,276	695	351	286	96	-
2000.....	4,905	46	401	669	1,253	1,147	628	374	267	120	-
1999.....	4,914	33	414	722	1,259	1,099	579	380	318	110	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
Non-Hispanic black, both sexes													
2003.....	1,918	43	438	498	409	266	132	77	43	12	-
2002.....	1,896	35	392	497	432	265	125	80	51	19	-
2001 3/.....	1,904	40	425	472	418	268	126	90	52	13	-
2000.....	1,923	39	471	469	427	266	101	85	53	12	-
1999.....	1,924	29	463	497	430	226	116	94	52	15	2
Non-Hispanic black, male													
2003.....	1,566	34	377	424	324	199	102	59	39	8	-
2002.....	1,595	29	342	441	345	212	97	66	46	17	-
2001 3/.....	1,578	31	388	406	329	196	102	73	44	9	-
2000.....	1,601	34	407	397	346	210	80	76	40	11	-
1999.....	1,630	22	408	432	353	183	95	78	45	12	2
Non-Hispanic black, female													
2003.....	352	9	61	74	85	67	30	18	4	4	-
2002.....	301	6	50	56	87	53	28	14	5	2	-
2001 3/.....	326	9	37	66	89	72	24	17	8	4	-
2000.....	322	5	64	72	81	56	21	9	13	1	-
1999.....	294	7	55	65	77	43	21	16	7	3	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Origin not stated 2/, both sexes													
2003.....	110	1	14	12	29	22	16	8	1	2	5
2002.....	158	1	19	24	31	33	21	14	5	6	4
2001 3/.....	125	-	12	20	25	21	19	12	5	3	8
2000.....	137	2	18	22	32	28	16	10	3	4	2
1999.....	125	2	10	18	29	29	11	10	8	3	5
Origin not stated 2/, male													
2003.....	86	1	10	10	22	16	14	5	1	2	5
2002.....	135	1	18	20	24	27	18	13	4	6	4
2001 3/.....	113	-	11	18	21	20	17	11	5	3	7
2000.....	112	1	14	15	27	24	15	9	3	2	2
1999.....	99	2	5	14	23	22	9	10	7	2	5
Origin not stated 2/, female													
2003.....	24	-	4	2	7	6	2	3	-	-	-
2002.....	23	-	1	4	7	6	3	1	1	-	-
2001 3/.....	12	-	1	2	4	1	2	1	-	-	1
2000.....	25	1	4	7	5	4	1	1	-	2	-
1999.....	26	-	5	4	6	7	2	-	1	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
All origins, both sexes													
2003.....	16,907	74	2,075	2,381	2,927	3,279	2,317	1,700	1,595	559	-
2002.....	17,108	86	2,088	2,399	3,157	3,136	2,234	1,776	1,680	550	2
2001.....	16,869	90	2,130	2,564	3,030	3,023	2,083	1,758	1,648	537	6
2000.....	16,586	110	2,267	2,453	3,191	2,858	1,833	1,700	1,611	558	5
1999.....	16,599	103	2,315	2,621	3,097	2,677	1,860	1,791	1,606	524	5
All origins, male													
2003.....	14,827	63	1,879	2,099	2,470	2,768	2,001	1,538	1,480	529	-
2002.....	15,045	68	1,897	2,078	2,672	2,677	1,930	1,620	1,576	525	2
2001.....	14,758	69	1,935	2,246	2,551	2,551	1,771	1,592	1,531	506	6
2000.....	14,454	90	2,043	2,123	2,657	2,409	1,568	1,528	1,505	526	5
1999.....	14,479	80	2,083	2,270	2,586	2,251	1,594	1,615	1,499	496	5
All origins, female													
2003.....	2,080	11	196	282	457	511	316	162	115	30	-
2002.....	2,063	18	191	321	485	459	304	156	104	25	-
2001.....	2,111	21	195	318	479	472	312	166	117	31	-
2000.....	2,132	20	224	330	534	449	265	172	106	32	-
1999.....	2,120	23	232	351	511	426	266	176	107	28	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
Hispanic, both sexes													
2003.....	835	9	228	208	145	113	57	38	26	11	-
2002.....	834	10	224	198	145	98	67	52	28	12	-
2001.....	798	5	193	224	135	102	59	45	26	8	1
2000.....	813	7	224	181	144	124	47	49	27	9	1
1999.....	794	13	219	182	148	96	48	37	39	11	1
Hispanic, male													
2003.....	774	7	210	194	133	102	55	38	25	10	-
2002.....	763	5	208	183	128	88	62	49	28	12	-
2001.....	731	4	172	203	126	90	57	44	26	8	1
2000.....	737	7	205	165	126	106	46	46	26	9	1
1999.....	705	7	196	164	127	81	45	35	38	11	1
Hispanic, female													
2003.....	61	2	18	14	12	11	2	-	1	1	-
2002.....	71	5	16	15	17	10	5	3	-	-	-
2001.....	67	1	21	21	9	12	2	1	-	-	-
2000.....	76	-	19	16	18	18	1	3	1	-	-
1999.....	89	6	23	18	21	15	3	2	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
Non-Hispanic 1/, both sexes													
2003.....	16,035	65	1,843	2,171	2,772	3,158	2,254	1,657	1,569	546	-
2002.....	16,213	76	1,855	2,195	3,002	3,024	2,159	1,716	1,649	536	1
2001.....	16,020	85	1,933	2,335	2,887	2,916	2,011	1,706	1,617	526	4
2000.....	15,717	103	2,035	2,264	3,036	2,722	1,779	1,645	1,583	547	3
1999.....	15,755	90	2,092	2,431	2,937	2,573	1,806	1,749	1,563	511	3
Non-Hispanic 1/, male													
2003.....	14,025	56	1,667	1,903	2,330	2,660	1,940	1,497	1,455	517	-
2002.....	14,226	63	1,680	1,890	2,536	2,576	1,861	1,563	1,545	511	1
2001.....	13,978	65	1,759	2,039	2,417	2,457	1,701	1,541	1,500	495	4
2000.....	13,664	83	1,831	1,951	2,520	2,291	1,515	1,476	1,478	516	3
1999.....	13,732	73	1,884	2,101	2,448	2,164	1,543	1,575	1,457	484	3
Non-Hispanic 1/, female													
2003.....	2,010	9	176	268	442	498	314	160	114	29	-
2002.....	1,987	13	175	305	466	448	298	153	104	25	-
2001.....	2,042	20	174	296	470	459	310	165	117	31	-
2000.....	2,053	20	204	313	516	431	264	169	105	31	-
1999.....	2,023	17	208	330	489	409	263	174	106	27	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
Non-Hispanic white, both sexes													
2003.....	14,737	56	1,503	1,813	2,564	2,996	2,158	1,589	1,520	538	-
2002.....	14,865	63	1,521	1,853	2,749	2,846	2,063	1,653	1,595	521	1
2001.....	14,648	69	1,565	2,001	2,629	2,760	1,919	1,628	1,560	514	3
2000.....	14,333	96	1,655	1,916	2,780	2,552	1,679	1,572	1,545	536	2
1999.....	14,316	81	1,684	2,071	2,676	2,413	1,703	1,672	1,514	499	3
Non-Hispanic white, male													
2003.....	12,865	49	1,361	1,573	2,145	2,521	1,857	1,438	1,411	510	-
2002.....	13,014	53	1,375	1,573	2,319	2,423	1,774	1,507	1,493	496	1
2001.....	12,771	55	1,414	1,739	2,199	2,333	1,619	1,474	1,451	484	3
2000.....	12,448	79	1,489	1,648	2,296	2,151	1,427	1,405	1,446	505	2
1999.....	12,441	65	1,514	1,776	2,213	2,029	1,451	1,505	1,412	473	3
Non-Hispanic white, female													
2003.....	1,872	7	142	240	419	475	301	151	109	28	-
2002.....	1,851	10	146	280	430	423	289	146	102	25	-
2001.....	1,877	14	151	262	430	427	300	154	109	30	-
2000.....	1,885	17	166	268	484	401	252	167	99	31	-
1999.....	1,875	16	170	295	463	384	252	167	102	26	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
Non-Hispanic black, both sexes													
2003.....	993	7	264	279	150	117	75	56	38	7	-
2002.....	1,041	9	250	272	199	130	71	54	41	15	-
2001.....	1,069	12	293	258	198	122	67	62	47	10	-
2000.....	1,073	4	304	268	193	131	71	62	32	8	-
1999.....	1,112	5	323	268	197	118	81	65	43	12	-
Non-Hispanic black, male													
2003.....	892	5	243	260	131	103	63	47	34	6	-
2002.....	945	8	229	258	170	115	63	48	39	15	-
2001.....	949	9	277	233	169	98	61	53	40	9	-
2000.....	949	4	277	237	168	107	61	60	27	8	-
1999.....	1,007	4	294	247	178	102	73	59	39	11	-
Non-Hispanic black, female													
2003.....	101	2	21	19	19	14	12	9	4	1	-
2002.....	96	1	21	14	29	15	8	6	2	-	-
2001.....	120	3	16	25	29	24	6	9	7	1	-
2000.....	124	-	27	31	25	24	10	2	5	-	-
1999.....	105	1	29	21	19	16	8	6	4	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Origin not stated 2/, both sexes													
2003.....	37	-	4	2	10	8	6	5	-	2	-
2002.....	61	-	9	6	10	14	8	8	3	2	1
2001.....	51	-	4	5	8	5	13	7	5	3	1
2000.....	56	-	8	8	11	12	7	6	1	2	1
1999.....	50	-	4	8	12	8	6	5	4	2	1
Origin not stated 2/, male													
2003.....	28	-	2	2	7	6	6	3	-	2	-
2002.....	56	-	9	5	8	13	7	8	3	2	1
2001.....	49	-	4	4	8	4	13	7	5	3	1
2000.....	53	-	7	7	11	12	7	6	1	1	1
1999.....	42	-	3	5	11	6	6	5	4	1	1
Origin not stated 2/, female													
2003.....	9	-	2	-	3	2	-	2	-	-	-
2002.....	5	-	-	1	2	1	1	-	-	-	-
2001.....	2	-	-	1	-	1	-	-	-	-	-
2000.....	3	-	1	1	-	-	-	-	-	1	-
1999.....	8	-	1	3	1	2	-	-	-	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)													
All origins, both sexes													
2003.....	14,577	176	1,913	2,684	3,675	3,202	1,526	635	520	239	7
2002.....	14,547	178	1,922	2,647	3,694	3,172	1,384	687	579	276	8
2001 3/.....	13,753	189	1,841	2,506	3,605	2,919	1,234	674	544	232	9
2000.....	12,764	197	1,727	2,339	3,371	2,579	1,112	592	570	275	2
1999.....	12,600	141	1,586	2,485	3,369	2,404	1,036	680	611	277	11
All origins, male													
2003.....	10,376	131	1,502	2,057	2,668	2,119	986	416	329	161	7
2002.....	10,364	131	1,535	2,057	2,633	2,119	907	433	362	179	8
2001 3/.....	9,914	145	1,474	1,953	2,629	1,953	792	448	354	158	8
2000.....	9,164	154	1,381	1,815	2,457	1,740	697	360	379	179	2
1999.....	8,979	114	1,243	1,924	2,433	1,603	661	436	369	185	11
All origins, female													
2003.....	4,201	45	411	627	1,007	1,083	540	219	191	78	-
2002.....	4,183	47	387	590	1,061	1,053	477	254	217	97	-
2001 3/.....	3,839	44	367	553	976	966	442	226	190	74	1
2000.....	3,600	43	346	524	914	839	415	232	191	96	-
1999.....	3,621	27	343	561	936	801	375	244	242	92	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)													
Hispanic, both sexes													
2003.....	1,172	34	263	291	257	162	88	42	25	10	-
2002.....	1,120	27	231	286	262	162	75	37	27	11	2
2001 3/.....	1,052	32	221	266	262	137	55	41	23	14	1
2000.....	974	22	223	239	241	120	55	35	32	7	-
1999.....	901	17	172	241	221	114	64	36	24	11	1
Hispanic, male													
2003.....	937	24	211	240	208	118	74	33	22	7	-
2002.....	888	16	180	243	212	118	61	28	19	9	2
2001 3/.....	845	22	171	216	224	102	43	33	19	14	1
2000.....	788	15	182	205	194	88	43	28	27	6	-
1999.....	725	14	142	200	180	88	49	28	16	7	1
Hispanic, female													
2003.....	235	10	52	51	49	44	14	9	3	3	-
2002.....	232	11	51	43	50	44	14	9	8	2	-
2001 3/.....	207	10	50	50	38	35	12	8	4	-	-
2000.....	186	7	41	34	47	32	12	7	5	1	-
1999.....	176	3	30	41	41	26	15	8	8	4	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)													
Non-Hispanic 1/, both sexes													
2003.....	13,332	141	1,640	2,383	3,399	3,026	1,428	590	494	229	2
2002.....	13,330	150	1,681	2,343	3,411	2,991	1,296	644	550	261	3
2001 3/.....	12,627	157	1,612	2,225	3,326	2,766	1,173	628	521	218	1
2000.....	11,709	173	1,494	2,086	3,109	2,443	1,048	553	536	266	1
1999.....	11,624	122	1,408	2,234	3,131	2,269	967	639	583	265	6
Non-Hispanic 1/, male													
2003.....	9,381	106	1,283	1,809	2,445	1,991	904	381	306	154	2
2002.....	9,397	114	1,346	1,799	2,405	1,987	835	400	342	166	3
2001 3/.....	9,005	123	1,296	1,723	2,392	1,835	745	411	335	144	1
2000.....	8,317	138	1,192	1,602	2,247	1,640	646	329	350	172	1
1999.....	8,197	98	1,099	1,715	2,241	1,499	609	403	350	177	6
Non-Hispanic 1/, female													
2003.....	3,951	35	357	574	954	1,035	524	209	188	75	-
2002.....	3,933	36	335	544	1,006	1,004	461	244	208	95	-
2001 3/.....	3,622	34	316	502	934	931	428	217	186	74	-
2000.....	3,392	35	302	484	862	803	402	224	186	94	-
1999.....	3,427	24	309	519	890	770	358	236	233	88	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)													
Non-Hispanic white, both sexes													
2003.....	11,697	95	1,329	1,998	2,999	2,778	1,304	521	460	211	2
2002.....	11,826	114	1,430	1,956	3,046	2,757	1,191	575	510	244	3
2001 3/.....	11,165	114	1,361	1,870	2,975	2,532	1,060	559	489	204	1
2000.....	10,283	129	1,213	1,742	2,760	2,222	980	503	483	250	1
1999.....	10,218	93	1,146	1,856	2,793	2,074	877	577	544	255	3
Non-Hispanic white, male													
2003.....	8,220	73	1,051	1,526	2,151	1,836	821	331	285	144	2
2002.....	8,309	85	1,152	1,492	2,143	1,829	772	358	319	156	3
2001 3/.....	7,939	89	1,108	1,451	2,130	1,683	665	362	312	138	1
2000.....	7,263	100	978	1,341	1,991	1,476	604	296	315	161	1
1999.....	7,179	76	902	1,429	1,997	1,359	550	364	328	171	3
Non-Hispanic white, female													
2003.....	3,477	22	278	472	848	942	483	190	175	67	-
2002.....	3,517	29	278	464	903	928	419	217	191	88	-
2001 3/.....	3,226	25	253	419	845	849	395	197	177	66	-
2000.....	3,020	29	235	401	769	746	376	207	168	89	-
1999.....	3,039	17	244	427	796	715	327	213	216	84	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)													
Non-Hispanic black, both sexes													
2003.....	925	36	174	219	259	149	57	21	5	5	-
2002.....	855	26	142	225	233	135	54	26	10	4	-
2001 3/.....	835	28	132	214	220	146	59	28	5	3	-
2000.....	850	35	167	201	234	135	30	23	21	4	-
1999.....	812	24	140	229	233	108	35	29	9	3	2
Non-Hispanic black, male													
2003.....	674	29	134	164	193	96	39	12	5	2	-
2002.....	650	21	113	183	175	97	34	18	7	2	-
2001 3/.....	629	22	111	173	160	98	41	20	4	-	-
2000.....	652	30	130	160	178	103	19	16	13	3	-
1999.....	623	18	114	185	175	81	22	19	6	1	2
Non-Hispanic black, female													
2003.....	251	7	40	55	66	53	18	9	-	3	-
2002.....	205	5	29	42	58	38	20	8	3	2	-
2001 3/.....	206	6	21	41	60	48	18	8	1	3	-
2000.....	198	5	37	41	56	32	11	7	8	1	-
1999.....	189	6	26	44	58	27	13	10	3	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Origin not stated 2/, both sexes

2003.....	73	1	10	10	19	14	10	3	1	-	5
2002.....	97	1	10	18	21	19	13	6	2	4	3
2001 3/.....	74	-	8	15	17	16	6	5	-	-	7
2000.....	81	2	10	14	21	16	9	4	2	2	1
1999.....	75	2	6	10	17	21	5	5	4	1	4

Origin not stated 2/, male

2003.....	58	1	8	8	15	10	8	2	1	-	5
2002.....	79	1	9	15	16	14	11	5	1	4	3
2001 3/.....	64	-	7	14	13	16	4	4	-	-	6
2000.....	59	1	7	8	16	12	8	3	2	1	1
1999.....	57	2	2	9	12	16	3	5	3	1	4

Origin not stated 2/, female

2003.....	15	-	2	2	4	4	2	1	-	-	-
2002.....	18	-	1	3	5	5	2	1	1	-	-
2001 3/.....	10	-	1	1	4	-	2	1	-	-	1
2000.....	22	1	3	6	5	4	1	1	-	1	-
1999.....	18	-	4	1	5	5	2	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
All origins, both sexes													
2003.....	17,732	341	376	324	5,368	4,516	3,110	2,017	786	432	317	104	41
2002.....	17,638	303	423	356	5,219	4,489	3,239	1,915	841	421	296	95	41
2001 3/.....	20,308	332	415	326	5,297	5,204	4,268	2,467	1,018	532	312	105	32
2000.....	16,765	349	356	371	4,939	4,164	3,219	1,755	738	443	301	100	30
1999.....	16,889	331	376	432	4,998	4,231	3,206	1,668	722	473	304	99	49
All origins, male													
2003.....	13,882	205	202	215	4,624	3,741	2,308	1,488	568	285	171	46	29
2002.....	13,640	164	234	197	4,474	3,669	2,386	1,394	614	277	158	41	32
2001 3/.....	15,555	197	238	193	4,541	4,157	3,122	1,832	723	332	155	42	23
2000.....	12,820	203	198	227	4,203	3,361	2,319	1,284	537	279	154	40	15
1999.....	12,785	187	194	250	4,191	3,346	2,307	1,229	541	304	162	41	33
All origins, female													
2003.....	3,850	136	174	109	744	775	802	529	218	147	146	58	12
2002.....	3,998	139	189	159	745	820	853	521	227	144	138	54	9
2001 3/.....	4,753	135	177	133	756	1,047	1,146	635	295	200	157	63	9
2000.....	3,945	146	158	144	736	803	900	471	201	164	147	60	15
1999.....	4,104	144	182	182	807	885	899	439	181	169	142	58	16

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Hispanic, both sexes													
2003.....	3,355	60	82	53	1,286	971	502	246	75	46	23	5	6
2002.....	3,129	52	81	64	1,200	903	464	216	87	29	19	3	11
2001 3/.....	3,331	55	69	56	1,222	960	567	248	82	45	16	3	8
2000.....	2,917	54	57	76	1,126	810	447	211	77	36	16	4	3
1999.....	2,864	53	52	89	1,119	801	432	197	69	33	11	3	5
Hispanic, male													
2003.....	2,809	35	36	36	1,139	852	401	194	59	37	13	2	5
2002.....	2,635	28	49	37	1,081	789	367	166	72	25	9	2	10
2001 3/.....	2,756	33	38	38	1,099	800	451	190	59	33	8	2	5
2000.....	2,431	26	30	56	1,014	697	359	158	57	22	8	2	2
1999.....	2,399	26	25	59	990	698	354	150	56	26	8	2	5
Hispanic, female													
2003.....	546	25	46	17	147	119	101	52	16	9	10	3	1
2002.....	494	24	32	27	119	114	97	50	15	4	10	1	1
2001 3/.....	575	22	31	18	123	160	116	58	23	12	8	1	3
2000.....	486	28	27	20	112	113	88	53	20	14	8	2	1
1999.....	465	27	27	30	129	103	78	47	13	7	3	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Non-Hispanic 1/, both sexes													
2003.....	14,245	278	293	271	4,045	3,520	2,588	1,757	705	381	291	97	19
2002.....	14,346	246	340	289	3,983	3,548	2,746	1,682	747	388	273	92	12
2001 3/.....	16,726	270	343	269	4,028	4,178	3,652	2,182	916	480	296	101	11
2000.....	13,717	291	299	293	3,776	3,337	2,749	1,526	652	402	285	95	12
1999.....	13,873	270	323	341	3,860	3,402	2,743	1,452	642	437	292	95	16
Non-Hispanic 1/, male													
2003.....	10,969	168	166	179	3,455	2,868	1,894	1,282	503	243	156	42	13
2002.....	10,881	131	184	157	3,362	2,852	2,002	1,213	536	248	148	39	9
2001 3/.....	12,608	161	199	154	3,401	3,309	2,632	1,612	651	294	147	39	9
2000.....	10,290	174	168	170	3,159	2,650	1,939	1,115	473	254	146	37	5
1999.....	10,275	158	168	191	3,188	2,624	1,930	1,063	476	275	153	39	10
Non-Hispanic 1/, female													
2003.....	3,276	110	127	92	590	652	694	475	202	138	135	55	6
2002.....	3,465	115	156	132	621	696	744	469	211	140	125	53	3
2001 3/.....	4,118	109	144	115	627	869	1,020	570	265	186	149	62	2
2000.....	3,427	117	131	123	617	687	810	411	179	148	139	58	7
1999.....	3,598	112	155	150	672	778	813	389	166	162	139	56	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Non-Hispanic white, both sexes													
2003.....	5,367	156	123	109	907	953	1,147	947	455	254	220	80	16
2002.....	5,571	117	169	133	958	977	1,268	898	496	275	197	73	10
2001 3/.....	7,928	139	160	164	1,058	1,642	2,081	1,373	652	342	225	85	7
2000.....	5,429	141	144	164	874	1,023	1,273	834	415	264	211	78	8
1999.....	5,748	143	156	176	962	1,135	1,327	814	418	319	214	76	8
Non-Hispanic white, male													
2003.....	3,550	94	61	72	665	648	747	645	303	153	115	36	11
2002.....	3,661	70	89	61	685	660	874	586	332	166	99	32	7
2001 3/.....	5,453	84	95	83	729	1,177	1,498	990	454	197	110	31	5
2000.....	3,510	96	86	93	588	677	836	560	288	153	101	28	4
1999.....	3,747	94	83	93	644	753	881	579	293	188	109	27	3
Non-Hispanic white, female													
2003.....	1,817	62	62	37	242	305	400	302	152	101	105	44	5
2002.....	1,910	47	80	72	273	317	394	312	164	109	98	41	3
2001 3/.....	2,475	55	65	81	329	465	583	383	198	145	115	54	2
2000.....	1,919	45	58	71	286	346	437	274	127	111	110	50	4
1999.....	2,001	49	73	83	318	382	446	235	125	131	105	49	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Non-Hispanic black, both sexes													
2003.....	8,276	106	155	147	2,955	2,443	1,335	731	218	105	63	16	2
2002.....	8,147	114	156	132	2,854	2,425	1,357	710	216	95	69	17	2
2001 3/.....	8,077	119	166	92	2,817	2,335	1,427	704	219	117	63	14	4
2000.....	7,748	139	143	122	2,771	2,193	1,359	615	201	123	64	14	4
1999.....	7,536	114	151	145	2,752	2,122	1,305	557	194	103	67	18	8
Non-Hispanic black, male													
2003.....	6,980	60	96	98	2,642	2,128	1,078	576	178	78	39	6	1
2002.....	6,780	54	89	84	2,540	2,092	1,042	572	180	72	47	6	2
2001 3/.....	6,658	68	100	66	2,551	1,998	1,036	544	165	85	34	7	4
2000.....	6,382	73	73	70	2,459	1,883	1,027	498	159	93	39	7	1
1999.....	6,131	58	74	89	2,433	1,771	980	431	159	79	39	11	7
Non-Hispanic black, female													
2003.....	1,296	46	59	49	313	315	257	155	40	27	24	10	1
2002.....	1,367	60	67	48	314	333	315	138	36	23	22	11	-
2001 3/.....	1,419	51	66	26	266	337	391	160	54	32	29	7	-
2000.....	1,366	66	70	52	312	310	332	117	42	30	25	7	3
1999.....	1,405	56	77	56	319	351	325	126	35	24	28	7	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Origin not stated 2/, both sexes													
2003.....	132	3	1	-	37	25	20	14	6	5	3	2	16
2002.....	163	5	2	3	36	38	29	17	7	4	4	-	18
2001 3/.....	251	7	3	1	47	66	49	37	20	7	-	1	13
2000.....	131	4	-	2	37	17	23	18	9	5	-	1	15
1999.....	152	8	1	2	19	28	31	19	11	3	1	1	28
Origin not stated 2/, male													
2003.....	104	2	-	-	30	21	13	12	6	5	2	2	11
2002.....	124	5	1	3	31	28	17	15	6	4	1	-	13
2001 3/.....	191	3	1	1	41	48	39	30	13	5	-	1	9
2000.....	99	3	-	1	30	14	21	11	7	3	-	1	8
1999.....	111	3	1	-	13	24	23	16	9	3	1	-	18
Origin not stated 2/, female													
2003.....	28	1	1	-	7	4	7	2	-	-	1	-	5
2002.....	39	-	1	-	5	10	12	2	1	-	3	-	5
2001 3/.....	60	4	2	-	6	18	10	7	7	2	-	-	4
2000.....	32	1	-	1	7	3	2	7	2	2	-	-	7
1999.....	41	5	-	2	6	4	8	3	2	-	-	1	10

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
All origins, both sexes													
2003.....	11,920	8	40	187	4,410	3,540	1,941	1,110	394	161	88	23	18
2002.....	11,829	9	49	205	4,317	3,465	2,042	1,025	409	161	115	19	13
2001.....	11,348	11	55	180	4,200	3,308	1,978	934	364	192	99	16	11
2000.....	10,801	12	28	187	3,963	3,074	1,906	959	367	184	81	28	12
1999.....	10,828	8	50	224	4,038	3,048	1,868	893	373	193	95	23	15
All origins, male													
2003.....	10,126	7	20	134	3,977	3,125	1,535	860	292	115	35	10	16
2002.....	9,899	3	30	127	3,851	3,017	1,623	761	301	101	64	9	12
2001.....	9,532	7	27	125	3,796	2,864	1,528	726	264	123	54	7	11
2000.....	9,006	9	17	138	3,548	2,650	1,487	730	266	108	37	9	7
1999.....	8,944	5	26	155	3,569	2,609	1,414	676	294	124	48	10	14
All origins, female													
2003.....	1,794	1	20	53	433	415	406	250	102	46	53	13	2
2002.....	1,930	6	19	78	466	448	419	264	108	60	51	10	1
2001.....	1,816	4	28	55	404	444	450	208	100	69	45	9	-
2000.....	1,795	3	11	49	415	424	419	229	101	76	44	19	5
1999.....	1,884	3	24	69	469	439	454	217	79	69	47	13	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Hispanic, both sexes													
2003.....	2,316	1	9	31	1,044	730	313	138	26	17	3	-	4
2002.....	2,168	2	12	40	976	679	294	116	31	9	4	1	4
2001.....	2,123	4	12	30	974	654	306	102	20	14	3	1	3
2000.....	1,958	1	5	41	904	581	249	126	34	14	1	-	2
1999.....	1,939	2	5	52	908	550	258	112	35	9	5	1	2
Hispanic, male													
2003.....	2,069	1	5	22	957	673	259	112	22	13	2	-	3
2002.....	1,942	-	8	29	909	619	245	91	26	7	3	1	4
2001.....	1,892	3	5	23	898	596	257	81	15	10	-	1	3
2000.....	1,738	-	3	34	837	524	207	99	26	7	-	-	1
1999.....	1,709	2	3	39	824	499	218	84	27	6	4	1	2
Hispanic, female													
2003.....	247	-	4	9	87	57	54	26	4	4	1	-	1
2002.....	226	2	4	11	67	60	49	25	5	2	1	-	-
2001.....	231	1	7	7	76	58	49	21	5	4	3	-	-
2000.....	220	1	2	7	67	57	42	27	8	7	1	-	1
1999.....	230	-	2	13	84	51	40	28	8	3	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Non-Hispanic 1/, both sexes													
2003.....	9,536	7	31	156	3,339	2,792	1,616	970	365	144	84	23	9
2002.....	9,575	7	37	164	3,309	2,760	1,735	902	376	151	110	18	6
2001.....	9,134	7	43	149	3,186	2,631	1,662	827	341	175	96	14	3
2000.....	8,767	11	23	144	3,032	2,480	1,640	828	330	168	80	28	3
1999.....	8,821	6	45	172	3,115	2,476	1,597	776	335	184	89	22	4
Non-Hispanic 1/, male													
2003.....	7,997	6	15	112	2,996	2,436	1,267	746	267	102	32	10	8
2002.....	7,884	3	22	97	2,913	2,378	1,367	664	274	93	60	8	5
2001.....	7,559	4	22	101	2,863	2,246	1,264	641	246	110	54	5	3
2000.....	7,205	9	14	103	2,686	2,115	1,264	628	237	101	37	9	2
1999.....	7,177	3	23	116	2,734	2,091	1,184	588	264	118	43	9	4
Non-Hispanic 1/, female													
2003.....	1,539	1	16	44	343	356	349	224	98	42	52	13	1
2002.....	1,691	4	15	67	396	382	368	238	102	58	50	10	1
2001.....	1,575	3	21	48	323	385	398	186	95	65	42	9	-
2000.....	1,562	2	9	41	346	365	376	200	93	67	43	19	1
1999.....	1,644	3	22	56	381	385	413	188	71	66	46	13	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Non-Hispanic white, both sexes													
2003.....	2,883	1	11	51	610	613	649	516	242	99	65	20	6
2002.....	3,052	1	22	66	642	636	735	483	253	107	87	16	4
2001.....	3,085	4	21	90	618	681	759	460	235	126	75	14	2
2000.....	2,861	4	12	72	544	624	729	445	224	118	63	24	2
1999.....	2,995	2	21	80	591	694	749	416	215	140	67	20	-
Non-Hispanic white, male													
2003.....	2,068	-	3	37	489	456	445	366	166	68	23	9	6
2002.....	2,139	1	12	34	488	465	531	321	174	58	44	8	3
2001.....	2,172	2	12	56	468	495	536	335	155	69	37	5	2
2000.....	1,967	2	6	47	401	444	510	311	149	63	25	8	1
1999.....	2,119	2	11	50	443	515	514	308	157	84	27	8	-
Non-Hispanic white, female													
2003.....	815	1	8	14	121	157	204	150	76	31	42	11	-
2002.....	913	-	10	32	154	171	204	162	79	49	43	8	1
2001.....	913	2	9	34	150	186	223	125	80	57	38	9	-
2000.....	894	2	6	25	143	180	219	134	75	55	38	16	1
1999.....	876	-	10	30	148	179	235	108	58	56	40	12	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Non-Hispanic black, both sexes													
2003.....	6,319	6	19	95	2,606	2,100	909	415	106	42	16	3	2
2002.....	6,181	6	14	80	2,547	2,048	931	385	107	39	20	2	2
2001.....	5,790	2	18	56	2,490	1,874	864	332	92	43	18	-	1
2000.....	5,622	7	8	68	2,396	1,790	854	347	85	45	17	4	1
1999.....	5,508	2	18	81	2,426	1,695	796	320	104	39	21	2	4
Non-Hispanic black, male													
2003.....	5,666	6	11	67	2,404	1,914	780	351	89	33	9	1	1
2002.....	5,482	2	9	54	2,325	1,854	784	317	88	32	15	-	2
2001.....	5,190	1	9	43	2,329	1,697	700	280	78	37	15	-	1
2000.....	5,011	7	5	52	2,206	1,616	715	290	69	37	12	1	1
1999.....	4,826	1	7	60	2,208	1,512	636	256	94	31	16	1	4
Non-Hispanic black, female													
2003.....	653	-	8	28	202	186	129	64	17	9	7	2	1
2002.....	699	4	5	26	222	194	147	68	19	7	5	2	-
2001.....	600	1	9	13	161	177	164	52	14	6	3	-	-
2000.....	611	-	3	16	190	174	139	57	16	8	5	3	-
1999.....	682	1	11	21	218	183	160	64	10	8	5	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Origin not stated 2/, both sexes													
2003.....	68	-	-	-	27	18	12	2	3	-	1	-	5
2002.....	86	-	-	1	32	26	13	7	2	1	1	-	3
2001.....	91	-	-	1	40	23	10	5	3	3	-	1	5
2000.....	76	-	-	2	27	13	17	5	3	2	-	-	7
1999.....	68	-	-	-	15	22	13	5	3	-	1	-	9
Origin not stated 2/, male													
2003.....	60	-	-	-	24	16	9	2	3	-	1	-	5
2002.....	73	-	-	1	29	20	11	6	1	1	1	-	3
2001.....	81	-	-	1	35	22	7	4	3	3	-	1	5
2000.....	63	-	-	1	25	11	16	3	3	-	-	-	4
1999.....	58	-	-	-	11	19	12	4	3	-	1	-	8
Origin not stated 2/, female													
2003.....	8	-	-	-	3	2	3	-	-	-	-	-	-
2002.....	13	-	-	-	3	6	2	1	1	-	-	-	-
2001.....	10	-	-	-	5	1	3	1	-	-	-	-	-
2000.....	13	-	-	1	2	2	1	2	-	2	-	-	3
1999.....	10	-	-	-	4	3	1	1	-	-	-	-	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All origins, both sexes													
2003.....	5,812	333	336	137	958	976	1,169	907	392	271	229	81	23
2002.....	5,809	294	374	151	902	1,024	1,197	890	432	260	181	76	28
2001 3/.....	8,960	321	360	146	1,097	1,896	2,290	1,533	654	340	213	89	21
2000.....	5,964	337	328	184	976	1,090	1,313	796	371	259	220	72	18
1999.....	6,061	323	326	208	960	1,183	1,338	775	349	280	209	76	34
All origins, male													
2003.....	3,756	198	182	81	647	616	773	628	276	170	136	36	13
2002.....	3,741	161	204	70	623	652	763	633	313	176	94	32	20
2001 3/.....	6,023	190	211	68	745	1,293	1,594	1,106	459	209	101	35	12
2000.....	3,814	194	181	89	655	711	832	554	271	171	117	31	8
1999.....	3,841	182	168	95	622	737	893	553	247	180	114	31	19
All origins, female													
2003.....	2,056	135	154	56	311	360	396	279	116	101	93	45	10
2002.....	2,068	133	170	81	279	372	434	257	119	84	87	44	8
2001 3/.....	2,937	131	149	78	352	603	696	427	195	131	112	54	9
2000.....	2,150	143	147	95	321	379	481	242	100	88	103	41	10
1999.....	2,220	141	158	113	338	446	445	222	102	100	95	45	15

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)													
Hispanic, both sexes													
2003.....	1,039	59	73	22	242	241	189	108	49	29	20	5	2
2002.....	961	50	69	24	224	224	170	100	56	20	15	2	7
2001 3/.....	1,208	51	57	26	248	306	261	146	62	31	13	2	5
2000.....	959	53	52	35	222	229	198	85	43	22	15	4	1
1999.....	925	51	47	37	211	251	174	85	34	24	6	2	3
Hispanic, male													
2003.....	740	34	31	14	182	179	142	82	37	24	11	2	2
2002.....	693	28	41	8	172	170	122	75	46	18	6	1	6
2001 3/.....	864	30	33	15	201	204	194	109	44	23	8	1	2
2000.....	693	26	27	22	177	173	152	59	31	15	8	2	1
1999.....	690	24	22	20	166	199	136	66	29	20	4	1	3
Hispanic, female													
2003.....	299	25	42	8	60	62	47	26	12	5	9	3	-
2002.....	268	22	28	16	52	54	48	25	10	2	9	1	1
2001 3/.....	344	21	24	11	47	102	67	37	18	8	5	1	3
2000.....	266	27	25	13	45	56	46	26	12	7	7	2	-
1999.....	235	27	25	17	45	52	38	19	5	4	2	1	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Non-Hispanic 1/, both sexes

2003.....	4,709	271	262	115	706	728	972	787	340	237	207	74	10
2002.....	4,771	239	303	125	674	788	1,011	780	371	237	163	74	6
2001 3/.....	7,592	263	300	120	842	1,547	1,990	1,355	575	305	200	87	8
2000.....	4,950	280	276	149	744	857	1,109	698	322	234	205	67	9
1999.....	5,052	264	278	169	745	926	1,146	676	307	253	203	73	12

Non-Hispanic 1/, male

2003.....	2,972	162	151	67	459	432	627	536	236	141	124	32	5
2002.....	2,997	128	162	60	449	474	635	549	262	155	88	31	4
2001 3/.....	5,049	157	177	53	538	1,063	1,368	971	405	184	93	34	6
2000.....	3,085	165	154	67	473	535	675	487	236	153	109	28	3
1999.....	3,098	155	145	75	454	533	746	475	212	157	110	30	6

Non-Hispanic 1/, female

2003.....	1,737	109	111	48	247	296	345	251	104	96	83	42	5
2002.....	1,774	111	141	65	225	314	376	231	109	82	75	43	2
2001 3/.....	2,543	106	123	67	304	484	622	384	170	121	107	53	2
2000.....	1,865	115	122	82	271	322	434	211	86	81	96	39	6
1999.....	1,954	109	133	94	291	393	400	201	95	96	93	43	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Non-Hispanic white, both sexes

2003.....	2,484	155	112	58	297	340	498	431	213	155	155	60	10
2002.....	2,519	116	147	67	316	341	533	415	243	168	110	57	6
2001 3/.....	4,843	135	139	74	440	961	1,322	913	417	216	150	71	5
2000.....	2,568	137	132	92	330	399	544	389	191	146	148	54	6
1999.....	2,753	141	135	96	371	441	578	398	203	179	147	56	8

Non-Hispanic white, male

2003.....	1,482	94	58	35	176	192	302	279	137	85	92	27	5
2002.....	1,522	69	77	27	197	195	343	265	158	108	55	24	4
2001 3/.....	3,281	82	83	27	261	682	962	655	299	128	73	26	3
2000.....	1,543	94	80	46	187	233	326	249	139	90	76	20	3
1999.....	1,628	92	72	43	201	238	367	271	136	104	82	19	3

Non-Hispanic white, female

2003.....	1,002	61	54	23	121	148	196	152	76	70	63	33	5
2002.....	997	47	70	40	119	146	190	150	85	60	55	33	2
2001 3/.....	1,562	53	56	47	179	279	360	258	118	88	77	45	2
2000.....	1,025	43	52	46	143	166	218	140	52	56	72	34	3
1999.....	1,125	49	63	53	170	203	211	127	67	75	65	37	5

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)													
Non-Hispanic black, both sexes													
2003.....	1,957	100	136	52	349	343	426	316	112	63	47	13	-
2002.....	1,966	108	142	52	307	377	426	325	109	56	49	15	-
2001 3/.....	2,287	117	148	36	327	461	563	372	127	74	45	14	3
2000.....	2,126	132	135	54	375	403	505	268	116	78	47	10	3
1999.....	2,028	112	133	64	326	427	509	237	90	64	46	16	4
Non-Hispanic black, male													
2003.....	1,314	54	85	31	238	214	298	225	89	45	30	5	-
2002.....	1,298	52	80	30	215	238	258	255	92	40	32	6	-
2001 3/.....	1,468	67	91	23	222	301	336	264	87	48	19	7	3
2000.....	1,371	66	68	18	253	267	312	208	90	56	27	6	-
1999.....	1,305	57	67	29	225	259	344	175	65	48	23	10	3
Non-Hispanic black, female													
2003.....	643	46	51	21	111	129	128	91	23	18	17	8	-
2002.....	668	56	62	22	92	139	168	70	17	16	17	9	-
2001 3/.....	819	50	57	13	105	160	227	108	40	26	26	7	-
2000.....	755	66	67	36	122	136	193	60	26	22	20	4	3
1999.....	723	55	66	35	101	168	165	62	25	16	23	6	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Origin not stated 2/, both sexes

2003.....	64	3	1	-	10	7	8	12	3	5	2	2	11
2002.....	77	5	2	2	4	12	16	10	5	3	3	-	15
2001 3/.....	160	7	3	-	7	43	39	32	17	4	-	-	8
2000.....	55	4	-	-	10	4	6	13	6	3	-	1	8
1999.....	84	8	1	2	4	6	18	14	8	3	-	1	19

Origin not stated 2/, male

2003.....	44	2	-	-	6	5	4	10	3	5	1	2	6
2002.....	51	5	1	2	2	8	6	9	5	3	-	-	10
2001 3/.....	110	3	1	-	6	26	32	26	10	2	-	-	4
2000.....	36	3	-	-	5	3	5	8	4	3	-	1	4
1999.....	53	3	1	-	2	5	11	12	6	3	-	-	10

Origin not stated 2/, female

2003.....	20	1	1	-	4	2	4	2	-	-	1	-	5
2002.....	26	-	1	-	2	4	10	1	-	-	3	-	5
2001 3/.....	50	4	2	-	1	17	7	6	7	2	-	-	4
2000.....	19	1	-	-	5	1	1	5	2	-	-	-	4
1999.....	31	5	-	2	2	1	7	2	2	-	-	1	9

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Legal intervention (Y35,Y89.0)													
All origins, both sexes													
2003.....	423	-	-	2	93	135	117	51	19	4	2	-	-
2002.....	384	-	-	3	91	102	109	56	16	6	1	-	-
2001.....	396	-	-	-	88	107	115	51	24	6	4	1	-
2000.....	359	-	-	3	73	110	101	55	10	5	1	1	-
1999.....	398	-	-	1	67	136	121	50	20	2	1	-	-
All origins, male													
2003.....	405	-	-	1	89	131	114	46	18	4	2	-	-
2002.....	369	-	-	3	87	99	105	52	16	6	1	-	-
2001.....	380	-	-	-	84	104	108	51	22	6	4	1	-
2000.....	348	-	-	3	71	108	96	54	9	5	1	1	-
1999.....	392	-	-	1	67	135	117	49	20	2	1	-	-
All origins, female													
2003.....	18	-	-	1	4	4	3	5	1	-	-	-	-
2002.....	15	-	-	-	4	3	4	4	-	-	-	-	-
2001.....	16	-	-	-	4	3	7	-	2	-	-	-	-
2000.....	11	-	-	-	2	2	5	1	1	-	-	-	-
1999.....	6	-	-	-	-	1	4	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Legal intervention (Y35,Y89.0)													
Non-Hispanic 1/, both sexes													
2003.....	347	-	-	1	64	103	107	49	18	3	2	-	-
2002.....	315	-	-	2	69	80	94	49	15	5	1	-	-
2001.....	330	-	-	-	65	83	102	46	23	6	4	1	-
2000.....	308	-	-	2	54	95	90	51	9	5	1	1	-
1999.....	338	-	-	1	54	112	106	45	18	1	1	-	-
Non-Hispanic 1/, male													
2003.....	330	-	-	1	60	99	104	44	17	3	2	-	-
2002.....	301	-	-	2	65	78	90	45	15	5	1	-	-
2001.....	314	-	-	-	61	80	95	46	21	6	4	1	-
2000.....	298	-	-	2	53	93	85	50	8	5	1	1	-
1999.....	332	-	-	1	54	111	102	44	18	1	1	-	-
Non-Hispanic 1/, female													
2003.....	17	-	-	-	4	4	3	5	1	-	-	-	-
2002.....	14	-	-	-	4	2	4	4	-	-	-	-	-
2001.....	16	-	-	-	4	3	7	-	2	-	-	-	-
2000.....	10	-	-	-	1	2	5	1	1	-	-	-	-
1999.....	6	-	-	-	-	1	4	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Legal intervention (Y35,Y89.0)													
Non-Hispanic white, both sexes													
2003.....	211	-	-	-	27	45	80	39	16	3	1	-	-
2002.....	191	-	-	-	28	48	64	34	12	4	1	-	-
2001.....	230	-	-	-	37	56	74	36	17	6	4	-	-
2000.....	187	-	-	-	26	57	52	39	8	3	1	1	-
1999.....	197	-	-	-	20	62	65	35	13	1	1	-	-
Non-Hispanic white, male													
2003.....	199	-	-	-	25	43	77	35	15	3	1	-	-
2002.....	181	-	-	-	25	47	62	30	12	4	1	-	-
2001.....	221	-	-	-	35	55	69	36	16	6	4	-	-
2000.....	182	-	-	-	25	55	51	39	7	3	1	1	-
1999.....	192	-	-	-	20	62	61	34	13	1	1	-	-
Non-Hispanic white, female													
2003.....	12	-	-	-	2	2	3	4	1	-	-	-	-
2002.....	10	-	-	-	3	1	2	4	-	-	-	-	-
2001.....	9	-	-	-	2	1	5	-	1	-	-	-	-
2000.....	5	-	-	-	1	2	1	-	1	-	-	-	-
1999.....	5	-	-	-	-	-	4	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Legal intervention (Y35,Y89.0)													
Non-Hispanic black, both sexes													
2003.....	117	-	-	1	33	52	22	7	1	-	1	-	-
2002.....	109	-	-	2	37	28	28	11	2	1	-	-	-
2001.....	82	-	-	-	26	21	22	9	3	-	-	1	-
2000.....	110	-	-	2	27	33	35	11	-	2	-	-	-
1999.....	125	-	-	1	31	42	36	10	5	-	-	-	-
Non-Hispanic black, male													
2003.....	114	-	-	1	31	51	22	7	1	-	1	-	-
2002.....	105	-	-	2	36	27	26	11	2	1	-	-	-
2001.....	77	-	-	-	24	20	20	9	3	-	-	1	-
2000.....	107	-	-	2	27	33	33	10	-	2	-	-	-
1999.....	124	-	-	1	31	41	36	10	5	-	-	-	-
Non-Hispanic black, female													
2003.....	3	-	-	-	2	1	-	-	-	-	-	-	-
2002.....	4	-	-	-	1	1	2	-	-	-	-	-	-
2001.....	5	-	-	-	2	1	2	-	-	-	-	-	-
2000.....	3	-	-	-	-	-	2	1	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
All origins, both sexes													
2003.....	5,072	75	31	55	563	835	1,581	1,267	368	131	96	52	18
2002.....	4,830	101	57	68	487	828	1,516	1,140	320	139	95	65	14
2001.....	4,198	83	44	51	412	728	1,396	990	225	99	108	44	18
2000.....	3,819	65	39	60	370	679	1,286	841	228	104	79	57	11
1999.....	3,917	55	43	49	430	743	1,295	793	222	114	96	58	19
All origins, male													
2003.....	3,295	50	17	41	412	580	1,013	811	222	65	49	21	14
2002.....	3,114	52	28	38	358	585	981	696	203	83	46	31	13
2001.....	2,833	49	19	38	316	515	931	662	146	64	57	20	16
2000.....	2,635	27	26	50	285	489	885	570	152	70	49	21	11
1999.....	2,692	33	21	38	345	523	887	532	148	72	55	25	13
All origins, female													
2003.....	1,777	25	14	14	151	255	568	456	146	66	47	31	4
2002.....	1,716	49	29	30	129	243	535	444	117	56	49	34	1
2001.....	1,365	34	25	13	96	213	465	328	79	35	51	24	2
2000.....	1,184	38	13	10	85	190	401	271	76	34	30	36	-
1999.....	1,225	22	22	11	85	220	408	261	74	42	41	33	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Hispanic, both sexes													
2003.....	339	6	6	13	51	85	80	66	19	6	5	2	-
2002.....	320	14	11	4	51	73	90	49	15	9	2	-	2
2001.....	283	7	6	4	34	63	92	54	9	6	2	4	2
2000.....	215	5	5	1	34	61	64	25	12	5	2	-	1
1999.....	249	6	9	4	52	69	61	28	13	3	3	-	1
Hispanic, male													
2003.....	258	1	4	9	36	69	65	50	13	5	4	2	-
2002.....	242	6	7	1	45	58	64	40	14	4	1	-	2
2001.....	216	4	3	4	26	52	73	36	7	5	1	3	2
2000.....	165	2	2	1	30	49	47	20	9	2	2	-	1
1999.....	200	3	6	3	46	54	49	23	11	2	2	-	1
Hispanic, female													
2003.....	81	5	2	4	15	16	15	16	6	1	1	-	-
2002.....	78	8	4	3	6	15	26	9	1	5	1	-	-
2001.....	67	3	3	-	8	11	19	18	2	1	1	1	-
2000.....	50	3	3	-	4	12	17	5	3	3	-	-	-
1999.....	49	3	3	1	6	15	12	5	2	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
Non-Hispanic 1/, both sexes													
2003.....	4,701	68	25	42	512	748	1,491	1,196	346	124	91	50	8
2002.....	4,470	87	44	63	432	749	1,420	1,086	301	128	93	62	5
2001.....	3,876	76	38	47	377	661	1,298	927	210	91	105	40	6
2000.....	3,571	58	34	59	336	614	1,214	812	211	96	77	56	4
1999.....	3,630	49	34	45	376	666	1,230	757	207	107	93	56	10
Non-Hispanic 1/, male													
2003.....	3,016	48	13	32	376	510	942	759	206	59	45	19	7
2002.....	2,836	46	21	36	310	521	912	651	185	77	45	28	4
2001.....	2,586	45	16	34	289	459	853	622	134	57	55	17	5
2000.....	2,445	25	24	49	255	436	834	546	140	65	47	20	4
1999.....	2,463	30	15	35	297	462	834	503	135	67	53	24	8
Non-Hispanic 1/, female													
2003.....	1,685	20	12	10	136	238	549	437	140	65	46	31	1
2002.....	1,634	41	23	27	122	228	508	435	116	51	48	34	1
2001.....	1,290	31	22	13	88	202	445	305	76	34	50	23	1
2000.....	1,126	33	10	10	81	178	380	266	71	31	30	36	-
1999.....	1,167	19	19	10	79	204	396	254	72	40	40	32	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
Non-Hispanic white, both sexes													
2003.....	3,898	45	15	24	439	635	1,247	986	277	96	80	48	6
2002.....	3,627	43	33	44	370	607	1,160	873	253	105	77	58	4
2001.....	3,155	40	25	35	318	549	1,054	749	176	75	93	37	4
2000.....	2,823	35	22	37	271	490	965	647	161	76	70	47	2
1999.....	2,848	30	15	38	290	531	960	578	172	92	86	50	6
Non-Hispanic white, male													
2003.....	2,441	28	7	20	316	422	780	604	155	45	41	18	5
2002.....	2,265	23	16	24	264	419	739	504	151	57	38	27	3
2001.....	2,054	25	10	25	238	377	673	485	106	45	51	16	3
2000.....	1,900	16	15	31	201	341	661	425	101	48	43	16	2
1999.....	1,887	19	8	30	232	357	646	363	101	57	47	22	5
Non-Hispanic white, female													
2003.....	1,457	17	8	4	123	213	467	382	122	51	39	30	1
2002.....	1,362	20	17	20	106	188	421	369	102	48	39	31	1
2001.....	1,101	15	15	10	80	172	381	264	70	30	42	21	1
2000.....	923	19	7	6	70	149	304	222	60	28	27	31	-
1999.....	961	11	7	8	58	174	314	215	71	35	39	28	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic black, both sexes													
2003.....	675	21	9	14	50	93	213	190	53	21	7	2	2
2002.....	738	37	7	15	52	118	240	195	39	16	16	3	-
2001.....	629	36	13	11	43	90	222	163	30	12	5	3	1
2000.....	654	14	11	21	50	103	230	153	42	15	6	7	2
1999.....	686	18	15	7	70	119	239	166	28	12	5	3	4
Non-Hispanic black, male													
2003.....	487	19	5	8	42	74	141	141	40	12	2	1	2
2002.....	504	18	3	11	39	84	161	136	30	15	7	-	-
2001.....	467	20	6	8	37	65	165	127	26	10	1	1	1
2000.....	478	3	8	17	45	78	160	113	32	13	4	3	2
1999.....	514	10	7	5	55	91	170	132	27	9	4	1	3
Non-Hispanic black, female													
2003.....	188	2	4	6	8	19	72	49	13	9	5	1	-
2002.....	234	19	4	4	13	34	79	59	9	1	9	3	-
2001.....	162	16	7	3	6	25	57	36	4	2	4	2	-
2000.....	176	11	3	4	5	25	70	40	10	2	2	4	-
1999.....	172	8	8	2	15	28	69	34	1	3	1	2	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Origin not stated 2/, both sexes

2003.....	32	1	-	-	-	2	10	5	3	1	-	-	10
2002.....	40	-	2	1	4	6	6	5	4	2	-	3	7
2001.....	39	-	-	-	1	4	6	9	6	2	1	-	10
2000.....	33	2	-	-	-	4	8	4	5	3	-	1	6
1999.....	38	-	-	-	2	8	4	8	2	4	-	2	8

Origin not stated 2/, male

2003.....	21	1	-	-	-	1	6	2	3	1	-	-	7
2002.....	36	-	-	1	3	6	5	5	4	2	-	3	7
2001.....	31	-	-	-	1	4	5	4	5	2	1	-	9
2000.....	25	-	-	-	-	4	4	4	3	3	-	1	6
1999.....	29	-	-	-	2	7	4	6	2	3	-	1	4

Origin not stated 2/, female

2003.....	11	-	-	-	-	1	4	3	-	-	-	-	3
2002.....	4	-	2	-	1	-	1	-	-	-	-	-	-
2001.....	8	-	-	-	-	-	1	5	1	-	-	-	1
2000.....	8	2	-	-	-	-	4	-	2	-	-	-	-
1999.....	9	-	-	-	-	1	-	2	-	1	-	1	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
All origins, both sexes													
2003.....	232	-	1	12	69	38	43	31	20	8	9	1	-
2002.....	243	-	1	6	79	46	50	29	12	15	5	-	-
2001.....	231	-	-	6	64	48	50	29	12	10	11	1	-
2000.....	230	-	-	10	71	52	39	23	15	11	6	2	1
1999.....	324	-	3	12	127	51	57	28	14	15	13	4	-
All origins, male													
2003.....	186	-	1	12	60	33	26	22	16	7	8	1	-
2002.....	199	-	1	5	71	39	39	23	7	10	4	-	-
2001.....	190	-	-	6	59	42	33	21	11	8	9	1	-
2000.....	190	-	-	9	62	43	30	18	11	9	5	2	1
1999.....	277	-	3	12	111	44	45	21	11	14	12	4	-
All origins, female													
2003.....	46	-	-	-	9	5	17	9	4	1	1	-	-
2002.....	44	-	-	1	8	7	11	6	5	5	1	-	-
2001.....	41	-	-	-	5	6	17	8	1	2	2	-	-
2000.....	40	-	-	1	9	9	9	5	4	2	1	-	-
1999.....	47	-	-	-	16	7	12	7	3	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
Hispanic, both sexes													
2003.....	31	-	-	3	13	6	5	3	-	1	-	-	-
2002.....	23	-	-	-	14	6	3	-	-	-	-	-	-
2001.....	21	-	-	1	8	4	4	3	1	-	-	-	-
2000.....	20	-	-	-	13	4	2	-	-	1	-	-	-
1999.....	31	-	1	1	18	3	3	2	1	2	-	-	-
Hispanic, male													
2003.....	26	-	-	3	9	6	5	2	-	1	-	-	-
2002.....	21	-	-	-	13	6	2	-	-	-	-	-	-
2001.....	16	-	-	1	7	3	2	2	1	-	-	-	-
2000.....	17	-	-	-	13	2	1	-	-	1	-	-	-
1999.....	26	-	1	1	16	2	2	1	1	2	-	-	-
Hispanic, female													
2003.....	5	-	-	-	4	-	-	1	-	-	-	-	-
2002.....	2	-	-	-	1	-	1	-	-	-	-	-	-
2001.....	5	-	-	-	1	1	2	1	-	-	-	-	-
2000.....	3	-	-	-	-	2	1	-	-	-	-	-	-
1999.....	5	-	-	-	2	1	1	1	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
Non-Hispanic 1/, both sexes													
2003.....	201	-	1	9	56	32	38	28	20	7	9	1	-
2002.....	215	-	1	5	63	39	47	29	11	15	5	-	-
2001.....	208	-	-	5	56	43	46	26	11	9	11	1	-
2000.....	209	-	-	10	58	48	37	23	15	10	6	2	-
1999.....	293	-	2	11	109	48	54	26	13	13	13	4	-
Non-Hispanic 1/, male													
2003.....	160	-	1	9	51	27	21	20	16	6	8	1	-
2002.....	173	-	1	4	56	32	37	23	6	10	4	-	-
2001.....	172	-	-	5	52	38	31	19	10	7	9	1	-
2000.....	172	-	-	9	49	41	29	18	11	8	5	2	-
1999.....	251	-	2	11	95	42	43	20	10	12	12	4	-
Non-Hispanic 1/, female													
2003.....	41	-	-	-	5	5	17	8	4	1	1	-	-
2002.....	42	-	-	1	7	7	10	6	5	5	1	-	-
2001.....	36	-	-	-	4	5	15	7	1	2	2	-	-
2000.....	37	-	-	1	9	7	8	5	4	2	1	-	-
1999.....	42	-	-	-	14	6	11	6	3	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
Non-Hispanic white, both sexes													
2003.....	152	-	-	4	34	21	33	26	17	7	9	1	-
2002.....	169	-	-	3	47	27	38	27	9	14	4	-	-
2001.....	166	-	-	3	36	35	41	23	8	9	10	1	-
2000.....	148	-	-	5	36	30	30	20	11	8	6	2	-
1999.....	217	-	1	10	67	33	47	21	11	11	12	4	-
Non-Hispanic white, male													
2003.....	118	-	-	4	31	17	17	19	15	6	8	1	-
2002.....	132	-	-	2	40	21	30	21	6	9	3	-	-
2001.....	133	-	-	3	33	31	27	16	7	7	8	1	-
2000.....	120	-	-	5	30	26	24	15	7	6	5	2	-
1999.....	187	-	1	10	60	29	39	15	8	10	11	4	-
Non-Hispanic white, female													
2003.....	34	-	-	-	3	4	16	7	2	1	1	-	-
2002.....	37	-	-	1	7	6	8	6	3	5	1	-	-
2001.....	33	-	-	-	3	4	14	7	1	2	2	-	-
2000.....	28	-	-	-	6	4	6	5	4	2	1	-	-
1999.....	30	-	-	-	7	4	8	6	3	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
Non-Hispanic black, both sexes													
2003.....	35	-	1	4	17	7	3	2	1	-	-	-	-
2002.....	37	-	1	2	14	10	6	1	1	1	1	-	-
2001.....	31	-	-	2	15	5	2	3	3	-	1	-	-
2000.....	53	-	-	5	19	15	7	2	3	2	-	-	-
1999.....	65	-	1	1	35	14	6	4	1	2	1	-	-
Non-Hispanic black, male													
2003.....	32	-	1	4	16	7	2	1	1	-	-	-	-
2002.....	35	-	1	2	14	10	5	1	1	1	1	-	-
2001.....	28	-	-	2	14	4	1	3	3	-	1	-	-
2000.....	46	-	-	4	18	12	5	2	3	2	-	-	-
1999.....	55	-	1	1	29	12	4	4	1	2	1	-	-
Non-Hispanic black, female													
2003.....	3	-	-	-	1	-	1	1	-	-	-	-	-
2002.....	2	-	-	-	-	-	1	-	1	-	-	-	-
2001.....	3	-	-	-	1	1	1	-	-	-	-	-	-
2000.....	7	-	-	1	1	3	2	-	-	-	-	-	-
1999.....	10	-	-	-	6	2	2	-	-	-	-	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All origins, both sexes

2003.....	4,840	75	30	43	494	797	1,538	1,236	348	123	87	51	18
2002.....	4,587	101	56	62	408	782	1,466	1,111	308	124	90	65	14
2001.....	3,967	83	44	45	348	680	1,346	961	213	89	97	43	18
2000.....	3,589	65	39	50	299	627	1,247	818	213	93	73	55	10
1999.....	3,593	55	40	37	303	692	1,238	765	208	99	83	54	19

All origins, male

2003.....	3,109	50	16	29	352	547	987	789	206	58	41	20	14
2002.....	2,915	52	27	33	287	546	942	673	196	73	42	31	13
2001.....	2,643	49	19	32	257	473	898	641	135	56	48	19	16
2000.....	2,445	27	26	41	223	446	855	552	141	61	44	19	10
1999.....	2,415	33	18	26	234	479	842	511	137	58	43	21	13

All origins, female

2003.....	1,731	25	14	14	142	250	551	447	142	65	46	31	4
2002.....	1,672	49	29	29	121	236	524	438	112	51	48	34	1
2001.....	1,324	34	25	13	91	207	448	320	78	33	49	24	2
2000.....	1,144	38	13	9	76	181	392	266	72	32	29	36	-
1999.....	1,178	22	22	11	69	213	396	254	71	41	40	33	6

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Hispanic, both sexes

2003.....	308	6	6	10	38	79	75	63	19	5	5	2	-
2002.....	297	14	11	4	37	67	87	49	15	9	2	-	2
2001.....	262	7	6	3	26	59	88	51	8	6	2	4	2
2000.....	195	5	5	1	21	57	62	25	12	4	2	-	1
1999.....	218	6	8	3	34	66	58	26	12	1	3	-	1

Hispanic, male

2003.....	232	1	4	6	27	63	60	48	13	4	4	2	-
2002.....	221	6	7	1	32	52	62	40	14	4	1	-	2
2001.....	200	4	3	3	19	49	71	34	6	5	1	3	2
2000.....	148	2	2	1	17	47	46	20	9	1	2	-	1
1999.....	174	3	5	2	30	52	47	22	10	-	2	-	1

Hispanic, female

2003.....	76	5	2	4	11	16	15	15	6	1	1	-	-
2002.....	76	8	4	3	5	15	25	9	1	5	1	-	-
2001.....	62	3	3	-	7	10	17	17	2	1	1	1	-
2000.....	47	3	3	-	4	10	16	5	3	3	-	-	-
1999.....	44	3	3	1	4	14	11	4	2	1	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic 1/, both sexes

2003.....	4,500	68	24	33	456	716	1,453	1,168	326	117	82	49	8
2002.....	4,255	87	43	58	369	710	1,373	1,057	290	113	88	62	5
2001.....	3,668	76	38	42	321	618	1,252	901	199	82	94	39	6
2000.....	3,362	58	34	49	278	566	1,177	789	196	86	71	54	4
1999.....	3,337	49	32	34	267	618	1,176	731	194	94	80	52	10

Non-Hispanic 1/, male

2003.....	2,856	48	12	23	325	483	921	739	190	53	37	18	7
2002.....	2,663	46	20	32	254	489	875	628	179	67	41	28	4
2001.....	2,414	45	16	29	237	421	822	603	124	50	46	16	5
2000.....	2,273	25	24	40	206	395	805	528	129	57	42	18	4
1999.....	2,212	30	13	24	202	420	791	483	125	55	41	20	8

Non-Hispanic 1/, female

2003.....	1,644	20	12	10	131	233	532	429	136	64	45	31	1
2002.....	1,592	41	23	26	115	221	498	429	111	46	47	34	1
2001.....	1,254	31	22	13	84	197	430	298	75	32	48	23	1
2000.....	1,089	33	10	9	72	171	372	261	67	29	29	36	-
1999.....	1,125	19	19	10	65	198	385	248	69	39	39	32	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic white, both sexes

2003.....	3,746	45	15	20	405	614	1,214	960	260	89	71	47	6
2002.....	3,458	43	33	41	323	580	1,122	846	244	91	73	58	4
2001.....	2,989	40	25	32	282	514	1,013	726	168	66	83	36	4
2000.....	2,675	35	22	32	235	460	935	627	150	68	64	45	2
1999.....	2,631	30	14	28	223	498	913	557	161	81	74	46	6

Non-Hispanic white, male

2003.....	2,323	28	7	16	285	405	763	585	140	39	33	17	5
2002.....	2,133	23	16	22	224	398	709	483	145	48	35	27	3
2001.....	1,921	25	10	22	205	346	646	469	99	38	43	15	3
2000.....	1,780	16	15	26	171	315	637	410	94	42	38	14	2
1999.....	1,700	19	7	20	172	328	607	348	93	47	36	18	5

Non-Hispanic white, female

2003.....	1,423	17	8	4	120	209	451	375	120	50	38	30	1
2002.....	1,325	20	17	19	99	182	413	363	99	43	38	31	1
2001.....	1,068	15	15	10	77	168	367	257	69	28	40	21	1
2000.....	895	19	7	6	64	145	298	217	56	26	26	31	-
1999.....	931	11	7	8	51	170	306	209	68	34	38	28	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic black, both sexes

2003.....	640	21	8	10	33	86	210	188	52	21	7	2	2
2002.....	701	37	6	13	38	108	234	194	38	15	15	3	-
2001.....	598	36	13	9	28	85	220	160	27	12	4	3	1
2000.....	601	14	11	16	31	88	223	151	39	13	6	7	2
1999.....	621	18	14	6	35	105	233	162	27	10	4	3	4

Non-Hispanic black, male

2003.....	455	19	4	4	26	67	139	140	39	12	2	1	2
2002.....	469	18	2	9	25	74	156	135	30	14	6	-	-
2001.....	439	20	6	6	23	61	164	124	23	10	-	1	1
2000.....	432	3	8	13	27	66	155	111	29	11	4	3	2
1999.....	459	10	6	4	26	79	166	128	26	7	3	1	3

Non-Hispanic black, female

2003.....	185	2	4	6	7	19	71	48	13	9	5	1	-
2002.....	232	19	4	4	13	34	78	59	8	1	9	3	-
2001.....	159	16	7	3	5	24	56	36	4	2	4	2	-
2000.....	169	11	3	3	4	22	68	40	10	2	2	4	-
1999.....	162	8	8	2	9	26	67	34	1	3	1	2	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Origin not stated 2/, both sexes

2003.....	32	1	-	-	-	2	10	5	3	1	-	-	10
2002.....	35	-	2	-	2	5	6	5	3	2	-	3	7
2001.....	37	-	-	-	1	3	6	9	6	1	1	-	10
2000.....	32	2	-	-	-	4	8	4	5	3	-	1	5
1999.....	38	-	-	-	2	8	4	8	2	4	-	2	8

Origin not stated 2/, male

2003.....	21	1	-	-	-	1	6	2	3	1	-	-	7
2002.....	31	-	-	-	1	5	5	5	3	2	-	3	7
2001.....	29	-	-	-	1	3	5	4	5	1	1	-	9
2000.....	24	-	-	-	-	4	4	4	3	3	-	1	5
1999.....	29	-	-	-	2	7	4	6	2	3	-	1	4

Origin not stated 2/, female

2003.....	11	-	-	-	-	1	4	3	-	-	-	-	3
2002.....	4	-	2	-	1	-	1	-	-	-	-	-	-
2001.....	8	-	-	-	-	-	1	5	1	-	-	-	1
2000.....	8	2	-	-	-	-	4	-	2	-	-	-	-
1999.....	9	-	-	-	-	1	-	2	-	1	-	1	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Operations of war and their sequelae (Y36,Y89.1)

All origins, both sexes

2003.....	14	-	-	-	1	-	-	1	5	2	5	-	-
2002.....	20	-	-	-	-	1	1	1	3	3	10	1	-
2001.....	17	-	-	-	1	-	1	2	-	4	8	1	-
2000.....	16	-	-	-	-	-	-	2	1	3	9	1	-
1999.....	23	-	-	-	-	-	1	2	1	8	8	3	-

All origins, male

2003.....	14	-	-	-	1	-	-	1	5	2	5	-	-
2002.....	20	-	-	-	-	1	1	1	3	3	10	1	-
2001.....	16	-	-	-	1	-	1	2	-	4	7	1	-
2000.....	16	-	-	-	-	-	-	2	1	3	9	1	-
1999.....	22	-	-	-	-	-	1	2	1	8	7	3	-

All origins, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Operations of war and their sequelae (Y36,Y89.1)

Non-Hispanic 1/, both sexes

2003.....	14	-	-	-	1	-	-	1	5	2	5	-	-
2002.....	19	-	-	-	-	-	1	1	3	3	10	1	-
2001.....	15	-	-	-	-	-	1	2	-	4	7	1	-
2000.....	15	-	-	-	-	-	-	2	1	2	9	1	-
1999.....	21	-	-	-	-	-	1	2	1	8	7	2	-

Non-Hispanic 1/, male

2003.....	14	-	-	-	1	-	-	1	5	2	5	-	-
2002.....	19	-	-	-	-	-	1	1	3	3	10	1	-
2001.....	14	-	-	-	-	-	1	2	-	4	6	1	-
2000.....	15	-	-	-	-	-	-	2	1	2	9	1	-
1999.....	20	-	-	-	-	-	1	2	1	8	6	2	-

Non-Hispanic 1/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Operations of war and their sequelae (Y36,Y89.1)													
Non-Hispanic white, both sexes													
2003.....	13	-	-	-	1	-	-	1	4	2	5	-	-
2002.....	14	-	-	-	-	-	-	-	2	2	9	1	-
2001.....	14	-	-	-	-	-	1	2	-	3	7	1	-
2000.....	14	-	-	-	-	-	-	1	1	2	9	1	-
1999.....	20	-	-	-	-	-	1	2	1	7	7	2	-
Non-Hispanic white, male													
2003.....	13	-	-	-	1	-	-	1	4	2	5	-	-
2002.....	14	-	-	-	-	-	-	-	2	2	9	1	-
2001.....	13	-	-	-	-	-	1	2	-	3	6	1	-
2000.....	14	-	-	-	-	-	-	1	1	2	9	1	-
1999.....	19	-	-	-	-	-	1	2	1	7	6	2	-
Non-Hispanic white, female													
2003.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	1	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
All origins, both sexes													
2003.....	2,855	15	24	12	44	69	179	273	380	534	794	530	1
2002.....	2,843	15	19	35	35	79	171	265	351	560	792	521	-
2001.....	3,021	26	24	14	43	79	172	283	384	567	864	564	1
2000.....	3,059	19	22	31	32	85	149	295	376	647	825	578	-
1999.....	2,823	54	30	18	36	78	152	214	315	586	811	529	-
All origins, male													
2003.....	1,333	8	16	4	26	36	96	146	195	264	355	186	1
2002.....	1,236	8	11	26	19	47	82	127	169	256	320	171	-
2001.....	1,357	15	14	8	19	42	83	143	196	265	383	189	-
2000.....	1,430	12	11	15	16	47	76	143	181	331	402	196	-
1999.....	1,308	31	12	9	21	43	78	98	165	279	375	197	-
All origins, female													
2003.....	1,522	7	8	8	18	33	83	127	185	270	439	344	-
2002.....	1,607	7	8	9	16	32	89	138	182	304	472	350	-
2001.....	1,664	11	10	6	24	37	89	140	188	302	481	375	1
2000.....	1,629	7	11	16	16	38	73	152	195	316	423	382	-
1999.....	1,515	23	18	9	15	35	74	116	150	307	436	332	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
Hispanic, both sexes													
2003.....	155	3	4	1	5	9	22	14	21	38	24	13	1
2002.....	137	2	7	5	6	8	13	14	19	28	29	6	-
2001.....	145	3	6	4	8	7	12	22	13	25	29	15	1
2000.....	136	3	4	6	7	4	11	13	11	35	26	16	-
1999.....	100	5	6	3	4	8	8	11	16	15	12	12	-
Hispanic, male													
2003.....	76	1	4	-	3	2	15	10	9	17	10	4	1
2002.....	57	2	4	4	3	1	5	8	6	11	13	-	-
2001.....	65	-	3	3	3	3	4	11	8	8	14	8	-
2000.....	70	2	2	4	6	2	5	4	8	16	15	6	-
1999.....	52	3	3	2	2	3	7	6	9	5	8	4	-
Hispanic, female													
2003.....	79	2	-	1	2	7	7	4	12	21	14	9	-
2002.....	80	-	3	1	3	7	8	6	13	17	16	6	-
2001.....	80	3	3	1	5	4	8	11	5	17	15	7	1
2000.....	66	1	2	2	1	2	6	9	3	19	11	10	-
1999.....	48	2	3	1	2	5	1	5	7	10	4	8	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
Non-Hispanic 1/, both sexes													
2003.....	2,693	12	19	11	38	60	156	257	359	496	769	516	-
2002.....	2,695	13	12	30	29	71	158	249	332	530	758	513	-
2001.....	2,861	23	18	10	34	72	158	261	369	537	831	548	-
2000.....	2,914	16	18	25	25	81	137	281	363	610	797	561	-
1999.....	2,714	49	24	15	32	70	141	202	298	570	798	515	-
Non-Hispanic 1/, male													
2003.....	1,252	7	11	4	22	34	80	135	186	247	344	182	-
2002.....	1,175	6	7	22	16	46	77	117	163	244	306	171	-
2001.....	1,286	15	11	5	15	39	78	132	188	254	368	181	-
2000.....	1,354	10	9	11	10	45	70	139	172	313	385	190	-
1999.....	1,253	28	9	7	19	40	70	91	156	274	366	193	-
Non-Hispanic 1/, female													
2003.....	1,441	5	8	7	16	26	76	122	173	249	425	334	-
2002.....	1,520	7	5	8	13	25	81	132	169	286	452	342	-
2001.....	1,575	8	7	5	19	33	80	129	181	283	463	367	-
2000.....	1,560	6	9	14	15	36	67	142	191	297	412	371	-
1999.....	1,461	21	15	8	13	30	71	111	142	296	432	322	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
Non-Hispanic white, both sexes													
2003.....	2,202	6	9	7	31	39	119	178	276	410	671	456	-
2002.....	2,188	6	7	16	21	49	107	169	252	440	664	457	-
2001.....	2,294	14	10	6	25	51	105	185	279	417	735	467	-
2000.....	2,378	12	8	17	17	56	103	218	272	498	691	486	-
1999.....	2,212	30	16	11	26	47	103	148	224	478	688	441	-
Non-Hispanic white, male													
2003.....	1,044	4	5	3	18	23	66	88	148	214	312	163	-
2002.....	966	3	5	12	13	34	60	85	129	200	276	149	-
2001.....	1,016	9	7	2	12	26	50	91	137	197	333	152	-
2000.....	1,135	9	6	7	7	31	58	111	135	255	347	169	-
1999.....	1,025	19	8	5	15	29	52	65	117	230	321	164	-
Non-Hispanic white, female													
2003.....	1,158	2	4	4	13	16	53	90	128	196	359	293	-
2002.....	1,222	3	2	4	8	15	47	84	123	240	388	308	-
2001.....	1,278	5	3	4	13	25	55	94	142	220	402	315	-
2000.....	1,243	3	2	10	10	25	45	107	137	243	344	317	-
1999.....	1,187	11	8	6	11	18	51	83	107	248	367	277	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
Non-Hispanic black, both sexes													
2003.....	425	5	9	3	6	16	33	66	74	76	86	51	-
2002.....	445	7	5	14	5	17	48	75	70	77	80	47	-
2001.....	515	9	7	3	7	21	48	70	84	104	86	76	-
2000.....	484	4	8	5	7	23	31	57	86	101	95	67	-
1999.....	442	18	8	4	6	18	35	51	61	84	94	63	-
Non-Hispanic black, male													
2003.....	174	3	5	1	3	8	11	40	35	27	27	14	-
2002.....	182	3	2	10	2	9	16	29	30	37	25	19	-
2001.....	247	6	4	2	2	13	26	38	47	53	30	26	-
2000.....	192	1	2	2	2	13	12	25	34	52	33	16	-
1999.....	195	8	1	2	4	10	16	26	30	40	38	20	-
Non-Hispanic black, female													
2003.....	251	2	4	2	3	8	22	26	39	49	59	37	-
2002.....	263	4	3	4	3	8	32	46	40	40	55	28	-
2001.....	268	3	3	1	5	8	22	32	37	51	56	50	-
2000.....	292	3	6	3	5	10	19	32	52	49	62	51	-
1999.....	247	10	7	2	2	8	19	25	31	44	56	43	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

Origin not stated 2/, both sexes

2003.....	7	-	1	-	1	-	1	2	-	-	1	1	-
2002.....	11	-	-	-	-	-	-	2	-	2	5	2	-
2001.....	15	-	-	-	1	-	2	-	2	5	4	1	-
2000.....	9	-	-	-	-	-	1	1	2	2	2	1	-
1999.....	9	-	-	-	-	-	3	1	1	1	1	2	-

Origin not stated 2/, male

2003.....	5	-	1	-	1	-	1	1	-	-	1	-	-
2002.....	4	-	-	-	-	-	-	2	-	1	1	-	-
2001.....	6	-	-	-	1	-	1	-	-	3	1	-	-
2000.....	6	-	-	-	-	-	1	-	1	2	2	-	-
1999.....	3	-	-	-	-	-	1	1	-	-	1	-	-

Origin not stated 2/, female

2003.....	2	-	-	-	-	-	-	1	-	-	-	1	-
2002.....	7	-	-	-	-	-	-	-	-	1	4	2	-
2001.....	9	-	-	-	-	-	1	-	2	2	3	1	-
2000.....	3	-	-	-	-	-	-	1	1	-	-	1	-
1999.....	6	-	-	-	-	-	2	-	1	1	-	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

All origins, both sexes

2003.....	20,687	-	-	5	124	596	3,572	7,063	5,168	2,771	1,183	194	11
2002 5/.....	20,218	1	2	3	105	601	3,636	6,748	4,888	2,835	1,190	201	8
2001 5/.....	20,114	2	-	2	106	645	3,746	6,727	4,599	2,846	1,204	221	16
2000 5/.....	19,643	3	1	4	95	650	3,837	6,143	4,542	2,910	1,223	227	8
1999 5/.....	19,469	1	-	2	100	641	3,829	6,002	4,439	2,926	1,291	230	8

All origins, male

2003.....	15,630	-	-	3	101	472	2,534	5,414	3,997	2,120	866	113	10
2002 5/.....	15,272	-	1	2	86	458	2,571	5,171	3,837	2,158	845	135	8
2001 5/.....	15,149	2	-	-	88	466	2,665	5,116	3,607	2,181	858	151	15
2000 5/.....	14,993	2	-	4	76	504	2,808	4,792	3,489	2,249	915	146	8
1999 5/.....	14,894	1	-	1	80	459	2,816	4,702	3,481	2,270	936	141	7

All origins, female

2003.....	5,057	-	-	2	23	124	1,038	1,649	1,171	651	317	81	1
2002 5/.....	4,946	1	1	1	19	143	1,065	1,577	1,051	677	345	66	-
2001 5/.....	4,965	-	-	2	18	179	1,081	1,611	992	665	346	70	1
2000 5/.....	4,650	1	1	-	19	146	1,029	1,351	1,053	661	308	81	-
1999 5/.....	4,575	-	-	1	20	182	1,013	1,300	958	656	355	89	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Hispanic, both sexes

2003.....	2,422	-	-	1	39	133	493	772	576	281	114	10	3
2002 5/.....	2,408	-	-	1	33	147	506	774	521	319	87	19	1
2001 5/.....	2,381	-	-	-	31	146	493	766	522	296	113	10	4
2000 5/.....	2,323	-	-	1	32	136	531	698	504	311	96	11	3
1999 5/.....	2,184	-	-	1	27	131	505	668	470	271	90	20	1

Hispanic, male

2003.....	2,048	-	-	-	34	113	410	673	500	219	91	6	2
2002 5/.....	2,065	-	-	-	30	134	431	661	454	276	65	13	1
2001 5/.....	2,026	-	-	-	25	122	417	652	450	262	85	9	4
2000 5/.....	2,024	-	-	1	29	129	445	606	440	278	86	7	3
1999 5/.....	1,864	-	-	1	25	114	426	575	406	233	69	14	1

Hispanic, female

2003.....	374	-	-	1	5	20	83	99	76	62	23	4	1
2002 5/.....	343	-	-	1	3	13	75	113	67	43	22	6	-
2001 5/.....	355	-	-	-	6	24	76	114	72	34	28	1	-
2000 5/.....	299	-	-	-	3	7	86	92	64	33	10	4	-
1999 5/.....	320	-	-	-	2	17	79	93	64	38	21	6	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Non-Hispanic 1/, both sexes

2003.....	18,160	-	-	4	84	460	3,066	6,259	4,556	2,475	1,067	184	5
2002 5/.....	17,661	1	2	2	71	450	3,102	5,921	4,330	2,500	1,097	181	4
2001 5/.....	17,593	2	-	2	75	495	3,234	5,909	4,041	2,530	1,087	211	7
2000 5/.....	17,177	3	1	3	62	509	3,280	5,389	4,012	2,579	1,120	215	4
1999 5/.....	17,143	1	-	1	71	505	3,291	5,288	3,938	2,637	1,196	210	5

Non-Hispanic 1/, male

2003.....	13,490	-	-	3	66	356	2,114	4,714	3,463	1,889	773	107	5
2002 5/.....	13,078	-	1	2	55	320	2,120	4,463	3,351	1,866	775	121	4
2001 5/.....	13,009	2	-	-	63	341	2,231	4,422	3,128	1,904	770	142	6
2000 5/.....	12,843	2	-	3	46	371	2,338	4,137	3,027	1,955	822	138	4
1999 5/.....	12,905	1	-	-	53	340	2,363	4,087	3,047	2,020	862	127	5

Non-Hispanic 1/, female

2003.....	4,670	-	-	1	18	104	952	1,545	1,093	586	294	77	-
2002 5/.....	4,583	1	1	-	16	130	982	1,458	979	634	322	60	-
2001 5/.....	4,584	-	-	2	12	154	1,003	1,487	913	626	317	69	1
2000 5/.....	4,334	1	1	-	16	138	942	1,252	985	624	298	77	-
1999 5/.....	4,238	-	-	1	18	165	928	1,201	891	617	334	83	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Non-Hispanic white, both sexes

2003.....	14,977	-	-	3	57	347	2,465	5,105	3,746	2,124	959	167	4
2002 5/.....	14,494	-	1	1	53	327	2,505	4,834	3,518	2,119	972	161	3
2001 5/.....	14,186	1	-	2	47	360	2,533	4,691	3,267	2,131	967	182	5
2000 5/.....	13,815	3	-	2	44	365	2,555	4,259	3,219	2,182	992	192	2
1999 5/.....	13,633	1	-	1	49	380	2,507	4,142	3,129	2,200	1,038	181	5

Non-Hispanic white, male

2003.....	11,133	-	-	3	48	267	1,718	3,837	2,845	1,628	687	96	4
2002 5/.....	10,783	-	1	1	43	236	1,730	3,678	2,736	1,564	687	104	3
2001 5/.....	10,497	1	-	-	41	254	1,740	3,553	2,512	1,585	685	122	4
2000 5/.....	10,408	2	-	2	34	269	1,840	3,317	2,444	1,648	729	121	2
1999 5/.....	10,337	1	-	-	37	261	1,830	3,243	2,431	1,669	751	109	5

Non-Hispanic white, female

2003.....	3,844	-	-	-	9	80	747	1,268	901	496	272	71	-
2002 5/.....	3,711	-	-	-	10	91	775	1,156	782	555	285	57	-
2001 5/.....	3,689	-	-	2	6	106	793	1,138	755	546	282	60	1
2000 5/.....	3,407	1	-	-	10	96	715	942	775	534	263	71	-
1999 5/.....	3,296	-	-	1	12	119	677	899	698	531	287	72	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Non-Hispanic black, both sexes

2003.....	2,367	-	-	1	7	60	388	867	657	285	87	14	1
2002 5/.....	2,396	1	1	-	10	70	384	853	653	303	104	16	1
2001 5/.....	2,677	1	-	-	17	70	496	986	645	337	104	19	2
2000 5/.....	2,672	-	1	1	8	92	516	946	639	343	104	20	2
1999 5/.....	2,794	-	-	-	10	72	559	933	687	377	132	24	-

Non-Hispanic black, male

2003.....	1,787	-	-	-	5	48	258	677	507	214	67	10	1
2002 5/.....	1,768	-	-	-	8	52	252	623	499	246	74	13	1
2001 5/.....	2,016	1	-	-	13	47	355	725	515	273	74	11	2
2000 5/.....	1,959	-	-	1	7	63	356	693	474	269	79	15	2
1999 5/.....	2,066	-	-	-	8	45	380	690	533	304	92	14	-

Non-Hispanic black, female

2003.....	580	-	-	1	2	12	130	190	150	71	20	4	-
2002 5/.....	628	1	1	-	2	18	132	230	154	57	30	3	-
2001 5/.....	661	-	-	-	4	23	141	261	130	64	30	8	-
2000 5/.....	713	-	1	-	1	29	160	253	165	74	25	5	-
1999 5/.....	728	-	-	-	2	27	179	243	154	73	40	10	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Origin not stated 2/, both sexes

2003.....	105	-	-	-	1	3	13	32	36	15	2	-	3
2002 5/.....	149	-	-	-	1	4	28	53	37	16	6	1	3
2001 5/.....	140	-	-	-	-	4	19	52	36	20	4	-	5
2000 5/.....	143	-	-	-	1	5	26	56	26	20	7	1	1
1999 5/.....	142	-	-	-	2	5	33	46	31	18	5	-	2

Origin not stated 2/, male

2003.....	92	-	-	-	1	3	10	27	34	12	2	-	3
2002 5/.....	129	-	-	-	1	4	20	47	32	16	5	1	3
2001 5/.....	114	-	-	-	-	3	17	42	29	15	3	-	5
2000 5/.....	126	-	-	-	1	4	25	49	22	16	7	1	1
1999 5/.....	125	-	-	-	2	5	27	40	28	17	5	-	1

Origin not stated 2/, female

2003.....	13	-	-	-	-	-	3	5	2	3	-	-	-
2002 5/.....	20	-	-	-	-	-	8	6	5	-	1	-	-
2001 5/.....	26	-	-	-	-	1	2	10	7	5	1	-	-
2000 5/.....	17	-	-	-	-	1	1	7	4	4	-	-	-
1999 5/.....	17	-	-	-	-	-	6	6	3	1	-	-	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

All origins, both sexes

2003.....	28,723	25	35	41	2,619	4,845	9,158	8,174	2,239	755	537	283	12
2002 5/.....	26,040	29	30	42	2,215	4,449	8,858	7,182	1,828	653	484	260	10
2001 5/.....	21,705	22	26	45	1,813	3,739	7,669	5,796	1,361	553	439	226	16
2000 5/.....	19,720	18	15	44	1,567	3,527	7,206	4,986	1,187	486	430	240	14
1999 5/.....	19,128	21	25	30	1,351	3,579	7,095	4,624	1,161	556	467	201	18

All origins, male

2003.....	18,426	17	21	20	1,955	3,361	5,873	5,153	1,266	384	253	112	11
2002 5/.....	16,734	20	9	24	1,622	3,139	5,728	4,517	1,040	311	215	102	7
2001 5/.....	14,253	14	21	22	1,350	2,640	5,079	3,793	760	293	191	74	16
2000 5/.....	13,137	12	13	25	1,166	2,515	4,890	3,358	631	235	187	92	13
1999 5/.....	12,885	14	12	16	981	2,579	4,907	3,133	688	286	184	74	11

All origins, female

2003.....	10,297	8	14	21	664	1,484	3,285	3,021	973	371	284	171	1
2002 5/.....	9,306	9	21	18	593	1,310	3,130	2,665	788	342	269	158	3
2001 5/.....	7,452	8	5	23	463	1,099	2,590	2,003	601	260	248	152	-
2000 5/.....	6,583	6	2	19	401	1,012	2,316	1,628	556	251	243	148	1
1999 5/.....	6,243	7	13	14	370	1,000	2,188	1,491	473	270	283	127	7

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

Hispanic, both sexes

2003.....	2,358	4	5	7	243	533	789	570	135	38	24	10	-
2002 5/.....	2,137	5	4	4	217	456	777	505	111	36	20	2	-
2001 5/.....	1,731	-	4	5	162	408	644	373	88	29	12	4	2
2000 5/.....	1,700	2	2	4	148	449	651	332	79	20	8	5	-
1999 5/.....	1,965	7	4	3	172	514	741	398	84	28	7	2	5

Hispanic, male

2003.....	1,800	3	3	1	194	421	606	422	104	28	13	5	-
2002 5/.....	1,647	4	-	1	172	371	601	385	81	18	13	1	-
2001 5/.....	1,335	-	3	1	119	320	504	291	69	18	6	2	2
2000 5/.....	1,348	2	2	-	121	367	521	265	52	11	3	4	-
1999 5/.....	1,605	4	4	2	142	431	606	320	69	21	1	1	4

Hispanic, female

2003.....	558	1	2	6	49	112	183	148	31	10	11	5	-
2002 5/.....	490	1	4	3	45	85	176	120	30	18	7	1	-
2001 5/.....	396	-	1	4	43	88	140	82	19	11	6	2	-
2000 5/.....	352	-	-	4	27	82	130	67	27	9	5	1	-
1999 5/.....	360	3	-	1	30	83	135	78	15	7	6	1	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

Non-Hispanic 1/, both sexes

2003.....	26,199	21	30	34	2,371	4,287	8,321	7,546	2,087	711	511	273	7
2002 5/.....	23,756	24	26	38	1,989	3,973	8,029	6,639	1,705	608	461	257	7
2001 5/.....	19,799	22	22	40	1,642	3,313	6,969	5,367	1,252	520	426	221	5
2000 5/.....	17,835	16	12	39	1,410	3,053	6,509	4,578	1,091	462	422	235	8
1999 5/.....	16,966	14	21	27	1,174	3,046	6,279	4,168	1,052	524	458	197	6

Non-Hispanic 1/, male

2003.....	16,497	14	18	19	1,758	2,921	5,231	4,689	1,146	350	238	107	6
2002 5/.....	14,978	16	9	23	1,445	2,751	5,089	4,107	948	285	200	100	5
2001 5/.....	12,778	14	18	21	1,224	2,304	4,532	3,459	673	272	184	72	5
2000 5/.....	11,656	10	10	24	1,038	2,133	4,334	3,041	565	222	184	88	7
1999 5/.....	11,136	10	8	14	835	2,132	4,249	2,768	600	261	183	72	4

Non-Hispanic 1/, female

2003.....	9,702	7	12	15	613	1,366	3,090	2,857	941	361	273	166	1
2002 5/.....	8,778	8	17	15	544	1,222	2,940	2,532	757	323	261	157	2
2001 5/.....	7,021	8	4	19	418	1,009	2,437	1,908	579	248	242	149	-
2000 5/.....	6,179	6	2	15	372	920	2,175	1,537	526	240	238	147	1
1999 5/.....	5,830	4	13	13	339	914	2,030	1,400	452	263	275	125	2

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/													
Non-Hispanic white, both sexes													
2003.....	22,245	12	17	25	2,208	3,739	6,994	6,158	1,730	612	488	258	4
2002 5/.....	19,949	9	14	32	1,819	3,429	6,722	5,328	1,385	542	428	236	5
2001 5/.....	16,367	10	13	33	1,484	2,835	5,730	4,190	1,019	456	394	198	5
2000 5/.....	14,585	9	5	33	1,266	2,554	5,264	3,532	900	410	399	209	4
1999 5/.....	13,644	6	10	22	1,019	2,493	5,001	3,179	840	461	433	178	2
Non-Hispanic white, male													
2003.....	13,959	11	11	16	1,651	2,578	4,414	3,767	900	286	225	97	3
2002 5/.....	12,478	7	3	19	1,325	2,393	4,276	3,230	708	241	179	94	3
2001 5/.....	10,465	4	12	18	1,112	1,992	3,750	2,614	498	232	164	64	5
2000 5/.....	9,439	5	3	21	944	1,790	3,525	2,284	428	185	175	76	3
1999 5/.....	8,831	5	4	12	727	1,750	3,401	2,040	431	223	169	68	1
Non-Hispanic white, female													
2003.....	8,286	1	6	9	557	1,161	2,580	2,391	830	326	263	161	1
2002 5/.....	7,471	2	11	13	494	1,036	2,446	2,098	677	301	249	142	2
2001 5/.....	5,902	6	1	15	372	843	1,980	1,576	521	224	230	134	-
2000 5/.....	5,146	4	2	12	322	764	1,739	1,248	472	225	224	133	1
1999 5/.....	4,813	1	6	10	292	743	1,600	1,139	409	238	264	110	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

Non-Hispanic black, both sexes

2003.....	3,466	9	13	8	112	431	1,181	1,278	315	82	21	13	3
2002 5/.....	3,404	13	10	5	127	451	1,181	1,229	301	48	25	13	1
2001 5/.....	3,099	11	8	5	112	404	1,140	1,108	214	52	25	20	-
2000 5/.....	2,977	6	5	4	113	440	1,143	990	178	48	22	24	4
1999 5/.....	3,030	8	11	4	123	488	1,170	940	190	55	20	17	4

Non-Hispanic black, male

2003.....	2,256	3	7	3	77	270	736	851	227	58	12	9	3
2002 5/.....	2,264	8	6	3	90	300	740	833	231	34	16	2	1
2001 5/.....	2,113	9	5	2	85	269	721	802	163	35	15	7	-
2000 5/.....	2,050	4	5	1	76	303	749	723	131	34	9	11	4
1999 5/.....	2,134	5	4	2	93	341	786	695	155	35	11	4	3

Non-Hispanic black, female

2003.....	1,210	6	6	5	35	161	445	427	88	24	9	4	-
2002 5/.....	1,140	5	4	2	37	151	441	396	70	14	9	11	-
2001 5/.....	986	2	3	3	27	135	419	306	51	17	10	13	-
2000 5/.....	927	2	-	3	37	137	394	267	47	14	13	13	-
1999 5/.....	896	3	7	2	30	147	384	245	35	20	9	13	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

Origin not stated 2/, both sexes

2003.....	166	-	-	-	5	25	48	58	17	6	2	-	5
2002 5/.....	147	-	-	-	9	20	52	38	12	9	3	1	3
2001 5/.....	175	-	-	-	9	18	56	56	21	4	1	1	9
2000 5/.....	185	-	1	1	9	25	46	76	17	4	-	-	6
1999 5/.....	197	-	-	-	5	19	75	58	25	4	2	2	7

Origin not stated 2/, male

2003.....	129	-	-	-	3	19	36	42	16	6	2	-	5
2002 5/.....	109	-	-	-	5	17	38	25	11	8	2	1	2
2001 5/.....	140	-	-	-	7	16	43	43	18	3	1	-	9
2000 5/.....	133	-	1	1	7	15	35	52	14	2	-	-	6
1999 5/.....	144	-	-	-	4	16	52	45	19	4	-	1	3

Origin not stated 2/, female

2003.....	37	-	-	-	2	6	12	16	1	-	-	-	-
2002 5/.....	38	-	-	-	4	3	14	13	1	1	1	-	1
2001 5/.....	35	-	-	-	2	2	13	13	3	1	-	1	-
2000 5/.....	52	-	-	-	2	10	11	24	3	2	-	-	-
1999 5/.....	53	-	-	-	1	3	23	13	6	-	2	1	4

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

All origins, both sexes

2003.....	30,136	8	48	324	6,846	6,188	5,118	4,550	2,804	1,912	1,731	589	18
2002.....	30,242	10	61	348	6,780	6,140	5,446	4,321	2,719	1,993	1,833	576	15
2001.....	29,573	11	70	333	6,687	6,137	5,288	4,120	2,544	1,998	1,803	563	19
2000.....	28,663	13	46	377	6,575	5,789	5,358	3,951	2,272	1,941	1,723	600	18
1999.....	28,874	8	65	416	6,795	5,975	5,234	3,722	2,306	2,034	1,738	561	20

All origins, male

2003.....	26,124	7	28	256	6,186	5,472	4,222	3,761	2,378	1,694	1,558	546	16
2002.....	26,098	4	38	242	6,096	5,347	4,503	3,579	2,296	1,768	1,671	540	14
2001.....	25,480	7	39	249	6,066	5,347	4,312	3,416	2,116	1,754	1,634	522	18
2000.....	24,582	9	28	302	5,906	5,003	4,363	3,258	1,896	1,684	1,571	549	13
1999.....	24,700	5	36	308	6,056	5,156	4,229	3,057	1,951	1,786	1,580	517	19

All origins, female

2003.....	4,012	1	20	68	660	716	896	789	426	218	173	43	2
2002.....	4,144	6	23	106	684	793	943	742	423	225	162	36	1
2001.....	4,093	4	31	84	621	790	976	704	428	244	169	41	1
2000.....	4,081	4	18	75	669	786	995	693	376	257	152	51	5
1999.....	4,174	3	29	108	739	819	1,005	665	355	248	158	44	1

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Hispanic, both sexes

2003.....	3,319	1	9	52	1,346	985	478	259	86	58	30	11	4
2002.....	3,143	2	15	56	1,260	920	459	220	99	62	33	13	4
2001.....	3,087	4	13	40	1,233	926	468	215	82	60	31	10	5
2000.....	2,891	1	7	55	1,184	788	414	255	83	64	28	9	3
1999.....	2,878	2	6	69	1,189	770	427	218	87	51	44	12	3

Hispanic, male

2003.....	2,998	1	5	38	1,234	914	411	220	80	54	28	10	3
2002.....	2,834	-	9	38	1,173	844	390	185	89	57	32	13	4
2001.....	2,774	3	6	31	1,132	841	408	180	75	55	28	10	5
2000.....	2,582	-	4	46	1,096	711	351	210	73	54	26	9	2
1999.....	2,549	2	4	50	1,078	699	364	173	76	46	42	12	3

Hispanic, female

2003.....	321	-	4	14	112	71	67	39	6	4	2	1	1
2002.....	309	2	6	18	87	76	69	35	10	5	1	-	-
2001.....	313	1	7	9	101	85	60	35	7	5	3	-	-
2000.....	309	1	3	9	88	77	63	45	10	10	2	-	1
1999.....	329	-	2	19	111	71	63	45	11	5	2	-	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Non-Hispanic 1/, both sexes

2003.....	26,710	7	39	272	5,468	5,183	4,617	4,281	2,709	1,849	1,700	576	9
2002.....	26,944	8	46	290	5,476	5,187	4,963	4,080	2,609	1,922	1,795	561	7
2001.....	26,341	7	57	292	5,409	5,182	4,802	3,895	2,446	1,927	1,767	549	8
2000.....	25,637	12	39	320	5,355	4,980	4,915	3,679	2,179	1,869	1,694	589	6
1999.....	25,877	6	59	347	5,587	5,174	4,782	3,491	2,210	1,978	1,689	547	7

Non-Hispanic 1/, male

2003.....	23,036	6	23	218	4,925	4,540	3,794	3,533	2,289	1,637	1,529	534	8
2002.....	23,127	4	29	202	4,882	4,477	4,093	3,375	2,198	1,702	1,634	525	6
2001.....	22,573	4	33	217	4,894	4,479	3,889	3,228	2,025	1,688	1,601	508	7
2000.....	21,881	9	24	255	4,777	4,274	3,984	3,033	1,813	1,624	1,544	539	5
1999.....	22,050	3	32	258	4,964	4,432	3,842	2,874	1,866	1,735	1,533	504	7

Non-Hispanic 1/, female

2003.....	3,674	1	16	54	543	643	823	748	420	212	171	42	1
2002.....	3,817	4	17	88	594	710	870	705	411	220	161	36	1
2001.....	3,768	3	24	75	515	703	913	667	421	239	166	41	1
2000.....	3,756	3	15	65	578	706	931	646	366	245	150	50	1
1999.....	3,827	3	27	89	623	742	940	617	344	243	156	43	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/													
Non-Hispanic white, both sexes													
2003.....	18,457	1	15	141	2,289	2,552	3,404	3,645	2,483	1,731	1,627	563	6
2002.....	18,762	2	28	163	2,356	2,639	3,662	3,457	2,380	1,812	1,714	544	5
2001.....	18,676	4	30	203	2,373	2,847	3,601	3,361	2,233	1,797	1,685	536	6
2000.....	18,042	5	21	219	2,377	2,702	3,694	3,106	1,965	1,738	1,637	574	4
1999.....	18,260	2	27	225	2,504	2,949	3,624	2,951	1,976	1,852	1,614	533	3
Non-Hispanic white, male													
2003.....	15,670	-	7	119	2,009	2,141	2,754	2,997	2,101	1,541	1,471	524	6
2002.....	15,881	2	16	114	2,038	2,169	2,998	2,855	2,005	1,608	1,562	510	4
2001.....	15,760	2	19	152	2,059	2,383	2,910	2,788	1,838	1,577	1,531	496	5
2000.....	15,160	2	11	175	2,050	2,232	2,965	2,560	1,629	1,508	1,498	527	3
1999.....	15,384	2	15	168	2,165	2,453	2,893	2,440	1,659	1,626	1,468	492	3
Non-Hispanic white, female													
2003.....	2,787	1	8	22	280	411	650	648	382	190	156	39	-
2002.....	2,881	-	12	49	318	470	664	602	375	204	152	34	1
2001.....	2,916	2	11	51	314	464	691	573	395	220	154	40	1
2000.....	2,882	3	10	44	327	470	729	546	336	230	139	47	1
1999.....	2,876	-	12	57	339	496	731	511	317	226	146	41	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Non-Hispanic black, both sexes

2003.....	7,566	6	23	116	2,968	2,460	1,088	544	186	103	58	12	2
2002.....	7,494	6	17	102	2,904	2,393	1,172	535	186	95	65	17	2
2001.....	7,063	2	23	78	2,872	2,172	1,092	464	171	108	69	11	1
2000.....	6,958	7	13	91	2,798	2,121	1,093	496	162	114	50	12	1
1999.....	6,933	2	26	105	2,882	2,034	1,037	457	195	109	68	14	4

Non-Hispanic black, male

2003.....	6,794	6	15	86	2,739	2,253	935	464	156	83	47	9	1
2002.....	6,681	2	12	74	2,657	2,182	991	450	156	82	58	15	2
2001.....	6,323	1	13	60	2,690	1,965	894	387	150	93	59	10	1
2000.....	6,193	7	8	74	2,574	1,910	920	411	136	103	40	9	1
1999.....	6,114	1	12	78	2,623	1,826	854	376	174	95	59	12	4

Non-Hispanic black, female

2003.....	772	-	8	30	229	207	153	80	30	20	11	3	1
2002.....	813	4	5	28	247	211	181	85	30	13	7	2	-
2001.....	740	1	10	18	182	207	198	77	21	15	10	1	-
2000.....	765	-	5	17	224	211	173	85	26	11	10	3	-
1999.....	819	1	14	27	259	208	183	81	21	14	9	2	-

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Origin not stated 2/, both sexes

2003.....	107	-	-	-	32	20	23	10	9	5	1	2	5
2002.....	155	-	-	2	44	33	24	21	11	9	5	2	4
2001.....	145	-	-	1	45	29	18	10	16	11	5	4	6
2000.....	135	-	-	2	36	21	29	17	10	8	1	2	9
1999.....	119	-	-	-	19	31	25	13	9	5	5	2	10

Origin not stated 2/, male

2003.....	90	-	-	-	27	18	17	8	9	3	1	2	5
2002.....	137	-	-	2	41	26	20	19	9	9	5	2	4
2001.....	133	-	-	1	40	27	15	8	16	11	5	4	6
2000.....	119	-	-	1	33	18	28	15	10	6	1	1	6
1999.....	101	-	-	-	14	25	23	10	9	5	5	1	9

Worktable Orig290F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these states were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age not stated
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------	----------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Origin not stated 2/, female

2003.....	17	-	-	-	5	2	6	2	-	2	-	-	-
2002.....	18	-	-	-	3	7	4	2	2	-	-	-	-
2001.....	12	-	-	-	5	2	3	2	-	-	-	-	-
2000.....	16	-	-	1	3	3	1	2	-	2	-	1	3
1999.....	18	-	-	-	5	6	2	3	-	-	-	1	1

- Quantity zero.

... Category not applicable.

1/ Includes races other than white and black.

2/ Includes deaths for which Hispanic origin was not reported on the death certificate.

3/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

4/ Included in selected categories.

5/ Figures may differ from those previously shown as they are based on a modified list of causes; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003."

Source: CDC/NCHS, National Vital Statistics System, Mortality.