

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All causes

All origins 3/, both sexes

2003.....	841.9	700.0	31.5	14.7	19.1	66.4	96.4	95.2	111.3	159.1	241.3
2002.....	847.3	695.0	31.2	15.2	19.5	67.8	95.2	94.7	111.6	161.3	242.6
2001.....	848.5	683.4	33.3	15.3	19.2	66.9	95.0	96.2	113.5	165.9	240.5
2000.....	854.0	736.7	32.4	15.8	20.3	67.1	93.6	91.2	111.0	159.2	239.1
1999.....	857.0	736.0	34.2	16.9	20.4	68.6	90.8	90.4	113.4	160.8	236.3

All origins 3/, male

2003.....	840.3	777.4	35.1	16.2	23.1	92.3	140.3	133.5	148.9	202.7	304.1
2002.....	846.6	761.5	35.2	16.7	23.1	94.0	140.8	134.4	149.4	206.6	306.4
2001.....	846.4	749.8	37.0	16.7	22.8	93.7	141.1	135.6	151.2	213.7	304.8
2000.....	853.0	806.5	35.9	17.6	24.2	93.3	138.0	128.8	147.9	205.4	305.9
1999.....	859.2	805.0	37.9	18.6	24.4	94.7	133.1	127.0	152.0	207.6	304.0

All origins 3/, female

2003.....	843.4	619.1	27.8	13.0	15.0	39.0	49.8	55.4	73.0	115.3	179.2
2002.....	848.0	625.3	27.0	13.5	15.9	40.1	47.3	53.6	73.3	116.1	179.7
2001.....	850.4	613.9	29.5	13.9	15.3	38.5	46.8	55.8	75.2	118.2	177.2
2000.....	855.0	663.4	28.7	14.0	16.1	39.3	47.1	52.8	73.6	113.2	173.4
1999.....	854.9	663.6	30.3	15.0	16.2	41.0	46.6	53.2	74.4	114.4	169.8

Hispanic, both sexes

2003.....	305.8	610.1	30.2	14.0	17.3	67.2	85.7	81.3	86.6	116.0	178.5
2002.....	302.2	592.7	29.8	14.4	16.6	65.2	87.9	74.7	84.2	115.5	181.8
2001.....	306.8	572.7	30.6	13.0	16.6	63.1	86.9	80.6	88.7	129.5	185.3
2000.....	303.8	596.3	29.6	14.1	17.5	61.6	83.3	76.7	93.2	129.4	192.2
1999.....	305.7	583.8	30.9	13.3	18.3	61.0	81.1	77.5	94.1	132.9	189.4

Hispanic, male

2003.....	330.7	665.5	33.8	16.0	20.9	98.5	124.6	115.7	120.7	153.8	232.5
2002.....	328.7	644.0	34.2	15.9	19.0	97.0	128.9	108.7	116.5	151.4	242.6
2001.....	332.9	624.4	33.8	14.9	18.6	92.0	128.2	115.2	121.1	175.5	249.0
2000.....	331.3	637.1	31.5	15.0	21.3	90.5	123.1	110.3	131.1	172.5	259.1
1999.....	332.6	623.4	32.9	14.9	21.2	87.8	119.1	111.1	130.8	183.0	255.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All causes

Hispanic, female

2003.....	279.3	552.4	26.3	12.0	13.6	33.5	37.4	39.1	47.1	73.4	120.5
2002.....	274.0	539.1	25.3	12.9	14.0	30.8	37.2	33.3	46.9	75.6	117.3
2001.....	279.0	518.9	27.2	11.0	14.5	31.1	36.1	39.4	51.3	79.0	118.3
2000.....	274.6	553.6	27.5	13.2	13.6	28.7	34.6	37.5	49.8	82.4	122.2
1999.....	277.2	542.3	28.7	11.6	15.2	30.5	35.3	38.0	52.1	78.3	120.4

Non-Hispanic 4/, both sexes

2003.....	924.4	716.2	31.7	14.7	19.5	65.9	98.3	98.2	116.0	165.7	248.7
2002.....	928.8	713.0	31.4	15.3	20.1	67.9	96.3	99.1	116.6	168.0	249.2
2001.....	926.2	703.5	33.8	15.8	19.6	67.3	96.2	99.2	117.5	170.6	245.8
2000.....	929.6	760.4	32.9	16.1	20.7	67.7	95.3	93.8	113.5	162.4	242.8
1999.....	929.9	760.7	34.8	17.5	20.7	69.7	92.4	92.5	116.1	163.7	239.9

Non-Hispanic 4/, male

2003.....	922.9	798.1	35.2	16.2	23.5	90.6	143.5	137.6	154.5	210.6	312.7
2002.....	928.0	782.6	35.4	16.8	23.7	93.0	142.8	140.6	155.7	215.0	313.3
2001.....	923.6	773.2	37.6	17.0	23.6	93.6	143.3	139.8	156.2	218.5	309.6
2000.....	928.1	836.8	36.8	18.1	24.7	93.4	140.9	132.2	150.3	208.8	309.0
1999.....	932.2	834.9	38.8	19.3	24.9	95.8	135.7	129.6	154.9	209.7	307.1

Non-Hispanic 4/, female

2003.....	925.9	630.7	28.0	13.2	15.2	39.9	52.0	58.8	77.8	121.6	186.1
2002.....	929.5	640.1	27.2	13.7	16.2	41.6	49.1	57.8	77.8	121.9	186.7
2001.....	928.6	630.4	29.9	14.4	15.4	39.7	48.7	58.8	79.1	123.4	183.5
2000.....	931.0	680.1	28.8	14.0	16.4	41.0	49.3	55.6	77.1	116.8	178.2
1999.....	927.8	682.9	30.6	15.7	16.3	42.5	48.6	55.7	77.7	118.5	174.2

Non-Hispanic White, both sexes

2003.....	993.6	576.5	27.6	13.3	17.6	62.9	88.9	86.2	104.4	149.5	227.3
2002.....	997.5	575.9	27.1	13.7	18.1	65.5	86.6	86.5	103.9	151.1	228.1
2001.....	991.1	555.4	30.1	14.5	18.0	63.4	85.4	87.2	104.4	153.1	220.9
2000.....	993.2	596.5	28.5	14.7	19.4	64.0	84.1	80.6	98.7	144.5	216.4
1999.....	990.7	593.1	30.1	15.6	19.1	65.3	80.2	80.3	101.3	144.5	211.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All causes

Non-Hispanic White, male

2003.....	979.1	647.2	30.2	14.7	21.2	84.3	128.3	119.2	139.4	192.2	289.4
2002.....	983.9	643.5	30.3	14.9	21.3	87.6	126.7	121.2	139.3	194.0	290.9
2001.....	975.6	611.6	33.7	15.2	21.7	86.1	125.6	122.4	139.3	197.9	282.4
2000.....	978.5	658.7	32.4	16.6	23.3	86.1	122.9	112.7	132.2	188.0	278.5
1999.....	979.6	658.1	33.4	17.0	22.7	87.1	116.1	112.1	135.7	187.1	272.8

Non-Hispanic White, female

2003.....	1,007.6	502.5	24.9	11.9	13.8	40.4	48.3	52.4	68.9	106.6	165.4
2002.....	1,010.6	504.8	23.8	12.3	14.8	42.0	45.6	51.3	68.1	108.0	165.4
2001.....	1,006.1	496.4	26.3	13.7	14.1	39.6	44.3	51.4	69.0	108.2	159.7
2000.....	1,007.3	530.9	24.4	12.6	15.2	41.0	44.5	48.0	64.7	100.8	154.5
1999.....	1,001.3	524.6	26.7	14.1	15.2	42.4	43.3	48.2	66.5	101.7	149.2

Non-Hispanic Black, both sexes

2003.....	788.8	1,290.0	48.0	20.6	26.7	82.3	152.2	175.4	201.8	285.1	421.1
2002.....	792.8	1,280.3	48.4	22.0	28.3	83.8	151.7	179.6	207.4	292.2	422.8
2001.....	798.1	1,354.4	49.5	21.4	26.6	88.7	156.7	174.9	207.1	294.6	438.1
2000.....	805.5	1,477.8	51.7	22.6	27.5	89.9	160.5	175.2	213.6	290.5	443.1
1999.....	812.1	1,510.2	54.6	26.2	29.0	94.8	160.6	169.8	212.0	301.4	457.4

Non-Hispanic Black, male

2003.....	840.6	1,427.7	55.1	22.8	32.6	124.3	232.0	259.9	273.1	357.4	520.3
2002.....	842.3	1,369.3	55.8	24.8	34.4	124.1	235.3	266.6	281.5	376.4	514.5
2001.....	849.7	1,484.2	55.1	24.6	32.5	133.9	244.7	255.7	279.6	375.1	538.9
2000.....	859.5	1,623.2	56.1	25.4	33.0	134.0	249.2	256.9	283.2	371.1	558.6
1999.....	872.8	1,650.8	61.5	30.1	36.3	142.3	249.7	247.5	288.0	382.5	583.2

Non-Hispanic Black, female

2003.....	741.6	1,146.9	40.7	18.4	20.6	39.1	73.3	97.5	137.5	220.8	333.8
2002.....	748.0	1,187.5	40.8	19.2	22.0	42.1	70.5	100.1	141.0	217.4	342.0
2001.....	751.2	1,217.8	43.6	18.1	20.5	42.0	72.2	101.4	142.2	223.2	349.0
2000.....	756.7	1,327.7	47.1	19.7	21.7	44.9	76.5	101.3	151.1	219.2	341.0
1999.....	757.3	1,365.4	47.4	22.1	21.5	46.3	76.1	99.5	144.0	229.5	346.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Meningococcal infection (A39)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.1	*	*	*	*	*	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	*	*	*	*	*
1999.....	0.1	0.8	0.2	*	*	0.2	*	*	*	*	*

Non-Hispanic 4/, male

2003.....	0.1	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.3	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, female

2003.....	0.1	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.3	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*	*

Non-Hispanic White, both sexes

2003.....	0.0	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	*	*	*	*	*
1999.....	0.1	1.0	0.3	*	*	0.2	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Septicemia (A40-A41)											
Hispanic, female											
2003.....	4.1	5.0	*	*	*	*	*	*	*	*	1.9
2002.....	4.1	7.6	*	*	*	*	*	*	*	*	*
2001.....	3.9	5.2	*	*	*	*	*	*	*	*	1.7
2000.....	3.5	7.7	*	*	*	*	*	*	*	*	1.9
1999.....	3.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	12.9	7.2	0.6	0.2	0.2	0.3	0.5	0.7	0.9	1.6	2.7
2002.....	12.9	7.1	0.5	0.2	0.3	0.2	0.4	0.7	1.0	1.6	2.4
2001.....	12.4	8.1	0.7	0.1	0.2	0.3	0.4	0.6	1.0	1.5	2.3
2000.....	12.1	6.9	0.7	0.2	0.1	0.2	0.3	0.5	0.9	1.5	2.5
1999.....	12.0	7.9	0.6	0.2	0.2	0.2	0.6	0.6	0.9	1.4	2.3
Non-Hispanic 4/, male											
2003.....	11.6	7.6	0.8	*	*	0.4	0.5	0.7	1.0	1.6	2.6
2002.....	11.6	7.3	0.5	0.2	0.3	*	0.4	0.7	1.0	1.5	2.5
2001.....	11.2	8.8	0.9	*	*	0.3	0.5	0.6	0.9	1.6	2.6
2000.....	10.7	6.7	0.7	0.2	*	0.2	0.4	0.7	1.0	1.5	2.6
1999.....	10.7	9.4	0.6	*	*	*	0.6	0.6	1.0	1.5	2.6
Non-Hispanic 4/, female											
2003.....	14.2	6.8	0.4	0.3	*	0.3	0.5	0.8	0.9	1.5	2.8
2002.....	14.1	6.8	0.5	*	0.3	0.3	0.3	0.7	0.9	1.6	2.4
2001.....	13.5	7.4	0.6	*	*	0.3	0.4	0.5	1.1	1.3	2.1
2000.....	13.5	7.0	0.6	*	*	*	*	0.4	0.9	1.5	2.4
1999.....	13.2	6.2	0.5	0.3	*	*	0.5	0.6	0.8	1.3	1.9
Non-Hispanic White, both sexes											
2003.....	12.9	4.2	0.5	0.2	0.2	0.2	0.4	0.5	0.7	1.2	2.1
2002.....	12.9	5.5	0.5	0.2	0.2	0.2	0.3	0.6	0.8	1.3	2.0
2001.....	12.3	5.1	0.7	*	*	0.2	0.3	0.4	0.7	1.1	1.8
2000.....	11.9	4.8	0.5	0.2	*	0.2	0.2	0.4	0.7	1.1	2.0
1999.....	11.7	5.3	0.5	0.2	*	0.2	0.4	0.5	0.7	1.0	1.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Septicemia (A40-A41)

Non-Hispanic White, male

2003.....	11.4	3.9	0.6	*	*	0.3	0.4	0.5	0.7	1.3	2.0
2002.....	11.5	5.6	0.5	*	*	*	0.4	0.6	0.7	1.2	2.0
2001.....	11.0	5.8	0.9	*	*	*	*	0.5	0.6	1.2	2.1
2000.....	10.4	4.7	0.5	*	*	*	0.3	0.4	0.7	1.2	2.1
1999.....	10.2	6.7	0.5	*	*	*	0.6	0.5	0.8	1.1	1.9

Non-Hispanic White, female

2003.....	14.3	4.5	*	*	*	*	0.5	0.6	0.7	1.1	2.2
2002.....	14.2	5.4	*	*	*	0.3	*	0.5	0.8	1.3	2.0
2001.....	13.6	4.4	0.5	*	*	*	*	0.4	0.8	0.9	1.5
2000.....	13.4	4.8	0.6	*	*	*	*	*	0.6	1.0	1.9
1999.....	13.1	3.8	0.5	*	*	*	*	0.4	0.6	0.9	1.5

Non-Hispanic Black, both sexes

2003.....	16.8	18.8	1.0	*	*	*	0.8	1.7	2.3	3.7	6.5
2002.....	16.8	13.1	*	*	*	*	*	1.6	2.3	3.5	5.6
2001.....	16.4	20.3	0.9	*	*	*	0.8	1.1	2.7	3.9	5.7
2000.....	16.7	16.6	1.1	*	*	*	0.8	1.5	2.6	3.8	5.9
1999.....	16.6	18.6	*	*	*	*	1.0	1.4	1.8	3.8	6.4

Non-Hispanic Black, male

2003.....	15.8	21.7	*	*	*	*	*	1.6	2.9	3.8	6.5
2002.....	15.5	13.9	*	*	*	*	*	*	2.5	3.8	6.1
2001.....	15.4	21.4	*	*	*	*	*	*	2.6	3.9	5.9
2000.....	15.3	16.4	*	*	*	*	*	2.2	2.6	3.6	6.0
1999.....	15.7	20.8	*	*	*	*	*	*	2.2	4.1	7.9

Non-Hispanic Black, female

2003.....	17.7	15.8	*	*	*	*	*	1.7	1.7	3.7	6.4
2002.....	18.0	12.3	*	*	*	*	*	2.0	2.1	3.1	5.2
2001.....	17.2	19.1	*	*	*	*	*	*	2.7	3.9	5.6
2000.....	17.9	16.9	*	*	*	*	*	*	2.6	4.1	5.8
1999.....	17.4	16.4	*	*	*	*	*	1.7	1.5	3.6	5.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Viral hepatitis (B15-B19)

All origins 3/, both sexes

2003.....	1.9	*	*	*	*	*	*	0.1	0.2	0.8	2.1
2002.....	2.0	*	*	*	*	*	*	0.2	0.3	1.0	2.3
2001.....	2.0	*	*	*	*	*	*	*	0.3	0.9	2.6
2000.....	1.9	*	*	*	*	*	*	0.1	0.3	1.0	2.5
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9	2.3

All origins 3/, male

2003.....	2.5	*	*	*	*	*	*	*	0.3	1.2	2.9
2002.....	2.7	*	*	*	*	*	*	0.2	0.4	1.4	3.2
2001.....	2.5	*	*	*	*	*	*	*	0.3	1.2	3.7
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.3	3.5
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.2	3.5

All origins 3/, female

2003.....	1.3	*	*	*	*	*	*	*	*	0.4	1.2
2002.....	1.4	*	*	*	*	*	*	*	*	0.6	1.4
2001.....	1.4	*	*	*	*	*	*	*	0.2	0.6	1.5
2000.....	1.4	*	*	*	*	*	*	*	0.2	0.6	1.5
1999.....	1.3	*	*	*	*	*	*	*	0.2	0.6	1.2

Hispanic, both sexes

2003.....	1.8	*	*	*	*	*	*	*	*	0.9	2.8
2002.....	2.0	*	*	*	*	*	*	*	*	1.0	2.7
2001.....	2.1	*	*	*	*	*	*	*	*	1.2	3.7
2000.....	2.0	*	*	*	*	*	*	*	*	1.4	3.8
1999.....	1.9	*	*	*	*	*	*	*	*	1.4	3.0

Hispanic, male

2003.....	2.3	*	*	*	*	*	*	*	*	1.2	4.2
2002.....	2.5	*	*	*	*	*	*	*	*	1.3	3.9
2001.....	2.6	*	*	*	*	*	*	*	*	1.5	5.2
2000.....	2.5	*	*	*	*	*	*	*	*	2.0	5.9
1999.....	2.3	*	*	*	*	*	*	*	*	2.1	4.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Viral hepatitis (B15-B19)

Hispanic, female

2003.....	1.3	*	*	*	*	*	*	*	*	*	*
2002.....	1.4	*	*	*	*	*	*	*	*	*	*
2001.....	1.7	*	*	*	*	*	*	*	*	*	2.1
2000.....	1.5	*	*	*	*	*	*	*	*	*	*
1999.....	1.5	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	1.9	*	*	*	*	*	*	*	0.2	0.8	2.0
2002.....	2.0	*	*	*	*	*	*	0.2	0.3	1.0	2.2
2001.....	1.9	*	*	*	*	*	*	*	0.3	0.9	2.5
2000.....	1.9	*	*	*	*	*	*	0.1	0.2	0.9	2.3
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.8	2.2

Non-Hispanic 4/, male

2003.....	2.5	*	*	*	*	*	*	*	0.3	1.2	2.8
2002.....	2.7	*	*	*	*	*	*	*	0.4	1.4	3.1
2001.....	2.5	*	*	*	*	*	*	*	0.4	1.1	3.5
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.2	3.2
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.1	3.3

Non-Hispanic 4/, female

2003.....	1.3	*	*	*	*	*	*	*	*	0.4	1.2
2002.....	1.4	*	*	*	*	*	*	*	*	0.6	1.4
2001.....	1.4	*	*	*	*	*	*	*	*	0.6	1.5
2000.....	1.4	*	*	*	*	*	*	*	*	0.6	1.5
1999.....	1.2	*	*	*	*	*	*	*	0.3	0.6	1.2

Non-Hispanic White, both sexes

2003.....	1.8	*	*	*	*	*	*	*	0.2	0.8	1.9
2002.....	1.9	*	*	*	*	*	*	*	0.2	1.0	2.1
2001.....	1.8	*	*	*	*	*	*	*	0.2	0.9	2.4
2000.....	1.8	*	*	*	*	*	*	*	0.2	0.9	2.2
1999.....	1.6	*	*	*	*	*	*	*	0.3	0.8	2.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Human immunodeficiency virus (HIV) disease (B20-B24)

All origins 3/, both sexes

2003.....	4.7	*	*	*	0.1	0.2	0.7	2.3	5.5	11.2	12.8
2002.....	4.9	*	*	*	0.1	0.2	0.7	2.6	6.4	11.9	13.4
2001.....	5.0	*	*	*	0.1	0.2	0.9	3.1	7.3	12.3	13.7
2000.....	5.1	*	*	*	0.1	0.2	0.8	3.3	8.7	12.8	13.5
1999.....	5.3	*	0.2	0.2	0.1	0.2	0.9	3.9	9.5	13.0	14.6

All origins 3/, male

2003.....	7.1	*	*	*	*	*	0.8	2.8	7.3	15.9	19.1
2002.....	7.4	*	*	*	*	*	0.7	3.0	8.5	17.7	19.8
2001.....	7.6	*	*	*	*	0.2	0.8	3.7	10.2	18.4	20.5
2000.....	7.9	*	*	*	*	*	0.8	3.8	11.9	19.2	20.4
1999.....	8.2	*	*	0.2	*	*	0.8	5.1	13.6	19.5	22.6

All origins 3/, female

2003.....	2.4	*	*	*	*	*	0.7	1.8	3.7	6.5	6.6
2002.....	2.5	*	*	*	*	0.2	0.6	2.3	4.2	6.2	7.2
2001.....	2.5	*	*	*	*	0.2	1.0	2.5	4.4	6.3	7.0
2000.....	2.5	*	*	*	*	*	0.7	2.9	5.5	6.4	6.7
1999.....	2.5	*	*	*	*	*	1.0	2.7	5.4	6.6	6.9

Hispanic, both sexes

2003.....	4.7	*	*	*	*	*	0.6	2.0	4.7	9.8	15.2
2002.....	4.8	*	*	*	*	*	0.6	2.1	5.1	11.2	16.1
2001.....	5.0	*	*	*	*	*	0.8	2.3	6.6	11.9	16.3
2000.....	5.4	*	*	*	*	*	*	2.9	8.5	13.5	16.5
1999.....	5.8	*	*	*	*	*	0.9	3.5	9.6	15.5	19.7

Hispanic, male

2003.....	7.1	*	*	*	*	*	*	2.6	6.7	14.2	21.9
2002.....	7.2	*	*	*	*	*	*	2.9	7.4	16.9	23.9
2001.....	7.6	*	*	*	*	*	*	2.9	9.4	17.5	24.9
2000.....	8.2	*	*	*	*	*	*	3.5	12.6	21.0	24.9
1999.....	8.7	*	*	*	*	*	*	4.9	13.9	23.8	29.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Human immunodeficiency virus (HIV) disease (B20-B24)

Hispanic, female

2003.....	2.2	*	*	*	*	*	*	1.2	2.4	4.8	7.9
2002.....	2.2	*	*	*	*	*	*	1.2	2.4	4.8	7.8
2001.....	2.3	*	*	*	*	*	*	1.5	3.5	5.8	7.3
2000.....	2.4	*	*	*	*	*	*	2.2	3.9	5.4	7.6
1999.....	2.6	*	*	*	*	*	*	1.9	4.6	6.4	9.7

Non-Hispanic 4/, both sexes

2003.....	4.6	*	*	*	0.1	0.1	0.7	2.4	5.7	11.3	12.3
2002.....	4.8	*	*	*	*	0.2	0.7	2.7	6.6	11.9	12.9
2001.....	4.9	*	*	*	*	0.2	1.0	3.2	7.4	12.3	13.2
2000.....	5.0	*	*	*	0.1	0.2	0.8	3.4	8.7	12.5	12.9
1999.....	5.2	*	*	0.2	0.1	0.2	0.9	3.9	9.5	12.6	13.9

Non-Hispanic 4/, male

2003.....	7.0	*	*	*	*	*	0.7	2.8	7.4	16.1	18.5
2002.....	7.3	*	*	*	*	*	0.7	3.0	8.7	17.6	19.0
2001.....	7.5	*	*	*	*	*	0.8	3.8	10.3	18.4	19.6
2000.....	7.7	*	*	*	*	*	0.8	3.8	11.7	18.6	19.5
1999.....	8.0	*	*	*	*	*	0.8	5.1	13.4	18.7	21.4

Non-Hispanic 4/, female

2003.....	2.4	*	*	*	*	*	0.7	2.0	3.9	6.6	6.4
2002.....	2.5	*	*	*	*	*	0.7	2.5	4.5	6.3	7.0
2001.....	2.5	*	*	*	*	*	1.1	2.6	4.4	6.3	6.9
2000.....	2.5	*	*	*	*	*	0.8	3.0	5.7	6.4	6.4
1999.....	2.5	*	*	*	*	*	1.0	2.8	5.5	6.6	6.5

Non-Hispanic White, both sexes

2003.....	2.1	*	*	*	*	*	0.2	0.5	2.4	5.1	6.0
2002.....	2.1	*	*	*	*	*	*	0.9	2.4	5.7	5.8
2001.....	2.2	*	*	*	*	*	0.2	0.9	3.0	6.0	6.1
2000.....	2.3	*	*	*	*	*	0.2	1.1	3.5	6.3	6.1
1999.....	2.4	*	*	*	*	*	0.2	1.5	4.4	6.5	6.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic White, male

2003.....	3.6	*	*	*	*	*	*	0.9	3.7	8.4	10.2
2002.....	3.7	*	*	*	*	*	*	1.3	3.8	9.6	9.9
2001.....	3.8	*	*	*	*	*	*	1.3	4.9	10.2	10.5
2000.....	4.0	*	*	*	*	*	*	1.5	5.6	10.6	10.2
1999.....	4.2	*	*	*	*	*	*	2.2	7.3	10.9	10.9

Non-Hispanic White, female

2003.....	0.6	*	*	*	*	*	*	*	1.0	1.8	1.8
2002.....	0.6	*	*	*	*	*	*	0.5	1.0	1.8	1.7
2001.....	0.6	*	*	*	*	*	*	0.5	1.2	1.7	1.7
2000.....	0.7	*	*	*	*	*	*	0.7	1.5	2.1	1.9
1999.....	0.7	*	*	*	*	*	*	0.8	1.5	2.0	1.8

Non-Hispanic Black, both sexes

2003.....	20.1	*	*	*	*	0.7	3.3	11.6	24.3	47.5	53.3
2002.....	21.3	*	*	*	*	0.8	3.4	12.2	30.2	49.3	58.6
2001.....	21.6	*	*	*	*	1.1	4.6	15.2	32.3	50.4	59.7
2000.....	22.0	*	*	*	*	0.7	3.9	15.8	38.4	51.0	58.2
1999.....	22.3	*	*	0.8	*	0.7	4.2	17.2	39.1	51.3	63.5

Non-Hispanic Black, male

2003.....	28.5	*	*	*	*	*	3.3	13.2	30.0	63.9	75.9
2002.....	30.3	*	*	*	*	*	3.5	12.2	38.0	69.9	81.3
2001.....	30.8	*	*	*	*	*	3.7	17.6	43.5	71.6	84.0
2000.....	31.7	*	*	*	*	*	3.8	17.1	49.4	73.0	85.3
1999.....	32.6	*	*	*	*	*	3.5	21.8	52.2	71.7	96.1

Non-Hispanic Black, female

2003.....	12.6	*	*	*	*	*	3.3	10.2	19.1	33.0	33.4
2002.....	13.2	*	*	*	*	*	3.4	12.3	23.2	31.1	38.6
2001.....	13.3	*	*	*	*	*	5.4	13.0	22.3	31.5	38.2
2000.....	13.1	*	*	*	*	*	4.0	14.6	28.5	31.5	34.3
1999.....	13.0	*	*	*	*	*	4.9	13.1	27.4	33.2	34.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Hispanic, female

2003.....	1.3	*	*	*	*	*	*	*	*	*	*
2002.....	1.2	*	*	*	*	*	*	*	*	*	*
2001.....	1.0	5.0	*	*	*	*	*	*	*	*	*
2000.....	1.0	*	*	*	*	*	*	*	*	*	*
1999.....	1.4	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	2.7	4.8	0.5	0.1	*	0.2	0.2	0.3	0.4	0.7	1.2
2002.....	2.4	5.7	0.4	0.2	0.2	0.2	0.3	0.3	0.4	0.8	1.2
2001.....	2.1	4.6	0.5	0.2	0.1	0.2	0.2	0.3	0.4	0.7	0.9
2000.....	2.1	4.7	0.4	0.1	0.2	0.2	0.2	0.3	0.4	0.7	0.8
1999.....	2.8	5.5	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.0	1.4

Non-Hispanic 4/, male

2003.....	2.9	4.8	0.5	*	*	0.3	0.2	0.3	0.5	0.8	1.5
2002.....	2.7	5.9	0.5	*	*	*	0.3	0.4	0.4	0.9	1.6
2001.....	2.2	4.8	0.5	*	*	*	0.3	0.3	0.5	0.8	1.2
2000.....	2.2	5.1	0.5	*	*	0.3	*	0.3	0.4	0.9	1.0
1999.....	2.8	6.2	0.6	*	0.2	0.2	0.3	0.4	0.8	1.2	1.6

Non-Hispanic 4/, female

2003.....	2.5	4.8	0.5	*	*	*	*	0.3	0.3	0.6	0.8
2002.....	2.2	5.4	*	0.3	*	*	*	*	0.4	0.6	0.8
2001.....	2.1	4.5	0.4	*	*	*	*	0.3	0.4	0.6	0.6
2000.....	2.0	4.3	0.4	*	*	*	*	0.3	0.4	0.5	0.6
1999.....	2.7	4.7	0.4	*	*	*	*	0.5	0.6	0.8	1.1

Non-Hispanic White, both sexes

2003.....	2.8	3.8	0.4	*	*	0.2	0.2	0.3	0.4	0.6	1.1
2002.....	2.5	4.8	0.4	0.2	0.2	0.2	0.2	0.2	0.3	0.6	1.0
2001.....	2.1	3.3	0.4	0.2	*	0.2	0.2	0.2	0.3	0.6	0.8
2000.....	2.1	3.5	0.4	*	0.2	0.2	0.2	0.2	0.3	0.5	0.7
1999.....	2.7	4.4	0.5	0.2	*	0.2	0.2	0.3	0.5	0.7	1.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic White, male

2003.....	2.9	3.6	*	*	*	*	*	*	0.4	0.7	1.4
2002.....	2.7	4.8	0.4	*	*	*	*	*	*	0.7	1.3
2001.....	2.2	3.3	0.4	*	*	*	*	*	0.4	0.6	1.0
2000.....	2.2	3.9	*	*	*	*	*	*	*	0.6	0.8
1999.....	2.7	5.5	0.6	*	*	*	*	*	0.5	1.0	1.3

Non-Hispanic White, female

2003.....	2.6	4.0	*	*	*	*	*	*	*	0.5	0.8
2002.....	2.3	4.9	*	*	*	*	*	*	0.3	0.4	0.6
2001.....	2.1	3.4	*	*	*	*	*	*	*	0.5	0.5
2000.....	2.0	3.1	*	*	*	*	*	*	0.3	0.4	0.5
1999.....	2.6	3.3	*	*	*	*	*	0.4	0.4	0.5	0.7

Non-Hispanic Black, both sexes

2003.....	2.6	8.4	0.9	*	*	*	*	*	0.8	1.5	2.1
2002.....	2.6	9.0	*	*	*	*	*	*	1.1	1.9	2.5
2001.....	2.3	10.0	*	*	*	*	*	*	1.1	1.5	1.8
2000.....	2.3	10.4	*	*	*	*	*	*	1.3	1.7	1.7
1999.....	3.6	10.0	1.0	*	*	*	*	1.2	2.2	2.6	3.6

Non-Hispanic Black, male

2003.....	3.0	9.3	*	*	*	*	*	*	*	1.6	2.7
2002.....	3.0	10.0	*	*	*	*	*	*	*	2.1	3.1
2001.....	2.5	11.5	*	*	*	*	*	*	*	1.7	2.6
2000.....	2.6	10.6	*	*	*	*	*	*	*	2.3	2.2
1999.....	3.7	10.0	*	*	*	*	*	*	2.7	2.7	3.6

Non-Hispanic Black, female

2003.....	2.2	7.4	*	*	*	*	*	*	*	*	1.5
2002.....	2.2	7.9	*	*	*	*	*	*	*	1.7	1.9
2001.....	2.1	8.4	*	*	*	*	*	*	*	*	*
2000.....	2.1	10.2	*	*	*	*	*	*	*	*	*
1999.....	3.5	10.0	*	*	*	*	*	*	1.7	2.5	3.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

All origins 3/, both sexes

2003.....	191.5	1.9	2.5	2.6	2.6	3.4	4.6	6.8	11.8	23.0	46.1
2002.....	193.2	1.8	2.6	2.7	2.5	3.5	5.0	6.6	12.5	23.5	47.5
2001.....	194.4	1.6	2.7	2.4	2.5	3.6	4.9	7.3	12.6	24.6	48.7
2000.....	196.5	2.4	2.7	2.4	2.6	3.7	5.1	7.0	12.5	24.5	48.8
1999.....	197.0	1.8	2.7	2.5	2.5	3.7	5.3	7.3	12.5	25.2	49.4

All origins 3/, male

2003.....	201.3	1.7	2.8	2.9	2.7	4.2	5.1	6.8	10.9	19.3	41.7
2002.....	203.8	2.0	2.7	2.9	2.8	4.0	5.7	7.0	11.3	19.8	42.8
2001.....	205.3	1.5	2.9	2.4	2.6	4.3	5.7	7.2	11.2	21.0	44.0
2000.....	207.2	2.6	3.0	2.6	2.9	4.3	5.9	7.1	11.3	21.2	44.4
1999.....	208.9	1.9	2.9	2.6	2.7	4.3	6.1	7.6	11.6	21.6	45.5

All origins 3/, female

2003.....	182.0	2.1	2.1	2.3	2.5	2.5	4.2	6.8	12.6	26.7	50.5
2002.....	183.0	1.6	2.4	2.5	2.2	3.0	4.2	6.2	13.7	27.2	52.2
2001.....	183.9	1.8	2.5	2.4	2.3	2.9	4.1	7.4	14.0	28.1	53.5
2000.....	186.2	2.3	2.5	2.2	2.2	3.0	4.2	7.0	13.7	27.8	53.2
1999.....	185.6	1.7	2.5	2.3	2.3	3.0	4.4	7.0	13.4	28.7	53.3

Hispanic, both sexes

2003.....	60.3	*	3.0	2.9	2.9	3.8	4.4	5.5	8.8	16.2	30.7
2002.....	59.7	*	2.3	2.8	3.0	3.8	5.1	4.7	9.1	15.8	31.9
2001.....	60.5	*	2.8	2.4	2.7	4.5	4.1	5.7	8.9	18.3	34.4
2000.....	59.9	*	2.5	2.5	3.4	3.8	4.5	5.8	9.0	18.2	33.9
1999.....	59.6	*	2.5	2.5	3.0	3.1	5.1	5.8	9.1	17.3	33.3

Hispanic, male

2003.....	61.5	*	3.6	3.6	2.8	5.4	4.6	5.5	8.2	12.7	24.8
2002.....	61.2	*	2.2	2.7	3.1	4.9	5.5	5.0	7.8	11.9	26.3
2001.....	62.2	*	2.9	2.6	2.4	5.7	5.4	5.2	7.5	15.1	29.9
2000.....	61.3	*	2.9	3.0	4.1	4.3	4.8	5.7	8.2	13.8	28.0
1999.....	61.2	*	2.3	2.5	3.1	3.4	5.8	5.6	7.6	13.9	27.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

Hispanic, female

2003.....	59.1	*	2.4	2.2	3.0	2.1	4.1	5.4	9.4	20.0	37.1
2002.....	58.1	*	2.4	2.9	2.9	2.6	4.5	4.4	10.6	20.2	37.9
2001.....	58.7	*	2.7	2.1	3.0	3.2	2.7	6.4	10.4	21.8	39.2
2000.....	58.5	*	2.2	2.0	2.6	3.4	4.1	5.8	9.9	23.0	40.0
1999.....	58.0	*	2.7	2.5	2.9	2.7	4.2	6.1	10.7	20.9	39.6

Non-Hispanic 4/, both sexes

2003.....	211.9	2.0	2.3	2.5	2.6	3.3	4.7	7.1	12.3	24.1	48.1
2002.....	213.4	1.7	2.6	2.7	2.4	3.5	4.9	7.1	13.1	24.7	49.4
2001.....	213.9	1.6	2.7	2.4	2.4	3.4	5.1	7.7	13.3	25.4	50.3
2000.....	215.6	2.5	2.7	2.3	2.4	3.6	5.2	7.2	13.1	25.3	50.3
1999.....	215.5	1.7	2.7	2.5	2.4	3.8	5.3	7.6	13.0	26.2	51.0

Non-Hispanic 4/, male

2003.....	224.3	1.9	2.6	2.7	2.7	3.9	5.2	7.1	11.4	20.5	44.0
2002.....	226.7	1.9	2.8	3.0	2.7	3.9	5.7	7.6	12.0	21.1	44.9
2001.....	227.2	1.6	2.9	2.4	2.7	4.0	5.8	7.7	11.9	21.8	45.5
2000.....	228.7	2.6	2.9	2.5	2.6	4.3	6.2	7.3	11.8	22.3	46.2
1999.....	229.9	1.6	3.0	2.7	2.6	4.5	6.1	8.1	12.3	22.7	47.4

Non-Hispanic 4/, female

2003.....	200.0	2.1	2.1	2.3	2.4	2.6	4.2	7.2	13.2	27.7	52.1
2002.....	200.8	1.5	2.4	2.4	2.1	3.1	4.1	6.6	14.2	28.2	53.8
2001.....	201.2	1.6	2.5	2.5	2.2	2.8	4.4	7.6	14.6	29.0	55.0
2000.....	203.1	2.4	2.6	2.2	2.2	3.0	4.2	7.2	14.3	28.3	54.4
1999.....	201.9	1.8	2.5	2.3	2.2	3.1	4.5	7.1	13.7	29.7	54.6

Non-Hispanic White, both sexes

2003.....	229.5	2.2	2.4	2.7	2.5	3.4	4.7	7.0	11.9	23.2	45.8
2002.....	230.9	1.6	2.6	2.8	2.4	3.5	5.1	6.9	12.7	23.9	47.2
2001.....	230.6	1.8	2.9	2.5	2.3	3.4	5.1	7.4	12.8	24.5	47.7
2000.....	232.2	2.6	2.8	2.4	2.4	3.5	5.1	7.3	12.7	24.0	46.9
1999.....	231.5	1.6	3.0	2.4	2.3	3.7	5.1	7.6	12.6	24.9	48.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

Non-Hispanic White, male

2003.....	241.8	2.1	2.6	2.9	2.5	4.0	5.2	7.3	11.2	20.6	42.0
2002.....	243.8	2.0	2.8	2.9	2.6	3.9	5.9	7.4	11.8	20.5	43.2
2001.....	243.4	1.7	3.1	2.3	2.4	4.0	5.8	7.7	12.0	21.2	43.2
2000.....	244.4	2.9	3.1	2.5	2.6	4.1	6.1	7.4	11.8	21.5	42.9
1999.....	245.1	1.7	3.3	2.7	2.5	4.2	5.8	8.1	12.0	22.1	44.8

Non-Hispanic White, female

2003.....	217.6	2.3	2.2	2.4	2.4	2.7	4.1	6.7	12.6	26.0	49.6
2002.....	218.5	*	2.3	2.5	2.2	3.0	4.3	6.5	13.5	27.2	51.2
2001.....	218.4	1.8	2.8	2.6	2.2	2.8	4.4	7.2	13.7	27.7	52.2
2000.....	220.6	2.4	2.4	2.4	2.2	2.9	4.1	7.1	13.5	26.5	50.9
1999.....	218.4	*	2.7	2.1	2.1	3.0	4.3	7.1	13.1	27.7	51.5

Non-Hispanic Black, both sexes

2003.....	170.0	*	2.3	2.4	3.0	3.0	5.2	8.9	16.2	32.1	68.2
2002.....	171.5	*	3.0	2.6	2.6	3.6	4.6	8.4	17.0	32.7	70.0
2001.....	172.8	*	2.0	2.4	2.9	3.7	5.3	9.6	17.0	33.7	72.2
2000.....	174.9	*	2.8	2.2	2.4	4.5	6.1	8.3	16.3	35.4	76.1
1999.....	176.9	*	2.0	2.5	3.1	4.6	6.9	8.8	16.4	36.9	74.9

Non-Hispanic Black, male

2003.....	184.8	*	2.4	2.6	3.3	3.6	5.8	7.3	13.9	22.5	62.7
2002.....	187.8	*	2.7	3.3	3.0	4.2	5.2	8.6	14.7	26.5	61.9
2001.....	190.8	*	2.4	2.6	3.4	4.3	6.1	8.9	12.9	27.6	64.3
2000.....	194.9	*	2.6	2.5	2.7	5.5	7.4	7.6	13.3	28.3	71.8
1999.....	197.4	*	2.0	2.4	3.4	5.6	7.7	9.2	14.5	28.6	69.9

Non-Hispanic Black, female

2003.....	156.5	*	2.1	2.2	2.6	2.4	4.5	10.4	18.4	40.7	73.0
2002.....	156.7	*	3.2	1.9	2.2	3.1	4.0	8.3	19.0	38.2	77.1
2001.....	156.5	*	*	2.2	2.3	3.0	4.5	10.3	20.8	39.2	79.2
2000.....	156.9	*	3.0	1.8	2.0	3.6	4.8	8.9	18.9	41.8	79.8
1999.....	158.3	*	2.0	2.7	2.7	3.5	6.1	8.4	18.1	44.3	79.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of esophagus (C15)

Hispanic, female

2003.....	0.5	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*
2001.....	0.5	*	*	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	4.9	*	*	*	*	*	*	*	*	0.4	1.0
2002.....	4.9	*	*	*	*	*	*	*	*	0.4	1.0
2001.....	4.9	*	*	*	*	*	*	*	0.1	0.3	1.1
2000.....	4.8	*	*	*	*	*	*	*	0.2	0.4	1.1
1999.....	4.7	*	*	*	*	*	*	*	0.1	0.3	1.0

Non-Hispanic 4/, male

2003.....	7.8	*	*	*	*	*	*	*	*	0.7	1.6
2002.....	7.8	*	*	*	*	*	*	*	*	0.7	1.7
2001.....	7.7	*	*	*	*	*	*	*	*	0.5	1.9
2000.....	7.5	*	*	*	*	*	*	*	0.3	0.7	1.9
1999.....	7.4	*	*	*	*	*	*	*	0.2	0.6	1.8

Non-Hispanic 4/, female

2003.....	2.2	*	*	*	*	*	*	*	*	*	0.4
2002.....	2.2	*	*	*	*	*	*	*	*	*	0.4
2001.....	2.2	*	*	*	*	*	*	*	*	*	0.4
2000.....	2.3	*	*	*	*	*	*	*	*	*	0.2
1999.....	2.2	*	*	*	*	*	*	*	*	*	0.3

Non-Hispanic White, both sexes

2003.....	5.3	*	*	*	*	*	*	*	*	0.5	1.0
2002.....	5.3	*	*	*	*	*	*	*	*	0.5	1.0
2001.....	5.1	*	*	*	*	*	*	*	*	0.4	1.1
2000.....	5.1	*	*	*	*	*	*	*	0.2	0.4	1.0
1999.....	4.9	*	*	*	*	*	*	*	0.2	0.4	0.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of stomach (C16)

Hispanic, female

2003.....	2.8	*	*	*	*	*	*	*	*	1.4	1.9
2002.....	2.7	*	*	*	*	*	*	*	*	2.0	1.8
2001.....	2.8	*	*	*	*	*	*	*	*	*	2.2
2000.....	2.7	*	*	*	*	*	*	*	*	1.6	2.8
1999.....	2.7	*	*	*	*	*	*	*	*	1.5	2.2

Non-Hispanic 4/, both sexes

2003.....	4.3	*	*	*	*	*	*	0.1	0.4	0.7	1.3
2002.....	4.4	*	*	*	*	*	*	0.2	0.4	0.6	1.1
2001.....	4.5	*	*	*	*	*	*	*	0.4	0.7	1.1
2000.....	4.7	*	*	*	*	*	*	0.2	0.4	0.6	1.3
1999.....	4.7	*	*	*	*	*	*	0.2	0.4	0.8	1.3

Non-Hispanic 4/, male

2003.....	5.2	*	*	*	*	*	*	*	0.4	0.8	1.5
2002.....	5.3	*	*	*	*	*	*	0.3	0.4	0.7	1.3
2001.....	5.4	*	*	*	*	*	*	*	0.5	0.8	1.2
2000.....	5.5	*	*	*	*	*	*	*	0.4	0.8	1.6
1999.....	5.7	*	*	*	*	*	*	*	0.4	0.9	1.6

Non-Hispanic 4/, female

2003.....	3.5	*	*	*	*	*	*	*	0.4	0.7	1.0
2002.....	3.5	*	*	*	*	*	*	*	0.4	0.5	0.9
2001.....	3.6	*	*	*	*	*	*	*	0.3	0.6	1.0
2000.....	3.8	*	*	*	*	*	*	*	0.4	0.4	1.0
1999.....	3.8	*	*	*	*	*	*	*	0.3	0.7	1.1

Non-Hispanic White, both sexes

2003.....	4.0	*	*	*	*	*	*	*	0.3	0.6	1.0
2002.....	4.1	*	*	*	*	*	*	0.2	0.3	0.5	0.8
2001.....	4.2	*	*	*	*	*	*	*	0.3	0.5	0.8
2000.....	4.4	*	*	*	*	*	*	*	0.3	0.4	0.9
1999.....	4.4	*	*	*	*	*	*	*	0.3	0.6	0.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Hispanic, female

2003.....	5.7	*	*	*	*	*	*	*	*	1.7	1.8
2002.....	5.7	*	*	*	*	*	*	*	*	1.4	3.0
2001.....	5.9	*	*	*	*	*	*	*	*	*	2.9
2000.....	5.8	*	*	*	*	*	*	*	*	2.0	3.5
1999.....	5.9	*	*	*	*	*	*	*	*	*	2.7

Non-Hispanic 4/, both sexes

2003.....	21.3	*	*	*	*	*	0.1	0.5	1.0	2.1	4.0
2002.....	21.7	*	*	*	*	*	0.2	0.5	1.1	2.1	4.4
2001.....	22.0	*	*	*	*	*	0.2	0.5	1.2	2.4	4.2
2000.....	22.4	*	*	*	*	*	0.1	0.4	1.0	2.2	4.0
1999.....	22.4	*	*	*	*	*	0.2	0.5	1.0	2.0	4.2

Non-Hispanic 4/, male

2003.....	21.7	*	*	*	*	*	*	0.6	1.1	2.3	4.4
2002.....	22.3	*	*	*	*	*	0.3	0.5	1.2	2.2	5.2
2001.....	22.3	*	*	*	*	*	*	0.4	1.3	2.7	4.4
2000.....	22.8	*	*	*	*	*	*	0.5	1.1	2.5	4.6
1999.....	22.7	*	*	*	*	*	*	0.7	1.1	2.1	4.3

Non-Hispanic 4/, female

2003.....	20.9	*	*	*	*	*	*	0.4	1.0	2.0	3.6
2002.....	21.2	*	*	*	*	*	*	0.4	0.9	2.0	3.7
2001.....	21.7	*	*	*	*	*	*	0.5	1.0	2.0	3.9
2000.....	22.1	*	*	*	*	*	*	0.4	0.8	1.8	3.5
1999.....	22.2	*	*	*	*	*	*	0.3	0.9	1.9	4.1

Non-Hispanic White, both sexes

2003.....	22.7	*	*	*	*	*	*	0.5	1.0	1.9	3.5
2002.....	23.2	*	*	*	*	*	0.2	0.5	1.1	2.0	4.0
2001.....	23.5	*	*	*	*	*	0.2	0.5	1.1	2.2	3.8
2000.....	24.0	*	*	*	*	*	*	0.4	0.9	1.9	3.6
1999.....	24.0	*	*	*	*	*	0.2	0.4	0.9	1.9	3.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic White, male

2003.....	23.2	*	*	*	*	*	*	0.6	1.0	2.1	3.9
2002.....	23.8	*	*	*	*	*	0.3	0.6	1.2	2.0	4.7
2001.....	23.9	*	*	*	*	*	*	0.4	1.3	2.6	4.0
2000.....	24.2	*	*	*	*	*	*	0.4	1.1	2.3	4.1
1999.....	24.3	*	*	*	*	*	*	0.6	1.0	2.1	4.1

Non-Hispanic White, female

2003.....	22.2	*	*	*	*	*	*	0.4	1.0	1.8	3.1
2002.....	22.7	*	*	*	*	*	*	0.4	1.0	1.9	3.3
2001.....	23.1	*	*	*	*	*	*	0.5	0.9	1.9	3.7
2000.....	23.8	*	*	*	*	*	*	0.4	0.6	1.5	3.2
1999.....	23.7	*	*	*	*	*	*	*	0.9	1.7	3.6

Non-Hispanic Black, both sexes

2003.....	18.8	*	*	*	*	*	*	*	1.2	3.5	6.8
2002.....	18.9	*	*	*	*	*	*	*	1.0	2.7	7.4
2001.....	19.2	*	*	*	*	*	*	*	1.6	3.4	6.6
2000.....	19.4	*	*	*	*	*	*	*	1.6	3.8	6.7
1999.....	19.3	*	*	*	*	*	*	0.8	1.3	3.2	6.7

Non-Hispanic Black, male

2003.....	19.0	*	*	*	*	*	*	*	*	3.4	7.4
2002.....	19.3	*	*	*	*	*	*	*	*	3.3	8.6
2001.....	19.0	*	*	*	*	*	*	*	1.9	4.0	7.6
2000.....	19.8	*	*	*	*	*	*	*	1.8	4.2	8.2
1999.....	19.2	*	*	*	*	*	*	*	*	3.1	6.5

Non-Hispanic Black, female

2003.....	18.7	*	*	*	*	*	*	*	*	3.6	6.3
2002.....	18.4	*	*	*	*	*	*	*	*	2.2	6.4
2001.....	19.4	*	*	*	*	*	*	*	*	3.0	5.7
2000.....	19.1	*	*	*	*	*	*	*	1.4	3.5	5.3
1999.....	19.5	*	*	*	*	*	*	*	*	3.2	6.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of pancreas (C25)

Hispanic, female

2003.....	3.8	*	*	*	*	*	*	*	*	*	*
2002.....	3.7	*	*	*	*	*	*	*	*	*	*
2001.....	3.9	*	*	*	*	*	*	*	*	*	*
2000.....	3.6	*	*	*	*	*	*	*	*	*	*
1999.....	3.5	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	11.7	*	*	*	*	*	*	*	0.2	0.8	1.8
2002.....	11.6	*	*	*	*	*	*	*	0.2	0.7	2.0
2001.....	11.5	*	*	*	*	*	*	*	0.2	0.8	1.8
2000.....	11.4	*	*	*	*	*	*	*	0.2	0.6	1.9
1999.....	11.4	*	*	*	*	*	*	*	0.2	0.8	1.9

Non-Hispanic 4/, male

2003.....	11.7	*	*	*	*	*	*	*	0.2	1.0	2.1
2002.....	11.7	*	*	*	*	*	*	*	0.2	0.8	2.5
2001.....	11.4	*	*	*	*	*	*	*	0.3	0.9	2.0
2000.....	11.3	*	*	*	*	*	*	*	0.2	0.8	2.2
1999.....	11.4	*	*	*	*	*	*	*	0.3	1.0	2.5

Non-Hispanic 4/, female

2003.....	11.6	*	*	*	*	*	*	*	*	0.5	1.5
2002.....	11.5	*	*	*	*	*	*	*	*	0.7	1.5
2001.....	11.5	*	*	*	*	*	*	*	*	0.6	1.7
2000.....	11.4	*	*	*	*	*	*	*	*	0.5	1.6
1999.....	11.4	*	*	*	*	*	*	*	*	0.6	1.4

Non-Hispanic White, both sexes

2003.....	12.5	*	*	*	*	*	*	*	0.2	0.7	1.6
2002.....	12.4	*	*	*	*	*	*	*	0.2	0.7	2.0
2001.....	12.3	*	*	*	*	*	*	*	0.2	0.8	1.8
2000.....	12.2	*	*	*	*	*	*	*	0.2	0.6	1.8
1999.....	12.1	*	*	*	*	*	*	*	0.2	0.7	1.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Hispanic, female

2003.....	7.8	*	*	*	*	*	*	*	*	*	2.1
2002.....	7.6	*	*	*	*	*	*	*	*	*	2.3
2001.....	7.4	*	*	*	*	*	*	*	*	*	2.7
2000.....	7.2	*	*	*	*	*	*	*	*	*	2.3
1999.....	7.1	*	*	*	*	*	*	*	*	*	2.3

Non-Hispanic 4/, both sexes

2003.....	61.2	*	*	*	*	*	*	0.1	0.6	2.7	9.3
2002.....	61.4	*	*	*	*	*	*	0.1	0.7	3.0	10.0
2001.....	61.3	*	*	*	*	*	*	0.2	0.6	3.2	10.2
2000.....	61.5	*	*	*	*	*	*	0.2	0.8	3.2	10.0
1999.....	60.5	*	*	*	*	*	*	0.2	0.7	3.2	9.9

Non-Hispanic 4/, male

2003.....	71.1	*	*	*	*	*	*	*	0.7	2.7	10.4
2002.....	71.8	*	*	*	*	*	*	*	0.7	3.0	10.5
2001.....	72.6	*	*	*	*	*	*	*	0.6	3.1	11.3
2000.....	73.2	*	*	*	*	*	*	0.3	0.8	3.6	11.3
1999.....	72.7	*	*	*	*	*	*	0.3	0.8	3.1	11.2

Non-Hispanic 4/, female

2003.....	51.8	*	*	*	*	*	*	*	0.6	2.7	8.2
2002.....	51.6	*	*	*	*	*	*	*	0.6	2.9	9.6
2001.....	50.5	*	*	*	*	*	*	*	0.7	3.3	9.0
2000.....	50.4	*	*	*	*	*	*	*	0.9	2.8	8.7
1999.....	48.8	*	*	*	*	*	*	*	0.6	3.2	8.7

Non-Hispanic White, both sexes

2003.....	67.5	*	*	*	*	*	*	*	0.7	2.7	9.1
2002.....	67.7	*	*	*	*	*	*	*	0.6	3.0	9.7
2001.....	67.3	*	*	*	*	*	*	0.2	0.7	3.2	9.8
2000.....	67.4	*	*	*	*	*	*	0.2	0.9	3.0	9.4
1999.....	65.9	*	*	*	*	*	*	0.2	0.7	3.0	9.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic White, male

2003.....	77.3	*	*	*	*	*	*	*	0.7	2.6	9.9
2002.....	77.8	*	*	*	*	*	*	*	0.7	2.9	9.8
2001.....	78.4	*	*	*	*	*	*	*	0.6	3.0	10.5
2000.....	78.9	*	*	*	*	*	*	*	0.8	3.2	10.0
1999.....	78.1	*	*	*	*	*	*	*	0.8	2.9	10.4

Non-Hispanic White, female

2003.....	58.1	*	*	*	*	*	*	*	0.6	2.7	8.3
2002.....	57.9	*	*	*	*	*	*	*	0.5	3.1	9.5
2001.....	56.6	*	*	*	*	*	*	*	0.7	3.4	9.1
2000.....	56.4	*	*	*	*	*	*	*	0.9	2.7	8.7
1999.....	54.3	*	*	*	*	*	*	*	0.6	3.1	8.6

Non-Hispanic Black, both sexes

2003.....	44.5	*	*	*	*	*	*	*	*	3.0	12.8
2002.....	44.7	*	*	*	*	*	*	*	*	3.5	14.5
2001.....	44.6	*	*	*	*	*	*	*	*	3.6	14.9
2000.....	45.4	*	*	*	*	*	*	*	0.9	5.0	16.1
1999.....	45.6	*	*	*	*	*	*	*	*	4.7	14.9

Non-Hispanic Black, male

2003.....	57.0	*	*	*	*	*	*	*	*	3.0	16.1
2002.....	58.1	*	*	*	*	*	*	*	*	4.4	17.5
2001.....	59.2	*	*	*	*	*	*	*	*	4.0	18.8
2000.....	60.3	*	*	*	*	*	*	*	*	6.0	21.9
1999.....	60.9	*	*	*	*	*	*	*	*	4.7	19.1

Non-Hispanic Black, female

2003.....	33.1	*	*	*	*	*	*	*	*	2.9	9.9
2002.....	32.6	*	*	*	*	*	*	*	*	2.7	11.9
2001.....	31.3	*	*	*	*	*	*	*	*	3.3	11.4
2000.....	31.9	*	*	*	*	*	*	*	*	4.1	10.9
1999.....	31.8	*	*	*	*	*	*	*	*	4.7	11.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Malignant melanoma of skin (C43)											
Hispanic, female											
2003.....	0.4	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	3.0	*	*	*	*	*	0.2	0.4	0.7	1.0	1.8
2002.....	2.9	*	*	*	*	*	0.2	0.4	0.7	1.1	1.8
2001.....	3.0	*	*	*	*	*	0.2	0.4	0.6	1.1	1.9
2000.....	3.0	*	*	*	*	*	0.2	0.4	0.7	1.1	1.9
1999.....	2.9	*	*	*	*	*	0.2	0.4	0.7	1.2	1.9
Non-Hispanic 4/, male											
2003.....	4.0	*	*	*	*	*	*	0.5	0.9	1.2	2.1
2002.....	3.8	*	*	*	*	*	*	0.5	0.9	1.2	2.3
2001.....	3.9	*	*	*	*	*	*	0.5	0.7	1.3	2.4
2000.....	3.7	*	*	*	*	*	*	0.4	0.9	1.1	2.3
1999.....	3.7	*	*	*	*	*	*	0.5	0.8	1.5	2.4
Non-Hispanic 4/, female											
2003.....	2.2	*	*	*	*	*	*	0.3	0.6	0.8	1.4
2002.....	2.1	*	*	*	*	*	*	0.4	0.5	1.0	1.3
2001.....	2.1	*	*	*	*	*	*	0.4	0.6	0.8	1.5
2000.....	2.2	*	*	*	*	*	*	0.3	0.6	1.1	1.5
1999.....	2.1	*	*	*	*	*	*	0.3	0.7	0.8	1.4
Non-Hispanic White, both sexes											
2003.....	3.8	*	*	*	*	*	0.2	0.5	0.9	1.3	2.2
2002.....	3.6	*	*	*	*	*	0.2	0.6	0.9	1.4	2.2
2001.....	3.7	*	*	*	*	*	0.2	0.6	0.8	1.3	2.4
2000.....	3.6	*	*	*	*	*	0.3	0.4	0.9	1.3	2.2
1999.....	3.5	*	*	*	*	*	0.2	0.5	0.9	1.4	2.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of breast (C50)

Hispanic, female

2003.....	9.8	*	*	*	*	*	*	*	2.7	5.8	12.9
2002.....	9.2	*	*	*	*	*	*	*	2.7	5.1	11.0
2001.....	9.7	*	*	*	*	*	*	*	2.5	7.3	11.5
2000.....	9.7	*	*	*	*	*	*	*	2.6	6.6	11.3
1999.....	9.3	*	*	*	*	*	*	*	2.4	5.7	11.4

Non-Hispanic 4/, both sexes

2003.....	15.9	*	*	*	*	*	*	0.3	1.8	4.5	8.1
2002.....	16.0	*	*	*	*	*	*	0.4	1.8	4.4	8.2
2001.....	16.1	*	*	*	*	*	*	0.5	1.9	4.4	8.5
2000.....	16.5	*	*	*	*	*	*	0.4	2.0	4.5	8.6
1999.....	16.3	*	*	*	*	*	*	0.4	1.8	4.5	8.1

Non-Hispanic 4/, male

2003.....	0.3	*	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, female

2003.....	30.8	*	*	*	*	*	*	0.6	3.6	8.9	15.9
2002.....	31.0	*	*	*	*	*	*	0.7	3.6	8.7	16.1
2001.....	31.1	*	*	*	*	*	*	0.9	3.8	8.8	16.7
2000.....	31.8	*	*	*	*	*	*	0.8	4.0	8.8	16.9
1999.....	31.4	*	*	*	*	*	*	0.9	3.6	8.9	16.0

Non-Hispanic White, both sexes

2003.....	16.8	*	*	*	*	*	*	*	1.5	4.0	7.2
2002.....	16.9	*	*	*	*	*	*	0.3	1.5	3.8	7.2
2001.....	17.0	*	*	*	*	*	*	0.4	1.6	3.9	7.5
2000.....	17.4	*	*	*	*	*	*	0.4	1.7	4.0	7.7
1999.....	17.1	*	*	*	*	*	*	0.4	1.5	3.9	7.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of breast (C50)

Non-Hispanic White, male

2003.....	0.3	*	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*

Non-Hispanic White, female

2003.....	32.6	*	*	*	*	*	*	*	3.1	8.0	14.3
2002.....	32.9	*	*	*	*	*	*	0.5	3.0	7.6	14.3
2001.....	32.9	*	*	*	*	*	*	0.7	3.2	7.8	15.0
2000.....	33.8	*	*	*	*	*	*	0.7	3.4	7.8	15.3
1999.....	33.3	*	*	*	*	*	*	0.7	3.1	7.8	14.3

Non-Hispanic Black, both sexes

2003.....	15.5	*	*	*	*	*	*	0.9	3.5	8.3	14.5
2002.....	15.4	*	*	*	*	*	*	0.9	3.8	8.4	15.6
2001.....	15.5	*	*	*	*	*	*	1.2	4.2	7.8	15.2
2000.....	15.3	*	*	*	*	*	*	0.8	3.8	8.0	15.2
1999.....	15.5	*	*	*	*	*	*	0.9	3.8	9.1	15.1

Non-Hispanic Black, male

2003.....	0.3	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*	*

Non-Hispanic Black, female

2003.....	29.4	*	*	*	*	*	*	1.7	6.7	15.6	26.9
2002.....	29.1	*	*	*	*	*	*	1.7	7.1	15.6	29.3
2001.....	29.2	*	*	*	*	*	*	2.3	7.8	14.7	28.3
2000.....	28.9	*	*	*	*	*	*	*	7.2	14.8	28.4
1999.....	29.2	*	*	*	*	*	*	1.8	7.1	16.8	28.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of cervix uteri (C53)

Hispanic, female

2003.....	2.0	*	*	*	*	*	*	*	*	2.5	3.6
2002.....	2.1	*	*	*	*	*	*	*	1.3	2.8	3.4
2001.....	2.3	*	*	*	*	*	*	*	1.7	3.0	5.0
2000.....	2.5	*	*	*	*	*	*	*	1.4	4.0	4.8
1999.....	2.3	*	*	*	*	*	*	*	1.7	2.6	4.9

Non-Hispanic 4/, both sexes

2003.....	1.4	*	*	*	*	*	*	0.3	0.7	1.2	1.7
2002.....	1.4	*	*	*	*	*	*	0.3	0.8	1.2	1.7
2001.....	1.5	*	*	*	*	*	*	0.3	0.8	1.3	1.8
2000.....	1.5	*	*	*	*	*	*	0.4	0.8	1.2	1.7
1999.....	1.6	*	*	*	*	*	*	0.3	0.7	1.3	1.8

Non-Hispanic 4/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2003.....	2.7	*	*	*	*	*	*	0.6	1.4	2.3	3.3
2002.....	2.8	*	*	*	*	*	*	0.5	1.6	2.4	3.4
2001.....	2.9	*	*	*	*	*	*	0.5	1.5	2.5	3.5
2000.....	3.0	*	*	*	*	*	*	0.8	1.5	2.4	3.4
1999.....	3.0	*	*	*	*	*	*	0.7	1.5	2.5	3.5

Non-Hispanic White, both sexes

2003.....	1.3	*	*	*	*	*	*	0.2	0.6	1.0	1.5
2002.....	1.3	*	*	*	*	*	*	0.2	0.8	1.1	1.5
2001.....	1.4	*	*	*	*	*	*	0.2	0.7	1.2	1.6
2000.....	1.4	*	*	*	*	*	*	0.4	0.7	1.1	1.4
1999.....	1.4	*	*	*	*	*	*	0.4	0.7	1.2	1.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic White, male											
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic White, female											
2003.....	2.5	*	*	*	*	*	*	0.5	1.3	2.0	3.1
2002.....	2.5	*	*	*	*	*	*	0.5	1.6	2.3	3.0
2001.....	2.7	*	*	*	*	*	*	0.5	1.4	2.4	3.2
2000.....	2.7	*	*	*	*	*	*	0.8	1.4	2.1	2.9
1999.....	2.8	*	*	*	*	*	*	0.8	1.5	2.3	3.2
Non-Hispanic Black, both sexes											
2003.....	2.2	*	*	*	*	*	*	*	1.2	2.1	2.5
2002.....	2.3	*	*	*	*	*	*	*	1.1	2.1	3.0
2001.....	2.2	*	*	*	*	*	*	*	1.2	1.9	3.0
2000.....	2.5	*	*	*	*	*	*	*	1.5	2.1	3.2
1999.....	2.4	*	*	*	*	*	*	*	1.0	2.1	3.0
Non-Hispanic Black, male											
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic Black, female											
2003.....	4.1	*	*	*	*	*	*	*	2.3	4.0	4.6
2002.....	4.3	*	*	*	*	*	*	*	2.1	4.0	5.7
2001.....	4.2	*	*	*	*	*	*	*	2.2	3.6	5.6
2000.....	4.7	*	*	*	*	*	*	*	2.8	4.0	6.0
1999.....	4.5	*	*	*	*	*	*	*	1.9	4.0	5.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasm of ovary (C56)

Hispanic, female

2003.....	3.3	*	*	*	*	*	*	*	*	*	*	2.3
2002.....	3.3	*	*	*	*	*	*	*	*	*	*	2.9
2001.....	3.4	*	*	*	*	*	*	*	*	*	*	2.5
2000.....	3.4	*	*	*	*	*	*	*	*	*	*	*
1999.....	2.9	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	5.6	*	*	*	*	*	*	0.2	0.3	0.6	1.4
2002.....	5.6	*	*	*	*	*	0.1	0.2	0.3	0.7	1.4
2001.....	5.6	*	*	*	*	*	*	0.2	0.4	0.7	1.4
2000.....	5.5	*	*	*	*	*	*	0.2	0.3	0.7	1.5
1999.....	5.3	*	*	*	*	*	*	0.1	0.3	0.7	1.6

Non-Hispanic 4/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2003.....	10.9	*	*	*	*	*	*	0.4	0.6	1.3	2.8
2002.....	11.0	*	*	*	*	*	0.3	0.4	0.6	1.5	2.7
2001.....	10.9	*	*	*	*	*	*	0.4	0.7	1.4	2.8
2000.....	10.7	*	*	*	*	*	*	0.4	0.7	1.3	3.0
1999.....	10.4	*	*	*	*	*	*	0.3	0.6	1.5	3.1

Non-Hispanic White, both sexes

2003.....	6.3	*	*	*	*	*	*	0.2	0.3	0.6	1.5
2002.....	6.3	*	*	*	*	*	0.2	0.2	0.3	0.8	1.4
2001.....	6.2	*	*	*	*	*	*	0.2	0.3	0.7	1.5
2000.....	6.2	*	*	*	*	*	*	0.2	0.3	0.7	1.6
1999.....	6.0	*	*	*	*	*	*	*	0.3	0.8	1.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of bladder (C67)

Hispanic, female

2003.....	0.7	*	*	*	*	*	*	*	*	*	*
2002.....	0.7	*	*	*	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	4.8	*	*	*	*	*	*	*	*	0.1	0.3
2002.....	4.9	*	*	*	*	*	*	*	*	0.2	0.4
2001.....	4.8	*	*	*	*	*	*	*	*	0.2	0.4
2000.....	4.7	*	*	*	*	*	*	*	*	0.2	0.3
1999.....	4.7	*	*	*	*	*	*	*	*	0.2	0.3

Non-Hispanic 4/, male

2003.....	6.8	*	*	*	*	*	*	*	*	*	0.5
2002.....	6.8	*	*	*	*	*	*	*	*	0.2	0.5
2001.....	6.7	*	*	*	*	*	*	*	*	*	0.5
2000.....	6.6	*	*	*	*	*	*	*	*	*	0.4
1999.....	6.6	*	*	*	*	*	*	*	*	*	0.4

Non-Hispanic 4/, female

2003.....	2.9	*	*	*	*	*	*	*	*	*	0.2
2002.....	3.1	*	*	*	*	*	*	*	*	*	0.3
2001.....	3.0	*	*	*	*	*	*	*	*	*	0.4
2000.....	3.0	*	*	*	*	*	*	*	*	*	0.3
1999.....	3.0	*	*	*	*	*	*	*	*	*	0.2

Non-Hispanic White, both sexes

2003.....	5.5	*	*	*	*	*	*	*	*	0.1	0.3
2002.....	5.6	*	*	*	*	*	*	*	*	0.2	0.4
2001.....	5.5	*	*	*	*	*	*	*	*	0.2	0.4
2000.....	5.4	*	*	*	*	*	*	*	*	0.2	0.4
1999.....	5.4	*	*	*	*	*	*	*	*	0.1	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Hispanic, female											
2003.....	1.6	*	*	*	*	*	*	*	*	*	*
2002.....	1.5	*	*	*	*	*	*	*	*	*	*
2001.....	1.6	*	*	*	*	*	*	*	*	*	*
2000.....	1.7	*	*	*	*	*	*	*	*	*	2.1
1999.....	1.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	4.9	*	0.7	0.9	0.8	0.6	0.6	0.9	1.3	2.1	3.1
2002.....	4.9	*	0.7	1.1	0.6	0.5	0.6	0.8	1.4	1.9	2.9
2001.....	4.8	*	0.6	0.9	0.7	0.5	0.7	0.8	1.5	1.9	2.9
2000.....	4.9	*	0.7	0.8	0.7	0.5	0.7	0.9	1.3	2.1	3.0
1999.....	5.0	*	0.7	0.9	0.6	0.7	0.6	1.0	1.5	2.0	2.9
Non-Hispanic 4/, male											
2003.....	5.5	*	0.7	0.8	0.8	0.7	0.7	1.0	1.5	2.5	3.7
2002.....	5.5	*	0.8	1.2	0.7	0.6	0.8	0.8	1.7	2.3	3.7
2001.....	5.5	*	0.6	0.8	0.7	0.6	0.7	1.0	1.9	2.3	3.7
2000.....	5.5	*	0.7	0.7	0.8	0.6	0.8	1.0	1.8	2.8	3.9
1999.....	5.6	*	0.7	0.9	0.6	0.8	0.7	1.1	1.9	2.5	3.5
Non-Hispanic 4/, female											
2003.....	4.3	*	0.6	0.9	0.8	0.5	0.6	0.9	1.0	1.7	2.5
2002.....	4.2	*	0.7	0.9	0.5	0.4	0.3	0.7	1.1	1.5	2.2
2001.....	4.2	*	0.6	1.0	0.6	0.4	0.6	0.5	1.1	1.4	2.1
2000.....	4.2	*	0.7	0.8	0.6	0.4	0.6	0.7	0.8	1.5	2.0
1999.....	4.4	*	0.7	0.9	0.7	0.5	0.4	0.8	1.0	1.6	2.4
Non-Hispanic White, both sexes											
2003.....	5.6	*	0.7	0.9	0.8	0.6	0.7	1.1	1.4	2.4	3.5
2002.....	5.6	*	0.6	1.1	0.6	0.6	0.6	0.8	1.6	2.2	3.3
2001.....	5.6	*	0.7	1.0	0.6	0.5	0.7	0.9	1.7	2.2	3.2
2000.....	5.6	*	0.7	0.8	0.7	0.5	0.7	1.0	1.4	2.4	3.2
1999.....	5.7	*	0.7	0.9	0.6	0.6	0.6	1.2	1.7	2.3	3.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All origins 3/, both sexes

2003.....	19.1	0.8	0.9	0.9	1.0	1.2	1.7	1.9	2.4	3.1	4.7
2002.....	19.5	0.7	0.8	0.8	1.0	1.5	1.9	1.9	2.6	3.2	4.6
2001.....	19.8	0.7	1.0	0.8	1.0	1.5	2.0	2.3	2.7	3.5	4.8
2000.....	20.1	1.0	0.9	0.8	1.0	1.6	2.1	2.1	2.5	3.4	4.8
1999.....	20.2	0.8	0.9	0.7	1.0	1.5	2.2	2.2	2.6	3.7	4.8

All origins 3/, male

2003.....	21.0	*	1.0	1.0	1.1	1.6	2.0	2.3	3.0	3.5	5.8
2002.....	21.4	*	0.8	0.9	1.2	1.8	2.2	2.3	3.0	3.7	5.5
2001.....	21.5	*	1.0	0.9	1.1	1.8	2.3	2.6	3.1	4.2	5.9
2000.....	21.6	1.1	1.0	0.9	1.2	1.9	2.6	2.3	3.0	3.9	5.7
1999.....	21.7	*	1.0	0.8	1.2	1.7	2.7	2.4	3.1	4.4	5.9

All origins 3/, female

2003.....	17.3	1.1	0.8	0.7	0.9	0.9	1.4	1.6	1.7	2.6	3.5
2002.....	17.7	*	0.7	0.8	0.8	1.2	1.7	1.5	2.2	2.6	3.7
2001.....	18.1	*	1.0	0.7	0.8	1.2	1.7	2.0	2.2	2.7	3.7
2000.....	18.6	*	0.9	0.7	0.8	1.2	1.5	1.9	1.9	3.0	3.9
1999.....	18.7	*	0.8	0.6	0.9	1.3	1.8	1.9	2.2	3.0	3.8

Hispanic, both sexes

2003.....	7.1	*	1.3	1.1	1.5	1.5	2.2	1.6	2.4	2.6	4.3
2002.....	7.3	*	0.9	1.2	1.3	1.9	2.3	1.7	2.2	3.3	4.5
2001.....	7.4	*	1.3	1.2	1.2	2.4	1.9	2.1	2.4	3.0	4.9
2000.....	7.5	*	1.0	0.8	1.4	1.7	2.2	1.9	1.9	3.0	5.0
1999.....	7.5	*	1.0	0.8	1.6	1.5	2.6	1.9	2.4	3.2	4.6

Hispanic, male

2003.....	7.4	*	1.5	1.6	1.3	2.3	2.4	1.9	2.9	3.3	5.3
2002.....	7.6	*	*	1.1	1.4	2.6	2.7	2.1	2.6	3.7	5.1
2001.....	8.3	*	1.4	1.5	1.2	3.1	2.4	2.3	3.1	3.8	6.5
2000.....	8.0	*	*	*	2.0	2.0	2.6	2.4	2.3	2.9	6.1
1999.....	7.6	*	*	*	1.5	1.4	2.8	1.9	2.5	4.0	4.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Hispanic, female

2003.....	6.8	*	*	*	1.7	*	2.0	1.2	1.9	1.9	3.3
2002.....	7.0	*	*	1.4	1.2	*	1.9	*	1.7	2.9	3.9
2001.....	6.5	*	*	*	1.3	1.7	1.3	1.8	1.6	2.2	3.2
2000.....	6.9	*	*	*	*	1.4	1.6	1.3	1.4	3.1	3.8
1999.....	7.4	*	*	*	1.6	1.5	2.2	1.9	2.2	2.4	4.7

Non-Hispanic 4/, both sexes

2003.....	21.0	0.8	0.8	0.8	0.9	1.2	1.6	2.0	2.4	3.1	4.7
2002.....	21.3	0.8	0.7	0.7	0.9	1.4	1.8	2.0	2.7	3.1	4.6
2001.....	21.6	0.6	0.9	0.7	0.9	1.4	2.0	2.3	2.7	3.5	4.8
2000.....	21.8	1.0	0.9	0.8	0.9	1.6	2.1	2.2	2.6	3.5	4.8
1999.....	21.9	0.7	0.9	0.7	0.9	1.5	2.2	2.2	2.6	3.7	4.9

Non-Hispanic 4/, male

2003.....	23.3	*	0.8	0.9	1.1	1.4	1.9	2.3	3.0	3.6	5.9
2002.....	23.6	*	0.8	0.8	1.2	1.6	2.1	2.4	3.1	3.7	5.5
2001.....	23.5	*	1.0	0.8	1.1	1.6	2.3	2.6	3.2	4.2	5.8
2000.....	23.7	*	1.0	1.0	1.0	1.9	2.6	2.3	3.2	4.0	5.7
1999.....	23.7	*	0.9	0.7	1.1	1.8	2.7	2.5	3.1	4.4	6.0

Non-Hispanic 4/, female

2003.....	18.9	*	0.7	0.8	0.8	0.9	1.3	1.7	1.7	2.7	3.6
2002.....	19.2	*	0.7	0.7	0.7	1.2	1.6	1.6	2.3	2.6	3.6
2001.....	19.7	*	0.9	0.6	0.7	1.1	1.7	2.0	2.3	2.8	3.7
2000.....	20.1	*	0.9	0.6	0.8	1.2	1.5	2.1	2.0	2.9	3.9
1999.....	20.1	*	0.8	0.6	0.8	1.2	1.7	1.9	2.1	3.0	3.7

Non-Hispanic White, both sexes

2003.....	23.4	1.0	0.8	0.9	0.9	1.2	1.6	2.0	2.2	3.0	4.6
2002.....	23.7	*	0.8	0.8	0.9	1.4	2.0	2.0	2.6	3.0	4.5
2001.....	23.8	*	1.1	0.7	0.9	1.4	2.0	2.1	2.6	3.4	4.5
2000.....	24.2	1.3	1.0	0.8	1.0	1.5	1.9	2.1	2.5	3.3	4.6
1999.....	24.1	*	1.0	0.7	0.9	1.4	2.1	2.2	2.5	3.7	4.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic White, male

2003.....	26.0	*	0.9	1.0	1.0	1.4	1.8	2.3	2.9	3.4	5.7
2002.....	26.2	*	0.9	0.8	1.1	1.7	2.2	2.4	2.9	3.5	5.5
2001.....	25.9	*	1.0	0.7	1.0	1.6	2.3	2.3	3.2	3.9	5.5
2000.....	26.2	*	1.1	1.0	1.1	1.8	2.5	2.2	3.0	3.9	5.7
1999.....	26.1	*	1.0	0.8	1.0	1.7	2.8	2.4	2.9	4.3	5.7

Non-Hispanic White, female

2003.....	20.8	*	0.7	0.7	0.8	0.9	1.3	1.7	1.6	2.6	3.4
2002.....	21.2	*	0.7	0.8	0.7	1.1	1.7	1.6	2.3	2.5	3.6
2001.....	21.8	*	1.1	0.6	0.8	1.1	1.7	1.8	2.1	2.9	3.5
2000.....	22.3	*	0.9	0.7	0.9	1.1	1.4	1.9	2.0	2.8	3.5
1999.....	22.2	*	0.9	0.6	0.7	1.2	1.5	1.9	2.1	3.0	3.7

Non-Hispanic Black, both sexes

2003.....	13.9	*	*	0.7	1.1	1.1	1.8	2.6	3.4	4.3	6.5
2002.....	14.1	*	*	*	1.0	1.4	1.5	2.5	3.5	3.8	5.8
2001.....	14.5	*	*	0.8	1.1	1.5	2.2	3.3	3.7	4.7	6.8
2000.....	14.3	*	*	0.6	0.7	2.0	2.7	2.9	3.2	4.5	6.4
1999.....	14.5	*	*	0.6	1.3	1.8	3.1	2.7	3.3	4.7	6.5

Non-Hispanic Black, male

2003.....	14.5	*	*	*	1.3	*	2.2	2.6	4.4	4.9	8.1
2002.....	15.1	*	*	*	1.2	1.6	1.4	2.9	4.6	5.1	7.3
2001.....	15.6	*	*	*	1.7	1.8	2.2	3.9	4.1	6.8	8.4
2000.....	15.2	*	*	*	*	2.4	3.3	2.7	4.6	4.8	6.7
1999.....	15.6	*	*	*	1.7	2.1	3.2	3.5	4.3	5.5	8.5

Non-Hispanic Black, female

2003.....	13.3	*	*	*	*	*	1.4	2.5	2.5	3.8	5.0
2002.....	13.1	*	*	*	*	*	1.6	2.1	2.6	2.6	4.4
2001.....	13.5	*	*	*	*	*	2.2	2.8	3.3	2.9	5.4
2000.....	13.5	*	*	*	*	1.5	2.1	3.0	1.9	4.3	6.2
1999.....	13.6	*	*	*	*	1.4	2.9	2.1	2.3	4.0	4.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Hodgkin's disease (C81)											
Hispanic, female											
2003.....	0.2	*	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4	0.4
2002.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.5	0.4	0.4
2000.....	0.5	*	*	*	*	*	0.3	0.3	0.5	0.4	0.3
1999.....	0.5	*	*	*	*	0.2	0.4	0.5	0.4	0.4	0.3
Non-Hispanic 4/, male											
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.6	0.4	0.5
2002.....	0.6	*	*	*	*	*	0.3	0.4	0.5	0.4	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.5	0.5	0.5	0.4
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4	0.4
1999.....	0.6	*	*	*	*	*	0.4	0.5	0.5	0.5	0.4
Non-Hispanic 4/, female											
2003.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4	0.2
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3	0.3
2001.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.3	0.3
2000.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3	0.3
1999.....	0.5	*	*	*	*	*	0.3	0.4	0.3	0.3	0.2
Non-Hispanic White, both sexes											
2003.....	0.6	*	*	*	*	*	0.2	0.4	0.5	0.4	0.4
2002.....	0.6	*	*	*	*	*	0.3	0.4	0.5	0.4	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4	0.4
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4	0.3
1999.....	0.6	*	*	*	*	0.2	0.4	0.5	0.4	0.5	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Non-Hodgkin's lymphoma (C82-C85)											
Hispanic, female											
2003.....	2.5	*	*	*	*	*	*	*	*	*	*
2002.....	2.8	*	*	*	*	*	*	*	*	*	1.6
2001.....	2.4	*	*	*	*	*	*	*	*	*	*
2000.....	2.7	*	*	*	*	*	*	*	*	*	2.0
1999.....	2.8	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	8.1	*	*	*	0.1	0.3	0.4	0.5	0.7	1.1	1.7
2002.....	8.3	*	*	*	0.2	0.2	0.5	0.6	0.9	1.2	1.9
2001.....	8.6	*	*	*	0.1	0.3	0.5	0.7	0.9	1.4	2.0
2000.....	8.8	*	*	0.1	0.1	0.3	0.7	0.7	0.9	1.3	2.0
1999.....	8.9	*	*	*	0.1	0.3	0.5	0.6	0.9	1.5	2.1
Non-Hispanic 4/, male											
2003.....	8.8	*	*	*	*	0.3	0.5	0.6	1.0	1.4	2.2
2002.....	9.0	*	*	*	0.3	0.3	0.6	0.8	1.1	1.5	2.5
2001.....	9.2	*	*	*	*	0.4	0.5	0.8	1.1	1.8	2.7
2000.....	9.4	*	*	*	*	0.5	0.9	0.7	1.1	1.8	2.6
1999.....	9.4	*	*	*	0.2	0.3	0.7	0.8	1.1	1.9	2.8
Non-Hispanic 4/, female											
2003.....	7.4	*	*	*	*	*	0.3	0.5	0.5	0.8	1.3
2002.....	7.7	*	*	*	*	*	0.4	0.5	0.6	0.8	1.3
2001.....	8.0	*	*	*	*	*	0.4	0.6	0.7	1.0	1.4
2000.....	8.3	*	*	*	*	*	0.5	0.6	0.6	0.9	1.5
1999.....	8.4	*	*	*	*	0.2	0.3	0.4	0.6	1.1	1.4
Non-Hispanic White, both sexes											
2003.....	9.3	*	*	*	*	0.2	0.4	0.4	0.6	1.0	1.7
2002.....	9.5	*	*	*	0.2	0.2	0.5	0.6	0.7	1.1	1.8
2001.....	9.7	*	*	*	*	0.3	0.4	0.6	0.8	1.3	2.0
2000.....	10.1	*	*	*	0.2	0.3	0.6	0.6	0.8	1.3	2.0
1999.....	10.1	*	*	*	*	0.2	0.5	0.6	0.8	1.4	2.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Non-Hodgkin's lymphoma (C82-C85)											
Non-Hispanic White, male											
2003.....	10.1	*	*	*	*	0.3	0.5	0.5	0.9	1.4	2.1
2002.....	10.2	*	*	*	0.3	*	0.6	0.8	0.9	1.4	2.4
2001.....	10.4	*	*	*	*	0.4	0.5	0.7	1.1	1.7	2.6
2000.....	10.6	*	*	*	*	0.5	0.8	0.6	1.0	1.7	2.7
1999.....	10.6	*	*	*	*	*	0.7	0.7	1.0	1.8	2.7
Non-Hispanic White, female											
2003.....	8.5	*	*	*	*	*	0.3	0.4	0.4	0.7	1.2
2002.....	8.8	*	*	*	*	*	0.3	0.4	0.5	0.8	1.3
2001.....	9.2	*	*	*	*	*	0.4	0.5	0.6	1.0	1.3
2000.....	9.5	*	*	*	*	*	0.4	0.5	0.6	0.8	1.4
1999.....	9.6	*	*	*	*	*	*	0.4	0.6	1.0	1.4
Non-Hispanic Black, both sexes											
2003.....	3.7	*	*	*	*	*	*	0.9	1.3	1.5	2.4
2002.....	4.0	*	*	*	*	*	*	0.9	1.5	1.3	2.7
2001.....	4.1	*	*	*	*	*	*	1.2	1.2	2.1	2.5
2000.....	3.9	*	*	*	*	*	1.0	1.2	1.0	1.9	2.4
1999.....	4.1	*	*	*	*	*	*	0.9	1.3	2.2	2.7
Non-Hispanic Black, male											
2003.....	3.9	*	*	*	*	*	*	*	1.9	1.7	3.1
2002.....	4.5	*	*	*	*	*	*	*	2.3	2.0	3.8
2001.....	4.7	*	*	*	*	*	*	*	*	3.2	3.5
2000.....	4.3	*	*	*	*	*	*	*	*	2.6	2.6
1999.....	4.7	*	*	*	*	*	*	*	1.8	2.7	4.0
Non-Hispanic Black, female											
2003.....	3.5	*	*	*	*	*	*	*	*	1.4	1.7
2002.....	3.6	*	*	*	*	*	*	*	*	*	1.7
2001.....	3.7	*	*	*	*	*	*	*	*	*	1.6
2000.....	3.5	*	*	*	*	*	*	*	*	*	2.1
1999.....	3.6	*	*	*	*	*	*	*	*	1.7	1.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Leukemia (C91-C95)											
All origins 3/, both sexes											
2003.....	7.4	0.7	0.8	0.8	0.9	0.9	1.1	1.1	1.1	1.5	2.1
2002.....	7.5	0.5	0.7	0.7	0.8	1.2	1.2	1.0	1.3	1.5	1.9
2001.....	7.5	0.6	1.0	0.8	0.8	1.1	1.3	1.2	1.4	1.6	1.9
2000.....	7.6	1.0	0.9	0.7	0.8	1.2	1.2	1.2	1.2	1.6	1.9
1999.....	7.5	0.7	0.8	0.6	0.9	1.1	1.4	1.2	1.3	1.6	1.9
All origins 3/, male											
2003.....	8.4	*	0.9	0.9	1.0	1.2	1.2	1.3	1.4	1.6	2.4
2002.....	8.5	*	0.7	0.7	0.9	1.4	1.4	1.2	1.5	1.6	2.2
2001.....	8.5	*	1.0	0.8	0.9	1.3	1.5	1.4	1.6	1.8	2.1
2000.....	8.5	1.1	0.9	0.8	1.0	1.4	1.5	1.3	1.5	1.6	2.1
1999.....	8.4	*	0.8	0.7	0.9	1.3	1.7	1.3	1.4	1.8	2.1
All origins 3/, female											
2003.....	6.4	1.1	0.8	0.7	0.8	0.6	0.9	0.9	0.8	1.4	1.7
2002.....	6.5	*	0.6	0.8	0.7	1.0	1.0	0.8	1.1	1.3	1.6
2001.....	6.6	*	1.0	0.7	0.7	0.9	1.0	1.0	1.2	1.3	1.7
2000.....	6.7	*	0.9	0.6	0.6	0.9	0.9	1.0	0.9	1.6	1.7
1999.....	6.7	*	0.7	0.5	0.8	0.9	1.1	1.1	1.2	1.5	1.7
Hispanic, both sexes											
2003.....	2.9	*	1.2	1.0	1.3	1.2	1.5	1.0	1.3	1.3	2.2
2002.....	2.8	*	0.8	1.1	1.2	1.7	1.5	1.0	1.2	2.0	2.2
2001.....	3.0	*	1.3	1.2	1.1	2.0	1.5	1.3	1.6	1.5	2.3
2000.....	2.9	*	0.9	0.8	1.4	1.5	1.5	1.1	1.1	1.8	2.0
1999.....	2.9	*	0.9	0.8	1.4	1.2	1.9	1.5	1.2	1.9	2.1
Hispanic, male											
2003.....	3.2	*	1.4	1.5	1.1	1.8	1.6	1.1	1.7	1.4	2.5
2002.....	3.0	*	*	*	1.3	2.2	1.8	1.2	1.7	2.1	2.5
2001.....	3.4	*	1.4	1.3	*	2.5	2.0	1.6	2.0	2.0	2.7
2000.....	3.3	*	*	*	2.0	1.7	1.6	1.6	1.3	1.6	2.4
1999.....	3.1	*	*	*	1.3	*	2.3	1.5	1.4	2.0	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Leukemia (C91-C95)											
Hispanic, female											
2003.....	2.6	*	*	*	1.5	*	1.4	*	*	*	1.9
2002.....	2.6	*	*	1.2	*	*	*	*	*	1.9	1.9
2001.....	2.6	*	*	*	1.3	1.3	*	*	*	*	1.9
2000.....	2.6	*	*	*	*	*	1.4	*	*	2.1	*
1999.....	2.8	*	*	*	1.5	1.5	1.6	1.5	*	1.8	2.4
Non-Hispanic 4/, both sexes											
2003.....	8.1	0.8	0.7	0.7	0.8	0.8	1.0	1.1	1.1	1.5	2.1
2002.....	8.2	*	0.6	0.7	0.7	1.1	1.1	1.0	1.4	1.4	1.9
2001.....	8.2	*	0.9	0.7	0.8	1.0	1.2	1.2	1.3	1.5	1.8
2000.....	8.2	1.0	0.9	0.7	0.7	1.1	1.1	1.2	1.2	1.6	1.9
1999.....	8.2	*	0.7	0.6	0.8	1.1	1.3	1.1	1.3	1.6	1.9
Non-Hispanic 4/, male											
2003.....	9.3	*	0.8	0.7	0.9	1.1	1.1	1.3	1.4	1.6	2.4
2002.....	9.4	*	0.7	0.7	0.8	1.2	1.2	1.2	1.5	1.5	2.2
2001.....	9.3	*	0.9	0.7	0.9	1.1	1.4	1.4	1.5	1.7	2.0
2000.....	9.3	*	0.9	0.8	0.9	1.4	1.4	1.3	1.5	1.6	2.1
1999.....	9.2	*	0.8	0.6	0.9	1.3	1.6	1.3	1.4	1.8	2.1
Non-Hispanic 4/, female											
2003.....	7.0	*	0.7	0.7	0.6	0.6	0.8	0.8	0.8	1.5	1.7
2002.....	7.0	*	0.5	0.6	0.6	0.9	0.9	0.8	1.2	1.3	1.6
2001.....	7.2	*	0.9	0.6	0.6	0.9	1.0	1.0	1.1	1.4	1.7
2000.....	7.2	*	0.9	0.5	0.6	0.9	0.7	1.1	0.9	1.5	1.7
1999.....	7.2	*	0.7	0.6	0.7	0.8	1.0	1.0	1.2	1.4	1.7
Non-Hispanic White, both sexes											
2003.....	9.1	1.0	0.8	0.8	0.7	0.8	0.9	1.1	1.0	1.4	2.0
2002.....	9.1	*	0.7	0.7	0.7	1.1	1.2	1.0	1.3	1.4	2.0
2001.....	9.1	*	1.0	0.6	0.7	1.0	1.2	1.0	1.3	1.5	1.8
2000.....	9.2	1.3	0.9	0.7	0.8	1.0	1.1	1.1	1.1	1.5	1.8
1999.....	9.1	*	0.8	0.6	0.7	1.0	1.2	1.2	1.2	1.6	1.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Leukemia (C91-C95)											
Non-Hispanic White, male											
2003.....	10.5	*	0.8	0.9	0.9	1.1	1.1	1.3	1.3	1.4	2.4
2002.....	10.5	*	0.8	0.7	0.8	1.2	1.3	1.2	1.5	1.5	2.3
2001.....	10.3	*	1.0	0.6	0.8	1.1	1.4	1.2	1.5	1.6	1.9
2000.....	10.4	*	1.0	0.8	0.9	1.2	1.4	1.2	1.3	1.6	2.1
1999.....	10.2	*	0.8	0.7	0.8	1.2	1.6	1.3	1.3	1.8	2.1
Non-Hispanic White, female											
2003.....	7.8	*	0.7	0.7	0.6	0.6	0.7	0.9	0.7	1.5	1.7
2002.....	7.8	*	0.5	0.7	0.6	0.9	1.0	0.7	1.2	1.3	1.7
2001.....	8.0	*	1.1	0.6	0.7	0.8	1.0	0.9	1.1	1.4	1.7
2000.....	8.1	*	0.9	0.6	0.7	0.8	0.7	1.0	0.8	1.5	1.6
1999.....	8.0	*	0.9	0.5	0.6	0.8	0.9	1.0	1.2	1.4	1.7
Non-Hispanic Black, both sexes											
2003.....	4.8	*	*	*	0.9	0.7	1.1	1.0	1.4	2.1	2.6
2002.....	4.8	*	*	*	0.7	1.0	0.8	1.1	1.5	1.5	1.7
2001.....	4.9	*	*	0.8	0.9	0.9	1.2	1.5	1.4	1.9	2.3
2000.....	4.9	*	*	*	0.6	1.5	1.3	1.5	1.9	1.7	2.5
1999.....	4.9	*	*	*	1.1	1.2	1.8	1.1	1.4	1.8	1.9
Non-Hispanic Black, male											
2003.....	5.2	*	*	*	*	*	1.4	*	1.8	2.6	3.0
2002.....	5.3	*	*	*	*	*	*	*	1.7	1.9	1.8
2001.....	5.4	*	*	*	1.5	*	*	1.8	1.8	2.4	2.7
2000.....	5.5	*	*	*	*	2.1	1.8	*	2.6	1.5	2.5
1999.....	5.4	*	*	*	1.5	1.5	1.7	*	1.8	1.9	2.3
Non-Hispanic Black, female											
2003.....	4.4	*	*	*	*	*	*	*	*	1.7	2.1
2002.....	4.4	*	*	*	*	*	*	*	*	*	1.6
2001.....	4.4	*	*	*	*	*	*	*	*	1.4	2.0
2000.....	4.4	*	*	*	*	*	*	1.6	*	1.9	2.4
1999.....	4.4	*	*	*	*	*	1.9	*	*	1.8	1.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All origins 3/, both sexes

2003.....	21.5	*	0.7	0.6	0.7	1.2	1.3	1.6	1.9	3.1	5.3
2002.....	21.8	0.6	0.8	0.7	0.7	1.1	1.5	1.5	2.1	3.3	5.8
2001.....	22.0	*	0.8	0.6	0.8	1.2	1.4	1.7	2.0	3.2	5.9
2000.....	22.4	0.6	0.8	0.6	0.7	1.2	1.4	1.5	2.0	3.4	5.9
1999.....	23.7	*	0.8	0.7	0.7	1.2	1.5	1.6	2.2	3.6	6.7

All origins 3/, male

2003.....	22.3	*	0.8	0.8	0.7	1.5	1.6	1.9	2.4	3.5	5.9
2002.....	22.7	*	0.8	0.8	0.7	1.3	1.9	2.0	2.5	3.7	6.2
2001.....	22.6	*	0.9	0.6	0.8	1.5	1.9	2.0	2.4	3.5	6.4
2000.....	23.0	*	1.0	0.6	0.7	1.4	1.8	1.9	2.4	3.7	6.4
1999.....	24.5	*	0.8	0.8	0.8	1.5	1.8	2.1	2.8	4.1	7.6

All origins 3/, female

2003.....	20.8	*	0.5	0.5	0.7	0.8	1.1	1.2	1.4	2.7	4.8
2002.....	21.0	*	0.8	0.6	0.8	0.9	1.0	1.0	1.7	2.8	5.4
2001.....	21.4	*	0.6	0.5	0.7	0.9	0.8	1.4	1.6	3.0	5.3
2000.....	21.9	*	0.7	0.6	0.6	1.0	1.0	1.1	1.6	3.0	5.3
1999.....	23.1	*	0.7	0.7	0.6	0.9	1.1	1.1	1.6	3.2	5.8

Hispanic, both sexes

2003.....	7.7	*	0.6	0.6	0.7	1.5	1.2	1.3	1.6	2.7	3.9
2002.....	7.8	*	*	0.6	0.9	1.2	1.9	1.1	1.8	2.2	4.0
2001.....	7.7	*	*	*	0.9	1.3	1.2	1.0	1.4	2.8	4.3
2000.....	7.8	*	0.8	*	0.9	1.4	1.1	1.4	1.7	2.1	3.6
1999.....	8.1	*	*	0.7	0.7	1.3	1.4	1.5	1.7	2.5	4.9

Hispanic, male

2003.....	7.9	*	*	*	*	2.1	1.2	1.9	2.3	2.7	4.0
2002.....	7.8	*	*	*	*	1.5	2.2	1.5	2.0	2.3	4.0
2001.....	7.7	*	*	*	*	1.5	2.0	1.1	2.0	2.7	4.2
2000.....	7.7	*	*	*	*	1.5	1.3	1.6	2.3	2.2	3.8
1999.....	8.0	*	*	*	*	1.7	2.0	1.9	1.8	2.5	4.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Hispanic, female											
2003.....	7.6	*	*	*	*	*	1.2	*	*	2.7	3.8
2002.....	7.7	*	*	*	*	*	1.5	*	1.7	2.1	3.9
2001.....	7.7	*	*	*	*	*	*	*	*	2.8	4.4
2000.....	7.9	*	*	*	*	*	*	*	*	1.9	3.4
1999.....	8.2	*	*	*	*	*	*	*	1.5	2.5	4.9
Non-Hispanic 4/, both sexes											
2003.....	23.7	*	0.7	0.6	0.7	1.1	1.3	1.6	1.9	3.2	5.5
2002.....	24.0	*	0.9	0.7	0.7	1.1	1.4	1.6	2.2	3.4	6.0
2001.....	24.1	*	0.8	0.6	0.8	1.2	1.4	1.8	2.1	3.3	6.0
2000.....	24.4	0.7	0.8	0.6	0.6	1.2	1.5	1.5	2.0	3.6	6.1
1999.....	25.8	*	0.8	0.7	0.7	1.2	1.5	1.6	2.3	3.8	6.8
Non-Hispanic 4/, male											
2003.....	24.6	*	0.8	0.8	0.7	1.4	1.7	1.8	2.4	3.6	6.1
2002.....	25.0	*	0.9	0.8	0.7	1.3	1.9	2.2	2.6	4.0	6.4
2001.....	24.9	*	1.0	0.7	0.8	1.4	1.9	2.2	2.5	3.6	6.6
2000.....	25.2	*	0.9	0.6	0.7	1.3	2.0	2.0	2.4	4.0	6.7
1999.....	26.8	*	1.0	0.8	0.8	1.5	1.7	2.1	2.9	4.3	7.9
Non-Hispanic 4/, female											
2003.....	22.8	*	0.6	0.5	0.7	0.8	1.0	1.4	1.5	2.7	4.9
2002.....	22.9	*	0.8	0.6	0.7	1.0	0.8	1.1	1.7	2.9	5.6
2001.....	23.3	*	0.6	0.6	0.7	0.9	0.9	1.4	1.7	3.0	5.4
2000.....	23.7	*	0.8	0.6	0.6	1.0	1.0	1.1	1.7	3.2	5.5
1999.....	25.0	*	0.7	0.7	0.6	0.9	1.2	1.1	1.6	3.3	5.8
Non-Hispanic White, both sexes											
2003.....	25.9	*	0.6	0.7	0.6	1.2	1.4	1.6	1.8	3.2	5.2
2002.....	26.0	*	0.8	0.7	0.7	1.2	1.3	1.6	2.1	3.4	5.7
2001.....	26.0	*	0.9	0.6	0.7	1.2	1.5	1.9	2.0	3.2	5.8
2000.....	26.5	*	0.8	0.7	0.6	1.1	1.6	1.7	2.0	3.4	5.6
1999.....	27.9	*	0.9	0.7	0.6	1.2	1.4	1.7	2.2	3.6	6.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic White, male

2003.....	26.8	*	0.7	0.9	0.6	1.5	1.7	2.0	2.2	3.8	5.7
2002.....	27.2	*	0.8	0.7	0.6	1.3	1.8	2.1	2.6	3.9	5.9
2001.....	26.8	*	1.1	0.7	0.8	1.4	2.0	2.3	2.4	3.5	6.4
2000.....	27.2	*	1.0	0.7	0.7	1.3	2.0	2.2	2.5	3.8	6.0
1999.....	28.8	*	1.1	0.8	0.7	1.4	1.6	2.3	2.8	4.2	7.4

Non-Hispanic White, female

2003.....	25.0	*	0.6	0.5	0.7	0.8	1.0	1.2	1.4	2.5	4.7
2002.....	24.9	*	0.7	0.6	0.8	1.0	0.9	1.1	1.6	2.9	5.5
2001.....	25.3	*	0.7	0.6	0.7	0.9	1.0	1.5	1.6	2.8	5.1
2000.....	25.8	*	0.6	0.7	0.6	0.9	1.1	1.1	1.6	2.9	5.2
1999.....	27.0	*	0.8	0.6	0.5	1.0	1.2	1.0	1.6	3.1	5.3

Non-Hispanic Black, both sexes

2003.....	17.9	*	0.9	0.7	0.9	0.9	1.4	1.9	2.8	3.8	7.9
2002.....	18.7	*	1.1	0.8	0.8	1.1	1.6	2.0	2.8	4.7	8.2
2001.....	18.9	*	*	*	0.8	1.2	1.4	2.1	2.7	4.3	7.8
2000.....	18.9	*	0.9	*	0.7	1.6	1.4	1.3	2.0	5.4	10.1
1999.....	20.2	*	*	0.7	0.9	1.3	1.7	1.6	2.5	5.1	10.4

Non-Hispanic Black, male

2003.....	18.8	*	*	*	*	*	1.8	1.7	3.1	3.7	9.4
2002.....	19.4	*	*	*	*	1.4	2.3	2.6	2.9	5.4	9.9
2001.....	20.0	*	*	*	*	1.5	2.1	2.6	3.0	4.4	8.3
2000.....	20.0	*	*	*	*	1.8	2.3	1.6	2.1	5.6	12.0
1999.....	21.3	*	*	*	*	1.9	2.1	1.6	3.2	5.2	11.9

Non-Hispanic Black, female

2003.....	17.1	*	*	*	*	*	*	2.1	2.4	4.0	6.6
2002.....	18.0	*	*	*	*	*	*	1.5	2.7	4.2	6.8
2001.....	17.8	*	*	*	*	*	*	1.6	2.4	4.3	7.4
2000.....	17.8	*	*	*	*	1.4	*	*	1.9	5.2	8.5
1999.....	19.2	*	*	*	*	*	*	1.5	1.8	4.9	9.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Hispanic, female

2003.....	1.6	*	*	*	*	*	*	*	*	*	*
2002.....	1.5	*	*	*	*	*	*	*	*	*	*
2001.....	1.5	*	*	*	*	*	*	*	*	*	*
2000.....	1.7	*	*	*	*	*	*	*	*	*	*
1999.....	1.6	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	5.2	1.5	0.4	0.2	0.2	0.2	0.2	0.4	0.5	0.7	0.9
2002.....	5.1	1.6	0.4	0.2	0.2	0.2	0.2	0.4	0.5	0.7	1.0
2001.....	5.3	1.4	0.4	0.2	0.3	0.3	0.3	0.3	0.5	0.7	0.9
2000.....	5.3	1.7	0.4	0.3	0.2	0.2	0.3	0.4	0.5	0.7	0.9
1999.....	5.2	1.6	0.4	0.3	0.2	0.3	0.3	0.3	0.6	0.7	0.9

Non-Hispanic 4/, male

2003.....	5.2	1.6	0.4	*	*	*	0.3	0.4	0.6	0.8	1.1
2002.....	5.1	1.3	0.4	*	*	*	0.3	0.4	0.5	0.8	1.1
2001.....	5.3	1.5	0.4	*	0.3	*	0.4	0.4	0.5	0.8	1.0
2000.....	5.3	1.8	0.5	0.3	*	0.3	0.4	0.4	0.5	0.8	0.9
1999.....	5.3	1.6	0.5	0.3	*	0.2	0.3	0.3	0.7	0.8	1.0

Non-Hispanic 4/, female

2003.....	5.1	1.3	*	*	*	*	*	0.3	0.5	0.6	0.7
2002.....	5.1	2.0	0.4	*	*	*	*	0.4	0.5	0.6	0.9
2001.....	5.2	1.4	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.7	0.9
2000.....	5.3	1.7	*	0.4	*	*	*	0.4	0.5	0.6	0.9
1999.....	5.1	1.7	0.4	0.3	*	0.3	*	0.3	0.5	0.6	0.8

Non-Hispanic White, both sexes

2003.....	5.8	1.4	0.3	*	0.2	0.2	0.2	0.4	0.5	0.6	0.9
2002.....	5.7	1.8	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.7	0.9
2001.....	5.9	1.3	0.4	0.2	0.2	0.2	0.3	0.4	0.5	0.7	0.9
2000.....	5.9	1.8	0.4	0.3	0.2	0.2	0.3	0.3	0.5	0.7	0.9
1999.....	5.7	1.5	0.4	0.3	0.2	0.2	0.2	0.2	0.5	0.6	0.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Anemias (D50-D64)

Non-Hispanic White, male

2003.....	1.3	*	*	*	*	*	*	*	*	*	*
2002.....	1.2	*	*	*	*	*	*	*	*	*	*
2001.....	1.3	*	*	*	*	*	*	*	*	*	*
2000.....	1.2	*	*	*	*	*	*	*	*	*	*
1999.....	1.2	*	*	*	*	*	*	*	*	*	*

Non-Hispanic White, female

2003.....	2.0	*	*	*	*	*	*	*	*	*	*
2002.....	2.1	*	*	*	*	*	*	*	*	*	*
2001.....	2.1	*	*	*	*	*	*	*	*	*	0.3
2000.....	2.1	*	*	*	*	*	*	*	*	*	*
1999.....	2.0	*	*	*	*	*	*	*	*	*	*

Non-Hispanic Black, both sexes

2003.....	2.8	*	*	*	*	*	2.0	2.1	2.3	2.2	2.6
2002.....	2.9	*	*	*	*	0.9	1.8	2.6	2.4	2.6	2.5
2001.....	2.8	*	*	*	*	0.9	1.6	2.0	2.5	2.2	3.3
2000.....	2.7	*	*	*	*	*	1.7	1.8	2.9	2.3	2.7
1999.....	2.9	*	0.9	*	*	0.8	2.4	2.1	3.2	2.3	2.8

Non-Hispanic Black, male

2003.....	2.8	*	*	*	*	*	1.8	2.3	2.4	2.2	3.0
2002.....	2.7	*	*	*	*	*	2.3	3.3	2.6	2.7	2.4
2001.....	2.6	*	*	*	*	*	1.7	2.6	3.4	2.3	2.8
2000.....	2.4	*	*	*	*	*	1.7	1.6	2.8	2.0	2.3
1999.....	2.7	*	*	*	*	*	2.9	2.4	3.4	2.6	3.0

Non-Hispanic Black, female

2003.....	2.9	*	*	*	*	*	2.2	1.8	2.3	2.3	2.1
2002.....	3.0	*	*	*	*	*	*	1.9	2.1	2.5	2.6
2001.....	2.9	*	*	*	*	*	1.5	*	1.8	2.1	3.8
2000.....	3.1	*	*	*	*	*	1.7	2.0	3.0	2.6	3.1
1999.....	3.1	*	*	*	*	*	1.8	1.9	2.9	2.1	2.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diabetes mellitus (E10-E14)

Hispanic, female

2003.....	16.6	*	*	*	*	*	*	*	*	1.5	4.1
2002.....	16.7	*	*	*	*	*	*	*	*	2.1	3.8
2001.....	17.1	*	*	*	*	*	*	*	*	2.0	2.4
2000.....	16.5	*	*	*	*	*	*	*	*	1.6	3.4
1999.....	17.3	*	*	*	*	*	*	*	*	1.9	4.4

Non-Hispanic 4/, both sexes

2003.....	27.0	*	*	*	*	0.2	0.6	1.3	2.3	3.6	5.9
2002.....	26.9	*	*	*	*	0.2	0.3	0.6	1.3	2.2	6.1
2001.....	26.4	*	*	*	*	0.2	0.2	0.6	1.3	2.0	5.5
2000.....	25.9	*	*	*	*	0.3	0.7	1.2	2.1	3.2	5.6
1999.....	25.7	*	*	*	*	0.3	0.6	1.1	2.1	3.3	5.5

Non-Hispanic 4/, male

2003.....	26.4	*	*	*	*	0.3	0.7	1.4	2.8	4.2	7.6
2002.....	25.8	*	*	*	*	0.3	0.7	1.4	2.8	4.4	7.9
2001.....	25.0	*	*	*	*	0.3	0.6	1.4	2.4	4.1	6.7
2000.....	24.2	*	*	*	*	0.3	0.8	1.3	2.6	3.7	6.9
1999.....	24.1	*	*	*	*	0.3	0.6	1.0	2.4	4.1	6.6

Non-Hispanic 4/, female

2003.....	27.6	*	*	*	*	*	0.5	1.2	1.8	3.0	4.3
2002.....	27.9	*	*	*	*	0.3	0.5	1.2	1.7	3.1	4.5
2001.....	27.8	*	*	*	*	*	0.5	1.1	1.6	2.8	4.4
2000.....	27.6	*	*	*	*	*	0.6	1.0	1.7	2.7	4.4
1999.....	27.3	*	*	*	*	0.3	0.6	1.1	1.7	2.6	4.6

Non-Hispanic White, both sexes

2003.....	26.6	*	*	*	*	0.2	0.5	0.9	2.0	3.0	5.0
2002.....	26.4	*	*	*	*	0.2	0.5	1.0	2.0	3.2	5.3
2001.....	26.0	*	*	*	*	0.2	0.4	0.9	1.7	2.9	4.8
2000.....	25.4	*	*	*	*	0.2	0.5	0.9	1.8	2.8	5.0
1999.....	25.0	*	*	*	*	0.3	0.5	1.0	1.8	2.9	4.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diabetes mellitus (E10-E14)

Non-Hispanic White, male

2003.....	26.6	*	*	*	*	*	0.6	0.8	2.5	3.6	6.6
2002.....	26.0	*	*	*	*	*	0.5	1.2	2.4	3.9	6.9
2001.....	25.1	*	*	*	*	*	0.5	0.9	1.9	3.4	6.2
2000.....	24.3	*	*	*	*	*	0.7	1.0	2.3	3.2	6.3
1999.....	24.0	*	*	*	*	0.3	0.5	1.0	2.0	3.6	5.6

Non-Hispanic White, female

2003.....	26.5	*	*	*	*	*	0.4	1.0	1.5	2.4	3.4
2002.....	26.8	*	*	*	*	*	0.5	0.9	1.5	2.6	3.7
2001.....	26.9	*	*	*	*	*	*	0.9	1.5	2.4	3.5
2000.....	26.5	*	*	*	*	*	0.4	0.8	1.3	2.4	3.7
1999.....	26.0	*	*	*	*	*	0.5	0.9	1.5	2.1	3.8

Non-Hispanic Black, both sexes

2003.....	35.0	*	*	*	*	*	1.0	3.4	4.6	7.7	12.1
2002.....	34.8	*	*	*	*	*	1.4	2.9	4.5	7.5	12.3
2001.....	34.1	*	*	*	*	*	1.4	3.3	4.4	7.2	11.0
2000.....	34.0	*	*	*	*	*	1.5	2.9	4.3	6.2	10.4
1999.....	34.1	*	*	*	*	*	1.2	1.9	4.2	6.8	11.6

Non-Hispanic Black, male

2003.....	30.9	*	*	*	*	*	*	4.4	5.5	8.8	14.9
2002.....	30.0	*	*	*	*	*	1.8	3.1	6.0	8.7	15.4
2001.....	29.4	*	*	*	*	*	1.5	4.1	5.8	9.0	12.0
2000.....	28.3	*	*	*	*	*	1.6	3.3	4.7	7.4	12.0
1999.....	28.6	*	*	*	*	*	*	1.7	5.1	8.0	13.5

Non-Hispanic Black, female

2003.....	38.7	*	*	*	*	*	*	2.4	3.9	6.6	9.6
2002.....	39.2	*	*	*	*	*	*	2.8	3.3	6.4	9.7
2001.....	38.4	*	*	*	*	*	*	2.6	3.0	5.6	10.1
2000.....	39.1	*	*	*	*	*	1.5	2.6	3.9	5.1	8.9
1999.....	39.0	*	*	*	*	*	*	2.0	3.4	5.6	9.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Meningitis (G00,G03)

Hispanic, female

2003.....	0.2	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.3	1.9	0.2	*	*	*	*	*	*	0.2	0.2
2002.....	0.3	2.1	*	*	0.1	*	*	*	*	*	0.2
2001.....	0.3	2.2	0.2	*	*	*	*	*	0.1	0.1	0.2
2000.....	0.3	2.7	0.3	*	*	*	*	*	0.1	0.2	0.2
1999.....	0.3	3.2	0.3	*	*	*	*	*	0.1	0.2	0.2

Non-Hispanic 4/, male

2003.....	0.3	1.6	*	*	*	*	*	*	*	0.2	*
2002.....	0.3	2.1	*	*	*	*	*	*	*	*	0.2
2001.....	0.3	2.1	*	*	*	*	*	*	*	*	*
2000.....	0.3	2.7	*	*	*	*	*	*	*	*	0.2
1999.....	0.3	3.1	0.3	*	*	*	*	*	*	*	0.2

Non-Hispanic 4/, female

2003.....	0.2	2.2	*	*	*	*	*	*	*	*	*
2002.....	0.2	2.0	*	*	*	*	*	*	*	*	*
2001.....	0.2	2.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	2.7	*	*	*	*	*	*	*	*	*
1999.....	0.3	3.3	*	*	*	*	*	*	*	*	*

Non-Hispanic White, both sexes

2003.....	0.2	1.5	*	*	*	*	*	*	*	*	0.1
2002.....	0.2	1.9	*	*	*	*	*	*	*	*	*
2001.....	0.2	1.6	*	*	*	*	*	*	*	*	0.1
2000.....	0.2	2.1	*	*	*	*	*	*	*	*	*
1999.....	0.3	2.2	0.3	*	*	*	*	*	*	0.1	0.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Major cardiovascular diseases (I00-I78)

All origins 3/, both sexes

2003.....	310.3	13.9	1.5	0.7	1.0	2.5	4.5	7.1	13.2	25.1	50.3
2002.....	318.6	15.9	1.4	0.6	1.1	2.4	3.9	7.1	12.6	25.3	49.7
2001.....	323.9	14.9	1.9	0.7	1.1	2.2	4.3	7.1	13.0	25.0	48.8
2000.....	332.9	16.7	1.5	0.7	1.1	2.4	4.3	6.7	12.2	24.2	49.6
1999.....	340.6	17.0	1.5	0.7	1.0	2.7	4.1	6.5	12.6	25.9	50.0

All origins 3/, male

2003.....	296.3	15.2	1.4	0.7	1.2	3.1	5.3	8.5	17.0	33.4	67.7
2002.....	303.3	16.6	1.5	0.6	1.3	3.1	5.0	9.0	16.1	32.9	67.9
2001.....	306.2	15.3	1.9	0.6	1.1	2.7	5.3	8.8	16.5	33.1	66.5
2000.....	315.8	17.5	1.6	0.7	1.2	2.8	5.0	8.2	15.3	31.3	68.0
1999.....	323.0	17.9	1.6	0.8	1.1	3.3	4.8	7.8	16.2	34.2	68.9

All origins 3/, female

2003.....	323.9	12.5	1.5	0.6	0.8	2.0	3.6	5.6	9.3	16.8	33.2
2002.....	333.3	15.1	1.3	0.7	0.9	1.7	2.8	5.1	9.0	17.7	31.6
2001.....	340.9	14.5	1.8	0.8	1.1	1.7	3.2	5.5	9.5	17.0	31.5
2000.....	349.5	15.9	1.4	0.6	1.0	2.0	3.5	5.2	9.0	17.1	31.5
1999.....	357.5	16.0	1.4	0.6	1.0	2.1	3.4	5.2	9.1	17.6	31.4

Hispanic, both sexes

2003.....	93.4	11.0	1.3	*	0.9	1.9	3.0	4.8	8.0	13.5	28.6
2002.....	94.3	13.1	1.2	*	1.0	1.8	3.3	4.7	7.5	13.8	28.2
2001.....	96.2	12.0	1.5	0.7	0.9	2.1	2.8	4.4	8.6	14.1	26.7
2000.....	96.5	13.7	1.3	0.6	0.8	1.9	3.2	4.8	8.2	14.5	29.7
1999.....	99.2	15.2	1.4	0.6	0.8	2.2	2.9	4.6	8.5	15.6	28.6

Hispanic, male

2003.....	92.6	12.2	1.5	*	*	2.4	4.0	5.7	10.2	18.3	39.2
2002.....	94.2	12.2	*	*	*	2.2	4.3	6.5	9.8	17.4	39.5
2001.....	95.4	12.7	1.4	*	*	2.4	3.5	5.6	11.2	19.0	36.5
2000.....	96.0	13.7	1.4	*	1.2	2.4	3.7	6.0	10.4	18.2	40.0
1999.....	99.2	12.7	1.4	*	*	2.5	3.5	6.1	11.4	20.8	40.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Major cardiovascular diseases (I00-I78)

Hispanic, female

2003.....	94.3	9.7	*	*	*	1.3	1.9	3.8	5.5	8.0	17.2
2002.....	94.4	14.0	1.3	*	1.2	1.4	2.0	2.5	4.9	9.8	16.2
2001.....	97.1	11.2	1.6	*	*	1.7	1.8	3.0	5.6	8.7	16.5
2000.....	96.9	13.8	*	*	*	*	2.6	3.4	5.7	10.5	18.9
1999.....	99.1	17.9	1.4	*	*	1.8	2.1	2.8	5.2	10.0	15.9

Non-Hispanic 4/, both sexes

2003.....	343.8	14.5	1.5	0.7	1.0	2.6	4.8	7.6	14.2	27.0	53.0
2002.....	352.2	16.5	1.5	0.7	1.1	2.5	4.0	7.7	13.6	27.1	52.2
2001.....	356.7	15.5	1.9	0.7	1.1	2.2	4.5	7.7	13.8	26.6	51.3
2000.....	365.6	17.3	1.6	0.7	1.1	2.5	4.5	7.0	12.8	25.4	51.5
1999.....	372.7	17.4	1.5	0.7	1.1	2.8	4.4	6.8	13.3	27.1	52.0

Non-Hispanic 4/, male

2003.....	329.5	15.9	1.4	0.8	1.2	3.2	5.7	9.2	18.5	36.1	71.3
2002.....	336.4	17.6	1.7	0.6	1.4	3.3	5.1	9.7	17.4	35.5	71.5
2001.....	338.1	15.8	2.0	0.6	1.2	2.7	5.6	9.5	17.4	35.2	69.8
2000.....	347.7	18.3	1.7	0.8	1.2	2.9	5.4	8.6	16.2	33.0	70.8
1999.....	354.2	19.1	1.7	0.8	1.1	3.5	5.1	8.1	17.0	35.9	71.6

Non-Hispanic 4/, female

2003.....	357.5	13.1	1.6	0.7	0.8	2.1	3.9	5.9	10.1	18.1	35.1
2002.....	367.2	15.3	1.3	0.8	0.9	1.7	2.9	5.7	9.8	18.9	33.4
2001.....	374.3	15.2	1.8	0.8	1.1	1.7	3.4	6.0	10.1	18.2	33.1
2000.....	382.6	16.3	1.5	0.6	1.1	2.1	3.7	5.5	9.5	18.0	32.7
1999.....	390.2	15.5	1.4	0.6	1.0	2.1	3.7	5.6	9.7	18.5	33.0

Non-Hispanic White, both sexes

2003.....	372.4	10.9	1.2	0.6	0.9	2.2	3.8	5.9	11.7	22.8	45.4
2002.....	381.9	13.4	1.4	0.6	0.9	2.2	3.2	6.0	10.8	22.9	45.6
2001.....	385.8	12.1	1.5	0.6	1.0	1.9	3.5	6.2	11.3	22.5	43.3
2000.....	395.5	14.6	1.4	0.6	0.9	2.1	3.5	5.6	10.4	21.4	43.9
1999.....	402.7	14.2	1.3	0.7	0.9	2.4	3.5	5.5	10.9	22.8	43.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic White, male

2003.....	354.1	11.3	0.9	0.6	1.1	2.7	4.5	7.3	15.5	31.4	63.5
2002.....	362.2	14.8	1.6	0.6	1.1	2.9	4.2	7.7	14.1	30.6	64.9
2001.....	363.0	13.0	1.6	0.6	1.1	2.2	4.5	7.8	14.6	30.4	61.8
2000.....	373.6	15.5	1.6	0.7	1.1	2.4	4.3	7.0	13.7	28.9	62.9
1999.....	379.8	16.1	1.5	0.8	0.9	2.9	4.0	6.5	14.3	30.9	62.8

Non-Hispanic White, female

2003.....	390.0	10.4	1.4	0.6	0.7	1.8	3.0	4.4	7.8	14.1	27.4
2002.....	400.8	12.0	1.1	0.7	0.7	1.5	2.2	4.2	7.5	15.2	26.4
2001.....	407.6	11.0	1.5	0.7	0.9	1.5	2.5	4.6	8.0	14.6	24.8
2000.....	416.5	13.6	1.1	0.4	0.8	1.7	2.8	4.2	7.0	13.8	25.0
1999.....	424.7	12.1	1.1	0.6	0.9	1.9	3.1	4.4	7.4	14.6	24.9

Non-Hispanic Black, both sexes

2003.....	282.2	26.6	2.9	1.3	1.7	4.5	9.8	16.9	30.3	54.7	107.0
2002.....	285.1	28.3	2.1	0.8	2.0	4.0	8.1	17.4	30.7	54.7	101.5
2001.....	289.7	30.1	3.7	1.0	1.6	4.1	9.8	17.1	29.5	53.3	108.4
2000.....	294.2	28.8	2.5	1.0	1.9	4.8	9.5	15.7	28.8	52.8	106.7
1999.....	298.5	30.2	2.6	1.1	1.9	5.0	9.3	15.1	28.0	56.8	112.2

Non-Hispanic Black, male

2003.....	277.5	31.7	3.3	1.5	1.7	5.8	11.7	20.6	37.4	68.5	133.4
2002.....	277.2	29.1	2.0	*	2.4	5.1	9.7	21.6	38.2	68.8	126.0
2001.....	281.2	29.7	3.8	*	1.6	5.2	11.7	21.0	36.4	67.9	132.7
2000.....	284.3	30.0	2.3	*	1.9	5.3	10.8	18.9	33.9	63.5	133.3
1999.....	291.1	31.5	2.7	*	2.2	6.5	11.2	18.6	34.5	71.6	141.1

Non-Hispanic Black, female

2003.....	286.5	21.3	2.4	*	1.6	3.2	8.0	13.5	23.8	42.5	83.8
2002.....	292.3	27.4	2.2	*	1.6	2.9	6.6	13.5	24.0	42.2	80.0
2001.....	297.4	30.6	3.6	1.3	1.5	2.9	7.9	13.6	23.3	40.4	86.9
2000.....	303.1	27.5	2.9	*	2.0	4.4	8.4	12.9	24.2	43.3	83.2
1999.....	305.1	28.9	2.4	*	1.6	3.3	7.4	11.9	22.2	43.8	86.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diseases of heart (I00-I09,I11,I13,I20-I51)

All origins 3/, both sexes

2003.....	235.6	11.0	1.2	0.5	0.8	1.9	3.6	5.6	10.5	20.1	40.5
2002.....	241.7	12.4	1.1	0.5	0.8	2.0	3.1	5.7	10.0	20.3	40.2
2001.....	245.8	11.9	1.5	0.5	0.8	1.7	3.3	5.6	10.2	19.9	39.1
2000.....	252.6	13.0	1.2	0.5	0.8	2.0	3.3	5.2	9.5	19.1	39.4
1999.....	259.9	13.8	1.2	0.6	0.8	2.3	3.3	5.1	10.1	20.6	40.1

All origins 3/, male

2003.....	235.0	12.1	1.1	0.6	0.9	2.4	4.4	6.9	13.9	27.8	56.9
2002.....	240.7	12.9	1.1	0.4	0.9	2.6	4.0	7.4	13.4	27.8	57.7
2001.....	242.5	11.8	1.5	0.4	0.9	2.1	4.2	7.0	13.4	27.4	55.8
2000.....	249.8	13.3	1.4	0.6	0.9	2.4	4.0	6.5	12.4	26.0	57.0
1999.....	257.0	13.9	1.3	0.6	0.9	2.9	3.9	6.3	13.2	28.5	58.1

All origins 3/, female

2003.....	236.2	9.8	1.3	0.5	0.6	1.4	2.7	4.2	7.1	12.5	24.3
2002.....	242.7	11.8	1.0	0.5	0.6	1.3	2.1	3.9	6.5	12.8	22.9
2001.....	249.0	12.0	1.4	0.6	0.8	1.3	2.4	4.1	6.9	12.4	22.6
2000.....	255.3	12.5	1.0	0.4	0.7	1.6	2.6	3.9	6.5	12.2	22.2
1999.....	262.6	13.7	1.1	0.5	0.7	1.7	2.6	3.9	6.9	12.6	22.4

Hispanic, both sexes

2003.....	70.9	8.7	1.1	*	0.6	1.4	2.4	3.5	6.0	10.3	20.1
2002.....	71.9	10.2	0.8	*	*	1.2	2.5	3.4	5.5	9.9	20.3
2001.....	73.3	9.7	1.1	*	0.6	1.4	1.9	3.2	6.0	10.1	18.1
2000.....	73.1	11.2	1.0	*	0.7	1.5	2.2	3.3	5.4	10.1	20.7
1999.....	76.2	12.7	1.2	*	*	1.7	2.0	2.9	6.3	11.7	20.0

Hispanic, male

2003.....	72.2	9.5	*	*	*	1.8	3.4	4.4	7.9	14.5	28.9
2002.....	74.0	9.6	*	*	*	1.6	3.4	4.8	7.5	12.9	30.0
2001.....	74.6	10.3	*	*	*	1.6	2.5	4.4	8.0	14.6	26.6
2000.....	74.7	11.2	1.4	*	*	2.1	2.7	4.3	7.2	13.4	29.9
1999.....	77.7	10.3	1.4	*	*	1.9	2.4	4.4	8.7	16.5	30.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Hispanic, female

2003.....	69.6	7.8	*	*	*	*	*	2.4	3.7	5.6	10.7
2002.....	69.7	10.8	*	*	*	*	1.3	1.7	3.1	6.6	10.0
2001.....	71.8	9.0	*	*	*	*	*	1.9	3.7	5.2	9.2
2000.....	71.5	11.2	*	*	*	*	1.6	2.1	3.3	6.4	11.0
1999.....	74.6	15.2	*	*	*	1.5	1.5	*	3.6	6.5	9.6

Non-Hispanic 4/, both sexes

2003.....	261.0	11.5	1.2	0.6	0.8	2.0	3.8	6.1	11.4	21.8	43.0
2002.....	267.1	12.9	1.1	0.5	0.8	2.1	3.2	6.2	10.8	21.9	42.6
2001.....	270.7	12.3	1.5	0.5	0.9	1.7	3.6	6.1	10.9	21.3	41.4
2000.....	277.3	13.3	1.2	0.5	0.8	2.1	3.5	5.6	10.2	20.3	41.3
1999.....	284.3	14.0	1.2	0.6	0.8	2.4	3.5	5.5	10.7	21.6	42.1

Non-Hispanic 4/, male

2003.....	261.4	12.8	1.1	0.6	0.9	2.5	4.6	7.6	15.2	30.1	60.5
2002.....	267.0	13.7	1.2	0.4	1.0	2.8	4.1	8.1	14.6	30.3	61.3
2001.....	268.0	12.0	1.7	0.4	0.9	2.2	4.6	7.6	14.4	29.3	59.2
2000.....	275.2	13.7	1.4	0.6	0.9	2.4	4.3	7.0	13.4	27.7	59.7
1999.....	282.0	14.8	1.3	0.6	0.9	3.0	4.2	6.6	14.0	30.0	60.9

Non-Hispanic 4/, female

2003.....	260.5	10.2	1.3	0.6	0.6	1.5	3.0	4.6	7.7	13.6	25.9
2002.....	267.2	12.0	1.1	0.6	0.6	1.4	2.2	4.3	7.1	13.7	24.4
2001.....	273.3	12.7	1.4	0.6	0.8	1.3	2.6	4.5	7.5	13.4	24.2
2000.....	279.3	12.8	1.1	0.4	0.7	1.7	2.7	4.2	7.0	12.9	23.3
1999.....	286.4	13.3	1.1	0.5	0.8	1.7	2.8	4.3	7.4	13.4	23.7

Non-Hispanic White, both sexes

2003.....	284.0	8.7	0.9	0.5	0.7	1.7	3.0	4.8	9.5	19.1	38.1
2002.....	290.8	10.7	1.0	0.4	0.7	1.8	2.4	4.8	8.7	19.2	38.5
2001.....	294.1	9.8	1.2	0.4	0.8	1.4	2.7	4.8	9.0	18.4	36.2
2000.....	301.3	11.4	1.1	0.4	0.7	1.7	2.7	4.3	8.3	17.6	36.5
1999.....	308.2	11.3	1.0	0.5	0.7	2.1	2.7	4.3	8.8	18.6	36.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic White, male

2003.....	282.9	9.4	0.7	0.5	0.9	2.0	3.5	6.1	13.1	27.2	55.5
2002.....	289.2	11.6	1.1	0.4	0.9	2.4	3.2	6.4	12.0	26.9	57.2
2001.....	289.5	10.3	1.3	0.4	0.8	1.7	3.5	6.3	12.1	26.0	53.9
2000.....	297.5	12.0	1.2	0.5	0.8	1.9	3.3	5.5	11.4	24.9	54.8
1999.....	304.0	12.4	1.1	0.5	0.7	2.5	3.2	5.3	11.9	26.4	54.9

Non-Hispanic White, female

2003.....	285.1	8.0	1.1	0.5	0.5	1.3	2.5	3.4	5.9	11.0	20.8
2002.....	292.3	9.6	0.8	0.5	0.5	1.2	1.6	3.3	5.4	11.4	19.8
2001.....	298.4	9.2	1.1	0.5	0.7	1.1	1.9	3.4	5.9	10.8	18.5
2000.....	304.9	10.8	0.9	0.3	0.5	1.4	2.0	3.1	5.2	10.2	18.2
1999.....	312.3	10.2	0.8	0.5	0.7	1.6	2.2	3.4	5.7	10.7	18.2

Non-Hispanic Black, both sexes

2003.....	209.4	20.4	2.3	1.1	1.3	3.5	8.1	13.6	23.7	41.2	81.1
2002.....	212.2	21.9	1.7	*	1.4	3.5	6.7	14.1	24.1	42.1	76.5
2001.....	215.5	22.9	3.1	0.8	1.3	3.5	7.8	13.5	23.5	41.3	81.6
2000.....	218.6	20.5	2.0	0.9	1.4	4.2	7.8	13.4	22.3	39.4	78.8
1999.....	223.8	24.8	2.1	1.0	1.4	4.3	8.0	12.3	22.5	43.8	84.7

Non-Hispanic Black, male

2003.....	212.6	24.5	2.5	*	1.4	4.6	10.4	16.6	29.5	53.3	105.4
2002.....	212.9	23.0	*	*	1.6	4.4	8.3	18.5	31.1	56.0	99.1
2001.....	215.7	20.7	3.4	*	1.4	4.5	10.0	16.7	30.3	54.1	104.2
2000.....	217.6	20.1	2.0	*	1.5	4.9	9.2	16.4	27.6	50.0	102.5
1999.....	224.9	24.6	2.3	*	1.7	6.0	9.7	15.6	28.4	57.9	111.5

Non-Hispanic Black, female

2003.....	206.5	16.1	2.0	*	*	2.4	5.7	10.8	18.6	30.3	59.6
2002.....	211.6	20.8	1.9	*	*	2.5	5.3	10.1	17.7	29.7	56.6
2001.....	215.4	25.2	2.9	*	*	2.4	5.8	10.7	17.5	30.0	61.6
2000.....	219.5	20.8	2.0	*	1.4	3.4	6.5	10.6	17.5	30.0	58.0
1999.....	222.8	25.0	1.9	*	*	2.6	6.5	9.4	17.2	31.4	61.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Hispanic, female

2003.....	0.5	*	*	*	*	*	*	*	*	*	*
2002.....	0.8	*	*	*	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*	*	*	*
2000.....	0.6	*	*	*	*	*	*	*	*	*	*
1999.....	0.8	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	1.4	*	*	*	*	*	*	*	*	0.2	0.3
2002.....	1.4	*	*	*	*	*	*	*	*	0.2	0.3
2001.....	1.3	*	*	*	*	*	*	*	0.1	0.2	0.4
2000.....	1.4	*	*	*	*	*	*	*	*	0.2	0.4
1999.....	1.4	*	*	*	*	*	*	*	0.2	0.2	0.3

Non-Hispanic 4/, male

2003.....	0.9	*	*	*	*	*	*	*	*	*	0.2
2002.....	0.8	*	*	*	*	*	*	*	*	*	0.2
2001.....	0.8	*	*	*	*	*	*	*	*	*	0.3
2000.....	0.9	*	*	*	*	*	*	*	*	0.2	0.3
1999.....	0.8	*	*	*	*	*	*	*	*	*	0.3

Non-Hispanic 4/, female

2003.....	1.8	*	*	*	*	*	*	*	*	0.2	0.3
2002.....	1.8	*	*	*	*	*	*	*	*	*	0.3
2001.....	1.8	*	*	*	*	*	*	*	*	0.2	0.4
2000.....	1.9	*	*	*	*	*	*	*	*	*	0.5
1999.....	2.0	*	*	*	*	*	*	*	*	0.2	0.4

Non-Hispanic White, both sexes

2003.....	1.5	*	*	*	*	*	*	*	*	0.2	0.2
2002.....	1.5	*	*	*	*	*	*	*	*	0.2	0.2
2001.....	1.5	*	*	*	*	*	*	*	*	0.2	0.3
2000.....	1.6	*	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.6	*	*	*	*	*	*	*	*	0.1	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Hypertensive heart disease (I11)											
Hispanic, female											
2003.....	3.7	*	*	*	*	*	*	*	*	*	*
2002.....	3.4	*	*	*	*	*	*	*	*	*	*
2001.....	3.1	*	*	*	*	*	*	*	*	*	*
2000.....	2.9	*	*	*	*	*	*	*	*	*	*
1999.....	2.9	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	10.6	*	*	*	*	*	0.2	0.5	1.2	2.0	4.1
2002.....	10.0	*	*	*	*	*	0.1	0.5	1.1	1.9	3.7
2001.....	9.4	*	*	*	*	*	*	0.4	1.0	1.6	3.4
2000.....	9.2	*	*	*	*	*	0.2	0.4	0.8	1.5	3.0
1999.....	8.8	*	*	*	*	*	0.2	0.3	0.8	1.6	2.9
Non-Hispanic 4/, male											
2003.....	9.5	*	*	*	*	*	0.3	0.7	1.5	2.8	5.3
2002.....	8.9	*	*	*	*	*	*	0.8	1.5	2.8	5.0
2001.....	8.1	*	*	*	*	*	*	0.4	1.4	2.1	4.6
2000.....	7.8	*	*	*	*	*	0.3	0.4	1.0	1.9	4.0
1999.....	7.4	*	*	*	*	*	*	0.4	1.1	2.1	4.1
Non-Hispanic 4/, female											
2003.....	11.7	*	*	*	*	*	*	0.3	0.8	1.3	2.8
2002.....	11.1	*	*	*	*	*	*	*	0.7	1.1	2.3
2001.....	10.7	*	*	*	*	*	*	0.3	0.6	1.1	2.1
2000.....	10.6	*	*	*	*	*	*	0.3	0.5	1.0	2.1
1999.....	10.2	*	*	*	*	*	*	*	0.6	1.1	1.8
Non-Hispanic White, both sexes											
2003.....	9.7	*	*	*	*	*	*	0.3	0.7	1.3	2.7
2002.....	9.1	*	*	*	*	*	*	0.3	0.6	1.3	2.5
2001.....	8.6	*	*	*	*	*	*	0.2	0.6	1.0	2.1
2000.....	8.3	*	*	*	*	*	*	0.2	0.4	0.9	2.0
1999.....	7.9	*	*	*	*	*	*	*	0.5	0.9	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Hypertensive heart disease (I11)											
Non-Hispanic White, male											
2003.....	8.2	*	*	*	*	*	*	0.6	1.0	2.0	3.9
2002.....	7.6	*	*	*	*	*	*	0.4	0.9	1.9	3.7
2001.....	6.9	*	*	*	*	*	*	*	0.9	1.5	3.1
2000.....	6.6	*	*	*	*	*	*	*	0.6	1.3	2.9
1999.....	6.2	*	*	*	*	*	*	*	0.8	1.2	2.7
Non-Hispanic White, female											
2003.....	11.1	*	*	*	*	*	*	*	0.4	0.7	1.4
2002.....	10.5	*	*	*	*	*	*	*	0.3	0.7	1.2
2001.....	10.2	*	*	*	*	*	*	*	*	0.6	1.0
2000.....	9.9	*	*	*	*	*	*	*	*	0.5	1.2
1999.....	9.6	*	*	*	*	*	*	*	*	0.5	0.8
Non-Hispanic Black, both sexes											
2003.....	18.5	*	*	*	*	*	0.7	1.5	3.6	6.2	13.2
2002.....	17.7	*	*	*	*	*	*	1.6	4.1	6.0	11.6
2001.....	16.5	*	*	*	*	*	*	1.1	3.3	5.2	11.8
2000.....	16.6	*	*	*	*	*	*	1.4	2.9	5.1	9.9
1999.....	16.0	*	*	*	*	*	*	1.0	2.7	5.9	10.4
Non-Hispanic Black, male											
2003.....	18.7	*	*	*	*	*	*	1.8	4.5	7.6	15.4
2002.....	18.0	*	*	*	*	*	*	2.5	5.6	8.7	14.7
2001.....	16.3	*	*	*	*	*	*	*	4.2	6.0	15.1
2000.....	16.1	*	*	*	*	*	*	1.6	3.7	6.0	12.6
1999.....	15.6	*	*	*	*	*	*	*	2.9	7.6	13.3
Non-Hispanic Black, female											
2003.....	18.4	*	*	*	*	*	*	*	2.8	4.9	11.3
2002.....	17.4	*	*	*	*	*	*	*	2.8	3.5	8.9
2001.....	16.7	*	*	*	*	*	*	*	2.4	4.6	8.9
2000.....	17.1	*	*	*	*	*	*	*	2.2	4.3	7.6
1999.....	16.4	*	*	*	*	*	*	*	2.6	4.5	7.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Ischemic heart diseases (I20-I25)

All origins 3/, both sexes

2003.....	165.1	0.5	*	*	*	0.2	0.6	1.4	3.9	10.1	23.5
2002.....	171.4	0.6	*	*	*	0.1	0.5	1.3	3.6	10.1	23.8
2001.....	176.3	0.6	*	*	*	0.2	0.7	1.5	4.0	9.9	23.4
2000.....	183.1	0.8	*	*	*	0.2	0.5	1.2	3.4	9.8	23.6
1999.....	189.8	0.7	*	*	*	0.3	0.5	1.3	3.7	10.4	24.2

All origins 3/, male

2003.....	172.2	*	*	*	*	0.2	0.8	2.0	5.9	15.2	35.5
2002.....	178.4	*	*	*	*	*	0.7	2.0	5.2	15.2	36.5
2001.....	181.7	*	*	*	*	0.2	0.9	2.2	5.9	15.0	35.8
2000.....	188.7	*	*	*	*	0.2	0.6	1.8	5.1	14.7	37.0
1999.....	195.4	*	*	*	*	0.3	0.6	1.7	5.5	15.8	37.9

All origins 3/, female

2003.....	158.1	*	*	*	*	*	0.4	0.8	1.9	4.9	11.6
2002.....	164.7	*	*	*	*	*	0.3	0.6	1.9	4.9	11.2
2001.....	171.2	*	*	*	*	*	0.4	0.7	2.1	4.9	11.1
2000.....	177.6	*	*	*	*	*	0.3	0.6	1.7	5.0	10.3
1999.....	184.5	*	*	*	*	*	0.3	0.8	1.9	5.1	10.7

Hispanic, both sexes

2003.....	52.1	*	*	*	*	*	*	0.8	2.0	4.8	10.8
2002.....	54.0	*	*	*	*	*	*	0.8	1.6	4.6	11.2
2001.....	55.9	*	*	*	*	*	*	0.9	2.3	4.3	9.7
2000.....	55.9	*	*	*	*	*	*	0.6	1.6	4.3	11.8
1999.....	58.7	*	*	*	*	*	*	0.7	2.1	4.7	10.8

Hispanic, male

2003.....	53.7	*	*	*	*	*	*	1.2	3.0	6.8	15.9
2002.....	56.5	*	*	*	*	*	*	1.3	2.6	6.4	17.5
2001.....	57.9	*	*	*	*	*	*	1.4	3.6	6.7	15.0
2000.....	58.2	*	*	*	*	*	*	*	2.6	6.6	19.0
1999.....	60.8	*	*	*	*	*	*	1.3	3.0	7.9	17.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Ischemic heart diseases (I20-I25)

Hispanic, female

2003.....	50.4	*	*	*	*	*	*	*	*	2.4	5.2
2002.....	51.4	*	*	*	*	*	*	*	*	2.5	4.6
2001.....	53.7	*	*	*	*	*	*	*	*	1.6	4.1
2000.....	53.6	*	*	*	*	*	*	*	*	1.7	4.2
1999.....	56.6	*	*	*	*	*	*	*	*	*	3.7

Non-Hispanic 4/, both sexes

2003.....	182.5	0.6	*	*	*	0.2	0.6	1.6	4.3	10.9	25.1
2002.....	189.0	*	*	*	*	0.2	0.5	1.4	3.9	11.0	25.3
2001.....	193.6	0.7	*	*	*	0.2	0.7	1.6	4.4	10.8	24.9
2000.....	200.5	0.8	*	*	*	0.2	0.5	1.3	3.7	10.6	24.7
1999.....	207.1	0.8	*	*	*	0.3	0.5	1.3	4.0	11.1	25.5

Non-Hispanic 4/, male

2003.....	191.5	*	*	*	*	*	0.8	2.2	6.5	16.7	38.1
2002.....	197.6	*	*	*	*	*	0.7	2.2	5.7	16.8	39.0
2001.....	200.4	*	*	*	*	0.2	1.0	2.4	6.4	16.3	38.2
2000.....	207.6	*	*	*	*	*	0.7	2.1	5.5	15.8	38.9
1999.....	214.1	*	*	*	*	0.4	0.7	1.8	6.0	16.8	40.0

Non-Hispanic 4/, female

2003.....	173.9	*	*	*	*	*	0.5	0.9	2.0	5.3	12.4
2002.....	180.7	*	*	*	*	*	0.3	0.7	2.1	5.3	12.0
2001.....	187.2	*	*	*	*	*	0.4	0.8	2.3	5.3	11.9
2000.....	193.8	*	*	*	*	*	0.3	0.6	1.9	5.4	10.9
1999.....	200.6	*	*	*	*	*	0.3	0.9	2.1	5.6	11.4

Non-Hispanic White, both sexes

2003.....	200.8	*	*	*	*	0.2	0.6	1.4	4.1	10.8	24.1
2002.....	208.0	*	*	*	*	0.2	0.4	1.3	3.7	10.6	24.5
2001.....	212.6	*	*	*	*	*	0.6	1.4	4.1	10.4	23.6
2000.....	220.2	*	*	*	*	0.2	0.5	1.1	3.5	10.2	23.4
1999.....	227.2	*	*	*	*	0.2	0.4	1.2	3.7	10.7	23.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Ischemic heart diseases (I20-I25)											
Non-Hispanic White, male											
2003.....	210.2	*	*	*	*	*	0.7	2.0	6.3	16.7	37.1
2002.....	217.2	*	*	*	*	*	0.6	2.1	5.5	16.3	38.5
2001.....	219.6	*	*	*	*	*	0.9	2.2	6.0	15.8	37.2
2000.....	227.6	*	*	*	*	*	0.6	1.8	5.3	15.5	37.8
1999.....	234.2	*	*	*	*	0.3	0.6	1.6	5.5	16.2	38.3
Non-Hispanic White, female											
2003.....	191.8	*	*	*	*	*	0.4	0.8	1.9	4.8	11.1
2002.....	199.1	*	*	*	*	*	*	0.6	1.9	4.8	10.5
2001.....	205.8	*	*	*	*	*	*	0.6	2.1	4.9	10.1
2000.....	213.1	*	*	*	*	*	*	0.5	1.6	4.9	9.0
1999.....	220.5	*	*	*	*	*	*	0.9	1.9	5.1	9.6
Non-Hispanic Black, both sexes											
2003.....	133.2	*	*	*	*	*	1.2	2.5	6.2	14.6	36.0
2002.....	137.0	*	*	*	*	*	0.9	2.2	5.8	15.5	36.1
2001.....	141.2	*	*	*	*	*	1.5	2.9	6.6	15.4	37.5
2000.....	144.4	*	*	*	*	*	0.8	2.6	6.2	15.0	37.7
1999.....	147.9	*	*	*	*	*	1.2	2.0	6.1	16.4	40.7
Non-Hispanic Black, male											
2003.....	137.6	*	*	*	*	*	1.5	3.3	9.5	20.4	51.5
2002.....	139.5	*	*	*	*	*	*	2.9	7.7	22.5	49.8
2001.....	143.2	*	*	*	*	*	2.0	4.0	9.5	22.5	52.0
2000.....	145.7	*	*	*	*	*	*	3.9	8.2	20.8	52.5
1999.....	150.0	*	*	*	*	*	*	3.1	8.8	23.8	58.1
Non-Hispanic Black, female											
2003.....	129.1	*	*	*	*	*	*	1.8	3.3	9.4	22.4
2002.....	134.8	*	*	*	*	*	*	1.6	4.1	9.3	23.9
2001.....	139.4	*	*	*	*	*	*	1.8	4.0	9.2	24.7
2000.....	143.2	*	*	*	*	*	*	*	4.4	9.9	24.7
1999.....	146.0	*	*	*	*	*	*	*	3.6	9.9	25.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Acute myocardial infarction (I21-I22)											
Hispanic, female											
2003.....	18.0	*	*	*	*	*	*	*	*	*	2.3
2002.....	18.7	*	*	*	*	*	*	*	*	1.3	2.4
2001.....	19.1	*	*	*	*	*	*	*	*	*	*
2000.....	19.1	*	*	*	*	*	*	*	*	*	*
1999.....	20.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	64.9	*	*	*	*	*	0.3	0.6	1.6	4.5	10.3
2002.....	68.7	*	*	*	*	*	0.3	0.5	1.6	4.4	10.6
2001.....	71.4	*	*	*	*	*	0.3	0.6	1.9	4.6	10.6
2000.....	75.3	*	*	*	*	*	0.2	0.6	1.4	4.3	10.5
1999.....	78.2	*	*	*	*	0.1	0.2	0.6	1.8	4.4	11.0
Non-Hispanic 4/, male											
2003.....	69.7	*	*	*	*	*	0.3	0.9	2.4	6.9	15.5
2002.....	73.5	*	*	*	*	*	0.3	0.8	2.3	6.6	15.9
2001.....	75.9	*	*	*	*	*	0.4	0.8	2.7	6.9	16.0
2000.....	80.3	*	*	*	*	*	*	0.9	2.0	6.3	16.4
1999.....	83.8	*	*	*	*	*	0.3	0.7	2.5	6.6	17.0
Non-Hispanic 4/, female											
2003.....	60.3	*	*	*	*	*	0.2	0.4	0.9	2.2	5.3
2002.....	64.1	*	*	*	*	*	*	0.3	0.9	2.2	5.4
2001.....	67.1	*	*	*	*	*	*	0.4	1.1	2.3	5.3
2000.....	70.6	*	*	*	*	*	*	0.3	0.8	2.3	4.7
1999.....	72.9	*	*	*	*	*	*	0.4	1.0	2.2	5.3
Non-Hispanic White, both sexes											
2003.....	71.5	*	*	*	*	*	0.2	0.6	1.6	4.6	10.1
2002.....	75.5	*	*	*	*	*	0.2	0.5	1.6	4.3	10.4
2001.....	78.4	*	*	*	*	*	0.2	0.5	1.9	4.5	10.3
2000.....	82.7	*	*	*	*	*	0.2	0.5	1.4	4.4	10.1
1999.....	86.0	*	*	*	*	*	*	0.5	1.7	4.4	10.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Acute myocardial infarction (I21-I22)											
Non-Hispanic White, male											
2003.....	77.0	*	*	*	*	*	*	0.9	2.3	7.1	15.3
2002.....	81.2	*	*	*	*	*	*	0.7	2.3	6.7	15.9
2001.....	83.7	*	*	*	*	*	0.3	0.6	2.7	6.8	16.1
2000.....	88.5	*	*	*	*	*	*	0.7	2.0	6.5	16.2
1999.....	92.4	*	*	*	*	*	*	0.7	2.3	6.7	16.8
Non-Hispanic White, female											
2003.....	66.2	*	*	*	*	*	*	*	0.9	2.0	4.8
2002.....	70.1	*	*	*	*	*	*	*	0.8	2.0	5.0
2001.....	73.4	*	*	*	*	*	*	0.4	1.1	2.2	4.5
2000.....	77.2	*	*	*	*	*	*	*	0.7	2.2	3.9
1999.....	79.8	*	*	*	*	*	*	0.4	1.0	2.0	4.6
Non-Hispanic Black, both sexes											
2003.....	47.3	*	*	*	*	*	*	0.9	2.4	5.9	14.8
2002.....	50.2	*	*	*	*	*	*	1.0	2.4	5.7	14.2
2001.....	51.6	*	*	*	*	*	*	1.1	2.6	6.0	14.6
2000.....	54.0	*	*	*	*	*	*	1.2	2.3	5.1	15.3
1999.....	55.1	*	*	*	*	*	*	0.9	2.5	5.6	15.8
Non-Hispanic Black, male											
2003.....	48.2	*	*	*	*	*	*	*	3.7	7.9	20.6
2002.....	50.1	*	*	*	*	*	*	*	3.1	7.8	19.6
2001.....	50.9	*	*	*	*	*	*	*	3.5	8.3	18.8
2000.....	53.7	*	*	*	*	*	*	1.7	2.7	6.7	20.9
1999.....	55.2	*	*	*	*	*	*	*	3.6	7.8	21.6
Non-Hispanic Black, female											
2003.....	46.5	*	*	*	*	*	*	*	*	4.0	9.6
2002.....	50.4	*	*	*	*	*	*	*	1.7	3.7	9.5
2001.....	52.3	*	*	*	*	*	*	*	1.8	3.9	10.9
2000.....	54.2	*	*	*	*	*	*	*	1.9	3.7	10.3
1999.....	54.9	*	*	*	*	*	*	*	1.5	3.6	10.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other forms of chronic ischemic heart disease (I20,I25)

All origins 3/, both sexes

2003.....	105.3	*	*	*	*	*	0.3	0.9	2.4	5.8	13.5
2002.....	108.0	*	*	*	*	*	0.3	0.8	2.0	6.0	13.6
2001.....	110.3	*	*	*	*	*	0.4	0.9	2.2	5.6	13.2
2000.....	113.3	*	*	*	*	*	0.3	0.6	2.0	5.7	13.4
1999.....	117.2	*	*	*	*	0.1	0.3	0.7	2.1	6.2	13.5

All origins 3/, male

2003.....	108.5	*	*	*	*	*	0.4	1.2	3.7	8.8	20.7
2002.....	110.9	*	*	*	*	*	0.4	1.3	3.0	9.2	21.4
2001.....	111.9	*	*	*	*	*	0.5	1.5	3.3	8.6	20.4
2000.....	114.8	*	*	*	*	*	0.4	1.1	3.1	8.7	21.1
1999.....	118.0	*	*	*	*	0.2	0.3	1.0	3.2	9.4	21.5

All origins 3/, female

2003.....	102.3	*	*	*	*	*	0.2	0.5	1.0	2.8	6.5
2002.....	105.2	*	*	*	*	*	*	0.4	1.0	2.8	6.0
2001.....	108.8	*	*	*	*	*	0.2	0.3	1.1	2.7	6.1
2000.....	111.9	*	*	*	*	*	*	0.2	1.0	2.8	5.8
1999.....	116.4	*	*	*	*	*	*	0.4	1.0	3.1	5.6

Hispanic, both sexes

2003.....	33.5	*	*	*	*	*	*	0.6	1.2	2.9	6.5
2002.....	34.2	*	*	*	*	*	*	*	1.0	2.7	6.7
2001.....	35.7	*	*	*	*	*	*	*	1.3	2.8	5.9
2000.....	35.6	*	*	*	*	*	*	*	1.0	2.6	7.1
1999.....	37.5	*	*	*	*	*	*	*	1.3	3.0	6.4

Hispanic, male

2003.....	34.6	*	*	*	*	*	*	*	1.9	4.3	9.9
2002.....	35.7	*	*	*	*	*	*	*	1.5	4.1	11.1
2001.....	36.8	*	*	*	*	*	*	*	2.0	4.6	8.9
2000.....	36.9	*	*	*	*	*	*	*	1.5	4.3	11.3
1999.....	38.7	*	*	*	*	*	*	*	2.0	5.2	10.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other forms of chronic ischemic heart disease (I20,I25)

Hispanic, female

2003.....	32.3	*	*	*	*	*	*	*	*	*	*	2.9
2002.....	32.6	*	*	*	*	*	*	*	*	*	*	2.2
2001.....	34.5	*	*	*	*	*	*	*	*	*	*	2.7
2000.....	34.2	*	*	*	*	*	*	*	*	*	*	2.7
1999.....	36.3	*	*	*	*	*	*	*	*	*	*	2.0

Non-Hispanic 4/, both sexes

2003.....	116.3	*	*	*	*	*	0.3	0.9	2.6	6.3	14.4
2002.....	118.9	*	*	*	*	*	0.3	0.9	2.2	6.5	14.5
2001.....	120.9	*	*	*	*	*	0.4	1.0	2.4	6.1	14.0
2000.....	123.9	*	*	*	*	*	0.3	0.7	2.2	6.1	14.0
1999.....	127.6	*	*	*	*	0.1	0.3	0.8	2.2	6.6	14.2

Non-Hispanic 4/, male

2003.....	120.4	*	*	*	*	*	0.4	1.3	4.1	9.6	22.1
2002.....	122.7	*	*	*	*	*	0.4	1.3	3.3	10.0	22.7
2001.....	123.1	*	*	*	*	*	0.6	1.6	3.6	9.3	21.8
2000.....	126.0	*	*	*	*	*	0.4	1.2	3.4	9.3	22.0
1999.....	128.9	*	*	*	*	*	0.4	1.1	3.4	10.0	22.6

Non-Hispanic 4/, female

2003.....	112.4	*	*	*	*	*	0.2	0.5	1.2	3.0	6.9
2002.....	115.4	*	*	*	*	*	*	0.4	1.1	3.1	6.4
2001.....	118.8	*	*	*	*	*	*	0.4	1.1	2.9	6.5
2000.....	122.0	*	*	*	*	*	*	*	1.1	3.0	6.1
1999.....	126.4	*	*	*	*	*	*	0.5	1.1	3.4	6.0

Non-Hispanic White, both sexes

2003.....	128.0	*	*	*	*	*	0.3	0.8	2.5	6.1	13.8
2002.....	131.0	*	*	*	*	*	0.2	0.9	2.1	6.1	13.8
2001.....	132.7	*	*	*	*	*	0.3	0.9	2.1	5.7	13.1
2000.....	136.1	*	*	*	*	*	0.2	0.6	2.0	5.7	13.1
1999.....	139.9	*	*	*	*	*	0.2	0.7	2.0	6.2	13.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic White, male

2003.....	131.9	*	*	*	*	*	0.4	1.1	3.9	9.3	21.3
2002.....	134.5	*	*	*	*	*	0.4	1.3	3.1	9.5	22.2
2001.....	134.4	*	*	*	*	*	0.5	1.6	3.2	8.8	20.7
2000.....	137.7	*	*	*	*	*	0.4	1.0	3.2	8.8	21.2
1999.....	140.4	*	*	*	*	*	*	0.9	3.1	9.3	21.1

Non-Hispanic White, female

2003.....	124.3	*	*	*	*	*	*	0.5	1.0	2.8	6.2
2002.....	127.7	*	*	*	*	*	*	0.4	1.0	2.8	5.5
2001.....	131.1	*	*	*	*	*	*	*	1.0	2.6	5.5
2000.....	134.6	*	*	*	*	*	*	*	0.9	2.6	4.9
1999.....	139.3	*	*	*	*	*	*	0.5	0.9	3.1	4.8

Non-Hispanic Black, both sexes

2003.....	84.7	*	*	*	*	*	*	1.5	3.8	8.5	20.6
2002.....	85.5	*	*	*	*	*	*	1.2	3.3	9.8	21.2
2001.....	88.2	*	*	*	*	*	0.8	1.8	3.9	9.2	22.2
2000.....	89.1	*	*	*	*	*	*	1.3	3.7	9.7	21.9
1999.....	91.6	*	*	*	*	*	*	1.2	3.5	10.4	24.4

Non-Hispanic Black, male

2003.....	87.9	*	*	*	*	*	*	2.2	5.7	12.2	30.1
2002.....	88.0	*	*	*	*	*	*	*	4.3	14.6	29.9
2001.....	90.8	*	*	*	*	*	*	2.5	6.0	13.8	32.1
2000.....	90.4	*	*	*	*	*	*	2.2	5.1	13.7	30.8
1999.....	93.3	*	*	*	*	*	*	1.8	5.2	15.3	35.6

Non-Hispanic Black, female

2003.....	81.7	*	*	*	*	*	*	*	2.0	5.1	12.2
2002.....	83.3	*	*	*	*	*	*	*	2.4	5.5	13.6
2001.....	86.0	*	*	*	*	*	*	*	2.0	5.0	13.4
2000.....	88.0	*	*	*	*	*	*	*	2.5	6.0	14.1
1999.....	90.1	*	*	*	*	*	*	*	2.0	6.0	14.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Atherosclerotic cardiovascular disease, so described (I25.0)											
All origins 3/, both sexes											
2003.....	23.1	*	*	*	*	*	*	0.3	0.8	1.9	4.7
2002.....	23.6	*	*	*	*	*	*	0.2	0.7	1.9	4.8
2001.....	23.9	*	*	*	*	*	*	0.2	0.7	1.8	4.6
2000.....	24.7	*	*	*	*	*	*	0.2	0.5	1.7	4.4
1999.....	25.8	*	*	*	*	*	*	0.2	0.6	2.0	4.7
All origins 3/, male											
2003.....	25.2	*	*	*	*	*	*	0.4	1.3	2.9	7.4
2002.....	25.4	*	*	*	*	*	*	0.3	1.1	3.0	7.8
2001.....	25.3	*	*	*	*	*	*	0.3	1.0	2.8	7.2
2000.....	26.0	*	*	*	*	*	*	0.3	0.9	2.8	7.1
1999.....	26.8	*	*	*	*	*	*	0.2	0.9	3.1	7.6
All origins 3/, female											
2003.....	21.1	*	*	*	*	*	*	*	0.2	0.9	2.0
2002.....	21.9	*	*	*	*	*	*	*	0.3	0.8	1.8
2001.....	22.5	*	*	*	*	*	*	*	0.4	0.8	2.1
2000.....	23.5	*	*	*	*	*	*	*	0.2	0.7	1.8
1999.....	24.8	*	*	*	*	*	*	*	0.3	0.9	1.8
Hispanic, both sexes											
2003.....	8.3	*	*	*	*	*	*	*	*	0.9	2.3
2002.....	8.2	*	*	*	*	*	*	*	*	1.0	2.9
2001.....	8.5	*	*	*	*	*	*	*	*	0.9	2.2
2000.....	8.4	*	*	*	*	*	*	*	*	1.0	2.4
1999.....	9.0	*	*	*	*	*	*	*	*	0.8	2.3
Hispanic, male											
2003.....	10.2	*	*	*	*	*	*	*	*	1.5	3.9
2002.....	9.7	*	*	*	*	*	*	*	*	1.3	4.9
2001.....	10.1	*	*	*	*	*	*	*	*	1.4	3.2
2000.....	9.9	*	*	*	*	*	*	*	*	1.7	3.9
1999.....	10.6	*	*	*	*	*	*	*	*	1.5	3.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Atherosclerotic cardiovascular disease, so described (I25.0)

Hispanic, female

2003.....	6.2	*	*	*	*	*	*	*	*	*	*	*
2002.....	6.5	*	*	*	*	*	*	*	*	*	*	*
2001.....	6.8	*	*	*	*	*	*	*	*	*	*	*
2000.....	6.8	*	*	*	*	*	*	*	*	*	*	*
1999.....	7.4	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	25.3	*	*	*	*	*	*	0.3	0.8	2.1	5.0
2002.....	25.9	*	*	*	*	*	*	0.2	0.7	2.0	5.0
2001.....	26.0	*	*	*	*	*	*	0.2	0.7	1.9	4.9
2000.....	26.9	*	*	*	*	*	*	0.2	0.6	1.8	4.6
1999.....	27.9	*	*	*	*	*	*	0.2	0.7	2.1	4.9

Non-Hispanic 4/, male

2003.....	27.5	*	*	*	*	*	*	0.4	1.5	3.2	7.8
2002.....	27.8	*	*	*	*	*	*	0.3	1.1	3.2	8.2
2001.....	27.5	*	*	*	*	*	*	0.3	1.0	3.1	7.6
2000.....	28.2	*	*	*	*	*	*	0.3	1.0	2.9	7.4
1999.....	28.9	*	*	*	*	*	*	*	1.0	3.3	8.0

Non-Hispanic 4/, female

2003.....	23.3	*	*	*	*	*	*	*	*	1.0	2.2
2002.....	24.0	*	*	*	*	*	*	*	0.3	0.8	2.0
2001.....	24.6	*	*	*	*	*	*	*	0.4	0.8	2.2
2000.....	25.6	*	*	*	*	*	*	*	0.3	0.7	1.9
1999.....	27.0	*	*	*	*	*	*	*	0.3	1.0	1.9

Non-Hispanic White, both sexes

2003.....	26.2	*	*	*	*	*	*	0.3	0.8	2.1	4.7
2002.....	26.8	*	*	*	*	*	*	0.2	0.6	1.8	4.6
2001.....	26.8	*	*	*	*	*	*	0.2	0.7	1.8	4.4
2000.....	27.6	*	*	*	*	*	*	*	0.6	1.6	4.2
1999.....	28.6	*	*	*	*	*	*	*	0.6	1.9	4.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Atherosclerotic cardiovascular disease, so described (I25.0)											
Non-Hispanic White, male											
2003.....	28.0	*	*	*	*	*	*	0.4	1.4	3.1	7.3
2002.....	28.2	*	*	*	*	*	*	*	1.0	2.9	7.7
2001.....	27.8	*	*	*	*	*	*	*	1.0	2.9	7.0
2000.....	28.4	*	*	*	*	*	*	*	0.9	2.7	6.9
1999.....	29.0	*	*	*	*	*	*	*	0.9	3.0	7.3
Non-Hispanic White, female											
2003.....	24.6	*	*	*	*	*	*	*	*	1.0	2.0
2002.....	25.3	*	*	*	*	*	*	*	*	0.7	1.6
2001.....	25.8	*	*	*	*	*	*	*	0.3	0.7	1.8
2000.....	26.9	*	*	*	*	*	*	*	*	0.6	1.5
1999.....	28.2	*	*	*	*	*	*	*	*	0.8	1.4
Non-Hispanic Black, both sexes											
2003.....	26.7	*	*	*	*	*	*	*	1.3	2.7	7.4
2002.....	27.7	*	*	*	*	*	*	*	1.3	3.6	8.3
2001.....	28.5	*	*	*	*	*	*	*	1.0	3.2	8.5
2000.....	29.3	*	*	*	*	*	*	*	0.9	3.0	7.6
1999.....	30.7	*	*	*	*	*	*	*	1.2	3.9	9.3
Non-Hispanic Black, male											
2003.....	31.2	*	*	*	*	*	*	*	2.3	4.0	11.6
2002.....	31.9	*	*	*	*	*	*	*	1.9	5.6	12.2
2001.....	32.3	*	*	*	*	*	*	*	*	4.8	12.8
2000.....	33.2	*	*	*	*	*	*	*	*	4.8	11.2
1999.....	34.5	*	*	*	*	*	*	*	1.8	6.0	13.9
Non-Hispanic Black, female											
2003.....	22.7	*	*	*	*	*	*	*	*	1.5	3.8
2002.....	24.0	*	*	*	*	*	*	*	*	1.8	5.0
2001.....	25.0	*	*	*	*	*	*	*	*	1.7	4.7
2000.....	25.8	*	*	*	*	*	*	*	*	1.5	4.4
1999.....	27.1	*	*	*	*	*	*	*	*	2.0	5.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All origins 3/, both sexes

2003.....	82.2	*	*	*	*	*	0.3	0.6	1.6	3.9	8.9
2002.....	84.4	*	*	*	*	*	0.2	0.6	1.3	4.1	8.8
2001.....	86.4	*	*	*	*	*	0.3	0.7	1.5	3.8	8.6
2000.....	88.6	*	*	*	*	*	0.2	0.5	1.5	4.0	9.0
1999.....	91.4	*	*	*	*	0.1	0.2	0.6	1.5	4.3	8.8

All origins 3/, male

2003.....	83.3	*	*	*	*	*	0.3	0.8	2.3	5.9	13.3
2002.....	85.5	*	*	*	*	*	0.3	0.9	1.9	6.2	13.6
2001.....	86.5	*	*	*	*	*	0.4	1.1	2.3	5.8	13.2
2000.....	88.8	*	*	*	*	*	0.2	0.8	2.2	5.9	14.0
1999.....	91.2	*	*	*	*	0.2	0.2	0.8	2.2	6.3	13.9

All origins 3/, female

2003.....	81.1	*	*	*	*	*		0.4	0.8	1.9	4.5
2002.....	83.3	*	*	*	*	*		0.3	0.8	2.0	4.2
2001.....	86.2	*	*	*	*	*		0.2	0.7	1.9	4.0
2000.....	88.4	*	*	*	*	*		*	0.8	2.1	4.0
1999.....	91.6	*	*	*	*	*		0.3	0.7	2.2	3.9

Hispanic, both sexes

2003.....	25.2	*	*	*	*	*	*	*	0.8	2.0	4.2
2002.....	26.0	*	*	*	*	*	*	*	*	1.7	3.8
2001.....	27.2	*	*	*	*	*	*	*	0.8	1.9	3.6
2000.....	27.2	*	*	*	*	*	*	*	0.8	1.6	4.7
1999.....	28.5	*	*	*	*	*	*	*	1.0	2.2	4.1

Hispanic, male

2003.....	24.4	*	*	*	*	*	*	*	1.1	2.9	6.0
2002.....	25.9	*	*	*	*	*	*	*	*	2.8	6.1
2001.....	26.7	*	*	*	*	*	*	*	1.3	3.1	5.7
2000.....	27.0	*	*	*	*	*	*	*	1.2	2.6	7.4
1999.....	28.1	*	*	*	*	*	*	*	1.4	3.7	6.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)											
Hispanic, female											
2003.....	26.1	*	*	*	*	*	*	*	*	*	2.3
2002.....	26.1	*	*	*	*	*	*	*	*	*	*
2001.....	27.7	*	*	*	*	*	*	*	*	*	*
2000.....	27.5	*	*	*	*	*	*	*	*	*	1.9
1999.....	28.9	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	91.0	*	*	*	*	*	0.3	0.6	1.8	4.2	9.5
2002.....	93.1	*	*	*	*	*	0.2	0.7	1.5	4.5	9.4
2001.....	94.9	*	*	*	*	*	0.3	0.8	1.6	4.1	9.2
2000.....	97.0	*	*	*	*	*	0.2	0.5	1.6	4.3	9.4
1999.....	99.7	*	*	*	*	0.1	0.2	0.6	1.6	4.5	9.3
Non-Hispanic 4/, male											
2003.....	92.9	*	*	*	*	*	0.3	0.9	2.6	6.4	14.3
2002.....	94.9	*	*	*	*	*	0.3	1.1	2.2	6.8	14.5
2001.....	95.6	*	*	*	*	*	0.4	1.3	2.5	6.2	14.1
2000.....	97.8	*	*	*	*	*	0.3	0.9	2.4	6.4	14.7
1999.....	100.0	*	*	*	*	*	*	0.8	2.4	6.6	14.6
Non-Hispanic 4/, female											
2003.....	89.2	*	*	*	*	*	*	0.4	0.9	2.0	4.7
2002.....	91.4	*	*	*	*	*	*	0.3	0.8	2.3	4.5
2001.....	94.2	*	*	*	*	*	*	0.3	0.8	2.1	4.3
2000.....	96.3	*	*	*	*	*	*	*	0.8	2.3	4.2
1999.....	99.4	*	*	*	*	*	*	0.4	0.7	2.4	4.1
Non-Hispanic White, both sexes											
2003.....	101.8	*	*	*	*	*	0.2	0.5	1.7	4.0	9.1
2002.....	104.3	*	*	*	*	*	0.2	0.7	1.4	4.3	9.2
2001.....	106.0	*	*	*	*	*	0.2	0.7	1.5	3.9	8.7
2000.....	108.4	*	*	*	*	*	*	0.5	1.5	4.0	8.9
1999.....	111.3	*	*	*	*	*	*	0.6	1.5	4.3	8.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic White, male

2003.....	103.9	*	*	*	*	*	*	0.7	2.5	6.2	14.0
2002.....	106.2	*	*	*	*	*	*	1.1	2.1	6.6	14.5
2001.....	106.6	*	*	*	*	*	0.4	1.3	2.3	5.9	13.7
2000.....	109.2	*	*	*	*	*	*	0.8	2.3	6.1	14.3
1999.....	111.4	*	*	*	*	*	*	0.7	2.3	6.3	13.8

Non-Hispanic White, female

2003.....	99.8	*	*	*	*	*	*	*	0.8	1.7	4.2
2002.....	102.4	*	*	*	*	*	*	*	0.7	2.1	4.0
2001.....	105.3	*	*	*	*	*	*	*	0.7	1.9	3.7
2000.....	107.7	*	*	*	*	*	*	*	0.7	2.0	3.5
1999.....	111.1	*	*	*	*	*	*	0.4	0.6	2.2	3.4

Non-Hispanic Black, both sexes

2003.....	57.9	*	*	*	*	*	*	1.1	2.5	5.8	13.1
2002.....	57.8	*	*	*	*	*	*	1.0	2.0	6.2	12.9
2001.....	59.7	*	*	*	*	*	*	1.3	2.9	6.0	13.7
2000.....	59.8	*	*	*	*	*	*	0.9	2.8	6.6	14.3
1999.....	61.0	*	*	*	*	*	*	0.9	2.3	6.5	15.1

Non-Hispanic Black, male

2003.....	56.7	*	*	*	*	*	*	*	3.4	8.2	18.5
2002.....	56.1	*	*	*	*	*	*	*	2.4	9.0	17.7
2001.....	58.4	*	*	*	*	*	*	1.9	4.6	9.0	19.4
2000.....	57.3	*	*	*	*	*	*	*	3.9	8.9	19.6
1999.....	58.8	*	*	*	*	*	*	*	3.4	9.3	21.6

Non-Hispanic Black, female

2003.....	59.1	*	*	*	*	*	*	*	1.6	3.7	8.4
2002.....	59.3	*	*	*	*	*	*	*	1.6	3.7	8.7
2001.....	60.9	*	*	*	*	*	*	*	*	3.4	8.7
2000.....	62.1	*	*	*	*	*	*	*	1.9	4.6	9.7
1999.....	63.0	*	*	*	*	*	*	*	*	4.0	9.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other heart diseases (I26-I51)

All origins 3/, both sexes

2003.....	58.5	10.4	1.1	0.5	0.7	1.7	2.7	3.6	5.4	7.8	12.6
2002.....	58.8	11.7	1.0	0.4	0.7	1.8	2.3	3.8	5.2	8.1	12.4
2001.....	58.6	11.1	1.4	0.5	0.7	1.5	2.5	3.6	5.1	8.0	11.9
2000.....	58.8	12.0	1.1	0.5	0.7	1.8	2.6	3.5	5.2	7.5	12.3
1999.....	59.4	12.9	1.2	0.5	0.7	2.0	2.6	3.5	5.3	8.2	12.5

All origins 3/, male

2003.....	52.3	11.4	1.0	0.5	0.8	2.1	3.2	4.1	6.5	9.5	15.6
2002.....	52.4	12.1	1.0	0.4	0.8	2.4	3.0	4.6	6.6	9.6	15.9
2001.....	51.8	10.8	1.5	0.4	0.8	1.9	3.1	4.3	6.0	9.9	14.9
2000.....	52.2	12.2	1.2	0.6	0.9	2.1	3.1	4.2	6.2	9.1	15.4
1999.....	53.1	13.1	1.3	0.5	0.8	2.4	3.0	4.0	6.4	10.3	15.6

All origins 3/, female

2003.....	64.5	9.3	1.2	0.4	0.5	1.3	2.1	3.1	4.3	6.1	9.5
2002.....	65.0	11.4	1.0	0.5	0.5	1.1	1.7	3.0	3.8	6.6	9.0
2001.....	65.2	11.5	1.3	0.5	0.7	1.1	1.9	2.9	4.1	6.0	8.9
2000.....	65.1	11.8	0.9	0.4	0.6	1.4	2.1	2.9	4.1	5.9	9.2
1999.....	65.5	12.7	1.0	0.4	0.7	1.5	2.2	2.9	4.2	6.1	9.4

Hispanic, both sexes

2003.....	14.1	8.3	1.0	*	0.5	1.2	1.6	2.3	3.4	4.3	6.3
2002.....	13.3	9.7	0.7	*	*	1.1	1.9	2.3	2.9	3.9	6.6
2001.....	13.3	9.2	1.0	*	*	1.2	1.3	2.1	3.2	4.1	5.9
2000.....	13.6	10.2	1.0	*	0.6	1.5	1.9	2.3	3.3	4.6	6.7
1999.....	13.8	12.2	1.2	*	*	1.5	1.7	1.9	3.7	5.4	7.4

Hispanic, male

2003.....	13.6	9.1	*	*	*	1.4	2.2	2.7	4.1	5.6	8.6
2002.....	12.9	9.1	*	*	*	1.5	2.6	3.1	3.7	4.6	8.6
2001.....	12.6	9.6	*	*	*	1.3	1.8	2.6	3.8	5.3	8.0
2000.....	13.2	9.9	1.3	*	*	2.0	2.2	3.0	4.1	5.4	8.3
1999.....	13.7	10.3	1.4	*	*	1.5	2.0	2.8	4.9	6.7	10.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Other heart diseases (I26-I51)											
Hispanic, female											
2003.....	14.5	7.6	*	*	*	*	*	1.9	2.5	2.8	3.9
2002.....	13.7	10.3	*	*	*	*	*	1.2	1.9	3.1	4.4
2001.....	13.9	8.7	*	*	*	*	*	1.4	2.5	2.8	3.6
2000.....	14.0	10.6	*	*	*	*	1.4	1.5	2.3	3.7	5.0
1999.....	13.9	14.1	*	*	*	1.4	*	*	2.4	4.0	4.5
Non-Hispanic 4/, both sexes											
2003.....	65.4	10.9	1.1	0.5	0.7	1.8	2.9	3.9	5.8	8.4	13.3
2002.....	65.7	12.2	1.1	0.5	0.7	1.9	2.4	4.2	5.6	8.7	13.1
2001.....	65.2	11.6	1.5	0.5	0.8	1.5	2.7	4.0	5.4	8.5	12.5
2000.....	65.1	12.4	1.1	0.5	0.8	1.8	2.8	3.8	5.5	7.9	12.8
1999.....	65.6	13.1	1.1	0.5	0.8	2.1	2.8	3.8	5.6	8.5	13.0
Non-Hispanic 4/, male											
2003.....	58.6	12.0	1.0	0.6	0.9	2.2	3.4	4.5	7.0	10.2	16.5
2002.....	58.7	12.8	1.1	0.4	0.9	2.6	3.1	5.0	7.1	10.4	16.7
2001.....	57.8	11.0	1.6	0.4	0.8	1.9	3.3	4.7	6.4	10.6	15.6
2000.....	58.0	12.7	1.2	0.6	0.9	2.1	3.3	4.4	6.6	9.6	16.1
1999.....	58.6	13.8	1.3	0.5	0.8	2.6	3.2	4.3	6.7	10.8	16.2
Non-Hispanic 4/, female											
2003.....	71.8	9.7	1.2	0.5	0.5	1.3	2.4	3.3	4.6	6.7	10.2
2002.....	72.3	11.6	1.0	0.5	0.5	1.2	1.8	3.3	4.1	7.1	9.5
2001.....	72.3	12.2	1.3	0.5	0.7	1.1	2.1	3.3	4.4	6.5	9.5
2000.....	71.8	12.0	1.0	0.4	0.7	1.5	2.3	3.2	4.4	6.2	9.6
1999.....	72.2	12.3	1.0	0.5	0.7	1.5	2.4	3.2	4.5	6.4	9.9
Non-Hispanic White, both sexes											
2003.....	71.0	8.1	0.8	0.4	0.6	1.5	2.3	3.0	4.6	6.8	11.1
2002.....	71.3	10.0	0.9	0.4	0.6	1.7	1.9	3.2	4.3	7.1	11.3
2001.....	70.6	9.1	1.1	0.4	0.7	1.3	2.1	3.1	4.3	6.8	10.1
2000.....	70.3	10.5	0.9	0.4	0.6	1.4	2.1	2.9	4.4	6.2	10.6
1999.....	70.4	10.5	0.9	0.4	0.6	1.8	2.2	2.9	4.4	6.8	10.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other heart diseases (I26-I51)

Non-Hispanic White, male

2003.....	62.6	8.7	0.7	0.4	0.8	1.9	2.7	3.4	5.7	8.3	14.2
2002.....	62.7	10.7	1.1	0.4	0.8	2.2	2.4	3.8	5.5	8.5	14.8
2001.....	61.4	9.4	1.2	0.3	0.8	1.5	2.6	3.7	5.0	8.5	13.2
2000.....	61.6	10.9	1.1	0.5	0.7	1.6	2.5	3.4	5.4	7.8	13.6
1999.....	61.7	11.6	1.1	0.5	0.6	2.2	2.5	3.3	5.4	8.7	13.6

Non-Hispanic White, female

2003.....	79.1	7.5	1.0	0.4	0.4	1.1	2.0	2.5	3.4	5.3	8.0
2002.....	79.6	9.3	0.8	0.5	0.5	1.1	1.3	2.5	3.1	5.7	7.8
2001.....	79.4	8.8	1.0	0.5	0.6	1.0	1.6	2.6	3.5	5.1	7.0
2000.....	78.7	10.1	0.8	*	0.5	1.2	1.6	2.4	3.4	4.7	7.6
1999.....	78.8	9.4	0.8	0.4	0.6	1.4	2.0	2.4	3.5	4.9	7.5

Non-Hispanic Black, both sexes

2003.....	54.4	19.8	2.0	1.0	1.1	3.1	6.1	9.3	13.2	18.9	29.7
2002.....	54.3	20.9	1.6	*	1.2	3.1	5.3	10.0	13.4	19.7	27.1
2001.....	54.7	21.3	3.0	0.8	1.2	3.0	5.8	9.1	12.8	19.1	30.0
2000.....	54.6	19.6	2.0	0.8	1.4	3.8	6.3	9.1	12.4	18.5	29.1
1999.....	56.2	23.2	2.0	0.8	1.3	3.7	5.9	9.0	13.0	20.1	30.8

Non-Hispanic Black, male

2003.....	53.4	23.6	2.2	*	1.3	4.0	7.7	11.2	14.8	23.5	35.6
2002.....	52.6	21.8	*	*	1.4	4.0	6.4	12.5	17.0	23.6	32.8
2001.....	53.5	18.8	3.3	*	1.3	3.9	7.3	11.1	15.5	24.1	34.6
2000.....	53.3	19.1	2.0	*	1.5	4.5	7.1	10.5	14.9	22.4	35.3
1999.....	56.2	22.9	2.3	*	1.5	5.1	7.1	10.8	15.9	25.1	37.0

Non-Hispanic Black, female

2003.....	55.4	15.8	1.8	*	*	2.1	4.5	7.6	11.8	14.9	24.5
2002.....	55.9	19.9	1.9	*	*	2.3	4.2	7.7	10.3	16.1	22.2
2001.....	55.8	23.8	2.7	*	*	2.1	4.3	7.2	10.4	14.7	26.0
2000.....	55.8	20.1	1.9	*	1.3	3.1	5.5	7.8	10.2	15.0	23.6
1999.....	56.3	23.6	1.8	*	*	2.2	4.7	7.4	10.4	15.7	25.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Hispanic, female

2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.3	0.9	*	*	*	*	0.2	*	0.1	0.2	0.3	
2002.....	0.3	0.7	0.2	*	*	*	*	*	0.2	0.2	0.2	
2001.....	0.3	0.9	0.2	*	*	*	0.1	*	0.1	0.2	0.2	
2000.....	0.3	*	*	*	*	*	*	0.1	0.1	0.2	0.2	
1999.....	0.3	0.8	0.2	*	*	*	*	*	0.1	0.2	0.3	

Non-Hispanic 4/, male

2003.....	0.3	*	*	*	*	*	*	*	*	*	0.3	
2002.....	0.3	*	*	*	*	*	*	*	0.2	*	0.3	
2001.....	0.3	*	*	*	*	*	*	*	*	0.2	*	
2000.....	0.3	*	*	*	*	*	*	*	*	0.3	0.3	
1999.....	0.3	*	*	*	*	*	*	*	*	0.3	0.3	

Non-Hispanic 4/, female

2003.....	0.3	*	*	*	*	*	*	*	*	*	0.2	
2002.....	0.3	*	*	*	*	*	*	*	*	*	*	
2001.....	0.3	*	*	*	*	*	*	*	*	*	0.3	
2000.....	0.3	*	*	*	*	*	*	*	*	*	*	
1999.....	0.3	*	*	*	*	*	*	*	*	0.2	0.3	

Non-Hispanic White, both sexes

2003.....	0.3	*	*	*	*	*	*	*	*	*	0.2	
2002.....	0.3	*	*	*	*	*	*	*	*	*	0.2	
2001.....	0.3	*	*	*	*	*	*	*	*	0.2	0.2	
2000.....	0.3	*	*	*	*	*	*	*	*	0.1	0.2	
1999.....	0.3	*	*	*	*	*	*	*	*	0.2	0.2	

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Heart failure (I50)											
Hispanic, female											
2003.....	4.6	*	*	*	*	*	*	*	*	*	*
2002.....	4.4	*	*	*	*	*	*	*	*	*	*
2001.....	4.5	*	*	*	*	*	*	*	*	*	*
2000.....	4.3	*	*	*	*	*	*	*	*	*	*
1999.....	4.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	22.2	1.0	*	*	*	*	*	0.2	0.4	0.5	1.1
2002.....	22.0	*	*	*	*	*	*	0.2	0.4	0.5	0.8
2001.....	22.4	*	*	*	*	*	*	0.2	0.3	0.5	0.9
2000.....	22.1	0.8	*	*	*	0.1	*	0.2	0.2	0.4	0.9
1999.....	21.8	*	*	*	*	*	*	0.2	0.2	0.4	0.9
Non-Hispanic 4/, male											
2003.....	17.7	1.4	*	*	*	*	*	*	0.5	0.5	1.3
2002.....	17.3	*	*	*	*	*	*	0.3	0.4	0.6	1.1
2001.....	17.4	*	*	*	*	*	*	0.3	0.3	0.7	1.0
2000.....	17.2	*	*	*	*	*	*	*	0.2	0.4	1.1
1999.....	16.9	*	*	*	*	*	*	*	*	0.5	1.1
Non-Hispanic 4/, female											
2003.....	26.5	*	*	*	*	*	*	*	0.3	0.4	0.8
2002.....	26.5	*	*	*	*	*	*	*	0.3	0.4	0.6
2001.....	27.1	*	*	*	*	*	*	*	*	0.4	0.8
2000.....	26.7	*	*	*	*	*	*	*	*	0.3	0.6
1999.....	26.5	*	*	*	*	*	*	*	*	0.3	0.7
Non-Hispanic White, both sexes											
2003.....	25.0	*	*	*	*	*	*	*	0.3	0.4	0.7
2002.....	24.7	*	*	*	*	*	*	*	0.3	0.4	0.6
2001.....	25.1	*	*	*	*	*	*	0.2	0.2	0.4	0.6
2000.....	24.7	*	*	*	*	*	*	*	0.2	0.3	0.7
1999.....	24.3	*	*	*	*	*	*	*	*	0.3	0.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)											
All origins 3/, both sexes											
2003.....	38.0	8.6	0.9	0.4	0.6	1.5	2.4	3.3	4.9	7.0	11.0
2002.....	38.5	10.5	0.8	0.3	0.6	1.6	2.1	3.5	4.6	7.3	11.2
2001.....	37.9	9.7	1.1	0.4	0.7	1.3	2.3	3.3	4.6	7.0	10.5
2000.....	38.3	10.6	0.9	0.4	0.6	1.5	2.4	3.2	4.8	6.8	10.9
1999.....	39.0	11.9	0.9	0.4	0.7	1.8	2.4	3.2	4.9	7.4	11.1
All origins 3/, male											
2003.....	35.8	9.2	0.9	0.4	0.7	1.8	2.9	3.7	5.8	8.6	13.8
2002.....	36.3	10.9	0.8	0.3	0.7	2.2	2.7	4.2	5.8	8.8	14.2
2001.....	35.6	9.1	1.2	0.3	0.7	1.7	2.8	3.8	5.4	8.8	13.4
2000.....	36.1	10.5	1.0	0.5	0.8	1.9	2.8	3.8	5.7	8.3	13.6
1999.....	37.1	12.1	1.0	0.4	0.8	2.2	2.7	3.7	5.9	9.3	13.9
All origins 3/, female											
2003.....	40.1	7.9	1.0	0.3	0.4	1.1	1.8	2.8	3.9	5.5	8.4
2002.....	40.6	10.0	0.8	0.4	0.4	1.0	1.5	2.7	3.4	5.9	8.2
2001.....	40.2	10.4	1.0	0.4	0.6	1.0	1.7	2.7	3.7	5.3	7.6
2000.....	40.4	10.7	0.7	0.3	0.5	1.2	2.0	2.6	3.8	5.3	8.2
1999.....	40.9	11.6	0.8	0.4	0.5	1.4	2.0	2.7	3.9	5.4	8.3
Hispanic, both sexes											
2003.....	9.7	6.8	0.9	*	*	1.0	1.4	2.1	3.1	4.0	5.7
2002.....	9.3	8.6	0.6	*	*	1.0	1.8	2.2	2.6	3.6	6.0
2001.....	9.3	8.0	0.8	*	*	1.2	1.2	1.8	2.7	3.6	5.1
2000.....	9.6	8.7	0.7	*	0.6	1.4	1.8	2.1	3.0	4.1	5.9
1999.....	9.8	11.5	0.9	*	*	1.3	1.4	1.8	3.3	4.7	6.5
Hispanic, male											
2003.....	9.8	7.0	*	*	*	*	2.0	2.4	3.6	5.3	7.6
2002.....	9.6	8.0	*	*	*	1.4	2.5	2.9	3.3	4.4	7.9
2001.....	9.3	7.9	*	*	*	1.3	1.6	2.2	3.3	4.4	7.3
2000.....	9.9	8.1	*	*	*	2.0	2.2	2.8	3.8	4.9	7.3
1999.....	10.3	9.8	*	*	*	1.3	1.7	2.5	4.3	5.8	8.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)											
Hispanic, female											
2003.....	9.6	6.6	*	*	*	*	*	1.6	2.4	2.6	3.6
2002.....	9.0	9.3	*	*	*	*	*	1.2	1.8	2.7	4.0
2001.....	9.2	8.0	*	*	*	*	*	1.3	1.9	2.7	2.9
2000.....	9.3	9.3	*	*	*	*	1.4	*	2.1	3.2	4.5
1999.....	9.2	13.3	*	*	*	*	*	*	2.1	3.5	4.0
Non-Hispanic 4/, both sexes											
2003.....	42.4	9.0	0.9	0.4	0.6	1.6	2.6	3.5	5.2	7.5	11.7
2002.....	42.9	10.9	0.8	0.4	0.6	1.7	2.2	3.8	4.9	7.9	11.8
2001.....	42.1	10.1	1.2	0.3	0.7	1.3	2.4	3.6	4.9	7.5	11.1
2000.....	42.3	11.0	0.9	0.4	0.6	1.6	2.5	3.4	5.0	7.2	11.4
1999.....	43.0	11.9	0.9	0.4	0.7	1.9	2.6	3.4	5.2	7.7	11.6
Non-Hispanic 4/, male											
2003.....	40.0	9.9	0.8	0.5	0.8	1.9	3.1	4.1	6.3	9.2	14.5
2002.....	40.6	11.6	0.8	0.3	0.8	2.3	2.8	4.6	6.3	9.5	14.9
2001.....	39.6	9.3	1.4	0.3	0.7	1.7	3.1	4.2	5.8	9.4	14.1
2000.....	40.0	10.9	1.0	0.5	0.7	1.9	3.0	3.9	6.1	8.8	14.3
1999.....	40.9	12.7	1.0	0.4	0.8	2.4	2.9	3.9	6.2	9.7	14.4
Non-Hispanic 4/, female											
2003.....	44.6	8.1	1.0	0.3	0.4	1.2	2.1	3.0	4.2	5.9	8.9
2002.....	45.1	10.1	0.8	0.4	0.4	1.1	1.6	3.0	3.7	6.4	8.7
2001.....	44.5	11.0	1.0	0.4	0.6	1.0	1.8	3.0	4.0	5.7	8.1
2000.....	44.5	11.0	0.8	0.2	0.5	1.3	2.1	2.9	4.0	5.6	8.6
1999.....	44.9	11.1	0.9	0.4	0.6	1.5	2.2	3.0	4.2	5.7	8.7
Non-Hispanic White, both sexes											
2003.....	45.2	7.1	0.7	0.3	0.6	1.3	2.0	2.7	4.2	6.2	9.9
2002.....	45.8	9.2	0.7	0.3	0.6	1.5	1.7	2.8	3.8	6.5	10.2
2001.....	44.7	8.1	0.9	0.3	0.6	1.1	1.9	2.8	3.9	6.1	9.1
2000.....	44.9	9.6	0.8	0.3	0.5	1.2	1.9	2.6	4.1	5.7	9.5
1999.....	45.4	9.8	0.8	0.4	0.6	1.7	2.1	2.6	4.1	6.1	9.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic White, male

2003.....	42.0	7.5	0.6	0.4	0.7	1.7	2.3	3.0	5.3	7.6	12.6
2002.....	42.7	9.8	0.8	0.3	0.7	2.1	2.2	3.4	4.8	7.8	13.2
2001.....	41.3	8.1	1.0	*	0.7	1.3	2.4	3.4	4.5	7.7	12.0
2000.....	41.7	9.6	0.8	0.4	0.6	1.4	2.3	3.1	5.0	7.1	12.0
1999.....	42.4	10.8	0.9	0.4	0.6	2.0	2.3	3.1	5.0	7.9	12.2

Non-Hispanic White, female

2003.....	48.3	6.7	0.8	*	0.4	1.0	1.7	2.3	3.1	4.8	7.2
2002.....	48.9	8.5	0.6	0.4	0.4	1.0	1.1	2.3	2.7	5.1	7.1
2001.....	47.9	8.1	0.7	*	0.5	0.8	1.4	2.2	3.2	4.4	6.1
2000.....	47.9	9.6	0.7	*	0.3	1.0	1.5	2.1	3.1	4.3	6.9
1999.....	48.2	8.7	0.6	0.4	0.6	1.4	1.9	2.2	3.3	4.3	6.7

Non-Hispanic Black, both sexes

2003.....	38.8	15.2	1.6	0.8	0.9	2.6	5.4	8.4	11.6	16.6	25.0
2002.....	39.1	17.8	1.2	*	1.0	2.9	4.9	9.1	11.8	17.7	24.0
2001.....	39.3	18.2	2.4	*	1.1	2.7	5.1	8.3	11.7	17.0	25.8
2000.....	39.5	16.1	1.5	0.7	1.1	3.4	5.7	8.3	11.3	16.5	25.7
1999.....	40.9	20.9	1.5	0.6	1.1	3.4	5.4	8.2	11.9	18.0	26.9

Non-Hispanic Black, male

2003.....	39.9	18.3	1.9	*	*	3.4	6.9	10.2	12.7	20.7	30.3
2002.....	39.6	19.7	*	*	1.2	3.6	5.9	11.5	14.9	21.3	28.8
2001.....	40.3	16.0	2.7	*	1.2	3.5	6.6	9.7	14.1	21.1	30.8
2000.....	40.5	15.0	1.7	*	1.2	4.0	6.4	9.5	13.7	20.0	31.2
1999.....	42.8	20.8	*	*	1.5	4.8	6.5	9.7	14.5	22.5	32.6

Non-Hispanic Black, female

2003.....	37.8	11.9	*	*	*	1.9	3.8	6.8	10.7	12.9	20.4
2002.....	38.7	15.8	*	*	*	2.1	4.0	7.0	9.1	14.5	19.8
2001.....	38.5	20.5	2.1	*	*	1.8	3.7	6.9	9.6	13.3	21.4
2000.....	38.6	17.3	*	*	*	2.7	5.1	7.1	9.0	13.4	20.9
1999.....	39.2	21.1	*	*	*	2.0	4.4	6.8	9.6	14.1	22.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Hispanic, female

2003.....	3.2	*	*	*	*	*	*	*	*	*	*
2002.....	3.0	*	*	*	*	*	*	*	*	*	*
2001.....	2.9	*	*	*	*	*	*	*	*	*	*
2000.....	2.5	*	*	*	*	*	*	*	*	*	*
1999.....	2.5	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	8.3	*	*	*	*	*	*	0.1	0.4	0.7	1.1
2002.....	7.7	*	*	*	*	*	*	0.2	0.3	0.6	1.0
2001.....	7.4	*	*	*	*	*	*	0.2	0.4	0.5	1.1
2000.....	7.0	*	*	*	*	*	*	*	0.3	0.5	1.1
1999.....	6.6	*	*	*	*	*	*	0.1	0.3	0.5	1.0

Non-Hispanic 4/, male

2003.....	6.3	*	*	*	*	*	*	*	0.5	0.8	1.3
2002.....	5.9	*	*	*	*	*	*	*	0.3	0.7	1.3
2001.....	5.7	*	*	*	*	*	*	*	0.4	0.6	1.3
2000.....	5.4	*	*	*	*	*	*	*	0.3	0.5	1.4
1999.....	5.1	*	*	*	*	*	*	*	0.4	0.7	1.2

Non-Hispanic 4/, female

2003.....	10.2	*	*	*	*	*	*	*	*	0.6	1.0
2002.....	9.4	*	*	*	*	*	*	*	0.2	0.5	0.7
2001.....	9.0	*	*	*	*	*	*	*	0.4	0.4	0.8
2000.....	8.5	*	*	*	*	*	*	*	0.2	0.4	0.8
1999.....	8.0	*	*	*	*	*	*	*	*	0.4	0.8

Non-Hispanic White, both sexes

2003.....	8.0	*	*	*	*	*	*	*	0.2	0.4	0.7
2002.....	7.4	*	*	*	*	*	*	*	*	0.2	0.6
2001.....	7.0	*	*	*	*	*	*	*	0.2	0.3	0.5
2000.....	6.5	*	*	*	*	*	*	*	*	0.2	0.6
1999.....	6.1	*	*	*	*	*	*	*	0.1	0.2	0.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Non-Hispanic White, male

2003.....	5.8	*	*	*	*	*	*	*	*	0.5	0.8
2002.....	5.5	*	*	*	*	*	*	*	*	*	0.8
2001.....	5.1	*	*	*	*	*	*	*	*	0.3	0.7
2000.....	4.9	*	*	*	*	*	*	*	*	0.3	0.8
1999.....	4.6	*	*	*	*	*	*	*	*	0.3	0.6

Non-Hispanic White, female

2003.....	10.0	*	*	*	*	*	*	*	*	*	0.6
2002.....	9.2	*	*	*	*	*	*	*	*	*	0.3
2001.....	8.8	*	*	*	*	*	*	*	*	*	0.3
2000.....	8.1	*	*	*	*	*	*	*	*	*	0.4
1999.....	7.6	*	*	*	*	*	*	*	*	*	0.3

Non-Hispanic Black, both sexes

2003.....	12.2	*	*	*	*	*	*	*	1.4	2.3	4.0
2002.....	11.5	*	*	*	*	*	*	*	1.0	2.8	3.9
2001.....	11.3	*	*	*	*	*	*	*	1.4	1.9	4.5
2000.....	11.0	*	*	*	*	*	*	*	1.1	2.1	4.2
1999.....	10.7	*	*	*	*	*	*	*	1.0	2.3	4.1

Non-Hispanic Black, male

2003.....	10.8	*	*	*	*	*	*	*	2.2	2.4	4.6
2002.....	9.9	*	*	*	*	*	*	*	*	3.7	4.9
2001.....	10.0	*	*	*	*	*	*	*	*	2.3	5.5
2000.....	9.5	*	*	*	*	*	*	*	*	2.2	5.4
1999.....	9.4	*	*	*	*	*	*	*	*	2.8	4.8

Non-Hispanic Black, female

2003.....	13.4	*	*	*	*	*	*	*	*	2.2	3.4
2002.....	13.0	*	*	*	*	*	*	*	*	2.1	3.1
2001.....	12.4	*	*	*	*	*	*	*	1.4	1.5	3.5
2000.....	12.4	*	*	*	*	*	*	*	*	2.1	3.2
1999.....	11.8	*	*	*	*	*	*	*	*	1.9	3.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Cerebrovascular diseases (I60-I69)

All origins 3/, both sexes

2003.....	54.2	2.5	0.3	0.1	0.2	0.4	0.6	1.0	1.9	3.6	7.4
2002.....	56.4	2.9	0.3	0.2	0.3	0.3	0.6	1.0	1.8	3.6	7.1
2001.....	57.4	2.7	0.4	0.2	0.2	0.3	0.7	1.1	1.9	3.8	7.2
2000.....	59.6	3.3	0.3	0.1	0.2	0.3	0.7	1.1	1.9	3.9	7.7
1999.....	60.0	2.7	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.0	7.5

All origins 3/, male

2003.....	42.9	2.8	0.3	*	0.2	0.5	0.6	1.0	2.0	3.8	7.6
2002.....	44.2	3.2	0.4	*	0.3	0.3	0.7	1.0	1.7	3.6	6.9
2001.....	45.2	3.1	0.3	0.2	0.2	0.3	0.7	1.1	2.0	3.9	7.4
2000.....	46.9	3.8	*	*	0.2	0.4	0.7	1.1	1.9	3.8	7.8
1999.....	47.1	3.3	0.3	*	*	0.3	0.6	1.0	1.9	3.9	7.8

All origins 3/, female

2003.....	65.1	2.2	0.3	*	*	0.4	0.7	0.9	1.8	3.3	7.2
2002.....	68.2	2.5	0.3	*	0.3	*	0.5	0.9	1.9	3.7	7.2
2001.....	69.2	2.3	0.4	*	0.2	0.3	0.6	1.0	1.9	3.6	7.1
2000.....	71.8	2.7	0.4	*	0.3	0.3	0.7	1.0	1.9	4.0	7.5
1999.....	72.3	2.1	0.3	*	0.2	0.3	0.6	1.0	1.8	4.1	7.1

Hispanic, both sexes

2003.....	16.7	*	*	*	*	*	*	0.9	1.5	2.5	6.8
2002.....	16.6	2.4	*	*	*	*	*	1.1	1.5	3.0	6.1
2001.....	17.4	*	*	*	*	*	0.8	1.0	2.0	3.4	7.0
2000.....	17.5	*	*	*	*	*	0.8	1.1	2.3	3.4	7.5
1999.....	17.4	*	*	*	*	*	0.7	1.4	1.8	3.3	6.9

Hispanic, male

2003.....	14.9	*	*	*	*	*	*	*	1.7	2.9	8.0
2002.....	15.0	*	*	*	*	*	*	1.4	1.7	3.4	7.1
2001.....	15.7	*	*	*	*	*	*	1.0	2.3	3.7	7.3
2000.....	15.8	*	*	*	*	*	*	1.2	2.6	3.6	7.8
1999.....	16.1	*	*	*	*	*	*	1.4	2.3	3.4	8.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cerebrovascular diseases (I60-I69)

Hispanic, female

2003.....	18.6	*	*	*	*	*	*	*	1.3	2.1	5.5
2002.....	18.4	*	*	*	*	*	*	*	1.3	2.5	5.1
2001.....	19.1	*	*	*	*	*	*	*	1.6	3.1	6.6
2000.....	19.4	*	*	*	*	*	*	*	1.9	3.1	7.1
1999.....	18.8	*	*	*	*	*	*	1.4	*	3.1	5.1

Non-Hispanic 4/, both sexes

2003.....	60.1	2.6	0.3	0.2	0.2	0.5	0.7	1.0	2.0	3.7	7.4
2002.....	62.4	3.0	0.3	0.2	0.2	0.3	0.6	0.9	1.9	3.7	7.2
2001.....	63.2	2.8	0.3	0.2	0.2	0.3	0.6	1.1	1.9	3.8	7.3
2000.....	65.4	3.6	0.3	0.1	0.3	0.4	0.7	1.1	1.8	3.9	7.6
1999.....	65.7	2.9	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.1	7.5

Non-Hispanic 4/, male

2003.....	47.6	2.8	*	*	*	0.5	0.7	1.1	2.1	4.0	7.6
2002.....	48.9	3.5	0.4	*	0.3	0.3	0.7	0.9	1.8	3.6	6.9
2001.....	49.7	3.3	0.3	*	*	0.3	0.6	1.1	1.9	3.9	7.3
2000.....	51.5	4.2	*	*	*	0.4	0.7	1.1	1.8	3.8	7.7
1999.....	51.5	3.7	0.3	*	*	0.3	0.6	0.9	1.8	4.0	7.7

Non-Hispanic 4/, female

2003.....	72.0	2.4	0.3	*	*	0.4	0.7	0.9	1.9	3.4	7.3
2002.....	75.3	2.5	*	*	*	*	0.5	0.9	2.0	3.9	7.5
2001.....	76.1	2.4	0.4	*	*	0.3	0.6	1.0	2.0	3.7	7.2
2000.....	78.7	2.8	0.4	*	0.3	0.3	0.7	1.0	1.9	4.1	7.6
1999.....	79.2	2.0	*	*	0.2	0.3	0.6	1.0	1.8	4.2	7.3

Non-Hispanic White, both sexes

2003.....	64.2	2.0	0.2	*	0.2	0.4	0.5	0.8	1.6	2.6	5.3
2002.....	67.0	2.2	0.3	0.2	0.2	0.2	0.6	0.7	1.5	2.8	5.4
2001.....	67.6	1.9	0.3	0.2	0.2	0.3	0.5	0.9	1.6	3.0	5.4
2000.....	70.0	2.6	0.3	*	0.2	0.4	0.6	1.0	1.4	2.9	5.6
1999.....	70.4	2.4	0.3	*	0.2	0.3	0.6	0.8	1.5	3.2	5.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Cerebrovascular diseases (I60-I69)

Non-Hispanic White, male

2003.....	49.7	1.9	*	*	*	0.4	0.7	0.8	1.7	2.7	5.5
2002.....	51.3	2.7	*	*	*	*	0.7	0.8	1.3	2.6	5.2
2001.....	52.0	2.2	*	*	*	0.3	0.6	0.8	1.7	2.9	5.6
2000.....	53.9	3.3	*	*	*	0.5	0.6	1.1	1.4	2.9	5.7
1999.....	53.9	3.1	*	*	*	0.3	0.5	0.8	1.5	3.1	5.7

Non-Hispanic White, female

2003.....	78.2	2.1	*	*	*	0.4	0.4	0.7	1.5	2.5	5.2
2002.....	82.1	*	*	*	*	*	0.5	0.7	1.7	3.0	5.5
2001.....	82.6	*	*	*	*	0.3	0.4	0.9	1.6	3.0	5.1
2000.....	85.5	1.9	*	*	*	*	0.6	0.9	1.4	3.0	5.5
1999.....	86.1	*	*	*	*	*	0.7	0.8	1.4	3.2	5.3

Non-Hispanic Black, both sexes

2003.....	51.1	4.9	*	*	*	0.7	1.3	2.1	4.3	9.1	19.2
2002.....	51.7	5.6	*	*	*	*	0.9	2.0	4.4	8.5	18.0
2001.....	52.9	6.9	*	*	*	*	1.4	2.3	3.6	8.3	18.9
2000.....	54.3	8.0	*	*	*	*	1.1	1.6	4.5	9.9	20.2
1999.....	53.9	4.7	*	*	*	*	0.8	1.9	3.8	9.4	20.3

Non-Hispanic Black, male

2003.....	44.9	*	*	*	*	*	*	2.6	4.3	10.1	19.5
2002.....	45.1	*	*	*	*	*	*	1.7	4.1	7.8	17.4
2001.....	46.2	8.3	*	*	*	*	*	2.8	3.4	8.9	18.2
2000.....	47.7	9.2	*	*	*	*	*	*	4.2	9.5	21.0
1999.....	47.5	*	*	*	*	*	*	2.0	3.8	9.2	21.1

Non-Hispanic Black, female

2003.....	56.7	*	*	*	*	*	1.7	1.6	4.3	8.3	18.9
2002.....	57.7	*	*	*	*	*	*	2.3	4.6	9.2	18.4
2001.....	59.0	*	*	*	*	*	1.8	1.8	3.9	7.7	19.6
2000.....	60.3	*	*	*	*	*	*	1.7	4.7	10.2	19.5
1999.....	59.8	*	*	*	*	*	*	1.8	3.7	9.6	19.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of circulatory system (I71-I78)

Hispanic, female

2003.....	1.9	*	*	*	*	*	*	*	*	*	*
2002.....	2.0	*	*	*	*	*	*	*	*	*	*
2001.....	1.8	*	*	*	*	*	*	*	*	*	*
2000.....	2.1	*	*	*	*	*	*	*	*	*	*
1999.....	1.9	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	9.5	*	*	*	*	0.1	0.2	0.3	0.4	0.8	1.3
2002.....	9.6	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.3
2001.....	9.9	*	*	*	*	*	0.2	0.4	0.5	0.9	1.4
2000.....	10.2	*	*	*	*	*	0.2	0.2	0.5	0.7	1.3
1999.....	10.2	*	*	*	*	*	0.2	0.3	0.5	0.8	1.4

Non-Hispanic 4/, male

2003.....	10.4	*	*	*	*	*	0.3	0.5	0.6	1.1	1.7
2002.....	10.5	*	*	*	*	*	0.2	0.3	0.7	0.8	1.9
2001.....	10.7	*	*	*	*	*	0.3	0.5	0.7	1.3	1.8
2000.....	11.3	*	*	*	*	*	0.3	0.4	0.7	0.9	1.7
1999.....	11.2	*	*	*	*	*	*	0.4	0.7	1.1	1.7

Non-Hispanic 4/, female

2003.....	8.6	*	*	*	*	*	*	*	0.3	0.5	0.8
2002.....	8.8	*	*	*	*	*	*	*	0.4	0.7	0.7
2001.....	9.1	*	*	*	*	*	*	0.3	0.3	0.6	0.9
2000.....	9.2	*	*	*	*	*	*	*	0.4	0.5	1.0
1999.....	9.3	*	*	*	*	*	*	*	0.2	0.6	1.1

Non-Hispanic White, both sexes

2003.....	10.4	*	*	*	*	*	0.2	0.3	0.4	0.6	1.1
2002.....	10.6	*	*	*	*	0.2	*	0.3	0.4	0.7	1.1
2001.....	10.8	*	*	*	*	*	0.2	0.4	0.5	0.8	1.1
2000.....	11.2	*	*	*	*	*	0.2	0.2	0.5	0.6	1.0
1999.....	11.3	*	*	*	*	*	0.2	0.3	0.5	0.8	1.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Aortic aneurysm and dissection (I71)

Hispanic, female

2003.....	0.9	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.8	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.7	*	*	*	*	*	*	*	*	*	*	*
2000.....	1.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.8	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	5.7	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9
2002.....	5.7	*	*	*	*	*	*	0.2	0.4	0.5	0.9
2001.....	6.0	*	*	*	*	*	0.2	0.3	0.4	0.7	0.9
2000.....	6.2	*	*	*	*	*	0.1	0.2	0.4	0.5	0.9
1999.....	6.3	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9

Non-Hispanic 4/, male

2003.....	7.0	*	*	*	*	*	0.2	0.4	0.5	1.0	1.4
2002.....	7.2	*	*	*	*	*	*	0.3	0.5	0.6	1.5
2001.....	7.3	*	*	*	*	*	*	0.4	0.5	1.0	1.4
2000.....	7.8	*	*	*	*	*	*	0.3	0.5	0.7	1.4
1999.....	7.8	*	*	*	*	*	*	0.3	0.6	0.9	1.3

Non-Hispanic 4/, female

2003.....	4.4	*	*	*	*	*	*	*	*	0.3	0.4
2002.....	4.4	*	*	*	*	*	*	*	*	0.4	0.3
2001.....	4.7	*	*	*	*	*	*	*	*	0.3	0.4
2000.....	4.7	*	*	*	*	*	*	*	*	0.3	0.5
1999.....	4.8	*	*	*	*	*	*	*	*	0.3	0.6

Non-Hispanic White, both sexes

2003.....	6.4	*	*	*	*	*	*	0.2	0.3	0.5	0.7
2002.....	6.5	*	*	*	*	*	*	*	0.3	0.5	0.7
2001.....	6.7	*	*	*	*	*	0.2	0.3	0.3	0.6	0.7
2000.....	7.0	*	*	*	*	*	*	*	0.3	0.4	0.7
1999.....	7.1	*	*	*	*	*	*	0.2	0.3	0.6	0.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other diseases of arteries, arterioles and capillaries (I72-I78)

Hispanic, female

2003.....	1.0	*	*	*	*	*	*	*	*	*	*
2002.....	1.3	*	*	*	*	*	*	*	*	*	*
2001.....	1.1	*	*	*	*	*	*	*	*	*	*
2000.....	1.1	*	*	*	*	*	*	*	*	*	*
1999.....	1.1	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	3.8	*	*	*	*	*	*	*	*	0.2	0.4
2002.....	3.9	*	*	*	*	*	*	*	0.1	0.2	0.4
2001.....	3.9	*	*	*	*	*	*	*	0.2	0.3	0.5
2000.....	4.0	*	*	*	*	*	*	*	0.2	0.2	0.4
1999.....	3.9	*	*	*	*	*	*	*	0.2	0.2	0.5

Non-Hispanic 4/, male

2003.....	3.3	*	*	*	*	*	*	*	*	*	0.4
2002.....	3.4	*	*	*	*	*	*	*	*	*	0.4
2001.....	3.4	*	*	*	*	*	*	*	*	0.3	0.4
2000.....	3.5	*	*	*	*	*	*	*	*	*	0.3
1999.....	3.4	*	*	*	*	*	*	*	*	*	0.4

Non-Hispanic 4/, female

2003.....	4.2	*	*	*	*	*	*	*	*	0.2	0.5
2002.....	4.3	*	*	*	*	*	*	*	*	0.3	0.4
2001.....	4.4	*	*	*	*	*	*	*	*	0.3	0.6
2000.....	4.4	*	*	*	*	*	*	*	*	0.3	0.5
1999.....	4.5	*	*	*	*	*	*	*	*	0.2	0.6

Non-Hispanic White, both sexes

2003.....	4.0	*	*	*	*	*	*	*	*	0.1	0.4
2002.....	4.1	*	*	*	*	*	*	*	*	0.2	0.3
2001.....	4.2	*	*	*	*	*	*	*	*	0.2	0.4
2000.....	4.3	*	*	*	*	*	*	*	0.2	0.2	0.3
1999.....	4.2	*	*	*	*	*	*	*	0.1	0.2	0.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other disorders of circulatory system (I80-I99)

Hispanic, female

2003.....	0.6	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.7	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	1.8	0.8	*	*	*	*	0.2	0.3	0.4	0.6	1.0
2002.....	1.8	0.6	*	*	*	*	0.1	0.2	0.5	0.6	0.8
2001.....	1.8	*	*	*	*	*	0.2	0.3	0.4	0.5	0.9
2000.....	1.8	0.7	*	*	*	*	0.2	0.4	0.4	0.6	0.9
1999.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.5	0.8

Non-Hispanic 4/, male

2003.....	1.5	*	*	*	*	*	*	0.3	0.4	0.6	1.0
2002.....	1.5	*	*	*	*	*	*	*	0.6	0.7	0.8
2001.....	1.5	*	*	*	*	*	*	*	0.4	0.5	0.9
2000.....	1.5	*	*	*	*	*	*	*	0.4	0.6	1.1
1999.....	1.4	*	*	*	*	*	*	*	0.3	0.5	1.0

Non-Hispanic 4/, female

2003.....	2.1	*	*	*	*	*	*	0.3	0.5	0.7	1.0
2002.....	2.1	*	*	*	*	*	*	0.3	0.5	0.6	0.8
2001.....	2.0	*	*	*	*	*	*	0.3	0.5	0.5	0.8
2000.....	2.0	*	*	*	*	*	*	0.5	0.4	0.5	0.6
1999.....	1.8	*	*	*	*	*	*	*	0.3	0.4	0.7

Non-Hispanic White, both sexes

2003.....	1.8	*	*	*	*	*	*	0.3	0.3	0.5	0.8
2002.....	1.8	*	*	*	*	*	*	*	0.4	0.5	0.7
2001.....	1.8	*	*	*	*	*	*	0.2	0.3	0.4	0.7
2000.....	1.8	*	*	*	*	*	*	0.3	0.3	0.5	0.8
1999.....	1.6	*	*	*	*	*	*	0.2	0.3	0.4	0.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Influenza and pneumonia (J10-J18)

All origins 3/, both sexes

2003.....	22.4	8.0	1.0	0.4	0.3	0.4	0.7	0.8	1.1	1.8	2.7
2002.....	22.8	6.5	0.7	0.2	0.3	0.4	0.5	0.7	1.0	1.7	2.6
2001.....	21.8	7.4	0.7	0.2	0.2	0.3	0.6	0.7	1.0	1.8	2.6
2000.....	23.2	7.6	0.7	0.2	0.2	0.3	0.7	0.7	1.1	1.8	3.0
1999.....	22.8	8.4	0.8	0.2	0.2	0.4	0.6	0.7	1.0	2.0	2.7

All origins 3/, male

2003.....	20.1	9.4	1.1	0.4	0.3	0.5	0.7	0.9	1.3	1.9	3.1
2002.....	20.4	7.8	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8	3.1
2001.....	19.6	8.1	0.7	0.2	0.2	0.3	0.7	0.7	1.2	2.0	3.0
2000.....	20.8	8.4	0.7	0.3	0.2	0.3	0.7	0.8	1.2	1.9	3.8
1999.....	20.3	8.8	0.8	0.3	0.3	0.3	0.6	0.8	1.2	2.2	3.1

All origins 3/, female

2003.....	24.6	6.6	0.9	0.4	0.4	0.3	0.7	0.6	0.8	1.6	2.3
2002.....	25.1	5.1	0.6	0.2	0.3	0.3	0.4	0.7	0.8	1.6	2.2
2001.....	23.9	6.7	0.7	0.2	0.2	0.3	0.5	0.6	0.9	1.6	2.2
2000.....	25.6	6.8	0.7	0.2	*	0.3	0.6	0.6	1.0	1.7	2.2
1999.....	25.3	8.0	0.9	*	0.2	0.4	0.5	0.6	0.8	1.8	2.3

Hispanic, both sexes

2003.....	7.4	7.6	1.1	*	*	*	0.7	0.6	0.8	0.9	1.7
2002.....	7.3	5.8	*	*	*	*	*	*	0.7	1.2	1.5
2001.....	7.4	7.0	1.1	*	*	*	*	*	*	1.1	1.7
2000.....	7.4	7.8	*	*	*	*	*	*	0.9	1.2	2.1
1999.....	6.6	8.5	*	*	*	*	*	*	*	1.7	1.3

Hispanic, male

2003.....	6.9	9.1	1.4	*	*	*	*	*	*	*	2.0
2002.....	7.0	7.7	*	*	*	*	*	*	*	1.4	2.1
2001.....	6.9	7.4	*	*	*	*	*	*	*	1.3	2.2
2000.....	7.2	7.6	*	*	*	*	*	*	*	1.4	3.0
1999.....	6.3	10.3	*	*	*	*	*	*	*	2.4	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Influenza and pneumonia (J10-J18)

Hispanic, female

2003.....	7.9	6.1	*	*	*	*	*	*	*	*	*
2002.....	7.6	*	*	*	*	*	*	*	*	*	*
2001.....	7.9	6.5	*	*	*	*	*	*	*	*	*
2000.....	7.7	8.0	*	*	*	*	*	*	*	*	*
1999.....	7.0	6.5	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	24.7	8.1	1.0	0.4	0.4	0.4	0.7	0.8	1.1	1.9	2.8
2002.....	25.1	6.7	0.7	0.2	0.2	0.4	0.5	0.8	1.1	1.7	2.8
2001.....	23.8	7.5	0.6	0.2	0.2	0.3	0.6	0.7	1.2	1.9	2.7
2000.....	25.4	7.5	0.7	0.2	0.2	0.4	0.7	0.8	1.1	1.8	3.0
1999.....	25.0	8.4	0.9	0.2	0.2	0.4	0.6	0.8	1.1	2.0	2.8

Non-Hispanic 4/, male

2003.....	22.3	9.4	1.1	0.4	0.3	0.5	0.7	1.0	1.4	2.2	3.2
2002.....	22.5	7.8	0.8	*	*	0.4	0.6	0.7	1.2	1.9	3.2
2001.....	21.5	8.2	0.6	0.2	0.2	0.3	0.8	0.7	1.4	2.1	3.1
2000.....	22.7	8.5	0.7	0.2	*	0.4	0.7	0.9	1.2	1.9	3.8
1999.....	22.2	8.4	0.9	0.3	0.3	0.3	0.6	0.9	1.3	2.1	3.2

Non-Hispanic 4/, female

2003.....	27.1	6.8	1.0	0.4	0.4	0.3	0.7	0.7	0.9	1.7	2.4
2002.....	27.5	5.4	0.7	*	0.3	0.4	0.4	0.8	0.9	1.6	2.4
2001.....	26.1	6.7	0.6	0.3	0.2	0.3	0.5	0.7	0.9	1.7	2.3
2000.....	27.9	6.5	0.6	*	*	0.3	0.6	0.6	1.0	1.8	2.2
1999.....	27.6	8.4	0.9	*	*	0.4	0.5	0.7	0.9	2.0	2.5

Non-Hispanic White, both sexes

2003.....	27.4	6.2	0.9	0.3	0.4	0.4	0.7	0.7	1.0	1.7	2.4
2002.....	27.9	4.9	0.5	0.2	0.2	0.3	0.5	0.7	0.9	1.5	2.4
2001.....	26.2	4.9	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.6	2.2
2000.....	27.9	5.0	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.7	2.5
1999.....	27.5	5.3	0.6	0.2	0.2	0.3	0.4	0.6	0.9	1.8	2.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Influenza and pneumonia (J10-J18)

Non-Hispanic White, male

2003.....	24.1	7.4	0.9	*	0.3	0.4	0.7	0.9	1.2	2.0	2.7
2002.....	24.6	6.1	0.5	*	*	0.4	0.6	0.7	1.1	1.5	2.8
2001.....	23.1	5.8	0.6	*	*	*	0.7	0.6	1.2	1.7	2.5
2000.....	24.5	5.8	0.5	*	*	0.4	0.7	0.7	1.0	1.7	3.2
1999.....	23.9	5.5	0.5	*	*	0.3	0.5	0.7	1.1	1.8	2.3

Non-Hispanic White, female

2003.....	30.6	5.0	0.8	*	0.4	*	0.7	0.5	0.8	1.5	2.2
2002.....	31.1	3.6	0.5	*	0.3	*	0.4	0.8	0.8	1.4	2.1
2001.....	29.2	3.9	0.5	*	*	0.3	0.4	0.7	0.8	1.5	2.0
2000.....	31.3	4.2	0.6	*	*	*	0.5	0.6	1.0	1.7	1.9
1999.....	31.0	5.1	0.8	*	*	0.4	0.4	0.5	0.7	1.7	2.1

Non-Hispanic Black, both sexes

2003.....	15.9	14.4	1.1	*	*	*	0.8	1.7	2.2	3.4	5.6
2002.....	16.1	13.3	1.7	*	*	*	0.7	1.1	2.0	3.6	5.3
2001.....	16.0	16.9	1.0	*	*	*	1.0	1.2	2.1	3.9	6.0
2000.....	16.9	17.5	1.2	*	*	*	1.1	1.4	2.1	2.9	6.3
1999.....	16.7	20.0	1.7	*	*	*	1.3	1.8	2.3	4.1	7.1

Non-Hispanic Black, male

2003.....	16.5	16.1	*	*	*	*	*	1.7	2.9	3.7	7.2
2002.....	15.9	14.2	1.8	*	*	*	*	*	2.4	4.1	6.2
2001.....	16.3	16.6	*	*	*	*	*	*	2.8	4.7	7.1
2000.....	17.3	18.8	1.7	*	*	*	*	2.1	2.6	3.2	8.6
1999.....	16.9	20.1	2.1	*	*	*	*	2.0	2.6	4.3	9.2

Non-Hispanic Black, female

2003.....	15.4	12.6	*	*	*	*	*	1.7	1.5	3.2	4.2
2002.....	16.2	12.3	*	*	*	*	*	*	1.6	3.2	4.5
2001.....	15.6	17.1	*	*	*	*	*	*	1.5	3.2	4.9
2000.....	16.5	16.2	*	*	*	*	*	*	1.7	2.7	4.3
1999.....	16.5	20.0	*	*	*	*	*	1.6	1.9	3.8	5.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Pneumonia (J12-J18)											
Hispanic, female											
2003.....	7.7	5.4	*	*	*	*	*	*	*	*	*
2002.....	7.5	*	*	*	*	*	*	*	*	*	*
2001.....	7.8	6.5	*	*	*	*	*	*	*	*	*
2000.....	7.6	7.4	*	*	*	*	*	*	*	*	*
1999.....	6.8	5.4	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	24.0	7.4	0.6	0.2	0.3	0.3	0.6	0.8	1.1	1.9	2.8
2002.....	24.8	6.5	0.7	0.2	0.2	0.3	0.5	0.7	1.1	1.7	2.7
2001.....	23.8	7.2	0.6	0.2	0.2	0.3	0.6	0.7	1.1	1.9	2.7
2000.....	24.7	7.3	0.6	0.2	0.1	0.3	0.7	0.7	1.1	1.8	2.9
1999.....	24.3	8.1	0.8	0.2	0.2	0.3	0.6	0.8	1.0	2.0	2.8
Non-Hispanic 4/, male											
2003.....	21.7	8.7	0.7	0.3	0.2	0.4	0.7	0.9	1.4	2.1	3.2
2002.....	22.3	7.6	0.8	*	*	0.4	0.6	0.7	1.2	1.9	3.1
2001.....	21.4	7.8	0.6	*	0.2	0.3	0.7	0.7	1.3	2.1	3.0
2000.....	22.2	8.2	0.7	*	*	0.4	0.7	0.8	1.2	1.9	3.7
1999.....	21.7	8.0	0.8	*	0.3	0.3	0.6	0.9	1.2	2.1	3.2
Non-Hispanic 4/, female											
2003.....	26.3	6.1	0.6	*	0.3	*	0.6	0.7	0.8	1.6	2.3
2002.....	27.2	5.3	0.6	*	*	0.3	0.4	0.8	0.9	1.6	2.3
2001.....	26.0	6.6	0.6	*	*	0.3	0.5	0.7	0.9	1.7	2.3
2000.....	27.1	6.4	0.5	*	*	0.3	0.6	0.6	1.0	1.8	2.2
1999.....	26.8	8.2	0.9	*	*	0.3	0.5	0.6	0.9	1.9	2.4
Non-Hispanic White, both sexes											
2003.....	26.6	5.7	0.6	0.2	0.3	0.3	0.6	0.7	0.9	1.7	2.4
2002.....	27.6	4.7	0.5	0.2	0.2	0.3	0.5	0.7	0.9	1.5	2.4
2001.....	26.1	4.6	0.5	0.2	0.2	0.3	0.5	0.6	1.0	1.6	2.2
2000.....	27.1	4.8	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.7	2.4
1999.....	26.8	5.2	0.6	0.2	0.2	0.3	0.4	0.6	0.9	1.7	2.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Pneumonia (J12-J18)											
Non-Hispanic White, male											
2003.....	23.5	6.8	0.6	*	*	0.4	0.7	0.8	1.1	1.9	2.6
2002.....	24.3	6.0	0.5	*	*	0.4	0.6	0.6	1.1	1.5	2.8
2001.....	23.0	5.3	0.6	*	*	*	0.7	0.5	1.1	1.7	2.5
2000.....	23.8	5.4	0.4	*	*	0.3	0.7	0.6	1.0	1.7	3.1
1999.....	23.4	5.4	*	*	*	0.3	0.5	0.7	1.0	1.8	2.3
Non-Hispanic White, female											
2003.....	29.6	4.5	0.5	*	0.3	*	0.6	0.5	0.7	1.4	2.1
2002.....	30.7	3.4	0.5	*	*	*	0.4	0.8	0.8	1.4	2.1
2001.....	29.1	3.8	0.4	*	*	0.3	0.4	0.7	0.8	1.5	1.9
2000.....	30.3	4.2	0.5	*	*	*	0.5	0.6	0.9	1.6	1.8
1999.....	30.0	4.9	0.7	*	*	*	*	0.5	0.7	1.7	2.0
Non-Hispanic Black, both sexes											
2003.....	15.6	13.3	0.9	*	*	*	0.7	1.6	2.2	3.4	5.6
2002.....	16.0	13.1	1.5	*	*	*	0.7	1.1	2.0	3.6	5.3
2001.....	15.9	16.9	1.0	*	*	*	1.0	1.2	2.1	3.9	5.9
2000.....	16.6	17.2	1.2	*	*	*	1.1	1.4	2.1	2.9	6.2
1999.....	16.5	19.3	1.7	*	*	*	1.3	1.8	2.3	4.0	7.0
Non-Hispanic Black, male											
2003.....	16.2	15.2	*	*	*	*	*	1.6	2.9	3.7	7.2
2002.....	15.8	13.9	1.6	*	*	*	*	*	2.4	4.1	6.2
2001.....	16.3	16.6	*	*	*	*	*	*	2.7	4.7	7.1
2000.....	17.1	18.8	1.7	*	*	*	*	2.1	2.6	3.2	8.3
1999.....	16.8	19.1	2.0	*	*	*	*	2.0	2.6	4.3	9.1
Non-Hispanic Black, female											
2003.....	15.0	11.3	*	*	*	*	*	1.6	1.4	3.2	4.2
2002.....	16.1	12.3	*	*	*	*	*	*	1.6	3.2	4.5
2001.....	15.6	17.1	*	*	*	*	*	*	1.5	3.2	4.8
2000.....	16.3	15.5	*	*	*	*	*	*	1.7	2.7	4.3
1999.....	16.3	19.6	*	*	*	*	*	1.5	1.9	3.7	5.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

Hispanic, female

2003.....	7.7	*	*	*	*	*	*	*	*	*	*
2002.....	7.6	*	*	*	*	*	*	*	*	*	2.0
2001.....	7.5	*	*	*	*	*	*	*	*	*	*
2000.....	7.2	*	*	*	*	*	*	*	*	*	1.9
1999.....	8.0	*	*	*	*	*	*	*	*	1.5	2.1

Non-Hispanic 4/, both sexes

2003.....	49.0	0.8	0.3	0.2	0.4	0.4	0.6	0.7	0.9	1.5	3.0
2002.....	48.6	1.1	0.4	0.2	0.5	0.5	0.5	0.7	0.9	1.7	3.0
2001.....	48.4	1.1	0.3	0.2	0.3	0.4	0.6	0.6	1.0	1.7	2.9
2000.....	48.3	1.0	0.3	0.2	0.5	0.5	0.6	0.7	0.9	1.4	2.9
1999.....	49.4	1.0	0.4	0.2	0.5	0.6	0.6	0.8	1.0	1.3	2.7

Non-Hispanic 4/, male

2003.....	48.1	*	0.5	0.3	0.5	0.5	0.7	0.8	1.0	1.4	2.7
2002.....	48.4	1.2	0.6	0.2	0.5	0.7	0.5	0.8	0.9	1.6	3.1
2001.....	48.0	*	0.4	0.3	0.4	0.4	0.7	0.6	1.0	1.7	2.7
2000.....	48.7	*	0.4	0.3	0.5	0.6	0.7	0.8	1.0	1.4	2.8
1999.....	50.8	*	0.4	0.3	0.6	0.6	0.8	0.8	0.9	1.3	2.5

Non-Hispanic 4/, female

2003.....	49.9	*	*	*	0.3	0.3	0.5	0.6	0.8	1.6	3.4
2002.....	48.9	*	*	*	0.5	0.3	0.5	0.6	0.9	1.8	2.9
2001.....	48.7	*	*	*	0.3	0.3	0.5	0.6	1.0	1.7	3.0
2000.....	48.0	*	*	*	0.4	0.3	0.5	0.6	0.8	1.5	3.0
1999.....	48.0	*	*	*	0.3	0.5	0.5	0.8	1.1	1.4	2.9

Non-Hispanic White, both sexes

2003.....	57.0	*	*	*	0.2	0.2	0.4	0.5	0.6	1.3	2.7
2002.....	56.4	0.9	0.2	*	0.2	0.3	0.4	0.5	0.7	1.4	2.6
2001.....	55.9	*	*	*	0.2	0.3	0.3	0.4	0.7	1.2	2.6
2000.....	55.7	*	*	*	0.2	0.3	0.4	0.5	0.6	1.1	2.4
1999.....	56.6	*	0.2	0.2	0.2	0.5	0.4	0.5	0.6	1.0	2.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic White, male

2003.....	54.9	*	*	*	*	*	0.5	0.4	0.6	1.2	2.3
2002.....	55.1	*	*	*	*	0.4	0.4	0.5	0.8	1.1	2.7
2001.....	54.6	*	*	*	*	*	0.4	0.4	0.6	1.3	2.4
2000.....	55.1	*	*	*	*	0.3	0.5	0.5	0.6	0.9	2.4
1999.....	57.3	*	*	*	0.3	0.5	0.5	0.4	0.6	1.0	2.0

Non-Hispanic White, female

2003.....	59.0	*	*	*	*	*	0.3	0.5	0.7	1.4	3.0
2002.....	57.7	*	*	*	*	*	0.5	0.5	0.7	1.6	2.5
2001.....	57.2	*	*	*	*	*	0.3	0.3	0.8	1.2	2.8
2000.....	56.2	*	*	*	*	*	0.4	0.5	0.7	1.3	2.5
1999.....	55.9	*	*	*	*	0.4	0.4	0.5	0.7	1.1	2.5

Non-Hispanic Black, both sexes

2003.....	20.9	*	1.0	0.7	1.2	1.3	1.4	1.8	2.2	3.0	5.8
2002.....	21.4	*	1.2	0.7	1.6	1.4	1.2	1.7	2.2	3.9	6.5
2001.....	21.1	*	0.9	0.7	0.9	1.0	1.7	1.9	3.1	4.7	5.1
2000.....	21.5	*	0.9	0.7	1.4	1.5	1.4	1.9	2.5	3.6	6.3
1999.....	22.6	*	*	*	1.5	1.2	1.6	2.6	2.8	3.3	6.2

Non-Hispanic Black, male

2003.....	24.2	*	*	*	1.6	1.6	1.6	2.5	3.3	3.0	5.2
2002.....	24.9	*	1.6	*	1.6	2.3	1.6	2.3	2.2	4.6	6.9
2001.....	24.5	*	*	*	*	1.4	1.9	2.2	3.4	4.4	5.1
2000.....	25.1	*	*	*	1.8	2.1	1.6	2.4	3.3	3.8	6.1
1999.....	27.0	*	*	*	2.1	1.5	2.3	2.8	2.6	3.3	6.2

Non-Hispanic Black, female

2003.....	17.9	*	*	*	*	*	*	*	*	3.0	6.4
2002.....	18.2	*	*	*	1.5	*	*	*	2.2	3.3	6.2
2001.....	18.0	*	*	*	*	*	1.5	1.6	2.7	4.9	5.2
2000.....	18.2	*	*	*	*	*	*	1.5	1.7	3.4	6.4
1999.....	18.6	*	*	*	*	*	*	2.4	2.9	3.3	6.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Asthma (J45-J46)											
Hispanic, female											
2003.....	0.9	*	*	*	*	*	*	*	*	*	*
2002.....	0.9	*	*	*	*	*	*	*	*	*	*
2001.....	1.0	*	*	*	*	*	*	*	*	*	*
2000.....	1.0	*	*	*	*	*	*	*	*	*	*
1999.....	1.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	1.5	*	0.2	0.2	0.4	0.4	0.5	0.6	0.7	0.8	1.2
2002.....	1.6	*	0.3	0.2	0.4	0.5	0.5	0.6	0.7	1.0	1.2
2001.....	1.6	*	0.2	0.2	0.3	0.3	0.5	0.5	0.8	1.0	1.1
2000.....	1.7	*	0.2	0.2	0.4	0.4	0.5	0.6	0.8	0.9	1.3
1999.....	1.8	*	0.2	0.2	0.4	0.5	0.5	0.7	0.7	0.9	1.1
Non-Hispanic 4/, male											
2003.....	1.1	*	0.4	0.3	0.4	0.4	0.5	0.7	0.8	0.6	0.9
2002.....	1.2	*	0.4	*	0.5	0.7	0.5	0.7	0.7	0.9	1.0
2001.....	1.1	*	*	0.3	0.3	0.4	0.5	0.6	0.7	0.9	1.0
2000.....	1.2	*	*	0.2	0.5	0.5	0.6	0.7	0.8	0.8	1.1
1999.....	1.3	*	*	0.3	0.6	0.6	0.7	0.7	0.7	0.9	0.8
Non-Hispanic 4/, female											
2003.....	1.9	*	*	*	0.3	0.3	0.4	0.5	0.6	0.9	1.5
2002.....	2.0	*	*	*	0.4	0.2	0.4	0.5	0.7	1.1	1.4
2001.....	2.1	*	*	*	0.3	0.3	0.4	0.5	0.8	1.1	1.3
2000.....	2.1	*	*	*	0.4	0.3	0.5	0.5	0.7	1.0	1.5
1999.....	2.2	*	*	*	0.3	0.5	0.4	0.6	0.8	1.0	1.4
Non-Hispanic White, both sexes											
2003.....	1.3	*	*	*	0.2	0.2	0.3	0.4	0.5	0.5	0.9
2002.....	1.4	*	*	*	0.2	0.3	0.3	0.4	0.5	0.7	0.7
2001.....	1.4	*	*	*	0.2	0.2	0.3	0.3	0.5	0.6	0.8
2000.....	1.4	*	*	*	0.2	0.2	0.4	0.4	0.5	0.5	0.9
1999.....	1.5	*	*	*	0.2	0.4	0.4	0.4	0.5	0.6	0.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Asthma (J45-J46)											
Non-Hispanic White, male											
2003.....	0.9	*	*	*	*	*	0.4	0.4	0.4	0.4	0.6
2002.....	0.9	*	*	*	*	0.3	0.3	0.4	0.5	0.5	0.6
2001.....	0.9	*	*	*	*	*	*	0.3	0.3	0.5	0.6
2000.....	1.0	*	*	*	*	*	0.4	0.4	0.5	0.4	0.7
1999.....	1.0	*	*	*	0.3	0.4	0.4	0.3	0.4	0.5	0.4
Non-Hispanic White, female											
2003.....	1.7	*	*	*	*	*	*	0.4	0.5	0.7	1.1
2002.....	1.8	*	*	*	*	*	0.4	0.4	0.6	0.9	0.9
2001.....	1.9	*	*	*	*	*	*	*	0.6	0.6	1.0
2000.....	1.9	*	*	*	*	*	0.3	*	0.5	0.7	1.0
1999.....	2.1	*	*	*	*	0.4	*	0.4	0.5	0.7	1.1
Non-Hispanic Black, both sexes											
2003.....	2.8	*	0.8	0.7	1.1	1.1	1.3	1.5	2.0	2.1	3.5
2002.....	3.0	*	0.9	*	1.6	1.3	1.2	1.6	1.6	3.0	4.1
2001.....	3.1	*	*	0.7	0.9	0.9	1.4	1.8	2.7	3.5	3.1
2000.....	3.3	*	*	0.7	1.4	1.3	1.2	1.8	2.1	2.7	3.7
1999.....	3.3	*	*	*	1.5	1.1	1.5	2.4	2.4	2.7	3.3
Non-Hispanic Black, male											
2003.....	2.5	*	*	*	1.5	1.3	1.4	2.2	2.9	2.0	2.7
2002.....	2.8	*	*	*	1.6	2.2	1.6	2.3	1.7	3.5	3.8
2001.....	2.7	*	*	*	*	*	1.6	2.0	3.0	3.1	2.9
2000.....	2.8	*	*	*	1.7	1.8	*	2.3	2.7	2.9	3.2
1999.....	2.9	*	*	*	2.0	1.3	2.1	2.8	2.6	2.7	2.9
Non-Hispanic Black, female											
2003.....	3.0	*	*	*	*	*	*	*	*	2.2	4.2
2002.....	3.1	*	*	*	1.5	*	*	*	1.5	2.4	4.3
2001.....	3.4	*	*	*	*	*	*	1.5	2.4	3.9	3.3
2000.....	3.7	*	*	*	*	*	*	*	1.5	2.7	4.1
1999.....	3.6	*	*	*	*	*	*	2.1	2.3	2.7	3.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other chronic lower respiratory diseases (J44,J47)

Hispanic, female

2003.....	6.0	*	*	*	*	*	*	*	*	*	*
2002.....	5.9	*	*	*	*	*	*	*	*	*	*
2001.....	5.8	*	*	*	*	*	*	*	*	*	*
2000.....	5.6	*	*	*	*	*	*	*	*	*	*
1999.....	5.9	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	41.4	*	*	*	*	*	*	*	0.1	0.6	1.5
2002.....	40.6	*	*	*	*	*	*	*	0.2	0.5	1.5
2001.....	40.0	*	*	*	*	*	*	*	0.2	0.6	1.4
2000.....	39.5	*	*	*	*	*	*	*	*	0.5	1.3
1999.....	40.0	*	*	*	*	*	*	*	0.1	0.3	1.2

Non-Hispanic 4/, male

2003.....	40.7	*	*	*	*	*	*	*	*	0.6	1.4
2002.....	40.4	*	*	*	*	*	*	*	*	0.4	1.6
2001.....	39.8	*	*	*	*	*	*	*	*	0.6	1.4
2000.....	40.0	*	*	*	*	*	*	*	*	0.4	1.3
1999.....	41.4	*	*	*	*	*	*	*	*	0.3	1.3

Non-Hispanic 4/, female

2003.....	42.0	*	*	*	*	*	*	*	*	0.6	1.5
2002.....	40.8	*	*	*	*	*	*	*	*	0.6	1.3
2001.....	40.1	*	*	*	*	*	*	*	*	0.5	1.5
2000.....	39.1	*	*	*	*	*	*	*	*	0.5	1.2
1999.....	38.7	*	*	*	*	*	*	*	*	0.3	1.1

Non-Hispanic White, both sexes

2003.....	48.6	*	*	*	*	*	*	*	*	0.6	1.5
2002.....	47.5	*	*	*	*	*	*	*	*	0.5	1.5
2001.....	46.6	*	*	*	*	*	*	*	0.2	0.5	1.5
2000.....	46.0	*	*	*	*	*	*	*	*	0.4	1.3
1999.....	46.3	*	*	*	*	*	*	*	*	0.3	1.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pneumonitis due to solids and liquids (J69)

Hispanic, female

2003.....	1.4	*	*	*	*	*	*	*	*	*	*	*
2002.....	1.4	*	*	*	*	*	*	*	*	*	*	*
2001.....	1.5	*	*	*	*	*	*	*	*	*	*	*
2000.....	1.4	*	*	*	*	*	*	*	*	*	*	*
1999.....	1.3	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	6.7	*	*	*	*	*	0.2	0.2	0.2	0.3	0.5
2002.....	6.8	*	*	*	*	*	0.1	0.2	0.2	0.4	0.6
2001.....	6.7	*	*	*	*	*	0.1	0.2	0.3	0.3	0.5
2000.....	6.5	*	*	*	*	*	0.1	*	0.2	0.3	0.5
1999.....	6.0	*	*	*	*	*	0.1	0.2	0.3	0.3	0.5

Non-Hispanic 4/, male

2003.....	6.9	*	*	*	*	*	0.2	*	0.3	0.4	0.5
2002.....	7.0	*	*	*	*	*	*	*	0.3	0.4	0.7
2001.....	6.8	*	*	*	*	*	*	*	0.3	0.4	0.7
2000.....	6.8	*	*	*	*	*	*	*	0.3	0.4	0.7
1999.....	6.3	*	*	*	*	*	*	*	0.3	0.3	0.6

Non-Hispanic 4/, female

2003.....	6.5	*	*	*	*	*	*	*	*	0.3	0.5
2002.....	6.6	*	*	*	*	*	*	*	*	0.3	0.4
2001.....	6.6	*	*	*	*	*	*	*	0.3	*	0.4
2000.....	6.3	*	*	*	*	*	*	*	*	*	0.4
1999.....	5.8	*	*	*	*	*	*	*	*	0.2	0.4

Non-Hispanic White, both sexes

2003.....	7.5	*	*	*	*	*	*	*	0.3	0.3	0.4
2002.....	7.7	*	*	*	*	*	*	0.2	0.2	0.3	0.5
2001.....	7.5	*	*	*	*	*	*	0.2	0.2	0.3	0.5
2000.....	7.3	*	*	*	*	*	*	*	0.2	0.2	0.6
1999.....	6.7	*	*	*	*	*	*	0.2	0.2	0.3	0.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Hispanic, female

2003.....	3.6	6.6	*	*	*	*	*	*	*	*	*
2002.....	3.4	5.4	*	*	*	*	*	*	*	*	*
2001.....	3.5	5.0	*	*	*	*	*	*	*	*	*
2000.....	3.6	5.3	*	*	*	*	*	*	*	*	*
1999.....	3.5	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	9.6	7.0	0.6	*	0.1	0.3	0.4	0.5	0.6	0.9	1.4
2002.....	9.5	7.3	0.5	0.2	0.2	0.3	0.4	0.4	0.6	0.9	1.5
2001.....	10.0	7.8	0.6	0.2	0.2	0.3	0.4	0.6	0.6	1.1	1.6
2000.....	9.8	7.5	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.0	1.8
1999.....	9.7	6.9	0.6	0.2	0.2	0.3	0.3	0.5	0.8	1.1	1.6

Non-Hispanic 4/, male

2003.....	9.5	8.3	0.6	*	*	0.3	0.4	0.6	0.7	0.9	1.4
2002.....	9.4	8.7	0.7	*	*	0.3	0.4	0.3	0.5	1.0	1.7
2001.....	9.8	10.0	0.6	*	0.2	0.3	0.5	0.5	0.6	1.0	1.8
2000.....	9.8	8.4	0.6	0.2	*	0.3	0.5	0.5	0.8	1.1	1.9
1999.....	9.6	7.9	0.8	0.2	*	0.4	0.4	0.5	0.9	1.2	1.7

Non-Hispanic 4/, female

2003.....	9.7	5.7	0.5	*	*	0.3	0.3	0.3	0.6	0.9	1.4
2002.....	9.6	5.9	0.4	*	0.3	0.2	0.4	0.4	0.6	0.9	1.4
2001.....	10.2	5.5	0.6	*	0.2	0.3	0.4	0.7	0.6	1.1	1.4
2000.....	9.9	6.6	0.5	*	*	0.3	0.3	0.6	0.7	0.9	1.6
1999.....	9.8	5.9	0.5	*	0.2	*	0.3	0.6	0.8	1.0	1.4

Non-Hispanic White, both sexes

2003.....	10.7	4.9	0.4	*	*	0.3	0.3	0.4	0.6	0.8	1.1
2002.....	10.5	4.7	0.4	0.2	0.2	0.2	0.3	0.3	0.5	0.8	1.4
2001.....	11.0	4.8	0.5	0.2	0.2	0.3	0.4	0.5	0.5	0.8	1.4
2000.....	10.8	5.4	0.5	0.2	0.2	0.3	0.3	0.5	0.7	0.8	1.6
1999.....	10.6	4.9	0.6	0.2	0.2	0.3	0.3	0.4	0.7	0.9	1.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic White, male

2003.....	10.6	5.5	0.4	*	*	*	0.3	0.5	0.6	0.8	1.2
2002.....	10.3	5.6	0.5	*	*	*	0.4	*	0.4	0.8	1.4
2001.....	10.8	6.2	0.4	*	*	0.3	0.4	0.4	0.4	0.8	1.7
2000.....	10.7	6.6	0.5	*	*	*	0.4	0.4	0.7	0.9	1.7
1999.....	10.5	5.6	0.6	*	*	0.4	0.4	0.5	0.8	1.0	1.5

Non-Hispanic White, female

2003.....	10.8	4.4	*	*	*	*	*	*	0.5	0.7	1.0
2002.....	10.7	3.7	*	*	*	*	*	0.4	0.6	0.8	1.3
2001.....	11.3	3.4	0.5	*	*	*	0.3	0.6	0.5	0.8	1.1
2000.....	11.0	4.1	0.5	*	*	*	*	0.5	0.6	0.7	1.5
1999.....	10.7	4.0	0.5	*	*	*	*	0.4	0.7	0.8	1.1

Non-Hispanic Black, both sexes

2003.....	6.0	15.3	1.1	*	*	*	0.9	0.8	1.2	1.9	3.1
2002.....	6.2	17.3	1.2	*	*	*	0.9	*	1.1	1.8	2.9
2001.....	6.9	19.0	1.1	*	*	*	0.8	1.2	1.4	2.6	3.3
2000.....	6.7	17.7	0.8	*	*	*	0.8	0.9	1.4	2.4	3.1
1999.....	6.7	15.8	1.1	*	*	*	*	1.1	1.7	2.4	3.4

Non-Hispanic Black, male

2003.....	6.2	18.6	*	*	*	*	*	*	*	2.0	2.6
2002.....	6.3	20.3	*	*	*	*	*	*	*	2.0	3.6
2001.....	6.6	24.3	*	*	*	*	*	*	1.6	2.0	3.2
2000.....	6.8	17.1	*	*	*	*	*	*	*	3.0	3.5
1999.....	6.6	18.0	*	*	*	*	*	*	1.8	2.5	3.2

Non-Hispanic Black, female

2003.....	5.9	11.9	*	*	*	*	*	*	*	1.8	3.6
2002.....	6.2	14.2	*	*	*	*	*	*	*	1.7	2.3
2001.....	7.2	13.4	*	*	*	*	*	*	*	3.0	3.4
2000.....	6.6	18.3	*	*	*	*	*	*	*	1.9	2.7
1999.....	6.9	13.6	*	*	*	*	*	1.8	1.6	2.3	3.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Peptic ulcer (K25-K28)

Hispanic, female

2003.....	0.4	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	1.5	*	*	*	*	*	*	*	*	0.2	0.3
2002.....	1.6	*	*	*	*	*	*	*	*	0.2	0.3
2001.....	1.7	*	*	*	*	*	*	*	*	0.2	0.4
2000.....	1.8	*	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.8	*	*	*	*	*	*	*	*	0.2	0.4

Non-Hispanic 4/, male

2003.....	1.4	*	*	*	*	*	*	*	*	*	0.4
2002.....	1.5	*	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.6	*	*	*	*	*	*	*	*	0.3	0.6
2000.....	1.8	*	*	*	*	*	*	*	*	0.3	0.4
1999.....	1.7	*	*	*	*	*	*	*	*	0.3	0.5

Non-Hispanic 4/, female

2003.....	1.5	*	*	*	*	*	*	*	*	*	0.3
2002.....	1.6	*	*	*	*	*	*	*	*	*	0.3
2001.....	1.8	*	*	*	*	*	*	*	*	*	0.2
2000.....	1.8	*	*	*	*	*	*	*	*	*	0.2
1999.....	1.9	*	*	*	*	*	*	*	*	*	0.3

Non-Hispanic White, both sexes

2003.....	1.6	*	*	*	*	*	*	*	*	0.1	0.3
2002.....	1.7	*	*	*	*	*	*	*	*	0.2	0.3
2001.....	1.9	*	*	*	*	*	*	*	*	*	0.4
2000.....	1.9	*	*	*	*	*	*	*	*	0.2	0.3
1999.....	2.0	*	*	*	*	*	*	*	*	0.1	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic liver disease and cirrhosis (K70,K73-K74)

All origins 3/, both sexes

2003.....	9.5	*	*	*	*	*	*	0.4	1.4	4.1	9.3
2002.....	9.5	*	*	*	*	*	0.1	0.4	1.4	4.2	9.7
2001.....	9.5	*	*	*	*	*	0.1	0.4	1.5	5.0	9.8
2000.....	9.4	*	*	*	*	*	0.2	0.5	1.6	4.8	10.2
1999.....	9.4	*	*	*	*	*	*	0.4	1.6	4.8	9.9

All origins 3/, male

2003.....	12.5	*	*	*	*	*	*	0.6	1.9	5.4	12.8
2002.....	12.3	*	*	*	*	*	*	0.5	1.8	5.5	12.9
2001.....	12.4	*	*	*	*	*	*	0.6	2.0	6.4	13.2
2000.....	12.5	*	*	*	*	*	*	0.6	2.1	6.3	14.4
1999.....	12.5	*	*	*	*	*	*	0.4	2.1	6.3	14.5

All origins 3/, female

2003.....	6.5	*	*	*	*	*	*	0.3	0.8	2.9	5.9
2002.....	6.7	*	*	*	*	*	*	0.3	1.0	3.0	6.5
2001.....	6.7	*	*	*	*	*	*	0.3	1.0	3.5	6.4
2000.....	6.5	*	*	*	*	*	*	0.3	1.1	3.2	6.1
1999.....	6.4	*	*	*	*	*	*	0.3	1.2	3.4	5.4

Hispanic, both sexes

2003.....	8.5	*	*	*	*	*	*	*	2.4	4.9	11.4
2002.....	8.8	*	*	*	*	*	*	*	1.8	5.1	13.0
2001.....	8.9	*	*	*	*	*	*	*	2.0	6.3	13.9
2000.....	9.0	*	*	*	*	*	*	*	2.4	6.0	14.4
1999.....	8.8	*	*	*	*	*	*	*	2.4	7.0	14.9

Hispanic, male

2003.....	11.6	*	*	*	*	*	*	*	3.5	7.9	17.4
2002.....	12.2	*	*	*	*	*	*	*	2.6	7.6	20.7
2001.....	12.7	*	*	*	*	*	*	*	2.9	9.6	21.7
2000.....	12.7	*	*	*	*	*	*	*	4.1	9.1	22.5
1999.....	12.3	*	*	*	*	*	*	*	3.4	10.8	23.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Chronic liver disease and cirrhosis (K70,K73-K74)

Hispanic, female

2003.....	5.2	*	*	*	*	*	*	*	1.2	1.5	5.0
2002.....	5.2	*	*	*	*	*	*	*	*	2.3	4.8
2001.....	5.0	*	*	*	*	*	*	*	*	2.6	5.6
2000.....	5.1	*	*	*	*	*	*	*	*	2.7	5.9
1999.....	5.0	*	*	*	*	*	*	*	*	2.8	5.7

Non-Hispanic 4/, both sexes

2003.....	9.6	*	*	*	*	*	*	0.4	1.1	4.0	9.0
2002.....	9.5	*	*	*	*	*	*	0.4	1.3	4.1	9.2
2001.....	9.5	*	*	*	*	*	*	0.4	1.4	4.8	9.2
2000.....	9.4	*	*	*	*	*	0.1	0.5	1.4	4.6	9.7
1999.....	9.5	*	*	*	*	*	*	0.4	1.5	4.5	9.3

Non-Hispanic 4/, male

2003.....	12.6	*	*	*	*	*	*	0.5	1.5	4.9	12.1
2002.....	12.2	*	*	*	*	*	*	0.4	1.6	5.1	11.8
2001.....	12.3	*	*	*	*	*	*	0.5	1.8	5.9	12.1
2000.....	12.4	*	*	*	*	*	*	0.6	1.7	5.9	13.3
1999.....	12.5	*	*	*	*	*	*	0.4	1.8	5.6	13.4

Non-Hispanic 4/, female

2003.....	6.7	*	*	*	*	*	*	0.3	0.7	3.1	6.0
2002.....	6.9	*	*	*	*	*	*	0.3	1.1	3.0	6.7
2001.....	6.9	*	*	*	*	*	*	0.3	1.0	3.7	6.5
2000.....	6.7	*	*	*	*	*	*	0.3	1.2	3.3	6.1
1999.....	6.6	*	*	*	*	*	*	0.3	1.2	3.4	5.3

Non-Hispanic White, both sexes

2003.....	10.3	*	*	*	*	*	*	0.3	1.1	4.0	9.2
2002.....	10.2	*	*	*	*	*	*	0.3	1.3	4.1	9.4
2001.....	10.1	*	*	*	*	*	*	0.3	1.4	4.8	9.0
2000.....	10.0	*	*	*	*	*	*	0.3	1.3	4.4	9.4
1999.....	9.9	*	*	*	*	*	*	0.4	1.4	4.4	8.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic White, male

2003.....	13.6	*	*	*	*	*	*	0.4	1.5	5.0	12.5
2002.....	13.2	*	*	*	*	*	*	0.4	1.6	5.1	12.2
2001.....	13.0	*	*	*	*	*	*	0.4	1.8	5.8	11.7
2000.....	13.1	*	*	*	*	*	*	0.5	1.6	5.7	13.1
1999.....	13.1	*	*	*	*	*	*	0.5	1.8	5.5	12.7

Non-Hispanic White, female

2003.....	7.2	*	*	*	*	*	*	*	0.7	3.1	6.0
2002.....	7.5	*	*	*	*	*	*	*	0.9	3.1	6.7
2001.....	7.4	*	*	*	*	*	*	*	1.0	3.8	6.3
2000.....	7.0	*	*	*	*	*	*	*	1.0	3.1	5.8
1999.....	6.9	*	*	*	*	*	*	*	1.1	3.3	5.0

Non-Hispanic Black, both sexes

2003.....	7.1	*	*	*	*	*	*	*	0.9	3.7	7.9
2002.....	7.0	*	*	*	*	*	*	*	1.4	3.0	8.2
2001.....	7.7	*	*	*	*	*	*	*	1.0	4.2	10.9
2000.....	7.7	*	*	*	*	*	*	0.8	1.4	5.0	11.2
1999.....	8.0	*	*	*	*	*	*	*	1.4	4.5	12.1

Non-Hispanic Black, male

2003.....	9.8	*	*	*	*	*	*	*	*	4.5	9.9
2002.....	9.3	*	*	*	*	*	*	*	*	3.8	10.0
2001.....	10.7	*	*	*	*	*	*	*	*	5.8	15.1
2000.....	10.3	*	*	*	*	*	*	*	*	6.8	14.8
1999.....	11.0	*	*	*	*	*	*	*	1.8	5.6	17.5

Non-Hispanic Black, female

2003.....	4.6	*	*	*	*	*	*	*	*	3.0	6.2
2002.....	5.0	*	*	*	*	*	*	*	*	2.4	6.6
2001.....	4.9	*	*	*	*	*	*	*	*	2.7	7.3
2000.....	5.4	*	*	*	*	*	*	*	1.5	3.5	8.0
1999.....	5.4	*	*	*	*	*	*	*	*	3.6	7.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcoholic liver disease (K70)

All origins 3/, both sexes

2003.....	4.3	*	*	*	*	*	*	0.3	0.9	2.9	5.9
2002.....	4.2	*	*	*	*	*	*	0.2	0.9	2.8	5.9
2001.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3	6.1
2000.....	4.3	*	*	*	*	*	*	0.3	1.1	3.3	6.6
1999.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3	6.4

All origins 3/, male

2003.....	6.4	*	*	*	*	*	*	0.4	1.4	3.7	8.0
2002.....	6.2	*	*	*	*	*	*	0.3	1.2	3.6	8.0
2001.....	6.4	*	*	*	*	*	*	0.3	1.4	4.4	8.4
2000.....	6.5	*	*	*	*	*	*	0.5	1.5	4.3	9.4
1999.....	6.5	*	*	*	*	*	*	0.2	1.4	4.3	9.4

All origins 3/, female

2003.....	2.2	*	*	*	*	*	*	*	0.5	2.0	3.8
2002.....	2.3	*	*	*	*	*	*	*	0.7	2.0	3.9
2001.....	2.3	*	*	*	*	*	*	*	0.7	2.3	3.9
2000.....	2.2	*	*	*	*	*	*	*	0.6	2.2	3.9
1999.....	2.1	*	*	*	*	*	*	*	0.8	2.4	3.5

Hispanic, both sexes

2003.....	4.3	*	*	*	*	*	*	*	1.6	3.8	7.7
2002.....	4.3	*	*	*	*	*	*	*	1.1	3.5	7.9
2001.....	4.7	*	*	*	*	*	*	*	1.5	4.1	8.8
2000.....	4.8	*	*	*	*	*	*	*	1.7	4.3	10.0
1999.....	4.6	*	*	*	*	*	*	*	1.5	4.7	9.7

Hispanic, male

2003.....	6.9	*	*	*	*	*	*	*	2.3	6.3	11.8
2002.....	7.1	*	*	*	*	*	*	*	1.8	5.7	13.0
2001.....	7.7	*	*	*	*	*	*	*	2.4	6.9	13.8
2000.....	7.9	*	*	*	*	*	*	*	3.0	6.4	16.0
1999.....	7.4	*	*	*	*	*	*	*	2.3	7.1	15.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Alcoholic liver disease (K70)

Hispanic, female

2003.....	1.5	*	*	*	*	*	*	*	*	*	3.4
2002.....	1.4	*	*	*	*	*	*	*	*	*	2.5
2001.....	1.5	*	*	*	*	*	*	*	*	*	3.6
2000.....	1.4	*	*	*	*	*	*	*	*	2.1	3.8
1999.....	1.5	*	*	*	*	*	*	*	*	2.1	3.6

Non-Hispanic 4/, both sexes

2003.....	4.2	*	*	*	*	*	*	0.3	0.8	2.7	5.6
2002.....	4.2	*	*	*	*	*	*	0.2	0.9	2.6	5.6
2001.....	4.2	*	*	*	*	*	*	0.2	1.0	3.2	5.8
2000.....	4.2	*	*	*	*	*	*	0.3	1.0	3.1	6.2
1999.....	4.2	*	*	*	*	*	*	0.2	1.0	3.1	6.0

Non-Hispanic 4/, male

2003.....	6.2	*	*	*	*	*	*	0.4	1.2	3.3	7.5
2002.....	6.0	*	*	*	*	*	*	0.3	1.1	3.2	7.3
2001.....	6.2	*	*	*	*	*	*	*	1.2	4.0	7.7
2000.....	6.2	*	*	*	*	*	*	0.4	1.2	4.0	8.6
1999.....	6.4	*	*	*	*	*	*	0.2	1.2	3.9	8.6

Non-Hispanic 4/, female

2003.....	2.3	*	*	*	*	*	*	*	0.4	2.1	3.9
2002.....	2.4	*	*	*	*	*	*	*	0.7	2.0	4.1
2001.....	2.4	*	*	*	*	*	*	*	0.7	2.4	3.9
2000.....	2.3	*	*	*	*	*	*	*	0.7	2.3	3.9
1999.....	2.2	*	*	*	*	*	*	*	0.8	2.4	3.5

Non-Hispanic White, both sexes

2003.....	4.5	*	*	*	*	*	*	0.2	0.8	2.7	5.8
2002.....	4.4	*	*	*	*	*	*	*	0.9	2.6	5.8
2001.....	4.4	*	*	*	*	*	*	*	1.0	3.2	5.7
2000.....	4.3	*	*	*	*	*	*	0.2	0.9	3.0	6.0
1999.....	4.3	*	*	*	*	*	*	0.2	0.9	3.0	5.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other chronic liver disease and cirrhosis (K73-K74)

Hispanic, female

2003.....	3.7	*	*	*	*	*	*	*	*	*	1.6
2002.....	3.8	*	*	*	*	*	*	*	*	*	2.3
2001.....	3.4	*	*	*	*	*	*	*	*	1.5	2.1
2000.....	3.7	*	*	*	*	*	*	*	*	*	2.0
1999.....	3.5	*	*	*	*	*	*	*	*	*	2.0

Non-Hispanic 4/, both sexes

2003.....	5.4	*	*	*	*	*	*	0.2	0.3	1.3	3.3
2002.....	5.4	*	*	*	*	*	*	0.2	0.4	1.4	3.6
2001.....	5.3	*	*	*	*	*	*	0.2	0.4	1.5	3.4
2000.....	5.2	*	*	*	*	*	*	0.2	0.5	1.5	3.5
1999.....	5.2	*	*	*	*	*	*	0.2	0.5	1.4	3.3

Non-Hispanic 4/, male

2003.....	6.4	*	*	*	*	*	*	*	0.4	1.6	4.6
2002.....	6.2	*	*	*	*	*	*	*	0.5	1.9	4.5
2001.....	6.2	*	*	*	*	*	*	0.3	0.6	1.9	4.4
2000.....	6.1	*	*	*	*	*	*	*	0.5	1.9	4.7
1999.....	6.1	*	*	*	*	*	*	*	0.6	1.7	4.8

Non-Hispanic 4/, female

2003.....	4.4	*	*	*	*	*	*	*	0.3	1.0	2.1
2002.....	4.6	*	*	*	*	*	*	*	0.3	1.0	2.6
2001.....	4.5	*	*	*	*	*	*	*	0.3	1.2	2.5
2000.....	4.4	*	*	*	*	*	*	*	0.4	1.0	2.2
1999.....	4.4	*	*	*	*	*	*	*	0.4	1.0	1.8

Non-Hispanic White, both sexes

2003.....	5.9	*	*	*	*	*	*	*	0.3	1.4	3.4
2002.....	5.9	*	*	*	*	*	*	*	0.4	1.5	3.6
2001.....	5.8	*	*	*	*	*	*	*	0.5	1.6	3.3
2000.....	5.7	*	*	*	*	*	*	*	0.4	1.4	3.4
1999.....	5.6	*	*	*	*	*	*	*	0.5	1.4	3.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Hispanic, female

2003.....	5.7	*	*	*	*	*	*	*	*	*	*
2002.....	5.1	*	*	*	*	*	*	*	*	*	*
2001.....	5.4	*	*	*	*	*	*	*	*	*	*
2000.....	4.9	*	*	*	*	*	*	*	*	*	*
1999.....	4.7	*	*	*	*	*	*	*	*	*	1.9

Non-Hispanic 4/, both sexes

2003.....	16.0	4.6	*	*	*	0.1	0.3	0.6	1.0	1.3	2.3
2002.....	15.6	4.3	*	*	*	0.1	0.3	0.6	0.8	1.3	2.1
2001.....	15.1	3.5	*	*	*	*	0.4	0.5	0.7	1.4	2.1
2000.....	14.4	4.4	*	*	*	0.1	0.3	0.5	0.8	1.3	2.1
1999.....	13.8	4.8	*	*	*	0.1	0.3	0.5	0.8	1.2	2.0

Non-Hispanic 4/, male

2003.....	15.8	5.4	*	*	*	*	0.4	0.6	1.0	1.5	2.8
2002.....	15.4	5.4	*	*	*	*	0.3	0.5	0.9	1.5	2.3
2001.....	14.8	3.7	*	*	*	*	0.3	0.5	0.7	1.5	2.2
2000.....	14.1	5.4	*	*	*	*	0.3	0.5	0.9	1.5	2.5
1999.....	13.5	4.8	*	*	*	*	0.3	0.5	0.9	1.4	2.3

Non-Hispanic 4/, female

2003.....	16.2	3.6	*	*	*	*	0.3	0.6	0.9	1.2	1.9
2002.....	15.8	3.2	*	*	*	*	0.3	0.7	0.7	1.2	1.9
2001.....	15.4	3.3	*	*	*	*	0.4	0.4	0.8	1.3	1.9
2000.....	14.7	3.4	*	*	*	*	0.4	0.4	0.7	1.2	1.7
1999.....	14.1	4.8	*	*	*	*	0.3	0.6	0.8	1.0	1.7

Non-Hispanic White, both sexes

2003.....	15.8	3.5	*	*	*	*	0.2	0.4	0.6	0.8	1.4
2002.....	15.4	3.3	*	*	*	*	0.2	0.4	0.6	0.9	1.4
2001.....	14.9	2.4	*	*	*	*	0.3	0.3	0.5	0.9	1.3
2000.....	14.2	2.7	*	*	*	*	0.3	0.4	0.5	0.8	1.4
1999.....	13.4	3.5	*	*	*	*	0.2	0.3	0.5	0.8	1.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic White, male

2003.....	15.7	4.4	*	*	*	*	*	0.4	0.6	0.8	1.8
2002.....	15.3	4.1	*	*	*	*	*	0.3	0.7	0.9	1.5
2001.....	14.8	2.5	*	*	*	*	*	*	0.4	1.2	1.5
2000.....	14.1	3.5	*	*	*	*	*	0.4	0.6	0.9	1.7
1999.....	13.3	3.9	*	*	*	*	*	0.3	0.6	0.9	1.4

Non-Hispanic White, female

2003.....	15.9	2.5	*	*	*	*	*	0.4	0.6	0.7	1.0
2002.....	15.6	2.5	*	*	*	*	*	0.4	0.5	0.8	1.3
2001.....	15.0	2.3	*	*	*	*	0.3	*	0.6	0.7	1.2
2000.....	14.2	1.8	*	*	*	*	*	*	0.4	0.7	1.2
1999.....	13.5	3.1	*	*	*	*	*	0.4	0.5	0.6	1.0

Non-Hispanic Black, both sexes

2003.....	21.4	9.5	*	*	*	*	0.8	1.5	3.1	4.6	8.2
2002.....	20.5	8.5	*	*	*	*	0.9	1.9	2.3	4.1	6.8
2001.....	20.2	8.5	*	*	*	*	0.8	1.4	2.2	4.1	6.9
2000.....	19.5	12.0	*	*	*	*	*	1.1	2.7	4.5	6.5
1999.....	19.2	10.2	*	*	*	*	0.8	1.5	2.4	4.2	7.2

Non-Hispanic Black, male

2003.....	20.8	9.9	*	*	*	*	*	*	3.7	5.6	9.1
2002.....	19.7	10.6	*	*	*	*	*	1.7	2.3	4.5	7.8
2001.....	18.6	8.9	*	*	*	*	*	1.8	2.6	3.4	7.2
2000.....	18.3	13.6	*	*	*	*	*	*	3.0	5.3	8.2
1999.....	18.1	8.7	*	*	*	*	*	*	2.4	5.0	8.5

Non-Hispanic Black, female

2003.....	21.9	9.0	*	*	*	*	*	*	2.5	3.7	7.4
2002.....	21.2	6.3	*	*	*	*	*	2.1	2.2	3.7	6.0
2001.....	21.7	8.1	*	*	*	*	*	*	1.9	4.6	6.7
2000.....	20.7	10.2	*	*	*	*	*	*	2.5	3.9	5.0
1999.....	20.2	11.8	*	*	*	*	*	1.7	2.4	3.4	6.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Renal failure (N17-N19)											
Hispanic, female											
2003.....	5.6	*	*	*	*	*	*	*	*	*	*
2002.....	5.0	*	*	*	*	*	*	*	*	*	*
2001.....	5.2	*	*	*	*	*	*	*	*	*	*
2000.....	4.8	*	*	*	*	*	*	*	*	*	*
1999.....	4.5	*	*	*	*	*	*	*	*	*	1.9
Non-Hispanic 4/, both sexes											
2003.....	15.7	4.3	*	*	*	*	0.3	0.5	0.9	1.3	2.3
2002.....	15.3	4.1	*	*	*	*	0.3	0.6	0.7	1.3	2.1
2001.....	14.9	3.3	*	*	*	*	0.3	0.4	0.7	1.3	2.0
2000.....	14.1	4.2	*	*	*	0.1	0.3	0.4	0.7	1.2	2.0
1999.....	13.5	4.7	*	*	*	0.1	0.2	0.5	0.8	1.1	1.9
Non-Hispanic 4/, male											
2003.....	15.5	5.0	*	*	*	*	0.3	0.5	1.0	1.4	2.7
2002.....	15.1	5.1	*	*	*	*	0.3	0.5	0.8	1.4	2.2
2001.....	14.6	3.5	*	*	*	*	0.3	0.5	0.7	1.4	2.2
2000.....	13.8	5.2	*	*	*	*	*	0.4	0.8	1.4	2.4
1999.....	13.2	4.6	*	*	*	*	*	0.4	0.8	1.3	2.2
Non-Hispanic 4/, female											
2003.....	15.9	3.5	*	*	*	*	0.3	0.5	0.8	1.2	1.9
2002.....	15.5	2.9	*	*	*	*	0.3	0.6	0.7	1.1	1.9
2001.....	15.2	3.1	*	*	*	*	0.3	0.4	0.7	1.3	1.8
2000.....	14.4	3.1	*	*	*	*	0.3	0.4	0.7	1.1	1.7
1999.....	13.8	4.8	*	*	*	*	0.3	0.5	0.7	0.9	1.6
Non-Hispanic White, both sexes											
2003.....	15.5	3.2	*	*	*	*	0.2	0.4	0.6	0.7	1.4
2002.....	15.2	3.1	*	*	*	*	0.2	0.3	0.5	0.8	1.3
2001.....	14.6	2.2	*	*	*	*	0.2	0.3	0.4	0.9	1.3
2000.....	13.9	2.5	*	*	*	*	0.2	0.3	0.4	0.8	1.3
1999.....	13.1	3.5	*	*	*	*	*	0.3	0.5	0.7	1.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Renal failure (N17-N19)											
Non-Hispanic White, male											
2003.....	15.4	4.0	*	*	*	*	*	0.4	0.6	0.8	1.7
2002.....	15.0	3.9	*	*	*	*	*	*	0.6	0.9	1.4
2001.....	14.5	2.3	*	*	*	*	*	*	0.4	1.1	1.5
2000.....	13.7	3.3	*	*	*	*	*	0.3	0.5	0.8	1.6
1999.....	13.0	3.8	*	*	*	*	*	*	0.5	0.8	1.4
Non-Hispanic White, female											
2003.....	15.6	2.3	*	*	*	*	*	0.4	0.6	0.7	1.0
2002.....	15.3	2.2	*	*	*	*	*	0.4	0.4	0.7	1.2
2001.....	14.7	2.1	*	*	*	*	*	*	0.5	0.7	1.1
2000.....	14.0	*	*	*	*	*	*	*	0.4	0.7	1.1
1999.....	13.2	3.1	*	*	*	*	*	*	0.5	0.6	1.0
Non-Hispanic Black, both sexes											
2003.....	21.1	9.3	*	*	*	*	0.7	1.4	2.9	4.6	8.1
2002.....	20.2	8.0	*	*	*	*	0.8	1.8	2.1	4.0	6.8
2001.....	20.0	8.5	*	*	*	*	0.8	1.3	2.2	3.9	6.8
2000.....	19.2	11.4	*	*	*	*	*	1.0	2.6	4.2	6.3
1999.....	18.8	10.2	*	*	*	*	*	1.5	2.3	3.9	6.9
Non-Hispanic Black, male											
2003.....	20.5	9.6	*	*	*	*	*	*	3.6	5.6	9.0
2002.....	19.4	10.0	*	*	*	*	*	*	2.1	4.5	7.6
2001.....	18.4	8.9	*	*	*	*	*	1.7	2.6	3.3	7.0
2000.....	17.9	13.6	*	*	*	*	*	*	2.8	5.1	8.0
1999.....	17.7	8.7	*	*	*	*	*	*	2.3	4.7	8.3
Non-Hispanic Black, female											
2003.....	21.7	9.0	*	*	*	*	*	*	2.4	3.7	7.2
2002.....	20.9	*	*	*	*	*	*	2.0	2.1	3.6	6.0
2001.....	21.4	8.1	*	*	*	*	*	*	1.8	4.5	6.6
2000.....	20.3	9.2	*	*	*	*	*	*	2.4	3.5	4.9
1999.....	19.8	11.8	*	*	*	*	*	1.7	2.3	3.3	5.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pregnancy, childbirth and the puerperium (000-099)

Hispanic, female

2003.....	0.5	*	*	*	1.8	*	*	*
2002.....	0.4	*	*	*	*	*	*	*
2001.....	0.5	*	*	*	1.3	*	*	*
2000.....	0.5	*	*	1.4	1.3	*	*	*
1999.....	0.4	*	*	*	*	*	1.6	*

Non-Hispanic 4/, both sexes

2003.....	0.2	*	*	0.4	0.5	0.6	0.5	0.2
2002.....	0.1	*	0.1	0.3	0.5	0.4	0.3	0.1
2001.....	0.1	*	0.2	0.3	0.5	0.5	0.3	0.1
2000.....	0.1	*	*	0.4	0.4	0.5	0.3	0.2
1999.....	0.1	*	0.2	0.3	0.4	0.5	0.3	0.1

Non-Hispanic 4/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2003.....	0.3	*	*	0.8	1.0	1.2	1.0	0.5
2002.....	0.2	*	0.3	0.6	0.9	0.8	0.7	0.2
2001.....	0.3	*	0.3	0.7	0.9	0.9	0.6	0.3
2000.....	0.3	*	*	0.7	0.8	0.9	0.7	0.3
1999.....	0.3	*	0.3	0.6	0.9	1.0	0.7	0.3

Non-Hispanic White, both sexes

2003.....	0.1	*	*	0.3	0.4	0.3	0.3	0.2
2002.....	0.1	*	*	*	0.3	0.3	0.2	*
2001.....	0.1	*	*	0.2	0.3	0.3	0.2	*
2000.....	0.1	*	*	0.2	0.3	0.3	0.2	*
1999.....	0.1	*	*	*	0.2	0.3	0.2	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic White, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2003.....	0.2	*	*	0.5	0.8	0.7	0.5	0.3	
2002.....	0.1	*	*	*	0.5	0.6	0.3	*	
2001.....	0.2	*	*	0.4	0.5	0.7	0.3	*	
2000.....	0.2	*	*	0.4	0.5	0.6	0.5	*	
1999.....	0.2	*	*	*	0.5	0.7	0.4	*	

Non-Hispanic Black, both sexes

2003.....	0.5	*	*	1.0	1.3	1.9	1.5	*	
2002.....	0.4	*	*	1.0	1.5	1.0	1.2	*	
2001.....	0.4	*	*	1.1	1.4	1.1	1.0	*	
2000.....	0.4	*	*	1.1	1.2	1.3	0.9	*	
1999.....	0.5	*	*	1.1	1.4	1.2	1.0	*	

Non-Hispanic Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2003.....	1.0	*	*	2.0	2.4	3.6	2.8	*	
2002.....	0.8	*	*	2.0	2.8	1.9	2.3	*	
2001.....	0.8	*	*	2.2	2.7	2.2	1.9	*	
2000.....	0.8	*	*	2.2	2.3	2.5	1.7	*	
1999.....	0.9	*	*	2.1	2.7	2.4	1.8	*	

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.0	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*

Non-Hispanic 4/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2003.....	0.0	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*

Non-Hispanic White, both sexes

2003.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

Non-Hispanic White, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2003.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Non-Hispanic Black, both sexes

2003.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Non-Hispanic Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2003.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Hispanic, female

2003.....	0.5	*	*	*	1.6	*	*	*
2002.....	0.4	*	*	*	*	*	*	*
2001.....	0.5	*	*	*	1.3	*	*	*
2000.....	0.5	*	*	1.3	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.2	*	*	0.4	0.5	0.5	0.5	0.2
2002.....	0.1	*	0.1	0.3	0.4	0.4	0.3	0.1
2001.....	0.1	*	0.1	0.3	0.4	0.4	0.3	0.1
2000.....	0.1	*	*	0.3	0.4	0.4	0.3	0.1
1999.....	0.1	*	0.2	0.3	0.4	0.4	0.3	0.1

Non-Hispanic 4/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2003.....	0.3	*	*	0.8	1.0	1.1	0.9	0.4
2002.....	0.2	*	0.3	0.5	0.9	0.8	0.6	0.2
2001.....	0.2	*	0.3	0.6	0.9	0.8	0.6	0.2
2000.....	0.2	*	*	0.7	0.7	0.8	0.6	0.3
1999.....	0.2	*	0.3	0.6	0.8	0.8	0.6	0.3

Non-Hispanic White, both sexes

2003.....	0.1	*	*	0.3	0.3	0.3	0.3	0.2
2002.....	0.1	*	*	*	0.3	0.3	0.2	*
2001.....	0.1	*	*	0.2	0.2	0.3	0.1	*
2000.....	0.1	*	*	0.2	0.2	0.3	0.2	*
1999.....	0.1	*	*	*	0.2	0.3	0.2	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other complications of pregnancy, childbirth and the puerperium (010-099)

Non-Hispanic White, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2003.....	0.2	*	*	0.5	0.7	0.6	0.5	0.3	
2002.....	0.1	*	*	*	0.5	0.5	0.3	*	
2001.....	0.1	*	*	0.3	0.5	0.6	0.3	*	
2000.....	0.1	*	*	0.4	0.5	0.6	0.4	*	
1999.....	0.1	*	*	*	0.5	0.6	0.4	*	

Non-Hispanic Black, both sexes

2003.....	0.5	*	*	0.9	1.3	1.7	1.4	*	
2002.....	0.4	*	*	0.9	1.4	0.9	1.1	*	
2001.....	0.4	*	*	1.0	1.3	1.0	0.9	*	
2000.....	0.3	*	*	1.0	1.0	1.1	0.7	*	
1999.....	0.4	*	*	1.0	1.3	1.1	0.8	*	

Non-Hispanic Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2003.....	0.9	*	*	1.8	2.4	3.2	2.6	*	
2002.....	0.7	*	*	1.9	2.7	1.8	2.0	*	
2001.....	0.7	*	*	2.0	2.5	1.9	1.6	*	
2000.....	0.7	*	*	1.9	1.9	2.2	1.4	*	
1999.....	0.8	*	*	2.0	2.4	2.1	1.6	*	

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All origins 3/, both sexes

2003.....	3.6	140.4	3.4	0.9	1.0	1.1	1.1	1.0	1.1	1.1	1.4
2002.....	3.7	139.4	3.4	1.0	1.0	1.2	1.2	1.1	1.3	1.2	1.3
2001.....	3.7	136.7	3.6	0.9	0.9	1.3	1.3	1.2	1.1	1.2	1.4
2000.....	3.8	150.9	3.2	1.0	1.0	1.1	1.1	1.1	1.3	1.2	1.3
1999.....	3.7	144.2	3.6	1.0	1.1	1.1	1.1	1.1	1.2	1.2	1.2

All origins 3/, male

2003.....	3.8	147.6	3.4	0.9	1.0	1.4	1.2	1.2	1.1	1.2	1.5
2002.....	3.9	142.6	3.5	1.0	1.1	1.4	1.3	1.4	1.4	1.2	1.4
2001.....	3.9	143.1	3.6	1.0	1.0	1.5	1.4	1.3	1.2	1.2	1.6
2000.....	4.0	155.4	3.3	1.0	1.0	1.3	1.4	1.2	1.5	1.4	1.5
1999.....	3.9	147.9	4.0	1.0	1.1	1.3	1.3	1.1	1.3	1.3	1.3

All origins 3/, female

2003.....	3.4	132.8	3.5	0.9	1.0	0.8	1.0	0.8	1.1	1.1	1.3
2002.....	3.5	136.1	3.3	1.0	1.0	1.0	1.1	0.9	1.1	1.2	1.2
2001.....	3.4	129.9	3.7	0.8	0.8	1.0	1.1	1.1	1.0	1.1	1.3
2000.....	3.5	146.2	3.1	0.9	1.0	0.9	0.9	0.9	1.1	1.1	1.1
1999.....	3.5	140.3	3.2	1.0	1.1	0.9	1.0	1.1	1.2	1.1	1.2

Hispanic, both sexes

2003.....	4.5	153.0	3.7	0.8	0.9	0.8	0.8	0.9	0.6	*	1.0
2002.....	4.4	150.7	3.5	1.1	0.8	1.2	0.8	0.6	0.8	0.7	*
2001.....	4.4	146.2	3.4	0.8	0.9	1.0	1.2	0.7	*	*	0.9
2000.....	4.6	153.3	3.8	0.9	*	0.8	0.9	*	0.9	0.8	*
1999.....	4.4	141.8	3.5	0.9	0.9	0.9	0.9	0.9	0.9	*	*

Hispanic, male

2003.....	4.5	156.9	3.7	*	*	*	*	1.1	*	*	*
2002.....	4.5	153.1	3.8	*	*	1.3	*	*	1.1	*	*
2001.....	4.7	153.4	3.5	*	*	*	1.1	*	*	*	*
2000.....	4.7	153.3	4.0	*	*	*	1.1	*	*	*	*
1999.....	4.3	138.1	4.3	*	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Hispanic, female

2003.....	4.4	148.9	3.7	*	1.2	*	*	*	*	*	*
2002.....	4.3	148.2	3.3	1.2	*	*	*	*	*	*	*
2001.....	4.1	138.8	3.3	*	*	*	*	*	*	*	*
2000.....	4.4	153.3	3.5	*	*	*	*	*	*	*	*
1999.....	4.5	145.8	2.7	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	3.5	136.0	3.4	0.9	1.0	1.1	1.2	1.1	1.2	1.2	1.4
2002.....	3.6	134.9	3.3	1.0	1.1	1.2	1.3	1.2	1.3	1.3	1.4
2001.....	3.5	132.8	3.7	0.9	0.9	1.3	1.3	1.3	1.2	1.2	1.5
2000.....	3.6	148.8	3.1	1.0	1.0	1.2	1.2	1.2	1.4	1.3	1.4
1999.....	3.6	143.1	3.6	1.0	1.1	1.1	1.2	1.1	1.3	1.3	1.3

Non-Hispanic 4/, male

2003.....	3.7	144.2	3.3	0.9	1.0	1.4	1.3	1.3	1.2	1.3	1.6
2002.....	3.8	138.1	3.5	1.0	1.1	1.4	1.4	1.6	1.5	1.3	1.5
2001.....	3.7	139.4	3.6	1.0	1.0	1.6	1.5	1.3	1.4	1.3	1.6
2000.....	3.9	154.4	3.1	1.0	1.0	1.4	1.4	1.3	1.7	1.4	1.5
1999.....	3.8	148.4	3.9	1.0	1.1	1.4	1.4	1.2	1.4	1.4	1.4

Non-Hispanic 4/, female

2003.....	3.3	127.6	3.4	0.9	0.9	0.8	1.0	0.8	1.2	1.2	1.3
2002.....	3.4	131.5	3.2	1.0	1.0	1.0	1.2	0.9	1.2	1.3	1.3
2001.....	3.3	125.8	3.7	0.9	0.9	0.9	1.1	1.2	1.1	1.2	1.4
2000.....	3.4	142.8	3.0	0.9	1.1	1.0	1.0	1.0	1.1	1.1	1.2
1999.....	3.4	137.4	3.3	1.1	1.1	0.9	1.0	1.1	1.2	1.2	1.2

Non-Hispanic White, both sexes

2003.....	3.3	129.8	3.0	0.8	0.9	1.1	1.1	1.1	1.2	1.3	1.5
2002.....	3.4	131.2	3.0	1.0	1.0	1.2	1.3	1.3	1.4	1.3	1.4
2001.....	3.4	127.4	3.1	0.9	0.9	1.2	1.2	1.3	1.3	1.2	1.5
2000.....	3.4	141.2	2.6	0.9	1.0	1.0	1.2	1.2	1.5	1.3	1.3
1999.....	3.4	136.8	3.0	0.9	1.0	1.1	1.1	1.2	1.3	1.3	1.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Non-Hispanic White, male

2003.....	3.4	136.1	2.9	0.8	1.0	1.4	1.3	1.3	1.2	1.4	1.7
2002.....	3.6	137.2	3.2	0.9	1.0	1.3	1.4	1.6	1.6	1.3	1.6
2001.....	3.6	135.5	3.1	1.0	1.0	1.5	1.3	1.5	1.5	1.3	1.6
2000.....	3.6	147.9	2.6	0.8	0.9	1.1	1.5	1.3	1.8	1.4	1.5
1999.....	3.6	144.4	3.2	0.8	1.0	1.4	1.4	1.3	1.4	1.5	1.4

Non-Hispanic White, female

2003.....	3.2	123.2	3.0	0.8	0.9	0.8	0.9	0.9	1.3	1.2	1.4
2002.....	3.3	125.0	2.9	1.0	1.1	1.0	1.2	1.0	1.2	1.3	1.3
2001.....	3.2	118.8	3.2	0.8	0.7	0.9	1.0	1.1	1.2	1.1	1.4
2000.....	3.2	134.2	2.6	0.9	1.0	0.9	1.0	1.0	1.2	1.1	1.2
1999.....	3.3	128.9	2.9	1.0	1.0	0.8	0.9	1.2	1.2	1.2	1.2

Non-Hispanic Black, both sexes

2003.....	4.6	160.5	4.8	1.2	1.1	1.3	1.8	1.3	1.6	1.3	1.2
2002.....	4.7	156.9	4.7	1.2	1.3	1.7	1.6	1.4	1.5	1.4	1.9
2001.....	4.7	163.5	5.4	1.0	1.2	2.0	1.9	1.4	1.2	1.5	1.7
2000.....	5.0	184.1	5.2	1.3	1.4	1.9	1.3	1.4	1.4	1.6	1.7
1999.....	4.8	174.8	5.8	1.5	1.8	1.4	1.5	1.1	1.2	1.5	1.3

Non-Hispanic Black, male

2003.....	5.2	173.1	5.0	1.3	*	1.7	1.9	*	*	*	*
2002.....	5.2	153.3	4.8	1.4	1.5	2.1	1.8	1.8	1.7	*	2.1
2001.....	5.1	167.5	5.5	*	*	2.7	2.3	*	*	*	2.2
2000.....	5.5	186.0	5.6	1.7	1.3	2.6	*	1.8	1.7	1.7	2.0
1999.....	5.3	179.5	6.6	1.4	1.8	1.7	1.9	*	*	*	1.8

Non-Hispanic Black, female

2003.....	4.0	147.3	4.7	*	*	*	1.6	*	1.6	*	*
2002.....	4.4	160.6	4.7	*	*	*	1.4	*	*	1.5	1.7
2001.....	4.4	159.2	5.4	*	1.4	1.4	1.5	1.8	*	1.7	*
2000.....	4.5	182.2	4.8	*	1.5	*	*	*	*	1.5	*
1999.....	4.4	170.0	5.0	1.5	1.8	*	*	*	*	1.5	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins 3/, both sexes

2003.....	10.8	82.9	1.6	0.4	0.5	1.5	2.6	3.2	3.4	4.8	5.9
2002.....	10.4	85.7	1.6	0.3	0.4	1.3	2.4	3.0	3.6	4.6	6.0
2001.....	11.4	86.5	1.8	0.4	0.5	1.8	3.0	3.5	4.2	6.2	7.7
2000.....	11.3	95.6	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.7	7.6
1999.....	9.6	93.2	1.5	0.3	0.4	1.1	2.0	2.3	3.1	4.1	5.1

All origins 3/, male

2003.....	10.2	96.5	1.8	0.3	0.6	1.9	3.4	4.4	4.5	6.4	7.4
2002.....	10.1	98.5	1.8	0.4	0.4	1.6	3.2	4.1	4.9	6.1	7.6
2001.....	11.3	98.9	2.1	0.4	0.5	2.4	4.0	4.8	5.6	8.0	9.9
2000.....	11.5	110.6	1.7	0.5	0.5	1.9	4.0	4.7	5.9	7.6	10.2
1999.....	9.4	106.6	1.7	0.3	0.4	1.5	2.8	2.9	4.2	5.3	6.7

All origins 3/, female

2003.....	11.4	68.6	1.5	0.5	0.4	1.0	1.7	1.9	2.3	3.3	4.4
2002.....	10.7	72.3	1.4	0.2	0.3	0.8	1.6	1.8	2.3	3.1	4.5
2001.....	11.4	73.6	1.5	0.4	0.5	1.1	1.9	2.2	2.8	4.4	5.5
2000.....	11.2	79.9	1.5	0.4	0.4	1.2	1.8	2.2	2.8	3.7	5.0
1999.....	9.7	79.2	1.3	0.3	0.3	0.8	1.2	1.6	2.1	3.0	3.4

Hispanic, both sexes

2003.....	4.1	56.8	0.9	*	*	1.4	1.4	2.4	2.2	3.0	3.2
2002.....	4.0	56.6	1.3	*	*	1.1	1.6	1.8	2.3	2.6	3.6
2001.....	5.9	60.7	1.6	*	0.6	2.3	3.2	3.7	4.7	5.9	6.1
2000.....	5.9	65.6	1.7	*	*	1.6	2.6	3.5	4.5	6.3	7.2
1999.....	3.8	54.9	1.0	*	*	1.0	1.6	1.8	2.6	2.9	3.1

Hispanic, male

2003.....	4.8	69.9	*	*	*	1.6	1.8	3.5	3.2	4.3	4.4
2002.....	4.6	66.4	1.6	*	*	1.2	2.1	2.5	3.4	3.6	4.4
2001.....	7.4	67.2	1.8	*	*	3.4	4.3	5.4	6.6	8.3	8.6
2000.....	7.5	78.6	2.1	*	*	1.7	3.9	5.1	6.5	9.3	10.6
1999.....	4.4	61.3	*	*	*	1.3	2.4	2.4	3.5	4.2	4.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Hispanic, female

2003.....	3.2	43.2	*	*	*	*	*	*	*	1.5	1.8
2002.....	3.3	46.4	*	*	*	*	*	*	*	1.4	2.8
2001.....	4.3	53.9	*	*	*	*	1.8	1.7	2.5	3.2	3.6
2000.....	4.1	52.1	1.4	*	*	1.4	*	1.6	2.3	3.0	3.6
1999.....	3.1	48.2	*	*	*	*	*	*	1.6	1.6	*

Non-Hispanic 4/, both sexes

2003.....	11.8	89.4	1.8	0.4	0.5	1.5	2.8	3.4	3.7	5.1	6.2
2002.....	11.3	92.8	1.7	0.3	0.4	1.3	2.5	3.2	3.9	4.9	6.3
2001.....	12.1	92.6	1.8	0.4	0.5	1.7	2.9	3.4	4.1	6.2	7.8
2000.....	12.0	102.4	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.5	7.5
1999.....	10.3	101.9	1.6	0.3	0.4	1.2	2.1	2.4	3.2	4.2	5.2

Non-Hispanic 4/, male

2003.....	11.0	102.9	2.0	0.3	0.6	1.9	3.8	4.7	4.8	6.7	7.7
2002.....	10.9	106.2	1.9	0.4	0.4	1.7	3.4	4.5	5.1	6.5	8.0
2001.....	11.8	106.2	2.1	0.4	0.5	2.2	3.9	4.6	5.3	7.9	9.9
2000.....	11.9	118.0	1.7	0.5	0.5	1.9	3.9	4.5	5.6	7.2	10.0
1999.....	10.0	116.7	1.9	0.3	0.4	1.5	2.9	3.0	4.3	5.3	6.9

Non-Hispanic 4/, female

2003.....	12.6	75.3	1.6	0.5	0.4	1.0	1.8	2.1	2.5	3.6	4.7
2002.....	11.7	78.7	1.4	0.3	0.3	0.8	1.7	2.0	2.6	3.4	4.6
2001.....	12.3	78.3	1.5	0.3	0.4	1.1	1.9	2.2	2.9	4.6	5.7
2000.....	12.1	86.1	1.5	0.3	0.4	1.2	1.9	2.2	2.9	3.8	5.1
1999.....	10.5	86.3	1.4	0.3	0.3	0.8	1.3	1.7	2.2	3.1	3.6

Non-Hispanic White, both sexes

2003.....	12.3	72.6	1.5	0.4	0.4	1.5	2.8	3.2	3.6	5.0	5.9
2002.....	11.6	76.0	1.3	0.2	0.3	1.2	2.5	3.1	3.7	4.7	6.0
2001.....	12.4	73.1	1.5	0.3	0.4	1.7	2.8	3.3	3.9	6.0	7.7
2000.....	12.2	79.7	1.4	0.3	0.4	1.5	2.8	3.1	4.0	5.4	7.2
1999.....	10.2	79.3	1.4	0.3	0.3	1.0	1.8	2.0	3.0	3.8	4.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic White, male

2003.....	11.0	87.4	1.6	0.3	0.4	2.0	3.9	4.4	4.8	6.4	7.3
2002.....	10.7	88.2	1.4	*	0.4	1.6	3.4	4.4	4.9	6.2	7.6
2001.....	11.8	85.2	1.7	*	0.5	2.1	3.8	4.6	5.1	7.5	9.9
2000.....	11.8	91.8	1.7	0.4	0.4	1.9	3.8	4.2	5.4	7.2	9.8
1999.....	9.6	92.3	1.7	0.3	0.4	1.3	2.6	2.7	3.8	4.9	6.0

Non-Hispanic White, female

2003.....	13.5	57.1	1.4	0.5	0.3	1.0	1.7	2.1	2.4	3.5	4.5
2002.....	12.5	63.2	1.3	*	*	0.8	1.5	1.8	2.5	3.2	4.4
2001.....	13.1	60.4	1.3	*	*	1.2	1.9	1.9	2.8	4.5	5.4
2000.....	12.6	66.9	1.2	*	0.4	1.0	1.7	2.0	2.6	3.5	4.7
1999.....	10.9	65.5	1.0	0.4	*	0.7	1.0	1.4	2.1	2.6	3.0

Non-Hispanic Black, both sexes

2003.....	12.7	161.6	3.3	*	1.1	1.5	3.5	5.1	5.2	7.3	9.9
2002.....	12.6	165.4	2.7	0.8	0.8	1.9	3.3	4.8	5.9	7.8	9.8
2001.....	12.9	176.9	3.0	0.6	0.7	1.7	3.8	4.8	5.5	8.5	10.1
2000.....	13.9	205.6	2.3	0.6	0.8	2.2	4.3	5.5	6.4	7.3	11.3
1999.....	13.0	202.0	2.9	*	*	1.8	3.8	4.4	5.5	7.9	10.5

Non-Hispanic Black, male

2003.....	13.8	171.9	3.6	*	1.3	1.8	4.4	7.6	6.4	9.7	12.6
2002.....	14.4	186.3	3.6	*	*	2.5	3.7	6.0	8.2	10.6	12.7
2001.....	14.2	195.9	3.7	*	*	2.4	5.3	5.6	6.6	10.9	12.0
2000.....	15.7	236.8	2.2	*	*	2.1	5.7	7.3	8.2	8.5	13.3
1999.....	14.9	228.0	2.9	*	*	2.5	4.9	5.5	8.4	9.5	13.8

Non-Hispanic Black, female

2003.....	11.6	150.9	3.0	*	*	*	2.7	2.9	4.0	5.1	7.5
2002.....	10.9	143.6	1.9	*	*	*	2.9	3.7	3.8	5.3	7.2
2001.....	11.7	156.8	2.2	*	*	*	2.4	4.1	4.5	6.4	8.4
2000.....	12.4	173.4	2.4	*	*	2.2	3.0	3.8	4.8	6.3	9.4
1999.....	11.2	175.3	2.8	*	*	*	2.7	3.5	2.9	6.5	7.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

All origins 3/, both sexes

2003.....	69.3	29.7	4.0	2.1	2.4	4.1	5.8	7.4	9.3	14.5	23.1
2002.....	67.5	28.4	4.2	1.9	2.3	3.5	5.6	6.6	9.0	14.1	23.5
2001.....	65.5	28.0	4.2	1.9	2.3	3.8	5.2	6.5	8.9	14.1	21.7
2000.....	63.4	29.0	4.0	1.8	2.5	3.6	5.4	6.5	9.0	13.9	22.1
1999.....	61.5	30.2	4.2	2.1	2.4	3.9	5.3	6.3	8.9	14.0	21.6

All origins 3/, male

2003.....	57.3	34.8	4.2	2.3	2.8	4.6	6.6	8.2	10.4	16.5	27.8
2002.....	55.8	33.4	4.6	2.2	2.5	4.0	6.3	7.4	10.6	16.4	28.2
2001.....	53.8	31.4	4.6	1.8	2.5	4.4	6.0	7.3	9.9	16.8	25.7
2000.....	52.3	32.3	4.3	1.9	2.6	4.2	6.5	7.5	10.4	16.5	26.3
1999.....	51.0	35.0	4.5	2.2	2.5	4.7	5.9	7.3	10.4	16.8	26.6

All origins 3/, female

2003.....	81.0	24.5	3.8	1.8	2.1	3.4	4.9	6.5	8.1	12.5	18.5
2002.....	78.8	23.0	3.9	1.7	2.1	3.0	4.9	5.8	7.3	11.9	18.9
2001.....	76.7	24.5	3.8	2.1	2.0	3.1	4.4	5.6	8.0	11.5	17.7
2000.....	74.0	25.6	3.6	1.8	2.3	3.1	4.4	5.4	7.5	11.3	17.9
1999.....	71.6	25.3	3.9	2.1	2.3	3.1	4.7	5.3	7.4	11.3	16.6

Hispanic, both sexes

2003.....	23.1	27.1	4.3	1.8	2.5	4.2	4.0	5.2	7.2	11.1	17.4
2002.....	21.6	25.1	3.8	2.0	2.2	3.1	4.2	4.6	6.6	9.7	17.0
2001.....	21.2	22.9	3.7	1.7	1.8	3.2	4.1	4.6	5.8	9.7	15.4
2000.....	20.7	25.4	3.7	1.6	1.8	2.6	3.9	5.0	6.8	11.0	17.1
1999.....	21.5	26.5	3.7	1.4	1.9	3.1	3.8	5.1	8.0	11.2	17.4

Hispanic, male

2003.....	22.2	32.6	4.5	1.9	2.7	4.5	4.3	6.1	8.7	12.6	21.6
2002.....	21.1	27.2	4.1	2.5	2.3	3.2	4.7	5.3	7.9	12.1	23.1
2001.....	20.2	25.4	4.3	1.6	2.0	3.9	4.2	5.0	7.1	12.6	20.0
2000.....	20.2	25.3	3.6	1.6	2.1	3.2	4.6	5.5	8.8	14.3	23.1
1999.....	21.0	32.1	3.4	1.6	1.7	3.6	3.9	5.9	10.6	15.3	23.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

Hispanic, female

2003.....	24.1	21.5	4.0	1.7	2.2	3.9	3.6	4.0	5.3	9.4	12.8
2002.....	22.1	22.8	3.5	1.5	2.0	3.0	3.5	3.8	5.1	7.1	10.4
2001.....	22.3	20.2	3.1	1.8	1.5	2.5	3.9	4.0	4.3	6.5	10.6
2000.....	21.4	25.5	3.8	1.6	1.4	1.9	3.0	4.4	4.6	7.4	10.9
1999.....	22.0	20.6	4.0	1.3	2.1	2.5	3.7	4.2	5.0	6.7	11.4

Non-Hispanic 4/, both sexes

2003.....	76.5	30.1	4.0	2.1	2.4	4.0	6.1	7.9	9.7	15.0	23.8
2002.....	74.4	29.0	4.3	1.9	2.3	3.6	5.9	7.1	9.4	14.8	24.3
2001.....	71.8	29.0	4.3	2.0	2.3	3.9	5.4	6.9	9.5	14.8	22.3
2000.....	69.3	29.6	4.0	1.9	2.6	3.8	5.8	6.8	9.3	14.3	22.5
1999.....	66.8	30.8	4.3	2.3	2.5	4.0	5.6	6.5	9.0	14.3	21.9

Non-Hispanic 4/, male

2003.....	63.0	35.0	4.2	2.4	2.8	4.6	7.1	8.7	10.7	17.1	28.5
2002.....	61.2	34.9	4.6	2.1	2.5	4.2	6.6	7.9	11.2	17.1	28.7
2001.....	58.8	32.7	4.6	1.8	2.6	4.5	6.4	7.8	10.4	17.4	26.2
2000.....	56.9	33.6	4.4	1.9	2.7	4.3	6.9	7.9	10.7	16.8	26.5
1999.....	55.2	35.1	4.7	2.3	2.6	4.9	6.3	7.6	10.3	16.8	26.7

Non-Hispanic 4/, female

2003.....	89.4	25.1	3.7	1.8	2.0	3.3	5.1	7.0	8.6	12.9	19.2
2002.....	86.8	22.7	4.0	1.8	2.1	3.0	5.2	6.2	7.7	12.5	19.9
2001.....	84.2	25.2	4.0	2.1	2.1	3.2	4.5	6.0	8.6	12.1	18.5
2000.....	81.0	25.4	3.5	1.8	2.4	3.3	4.6	5.6	7.9	11.8	18.6
1999.....	77.9	26.2	3.9	2.2	2.4	3.2	4.8	5.6	7.7	11.8	17.1

Non-Hispanic White, both sexes

2003.....	83.0	24.6	3.5	2.0	2.2	3.8	5.8	7.0	8.4	13.2	21.8
2002.....	80.5	24.7	3.8	1.8	2.2	3.5	5.2	6.3	8.3	13.2	22.1
2001.....	77.3	23.2	3.9	2.0	2.2	3.6	5.0	6.0	8.2	13.0	19.6
2000.....	74.3	24.0	3.4	1.7	2.4	3.7	5.4	5.8	8.1	12.3	19.9
1999.....	71.5	24.2	4.0	2.1	2.3	3.9	5.0	6.0	8.0	12.1	19.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

Non-Hispanic White, male

2003.....	66.9	30.4	3.6	2.5	2.5	4.7	7.3	8.2	9.7	15.5	26.9
2002.....	64.8	31.4	4.0	2.0	2.3	4.2	6.2	7.3	10.3	16.0	27.0
2001.....	61.9	25.3	4.4	1.9	2.4	4.4	6.2	7.0	9.3	15.8	23.5
2000.....	59.6	26.7	3.8	1.9	2.7	4.3	6.6	7.3	9.7	14.9	23.9
1999.....	57.6	27.4	4.5	2.1	2.3	5.0	5.9	7.2	9.4	14.8	24.0

Non-Hispanic White, female

2003.....	98.4	18.6	3.5	1.6	1.9	2.9	4.3	5.8	7.1	10.8	16.7
2002.....	95.5	17.7	3.6	1.7	2.0	2.6	4.2	5.3	6.3	10.3	17.1
2001.....	92.1	21.0	3.4	2.1	2.0	2.8	3.8	5.0	7.2	10.2	15.8
2000.....	88.3	21.2	2.9	1.6	2.1	3.0	4.1	4.3	6.6	9.7	15.9
1999.....	84.9	20.9	3.5	2.1	2.3	2.9	4.1	4.7	6.5	9.4	14.0

Non-Hispanic Black, both sexes

2003.....	63.0	52.6	5.5	2.3	3.3	5.0	8.3	13.8	17.5	28.1	39.8
2002.....	61.7	46.1	6.1	2.3	3.0	4.4	9.8	12.7	17.2	26.5	42.8
2001.....	61.1	53.8	5.6	2.0	3.0	5.4	8.0	12.0	17.3	27.4	42.3
2000.....	60.1	51.3	6.1	2.4	3.5	4.9	8.6	13.4	17.8	27.8	42.7
1999.....	58.0	58.4	5.5	3.0	3.2	4.9	9.3	10.9	16.0	29.5	43.4

Non-Hispanic Black, male

2003.....	57.6	56.8	6.3	1.9	4.1	4.8	7.3	13.4	17.8	29.9	43.4
2002.....	56.6	50.6	6.6	2.3	3.4	4.3	9.8	13.0	18.5	27.2	45.2
2001.....	56.4	61.3	5.2	1.6	3.4	5.6	8.1	12.9	18.2	30.5	48.2
2000.....	56.1	61.4	6.4	2.1	3.2	5.1	9.4	13.6	18.9	30.9	48.8
1999.....	54.4	66.5	5.6	3.3	3.9	5.0	10.1	11.0	16.7	32.5	48.5

Non-Hispanic Black, female

2003.....	67.9	48.4	4.6	2.7	2.6	5.3	9.4	14.1	17.3	26.5	36.7
2002.....	66.3	41.3	5.6	2.3	2.5	4.5	9.9	12.4	16.0	25.9	40.7
2001.....	65.2	46.0	6.0	2.5	2.7	5.2	8.0	11.2	16.5	24.6	37.2
2000.....	63.7	40.9	5.8	2.7	3.7	4.8	7.8	13.2	16.8	25.0	37.3
1999.....	61.3	50.0	5.4	2.7	2.5	4.7	8.6	10.8	15.3	26.9	38.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All origins 3/, both sexes

2003.....	37.6	23.6	10.9	5.5	7.2	33.0	41.1	32.8	30.2	35.0	40.4
2002.....	37.0	23.5	10.5	5.9	7.3	35.0	40.9	32.3	30.8	34.9	39.4
2001.....	35.7	24.2	11.2	6.4	7.4	32.8	39.5	31.4	28.5	34.1	36.7
2000.....	34.8	23.1	11.9	6.8	7.7	33.4	38.8	30.4	28.6	32.9	35.4
1999.....	35.1	22.3	12.4	7.1	8.1	33.3	37.5	30.0	29.2	32.2	35.4

All origins 3/, male

2003.....	49.3	26.4	13.0	6.4	9.3	44.7	61.8	49.4	45.1	50.2	58.0
2002.....	48.9	26.8	12.9	7.1	9.2	47.6	62.4	49.5	46.1	50.5	56.8
2001.....	47.2	26.2	13.3	7.6	9.5	45.3	60.7	48.0	43.1	49.5	54.1
2000.....	46.2	25.7	14.0	7.8	9.9	45.8	58.9	47.0	43.1	48.9	53.1
1999.....	46.4	25.4	14.4	8.5	10.4	44.8	57.1	46.1	43.9	47.4	53.1

All origins 3/, female

2003.....	26.2	20.7	8.7	4.6	4.9	20.6	19.2	15.5	15.1	19.7	23.0
2002.....	25.6	20.0	8.1	4.7	5.3	21.8	18.4	14.5	15.2	19.2	22.4
2001.....	24.5	22.1	8.9	5.0	5.3	19.5	17.2	14.3	13.8	18.7	19.6
2000.....	23.8	20.5	9.7	5.7	5.5	20.3	17.8	13.5	13.9	17.0	18.0
1999.....	24.1	19.0	10.3	5.6	5.6	21.1	17.1	13.7	14.3	17.0	18.1

Hispanic, both sexes

2003.....	26.1	15.1	9.9	5.7	5.8	31.0	36.3	31.8	26.8	29.9	33.9
2002.....	26.1	15.2	10.9	5.3	5.4	30.8	39.6	28.8	26.4	28.5	35.6
2001.....	25.8	18.8	10.3	5.0	5.7	27.8	37.0	30.2	26.1	31.6	33.1
2000.....	25.0	15.3	10.4	5.7	5.6	27.9	35.5	27.9	26.6	30.3	33.1
1999.....	25.5	17.7	11.9	4.9	6.4	26.1	35.3	28.7	28.0	31.0	34.6

Hispanic, male

2003.....	38.3	17.4	11.9	7.0	8.1	44.9	55.4	48.8	41.9	45.4	51.1
2002.....	38.5	19.5	13.5	6.3	6.8	45.8	60.0	44.5	41.3	42.9	55.3
2001.....	37.6	18.0	12.7	6.0	6.4	40.2	56.9	47.0	40.2	48.9	51.7
2000.....	36.9	17.0	11.6	5.7	6.5	40.8	54.8	44.0	41.7	46.3	53.9
1999.....	37.7	20.7	13.5	6.0	7.8	37.3	54.5	44.7	43.4	48.7	54.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Hispanic, female

2003.....	13.1	12.7	7.8	4.3	3.5	15.9	12.7	10.9	9.3	12.4	15.4
2002.....	12.8	10.8	8.2	4.3	3.8	14.4	14.5	9.7	9.2	12.4	14.5
2001.....	13.2	19.7	7.8	4.0	5.0	14.0	12.6	10.2	10.0	12.6	13.5
2000.....	12.4	13.5	9.2	5.6	4.7	13.1	11.9	9.0	9.4	12.9	11.3
1999.....	12.6	14.6	10.3	3.7	4.9	13.4	12.2	9.9	10.3	11.6	13.7

Non-Hispanic 4/, both sexes

2003.....	39.2	25.9	11.1	5.5	7.5	33.3	42.0	32.9	30.8	35.7	41.1
2002.....	38.5	25.5	10.4	6.0	7.6	35.7	41.1	33.0	31.5	35.7	39.7
2001.....	36.9	25.4	11.3	6.6	7.7	33.6	39.8	31.5	28.8	34.3	36.9
2000.....	36.0	25.0	12.2	7.0	8.1	34.3	39.3	30.8	28.8	33.1	35.4
1999.....	36.2	23.4	12.4	7.5	8.3	34.5	37.8	30.1	29.2	32.2	35.3

Non-Hispanic 4/, male

2003.....	50.9	28.7	13.2	6.2	9.6	44.6	63.1	49.3	45.6	50.9	58.6
2002.....	50.3	28.6	12.7	7.2	9.6	47.8	62.7	50.6	46.8	51.5	56.7
2001.....	48.5	28.0	13.4	7.9	10.0	46.1	61.4	48.1	43.4	49.3	54.0
2000.....	47.3	27.7	14.5	8.3	10.5	46.6	59.6	47.4	43.1	49.0	52.5
1999.....	47.4	26.6	14.5	9.0	10.8	46.1	57.3	46.1	43.6	46.9	52.5

Non-Hispanic 4/, female

2003.....	28.1	22.9	8.9	4.7	5.2	21.4	20.4	16.5	16.2	20.8	23.9
2002.....	27.3	22.3	8.1	4.8	5.6	23.0	19.2	15.5	16.3	20.2	23.2
2001.....	26.0	22.6	9.2	5.3	5.3	20.5	18.1	15.1	14.4	19.6	20.2
2000.....	25.2	22.2	9.7	5.6	5.6	21.5	18.9	14.3	14.6	17.5	18.6
1999.....	25.5	20.0	10.2	6.0	5.8	22.4	17.9	14.3	14.9	17.6	18.4

Non-Hispanic White, both sexes

2003.....	41.6	21.6	10.7	5.0	7.2	36.5	45.7	34.9	32.3	37.0	42.3
2002.....	40.6	19.8	9.4	5.1	7.3	39.0	44.2	34.3	33.2	36.4	40.8
2001.....	38.5	20.9	10.9	5.9	7.3	36.0	42.2	32.9	29.6	34.9	36.8
2000.....	37.3	20.1	11.1	6.3	8.0	37.3	41.4	31.1	29.1	33.4	35.0
1999.....	37.2	19.0	11.1	6.6	8.1	37.2	39.1	30.8	29.1	32.1	34.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic White, male

2003.....	53.1	24.1	12.9	5.6	9.2	48.4	68.2	52.0	47.4	51.9	60.2
2002.....	52.0	23.3	11.1	6.1	9.2	51.6	67.2	51.9	48.9	51.9	57.9
2001.....	49.7	22.9	13.2	6.7	9.5	48.8	65.0	49.7	44.2	49.7	53.7
2000.....	48.1	22.9	13.4	7.5	10.2	50.0	62.6	47.2	43.4	49.3	51.5
1999.....	47.8	22.0	12.9	7.7	10.4	48.8	59.1	46.9	43.0	46.6	51.5

Non-Hispanic White, female

2003.....	30.5	19.0	8.4	4.4	5.1	24.0	22.4	17.4	17.1	21.9	24.4
2002.....	29.6	16.0	7.5	4.1	5.4	25.7	20.7	16.4	17.2	20.9	23.8
2001.....	27.8	18.7	8.5	5.0	5.1	22.5	18.9	15.8	14.7	20.0	19.9
2000.....	26.9	17.1	8.7	5.0	5.6	24.0	19.6	14.7	14.6	17.5	18.5
1999.....	27.0	15.9	9.3	5.4	5.7	25.1	18.6	14.5	15.0	17.6	18.2

Non-Hispanic Black, both sexes

2003.....	33.3	44.7	13.7	7.7	8.4	21.3	31.7	30.6	31.1	36.7	42.7
2002.....	34.0	50.1	14.6	9.6	9.2	24.4	32.7	34.1	30.9	39.1	40.8
2001.....	34.4	44.5	14.2	9.8	9.7	25.3	34.2	31.3	30.9	36.5	45.3
2000.....	34.4	46.5	16.8	10.0	9.1	24.3	35.5	34.2	32.8	36.7	44.6
1999.....	36.1	44.1	18.2	11.8	9.9	25.8	35.8	32.5	34.0	37.4	44.3

Non-Hispanic Black, male

2003.....	47.4	49.0	15.9	9.1	11.0	30.9	49.3	47.9	48.9	56.7	62.0
2002.....	49.0	52.1	18.9	11.4	12.4	35.1	50.7	55.8	48.8	60.3	59.1
2001.....	49.5	50.4	15.8	13.0	13.2	37.0	52.9	49.5	48.0	54.3	68.4
2000.....	50.3	49.5	19.4	12.2	12.5	35.3	54.5	55.9	51.1	56.9	69.9
1999.....	52.3	48.5	21.8	15.0	13.5	37.9	56.5	52.7	53.4	56.5	70.0

Non-Hispanic Black, female

2003.....	20.5	40.3	11.4	6.1	5.7	11.5	14.2	14.7	15.1	18.9	25.7
2002.....	20.3	48.0	10.1	7.7	5.9	13.3	15.2	14.4	14.9	20.3	24.6
2001.....	20.6	38.3	12.4	6.4	6.0	13.3	16.3	14.6	15.5	20.7	24.9
2000.....	20.1	43.3	14.2	7.8	5.5	13.0	17.5	14.6	16.5	18.8	22.3
1999.....	21.4	39.6	14.5	8.4	6.3	13.4	16.3	14.3	16.6	20.5	21.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Transport accidents (V01-V99,Y85)

All origins 3/, both sexes

2003.....	16.5	3.7	4.1	3.5	5.1	26.7	28.8	20.1	16.5	16.7	17.5
2002.....	16.8	3.1	4.1	3.5	4.9	28.4	29.9	20.5	17.1	17.1	17.1
2001.....	16.6	3.7	4.4	3.8	5.0	26.6	29.2	20.9	16.9	17.4	17.3
2000.....	16.6	4.6	4.5	4.1	5.3	27.1	29.3	20.4	16.8	17.1	16.8
1999.....	16.6	5.0	4.6	4.4	5.3	27.1	28.2	20.0	17.0	16.9	16.6

All origins 3/, male

2003.....	23.3	4.0	4.5	4.0	6.3	34.7	42.4	29.8	24.8	24.1	25.5
2002.....	23.7	3.5	4.8	4.0	5.9	37.0	44.4	31.0	25.4	24.7	24.5
2001.....	23.5	3.3	5.0	4.3	6.2	35.4	43.9	31.4	25.4	24.7	25.4
2000.....	23.3	4.8	5.0	4.4	6.4	35.5	43.3	30.8	24.7	24.8	24.8
1999.....	23.1	5.2	5.1	5.0	6.5	34.8	41.8	29.9	24.7	24.0	24.0

All origins 3/, female

2003.....	10.0	3.4	3.7	3.0	3.8	18.3	14.3	10.0	8.1	9.3	9.5
2002.....	10.1	2.8	3.4	3.1	3.9	19.3	14.6	9.7	8.7	9.6	9.9
2001.....	10.0	4.2	3.8	3.3	3.8	17.3	13.8	10.1	8.3	10.2	9.4
2000.....	10.2	4.3	4.0	3.8	4.1	18.1	14.6	9.8	8.7	9.5	9.0
1999.....	10.4	4.8	4.2	3.6	4.1	19.0	14.0	10.0	9.3	9.8	9.4

Hispanic, both sexes

2003.....	15.2	4.5	4.2	3.8	4.4	25.2	27.1	21.6	16.5	15.9	17.1
2002.....	15.3	3.1	5.8	3.7	3.9	24.6	30.2	20.2	16.2	15.2	17.0
2001.....	15.6	5.3	5.3	3.3	4.3	22.7	29.0	21.6	15.9	17.3	17.3
2000.....	14.8	5.4	5.2	4.0	4.3	21.9	27.0	19.4	15.0	15.6	16.9
1999.....	14.4	5.3	5.5	3.1	4.7	20.0	25.9	19.6	15.9	14.4	16.6

Hispanic, male

2003.....	22.2	5.7	4.6	4.8	6.1	35.1	40.7	32.7	25.5	23.2	24.7
2002.....	22.4	*	7.1	4.2	5.1	35.4	44.7	30.3	24.5	22.4	25.6
2001.....	22.8	*	6.0	3.8	4.9	31.7	44.1	33.2	24.1	25.8	26.0
2000.....	21.6	6.1	5.6	3.8	5.2	31.3	40.4	30.2	22.1	22.9	25.6
1999.....	20.8	5.7	5.8	3.6	5.6	27.7	38.5	29.7	23.5	21.6	25.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Transport accidents (V01-V99,Y85)

Hispanic, female

2003.....	7.6	*	3.8	2.8	2.6	14.4	10.3	8.0	6.1	7.7	8.9
2002.....	7.7	*	4.5	3.2	2.7	12.8	12.4	8.0	6.6	7.2	7.8
2001.....	8.0	6.7	4.6	2.7	3.7	12.7	10.5	7.8	6.4	7.9	8.2
2000.....	7.6	*	4.8	4.1	3.3	11.3	10.7	6.8	6.8	7.8	7.7
1999.....	7.6	*	5.2	2.5	3.8	11.3	10.7	7.9	7.2	6.5	7.7

Non-Hispanic 4/, both sexes

2003.....	16.7	3.5	4.0	3.4	5.2	26.9	29.0	19.7	16.5	16.8	17.5
2002.....	16.9	3.2	3.7	3.5	5.1	29.0	29.7	20.5	17.3	17.3	17.1
2001.....	16.7	3.4	4.2	3.9	5.2	27.2	29.2	20.7	17.1	17.4	17.2
2000.....	16.8	4.3	4.3	4.1	5.4	27.9	29.7	20.5	17.0	17.3	16.7
1999.....	16.9	4.9	4.4	4.6	5.4	28.3	28.5	20.0	17.2	17.2	16.5

Non-Hispanic 4/, male

2003.....	23.4	3.5	4.5	3.7	6.3	34.5	42.6	28.9	24.6	24.1	25.6
2002.....	23.8	3.4	4.2	3.9	6.0	37.2	44.1	31.1	25.5	24.9	24.3
2001.....	23.5	3.2	4.7	4.4	6.5	36.0	43.7	30.9	25.6	24.4	25.2
2000.....	23.4	4.5	4.8	4.5	6.6	36.2	43.9	30.8	25.1	24.9	24.5
1999.....	23.3	5.1	4.9	5.3	6.6	36.0	42.3	29.7	24.8	24.3	23.7

Non-Hispanic 4/, female

2003.....	10.3	3.5	3.6	3.1	4.0	18.9	15.1	10.4	8.4	9.5	9.6
2002.....	10.4	2.9	3.1	3.1	4.1	20.4	15.0	10.0	9.1	9.9	10.1
2001.....	10.2	3.6	3.6	3.4	3.8	18.0	14.4	10.5	8.6	10.5	9.5
2000.....	10.5	4.2	3.8	3.7	4.2	19.3	15.3	10.4	9.0	9.8	9.1
1999.....	10.7	4.7	3.9	3.9	4.2	20.3	14.5	10.3	9.6	10.2	9.5

Non-Hispanic White, both sexes

2003.....	17.2	3.2	3.8	3.3	5.4	29.7	30.7	20.1	16.6	17.0	17.7
2002.....	17.5	2.5	3.3	3.3	5.2	32.1	31.4	20.5	17.7	17.3	17.4
2001.....	17.0	3.1	3.8	3.6	5.2	29.6	30.3	21.3	17.1	17.5	17.3
2000.....	17.1	3.5	3.8	4.0	5.6	30.9	30.9	20.2	17.0	17.4	16.6
1999.....	17.1	4.3	3.7	4.4	5.6	31.1	29.5	20.2	17.2	16.9	16.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Transport accidents (V01-V99,Y85)

Non-Hispanic White, male

2003.....	23.9	3.3	4.4	3.5	6.5	37.6	44.7	29.2	24.6	24.0	25.7
2002.....	24.3	2.9	3.6	3.7	6.1	40.8	46.6	30.7	25.8	24.6	24.5
2001.....	23.8	2.9	4.4	4.0	6.4	38.6	45.5	31.5	25.4	24.2	25.0
2000.....	23.7	3.8	4.3	4.5	6.8	39.6	45.6	29.7	24.8	24.8	24.1
1999.....	23.5	4.9	4.0	4.9	6.8	39.0	43.7	29.8	24.3	23.7	23.5

Non-Hispanic White, female

2003.....	10.7	3.0	3.2	3.1	4.2	21.3	16.4	10.7	8.5	9.9	9.6
2002.....	10.9	2.1	2.9	2.8	4.2	22.9	15.8	10.2	9.5	10.0	10.3
2001.....	10.6	3.3	3.2	3.2	3.8	20.0	14.8	10.9	8.7	10.8	9.6
2000.....	10.9	3.2	3.4	3.5	4.4	21.8	15.9	10.5	9.0	10.0	9.1
1999.....	11.0	3.6	3.4	3.7	4.4	22.8	15.0	10.5	9.9	10.2	9.6

Non-Hispanic Black, both sexes

2003.....	15.6	4.4	5.1	3.8	4.7	16.7	24.0	21.0	19.1	18.5	19.2
2002.....	15.7	5.6	5.0	4.7	4.7	18.3	25.1	23.9	18.4	19.7	17.3
2001.....	16.3	*	5.4	5.5	5.4	19.3	26.7	21.6	19.6	18.5	19.8
2000.....	16.4	6.9	5.9	4.6	5.0	17.8	27.7	24.4	20.0	18.4	19.7
1999.....	16.7	7.7	7.1	6.0	5.2	19.2	26.6	22.0	19.2	20.5	18.4

Non-Hispanic Black, male

2003.....	23.3	*	5.1	4.7	5.8	23.4	37.6	32.9	30.7	29.4	28.9
2002.....	23.6	*	6.5	5.0	5.8	24.7	37.8	39.0	29.5	30.5	25.6
2001.....	24.5	*	5.8	6.9	7.3	27.2	40.5	34.1	31.1	27.9	31.5
2000.....	24.8	7.2	6.4	5.0	6.7	24.6	41.7	39.5	31.7	29.5	31.6
1999.....	24.9	*	8.0	7.3	6.6	26.7	40.3	35.5	30.5	30.7	28.8

Non-Hispanic Black, female

2003.....	8.5	*	5.2	2.9	3.5	9.7	10.6	10.0	8.7	8.8	10.7
2002.....	8.5	*	3.6	4.3	3.6	11.6	12.7	10.2	8.5	10.2	10.1
2001.....	8.9	*	5.0	4.1	3.4	11.1	13.5	10.1	9.2	10.2	9.5
2000.....	8.8	*	5.4	4.2	3.2	10.8	14.4	10.7	9.6	8.6	9.2
1999.....	9.4	9.3	6.2	4.6	3.7	11.7	13.7	9.8	9.0	11.5	9.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All origins 3/, both sexes

2003.....	15.4	3.6	3.9	3.3	4.7	25.7	27.5	19.0	15.4	15.4	16.0
2002.....	15.7	3.0	3.9	3.3	4.5	27.6	28.8	19.7	16.0	15.8	15.7
2001.....	15.4	3.6	4.1	3.6	4.6	25.7	28.0	19.6	15.6	15.9	15.5
2000.....	15.4	4.4	4.2	3.8	4.8	26.0	28.0	19.2	15.5	15.6	15.1
1999.....	15.2	4.8	4.2	3.9	4.8	25.9	26.5	18.4	15.4	15.1	14.8

All origins 3/, male

2003.....	21.4	3.9	4.4	3.7	5.9	33.3	40.4	28.2	23.0	21.9	23.1
2002.....	21.9	3.3	4.5	3.8	5.3	36.0	42.6	29.6	23.6	22.4	22.1
2001.....	21.5	3.2	4.5	4.0	5.7	34.0	42.1	29.5	23.1	22.3	22.5
2000.....	21.3	4.6	4.7	4.1	5.8	33.9	41.2	28.6	22.6	22.1	21.9
1999.....	20.9	5.0	4.7	4.4	5.7	33.1	39.1	27.2	22.2	21.2	21.0

All origins 3/, female

2003.....	9.5	3.3	3.5	2.8	3.5	17.8	13.9	9.5	7.7	8.8	9.0
2002.....	9.8	2.8	3.3	2.9	3.6	18.9	14.3	9.4	8.3	9.2	9.4
2001.....	9.5	4.0	3.6	3.0	3.5	16.9	13.4	9.5	8.0	9.6	8.7
2000.....	9.7	4.2	3.8	3.5	3.8	17.5	14.2	9.5	8.2	9.1	8.4
1999.....	9.7	4.7	3.8	3.3	3.8	18.3	13.4	9.4	8.5	9.0	8.6

Hispanic, both sexes

2003.....	14.4	4.5	4.1	3.7	4.2	24.6	26.2	20.3	15.6	15.0	15.9
2002.....	14.6	3.1	5.6	3.5	3.7	23.9	29.0	19.6	15.6	14.5	16.1
2001.....	14.3	4.9	4.7	3.0	4.0	22.1	27.7	20.2	14.5	15.4	15.2
2000.....	13.9	5.2	4.9	3.7	4.0	21.2	26.0	18.5	14.0	14.4	15.2
1999.....	13.3	5.3	5.2	2.9	4.4	19.1	24.5	18.2	14.5	13.0	15.0

Hispanic, male

2003.....	21.1	5.7	4.3	4.6	5.7	34.3	39.5	30.9	24.1	21.6	22.6
2002.....	21.3	*	6.8	3.9	4.9	34.3	42.7	29.3	23.4	21.3	24.2
2001.....	21.0	*	5.3	3.6	4.5	30.8	42.1	31.1	22.2	22.9	23.4
2000.....	20.1	6.1	5.2	3.5	4.8	30.2	38.7	28.6	20.7	20.6	22.9
1999.....	19.1	5.7	5.6	3.3	5.0	26.4	36.4	27.3	21.4	19.4	22.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Hispanic, female

2003.....	7.3	*	3.8	2.7	2.6	14.1	9.8	7.4	5.9	7.4	8.7
2002.....	7.4	*	4.4	3.0	2.4	12.5	12.2	7.8	6.6	7.0	7.5
2001.....	7.2	6.2	4.1	2.5	3.5	12.5	10.1	7.1	5.8	7.1	6.6
2000.....	7.2	*	4.6	4.0	3.1	10.9	10.4	6.7	6.4	7.6	7.1
1999.....	7.2	*	4.7	2.4	3.7	10.9	10.3	7.5	6.6	6.0	7.0

Non-Hispanic 4/, both sexes

2003.....	15.5	3.4	3.9	3.2	4.8	25.9	27.7	18.6	15.3	15.4	16.0
2002.....	15.9	3.0	3.5	3.3	4.6	28.3	28.7	19.6	16.1	15.9	15.6
2001.....	15.5	3.2	3.9	3.7	4.7	26.3	28.0	19.4	15.7	16.0	15.5
2000.....	15.6	4.2	4.1	3.8	5.0	26.8	28.3	19.2	15.6	15.7	15.0
1999.....	15.4	4.7	4.0	4.1	4.9	27.0	26.8	18.3	15.5	15.3	14.7

Non-Hispanic 4/, male

2003.....	21.4	3.4	4.3	3.4	5.9	33.0	40.5	27.3	22.7	21.9	23.1
2002.....	21.9	3.2	4.0	3.7	5.4	36.2	42.4	29.6	23.5	22.5	21.8
2001.....	21.5	3.1	4.3	4.1	5.9	34.5	41.9	29.0	23.2	22.1	22.3
2000.....	21.4	4.2	4.5	4.2	6.0	34.5	41.6	28.5	22.8	22.3	21.6
1999.....	21.0	4.8	4.4	4.6	5.8	34.2	39.6	27.1	22.3	21.4	20.7

Non-Hispanic 4/, female

2003.....	9.8	3.3	3.4	2.9	3.7	18.4	14.6	9.9	8.0	9.0	9.0
2002.....	10.1	2.9	3.0	2.8	3.8	20.0	14.7	9.7	8.7	9.5	9.6
2001.....	9.8	3.4	3.5	3.2	3.5	17.6	14.0	9.9	8.3	9.9	8.9
2000.....	10.0	4.2	3.6	3.4	3.9	18.7	14.9	10.0	8.5	9.3	8.5
1999.....	10.0	4.7	3.6	3.5	3.8	19.5	13.9	9.7	8.8	9.4	8.8

Non-Hispanic White, both sexes

2003.....	15.9	3.1	3.6	3.0	4.9	28.5	29.3	18.9	15.3	15.4	16.0
2002.....	16.3	2.4	3.0	3.1	4.8	31.3	30.2	19.5	16.4	15.8	15.9
2001.....	15.8	3.0	3.5	3.4	4.7	28.5	29.0	19.9	15.7	16.0	15.5
2000.....	15.8	3.3	3.7	3.7	5.1	29.7	29.4	18.7	15.5	15.8	14.8
1999.....	15.6	4.1	3.4	3.8	5.0	29.7	27.7	18.4	15.4	15.0	14.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Non-Hispanic White, male

2003.....	21.8	3.2	4.2	3.2	6.0	35.9	42.4	27.5	22.6	21.6	23.1
2002.....	22.3	2.7	3.3	3.5	5.6	39.7	44.6	29.0	23.6	22.1	22.0
2001.....	21.7	2.9	3.9	3.7	5.9	36.9	43.5	29.3	22.9	21.8	22.0
2000.....	21.5	3.4	4.1	4.2	6.1	37.9	43.0	27.2	22.4	22.1	21.1
1999.....	21.1	4.7	3.6	4.3	5.9	37.0	40.8	26.9	21.8	20.7	20.4

Non-Hispanic White, female

2003.....	10.2	2.9	3.0	2.9	3.7	20.7	15.8	10.1	8.0	9.2	9.0
2002.....	10.5	2.1	2.8	2.6	3.9	22.5	15.5	9.8	9.0	9.5	9.8
2001.....	10.1	3.1	3.1	3.0	3.5	19.6	14.2	10.3	8.4	10.2	8.9
2000.....	10.3	3.2	3.2	3.2	4.1	21.1	15.4	10.1	8.5	9.5	8.5
1999.....	10.3	3.5	3.1	3.3	4.0	22.0	14.3	9.8	9.0	9.3	8.7

Non-Hispanic Black, both sexes

2003.....	14.7	4.4	5.0	3.6	4.5	15.9	23.1	20.2	18.1	17.3	17.9
2002.....	14.8	5.4	4.9	4.3	4.3	17.7	24.5	23.3	17.6	18.4	16.1
2001.....	15.2	*	5.3	5.1	4.9	18.6	26.1	20.7	18.5	17.2	18.1
2000.....	15.5	6.9	5.6	4.3	4.7	16.9	26.9	23.7	18.6	16.9	18.3
1999.....	15.5	7.6	6.5	5.5	4.8	18.3	25.6	20.9	17.6	18.8	16.9

Non-Hispanic Black, male

2003.....	21.7	*	5.0	4.3	5.4	22.2	35.8	31.6	28.9	27.3	26.9
2002.....	22.0	*	6.3	4.8	5.1	23.9	36.9	37.7	28.0	28.2	23.3
2001.....	22.7	*	5.6	6.5	6.5	26.1	39.4	32.7	29.3	25.6	28.2
2000.....	23.1	7.2	6.0	4.5	6.4	23.2	40.3	38.0	29.3	26.9	29.0
1999.....	22.8	*	7.0	6.5	6.0	25.3	38.8	33.6	28.0	27.8	26.0

Non-Hispanic Black, female

2003.....	8.3	*	4.9	2.9	3.5	9.4	10.5	9.6	8.4	8.5	10.1
2002.....	8.3	*	3.5	3.9	3.4	11.4	12.5	10.1	8.2	9.7	9.7
2001.....	8.5	*	4.9	3.7	3.2	10.8	13.4	9.7	8.8	9.7	9.1
2000.....	8.5	*	5.2	4.1	3.1	10.5	14.2	10.6	9.0	8.2	8.8
1999.....	8.9	9.3	5.9	4.5	3.6	11.2	13.1	9.4	8.4	10.8	8.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All origins 3/, both sexes

2003.....	0.5	*	*	0.1	0.2	0.6	0.7	0.5	0.5	0.5	0.6
2002.....	0.4	*	*	0.1	0.2	0.3	0.5	0.3	0.4	0.5	0.5
2001.....	0.5	*	0.2	0.1	0.3	0.5	0.5	0.4	0.4	0.6	0.6
2000.....	0.5	*	0.2	0.2	0.3	0.6	0.6	0.5	0.6	0.7	0.7
1999.....	0.8	*	0.3	0.3	0.3	0.9	1.0	0.8	0.8	0.9	0.9

All origins 3/, male

2003.....	0.7	*	*	*	0.2	0.9	1.0	0.8	0.8	0.8	0.9
2002.....	0.6	*	*	*	0.3	0.5	0.8	0.5	0.6	0.8	0.9
2001.....	0.8	*	0.3	*	0.3	0.7	0.9	0.7	0.8	1.1	1.0
2000.....	0.8	*	*	0.2	0.4	1.0	1.0	0.8	0.9	1.1	1.2
1999.....	1.1	*	0.3	0.4	0.5	1.2	1.7	1.3	1.1	1.4	1.5

All origins 3/, female

2003.....	0.2	*	*	*	*	0.3	0.2	0.2	*	0.2	0.3
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.2
2001.....	0.2	*	*	*	*	0.2	*	*	*	0.2	0.2
2000.....	0.3	*	*	*	*	0.3	0.2	*	0.3	0.2	0.3
1999.....	0.4	*	0.3	*	*	0.5	0.3	0.3	0.5	0.4	0.4

Hispanic, both sexes

2003.....	0.4	*	*	*	*	*	0.6	0.8	*	*	*
2002.....	0.4	*	*	*	*	*	0.8	*	*	*	*
2001.....	0.5	*	*	*	*	*	0.7	0.7	*	0.7	0.9
2000.....	0.6	*	*	*	*	*	*	*	0.7	*	1.1
1999.....	0.8	*	*	*	*	0.7	1.1	0.9	1.0	1.0	1.3

Hispanic, male

2003.....	0.7	*	*	*	*	*	*	1.2	*	*	*
2002.....	0.6	*	*	*	*	*	1.3	*	*	*	*
2001.....	0.9	*	*	*	*	*	1.3	1.2	*	*	1.6
2000.....	0.9	*	*	*	*	*	*	*	*	*	*
1999.....	1.3	*	*	*	*	*	1.6	1.6	1.5	1.7	2.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)											
Hispanic, female											
2003.....	0.1	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	0.5	*	*	*	0.2	0.7	0.7	0.5	0.5	0.5	0.6
2002.....	0.4	*	*	*	0.2	0.3	0.4	0.3	0.4	0.5	0.5
2001.....	0.5	*	*	0.1	0.3	0.5	0.5	0.4	0.4	0.6	0.6
2000.....	0.5	*	*	0.2	0.3	0.7	0.6	0.5	0.6	0.6	0.7
1999.....	0.7	*	0.3	0.3	0.3	0.9	1.0	0.7	0.7	0.9	0.9
Non-Hispanic 4/, male											
2003.....	0.7	*	*	*	0.2	1.0	1.1	0.7	0.8	0.8	0.8
2002.....	0.6	*	*	*	0.4	0.4	0.7	0.5	0.6	0.8	0.9
2001.....	0.7	*	*	*	0.4	0.7	0.8	0.6	0.8	1.0	1.0
2000.....	0.8	*	*	*	0.4	1.0	1.0	0.8	0.8	1.0	1.2
1999.....	1.1	*	0.4	0.4	0.5	1.2	1.7	1.2	1.0	1.3	1.4
Non-Hispanic 4/, female											
2003.....	0.2	*	*	*	*	0.4	0.3	*	*	0.2	0.3
2002.....	0.1	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	0.3	*	*	*	0.2	0.2
2000.....	0.3	*	*	*	*	0.3	*	*	0.3	0.3	0.2
1999.....	0.4	*	*	*	*	0.6	0.3	0.3	0.5	0.4	0.3
Non-Hispanic White, both sexes											
2003.....	0.5	*	*	*	0.2	0.8	0.7	0.5	0.5	0.5	0.5
2002.....	0.4	*	*	*	0.2	0.3	0.4	0.3	0.4	0.5	0.4
2001.....	0.5	*	*	*	0.3	0.5	0.5	0.4	0.4	0.5	0.5
2000.....	0.5	*	*	0.2	0.3	0.7	0.6	0.5	0.5	0.6	0.6
1999.....	0.8	*	0.3	0.3	0.4	0.9	1.0	0.8	0.7	0.9	0.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Hispanic, female

2003.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.7	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.7	*	*	0.2	0.2	0.4	0.6	0.6	0.7	0.9	1.0	
2002.....	0.7	*	*	*	0.2	0.4	0.6	0.6	0.8	0.9	0.9	
2001.....	0.7	*	*	0.1	0.1	0.5	0.7	0.8	0.9	0.8	1.1	
2000.....	0.7	*	*	0.1	0.2	0.5	0.7	0.8	0.9	0.9	1.0	
1999.....	0.7	*	*	0.2	0.2	0.4	0.7	0.9	0.9	1.0	1.0	

Non-Hispanic 4/, male

2003.....	1.2	*	*	*	*	0.6	1.1	0.9	1.1	1.5	1.6	
2002.....	1.2	*	*	*	*	0.6	1.0	1.0	1.3	1.6	1.6	
2001.....	1.2	*	*	*	*	0.8	1.0	1.3	1.5	1.3	1.9	
2000.....	1.2	*	*	*	*	0.7	1.3	1.5	1.5	1.6	1.8	
1999.....	1.2	*	*	*	0.4	0.7	1.0	1.5	1.5	1.5	1.6	

Non-Hispanic 4/, female

2003.....	0.3	*	*	*	*	*	*	0.3	0.3	0.3	0.3	
2002.....	0.2	*	*	*	*	0.3	*	*	0.3	0.3	0.3	
2001.....	0.2	*	*	*	*	*	0.3	0.4	*	0.3	0.4	
2000.....	0.2	*	*	*	*	0.3	*	*	0.2	0.2	0.3	
1999.....	0.3	*	*	*	*	*	0.3	0.3	0.4	0.4	0.4	

Non-Hispanic White, both sexes

2003.....	0.8	*	*	0.2	0.2	0.4	0.7	0.7	0.8	1.0	1.1	
2002.....	0.8	*	*	*	0.2	0.5	0.7	0.7	0.9	1.0	1.1	
2001.....	0.8	*	*	*	0.2	0.6	0.8	1.0	1.0	0.9	1.3	
2000.....	0.8	*	*	*	0.2	0.5	0.9	1.0	0.9	1.0	1.2	
1999.....	0.8	*	*	0.2	0.3	0.5	0.8	1.0	1.0	1.1	1.1	

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Nontransport accidents (W00-X59,Y86)

All origins 3/, both sexes

2003.....	21.0	19.9	6.8	2.0	2.1	6.3	12.3	12.7	13.7	18.3	22.9
2002.....	20.2	20.3	6.4	2.4	2.4	6.6	11.1	11.8	13.6	17.8	22.3
2001.....	19.0	20.5	6.8	2.5	2.4	6.2	10.3	10.5	11.6	16.7	19.4
2000.....	18.2	18.6	7.4	2.7	2.5	6.3	9.5	10.0	11.8	15.8	18.6
1999.....	18.4	17.3	7.7	2.7	2.7	6.2	9.3	10.0	12.1	15.3	18.8

All origins 3/, male

2003.....	26.0	22.4	8.5	2.5	3.0	10.0	19.4	19.6	20.2	26.2	32.4
2002.....	25.2	23.3	8.1	3.1	3.3	10.6	18.0	18.4	20.6	25.9	32.3
2001.....	23.8	22.9	8.3	3.3	3.3	9.9	16.8	16.6	17.7	24.8	28.7
2000.....	22.9	20.9	9.0	3.4	3.5	10.2	15.5	16.2	18.4	24.2	28.3
1999.....	23.3	20.2	9.3	3.5	3.9	10.0	15.3	16.2	19.2	23.4	29.1

All origins 3/, female

2003.....	16.2	17.3	5.0	1.6	1.2	2.3	4.9	5.5	7.0	10.4	13.5
2002.....	15.4	17.2	4.7	1.6	1.5	2.5	3.9	4.9	6.5	9.7	12.5
2001.....	14.5	17.9	5.1	1.8	1.5	2.3	3.4	4.2	5.5	8.5	10.2
2000.....	13.6	16.2	5.7	1.9	1.4	2.2	3.2	3.7	5.2	7.4	9.0
1999.....	13.7	14.2	6.1	2.0	1.5	2.1	3.1	3.7	5.0	7.2	8.7

Hispanic, both sexes

2003.....	11.0	10.6	5.6	1.9	1.4	5.8	9.2	10.2	10.3	14.0	16.8
2002.....	10.8	12.1	5.1	1.6	1.4	6.2	9.4	8.6	10.2	13.2	18.6
2001.....	10.1	13.6	5.0	1.8	1.4	5.1	8.0	8.6	10.3	14.4	15.8
2000.....	10.2	9.9	5.2	1.7	1.4	6.0	8.5	8.4	11.6	14.7	16.2
1999.....	11.1	12.4	6.4	1.8	1.7	6.0	9.4	9.1	12.1	16.6	18.0

Hispanic, male

2003.....	16.1	11.8	7.3	2.2	2.0	9.7	14.7	16.1	16.4	22.2	26.4
2002.....	16.1	15.7	6.4	2.1	1.7	10.5	15.3	14.2	16.8	20.5	29.7
2001.....	14.8	14.2	6.7	2.3	1.5	8.5	12.8	13.8	16.1	23.1	25.7
2000.....	15.3	10.9	5.9	1.8	1.3	9.5	14.5	13.8	19.5	23.4	28.3
1999.....	16.9	15.0	7.7	2.4	2.2	9.6	15.9	15.1	20.0	27.1	29.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Nontransport accidents (W00-X59,Y86)

Hispanic, female

2003.....	5.4	9.4	3.9	1.5	*	1.5	2.4	2.9	3.2	4.7	6.5
2002.....	5.2	8.3	3.7	1.2	1.2	1.6	2.1	1.7	2.6	5.2	6.8
2001.....	5.2	13.0	3.3	1.3	1.3	*	2.1	2.3	3.6	4.7	5.3
2000.....	4.9	8.8	4.4	1.5	1.4	1.9	*	2.2	2.6	5.1	3.6
1999.....	5.0	9.8	5.2	1.2	*	2.0	1.4	2.1	3.1	5.1	6.0

Non-Hispanic 4/, both sexes

2003.....	22.5	22.4	7.1	2.1	2.3	6.3	13.0	13.2	14.3	18.9	23.6
2002.....	21.6	22.3	6.7	2.5	2.6	6.7	11.4	12.5	14.2	18.4	22.7
2001.....	20.2	22.0	7.2	2.7	2.6	6.4	10.7	10.9	11.8	16.9	19.7
2000.....	19.2	20.7	7.9	2.9	2.7	6.4	9.7	10.3	11.8	15.8	18.7
1999.....	19.3	18.5	8.0	2.9	2.9	6.1	9.3	10.1	12.0	15.0	18.7

Non-Hispanic 4/, male

2003.....	27.5	25.2	8.7	2.5	3.2	10.0	20.5	20.4	20.9	26.7	33.1
2002.....	26.5	25.2	8.5	3.3	3.6	10.5	18.5	19.5	21.3	26.6	32.4
2001.....	25.0	24.9	8.7	3.5	3.6	10.1	17.7	17.2	17.8	24.8	28.8
2000.....	23.9	23.2	9.7	3.7	3.8	10.4	15.7	16.7	18.0	24.1	28.0
1999.....	24.0	21.5	9.6	3.7	4.2	10.0	15.1	16.4	18.8	22.7	28.8

Non-Hispanic 4/, female

2003.....	17.8	19.5	5.3	1.6	1.3	2.5	5.3	6.0	7.7	11.3	14.3
2002.....	16.9	19.4	4.9	1.7	1.5	2.7	4.2	5.5	7.2	10.3	13.2
2001.....	15.7	19.1	5.6	1.9	1.5	2.5	3.6	4.6	5.8	9.1	10.7
2000.....	14.7	18.0	6.0	1.9	1.4	2.2	3.6	3.9	5.6	7.7	9.6
1999.....	14.8	15.2	6.3	2.1	1.6	2.1	3.4	3.9	5.3	7.4	8.9

Non-Hispanic White, both sexes

2003.....	24.4	18.4	6.9	1.7	1.8	6.8	14.9	14.8	15.7	20.0	24.6
2002.....	23.1	17.2	6.1	1.9	2.1	6.9	12.8	13.8	15.5	19.1	23.4
2001.....	21.5	17.8	7.1	2.3	2.2	6.5	11.9	11.6	12.5	17.4	19.5
2000.....	20.1	16.6	7.3	2.3	2.3	6.4	10.5	10.9	12.1	16.0	18.4
1999.....	20.0	14.7	7.4	2.2	2.5	6.1	9.6	10.6	11.9	15.2	18.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Nontransport accidents (W00-X59,Y86)											
Non-Hispanic White, male											
2003.....	29.2	20.8	8.5	2.1	2.7	10.8	23.5	22.7	22.8	27.8	34.5
2002.....	27.8	20.4	7.6	2.4	3.0	10.8	20.6	21.2	23.1	27.2	33.4
2001.....	25.9	20.0	8.8	2.8	3.0	10.2	19.6	18.2	18.9	25.6	28.7
2000.....	24.5	19.1	9.1	3.0	3.4	10.4	17.0	17.5	18.6	24.5	27.4
1999.....	24.3	17.1	8.9	2.7	3.6	9.8	15.4	17.1	18.7	22.9	28.0
Non-Hispanic White, female											
2003.....	19.7	16.0	5.1	1.3	1.0	2.7	6.1	6.7	8.6	12.1	14.7
2002.....	18.6	13.9	4.6	1.2	1.2	2.8	4.8	6.2	7.8	10.9	13.5
2001.....	17.2	15.5	5.3	1.8	1.2	2.5	4.1	4.9	6.0	9.2	10.3
2000.....	16.0	14.0	5.4	1.5	1.2	2.3	3.8	4.2	5.6	7.5	9.3
1999.....	16.0	12.3	5.9	1.7	1.4	2.2	3.6	4.0	5.1	7.4	8.6
Non-Hispanic Black, both sexes											
2003.....	17.7	40.3	8.5	3.8	3.7	4.7	7.6	9.7	12.0	18.2	23.5
2002.....	18.3	44.5	9.5	4.9	4.5	6.1	7.6	10.2	12.5	19.4	23.4
2001.....	18.1	41.6	8.7	4.3	4.3	6.1	7.5	9.7	11.3	18.0	25.5
2000.....	18.0	39.5	10.9	5.4	4.1	6.5	7.8	9.8	12.8	18.3	24.9
1999.....	19.4	36.4	11.1	5.8	4.8	6.5	9.2	10.5	14.8	16.9	25.9
Non-Hispanic Black, male											
2003.....	24.1	45.3	10.8	4.4	5.2	7.5	11.6	15.0	18.2	27.3	33.1
2002.....	25.5	46.7	12.5	6.4	6.7	10.4	12.9	16.8	19.2	29.8	33.5
2001.....	25.0	48.2	10.0	6.2	5.9	9.8	12.4	15.4	16.9	26.4	36.9
2000.....	25.5	42.3	12.9	7.2	5.8	10.7	12.8	16.4	19.4	27.5	38.2
1999.....	27.5	42.3	13.7	7.7	6.8	11.2	16.2	17.2	22.8	25.9	41.2
Non-Hispanic Black, female											
2003.....	12.0	35.1	6.2	3.2	2.2	1.8	3.6	4.7	6.5	10.2	15.1
2002.....	11.8	42.3	6.5	3.4	2.4	1.7	2.5	4.2	6.4	10.1	14.6
2001.....	11.8	34.6	7.4	2.3	2.6	2.2	2.8	4.5	6.3	10.5	15.4
2000.....	11.3	36.6	8.8	3.6	2.4	2.2	3.1	3.9	6.9	10.2	13.0
1999.....	12.0	30.3	8.3	3.9	2.6	1.7	2.6	4.5	7.6	9.0	12.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Falls (W00-W19)											
All origins 3/, both sexes											
2003.....	5.9	*	0.3	*	0.1	0.4	0.7	0.7	0.7	1.2	1.7
2002.....	5.6	*	0.2	*	0.1	0.4	0.8	0.7	0.8	1.2	1.8
2001.....	5.3	0.6	0.2	0.2	0.2	0.4	0.9	0.8	0.9	1.2	1.6
2000.....	4.7	*	0.2	*	0.1	0.5	0.7	0.7	0.8	1.0	1.6
1999.....	4.7	*	0.4	0.1	0.1	0.6	0.7	0.8	0.9	1.2	1.7
All origins 3/, male											
2003.....	6.2	*	0.5	*	0.2	0.7	1.1	1.3	1.1	2.0	2.7
2002.....	6.0	*	0.4	*	*	0.6	1.4	1.2	1.4	1.7	2.9
2001.....	5.8	*	0.3	0.2	0.2	0.8	1.5	1.5	1.5	2.1	2.8
2000.....	5.2	*	0.3	*	*	0.8	1.2	1.2	1.5	1.8	2.7
1999.....	5.2	*	0.5	*	0.2	0.9	1.2	1.4	1.5	2.0	2.8
All origins 3/, female											
2003.....	5.6	*	*	*	*	*	0.2	*	0.3	0.3	0.6
2002.....	5.3	*	*	*	*	*	*	*	0.3	0.6	0.7
2001.....	4.8	*	*	*	*	*	*	*	0.3	0.4	0.6
2000.....	4.3	*	*	*	*	*	0.2	*	*	0.3	0.6
1999.....	4.3	*	*	*	*	*	*	*	0.3	0.4	0.6
Hispanic, both sexes											
2003.....	2.1	*	*	*	*	*	0.7	0.9	0.7	1.0	1.8
2002.....	2.1	*	*	*	*	*	1.1	0.9	1.0	1.3	2.0
2001.....	2.0	*	*	*	*	*	0.8	1.0	1.1	1.5	2.3
2000.....	2.0	*	*	*	*	*	0.8	1.1	1.2	1.1	1.5
1999.....	1.9	*	*	*	*	*	1.0	1.1	1.1	1.2	1.7
Hispanic, male											
2003.....	2.7	*	*	*	*	*	1.2	1.6	1.1	1.8	3.5
2002.....	2.8	*	*	*	*	*	1.9	1.5	1.8	1.9	3.7
2001.....	2.8	*	*	*	*	*	1.5	1.8	2.1	2.6	4.2
2000.....	2.7	*	*	*	*	*	1.4	1.9	2.2	1.8	3.0
1999.....	2.7	*	*	*	*	*	1.7	1.9	1.9	2.1	2.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Falls (W00-W19)											
Hispanic, female											
2003.....	1.5	*	*	*	*	*	*	*	*	*	*
2002.....	1.4	*	*	*	*	*	*	*	*	*	*
2001.....	1.3	*	*	*	*	*	*	*	*	*	*
2000.....	1.2	*	*	*	*	*	*	*	*	*	*
1999.....	1.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	6.5	*	0.3	*	0.1	0.5	0.7	0.7	0.7	1.2	1.6
2002.....	6.2	*	0.2	*	0.1	0.4	0.8	0.6	0.8	1.1	1.7
2001.....	5.7	*	0.2	0.1	0.2	0.4	0.9	0.8	0.8	1.2	1.6
2000.....	5.1	*	0.2	*	*	0.5	0.7	0.6	0.7	1.0	1.6
1999.....	5.1	*	0.4	0.1	0.2	0.6	0.6	0.7	0.9	1.2	1.7
Non-Hispanic 4/, male											
2003.....	6.8	*	0.5	*	*	0.7	1.1	1.1	1.1	2.1	2.5
2002.....	6.5	*	0.3	*	*	0.6	1.3	1.2	1.3	1.7	2.8
2001.....	6.2	*	0.3	*	0.2	0.7	1.5	1.4	1.3	2.0	2.6
2000.....	5.5	*	*	*	*	0.8	1.1	1.1	1.3	1.7	2.6
1999.....	5.5	*	0.5	*	*	1.0	1.1	1.3	1.4	1.9	2.7
Non-Hispanic 4/, female											
2003.....	6.2	*	*	*	*	*	0.3	*	0.3	0.4	0.7
2002.....	5.9	*	*	*	*	*	*	*	0.3	0.6	0.7
2001.....	5.3	*	*	*	*	*	*	*	0.3	0.4	0.6
2000.....	4.7	*	*	*	*	*	0.3	*	*	0.3	0.7
1999.....	4.6	*	*	*	*	*	*	*	0.3	0.5	0.6
Non-Hispanic White, both sexes											
2003.....	7.5	*	0.3	*	0.2	0.5	0.8	0.8	0.8	1.2	1.7
2002.....	7.1	*	*	*	*	0.5	0.9	0.8	0.9	1.2	1.8
2001.....	6.6	*	0.2	*	0.2	0.5	1.0	0.9	0.9	1.2	1.5
2000.....	5.8	*	*	*	*	0.6	0.8	0.7	0.8	1.0	1.7
1999.....	5.7	*	0.4	*	0.2	0.7	0.7	0.8	0.8	1.3	1.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental discharge of firearms (W32-W34)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.2	0.3	0.3
2002.....	0.3	*	*	*	0.2	0.5	0.6	0.4	0.4	0.3	0.3
2001.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.2	0.3	0.3
2000.....	0.3	*	*	*	0.3	0.6	0.5	0.4	0.3	0.3	0.4
1999.....	0.3	*	*	*	0.3	0.7	0.7	0.4	0.3	0.3	0.3

Non-Hispanic 4/, male

2003.....	0.5	*	*	*	0.3	0.8	0.9	0.8	0.4	0.5	0.5
2002.....	0.5	*	*	*	0.3	1.0	1.0	0.8	0.6	0.4	0.6
2001.....	0.5	*	*	*	0.4	1.0	0.9	0.7	0.4	0.5	0.6
2000.....	0.5	*	*	*	0.5	1.0	0.9	0.7	0.5	0.6	0.6
1999.....	0.5	*	*	*	0.5	1.2	1.3	0.7	0.6	0.5	0.5

Non-Hispanic 4/, female

2003.....	0.1	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*	*

Non-Hispanic White, both sexes

2003.....	0.3	*	*	*	0.2	0.4	0.5	0.3	0.2	0.3	0.3
2002.....	0.3	*	*	*	0.2	0.4	0.5	0.3	0.3	0.2	0.4
2001.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.4	0.4
2000.....	0.3	*	*	*	0.3	0.5	0.4	0.4	0.3	0.3	0.4
1999.....	0.3	*	*	*	0.3	0.6	0.6	0.4	0.3	0.3	0.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Accidental drowning and submersion (W65-W74)											
All origins 3/, both sexes											
2003.....	1.1	1.4	2.9	0.6	0.7	1.4	1.4	1.0	0.8	1.0	1.1
2002.....	1.2	1.6	2.9	0.8	0.8	1.6	1.5	1.2	1.0	1.2	1.1
2001.....	1.2	1.7	3.0	0.8	0.8	1.6	1.4	1.1	0.8	1.0	1.1
2000.....	1.2	2.0	3.2	1.0	0.8	1.8	1.5	1.1	1.0	1.1	1.0
1999.....	1.3	1.8	3.2	0.9	0.9	1.8	1.5	1.1	1.1	1.1	1.0
All origins 3/, male											
2003.....	1.8	1.7	3.7	0.9	1.0	2.5	2.5	1.8	1.4	1.7	1.8
2002.....	1.9	1.9	3.9	1.2	1.1	2.8	2.8	2.1	1.7	2.0	1.9
2001.....	1.8	1.5	3.7	1.2	1.1	2.8	2.5	1.9	1.4	1.6	1.7
2000.....	2.0	2.2	4.0	1.3	1.2	3.3	2.5	1.9	1.7	1.9	1.7
1999.....	2.0	2.0	4.0	1.3	1.3	3.2	2.8	2.0	2.0	1.9	1.7
All origins 3/, female											
2003.....	0.5	1.2	2.0	0.4	0.3	*	0.2	*	*	0.3	0.3
2002.....	0.5	1.2	1.9	0.3	0.4	0.3	*	0.3	0.3	0.4	0.4
2001.....	0.5	1.9	2.2	0.5	0.5	0.3	0.3	0.3	0.2	0.4	0.4
2000.....	0.5	1.7	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3	0.3
1999.....	0.5	1.6	2.4	0.5	0.4	0.3	0.3	*	0.3	0.3	0.3
Hispanic, both sexes											
2003.....	1.1	*	2.5	0.6	0.6	1.4	1.8	1.4	0.9	0.9	0.8
2002.....	1.2	*	2.1	*	*	2.2	2.1	1.2	1.0	0.9	1.0
2001.....	1.1	*	2.6	0.7	*	1.3	1.7	1.1	1.1	1.2	0.8
2000.....	1.2	*	2.5	0.8	*	2.3	2.2	1.0	0.9	0.9	*
1999.....	1.4	*	2.7	0.7	0.7	2.2	2.3	1.2	1.3	1.0	*
Hispanic, male											
2003.....	1.9	*	4.0	*	*	2.7	3.1	2.4	1.5	1.5	1.5
2002.....	2.0	*	3.0	*	*	3.8	3.7	2.1	1.8	1.4	1.8
2001.....	1.8	*	3.5	1.1	*	2.5	2.9	1.7	1.8	1.9	*
2000.....	1.9	*	2.9	*	*	3.8	3.7	1.8	1.7	1.4	*
1999.....	2.3	*	2.9	*	*	4.1	4.1	2.1	2.3	1.7	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental drowning and submersion (W65-W74)

Hispanic, female

2003.....	0.3	*	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	1.7	*	*	*	*	*	*	*	*
2000.....	0.5	*	2.2	*	*	*	*	*	*	*	*
1999.....	0.5	*	2.5	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	1.1	1.5	3.0	0.6	0.7	1.3	1.3	0.9	0.8	1.0	1.1
2002.....	1.2	1.4	3.1	0.9	0.8	1.5	1.4	1.2	1.0	1.2	1.2
2001.....	1.1	1.6	3.1	0.8	0.9	1.6	1.3	1.1	0.7	1.0	1.1
2000.....	1.2	2.1	3.4	1.0	0.9	1.7	1.3	1.1	1.0	1.1	1.1
1999.....	1.2	1.7	3.3	1.0	0.9	1.7	1.4	1.1	1.1	1.1	1.0

Non-Hispanic 4/, male

2003.....	1.8	1.7	3.7	0.9	1.0	2.4	2.3	1.6	1.4	1.8	1.8
2002.....	1.9	1.6	4.1	1.3	1.2	2.5	2.6	2.0	1.7	2.1	1.9
2001.....	1.8	1.3	3.8	1.1	1.2	2.9	2.4	2.0	1.3	1.6	1.8
2000.....	2.0	2.4	4.3	1.4	1.4	3.1	2.2	1.9	1.8	1.9	1.8
1999.....	2.0	1.7	4.2	1.4	1.3	3.0	2.4	1.9	1.9	1.9	1.8

Non-Hispanic 4/, female

2003.....	0.5	*	2.3	0.4	0.3	*	0.3	*	*	0.3	0.4
2002.....	0.5	*	2.1	0.4	0.5	0.3	*	0.3	0.3	0.4	0.4
2001.....	0.5	1.9	2.4	0.5	0.5	0.3	0.3	0.3	*	0.3	0.4
2000.....	0.5	1.9	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3	0.3
1999.....	0.5	1.7	2.4	0.6	0.4	0.3	0.3	*	0.3	0.3	0.3

Non-Hispanic White, both sexes

2003.....	1.1	1.4	3.3	0.5	0.4	1.2	1.2	0.8	0.7	1.0	1.1
2002.....	1.1	1.4	3.3	0.6	0.5	1.2	1.3	1.0	1.0	1.2	1.2
2001.....	1.1	1.5	3.5	0.6	0.5	1.3	1.3	1.0	0.7	0.9	1.0
2000.....	1.1	1.7	3.7	0.8	0.6	1.4	1.1	1.0	1.0	1.1	1.0
1999.....	1.1	1.3	3.6	0.6	0.5	1.4	1.2	0.9	1.0	1.1	1.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Accidental exposure to smoke, fire and flames (X00-X09)											
Hispanic, female											
2003.....	0.4	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	1.3	0.8	1.6	0.7	0.4	0.4	0.7	0.7	0.7	0.8	1.1
2002.....	1.2	1.0	1.5	0.8	0.5	0.4	0.6	0.7	0.8	0.7	1.0
2001.....	1.3	1.3	1.7	0.9	0.5	0.4	0.7	0.7	0.7	0.9	1.1
2000.....	1.3	1.0	2.2	1.0	0.5	0.4	0.7	0.7	0.7	0.9	1.0
1999.....	1.3	1.2	2.1	0.9	0.5	0.4	0.7	0.8	0.7	0.9	1.0
Non-Hispanic 4/, male											
2003.....	1.5	*	1.9	0.8	0.4	0.4	0.8	0.8	0.9	1.2	1.3
2002.....	1.5	*	1.7	1.0	0.6	0.6	0.7	0.9	1.0	0.9	1.4
2001.....	1.6	1.5	2.0	1.1	0.5	0.5	1.0	0.8	0.9	1.3	1.6
2000.....	1.6	*	2.6	1.2	0.4	0.4	0.9	0.9	0.9	1.3	1.2
1999.....	1.6	*	2.3	1.0	0.6	0.6	0.8	1.1	1.1	1.3	1.5
Non-Hispanic 4/, female											
2003.....	1.0	1.3	1.3	0.7	0.4	0.3	0.5	0.6	0.5	0.5	0.8
2002.....	0.9	*	1.2	0.7	0.4	0.3	0.4	0.4	0.6	0.5	0.6
2001.....	1.0	*	1.3	0.6	0.4	0.4	0.4	0.5	0.5	0.6	0.7
2000.....	1.0	*	1.7	0.8	0.5	0.3	0.5	0.5	0.5	0.5	0.7
1999.....	1.0	*	1.8	0.8	0.4	0.3	0.5	0.6	0.4	0.6	0.6
Non-Hispanic White, both sexes											
2003.....	1.2	*	1.3	0.5	0.3	0.4	0.7	0.7	0.7	0.9	1.0
2002.....	1.0	*	1.0	0.5	0.4	0.4	0.5	0.6	0.8	0.7	1.0
2001.....	1.1	1.1	1.3	0.7	0.4	0.4	0.7	0.5	0.7	0.8	1.0
2000.....	1.1	*	1.5	0.7	0.4	0.4	0.6	0.6	0.6	0.8	0.9
1999.....	1.1	0.9	1.5	0.6	0.4	0.4	0.6	0.8	0.7	0.8	1.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Accidental exposure to smoke, fire and flames (X00-X09)											
Non-Hispanic White, male											
2003.....	1.4	*	1.6	0.5	*	0.4	1.0	0.8	0.9	1.2	1.4
2002.....	1.3	*	1.1	0.6	0.5	0.5	0.7	0.8	1.0	0.8	1.4
2001.....	1.4	*	1.7	0.9	0.4	0.5	1.0	0.8	0.9	1.1	1.5
2000.....	1.4	*	1.8	0.8	0.3	0.4	0.9	0.7	0.8	1.1	1.1
1999.....	1.4	*	1.7	0.6	0.5	0.6	0.8	1.1	1.1	1.1	1.4
Non-Hispanic White, female											
2003.....	1.0	*	1.0	0.5	*	*	0.5	0.6	0.6	0.5	0.7
2002.....	0.8	*	0.8	0.4	*	*	0.3	*	0.6	0.5	0.6
2001.....	0.9	*	0.9	0.5	0.4	0.3	0.3	*	0.5	0.5	0.6
2000.....	0.9	*	1.1	0.6	0.4	*	0.4	0.4	0.4	0.4	0.6
1999.....	0.8	*	1.3	0.6	0.3	*	0.4	0.5	0.3	0.5	0.5
Non-Hispanic Black, both sexes											
2003.....	2.1	*	3.1	2.0	1.0	*	*	1.0	0.8	1.0	1.4
2002.....	2.1	*	3.7	2.1	0.9	0.7	0.9	1.2	1.1	1.0	1.1
2001.....	2.3	*	3.4	1.8	0.8	*	1.0	1.4	1.0	1.6	2.1
2000.....	2.6	*	5.1	2.2	0.8	*	1.0	1.2	1.3	1.6	2.0
1999.....	2.6	*	4.7	2.2	0.9	*	0.9	1.1	0.9	1.9	1.7
Non-Hispanic Black, male											
2003.....	2.5	*	3.4	2.0	*	*	*	*	*	*	1.6
2002.....	2.6	*	4.3	2.3	*	*	*	*	*	1.6	1.8
2001.....	3.0	*	3.7	2.3	*	*	*	*	*	2.6	3.0
2000.....	3.2	*	5.8	2.7	*	*	*	*	1.6	2.3	2.6
1999.....	3.1	*	5.3	2.5	*	*	*	*	*	2.8	2.5
Non-Hispanic Black, female											
2003.....	1.8	*	2.8	1.9	*	*	*	*	*	*	*
2002.....	1.6	*	3.0	1.8	*	*	*	*	*	*	*
2001.....	1.8	*	3.1	1.3	*	*	*	1.5	*	*	*
2000.....	2.0	*	4.3	1.7	*	*	*	*	*	*	1.4
1999.....	2.1	*	4.1	1.9	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental poisoning and exposure to noxious substances (X40-X49)

All origins 3/, both sexes

2003.....	6.7	0.5	0.3	*	0.2	2.5	7.1	7.9	9.3	12.5	15.5
2002.....	6.1	0.6	0.2	*	0.1	2.4	5.9	6.9	8.6	11.9	14.8
2001.....	4.9	*	0.2	*	0.2	2.0	4.9	5.7	6.9	10.4	12.0
2000.....	4.5	*	0.2	*	0.1	1.7	4.3	5.1	6.8	9.5	11.2
1999.....	4.4	*	0.2	*	0.1	1.3	3.8	4.9	6.8	9.1	11.1

All origins 3/, male

2003.....	9.2	*	0.3	*	0.2	3.8	10.9	11.7	13.1	16.8	20.7
2002.....	8.5	1.0	*	*	*	3.5	9.1	10.4	12.5	16.5	20.3
2001.....	7.1	*	0.3	*	*	3.0	7.6	8.7	9.9	14.6	16.8
2000.....	6.6	*	0.3	*	*	2.6	6.7	7.9	10.1	13.7	16.4
1999.....	6.5	*	*	*	*	1.9	5.9	7.7	10.3	13.3	16.6

All origins 3/, female

2003.....	4.3	*	0.3	*	*	1.2	3.1	4.0	5.4	8.1	10.3
2002.....	3.7	*	*	*	*	1.2	2.6	3.3	4.7	7.2	9.4
2001.....	2.9	*	*	*	*	1.0	1.9	2.6	3.9	6.1	7.3
2000.....	2.5	*	*	*	*	0.8	1.7	2.1	3.5	5.3	6.0
1999.....	2.3	*	*	*	*	0.7	1.6	2.0	3.2	4.9	5.7

Hispanic, both sexes

2003.....	4.5	*	*	*	*	2.1	4.0	5.2	6.3	9.7	11.1
2002.....	4.3	*	*	*	*	1.8	3.8	4.3	5.7	8.3	12.2
2001.....	3.6	*	*	*	*	1.5	2.7	4.3	5.5	8.4	9.1
2000.....	3.7	*	*	*	*	1.2	2.8	3.9	6.8	9.5	9.8
1999.....	4.4	*	*	*	*	1.3	3.2	4.8	7.5	11.4	12.4

Hispanic, male

2003.....	6.9	*	*	*	*	3.3	6.0	7.7	9.7	14.9	16.3
2002.....	6.6	*	*	*	*	2.6	5.9	6.8	8.9	12.4	19.0
2001.....	5.6	*	*	*	*	2.1	3.9	6.8	8.1	13.1	14.2
2000.....	6.0	*	*	*	*	1.8	4.4	6.1	11.0	14.5	17.1
1999.....	7.3	*	*	*	*	1.7	5.3	7.6	12.2	18.4	20.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/ years	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Accidental poisoning and exposure to noxious substances (X40-X49)											
Hispanic, female											
2003.....	1.9	*	*	*	*	*	1.6	2.1	2.4	3.8	5.4
2002.....	1.7	*	*	*	*	*	1.3	*	2.0	3.7	5.0
2001.....	1.5	*	*	*	*	*	1.3	1.4	2.5	3.2	3.7
2000.....	1.3	*	*	*	*	*	*	1.3	1.9	3.9	2.2
1999.....	1.4	*	*	*	*	*	*	1.5	2.0	3.8	4.0
Non-Hispanic 4/, both sexes											
2003.....	7.0	*	0.3	*	0.2	2.6	7.8	8.5	9.8	12.9	16.0
2002.....	6.3	0.6	0.2	*	0.1	2.5	6.3	7.5	9.2	12.4	15.1
2001.....	5.1	*	0.2	*	0.1	2.1	5.3	6.0	7.1	10.6	12.2
2000.....	4.6	*	0.2	*	0.1	1.8	4.6	5.3	6.8	9.4	11.2
1999.....	4.3	*	0.2	*	0.1	1.3	3.9	4.8	6.6	8.7	10.8
Non-Hispanic 4/, male											
2003.....	9.5	*	0.3	*	0.3	3.9	12.1	12.7	13.8	17.0	21.2
2002.....	8.8	*	*	*	*	3.7	9.8	11.3	13.1	17.1	20.3
2001.....	7.2	*	*	*	*	3.1	8.5	9.1	10.1	14.7	17.0
2000.....	6.6	*	*	*	*	2.8	7.2	8.3	9.8	13.5	16.2
1999.....	6.3	*	*	*	*	1.9	6.0	7.6	9.9	12.5	15.9
Non-Hispanic 4/, female											
2003.....	4.6	*	*	*	*	1.3	3.4	4.4	5.9	8.8	10.9
2002.....	4.0	*	*	*	*	1.3	2.8	3.8	5.2	7.7	9.9
2001.....	3.1	*	*	*	*	1.0	2.1	2.9	4.1	6.5	7.6
2000.....	2.7	*	*	*	*	0.8	1.9	2.2	3.8	5.5	6.3
1999.....	2.4	*	*	*	*	0.6	1.8	2.1	3.4	5.0	5.8
Non-Hispanic White, both sexes											
2003.....	7.4	*	*	*	0.2	3.2	9.7	10.1	11.1	13.8	16.9
2002.....	6.6	*	*	*	0.2	3.0	7.6	8.8	10.3	13.1	15.7
2001.....	5.2	*	*	*	*	2.4	6.2	6.9	7.7	11.0	12.3
2000.....	4.6	*	*	*	*	2.2	5.4	5.9	7.2	9.7	11.0
1999.....	4.2	*	*	*	*	1.5	4.3	5.3	6.7	8.8	10.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic White, male

2003.....	10.1	*	*	*	*	4.8	14.9	15.1	15.4	18.0	22.4
2002.....	9.1	*	*	*	*	4.5	11.8	13.0	14.7	18.1	21.0
2001.....	7.3	*	*	*	*	3.8	9.9	10.3	11.2	15.3	17.2
2000.....	6.6	*	*	*	*	3.3	8.6	9.2	10.4	14.0	15.8
1999.....	6.1	*	*	*	*	2.2	6.6	8.3	10.0	12.6	15.3

Non-Hispanic White, female

2003.....	4.9	*	*	*	*	1.5	4.3	5.1	6.6	9.5	11.5
2002.....	4.3	*	*	*	*	1.5	3.4	4.5	5.8	8.2	10.4
2001.....	3.1	*	*	*	*	1.0	2.5	3.4	4.2	6.7	7.4
2000.....	2.7	*	*	*	*	1.0	2.0	2.5	3.9	5.4	6.2
1999.....	2.4	*	*	*	*	0.7	2.0	2.3	3.3	4.9	5.4

Non-Hispanic Black, both sexes

2003.....	6.4	*	*	*	*	*	2.3	4.1	7.5	12.1	15.7
2002.....	6.4	*	*	*	*	*	2.3	4.5	7.0	12.1	16.1
2001.....	6.0	*	*	*	*	0.9	2.4	4.1	6.4	11.2	15.6
2000.....	5.7	*	*	*	*	*	2.2	4.3	7.0	10.9	15.6
1999.....	5.8	*	*	*	*	*	2.9	4.3	8.2	10.4	16.3

Non-Hispanic Black, male

2003.....	9.1	*	*	*	*	*	3.7	5.8	10.7	17.1	21.2
2002.....	9.3	*	*	*	*	*	3.7	7.2	10.1	17.3	22.2
2001.....	8.6	*	*	*	*	*	3.9	6.8	8.4	15.5	20.9
2000.....	8.4	*	*	*	*	*	2.8	7.1	9.8	15.0	23.2
1999.....	8.8	*	*	*	*	*	4.8	6.9	11.7	14.9	24.4

Non-Hispanic Black, female

2003.....	4.1	*	*	*	*	*	*	2.5	4.7	7.7	10.9
2002.....	3.8	*	*	*	*	*	*	2.1	4.2	7.5	10.7
2001.....	3.6	*	*	*	*	*	*	1.7	4.7	7.4	10.9
2000.....	3.2	*	*	*	*	*	1.5	1.8	4.4	7.2	8.8
1999.....	3.1	*	*	*	*	*	*	2.0	5.0	6.4	9.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All origins 3/, both sexes

2003.....	5.9	16.9	1.8	0.5	0.6	1.1	2.0	2.0	2.0	2.6	3.4
2002.....	6.0	16.8	1.6	0.6	0.8	1.3	1.8	2.0	2.1	2.6	3.3
2001.....	6.2	16.6	1.8	0.6	0.7	1.2	2.0	1.8	2.2	2.9	3.2
2000.....	6.2	15.0	1.7	0.6	0.7	1.4	2.0	2.1	2.3	3.0	3.4
1999.....	6.6	13.8	1.9	0.7	0.9	1.4	2.0	2.2	2.3	2.7	3.7

All origins 3/, male

2003.....	6.9	19.5	2.2	0.6	0.9	1.8	3.3	3.3	3.4	4.0	5.5
2002.....	6.9	18.8	2.0	0.7	1.1	2.1	3.1	3.2	3.6	4.3	5.4
2001.....	7.2	18.9	2.1	0.7	1.0	1.9	3.4	3.1	3.7	4.8	5.3
2000.....	7.2	16.8	2.1	0.7	1.1	2.1	3.5	3.7	3.9	5.1	5.6
1999.....	7.6	16.2	2.3	0.9	1.2	2.3	3.4	3.7	3.7	4.5	6.1

All origins 3/, female

2003.....	4.9	14.1	1.3	0.3	0.2	0.3	0.7	0.7	0.6	1.1	1.4
2002.....	5.0	14.6	1.2	0.4	0.4	0.5	0.5	0.7	0.6	1.0	1.3
2001.....	5.3	14.2	1.5	0.4	0.3	0.6	0.6	0.6	0.6	1.1	1.2
2000.....	5.2	13.2	1.4	0.4	0.3	0.5	0.4	0.6	0.7	0.8	1.3
1999.....	5.6	11.3	1.5	0.5	0.4	0.5	0.5	0.6	0.8	0.9	1.4

Hispanic, both sexes

2003.....	2.5	8.2	1.6	0.5	*	1.2	2.0	2.1	2.0	1.9	2.5
2002.....	2.5	8.2	1.3	*	*	1.2	1.7	1.8	2.0	2.2	2.8
2001.....	2.7	9.5	1.4	*	*	1.1	1.8	1.5	2.0	2.6	2.8
2000.....	2.7	7.5	1.4	*	*	1.1	2.2	1.9	2.5	2.7	3.2
1999.....	2.6	8.6	1.9	*	*	1.1	1.9	1.7	1.5	2.5	2.4

Hispanic, male

2003.....	3.6	9.1	1.9	*	*	2.0	3.6	3.5	3.3	3.2	4.2
2002.....	3.6	10.1	1.8	*	*	2.0	2.9	3.1	3.6	3.7	4.8
2001.....	3.7	9.6	1.8	*	*	1.9	3.2	2.6	3.4	4.5	4.7
2000.....	3.8	7.9	1.5	*	*	1.8	3.9	3.2	4.3	5.0	5.6
1999.....	3.6	9.0	2.5	*	*	1.7	3.4	2.9	2.4	4.2	3.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)											
Hispanic, female											
2003.....	1.3	7.3	1.3	*	*	*	*	*	*	*	*
2002.....	1.2	6.1	*	*	*	*	*	*	*	*	*
2001.....	1.6	9.5	*	*	*	*	*	*	*	*	*
2000.....	1.4	7.2	*	*	*	*	*	*	*	*	*
1999.....	1.6	8.1	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	6.4	19.2	1.8	0.5	0.6	1.1	2.0	2.0	2.0	2.7	3.5
2002.....	6.5	18.9	1.6	0.6	0.8	1.4	1.8	2.0	2.1	2.7	3.4
2001.....	6.7	18.3	1.9	0.6	0.7	1.3	2.0	1.9	2.2	2.9	3.3
2000.....	6.7	16.9	1.8	0.6	0.8	1.4	1.9	2.2	2.3	3.0	3.4
1999.....	7.1	15.0	1.9	0.7	0.9	1.5	2.0	2.2	2.4	2.7	3.9
Non-Hispanic 4/, male											
2003.....	7.5	22.2	2.3	0.6	1.0	1.8	3.2	3.3	3.4	4.2	5.6
2002.....	7.5	21.0	2.0	0.7	1.2	2.1	3.1	3.3	3.6	4.3	5.4
2001.....	7.7	21.1	2.1	0.8	1.1	1.9	3.4	3.2	3.8	4.8	5.3
2000.....	7.7	19.0	2.2	0.8	1.3	2.2	3.4	3.8	3.8	5.1	5.6
1999.....	8.1	17.9	2.3	0.9	1.4	2.4	3.5	3.8	4.0	4.5	6.3
Non-Hispanic 4/, female											
2003.....	5.4	16.0	1.3	0.3	0.3	0.3	0.8	0.7	0.7	1.2	1.4
2002.....	5.6	16.8	1.3	0.5	0.4	0.5	0.5	0.8	0.7	1.1	1.4
2001.....	5.8	15.4	1.6	0.4	0.3	0.6	0.6	0.6	0.6	1.2	1.3
2000.....	5.7	14.7	1.4	0.4	0.3	0.6	0.5	0.6	0.8	0.9	1.3
1999.....	6.2	12.0	1.5	0.5	0.5	0.5	0.5	0.7	0.8	1.0	1.5
Non-Hispanic White, both sexes											
2003.....	6.9	15.5	1.7	0.5	0.7	1.1	2.1	2.1	2.2	2.7	3.6
2002.....	7.0	14.5	1.5	0.5	0.8	1.5	2.0	2.3	2.3	2.7	3.5
2001.....	7.2	14.5	1.7	0.7	0.7	1.4	2.2	2.0	2.2	3.1	3.2
2000.....	7.1	13.7	1.7	0.6	0.8	1.4	2.0	2.3	2.4	3.0	3.4
1999.....	7.6	12.1	1.7	0.8	1.0	1.6	2.2	2.3	2.4	2.9	3.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All origins 3/, both sexes

2003.....	10.8	*	1.2	7.3	12.1	11.9	13.5	13.8	15.9
2002.....	11.0	*	1.2	7.4	12.4	12.8	12.5	14.3	16.1
2001 5/.....	10.8	*	1.3	7.9	12.0	12.6	13.0	14.3	15.2
2000.....	10.4	*	1.5	8.0	12.5	12.1	12.0	13.7	15.4
1999.....	10.5	*	1.2	8.0	12.3	12.3	13.1	14.4	14.3

All origins 3/, male

2003.....	17.6	*	1.7	11.6	20.2	19.5	21.6	21.9	24.5
2002.....	17.9	*	1.8	12.2	20.8	20.8	20.2	22.7	24.7
2001 5/.....	17.6	*	1.9	12.9	20.5	21.0	21.0	22.7	23.6
2000.....	17.1	*	2.3	13.0	21.4	20.0	19.2	21.7	23.9
1999.....	17.1	*	1.9	13.0	20.8	20.3	21.1	22.7	22.1

All origins 3/, female

2003.....	4.3	*	0.5	2.7	3.4	4.0	5.2	5.7	7.4
2002.....	4.3	*	0.6	2.4	3.5	4.5	4.7	6.0	7.7
2001 5/.....	4.1	*	0.6	2.7	3.1	4.0	4.8	5.9	6.9
2000.....	4.0	*	0.6	2.7	3.2	4.0	4.6	5.8	7.0
1999.....	4.0	*	0.5	2.7	3.4	4.1	5.1	6.1	6.7

Hispanic, both sexes

2003.....	5.0	*	1.2	5.9	8.0	6.6	6.9	6.4	7.3
2002.....	5.0	*	1.0	5.7	7.6	6.6	6.6	7.0	7.0
2001 5/.....	5.0	*	1.1	5.2	7.0	7.3	7.0	7.5	7.0
2000.....	5.1	*	0.9	5.6	7.9	6.7	6.2	6.9	8.3
1999.....	5.0	*	1.0	5.7	6.7	6.7	6.7	8.4	6.4

Hispanic, male

2003.....	8.3	*	1.7	9.2	12.8	10.5	11.1	10.3	11.8
2002.....	8.3	*	1.1	9.1	11.9	10.8	10.6	11.3	11.1
2001 5/.....	8.3	*	1.5	7.8	11.0	11.6	11.1	12.8	11.8
2000.....	8.4	*	1.2	8.5	13.0	11.1	10.0	10.7	13.7
1999.....	8.2	*	1.4	8.9	10.8	10.6	10.9	13.2	10.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Hispanic, female

2003.....	1.5	*	*	2.4	2.0	1.8	2.1	2.0	2.3
2002.....	1.6	*	*	2.0	2.3	1.5	1.9	2.3	2.6
2001 5/.....	1.5	*	*	2.5	2.2	2.2	2.3	1.7	1.9
2000.....	1.5	*	*	2.4	1.6	1.5	1.8	2.7	2.6
1999.....	1.6	*	*	2.0	1.6	2.2	1.8	3.1	2.0

Non-Hispanic 4/, both sexes

2003.....	11.7	*	1.1	7.5	12.9	13.1	14.8	15.0	17.0
2002.....	11.8	*	1.3	7.7	13.3	14.2	13.7	15.5	17.3
2001 5/.....	11.6	*	1.3	8.4	13.0	13.8	14.1	15.3	16.1
2000.....	11.1	*	1.6	8.4	13.5	13.1	12.9	14.6	16.1
1999.....	11.2	*	1.2	8.4	13.5	13.3	14.2	15.1	15.1

Non-Hispanic 4/, male

2003.....	19.1	*	1.7	12.1	21.9	21.8	23.9	23.9	26.2
2002.....	19.4	*	1.9	12.7	22.8	23.4	22.2	24.6	26.5
2001 5/.....	19.0	*	2.0	13.8	22.7	23.3	22.9	24.2	24.9
2000.....	18.3	*	2.4	13.8	23.3	21.9	20.9	23.2	24.9
1999.....	18.4	*	1.9	13.8	23.1	22.3	23.0	24.0	23.2

Non-Hispanic 4/, female

2003.....	4.6	*	0.5	2.7	3.6	4.4	5.8	6.3	8.0
2002.....	4.6	*	0.6	2.4	3.7	5.1	5.2	6.5	8.3
2001 5/.....	4.5	*	0.6	2.7	3.3	4.4	5.3	6.5	7.5
2000.....	4.3	*	0.6	2.8	3.5	4.4	5.0	6.1	7.5
1999.....	4.3	*	0.5	2.8	3.7	4.4	5.6	6.4	7.2

Non-Hispanic White, both sexes

2003.....	13.3	*	1.1	8.3	13.6	14.3	16.6	17.0	19.4
2002.....	13.4	*	1.3	8.6	14.5	15.5	15.3	17.5	19.7
2001 5/.....	13.0	*	1.3	9.3	13.9	15.1	15.7	17.2	18.2
2000.....	12.5	*	1.7	9.0	14.3	14.0	14.4	16.4	18.0
1999.....	12.4	*	1.3	9.0	14.2	14.2	15.6	16.7	17.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Non-Hispanic White, male

2003.....	21.6	*	1.7	13.3	23.3	23.4	26.4	26.6	29.7
2002.....	21.9	*	2.0	14.2	24.7	24.7	24.4	27.3	30.0
2001 5/.....	21.4	*	2.1	15.4	24.1	25.3	25.2	26.8	27.9
2000.....	20.4	*	2.6	14.8	24.8	23.2	22.9	25.7	27.6
1999.....	20.3	*	2.1	14.7	24.3	23.5	25.0	25.9	26.0

Non-Hispanic White, female

2003.....	5.3	*	0.5	3.0	3.6	5.0	6.6	7.3	9.2
2002.....	5.3	*	0.6	2.7	4.1	6.1	6.0	7.6	9.5
2001 5/.....	5.0	*	0.6	3.0	3.6	4.8	6.1	7.6	8.5
2000.....	4.9	*	0.7	3.0	3.7	4.7	5.7	7.1	8.5
1999.....	4.9	*	0.5	3.0	3.9	4.8	6.2	7.5	8.1

Non-Hispanic Black, both sexes

2003.....	5.3	*	1.2	3.9	10.8	9.8	9.4	7.7	7.1
2002.....	5.2	*	1.0	4.0	9.4	10.8	8.5	8.3	7.2
2001 5/.....	5.3	*	1.2	4.5	10.5	9.7	8.7	8.1	6.9
2000.....	5.5	*	1.2	5.7	11.4	10.2	7.9	7.2	8.2
1999.....	5.5	*	0.9	5.9	11.1	10.0	9.0	8.6	7.0

Non-Hispanic Black, male

2003.....	9.0	*	1.9	6.7	18.5	18.1	16.2	13.4	11.6
2002.....	9.3	*	1.6	6.9	16.6	20.5	15.6	14.7	11.7
2001 5/.....	9.3	*	1.7	7.5	20.0	17.7	15.5	14.2	10.8
2000.....	9.6	*	2.1	9.9	20.1	18.2	14.1	12.9	13.7
1999.....	9.9	*	1.4	10.1	20.3	18.7	16.3	16.1	11.1

Non-Hispanic Black, female

2003.....	1.8	*	*	*	3.2	2.2	3.2	2.6	3.1
2002.....	1.6	*	*	*	2.4	2.1	2.1	2.6	3.3
2001 5/.....	1.7	*	*	*	*	2.4	2.5	2.6	3.4
2000.....	1.7	*	*	1.5	3.1	3.0	2.3	2.2	3.3
1999.....	1.6	*	*	1.6	2.3	2.1	2.6	1.9	3.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All origins 3/, both sexes

2003.....	5.8	*	0.3	3.6	6.5	5.8	6.1	6.0	7.2
2002.....	5.9	*	0.4	3.6	6.7	6.4	5.6	6.6	7.4
2001.....	5.9	*	0.4	4.1	6.6	6.7	6.3	6.4	7.0
2000.....	5.9	*	0.5	4.4	7.2	6.3	6.0	6.7	7.4
1999.....	5.9	*	0.5	4.9	7.2	6.6	6.5	7.1	6.7

All origins 3/, male

2003.....	10.4	*	0.6	6.3	11.4	10.1	10.6	10.2	12.1
2002.....	10.6	*	0.6	6.4	11.9	11.1	9.6	11.3	12.6
2001.....	10.6	*	0.6	7.1	11.8	11.7	10.8	10.8	12.0
2000.....	10.5	*	0.9	7.6	12.9	11.1	10.1	11.1	12.6
1999.....	10.6	*	0.8	8.4	12.8	11.5	10.9	11.8	11.2

All origins 3/, female

2003.....	1.4	*	*	0.8	1.2	1.3	1.5	1.8	2.3
2002.....	1.4	*	*	0.7	1.2	1.6	1.7	2.0	2.3
2001.....	1.5	*	0.2	1.0	1.0	1.5	1.7	2.1	2.2
2000.....	1.5	*	0.2	1.1	1.3	1.5	1.9	2.3	2.4
1999.....	1.5	*	0.2	1.1	1.4	1.6	1.9	2.3	2.2

Hispanic, both sexes

2003.....	2.1	*	*	2.6	3.9	3.2	2.4	2.4	2.5
2002.....	2.2	*	*	2.8	3.8	3.0	2.4	2.5	2.5
2001.....	2.2	*	*	2.1	3.6	3.6	2.9	2.5	2.4
2000.....	2.3	*	*	2.8	4.0	3.1	2.5	2.3	3.4
1999.....	2.3	*	*	3.1	3.8	3.2	2.6	3.3	2.6

Hispanic, male

2003.....	3.8	*	*	4.3	6.6	5.4	4.2	4.0	4.6
2002.....	3.8	*	*	4.9	6.4	4.9	4.3	4.3	4.1
2001.....	3.8	*	*	3.5	5.9	6.0	4.9	4.5	4.4
2000.....	4.1	*	*	4.6	6.8	5.3	4.1	3.7	6.0
1999.....	4.0	*	*	5.0	6.5	5.2	4.4	5.3	4.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Hispanic, female

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	6.4	*	0.4	3.8	7.0	6.5	6.9	6.6	7.8
2002.....	6.5	*	0.4	3.8	7.3	7.2	6.3	7.3	8.1
2001.....	6.5	*	0.5	4.5	7.2	7.4	6.9	7.0	7.6
2000.....	6.4	*	0.6	4.7	7.9	7.0	6.6	7.3	7.9
1999.....	6.4	*	0.5	5.2	7.9	7.3	7.1	7.5	7.1

Non-Hispanic 4/, male

2003.....	11.5	*	0.6	6.6	12.6	11.4	12.0	11.3	13.1
2002.....	11.7	*	0.7	6.6	13.1	12.7	10.7	12.4	13.7
2001.....	11.6	*	0.7	7.8	13.2	13.1	12.0	11.8	12.9
2000.....	11.4	*	0.9	8.2	14.3	12.3	11.2	12.1	13.3
1999.....	11.5	*	0.8	9.0	14.3	12.8	12.1	12.7	12.0

Non-Hispanic 4/, female

2003.....	1.6	*	*	0.8	1.3	1.5	1.8	2.0	2.6
2002.....	1.6	*	*	0.8	1.3	1.8	1.9	2.2	2.5
2001.....	1.6	*	0.2	1.0	1.1	1.7	1.9	2.3	2.4
2000.....	1.6	*	0.2	1.1	1.4	1.6	2.1	2.6	2.6
1999.....	1.6	*	*	1.1	1.5	1.7	2.1	2.5	2.3

Non-Hispanic White, both sexes

2003.....	7.4	*	0.4	4.3	7.4	7.0	7.7	7.6	9.1
2002.....	7.5	*	0.5	4.1	7.8	7.8	7.2	8.3	9.4
2001.....	7.4	*	0.5	5.0	7.5	8.1	7.9	8.0	8.6
2000.....	7.3	*	0.7	5.0	8.5	7.4	7.5	8.3	9.0
1999.....	7.3	*	0.6	5.6	8.2	7.8	8.0	8.4	8.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Non-Hispanic White, male

2003.....	13.2	*	0.7	7.4	13.3	12.1	13.3	12.8	15.2
2002.....	13.4	*	0.8	7.2	14.0	13.3	12.0	14.0	15.8
2001.....	13.2	*	0.8	8.6	13.6	14.2	13.4	13.2	14.5
2000.....	12.9	*	1.2	8.6	15.3	12.9	12.5	13.5	15.0
1999.....	12.9	*	1.0	9.7	14.7	13.6	13.3	13.8	13.5

Non-Hispanic White, female

2003.....	1.8	*	*	0.9	1.3	1.8	2.1	2.4	3.0
2002.....	1.8	*	*	0.9	1.5	2.3	2.3	2.6	2.9
2001.....	1.9	*	*	1.2	1.2	1.9	2.3	2.7	2.7
2000.....	1.9	*	*	1.2	1.5	1.8	2.3	3.0	3.0
1999.....	1.9	*	*	1.3	1.5	2.0	2.5	2.9	2.8

Non-Hispanic Black, both sexes

2003.....	2.7	*	*	2.1	6.7	5.8	5.0	2.9	2.5
2002.....	2.9	*	*	2.5	6.0	6.5	4.1	4.1	3.0
2001.....	3.0	*	*	2.9	7.4	5.9	4.2	3.8	3.3
2000.....	3.1	*	*	3.9	7.1	6.2	4.1	3.5	3.4
1999.....	3.2	*	*	4.0	7.9	5.9	4.4	4.3	2.8

Non-Hispanic Black, male

2003.....	5.1	*	*	4.0	12.3	11.6	9.5	5.8	4.3
2002.....	5.5	*	*	4.4	11.4	13.2	8.0	7.7	5.3
2001.....	5.6	*	*	5.2	14.5	11.0	8.2	7.3	5.6
2000.....	5.7	*	*	7.3	13.0	11.8	7.5	6.6	6.3
1999.....	6.1	*	*	7.1	14.8	11.6	8.4	8.4	5.3

Non-Hispanic Black, female

2003.....	0.5	*	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
All origins 3/, both sexes											
2003.....	5.0	*	0.8	3.7	5.6	6.1	7.4	7.8	8.7
2002.....	5.0	*	0.8	3.8	5.7	6.4	6.9	7.7	8.7
2001 5/.....	4.8	*	0.9	3.8	5.4	5.9	6.7	7.9	8.2
2000.....	4.5	*	0.9	3.6	5.3	5.8	6.0	7.0	7.9
1999.....	4.5	*	0.7	3.2	5.1	5.7	6.7	7.3	7.7
All origins 3/, male											
2003.....	7.3	*	1.2	5.4	8.8	9.3	11.0	11.7	12.4
2002.....	7.3	*	1.2	5.8	8.9	9.7	10.6	11.4	12.1
2001 5/.....	7.1	*	1.3	5.8	8.7	9.3	10.2	11.9	11.6
2000.....	6.6	*	1.4	5.4	8.5	8.9	9.1	10.6	11.3
1999.....	6.6	*	1.1	4.6	8.0	8.8	10.2	10.9	10.8
All origins 3/, female											
2003.....	2.8	*	0.4	1.9	2.2	2.7	3.7	4.0	5.1
2002.....	2.9	*	0.4	1.6	2.3	2.9	3.1	4.0	5.4
2001 5/.....	2.6	*	0.4	1.7	2.0	2.5	3.1	3.9	4.8
2000.....	2.5	*	0.4	1.7	1.9	2.6	2.7	3.4	4.6
1999.....	2.5	*	0.3	1.6	2.0	2.5	3.2	3.7	4.5
Hispanic, both sexes											
2003.....	2.9	*	0.9	3.4	4.1	3.4	4.5	4.0	4.7
2002.....	2.9	*	0.7	2.9	3.8	3.6	4.2	4.5	4.5
2001 5/.....	2.8	*	1.0	3.1	3.5	3.7	4.1	5.0	4.6
2000.....	2.8	*	0.6	2.8	3.9	3.6	3.7	4.6	4.9
1999.....	2.7	*	*	2.6	2.8	3.6	4.1	5.0	3.8
Hispanic, male											
2003.....	4.5	*	1.3	4.9	6.2	5.1	6.9	6.2	7.3
2002.....	4.4	*	*	4.2	5.5	5.9	6.4	7.0	7.0
2001 5/.....	4.4	*	1.3	4.3	5.1	5.5	6.2	8.3	7.4
2000.....	4.3	*	*	3.8	6.2	5.9	5.9	7.0	7.7
1999.....	4.2	*	*	4.0	4.3	5.4	6.5	7.9	6.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Hispanic, female											
2003.....	1.2	*	*	1.7	1.5	1.3	1.7	1.5	2.0
2002.....	1.2	*	*	1.5	1.7	*	1.7	1.7	1.9
2001 5/.....	1.2	*	*	1.9	1.4	1.4	1.8	*	*
2000.....	1.1	*	*	1.7	*	*	*	1.9	1.9
1999.....	1.1	*	*	*	*	1.4	1.4	1.9	*
Non-Hispanic 4/, both sexes											
2003.....	5.3	*	0.8	3.7	5.9	6.7	7.9	8.4	9.2
2002.....	5.3	*	0.8	3.9	6.1	7.0	7.4	8.2	9.2
2001 5/.....	5.1	*	0.9	3.9	5.8	6.4	7.1	8.3	8.5
2000.....	4.8	*	1.0	3.7	5.5	6.2	6.3	7.3	8.2
1999.....	4.7	*	0.7	3.3	5.6	6.0	7.1	7.6	8.0
Non-Hispanic 4/, male											
2003.....	7.7	*	1.1	5.4	9.3	10.4	11.8	12.6	13.1
2002.....	7.7	*	1.2	6.1	9.7	10.7	11.5	12.1	12.7
2001 5/.....	7.5	*	1.3	6.0	9.5	10.2	10.9	12.4	12.0
2000.....	6.9	*	1.5	5.6	9.0	9.6	9.7	11.1	11.6
1999.....	6.9	*	1.1	4.8	8.9	9.4	10.9	11.3	11.3
Non-Hispanic 4/, female											
2003.....	3.1	*	0.4	1.9	2.3	2.9	4.0	4.3	5.4
2002.....	3.1	*	0.4	1.6	2.4	3.3	3.3	4.4	5.8
2001 5/.....	2.9	*	0.4	1.7	2.2	2.7	3.4	4.2	5.1
2000.....	2.7	*	0.4	1.7	2.1	2.8	2.9	3.6	4.9
1999.....	2.7	*	0.3	1.7	2.2	2.6	3.4	3.9	4.8
Non-Hispanic White, both sexes											
2003.....	5.9	*	0.7	4.1	6.2	7.3	8.9	9.3	10.3
2002.....	6.0	*	0.9	4.5	6.7	7.6	8.1	9.2	10.4
2001 5/.....	5.6	*	0.8	4.4	6.5	7.0	7.8	9.2	9.6
2000.....	5.2	*	0.9	4.0	5.8	6.6	6.9	8.1	9.1
1999.....	5.2	*	0.7	3.4	6.0	6.4	7.7	8.3	8.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Non-Hispanic White, male											
2003.....	8.4	*	1.0	5.9	10.0	11.4	13.2	13.8	14.4
2002.....	8.5	*	1.2	7.0	10.7	11.4	12.4	13.4	14.2
2001 5/.....	8.2	*	1.2	6.8	10.5	11.0	11.9	13.6	13.4
2000.....	7.5	*	1.4	6.2	9.5	10.3	10.4	12.1	12.6
1999.....	7.4	*	1.1	5.0	9.6	9.9	11.6	12.1	12.6
Non-Hispanic White, female											
2003.....	3.4	*	0.3	2.1	2.3	3.2	4.5	4.9	6.2
2002.....	3.5	*	0.5	1.8	2.6	3.8	3.8	5.0	6.6
2001 5/.....	3.2	*	0.4	1.7	2.4	2.9	3.8	4.8	5.8
2000.....	3.0	*	0.4	1.7	2.1	2.9	3.4	4.0	5.5
1999.....	3.0	*	*	1.7	2.3	2.8	3.7	4.5	5.3
Non-Hispanic Black, both sexes											
2003.....	2.5	*	1.0	1.8	4.1	4.0	4.4	4.7	4.6
2002.....	2.4	*	0.7	1.5	3.4	4.4	4.4	4.1	4.2
2001 5/.....	2.3	*	0.8	1.5	3.1	3.8	4.5	4.3	3.6
2000.....	2.4	*	1.1	1.8	4.3	4.0	3.7	3.7	4.8
1999.....	2.3	*	0.8	2.0	3.2	4.1	4.7	4.3	4.2
Non-Hispanic Black, male											
2003.....	3.9	*	1.6	2.8	6.2	6.5	6.8	7.6	7.3
2002.....	3.8	*	1.2	2.5	5.2	7.3	7.6	7.0	6.4
2001 5/.....	3.7	*	1.2	2.3	5.6	6.8	7.4	7.0	5.2
2000.....	3.9	*	1.9	2.6	7.1	6.4	6.6	6.3	7.4
1999.....	3.8	*	*	3.0	5.5	7.1	7.9	7.7	5.8
Non-Hispanic Black, female											
2003.....	1.3	*	*	*	2.0	1.7	2.3	2.2	2.3
2002.....	1.1	*	*	*	1.6	1.7	1.4	1.6	2.3
2001 5/.....	1.1	*	*	*	*	*	1.9	1.9	2.2
2000.....	1.1	*	*	*	1.7	1.9	*	1.4	2.4
1999.....	1.0	*	*	*	*	*	1.9	*	2.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All origins 3/, both sexes

2003.....	6.1	8.5	2.4	0.6	1.0	9.5	16.5	13.4	9.4	7.6	6.4
2002.....	6.1	7.5	2.7	0.7	1.0	9.3	16.5	13.3	9.4	7.9	6.5
2001 5/.....	7.1	8.2	2.7	0.7	0.9	9.4	17.3	14.9	11.5	10.4	8.6
2000.....	6.0	9.2	2.3	0.7	1.1	9.5	16.0	11.9	9.0	7.6	6.6
1999.....	6.1	8.7	2.5	0.9	1.2	10.4	15.6	12.1	9.1	7.7	6.6

All origins 3/, male

2003.....	9.7	10.0	2.5	0.7	1.3	15.9	27.6	22.4	14.8	11.4	9.5
2002.....	9.6	7.9	2.9	0.6	1.2	15.3	27.7	22.0	14.7	11.8	9.6
2001 5/.....	11.1	9.5	3.0	0.7	1.1	15.7	28.9	24.1	17.7	15.3	12.6
2000.....	9.3	10.4	2.5	0.7	1.4	15.5	26.7	19.6	14.0	10.9	9.7
1999.....	9.3	9.6	2.5	0.9	1.5	16.9	25.7	19.5	13.8	11.0	9.5

All origins 3/, female

2003.....	2.6	6.9	2.3	0.5	0.6	2.6	4.8	4.0	3.9	3.9	3.4
2002.....	2.7	7.1	2.5	0.8	0.8	2.9	4.6	4.2	4.1	4.1	3.5
2001 5/.....	3.3	6.9	2.4	0.6	0.7	2.7	5.1	5.5	5.2	5.5	4.7
2000.....	2.8	7.9	2.1	0.6	0.8	3.1	4.7	4.1	4.0	4.4	3.6
1999.....	2.9	7.8	2.4	0.9	0.9	3.5	5.1	4.5	4.3	4.3	3.6

Hispanic, both sexes

2003.....	8.4	6.9	2.5	*	1.0	14.5	21.8	15.6	10.6	9.3	7.5
2002.....	8.1	6.2	2.6	0.8	1.0	14.8	20.2	14.8	9.8	8.7	7.2
2001 5/.....	9.0	6.7	2.3	0.5	1.1	13.8	22.3	16.1	11.7	11.0	9.6
2000.....	8.3	7.0	1.9	0.7	1.6	15.0	19.1	13.7	11.1	8.8	8.6
1999.....	8.4	7.0	1.8	0.8	2.0	15.7	19.6	14.8	10.5	9.1	8.4

Hispanic, male

2003.....	13.6	7.9	2.1	*	1.5	25.1	34.5	25.2	16.9	14.6	11.0
2002.....	13.2	6.6	3.1	*	1.2	25.5	32.9	23.9	15.4	13.0	11.1
2001 5/.....	14.5	7.9	2.5	*	1.5	23.9	36.1	25.3	17.6	16.5	15.2
2000.....	13.4	6.6	2.0	*	2.3	25.7	30.9	22.2	17.5	13.5	13.6
1999.....	13.8	6.7	1.7	*	2.8	26.0	31.8	24.3	16.6	14.1	13.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Hispanic, female

2003.....	2.8	5.9	2.9	*	*	3.1	5.9	3.8	3.3	3.4	3.8
2002.....	2.6	5.9	2.1	*	*	3.1	4.5	3.6	3.2	3.8	3.1
2001 5/.....	3.2	5.5	2.1	*	*	2.6	5.3	5.2	4.9	5.0	3.7
2000.....	2.8	7.4	1.9	*	*	2.8	4.6	3.7	3.8	3.6	3.5
1999.....	2.8	7.3	1.9	*	*	4.0	4.9	3.6	3.4	3.5	3.0

Non-Hispanic 4/, both sexes

2003.....	5.7	8.9	2.3	0.7	0.9	8.4	15.3	12.8	9.1	7.3	6.2
2002.....	5.7	7.7	2.7	0.7	1.0	8.2	15.5	12.7	9.3	7.7	6.4
2001 5/.....	6.7	8.4	2.8	0.7	0.9	8.5	16.0	14.4	11.3	10.1	8.4
2000.....	5.6	9.6	2.4	0.7	1.0	8.4	15.1	11.5	8.6	7.4	6.4
1999.....	5.7	8.9	2.6	0.9	1.1	9.4	14.7	11.5	8.7	7.4	6.3

Non-Hispanic 4/, male

2003.....	9.0	10.5	2.6	0.8	1.3	14.0	25.8	21.5	14.3	10.7	9.2
2002.....	8.9	8.0	2.9	0.6	1.2	13.3	26.2	21.3	14.3	11.5	9.3
2001 5/.....	10.4	9.8	3.2	0.7	1.0	14.0	26.8	23.4	17.5	14.9	12.1
2000.....	8.6	11.2	2.6	0.7	1.3	13.4	25.5	18.9	13.2	10.4	9.2
1999.....	8.6	10.1	2.6	0.9	1.3	15.1	24.3	18.3	13.1	10.4	9.0

Non-Hispanic 4/, female

2003.....	2.5	7.2	2.1	0.6	0.6	2.5	4.5	4.0	4.0	3.9	3.3
2002.....	2.7	7.4	2.6	0.8	0.8	2.8	4.6	4.3	4.2	4.1	3.5
2001 5/.....	3.2	6.9	2.4	0.7	0.7	2.7	5.0	5.4	5.2	5.5	4.7
2000.....	2.7	7.9	2.2	0.7	0.8	3.0	4.7	4.2	4.0	4.5	3.6
1999.....	2.9	7.6	2.5	0.9	0.9	3.4	5.1	4.7	4.4	4.4	3.6

Non-Hispanic White, both sexes

2003.....	2.7	6.8	1.3	0.4	0.5	2.5	4.5	4.0	3.8	3.9	3.7
2002.....	2.8	5.0	1.8	0.5	0.6	2.8	4.7	4.1	3.8	4.2	4.0
2001 5/.....	4.0	5.8	1.7	0.7	0.6	2.9	5.5	6.5	6.6	7.0	6.2
2000.....	2.8	6.2	1.6	0.6	0.7	2.7	4.4	4.0	3.9	4.1	3.8
1999.....	2.9	6.3	1.7	0.6	0.7	3.1	4.7	4.2	4.4	4.3	3.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Non-Hispanic White, male

2003.....	3.6	8.0	1.3	0.4	0.7	3.6	6.5	5.4	5.1	5.0	4.8
2002.....	3.8	5.8	1.9	0.3	0.6	3.9	6.6	5.6	5.0	5.6	5.7
2001 5/.....	5.6	6.8	2.0	0.6	0.6	3.9	7.5	9.3	9.4	10.3	8.7
2000.....	3.6	8.3	1.8	0.6	0.8	3.5	6.0	5.2	5.2	5.1	5.3
1999.....	3.9	8.0	1.7	0.6	0.8	4.2	6.1	5.7	5.7	5.6	5.2

Non-Hispanic White, female

2003.....	1.8	5.5	1.4	*	*	1.4	2.4	2.6	2.4	2.8	2.5
2002.....	1.9	4.1	1.8	0.6	0.6	1.6	2.7	2.5	2.7	2.8	2.3
2001 5/.....	2.4	4.7	1.5	0.7	0.6	1.9	3.5	3.6	3.8	3.7	3.7
2000.....	1.9	4.1	1.3	0.5	0.6	1.9	2.8	2.8	2.6	3.1	2.3
1999.....	2.0	4.4	1.6	0.6	0.7	2.0	3.3	2.8	3.1	3.0	2.5

Non-Hispanic Black, both sexes

2003.....	22.7	16.8	6.4	1.8	2.8	34.2	64.5	56.9	37.7	26.3	22.1
2002.....	22.5	17.6	6.5	1.4	2.6	31.9	65.6	56.5	38.1	27.8	20.9
2001 5/.....	22.7	19.5	7.1	0.9	1.9	33.6	64.9	54.6	36.5	29.0	22.0
2000.....	22.1	24.1	6.1	1.2	2.6	34.1	66.4	50.4	34.3	26.6	22.3
1999.....	21.7	20.0	6.4	2.0	2.6	37.7	63.0	48.8	32.6	25.7	21.5

Non-Hispanic Black, male

2003.....	40.1	18.6	7.8	2.1	3.8	60.7	115.5	106.2	66.6	45.4	37.9
2002.....	39.4	16.4	7.3	1.5	3.5	54.8	119.6	104.5	67.2	46.6	33.1
2001 5/.....	39.3	21.7	8.4	*	2.8	59.7	120.3	99.7	64.5	44.2	34.7
2000.....	38.3	24.9	6.1	*	3.3	59.0	122.1	93.5	59.9	43.8	34.8
1999.....	37.2	20.1	6.2	2.0	3.5	64.6	116.0	87.2	56.2	41.1	34.6

Non-Hispanic Black, female

2003.....	6.8	14.8	4.9	1.4	1.7	7.0	14.0	11.4	11.8	9.4	8.2
2002.....	7.2	18.9	5.7	1.4	1.6	8.2	13.2	12.7	11.9	11.1	10.2
2001 5/.....	7.6	17.1	5.7	*	*	6.6	11.8	13.6	11.4	15.6	10.7
2000.....	7.4	23.3	6.0	1.5	1.8	8.6	13.7	11.4	11.3	11.3	11.2
1999.....	7.7	20.0	6.7	1.9	1.6	10.1	12.8	14.1	11.4	12.1	10.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All origins 3/, both sexes

2003.....	4.1	*	0.3	0.2	0.7	7.7	13.6	10.8	7.1	5.0	3.8
2002.....	4.1	*	0.3	0.3	0.7	7.7	13.6	10.6	6.9	5.2	3.9
2001.....	4.0	*	0.4	0.3	0.6	7.5	13.6	10.3	6.6	5.1	3.7
2000.....	3.8	*	0.2	0.2	0.7	7.7	12.7	9.2	6.3	4.7	3.8
1999.....	3.9	*	0.3	0.3	0.8	8.5	12.5	9.2	6.1	4.6	3.7

All origins 3/, male

2003.....	7.1	*	0.2	0.3	1.0	13.7	23.8	19.1	12.0	8.0	5.9
2002.....	7.0	*	0.4	0.3	0.9	13.2	23.8	18.6	11.6	8.4	6.2
2001.....	6.8	*	0.3	0.3	0.8	13.2	24.0	17.8	11.1	8.0	5.6
2000.....	6.5	*	*	0.3	1.0	13.3	22.4	16.1	10.4	7.4	5.8
1999.....	6.5	*	0.3	0.4	1.1	14.5	21.8	15.9	10.1	7.1	5.5

All origins 3/, female

2003.....	1.2	*	0.3	0.2	0.3	1.5	2.8	2.2	2.1	1.9	1.7
2002.....	1.3	*	*	0.3	0.5	1.8	2.9	2.4	2.1	2.0	1.7
2001.....	1.3	*	0.4	0.2	0.3	1.5	2.7	2.5	2.1	2.2	1.7
2000.....	1.3	*	*	0.2	0.3	1.7	2.7	2.2	2.1	1.9	1.7
1999.....	1.3	*	0.3	0.2	0.5	2.2	2.8	2.3	2.1	2.2	1.8

Hispanic, both sexes

2003.....	5.8	*	*	*	0.7	11.9	17.6	11.9	7.7	6.1	4.4
2002.....	5.6	*	*	*	0.7	12.6	15.9	11.6	6.8	5.4	4.6
2001.....	5.7	*	*	*	0.7	11.7	17.1	11.7	7.2	5.8	5.4
2000.....	5.5	*	*	*	1.0	12.6	14.8	10.3	7.4	4.8	4.9
1999.....	5.7	*	*	*	1.4	13.4	15.3	10.5	6.8	5.4	5.0

Hispanic, male

2003.....	10.0	*	*	*	1.1	21.1	29.0	20.1	13.1	9.7	6.7
2002.....	9.7	*	*	*	*	22.3	27.0	19.3	11.6	8.7	7.3
2001.....	9.9	*	*	*	*	20.4	28.8	19.8	12.0	9.4	8.8
2000.....	9.6	*	*	*	1.7	21.9	24.9	17.5	12.3	7.5	8.1
1999.....	9.8	*	*	*	2.1	22.9	25.4	18.1	11.1	9.1	7.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Hispanic, female

2003.....	1.3	*	*	*	*	2.0	3.3	1.9	1.5	1.9	1.9
2002.....	1.2	*	*	*	*	2.0	2.2	2.2	1.4	1.7	1.8
2001.....	1.3	*	*	*	*	2.1	2.8	2.0	1.6	1.8	1.9
2000.....	1.3	*	*	*	*	2.0	2.5	2.0	1.8	1.9	*
1999.....	1.4	*	*	*	*	2.6	3.2	1.7	1.8	*	1.9

Non-Hispanic 4/, both sexes

2003.....	3.8	*	0.2	0.3	0.7	6.9	12.7	10.5	6.9	4.7	3.7
2002.....	3.8	*	0.3	0.3	0.7	6.7	13.0	10.3	6.9	5.1	3.8
2001.....	3.7	*	0.3	0.3	0.5	6.7	12.7	9.8	6.5	5.0	3.4
2000.....	3.6	*	0.2	0.2	0.6	6.7	12.2	9.0	6.0	4.6	3.6
1999.....	3.6	*	0.4	0.3	0.7	7.6	11.9	8.8	5.9	4.5	3.5

Non-Hispanic 4/, male

2003.....	6.5	*	*	0.3	1.0	12.1	22.4	18.7	11.7	7.7	5.8
2002.....	6.5	*	0.3	*	0.9	11.4	22.9	18.2	11.5	8.3	6.0
2001.....	6.3	*	0.3	0.4	0.8	11.7	22.6	17.1	10.8	7.8	5.2
2000.....	6.0	*	*	0.2	0.9	11.5	21.6	15.7	10.0	7.3	5.5
1999.....	6.0	*	0.4	0.4	1.0	12.9	20.9	15.3	9.8	6.7	5.2

Non-Hispanic 4/, female

2003.....	1.2	*	*	*	0.3	1.4	2.7	2.2	2.2	1.9	1.7
2002.....	1.3	*	*	0.3	0.5	1.8	3.0	2.4	2.2	2.0	1.7
2001.....	1.2	*	0.3	0.3	0.3	1.4	2.6	2.6	2.1	2.3	1.7
2000.....	1.2	*	*	*	0.3	1.7	2.7	2.3	2.1	2.0	1.8
1999.....	1.3	*	0.4	*	0.4	2.0	2.8	2.5	2.1	2.2	1.8

Non-Hispanic White, both sexes

2003.....	1.4	*	*	*	0.3	1.7	3.0	2.6	2.4	2.3	2.0
2002.....	1.5	*	0.2	0.2	0.3	1.9	3.1	2.7	2.5	2.5	2.3
2001.....	1.6	*	0.2	0.3	0.4	1.8	3.2	2.8	2.6	2.7	2.1
2000.....	1.4	*	*	0.2	0.3	1.7	2.7	2.5	2.4	2.4	2.1
1999.....	1.5	*	0.2	0.2	0.4	2.0	2.9	2.7	2.6	2.5	2.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Non-Hispanic White, male

2003.....	2.1	*	*	*	0.4	2.6	4.8	3.9	3.5	3.1	2.8
2002.....	2.2	*	*	*	0.4	2.8	4.7	4.0	3.5	3.5	3.4
2001.....	2.2	*	*	0.4	0.5	2.6	4.7	4.3	3.6	3.9	2.9
2000.....	2.0	*	*	*	0.5	2.5	4.0	3.5	3.4	3.4	3.0
1999.....	2.2	*	*	*	0.5	3.0	4.2	4.0	3.8	3.4	2.9

Non-Hispanic White, female

2003.....	0.8	*	*	*	*	0.7	1.2	1.2	1.4	1.5	1.2
2002.....	0.9	*	*	*	*	1.0	1.5	1.4	1.4	1.5	1.2
2001.....	0.9	*	*	*	*	0.9	1.6	1.4	1.6	1.5	1.3
2000.....	0.9	*	*	*	*	0.9	1.5	1.4	1.4	1.5	1.3
1999.....	0.9	*	*	*	0.3	0.9	1.5	1.3	1.4	1.6	1.3

Non-Hispanic Black, both sexes

2003.....	17.3	*	*	0.8	2.1	30.0	57.1	50.1	31.3	18.8	14.1
2002.....	17.1	*	*	*	2.0	28.1	59.0	48.9	31.0	20.1	13.3
2001.....	16.3	*	*	*	1.4	29.5	57.6	45.3	27.9	19.1	11.8
2000.....	16.0	*	*	*	1.8	29.5	57.3	42.5	26.6	17.8	12.9
1999.....	15.9	*	*	0.7	1.9	33.0	55.8	40.7	24.3	16.3	12.5

Non-Hispanic Black, male

2003.....	32.6	*	*	*	3.0	55.0	105.3	96.7	58.8	35.3	25.1
2002.....	31.9	*	*	*	2.8	49.8	109.9	94.2	58.1	36.9	23.1
2001.....	30.6	*	*	*	2.2	54.2	110.3	86.3	53.2	32.6	20.7
2000.....	30.1	*	*	*	2.8	53.3	109.1	82.3	49.4	32.6	22.0
1999.....	29.3	*	*	*	2.9	58.3	105.7	77.2	45.3	28.0	21.0

Non-Hispanic Black, female

2003.....	3.4	*	*	*	1.2	4.2	9.4	7.2	6.5	4.2	4.6
2002.....	3.7	*	*	*	*	5.6	9.5	7.6	6.8	5.2	4.7
2001.....	3.2	*	*	*	*	4.0	7.2	8.0	5.2	7.0	4.0
2000.....	3.3	*	*	*	*	5.2	8.4	6.5	6.2	4.6	4.9
1999.....	3.7	*	*	*	*	7.2	8.5	7.8	5.5	5.9	5.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1)

All origins 3/, both sexes

2003.....	2.0	8.3	2.1	0.4	0.3	1.7	2.9	2.6	2.3	2.7	2.6
2002.....	2.0	7.3	2.4	0.4	0.3	1.6	2.9	2.6	2.5	2.7	2.6
2001 5/.....	3.1	8.0	2.3	0.4	0.3	1.8	3.7	4.6	4.9	5.2	4.9
2000.....	2.1	8.9	2.1	0.4	0.5	1.8	3.2	2.7	2.8	2.9	2.9
1999.....	2.2	8.5	2.1	0.6	0.4	1.9	3.1	2.9	3.0	3.0	2.9

All origins 3/, male

2003.....	2.6	9.7	2.3	0.4	0.4	2.3	3.8	3.3	2.8	3.4	3.6
2002.....	2.6	7.8	2.6	0.4	0.3	2.1	3.9	3.4	3.0	3.4	3.4
2001 5/.....	4.3	9.2	2.7	0.4	0.3	2.4	4.9	6.3	6.7	7.3	7.0
2000.....	2.8	10.0	2.3	0.5	0.4	2.2	4.4	3.5	3.6	3.5	3.9
1999.....	2.8	9.4	2.1	0.5	0.4	2.4	3.9	3.6	3.7	3.9	4.0

All origins 3/, female

2003.....	1.4	6.9	2.0	0.3	0.2	1.1	2.0	1.8	1.8	1.9	1.6
2002.....	1.4	6.8	2.2	0.5	0.3	1.1	1.8	1.8	2.0	2.1	1.8
2001 5/.....	2.0	6.7	2.0	0.4	0.4	1.2	2.4	3.0	3.2	3.2	2.9
2000.....	1.5	7.7	2.0	0.4	0.5	1.4	2.0	1.9	2.0	2.4	1.8
1999.....	1.6	7.6	2.1	0.7	0.5	1.4	2.2	2.2	2.3	2.2	1.8

Hispanic, both sexes

2003.....	2.6	6.8	2.2	*	*	2.6	4.2	3.7	2.8	3.2	3.1
2002.....	2.5	6.0	2.2	*	*	2.2	4.3	3.2	2.9	3.2	2.6
2001 5/.....	3.3	6.2	1.9	*	*	2.1	5.2	4.4	4.5	5.3	4.2
2000.....	2.7	6.9	1.8	*	*	2.4	4.3	3.3	3.7	4.0	3.7
1999.....	2.7	6.7	1.6	*	*	2.3	4.3	4.2	3.7	3.6	3.4

Hispanic, male

2003.....	3.6	7.7	1.8	*	*	4.0	5.5	5.1	3.8	4.8	4.3
2002.....	3.5	6.6	2.6	*	*	3.2	5.9	4.7	3.9	4.3	3.7
2001 5/.....	4.5	7.2	2.2	*	*	3.5	7.4	5.4	5.6	7.2	6.5
2000.....	3.8	6.6	1.8	*	*	3.8	6.0	4.7	5.2	6.0	5.4
1999.....	4.0	6.2	1.5	*	*	3.2	6.4	6.2	5.5	5.1	5.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1)

Hispanic, female

2003.....	1.5	5.9	2.6	*	*	*	2.6	1.9	1.8	1.5	1.9
2002.....	1.4	5.4	1.8	*	*	*	2.2	1.4	1.8	2.1	*
2001 5/.....	1.9	5.2	1.6	*	*	*	2.5	3.2	3.2	3.2	1.8
2000.....	1.6	7.2	1.7	*	*	*	2.1	1.7	2.0	1.8	2.0
1999.....	1.4	7.3	1.8	*	*	1.4	1.7	1.9	1.6	2.1	*

Non-Hispanic 4/, both sexes

2003.....	1.9	8.6	2.1	0.4	0.3	1.5	2.6	2.3	2.2	2.5	2.5
2002.....	1.9	7.5	2.4	0.4	0.3	1.5	2.5	2.4	2.4	2.6	2.5
2001 5/.....	3.1	8.2	2.4	0.4	0.3	1.8	3.3	4.6	4.9	5.1	4.9
2000.....	2.0	9.2	2.2	0.4	0.4	1.7	3.0	2.6	2.6	2.8	2.8
1999.....	2.1	8.7	2.2	0.6	0.4	1.8	2.8	2.6	2.8	2.9	2.8

Non-Hispanic 4/, male

2003.....	2.4	10.1	2.4	0.5	0.3	1.9	3.4	2.8	2.5	3.1	3.5
2002.....	2.5	7.8	2.5	0.4	0.3	1.9	3.4	3.1	2.8	3.2	3.4
2001 5/.....	4.2	9.5	2.8	0.4	0.2	2.2	4.2	6.3	6.7	7.1	6.9
2000.....	2.6	10.6	2.4	0.4	0.3	1.9	3.9	3.2	3.2	3.1	3.8
1999.....	2.6	10.0	2.3	0.5	0.3	2.3	3.3	3.0	3.3	3.7	3.8

Non-Hispanic 4/, female

2003.....	1.4	7.1	1.8	0.3	0.3	1.1	1.8	1.8	1.8	2.0	1.6
2002.....	1.4	7.1	2.3	0.5	0.3	1.1	1.7	1.8	2.0	2.1	1.7
2001 5/.....	2.0	6.8	2.0	0.4	0.4	1.3	2.4	2.8	3.1	3.2	3.0
2000.....	1.5	7.8	2.0	0.5	0.5	1.4	2.0	1.9	1.9	2.5	1.8
1999.....	1.6	7.3	2.2	0.7	0.4	1.4	2.3	2.2	2.4	2.1	1.8

Non-Hispanic White, both sexes

2003.....	1.2	6.8	1.2	0.2	0.2	0.8	1.5	1.4	1.4	1.6	1.6
2002.....	1.3	5.0	1.6	0.3	0.2	0.9	1.6	1.4	1.4	1.7	1.7
2001 5/.....	2.4	5.6	1.5	0.3	0.3	1.2	2.4	3.6	4.0	4.3	4.1
2000.....	1.3	6.0	1.4	0.3	0.4	1.0	1.7	1.6	1.5	1.7	1.7
1999.....	1.4	6.2	1.4	0.4	0.3	1.2	1.9	1.6	1.8	1.8	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1)

Non-Hispanic White, male

2003.....	1.5	8.0	1.2	*	*	1.0	1.7	1.4	1.6	1.9	2.1
2002.....	1.6	5.8	1.6	*	*	1.1	2.0	1.6	1.5	2.1	2.3
2001 5/.....	3.4	6.7	1.8	*	*	1.3	2.8	5.0	5.7	6.4	5.8
2000.....	1.6	8.1	1.7	0.4	0.3	1.0	2.0	1.7	1.8	1.8	2.3
1999.....	1.7	7.8	1.5	0.4	*	1.3	1.9	1.7	1.9	2.2	2.3

Non-Hispanic White, female

2003.....	1.0	5.4	1.2	*	*	0.7	1.2	1.4	1.1	1.4	1.2
2002.....	1.0	4.1	1.6	0.4	*	0.6	1.3	1.1	1.3	1.4	1.1
2001 5/.....	1.5	4.5	1.2	0.4	0.4	1.0	1.9	2.2	2.3	2.2	2.4
2000.....	1.0	3.9	1.2	*	0.4	1.0	1.4	1.4	1.2	1.7	1.0
1999.....	1.1	4.4	1.4	0.5	0.4	1.1	1.8	1.4	1.7	1.4	1.2

Non-Hispanic Black, both sexes

2003.....	5.4	15.8	5.6	1.0	0.6	4.3	7.4	6.8	6.5	7.5	7.9
2002.....	5.4	16.7	5.9	1.0	0.6	3.8	6.6	7.6	7.0	7.6	7.6
2001 5/.....	6.4	19.1	6.3	*	*	4.1	7.3	9.3	8.6	10.0	10.2
2000.....	6.1	22.9	5.7	0.9	0.8	4.6	9.0	7.8	7.7	8.8	9.4
1999.....	5.8	19.7	5.7	1.3	0.7	4.7	7.2	8.1	8.3	9.4	9.1

Non-Hispanic Black, male

2003.....	7.6	16.8	6.9	*	*	5.6	10.2	9.5	7.8	10.1	12.9
2002.....	7.6	15.8	6.6	*	*	5.0	9.7	10.3	9.1	9.7	10.1
2001 5/.....	8.7	21.4	7.7	*	*	5.6	10.0	13.4	11.3	11.6	14.1
2000.....	8.2	22.5	5.7	*	*	5.7	13.0	11.2	10.5	11.2	12.8
1999.....	7.9	19.7	5.6	1.2	*	6.3	10.3	10.0	10.9	13.0	13.6

Non-Hispanic Black, female

2003.....	3.4	14.8	4.3	*	*	2.8	4.6	4.2	5.2	5.1	3.6
2002.....	3.5	17.7	5.2	*	*	2.6	3.6	5.1	5.1	5.9	5.5
2001 5/.....	4.4	16.8	5.0	*	*	2.6	4.6	5.6	6.2	8.6	6.7
2000.....	4.1	23.3	5.8	*	*	3.4	5.3	4.8	5.1	6.7	6.4
1999.....	4.0	19.6	5.7	1.4	*	2.9	4.4	6.3	5.9	6.2	5.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Legal intervention (Y35,Y89.0)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.1	*	*	*	*	*	0.3	0.3	0.3	0.3	0.2
2002.....	0.1	*	*	*	*	*	0.3	0.2	0.2	0.3	0.2
2001.....	0.1	*	*	*	*	*	0.3	0.2	0.3	0.3	0.2
2000.....	0.1	*	*	*	*	0.1	0.2	0.3	0.2	0.3	0.2
1999.....	0.1	*	*	*	*	*	0.3	0.3	0.3	0.3	0.2

Non-Hispanic 4/, male

2003.....	0.3	*	*	*	*	*	0.5	0.6	0.6	0.7	0.5
2002.....	0.2	*	*	*	*	*	0.6	0.5	0.5	0.5	0.4
2001.....	0.3	*	*	*	*	*	0.5	0.4	0.6	0.6	0.4
2000.....	0.2	*	*	*	*	0.3	0.4	0.7	0.5	0.5	0.3
1999.....	0.3	*	*	*	*	*	0.5	0.7	0.6	0.6	0.4

Non-Hispanic 4/, female

2003.....	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic White, both sexes

2003.....	0.1	*	*	*	*	*	0.2	*	0.2	0.3	0.2
2002.....	0.1	*	*	*	*	*	*	*	0.2	0.2	0.2
2001.....	0.1	*	*	*	*	*	0.2	0.2	0.3	0.3	0.2
2000.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2	0.1
1999.....	0.1	*	*	*	*	*	*	0.2	0.2	0.2	0.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All origins 3/, both sexes

2003.....	1.7	1.9	0.2	*	0.2	0.8	1.9	1.8	2.3	3.2	3.9
2002.....	1.7	2.5	0.4	0.1	0.2	0.7	1.7	1.9	2.3	3.3	3.4
2001.....	1.5	2.1	0.3	*	0.2	0.7	1.4	1.6	2.0	3.0	3.2
2000.....	1.4	1.7	0.3	*	0.2	0.6	1.3	1.6	1.8	2.6	3.1
1999.....	1.4	1.4	0.3	*	0.2	0.8	1.5	1.6	2.1	2.8	2.9

All origins 3/, male

2003.....	2.3	2.4	*	*	0.3	1.2	2.7	2.5	3.2	4.2	4.9
2002.....	2.2	2.5	0.4	*	0.2	1.0	2.4	2.7	3.1	4.5	4.3
2001.....	2.0	2.4	*	*	0.3	1.0	2.1	2.4	2.7	3.9	4.4
2000.....	1.9	1.4	0.3	*	0.4	0.9	2.0	2.3	2.6	3.7	4.2
1999.....	2.0	1.7	0.3	*	0.3	1.3	2.2	2.3	2.8	3.9	4.0

All origins 3/, female

2003.....	1.2	1.3	*	*	*	0.5	1.1	1.1	1.5	2.2	2.9
2002.....	1.2	2.5	0.4	*	*	0.4	1.0	1.0	1.4	2.1	2.6
2001.....	0.9	1.7	0.3	*	*	0.3	0.7	0.9	1.3	2.1	2.0
2000.....	0.8	2.0	*	*	*	0.3	0.6	0.9	1.1	1.6	1.9
1999.....	0.9	1.2	0.3	*	*	0.3	0.6	0.9	1.3	1.7	1.9

Hispanic, both sexes

2003.....	0.8	*	*	*	*	*	1.0	0.9	1.4	1.3	1.4
2002.....	0.8	*	*	*	*	*	1.1	0.8	1.2	1.7	1.4
2001.....	0.8	*	*	*	*	*	0.6	0.9	1.0	1.6	1.7
2000.....	0.6	*	*	*	*	*	0.7	0.9	1.0	1.5	1.0
1999.....	0.7	*	*	*	*	0.7	1.0	1.0	1.2	1.3	1.1

Hispanic, male

2003.....	1.3	*	*	*	*	*	1.3	1.4	2.1	1.8	2.4
2002.....	1.2	*	*	*	*	*	1.7	1.1	1.8	2.4	1.8
2001.....	1.1	*	*	*	*	*	*	1.3	1.5	2.2	3.0
2000.....	0.9	*	*	*	*	*	1.1	1.2	1.6	2.1	*
1999.....	1.1	*	*	*	*	*	1.6	1.5	1.7	2.0	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Hispanic, female

2003.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	1.9	2.2	0.2	*	0.2	0.9	2.1	2.0	2.5	3.5	4.2
2002.....	1.8	2.7	0.4	0.1	0.2	0.7	1.8	2.1	2.5	3.6	3.7
2001.....	1.6	2.4	0.3	*	0.2	0.7	1.6	1.8	2.2	3.2	3.3
2000.....	1.5	1.9	0.3	*	0.2	0.6	1.5	1.7	1.9	2.8	3.3
1999.....	1.5	1.6	0.3	*	0.2	0.8	1.5	1.7	2.2	3.0	3.1

Non-Hispanic 4/, male

2003.....	2.5	3.0	*	*	0.3	1.3	3.1	2.9	3.4	4.6	5.3
2002.....	2.3	2.8	0.3	*	0.3	1.1	2.6	3.0	3.4	4.9	4.6
2001.....	2.1	2.7	*	*	0.3	1.0	2.5	2.7	2.9	4.2	4.5
2000.....	2.0	1.6	0.4	*	0.4	1.0	2.2	2.5	2.7	3.9	4.5
1999.....	2.1	1.9	*	*	0.3	1.3	2.4	2.5	3.0	4.2	4.2

Non-Hispanic 4/, female

2003.....	1.3	1.3	*	*	*	0.5	1.1	1.2	1.7	2.4	3.2
2002.....	1.3	2.6	0.4	*	*	0.4	1.1	1.2	1.6	2.3	2.8
2001.....	1.0	2.0	0.4	*	*	0.3	0.7	1.0	1.5	2.3	2.2
2000.....	0.9	2.2	*	*	*	0.3	0.7	0.9	1.2	1.7	2.1
1999.....	0.9	*	*	*	*	0.3	0.7	0.9	1.5	1.9	2.0

Non-Hispanic White, both sexes

2003.....	2.0	2.0	*	*	0.2	1.0	2.4	2.3	2.8	3.7	4.4
2002.....	1.8	1.8	0.4	*	0.2	0.8	2.1	2.3	2.6	3.7	3.8
2001.....	1.6	1.7	0.3	*	0.2	0.8	1.8	2.0	2.4	3.3	3.4
2000.....	1.4	1.5	0.2	*	0.2	0.7	1.5	1.8	2.0	2.8	3.2
1999.....	1.4	1.3	*	*	0.2	0.8	1.6	1.7	2.3	3.0	3.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic White, male

2003.....	2.5	2.4	*	*	*	1.5	3.3	3.1	3.7	4.9	5.4
2002.....	2.3	1.9	*	*	0.3	1.1	2.9	3.2	3.5	4.9	4.7
2001.....	2.1	2.0	*	*	0.3	1.2	2.6	2.8	3.1	4.1	4.4
2000.....	2.0	*	*	*	0.4	1.1	2.2	2.6	2.7	3.9	4.3
1999.....	2.0	*	*	*	0.4	1.4	2.4	2.4	3.0	3.9	4.0

Non-Hispanic White, female

2003.....	1.4	*	*	*	*	0.6	1.4	1.5	2.0	2.6	3.5
2002.....	1.3	1.8	*	*	*	0.5	1.2	1.3	1.7	2.5	2.9
2001.....	1.1	*	*	*	*	0.4	0.9	1.1	1.6	2.5	2.4
2000.....	0.9	*	*	*	*	0.4	0.8	1.1	1.3	1.8	2.0
1999.....	1.0	*	*	*	*	*	0.7	1.1	1.6	2.0	1.9

Non-Hispanic Black, both sexes

2003.....	1.8	3.3	*	*	*	*	1.2	1.4	2.2	3.3	4.3
2002.....	2.0	5.7	*	*	*	*	1.3	2.0	2.5	4.1	4.5
2001.....	1.8	5.9	*	*	*	*	1.1	1.5	2.0	3.9	4.0
2000.....	1.9	*	*	*	*	*	1.4	1.7	2.2	3.4	4.9
1999.....	2.0	*	*	*	*	0.8	1.7	2.0	2.6	4.0	4.7

Non-Hispanic Black, male

2003.....	2.8	*	*	*	*	*	2.0	2.5	3.5	4.5	6.3
2002.....	2.9	*	*	*	*	*	1.9	3.2	3.7	6.7	5.6
2001.....	2.8	6.4	*	*	*	*	2.0	2.5	2.8	6.0	6.6
2000.....	2.9	*	*	*	*	*	2.6	2.8	3.5	5.0	7.3
1999.....	3.1	*	*	*	*	1.3	2.8	3.6	3.8	6.4	6.7

Non-Hispanic Black, female

2003.....	1.0	*	*	*	*	*	*	*	*	2.3	2.6
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.9	3.4
2001.....	0.9	*	*	*	*	*	*	*	*	2.1	1.8
2000.....	1.0	*	*	*	*	*	*	*	*	2.0	2.8
1999.....	0.9	*	*	*	*	*	*	*	1.5	1.8	3.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Discharge of firearms, undetermined intent (Y22-Y24)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.1	*	*	*	*	0.1	0.2	*	*	*	*
2002.....	0.1	*	*	*	*	0.2	0.2	0.1	*	0.1	*
2001.....	0.1	*	*	*	*	0.1	0.2	0.2	*	0.2	*
2000.....	0.1	*	*	*	*	*	0.3	0.2	*	*	0.1
1999.....	0.1	*	*	*	*	0.3	0.3	0.2	*	0.1	0.1

Non-Hispanic 4/, male

2003.....	0.1	*	*	*	*	0.3	0.3	*	*	*	*
2002.....	0.1	*	*	*	*	0.3	0.4	*	*	0.2	*
2001.....	0.1	*	*	*	*	*	0.4	0.3	*	*	*
2000.....	0.1	*	*	*	*	*	0.4	0.3	*	*	*
1999.....	0.2	*	*	*	*	0.6	0.5	0.3	*	0.2	0.2

Non-Hispanic 4/, female

2003.....	0.0	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*

Non-Hispanic White, both sexes

2003.....	0.1	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.2	*	*	0.1	*
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	0.2	*
2000.....	0.1	*	*	*	*	*	0.2	0.2	*	*	*
1999.....	0.1	*	*	*	*	0.3	0.3	0.2	*	0.1	0.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All origins 3/, both sexes

2003.....	1.7	1.9	0.2	*	0.1	0.7	1.7	1.7	2.3	3.1	3.8
2002.....	1.6	2.5	0.4	0.1	0.2	0.5	1.5	1.7	2.2	3.2	3.3
2001.....	1.4	2.1	0.3	*	0.1	0.5	1.2	1.5	1.9	2.9	3.1
2000.....	1.3	1.7	0.3	*	0.2	0.5	1.1	1.4	1.7	2.6	3.0
1999.....	1.3	1.4	0.3	*	0.1	0.5	1.1	1.4	2.0	2.7	2.8

All origins 3/, male

2003.....	2.2	2.4	*	*	0.2	0.9	2.4	2.4	3.0	4.1	4.8
2002.....	2.1	2.5	0.3	*	0.2	0.7	2.1	2.4	3.0	4.3	4.1
2001.....	1.9	2.4	*	*	0.2	0.8	1.8	2.1	2.6	3.8	4.2
2000.....	1.8	1.4	0.3	*	0.3	0.7	1.6	2.0	2.4	3.6	4.1
1999.....	1.8	1.7	*	*	0.2	0.7	1.7	2.0	2.7	3.7	3.8

All origins 3/, female

2003.....	1.2	1.3	*	*	*	0.4	1.0	1.1	1.5	2.1	2.8
2002.....	1.1	2.5	0.4	*	*	0.3	0.9	1.0	1.4	2.0	2.6
2001.....	0.9	1.7	0.3	*	*	0.3	0.6	0.8	1.3	2.0	2.0
2000.....	0.8	2.0	*	*	*	0.3	0.5	0.8	1.0	1.6	1.9
1999.....	0.8	1.2	0.3	*	*	0.2	0.5	0.8	1.3	1.7	1.8

Hispanic, both sexes

2003.....	0.8	*	*	*	*	*	0.8	0.8	1.3	1.3	1.3
2002.....	0.8	*	*	*	*	*	0.8	0.7	1.2	1.6	1.4
2001.....	0.7	*	*	*	*	*	*	0.8	0.9	1.5	1.7
2000.....	0.6	*	*	*	*	*	*	0.8	1.0	1.4	1.0
1999.....	0.6	*	*	*	*	*	0.7	0.9	1.2	1.2	1.1

Hispanic, male

2003.....	1.1	*	*	*	*	*	1.0	1.2	2.0	1.7	2.2
2002.....	1.1	*	*	*	*	*	1.3	*	1.7	2.3	1.8
2001.....	1.1	*	*	*	*	*	*	1.2	1.5	2.1	3.0
2000.....	0.8	*	*	*	*	*	*	1.1	1.6	2.0	*
1999.....	1.0	*	*	*	*	*	1.2	1.4	1.7	1.9	1.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Hispanic, female											
2003.....	0.4	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes											
2003.....	1.8	2.2	0.2	*	0.1	0.7	1.9	1.9	2.4	3.4	4.1
2002.....	1.7	2.7	0.3	0.1	0.2	0.6	1.6	2.0	2.4	3.4	3.6
2001.....	1.5	2.4	0.3	*	0.2	0.6	1.4	1.6	2.1	3.1	3.2
2000.....	1.4	1.9	0.3	*	0.2	0.5	1.2	1.5	1.8	2.7	3.2
1999.....	1.4	1.6	0.3	*	0.1	0.5	1.2	1.5	2.1	2.9	3.0
Non-Hispanic 4/, male											
2003.....	2.3	3.0	*	*	*	1.0	2.7	2.7	3.3	4.5	5.2
2002.....	2.2	2.8	0.3	*	0.3	0.8	2.2	2.8	3.2	4.6	4.4
2001.....	2.0	2.7	*	*	0.2	0.8	2.0	2.3	2.8	4.0	4.4
2000.....	1.9	1.6	0.4	*	0.3	0.7	1.8	2.2	2.5	3.8	4.4
1999.....	1.9	1.9	*	*	*	0.7	1.8	2.1	2.8	3.9	4.0
Non-Hispanic 4/, female											
2003.....	1.3	1.3	*	*	*	0.5	1.1	1.2	1.6	2.3	3.1
2002.....	1.2	2.6	0.4	*	*	0.4	1.0	1.1	1.5	2.2	2.8
2001.....	1.0	2.0	0.4	*	*	0.3	0.7	0.9	1.5	2.2	2.1
2000.....	0.9	2.2	*	*	*	0.3	0.6	0.9	1.1	1.7	2.0
1999.....	0.9	*	*	*	*	*	0.6	0.9	1.4	1.8	2.0
Non-Hispanic White, both sexes											
2003.....	1.9	2.0	*	*	*	0.9	2.2	2.2	2.7	3.6	4.3
2002.....	1.7	1.8	0.4	*	0.2	0.7	1.9	2.2	2.5	3.6	3.7
2001.....	1.5	1.7	0.3	*	0.2	0.7	1.6	1.8	2.3	3.1	3.3
2000.....	1.4	1.5	0.2	*	0.2	0.6	1.3	1.7	1.9	2.7	3.1
1999.....	1.3	1.3	*	*	*	0.6	1.3	1.6	2.2	2.8	2.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Non-Hispanic White, male											
2003.....	2.4	2.4	*	*	*	1.2	3.1	3.0	3.5	4.7	5.2
2002.....	2.2	1.9	*	*	*	0.9	2.6	3.0	3.3	4.7	4.5
2001.....	2.0	2.0	*	*	*	1.0	2.2	2.5	3.0	3.9	4.3
2000.....	1.8	*	*	*	0.3	0.9	1.9	2.3	2.5	3.8	4.1
1999.....	1.8	*	*	*	*	0.9	1.9	2.1	2.8	3.7	3.8
Non-Hispanic White, female											
2003.....	1.4	*	*	*	*	0.6	1.3	1.5	1.9	2.5	3.4
2002.....	1.3	1.8	*	*	*	0.5	1.1	1.3	1.7	2.5	2.8
2001.....	1.1	*	*	*	*	0.4	0.9	1.1	1.6	2.3	2.3
2000.....	0.9	*	*	*	*	0.3	0.7	1.0	1.3	1.7	2.0
1999.....	0.9	*	*	*	*	*	0.6	1.1	1.5	1.9	1.8
Non-Hispanic Black, both sexes											
2003.....	1.8	3.3	*	*	*	*	0.8	1.3	2.0	3.3	4.3
2002.....	1.9	5.7	*	*	*	*	1.1	1.8	2.4	4.0	4.4
2001.....	1.7	5.9	*	*	*	*	0.8	1.3	2.0	3.9	4.0
2000.....	1.7	*	*	*	*	*	0.9	1.4	2.0	3.3	4.8
1999.....	1.8	*	*	*	*	*	1.2	1.6	2.4	3.9	4.6
Non-Hispanic Black, male											
2003.....	2.6	*	*	*	*	*	1.4	2.1	3.3	4.5	6.2
2002.....	2.7	*	*	*	*	*	1.4	2.7	3.3	6.4	5.5
2001.....	2.6	6.4	*	*	*	*	1.5	2.3	2.7	6.0	6.5
2000.....	2.6	*	*	*	*	*	1.6	2.2	3.1	4.7	7.2
1999.....	2.8	*	*	*	*	*	1.8	2.9	3.5	6.2	6.6
Non-Hispanic Black, female											
2003.....	1.0	*	*	*	*	*	*	*	*	2.2	2.6
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.8	3.4
2001.....	0.9	*	*	*	*	*	*	*	*	2.1	1.7
2000.....	0.9	*	*	*	*	*	*	*	*	2.0	2.6
1999.....	0.9	*	*	*	*	*	*	*	1.4	1.8	2.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Complications of medical and surgical care (Y40-Y84,Y88)

Hispanic, female

2003.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	1.1	*	*	*	*	*	0.1	0.1	0.2	0.4	0.5
2002.....	1.1	*	*	*	*	*	*	0.2	0.3	0.4	0.5
2001.....	1.2	0.7	*	*	*	*	0.1	*	0.3	0.3	0.4
2000.....	1.2	*	*	*	*	*	*	0.2	0.3	0.3	0.4
1999.....	1.1	1.6	0.2	*	*	*	*	0.2	0.2	0.3	0.4

Non-Hispanic 4/, male

2003.....	1.0	*	*	*	*	*	*	*	0.3	0.3	0.5
2002.....	1.0	*	*	*	*	*	*	*	0.4	0.3	0.4
2001.....	1.1	*	*	*	*	*	*	*	0.3	0.4	0.4
2000.....	1.1	*	*	*	*	*	*	0.3	0.3	0.3	0.4
1999.....	1.0	1.8	*	*	*	*	*	*	0.4	0.3	0.4

Non-Hispanic 4/, female

2003.....	1.1	*	*	*	*	*	*	*	*	0.4	0.4
2002.....	1.2	*	*	*	*	*	*	*	*	0.4	0.5
2001.....	1.2	*	*	*	*	*	*	*	0.3	0.3	0.5
2000.....	1.2	*	*	*	*	*	*	*	0.3	0.3	0.4
1999.....	1.2	1.4	*	*	*	*	*	*	*	0.3	0.4

Non-Hispanic White, both sexes

2003.....	1.1	*	*	*	*	*	*	*	0.2	0.3	0.5
2002.....	1.1	*	*	*	*	*	*	*	0.2	0.3	0.4
2001.....	1.2	*	*	*	*	*	*	*	0.3	0.3	0.4
2000.....	1.2	*	*	*	*	*	*	0.2	0.3	0.3	0.3
1999.....	1.1	1.3	*	*	*	*	*	*	0.2	0.3	0.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/

All origins 3/, both sexes

2003.....	7.1	*	*	*	*	0.2	0.4	0.9	2.1	5.3	10.6
2002 7/.....	7.0	*	*	*	*	0.2	0.4	0.8	2.1	5.4	10.6
2001 7/.....	7.1	*	*	*	*	0.2	0.3	0.9	2.3	6.0	10.6
2000 7/.....	7.0	*	*	*	*	0.2	0.3	0.9	2.3	6.0	11.1
1999 7/.....	7.0	*	*	*	*	0.1	0.4	0.8	2.4	6.2	10.9

All origins 3/, male

2003.....	10.9	*	*	*	*	0.4	0.6	1.4	3.2	7.4	15.2
2002 7/.....	10.8	*	*	*	*	0.3	0.6	1.4	3.1	7.6	15.3
2001 7/.....	10.8	*	*	*	*	0.3	0.6	1.3	3.3	8.5	15.3
2000 7/.....	10.9	*	*	*	*	0.3	0.5	1.5	3.5	8.7	16.4
1999 7/.....	10.9	*	*	*	*	0.2	0.6	1.2	3.3	8.7	16.6

All origins 3/, female

2003.....	3.4	*	*	*	*	*	*	0.3	0.9	3.2	6.0
2002 7/.....	3.4	*	*	*	*	*	*	0.3	1.1	3.3	6.1
2001 7/.....	3.4	*	*	*	*	*	*	0.5	1.3	3.5	6.0
2000 7/.....	3.2	*	*	*	*	*	*	0.3	1.1	3.2	5.8
1999 7/.....	3.2	*	*	*	*	*	*	0.3	1.5	3.7	5.3

Hispanic, both sexes

2003.....	6.1	*	*	*	*	*	0.6	1.0	2.7	6.1	10.9
2002 7/.....	6.2	*	*	*	*	*	0.6	1.3	2.7	5.9	12.0
2001 7/.....	6.4	*	*	*	*	*	*	1.5	2.8	6.3	12.4
2000 7/.....	6.6	*	*	*	*	*	0.7	1.3	3.0	7.3	14.1
1999 7/.....	6.4	*	*	*	*	*	0.6	1.0	3.2	7.6	13.5

Hispanic, male

2003.....	9.9	*	*	*	*	*	1.0	1.7	4.1	10.0	17.0
2002 7/.....	10.3	*	*	*	*	*	1.1	2.3	4.5	9.5	20.1
2001 7/.....	10.7	*	*	*	*	*	*	2.2	4.5	10.6	19.8
2000 7/.....	11.1	*	*	*	*	*	1.2	2.2	5.3	11.7	23.2
1999 7/.....	10.7	*	*	*	*	*	1.2	1.8	5.1	12.2	22.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/

Hispanic, female

2003.....	1.9	*	*	*	*	*	*	*	*	1.7	4.3
2002 7/.....	1.8	*	*	*	*	*	*	*	*	2.0	3.5
2001 7/.....	2.0	*	*	*	*	*	*	*	*	1.5	4.6
2000 7/.....	1.7	*	*	*	*	*	*	*	*	2.5	4.6
1999 7/.....	1.9	*	*	*	*	*	*	*	*	2.6	4.2

Non-Hispanic 4/, both sexes

2003.....	7.2	*	*	*	*	0.2	0.3	0.8	1.9	5.2	10.5
2002 7/.....	7.1	*	*	*	*	0.1	0.3	0.7	2.0	5.3	10.4
2001 7/.....	7.1	*	*	*	*	0.2	0.3	0.7	2.2	5.9	10.3
2000 7/.....	7.0	*	*	*	*	0.2	0.2	0.8	2.2	5.7	10.6
1999 7/.....	7.0	*	*	*	*	*	0.3	0.7	2.2	5.9	10.5

Non-Hispanic 4/, male

2003.....	11.0	*	*	*	*	0.3	0.5	1.3	3.0	6.9	14.9
2002 7/.....	10.7	*	*	*	*	0.2	0.4	1.1	2.7	7.1	14.5
2001 7/.....	10.8	*	*	*	*	0.2	0.5	1.1	3.0	8.1	14.6
2000 7/.....	10.7	*	*	*	*	0.2	0.3	1.3	3.1	8.2	15.4
1999 7/.....	10.8	*	*	*	*	*	0.5	1.1	2.9	8.1	15.7

Non-Hispanic 4/, female

2003.....	3.6	*	*	*	*	*	*	0.3	0.9	3.5	6.2
2002 7/.....	3.6	*	*	*	*	*	*	0.3	1.2	3.5	6.4
2001 7/.....	3.6	*	*	*	*	*	*	0.4	1.4	3.8	6.2
2000 7/.....	3.4	*	*	*	*	*	*	0.3	1.3	3.3	6.0
1999 7/.....	3.4	*	*	*	*	*	*	0.3	1.6	3.8	5.4

Non-Hispanic White, both sexes

2003.....	7.5	*	*	*	*	0.2	0.3	0.8	1.9	5.2	10.7
2002 7/.....	7.3	*	*	*	*	0.2	0.3	0.7	1.9	5.2	10.6
2001 7/.....	7.2	*	*	*	*	*	0.2	0.6	2.1	5.9	10.0
2000 7/.....	7.0	*	*	*	*	0.2	0.2	0.8	2.0	5.6	10.2
1999 7/.....	6.9	*	*	*	*	*	0.3	0.7	2.1	5.6	9.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/

Non-Hispanic White, male

2003.....	11.4	*	*	*	*	*	0.5	1.3	2.9	6.9	15.2
2002 7/.....	11.1	*	*	*	*	*	0.4	1.0	2.6	7.0	14.8
2001 7/.....	10.8	*	*	*	*	*	0.4	1.0	2.9	7.8	13.9
2000 7/.....	10.8	*	*	*	*	*	*	1.2	2.8	8.0	14.8
1999 7/.....	10.7	*	*	*	*	*	0.5	1.1	2.7	7.6	14.9

Non-Hispanic White, female

2003.....	3.8	*	*	*	*	*	*	*	1.0	3.4	6.2
2002 7/.....	3.7	*	*	*	*	*	*	*	1.1	3.4	6.4
2001 7/.....	3.6	*	*	*	*	*	*	*	1.3	3.9	6.1
2000 7/.....	3.4	*	*	*	*	*	*	*	1.2	3.2	5.7
1999 7/.....	3.3	*	*	*	*	*	*	*	1.5	3.5	4.8

Non-Hispanic Black, both sexes

2003.....	6.5	*	*	*	*	*	*	0.9	1.4	5.0	9.0
2002 7/.....	6.6	*	*	*	*	*	*	*	2.0	4.5	9.2
2001 7/.....	7.5	*	*	*	*	*	*	0.8	1.9	5.3	12.5
2000 7/.....	7.6	*	*	*	*	*	*	0.9	2.6	5.6	13.1
1999 7/.....	8.0	*	*	*	*	*	*	*	2.2	7.3	13.2

Non-Hispanic Black, male

2003.....	10.3	*	*	*	*	*	*	*	2.3	6.8	13.0
2002 7/.....	10.3	*	*	*	*	*	*	*	3.0	6.3	12.9
2001 7/.....	11.9	*	*	*	*	*	*	*	2.6	8.5	18.6
2000 7/.....	11.7	*	*	*	*	*	*	1.6	3.5	8.3	19.3
1999 7/.....	12.5	*	*	*	*	*	*	*	2.9	10.5	19.2

Non-Hispanic Black, female

2003.....	3.0	*	*	*	*	*	*	*	*	3.3	5.5
2002 7/.....	3.3	*	*	*	*	*	*	*	*	2.9	6.0
2001 7/.....	3.5	*	*	*	*	*	*	*	*	2.5	7.1
2000 7/.....	3.9	*	*	*	*	*	*	*	1.8	3.3	7.6
1999 7/.....	4.0	*	*	*	*	*	*	*	1.6	4.5	7.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

All origins 3/, both sexes

2003.....	9.9	0.6	0.2	*	0.2	3.3	9.4	10.9	13.3	18.3	22.8
2002 7/.....	9.0	0.7	0.2	*	0.2	3.1	7.9	9.8	12.4	17.5	21.8
2001 7/.....	7.6	0.5	0.2	*	0.2	2.7	6.5	8.2	10.6	15.5	18.5
2000 7/.....	7.0	*	*	*	0.2	2.2	5.9	7.5	10.1	14.4	17.5
1999 7/.....	6.9	0.6	0.2	*	0.1	1.8	5.3	7.4	10.3	14.2	17.4

All origins 3/, male

2003.....	12.9	*	0.3	*	*	4.5	13.9	15.4	17.7	23.8	29.1
2002 7/.....	11.8	1.0	*	*	*	4.3	11.4	14.0	17.0	23.1	28.0
2001 7/.....	10.2	*	0.3	*	*	3.7	9.6	11.9	14.4	20.7	24.6
2000 7/.....	9.5	*	*	*	*	3.1	8.8	10.8	14.1	19.8	23.8
1999 7/.....	9.4	*	*	*	*	2.4	7.7	10.8	14.6	19.6	24.3

All origins 3/, female

2003.....	7.0	*	*	*	*	1.9	4.7	6.2	8.8	12.8	16.6
2002 7/.....	6.3	*	0.3	*	*	1.8	4.2	5.4	7.8	11.9	15.8
2001 7/.....	5.1	*	*	*	*	1.6	3.2	4.4	6.7	10.3	12.6
2000 7/.....	4.6	*	*	*	*	1.4	2.9	4.1	6.1	9.0	11.4
1999 7/.....	4.4	*	*	*	*	1.2	2.8	4.0	6.0	8.8	10.6

Hispanic, both sexes

2003.....	5.9	*	*	*	*	2.1	4.7	6.0	8.5	12.0	14.9
2002 7/.....	5.5	*	*	*	*	1.8	4.4	5.0	7.5	11.3	15.8
2001 7/.....	4.7	*	*	*	*	1.7	3.1	5.0	7.0	11.1	12.7
2000 7/.....	4.8	*	*	*	*	1.3	3.2	5.0	8.9	12.2	13.3
1999 7/.....	5.8	*	*	*	*	1.6	3.8	6.5	9.9	14.7	15.5

Hispanic, male

2003.....	8.7	*	*	*	*	3.0	6.9	8.8	12.3	17.7	21.8
2002 7/.....	8.2	*	*	*	*	2.6	6.4	7.8	11.0	16.4	24.1
2001 7/.....	7.0	*	*	*	*	2.3	4.1	7.6	9.8	16.6	19.3
2000 7/.....	7.4	*	*	*	*	1.8	4.9	7.1	14.2	17.9	21.8
1999 7/.....	9.2	*	*	*	*	2.1	6.1	9.9	15.8	22.5	25.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

Hispanic, female

2003.....	2.9	*	*	*	*	*	1.9	2.5	4.2	5.6	7.6
2002 7/.....	2.6	*	*	*	*	*	1.9	1.7	3.4	5.8	7.0
2001 7/.....	2.2	*	*	*	*	*	1.8	1.8	3.8	5.0	5.7
2000 7/.....	2.1	*	*	*	*	*	*	2.6	2.9	6.0	4.4
1999 7/.....	2.2	*	*	*	*	*	*	2.5	3.2	6.1	5.1

Non-Hispanic 4/, both sexes

2003.....	10.4	0.7	0.2	*	0.1	3.5	10.4	12.0	14.3	19.3	23.8
2002 7/.....	9.5	0.8	0.2	*	0.2	3.3	8.6	10.9	13.3	18.4	22.5
2001 7/.....	8.0	0.7	0.2	*	0.2	2.8	7.2	8.9	11.2	16.0	19.1
2000 7/.....	7.2	*	*	*	0.2	2.4	6.4	7.9	10.2	14.6	17.9
1999 7/.....	6.9	*	0.2	*	0.1	1.9	5.6	7.6	10.3	14.0	17.3

Non-Hispanic 4/, male

2003.....	13.5	*	*	*	*	4.8	15.5	17.1	18.8	24.7	30.0
2002 7/.....	12.3	*	*	*	*	4.6	12.5	15.6	18.2	24.2	28.3
2001 7/.....	10.6	*	*	*	*	3.9	10.8	12.9	15.2	21.2	25.0
2000 7/.....	9.7	*	*	*	*	3.3	9.6	11.6	14.0	20.0	23.8
1999 7/.....	9.3	*	*	*	*	2.5	8.0	10.9	14.3	19.0	23.8

Non-Hispanic 4/, female

2003.....	7.6	*	*	*	*	2.1	5.2	6.9	9.7	14.0	17.7
2002 7/.....	6.9	*	*	*	*	1.9	4.6	6.2	8.6	12.8	16.8
2001 7/.....	5.5	*	*	*	*	1.6	3.5	4.9	7.2	11.0	13.3
2000 7/.....	4.9	*	*	*	*	1.5	3.2	4.3	6.6	9.4	12.1
1999 7/.....	4.6	*	*	*	*	1.2	3.1	4.2	6.4	9.0	11.0

Non-Hispanic White, both sexes

2003.....	11.2	*	*	*	*	4.3	12.8	14.2	16.1	20.7	25.0
2002 7/.....	10.0	*	*	*	0.2	3.9	10.4	12.6	14.9	19.7	23.4
2001 7/.....	8.3	*	*	*	0.2	3.3	8.6	10.3	12.2	16.8	19.3
2000 7/.....	7.4	*	*	*	0.2	2.9	7.5	8.9	10.9	15.0	17.6
1999 7/.....	6.9	*	*	*	*	2.2	6.3	8.2	10.7	14.2	16.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

Non-Hispanic White, male

2003.....	14.3	*	*	*	*	6.0	19.1	20.4	21.2	26.3	31.4
2002 7/.....	12.8	*	*	*	*	5.5	15.0	17.9	20.3	25.6	29.3
2001 7/.....	10.8	*	*	*	*	4.8	12.7	14.7	16.7	22.1	25.2
2000 7/.....	9.8	*	*	*	*	4.0	11.3	12.8	14.8	20.6	23.2
1999 7/.....	9.2	*	*	*	*	2.9	9.0	11.5	14.8	19.1	22.8

Non-Hispanic White, female

2003.....	8.2	*	*	*	*	2.5	6.3	7.9	10.9	15.0	18.7
2002 7/.....	7.4	*	*	*	*	2.3	5.6	7.3	9.4	13.8	17.5
2001 7/.....	5.8	*	*	*	*	1.8	4.3	5.8	7.6	11.5	13.5
2000 7/.....	5.1	*	*	*	*	1.7	3.7	4.9	6.9	9.5	12.1
1999 7/.....	4.8	*	*	*	*	1.4	3.5	4.7	6.6	9.2	10.6

Non-Hispanic Black, both sexes

2003.....	9.5	*	*	*	*	0.7	3.0	5.8	10.7	18.4	24.2
2002 7/.....	9.4	*	*	*	*	1.0	3.3	6.8	10.6	17.7	24.6
2001 7/.....	8.7	*	*	*	*	0.9	3.1	5.9	9.7	17.0	23.9
2000 7/.....	8.5	*	*	*	*	0.9	3.2	6.2	10.7	16.9	24.5
1999 7/.....	8.7	*	*	*	*	0.8	3.9	7.1	11.5	16.7	26.1

Non-Hispanic Black, male

2003.....	13.0	*	*	*	*	*	4.6	7.7	14.0	24.9	31.8
2002 7/.....	13.2	*	*	*	*	1.4	4.8	10.1	14.3	24.7	31.8
2001 7/.....	12.5	*	*	*	*	*	5.2	9.4	12.6	22.7	32.4
2000 7/.....	12.3	*	*	*	*	*	4.7	9.9	14.7	23.2	34.5
1999 7/.....	12.9	*	*	*	*	*	5.9	11.4	16.1	24.1	37.2

Non-Hispanic Black, female

2003.....	6.3	*	*	*	*	*	1.5	4.1	7.7	12.7	17.6
2002 7/.....	6.0	*	*	*	*	*	1.9	3.8	7.2	11.4	18.3
2001 7/.....	5.3	*	*	*	*	*	*	2.7	7.2	11.9	16.4
2000 7/.....	5.0	*	*	*	*	*	1.9	3.0	7.0	11.3	15.5
1999 7/.....	4.9	*	*	*	*	*	2.0	3.2	7.5	10.2	16.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

All origins 3/, both sexes

2003.....	10.4	*	0.3	0.3	1.2	12.1	21.1	17.4	13.8	11.6	11.5
2002.....	10.5	*	0.4	0.4	1.3	12.1	21.3	17.8	13.2	12.4	11.9
2001.....	10.4	*	0.5	0.4	1.2	12.4	21.2	17.7	13.5	12.2	11.3
2000.....	10.2	*	0.3	0.3	1.5	12.9	20.9	16.3	12.8	12.0	11.8
1999.....	10.3	*	0.4	0.4	1.7	14.4	21.0	16.6	13.2	12.3	10.9

All origins 3/, male

2003.....	18.3	*	0.3	0.4	2.0	21.2	37.1	30.7	23.7	19.3	18.9
2002.....	18.4	*	0.5	0.4	1.9	21.1	37.6	31.1	22.3	20.6	19.7
2001.....	18.2	*	0.5	0.5	1.9	21.8	37.7	30.9	22.9	20.0	18.6
2000.....	17.8	*	0.4	0.4	2.4	22.4	37.0	28.5	21.4	19.5	19.4
1999.....	18.1	*	0.5	0.5	2.5	24.8	36.8	28.8	22.3	20.0	17.7

All origins 3/, female

2003.....	2.7	*	0.3	0.2	0.4	2.4	4.2	3.6	3.7	3.8	4.2
2002.....	2.8	*	0.3	0.3	0.7	2.7	4.2	4.1	3.9	4.2	4.2
2001.....	2.8	*	0.4	0.3	0.5	2.6	3.8	4.1	3.9	4.6	4.1
2000.....	2.8	*	*	0.2	0.5	2.9	4.2	3.9	4.1	4.5	4.3
1999.....	2.9	*	0.4	0.3	0.8	3.4	4.4	4.1	4.1	4.6	4.2

Hispanic, both sexes

2003.....	8.3	*	*	*	1.1	15.3	22.6	15.8	10.8	8.8	7.3
2002.....	8.1	*	*	*	1.1	16.1	20.7	15.2	9.7	8.5	7.2
2001.....	8.3	*	*	*	1.0	14.8	21.7	16.2	10.6	8.9	8.2
2000.....	8.2	*	*	*	1.4	16.1	19.7	13.9	10.2	7.5	8.7
1999.....	8.5	*	*	*	1.9	17.4	20.2	14.3	10.0	9.2	8.0

Hispanic, male

2003.....	14.6	*	*	*	1.8	27.0	37.6	26.7	18.5	14.3	12.0
2002.....	14.2	*	*	*	1.4	28.5	35.1	25.4	16.7	13.9	11.6
2001.....	14.6	*	*	*	1.5	25.5	36.4	27.3	17.8	14.8	14.0
2000.....	14.2	*	*	*	2.4	27.9	33.3	23.5	16.9	11.9	14.9
1999.....	14.6	*	*	*	2.9	29.5	33.6	24.3	16.7	15.2	13.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
Hispanic, female											
2003.....	1.7	*	*	*	*	2.8	4.1	2.3	1.9	2.5	2.2
2002.....	1.6	*	*	*	*	2.6	2.9	2.8	1.7	2.4	2.6
2001.....	1.7	*	*	*	*	2.9	3.7	3.0	2.3	2.3	2.2
2000.....	1.8	*	*	*	*	2.7	3.1	2.6	2.5	2.8	2.2
1999.....	2.0	*	*	*	*	3.6	4.0	2.6	2.2	2.7	2.6
Non-Hispanic 4/, both sexes											
2003.....	10.6	*	0.3	0.3	1.2	11.4	20.7	17.7	14.3	12.0	12.0
2002.....	10.8	*	0.4	0.3	1.3	11.3	21.3	18.3	13.8	13.0	12.4
2001.....	10.6	*	0.5	0.4	1.3	11.9	20.8	17.9	14.0	12.7	11.6
2000.....	10.4	*	0.3	0.3	1.5	12.2	21.0	16.8	13.2	12.5	12.0
1999.....	10.6	*	0.5	0.4	1.6	13.8	21.1	17.0	13.7	12.6	11.2
Non-Hispanic 4/, male											
2003.....	18.8	*	0.4	0.4	2.0	20.0	36.8	31.6	24.8	20.1	19.8
2002.....	19.0	*	0.5	0.3	2.0	19.5	37.9	32.4	23.4	21.7	20.7
2001.....	18.7	*	0.5	0.5	1.9	20.9	37.7	31.6	23.8	20.7	19.0
2000.....	18.3	*	0.4	0.4	2.4	21.1	37.6	29.6	22.1	20.5	19.8
1999.....	18.5	*	0.5	0.5	2.4	23.8	37.5	29.7	23.2	20.5	18.1
Non-Hispanic 4/, female											
2003.....	2.9	*	*	*	0.4	2.3	4.2	3.9	4.0	4.0	4.4
2002.....	3.0	*	*	0.4	0.7	2.7	4.5	4.4	4.3	4.4	4.4
2001.....	3.0	*	0.4	0.3	0.6	2.5	3.8	4.4	4.2	4.9	4.3
2000.....	3.0	*	*	0.2	0.5	2.9	4.4	4.1	4.3	4.7	4.5
1999.....	3.0	*	0.4	0.3	0.8	3.4	4.5	4.4	4.3	4.9	4.4
Non-Hispanic White, both sexes											
2003.....	9.3	*	*	0.2	0.9	6.5	11.2	10.1	10.6	10.6	11.7
2002.....	9.4	*	0.3	0.2	1.0	6.7	11.8	11.2	10.2	11.3	12.2
2001.....	9.4	*	0.3	0.4	1.1	7.4	11.5	11.7	11.1	11.5	11.3
2000.....	9.1	*	0.2	0.3	1.4	7.4	12.0	10.7	10.4	11.3	11.7
1999.....	9.3	*	0.3	0.3	1.5	8.4	12.1	11.3	11.2	11.5	10.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
ages 1/ 1 year 2/ years years years years years years years years years

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Non-Hispanic White, male

2003.....	16.0	*	*	*	1.5	11.1	19.4	17.0	17.6	17.1	18.9
2002.....	16.3	*	*	*	1.5	11.2	20.2	18.3	16.6	18.3	20.2
2001.....	16.3	*	*	0.5	1.7	12.3	19.9	19.7	18.1	18.3	18.5
2000.....	15.7	*	*	0.4	2.2	12.3	20.6	17.7	16.8	17.8	19.0
1999.....	15.9	*	*	*	2.2	14.2	20.6	18.9	18.2	18.3	17.4

Non-Hispanic White, female

2003.....	2.7	*	*	*	*	1.7	2.7	3.1	3.6	4.1	4.4
2002.....	2.8	*	*	*	0.5	1.9	3.2	3.8	3.8	4.2	4.3
2001.....	2.9	*	*	0.4	0.5	2.2	2.9	3.5	4.0	4.6	4.2
2000.....	2.9	*	*	*	0.5	2.2	3.2	3.5	3.9	4.7	4.4
1999.....	2.9	*	*	*	0.7	2.3	3.3	3.5	4.0	4.7	4.3

Non-Hispanic Black, both sexes

2003.....	20.7	*	0.9	0.9	2.6	33.2	66.0	58.1	37.2	22.3	17.1
2002.....	20.7	*	*	*	2.5	32.3	67.0	57.1	36.3	25.2	16.9
2001.....	19.8	*	1.0	*	1.9	33.8	66.7	52.3	32.6	23.3	15.7
2000.....	19.8	*	*	*	2.3	35.0	66.3	50.2	31.7	22.3	16.9
1999.....	20.0	*	1.1	0.9	2.4	39.2	66.3	48.2	29.8	21.5	16.0

Non-Hispanic Black, male

2003.....	39.1	*	*	*	3.9	61.2	121.6	112.6	70.3	42.0	30.3
2002.....	38.9	*	*	*	3.7	57.5	125.0	111.0	68.1	46.3	29.6
2001.....	37.3	*	*	*	3.1	62.1	127.9	99.4	62.2	40.8	27.2
2000.....	37.1	*	*	*	3.8	63.5	125.7	97.1	58.5	41.1	29.2
1999.....	37.1	*	*	1.3	3.5	69.4	125.3	91.9	55.9	38.1	27.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Non-Hispanic Black, female											
2003.....	4.0	*	*	*	1.3	4.4	10.9	7.8	7.5	4.9	5.6
2002.....	4.3	*	*	*	1.3	6.2	10.6	8.0	7.6	6.5	5.7
2001.....	4.0	*	*	*	*	4.6	8.0	9.4	6.0	7.8	5.6
2000.....	4.1	*	*	*	*	5.9	10.1	7.8	7.7	5.6	6.1
1999.....	4.5	*	*	*	*	8.3	10.3	8.7	6.5	6.7	5.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes									
All origins 3/, both sexes									
2003.....	359.2	517.8	755.6	1,182.7	1,793.7	2,778.3	4,363.4	6,976.2	14,593.3
2002.....	357.1	512.8	770.1	1,187.7	1,832.7	2,845.9	4,449.1	7,103.8	14,828.3
2001.....	354.8	512.4	771.8	1,210.7	1,869.7	2,878.3	4,494.0	7,151.9	15,112.8
2000.....	352.5	509.0	792.6	1,241.0	1,906.3	2,929.4	4,553.8	7,335.0	15,524.4
1999.....	346.6	501.4	801.6	1,254.0	1,956.7	2,995.3	4,608.0	7,378.5	15,554.6
All origins 3/, male									
2003.....	457.9	660.8	942.3	1,462.2	2,216.6	3,428.7	5,400.3	8,513.7	15,794.0
2002.....	454.7	653.5	961.8	1,476.0	2,265.9	3,533.0	5,522.7	8,652.3	16,254.5
2001.....	455.4	646.8	958.4	1,496.8	2,315.6	3,587.4	5,603.2	8,781.9	16,744.8
2000.....	456.3	642.2	989.8	1,535.9	2,362.0	3,676.1	5,693.2	9,095.4	17,501.4
1999.....	448.0	632.4	1,003.8	1,560.8	2,444.1	3,785.1	5,806.4	9,216.8	17,597.2
All origins 3/, female									
2003.....	263.2	380.9	579.7	928.8	1,427.1	2,256.6	3,625.3	6,036.0	14,062.5
2002.....	262.4	378.2	590.0	926.0	1,458.7	2,296.7	3,690.9	6,162.2	14,209.6
2001.....	257.2	383.7	596.9	951.3	1,485.9	2,314.8	3,717.6	6,175.1	14,429.9
2000.....	252.0	381.3	608.1	973.7	1,515.7	2,341.2	3,760.2	6,296.6	14,719.2
1999.....	248.5	375.9	612.6	976.4	1,539.6	2,378.2	3,774.8	6,302.8	14,731.3
Hispanic, both sexes									
2003.....	272.7	400.3	590.7	886.5	1,352.1	2,107.8	3,380.8	5,251.8	10,431.5
2002.....	266.6	393.2	583.0	905.8	1,378.9	2,179.7	3,411.7	5,365.5	10,679.0
2001.....	280.9	403.9	580.6	910.1	1,412.8	2,229.5	3,556.3	5,501.5	11,821.8
2000.....	281.1	394.3	612.1	913.8	1,425.9	2,278.4	3,543.6	5,777.6	11,822.9
1999.....	283.6	398.1	602.7	940.8	1,527.8	2,264.3	3,577.0	5,774.4	12,177.9
Hispanic, male									
2003.....	365.3	532.3	763.9	1,146.5	1,701.1	2,660.5	4,124.7	6,357.9	11,029.2
2002.....	353.7	516.8	779.4	1,158.2	1,765.5	2,747.9	4,258.0	6,447.7	11,674.1
2001.....	379.7	529.6	751.2	1,158.6	1,817.5	2,791.5	4,456.8	6,783.5	12,888.3
2000.....	378.6	519.5	801.5	1,181.3	1,804.0	2,918.9	4,480.2	7,220.4	13,086.2
1999.....	383.9	525.5	788.0	1,215.5	1,942.0	2,931.5	4,571.8	7,005.8	13,485.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes									
Hispanic, female									
2003.....	177.9	271.9	430.3	659.9	1,065.6	1,683.6	2,853.0	4,531.9	10,128.1
2002.....	178.1	273.2	402.6	685.7	1,063.3	1,742.1	2,807.4	4,674.8	10,186.0
2001.....	181.7	283.8	426.0	695.1	1,085.0	1,800.8	2,917.0	4,707.4	11,300.5
2000.....	183.8	274.9	440.6	683.1	1,119.7	1,793.2	2,880.3	4,899.7	11,202.8
1999.....	184.2	277.2	435.3	704.0	1,192.9	1,762.5	2,874.8	5,039.8	11,547.3
Non-Hispanic 4/, both sexes									
2003.....	367.2	526.6	766.2	1,201.4	1,821.5	2,815.4	4,408.5	7,042.5	14,740.6
2002.....	365.0	521.3	782.3	1,204.9	1,859.4	2,879.6	4,490.9	7,161.0	14,958.7
2001.....	360.4	519.3	783.7	1,228.7	1,896.2	2,908.7	4,528.0	7,201.4	15,199.0
2000.....	357.3	515.8	802.9	1,259.9	1,931.9	2,956.6	4,586.6	7,375.0	15,619.8
1999.....	350.4	507.2	812.7	1,271.3	1,977.5	3,025.1	4,639.4	7,416.1	15,631.0
Non-Hispanic 4/, male									
2003.....	466.2	669.7	952.7	1,479.9	2,246.3	3,468.2	5,458.2	8,599.8	15,984.2
2002.....	463.0	662.5	972.1	1,493.6	2,292.2	3,569.8	5,572.1	8,727.3	16,418.0
2001.....	460.5	653.3	969.6	1,514.1	2,341.5	3,621.9	5,644.0	8,842.8	16,866.3
2000.....	460.9	648.3	998.9	1,554.1	2,388.4	3,704.3	5,730.8	9,144.1	17,639.8
1999.....	450.9	637.0	1,014.4	1,577.2	2,466.2	3,815.8	5,842.6	9,272.9	17,710.4
Non-Hispanic 4/, female									
2003.....	271.5	389.8	590.4	947.5	1,451.7	2,290.3	3,661.1	6,092.9	14,194.2
2002.....	270.3	386.3	603.7	942.0	1,484.2	2,326.6	3,727.8	6,210.5	14,329.1
2001.....	263.6	391.0	608.9	968.9	1,511.2	2,340.5	3,747.4	6,219.1	14,505.6
2000.....	257.3	388.8	619.1	992.4	1,539.1	2,366.3	3,790.4	6,332.6	14,802.9
1999.....	253.4	382.7	623.8	993.6	1,557.9	2,406.0	3,803.4	6,330.3	14,798.4
Non-Hispanic White, both sexes									
2003.....	333.6	478.6	716.6	1,142.1	1,762.4	2,761.7	4,371.2	7,056.2	14,944.8
2002.....	331.7	475.5	731.3	1,145.4	1,800.0	2,825.5	4,446.2	7,166.0	15,149.1
2001.....	324.7	474.0	730.8	1,169.7	1,827.4	2,849.2	4,476.8	7,196.1	15,371.2
2000.....	318.7	470.1	749.9	1,199.4	1,865.4	2,889.5	4,525.0	7,369.2	15,799.7
1999.....	310.1	461.9	754.0	1,206.2	1,911.7	2,949.5	4,570.3	7,406.6	15,795.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes									
Non-Hispanic White, male									
2003.....	426.7	609.7	888.4	1,402.1	2,169.8	3,393.7	5,403.0	8,607.0	16,234.4
2002.....	424.3	604.5	906.2	1,413.9	2,216.9	3,500.6	5,514.6	8,733.5	16,641.9
2001.....	417.3	596.5	901.6	1,439.8	2,256.8	3,543.1	5,580.2	8,828.0	17,055.3
2000.....	415.7	588.9	928.4	1,477.3	2,305.7	3,615.5	5,653.4	9,137.4	17,853.2
1999.....	399.6	578.8	939.4	1,494.4	2,384.2	3,724.5	5,759.4	9,266.7	17,871.2
Non-Hispanic White, female									
2003.....	241.4	350.2	551.1	900.2	1,400.2	2,242.3	3,624.5	6,103.0	14,377.2
2002.....	240.1	349.1	563.3	896.1	1,430.8	2,273.2	3,681.6	6,207.1	14,504.3
2001.....	233.1	353.9	567.1	919.2	1,448.1	2,285.2	3,695.5	6,211.0	14,670.6
2000.....	222.8	353.6	579.3	941.7	1,478.3	2,305.2	3,731.9	6,318.4	14,971.7
1999.....	221.7	347.5	576.9	939.4	1,497.5	2,331.8	3,735.9	6,309.1	14,967.5
Non-Hispanic Black, both sexes									
2003.....	647.8	959.3	1,321.4	1,920.6	2,640.7	3,810.9	5,490.4	7,836.9	14,107.7
2002.....	648.6	943.7	1,346.7	1,911.2	2,670.7	3,877.8	5,630.1	8,036.6	14,386.7
2001.....	656.8	946.1	1,351.9	1,920.5	2,777.9	3,951.1	5,712.5	8,098.0	14,660.8
2000.....	679.7	956.6	1,368.9	1,959.9	2,809.4	4,065.0	5,837.2	8,234.9	14,837.5
1999.....	681.6	945.4	1,422.8	1,998.3	2,851.2	4,191.4	5,927.6	8,223.1	14,839.2
Non-Hispanic Black, male									
2003.....	819.8	1,251.2	1,722.5	2,493.7	3,390.4	4,907.6	7,066.5	9,941.1	15,082.2
2002.....	818.5	1,229.3	1,751.3	2,503.4	3,402.3	4,983.1	7,157.7	10,054.7	15,776.9
2001.....	846.0	1,222.7	1,746.2	2,456.8	3,547.9	5,105.6	7,313.8	10,346.7	16,576.2
2000.....	869.4	1,248.2	1,802.9	2,498.8	3,559.1	5,258.8	7,462.1	10,491.3	16,944.8
1999.....	902.1	1,238.5	1,851.6	2,583.0	3,644.4	5,379.6	7,591.6	10,490.8	17,385.3
Non-Hispanic Black, female									
2003.....	498.1	712.7	992.0	1,474.3	2,084.8	3,071.6	4,554.5	6,735.8	13,730.3
2002.....	501.4	702.1	1,014.9	1,448.4	2,128.3	3,136.6	4,718.0	6,986.6	13,855.4
2001.....	493.1	710.9	1,028.9	1,498.8	2,209.4	3,182.0	4,756.8	6,952.7	13,951.4
2000.....	515.5	708.3	1,013.7	1,534.7	2,256.3	3,272.3	4,857.7	7,099.0	14,054.9
1999.....	491.7	697.1	1,073.0	1,539.8	2,263.4	3,402.8	4,925.0	7,099.0	13,887.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Certain other intestinal infections (A04,A07-A09)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	*	0.2	0.5	0.9	1.7	3.4	7.2	12.7	25.2
2002.....	0.1	0.2	0.5	0.7	1.4	2.7	5.1	9.8	21.4
2001.....	*	0.2	0.2	0.5	0.8	1.9	3.2	6.4	12.8
2000.....	*	*	0.2	0.3	0.9	1.6	2.8	5.3	11.9
1999.....	*	*	*	0.3	0.7	1.3	2.3	4.7	10.4

Non-Hispanic 4/, male

2003.....	*	*	0.5	0.8	1.5	3.2	7.0	13.3	21.4
2002.....	*	*	0.4	0.7	1.6	3.2	5.1	10.2	21.0
2001.....	*	*	*	0.4	0.8	2.0	3.5	6.4	12.0
2000.....	*	*	*	*	1.0	1.5	2.9	5.9	11.5
1999.....	*	*	*	*	0.6	1.2	2.7	5.1	9.6

Non-Hispanic 4/, female

2003.....	*	0.2	0.5	0.9	1.9	3.6	7.3	12.3	26.8
2002.....	*	*	0.5	0.6	1.3	2.4	5.0	9.6	21.6
2001.....	*	0.2	*	0.6	0.8	1.8	3.0	6.4	13.2
2000.....	*	*	*	*	0.8	1.6	2.7	5.0	12.1
1999.....	*	*	*	*	0.8	1.3	2.1	4.4	10.8

Non-Hispanic White, both sexes

2003.....	*	0.2	0.5	0.8	1.8	3.7	7.3	13.6	26.5
2002.....	0.1	0.2	0.4	0.6	1.6	2.8	5.4	10.4	22.5
2001.....	*	0.2	0.2	0.5	0.7	2.0	3.4	6.6	13.6
2000.....	*	*	0.2	0.3	0.9	1.6	2.9	5.7	12.3
1999.....	*	*	*	0.3	0.7	1.3	2.4	4.9	11.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Tuberculosis (A16-A19)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.2	0.3	0.3	0.3	0.5	0.7	1.3	2.0	2.2
2002.....	0.2	0.2	0.3	0.4	0.8	0.8	1.5	2.1	2.5
2001.....	0.2	0.2	0.4	0.4	0.6	1.0	1.3	1.9	2.7
2000.....	0.2	0.3	0.2	0.4	0.7	0.9	1.6	2.3	2.5
1999.....	0.2	0.3	0.5	0.5	0.7	1.1	1.6	2.2	3.9

Non-Hispanic 4/, male

2003.....	0.4	0.4	0.5	0.4	0.6	0.9	1.9	2.6	3.0
2002.....	*	0.3	0.5	0.6	1.1	1.2	1.8	3.0	3.8
2001.....	0.3	0.3	0.6	0.7	0.8	1.4	1.7	2.2	4.3
2000.....	0.3	0.5	0.4	0.6	1.0	1.0	2.0	3.6	4.3
1999.....	0.4	0.4	0.7	0.8	1.0	1.6	2.1	3.5	6.8

Non-Hispanic 4/, female

2003.....	*	*	*	*	*	0.5	1.0	1.6	1.8
2002.....	*	*	*	*	0.5	0.6	1.4	1.5	1.9
2001.....	*	*	*	*	*	0.7	1.0	1.7	2.1
2000.....	*	*	*	*	0.5	0.9	1.4	1.6	1.8
1999.....	*	*	*	*	*	0.7	1.3	1.5	2.8

Non-Hispanic White, both sexes

2003.....	*	*	0.2	*	*	0.6	0.9	1.6	1.7
2002.....	*	*	*	0.3	0.4	0.5	1.2	1.6	1.9
2001.....	*	*	0.2	0.3	0.4	0.6	1.0	1.3	2.0
2000.....	*	0.2	*	*	0.4	0.5	1.0	1.6	2.0
1999.....	*	*	0.3	0.3	0.5	0.6	1.1	1.7	3.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Respiratory tuberculosis (A16)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.2	0.2	0.3	0.2	0.4	0.5	1.1	1.5	1.7
2002.....	0.1	0.2	0.2	0.3	0.6	0.6	1.2	1.8	2.2
2001.....	0.1	0.2	0.3	0.4	0.4	0.7	1.1	1.5	2.4
2000.....	0.2	0.2	0.2	0.3	0.5	0.7	1.3	2.0	2.2
1999.....	0.2	0.2	0.4	0.5	0.6	0.8	1.4	2.0	3.5

Non-Hispanic 4/, male

2003.....	0.3	0.3	0.4	*	0.5	0.8	1.6	2.0	2.4
2002.....	*	0.3	0.4	0.5	0.9	0.9	1.3	2.5	3.3
2001.....	*	0.3	0.5	0.7	0.6	1.0	1.4	1.7	3.8
2000.....	0.3	0.4	0.3	0.5	0.8	0.8	1.7	3.0	3.8
1999.....	0.3	0.3	0.6	0.8	0.9	1.3	1.8	3.2	6.1

Non-Hispanic 4/, female

2003.....	*	*	*	*	*	*	0.8	1.2	1.5
2002.....	*	*	*	*	*	*	1.1	1.4	1.7
2001.....	*	*	*	*	*	0.5	0.8	1.4	1.8
2000.....	*	*	*	*	*	0.6	1.1	1.4	1.5
1999.....	*	*	*	*	*	0.5	1.1	1.3	2.4

Non-Hispanic White, both sexes

2003.....	*	*	0.2	*	*	0.4	0.8	1.2	1.3
2002.....	*	*	*	*	0.3	0.4	0.9	1.3	1.6
2001.....	*	*	0.2	0.2	0.3	0.4	0.8	1.0	1.7
2000.....	*	0.2	*	*	0.3	0.4	0.9	1.3	1.8
1999.....	*	*	0.2	0.3	0.4	0.5	0.9	1.5	2.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)									
All origins 3/, both sexes									
2003.....	4.2	6.5	10.3	16.7	25.2	40.9	68.4	107.8	202.5
2002.....	4.2	6.2	10.0	16.0	27.0	43.1	70.0	109.4	203.0
2001.....	4.0	6.2	9.8	15.5	25.3	40.9	67.3	103.9	205.9
2000.....	3.9	6.1	9.4	15.2	24.2	38.4	64.0	105.0	215.7
1999.....	3.7	5.6	9.0	14.4	24.7	38.3	63.8	102.9	220.7
All origins 3/, male									
2003.....	4.6	7.1	11.3	17.7	28.7	45.1	76.7	122.5	213.8
2002.....	4.7	6.8	10.6	17.8	29.1	48.2	80.6	127.0	213.9
2001.....	4.3	6.7	10.6	16.6	27.8	46.7	78.6	122.3	224.3
2000.....	4.4	7.0	10.1	16.7	26.9	43.2	73.9	121.7	223.0
1999.....	4.2	6.2	9.7	15.7	27.7	43.7	74.8	120.6	233.5
All origins 3/, female									
2003.....	3.8	6.0	9.4	15.7	22.2	37.5	62.5	98.8	197.5
2002.....	3.8	5.6	9.5	14.5	25.2	39.0	62.5	98.7	198.3
2001.....	3.8	5.7	9.1	14.4	23.1	36.2	59.5	92.9	198.3
2000.....	3.4	5.2	8.7	13.8	21.8	34.6	57.1	95.2	212.7
1999.....	3.3	5.1	8.4	13.1	22.1	34.1	56.2	92.5	215.5
Hispanic, both sexes									
2003.....	3.4	4.5	7.4	11.5	19.0	26.0	50.2	81.2	151.8
2002.....	2.7	5.1	6.2	11.2	20.1	39.9	54.9	78.4	156.3
2001.....	3.6	4.3	5.4	11.8	20.5	31.5	54.1	77.0	171.3
2000.....	2.9	5.2	6.4	11.2	16.2	28.9	48.1	79.6	144.7
1999.....	2.1	4.4	6.9	9.4	20.6	31.9	52.3	78.7	180.3
Hispanic, male									
2003.....	4.4	5.7	9.1	10.3	22.7	28.6	56.5	85.2	160.1
2002.....	3.8	6.9	6.4	12.6	20.7	43.9	61.4	91.5	164.0
2001.....	3.8	6.0	6.4	14.7	24.5	36.6	68.9	89.6	160.0
2000.....	4.1	6.3	8.1	12.4	22.4	32.1	58.3	94.2	157.2
1999.....	2.6	5.4	8.6	12.2	25.9	34.1	70.2	84.9	181.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)									
Hispanic, female									
2003.....	2.4	3.3	5.9	12.4	16.0	23.9	45.7	78.5	147.7
2002.....	*	3.2	6.1	10.1	19.6	36.8	50.3	70.1	152.5
2001.....	3.4	2.6	4.5	9.4	17.2	27.6	43.6	69.2	176.9
2000.....	*	4.2	4.8	10.2	11.2	26.5	40.8	70.8	138.5
1999.....	*	3.5	5.4	6.9	16.3	30.3	39.6	75.0	179.9
Non-Hispanic 4/, both sexes									
2003.....	4.3	6.7	10.5	17.0	25.6	41.7	69.3	108.7	204.1
2002.....	4.4	6.3	10.3	16.4	27.4	43.2	70.6	110.5	204.2
2001.....	4.1	6.3	10.1	15.6	25.6	41.3	67.8	104.7	206.8
2000.....	4.0	6.1	9.5	15.4	24.6	38.8	64.4	105.6	217.5
1999.....	3.9	5.7	9.1	14.7	24.8	38.5	64.1	103.4	221.2
Non-Hispanic 4/, male									
2003.....	4.7	7.2	11.4	18.2	29.0	46.0	77.7	124.0	215.9
2002.....	4.8	6.8	10.8	18.1	29.5	48.3	81.2	128.1	215.6
2001.....	4.3	6.7	10.9	16.6	28.0	47.1	78.8	123.1	226.8
2000.....	4.5	7.0	10.2	16.9	26.9	43.6	74.3	122.0	224.8
1999.....	4.3	6.2	9.7	16.0	27.7	44.1	74.5	121.5	234.5
Non-Hispanic 4/, female									
2003.....	3.9	6.2	9.7	15.9	22.6	38.3	63.3	99.4	199.0
2002.....	4.0	5.8	9.7	14.8	25.5	39.1	63.0	99.7	199.3
2001.....	3.8	6.0	9.3	14.8	23.5	36.6	60.2	93.6	198.5
2000.....	3.5	5.2	9.0	14.0	22.5	35.0	57.6	95.9	214.5
1999.....	3.4	5.2	8.5	13.5	22.3	34.1	56.8	92.9	215.9
Non-Hispanic White, both sexes									
2003.....	3.4	5.2	8.8	14.5	21.8	37.5	64.3	103.1	192.4
2002.....	3.6	4.9	8.5	13.6	23.7	38.9	64.6	104.0	192.6
2001.....	3.1	5.0	8.2	13.2	22.3	36.9	62.4	99.1	195.6
2000.....	3.0	4.8	7.9	12.7	20.9	34.8	58.2	97.7	204.3
1999.....	2.8	4.5	7.2	12.3	21.2	33.5	57.6	95.7	206.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)									
Non-Hispanic White, male									
2003.....	3.8	5.6	9.3	15.6	24.6	41.5	71.9	118.1	204.8
2002.....	4.0	5.2	9.1	15.4	25.2	43.8	73.6	119.9	203.9
2001.....	3.3	5.2	8.6	14.2	24.2	41.6	72.4	116.6	213.6
2000.....	3.4	5.3	8.4	14.1	22.9	39.7	66.7	114.0	211.2
1999.....	3.2	4.9	7.5	12.8	23.4	38.3	67.8	112.1	218.1
Non-Hispanic White, female									
2003.....	3.1	4.8	8.4	13.4	19.4	34.2	58.8	93.9	187.0
2002.....	3.3	4.6	7.9	11.9	22.3	34.9	58.2	94.2	187.7
2001.....	2.9	4.7	7.9	12.3	20.6	33.1	55.2	88.6	188.1
2000.....	2.5	4.2	7.3	11.4	19.1	30.8	52.1	88.1	201.6
1999.....	2.5	4.2	6.9	11.8	19.4	29.7	50.4	86.1	201.2
Non-Hispanic Black, both sexes									
2003.....	10.8	18.0	25.8	42.2	62.0	89.2	135.4	196.5	373.9
2002.....	9.6	17.4	27.6	41.8	63.3	92.0	148.3	210.3	370.8
2001.....	10.9	17.1	27.6	38.9	58.2	90.9	141.2	194.2	368.6
2000.....	11.3	17.3	24.7	40.5	60.8	86.7	148.0	218.1	395.9
1999.....	11.3	15.5	25.8	37.0	58.8	95.6	143.9	211.6	414.2
Non-Hispanic Black, male									
2003.....	12.1	20.3	32.7	46.8	76.2	101.1	162.6	237.7	409.1
2002.....	10.5	20.6	29.3	46.2	73.2	106.7	185.4	260.5	399.3
2001.....	12.0	19.1	34.0	41.8	67.5	112.6	173.0	237.9	430.8
2000.....	12.8	21.9	27.9	45.9	69.8	96.7	185.0	251.5	432.1
1999.....	13.2	18.1	30.1	47.7	72.8	116.0	165.8	270.1	451.6
Non-Hispanic Black, female									
2003.....	9.6	15.9	20.2	38.5	51.4	81.1	119.3	174.9	360.3
2002.....	8.9	14.8	26.3	38.3	56.0	82.1	126.1	184.2	359.9
2001.....	10.0	15.5	22.4	36.6	51.4	76.4	122.2	172.0	345.6
2000.....	10.0	13.4	22.0	36.2	54.1	80.1	125.7	201.3	382.5
1999.....	9.6	13.3	22.2	28.7	48.4	82.0	130.8	182.6	400.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Viral hepatitis (B15-B19)

All origins 3/, both sexes

2003.....	4.8	6.3	4.5	3.1	3.4	4.3	4.3	4.1	2.4
2002.....	5.6	6.1	4.3	3.7	3.9	4.5	4.7	4.4	3.9
2001.....	5.3	5.2	4.1	3.8	4.3	4.9	5.1	4.6	3.8
2000.....	5.2	4.8	3.7	3.8	4.3	4.9	5.4	5.2	4.0
1999.....	4.4	3.8	3.3	3.3	4.5	5.1	5.3	5.4	4.7

All origins 3/, male

2003.....	7.1	9.7	6.6	3.9	3.7	4.8	4.7	5.0	2.8
2002.....	8.5	9.5	6.2	4.6	4.3	4.7	5.7	4.8	5.0
2001.....	8.1	8.2	5.9	4.5	4.7	4.9	5.1	5.5	4.4
2000.....	8.0	7.4	5.2	4.7	4.6	5.0	6.5	6.2	5.7
1999.....	6.7	5.9	4.7	3.8	4.9	5.4	6.1	7.3	6.9

All origins 3/, female

2003.....	2.6	3.1	2.5	2.4	3.1	3.9	4.0	3.5	2.3
2002.....	2.8	2.8	2.5	2.9	3.4	4.4	4.1	4.2	3.3
2001.....	2.5	2.4	2.5	3.3	3.9	4.9	5.0	4.1	3.5
2000.....	2.5	2.4	2.2	3.1	4.1	4.9	4.6	4.7	3.3
1999.....	2.2	1.7	2.1	2.8	4.1	4.9	4.7	4.2	3.9

Hispanic, both sexes

2003.....	5.7	8.6	7.7	6.6	8.6	8.7	9.8	8.2	*
2002.....	6.8	9.4	8.1	8.6	8.8	8.4	11.5	8.9	*
2001.....	8.1	7.5	11.1	9.5	8.2	12.1	12.2	*	*
2000.....	6.7	7.3	7.0	10.7	10.3	12.2	12.9	13.4	*
1999.....	5.9	8.1	7.6	8.0	11.7	13.3	13.6	13.5	*

Hispanic, male

2003.....	8.8	13.5	11.5	8.9	9.1	*	*	*	*
2002.....	10.8	13.5	11.9	11.0	7.4	*	12.8	*	*
2001.....	12.4	12.1	17.0	10.3	*	*	*	*	*
2000.....	10.3	11.1	9.4	14.7	10.4	11.7	*	*	*
1999.....	9.1	12.3	9.9	9.5	12.4	13.7	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)									
Hispanic, female									
2003.....	2.5	3.8	4.1	4.7	8.3	10.8	10.8	*	*
2002.....	2.8	5.5	4.6	6.5	10.0	8.8	10.5	*	*
2001.....	3.8	3.2	5.8	8.7	10.0	15.0	13.6	*	*
2000.....	3.2	3.6	4.8	7.2	10.3	12.5	13.1	*	*
1999.....	2.7	4.1	5.4	6.7	11.3	13.0	15.4	*	*
Non-Hispanic 4/, both sexes									
2003.....	4.7	6.1	4.2	2.8	3.0	3.9	4.0	3.9	2.3
2002.....	5.4	5.7	3.9	3.3	3.5	4.2	4.4	4.2	3.7
2001.....	5.0	5.0	3.5	3.4	4.0	4.4	4.7	4.4	3.6
2000.....	5.0	4.6	3.4	3.3	3.9	4.5	5.1	4.9	3.8
1999.....	4.3	3.4	3.0	2.9	4.0	4.7	4.9	5.1	4.7
Non-Hispanic 4/, male									
2003.....	6.9	9.3	6.2	3.5	3.3	4.7	4.5	4.9	2.7
2002.....	8.1	9.1	5.6	4.1	4.1	4.5	5.3	4.7	4.8
2001.....	7.7	7.8	5.0	4.0	4.6	4.7	4.8	5.3	4.3
2000.....	7.7	7.0	4.8	3.9	4.2	4.6	6.2	5.9	5.4
1999.....	6.5	5.3	4.3	3.4	4.4	4.9	5.9	7.4	6.8
Non-Hispanic 4/, female									
2003.....	2.6	3.0	2.3	2.2	2.7	3.4	3.7	3.2	2.2
2002.....	2.8	2.6	2.3	2.6	2.9	4.1	3.8	3.9	3.2
2001.....	2.4	2.3	2.2	2.8	3.4	4.3	4.6	3.9	3.3
2000.....	2.4	2.3	2.0	2.7	3.6	4.4	4.3	4.3	3.2
1999.....	2.1	1.5	1.8	2.5	3.6	4.5	4.2	3.7	3.9
Non-Hispanic White, both sexes									
2003.....	4.7	5.7	3.4	2.3	2.4	3.5	3.6	3.6	2.2
2002.....	5.2	5.4	3.3	2.7	2.9	3.8	4.1	4.0	3.5
2001.....	4.8	4.4	3.0	2.9	3.4	3.9	4.4	4.2	3.3
2000.....	4.7	3.9	2.7	2.7	3.3	4.1	4.9	4.6	3.7
1999.....	3.9	2.9	2.4	2.5	3.4	4.1	4.5	5.0	4.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Viral hepatitis (B15-B19)

Non-Hispanic White, male

2003.....	6.9	8.6	5.1	2.9	2.8	4.1	3.9	4.7	2.7
2002.....	7.9	8.5	4.7	3.3	3.7	4.1	5.1	4.7	4.6
2001.....	7.4	7.0	4.3	3.6	4.3	4.4	4.9	5.4	4.4
2000.....	7.3	6.1	3.9	3.3	3.6	4.2	6.0	5.6	5.4
1999.....	5.7	4.5	3.6	3.1	4.0	4.6	5.6	7.7	6.5

Non-Hispanic White, female

2003.....	2.5	2.8	1.8	1.7	2.1	2.9	3.4	2.9	2.0
2002.....	2.6	2.4	2.0	2.2	2.2	3.5	3.4	3.6	3.1
2001.....	2.2	1.8	1.7	2.3	2.7	3.4	4.1	3.5	2.9
2000.....	2.1	1.8	1.5	2.3	3.2	4.0	4.0	4.0	3.1
1999.....	2.1	1.3	1.4	1.9	2.9	3.7	3.8	3.4	3.7

Non-Hispanic Black, both sexes

2003.....	5.3	10.3	9.6	6.4	5.2	5.6	6.3	5.4	*
2002.....	7.3	9.2	7.8	6.3	6.1	5.8	6.0	*	*
2001.....	6.8	9.6	7.3	6.2	5.1	6.2	6.2	*	*
2000.....	7.4	8.8	6.9	4.5	5.3	6.3	5.8	7.2	*
1999.....	7.2	6.8	5.7	4.3	6.7	6.9	6.5	*	*

Non-Hispanic Black, male

2003.....	7.8	16.5	15.3	7.7	5.6	*	*	*	*
2002.....	10.6	15.1	12.6	8.1	7.9	*	*	*	*
2001.....	10.1	14.3	10.1	7.5	*	*	*	*	*
2000.....	11.2	13.8	10.3	5.3	6.9	7.5	*	*	*
1999.....	12.2	11.4	7.7	4.6	5.6	*	*	*	*

Non-Hispanic Black, female

2003.....	3.2	5.1	4.9	5.4	4.9	5.0	5.5	*	*
2002.....	4.5	4.3	3.9	4.9	4.7	5.7	5.6	*	*
2001.....	3.9	5.6	4.9	5.1	5.4	6.7	7.4	*	*
2000.....	4.2	4.5	4.0	3.9	4.1	5.4	*	*	*
1999.....	2.9	2.9	4.1	4.1	7.5	7.6	6.7	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All origins 3/, both sexes

2003.....	12.1	9.5	6.4	4.1	2.8	1.9	1.0	0.4	*
2002.....	12.6	9.6	5.9	4.0	2.9	1.4	1.0	0.5	*
2001.....	12.1	8.7	5.8	4.5	2.7	1.4	0.9	0.4	*
2000.....	12.7	9.0	5.9	4.1	2.6	1.7	0.9	*	*
1999.....	12.4	8.7	5.9	3.5	2.7	1.5	0.7	*	*

All origins 3/, male

2003.....	18.8	15.3	10.7	6.9	4.6	3.3	1.8	1.0	*
2002.....	19.4	15.7	9.8	6.8	5.0	2.6	1.7	*	*
2001.....	19.0	14.4	9.8	7.1	4.2	2.7	1.8	*	*
2000.....	20.3	15.0	10.1	7.0	4.5	2.9	1.8	*	*
1999.....	20.0	14.5	10.2	5.9	4.8	2.7	1.3	*	*

All origins 3/, female

2003.....	5.6	3.9	2.5	1.6	1.3	0.7	*	*	*
2002.....	5.9	3.6	2.3	1.5	1.0	0.5	*	*	*
2001.....	5.4	3.3	2.0	2.0	1.4	0.4	*	*	*
2000.....	5.3	3.4	2.0	1.5	1.0	0.7	*	*	*
1999.....	5.0	3.1	1.9	1.2	1.0	0.6	*	*	*

Hispanic, both sexes

2003.....	17.0	14.4	12.3	6.2	6.2	4.1	*	*	*
2002.....	15.4	13.7	10.8	7.8	4.3	*	*	*	*
2001.....	16.4	14.2	10.5	7.3	6.0	*	*	*	*
2000.....	18.1	15.4	10.0	8.9	4.7	*	*	*	*
1999.....	17.1	13.1	9.8	6.6	6.0	*	*	*	*

Hispanic, male

2003.....	27.9	23.7	20.2	8.6	10.1	*	*	*	*
2002.....	24.0	21.8	17.4	11.5	8.5	*	*	*	*
2001.....	25.0	24.8	17.6	11.1	11.3	*	*	*	*
2000.....	28.4	26.2	17.3	12.7	8.2	*	*	*	*
1999.....	26.8	20.8	17.6	12.5	10.8	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Hispanic, female

2003.....	5.8	5.4	5.1	*	*	*	*	*	*
2002.....	6.7	5.9	4.7	4.5	*	*	*	*	*
2001.....	7.8	4.1	4.2	*	*	*	*	*	*
2000.....	7.9	5.0	*	5.7	*	*	*	*	*
1999.....	7.5	5.8	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	11.5	8.9	5.9	3.9	2.5	1.7	0.9	0.4	*
2002.....	12.1	9.0	5.5	3.6	2.7	1.4	0.9	0.4	*
2001.....	11.5	8.1	5.3	4.1	2.4	1.3	0.8	*	*
2000.....	12.0	8.4	5.5	3.7	2.4	1.6	0.8	*	*
1999.....	11.7	8.1	5.5	3.1	2.4	1.4	0.7	*	*

Non-Hispanic 4/, male

2003.....	17.5	14.3	9.7	6.7	4.1	3.0	1.7	*	*
2002.....	18.6	14.9	9.1	6.3	4.8	2.4	1.7	*	*
2001.....	18.0	13.2	9.1	6.6	3.6	2.5	1.6	*	*
2000.....	19.2	13.8	9.3	6.5	4.1	2.7	1.6	*	*
1999.....	18.9	13.6	9.5	5.2	4.3	2.5	1.2	*	*

Non-Hispanic 4/, female

2003.....	5.6	3.7	2.2	1.4	1.2	0.7	*	*	*
2002.....	5.8	3.4	2.0	1.2	1.0	0.6	*	*	*
2001.....	5.1	3.2	1.8	1.8	1.4	*	*	*	*
2000.....	5.1	3.2	1.9	1.2	0.9	0.7	*	*	*
1999.....	4.8	2.8	1.9	1.2	0.9	0.6	*	*	*

Non-Hispanic White, both sexes

2003.....	5.2	3.7	2.6	1.7	0.9	0.8	0.5	*	*
2002.....	5.5	3.5	2.6	1.7	1.0	0.5	0.4	*	*
2001.....	5.0	3.4	2.6	1.5	0.8	0.6	0.3	*	*
2000.....	5.5	3.7	2.7	1.5	1.0	0.6	0.4	*	*
1999.....	5.2	3.9	2.4	1.4	1.1	0.5	0.3	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic White, male

2003.....	8.9	6.6	4.6	3.2	1.5	1.5	1.1	*	*
2002.....	9.5	6.3	4.7	3.2	1.9	1.0	*	*	*
2001.....	8.8	6.2	4.8	2.6	1.4	1.4	*	*	*
2000.....	9.6	6.5	4.9	2.8	2.0	0.9	0.8	*	*
1999.....	9.0	7.2	4.4	2.6	2.1	0.9	*	*	*

Non-Hispanic White, female

2003.....	1.6	0.9	0.7	*	*	*	*	*	*
2002.....	1.6	0.8	0.6	*	*	*	*	*	*
2001.....	1.2	0.6	0.6	0.4	*	*	*	*	*
2000.....	1.4	0.8	0.7	*	*	*	*	*	*
1999.....	1.3	0.8	0.5	*	*	*	*	*	*

Non-Hispanic Black, both sexes

2003.....	54.5	47.5	34.4	23.9	17.1	10.9	4.9	*	*
2002.....	58.4	50.5	30.8	20.5	17.8	10.6	7.2	*	*
2001.....	57.6	45.0	29.3	26.3	16.2	8.5	6.4	*	*
2000.....	59.5	46.9	29.6	22.2	14.7	11.5	6.0	*	*
1999.....	59.6	42.8	33.0	18.4	14.9	11.2	4.9	*	*

Non-Hispanic Black, male

2003.....	81.2	76.3	58.5	41.7	29.6	19.5	*	*	*
2002.....	87.2	84.9	50.9	36.7	32.0	19.5	15.0	*	*
2001.....	88.7	72.3	50.7	43.7	25.8	15.5	12.8	*	*
2000.....	93.8	77.5	51.3	40.2	25.3	21.9	11.6	*	*
1999.....	96.6	71.1	57.7	30.5	26.3	21.1	*	*	*

Non-Hispanic Black, female

2003.....	31.3	23.2	14.6	10.0	7.9	5.0	*	*	*
2002.....	33.4	21.4	14.2	8.0	7.2	4.6	*	*	*
2001.....	30.6	21.7	11.7	12.7	9.0	*	*	*	*
2000.....	29.8	20.8	11.8	8.1	6.9	4.5	*	*	*
1999.....	27.7	18.8	12.8	8.9	6.5	4.6	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All origins 3/, both sexes

2003.....	2.4	3.5	3.7	4.5	5.9	8.9	12.3	14.8	22.5
2002.....	1.9	2.7	3.0	3.8	5.4	8.1	11.6	13.8	21.7
2001.....	1.5	1.9	2.4	3.9	5.6	6.9	9.4	12.3	20.1
2000.....	1.4	1.8	2.4	3.6	4.7	7.0	10.0	12.9	21.2
1999.....	2.0	2.6	3.3	4.4	6.8	8.9	12.6	16.6	27.0

All origins 3/, male

2003.....	3.3	4.7	4.9	5.4	7.0	10.3	13.9	18.8	26.5
2002.....	2.6	3.8	3.8	4.7	6.5	9.7	14.0	17.7	26.8
2001.....	2.1	2.4	3.0	4.5	6.2	7.5	11.4	15.9	23.6
2000.....	1.7	2.3	2.8	4.0	5.7	8.8	12.1	16.2	27.6
1999.....	2.4	3.0	3.5	5.2	8.0	10.5	15.3	20.4	31.9

All origins 3/, female

2003.....	1.6	2.2	2.6	3.7	5.0	7.8	11.1	12.4	20.8
2002.....	1.2	1.7	2.2	3.0	4.5	6.8	9.9	11.5	19.5
2001.....	0.9	1.5	1.8	3.3	5.1	6.5	8.0	10.1	18.7
2000.....	1.1	1.3	2.1	3.2	3.9	5.6	8.5	10.9	18.6
1999.....	1.6	2.3	3.1	3.7	5.9	7.7	10.8	14.4	25.1

Hispanic, both sexes

2003.....	3.3	5.9	4.7	6.8	8.4	8.5	15.1	15.6	13.0
2002.....	2.4	4.9	4.6	3.9	6.2	6.5	8.8	15.6	17.7
2001.....	2.1	2.0	3.2	2.8	4.9	7.1	10.5	9.9	15.0
2000.....	1.7	2.3	3.1	3.2	4.5	7.3	9.5	14.5	19.2
1999.....	2.0	2.7	3.0	4.8	8.1	7.2	13.9	14.7	26.0

Hispanic, male

2003.....	4.9	9.4	7.5	7.9	12.0	*	16.4	*	*
2002.....	3.4	7.6	6.2	*	8.1	*	*	*	*
2001.....	2.7	2.9	5.0	*	*	*	*	*	*
2000.....	*	*	4.6	*	*	*	*	*	*
1999.....	*	3.7	*	6.0	9.3	*	16.4	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Hispanic, female

2003.....	*	2.6	*	5.8	5.3	9.8	14.2	14.9	*
2002.....	*	*	*	*	*	*	*	14.6	*
2001.....	*	*	*	*	*	*	9.7	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	7.2	8.0	12.1	*	28.1

Non-Hispanic 4/, both sexes

2003.....	2.3	3.2	3.6	4.3	5.7	8.9	12.1	14.8	22.9
2002.....	1.8	2.5	2.8	3.8	5.4	8.2	11.7	13.7	21.8
2001.....	1.4	1.9	2.3	4.0	5.7	6.9	9.3	12.4	20.3
2000.....	1.3	1.7	2.3	3.6	4.7	7.0	9.9	12.8	21.2
1999.....	2.0	2.6	3.3	4.4	6.7	9.0	12.5	16.7	27.0

Non-Hispanic 4/, male

2003.....	3.1	4.3	4.7	5.2	6.6	10.5	13.7	18.9	27.0
2002.....	2.6	3.4	3.6	4.7	6.4	9.8	14.1	17.7	26.8
2001.....	2.0	2.3	2.9	4.6	6.2	7.3	11.4	16.0	24.0
2000.....	1.6	2.2	2.6	4.0	5.9	8.8	12.1	15.9	27.8
1999.....	2.4	2.9	3.5	5.1	7.9	10.7	15.2	20.6	32.3

Non-Hispanic 4/, female

2003.....	1.6	2.2	2.7	3.5	5.0	7.6	10.9	12.2	21.2
2002.....	1.1	1.6	2.1	3.0	4.5	6.9	10.0	11.3	19.7
2001.....	0.9	1.5	1.8	3.4	5.2	6.5	7.9	10.2	18.8
2000.....	1.1	1.2	2.1	3.2	3.8	5.6	8.4	10.9	18.6
1999.....	1.6	2.3	3.0	3.7	5.7	7.6	10.7	14.4	24.9

Non-Hispanic White, both sexes

2003.....	2.2	3.0	3.3	4.0	5.6	8.9	11.5	14.6	23.4
2002.....	1.7	2.2	2.4	3.5	5.3	8.0	11.4	13.7	22.2
2001.....	1.3	1.7	2.0	3.7	5.5	6.6	9.2	12.2	20.2
2000.....	1.2	1.5	2.2	3.2	4.6	6.8	9.8	12.6	21.0
1999.....	1.5	2.1	2.9	3.9	6.3	8.6	11.9	16.1	26.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic White, male

2003.....	3.0	3.9	4.3	4.8	6.4	10.2	13.2	18.6	27.6
2002.....	2.4	3.0	2.9	4.3	6.4	9.6	13.8	17.5	27.2
2001.....	1.8	2.1	2.5	4.3	5.9	7.1	11.3	16.0	24.2
2000.....	1.5	1.9	2.5	3.6	5.6	8.6	11.6	15.8	27.0
1999.....	2.0	2.4	3.3	4.4	7.3	10.4	14.7	18.9	30.8

Non-Hispanic White, female

2003.....	1.4	2.0	2.4	3.3	4.9	7.8	10.3	12.2	21.6
2002.....	1.0	1.4	1.9	2.8	4.3	6.7	9.8	11.4	20.0
2001.....	0.7	1.4	1.5	3.1	5.1	6.2	7.7	10.0	18.6
2000.....	1.0	1.0	1.9	2.8	3.7	5.3	8.5	10.8	18.6
1999.....	1.1	1.9	2.5	3.4	5.4	7.2	10.0	14.4	24.7

Non-Hispanic Black, both sexes

2003.....	3.6	5.3	6.1	6.3	5.8	8.4	16.5	16.1	19.9
2002.....	3.2	4.9	6.2	5.5	5.4	9.9	15.3	11.5	21.8
2001.....	2.6	3.5	4.5	6.2	7.1	9.8	9.9	13.5	20.9
2000.....	2.7	3.5	4.0	6.0	6.2	10.1	11.6	16.1	24.2
1999.....	5.2	6.0	6.5	8.2	10.9	13.3	19.1	23.8	34.3

Non-Hispanic Black, male

2003.....	4.7	7.1	7.8	8.5	8.6	13.1	16.8	22.6	23.4
2002.....	4.5	7.0	9.5	7.5	6.7	12.3	18.3	*	28.4
2001.....	3.7	4.4	5.5	7.5	9.6	10.0	10.4	16.4	*
2000.....	3.3	4.8	4.6	7.4	8.8	12.6	16.9	19.7	31.7
1999.....	6.0	6.6	5.9	10.4	12.5	16.3	22.2	42.5	46.4

Non-Hispanic Black, female

2003.....	2.6	3.7	4.6	4.5	3.8	5.2	16.4	12.6	18.6
2002.....	2.0	3.2	3.4	4.0	4.6	8.3	13.5	9.7	19.2
2001.....	1.6	2.8	3.7	5.1	5.2	9.6	9.6	12.1	20.5
2000.....	2.1	2.3	3.5	4.9	4.3	8.4	8.4	14.3	21.4
1999.....	4.5	5.5	7.0	6.5	9.6	11.3	17.2	14.5	29.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)									
All origins 3/, both sexes									
2003.....	89.8	159.1	267.9	440.9	647.9	909.2	1,195.4	1,450.0	1,698.2
2002.....	90.6	161.5	276.8	447.0	663.8	933.6	1,207.0	1,458.3	1,723.9
2001.....	92.4	164.8	280.7	453.3	673.2	943.5	1,211.7	1,465.9	1,765.6
2000.....	93.6	166.2	289.0	463.4	690.6	951.6	1,222.5	1,505.3	1,819.4
1999.....	93.7	167.0	295.3	471.6	700.2	963.4	1,224.0	1,493.1	1,805.8
All origins 3/, male									
2003.....	90.6	169.8	295.9	507.6	774.4	1,117.9	1,533.3	1,939.7	2,413.8
2002.....	89.8	171.7	308.6	519.7	796.8	1,157.8	1,554.1	1,951.5	2,491.1
2001.....	91.7	174.2	311.6	526.7	814.1	1,178.0	1,568.4	1,978.7	2,613.6
2000.....	93.5	173.9	321.3	535.7	834.7	1,190.4	1,586.0	2,050.4	2,710.7
1999.....	94.7	174.9	330.6	550.1	848.2	1,212.9	1,600.1	2,046.0	2,722.2
All origins 3/, female									
2003.....	89.1	148.8	241.5	380.3	538.1	741.8	954.8	1,150.5	1,381.9
2002.....	91.4	151.8	246.8	381.1	548.9	754.4	961.9	1,158.4	1,391.1
2001.....	93.2	155.9	251.7	386.7	551.9	757.2	962.0	1,158.6	1,410.7
2000.....	93.7	158.8	258.9	398.0	567.2	763.5	969.3	1,183.8	1,456.4
1999.....	92.7	159.5	262.3	400.6	573.7	768.4	962.5	1,169.5	1,436.5
Hispanic, both sexes									
2003.....	60.7	107.1	177.1	267.0	408.6	582.3	819.1	1,005.4	1,234.9
2002.....	57.4	105.5	173.2	271.9	412.8	600.5	831.8	1,000.4	1,308.8
2001.....	61.7	110.3	172.3	291.3	413.5	616.9	846.0	1,033.1	1,363.7
2000.....	64.3	105.4	180.7	272.5	422.5	633.8	851.8	1,138.5	1,383.5
1999.....	65.0	110.9	179.3	282.8	448.9	611.2	834.8	1,094.3	1,399.6
Hispanic, male									
2003.....	58.9	111.1	194.1	321.9	509.5	759.7	1,064.0	1,334.8	1,668.6
2002.....	54.2	111.0	204.8	323.5	518.7	756.5	1,111.5	1,332.6	1,869.0
2001.....	60.1	109.9	189.4	340.6	519.1	802.4	1,171.7	1,423.2	1,967.4
2000.....	60.9	104.2	199.7	323.3	521.3	820.6	1,140.5	1,637.2	2,049.7
1999.....	64.4	111.7	208.3	345.2	567.6	802.5	1,120.3	1,520.0	1,945.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)									
Hispanic, female									
2003.....	62.4	103.1	161.4	219.2	325.7	446.1	645.4	791.1	1,014.8
2002.....	60.7	100.3	144.2	227.0	326.2	480.3	632.0	788.4	1,031.2
2001.....	63.3	110.7	156.7	248.7	328.0	475.3	614.9	791.4	1,068.6
2000.....	67.6	106.5	163.5	228.8	342.4	492.3	647.3	835.0	1,056.5
1999.....	65.7	110.1	153.2	228.9	352.8	467.3	633.2	840.3	1,136.5
Non-Hispanic 4/, both sexes									
2003.....	92.8	163.5	274.6	453.4	664.3	929.0	1,214.2	1,468.1	1,714.7
2002.....	93.8	166.2	284.5	459.2	680.4	952.6	1,224.0	1,474.9	1,736.8
2001.....	95.3	169.1	288.6	464.3	690.0	960.9	1,226.9	1,480.7	1,776.6
2000.....	96.1	170.7	296.5	476.3	706.7	967.3	1,236.6	1,515.9	1,831.1
1999.....	96.1	171.2	303.1	484.0	714.7	980.0	1,237.8	1,503.6	1,816.1
Non-Hispanic 4/, male									
2003.....	94.0	174.7	303.4	520.4	791.7	1,138.5	1,556.5	1,965.5	2,444.0
2002.....	93.3	176.8	316.2	532.8	814.2	1,180.0	1,573.7	1,974.9	2,513.2
2001.....	94.7	179.2	320.0	538.6	832.2	1,196.9	1,584.2	1,998.0	2,634.8
2000.....	96.4	179.1	329.4	549.5	852.4	1,207.8	1,602.6	2,061.5	2,732.4
1999.....	97.2	179.5	338.6	563.0	863.4	1,231.0	1,616.8	2,060.5	2,747.5
Non-Hispanic 4/, female									
2003.....	91.7	152.7	247.5	392.3	553.5	760.5	970.5	1,164.9	1,394.3
2002.....	94.3	156.0	254.6	392.3	564.4	770.5	977.1	1,171.5	1,401.8
2001.....	95.8	159.5	259.0	396.6	567.0	773.0	976.9	1,171.2	1,419.6
2000.....	95.8	162.7	265.6	409.7	581.3	777.5	981.8	1,194.4	1,466.6
1999.....	95.0	163.1	269.9	412.2	587.0	783.4	974.5	1,178.0	1,443.2
Non-Hispanic White, both sexes									
2003.....	87.3	155.1	267.4	447.1	661.9	928.9	1,213.9	1,468.8	1,717.8
2002.....	89.0	158.8	276.0	451.8	676.5	950.9	1,222.3	1,473.3	1,737.9
2001.....	90.2	162.2	279.7	458.2	682.7	955.9	1,222.7	1,473.0	1,775.2
2000.....	90.3	163.6	288.2	470.0	701.3	959.9	1,227.7	1,512.6	1,826.6
1999.....	90.4	163.5	293.3	474.6	707.7	971.9	1,226.4	1,498.7	1,808.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)									
Non-Hispanic White, male									
2003.....	87.9	163.8	292.1	506.7	780.1	1,128.6	1,541.2	1,955.2	2,441.7
2002.....	88.6	167.2	303.2	515.9	803.1	1,167.1	1,556.2	1,957.1	2,507.7
2001.....	89.1	169.4	305.9	525.6	815.0	1,178.6	1,562.9	1,968.7	2,616.8
2000.....	90.1	168.5	314.5	535.3	837.1	1,182.9	1,574.8	2,036.1	2,714.1
1999.....	89.9	168.3	323.3	544.9	846.1	1,209.4	1,586.9	2,037.9	2,721.3
Non-Hispanic White, female									
2003.....	86.7	146.5	243.7	391.6	556.9	764.8	977.1	1,169.9	1,399.1
2002.....	89.5	150.7	249.8	392.4	564.4	774.0	983.3	1,177.4	1,405.4
2001.....	91.3	155.1	254.6	395.8	565.8	775.0	981.8	1,173.8	1,425.1
2000.....	90.5	158.8	263.2	409.5	582.0	780.3	983.7	1,201.6	1,468.7
1999.....	91.0	158.8	264.6	409.6	586.3	782.7	973.5	1,180.6	1,444.2
Non-Hispanic Black, both sexes									
2003.....	141.5	248.2	390.9	600.8	816.7	1,097.7	1,405.0	1,662.0	1,867.5
2002.....	138.6	243.5	407.7	606.9	845.5	1,143.2	1,443.0	1,680.9	1,904.3
2001.....	141.4	246.5	412.8	598.1	871.7	1,172.3	1,446.2	1,735.8	1,947.5
2000.....	148.6	257.2	415.1	616.2	874.9	1,193.3	1,497.2	1,718.5	2,015.1
1999.....	147.7	260.2	437.3	644.3	891.9	1,206.6	1,516.9	1,715.6	2,024.0
Non-Hispanic Black, male									
2003.....	149.1	288.5	475.6	763.8	1,065.1	1,477.8	2,004.0	2,453.4	2,859.6
2002.....	140.8	277.8	494.5	790.8	1,094.5	1,559.5	2,071.8	2,539.6	3,008.4
2001.....	147.2	286.7	504.5	756.4	1,154.2	1,619.6	2,100.9	2,672.1	3,238.5
2000.....	155.3	299.9	521.3	783.3	1,154.5	1,675.4	2,183.1	2,709.5	3,290.1
1999.....	164.2	305.9	543.1	830.7	1,174.9	1,663.5	2,224.9	2,656.8	3,361.1
Non-Hispanic Black, female									
2003.....	134.9	214.2	321.3	473.8	632.4	841.5	1,049.3	1,247.9	1,483.2
2002.....	136.7	214.5	336.6	463.2	660.9	863.9	1,067.5	1,234.1	1,482.4
2001.....	136.3	212.4	337.8	473.7	663.1	874.2	1,055.5	1,258.9	1,469.3
2000.....	142.8	220.9	328.1	484.4	668.7	873.2	1,083.7	1,219.7	1,541.6
1999.....	133.4	221.4	351.1	498.2	682.1	903.4	1,090.3	1,249.0	1,524.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)									
All origins 3/, both sexes									
2003.....	2.0	3.6	5.5	7.9	9.7	10.8	12.8	13.6	18.6
2002.....	2.0	3.6	5.8	8.1	8.8	12.1	13.1	14.1	18.7
2001.....	2.1	3.5	6.1	7.9	9.4	11.6	13.5	15.2	18.5
2000.....	2.0	3.8	5.7	8.1	9.9	11.5	12.6	14.0	19.4
1999.....	2.3	3.6	5.8	8.1	9.3	11.3	13.5	15.4	19.3
All origins 3/, male									
2003.....	3.2	6.2	9.1	12.6	15.0	16.4	19.0	19.9	26.9
2002.....	3.2	5.8	9.4	12.9	13.9	18.1	19.2	20.3	26.9
2001.....	3.2	5.8	10.0	13.0	14.3	17.9	19.6	22.1	25.8
2000.....	3.1	6.0	9.5	12.7	15.2	16.9	18.7	20.2	27.1
1999.....	3.6	5.7	9.0	13.1	14.3	17.4	20.1	22.3	27.1
All origins 3/, female									
2003.....	0.9	1.1	2.1	3.6	5.2	6.3	8.4	9.7	15.0
2002.....	0.9	1.4	2.5	3.7	4.4	7.3	8.8	10.3	15.1
2001.....	1.0	1.4	2.4	3.3	5.1	6.6	9.3	11.1	15.4
2000.....	1.0	1.6	2.2	3.9	5.4	7.2	8.3	10.4	16.3
1999.....	1.0	1.6	2.7	3.6	5.1	6.6	9.0	11.3	16.1
Hispanic, both sexes									
2003.....	1.2	1.5	3.1	4.2	6.7	6.3	9.6	10.2	15.0
2002.....	*	2.0	3.3	3.3	4.6	6.1	9.1	8.9	16.1
2001.....	1.3	2.1	2.8	5.0	6.5	7.3	11.0	*	19.4
2000.....	*	2.0	4.2	4.7	6.7	6.9	8.3	11.7	18.6
1999.....	*	1.7	3.2	5.1	6.4	7.7	11.6	15.9	14.0
Hispanic, male									
2003.....	*	2.8	5.6	7.6	11.4	9.0	17.0	*	*
2002.....	*	3.2	5.9	5.4	7.8	12.3	16.0	*	*
2001.....	*	3.3	4.6	8.7	11.0	12.4	17.7	*	*
2000.....	*	3.2	7.5	8.1	11.9	12.6	*	*	40.3
1999.....	*	*	6.2	8.3	12.8	14.8	21.1	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	2.1	3.8	5.7	8.2	10.0	11.0	13.0	13.7	18.7
2002.....	2.1	3.7	6.0	8.4	9.1	12.4	13.3	14.3	18.7
2001.....	2.1	3.6	6.3	8.1	9.5	11.8	13.6	15.4	18.4
2000.....	2.1	3.9	5.8	8.3	10.0	11.7	12.8	14.1	19.5
1999.....	2.4	3.8	5.9	8.2	9.5	11.5	13.6	15.4	19.4

Non-Hispanic 4/, male

2003.....	3.4	6.5	9.3	13.0	15.2	16.8	19.1	20.2	27.1
2002.....	3.4	6.0	9.7	13.4	14.4	18.3	19.3	20.6	26.7
2001.....	3.3	6.0	10.4	13.3	14.5	18.2	19.5	22.4	25.8
2000.....	3.3	6.2	9.6	13.0	15.3	17.1	19.0	20.1	26.5
1999.....	3.7	6.0	9.1	13.4	14.3	17.5	20.0	22.7	27.3

Non-Hispanic 4/, female

2003.....	0.9	1.2	2.2	3.8	5.4	6.4	8.6	9.8	15.0
2002.....	1.0	1.5	2.6	3.8	4.6	7.7	9.1	10.5	15.3
2001.....	1.0	1.4	2.5	3.4	5.3	6.8	9.4	11.3	15.3
2000.....	1.0	1.7	2.2	4.0	5.5	7.5	8.4	10.5	16.6
1999.....	1.1	1.7	2.9	3.6	5.3	6.8	9.1	11.1	16.2

Non-Hispanic White, both sexes

2003.....	1.9	3.3	5.4	7.5	9.5	10.7	12.7	13.5	19.1
2002.....	1.8	3.2	5.5	7.8	8.6	12.7	13.5	14.4	18.9
2001.....	1.8	3.3	5.7	7.5	8.9	11.6	13.6	15.7	18.9
2000.....	1.8	3.3	5.1	7.4	9.5	11.3	12.7	14.1	19.8
1999.....	1.8	3.1	5.1	7.6	9.0	11.1	13.3	15.5	19.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)									
Non-Hispanic White, male									
2003.....	2.8	5.8	8.7	11.5	14.3	15.7	18.2	19.4	27.4
2002.....	2.9	5.1	8.8	12.2	13.3	18.4	19.5	20.5	26.8
2001.....	2.7	5.4	9.2	11.9	13.4	17.4	19.2	22.5	26.4
2000.....	2.8	5.0	8.3	11.4	14.1	16.1	18.3	19.7	27.4
1999.....	2.8	4.9	7.7	12.3	13.5	16.9	19.1	22.3	27.5
Non-Hispanic White, female									
2003.....	0.9	0.9	2.2	3.9	5.3	6.6	8.7	9.9	15.4
2002.....	0.8	1.3	2.3	3.7	4.5	8.0	9.1	10.7	15.5
2001.....	0.8	1.2	2.3	3.3	5.0	6.9	9.7	11.6	15.7
2000.....	0.9	1.6	2.0	3.7	5.5	7.4	8.7	10.7	16.8
1999.....	0.9	1.3	2.6	3.4	5.1	6.6	9.2	11.4	16.8
Non-Hispanic Black, both sexes									
2003.....	4.1	6.6	9.1	14.8	15.8	15.5	15.1	16.3	15.5
2002.....	4.3	7.3	10.4	15.0	13.5	12.2	11.6	12.8	16.0
2001.....	4.4	6.4	12.5	13.4	15.5	16.0	14.2	13.5	13.1
2000.....	4.1	8.5	12.3	15.6	15.0	16.5	14.5	16.1	14.9
1999.....	6.0	9.0	12.7	14.7	14.5	16.6	17.4	15.8	14.5
Non-Hispanic Black, male									
2003.....	7.3	11.6	16.8	28.6	28.1	30.3	27.1	30.9	25.6
2002.....	7.0	13.2	18.7	27.5	24.6	20.9	16.9	21.8	26.3
2001.....	7.1	11.2	22.7	25.5	25.5	29.6	24.6	21.3	*
2000.....	6.3	16.3	22.0	28.1	27.7	29.7	28.9	29.2	*
1999.....	10.4	14.8	21.8	25.7	24.2	27.0	33.4	29.2	25.0
Non-Hispanic Black, female									
2003.....	*	2.4	2.7	4.0	6.8	5.5	8.0	8.7	11.6
2002.....	2.0	2.3	3.5	5.3	5.3	6.4	8.4	8.1	12.1
2001.....	2.0	2.3	4.1	3.9	8.1	6.9	7.9	9.6	10.7
2000.....	2.2	*	4.3	5.7	5.7	7.7	5.8	9.5	14.4
1999.....	2.2	4.0	5.3	6.1	7.3	9.6	7.9	9.2	10.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of esophagus (C15)									
All origins 3/, both sexes									
2003.....	2.4	4.6	8.8	12.8	17.6	22.2	25.4	27.6	25.0
2002.....	2.4	4.7	8.6	12.7	18.1	22.5	25.7	27.8	24.6
2001.....	2.4	5.0	7.9	13.0	18.5	22.9	26.1	26.5	25.8
2000.....	2.4	4.9	8.8	13.2	17.9	22.1	24.5	27.2	25.9
1999.....	2.4	4.7	8.7	13.0	18.2	22.0	24.3	24.6	26.8
All origins 3/, male									
2003.....	4.1	8.1	15.3	22.7	30.6	38.3	45.7	51.0	45.3
2002.....	4.1	8.3	15.1	22.1	31.4	40.3	46.8	50.6	43.4
2001.....	4.1	8.7	13.7	22.6	32.6	39.9	46.5	48.8	48.1
2000.....	4.2	8.7	15.2	23.0	30.7	38.0	43.9	48.8	47.8
1999.....	4.2	8.3	15.3	21.9	31.7	39.3	44.1	44.8	48.0
All origins 3/, female									
2003.....	0.7	1.3	2.6	3.8	6.3	9.2	11.0	13.3	16.1
2002.....	0.7	1.3	2.4	4.2	6.7	8.2	10.9	14.0	16.5
2001.....	0.7	1.4	2.4	4.2	6.4	9.4	11.9	13.2	16.5
2000.....	0.6	1.4	2.8	4.4	7.1	9.6	11.0	14.5	17.1
1999.....	0.6	1.3	2.5	5.0	6.7	8.5	10.5	12.8	18.3
Hispanic, both sexes									
2003.....	0.9	1.8	4.7	5.5	8.6	11.7	12.6	17.2	16.1
2002.....	1.1	2.0	3.2	5.3	8.8	11.8	14.4	16.5	23.3
2001.....	1.1	2.8	3.2	7.8	5.9	11.5	13.9	25.8	20.0
2000.....	*	1.7	3.9	6.5	9.3	13.4	19.0	20.1	19.9
1999.....	1.3	1.7	3.9	6.8	9.7	8.5	17.4	19.4	26.7
Hispanic, male									
2003.....	*	3.2	7.5	8.6	15.3	19.6	24.9	34.3	*
2002.....	2.1	3.5	6.4	10.3	15.9	21.5	26.2	28.6	36.8
2001.....	*	5.8	5.0	13.3	11.7	21.5	22.5	45.4	41.9
2000.....	*	3.2	5.7	10.9	16.4	23.3	35.4	35.3	40.3
1999.....	2.4	*	7.9	13.1	16.2	16.3	32.8	33.0	49.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of esophagus (C15)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	16.6
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	2.6	4.9	9.1	13.4	18.2	22.8	26.1	28.1	25.4
2002.....	2.5	5.0	9.0	13.3	18.8	23.1	26.3	28.4	24.5
2001.....	2.5	5.1	8.2	13.3	19.3	23.5	26.7	26.5	26.0
2000.....	2.5	5.2	9.1	13.7	18.5	22.5	24.7	27.4	26.1
1999.....	2.5	4.9	9.0	13.4	18.7	22.7	24.5	24.8	26.8

Non-Hispanic 4/, male

2003.....	4.4	8.6	15.9	23.7	31.7	39.5	46.8	51.9	46.3
2002.....	4.3	8.7	15.8	22.9	32.4	41.4	47.8	51.6	43.5
2001.....	4.3	9.0	14.3	23.2	33.8	40.8	47.6	48.9	48.4
2000.....	4.5	9.1	15.9	23.8	31.5	38.6	44.2	49.3	47.8
1999.....	4.4	8.7	15.8	22.4	32.6	40.4	44.4	45.3	48.0

Non-Hispanic 4/, female

2003.....	0.8	1.3	2.7	3.9	6.5	9.4	11.4	13.7	16.1
2002.....	0.8	1.4	2.6	4.5	6.9	8.4	11.1	14.2	16.3
2001.....	0.8	1.5	2.4	4.3	6.8	9.7	12.1	13.1	16.7
2000.....	0.7	1.5	2.8	4.5	7.3	9.8	11.1	14.5	17.3
1999.....	0.6	1.3	2.7	5.3	6.8	8.9	10.7	12.8	18.3

Non-Hispanic White, both sexes

2003.....	2.4	4.6	8.8	12.9	17.7	23.2	26.5	28.3	26.1
2002.....	2.4	4.9	8.5	12.8	18.2	23.0	26.3	28.7	25.1
2001.....	2.4	4.8	7.4	12.6	18.5	22.7	26.8	26.8	26.5
2000.....	2.4	4.8	8.4	12.9	17.9	22.2	24.2	27.4	26.7
1999.....	2.3	4.5	8.3	12.5	17.9	22.2	23.9	25.0	27.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of esophagus (C15)

Non-Hispanic White, male

2003.....	4.2	8.1	15.5	23.1	30.6	40.2	47.7	53.2	47.9
2002.....	4.3	8.6	15.0	22.3	31.8	41.3	48.3	52.3	44.7
2001.....	4.1	8.3	13.1	22.1	32.5	39.8	48.1	49.5	49.0
2000.....	4.4	8.4	14.7	22.8	30.8	38.2	43.4	50.1	49.4
1999.....	4.1	8.0	14.6	21.0	31.7	39.5	43.4	46.1	48.0

Non-Hispanic White, female

2003.....	0.7	1.1	2.3	3.4	6.2	9.2	11.2	13.0	16.5
2002.....	0.6	1.2	2.3	4.0	6.1	8.0	10.5	14.3	16.6
2001.....	0.6	1.3	1.9	3.7	6.2	8.8	11.8	13.2	17.1
2000.....	0.4	1.2	2.4	3.7	6.6	9.3	10.7	13.8	17.5
1999.....	0.5	1.1	2.3	4.6	5.9	8.3	10.3	12.6	18.9

Non-Hispanic Black, both sexes

2003.....	4.2	8.5	14.3	21.2	26.8	25.2	27.3	30.1	21.4
2002.....	3.7	7.0	15.3	20.8	28.1	29.8	31.9	26.7	20.6
2001.....	3.9	8.8	16.3	22.2	30.1	36.1	31.2	28.1	24.0
2000.....	4.2	10.0	17.4	23.7	27.5	29.9	36.4	30.5	22.6
1999.....	4.5	9.4	17.4	23.9	29.1	31.6	36.0	24.1	23.6

Non-Hispanic Black, male

2003.....	7.3	14.2	23.7	37.1	47.8	41.8	48.1	44.5	38.4
2002.....	5.8	11.7	27.5	35.0	46.3	54.4	54.0	49.1	38.3
2001.....	6.2	15.7	28.3	39.1	52.6	60.9	55.9	54.8	47.4
2000.....	6.5	17.8	31.3	39.1	45.6	50.2	66.0	46.3	38.7
1999.....	7.7	16.5	30.8	40.9	46.8	56.8	68.2	37.2	49.9

Non-Hispanic Black, female

2003.....	1.6	3.6	6.5	8.8	11.3	14.1	15.0	22.5	14.9
2002.....	1.9	3.1	5.3	9.7	14.6	13.3	18.8	15.0	13.8
2001.....	1.8	2.9	6.5	8.9	13.5	19.5	16.4	14.6	15.4
2000.....	2.2	3.4	6.0	11.6	14.2	16.3	18.6	22.5	16.6
1999.....	1.7	3.4	6.4	10.6	15.9	14.9	16.6	17.6	13.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of stomach (C16)

All origins 3/, both sexes

2003.....	2.3	3.8	5.3	8.6	12.4	18.4	24.2	32.2	43.6
2002.....	2.1	3.5	5.6	8.5	12.3	18.8	25.6	33.0	47.6
2001.....	2.2	3.7	5.3	9.2	13.7	19.5	27.0	34.7	45.4
2000.....	2.3	3.6	5.6	9.7	13.8	19.6	27.1	36.9	52.7
1999.....	2.3	3.8	6.2	9.8	14.2	20.2	28.1	36.8	50.1

All origins 3/, male

2003.....	2.8	5.0	7.5	12.3	17.9	26.6	34.2	44.7	60.0
2002.....	2.7	4.8	7.7	11.9	18.1	27.5	36.4	47.1	67.2
2001.....	2.8	4.9	7.5	13.7	19.5	27.5	39.1	49.4	62.6
2000.....	2.9	4.8	7.8	14.1	19.5	28.3	39.4	51.4	75.6
1999.....	3.1	5.4	8.8	13.7	20.3	28.6	41.5	54.9	71.3

All origins 3/, female

2003.....	1.7	2.6	3.1	5.3	7.7	11.9	17.1	24.5	36.4
2002.....	1.5	2.3	3.7	5.4	7.3	11.9	18.1	24.4	39.1
2001.....	1.6	2.5	3.2	5.1	8.8	13.1	18.5	25.9	38.3
2000.....	1.7	2.4	3.5	5.8	8.9	12.7	18.5	28.3	43.4
1999.....	1.5	2.4	3.7	6.4	9.1	13.7	18.8	26.2	41.6

Hispanic, both sexes

2003.....	4.1	6.4	9.3	14.4	19.8	28.9	42.9	44.7	62.7
2002.....	3.4	5.2	8.5	14.5	16.9	28.3	43.2	53.9	72.1
2001.....	4.1	7.0	8.8	14.6	21.4	32.7	40.2	55.1	63.8
2000.....	4.1	5.7	7.1	13.2	23.2	35.0	41.0	66.8	65.7
1999.....	3.0	5.5	9.1	15.0	23.2	29.9	44.2	59.9	73.7

Hispanic, male

2003.....	4.6	7.0	11.5	18.0	25.0	44.0	62.0	66.5	73.9
2002.....	4.0	5.5	10.8	17.7	24.0	44.3	55.7	72.1	107.1
2001.....	4.5	8.5	10.2	21.5	30.5	45.7	51.8	76.6	95.2
2000.....	3.9	6.5	9.4	19.3	34.7	54.5	53.9	104.5	98.8
1999.....	4.2	6.6	12.7	21.1	33.2	38.7	67.9	84.9	75.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of stomach (C16)									
Hispanic, female									
2003.....	3.5	5.7	7.3	11.4	15.5	17.4	29.3	30.5	57.0
2002.....	2.8	5.0	6.4	11.6	11.1	15.9	34.3	42.3	54.7
2001.....	3.7	5.7	7.6	8.7	14.1	22.7	32.0	41.8	48.4
2000.....	4.3	5.0	5.0	7.9	13.9	20.3	31.9	43.9	49.5
1999.....	*	4.6	5.8	9.8	15.0	23.4	27.5	45.0	72.8
Non-Hispanic 4/, both sexes									
2003.....	2.1	3.5	4.9	8.1	11.8	17.7	23.1	31.5	42.7
2002.....	1.9	3.4	5.4	8.0	11.9	18.2	24.6	32.0	46.3
2001.....	2.0	3.4	5.0	8.7	13.2	18.6	26.2	33.9	44.7
2000.....	2.2	3.4	5.4	9.4	13.1	18.7	26.4	35.6	52.0
1999.....	2.2	3.7	5.9	9.4	13.6	19.5	27.4	35.9	49.2
Non-Hispanic 4/, male									
2003.....	2.6	4.7	7.2	11.8	17.3	25.4	32.6	43.4	59.2
2002.....	2.6	4.7	7.4	11.5	17.6	26.4	35.3	45.8	65.0
2001.....	2.6	4.6	7.2	13.1	18.7	26.3	38.3	48.2	61.0
2000.....	2.8	4.6	7.6	13.6	18.4	26.8	38.6	49.3	74.5
1999.....	3.0	5.2	8.5	13.1	19.4	27.8	40.3	53.7	71.0
Non-Hispanic 4/, female									
2003.....	1.5	2.3	2.8	4.8	7.1	11.5	16.3	24.2	35.5
2002.....	1.3	2.1	3.5	4.9	7.0	11.6	17.1	23.6	38.3
2001.....	1.3	2.2	2.9	4.8	8.4	12.5	17.8	25.3	37.9
2000.....	1.5	2.2	3.4	5.6	8.6	12.3	17.8	27.6	42.9
1999.....	1.4	2.2	3.5	6.1	8.6	13.1	18.4	25.5	40.5
Non-Hispanic White, both sexes									
2003.....	1.5	2.8	4.1	6.9	10.2	15.3	20.0	27.8	37.5
2002.....	1.6	2.6	4.5	6.7	9.9	15.5	21.3	28.0	40.7
2001.....	1.6	2.6	4.3	7.3	10.8	16.3	23.0	29.8	40.0
2000.....	1.6	2.8	4.6	8.1	11.0	16.0	22.6	31.2	46.3
1999.....	1.8	2.9	4.8	7.7	12.0	16.8	23.8	31.8	43.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of stomach (C16)

Non-Hispanic White, male

2003.....	2.0	3.7	6.1	9.9	14.6	21.9	28.2	38.7	52.3
2002.....	2.1	3.7	6.4	9.6	14.9	22.7	31.2	40.7	58.0
2001.....	2.1	3.6	6.1	10.9	15.1	23.0	34.2	42.4	55.7
2000.....	2.1	4.0	6.5	11.9	15.7	23.2	33.5	43.8	67.3
1999.....	2.5	4.1	7.1	10.7	17.3	24.3	35.2	47.6	62.6

Non-Hispanic White, female

2003.....	1.1	1.8	2.2	4.2	6.2	9.9	14.0	21.2	31.0
2002.....	1.0	1.5	2.7	3.9	5.4	9.6	14.2	20.2	33.3
2001.....	1.0	1.7	2.4	3.9	7.1	10.9	15.0	22.2	33.5
2000.....	1.1	1.7	2.8	4.5	6.9	10.2	15.0	23.8	37.9
1999.....	1.1	1.6	2.6	4.9	7.4	10.9	15.7	22.6	35.9

Non-Hispanic Black, both sexes

2003.....	4.5	7.5	9.9	17.0	22.6	33.7	49.6	63.0	86.0
2002.....	3.7	7.7	10.7	16.4	25.0	39.7	53.7	63.8	100.1
2001.....	3.8	7.5	9.3	18.0	27.8	35.0	51.7	70.6	89.3
2000.....	4.8	7.2	11.2	17.6	27.0	38.7	57.3	75.6	102.7
1999.....	4.0	8.6	13.3	19.4	24.0	40.7	59.2	74.6	103.1

Non-Hispanic Black, male

2003.....	6.4	11.2	15.2	26.6	37.0	55.3	76.6	86.8	107.6
2002.....	5.2	11.3	15.7	24.6	37.1	60.9	74.6	91.3	136.7
2001.....	5.2	11.6	13.8	28.0	44.8	55.1	76.8	107.1	106.3
2000.....	6.7	9.9	16.6	26.4	38.6	57.7	83.3	104.7	138.6
1999.....	5.8	12.4	19.0	28.7	35.6	60.6	90.9	121.3	149.8

Non-Hispanic Black, female

2003.....	2.8	4.4	5.5	9.6	12.0	19.1	33.6	50.5	77.6
2002.....	2.4	4.6	6.7	10.0	15.9	25.5	41.2	49.5	86.1
2001.....	2.6	4.1	5.5	10.2	15.2	21.6	36.8	52.1	83.0
2000.....	3.1	4.8	6.8	10.7	18.5	26.1	41.5	60.9	89.4
1999.....	2.4	5.4	8.7	12.1	15.4	27.6	40.2	51.4	85.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)									
All origins 3/, both sexes									
2003.....	7.9	14.3	23.0	38.6	59.3	83.3	117.0	159.0	239.7
2002.....	8.2	15.0	24.6	40.7	61.6	86.2	118.9	163.2	241.4
2001.....	8.4	15.2	25.4	39.3	62.9	88.8	121.1	165.8	255.7
2000.....	8.2	14.7	25.2	42.0	62.0	92.4	126.9	174.8	270.4
1999.....	8.3	14.4	25.3	43.6	63.9	91.9	129.0	173.2	271.7
All origins 3/, male									
2003.....	9.0	16.6	27.6	47.1	75.0	103.4	143.0	196.5	273.3
2002.....	9.1	17.5	30.1	51.6	77.8	106.8	147.7	197.4	275.1
2001.....	9.5	17.1	30.8	48.0	78.9	110.4	149.4	201.3	298.3
2000.....	9.0	17.0	30.6	51.0	77.4	115.7	159.4	213.0	315.6
1999.....	9.1	16.6	30.6	52.9	78.0	114.3	161.3	216.2	324.0
All origins 3/, female									
2003.....	6.9	12.1	18.6	30.9	45.6	67.3	98.5	136.1	224.8
2002.....	7.4	12.6	19.5	30.8	47.6	69.7	98.6	142.3	226.8
2001.....	7.4	13.4	20.4	31.5	49.2	71.5	101.2	144.6	237.8
2000.....	7.5	12.4	20.2	33.9	48.8	74.1	104.3	152.2	251.9
1999.....	7.5	12.3	20.4	35.1	51.8	74.3	106.6	148.0	250.6
Hispanic, both sexes									
2003.....	6.1	11.0	20.6	27.8	41.9	59.5	83.2	115.6	144.6
2002.....	5.0	11.3	17.3	30.7	41.9	63.0	90.1	104.7	165.7
2001.....	4.6	10.6	16.7	30.1	40.4	66.7	89.8	108.3	193.8
2000.....	6.3	11.2	19.2	28.1	40.4	60.3	85.4	121.4	181.1
1999.....	6.7	10.3	14.5	29.2	46.8	67.8	88.4	123.9	176.8
Hispanic, male									
2003.....	7.3	13.7	24.6	37.2	54.3	78.6	102.1	151.8	172.4
2002.....	6.3	13.4	24.8	40.0	56.9	80.3	118.4	125.8	174.0
2001.....	4.6	11.1	19.8	39.7	53.2	83.7	113.9	122.1	257.1
2000.....	6.0	13.1	22.4	36.0	54.8	74.9	109.3	175.2	235.8
1999.....	7.0	11.8	16.6	33.7	58.7	91.1	114.7	172.9	211.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)									
Hispanic, female									
2003.....	4.9	8.5	16.8	19.7	31.7	44.9	69.8	92.0	130.5
2002.....	3.7	9.2	10.5	22.6	29.6	49.7	69.9	91.2	161.6
2001.....	4.6	10.2	13.9	21.9	30.1	53.7	72.6	99.7	162.9
2000.....	6.5	9.3	16.3	21.3	28.7	49.3	68.5	88.7	154.3
1999.....	6.4	8.8	12.5	25.4	37.2	50.2	69.9	94.7	160.1
Non-Hispanic 4/, both sexes									
2003.....	8.1	14.5	23.1	39.3	60.4	84.8	118.7	160.7	243.2
2002.....	8.6	15.3	25.2	41.4	62.9	87.4	120.2	165.3	244.0
2001.....	8.8	15.6	26.0	39.9	64.4	89.9	122.4	167.9	257.4
2000.....	8.4	14.9	25.6	43.0	63.3	94.1	128.6	176.5	273.1
1999.....	8.4	14.7	26.0	44.5	64.8	93.0	130.5	174.2	274.3
Non-Hispanic 4/, male									
2003.....	9.2	16.8	27.8	47.7	76.3	104.8	145.0	198.2	277.3
2002.....	9.4	17.8	30.5	52.3	79.1	108.1	148.8	200.3	279.2
2001.....	9.9	17.6	31.5	48.4	80.5	111.7	151.0	204.3	299.6
2000.....	9.3	17.3	31.1	52.0	78.7	117.9	161.4	214.0	318.6
1999.....	9.2	16.9	31.5	54.2	79.0	115.4	163.0	216.7	327.5
Non-Hispanic 4/, female									
2003.....	7.1	12.4	18.8	31.7	46.6	68.7	100.0	137.8	228.1
2002.....	7.8	12.9	20.2	31.5	48.9	70.9	99.9	144.1	228.8
2001.....	7.6	13.6	20.9	32.2	50.5	72.6	102.4	146.1	239.9
2000.....	7.5	12.6	20.4	34.8	50.1	75.4	105.7	154.4	254.7
1999.....	7.5	12.5	20.9	35.7	52.7	75.4	108.0	149.4	253.0
Non-Hispanic White, both sexes									
2003.....	7.4	13.1	21.5	37.6	59.0	81.9	116.6	159.8	242.1
2002.....	7.7	14.5	23.7	39.4	61.1	85.8	118.0	162.3	243.2
2001.....	8.0	14.5	24.4	38.4	62.2	87.8	119.2	165.7	256.9
2000.....	7.5	13.9	23.9	41.4	61.5	92.3	126.6	173.7	271.9
1999.....	7.8	13.7	24.5	42.8	62.9	91.0	128.3	171.9	273.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic White, male									
2003.....	8.4	15.2	26.2	45.4	74.7	102.1	142.3	197.5	276.8
2002.....	8.6	16.8	28.9	49.8	77.2	106.6	145.8	195.9	277.6
2001.....	9.1	16.8	29.9	47.0	78.0	109.8	148.2	201.9	298.2
2000.....	8.5	16.3	29.0	49.9	76.3	115.6	159.5	209.2	315.8
1999.....	8.6	16.0	29.6	52.3	77.1	114.2	159.7	214.9	328.2
Non-Hispanic White, female									
2003.....	6.4	10.9	16.9	30.3	45.0	65.4	97.9	136.7	226.8
2002.....	6.9	12.2	18.8	29.8	46.9	68.7	98.2	141.8	228.3
2001.....	6.9	12.3	19.2	30.4	48.2	69.9	98.6	143.8	239.8
2000.....	6.6	11.7	19.1	33.5	48.4	73.5	103.5	152.6	254.1
1999.....	6.9	11.4	19.6	34.1	50.4	72.5	106.3	146.5	251.5
Non-Hispanic Black, both sexes									
2003.....	13.5	26.0	40.1	60.7	82.5	127.2	158.1	200.1	282.9
2002.....	14.9	24.1	40.9	63.1	88.2	116.3	162.7	221.3	278.3
2001.....	14.8	25.2	44.0	59.0	92.8	122.1	174.5	213.3	289.0
2000.....	15.6	25.4	40.7	63.9	88.5	127.9	169.6	228.8	312.9
1999.....	13.6	23.9	42.8	67.1	91.4	126.5	174.1	227.5	308.8
Non-Hispanic Black, male									
2003.....	15.6	30.8	46.7	78.0	104.8	159.4	199.5	245.3	321.8
2002.....	16.1	28.7	48.6	81.5	110.8	139.2	212.2	288.6	332.6
2001.....	16.8	27.0	52.3	68.3	117.0	148.7	209.5	264.9	352.3
2000.....	17.0	29.9	52.3	78.0	111.1	162.6	211.9	302.1	381.6
1999.....	15.4	27.4	54.9	82.1	111.6	148.9	229.6	275.4	351.8
Non-Hispanic Black, female									
2003.....	11.7	22.0	34.7	47.2	65.9	105.4	133.4	176.5	267.8
2002.....	13.8	20.1	34.5	48.7	71.3	100.9	133.2	186.3	257.5
2001.....	13.1	23.7	37.2	51.6	75.0	104.4	153.6	187.0	265.6
2000.....	14.4	21.6	31.2	52.7	71.8	104.8	144.0	191.8	287.4
1999.....	12.0	20.9	33.0	55.3	76.4	111.7	140.7	203.7	292.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All origins 3/, both sexes

2003.....	3.7	7.4	8.5	11.7	16.5	23.5	29.6	32.1	33.4
2002.....	3.4	6.6	8.1	11.6	16.9	22.6	27.6	31.6	35.0
2001.....	3.8	5.8	7.5	11.6	16.3	21.5	27.0	32.0	34.1
2000.....	3.7	5.4	7.7	11.3	16.0	21.6	27.1	31.3	32.7
1999.....	3.4	4.9	7.1	10.9	16.3	21.8	26.1	30.9	30.6

All origins 3/, male

2003.....	6.0	12.2	13.0	17.3	23.3	33.4	43.7	46.1	47.5
2002.....	5.7	11.1	12.4	17.4	24.8	32.8	39.5	44.8	46.2
2001.....	6.0	9.7	11.4	17.2	23.5	30.6	39.2	43.9	50.3
2000.....	6.0	8.5	11.7	16.5	23.6	30.9	39.3	46.7	44.7
1999.....	5.5	7.6	10.6	15.6	23.7	31.9	36.6	42.9	40.1

All origins 3/, female

2003.....	1.6	2.8	4.3	6.6	10.7	15.6	19.6	23.6	27.2
2002.....	1.2	2.4	4.0	6.4	10.1	14.4	19.2	23.6	30.2
2001.....	1.7	2.1	3.9	6.5	10.2	14.2	18.4	24.8	27.2
2000.....	1.4	2.4	4.0	6.5	9.5	14.2	18.6	22.2	27.8
1999.....	1.3	2.3	3.8	6.6	10.1	13.8	18.7	23.9	26.8

Hispanic, both sexes

2003.....	5.1	10.7	14.4	18.6	30.0	35.6	48.7	51.6	51.8
2002.....	4.6	7.7	12.8	19.6	30.1	39.1	44.0	44.6	59.9
2001.....	5.6	8.1	12.1	16.9	24.9	33.7	50.7	54.6	52.5
2000.....	5.2	6.2	11.6	16.4	24.2	39.8	45.9	62.4	58.4
1999.....	4.5	7.3	9.8	17.2	29.2	33.9	43.6	61.1	53.3

Hispanic, male

2003.....	8.2	18.3	21.1	29.0	38.7	53.8	69.3	60.3	55.4
2002.....	7.1	13.5	20.9	28.5	44.0	55.3	62.1	56.0	75.3
2001.....	10.0	12.6	19.2	24.7	35.4	48.5	68.2	67.5	64.8
2000.....	8.1	10.4	18.0	23.9	35.4	54.9	65.0	94.2	72.6
1999.....	6.9	13.3	16.0	26.8	36.3	51.4	56.2	77.0	66.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)									
Hispanic, female									
2003.....	2.0	3.2	8.2	9.4	22.9	21.6	34.1	46.0	50.0
2002.....	2.0	*	5.4	11.9	18.8	26.7	31.1	37.2	52.2
2001.....	*	3.7	5.6	10.1	16.4	22.3	38.2	46.6	46.5
2000.....	2.3	*	5.8	9.9	15.1	28.4	32.4	43.0	51.4
1999.....	*	*	4.2	9.0	23.4	20.7	34.7	51.6	46.8
Non-Hispanic 4/, both sexes									
2003.....	3.6	7.1	8.0	11.1	15.4	22.7	28.5	31.2	32.6
2002.....	3.3	6.5	7.6	10.9	15.9	21.5	26.7	30.9	33.9
2001.....	3.6	5.6	7.1	11.1	15.7	20.6	25.7	31.0	33.2
2000.....	3.5	5.3	7.4	10.8	15.4	20.5	26.2	30.0	31.7
1999.....	3.3	4.7	6.8	10.4	15.5	21.1	25.3	29.8	29.7
Non-Hispanic 4/, male									
2003.....	5.7	11.6	12.4	16.3	22.0	32.0	42.3	45.3	47.1
2002.....	5.5	10.8	11.6	16.3	23.3	31.3	38.1	44.0	44.8
2001.....	5.6	9.4	10.6	16.6	22.6	29.4	37.6	42.9	49.7
2000.....	5.8	8.3	11.2	15.9	22.7	29.5	38.0	44.6	43.4
1999.....	5.4	7.1	10.2	14.8	22.8	30.7	35.7	41.6	38.7
Non-Hispanic 4/, female									
2003.....	1.5	2.8	3.9	6.4	9.7	15.1	18.8	22.6	26.2
2002.....	1.2	2.4	3.9	6.0	9.4	13.6	18.6	23.0	29.2
2001.....	1.7	2.0	3.7	6.1	9.7	13.7	17.4	23.9	26.4
2000.....	1.4	2.4	3.9	6.2	9.1	13.4	18.0	21.4	27.0
1999.....	1.2	2.3	3.7	6.4	9.1	13.5	18.0	22.9	26.1
Non-Hispanic White, both sexes									
2003.....	2.8	5.7	6.8	9.5	13.8	20.6	27.0	29.8	31.1
2002.....	2.7	5.2	6.2	9.7	14.4	19.7	25.6	29.8	32.6
2001.....	2.9	4.6	5.7	9.9	14.2	19.2	24.6	29.2	32.4
2000.....	2.8	4.2	6.1	9.4	13.8	18.8	24.7	29.2	30.8
1999.....	2.5	3.9	5.5	9.0	14.1	19.6	24.2	28.8	28.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)									
Non-Hispanic White, male									
2003.....	4.5	9.2	10.4	13.9	19.8	29.5	40.1	43.8	45.0
2002.....	4.5	8.5	9.3	14.3	21.2	28.4	36.5	41.9	43.6
2001.....	4.3	7.5	8.4	14.9	21.0	27.6	35.9	41.3	48.9
2000.....	4.4	6.4	8.9	13.6	20.3	27.1	36.5	43.5	41.6
1999.....	4.0	5.6	8.1	12.8	20.8	28.8	34.1	40.8	39.0
Non-Hispanic White, female									
2003.....	1.2	2.4	3.3	5.3	8.4	13.2	17.5	21.2	25.0
2002.....	0.9	2.0	3.2	5.3	8.4	12.5	17.8	22.5	27.8
2001.....	1.4	1.7	3.2	5.2	8.2	12.3	16.6	22.0	25.5
2000.....	1.2	2.0	3.4	5.5	8.0	12.1	16.3	20.8	26.5
1999.....	1.0	2.1	3.0	5.6	8.1	12.3	17.2	21.7	24.8
Non-Hispanic Black, both sexes									
2003.....	6.2	14.5	14.9	18.7	18.9	28.5	30.5	33.4	41.9
2002.....	5.8	12.6	15.7	14.1	19.6	28.0	30.0	32.3	38.1
2001.....	7.3	10.9	15.6	14.5	20.1	24.1	26.2	41.7	35.3
2000.....	7.1	10.9	13.0	16.2	20.7	27.3	29.2	29.0	31.8
1999.....	6.7	8.3	12.2	14.9	21.4	28.4	26.9	34.3	34.9
Non-Hispanic Black, male									
2003.....	10.2	26.1	24.5	29.2	26.8	43.1	43.9	46.0	63.9
2002.....	9.9	22.3	25.6	22.8	32.3	46.2	43.2	51.5	41.6
2001.....	11.8	19.1	26.1	21.7	28.1	33.1	38.4	49.9	42.7
2000.....	12.6	18.1	22.7	25.9	33.6	42.0	37.1	46.3	48.1
1999.....	12.1	14.5	19.0	22.0	32.4	42.6	40.6	43.4	30.9
Non-Hispanic Black, female									
2003.....	2.8	4.7	7.1	10.5	13.0	18.6	22.5	26.9	33.4
2002.....	2.3	4.4	7.5	7.3	10.2	15.8	22.1	22.3	36.8
2001.....	3.3	3.9	7.0	8.9	14.2	18.1	19.0	37.5	32.5
2000.....	2.3	4.8	5.1	8.6	11.2	17.5	24.4	20.3	25.7
1999.....	2.1	3.0	6.7	9.4	13.2	19.1	18.6	29.9	36.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of pancreas (C25)									
All origins 3/, both sexes									
2003.....	4.1	8.7	14.7	24.7	36.4	53.0	67.6	86.2	95.2
2002.....	4.2	8.2	15.5	24.0	37.2	53.2	70.2	81.8	94.5
2001.....	4.0	8.6	14.9	24.8	37.0	52.8	67.2	84.2	100.1
2000.....	4.2	8.7	15.1	24.8	37.2	51.4	70.4	85.7	96.4
1999.....	4.5	8.7	15.6	25.0	38.1	52.5	67.2	82.0	99.7
All origins 3/, male									
2003.....	5.2	10.7	18.2	29.8	42.5	60.9	76.1	99.5	95.6
2002.....	5.3	10.6	19.2	29.3	43.3	60.8	78.9	91.9	100.9
2001.....	4.9	10.7	18.5	30.3	43.6	59.0	77.2	94.9	105.1
2000.....	5.4	11.1	18.5	30.0	44.7	57.0	80.4	94.9	102.9
1999.....	5.9	11.0	19.1	29.9	44.9	60.9	75.7	92.0	107.5
All origins 3/, female									
2003.....	3.0	6.8	11.5	20.2	31.1	46.7	61.6	78.1	95.0
2002.....	3.1	6.0	12.0	19.2	31.9	47.1	64.1	75.6	91.7
2001.....	3.1	6.5	11.5	19.8	31.3	47.8	60.1	77.8	98.1
2000.....	3.0	6.4	11.8	20.0	30.8	47.1	63.5	80.2	93.8
1999.....	3.1	6.5	12.3	20.5	32.2	46.0	61.2	76.1	96.6
Hispanic, both sexes									
2003.....	2.9	5.6	11.5	18.3	29.2	42.7	55.7	67.6	79.8
2002.....	2.8	5.6	11.2	17.4	26.8	44.7	53.9	58.4	79.3
2001.....	2.4	5.4	11.6	17.8	29.2	46.5	55.5	71.5	83.8
2000.....	3.0	5.5	9.6	18.9	29.4	46.3	56.6	71.3	82.9
1999.....	3.7	5.8	10.4	17.0	26.8	40.0	55.5	58.7	84.2
Hispanic, male									
2003.....	3.6	6.9	15.1	21.7	33.1	49.1	63.8	72.8	70.8
2002.....	3.7	7.3	13.0	19.5	32.8	51.3	54.4	58.3	60.2
2001.....	2.9	6.2	13.6	19.8	36.1	50.3	64.1	77.9	62.9
2000.....	4.2	6.5	12.7	21.6	39.5	52.5	57.6	73.6	80.6
1999.....	5.1	6.6	13.4	17.6	32.8	49.4	53.1	62.9	86.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of pancreas (C25)									
Hispanic, female									
2003.....	2.3	4.4	8.2	15.4	25.9	37.7	50.0	64.3	84.4
2002.....	1.9	4.0	9.4	15.7	21.8	39.5	53.5	58.4	88.7
2001.....	*	4.6	9.8	16.0	23.5	43.6	49.4	67.6	94.0
2000.....	*	4.6	6.7	16.6	21.1	41.6	55.9	69.9	84.1
1999.....	2.4	5.0	7.7	16.4	21.9	32.9	57.3	56.3	83.2
Non-Hispanic 4/, both sexes									
2003.....	4.2	8.9	15.0	25.1	36.9	53.6	68.1	86.9	95.7
2002.....	4.3	8.4	15.8	24.4	37.9	53.7	70.9	82.6	94.9
2001.....	4.2	8.8	15.1	25.3	37.4	53.0	67.6	84.6	100.6
2000.....	4.2	8.9	15.5	25.2	37.7	51.6	70.9	85.9	96.6
1999.....	4.6	8.9	16.0	25.5	38.7	53.0	67.5	82.6	100.1
Non-Hispanic 4/, male									
2003.....	5.4	11.0	18.4	30.3	43.2	61.5	76.6	100.5	96.5
2002.....	5.5	10.8	19.6	30.0	44.0	61.3	80.1	93.2	102.4
2001.....	5.1	11.1	18.9	31.1	43.9	59.3	77.7	95.5	106.4
2000.....	5.5	11.4	18.9	30.6	44.9	57.2	81.4	95.5	103.5
1999.....	6.0	11.3	19.4	30.6	45.6	61.2	76.5	92.9	108.2
Non-Hispanic 4/, female									
2003.....	3.1	6.9	11.8	20.4	31.5	47.3	62.1	78.6	95.3
2002.....	3.2	6.1	12.2	19.4	32.5	47.6	64.5	76.1	91.7
2001.....	3.2	6.6	11.6	20.0	31.8	48.0	60.5	78.0	98.2
2000.....	3.1	6.5	12.2	20.3	31.4	47.3	63.7	80.3	93.8
1999.....	3.2	6.6	12.7	20.8	32.8	46.5	61.2	76.5	96.8
Non-Hispanic White, both sexes									
2003.....	4.0	8.3	14.5	24.2	36.4	52.7	67.6	85.8	94.9
2002.....	4.2	7.9	15.3	23.8	36.7	52.3	69.8	82.2	94.6
2001.....	3.9	8.6	14.6	25.3	36.7	51.8	66.7	83.6	99.4
2000.....	4.1	8.6	14.9	24.6	36.6	50.5	69.6	84.7	94.7
1999.....	4.5	8.4	15.1	24.8	37.8	51.9	66.0	81.1	98.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of pancreas (C25)									
Non-Hispanic White, male									
2003.....	5.0	10.3	17.8	29.4	43.3	60.6	76.5	100.2	96.8
2002.....	5.3	10.3	19.0	29.5	43.3	60.1	80.0	93.6	102.9
2001.....	4.9	10.8	18.6	31.0	43.1	58.3	77.6	95.1	108.0
2000.....	5.3	11.0	18.2	30.1	44.5	56.0	80.9	94.5	103.0
1999.....	5.8	10.5	18.2	29.9	44.6	60.4	75.3	91.7	108.1
Non-Hispanic White, female									
2003.....	3.0	6.4	11.2	19.4	30.2	46.2	61.2	76.9	94.0
2002.....	3.1	5.6	11.8	18.4	30.9	45.9	62.6	75.2	91.0
2001.....	3.0	6.4	10.9	19.9	31.0	46.6	58.9	76.7	95.8
2000.....	3.0	6.3	11.7	19.6	29.8	46.0	61.6	78.9	91.4
1999.....	3.1	6.4	12.1	20.1	31.8	45.1	59.5	74.9	95.2
Non-Hispanic Black, both sexes									
2003.....	6.6	14.8	21.1	36.5	48.2	69.3	84.3	108.6	113.3
2002.....	5.8	12.6	22.9	35.2	51.3	74.8	89.3	93.2	108.3
2001.....	6.5	12.5	21.1	31.9	48.9	71.5	85.2	106.0	122.0
2000.....	5.9	13.5	23.0	33.9	51.1	70.6	94.9	105.2	121.2
1999.....	6.4	14.4	25.6	32.7	51.9	70.8	88.3	105.7	119.0
Non-Hispanic Black, male									
2003.....	9.2	19.1	26.8	43.3	51.4	82.6	89.7	114.0	106.5
2002.....	7.4	16.8	29.6	42.0	57.6	85.5	92.0	106.1	102.8
2001.....	8.0	15.6	24.6	38.9	60.2	78.8	94.8	110.4	100.5
2000.....	7.4	18.0	29.5	41.0	56.8	80.0	102.6	117.6	110.4
1999.....	8.6	20.0	34.8	40.5	61.1	78.6	101.5	118.7	115.3
Non-Hispanic Black, female									
2003.....	4.3	11.2	16.4	31.2	45.8	60.4	81.0	105.8	116.0
2002.....	4.3	9.1	17.5	29.9	46.7	67.7	87.7	86.4	110.4
2001.....	5.2	9.8	18.2	26.5	40.6	66.5	79.5	103.7	130.0
2000.....	4.5	9.7	17.7	28.3	47.0	64.4	90.3	98.9	125.2
1999.....	4.4	9.6	18.1	26.6	45.1	65.7	80.4	99.2	120.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of larynx (C32)									
All origins 3/, both sexes									
2003.....	0.7	1.4	2.6	4.6	5.8	6.6	6.6	7.4	6.5
2002.....	0.7	1.4	2.7	4.2	5.3	6.6	7.9	7.2	5.5
2001.....	0.7	1.5	2.9	4.5	5.6	7.3	7.4	6.9	6.7
2000.....	0.8	1.5	2.9	4.5	6.0	7.3	8.0	8.1	6.5
1999.....	0.8	1.5	3.1	4.6	6.1	7.4	7.2	7.4	7.0
All origins 3/, male									
2003.....	1.2	2.4	4.2	7.9	10.3	11.4	12.6	15.4	15.3
2002.....	1.2	2.4	4.7	7.2	8.9	11.3	14.8	14.9	13.2
2001.....	1.2	2.5	4.9	7.5	9.5	12.9	14.5	13.9	18.3
2000.....	1.2	2.5	4.9	7.9	10.0	12.9	14.8	17.1	17.4
1999.....	1.3	2.6	5.4	7.8	10.4	12.6	13.2	15.8	17.8
All origins 3/, female									
2003.....	0.3	0.4	1.0	1.6	2.0	2.8	2.3	2.5	2.7
2002.....	0.2	0.5	0.9	1.5	2.2	2.9	3.1	2.5	2.2
2001.....	0.2	0.5	1.0	1.7	2.2	2.8	2.5	2.8	1.9
2000.....	0.3	0.6	0.9	1.4	2.6	2.8	3.3	2.8	2.1
1999.....	0.2	0.4	1.0	1.7	2.5	3.3	3.0	2.6	2.6
Hispanic, both sexes									
2003.....	*	*	1.6	2.8	3.7	*	5.0	*	*
2002.....	*	*	1.8	2.6	3.2	5.2	6.4	*	*
2001.....	*	*	*	2.5	5.5	4.9	5.7	*	*
2000.....	*	*	*	*	4.5	*	*	11.1	*
1999.....	*	*	*	*	5.2	5.5	7.7	*	*
Hispanic, male									
2003.....	*	*	3.4	5.7	7.5	*	*	*	*
2002.....	*	*	3.9	*	*	10.1	*	*	*
2001.....	*	*	*	*	10.3	11.0	*	*	*
2000.....	*	*	*	*	8.6	*	*	*	*
1999.....	*	*	*	*	10.0	11.2	15.6	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of larynx (C32)									
Hispanic, female									
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes									
2003.....	0.8	1.4	2.6	4.7	6.0	6.8	6.6	7.3	6.5
2002.....	0.8	1.5	2.8	4.4	5.4	6.7	8.0	7.3	5.4
2001.....	0.7	1.6	3.0	4.6	5.5	7.4	7.5	7.0	6.7
2000.....	0.8	1.6	3.0	4.7	6.1	7.4	8.2	8.0	6.5
1999.....	0.8	1.5	3.2	4.8	6.2	7.5	7.2	7.3	6.8
Non-Hispanic 4/, male									
2003.....	1.2	2.5	4.3	8.0	10.4	11.6	12.7	15.3	15.0
2002.....	1.3	2.5	4.7	7.4	9.1	11.3	14.9	15.1	13.2
2001.....	1.2	2.6	5.0	7.6	9.4	13.0	14.5	14.0	18.4
2000.....	1.3	2.6	5.0	8.1	10.0	13.1	15.1	16.7	17.3
1999.....	1.3	2.7	5.5	8.1	10.4	12.6	13.0	15.6	17.3
Non-Hispanic 4/, female									
2003.....	0.3	0.4	1.1	1.7	2.1	2.9	2.3	2.5	2.7
2002.....	0.3	0.5	1.0	1.6	2.3	3.0	3.1	2.5	2.1
2001.....	0.3	0.5	1.1	1.8	2.3	3.0	2.6	2.8	1.8
2000.....	0.4	0.6	1.0	1.5	2.7	2.9	3.4	2.8	2.1
1999.....	0.2	0.5	1.1	1.8	2.5	3.4	3.1	2.5	2.6
Non-Hispanic White, both sexes									
2003.....	0.6	1.2	2.4	4.2	5.6	6.4	6.4	7.0	6.1
2002.....	0.6	1.3	2.5	3.9	5.2	6.4	7.5	7.1	5.5
2001.....	0.5	1.3	2.6	4.2	5.2	6.8	7.0	6.8	6.5
2000.....	0.7	1.3	2.5	4.0	5.7	6.8	7.6	7.8	6.5
1999.....	0.6	1.3	2.8	4.3	5.7	7.2	7.0	7.2	6.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of larynx (C32)									
Non-Hispanic White, male									
2003.....	1.1	2.0	3.7	7.2	9.4	10.6	11.9	14.3	14.1
2002.....	1.1	2.1	4.1	6.4	8.5	10.6	13.8	14.8	13.3
2001.....	0.9	2.1	4.3	7.0	8.6	11.9	13.3	13.3	18.3
2000.....	1.1	2.1	4.1	6.9	9.2	11.9	14.1	16.3	17.4
1999.....	1.0	2.3	4.7	7.3	9.5	11.9	12.7	15.0	16.0
Non-Hispanic White, female									
2003.....	*	0.4	1.0	1.5	2.2	2.9	2.4	2.5	2.6
2002.....	*	0.4	1.0	1.6	2.3	2.9	3.0	2.3	2.2
2001.....	*	0.5	0.9	1.6	2.1	2.7	2.6	2.9	1.6
2000.....	0.3	0.6	0.9	1.4	2.6	2.7	3.1	2.8	2.1
1999.....	*	0.3	1.0	1.6	2.4	3.4	3.1	2.6	2.7
Non-Hispanic Black, both sexes									
2003.....	1.7	3.6	5.9	10.0	10.7	12.8	11.3	12.7	11.3
2002.....	1.9	3.7	6.3	9.6	9.4	12.1	15.6	11.2	*
2001.....	2.0	3.9	7.1	8.9	9.6	13.9	13.6	9.4	8.4
2000.....	2.1	4.1	7.7	11.0	10.7	15.5	15.9	11.8	7.0
1999.....	2.2	4.0	7.7	10.0	11.5	11.9	10.4	9.4	8.4
Non-Hispanic Black, male									
2003.....	2.8	6.9	11.0	18.1	22.8	25.6	26.2	31.7	27.7
2002.....	3.3	6.2	11.8	19.3	17.4	22.9	33.8	22.6	*
2001.....	3.7	7.1	12.9	15.9	17.7	26.2	31.3	22.9	*
2000.....	3.3	7.7	14.7	21.0	20.0	30.4	29.9	27.5	*
1999.....	4.0	7.1	14.8	18.1	21.3	23.2	20.3	26.6	25.0
Non-Hispanic Black, female									
2003.....	*	*	*	3.7	*	*	*	*	*
2002.....	*	*	*	*	*	4.8	*	*	*
2001.....	*	*	*	3.5	*	5.7	*	*	*
2000.....	*	*	*	*	3.9	5.7	7.6	*	*
1999.....	*	*	*	3.8	4.3	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)									
All origins 3/, both sexes									
2003.....	21.2	40.7	81.8	149.0	230.0	314.0	376.1	380.1	298.9
2002.....	20.4	41.6	86.3	152.9	233.4	320.9	376.9	378.5	297.2
2001.....	20.5	42.3	88.8	154.5	239.7	323.2	373.4	368.6	302.7
2000.....	21.0	43.8	92.5	159.8	245.8	325.4	368.2	374.7	302.1
1999.....	19.9	44.4	93.0	160.7	243.6	322.3	361.7	362.6	297.0
All origins 3/, male									
2003.....	24.8	50.0	101.3	183.7	292.1	411.0	514.2	541.5	475.1
2002.....	23.6	51.4	107.0	192.7	297.5	423.0	515.5	546.8	482.2
2001.....	24.3	52.2	109.6	195.7	312.9	434.7	522.3	541.0	512.4
2000.....	25.7	53.2	115.9	202.3	322.5	440.5	516.5	558.1	521.2
1999.....	24.3	54.4	118.1	207.7	322.4	439.2	513.4	551.3	529.2
All origins 3/, female									
2003.....	17.8	31.9	63.5	117.4	176.2	236.1	277.9	281.4	221.0
2002.....	17.3	32.2	66.8	116.8	177.9	239.3	279.1	276.1	217.0
2001.....	16.8	32.8	69.2	117.2	176.7	234.5	269.1	265.2	215.0
2000.....	16.5	34.7	70.5	121.3	180.2	234.7	264.9	266.6	212.8
1999.....	15.7	34.9	69.5	118.1	176.1	230.9	256.2	252.2	203.5
Hispanic, both sexes									
2003.....	6.6	15.0	27.2	53.2	88.6	127.6	177.3	194.3	189.2
2002.....	6.0	15.0	32.0	53.1	87.4	139.2	181.6	187.2	189.0
2001.....	6.6	16.0	30.9	54.6	89.4	131.8	177.2	191.2	200.7
2000.....	7.0	16.0	31.6	49.2	90.9	135.4	182.1	228.9	213.7
1999.....	6.6	15.4	33.2	58.0	96.1	134.2	178.5	204.4	227.3
Hispanic, male									
2003.....	7.0	20.1	39.9	77.8	137.5	193.9	272.2	285.9	290.9
2002.....	6.8	20.3	47.0	79.8	133.1	198.8	296.3	304.3	307.9
2001.....	8.6	21.6	43.8	80.9	130.8	206.3	281.7	316.8	335.2
2000.....	9.1	22.3	46.0	75.1	130.9	214.9	304.1	374.0	368.8
1999.....	8.8	21.4	50.6	90.5	152.2	205.7	294.1	360.0	362.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)									
Hispanic, female									
2003.....	6.2	10.0	15.5	31.7	48.5	76.7	109.9	134.7	137.5
2002.....	5.2	10.0	18.2	29.7	50.0	93.3	99.6	112.4	130.1
2001.....	4.6	10.7	19.2	31.7	56.0	75.0	103.1	113.4	135.0
2000.....	4.8	10.0	18.5	26.8	58.6	75.1	95.7	140.7	137.5
1999.....	4.4	9.6	17.5	30.0	50.6	80.4	96.9	111.6	162.2
Non-Hispanic 4/, both sexes									
2003.....	22.8	43.1	86.1	156.2	240.0	325.7	386.5	388.2	303.0
2002.....	21.9	43.9	90.5	160.2	243.4	331.9	386.3	386.0	301.0
2001.....	21.9	44.4	93.2	161.7	249.8	334.1	382.4	375.1	305.8
2000.....	22.3	45.9	96.8	167.6	255.5	335.4	376.0	379.5	304.5
1999.....	21.1	46.6	97.2	167.7	252.4	331.5	368.7	367.5	298.8
Non-Hispanic 4/, male									
2003.....	26.7	52.7	106.1	191.5	302.3	424.0	526.5	553.1	483.1
2002.....	25.3	54.1	111.7	200.8	308.3	436.2	525.7	556.7	488.9
2001.....	25.9	54.6	114.3	203.5	324.7	447.4	533.1	549.4	518.4
2000.....	27.2	55.6	120.7	210.9	334.0	451.9	525.2	564.0	526.2
1999.....	25.7	56.8	122.7	215.3	332.1	450.2	521.5	557.1	534.6
Non-Hispanic 4/, female									
2003.....	18.9	33.9	67.3	124.0	185.8	246.6	286.8	287.7	223.9
2002.....	18.5	34.1	70.7	123.3	187.2	248.2	287.9	282.4	220.0
2001.....	17.9	34.6	73.3	123.7	185.0	243.9	276.9	270.8	217.4
2000.....	17.5	36.6	74.4	128.2	188.0	243.4	272.1	270.8	214.8
1999.....	16.7	36.8	73.3	124.5	183.9	238.6	262.5	256.5	204.4
Non-Hispanic White, both sexes									
2003.....	21.5	41.6	85.3	157.8	244.7	332.7	392.4	392.9	306.7
2002.....	21.0	42.4	89.4	160.6	247.9	339.0	392.3	392.1	304.8
2001.....	20.7	43.2	92.7	163.9	253.2	339.3	386.7	377.9	308.6
2000.....	20.8	44.4	96.3	169.9	258.9	339.0	378.9	385.7	306.8
1999.....	19.6	45.0	96.5	168.4	254.2	334.6	370.3	370.4	300.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)									
Non-Hispanic White, male									
2003.....	24.8	49.5	102.1	188.4	302.0	426.0	525.8	553.6	486.8
2002.....	23.5	50.8	107.5	195.6	307.6	438.5	525.3	557.9	493.3
2001.....	23.8	51.4	110.5	201.8	322.2	446.0	530.5	545.9	520.0
2000.....	25.0	51.8	116.5	209.8	331.3	448.7	520.4	565.8	527.3
1999.....	23.2	52.9	118.9	211.0	328.0	446.4	517.2	553.9	536.9
Non-Hispanic White, female									
2003.....	18.3	33.8	69.1	129.4	193.8	255.9	295.8	294.2	227.4
2002.....	18.5	34.2	72.1	128.1	194.9	257.7	297.2	290.6	223.4
2001.....	17.7	35.1	75.5	128.8	192.3	252.7	284.9	276.5	220.6
2000.....	16.7	37.2	77.0	133.0	195.2	250.6	279.5	278.7	218.0
1999.....	16.1	37.2	75.1	128.9	189.4	245.5	267.2	262.1	206.2
Non-Hispanic Black, both sexes									
2003.....	36.2	63.4	117.7	186.9	261.7	334.4	393.2	383.7	283.5
2002.....	33.3	66.0	123.3	197.4	270.6	336.0	395.6	370.2	274.9
2001.....	33.6	64.6	121.4	185.0	277.6	353.4	401.1	375.0	281.6
2000.....	36.4	70.5	124.8	192.2	284.2	363.5	399.7	347.2	291.0
1999.....	35.8	71.8	127.4	202.3	286.3	356.2	406.5	365.7	293.3
Non-Hispanic Black, male									
2003.....	46.3	88.7	172.0	274.1	383.7	503.2	627.5	640.7	509.3
2002.....	44.0	93.7	175.7	297.6	403.1	515.9	638.5	626.3	490.1
2001.....	46.2	93.9	177.0	271.9	426.0	561.5	668.4	667.2	550.9
2000.....	48.8	102.3	187.8	276.6	438.7	568.2	664.3	623.1	567.1
1999.....	49.8	104.8	189.6	307.4	434.8	564.3	662.7	669.5	580.0
Non-Hispanic Black, female									
2003.....	27.4	42.1	73.1	119.0	171.2	220.6	254.1	249.2	196.0
2002.....	24.1	42.5	80.4	119.2	172.3	215.4	250.6	237.0	192.7
2001.....	22.6	39.8	75.8	116.7	168.1	214.7	241.6	226.1	181.8
2000.....	25.7	43.4	73.3	125.6	170.3	227.5	240.2	208.2	188.4
1999.....	23.7	43.9	76.8	120.0	176.2	218.0	252.2	215.1	186.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant melanoma of skin (C43)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	2.7	3.5	5.2	6.7	7.7	10.7	14.5	16.0	18.1
2002.....	2.5	3.6	4.7	6.3	8.3	10.5	13.3	15.3	18.2
2001.....	2.8	3.9	4.9	6.5	8.2	11.2	12.7	15.1	18.3
2000.....	2.8	4.0	4.9	6.6	8.3	10.3	12.3	15.3	19.3
1999.....	2.8	3.8	4.9	6.4	8.6	10.1	13.2	14.6	16.7

Non-Hispanic 4/, male

2003.....	3.4	4.8	7.3	9.5	10.9	15.8	23.4	25.4	30.9
2002.....	3.5	4.7	6.5	8.9	12.0	15.6	20.6	24.0	30.2
2001.....	3.5	5.1	6.6	9.3	12.5	17.0	20.2	24.8	31.0
2000.....	3.4	5.5	6.7	9.2	11.9	15.2	19.0	25.2	32.6
1999.....	3.7	4.8	7.0	8.8	12.4	15.2	20.4	23.3	28.2

Non-Hispanic 4/, female

2003.....	2.0	2.3	3.2	4.2	4.8	6.6	8.1	10.2	12.4
2002.....	1.5	2.6	3.0	3.8	5.1	6.4	8.1	10.0	12.9
2001.....	2.1	2.8	3.2	3.9	4.6	6.6	7.5	9.2	12.9
2000.....	2.2	2.6	3.1	4.3	5.2	6.5	7.7	9.4	13.9
1999.....	2.0	2.7	2.9	4.2	5.4	6.1	8.1	9.5	12.0

Non-Hispanic White, both sexes

2003.....	3.2	4.2	6.0	7.7	8.8	12.1	15.9	17.5	19.6
2002.....	3.0	4.3	5.4	7.2	9.5	11.7	14.7	16.8	19.6
2001.....	3.4	4.6	5.7	7.5	9.3	12.5	14.0	16.3	19.9
2000.....	3.4	4.7	5.6	7.6	9.4	11.4	13.5	16.4	20.8
1999.....	3.4	4.4	5.7	7.3	9.8	11.2	14.4	15.7	17.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)									
Hispanic, female									
2003.....	16.9	26.2	37.9	40.1	50.1	46.2	72.9	67.7	114.8
2002.....	17.9	26.4	30.5	37.3	47.6	50.0	63.1	89.1	105.3
2001.....	19.2	29.5	39.1	44.7	44.2	55.5	64.9	64.3	102.4
2000.....	20.7	28.0	39.9	38.2	54.7	55.2	69.5	84.2	105.8
1999.....	18.8	30.4	33.6	40.3	45.0	54.0	62.8	87.2	127.9
Non-Hispanic 4/, both sexes									
2003.....	13.0	19.6	27.1	34.4	40.7	53.3	67.1	89.2	134.7
2002.....	13.3	20.2	26.9	34.7	42.2	53.7	68.6	91.0	136.8
2001.....	13.8	21.1	27.6	34.7	42.7	55.3	70.4	89.8	136.4
2000.....	13.9	21.3	28.6	35.9	44.1	56.4	72.0	92.1	149.7
1999.....	13.6	21.4	28.8	35.6	44.5	57.1	71.3	93.5	146.2
Non-Hispanic 4/, male									
2003.....	*	0.4	0.4	0.8	1.0	1.3	1.6	3.0	3.8
2002.....	*	0.3	0.4	0.7	0.8	1.3	2.0	2.8	3.5
2001.....	*	0.4	0.5	0.8	1.2	1.8	2.3	2.4	3.7
2000.....	*	0.3	0.7	0.8	1.1	1.7	2.2	2.4	4.2
1999.....	*	0.4	0.6	0.7	1.1	1.4	1.9	2.4	4.7
Non-Hispanic 4/, female									
2003.....	25.5	37.9	52.3	65.1	75.3	95.2	113.7	141.7	192.2
2002.....	26.0	39.3	51.9	65.7	78.2	95.8	115.6	144.6	194.4
2001.....	27.0	40.9	53.0	65.7	78.7	97.9	118.0	142.1	191.6
2000.....	27.1	41.3	54.8	67.8	81.1	99.6	120.6	145.0	208.6
1999.....	26.7	41.6	55.3	67.3	81.8	100.8	119.4	146.9	202.8
Non-Hispanic White, both sexes									
2003.....	11.8	18.1	25.6	33.4	40.1	53.2	67.2	89.6	134.7
2002.....	12.0	18.8	25.7	34.1	41.8	53.4	69.1	90.3	137.7
2001.....	12.6	19.8	26.2	33.9	41.9	54.9	70.8	89.9	137.1
2000.....	12.5	20.1	27.2	35.3	44.0	56.8	72.3	92.9	149.6
1999.....	12.3	20.1	27.4	34.6	43.8	56.8	71.0	94.0	147.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of breast (C50)

Non-Hispanic White, male

2003.....	*	0.4	0.4	0.8	1.0	1.2	1.6	3.0	3.9
2002.....	*	*	0.4	0.6	0.6	1.3	1.8	2.7	3.4
2001.....	*	0.3	0.5	0.8	1.1	1.7	2.1	2.4	3.5
2000.....	*	0.3	0.6	0.8	1.1	1.7	2.2	2.2	4.0
1999.....	*	0.3	0.5	0.6	1.0	1.2	1.8	2.3	4.5

Non-Hispanic White, female

2003.....	23.3	35.5	50.0	63.7	74.8	96.0	114.6	142.8	192.3
2002.....	23.7	37.0	50.0	65.2	78.3	96.0	117.2	143.9	195.8
2001.....	24.9	38.8	50.8	64.6	78.0	98.1	119.4	142.8	192.6
2000.....	24.7	39.4	52.6	67.2	81.7	101.1	121.5	146.9	208.3
1999.....	24.3	39.5	53.0	66.0	81.3	101.1	119.5	148.1	204.9

Non-Hispanic Black, both sexes

2003.....	23.1	32.0	44.2	48.7	55.3	66.2	81.1	106.5	163.3
2002.....	23.4	32.9	41.2	47.0	55.4	68.2	79.9	118.8	154.0
2001.....	23.2	33.7	43.4	48.5	58.9	71.7	83.8	107.3	153.9
2000.....	24.3	33.7	44.9	48.2	53.5	66.5	85.5	101.4	178.7
1999.....	24.6	35.2	45.2	49.6	59.0	71.1	89.0	106.3	149.1

Non-Hispanic Black, male

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic Black, female

2003.....	42.9	58.7	79.9	85.6	95.2	109.3	128.5	160.7	224.5
2002.....	43.4	60.1	74.3	82.3	94.3	112.9	125.3	178.6	210.7
2001.....	42.9	61.5	78.3	85.6	101.0	117.2	131.3	160.3	208.3
2000.....	45.0	61.9	80.5	84.9	92.4	108.9	135.6	150.4	241.6
1999.....	45.6	64.2	81.0	86.9	101.6	116.0	140.9	156.7	201.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

Hispanic, female

2003.....	5.0	6.6	6.7	4.9	5.6	7.2	9.5	13.5	*
2002.....	4.1	5.6	6.7	6.9	6.6	10.5	11.0	*	*
2001.....	6.3	5.6	7.4	8.0	*	8.0	10.2	*	*
2000.....	5.3	5.7	7.9	9.4	9.1	11.0	12.5	*	20.8
1999.....	6.5	6.7	7.1	7.7	8.4	9.2	11.0	*	*

Non-Hispanic 4/, both sexes

2003.....	2.0	2.3	2.8	2.9	3.2	2.9	3.6	5.0	5.7
2002.....	2.1	2.4	2.6	2.7	3.3	3.5	4.0	4.0	6.1
2001.....	2.2	2.6	2.5	3.1	3.0	3.4	3.9	5.4	7.0
2000.....	2.2	2.6	3.0	2.9	3.5	3.7	4.1	5.5	7.1
1999.....	2.3	2.6	2.9	3.5	3.6	3.8	4.6	4.5	7.9

Non-Hispanic 4/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2003.....	3.8	4.4	5.4	5.5	5.9	5.2	6.1	8.0	8.2
2002.....	4.0	4.7	5.0	5.2	6.1	6.2	6.8	6.5	8.7
2001.....	4.3	5.0	4.9	5.9	5.5	6.1	6.7	8.6	10.0
2000.....	4.3	5.1	5.8	5.5	6.5	6.7	7.0	8.7	9.9
1999.....	4.5	5.1	5.7	6.6	6.7	6.8	7.8	7.1	11.0

Non-Hispanic White, both sexes

2003.....	1.8	2.0	2.5	2.5	2.7	2.3	2.7	4.4	4.8
2002.....	1.8	2.1	2.4	2.2	2.7	2.7	3.5	3.4	4.9
2001.....	2.0	2.3	2.2	2.7	2.4	2.9	3.3	4.5	6.1
2000.....	1.9	2.3	2.6	2.4	2.8	3.1	3.6	4.8	6.0
1999.....	2.1	2.4	2.5	2.9	3.0	2.9	3.9	4.0	6.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic White, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2003.....	3.5	3.9	4.9	4.9	5.1	4.2	4.6	7.1	7.0
2002.....	3.6	4.2	4.6	4.3	5.0	5.0	5.9	5.4	7.0
2001.....	4.0	4.5	4.2	5.2	4.4	5.3	5.6	7.3	8.6
2000.....	3.8	4.5	5.1	4.6	5.2	5.5	6.1	7.6	8.4
1999.....	4.2	4.7	5.0	5.5	5.7	5.3	6.6	6.3	9.3

Non-Hispanic Black, both sexes

2003.....	3.6	3.9	5.2	6.3	7.2	7.9	11.5	11.7	14.9
2002.....	3.9	4.6	4.3	6.2	8.4	10.3	9.5	12.5	17.5
2001.....	3.3	4.5	4.7	6.1	8.0	7.8	9.7	15.5	19.0
2000.....	3.8	5.0	6.2	6.6	9.6	10.0	10.9	13.8	19.7
1999.....	3.1	4.2	6.3	8.3	8.7	13.3	12.4	9.7	21.7

Non-Hispanic Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2003.....	6.7	7.2	9.5	11.1	12.6	13.2	18.3	17.8	20.6
2002.....	7.2	8.4	7.8	11.0	14.6	17.2	15.1	19.1	24.2
2001.....	6.1	8.3	8.6	10.8	13.8	13.1	15.6	23.3	26.1
2000.....	7.0	9.3	11.4	11.7	16.7	16.6	17.4	20.7	27.0
1999.....	5.7	7.7	11.4	14.9	15.2	22.1	19.8	14.5	29.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Hispanic, female

2003.....	*	3.3	4.8	6.6	11.5	13.8	15.1	18.9	31.3
2002.....	*	2.6	6.4	5.8	14.9	17.6	18.3	19.0	30.7
2001.....	*	4.2	5.8	9.2	11.2	16.1	16.5	24.9	27.9
2000.....	*	3.4	6.2	7.7	10.6	14.7	18.3	*	29.7
1999.....	*	3.5	5.8	8.5	14.1	15.7	16.5	19.7	22.9

Non-Hispanic 4/, both sexes

2003.....	0.8	1.9	3.6	6.3	9.2	11.5	13.9	17.3	22.9
2002.....	0.9	2.1	3.5	5.9	8.4	11.9	14.3	17.2	24.6
2001.....	0.9	2.0	3.2	5.9	8.4	11.5	14.7	19.3	24.8
2000.....	1.0	1.8	3.6	6.1	8.6	11.2	14.2	18.8	24.9
1999.....	1.0	1.7	3.3	5.6	8.8	11.0	14.6	19.6	24.3

Non-Hispanic 4/, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, female

2003.....	1.5	3.8	7.0	12.1	17.2	20.8	23.9	27.9	33.0
2002.....	1.7	4.0	6.8	11.3	15.8	21.4	24.4	27.7	35.3
2001.....	1.8	3.9	6.2	11.2	15.6	20.6	25.0	30.9	35.1
2000.....	1.9	3.5	6.9	11.6	15.9	20.0	24.0	29.9	34.9
1999.....	1.9	3.3	6.3	10.7	16.4	19.7	24.8	31.1	34.1

Non-Hispanic White, both sexes

2003.....	0.7	1.9	3.5	5.5	8.0	10.7	13.2	16.4	22.8
2002.....	0.8	1.9	3.1	5.4	7.1	10.8	13.2	16.5	23.8
2001.....	0.9	1.9	3.1	5.2	7.3	10.5	13.9	18.7	24.5
2000.....	1.0	1.7	3.4	5.4	7.4	9.9	13.1	17.7	24.5
1999.....	0.9	1.6	3.0	5.2	7.6	10.3	13.9	19.0	23.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic White, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2003.....	1.5	3.7	6.8	10.7	15.1	19.4	22.7	26.5	32.8
2002.....	1.6	3.8	6.2	10.3	13.4	19.7	22.6	26.6	34.1
2001.....	1.7	3.7	6.1	10.0	13.7	19.0	23.8	30.0	34.7
2000.....	1.9	3.4	6.7	10.5	13.8	17.9	22.3	28.2	34.3
1999.....	1.9	3.2	5.9	10.0	14.2	18.5	23.6	30.3	33.1

Non-Hispanic Black, both sexes

2003.....	1.0	2.6	5.3	14.3	20.4	22.8	25.4	31.9	30.3
2002.....	1.1	3.0	6.8	11.0	20.4	25.5	28.1	28.8	39.6
2001.....	1.4	3.0	4.6	11.7	19.1	24.6	26.6	31.7	32.8
2000.....	1.0	2.4	5.3	12.3	20.0	26.5	28.3	35.6	33.7
1999.....	*	2.3	5.9	10.8	21.1	22.0	25.8	30.2	35.9

Non-Hispanic Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2003.....	1.9	4.8	9.6	25.4	35.5	38.2	40.5	48.6	42.1
2002.....	2.0	5.5	12.3	19.6	35.5	42.7	44.9	43.8	54.8
2001.....	2.7	5.6	8.3	20.9	33.3	41.1	42.4	47.9	44.9
2000.....	2.0	4.5	9.7	22.0	34.8	44.0	45.3	53.6	46.2
1999.....	*	4.2	10.8	19.3	36.8	36.5	41.4	45.2	49.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of ovary (C56)									
Hispanic, female									
2003.....	4.8	6.3	11.1	13.7	19.7	26.6	35.8	39.2	43.0
2002.....	4.4	8.8	7.9	14.3	19.6	25.0	41.1	40.1	35.6
2001.....	4.1	6.9	10.3	17.4	25.5	26.2	28.1	42.6	45.6
2000.....	4.7	7.5	10.9	13.9	22.6	29.1	41.8	41.2	48.5
1999.....	3.4	7.7	8.6	15.1	15.9	28.7	32.5	38.4	35.4
Non-Hispanic 4/, both sexes									
2003.....	2.9	5.4	8.5	13.4	17.8	22.3	29.9	37.5	38.8
2002.....	3.3	5.6	8.7	13.3	17.7	22.8	30.6	38.4	38.7
2001.....	3.0	5.7	8.5	13.2	17.8	23.0	30.9	37.8	39.8
2000.....	3.1	5.6	8.5	12.6	17.8	22.1	31.2	36.8	41.2
1999.....	3.3	5.6	8.5	12.9	17.5	22.6	29.5	35.7	38.1
Non-Hispanic 4/, male									
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic 4/, female									
2003.....	5.8	10.6	16.5	25.7	33.2	40.3	51.2	60.4	55.9
2002.....	6.5	10.9	16.8	25.4	33.1	41.1	52.2	61.6	55.3
2001.....	5.9	11.1	16.4	25.2	33.3	41.4	52.4	60.4	56.4
2000.....	6.1	11.0	16.5	24.0	33.1	39.5	52.9	58.4	57.8
1999.....	6.4	11.0	16.5	24.5	32.6	40.3	50.0	56.6	53.3
Non-Hispanic White, both sexes									
2003.....	3.0	5.6	8.9	13.9	18.4	22.8	30.8	38.7	39.5
2002.....	3.4	5.9	9.0	13.9	18.5	23.5	31.4	39.3	39.6
2001.....	3.2	5.9	8.7	13.4	18.3	23.8	31.8	39.0	40.6
2000.....	3.3	5.8	8.9	13.1	18.3	22.7	32.3	37.9	42.2
1999.....	3.4	5.8	8.9	13.4	18.1	23.1	30.0	36.7	39.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of ovary (C56)

Non-Hispanic White, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, female

2003.....	6.0	11.0	17.5	26.9	34.8	41.6	53.0	62.6	56.8
2002.....	6.8	11.6	17.6	26.9	34.8	42.6	53.9	63.4	56.7
2001.....	6.4	11.8	17.1	25.9	34.4	43.2	54.3	62.5	57.5
2000.....	6.6	11.4	17.5	25.2	34.4	41.0	54.9	60.4	59.2
1999.....	6.8	11.4	17.3	25.8	33.9	41.5	51.0	58.3	54.9

Non-Hispanic Black, both sexes

2003.....	2.8	5.2	6.9	11.3	15.4	20.9	26.8	29.3	37.2
2002.....	2.9	4.2	7.8	11.1	15.1	20.3	26.9	33.1	35.4
2001.....	2.0	4.5	8.1	12.8	16.6	20.8	26.4	28.4	36.5
2000.....	2.2	4.8	6.6	11.2	16.9	19.9	26.6	27.9	34.7
1999.....	2.4	5.2	7.4	11.2	15.3	22.9	28.9	28.5	29.1

Non-Hispanic Black, male

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic Black, female

2003.....	5.3	9.6	12.6	20.1	26.8	35.0	42.7	44.6	51.6
2002.....	5.4	7.8	14.2	19.7	26.4	34.0	42.9	50.3	48.9
2001.....	3.7	8.3	14.8	22.8	28.9	34.6	42.2	42.9	50.0
2000.....	4.1	9.0	11.9	20.0	29.3	33.1	42.7	41.9	47.5
1999.....	4.5	9.6	13.4	20.0	26.6	38.1	46.3	42.6	40.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of prostate (C61)

All origins 3/, both sexes

2003.....	0.5	1.7	4.2	11.6	23.5	43.6	76.1	122.1	184.9
2002.....	0.4	1.6	4.6	11.4	24.2	45.8	79.7	128.7	195.0
2001.....	0.5	1.7	4.6	11.3	25.0	47.4	82.4	133.2	200.9
2000.....	0.5	1.5	4.6	12.1	25.9	49.0	85.1	136.6	209.3
1999.....	0.5	1.6	4.9	12.3	28.4	51.9	89.0	141.4	207.5

All origins 3/, male

2003.....	0.9	3.4	8.7	24.4	50.6	97.9	183.0	321.8	603.3
2002.....	0.8	3.3	9.6	24.0	52.2	103.2	192.6	340.3	644.5
2001.....	1.0	3.5	9.4	23.8	53.9	107.1	200.1	355.4	681.2
2000.....	1.1	3.1	9.6	25.4	56.0	111.3	207.4	368.2	723.1
1999.....	1.0	3.3	10.1	25.8	61.6	118.3	216.9	383.0	722.4

All origins 3/, female

2003.....
2002.....
2001.....
2000.....
1999.....

Hispanic, both sexes

2003.....	*	*	3.1	8.4	18.6	34.3	58.0	101.2	151.8
2002.....	*	1.4	3.2	8.4	19.9	30.7	62.7	105.6	170.7
2001.....	*	1.5	3.8	8.6	22.5	37.6	78.7	103.8	170.1
2000.....	*	*	2.2	7.6	15.9	38.1	63.7	107.5	165.9
1999.....	*	1.6	4.4	7.2	23.3	35.0	67.5	104.0	176.1

Hispanic, male

2003.....	*	*	6.5	18.0	41.3	79.0	139.8	256.8	451.0
2002.....	*	2.8	6.6	18.0	44.4	70.7	150.4	271.1	515.4
2001.....	*	3.0	8.0	18.5	50.3	86.9	189.6	271.4	518.0
2000.....	*	*	4.6	16.4	35.4	88.5	153.5	284.2	503.9
1999.....	*	3.2	9.2	15.5	52.2	81.5	163.1	278.2	541.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of prostate (C61)									
Hispanic, female									
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic 4/, both sexes									
2003.....	0.5	1.7	4.3	11.8	23.8	44.1	76.9	122.9	186.0
2002.....	0.4	1.6	4.7	11.6	24.4	46.6	80.4	129.4	195.5
2001.....	0.5	1.7	4.6	11.5	25.1	47.9	82.3	134.0	201.6
2000.....	0.6	1.5	4.8	12.4	26.5	49.5	85.8	137.4	210.4
1999.....	0.5	1.6	4.9	12.6	28.7	52.6	89.7	142.5	208.2
Non-Hispanic 4/, male									
2003.....	1.0	3.5	8.9	24.8	51.2	98.8	185.0	324.6	609.4
2002.....	0.8	3.3	9.8	24.3	52.5	104.7	194.4	342.5	648.7
2001.....	1.0	3.5	9.5	24.1	54.1	108.1	199.9	357.9	686.3
2000.....	1.1	3.1	9.9	26.0	57.3	112.1	209.2	370.6	730.7
1999.....	1.1	3.3	10.1	26.4	62.1	119.8	218.8	386.3	728.2
Non-Hispanic 4/, female									
2003.....
2002.....
2001.....
2000.....
1999.....
Non-Hispanic White, both sexes									
2003.....	0.4	1.4	3.5	10.0	20.6	39.8	70.2	114.6	178.2
2002.....	0.3	1.3	4.0	9.7	21.2	41.3	73.0	120.6	186.6
2001.....	0.4	1.4	3.9	9.7	22.0	42.4	74.8	123.6	189.5
2000.....	0.4	1.3	4.0	10.6	22.8	43.3	78.1	126.3	199.3
1999.....	0.4	1.4	4.1	10.3	24.9	46.8	81.5	133.8	195.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)									
All origins 3/, both sexes									
2003.....	2.0	4.0	7.1	11.4	14.5	19.6	24.8	30.0	33.2
2002.....	2.4	4.3	7.2	10.9	16.0	19.6	24.6	28.7	32.5
2001.....	2.4	4.4	7.1	11.9	14.6	20.8	24.4	29.1	32.3
2000.....	2.2	4.5	7.6	10.8	15.0	19.4	24.5	29.6	33.8
1999.....	2.2	4.3	7.3	10.8	14.4	19.2	23.1	26.9	31.4
All origins 3/, male									
2003.....	3.0	5.7	10.4	17.2	20.2	27.7	35.9	44.9	48.7
2002.....	3.4	6.4	10.6	16.1	23.0	27.6	36.6	39.9	50.1
2001.....	3.5	6.5	10.5	17.2	21.0	30.0	34.4	42.2	52.9
2000.....	3.1	6.6	11.2	15.6	21.4	27.4	34.6	44.2	55.1
1999.....	3.2	6.1	10.6	15.8	20.3	27.3	33.7	38.9	49.4
All origins 3/, female									
2003.....	1.0	2.3	4.0	6.1	9.4	13.0	16.9	20.9	26.4
2002.....	1.5	2.3	3.9	6.2	10.0	13.2	16.2	21.9	24.9
2001.....	1.4	2.4	3.9	7.0	9.1	13.5	17.4	21.2	23.7
2000.....	1.4	2.4	4.3	6.5	9.5	13.0	17.5	21.0	25.1
1999.....	1.2	2.5	4.3	6.2	9.3	12.8	15.7	19.9	24.1
Hispanic, both sexes									
2003.....	1.7	2.7	6.3	10.8	12.9	20.4	24.2	22.5	26.9
2002.....	2.1	3.8	6.6	10.2	11.9	14.1	23.5	25.8	26.1
2001.....	2.6	4.3	4.6	10.0	12.2	17.0	24.4	21.4	31.9
2000.....	2.4	3.4	6.2	10.0	10.8	14.0	21.4	30.6	25.9
1999.....	1.9	4.1	6.7	9.4	13.3	14.6	21.0	25.3	28.8
Hispanic, male									
2003.....	2.6	3.2	8.2	15.0	18.8	26.9	34.6	28.1	38.5
2002.....	2.6	5.8	9.9	13.3	16.9	20.6	41.0	35.5	40.2
2001.....	3.8	6.0	6.6	14.4	17.7	26.1	35.5	35.1	57.1
2000.....	3.4	4.5	10.1	13.2	14.9	21.9	34.7	45.6	40.3
1999.....	3.3	5.6	8.8	15.8	18.9	23.9	29.6	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Hispanic, female

2003.....	*	*	4.4	7.1	8.0	15.4	16.8	18.9	21.1
2002.....	*	*	3.5	7.4	7.7	9.1	11.0	19.7	19.1
2001.....	*	2.6	*	6.1	7.7	10.1	16.5	*	19.5
2000.....	*	*	*	7.2	7.5	8.1	12.0	21.5	*
1999.....	*	*	4.8	*	8.8	7.7	14.9	22.5	23.9

Non-Hispanic 4/, both sexes

2003.....	2.0	4.1	7.2	11.4	14.6	19.4	24.8	30.3	33.4
2002.....	2.4	4.3	7.2	10.9	16.3	19.9	24.6	28.8	32.7
2001.....	2.4	4.4	7.3	12.0	14.8	21.0	24.4	29.3	32.3
2000.....	2.2	4.6	7.7	10.9	15.2	19.5	24.6	29.5	33.9
1999.....	2.2	4.3	7.4	10.9	14.4	19.4	23.1	26.9	31.4

Non-Hispanic 4/, male

2003.....	3.1	6.0	10.6	17.3	20.3	27.6	36.0	45.6	49.0
2002.....	3.5	6.5	10.6	16.2	23.3	28.0	36.3	40.0	50.4
2001.....	3.4	6.5	10.8	17.4	21.2	30.1	34.3	42.4	52.6
2000.....	3.1	6.8	11.2	15.8	21.7	27.5	34.5	44.1	55.4
1999.....	3.2	6.1	10.8	15.8	20.4	27.5	33.8	39.2	49.8

Non-Hispanic 4/, female

2003.....	1.1	2.3	3.9	6.0	9.6	12.8	16.9	21.0	26.6
2002.....	1.5	2.3	4.0	6.1	10.1	13.5	16.4	22.0	25.1
2001.....	1.4	2.4	4.0	7.1	9.2	13.7	17.4	21.5	23.8
2000.....	1.4	2.4	4.4	6.4	9.6	13.3	17.7	20.9	25.2
1999.....	1.2	2.5	4.2	6.4	9.4	13.1	15.6	19.7	24.1

Non-Hispanic White, both sexes

2003.....	2.0	4.3	7.2	11.6	15.1	20.1	25.4	31.1	34.0
2002.....	2.5	4.4	7.4	11.2	16.9	20.5	25.0	29.3	33.5
2001.....	2.4	4.4	7.5	12.2	15.4	21.5	24.7	29.7	32.8
2000.....	2.3	4.7	7.9	11.2	15.5	19.9	24.9	29.9	34.6
1999.....	2.1	4.3	7.5	11.3	14.8	19.7	23.3	27.3	31.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)									
Non-Hispanic White, male									
2003.....	3.1	6.1	10.8	17.6	20.9	28.2	36.5	46.9	50.4
2002.....	3.6	6.6	10.8	16.6	24.2	28.6	37.1	40.3	52.4
2001.....	3.6	6.5	11.1	17.3	22.0	30.8	34.3	42.9	54.3
2000.....	3.2	6.9	11.4	16.2	21.7	27.8	34.9	44.5	56.9
1999.....	3.1	6.0	10.9	16.4	20.7	27.7	34.1	40.4	50.2
Non-Hispanic White, female									
2003.....	0.9	2.5	3.8	6.2	9.8	13.5	17.4	21.3	26.8
2002.....	1.4	2.2	4.1	6.2	10.4	13.9	16.3	22.6	25.3
2001.....	1.3	2.4	4.0	7.5	9.6	14.0	17.9	21.7	23.8
2000.....	1.4	2.5	4.6	6.4	10.1	13.5	17.9	21.3	25.7
1999.....	1.2	2.6	4.4	6.5	9.7	13.3	15.7	19.5	24.1
Non-Hispanic Black, both sexes									
2003.....	2.6	4.2	8.4	11.5	13.6	16.3	22.5	26.7	34.2
2002.....	2.9	4.5	7.2	10.9	14.7	17.2	26.2	27.0	28.1
2001.....	2.6	5.1	7.5	12.5	12.0	19.4	24.3	28.7	30.0
2000.....	2.5	4.6	7.5	10.7	15.2	18.7	23.9	29.0	29.6
1999.....	3.0	5.1	7.7	9.9	13.8	19.5	23.4	24.7	31.7
Non-Hispanic Black, male									
2003.....	3.7	6.6	11.7	17.5	19.5	25.9	35.5	40.0	44.7
2002.....	3.6	6.6	12.1	17.3	20.5	24.3	35.2	39.0	39.4
2001.....	3.5	8.0	10.8	21.1	17.5	29.6	39.3	43.3	46.2
2000.....	3.3	7.1	11.5	15.7	25.6	28.4	32.8	48.1	49.3
1999.....	4.3	8.2	12.5	14.1	20.5	28.4	34.8	28.3	49.9
Non-Hispanic Black, female									
2003.....	1.5	2.2	5.7	6.8	9.2	9.8	14.7	19.7	30.1
2002.....	2.3	2.6	3.1	5.9	10.4	12.4	20.7	20.7	23.8
2001.....	1.9	2.7	4.9	5.8	7.9	12.6	15.3	21.2	24.0
2000.....	1.9	2.5	4.3	6.7	7.5	12.3	18.6	19.4	22.2
1999.....	1.9	2.4	3.8	6.5	8.9	13.6	16.6	22.8	24.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of bladder (C67)									
All origins 3/, both sexes									
2003.....	0.8	1.8	3.5	6.9	11.8	18.5	30.0	45.1	64.8
2002.....	0.9	1.7	3.8	6.4	12.0	20.2	31.9	44.1	66.6
2001.....	0.8	1.7	3.4	7.0	12.0	19.0	29.3	44.8	68.9
2000.....	0.8	1.6	3.6	6.8	11.7	19.6	29.1	45.5	69.6
1999.....	0.7	1.9	3.8	7.0	11.8	20.2	29.7	43.8	68.5
All origins 3/, male									
2003.....	1.4	2.7	5.5	10.9	19.1	29.6	51.7	80.3	125.1
2002.....	1.2	2.4	5.9	10.1	19.0	32.2	54.0	76.6	128.9
2001.....	1.2	2.6	5.2	11.3	19.5	31.6	50.8	81.0	134.8
2000.....	1.1	2.5	5.7	10.6	19.0	32.2	50.6	82.4	139.4
1999.....	1.1	2.9	5.9	11.0	19.3	32.9	52.3	79.5	134.8
All origins 3/, female									
2003.....	0.3	0.9	1.7	3.3	5.4	9.5	14.6	23.5	38.1
2002.....	0.5	1.0	1.9	2.9	6.0	10.6	16.3	24.2	39.6
2001.....	0.4	0.9	1.7	3.2	5.6	9.0	14.3	23.2	41.4
2000.....	0.5	0.8	1.7	3.4	5.5	9.7	14.0	23.8	41.2
1999.....	0.3	0.9	1.7	3.4	5.4	10.3	13.9	22.9	41.8
Hispanic, both sexes									
2003.....	*	*	2.0	3.4	6.3	9.8	16.4	27.9	42.0
2002.....	*	*	2.4	3.6	6.1	9.7	14.1	25.4	54.9
2001.....	*	*	*	3.2	6.3	9.9	14.4	26.3	46.3
2000.....	*	*	2.6	3.3	3.5	10.5	14.1	24.0	47.1
1999.....	*	*	2.3	4.1	6.0	11.6	13.2	22.3	38.6
Hispanic, male									
2003.....	*	*	*	5.7	12.0	14.5	24.3	48.9	77.0
2002.....	*	*	*	6.2	10.5	15.8	21.8	38.9	100.4
2001.....	*	*	*	6.0	9.6	14.6	21.1	46.7	87.6
2000.....	*	*	4.6	*	*	16.5	28.8	44.2	84.6
1999.....	*	*	*	7.4	11.6	19.9	21.8	37.7	77.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of bladder (C67)									
Hispanic, female									
2003.....	*	*	*	*	*	*	10.8	14.2	24.2
2002.....	*	*	*	*	*	*	*	16.8	32.3
2001.....	*	*	*	*	*	*	9.7	*	26.1
2000.....	*	*	*	*	*	*	*	*	28.7
1999.....	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes									
2003.....	0.9	1.8	3.6	7.2	12.2	19.0	30.8	45.7	65.6
2002.....	0.9	1.8	3.9	6.5	12.4	20.8	32.8	44.8	66.9
2001.....	0.8	1.8	3.6	7.3	12.4	19.5	30.0	45.5	69.6
2000.....	0.8	1.7	3.7	7.1	12.3	20.1	29.7	46.2	70.3
1999.....	0.8	1.9	3.9	7.2	12.1	20.7	30.3	44.5	69.3
Non-Hispanic 4/, male									
2003.....	1.5	2.8	5.7	11.3	19.6	30.6	53.1	81.6	127.2
2002.....	1.3	2.5	6.1	10.4	19.6	33.2	55.7	78.2	129.8
2001.....	1.2	2.8	5.4	11.6	20.1	32.6	52.2	82.2	136.4
2000.....	1.2	2.6	5.7	11.1	19.9	33.0	51.5	83.5	141.3
1999.....	1.2	3.0	6.0	11.2	19.7	33.6	53.5	81.0	136.6
Non-Hispanic 4/, female									
2003.....	0.3	0.9	1.7	3.4	5.7	9.7	14.8	23.8	38.6
2002.....	0.5	1.1	1.9	3.1	6.2	10.9	16.6	24.5	39.8
2001.....	0.4	0.9	1.8	3.3	5.7	9.1	14.5	23.6	41.9
2000.....	0.5	0.8	1.7	3.4	5.7	9.9	14.4	24.2	41.6
1999.....	0.3	0.9	1.8	3.6	5.6	10.6	14.1	23.2	42.3
Non-Hispanic White, both sexes									
2003.....	0.9	1.9	3.8	7.6	12.7	20.2	31.8	46.9	67.6
2002.....	1.0	1.9	4.0	7.0	13.0	21.6	34.0	46.0	68.0
2001.....	0.8	1.9	3.7	7.6	13.0	20.4	31.0	46.4	71.5
2000.....	0.9	1.8	3.9	7.5	12.9	20.8	30.4	47.1	72.8
1999.....	0.7	2.0	3.9	7.5	12.5	21.3	31.2	45.8	70.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of bladder (C67)									
Non-Hispanic White, male									
2003.....	1.5	3.0	5.9	12.2	20.4	32.7	55.4	84.9	133.6
2002.....	1.4	2.7	6.5	11.1	20.7	34.9	57.9	81.1	135.2
2001.....	1.2	2.9	5.6	12.4	21.4	34.3	54.7	85.0	143.0
2000.....	1.2	2.8	6.0	11.9	21.1	34.5	53.6	85.9	148.0
1999.....	1.1	3.0	6.3	11.7	20.6	35.2	55.6	83.7	143.0
Non-Hispanic White, female									
2003.....	0.3	0.8	1.7	3.4	5.8	10.0	14.8	23.6	38.6
2002.....	0.5	1.0	1.7	3.1	6.1	10.7	16.9	24.5	39.0
2001.....	0.3	0.9	1.8	3.2	5.6	9.2	14.3	23.1	41.7
2000.....	0.5	0.8	1.8	3.4	5.8	9.8	14.0	24.1	42.5
1999.....	0.3	1.0	1.6	3.5	5.4	10.3	14.0	23.5	41.9
Non-Hispanic Black, both sexes									
2003.....	1.2	2.2	3.3	5.1	11.7	12.9	27.0	41.5	53.8
2002.....	*	1.6	4.1	5.1	11.2	19.1	27.0	37.6	64.1
2001.....	0.8	1.8	3.5	7.1	10.3	14.3	24.8	41.7	55.6
2000.....	0.9	1.3	3.0	5.5	9.7	18.0	28.8	44.5	49.6
1999.....	1.0	2.2	4.6	6.8	11.9	19.2	25.4	35.8	60.8
Non-Hispanic Black, male									
2003.....	1.8	2.8	4.6	6.7	18.7	16.5	41.6	62.6	79.9
2002.....	*	*	4.4	6.9	14.6	24.3	42.3	57.7	91.9
2001.....	*	2.9	4.3	9.4	13.0	21.0	34.1	58.9	78.5
2000.....	*	*	4.7	7.4	12.8	24.3	41.4	73.0	88.1
1999.....	*	3.8	5.6	9.6	15.7	23.9	39.2	60.2	86.8
Non-Hispanic Black, female									
2003.....	*	1.8	*	3.9	6.6	10.5	18.3	30.4	43.7
2002.....	*	*	3.8	3.7	8.7	15.6	17.9	27.2	53.5
2001.....	*	*	2.8	5.3	8.3	9.9	19.2	32.9	47.1
2000.....	*	*	*	4.0	7.5	13.8	21.2	30.2	35.3
1999.....	*	*	3.8	4.6	9.1	16.1	17.2	23.7	51.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All origins 3/, both sexes

2003.....	3.9	5.8	8.5	11.2	13.8	17.3	19.4	19.6	15.2
2002.....	4.2	6.2	8.5	11.1	14.7	17.8	19.6	18.5	16.4
2001.....	4.0	6.1	8.3	11.6	14.2	16.7	20.3	20.6	16.4
2000.....	4.1	6.3	8.2	11.4	14.8	18.3	19.4	21.1	16.9
1999.....	4.3	6.1	9.2	11.7	15.7	18.3	21.0	19.4	16.0

All origins 3/, male

2003.....	5.1	7.4	9.7	13.7	16.4	21.3	24.0	25.4	20.0
2002.....	5.3	7.2	10.6	13.1	18.4	22.2	24.4	22.9	20.1
2001.....	4.9	7.5	10.7	13.9	17.3	21.2	25.4	26.5	20.1
2000.....	4.9	7.7	10.1	14.0	17.9	22.8	23.2	26.3	22.4
1999.....	5.2	7.4	10.6	14.7	19.5	22.3	26.1	24.5	20.7

All origins 3/, female

2003.....	2.8	4.3	7.3	9.0	11.6	14.1	16.2	16.0	13.1
2002.....	3.1	5.1	6.5	9.2	11.4	14.2	16.2	15.8	14.8
2001.....	3.1	4.8	6.0	9.5	11.5	13.1	16.7	17.1	14.8
2000.....	3.3	4.9	6.5	9.0	12.1	14.8	16.7	18.0	14.7
1999.....	3.4	4.9	7.9	8.9	12.4	15.1	17.5	16.5	14.1

Hispanic, both sexes

2003.....	2.0	3.2	4.7	7.6	8.2	11.1	16.1	13.9	*
2002.....	2.4	4.6	4.7	7.4	9.1	11.1	14.4	12.5	13.3
2001.....	2.2	3.0	4.5	7.8	9.5	11.3	13.3	17.4	*
2000.....	2.3	3.7	3.7	9.3	9.7	9.8	16.2	*	15.9
1999.....	2.2	4.0	5.2	7.2	8.1	12.5	12.3	17.0	*

Hispanic, male

2003.....	2.7	3.8	4.8	8.4	10.1	14.1	15.2	*	*
2002.....	2.9	6.4	5.0	9.0	12.2	12.7	17.3	*	*
2001.....	3.0	2.9	6.2	7.6	11.3	18.3	16.4	*	*
2000.....	2.6	*	4.4	12.1	11.9	13.1	19.9	*	*
1999.....	*	3.8	5.5	10.7	9.7	11.7	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Hispanic, female

2003.....	*	2.6	4.6	6.9	6.7	8.9	16.8	*	*
2002.....	1.9	2.9	4.4	6.0	6.6	9.8	12.3	*	*
2001.....	*	3.2	*	8.0	8.0	*	11.1	*	*
2000.....	*	4.7	*	6.9	7.9	7.4	13.6	*	*
1999.....	*	4.1	4.8	*	6.9	13.0	*	*	*

Non-Hispanic 4/, both sexes

2003.....	4.1	6.0	8.8	11.5	14.2	17.7	19.6	19.8	15.4
2002.....	4.4	6.3	8.8	11.3	15.1	18.2	19.8	18.7	16.5
2001.....	4.2	6.3	8.6	11.8	14.5	17.1	20.6	20.7	16.5
2000.....	4.2	6.5	8.6	11.5	15.1	18.8	19.5	21.3	16.9
1999.....	4.5	6.3	9.5	11.9	16.2	18.5	21.3	19.5	16.1

Non-Hispanic 4/, male

2003.....	5.3	7.7	10.1	14.1	16.8	21.8	24.5	25.7	20.1
2002.....	5.6	7.3	11.1	13.3	18.9	22.8	24.7	23.3	20.2
2001.....	5.1	7.9	11.0	14.3	17.7	21.4	25.8	26.7	20.5
2000.....	5.1	8.1	10.6	14.1	18.2	23.3	23.3	26.5	22.3
1999.....	5.5	7.7	10.9	14.9	20.1	22.9	26.5	24.6	21.3

Non-Hispanic 4/, female

2003.....	3.0	4.4	7.6	9.2	12.0	14.4	16.1	16.2	13.4
2002.....	3.3	5.3	6.7	9.5	11.8	14.5	16.4	15.9	14.9
2001.....	3.3	4.9	6.3	9.5	11.7	13.6	16.9	17.1	14.9
2000.....	3.4	4.9	6.7	9.1	12.4	15.2	16.8	18.3	14.7
1999.....	3.4	5.0	8.2	9.2	12.8	15.1	17.8	16.5	14.0

Non-Hispanic White, both sexes

2003.....	4.5	6.6	9.5	12.6	15.4	19.0	20.9	20.9	15.9
2002.....	4.9	6.9	9.5	12.3	16.4	19.5	20.9	19.6	17.3
2001.....	4.6	7.0	9.2	12.9	15.6	18.2	22.0	21.8	17.2
2000.....	4.7	7.2	9.2	12.5	16.4	20.3	20.6	22.1	17.5
1999.....	5.0	6.8	10.4	12.9	17.4	20.0	22.6	20.4	16.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic White, male

2003.....	5.9	8.4	10.8	15.4	17.9	23.2	25.8	26.9	21.0
2002.....	6.2	8.0	11.7	14.4	20.5	24.3	25.9	24.2	21.4
2001.....	5.6	8.6	11.8	15.4	19.1	22.8	27.2	28.0	20.9
2000.....	5.7	8.7	11.4	15.3	19.8	24.9	24.5	27.7	23.5
1999.....	6.0	8.2	11.8	16.0	21.6	24.6	27.9	25.9	22.4

Non-Hispanic White, female

2003.....	3.2	5.0	8.2	10.0	13.1	15.6	17.3	17.2	13.7
2002.....	3.7	5.8	7.3	10.5	12.8	15.6	17.4	16.9	15.5
2001.....	3.6	5.4	6.7	10.6	12.4	14.5	18.3	18.0	15.7
2000.....	3.7	5.6	7.2	9.9	13.4	16.6	17.8	18.8	15.1
1999.....	3.9	5.5	9.0	10.1	13.8	16.4	18.8	17.1	14.4

Non-Hispanic Black, both sexes

2003.....	2.7	3.7	5.7	5.6	8.3	9.8	10.6	10.6	10.7
2002.....	2.0	3.7	5.7	5.5	7.8	9.6	12.1	11.5	9.7
2001.....	2.7	3.5	5.5	6.3	9.2	9.6	9.2	11.0	11.9
2000.....	2.2	3.7	5.5	6.2	7.6	9.0	10.5	14.7	10.5
1999.....	2.7	4.0	5.5	6.9	8.7	7.6	12.0	11.2	10.0

Non-Hispanic Black, male

2003.....	3.1	5.3	6.8	6.7	11.6	12.1	16.4	*	*
2002.....	2.2	3.3	8.3	7.7	9.0	12.0	15.0	17.9	*
2001.....	3.1	4.6	7.1	9.0	9.4	11.7	11.9	*	*
2000.....	2.2	5.7	6.6	7.9	7.7	12.6	12.0	*	*
1999.....	3.8	5.3	6.1	8.9	11.2	9.0	14.5	*	*

Non-Hispanic Black, female

2003.....	2.3	2.3	4.8	4.6	5.8	8.2	7.2	8.7	12.0
2002.....	1.8	4.1	3.5	3.8	7.0	8.0	10.4	8.1	10.5
2001.....	2.3	2.7	4.2	4.3	9.0	8.3	7.6	10.4	9.8
2000.....	2.3	2.0	4.6	4.9	7.5	6.6	9.6	14.3	10.5
1999.....	1.7	2.9	5.0	5.3	6.9	6.7	10.5	12.3	9.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All origins 3/, both sexes

2003.....	7.2	11.7	20.1	35.5	55.2	87.3	127.4	165.2	198.0
2002.....	7.5	12.6	21.5	36.9	59.2	90.1	125.6	168.8	197.5
2001.....	8.1	13.0	22.7	38.1	58.8	91.8	129.0	168.5	199.9
2000.....	8.3	13.4	23.4	38.2	62.2	91.6	131.4	176.0	204.6
1999.....	8.4	13.6	25.6	39.4	62.8	93.9	130.7	174.0	199.4

All origins 3/, male

2003.....	8.8	14.5	25.2	45.6	71.9	112.3	163.5	216.8	268.7
2002.....	9.1	15.1	26.8	45.7	76.4	116.8	164.3	224.2	269.1
2001.....	9.8	16.0	27.7	47.5	75.2	116.9	165.7	220.9	275.3
2000.....	10.3	16.1	28.4	45.8	78.3	117.7	169.8	230.6	281.8
1999.....	10.2	16.7	31.0	48.6	78.1	118.9	169.4	221.8	279.0

All origins 3/, female

2003.....	5.5	9.1	15.3	26.4	40.8	67.3	101.7	133.6	166.8
2002.....	6.0	10.1	16.4	28.9	44.4	68.8	98.2	135.1	166.5
2001.....	6.4	10.1	18.1	29.5	44.6	71.8	103.4	137.1	168.4
2000.....	6.5	10.8	18.8	31.3	48.4	71.0	104.7	143.8	173.1
1999.....	6.6	10.6	20.5	31.0	49.8	74.3	103.8	145.9	167.3

Hispanic, both sexes

2003.....	7.3	9.9	17.1	24.9	38.5	58.0	91.3	111.9	129.6
2002.....	6.5	9.3	18.1	27.6	47.3	61.7	89.8	119.0	132.5
2001.....	6.1	10.9	17.0	33.1	43.0	64.5	98.0	112.7	134.4
2000.....	7.8	10.4	17.9	34.0	45.7	75.2	97.0	114.7	143.3
1999.....	8.4	12.3	20.0	29.5	49.6	69.7	88.1	124.5	131.9

Hispanic, male

2003.....	8.3	11.2	19.9	30.8	46.5	71.7	107.6	126.8	149.3
2002.....	7.3	11.2	21.3	33.1	56.5	72.0	108.1	138.4	180.7
2001.....	7.5	12.7	21.6	38.8	56.3	81.9	139.8	146.7	180.9
2000.....	10.7	11.6	23.2	40.6	58.2	88.5	114.4	151.7	197.5
1999.....	10.0	14.6	23.6	36.9	56.0	85.0	106.9	139.9	135.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)									
Hispanic, female									
2003.....	6.3	8.7	14.6	19.7	32.0	47.5	79.8	102.2	119.5
2002.....	5.7	7.5	15.2	22.8	39.8	53.7	76.8	106.6	108.6
2001.....	4.8	9.1	12.8	28.2	32.1	51.3	68.3	91.7	111.7
2000.....	4.8	9.3	13.1	28.3	35.6	65.2	84.7	92.3	116.7
1999.....	6.8	10.2	16.7	23.1	44.4	58.2	74.9	115.3	130.0
Non-Hispanic 4/, both sexes									
2003.....	7.1	11.8	20.3	36.3	56.4	89.1	129.2	167.4	200.5
2002.....	7.6	12.8	21.7	37.5	59.9	91.7	127.2	170.5	199.8
2001.....	8.2	13.1	23.1	38.3	59.7	93.2	130.3	170.5	202.1
2000.....	8.4	13.6	23.8	38.4	63.2	92.3	132.7	178.1	206.5
1999.....	8.4	13.7	25.9	40.0	63.6	95.1	132.2	175.3	201.4
Non-Hispanic 4/, male									
2003.....	8.9	14.7	25.5	46.6	73.6	114.9	166.3	220.9	273.6
2002.....	9.2	15.4	27.2	46.6	77.5	119.2	166.8	227.5	272.8
2001.....	10.0	16.2	28.1	48.0	76.3	118.7	166.4	223.6	278.8
2000.....	10.2	16.4	28.7	46.0	79.4	119.1	172.1	233.3	285.0
1999.....	10.2	16.9	31.4	49.4	79.3	120.5	171.8	224.3	284.3
Non-Hispanic 4/, female									
2003.....	5.4	9.1	15.3	26.8	41.4	68.4	102.8	134.8	168.4
2002.....	6.0	10.3	16.5	29.3	44.6	69.7	99.2	135.9	168.3
2001.....	6.5	10.2	18.5	29.5	45.4	72.9	105.0	138.7	170.2
2000.....	6.6	10.9	19.2	31.4	49.2	71.2	105.3	145.6	174.8
1999.....	6.6	10.6	20.8	31.4	50.1	75.2	104.7	146.7	168.2
Non-Hispanic White, both sexes									
2003.....	6.9	11.4	20.3	36.5	57.3	90.7	132.3	172.1	207.2
2002.....	7.3	12.3	21.2	37.6	61.0	93.5	130.4	175.5	206.5
2001.....	8.0	12.9	22.6	38.2	60.0	95.2	133.3	174.9	208.2
2000.....	8.0	13.3	23.6	38.6	64.5	94.1	135.7	183.7	210.7
1999.....	8.2	13.4	25.6	39.8	64.6	96.9	134.7	179.9	206.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)									
Non-Hispanic White, male									
2003.....	8.6	14.1	25.4	47.3	75.4	118.0	171.9	228.8	284.6
2002.....	8.9	15.0	26.8	46.6	79.1	121.8	171.7	234.5	283.5
2001.....	9.9	16.0	27.4	47.9	76.7	121.2	170.6	229.3	288.7
2000.....	9.8	15.9	28.4	46.2	81.5	120.4	176.5	240.8	293.8
1999.....	9.9	16.6	31.0	49.4	80.6	122.7	174.6	231.2	291.8
Non-Hispanic White, female									
2003.....	5.2	8.7	15.3	26.5	41.2	68.2	103.7	137.2	173.1
2002.....	5.6	9.7	15.8	29.2	45.0	70.3	100.9	139.5	173.2
2001.....	6.2	9.8	18.0	29.3	45.2	74.0	106.8	142.1	174.6
2000.....	6.2	10.8	19.0	31.5	49.5	73.0	107.0	149.8	177.2
1999.....	6.5	10.2	20.5	30.9	50.5	76.3	106.6	149.7	172.4
Non-Hispanic Black, both sexes									
2003.....	9.8	17.7	24.0	42.2	60.5	91.7	118.4	140.8	152.3
2002.....	10.8	17.5	30.3	44.3	63.1	94.6	113.6	138.5	153.7
2001.....	10.7	16.9	31.0	44.9	68.4	92.2	117.1	144.0	157.3
2000.....	12.6	18.0	29.7	44.8	63.5	91.5	115.8	132.8	177.1
1999.....	10.6	18.5	33.7	46.5	65.8	91.8	120.8	140.0	162.3
Non-Hispanic Black, male									
2003.....	12.2	23.1	32.7	52.6	74.9	106.8	138.3	169.8	188.6
2002.....	13.5	20.0	36.8	56.4	80.1	125.2	140.4	180.2	203.5
2001.....	12.1	21.4	38.5	57.1	89.4	115.7	142.2	194.6	206.7
2000.....	15.0	22.4	37.6	55.1	74.9	127.1	149.8	170.8	226.6
1999.....	13.4	22.8	42.2	56.2	82.1	116.3	161.0	166.5	221.1
Non-Hispanic Black, female									
2003.....	7.6	13.2	16.9	34.0	49.7	81.6	106.5	125.6	138.3
2002.....	8.5	15.4	25.0	34.8	50.5	74.1	97.6	116.9	134.6
2001.....	9.5	13.0	24.8	35.3	52.9	76.6	102.1	118.3	139.0
2000.....	10.5	14.3	23.3	36.7	55.0	67.8	95.2	113.6	158.7
1999.....	8.2	14.8	26.9	38.9	53.7	75.6	96.7	126.9	140.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hodgkin's disease (C81)									
Hispanic, female									
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes									
2003.....	0.3	0.4	0.6	0.9	0.8	1.5	2.0	2.3	2.3
2002.....	0.4	0.5	0.5	0.8	1.0	1.6	1.7	2.2	2.8
2001.....	0.5	0.5	0.4	0.7	1.0	1.2	2.1	2.5	2.2
2000.....	0.4	0.5	0.5	0.8	1.0	1.6	1.7	2.3	2.4
1999.....	0.4	0.5	0.6	0.8	1.2	1.8	2.0	2.2	2.3
Non-Hispanic 4/, male									
2003.....	0.4	0.5	0.8	1.1	1.1	1.9	2.3	2.9	2.7
2002.....	0.5	0.5	0.7	1.0	1.4	2.2	2.0	3.1	3.7
2001.....	0.6	0.6	0.5	0.9	1.4	1.6	2.5	2.9	2.5
2000.....	0.4	0.6	0.5	1.0	1.3	1.9	2.1	2.6	3.7
1999.....	0.5	0.5	0.8	1.2	1.6	2.2	2.2	2.5	3.7
Non-Hispanic 4/, female									
2003.....	0.2	0.4	0.5	0.6	0.6	1.2	1.9	2.0	2.2
2002.....	0.3	0.5	*	0.7	0.6	1.1	1.5	1.6	2.4
2001.....	0.4	0.4	0.3	0.6	0.7	0.9	1.8	2.2	2.0
2000.....	0.3	0.4	0.5	0.6	0.7	1.3	1.4	2.1	2.0
1999.....	0.3	0.4	0.5	0.5	0.9	1.5	1.8	2.0	1.7
Non-Hispanic White, both sexes									
2003.....	0.4	0.5	0.6	0.9	0.8	1.6	2.1	2.4	2.5
2002.....	0.4	0.5	0.5	0.9	1.0	1.7	1.8	2.3	2.9
2001.....	0.5	0.5	0.4	0.8	1.2	1.2	2.1	2.6	2.3
2000.....	0.4	0.5	0.5	0.8	1.1	1.6	1.9	2.4	2.6
1999.....	0.4	0.5	0.7	0.9	1.2	1.9	2.1	2.3	2.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)									
All origins 3/, both sexes									
2003.....	3.0	4.7	8.1	14.3	21.6	32.8	49.3	65.4	78.8
2002.....	3.0	5.0	8.8	14.9	23.2	35.0	50.2	67.0	77.3
2001.....	3.4	5.5	9.6	15.6	23.0	36.0	50.9	69.8	79.6
2000.....	3.6	5.8	10.6	16.1	25.0	36.3	53.7	71.8	80.2
1999.....	3.5	6.2	10.8	16.3	25.6	38.4	54.6	72.6	77.7
All origins 3/, male									
2003.....	3.9	6.0	10.1	18.7	27.8	41.0	60.7	81.8	103.6
2002.....	3.9	6.2	11.3	18.4	30.0	42.8	61.8	84.4	99.7
2001.....	4.4	7.4	11.9	19.1	29.2	44.7	61.3	87.3	106.6
2000.....	4.7	7.5	13.3	19.2	30.5	45.2	66.3	88.1	105.9
1999.....	4.7	7.9	13.5	19.6	30.8	46.8	68.1	85.7	104.4
All origins 3/, female									
2003.....	2.1	3.5	6.1	10.4	16.2	26.1	41.2	55.4	67.8
2002.....	2.2	3.8	6.4	11.7	17.3	28.7	41.9	56.4	67.6
2001.....	2.5	3.8	7.5	12.4	17.7	29.1	43.6	59.3	68.3
2000.....	2.6	4.1	8.2	13.3	20.3	29.2	45.0	62.2	69.8
1999.....	2.4	4.5	8.3	13.3	21.1	31.8	45.2	65.0	66.9
Hispanic, both sexes									
2003.....	2.8	3.4	7.7	9.4	15.4	23.7	35.8	48.0	54.4
2002.....	2.5	3.5	6.7	13.1	18.6	27.9	41.3	47.7	50.4
2001.....	2.9	4.5	7.9	14.9	14.4	23.2	41.6	46.2	59.4
2000.....	2.9	4.6	8.1	14.9	18.5	34.8	39.2	45.7	59.7
1999.....	3.8	5.2	6.8	12.5	20.2	31.5	37.8	51.7	56.1
Hispanic, male									
2003.....	3.1	3.8	9.4	12.1	18.8	30.3	40.7	53.0	57.0
2002.....	3.3	4.0	8.4	16.2	23.4	29.4	42.2	51.5	60.2
2001.....	3.6	5.4	10.0	17.7	19.5	30.7	55.9	63.6	68.6
2000.....	3.8	5.7	10.5	18.4	22.4	40.4	44.3	55.9	82.6
1999.....	5.0	6.7	8.8	15.8	21.6	35.1	46.0	51.9	60.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)									
Hispanic, female									
2003.....	2.4	3.0	6.2	7.1	12.5	18.7	32.3	44.7	53.1
2002.....	*	3.1	5.1	10.5	14.7	26.7	40.7	45.3	45.6
2001.....	2.1	3.7	6.0	12.5	10.3	17.4	31.5	35.4	54.9
2000.....	*	3.4	6.0	11.9	15.4	30.6	35.6	39.4	48.5
1999.....	2.6	3.8	5.0	9.8	19.1	28.7	31.9	51.6	54.1
Non-Hispanic 4/, both sexes									
2003.....	3.0	4.8	8.1	14.7	22.0	33.3	50.0	66.1	79.6
2002.....	3.0	5.1	9.0	15.0	23.5	35.4	50.5	67.5	78.2
2001.....	3.5	5.6	9.8	15.6	23.5	36.7	51.2	70.6	80.3
2000.....	3.7	5.8	10.8	16.1	25.4	36.3	54.3	72.7	80.9
1999.....	3.5	6.2	11.1	16.5	25.9	38.7	55.2	73.2	78.3
Non-Hispanic 4/, male									
2003.....	3.9	6.2	10.1	19.1	28.5	41.7	61.8	83.1	105.6
2002.....	3.9	6.4	11.5	18.5	30.4	43.5	62.7	85.5	101.4
2001.....	4.5	7.5	12.1	19.1	29.7	45.5	61.3	88.2	108.1
2000.....	4.8	7.6	13.4	19.1	31.0	45.3	67.2	89.3	106.7
1999.....	4.7	8.0	13.8	19.9	31.3	47.3	68.9	86.5	106.0
Non-Hispanic 4/, female									
2003.....	2.1	3.5	6.1	10.6	16.4	26.6	41.6	55.8	68.2
2002.....	2.2	3.9	6.5	11.8	17.5	28.8	42.0	56.6	68.2
2001.....	2.5	3.8	7.6	12.4	18.1	29.8	44.1	60.1	68.7
2000.....	2.6	4.2	8.3	13.4	20.7	29.1	45.3	62.9	70.4
1999.....	2.3	4.6	8.6	13.5	21.2	32.0	45.6	65.3	67.2
Non-Hispanic White, both sexes									
2003.....	2.9	4.8	8.5	15.2	23.3	34.9	52.6	69.9	84.3
2002.....	3.1	5.0	9.1	15.8	24.7	36.9	53.4	71.1	83.1
2001.....	3.4	5.7	9.9	16.3	24.5	38.7	54.0	74.0	84.1
2000.....	3.7	6.0	11.4	16.9	26.8	38.1	57.1	76.9	85.1
1999.....	3.5	6.5	11.5	17.4	27.5	40.6	57.6	77.3	82.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)									
Non-Hispanic White, male									
2003.....	3.8	6.2	10.6	19.8	30.3	43.9	65.2	88.1	112.0
2002.....	4.0	6.4	11.8	19.2	31.9	45.3	66.1	89.8	107.6
2001.....	4.5	7.6	12.1	19.8	30.9	47.5	64.3	92.2	113.1
2000.....	4.7	7.6	13.9	19.9	32.8	46.9	70.8	94.1	113.0
1999.....	4.7	8.2	14.1	20.6	32.9	49.5	71.1	91.8	110.0
Non-Hispanic White, female									
2003.....	2.1	3.5	6.4	10.9	17.0	27.5	43.6	58.7	72.2
2002.....	2.1	3.7	6.6	12.5	18.4	30.0	44.3	59.7	72.6
2001.....	2.4	3.9	7.8	13.1	18.8	31.6	46.7	63.0	72.0
2000.....	2.6	4.4	9.0	14.1	21.6	31.1	47.6	66.6	73.8
1999.....	2.3	4.8	9.0	14.4	22.7	33.4	48.2	68.8	70.8
Non-Hispanic Black, both sexes									
2003.....	4.0	6.1	6.2	12.9	14.9	22.9	27.0	31.1	31.2
2002.....	3.6	6.2	9.4	12.6	18.0	26.2	26.9	35.0	31.2
2001.....	4.3	6.1	9.9	12.1	19.3	24.0	25.9	38.1	40.0
2000.....	4.5	5.6	9.1	12.8	17.0	22.9	26.3	28.5	36.6
1999.....	4.1	5.6	10.8	11.6	17.3	24.4	32.5	31.4	40.4
Non-Hispanic Black, male									
2003.....	5.8	7.7	8.0	16.5	17.7	24.6	29.0	34.7	35.2
2002.....	4.2	7.6	11.7	16.9	23.3	34.9	31.9	42.1	43.8
2001.....	5.1	8.8	12.4	15.5	25.0	32.4	30.8	48.2	55.4
2000.....	6.2	8.0	13.9	15.9	19.7	32.5	31.3	32.6	43.4
1999.....	5.6	7.9	14.1	15.7	23.9	29.1	47.9	28.3	59.4
Non-Hispanic Black, female									
2003.....	2.4	4.7	4.8	10.2	12.8	21.8	25.8	29.2	29.7
2002.....	3.1	5.0	7.5	9.2	14.0	20.4	23.8	31.2	26.3
2001.....	3.7	3.7	7.9	9.5	15.0	18.4	22.9	32.9	34.2
2000.....	3.1	3.6	5.1	10.4	14.9	16.6	23.2	26.4	34.0
1999.....	2.9	3.8	8.1	8.4	12.4	21.4	23.3	32.9	33.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)									
All origins 3/, both sexes									
2003.....	2.7	4.0	7.0	12.4	19.6	32.5	47.4	62.9	80.3
2002.....	2.8	4.5	7.4	12.9	20.9	32.4	45.7	63.4	80.5
2001.....	3.0	4.6	7.6	13.1	20.6	33.3	47.7	61.3	81.4
2000.....	3.1	4.5	7.7	12.7	21.7	33.2	47.4	65.1	83.6
1999.....	3.2	4.4	8.5	13.1	21.8	33.1	46.3	62.7	82.4
All origins 3/, male									
2003.....	3.2	4.9	8.9	15.8	26.8	44.9	66.3	88.8	111.8
2002.....	3.2	5.3	9.2	16.1	28.0	45.8	65.0	90.0	114.5
2001.....	3.4	5.2	9.2	17.4	27.6	44.9	66.2	86.9	117.5
2000.....	3.6	5.1	9.7	16.0	29.2	46.2	65.1	93.4	119.8
1999.....	3.6	5.2	10.1	16.8	29.0	45.7	64.6	87.6	119.2
All origins 3/, female									
2003.....	2.2	3.2	5.2	9.4	13.3	22.7	33.9	47.0	66.4
2002.....	2.5	3.8	5.7	10.0	14.7	21.8	32.1	47.3	65.7
2001.....	2.7	4.0	6.0	9.3	14.6	24.1	34.8	45.9	66.3
2000.....	2.6	3.9	5.9	9.7	15.3	22.9	35.1	48.4	68.8
1999.....	2.8	3.6	7.0	9.8	15.6	23.2	33.6	48.1	67.6
Hispanic, both sexes									
2003.....	3.5	3.8	4.7	8.6	11.9	16.5	29.0	39.3	47.2
2002.....	2.5	2.7	6.5	8.2	13.4	18.1	21.6	37.0	47.1
2001.....	2.6	3.5	4.5	9.2	14.3	21.2	32.0	35.8	42.5
2000.....	3.3	3.8	6.4	9.6	13.5	21.2	29.1	39.0	49.1
1999.....	3.0	3.9	6.8	7.3	14.9	20.3	27.4	37.6	44.2
Hispanic, male									
2003.....	4.2	3.9	5.0	11.3	14.6	23.1	35.9	48.9	61.6
2002.....	2.1	3.7	6.6	8.0	15.2	22.8	32.0	50.3	65.3
2001.....	3.1	3.6	5.2	12.8	16.3	23.8	49.1	39.0	68.6
2000.....	4.3	3.9	9.0	11.8	16.8	25.8	35.4	57.4	76.6
1999.....	3.6	3.5	7.2	10.4	17.4	31.6	35.1	47.2	49.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)									
Hispanic, female									
2003.....	2.9	3.6	4.4	6.2	9.6	11.5	24.1	33.2	39.8
2002.....	2.8	*	6.4	8.5	11.9	14.5	14.2	28.5	38.1
2001.....	*	3.3	3.8	6.1	12.6	19.2	19.8	33.8	29.8
2000.....	2.3	3.7	4.0	7.7	10.9	17.7	24.6	27.8	35.6
1999.....	2.4	4.3	6.5	*	12.8	11.9	22.0	31.9	41.6
Non-Hispanic 4/, both sexes									
2003.....	2.6	4.0	7.1	12.7	20.1	33.5	48.4	63.9	81.6
2002.....	2.9	4.7	7.5	13.3	21.4	33.2	46.9	64.5	81.7
2001.....	3.1	4.7	7.8	13.4	21.0	33.9	48.4	62.2	82.7
2000.....	3.1	4.5	7.8	12.9	22.2	33.8	48.1	66.0	84.7
1999.....	3.2	4.4	8.6	13.5	22.2	33.7	47.0	63.5	83.7
Non-Hispanic 4/, male									
2003.....	3.1	4.9	9.2	16.1	27.7	46.2	67.9	90.7	113.7
2002.....	3.3	5.4	9.4	16.7	28.8	47.0	66.5	91.7	116.5
2001.....	3.4	5.3	9.5	17.7	28.3	46.0	66.9	88.7	119.3
2000.....	3.5	5.1	9.6	16.3	30.0	47.3	66.4	94.5	121.5
1999.....	3.6	5.3	10.2	17.2	29.6	46.3	65.8	89.0	122.1
Non-Hispanic 4/, female									
2003.....	2.2	3.2	5.2	9.6	13.6	23.4	34.5	47.6	67.4
2002.....	2.5	4.0	5.7	10.1	14.9	22.2	33.0	48.0	66.6
2001.....	2.8	4.1	6.2	9.5	14.7	24.3	35.5	46.4	67.5
2000.....	2.6	4.0	6.0	9.8	15.5	23.1	35.4	49.2	69.8
1999.....	2.8	3.5	7.1	10.1	15.8	23.8	34.0	48.5	68.3
Non-Hispanic White, both sexes									
2003.....	2.7	4.0	7.3	13.0	21.0	34.8	50.4	66.2	84.7
2002.....	2.7	4.7	7.4	13.4	22.4	34.5	48.3	66.9	84.4
2001.....	3.1	4.6	7.9	13.7	21.4	35.2	49.8	64.8	86.1
2000.....	2.9	4.5	7.8	13.1	23.2	35.1	49.9	68.5	87.3
1999.....	3.2	4.3	8.8	13.7	22.6	34.9	48.8	65.0	86.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)									
Non-Hispanic White, male									
2003.....	3.3	4.8	9.3	16.8	28.7	48.0	71.1	94.8	119.0
2002.....	3.1	5.5	9.4	16.9	29.9	48.5	68.7	95.1	121.3
2001.....	3.5	5.2	9.7	18.1	28.6	47.8	69.0	92.0	125.3
2000.....	3.4	5.0	9.7	16.5	31.3	48.6	68.9	97.9	125.5
1999.....	3.5	5.2	10.5	17.5	30.2	47.9	68.0	91.7	127.2
Non-Hispanic White, female									
2003.....	2.2	3.2	5.4	9.6	14.0	23.9	35.4	48.6	69.6
2002.....	2.4	4.0	5.4	10.2	15.6	23.1	33.8	49.6	68.4
2001.....	2.7	3.9	6.2	9.7	15.0	25.0	36.2	48.4	69.8
2000.....	2.5	4.0	6.0	9.9	16.0	24.2	36.6	51.1	72.0
1999.....	2.9	3.4	7.2	10.1	15.9	24.6	35.3	49.2	70.8
Non-Hispanic Black, both sexes									
2003.....	2.6	5.2	7.2	13.1	18.2	29.9	37.6	48.7	60.4
2002.....	3.8	5.1	9.4	14.2	17.4	29.1	40.0	47.2	65.3
2001.....	3.1	5.7	8.1	12.5	22.4	28.4	41.8	42.5	57.1
2000.....	4.7	5.3	8.1	14.3	18.1	29.0	35.3	44.3	66.1
1999.....	3.6	5.5	8.3	14.0	22.6	28.3	35.6	51.4	55.9
Non-Hispanic Black, male									
2003.....	2.8	6.9	10.9	14.9	25.1	40.1	50.5	59.6	78.8
2002.....	5.0	5.6	11.4	17.7	23.8	43.4	53.5	65.5	87.5
2001.....	2.9	6.8	10.1	17.6	32.6	36.8	52.6	62.1	75.1
2000.....	5.1	6.3	10.2	18.9	22.9	44.4	50.6	65.2	98.6
1999.....	4.6	7.1	10.6	17.6	29.5	38.4	52.2	62.0	78.4
Non-Hispanic Black, female									
2003.....	2.5	3.8	4.3	11.7	13.1	22.9	30.0	43.0	53.2
2002.....	2.7	4.7	7.8	11.5	12.7	19.5	32.0	37.7	56.9
2001.....	3.4	4.8	6.5	8.5	14.8	22.7	35.4	32.5	50.5
2000.....	4.3	4.5	6.4	10.6	14.5	18.8	26.1	33.7	54.1
1999.....	2.7	4.2	6.4	11.1	17.5	21.6	25.6	46.1	47.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)									
All origins 3/, both sexes									
2003.....	1.1	2.5	4.4	7.9	13.1	20.4	28.5	34.5	36.3
2002.....	1.3	2.5	4.7	8.3	14.1	21.0	27.9	35.8	36.7
2001.....	1.2	2.3	5.1	8.5	14.0	21.1	28.3	34.7	36.3
2000.....	1.2	2.6	4.5	8.5	14.4	20.4	28.3	36.4	37.8
1999.....	1.2	2.5	5.6	9.0	14.1	20.6	27.6	36.2	36.6
All origins 3/, male									
2003.....	1.3	3.1	5.4	10.1	16.0	24.3	33.9	43.3	50.4
2002.....	1.5	3.0	5.6	10.2	16.8	25.8	35.5	46.2	50.8
2001.....	1.5	2.8	6.0	10.1	16.8	25.3	35.6	43.4	48.0
2000.....	1.4	2.9	4.9	9.5	17.1	24.2	35.7	46.1	51.8
1999.....	1.4	3.0	6.5	11.0	16.6	24.1	34.1	45.5	51.3
All origins 3/, female									
2003.....	0.9	1.9	3.5	6.0	10.6	17.2	24.7	29.2	30.1
2002.....	1.0	2.0	4.0	6.5	11.7	17.2	22.5	29.5	30.6
2001.....	0.9	1.9	4.2	7.1	11.5	17.7	23.2	29.4	31.3
2000.....	1.0	2.3	4.2	7.6	12.1	17.3	23.1	30.7	32.2
1999.....	1.1	2.1	4.7	7.3	12.0	17.8	23.0	30.7	30.6
Hispanic, both sexes									
2003.....	*	2.3	4.4	6.4	10.1	15.8	24.2	23.4	24.9
2002.....	1.2	2.7	4.3	5.9	14.3	14.3	25.1	29.4	33.3
2001.....	*	2.2	3.9	8.3	12.2	18.2	22.7	28.3	29.4
2000.....	1.2	1.5	3.2	7.9	13.0	17.4	25.1	23.4	31.2
1999.....	1.2	2.3	5.6	8.3	12.1	15.1	19.4	30.0	28.1
Hispanic, male									
2003.....	*	2.8	5.1	6.6	11.7	15.4	27.3	22.9	*
2002.....	*	2.8	5.5	8.2	17.3	18.0	32.6	29.7	53.5
2001.....	*	4.8	7.6	17.4	24.2	32.7	41.6	38.1	
2000.....	*	*	*	8.1	17.5	19.4	28.1	33.9	*
1999.....	*	*	6.7	9.2	14.3	15.3	21.1	31.4	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Hispanic, female

2003.....	*	*	3.6	6.2	8.8	16.1	22.0	23.7	22.7
2002.....	*	2.5	*	*	11.9	11.5	19.7	29.2	23.2
2001.....	*	*	*	8.9	8.0	13.6	15.5	20.1	25.1
2000.....	*	*	*	7.7	9.4	15.8	23.0	*	30.7
1999.....	*	*	4.6	7.4	10.3	14.9	18.2	29.1	31.2

Non-Hispanic 4/, both sexes

2003.....	1.1	2.5	4.4	8.0	13.3	20.6	28.7	35.0	36.7
2002.....	1.3	2.5	4.8	8.4	14.0	21.4	27.9	36.1	36.7
2001.....	1.2	2.3	5.2	8.5	14.1	21.2	28.5	34.8	36.5
2000.....	1.2	2.7	4.6	8.5	14.5	20.5	28.4	36.8	38.0
1999.....	1.2	2.5	5.5	9.1	14.2	20.8	27.9	36.3	36.8

Non-Hispanic 4/, male

2003.....	1.4	3.1	5.4	10.3	16.3	24.9	34.1	44.2	51.2
2002.....	1.5	3.0	5.5	10.3	16.8	26.3	35.5	46.8	50.6
2001.....	1.6	2.8	6.1	10.2	16.8	25.3	35.6	43.4	48.4
2000.....	1.4	3.0	5.0	9.5	17.0	24.5	36.0	46.5	52.5
1999.....	1.4	3.0	6.5	11.1	16.7	24.6	34.6	46.0	52.2

Non-Hispanic 4/, female

2003.....	0.9	1.9	3.5	6.0	10.7	17.2	24.8	29.4	30.3
2002.....	1.1	2.0	4.0	6.7	11.6	17.5	22.5	29.5	30.8
2001.....	0.9	1.9	4.3	6.9	11.8	17.9	23.5	29.7	31.5
2000.....	1.0	2.3	4.2	7.6	12.3	17.4	23.1	31.1	32.1
1999.....	1.1	2.1	4.7	7.3	12.1	17.9	23.2	30.7	30.6

Non-Hispanic White, both sexes

2003.....	0.9	2.0	3.9	7.4	12.2	19.2	27.0	33.5	35.3
2002.....	1.0	2.0	4.1	7.6	12.8	20.2	26.8	35.1	35.7
2001.....	1.0	2.0	4.4	7.4	12.9	19.9	27.2	33.1	35.1
2000.....	1.0	2.3	3.9	7.7	13.3	19.2	26.5	35.6	35.2
1999.....	1.1	2.1	4.6	7.8	13.2	19.4	26.0	35.1	35.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)									
Non-Hispanic White, male									
2003.....	1.1	2.5	4.8	9.5	15.3	23.9	33.0	42.8	50.5
2002.....	1.3	2.7	4.9	9.5	15.6	25.3	34.6	46.0	50.1
2001.....	1.4	2.5	5.2	9.1	15.5	23.9	34.6	41.6	47.3
2000.....	1.2	2.6	4.3	8.8	15.8	22.9	34.1	45.7	50.9
1999.....	1.3	2.6	5.5	10.0	15.8	22.9	32.9	44.7	50.6
Non-Hispanic White, female									
2003.....	0.7	1.6	3.0	5.3	9.5	15.4	22.6	27.8	28.7
2002.....	0.8	1.4	3.4	5.8	10.3	16.0	21.2	28.4	29.5
2001.....	0.7	1.6	3.7	5.8	10.6	16.6	22.0	28.0	30.0
2000.....	0.8	2.0	3.5	6.7	11.1	16.2	21.2	29.7	28.9
1999.....	0.9	1.5	3.8	5.8	10.9	16.6	21.1	29.4	28.7
Non-Hispanic Black, both sexes									
2003.....	2.9	6.1	9.6	15.6	26.1	38.1	52.7	60.4	60.4
2002.....	3.0	5.7	10.9	16.6	26.9	38.0	45.5	55.0	55.7
2001.....	2.7	4.8	12.1	19.3	26.2	38.8	47.7	61.8	59.3
2000.....	3.0	6.5	11.9	16.9	27.9	38.0	53.8	58.3	72.8
1999.....	2.5	6.8	13.9	20.5	24.6	38.1	51.6	56.9	63.4
Non-Hispanic Black, male									
2003.....	3.3	7.8	12.4	20.2	30.1	40.4	57.5	74.7	74.6
2002.....	3.7	6.1	12.6	20.4	32.0	45.2	53.5	71.0	71.1
2001.....	3.5	5.4	15.0	23.0	31.0	45.4	56.9	81.8	72.8
2000.....	3.1	7.4	12.7	18.7	31.7	48.5	67.4	70.4	81.0
1999.....	2.6	6.7	16.4	22.2	26.6	47.4	59.5	75.3	80.8
Non-Hispanic Black, female									
2003.....	2.5	4.6	7.3	11.9	23.1	36.6	49.9	52.9	54.9
2002.....	2.5	5.4	9.4	13.7	23.2	33.3	40.6	46.7	49.7
2001.....	2.0	4.4	9.8	16.4	22.7	34.4	42.2	51.6	54.3
2000.....	2.9	5.7	11.2	15.4	25.2	31.1	45.6	52.3	69.8
1999.....	2.3	6.8	11.8	19.1	23.2	31.9	46.9	47.9	56.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)									
All origins 3/, both sexes									
2003.....	10.3	17.7	28.1	44.4	64.6	92.9	131.5	167.7	224.0
2002.....	10.4	17.5	28.8	46.5	66.0	97.0	131.0	168.7	230.1
2001.....	10.5	18.0	30.1	46.4	67.2	98.1	133.4	170.9	234.5
2000.....	10.5	18.2	30.4	48.2	71.5	100.4	136.0	177.4	239.8
1999.....	11.4	19.1	32.5	52.5	76.1	107.8	142.3	188.5	250.4
All origins 3/, male									
2003.....	11.9	20.1	33.0	52.9	78.3	111.3	162.7	208.2	275.0
2002.....	11.7	20.8	33.0	56.5	79.9	118.9	161.8	207.8	290.8
2001.....	12.0	21.5	34.9	55.4	79.4	120.3	162.3	211.0	295.2
2000.....	12.0	20.6	35.1	57.2	86.2	122.5	167.5	221.4	300.7
1999.....	13.2	22.0	38.2	62.6	90.6	132.8	174.0	233.0	318.9
All origins 3/, female									
2003.....	8.7	15.5	23.6	36.7	52.8	78.1	109.2	142.9	201.4
2002.....	9.1	14.3	24.9	37.5	53.9	79.5	109.2	144.9	203.8
2001.....	9.1	14.7	25.7	38.3	56.7	80.5	113.1	146.8	209.1
2000.....	9.1	15.8	25.9	40.0	59.0	83.1	114.1	151.4	215.0
1999.....	9.7	16.3	27.1	43.3	63.8	88.2	120.3	162.5	222.8
Hispanic, both sexes									
2003.....	7.1	13.3	18.6	30.7	45.4	75.5	97.3	129.1	168.9
2002.....	6.9	12.8	20.4	32.3	48.4	75.2	102.6	129.7	164.1
2001.....	7.4	12.9	19.0	35.4	46.8	78.2	98.3	133.1	193.8
2000.....	6.6	12.6	23.5	30.5	52.9	78.2	109.3	152.6	192.4
1999.....	8.6	13.2	23.7	35.2	56.2	75.6	108.5	147.4	202.1
Hispanic, male									
2003.....	9.0	14.2	21.2	36.9	54.9	92.2	117.9	143.5	194.0
2002.....	7.3	14.5	25.1	35.9	55.5	86.9	127.3	159.0	204.1
2001.....	8.1	14.3	23.0	39.9	50.0	90.6	130.3	161.0	213.3
2000.....	6.8	13.2	24.8	35.1	60.8	90.4	138.0	179.6	231.8
1999.....	9.0	13.9	26.6	42.0	62.2	96.7	124.0	187.1	235.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)									
Hispanic, female									
2003.....	5.2	12.6	16.2	25.3	37.6	62.6	82.8	119.8	156.3
2002.....	6.5	11.2	16.0	29.1	42.6	66.2	85.0	111.0	144.2
2001.....	6.8	11.7	15.4	31.5	44.2	68.8	75.5	115.8	184.3
2000.....	6.5	12.1	22.4	26.6	46.5	68.9	88.9	136.2	173.1
1999.....	8.2	12.6	21.1	29.3	51.3	59.8	97.5	123.8	186.1
Non-Hispanic 4/, both sexes									
2003.....	10.6	18.1	28.8	45.3	65.9	93.8	133.2	169.3	225.8
2002.....	10.7	17.9	29.4	47.5	67.0	98.2	132.1	170.1	232.1
2001.....	10.8	18.4	30.9	47.2	68.5	98.8	134.8	172.0	235.3
2000.....	10.9	18.5	30.8	49.3	72.6	101.4	136.7	177.9	241.0
1999.....	11.7	19.5	33.0	53.6	77.3	109.2	143.5	189.5	251.6
Non-Hispanic 4/, male									
2003.....	12.1	20.5	33.8	53.9	79.8	112.3	165.0	211.0	278.2
2002.....	12.1	21.3	33.6	57.8	81.3	120.7	163.2	209.4	294.0
2001.....	12.4	22.0	35.7	56.3	81.2	121.3	163.7	212.5	297.9
2000.....	12.5	21.2	35.8	58.6	87.5	123.8	168.2	222.5	303.0
1999.....	13.5	22.7	39.0	63.9	92.2	134.2	175.7	233.9	321.9
Non-Hispanic 4/, female									
2003.....	9.1	15.7	24.1	37.5	53.9	79.0	110.5	143.8	202.7
2002.....	9.4	14.6	25.5	38.1	54.6	80.2	110.2	146.2	205.4
2001.....	9.3	15.0	26.4	38.8	57.5	81.0	114.6	147.7	209.3
2000.....	9.3	16.0	26.1	40.9	59.7	83.6	114.9	151.7	215.9
1999.....	9.9	16.5	27.5	44.2	64.5	89.6	121.1	163.5	223.5
Non-Hispanic White, both sexes									
2003.....	10.0	17.3	28.0	44.9	66.0	94.4	134.5	171.7	229.9
2002.....	10.1	17.1	28.7	46.6	66.5	98.0	132.8	171.4	235.2
2001.....	10.1	17.5	29.7	45.9	67.7	98.1	135.5	172.5	238.9
2000.....	10.2	17.5	30.1	48.3	72.4	100.8	136.2	179.9	244.9
1999.....	11.0	18.6	31.6	52.3	77.5	108.5	143.3	190.3	253.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic White, male

2003.....	11.2	19.6	32.5	53.2	79.4	112.5	166.5	215.2	284.6
2002.....	11.4	20.4	32.5	56.5	81.6	120.4	163.5	212.7	300.2
2001.....	11.6	20.5	33.8	54.3	79.8	120.8	164.4	213.7	302.9
2000.....	11.5	20.0	34.7	56.1	87.6	123.1	166.3	226.2	309.5
1999.....	12.4	21.3	37.2	62.1	91.7	133.2	176.2	236.1	326.9

Non-Hispanic White, female

2003.....	8.7	15.0	23.6	37.2	54.0	79.6	111.3	144.9	205.8
2002.....	8.8	13.9	25.0	37.4	53.2	79.7	110.7	146.2	207.0
2001.....	8.6	14.4	25.8	38.1	57.1	79.6	115.1	147.6	212.3
2000.....	8.8	14.9	25.7	41.0	59.1	82.9	115.1	152.4	218.9
1999.....	9.5	16.0	26.2	43.3	65.0	88.8	120.2	163.2	224.3

Non-Hispanic Black, both sexes

2003.....	16.3	26.9	41.9	58.6	78.5	104.5	141.2	167.7	206.4
2002.....	16.1	25.8	42.4	64.0	83.8	115.8	149.2	175.4	223.2
2001.....	17.2	28.4	45.2	65.3	86.5	120.7	143.0	182.4	218.5
2000.....	17.1	29.5	42.1	65.9	86.8	118.5	156.7	174.4	211.1
1999.....	18.7	29.8	49.2	74.4	86.6	129.5	159.2	194.6	244.8

Non-Hispanic Black, male

2003.....	20.2	31.6	54.4	72.8	100.2	131.4	179.0	207.5	239.7
2002.....	18.7	31.6	51.4	82.3	99.1	145.7	190.1	210.6	265.8
2001.....	19.4	36.6	58.6	82.6	111.8	150.5	181.6	218.3	287.6
2000.....	21.4	33.9	52.1	90.4	103.1	150.3	211.9	212.0	253.6
1999.....	23.8	37.4	62.9	95.8	111.0	165.1	193.8	235.6	293.6

Non-Hispanic Black, female

2003.....	12.9	22.9	31.6	47.5	62.3	86.3	118.7	146.9	193.6
2002.....	13.9	20.9	35.0	49.6	72.5	95.7	124.7	157.1	206.9
2001.....	15.2	21.5	34.2	51.8	67.9	100.9	120.0	164.1	192.9
2000.....	13.5	25.7	33.8	46.6	74.7	97.3	123.4	155.5	195.4
1999.....	14.4	23.3	38.1	57.7	68.5	105.9	138.3	174.3	226.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All origins 3/, both sexes

2003.....	1.4	2.0	3.2	6.0	9.8	16.3	28.5	47.4	80.5
2002.....	1.5	2.1	3.7	5.4	9.7	16.3	28.8	46.3	78.7
2001.....	1.5	2.3	3.9	6.3	10.5	17.9	29.5	47.8	82.7
2000.....	1.5	2.3	3.8	6.6	10.6	17.4	28.9	50.6	84.9
1999.....	1.4	2.3	3.5	6.3	10.8	18.6	29.3	47.5	82.9

All origins 3/, male

2003.....	1.4	2.3	3.6	7.0	11.6	21.2	35.9	61.5	101.7
2002.....	1.6	2.1	4.5	6.3	12.2	20.0	37.3	59.1	101.1
2001.....	1.7	2.4	4.1	7.5	12.8	22.5	39.2	62.4	107.6
2000.....	1.6	2.6	4.1	7.5	13.3	21.2	36.8	65.3	108.9
1999.....	1.5	2.7	4.0	7.9	13.4	23.9	37.2	61.0	109.2

All origins 3/, female

2003.....	1.3	1.7	2.8	5.1	8.3	12.4	23.2	38.7	71.1
2002.....	1.5	2.0	3.0	4.6	7.5	13.4	22.8	38.5	68.9
2001.....	1.2	2.1	3.7	5.2	8.5	14.3	22.7	39.1	72.3
2000.....	1.3	1.9	3.5	5.7	8.2	14.4	23.4	41.9	75.2
1999.....	1.3	1.9	3.0	4.8	8.5	14.5	23.8	39.7	72.3

Hispanic, both sexes

2003.....	1.1	*	2.2	4.2	7.2	10.6	22.2	39.8	48.2
2002.....	1.4	1.5	1.8	4.8	6.2	12.2	17.3	37.0	48.8
2001.....	1.1	1.9	2.4	5.3	7.5	10.5	19.0	30.3	57.5
2000.....	1.1	1.7	2.4	4.0	6.7	12.2	20.5	31.2	70.3
1999.....	*	1.9	3.4	4.0	6.7	15.3	18.4	31.7	51.9

Hispanic, male

2003.....	*	*	*	5.4	8.1	12.4	26.1	46.8	52.3
2002.....	*	*	*	5.4	7.8	13.6	19.8	52.6	56.9
2001.....	*	*	*	6.8	7.4	13.3	19.8	49.3	70.5
2000.....	*	*	*	*	*	14.6	26.6	*	86.7
1999.....	*	*	*	*	*	17.3	18.7	36.2	49.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Hispanic, female

2003.....	*	*	*	*	6.4	9.2	19.4	35.2	46.1
2002.....	*	*	*	*	*	11.2	15.5	27.0	44.8
2001.....	*	*	*	*	7.5	8.4	18.4	18.5	51.2
2000.....	*	*	*	*	7.2	10.3	16.2	34.0	62.3
1999.....	*	*	*	*	8.1	13.8	18.2	29.1	53.0

Non-Hispanic 4/, both sexes

2003.....	1.4	2.1	3.3	6.2	10.0	16.6	28.8	47.7	81.8
2002.....	1.5	2.1	3.9	5.5	9.9	16.6	29.3	46.6	79.8
2001.....	1.5	2.3	4.0	6.3	10.7	18.3	30.0	48.3	83.5
2000.....	1.5	2.3	3.9	6.8	10.8	17.6	29.2	51.2	85.2
1999.....	1.5	2.3	3.5	6.4	11.0	18.8	29.7	48.0	83.8

Non-Hispanic 4/, male

2003.....	1.4	2.4	3.7	7.1	11.8	21.6	36.4	62.2	103.9
2002.....	1.6	2.2	4.7	6.4	12.5	20.4	38.0	59.2	102.9
2001.....	1.7	2.4	4.2	7.5	13.1	23.0	40.1	62.7	109.0
2000.....	1.7	2.7	4.2	7.8	13.8	21.6	37.2	66.6	109.4
1999.....	1.5	2.6	4.1	8.1	13.9	24.3	37.9	61.8	111.4

Non-Hispanic 4/, female

2003.....	1.3	1.8	2.9	5.2	8.4	12.6	23.4	38.9	72.1
2002.....	1.5	2.0	3.1	4.7	7.7	13.5	23.2	38.9	69.8
2001.....	1.3	2.1	3.8	5.3	8.6	14.6	22.9	39.7	72.9
2000.....	1.3	1.9	3.6	5.8	8.2	14.6	23.7	42.1	75.4
1999.....	1.4	2.0	3.0	4.9	8.5	14.5	24.0	39.9	72.8

Non-Hispanic White, both sexes

2003.....	1.3	2.1	3.2	6.3	10.0	17.1	29.9	49.1	84.4
2002.....	1.6	2.0	3.8	5.6	10.1	17.0	30.2	47.4	81.8
2001.....	1.5	2.3	4.0	6.3	11.0	18.6	30.9	49.4	84.8
2000.....	1.4	2.3	4.0	6.8	11.0	17.6	29.8	52.8	87.6
1999.....	1.3	2.2	3.4	6.4	10.9	19.2	30.1	49.2	86.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic White, male

2003.....	1.4	2.4	3.5	7.2	12.0	22.3	37.9	64.0	107.9
2002.....	1.7	2.2	4.5	6.5	12.8	21.1	39.2	60.2	106.0
2001.....	1.8	2.5	4.3	7.4	13.5	23.6	41.3	63.9	111.4
2000.....	1.6	2.7	4.4	7.5	13.9	21.3	38.0	69.4	113.1
1999.....	1.5	2.6	3.9	8.2	13.9	24.9	38.6	63.8	116.8

Non-Hispanic White, female

2003.....	1.2	1.8	2.9	5.4	8.1	12.9	24.2	39.9	74.0
2002.....	1.4	1.8	3.1	4.8	7.8	13.6	23.8	39.6	71.4
2001.....	1.3	2.1	3.8	5.3	8.7	14.5	23.5	40.6	73.7
2000.....	1.2	1.8	3.6	6.1	8.5	14.6	24.2	43.0	77.3
1999.....	1.2	1.9	3.0	4.7	8.3	14.7	24.2	40.6	74.1

Non-Hispanic Black, both sexes

2003.....	2.3	2.8	4.3	6.7	11.5	15.6	22.5	40.2	58.9
2002.....	2.0	3.1	5.3	6.0	10.8	15.4	24.7	42.4	64.4
2001.....	1.5	2.8	4.6	7.7	11.0	18.9	25.0	42.8	72.7
2000.....	2.4	3.2	4.2	7.3	10.9	20.3	27.0	38.2	64.2
1999.....	2.3	3.1	5.0	8.0	14.0	18.2	29.4	39.0	62.7

Non-Hispanic Black, male

2003.....	2.1	3.0	5.6	8.5	12.7	21.6	26.6	52.8	68.2
2002.....	*	2.9	7.7	6.7	12.8	16.4	30.0	54.6	81.0
2001.....	*	2.5	4.5	9.4	13.3	23.4	32.2	59.7	87.8
2000.....	2.2	3.7	4.2	10.4	16.0	23.9	34.7	39.5	76.3
1999.....	*	3.2	6.3	8.5	17.3	21.8	36.3	43.4	64.2

Non-Hispanic Black, female

2003.....	2.4	2.6	3.2	5.3	10.7	11.6	20.0	33.6	55.3
2002.....	2.3	3.3	3.4	5.4	9.3	14.7	21.6	36.1	58.1
2001.....	*	3.0	4.6	6.4	9.2	15.8	20.7	34.1	67.2
2000.....	2.5	2.8	4.2	4.9	7.1	17.9	22.4	37.6	59.7
1999.....	2.6	3.1	4.0	7.7	11.6	15.9	25.3	36.9	62.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Anemias (D50-D64)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	28.1
2002.....	*	*	*	*	*	*	*	*	24.9
2001.....	*	*	*	*	*	*	*	*	32.6
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	32.2

Non-Hispanic 4/, both sexes

2003.....	0.7	0.6	0.7	1.3	1.9	3.5	6.1	12.7	41.2
2002.....	0.6	0.7	0.9	1.2	1.9	3.7	6.5	12.9	40.5
2001.....	0.6	0.7	0.8	1.2	1.9	3.6	7.1	13.7	42.5
2000.....	0.6	0.7	1.0	1.2	2.2	3.4	6.6	13.9	43.2
1999.....	0.7	0.6	0.8	1.2	1.8	3.4	6.8	13.7	44.0

Non-Hispanic 4/, male

2003.....	0.7	0.5	0.8	1.6	2.3	3.9	6.6	13.4	38.3
2002.....	0.6	0.6	0.8	1.3	1.7	4.3	6.7	12.5	37.6
2001.....	0.6	0.7	0.8	1.1	2.0	3.9	7.1	14.4	40.8
2000.....	0.6	0.7	1.1	1.3	2.2	3.5	6.6	14.4	41.0
1999.....	0.7	0.6	0.7	1.2	2.1	3.2	7.2	14.7	43.3

Non-Hispanic 4/, female

2003.....	0.6	0.6	0.7	1.1	1.5	3.2	5.7	12.3	42.4
2002.....	0.7	0.7	0.9	1.1	2.1	3.2	6.3	13.2	41.8
2001.....	0.5	0.7	0.9	1.3	1.9	3.3	7.0	13.3	43.1
2000.....	0.6	0.7	0.9	1.2	2.3	3.4	6.6	13.6	44.1
1999.....	0.7	0.6	0.9	1.2	1.6	3.6	6.5	13.2	44.2

Non-Hispanic White, both sexes

2003.....	0.3	0.3	0.4	1.0	1.6	3.2	5.8	12.5	41.0
2002.....	0.3	0.3	0.6	0.9	1.7	3.3	5.9	12.6	39.9
2001.....	0.2	0.4	0.5	1.1	1.7	3.3	6.7	13.2	42.1
2000.....	0.2	0.4	0.7	0.9	2.0	3.2	6.3	13.8	42.9
1999.....	0.3	0.3	0.4	0.8	1.6	3.1	6.5	13.7	43.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)									
Non-Hispanic White, male									
2003.....	0.3	0.3	0.4	1.1	2.0	3.3	6.4	13.5	38.1
2002.....	0.3	0.4	0.6	1.0	1.5	3.6	6.3	11.9	36.3
2001.....	*	0.4	0.5	1.0	1.8	3.4	6.9	14.2	41.2
2000.....	*	0.4	0.8	0.9	2.1	3.3	6.3	14.1	39.9
1999.....	0.4	0.3	*	0.7	1.9	2.8	7.0	14.7	43.4
Non-Hispanic White, female									
2003.....	*	*	0.3	0.8	1.2	3.1	5.4	11.8	42.2
2002.....	0.3	0.3	0.6	0.8	1.8	3.0	5.6	13.0	41.4
2001.....	*	0.4	0.5	1.1	1.7	3.1	6.5	12.7	42.4
2000.....	*	0.4	0.6	0.9	1.8	3.1	6.3	13.5	44.1
1999.....	0.3	*	0.5	1.0	1.4	3.4	6.1	13.1	43.1
Non-Hispanic Black, both sexes									
2003.....	3.4	2.7	4.2	4.5	4.3	7.3	11.0	19.7	50.3
2002.....	3.2	3.1	3.2	4.2	4.5	8.1	14.0	19.5	52.9
2001.....	3.3	2.9	3.3	2.7	4.0	7.4	12.6	21.5	49.3
2000.....	3.0	3.2	3.0	4.0	4.9	6.3	10.5	18.1	51.8
1999.....	3.3	3.0	4.3	4.3	3.7	6.6	11.4	16.4	53.0
Non-Hispanic Black, male									
2003.....	4.1	2.6	5.1	6.4	5.6	11.1	13.1	18.9	50.1
2002.....	2.7	2.6	3.1	5.3	*	12.0	13.1	23.4	58.0
2001.....	4.0	3.2	3.4	*	*	10.7	12.3	20.4	42.7
2000.....	3.3	3.5	*	*	*	*	11.1	18.9	56.4
1999.....	3.4	3.3	4.4	6.1	*	8.0	10.6	*	44.0
Non-Hispanic Black, female									
2003.....	2.8	2.8	3.4	3.1	*	4.8	9.7	20.1	50.3
2002.....	3.6	3.5	3.4	3.3	4.7	5.5	14.6	17.5	51.0
2001.....	2.7	2.7	3.3	*	4.0	5.3	12.7	22.1	51.8
2000.....	2.8	3.1	3.0	4.0	5.9	6.4	10.2	17.7	50.2
1999.....	3.2	2.8	4.3	*	*	5.7	11.9	16.2	56.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diabetes mellitus (E10-E14)

All origins 3/, both sexes

2003.....	10.3	17.9	30.4	48.9	75.0	108.8	155.3	216.5	317.5
2002.....	9.9	18.0	30.1	47.5	75.7	108.8	157.7	217.9	320.6
2001.....	9.9	17.8	29.5	48.4	75.5	108.6	155.5	218.7	321.8
2000.....	9.5	17.3	29.6	48.1	74.0	108.6	156.5	213.9	319.7
1999.....	9.9	16.5	29.5	48.9	75.6	109.2	153.5	214.8	317.2

All origins 3/, male

2003.....	12.6	21.3	35.4	57.8	86.8	124.4	179.0	249.7	340.7
2002.....	11.9	21.5	36.0	55.5	84.3	121.1	175.9	243.5	348.5
2001.....	12.1	21.3	34.4	54.4	84.7	121.0	171.5	242.3	343.1
2000.....	11.4	19.7	33.4	53.9	80.8	119.7	173.7	242.5	345.1
1999.....	12.3	19.4	33.5	53.9	82.5	119.4	172.3	241.3	345.7

All origins 3/, female

2003.....	8.1	14.7	25.7	40.9	64.8	96.2	138.4	196.2	307.2
2002.....	7.9	14.7	24.5	40.2	68.2	99.0	144.9	202.3	308.4
2001.....	7.9	14.5	24.9	42.9	67.6	98.8	144.3	204.5	312.9
2000.....	7.8	14.9	26.1	42.8	68.1	99.9	144.6	197.0	309.4
1999.....	7.5	13.8	25.8	44.5	69.7	101.2	140.4	199.3	305.8

Hispanic, both sexes

2003.....	10.1	22.7	40.5	72.1	105.3	165.4	238.3	327.1	428.1
2002.....	9.2	20.4	38.0	71.3	108.2	168.0	257.0	333.3	437.9
2001.....	11.9	18.8	38.4	72.8	121.1	166.3	254.5	322.3	489.6
2000.....	9.8	19.5	45.4	68.5	114.1	190.3	250.1	330.3	465.1
1999.....	10.1	19.0	40.4	77.1	124.6	181.4	252.4	344.8	493.9

Hispanic, male

2003.....	11.6	26.8	45.2	87.1	116.7	185.3	254.6	370.1	426.4
2002.....	10.2	25.4	43.3	80.8	110.7	186.5	282.8	350.0	425.0
2001.....	14.4	23.8	43.4	79.6	132.5	175.7	262.6	328.5	422.8
2000.....	11.1	21.4	52.8	80.3	126.0	206.1	267.2	363.7	449.4
1999.....	12.5	23.5	46.7	82.8	127.9	192.5	262.1	355.2	468.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)									
Hispanic, female									
2003.....	8.6	18.7	36.1	59.0	96.0	150.1	226.8	299.1	428.9
2002.....	8.2	15.4	33.1	63.1	106.2	153.8	238.6	322.7	444.3
2001.....	9.4	14.0	33.8	67.0	111.9	159.1	248.9	318.5	522.2
2000.....	8.6	17.7	38.7	58.3	104.5	178.2	237.9	310.0	472.8
1999.....	7.6	14.7	34.6	72.1	121.9	173.1	245.6	338.5	506.4
Non-Hispanic 4/, both sexes									
2003.....	10.3	17.4	29.5	47.0	72.5	104.8	150.2	210.7	312.0
2002.....	9.9	17.8	29.3	45.4	73.0	104.7	151.8	212.2	315.0
2001.....	9.7	17.6	28.7	46.4	71.9	104.7	150.1	213.9	314.8
2000.....	9.5	17.0	28.3	46.4	71.1	103.6	151.7	209.0	313.4
1999.....	9.8	16.2	28.6	46.7	72.3	105.0	148.7	209.5	310.0
Non-Hispanic 4/, male									
2003.....	12.7	20.8	34.5	55.4	84.4	120.2	174.2	243.1	335.9
2002.....	12.0	21.1	35.4	53.3	82.0	116.7	169.5	237.7	344.0
2001.....	11.7	21.0	33.6	52.4	81.0	117.1	166.4	238.1	338.9
2000.....	11.4	19.5	31.8	51.9	77.6	114.4	168.6	237.2	339.7
1999.....	12.3	18.9	32.5	51.6	79.4	115.2	167.9	236.6	339.5
Non-Hispanic 4/, female									
2003.....	8.0	14.2	24.8	39.4	62.2	92.4	133.1	190.9	301.5
2002.....	7.9	14.6	23.7	38.2	65.1	95.2	139.4	196.7	302.5
2001.....	7.7	14.4	24.1	41.0	64.1	94.7	138.7	199.4	304.7
2000.....	7.7	14.6	25.1	41.4	65.5	95.1	139.9	192.4	302.8
1999.....	7.5	13.7	25.0	42.2	66.1	97.0	135.3	193.6	298.2
Non-Hispanic White, both sexes									
2003.....	8.6	14.2	24.8	39.8	62.6	92.5	136.6	195.4	293.8
2002.....	8.2	14.7	24.7	38.3	61.9	92.7	138.0	197.8	296.4
2001.....	8.1	14.7	23.9	39.4	62.0	92.7	136.3	199.9	297.4
2000.....	7.8	14.0	23.7	39.7	61.2	91.7	137.5	193.0	295.5
1999.....	8.1	13.2	23.0	38.7	62.2	92.0	135.2	194.3	290.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diabetes mellitus (E10-E14)

Non-Hispanic White, male

2003.....	11.0	17.7	30.1	48.1	75.3	109.7	164.0	232.9	327.3
2002.....	10.2	17.9	30.7	46.4	72.5	107.2	160.1	228.4	338.4
2001.....	9.9	17.9	29.0	45.9	72.8	108.6	155.6	226.9	332.1
2000.....	9.5	16.7	27.3	45.7	69.6	105.1	157.9	227.9	330.4
1999.....	10.3	16.0	27.2	44.5	70.9	106.0	158.4	227.5	326.0

Non-Hispanic White, female

2003.....	6.2	10.9	19.8	32.1	51.4	78.3	116.7	172.3	279.0
2002.....	6.3	11.6	18.8	30.7	52.5	80.8	122.1	179.0	278.3
2001.....	6.3	11.6	18.9	33.3	52.5	79.7	122.6	183.7	283.0
2000.....	6.1	11.3	20.4	34.2	53.7	80.9	123.2	172.2	281.4
1999.....	5.9	10.5	18.9	33.3	54.5	80.8	119.0	174.7	275.9

Non-Hispanic Black, both sexes

2003.....	22.7	40.7	71.3	109.1	159.9	225.4	303.4	394.6	532.2
2002.....	22.0	41.8	69.3	107.3	164.9	225.1	303.3	396.4	544.7
2001.....	21.3	41.0	70.6	106.6	159.4	229.9	307.4	385.5	533.8
2000.....	22.4	41.7	67.7	103.7	160.7	226.7	320.5	409.2	530.1
1999.....	23.1	41.1	74.2	113.7	162.6	232.3	297.1	399.2	530.9

Non-Hispanic Black, male

2003.....	26.2	45.0	78.9	123.9	174.9	236.2	311.6	393.2	457.1
2002.....	26.1	47.7	78.6	120.6	168.4	226.5	297.7	390.8	423.4
2001.....	25.8	46.4	77.8	114.5	163.9	223.4	310.5	404.7	452.7
2000.....	26.1	44.4	73.5	110.1	155.4	227.2	314.1	384.5	456.8
1999.....	28.2	44.5	79.6	115.2	167.3	218.8	284.2	379.0	506.3

Non-Hispanic Black, female

2003.....	19.7	37.2	65.0	97.6	148.9	218.1	298.5	395.3	561.3
2002.....	18.5	36.7	61.8	96.9	162.2	224.1	306.7	399.3	591.1
2001.....	17.4	36.4	64.8	100.4	156.0	234.2	305.5	375.6	563.8
2000.....	19.3	39.4	63.0	98.6	164.6	226.4	324.4	421.7	557.3
1999.....	18.7	38.2	69.8	112.5	159.1	241.2	304.9	409.2	540.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nutritional deficiencies (E40-E64)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	28.9
2002.....	*	*	*	*	*	*	*	*	29.0
2001.....	*	*	*	*	*	*	*	*	36.3
2000.....	*	*	*	*	*	*	*	*	41.5
1999.....	*	*	*	*	*	*	*	*	51.0

Non-Hispanic 4/, both sexes

2003.....	0.2	0.3	0.5	0.8	1.3	2.5	5.8	10.5	35.7
2002.....	0.2	0.3	0.5	0.8	1.6	2.6	6.0	12.9	41.8
2001.....	0.2	0.4	0.5	0.8	1.4	2.9	5.9	12.9	43.0
2000.....	0.2	0.3	0.5	1.0	2.0	3.1	6.8	15.8	50.3
1999.....	0.2	0.4	0.6	0.9	1.6	3.2	6.5	16.1	53.9

Non-Hispanic 4/, male

2003.....	*	0.3	0.5	0.7	1.4	2.6	5.8	11.0	30.7
2002.....	*	0.3	0.6	1.0	1.6	3.0	6.2	12.0	37.1
2001.....	0.3	0.3	0.5	0.9	1.5	3.0	5.5	13.2	34.8
2000.....	0.3	*	0.7	1.1	2.0	3.4	6.7	14.3	43.9
1999.....	0.2	0.3	0.7	1.2	1.7	3.8	6.7	16.4	48.1

Non-Hispanic 4/, female

2003.....	*	0.4	0.5	0.8	1.3	2.3	5.8	10.1	37.9
2002.....	*	0.4	0.5	0.7	1.6	2.4	5.8	13.4	43.9
2001.....	0.2	0.4	0.4	0.8	1.3	2.7	6.2	12.7	46.4
2000.....	0.2	0.4	0.4	0.9	2.0	2.8	6.9	16.6	52.8
1999.....	0.3	0.4	0.5	0.7	1.5	2.6	6.3	16.0	56.2

Non-Hispanic White, both sexes

2003.....	0.2	0.3	0.5	0.7	1.3	2.3	5.6	9.9	34.4
2002.....	0.2	0.3	0.5	0.7	1.4	2.4	5.7	12.3	40.6
2001.....	0.2	0.3	0.4	0.8	1.2	2.7	5.4	12.0	42.2
2000.....	0.2	0.2	0.5	0.9	1.9	2.7	6.4	15.3	49.0
1999.....	0.2	0.3	0.4	0.8	1.4	2.6	6.0	15.1	52.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nutritional deficiencies (E40-E64)

Non-Hispanic White, male

2003.....	*	*	0.4	0.6	1.4	2.2	5.3	10.1	29.6
2002.....	*	*	0.5	0.7	1.4	2.7	6.0	11.0	36.5
2001.....	*	*	0.4	0.8	1.2	2.7	5.1	12.3	33.3
2000.....	*	*	0.5	0.9	1.8	2.8	6.1	13.8	43.0
1999.....	*	*	*	1.0	1.4	3.0	5.9	15.0	46.2

Non-Hispanic White, female

2003.....	*	*	0.5	0.8	1.1	2.3	5.9	9.8	36.5
2002.....	*	0.4	0.5	0.7	1.4	2.2	5.5	13.2	42.3
2001.....	*	0.3	0.4	0.8	1.3	2.7	5.7	11.8	45.9
2000.....	*	0.3	0.4	0.9	1.9	2.6	6.5	16.1	51.5
1999.....	*	0.4	0.4	0.6	1.5	2.3	6.0	15.1	55.4

Non-Hispanic Black, both sexes

2003.....	*	1.0	*	1.7	*	4.7	9.2	19.2	55.9
2002.....	*	*	1.6	2.1	3.6	5.1	10.7	20.8	60.2
2001.....	*	1.1	*	*	3.1	4.5	11.0	24.0	55.2
2000.....	*	*	*	2.2	3.3	6.7	12.1	23.0	67.4
1999.....	*	*	2.0	2.0	3.3	8.4	12.5	31.1	70.8

Non-Hispanic Black, male

2003.....	*	*	*	*	*	7.4	14.0	24.1	49.0
2002.....	*	*	*	*	*	9.9	27.3	45.9	
2001.....	*	*	*	*	*	6.9	11.9	24.5	58.9
2000.....	*	*	*	*	*	9.9	14.9	23.2	64.6
1999.....	*	*	*	*	*	13.2	16.4	35.4	76.1

Non-Hispanic Black, female

2003.....	*	*	*	*	*	*	6.4	16.6	58.6
2002.....	*	*	*	*	*	4.6	11.2	17.5	65.6
2001.....	*	*	*	*	*	*	10.5	23.7	53.9
2000.....	*	*	*	*	*	4.5	10.5	22.9	68.5
1999.....	*	*	*	*	*	5.3	10.2	29.0	68.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)									
Hispanic, female									
2003.....	*	*	*	*	*	*	*	*	26.6
2002.....	*	*	*	*	*	*	*	*	28.2
2001.....	*	*	*	*	*	*	*	*	36.3
2000.....	*	*	*	*	*	*	*	*	41.5
1999.....	*	*	*	*	*	*	*	*	48.9
Non-Hispanic 4/, both sexes									
2003.....	0.2	0.3	0.5	0.7	1.3	2.3	5.6	10.1	33.7
2002.....	0.2	0.3	0.5	0.7	1.4	2.5	5.5	12.2	38.7
2001.....	0.2	0.3	0.5	0.8	1.3	2.6	5.4	12.1	40.2
2000.....	0.2	0.3	0.5	0.9	1.9	2.8	6.4	14.6	46.5
1999.....	0.2	0.3	0.5	0.9	1.5	3.0	6.0	14.9	50.0
Non-Hispanic 4/, male									
2003.....	*	0.3	0.5	0.7	1.3	2.5	5.5	10.5	28.4
2002.....	*	0.3	0.6	0.9	1.4	2.8	5.7	11.4	35.3
2001.....	0.2	0.3	0.5	0.8	1.4	2.7	4.9	12.7	33.2
2000.....	0.2	*	0.6	1.0	1.8	3.2	6.3	13.0	40.9
1999.....	0.2	0.3	0.6	1.1	1.6	3.6	6.1	15.1	44.3
Non-Hispanic 4/, female									
2003.....	*	0.3	0.5	0.8	1.2	2.2	5.8	9.8	36.0
2002.....	*	0.3	0.5	0.6	1.5	2.2	5.4	12.7	40.1
2001.....	*	0.4	0.4	0.8	1.3	2.6	5.7	11.7	43.2
2000.....	0.2	0.3	0.3	0.9	1.9	2.5	6.6	15.5	48.7
1999.....	0.2	0.4	0.4	0.6	1.5	2.5	6.0	14.9	52.2
Non-Hispanic White, both sexes									
2003.....	0.1	0.2	0.4	0.6	1.2	2.1	5.4	9.5	32.3
2002.....	0.2	0.3	0.4	0.6	1.3	2.3	5.2	11.7	37.4
2001.....	0.2	0.2	0.4	0.8	1.2	2.5	5.0	11.2	39.4
2000.....	0.2	0.2	0.4	0.8	1.7	2.4	6.1	14.1	45.3
1999.....	0.2	0.3	0.3	0.8	1.3	2.5	5.6	13.9	48.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malnutrition (E40-E46)

Non-Hispanic White, male

2003.....	*	*	0.3	0.6	1.3	2.1	5.0	9.6	27.3
2002.....	*	*	0.4	0.6	1.3	2.6	5.4	10.5	34.5
2001.....	*	*	0.4	0.7	1.1	2.5	4.5	11.8	31.7
2000.....	*	*	0.4	0.8	1.7	2.6	5.8	12.3	39.8
1999.....	*	*	*	1.0	1.3	2.8	5.5	13.7	42.0

Non-Hispanic White, female

2003.....	*	*	0.5	0.7	1.1	2.2	5.8	9.4	34.5
2002.....	*	0.3	0.5	0.6	1.3	2.1	5.1	12.5	38.6
2001.....	*	0.3	0.4	0.8	1.2	2.5	5.3	10.9	42.7
2000.....	*	*	*	0.9	1.8	2.2	6.3	15.1	47.4
1999.....	*	0.4	*	0.5	1.4	2.2	5.6	14.1	51.6

Non-Hispanic Black, both sexes

2003.....	*	1.0	*	*	*	4.5	9.1	18.9	54.1
2002.....	*	*	1.5	2.0	3.1	4.4	10.4	20.0	57.2
2001.....	*	*	*	*	2.9	4.1	10.3	22.3	52.1
2000.....	*	*	*	2.0	3.2	6.7	11.2	21.3	64.2
1999.....	*	*	1.6	2.0	3.3	8.1	11.3	29.1	66.0

Non-Hispanic Black, male

2003.....	*	*	*	*	*	7.1	13.5	23.4	46.9
2002.....	*	*	*	*	*	*	9.4	25.0	44.9
2001.....	*	*	*	*	*	*	11.4	23.7	56.6
2000.....	*	*	*	*	*	9.9	14.0	23.2	62.2
1999.....	*	*	*	*	*	12.5	13.5	33.7	74.9

Non-Hispanic Black, female

2003.....	*	*	*	*	*	*	6.4	16.6	57.0
2002.....	*	*	*	*	*	*	10.9	17.5	61.9
2001.....	*	*	*	*	*	*	9.6	21.7	50.5
2000.....	*	*	*	*	*	4.5	9.6	20.3	65.0
1999.....	*	*	*	*	*	5.3	9.9	26.8	62.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other nutritional deficiencies (E50-E64)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	*	*	*	*	*	*	*	0.4	2.1
2002.....	*	*	*	*	*	*	0.5	0.6	3.2
2001.....	*	*	*	*	*	*	0.5	0.8	2.7
2000.....	*	*	*	*	*	0.3	0.4	1.2	3.8
1999.....	*	*	*	*	*	*	0.5	1.2	3.9

Non-Hispanic 4/, male

2003.....	*	*	*	*	*	*	*	*	2.3
2002.....	*	*	*	*	*	*	*	*	1.9
2001.....	*	*	*	*	*	*	*	*	1.6
2000.....	*	*	*	*	*	*	*	1.4	3.0
1999.....	*	*	*	*	*	*	*	1.3	3.8

Non-Hispanic 4/, female

2003.....	*	*	*	*	*	*	*	*	1.9
2002.....	*	*	*	*	*	*	*	0.7	3.7
2001.....	*	*	*	*	*	*	*	1.0	3.2
2000.....	*	*	*	*	*	*	*	1.1	4.1
1999.....	*	*	*	*	*	*	*	1.1	4.0

Non-Hispanic White, both sexes

2003.....	*	*	*	*	*	*	*	0.5	2.1
2002.....	*	*	*	*	*	*	0.5	0.6	3.2
2001.....	*	*	*	*	*	*	0.4	0.8	2.8
2000.....	*	*	*	*	*	0.3	*	1.2	3.8
1999.....	*	*	*	*	*	*	0.4	1.1	3.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Meningitis (G00,G03)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.3	0.3	0.3	0.4	0.6	0.5	0.8	0.7	1.1
2002.....	0.3	0.3	0.3	0.4	0.5	0.7	0.6	0.6	1.2
2001.....	0.3	0.3	0.4	0.5	0.5	0.5	0.6	0.9	0.9
2000.....	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1	1.2
1999.....	0.3	0.3	0.3	0.5	0.5	0.7	0.9	1.2	1.4

Non-Hispanic 4/, male

2003.....	0.3	0.4	0.4	0.5	0.6	*	0.9	*	*
2002.....	0.4	0.3	0.4	*	0.6	0.9	*	*	*
2001.....	0.3	0.4	0.4	0.6	0.5	0.7	0.9	*	*
2000.....	0.3	0.3	0.4	0.5	0.6	0.9	1.3	1.1	*
1999.....	0.4	0.4	0.3	0.4	0.6	0.8	0.9	1.3	*

Non-Hispanic 4/, female

2003.....	0.2	*	0.3	*	0.6	0.4	0.7	0.7	1.0
2002.....	*	0.3	*	*	*	0.6	0.5	*	1.2
2001.....	0.2	*	0.4	0.4	0.5	0.4	*	0.9	0.7
2000.....	0.2	0.3	*	*	0.4	0.5	0.6	1.1	1.1
1999.....	*	*	*	0.5	*	0.7	0.8	1.1	1.4

Non-Hispanic White, both sexes

2003.....	0.2	0.2	0.3	0.3	0.5	0.4	0.8	0.6	1.0
2002.....	0.2	0.2	0.2	0.3	0.4	0.6	0.5	0.6	1.2
2001.....	0.2	0.2	0.3	0.5	0.4	0.6	0.6	0.8	0.7
2000.....	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1	1.1
1999.....	0.2	0.2	0.3	0.4	0.4	0.6	0.7	1.1	1.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

All origins 3/, both sexes

2003.....	*	0.2	0.7	2.0	6.4	19.9	49.3	93.3	138.2
2002.....	*	0.2	0.6	1.9	6.5	18.4	45.4	89.8	135.2
2001.....	*	0.2	0.7	1.8	5.8	18.1	46.4	90.9	134.2
2000.....	*	0.2	0.6	1.7	6.6	16.7	44.8	87.6	131.9
1999.....	*	0.2	0.6	1.6	5.6	16.7	41.8	82.8	124.4

All origins 3/, male

2003.....	*	0.2	0.8	2.7	8.9	29.2	74.2	140.7	222.4
2002.....	*	0.3	0.8	2.5	9.8	27.3	67.2	136.6	220.1
2001.....	*	0.3	1.0	2.7	8.2	26.5	68.8	141.7	224.4
2000.....	*	*	0.8	2.2	9.0	24.1	67.4	135.0	222.3
1999.....	*	0.3	0.7	2.3	7.7	24.8	64.4	131.3	205.6

All origins 3/, female

2003.....	*	0.2	0.6	1.4	4.3	12.4	31.5	64.3	101.0
2002.....	*	*	0.5	1.4	3.8	11.3	30.0	61.3	98.4
2001.....	*	*	0.4	1.0	3.7	11.5	30.7	60.4	96.4
2000.....	*	*	0.4	1.3	4.4	10.9	29.1	59.5	95.0
1999.....	*	*	0.4	0.9	3.9	10.4	26.0	54.4	91.7

Hispanic, both sexes

2003.....	*	*	*	*	3.7	13.2	28.2	59.8	77.7
2002.....	*	*	*	*	5.9	12.0	27.7	53.0	76.5
2001.....	*	*	*	*	4.8	13.5	30.6	60.6	86.3
2000.....	*	*	*	*	5.7	10.5	27.2	56.8	86.3
1999.....	*	*	*	*	4.1	13.1	31.3	51.1	77.2

Hispanic, male

2003.....	*	*	*	*	6.5	20.1	38.9	91.5	129.3
2002.....	*	*	*	*	8.8	19.7	40.3	67.5	113.8
2001.....	*	*	*	*	*	16.9	43.0	101.3	127.6
2000.....	*	*	*	*	*	13.1	39.1	86.9	143.1
1999.....	*	*	*	*	*	16.8	46.8	83.3	105.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

Hispanic, female

2003.....	*	*	*	*	*	7.9	20.7	39.2	51.6
2002.....	*	*	*	*	*	*	18.7	43.8	58.0
2001.....	*	*	*	*	*	10.8	21.8	35.4	66.1
2000.....	*	*	*	*	*	8.5	18.8	38.5	58.4
1999.....	*	*	*	*	*	10.3	20.4	31.9	63.4

Non-Hispanic 4/, both sexes

2003.....	*	0.3	0.7	2.1	6.6	20.3	50.3	94.6	140.6
2002.....	*	0.2	0.6	1.9	6.6	18.8	46.2	91.2	137.4
2001.....	*	0.2	0.7	1.9	5.9	18.4	47.1	91.8	135.6
2000.....	*	0.2	0.6	1.8	6.6	17.1	45.5	88.5	133.3
1999.....	*	0.2	0.6	1.6	5.7	16.9	42.2	83.8	125.8

Non-Hispanic 4/, male

2003.....	*	0.3	0.9	2.8	9.0	29.7	76.0	142.7	226.5
2002.....	*	0.3	0.8	2.4	9.8	27.7	68.5	139.5	224.4
2001.....	*	0.3	1.1	2.8	8.3	27.1	69.9	142.9	227.8
2000.....	*	*	0.9	2.3	9.1	24.7	68.5	136.7	225.2
1999.....	*	0.3	0.7	2.4	7.7	25.2	65.1	132.9	209.4

Non-Hispanic 4/, female

2003.....	*	0.2	0.6	1.4	4.5	12.7	32.0	65.3	102.8
2002.....	*	*	0.5	1.5	3.8	11.6	30.5	61.9	99.8
2001.....	*	*	0.4	1.0	3.8	11.5	31.1	61.3	97.3
2000.....	*	*	0.4	1.3	4.4	11.0	29.4	60.2	96.1
1999.....	*	*	0.4	1.0	4.0	10.4	26.2	55.1	92.3

Non-Hispanic White, both sexes

2003.....	*	0.3	0.7	2.1	7.0	21.7	53.6	100.6	148.8
2002.....	*	0.3	0.6	2.1	7.0	20.0	49.3	97.4	145.8
2001.....	*	0.2	0.8	2.0	6.3	19.7	50.4	97.5	143.5
2000.....	*	0.2	0.7	1.9	7.1	18.2	48.4	93.7	141.2
1999.....	*	0.2	0.6	1.7	6.1	18.0	44.6	88.6	132.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

Non-Hispanic White, male

2003.....	*	*	0.9	2.8	9.5	31.5	80.4	151.4	237.9
2002.....	*	0.4	0.8	2.7	10.3	29.3	72.7	148.2	237.8
2001.....	*	0.3	1.1	3.1	8.9	28.7	74.7	151.0	241.7
2000.....	*	*	0.9	2.5	9.8	26.2	72.0	143.4	238.6
1999.....	*	0.3	0.8	2.5	8.1	26.9	68.5	140.4	221.0

Non-Hispanic White, female

2003.....	*	*	0.6	1.5	4.7	13.7	34.3	69.4	109.6
2002.....	*	*	0.5	1.6	4.0	12.4	32.6	66.3	106.0
2001.....	*	*	0.4	0.9	4.0	12.4	33.1	65.2	102.7
2000.....	*	*	0.5	1.4	4.8	11.8	31.8	64.1	101.9
1999.....	*	*	0.4	1.0	4.3	10.8	27.8	58.0	97.7

Non-Hispanic Black, both sexes

2003.....	*	*	*	1.9	4.2	11.0	23.3	38.4	53.5
2002.....	*	*	*	*	4.5	9.6	21.1	32.8	51.7
2001.....	*	*	*	*	2.8	9.1	19.3	35.6	59.0
2000.....	*	*	*	*	3.2	8.9	21.0	37.4	52.5
1999.....	*	*	*	*	3.6	8.7	19.8	32.9	51.7

Non-Hispanic Black, male

2003.....	*	*	*	*	5.6	17.2	37.8	58.1	99.1
2002.....	*	*	*	*	7.4	16.1	31.0	56.2	87.5
2001.....	*	*	*	*	*	15.1	26.1	58.9	99.3
2000.....	*	*	*	*	*	14.0	37.6	65.2	88.1
1999.....	*	*	*	*	5.8	12.5	32.4	49.6	87.9

Non-Hispanic Black, female

2003.....	*	*	*	*	*	6.8	14.7	28.0	35.9
2002.....	*	*	*	*	*	5.3	15.1	20.7	38.0
2001.....	*	*	*	*	*	5.0	15.3	23.7	44.1
2000.....	*	*	*	*	*	5.4	11.0	23.3	39.2
1999.....	*	*	*	*	*	6.2	12.2	24.6	38.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)									
All origins 3/, both sexes									
2003.....	0.1	0.3	1.0	3.3	10.0	33.3	98.6	255.0	802.4
2002.....	*	0.2	0.9	3.2	9.8	30.6	95.4	245.7	752.3
2001.....	*	0.3	1.2	3.3	9.0	29.2	88.5	232.5	710.3
2000.....	*	0.3	1.1	3.2	9.1	28.9	85.2	221.1	667.7
1999.....	0.1	0.2	1.0	3.0	8.7	26.7	78.6	206.2	601.3
All origins 3/, male									
2003.....	*	0.3	1.0	2.9	10.1	32.1	92.1	231.2	606.5
2002.....	*	0.2	0.8	3.1	9.2	29.5	90.9	220.6	572.0
2001.....	*	0.3	0.9	3.2	9.0	27.7	85.6	216.6	556.1
2000.....	*	0.3	0.9	2.9	9.5	29.1	79.9	206.8	522.1
1999.....	*	*	1.1	3.0	8.8	26.9	77.4	198.2	485.7
All origins 3/, female									
2003.....	*	0.3	0.9	3.7	10.0	34.2	103.2	269.5	889.0
2002.....	*	*	1.0	3.3	10.3	31.5	98.5	260.9	830.5
2001.....	*	0.3	1.4	3.4	9.0	30.4	90.5	242.1	774.9
2000.....	*	0.3	1.2	3.5	8.7	28.8	88.9	229.5	727.0
1999.....	*	0.3	0.8	3.0	8.6	26.6	79.4	210.9	648.0
Hispanic, both sexes									
2003.....	*	*	*	2.5	6.4	25.0	67.6	162.3	493.4
2002.....	*	*	*	*	5.9	16.8	56.3	144.4	438.5
2001.....	*	*	*	*	5.1	15.8	54.9	134.6	399.5
2000.....	*	*	*	*	7.2	17.8	55.1	123.1	380.2
1999.....	*	*	*	*	4.3	16.4	53.6	115.1	351.5
Hispanic, male									
2003.....	*	*	*	*	*	24.3	69.3	155.9	426.4
2002.....	*	*	*	*	*	16.2	54.4	127.0	341.3
2001.....	*	*	*	*	*	12.8	60.7	126.0	306.6
2000.....	*	*	*	*	*	22.8	55.4	134.0	284.2
1999.....	*	*	*	*	*	17.3	52.3	95.9	278.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alzheimer's disease (G30)

Hispanic, female

2003.....	*	*	*	*	6.9	25.6	66.4	166.5	527.3
2002.....	*	*	*	*	5.8	17.2	57.6	155.5	486.6
2001.....	*	*	*	*	*	18.1	50.8	139.9	444.9
2000.....	*	*	*	*	8.2	14.0	54.9	116.5	427.3
1999.....	*	*	*	*	*	15.7	54.5	126.6	386.8

Non-Hispanic 4/, both sexes

2003.....	*	0.3	1.0	3.4	10.3	33.7	100.2	258.9	814.0
2002.....	*	0.2	1.0	3.3	10.0	31.4	97.2	249.6	763.2
2001.....	*	0.3	1.2	3.4	9.3	30.0	90.0	236.3	720.5
2000.....	*	0.3	1.1	3.3	9.2	29.5	86.5	224.5	676.7
1999.....	0.1	0.2	1.0	3.1	9.0	27.3	79.5	209.1	608.9

Non-Hispanic 4/, male

2003.....	*	0.2	1.0	3.0	10.3	32.5	93.4	234.4	613.6
2002.....	*	0.3	0.8	3.2	9.4	30.2	92.6	224.5	581.4
2001.....	*	0.3	1.0	3.3	9.2	28.5	86.6	220.3	565.2
2000.....	*	0.3	1.0	3.0	9.8	29.3	81.0	209.5	531.0
1999.....	*	*	1.2	3.1	9.0	27.4	78.5	201.6	492.8

Non-Hispanic 4/, female

2003.....	*	0.3	1.0	3.7	10.2	34.7	105.0	273.8	902.1
2002.....	*	*	1.1	3.4	10.6	32.4	100.5	264.8	841.6
2001.....	*	0.3	1.4	3.5	9.3	31.1	92.3	245.8	785.1
2000.....	*	0.3	1.2	3.6	8.7	29.6	90.3	233.3	735.6
1999.....	*	*	0.9	3.1	9.0	27.2	80.3	213.5	655.4

Non-Hispanic White, both sexes

2003.....	*	0.3	1.1	3.7	10.5	35.1	103.5	268.7	846.9
2002.....	*	0.3	1.1	3.5	10.5	32.8	101.2	259.6	793.9
2001.....	*	0.4	1.4	3.7	9.6	31.5	92.5	244.5	749.2
2000.....	*	0.3	1.2	3.5	9.7	30.5	89.5	232.7	705.1
1999.....	*	0.3	1.1	3.4	9.4	28.1	82.4	216.8	636.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alzheimer's disease (G30)

Non-Hispanic White, male

2003.....	*	0.3	1.1	3.1	10.6	33.5	96.2	241.9	638.1
2002.....	*	*	0.9	3.4	9.6	31.4	96.2	232.0	605.4
2001.....	*	0.3	1.0	3.5	9.4	29.7	89.2	226.7	587.2
2000.....	*	*	1.0	3.2	10.3	30.2	83.8	217.3	551.7
1999.....	*	*	1.2	3.4	9.3	28.0	81.3	209.4	515.1

Non-Hispanic White, female

2003.....	*	0.3	1.1	4.2	10.5	36.3	108.8	285.2	938.8
2002.....	*	*	1.2	3.6	11.4	34.0	104.8	276.6	875.3
2001.....	*	0.4	1.7	3.9	9.8	32.9	94.9	255.3	816.6
2000.....	*	*	1.4	3.9	9.2	30.8	93.4	241.9	767.0
1999.....	*	*	0.9	3.4	9.5	28.1	83.2	221.1	685.0

Non-Hispanic Black, both sexes

2003.....	*	*	*	1.8	10.8	30.7	93.1	210.5	591.1
2002.....	*	*	*	2.5	8.4	27.5	82.1	188.2	548.7
2001.....	*	*	*	1.8	8.9	23.8	88.4	186.8	513.2
2000.....	*	*	*	2.2	7.6	26.5	72.3	173.9	450.9
1999.....	*	*	*	*	7.0	25.8	65.1	154.1	374.7

Non-Hispanic Black, male

2003.....	*	*	*	*	11.4	32.7	94.8	209.8	465.6
2002.....	*	*	*	*	9.2	29.4	82.2	189.5	414.6
2001.....	*	*	*	*	9.9	24.8	86.3	188.1	420.4
2000.....	*	*	*	*	8.0	28.4	65.5	157.9	378.1
1999.....	*	*	*	*	8.2	28.0	62.8	141.7	320.9

Non-Hispanic Black, female

2003.....	*	*	*	*	10.3	29.3	92.1	210.9	639.7
2002.....	*	*	*	*	7.8	26.2	82.1	187.5	599.9
2001.....	*	*	*	*	8.1	23.2	89.7	186.2	547.5
2000.....	*	*	*	*	7.3	25.2	76.4	181.9	478.0
1999.....	*	*	*	*	6.1	24.4	66.4	160.2	394.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

All origins 3/, both sexes									
2003.....	87.7	142.8	224.5	363.7	574.0	946.1	1,627.3	2,918.8	7,138.0
2002.....	87.9	144.2	233.5	374.0	602.6	993.2	1,700.2	3,034.4	7,389.6
2001.....	85.6	144.9	234.9	385.7	619.6	1,017.5	1,759.4	3,133.9	7,673.7
2000.....	88.5	146.7	248.4	407.5	647.9	1,066.0	1,824.4	3,253.7	8,030.3
1999.....	88.3	148.7	254.8	416.0	678.9	1,111.3	1,888.7	3,356.4	8,197.1
All origins 3/, male									
2003.....	123.1	204.5	314.7	497.9	765.9	1,225.3	2,051.8	3,517.1	7,329.3
2002.....	123.9	206.1	326.7	512.6	802.0	1,287.4	2,140.7	3,633.4	7,638.8
2001.....	121.4	204.7	327.6	526.3	821.4	1,325.6	2,223.2	3,774.2	7,936.0
2000.....	126.8	209.0	350.6	556.9	857.2	1,403.8	2,305.7	3,956.7	8,448.1
1999.....	125.7	212.5	358.4	568.4	908.7	1,461.9	2,392.7	4,078.9	8,603.2
All origins 3/, female									
2003.....	53.3	83.7	139.7	241.8	407.6	722.2	1,325.1	2,552.9	7,053.5
2002.....	52.9	85.0	146.0	248.2	430.4	758.0	1,389.0	2,670.1	7,281.5
2001.....	50.7	87.6	148.1	258.3	445.9	772.7	1,434.7	2,750.2	7,563.9
2000.....	51.4	87.0	152.8	272.0	468.5	799.9	1,489.2	2,839.0	7,860.1
1999.....	52.1	87.5	158.0	278.2	482.2	837.3	1,538.3	2,933.7	8,033.5
Hispanic, both sexes									
2003.....	54.6	94.7	165.1	277.9	457.5	768.9	1,336.3	2,247.8	5,176.6
2002.....	56.4	96.2	171.9	299.2	471.4	810.5	1,353.8	2,354.6	5,369.7
2001.....	54.0	103.2	165.7	283.6	490.6	837.6	1,459.3	2,475.8	6,036.2
2000.....	56.1	99.6	178.2	310.5	501.0	861.4	1,451.4	2,583.3	6,168.2
1999.....	59.1	108.5	186.0	313.9	543.7	875.1	1,519.7	2,691.9	6,335.9
Hispanic, male									
2003.....	76.4	138.4	233.3	385.1	608.6	1,004.0	1,642.5	2,633.2	5,079.7
2002.....	79.0	135.0	247.0	413.5	626.1	1,077.3	1,686.2	2,749.8	5,494.9
2001.....	77.7	146.6	233.8	383.5	672.6	1,084.7	1,808.6	3,004.8	6,155.6
2000.....	80.2	144.5	257.4	426.9	658.5	1,145.9	1,849.2	3,116.9	6,304.3
1999.....	83.2	160.7	259.8	427.7	730.3	1,182.4	1,968.4	3,137.4	6,595.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Major cardiovascular diseases (I00-I78)									
Hispanic, female									
2003.....	32.3	52.3	101.9	184.4	333.4	588.4	1,119.1	1,996.9	5,225.8
2002.....	33.4	58.5	102.9	199.5	345.0	605.0	1,116.5	2,102.3	5,307.7
2001.....	30.2	61.7	104.1	197.2	343.2	649.1	1,211.3	2,148.2	5,977.9
2000.....	32.0	56.8	106.6	210.2	373.5	645.9	1,169.6	2,258.6	6,101.4
1999.....	35.3	58.9	119.4	215.9	392.8	643.9	1,203.1	2,426.1	6,210.9
Non-Hispanic 4/, both sexes									
2003.....	91.0	146.7	228.4	368.7	580.8	954.6	1,639.3	2,943.4	7,206.9
2002.....	90.9	147.9	237.6	378.1	609.6	1,001.0	1,713.0	3,056.0	7,451.4
2001.....	88.3	147.8	239.3	391.7	626.6	1,025.2	1,768.6	3,152.1	7,715.2
2000.....	91.1	149.9	252.6	412.4	655.5	1,073.4	1,835.5	3,270.0	8,076.4
1999.....	90.6	151.3	258.7	421.2	684.9	1,119.5	1,898.7	3,370.5	8,239.8
Non-Hispanic 4/, male									
2003.....	127.8	210.0	319.7	503.9	774.5	1,235.1	2,068.8	3,551.2	7,419.6
2002.....	128.4	211.5	331.6	517.8	810.8	1,295.2	2,157.3	3,663.2	7,713.8
2001.....	125.4	208.7	333.1	533.8	828.4	1,335.1	2,236.3	3,795.4	7,990.6
2000.....	130.7	213.2	355.8	563.0	866.4	1,411.9	2,318.5	3,979.0	8,513.1
1999.....	129.0	215.8	363.7	574.7	916.0	1,469.7	2,402.8	4,101.3	8,657.6
Non-Hispanic 4/, female									
2003.....	55.4	86.3	142.4	245.5	412.2	728.9	1,333.5	2,572.8	7,113.4
2002.....	54.7	87.1	149.1	251.0	435.1	765.3	1,399.4	2,687.5	7,338.2
2001.....	52.6	89.5	151.1	262.3	452.1	778.4	1,441.4	2,767.2	7,600.7
2000.....	53.0	89.2	155.8	275.5	473.9	806.2	1,499.3	2,852.2	7,899.8
1999.....	53.4	89.4	160.3	281.8	486.4	845.2	1,548.4	2,943.2	8,072.5
Non-Hispanic White, both sexes									
2003.....	78.3	127.1	205.3	335.9	541.5	912.5	1,603.0	2,932.8	7,304.7
2002.....	78.6	129.6	214.1	346.2	572.1	961.8	1,673.1	3,043.0	7,546.1
2001.....	76.0	129.1	215.0	359.1	584.0	984.9	1,727.4	3,139.5	7,811.4
2000.....	78.1	132.4	228.1	378.9	617.5	1,032.1	1,791.9	3,258.1	8,179.0
1999.....	77.8	133.6	232.2	388.2	648.2	1,074.2	1,856.0	3,358.3	8,342.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic White, male

2003.....	113.8	187.0	292.8	467.4	733.0	1,190.2	2,037.8	3,555.0	7,561.6
2002.....	115.3	190.6	304.1	481.2	771.3	1,257.4	2,127.0	3,670.7	7,845.8
2001.....	111.7	188.0	305.3	498.8	785.3	1,292.9	2,203.7	3,797.2	8,116.3
2000.....	117.3	193.1	326.7	527.5	829.2	1,372.2	2,284.2	3,984.9	8,650.5
1999.....	115.3	195.1	333.2	537.4	879.0	1,426.0	2,370.6	4,111.9	8,776.1

Non-Hispanic White, female

2003.....	43.1	68.4	121.1	213.6	371.2	684.2	1,288.4	2,550.3	7,191.6
2002.....	42.2	69.9	127.6	220.8	395.7	719.9	1,348.3	2,659.1	7,416.6
2001.....	40.6	71.4	128.4	229.5	406.1	734.6	1,390.3	2,742.5	7,684.6
2000.....	39.4	73.0	133.8	241.2	431.4	758.3	1,445.9	2,826.1	7,988.9
1999.....	40.8	73.5	135.8	249.9	445.8	793.8	1,495.0	2,913.7	8,170.1

Non-Hispanic Black, both sexes

2003.....	188.3	313.0	460.6	705.3	999.0	1,508.1	2,276.6	3,444.0	6,923.9
2002.....	188.5	308.2	477.1	708.5	1,015.7	1,529.7	2,367.5	3,588.7	7,136.9
2001.....	185.9	313.3	478.6	725.1	1,073.1	1,567.7	2,444.2	3,650.2	7,366.6
2000.....	195.9	310.0	497.1	748.1	1,078.7	1,639.8	2,519.2	3,750.0	7,576.5
1999.....	192.5	317.0	517.6	753.4	1,106.8	1,719.8	2,571.4	3,809.5	7,624.4

Non-Hispanic Black, male

2003.....	245.3	422.8	621.2	935.5	1,291.9	1,922.6	2,820.7	4,073.6	6,625.9
2002.....	242.6	413.5	644.4	949.5	1,310.3	1,933.1	2,831.4	4,127.6	6,982.7
2001.....	242.7	413.4	637.0	947.4	1,361.6	2,006.9	2,966.0	4,316.4	7,413.8
2000.....	252.1	415.3	680.4	967.2	1,346.6	2,057.4	3,018.8	4,434.6	7,755.5
1999.....	251.7	431.3	697.1	1,003.1	1,415.5	2,159.6	3,060.6	4,425.3	7,952.2

Non-Hispanic Black, female

2003.....	138.7	220.2	328.7	525.9	781.8	1,228.6	1,953.5	3,114.6	7,039.3
2002.....	141.6	219.1	339.9	520.3	797.4	1,259.2	2,090.4	3,308.3	7,195.9
2001.....	136.8	228.2	348.9	550.3	860.0	1,275.0	2,132.8	3,310.8	7,349.1
2000.....	147.3	220.3	347.1	575.2	881.0	1,362.6	2,218.0	3,405.4	7,510.0
1999.....	141.5	220.2	371.1	557.6	878.0	1,427.9	2,276.7	3,504.3	7,501.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)									
All origins 3/, both sexes									
2003.....	71.3	116.6	185.0	296.1	456.3	730.9	1,218.3	2,151.6	5,278.4
2002.....	71.3	119.2	192.8	304.4	479.6	766.1	1,278.2	2,234.3	5,446.8
2001.....	69.3	119.5	194.2	314.1	494.4	787.7	1,321.2	2,309.0	5,664.2
2000.....	71.2	120.4	205.5	330.6	515.8	826.7	1,372.3	2,392.1	5,926.1
1999.....	71.7	123.5	211.0	341.9	547.1	867.9	1,427.8	2,484.7	6,063.0
All origins 3/, male									
2003.....	104.0	173.3	268.1	416.1	624.7	975.4	1,589.0	2,695.8	5,621.5
2002.....	104.7	177.0	278.2	429.1	654.3	1,025.8	1,667.3	2,786.9	5,823.5
2001.....	102.6	175.2	279.7	440.9	672.2	1,054.4	1,732.2	2,882.7	6,040.5
2000.....	107.1	178.1	298.9	463.9	702.0	1,119.7	1,787.5	3,012.3	6,430.0
1999.....	107.0	182.9	306.5	478.2	752.8	1,174.1	1,865.5	3,123.3	6,561.6
All origins 3/, female									
2003.....	39.5	62.4	106.7	187.0	310.3	534.8	954.5	1,818.9	5,126.7
2002.....	38.8	63.9	112.5	191.2	328.8	558.5	1,003.3	1,898.3	5,283.3
2001.....	37.0	66.1	114.2	199.0	341.4	575.8	1,033.5	1,965.3	5,506.8
2000.....	36.4	65.0	118.1	209.8	356.3	596.0	1,083.0	2,026.2	5,720.9
1999.....	37.5	66.6	121.7	218.6	371.2	628.7	1,123.5	2,111.0	5,862.1
Hispanic, both sexes									
2003.....	39.9	71.0	126.8	219.2	353.0	598.1	1,000.2	1,685.0	3,932.9
2002.....	41.0	73.9	134.3	234.7	376.2	620.0	1,028.8	1,746.8	4,121.9
2001.....	38.3	77.1	129.3	225.5	391.0	646.2	1,097.2	1,866.0	4,607.5
2000.....	39.9	76.8	141.1	237.3	388.4	654.8	1,096.6	1,940.5	4,701.8
1999.....	42.6	84.9	146.0	251.5	425.7	676.2	1,167.9	2,013.5	4,888.6
Hispanic, male									
2003.....	58.0	108.6	186.7	305.9	487.8	805.4	1,260.2	2,028.2	3,874.5
2002.....	59.4	108.7	199.5	335.0	511.6	846.9	1,314.4	2,109.2	4,301.8
2001.....	58.7	115.3	191.0	311.8	547.1	860.1	1,385.1	2,307.5	4,784.3
2000.....	60.8	115.6	208.7	338.5	525.4	892.6	1,411.5	2,374.9	4,897.5
1999.....	63.4	130.7	213.6	347.6	588.1	933.4	1,526.8	2,398.7	5,012.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)									
Hispanic, female									
2003.....	21.3	34.3	71.3	143.7	242.5	438.9	815.6	1,461.7	3,962.5
2002.....	22.4	40.2	74.4	147.2	265.7	445.1	824.9	1,515.4	4,032.8
2001.....	17.8	40.7	73.5	150.9	264.6	483.0	892.8	1,592.5	4,521.1
2000.....	19.1	39.8	80.0	150.1	277.5	474.6	873.7	1,676.2	4,605.8
1999.....	21.9	41.5	84.9	168.6	294.4	482.7	914.6	1,783.7	4,828.9
Non-Hispanic 4/, both sexes									
2003.....	74.5	120.5	188.9	300.6	462.4	737.2	1,227.0	2,168.1	5,324.5
2002.....	74.2	122.7	196.7	308.3	484.9	772.1	1,287.0	2,249.3	5,484.9
2001.....	72.1	122.6	198.4	319.3	499.9	793.7	1,327.9	2,320.3	5,688.6
2000.....	73.9	123.4	209.4	335.7	522.3	833.0	1,380.1	2,402.3	5,953.6
1999.....	74.1	126.1	214.7	346.6	552.7	874.4	1,434.0	2,493.7	6,086.3
Non-Hispanic 4/, male									
2003.....	108.7	178.7	273.2	422.3	632.2	982.7	1,602.4	2,720.7	5,690.6
2002.....	109.3	182.2	283.1	434.2	661.3	1,032.2	1,679.9	2,809.3	5,874.4
2001.....	106.6	179.5	285.1	448.0	678.1	1,061.9	1,743.4	2,898.2	6,077.8
2000.....	111.1	182.4	304.2	470.1	710.1	1,127.0	1,798.4	3,028.8	6,474.6
1999.....	110.6	186.3	311.6	484.3	759.8	1,180.7	1,873.1	3,140.2	6,602.0
Non-Hispanic 4/, female									
2003.....	41.3	64.8	109.3	189.7	314.5	539.6	959.7	1,831.1	5,163.6
2002.....	40.4	65.8	115.3	193.8	332.0	563.7	1,009.7	1,909.4	5,316.8
2001.....	38.7	68.0	117.0	202.2	345.9	580.0	1,037.2	1,974.4	5,526.8
2000.....	37.9	66.9	120.5	213.4	360.8	600.9	1,089.1	2,033.1	5,742.9
1999.....	38.8	68.3	124.0	221.4	374.9	634.5	1,129.0	2,115.7	5,879.8
Non-Hispanic White, both sexes									
2003.....	66.2	107.6	173.7	278.8	436.3	710.7	1,206.9	2,168.5	5,412.2
2002.....	66.3	110.8	180.7	286.8	460.5	748.2	1,262.7	2,248.0	5,568.4
2001.....	64.0	110.9	181.6	298.4	471.8	769.0	1,302.6	2,318.8	5,776.0
2000.....	65.8	112.1	192.8	313.1	497.3	805.7	1,353.2	2,402.0	6,047.9
1999.....	65.8	114.4	196.2	324.5	528.0	844.9	1,407.9	2,492.7	6,176.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)									
Non-Hispanic White, male									
2003.....	99.5	163.7	255.9	398.9	604.6	954.8	1,588.2	2,735.5	5,821.0
2002.....	100.8	168.9	264.7	409.8	636.5	1,010.8	1,665.4	2,827.0	5,994.1
2001.....	97.9	166.9	265.9	426.1	650.7	1,037.0	1,726.7	2,911.3	6,195.4
2000.....	102.8	169.3	284.7	446.8	686.5	1,102.9	1,780.9	3,046.4	6,606.6
1999.....	101.8	172.6	289.8	459.4	735.6	1,154.2	1,858.9	3,162.2	6,713.7
Non-Hispanic White, female									
2003.....	33.2	52.6	94.6	167.1	286.6	510.1	930.9	1,820.0	5,232.2
2002.....	32.2	54.0	99.9	172.5	304.7	533.4	974.5	1,893.8	5,384.5
2001.....	30.5	56.0	100.8	180.0	313.7	551.2	1,002.4	1,961.1	5,601.6
2000.....	29.1	56.0	105.0	189.2	330.9	566.6	1,052.6	2,019.1	5,822.7
1999.....	30.3	57.3	106.9	199.5	346.0	598.4	1,091.5	2,097.7	5,962.4
Non-Hispanic Black, both sexes									
2003.....	143.0	238.0	355.0	540.9	764.5	1,128.5	1,655.8	2,471.6	4,977.8
2002.....	142.1	236.4	372.5	550.7	774.7	1,140.0	1,731.2	2,577.8	5,151.0
2001.....	140.3	236.6	375.4	554.8	820.9	1,173.2	1,795.1	2,634.8	5,307.7
2000.....	145.2	236.1	386.8	579.6	826.1	1,242.2	1,852.8	2,693.6	5,424.6
1999.....	145.9	244.9	407.2	587.8	861.8	1,297.7	1,889.6	2,762.0	5,514.7
Non-Hispanic Black, male									
2003.....	192.4	329.9	489.3	726.1	1,012.2	1,475.4	2,104.0	3,007.3	4,897.8
2002.....	190.7	324.8	512.2	751.5	1,014.9	1,464.8	2,114.1	3,061.4	5,161.2
2001.....	188.8	320.2	510.4	740.5	1,053.7	1,531.1	2,230.3	3,202.0	5,495.4
2000.....	195.0	325.5	540.6	763.9	1,047.1	1,582.4	2,260.1	3,247.6	5,677.2
1999.....	197.0	341.6	565.4	797.1	1,122.5	1,658.3	2,283.9	3,268.7	5,877.1
Non-Hispanic Black, female									
2003.....	100.0	160.3	244.6	396.7	580.8	894.7	1,389.7	2,191.3	5,008.9
2002.....	100.0	161.6	258.0	393.8	596.6	922.2	1,502.6	2,326.2	5,147.0
2001.....	98.4	165.5	264.8	408.8	649.0	934.8	1,535.3	2,345.9	5,238.2
2000.....	102.0	159.9	260.9	434.1	663.2	1,016.4	1,607.3	2,414.7	5,330.8
1999.....	101.9	163.0	278.2	423.7	668.6	1,058.5	1,652.1	2,510.8	5,379.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	8.1	9.1	*	*
2001.....	*	*	*	*	*	7.3	*	*	*
2000.....	*	*	*	*	*	7.4	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.5	0.6	1.0	1.8	3.0	4.7	8.2	11.8	20.9
2002.....	0.4	0.7	1.3	2.1	2.8	4.9	8.5	11.5	18.1
2001.....	0.5	0.7	1.2	1.7	3.2	5.0	8.2	11.6	18.0
2000.....	0.6	0.8	1.4	2.2	3.4	5.5	8.3	11.8	17.1
1999.....	0.5	0.8	1.4	2.1	3.5	5.9	8.4	12.7	18.3

Non-Hispanic 4/, male

2003.....	0.4	0.4	0.8	1.5	2.3	3.7	6.5	9.3	16.6
2002.....	0.4	0.6	1.0	1.4	1.9	4.2	6.0	8.4	12.0
2001.....	0.5	0.5	1.1	1.2	2.3	3.3	6.0	7.8	13.1
2000.....	0.5	0.6	1.1	1.8	2.3	3.5	5.9	8.1	12.5
1999.....	0.4	0.6	1.0	1.3	2.4	4.8	5.6	9.1	10.7

Non-Hispanic 4/, female

2003.....	0.6	0.8	1.1	2.1	3.6	5.6	9.4	13.3	22.8
2002.....	0.5	0.9	1.5	2.6	3.6	5.4	10.2	13.4	20.8
2001.....	0.6	0.9	1.3	2.1	4.0	6.3	9.8	13.9	20.0
2000.....	0.7	0.9	1.6	2.5	4.5	7.1	10.0	13.9	19.0
1999.....	0.6	1.0	1.8	2.8	4.4	6.7	10.3	14.8	21.3

Non-Hispanic White, both sexes

2003.....	0.4	0.5	0.8	1.6	3.1	4.8	8.7	12.6	22.3
2002.....	0.4	0.7	1.1	2.0	2.7	5.1	8.7	12.3	19.3
2001.....	0.4	0.6	1.1	1.7	3.1	5.0	8.6	12.4	19.0
2000.....	0.5	0.7	1.3	2.2	3.6	5.6	8.9	12.5	18.0
1999.....	0.5	0.7	1.4	2.0	3.6	6.1	8.8	13.3	19.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Non-Hispanic White, male

2003.....	0.3	0.3	0.6	1.3	2.3	3.8	7.1	10.1	17.7
2002.....	0.3	0.5	0.9	1.4	1.8	4.4	6.2	9.1	12.9
2001.....	0.4	0.4	1.0	1.2	2.2	3.4	6.3	8.2	13.7
2000.....	0.5	0.6	1.1	1.9	2.3	3.5	6.3	8.7	13.0
1999.....	0.4	0.6	1.0	1.3	2.6	5.0	5.7	9.7	11.6

Non-Hispanic White, female

2003.....	0.5	0.6	1.0	1.9	3.8	5.6	9.9	14.1	24.4
2002.....	0.4	0.9	1.4	2.5	3.6	5.7	10.6	14.2	22.0
2001.....	0.4	0.7	1.1	2.1	4.0	6.3	10.2	14.9	21.2
2000.....	0.6	0.8	1.4	2.4	4.7	7.3	10.7	14.8	20.0
1999.....	0.5	0.9	1.8	2.7	4.4	7.0	11.0	15.5	22.3

Non-Hispanic Black, both sexes

2003.....	1.3	1.5	2.2	3.3	3.2	4.9	4.0	*	7.1
2002.....	0.9	1.1	2.1	2.9	3.2	*	4.9	*	*
2001.....	1.1	1.7	2.2	2.0	3.4	5.2	5.0	*	6.9
2000.....	1.0	1.1	2.0	2.1	3.5	3.7	3.8	*	8.6
1999.....	0.9	1.7	2.3	2.1	2.7	4.1	4.2	*	8.4

Non-Hispanic Black, male

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic Black, female

2003.....	1.7	1.8	*	3.4	*	6.4	*	*	*
2002.....	*	*	*	3.7	*	*	5.6	*	*
2001.....	*	2.3	2.8	*	3.8	7.3	6.5	*	*
2000.....	*	*	*	*	4.5	*	*	*	*
1999.....	*	*	2.8	*	*	4.8	*	*	11.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart disease (I11)

All origins 3/, both sexes

2003.....	6.6	9.4	11.7	14.9	19.4	25.1	39.5	66.8	195.0
2002.....	5.9	8.6	11.1	14.3	18.9	24.0	37.9	66.0	187.0
2001.....	5.1	7.7	9.7	12.5	16.3	23.5	36.6	66.0	189.5
2000.....	4.7	6.6	9.3	13.2	16.5	23.0	36.3	66.1	193.1
1999.....	4.5	6.2	9.6	11.9	16.3	22.1	36.4	64.7	184.9

All origins 3/, male

2003.....	9.0	12.8	16.0	18.8	25.2	29.4	42.8	63.3	142.2
2002.....	8.0	11.9	15.2	19.0	23.6	27.0	41.0	62.2	134.6
2001.....	7.0	10.3	12.8	16.5	20.5	27.1	38.9	63.5	134.4
2000.....	6.3	9.4	12.1	17.4	20.7	27.2	37.3	61.8	139.1
1999.....	6.1	8.1	12.7	15.0	19.5	24.6	39.4	59.2	133.6

All origins 3/, female

2003.....	4.4	6.1	7.7	11.2	14.4	21.6	37.2	69.0	218.3
2002.....	3.9	5.4	7.2	10.0	14.8	21.6	35.8	68.2	209.7
2001.....	3.2	5.1	6.8	8.9	12.7	20.6	35.1	67.5	212.6
2000.....	3.2	3.8	6.6	9.3	12.9	19.6	35.7	68.6	215.1
1999.....	3.0	4.3	6.8	9.2	13.5	20.2	34.3	67.9	205.5

Hispanic, both sexes

2003.....	5.0	6.6	9.8	14.7	19.8	26.9	42.6	73.0	179.3
2002.....	4.0	6.1	10.7	14.2	19.8	25.6	43.2	65.1	165.7
2001.....	3.1	7.2	8.5	11.8	16.8	26.3	37.7	63.1	151.9
2000.....	3.1	4.6	5.8	11.3	13.3	24.7	35.5	65.2	143.3
1999.....	2.6	4.4	8.4	9.5	16.4	17.3	33.2	65.8	152.2

Hispanic, male

2003.....	7.2	9.7	14.4	20.7	25.7	37.6	51.6	84.2	144.7
2002.....	6.2	9.0	15.4	19.2	24.4	33.8	51.8	77.8	137.2
2001.....	4.5	10.4	11.4	17.1	22.3	31.6	38.9	66.2	144.7
2000.....	4.7	6.3	8.1	14.1	16.4	30.1	43.6	64.8	112.9
1999.....	4.1	5.6	11.8	13.4	20.1	21.9	34.3	59.7	107.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)									
Hispanic, female									
2003.....	2.8	3.6	5.5	9.4	14.9	18.7	36.2	65.6	196.9
2002.....	*	3.4	6.4	9.8	16.0	19.3	37.0	56.9	179.9
2001.....	*	4.1	6.0	7.3	12.3	22.3	36.8	61.1	155.5
2000.....	*	3.0	*	8.9	10.9	20.6	29.8	65.4	158.3
1999.....	*	3.2	5.4	6.2	13.4	13.8	32.5	69.4	173.7
Non-Hispanic 4/, both sexes									
2003.....	6.8	9.6	11.8	14.8	19.3	24.8	39.3	66.3	195.0
2002.....	6.1	8.7	11.0	14.2	18.5	23.8	37.5	65.7	187.1
2001.....	5.2	7.6	9.7	12.4	16.3	23.2	36.4	66.0	190.3
2000.....	4.8	6.7	9.5	13.3	16.5	22.7	36.1	66.0	194.3
1999.....	4.7	6.3	9.6	12.0	16.1	22.2	36.4	64.3	185.3
Non-Hispanic 4/, male									
2003.....	9.1	13.0	15.9	18.6	25.0	28.6	42.2	62.0	141.7
2002.....	8.2	12.1	15.1	18.9	23.1	26.4	40.1	61.1	133.7
2001.....	7.2	10.2	12.7	16.2	20.4	26.5	38.6	63.4	133.4
2000.....	6.4	9.6	12.3	17.6	20.6	26.7	36.6	61.5	139.6
1999.....	6.2	8.3	12.6	14.8	19.3	24.4	39.4	58.8	134.2
Non-Hispanic 4/, female									
2003.....	4.5	6.3	7.9	11.3	14.3	21.7	37.2	68.9	218.5
2002.....	4.1	5.6	7.2	9.9	14.5	21.6	35.6	68.5	210.2
2001.....	3.3	5.2	6.8	9.0	12.7	20.5	34.8	67.6	214.0
2000.....	3.3	3.8	6.8	9.3	13.0	19.5	35.8	68.6	216.5
1999.....	3.2	4.3	6.8	9.4	13.4	20.4	34.3	67.6	205.8
Non-Hispanic White, both sexes									
2003.....	4.2	6.5	8.4	11.1	14.2	19.5	32.4	60.3	184.9
2002.....	4.2	6.0	7.8	10.0	13.7	18.7	31.0	57.5	177.0
2001.....	3.3	5.0	6.7	8.8	11.8	17.9	30.4	58.9	180.4
2000.....	3.1	4.3	6.2	9.1	12.2	17.0	29.7	58.0	182.5
1999.....	2.9	4.0	6.3	8.5	11.7	16.7	29.6	56.8	175.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)									
Non-Hispanic White, male									
2003.....	6.2	9.6	11.9	14.4	18.7	22.4	34.9	56.2	130.8
2002.....	6.0	8.8	11.2	13.5	17.3	21.0	33.2	53.3	123.5
2001.....	4.8	7.1	9.4	12.0	15.4	21.1	32.5	55.5	122.0
2000.....	4.5	6.5	8.5	12.4	15.6	20.7	30.2	52.9	129.5
1999.....	4.1	5.7	8.6	10.9	13.9	18.7	32.1	51.2	123.8
Non-Hispanic White, female									
2003.....	2.3	3.5	5.0	7.9	10.1	17.2	30.6	62.8	208.7
2002.....	2.5	3.3	4.5	6.8	10.5	16.8	29.5	60.0	200.1
2001.....	1.8	2.9	4.2	5.8	8.6	15.3	28.9	60.9	204.7
2000.....	1.7	2.1	4.0	6.0	9.2	14.0	29.4	61.0	203.9
1999.....	1.6	2.4	4.0	6.3	9.7	15.1	27.9	60.1	195.7
Non-Hispanic Black, both sexes									
2003.....	24.3	33.6	42.5	48.6	64.0	78.2	115.8	147.2	334.9
2002.....	19.8	30.8	40.4	51.2	61.1	77.3	111.6	168.4	322.1
2001.....	18.9	28.5	35.7	44.1	56.4	78.2	106.1	155.9	328.7
2000.....	18.1	25.9	38.4	48.9	55.6	82.1	111.8	165.8	344.1
1999.....	18.2	25.3	38.4	42.2	57.4	79.5	114.9	161.4	308.5
Non-Hispanic Black, male									
2003.....	30.0	42.4	55.6	60.1	87.6	101.1	132.2	150.2	291.9
2002.....	24.9	40.4	54.2	72.1	78.8	91.0	126.8	166.9	264.7
2001.....	25.5	36.5	45.2	56.4	71.1	91.2	116.6	173.3	288.7
2000.....	22.0	36.5	48.8	65.1	71.2	95.7	117.1	179.4	275.9
1999.....	22.9	32.0	50.0	52.5	72.8	90.7	130.5	166.5	266.2
Non-Hispanic Black, female									
2003.....	19.2	26.2	31.7	39.6	46.6	62.7	106.0	145.7	351.6
2002.....	15.4	22.6	29.2	34.8	48.0	68.1	102.6	169.2	344.1
2001.....	13.2	21.7	28.0	34.3	45.6	69.5	99.9	147.0	343.5
2000.....	14.8	16.9	29.9	36.1	44.0	73.1	108.6	159.0	369.4
1999.....	14.1	19.7	29.0	34.1	46.1	72.1	105.4	158.9	324.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)									
Hispanic, female									
2003.....	*	*	*	*	*	*	*	*	20.3
2002.....	*	*	*	*	*	*	*	*	21.6
2001.....	*	*	*	*	*	*	*	*	24.2
2000.....	*	*	*	*	*	*	*	*	26.7
1999.....	*	*	*	*	*	*	*	*	20.8
Non-Hispanic 4/, both sexes									
2003.....	0.4	0.6	0.9	1.1	1.6	2.8	5.1	9.7	25.6
2002.....	0.4	0.5	0.9	1.1	1.6	2.8	4.8	9.4	23.5
2001.....	0.3	0.4	0.8	1.2	1.7	2.7	5.0	9.2	23.1
2000.....	0.3	0.4	0.8	1.0	1.6	3.0	5.0	9.8	24.2
1999.....	0.4	0.5	0.9	1.4	2.0	3.0	6.6	12.3	29.3
Non-Hispanic 4/, male									
2003.....	0.6	0.8	1.1	1.3	1.7	3.0	5.6	10.1	26.3
2002.....	0.5	0.6	1.1	1.3	1.5	3.2	5.1	9.7	22.9
2001.....	0.5	0.6	0.9	1.1	2.0	2.8	5.3	9.0	23.8
2000.....	0.4	0.5	0.9	1.0	1.7	3.1	4.8	10.7	27.9
1999.....	0.5	0.6	1.0	1.5	2.2	3.1	7.4	13.4	31.4
Non-Hispanic 4/, female									
2003.....	0.2	0.5	0.6	0.9	1.5	2.6	4.8	9.4	25.3
2002.....	0.2	0.4	0.7	1.0	1.6	2.4	4.5	9.1	23.7
2001.....	*	0.3	0.7	1.3	1.4	2.7	4.8	9.3	22.8
2000.....	0.2	0.4	0.7	0.9	1.5	2.8	5.2	9.3	22.8
1999.....	0.3	0.3	0.7	1.2	1.9	2.9	6.1	11.7	28.4
Non-Hispanic White, both sexes									
2003.....	*	0.2	0.3	0.6	0.8	1.9	4.1	8.4	23.2
2002.....	*	*	0.3	0.5	0.7	1.8	3.6	8.2	20.8
2001.....	*	0.2	0.3	0.6	0.9	1.8	3.8	8.1	20.7
2000.....	*	0.2	0.4	0.4	0.8	2.1	3.9	8.0	21.8
1999.....	*	0.2	0.4	0.6	1.1	1.8	5.1	10.4	26.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)									
Non-Hispanic White, male									
2003.....	*	*	0.4	0.7	1.0	2.2	4.8	9.2	23.6
2002.....	*	*	0.4	0.5	0.7	2.2	4.1	8.6	20.2
2001.....	*	*	0.4	0.5	1.1	1.9	4.2	8.3	22.2
2000.....	*	*	0.4	*	1.0	2.2	4.2	9.0	24.3
1999.....	*	*	0.4	0.7	1.3	1.8	5.9	12.2	28.7
Non-Hispanic White, female									
2003.....	*	*	*	0.4	0.7	1.6	3.6	7.9	23.0
2002.....	*	*	*	0.5	0.8	1.5	3.3	8.0	21.0
2001.....	*	*	*	0.7	0.8	1.8	3.6	7.9	20.1
2000.....	*	*	*	*	0.7	2.0	3.7	7.4	20.8
1999.....	*	*	*	0.6	0.9	1.7	4.6	9.3	25.1
Non-Hispanic Black, both sexes									
2003.....	2.4	3.9	5.5	5.6	7.7	11.3	16.9	24.4	53.8
2002.....	2.3	3.3	6.1	6.3	8.1	12.5	17.6	23.2	57.2
2001.....	1.9	2.4	4.4	5.9	8.6	11.8	18.4	22.9	50.6
2000.....	2.0	2.5	4.1	5.4	8.4	11.9	17.0	31.3	50.9
1999.....	2.3	2.3	5.2	7.4	10.3	15.7	22.0	35.8	66.0
Non-Hispanic Black, male									
2003.....	3.5	4.8	7.8	6.4	6.8	12.1	15.9	24.1	51.1
2002.....	3.1	4.7	7.8	7.5	7.4	14.0	17.4	22.6	62.4
2001.....	3.0	3.6	5.6	6.5	10.9	13.4	18.5	18.8	45.0
2000.....	2.8	3.0	5.1	5.7	9.3	13.3	12.5	33.5	64.6
1999.....	3.1	3.6	6.4	9.6	10.9	17.7	23.7	30.1	60.6
Non-Hispanic Black, female									
2003.....	*	3.2	3.6	4.9	8.3	10.7	17.5	24.5	54.9
2002.....	1.5	2.2	4.7	5.4	8.5	11.5	17.7	23.5	55.2
2001.....	*	*	3.4	5.4	6.9	10.8	18.4	25.0	52.6
2000.....	*	2.1	3.3	5.2	7.7	10.9	19.7	30.2	45.8
1999.....	1.7	*	4.1	5.6	9.8	14.5	21.0	38.6	68.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)									
All origins 3/, both sexes									
2003.....	45.9	80.7	132.4	218.7	335.0	538.1	886.4	1,538.2	3,610.2
2002.....	46.4	83.5	140.1	226.2	358.4	569.8	940.2	1,611.2	3,775.0
2001.....	45.6	84.8	142.8	236.6	372.4	591.8	984.3	1,685.2	3,958.2
2000.....	48.0	86.7	153.9	250.0	390.8	628.0	1,032.5	1,761.7	4,182.5
1999.....	48.5	90.1	157.2	260.0	420.1	661.9	1,081.1	1,841.7	4,318.1
All origins 3/, male									
2003.....	71.4	126.9	200.3	319.7	476.1	744.1	1,200.1	2,013.5	4,005.2
2002.....	72.3	130.3	211.4	331.4	508.5	791.5	1,274.8	2,094.2	4,205.3
2001.....	71.6	131.1	215.5	344.4	525.7	820.3	1,337.5	2,180.0	4,395.8
2000.....	76.9	134.2	232.8	362.9	550.5	879.6	1,394.4	2,305.0	4,701.7
1999.....	77.0	140.3	237.7	376.1	595.2	924.6	1,462.5	2,400.4	4,837.5
All origins 3/, female									
2003.....	21.1	36.5	68.4	126.9	212.6	373.0	663.1	1,247.5	3,435.5
2002.....	21.4	38.7	73.1	130.7	228.8	392.6	704.0	1,317.5	3,588.3
2001.....	20.3	40.3	74.8	138.8	240.5	410.2	737.0	1,388.7	3,775.1
2000.....	20.0	41.3	80.0	147.7	253.9	429.9	780.5	1,441.2	3,971.0
1999.....	20.8	42.1	82.1	155.0	270.2	456.7	815.9	1,514.7	4,108.8
Hispanic, both sexes									
2003.....	23.8	50.6	91.5	164.3	267.1	466.2	771.2	1,301.0	2,970.0
2002.....	26.9	52.6	99.9	184.7	293.4	493.0	805.7	1,353.7	3,182.9
2001.....	25.8	55.5	98.1	172.8	306.5	518.7	881.4	1,481.0	3,620.2
2000.....	27.4	55.4	109.9	186.8	312.5	521.9	873.5	1,544.0	3,697.2
1999.....	29.9	64.4	111.6	199.2	340.5	551.1	948.1	1,596.5	3,876.2
Hispanic, male									
2003.....	36.0	82.6	141.2	235.0	382.1	642.3	995.3	1,590.5	2,933.9
2002.....	41.4	80.3	155.3	272.2	413.4	681.9	1,053.3	1,662.0	3,374.9
2001.....	41.3	85.6	152.6	245.2	441.9	712.3	1,147.8	1,860.8	3,828.2
2000.....	44.8	86.7	167.7	276.6	435.1	726.3	1,161.2	1,936.1	3,865.6
1999.....	46.7	104.2	171.1	279.1	483.8	773.0	1,270.9	1,949.1	3,987.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)									
Hispanic, female									
2003.....	11.4	19.6	45.5	102.7	172.6	331.1	612.2	1,112.5	2,988.3
2002.....	12.1	25.8	49.1	108.5	195.5	347.5	628.8	1,157.0	3,087.8
2001.....	10.1	26.6	48.7	110.0	196.9	371.0	692.3	1,245.8	3,518.6
2000.....	10.0	25.5	57.6	109.4	213.2	367.1	669.8	1,305.3	3,614.6
1999.....	13.2	26.6	58.0	130.4	224.7	384.2	720.2	1,386.1	3,822.4
Non-Hispanic 4/, both sexes									
2003.....	48.1	83.3	135.1	221.8	338.7	540.9	889.9	1,545.2	3,629.3
2002.....	48.4	85.9	142.7	228.2	361.5	572.0	943.8	1,617.7	3,787.0
2001.....	47.4	86.9	145.7	240.2	375.6	594.1	985.9	1,688.1	3,959.4
2000.....	49.7	88.9	156.4	253.1	394.4	631.3	1,036.0	1,764.3	4,187.0
1999.....	50.0	91.8	159.8	262.9	423.4	664.7	1,082.6	1,844.1	4,319.6
Non-Hispanic 4/, male									
2003.....	75.1	130.6	203.9	324.5	480.9	747.7	1,207.3	2,028.4	4,045.5
2002.....	75.4	134.1	214.8	334.2	512.8	794.4	1,281.4	2,107.5	4,228.4
2001.....	74.5	134.5	219.2	349.7	529.1	823.6	1,342.0	2,185.7	4,408.5
2000.....	79.6	137.3	236.5	366.8	555.3	883.8	1,399.7	2,312.5	4,721.1
1999.....	79.5	142.5	241.2	380.4	599.4	927.6	1,464.8	2,409.4	4,854.5
Non-Hispanic 4/, female									
2003.....	22.0	38.0	70.1	128.2	215.0	374.6	664.0	1,250.5	3,446.4
2002.....	22.2	39.7	74.9	131.8	230.2	393.9	705.5	1,320.5	3,596.5
2001.....	21.2	41.4	76.6	140.6	242.9	411.4	736.7	1,390.3	3,772.7
2000.....	20.8	42.4	81.4	149.8	255.9	431.9	782.9	1,441.2	3,971.0
1999.....	21.4	43.1	83.6	156.2	272.3	459.0	817.1	1,513.5	4,105.4
Non-Hispanic White, both sexes									
2003.....	45.7	78.2	128.7	210.6	325.4	527.9	879.9	1,548.1	3,691.6
2002.....	46.0	81.2	135.4	217.1	348.4	560.1	931.0	1,622.0	3,848.2
2001.....	44.8	82.1	137.8	228.7	359.7	581.5	973.2	1,692.1	4,022.8
2000.....	46.9	84.3	148.3	240.8	380.8	617.2	1,021.9	1,770.0	4,261.3
1999.....	47.3	86.6	151.0	251.6	410.9	649.3	1,071.3	1,851.7	4,390.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)									
Non-Hispanic White, male									
2003.....	72.5	124.9	197.4	312.8	468.4	733.9	1,202.8	2,043.6	4,149.5
2002.....	73.5	129.4	206.7	322.1	501.2	786.3	1,278.6	2,128.1	4,324.5
2001.....	72.2	129.4	210.4	338.5	514.8	812.4	1,338.8	2,207.1	4,506.9
2000.....	77.2	132.8	226.8	355.2	545.0	873.4	1,394.6	2,335.6	4,833.4
1999.....	77.1	136.7	231.0	368.1	589.6	915.9	1,466.0	2,439.8	4,954.1
Non-Hispanic White, female									
2003.....	19.2	32.6	62.5	115.5	198.3	358.6	646.2	1,243.6	3,490.0
2002.....	18.8	34.2	66.9	119.6	213.1	375.2	682.3	1,312.3	3,642.5
2001.....	17.7	35.8	68.1	126.7	222.7	393.8	714.3	1,381.2	3,821.4
2000.....	17.0	36.7	73.2	134.8	236.5	410.9	759.9	1,433.9	4,030.7
1999.....	17.9	37.6	74.7	143.8	254.3	436.8	794.3	1,504.7	4,165.3
Non-Hispanic Black, both sexes									
2003.....	73.2	136.6	217.2	357.2	509.6	764.3	1,148.6	1,713.3	3,328.7
2002.....	74.4	139.0	234.9	366.7	530.0	787.4	1,206.0	1,780.5	3,481.1
2001.....	74.3	141.7	239.4	381.7	570.0	820.0	1,257.4	1,844.5	3,628.7
2000.....	79.8	142.9	254.5	398.2	573.6	874.6	1,307.5	1,894.4	3,701.7
1999.....	78.7	152.1	262.9	400.2	602.8	911.2	1,327.7	1,932.5	3,817.9
Non-Hispanic Black, male									
2003.....	105.8	200.7	306.9	496.8	685.4	1,036.6	1,511.0	2,158.3	3,309.2
2002.....	103.8	199.8	335.4	516.5	713.6	1,034.1	1,508.4	2,183.9	3,572.8
2001.....	104.5	203.8	337.8	522.1	752.5	1,102.1	1,591.8	2,254.3	3,791.8
2000.....	113.9	201.0	371.1	536.0	735.3	1,148.8	1,639.7	2,329.3	3,911.2
1999.....	112.9	221.3	378.6	558.1	795.8	1,191.3	1,633.7	2,312.3	4,082.5
Non-Hispanic Black, female									
2003.....	44.9	82.5	143.6	248.4	379.2	580.7	933.3	1,480.5	3,336.2
2002.....	48.9	87.6	152.6	249.7	393.9	621.9	1,025.5	1,570.5	3,446.0
2001.....	48.2	88.8	158.8	271.3	435.3	632.0	1,057.8	1,635.8	3,568.3
2000.....	50.2	93.4	159.1	289.5	454.4	692.6	1,107.3	1,675.5	3,623.9
1999.....	49.3	93.5	168.6	276.3	459.7	725.3	1,143.3	1,744.2	3,719.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)									
All origins 3/, both sexes									
2003.....	18.8	33.2	55.5	90.8	134.5	207.4	325.6	532.5	1,118.8
2002.....	19.0	34.9	58.4	96.2	148.5	227.5	352.5	572.7	1,191.3
2001.....	19.7	36.5	61.3	100.1	154.1	235.6	374.0	611.5	1,265.4
2000.....	21.1	37.8	67.0	110.9	166.8	254.7	405.3	650.1	1,338.5
1999.....	21.5	39.0	69.7	114.9	181.7	271.6	422.9	689.4	1,379.3
All origins 3/, male									
2003.....	29.2	52.1	83.4	131.6	188.5	281.4	430.0	682.4	1,263.5
2002.....	29.2	53.7	86.6	139.0	207.6	312.0	466.1	726.6	1,366.4
2001.....	30.8	55.6	91.3	143.5	214.2	322.0	495.6	776.1	1,461.3
2000.....	33.6	57.5	99.7	158.6	231.3	349.6	535.2	835.8	1,568.3
1999.....	33.7	60.9	105.3	164.9	253.5	373.8	564.2	885.1	1,622.1
All origins 3/, female									
2003.....	8.7	15.1	29.3	53.8	87.7	148.0	251.3	440.9	1,054.8
2002.....	9.1	17.0	32.0	57.4	97.6	159.8	272.2	479.1	1,115.3
2001.....	9.0	18.2	33.2	60.7	102.4	167.0	288.8	512.9	1,183.5
2000.....	8.9	18.9	36.5	67.7	111.5	179.9	314.9	540.6	1,244.8
1999.....	9.7	18.0	36.5	69.6	120.2	191.7	324.7	574.9	1,281.5
Hispanic, both sexes									
2003.....	8.3	19.9	35.7	56.7	104.7	179.5	272.6	453.7	952.0
2002.....	9.5	20.2	40.1	75.6	112.2	188.8	297.8	475.5	1,048.2
2001.....	10.5	20.5	36.2	67.2	113.4	195.2	328.7	511.1	1,185.5
2000.....	11.0	21.3	41.7	72.1	118.8	198.8	331.8	537.5	1,177.1
1999.....	12.9	24.3	38.6	78.2	129.6	207.5	365.1	558.6	1,227.8
Hispanic, male									
2003.....	12.2	32.0	53.1	80.0	145.3	243.0	333.6	538.5	963.6
2002.....	14.5	29.7	60.9	108.5	155.8	255.8	383.9	595.9	1,107.7
2001.....	17.2	30.1	53.3	91.3	162.3	264.4	408.5	641.5	1,331.3
2000.....	17.6	31.2	62.7	102.8	162.9	265.9	435.5	672.9	1,273.8
1999.....	19.0	37.1	57.3	106.9	181.6	284.1	511.0	655.5	1,276.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)									
Hispanic, female									
2003.....	4.3	8.1	19.7	36.5	71.5	130.8	229.3	398.6	946.1
2002.....	4.6	11.1	20.9	47.0	76.6	137.2	236.3	398.6	1,018.8
2001.....	3.7	11.3	20.6	46.3	73.7	142.4	272.1	430.3	1,114.2
2000.....	4.5	11.9	22.6	45.7	83.0	148.0	258.3	455.1	1,129.7
1999.....	6.8	12.2	21.7	53.4	87.5	149.8	262.1	500.8	1,204.1
Non-Hispanic 4/, both sexes									
2003.....	19.9	34.4	57.0	93.3	136.3	208.7	327.8	535.2	1,123.8
2002.....	20.0	36.2	59.8	97.6	150.8	229.1	354.6	575.7	1,194.0
2001.....	20.6	37.8	63.2	102.1	156.7	237.5	375.3	614.1	1,265.3
2000.....	22.0	39.1	68.7	113.4	169.5	257.1	407.8	652.8	1,340.8
1999.....	22.2	40.1	71.7	117.2	184.5	274.1	424.3	692.4	1,381.0
Non-Hispanic 4/, male									
2003.....	31.0	53.9	85.5	135.2	191.2	283.1	434.3	687.7	1,274.5
2002.....	30.8	55.8	88.5	141.0	210.7	314.5	469.4	731.0	1,374.2
2001.....	32.1	57.7	94.0	146.7	217.3	324.5	498.6	779.7	1,463.9
2000.....	35.1	59.5	102.2	162.2	235.0	353.2	538.5	840.4	1,576.2
1999.....	35.0	62.6	108.3	168.4	257.4	377.0	564.9	891.3	1,631.5
Non-Hispanic 4/, female									
2003.....	9.2	15.8	30.1	55.0	88.6	148.8	251.9	442.2	1,057.5
2002.....	9.6	17.5	32.8	58.1	98.8	160.7	273.6	481.4	1,116.3
2001.....	9.5	18.7	34.2	61.6	104.3	168.3	288.9	515.1	1,182.7
2000.....	9.3	19.4	37.4	69.0	113.2	181.3	316.8	542.3	1,245.6
1999.....	9.9	18.4	37.5	70.6	121.9	193.5	326.6	576.1	1,280.8
Non-Hispanic White, both sexes									
2003.....	19.3	33.2	55.8	91.5	133.9	206.3	326.2	534.8	1,137.4
2002.....	19.4	35.4	58.6	95.4	148.4	226.2	351.1	575.3	1,202.8
2001.....	20.2	36.7	61.6	100.5	153.1	234.9	372.6	614.8	1,276.8
2000.....	21.6	37.9	67.3	111.3	166.6	253.9	403.8	654.9	1,355.3
1999.....	21.8	39.3	70.2	115.6	183.3	270.8	422.0	695.3	1,394.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)									
Non-Hispanic White, male									
2003.....	30.7	52.8	85.2	134.4	190.9	281.3	434.8	690.7	1,300.9
2002.....	30.7	55.5	88.0	140.6	210.8	315.1	469.8	734.8	1,393.0
2001.....	32.4	57.3	93.3	147.1	216.2	324.4	500.3	786.9	1,483.2
2000.....	35.3	59.3	101.2	161.8	234.8	352.6	539.5	847.9	1,600.7
1999.....	35.2	62.6	107.0	168.7	258.1	377.4	569.2	903.1	1,653.8
Non-Hispanic White, female									
2003.....	8.1	13.9	27.4	51.5	83.3	144.6	247.6	439.0	1,065.3
2002.....	8.3	15.8	30.3	53.5	93.2	153.4	266.1	477.8	1,120.7
2001.....	8.1	16.5	31.1	57.2	97.4	162.3	282.2	510.9	1,190.9
2000.....	8.0	17.0	34.9	64.5	106.7	174.5	308.4	540.2	1,256.3
1999.....	8.6	16.4	35.0	66.5	117.7	185.8	318.7	572.7	1,290.7
Non-Hispanic Black, both sexes									
2003.....	28.3	51.0	81.0	126.6	183.4	273.0	406.5	605.6	1,103.6
2002.....	28.5	51.8	86.3	138.6	197.2	299.8	440.2	648.9	1,212.9
2001.....	28.6	54.3	90.9	137.8	214.1	308.1	458.0	677.1	1,247.0
2000.....	30.2	56.7	96.2	154.4	222.8	333.5	495.3	705.5	1,281.8
1999.....	30.9	56.3	98.8	153.1	227.4	349.5	499.9	735.3	1,328.6
Non-Hispanic Black, male									
2003.....	39.8	73.1	110.1	171.3	234.1	357.6	522.8	742.6	1,121.9
2002.....	38.0	72.3	117.2	180.6	252.6	375.9	539.9	783.1	1,296.3
2001.....	37.5	72.9	122.5	176.6	267.8	390.8	556.0	802.9	1,389.4
2000.....	41.9	74.6	133.6	200.3	275.0	426.1	596.8	865.1	1,426.6
1999.....	42.3	78.8	141.5	202.7	295.6	436.9	597.4	863.5	1,509.4
Non-Hispanic Black, female									
2003.....	18.4	32.3	57.2	91.8	145.9	216.1	337.4	533.9	1,096.5
2002.....	20.2	34.5	60.9	105.8	156.2	248.7	380.7	579.1	1,181.0
2001.....	20.8	38.5	65.0	107.3	174.5	253.0	399.5	613.0	1,194.3
2000.....	20.1	41.4	65.7	118.3	184.2	272.0	434.1	625.2	1,228.0
1999.....	21.0	37.3	64.0	114.2	176.8	291.6	441.2	671.7	1,261.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)									
Hispanic, female									
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes									
2003.....	0.5	1.0	1.3	1.8	2.9	3.8	5.4	8.6	20.6
2002.....	0.6	1.3	1.5	2.2	3.2	4.1	5.8	9.3	21.8
2001.....	0.6	1.1	1.5	2.5	3.0	4.3	5.5	9.5	22.0
2000.....	0.5	1.1	1.7	2.6	3.2	4.0	6.1	9.6	22.2
1999.....	0.5	1.0	1.7	2.3	3.2	4.5	5.6	9.5	22.2
Non-Hispanic 4/, male									
2003.....	0.8	1.4	1.9	2.6	4.0	5.1	7.4	9.8	20.2
2002.....	0.9	1.9	2.2	3.1	4.6	5.2	7.3	11.7	22.1
2001.....	1.0	1.7	2.0	3.7	4.2	5.6	6.9	11.6	20.9
2000.....	0.8	1.7	2.6	3.7	4.6	5.3	7.8	11.9	21.0
1999.....	0.8	1.4	2.6	3.2	4.5	5.9	7.7	11.0	22.9
Non-Hispanic 4/, female									
2003.....	0.3	0.5	0.8	1.1	1.9	2.7	4.0	7.8	20.7
2002.....	0.3	0.7	0.8	1.4	1.9	3.2	4.8	7.8	21.6
2001.....	0.3	0.5	1.0	1.4	2.1	3.2	4.5	8.2	22.4
2000.....	*	0.6	0.8	1.6	2.1	3.0	4.9	8.2	22.7
1999.....	0.3	0.6	0.8	1.5	2.1	3.3	4.2	8.5	21.9
Non-Hispanic White, both sexes									
2003.....	0.4	0.9	1.2	1.6	2.7	3.6	5.2	8.4	20.9
2002.....	0.5	1.2	1.4	2.0	3.1	3.9	5.8	9.2	22.0
2001.....	0.6	0.9	1.3	2.2	2.7	4.2	5.3	9.4	22.0
2000.....	0.4	1.0	1.4	2.4	3.0	3.8	5.7	9.5	22.5
1999.....	0.4	0.9	1.5	2.2	3.1	4.2	5.4	9.2	22.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)									
Non-Hispanic White, male									
2003.....	0.6	1.4	1.7	2.3	3.8	5.0	6.9	9.3	20.4
2002.....	0.8	1.7	2.2	2.9	4.4	4.8	7.3	11.5	22.7
2001.....	0.9	1.4	1.9	3.3	3.6	5.5	6.7	11.9	20.9
2000.....	0.8	1.6	2.3	3.4	4.2	5.0	7.4	11.8	21.3
1999.....	0.6	1.3	2.4	2.9	4.3	5.5	7.7	10.5	23.6
Non-Hispanic White, female									
2003.....	0.2	0.5	0.7	1.0	1.7	2.5	3.9	7.9	21.1
2002.....	0.3	0.7	0.8	1.2	1.9	3.1	4.7	7.8	21.8
2001.....	*	0.5	0.8	1.2	2.0	3.1	4.4	7.9	22.4
2000.....	*	0.5	0.6	1.4	2.0	2.9	4.6	8.2	23.0
1999.....	*	0.5	0.7	1.5	2.0	3.1	3.9	8.4	22.4
Non-Hispanic Black, both sexes									
2003.....	1.2	1.5	2.7	3.4	5.0	6.1	8.9	11.9	20.2
2002.....	1.1	2.5	2.1	4.2	5.1	7.3	7.7	12.0	21.5
2001.....	1.1	2.2	2.9	5.3	5.4	6.3	6.9	11.0	24.3
2000.....	1.0	2.1	3.9	4.9	5.2	6.1	10.1	11.8	21.9
1999.....	1.5	2.3	3.3	3.9	4.4	7.5	8.9	14.1	17.5
Non-Hispanic Black, male									
2003.....	1.8	2.3	4.6	5.2	7.3	8.1	14.5	19.6	24.5
2002.....	2.0	3.8	3.2	5.9	8.2	10.9	*	16.4	21.9
2001.....	*	3.7	3.5	8.4	9.4	7.9	11.4	*	24.3
2000.....	*	3.5	5.9	6.8	8.3	9.2	13.5	*	6.8
1999.....	2.5	2.9	5.7	6.5	6.9	10.4	10.2	17.7	*
Non-Hispanic Black, female									
2003.....	*	*	*	*	*	4.8	5.5	7.9	18.6
2002.....	*	*	*	*	*	4.8	7.0	9.7	21.3
2001.....	*	*	*	*	*	5.3	*	12.1	24.4
2000.....	*	*	*	3.4	*	*	8.1	9.9	22.2
1999.....	*	*	*	*	*	5.5	8.2	12.3	16.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)									
All origins 3/, both sexes									
2003.....	26.6	46.6	75.6	126.1	197.8	327.1	555.6	997.3	2,471.4
2002.....	26.8	47.3	80.2	127.8	206.8	338.4	582.1	1,029.5	2,562.5
2001.....	25.2	47.3	80.1	134.1	215.4	352.0	605.0	1,064.5	2,671.3
2000.....	26.5	47.9	85.2	136.6	221.0	369.4	621.2	1,102.2	2,822.1
1999.....	26.5	50.2	85.9	142.9	235.3	386.1	652.6	1,143.1	2,917.0
All origins 3/, male									
2003.....	41.5	73.5	115.1	185.6	283.8	457.8	763.1	1,321.6	2,722.2
2002.....	42.2	74.9	122.7	189.4	296.6	474.3	801.6	1,356.2	2,817.4
2001.....	40.0	74.0	122.2	197.4	307.4	492.9	835.3	1,392.6	2,913.8
2000.....	42.5	75.1	130.6	200.8	314.8	524.7	851.5	1,457.6	3,112.7
1999.....	42.6	78.1	129.9	208.1	337.3	545.2	890.7	1,504.6	3,193.1
All origins 3/, female									
2003.....	12.1	20.9	38.4	72.0	123.2	222.4	408.0	799.1	2,360.5
2002.....	11.9	21.0	40.3	72.0	129.4	229.7	427.1	830.8	2,451.9
2001.....	11.0	21.7	40.6	76.8	136.2	240.1	443.8	867.9	2,569.8
2000.....	10.9	21.8	42.7	78.5	140.5	247.0	460.8	892.5	2,703.8
1999.....	10.9	23.4	44.9	83.9	148.1	261.7	487.1	931.6	2,805.7
Hispanic, both sexes									
2003.....	15.4	30.6	55.6	107.1	161.9	285.4	496.6	845.6	2,012.8
2002.....	17.1	32.1	59.5	108.1	180.7	302.3	505.5	875.6	2,129.7
2001.....	15.3	34.8	61.3	104.8	191.7	321.5	549.9	968.0	2,426.6
2000.....	16.1	33.9	67.6	114.2	193.0	320.8	539.0	1,002.0	2,508.8
1999.....	17.0	39.6	72.6	119.8	210.1	343.0	579.7	1,036.7	2,640.7
Hispanic, male									
2003.....	23.7	50.4	88.1	154.3	236.2	398.0	659.9	1,051.0	1,965.7
2002.....	26.6	50.1	93.8	162.1	257.3	422.2	666.8	1,062.6	2,262.2
2001.....	24.1	55.3	98.3	152.9	277.5	444.2	735.9	1,216.7	2,479.8
2000.....	27.0	55.4	104.3	173.3	270.7	457.0	722.0	1,257.4	2,577.7
1999.....	27.6	66.6	113.3	171.0	301.0	488.3	754.4	1,292.1	2,706.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)									
Hispanic, female									
2003.....	6.9	11.4	25.5	66.1	100.8	199.0	380.7	711.9	2,036.7
2002.....	7.4	14.6	28.0	61.1	118.1	209.9	390.3	756.3	2,064.1
2001.....	6.4	15.2	27.9	63.2	122.2	227.9	417.8	813.9	2,400.7
2000.....	5.3	13.5	34.3	63.3	130.1	217.6	409.4	846.6	2,475.0
1999.....	6.4	14.0	35.9	75.7	136.6	233.7	456.5	884.3	2,608.9
Non-Hispanic 4/, both sexes									
2003.....	27.7	47.9	76.7	126.8	199.5	328.5	556.8	1,001.4	2,484.9
2002.....	27.8	48.4	81.4	128.4	207.5	338.9	583.4	1,032.8	2,571.2
2001.....	26.1	48.0	81.1	135.6	215.9	352.3	605.2	1,064.5	2,672.2
2000.....	27.2	48.7	86.0	137.1	221.6	370.1	622.1	1,101.9	2,824.0
1999.....	27.2	50.7	86.5	143.5	235.7	386.2	652.7	1,142.2	2,916.3
Non-Hispanic 4/, male									
2003.....	43.3	75.3	116.5	186.7	285.7	459.5	765.6	1,330.9	2,750.7
2002.....	43.7	76.5	124.1	190.0	297.4	474.7	804.7	1,364.8	2,832.2
2001.....	41.4	75.1	123.2	199.4	307.7	493.5	836.5	1,394.4	2,923.7
2000.....	43.7	76.1	131.7	200.9	315.7	525.3	853.4	1,460.2	3,123.9
1999.....	43.7	78.5	130.3	208.8	337.5	544.7	892.2	1,507.1	3,200.1
Non-Hispanic 4/, female									
2003.....	12.6	21.7	39.2	72.1	124.5	223.1	408.1	800.5	2,368.1
2002.....	12.3	21.5	41.2	72.3	129.5	230.0	427.1	831.3	2,458.6
2001.....	11.4	22.1	41.5	77.6	136.6	239.9	443.3	867.0	2,567.6
2000.....	11.3	22.4	43.2	79.1	140.7	247.6	461.2	890.7	2,702.7
1999.....	11.3	24.1	45.4	84.1	148.3	262.1	486.4	928.9	2,802.7
Non-Hispanic White, both sexes									
2003.....	25.9	44.2	71.7	117.5	188.8	318.0	548.5	1,004.8	2,533.4
2002.....	26.0	44.7	75.4	119.7	196.9	330.1	574.2	1,037.5	2,623.4
2001.....	24.0	44.5	74.9	126.0	203.8	342.4	595.3	1,067.8	2,724.1
2000.....	24.9	45.3	79.5	127.1	211.2	359.4	612.4	1,105.6	2,883.6
1999.....	25.2	46.5	79.3	133.8	224.5	374.3	643.9	1,147.2	2,973.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)									
Non-Hispanic White, male									
2003.....	41.1	70.7	110.5	176.0	273.7	447.6	761.1	1,343.5	2,828.2
2002.....	42.1	72.2	116.6	178.6	285.9	466.3	801.5	1,381.9	2,908.8
2001.....	39.0	70.6	115.2	188.2	295.0	482.6	831.9	1,408.3	3,002.8
2000.....	41.1	72.0	123.2	189.9	306.1	515.8	847.8	1,475.9	3,211.3
1999.....	41.4	72.8	121.6	196.5	327.2	533.0	889.2	1,526.1	3,276.7
Non-Hispanic White, female									
2003.....	10.9	18.2	34.3	63.0	113.3	211.5	394.7	796.6	2,403.6
2002.....	10.2	17.7	35.8	64.9	118.0	218.6	411.5	826.8	2,500.1
2001.....	9.3	18.8	36.2	68.2	123.3	228.4	427.7	862.3	2,608.1
2000.....	8.9	19.2	37.8	68.9	127.8	233.5	447.0	885.5	2,751.4
1999.....	9.2	20.7	38.9	75.8	134.6	247.9	471.7	923.6	2,852.2
Non-Hispanic Black, both sexes									
2003.....	43.7	84.1	133.4	227.2	321.2	485.2	733.2	1,095.8	2,204.8
2002.....	44.8	84.7	146.5	223.9	327.7	480.3	758.1	1,119.5	2,246.7
2001.....	44.6	85.2	145.6	238.5	350.5	505.5	792.5	1,156.4	2,357.3
2000.....	48.5	84.1	154.4	238.9	345.6	535.0	802.0	1,177.1	2,398.0
1999.....	46.4	93.5	160.9	243.2	371.0	554.2	818.9	1,183.2	2,471.9
Non-Hispanic Black, male									
2003.....	64.3	125.3	192.1	320.4	444.0	671.0	973.7	1,396.1	2,162.8
2002.....	63.8	123.8	215.0	329.9	452.8	647.2	959.6	1,384.5	2,254.6
2001.....	65.5	127.3	211.7	337.1	475.2	703.4	1,024.4	1,442.4	2,378.1
2000.....	70.5	122.9	231.6	328.9	452.0	713.5	1,029.4	1,448.7	2,463.5
1999.....	68.1	139.6	231.4	348.9	493.3	744.0	1,026.2	1,431.1	2,552.9
Non-Hispanic Black, female									
2003.....	25.8	49.3	85.2	154.6	230.1	359.9	590.4	938.7	2,221.1
2002.....	28.4	51.7	90.4	141.0	234.9	368.4	637.8	981.7	2,243.7
2001.....	26.6	49.4	91.4	161.0	258.3	373.7	654.1	1,010.7	2,349.6
2000.....	29.5	51.1	91.2	167.9	267.2	416.5	665.0	1,040.4	2,373.7
1999.....	27.6	54.5	103.3	160.3	280.3	428.3	694.0	1,060.2	2,441.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)									
All origins 3/, both sexes									
2003.....	9.9	18.6	26.6	40.5	53.3	73.3	110.7	186.2	442.9
2002.....	9.8	17.5	27.9	39.7	53.9	76.5	116.6	187.6	474.5
2001.....	8.8	17.4	27.7	40.6	55.3	76.7	121.8	197.2	493.6
2000.....	9.4	17.5	28.5	40.7	55.7	81.9	125.0	207.8	534.6
1999.....	9.1	17.4	28.6	43.4	59.6	86.8	131.6	218.2	563.4
All origins 3/, male									
2003.....	16.0	30.5	41.8	60.6	77.1	102.8	146.6	233.1	462.1
2002.....	15.6	28.4	44.8	59.7	78.8	106.7	154.5	232.4	491.8
2001.....	14.6	28.3	43.4	61.8	79.0	107.6	160.2	247.4	502.0
2000.....	15.5	28.4	45.8	61.3	80.8	115.6	167.1	252.8	550.8
1999.....	14.9	28.0	45.0	64.6	87.3	120.8	172.7	264.3	578.6
All origins 3/, female									
2003.....	4.0	7.2	12.4	22.2	32.6	49.7	85.2	157.6	434.4
2002.....	4.1	7.1	12.1	21.4	32.4	52.5	89.7	160.4	467.1
2001.....	3.3	7.0	12.9	21.4	34.9	52.1	94.9	167.1	490.1
2000.....	3.5	7.2	12.4	22.1	34.1	55.4	95.7	181.2	528.0
1999.....	3.5	7.3	13.4	24.1	36.0	60.2	103.1	191.3	557.2
Hispanic, both sexes									
2003.....	7.1	13.3	22.1	39.5	46.6	70.8	105.2	156.2	357.1
2002.....	6.8	11.8	22.7	39.2	49.8	75.4	105.3	155.5	358.6
2001.....	6.2	15.1	23.5	37.9	54.9	79.6	112.1	168.4	417.0
2000.....	6.3	13.8	24.0	38.9	54.1	80.2	101.3	177.1	452.5
1999.....	6.5	15.4	27.1	40.4	62.9	83.7	124.0	190.3	469.4
Hispanic, male									
2003.....	12.0	24.0	38.2	63.7	74.1	107.6	148.9	204.8	397.1
2002.....	10.7	20.3	40.4	61.8	74.8	111.5	140.1	178.4	415.0
2001.....	10.0	26.6	39.2	59.7	87.2	117.1	159.6	222.0	440.0
2000.....	10.9	24.1	39.9	64.5	74.9	115.2	158.7	225.3	493.8
1999.....	10.6	27.4	42.5	59.6	103.6	131.9	171.6	238.9	485.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)									
Hispanic, female									
2003.....	2.1	2.9	7.3	18.4	24.0	42.6	74.1	124.5	336.7
2002.....	2.8	3.5	6.4	19.5	29.3	47.7	80.4	140.9	330.7
2001.....	2.5	4.1	9.2	19.0	28.7	51.0	78.4	135.1	405.8
2000.....	*	4.0	9.5	16.9	37.1	53.8	60.6	147.8	432.3
1999.....	2.5	4.1	13.1	23.9	30.0	47.5	90.3	161.3	461.7
Non-Hispanic 4/, both sexes									
2003.....	10.1	18.9	26.7	40.1	53.5	72.9	110.2	186.8	445.3
2002.....	10.0	17.8	28.1	39.2	53.7	75.9	116.2	188.3	477.8
2001.....	9.1	17.5	27.7	40.5	54.8	75.9	121.3	197.5	494.8
2000.....	9.6	17.7	28.6	40.3	55.3	81.3	125.0	207.9	535.9
1999.....	9.3	17.4	28.5	43.0	58.9	86.0	130.8	217.9	564.9
Non-Hispanic 4/, male									
2003.....	16.3	30.9	41.6	59.6	76.8	101.5	145.1	233.3	463.3
2002.....	16.0	28.7	44.5	58.7	78.4	105.3	153.9	233.6	493.3
2001.....	14.9	28.2	43.2	61.4	77.6	106.0	158.8	247.2	503.0
2000.....	15.8	28.4	45.7	60.2	80.4	114.3	165.7	252.5	551.0
1999.....	15.2	27.7	44.8	63.8	85.4	118.6	170.8	263.4	580.2
Non-Hispanic 4/, female									
2003.....	4.1	7.5	12.6	22.4	33.2	49.8	85.3	158.4	437.4
2002.....	4.2	7.3	12.6	21.4	32.3	52.3	89.7	160.7	471.1
2001.....	3.4	7.2	13.1	21.5	35.1	51.9	95.0	167.7	491.4
2000.....	3.7	7.4	12.5	22.3	33.8	55.2	96.7	181.6	529.8
1999.....	3.6	7.5	13.3	24.0	36.1	60.5	103.0	191.3	558.8
Non-Hispanic White, both sexes									
2003.....	9.0	16.9	24.0	34.6	47.1	65.6	101.9	179.7	441.5
2002.....	9.3	15.8	24.6	34.4	46.6	68.5	107.0	180.7	473.5
2001.....	8.1	15.4	23.6	34.4	46.7	68.0	111.2	190.8	490.7
2000.....	8.5	15.7	24.5	34.3	48.6	72.3	115.5	198.2	530.2
1999.....	8.0	14.9	24.2	36.6	51.0	76.8	120.3	209.3	558.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)									
Non-Hispanic White, male									
2003.....	14.9	27.9	37.9	51.8	68.1	90.5	134.7	224.0	459.5
2002.....	15.1	26.0	39.3	52.0	68.2	96.0	142.1	222.9	489.2
2001.....	13.6	25.3	37.7	52.9	66.9	95.2	146.4	239.6	499.4
2000.....	14.2	25.6	39.5	52.4	71.8	102.1	152.6	240.8	544.7
1999.....	13.4	24.2	38.7	55.0	75.0	106.5	157.5	252.7	572.8
Non-Hispanic White, female									
2003.....	3.3	6.2	10.7	18.6	28.5	45.1	78.1	152.5	433.5
2002.....	3.5	5.8	10.5	18.0	27.5	46.0	81.9	154.9	466.8
2001.....	2.6	5.7	10.1	17.3	28.8	45.9	86.3	161.3	487.0
2000.....	2.8	5.9	10.2	17.4	28.2	48.3	89.5	172.9	524.4
1999.....	2.8	5.9	10.3	19.7	30.0	53.1	94.2	183.6	553.2
Non-Hispanic Black, both sexes									
2003.....	17.7	36.3	53.6	92.0	113.3	150.1	215.7	286.5	546.7
2002.....	17.3	35.9	60.8	86.1	120.6	158.1	233.1	303.8	580.4
2001.....	17.1	36.3	64.5	95.5	129.4	164.1	244.8	300.8	595.9
2000.....	19.7	36.9	66.3	94.4	120.8	178.9	240.8	342.8	652.2
1999.....	19.5	39.1	68.6	98.4	135.0	187.7	260.8	339.9	670.3
Non-Hispanic Black, male									
2003.....	26.8	58.7	81.8	139.3	163.8	229.2	302.8	384.9	588.1
2002.....	26.0	54.5	97.1	131.4	182.2	220.6	321.1	420.4	603.9
2001.....	26.3	56.7	97.3	145.3	185.0	241.7	332.8	385.9	609.8
2000.....	30.5	57.2	107.3	137.8	170.0	259.7	339.1	429.1	711.6
1999.....	30.5	60.8	105.2	146.7	198.2	267.3	346.6	436.6	714.3
Non-Hispanic Black, female									
2003.....	9.8	17.4	30.4	55.2	75.8	96.8	164.0	235.0	530.7
2002.....	9.7	20.1	31.1	50.8	75.0	116.1	180.5	243.1	571.5
2001.....	9.2	18.9	37.6	56.4	88.3	112.4	192.4	257.4	590.7
2000.....	10.4	19.5	32.7	60.2	84.5	125.2	181.5	299.4	630.2
1999.....	10.1	20.7	38.7	60.4	88.2	134.9	209.1	291.9	653.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)									
All origins 3/, both sexes									
2003.....	16.7	28.0	49.0	85.6	144.5	253.8	444.9	811.1	2,028.5
2002.....	17.1	29.9	52.3	88.2	152.9	261.8	465.6	841.9	2,088.0
2001.....	16.4	29.8	52.4	93.5	160.1	275.3	483.2	867.3	2,177.7
2000.....	17.0	30.3	56.7	95.9	165.3	287.4	496.2	894.4	2,287.5
1999.....	17.4	32.8	57.3	99.5	175.7	299.3	521.0	924.9	2,353.6
All origins 3/, male									
2003.....	25.5	43.0	73.3	125.0	206.7	355.0	616.5	1,088.5	2,260.1
2002.....	26.6	46.5	78.0	129.6	217.8	367.6	647.1	1,123.8	2,325.6
2001.....	25.4	45.7	78.8	135.6	228.4	385.2	675.1	1,145.1	2,411.8
2000.....	27.0	46.7	84.9	139.5	234.0	409.1	684.5	1,204.8	2,561.9
1999.....	27.7	50.2	84.9	143.5	250.0	424.4	718.0	1,240.3	2,614.4
All origins 3/, female									
2003.....	8.2	13.7	26.0	49.8	90.5	172.7	322.7	641.5	1,926.2
2002.....	7.8	13.9	28.2	50.5	97.0	177.3	337.3	670.4	1,984.9
2001.....	7.6	14.7	27.7	55.4	101.2	188.0	348.9	700.7	2,079.8
2000.....	7.3	14.6	30.3	56.4	106.3	191.6	365.1	711.3	2,175.8
1999.....	7.4	16.1	31.5	59.7	112.1	201.5	384.0	740.3	2,248.4
Hispanic, both sexes									
2003.....	8.2	17.3	33.5	67.6	115.3	214.6	391.4	689.4	1,655.7
2002.....	10.3	20.3	36.8	68.9	130.9	226.8	400.2	720.1	1,771.1
2001.....	9.0	19.7	37.9	66.9	136.8	241.9	437.7	799.6	2,009.6
2000.....	9.9	20.1	43.6	75.3	139.0	240.5	437.7	824.9	2,056.3
1999.....	10.4	24.2	45.5	79.4	147.2	259.3	455.8	846.4	2,171.4
Hispanic, male									
2003.....	11.7	26.5	50.0	90.6	162.2	290.4	511.0	846.2	1,568.6
2002.....	15.9	29.8	53.4	100.3	182.5	310.7	526.6	884.2	1,847.2
2001.....	14.1	28.7	59.1	93.2	190.3	327.1	576.3	994.7	2,039.8
2000.....	16.0	31.3	64.5	108.8	195.8	341.8	563.3	1,032.1	2,084.0
1999.....	17.0	39.2	70.7	111.4	197.4	356.4	582.8	1,053.1	2,221.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)									
Hispanic, female									
2003.....	4.8	8.5	18.2	47.6	76.8	156.4	306.5	587.4	1,700.0
2002.....	4.6	11.1	21.6	41.6	88.7	162.2	309.9	615.4	1,733.3
2001.....	3.9	11.1	18.6	44.2	93.5	176.9	339.4	678.8	1,994.8
2000.....	3.7	9.5	24.8	46.4	93.0	163.9	348.7	698.8	2,042.7
1999.....	3.9	9.9	22.7	51.8	106.6	186.2	366.2	723.0	2,147.2
Non-Hispanic 4/, both sexes									
2003.....	17.6	28.9	50.0	86.6	146.0	255.6	446.6	814.6	2,039.6
2002.....	17.7	30.6	53.3	89.2	153.8	263.0	467.1	844.5	2,093.4
2001.....	17.1	30.6	53.4	95.1	161.1	276.4	483.9	867.0	2,177.4
2000.....	17.6	31.0	57.4	96.8	166.3	288.8	497.1	894.0	2,288.1
1999.....	17.9	33.3	57.9	100.5	176.8	300.2	522.0	924.3	2,351.4
Non-Hispanic 4/, male									
2003.....	27.1	44.4	74.8	127.1	208.9	358.0	620.5	1,097.6	2,287.4
2002.....	27.7	47.8	79.6	131.3	219.0	369.4	650.8	1,131.1	2,338.9
2001.....	26.5	46.9	79.9	138.0	230.1	387.4	677.7	1,147.3	2,420.7
2000.....	27.9	47.7	86.0	140.8	235.3	411.0	687.7	1,207.7	2,572.9
1999.....	28.6	50.8	85.6	145.0	252.0	426.1	721.4	1,243.7	2,619.9
Non-Hispanic 4/, female									
2003.....	8.5	14.2	26.6	49.6	91.3	173.2	322.7	642.1	1,930.7
2002.....	8.1	14.1	28.7	51.0	97.2	177.8	337.4	670.6	1,987.5
2001.....	8.0	14.9	28.4	56.1	101.5	188.0	348.3	699.3	2,076.2
2000.....	7.6	15.0	30.7	56.9	106.9	192.4	364.5	709.1	2,172.9
1999.....	7.7	16.6	32.1	60.1	112.2	201.6	383.4	737.6	2,243.9
Non-Hispanic White, both sexes									
2003.....	16.9	27.2	47.7	82.9	141.7	252.4	446.6	825.1	2,091.9
2002.....	16.7	28.9	50.8	85.3	150.3	261.5	467.2	856.7	2,149.8
2001.....	16.0	29.1	51.3	91.6	157.1	274.4	484.0	877.0	2,233.4
2000.....	16.4	29.6	55.0	92.8	162.6	287.1	496.9	907.4	2,353.3
1999.....	17.1	31.5	55.1	97.1	173.5	297.5	523.6	937.9	2,414.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)									
Non-Hispanic White, male									
2003.....	26.3	42.8	72.7	124.2	205.6	357.1	626.4	1,119.6	2,368.6
2002.....	27.0	46.2	77.2	126.6	217.7	370.3	659.5	1,158.9	2,419.7
2001.....	25.4	45.4	77.5	135.4	228.0	387.4	685.5	1,168.6	2,503.4
2000.....	26.9	46.3	83.7	137.5	234.2	413.6	695.2	1,235.1	2,666.6
1999.....	28.0	48.6	82.9	141.5	252.2	426.5	731.7	1,273.5	2,703.9
Non-Hispanic White, female									
2003.....	7.6	12.0	23.6	44.4	84.8	166.4	316.6	644.1	1,970.1
2002.....	6.7	12.0	25.3	46.9	90.6	172.6	329.6	671.9	2,033.3
2001.....	6.7	13.1	26.2	51.0	94.5	182.5	341.4	701.0	2,121.0
2000.....	6.1	13.2	27.6	51.5	99.6	185.3	357.5	712.6	2,227.0
1999.....	6.4	14.8	28.6	56.1	104.6	194.7	377.5	739.9	2,299.0
Non-Hispanic Black, both sexes									
2003.....	26.0	47.8	79.9	135.1	207.9	335.1	517.5	809.4	1,658.1
2002.....	27.6	48.9	85.7	137.7	207.0	322.3	525.0	815.8	1,666.3
2001.....	27.5	48.9	81.1	143.0	221.1	341.4	547.6	855.6	1,761.4
2000.....	28.8	47.3	88.1	144.5	224.8	356.1	561.3	834.3	1,745.8
1999.....	26.8	54.4	92.3	144.8	236.0	366.5	558.1	843.3	1,801.6
Non-Hispanic Black, male									
2003.....	37.5	66.6	110.4	181.0	280.2	441.8	671.0	1,011.2	1,574.7
2002.....	37.8	69.4	117.8	198.6	270.6	426.6	638.5	964.1	1,650.8
2001.....	39.2	70.5	114.5	191.9	290.3	461.7	691.6	1,056.4	1,768.3
2000.....	40.0	65.6	124.3	191.1	282.0	453.8	690.3	1,019.6	1,751.9
1999.....	37.7	78.8	126.1	202.2	295.1	476.7	679.6	994.5	1,838.6
Non-Hispanic Black, female									
2003.....	15.9	31.9	54.8	99.4	154.3	263.1	426.4	703.7	1,690.4
2002.....	18.7	31.6	59.4	90.2	160.0	252.3	457.2	738.6	1,672.2
2001.....	17.4	30.5	53.8	104.7	170.0	261.3	461.7	753.4	1,758.9
2000.....	19.2	31.6	58.5	107.7	182.6	291.3	483.5	741.0	1,743.5
1999.....	17.5	33.7	64.7	99.9	192.1	293.4	484.9	768.3	1,787.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)									
All origins 3/, both sexes									
2003.....	17.9	25.3	39.1	59.7	97.3	160.2	279.1	525.5	1,427.2
2002.....	18.1	25.9	39.4	60.6	97.9	164.5	286.7	536.3	1,443.5
2001.....	17.9	25.9	39.7	62.0	100.7	164.7	287.2	537.0	1,475.5
2000.....	17.6	25.9	40.2	64.3	103.6	167.1	290.2	542.8	1,509.4
1999.....	17.8	25.9	41.8	66.5	105.2	175.0	295.4	553.5	1,512.9
All origins 3/, male									
2003.....	22.7	32.4	49.9	74.9	119.4	195.4	334.0	599.9	1,431.3
2002.....	23.5	33.6	49.4	75.7	118.8	199.8	340.7	612.4	1,448.6
2001.....	23.1	32.7	49.4	77.7	121.5	200.9	344.7	622.1	1,473.4
2000.....	23.0	33.4	52.0	80.8	126.9	205.9	345.1	626.8	1,549.1
1999.....	23.0	33.3	54.1	84.3	133.4	217.0	350.9	641.4	1,548.9
All origins 3/, female									
2003.....	13.2	18.6	28.8	45.9	78.1	131.9	240.0	480.0	1,425.4
2002.....	12.8	18.5	30.1	46.8	79.8	136.3	248.6	490.0	1,441.3
2001.....	12.8	19.5	30.6	47.8	82.8	135.9	247.0	486.0	1,476.4
2000.....	12.3	18.7	29.2	49.3	83.6	136.6	252.1	493.2	1,493.2
1999.....	12.8	18.9	30.2	50.5	81.0	142.2	256.8	502.0	1,498.5
Hispanic, both sexes									
2003.....	10.2	12.8	24.0	37.7	61.4	97.5	175.0	296.3	749.9
2002.....	9.2	14.2	21.4	31.4	57.4	91.6	167.4	307.5	740.0
2001.....	8.9	13.5	21.3	36.8	61.7	92.5	168.2	304.0	798.5
2000.....	8.5	15.6	23.3	35.8	58.2	98.1	178.7	314.7	829.4
1999.....	9.1	15.0	23.6	39.4	61.5	100.3	173.0	332.5	827.9
Hispanic, male									
2003.....	13.9	15.7	29.3	48.2	75.4	120.9	202.3	341.0	752.7
2002.....	11.2	18.4	26.6	38.7	69.8	122.0	202.2	354.6	758.0
2001.....	12.5	18.3	25.4	45.9	75.5	111.2	190.3	359.7	775.2
2000.....	10.7	20.9	31.3	44.9	70.8	124.5	194.9	362.2	890.8
1999.....	11.6	20.5	29.1	51.2	79.6	132.4	213.0	377.2	888.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)									
Hispanic, female									
2003.....	6.4	9.9	19.2	28.5	49.9	79.7	155.6	267.3	748.4
2002.....	7.2	10.2	16.7	25.0	47.3	68.3	142.6	277.4	731.1
2001.....	5.3	8.9	17.6	28.9	50.5	78.2	152.5	269.4	809.8
2000.....	6.3	10.5	16.1	28.0	48.0	78.1	167.3	285.8	799.3
1999.....	6.6	9.7	18.6	29.3	46.9	76.2	144.8	305.7	798.6
Non-Hispanic 4/, both sexes									
2003.....	18.7	26.4	40.2	61.2	99.8	164.0	284.5	535.2	1,453.7
2002.....	19.0	26.8	40.8	62.7	100.6	168.7	292.4	545.0	1,469.2
2001.....	18.7	26.9	41.1	63.8	103.1	168.7	292.4	545.3	1,497.8
2000.....	18.4	26.7	41.3	66.1	106.4	170.6	294.7	550.4	1,530.9
1999.....	18.6	26.7	43.0	68.2	107.7	178.5	300.0	560.2	1,533.9
Non-Hispanic 4/, male									
2003.....	23.6	33.8	51.5	76.5	122.4	199.8	340.8	610.9	1,460.5
2002.....	24.8	34.8	51.1	78.4	122.0	204.1	347.2	622.6	1,477.3
2001.....	24.0	33.8	51.2	79.7	124.2	205.6	351.5	632.2	1,499.0
2000.....	24.1	34.3	53.3	83.0	130.2	209.8	351.4	636.0	1,573.6
1999.....	24.0	34.2	55.7	86.2	136.5	220.7	356.0	649.4	1,571.2
Non-Hispanic 4/, female									
2003.....	13.9	19.3	29.6	47.2	80.1	135.2	244.3	489.0	1,450.7
2002.....	13.3	19.2	31.1	48.4	82.1	140.3	253.8	497.9	1,465.7
2001.....	13.4	20.3	31.5	49.3	84.9	139.2	251.1	493.4	1,497.3
2000.....	12.8	19.3	30.1	50.8	86.0	139.6	255.3	500.0	1,513.7
1999.....	13.3	19.6	31.0	51.9	83.0	145.5	261.1	508.0	1,519.0
Non-Hispanic White, both sexes									
2003.....	15.8	22.2	35.6	55.0	92.8	156.6	281.8	539.1	1,490.1
2002.....	15.6	22.8	36.0	57.1	94.9	162.5	288.2	548.1	1,503.1
2001.....	15.4	23.0	35.7	58.7	96.3	162.8	286.7	547.4	1,533.2
2000.....	15.2	22.6	36.7	60.6	99.8	163.9	288.8	553.5	1,564.2
1999.....	15.1	22.8	37.2	61.7	100.8	171.0	293.1	560.5	1,565.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)									
Non-Hispanic White, male									
2003.....	20.3	28.7	45.6	69.7	114.2	192.5	338.7	616.3	1,499.3
2002.....	20.8	30.1	45.5	72.3	115.5	197.0	343.3	627.9	1,512.9
2001.....	20.3	29.7	44.8	73.9	117.3	198.2	344.9	632.2	1,530.7
2000.....	20.5	29.2	47.9	76.8	122.7	203.1	345.6	640.2	1,606.4
1999.....	20.1	29.4	48.8	78.5	128.2	212.7	349.2	649.4	1,595.5
Non-Hispanic White, female									
2003.....	11.3	15.8	25.9	41.4	73.7	127.1	240.5	491.6	1,486.1
2002.....	10.5	15.6	27.0	43.1	76.7	134.2	248.8	499.2	1,498.8
2001.....	10.6	16.5	27.1	44.6	77.7	134.0	245.4	496.2	1,534.2
2000.....	9.9	16.2	26.1	45.6	79.8	132.3	248.8	502.0	1,547.2
1999.....	10.1	16.3	26.2	46.2	76.7	137.8	253.7	508.0	1,554.1
Non-Hispanic Black, both sexes									
2003.....	41.9	62.4	87.6	126.3	180.0	269.9	370.6	582.3	1,253.2
2002.....	44.8	62.1	89.0	123.6	172.3	260.2	391.1	601.9	1,285.5
2001.....	44.0	62.3	93.6	121.1	182.4	258.0	408.2	608.1	1,292.9
2000.....	44.3	63.6	87.8	125.0	185.1	270.0	412.7	597.8	1,319.4
1999.....	45.8	63.5	98.4	135.9	188.6	287.1	420.9	627.5	1,314.0
Non-Hispanic Black, male									
2003.....	52.2	80.9	116.6	159.6	229.1	322.9	442.0	673.2	1,239.1
2002.....	58.3	78.8	112.9	153.6	211.9	322.9	457.7	685.6	1,258.0
2001.....	54.7	75.2	120.1	154.0	216.3	322.3	501.0	753.1	1,362.9
2000.....	55.6	84.4	114.4	156.1	229.2	321.9	487.5	704.6	1,416.1
1999.....	57.3	83.5	128.8	175.5	241.5	355.6	491.1	756.3	1,466.6
Non-Hispanic Black, female									
2003.....	32.9	46.7	63.7	100.3	143.6	234.2	328.2	534.7	1,258.7
2002.....	33.1	48.1	69.3	100.2	142.9	218.2	351.3	558.4	1,296.0
2001.....	34.8	51.3	71.9	95.3	157.3	215.2	352.8	534.3	1,267.0
2000.....	34.5	45.9	66.0	100.5	152.6	235.5	367.6	544.0	1,283.4
1999.....	35.8	46.5	73.6	105.0	149.4	241.7	378.6	563.7	1,257.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and subacute endocarditis (I33)									
Hispanic, female									
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes									
2003.....	0.4	0.5	0.6	1.0	1.0	1.6	2.3	2.7	3.0
2002.....	0.4	0.5	0.5	0.8	1.1	1.5	2.2	2.7	2.7
2001.....	0.5	0.5	0.5	0.8	1.1	1.7	2.2	3.1	2.6
2000.....	0.5	0.5	0.7	0.7	1.1	1.3	2.2	2.8	2.9
1999.....	0.4	0.5	0.7	0.7	1.1	1.5	2.1	2.8	2.9
Non-Hispanic 4/, male									
2003.....	0.6	0.7	0.8	1.2	1.1	1.7	3.0	3.9	4.1
2002.....	0.5	0.7	0.6	0.9	1.5	1.7	2.6	3.7	3.6
2001.....	0.7	0.7	0.6	1.1	1.2	2.2	3.1	4.5	3.9
2000.....	0.6	0.7	0.8	0.9	1.3	1.5	3.2	4.1	4.2
1999.....	0.5	0.7	1.0	0.8	1.2	1.6	2.7	4.0	4.7
Non-Hispanic 4/, female									
2003.....	0.3	0.3	0.4	0.8	0.8	1.5	1.9	2.0	2.5
2002.....	0.3	0.3	0.4	0.8	0.8	1.4	1.9	2.1	2.3
2001.....	0.3	0.3	0.4	0.6	1.1	1.3	1.5	2.2	2.0
2000.....	0.3	0.4	0.5	0.6	0.9	1.1	1.6	2.0	2.3
1999.....	*	0.3	0.5	0.7	1.0	1.4	1.7	2.1	2.2
Non-Hispanic White, both sexes									
2003.....	0.3	0.4	0.5	0.8	0.9	1.4	2.3	2.8	3.2
2002.....	0.2	0.4	0.3	0.8	1.0	1.4	2.2	2.9	2.7
2001.....	0.4	0.4	0.4	0.7	0.9	1.5	2.0	3.1	2.7
2000.....	0.4	0.4	0.5	0.7	1.1	1.2	2.2	2.8	2.8
1999.....	0.3	0.3	0.5	0.7	1.0	1.4	2.1	2.8	3.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)									
Hispanic, female									
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes									
2003.....	0.3	0.3	0.4	0.4	0.8	0.9	1.0	1.5	1.7
2002.....	0.3	0.3	0.4	0.4	0.6	1.0	1.1	1.4	2.1
2001.....	0.3	0.3	0.4	0.5	0.5	0.8	1.2	1.3	1.9
2000.....	0.2	0.2	0.3	0.4	0.5	0.9	1.1	1.2	1.8
1999.....	0.3	0.3	0.5	0.6	0.7	0.9	1.2	1.4	2.1
Non-Hispanic 4/, male									
2003.....	0.4	0.3	0.4	0.4	0.9	1.1	1.1	1.6	1.6
2002.....	0.4	0.3	0.4	0.6	0.7	1.1	1.2	1.3	2.3
2001.....	0.3	0.3	0.4	0.6	0.7	0.8	1.3	1.2	1.8
2000.....	0.3	*	0.5	*	0.6	1.2	1.4	1.5	2.1
1999.....	0.3	0.3	0.7	0.6	0.8	0.9	1.3	1.5	2.6
Non-Hispanic 4/, female									
2003.....	0.2	*	0.3	0.3	0.7	0.7	0.8	1.5	1.7
2002.....	*	0.2	0.4	*	0.5	0.9	1.1	1.5	2.0
2001.....	0.3	0.3	0.3	*	*	0.9	1.1	1.4	1.9
2000.....	*	0.3	*	*	0.5	0.7	0.9	1.1	1.7
1999.....	0.2	0.3	0.4	0.6	0.7	0.8	1.1	1.4	1.9
Non-Hispanic White, both sexes									
2003.....	0.3	0.2	0.4	0.4	0.8	0.9	1.0	1.5	1.7
2002.....	0.2	0.2	0.3	0.4	0.6	1.0	1.0	1.4	2.1
2001.....	0.3	0.2	0.3	0.5	0.4	0.9	1.2	1.3	1.9
2000.....	0.2	0.2	0.3	0.3	0.5	0.9	1.0	1.3	1.8
1999.....	0.3	0.3	0.5	0.6	0.7	0.8	1.2	1.4	2.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)									
All origins 3/, both sexes									
2003.....	1.6	3.3	6.4	12.1	23.6	44.7	89.9	192.7	641.1
2002.....	1.7	3.4	6.0	12.4	22.9	45.7	89.0	194.0	645.0
2001.....	1.7	3.3	6.5	13.0	23.8	45.6	92.6	199.2	674.6
2000.....	1.4	3.0	6.1	12.8	24.0	46.5	93.5	199.9	684.9
1999.....	1.5	2.9	6.5	11.7	23.5	48.3	94.3	201.6	681.9
All origins 3/, male									
2003.....	2.1	3.9	7.9	14.7	28.3	53.1	105.3	213.9	629.8
2002.....	2.0	4.4	7.2	14.8	27.0	53.2	103.9	218.0	621.4
2001.....	2.2	4.0	8.1	15.0	28.0	54.2	107.9	226.6	646.4
2000.....	1.9	3.5	7.4	15.3	28.2	55.6	107.9	229.5	672.2
1999.....	1.9	3.6	8.0	14.2	28.0	58.9	108.4	224.9	667.0
All origins 3/, female									
2003.....	1.2	2.7	4.9	9.7	19.6	37.8	78.9	179.8	646.1
2002.....	1.3	2.4	4.9	10.1	19.3	39.8	78.4	179.3	655.3
2001.....	1.2	2.6	5.1	11.3	20.2	38.7	81.9	182.7	686.4
2000.....	0.9	2.5	5.0	10.5	20.3	39.3	83.5	182.4	690.0
1999.....	1.1	2.1	5.1	9.5	19.6	39.9	84.5	188.0	687.9
Hispanic, both sexes									
2003.....	1.0	1.6	4.1	7.8	16.8	29.7	55.7	111.9	330.1
2002.....	*	1.8	3.1	6.7	14.2	27.1	53.3	111.8	316.0
2001.....	*	1.7	2.9	9.1	14.3	24.4	50.4	112.2	349.5
2000.....	*	*	4.0	8.3	12.2	27.2	52.3	108.1	369.6
1999.....	*	*	3.5	7.6	12.3	28.6	53.9	109.3	377.4
Hispanic, male									
2003.....	*	*	5.3	11.3	18.2	36.3	68.7	134.1	312.5
2002.....	*	*	4.0	8.2	17.3	30.7	64.0	124.7	296.2
2001.....	*	*	*	9.5	19.5	24.7	47.1	129.9	333.3
2000.....	*	*	5.3	10.9	16.4	31.6	52.4	117.8	380.9
1999.....	*	*	*	8.3	15.5	36.7	67.9	113.2	358.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)									
Hispanic, female									
2003.....	*	*	*	4.7	15.7	24.6	46.6	97.4	339.1
2002.....	*	*	*	5.4	11.6	24.3	45.7	103.7	325.8
2001.....	*	*	*	8.7	10.0	24.1	52.8	101.3	357.4
2000.....	*	*	*	6.0	8.8	23.9	52.3	102.1	364.0
1999.....	*	*	*	6.9	9.7	22.6	44.0	106.9	386.8
Non-Hispanic 4/, both sexes									
2003.....	1.7	3.4	6.6	12.4	24.1	45.5	91.7	196.1	653.2
2002.....	1.8	3.5	6.3	12.8	23.5	46.7	90.7	197.0	656.8
2001.....	1.8	3.4	6.8	13.3	24.4	46.8	94.5	202.2	685.6
2000.....	1.5	3.2	6.3	13.1	24.7	47.5	95.2	202.8	694.8
1999.....	1.6	3.0	6.7	11.9	24.1	49.2	95.8	204.5	691.0
Non-Hispanic 4/, male									
2003.....	2.2	4.2	8.1	14.9	29.0	54.0	107.2	217.1	643.2
2002.....	2.2	4.6	7.5	15.3	27.7	54.4	105.9	221.4	634.7
2001.....	2.3	4.2	8.4	15.3	28.5	55.8	110.7	230.4	658.0
2000.....	2.1	3.8	7.5	15.5	28.9	56.8	110.3	233.4	683.3
1999.....	2.0	3.8	8.2	14.5	28.7	60.0	109.9	228.4	677.1
Non-Hispanic 4/, female									
2003.....	1.3	2.7	5.1	10.1	19.8	38.6	80.6	183.3	657.6
2002.....	1.4	2.5	5.1	10.5	19.8	40.6	79.9	182.2	666.4
2001.....	1.3	2.7	5.2	11.5	20.8	39.6	83.2	185.4	697.0
2000.....	0.9	2.6	5.2	10.9	21.1	40.1	84.7	184.8	699.5
1999.....	1.2	2.2	5.3	9.6	20.2	40.8	86.1	190.5	696.5
Non-Hispanic White, both sexes									
2003.....	1.3	2.7	5.7	10.9	22.7	44.0	91.3	198.9	673.5
2002.....	1.2	2.8	5.5	11.7	22.2	45.1	89.8	198.8	677.3
2001.....	1.4	2.8	5.9	12.2	22.9	45.5	93.9	204.3	706.8
2000.....	1.1	2.7	5.5	12.0	23.3	45.8	94.5	204.9	714.3
1999.....	1.2	2.3	5.7	11.1	22.6	47.7	94.0	205.5	709.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)									
Non-Hispanic White, male									
2003.....	1.7	3.2	7.1	13.1	27.5	52.9	107.8	220.6	665.3
2002.....	1.6	3.7	6.6	14.0	26.1	52.5	105.5	224.9	656.2
2001.....	1.9	3.5	7.5	14.4	26.9	54.4	110.5	232.1	679.1
2000.....	1.7	3.2	6.9	14.2	27.8	55.5	110.1	235.8	701.6
1999.....	1.5	3.0	7.3	13.5	26.8	59.1	108.8	229.9	690.8
Non-Hispanic White, female									
2003.....	0.9	2.1	4.4	8.8	18.4	36.6	79.3	185.5	677.2
2002.....	0.9	1.9	4.4	9.6	18.9	39.0	78.6	182.8	686.4
2001.....	0.8	2.1	4.3	10.2	19.4	38.2	82.2	187.4	718.4
2000.....	0.6	2.1	4.2	9.9	19.3	38.1	83.6	186.6	719.4
1999.....	0.8	1.7	4.2	8.8	18.9	38.7	83.6	191.1	716.4
Non-Hispanic Black, both sexes									
2003.....	4.9	9.7	15.6	28.2	42.2	71.8	116.8	206.1	538.7
2002.....	5.6	9.4	14.1	25.3	39.3	72.8	119.4	217.6	534.5
2001.....	4.7	8.9	16.0	25.1	42.1	72.3	122.2	216.6	551.2
2000.....	4.3	7.9	14.4	24.8	42.0	73.4	121.4	211.8	564.8
1999.....	4.9	8.6	16.6	22.4	43.0	74.9	131.6	218.4	557.4
Non-Hispanic Black, male									
2003.....	5.7	12.0	20.0	36.1	52.4	80.2	127.1	228.7	521.0
2002.....	6.7	12.3	16.6	32.2	48.6	88.9	138.0	240.2	489.0
2001.....	5.2	10.7	19.3	26.8	50.5	87.1	141.7	260.8	547.5
2000.....	5.7	9.3	15.2	30.0	46.4	84.1	137.3	246.3	589.4
1999.....	6.5	11.3	19.3	27.6	52.6	84.5	146.9	241.8	609.7
Non-Hispanic Black, female									
2003.....	4.1	7.8	12.0	22.0	34.7	66.1	110.7	194.3	545.6
2002.....	4.7	7.0	12.1	19.9	32.4	61.9	108.2	205.8	551.8
2001.....	4.2	7.3	13.3	23.8	35.8	62.4	110.6	194.1	552.6
2000.....	3.1	6.6	13.7	20.6	38.7	66.2	111.8	194.4	555.6
1999.....	3.5	6.3	14.4	18.3	35.8	68.5	122.3	206.8	537.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)									
All origins 3/, both sexes									
2003.....	15.5	21.3	31.7	46.3	71.8	113.2	186.0	328.5	781.5
2002.....	15.7	21.7	32.5	46.9	73.3	116.3	194.4	338.2	793.7
2001.....	15.4	21.9	32.3	47.8	75.3	116.6	191.2	333.6	796.5
2000.....	15.5	22.1	33.1	50.4	78.0	118.4	193.4	338.9	820.0
1999.....	15.7	22.3	34.1	53.5	79.9	124.4	197.7	347.6	826.1
All origins 3/, male									
2003.....	19.7	27.5	40.8	58.6	88.8	139.6	224.8	380.7	795.9
2002.....	20.6	28.1	41.1	59.4	89.7	143.8	233.1	389.4	821.3
2001.....	19.9	27.7	40.3	61.1	91.7	143.7	232.4	390.0	821.3
2000.....	20.2	28.9	43.4	64.2	96.8	147.6	232.6	391.5	870.7
1999.....	20.2	28.7	44.5	68.7	103.5	155.5	238.5	411.1	874.7
All origins 3/, female									
2003.....	11.5	15.4	23.1	35.1	57.0	92.0	158.4	296.6	775.1
2002.....	11.0	15.6	24.4	35.6	59.2	94.3	167.1	307.1	781.7
2001.....	11.1	16.3	24.8	35.7	61.2	95.0	162.4	299.7	786.1
2000.....	11.0	15.6	23.4	37.8	61.9	95.4	166.0	307.9	799.3
1999.....	11.2	16.2	24.3	39.8	59.7	100.1	169.4	310.5	806.5
Hispanic, both sexes									
2003.....	8.5	10.7	18.9	29.1	41.8	67.1	117.0	180.8	418.2
2002.....	7.8	11.9	17.2	23.2	41.7	62.4	111.4	192.1	419.6
2001.....	7.5	11.2	17.8	26.6	46.5	66.3	114.7	188.7	444.6
2000.....	7.8	13.4	18.4	26.5	44.2	68.3	123.7	203.9	457.2
1999.....	7.9	13.1	19.3	31.2	47.7	69.3	114.9	219.1	447.6
Hispanic, male									
2003.....	11.7	14.0	22.9	35.9	52.6	84.1	131.9	205.8	437.2
2002.....	9.8	15.4	20.9	28.7	50.5	88.6	136.3	225.3	455.1
2001.....	10.4	15.5	22.0	34.5	55.3	83.7	139.1	229.8	438.1
2000.....	10.0	18.7	25.0	33.4	52.6	89.9	140.3	237.0	505.9
1999.....	10.5	18.1	23.6	42.6	63.0	92.2	139.7	259.4	526.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)									
Hispanic, female									
2003.....	5.3	7.5	15.2	23.2	32.8	54.1	106.5	164.4	408.6
2002.....	5.8	8.5	13.8	18.3	34.6	42.3	93.7	170.8	402.0
2001.....	4.7	7.2	13.9	19.7	39.3	53.0	97.3	163.3	447.7
2000.....	5.5	8.5	12.5	20.6	37.4	51.9	111.9	183.7	433.3
1999.....	5.2	8.4	15.4	21.3	35.3	52.1	97.5	195.1	409.7
Non-Hispanic 4/, both sexes									
2003.....	16.2	22.2	32.7	47.5	73.9	116.0	189.5	334.8	795.8
2002.....	16.6	22.6	33.6	48.7	75.4	119.4	198.4	343.8	807.6
2001.....	16.2	22.7	33.4	49.2	77.1	119.3	194.6	338.8	807.8
2000.....	16.2	22.7	34.0	52.0	80.1	120.9	196.2	343.5	831.4
1999.....	16.3	23.0	35.0	54.9	81.7	126.9	200.9	351.5	837.9
Non-Hispanic 4/, male									
2003.....	20.5	28.6	42.1	60.0	91.3	142.9	229.5	388.3	811.6
2002.....	21.8	29.2	42.6	61.6	92.2	146.8	237.6	396.1	836.7
2001.....	20.8	28.6	41.7	62.8	93.9	146.8	236.4	396.2	835.3
2000.....	21.1	29.7	44.5	66.2	99.4	150.3	236.4	396.9	883.9
1999.....	21.1	29.5	45.8	70.3	105.8	158.2	242.1	415.5	886.8
Non-Hispanic 4/, female									
2003.....	12.1	16.1	23.7	36.0	58.7	94.4	160.9	302.1	788.8
2002.....	11.5	16.2	25.1	36.9	60.9	97.4	170.8	312.1	795.0
2001.....	11.7	17.1	25.6	36.9	62.6	97.4	165.3	304.4	796.3
2000.....	11.4	16.1	24.2	39.0	63.5	97.7	168.2	312.1	810.2
1999.....	11.7	16.9	24.9	41.0	61.1	102.4	172.3	314.0	818.4
Non-Hispanic White, both sexes									
2003.....	13.9	18.9	29.0	42.9	68.3	110.3	187.2	335.9	811.6
2002.....	13.9	19.4	29.9	44.2	71.1	114.9	195.2	345.0	820.9
2001.....	13.4	19.6	29.1	45.3	72.0	114.9	189.5	338.8	821.8
2000.....	13.5	19.3	30.4	47.6	75.0	115.9	191.0	344.5	845.3
1999.....	13.4	19.9	30.5	49.4	76.5	121.0	195.8	350.7	851.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)									
Non-Hispanic White, male									
2003.....	17.9	24.6	37.5	55.0	84.6	136.9	226.8	390.1	827.7
2002.....	18.5	25.6	38.1	56.8	87.4	141.7	234.0	397.9	850.7
2001.....	17.6	25.4	36.4	58.0	88.9	141.1	230.1	394.4	845.6
2000.....	18.2	25.2	39.8	61.4	93.3	144.9	231.0	398.7	898.3
1999.....	17.9	25.7	40.1	63.6	99.5	151.1	236.2	413.9	897.0
Non-Hispanic White, female									
2003.....	10.0	13.4	20.9	31.7	53.8	88.5	158.6	302.7	804.5
2002.....	9.3	13.3	21.9	32.6	56.6	93.1	167.4	312.7	808.1
2001.....	9.2	13.9	22.1	33.5	57.0	93.6	160.7	305.2	811.9
2000.....	8.9	13.6	21.4	34.9	58.9	92.6	162.9	312.3	824.0
1999.....	9.0	14.2	21.3	36.2	56.3	97.1	167.5	313.5	833.7
Non-Hispanic Black, both sexes									
2003.....	35.3	50.8	69.8	95.4	135.3	194.1	249.6	371.5	711.8
2002.....	37.5	50.6	72.2	96.2	129.8	183.6	266.8	381.5	747.1
2001.....	37.7	51.0	75.4	93.3	137.0	182.1	280.5	386.8	737.9
2000.....	38.1	53.9	70.9	98.3	140.3	193.5	287.0	382.5	748.9
1999.....	39.3	52.9	78.7	111.2	142.1	208.0	285.7	405.3	752.7
Non-Hispanic Black, male									
2003.....	44.1	66.1	93.3	121.3	174.4	238.3	310.2	440.0	718.1
2002.....	49.1	64.1	92.7	119.1	159.5	230.2	316.4	440.7	763.6
2001.....	46.8	61.6	98.2	123.3	161.0	229.6	352.7	484.9	813.1
2000.....	47.4	72.9	96.8	124.0	177.8	235.1	344.9	454.0	821.9
1999.....	48.9	68.9	105.2	145.4	185.7	266.2	342.7	509.2	855.7
Non-Hispanic Black, female									
2003.....	27.7	37.8	50.4	75.3	106.2	164.3	213.6	335.7	709.4
2002.....	27.5	39.3	55.3	78.3	107.8	152.3	237.2	350.6	740.8
2001.....	29.7	42.1	56.7	69.7	119.3	150.5	237.4	336.9	710.1
2000.....	30.0	37.6	49.6	78.0	112.7	165.8	252.1	346.5	721.7
1999.....	31.1	39.3	57.1	84.3	109.9	169.3	251.3	353.9	714.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All origins 3/, both sexes

2003.....	1.8	3.3	5.0	8.0	13.1	21.3	38.4	69.9	188.9
2002.....	1.8	2.9	4.5	7.3	11.8	20.7	34.9	66.7	180.4
2001.....	1.9	3.0	4.4	7.5	12.2	19.0	35.5	65.2	171.9
2000.....	1.8	3.0	4.4	7.8	11.5	19.1	33.8	63.0	162.9
1999.....	1.7	2.8	4.1	7.3	12.0	18.6	32.7	59.9	152.1

All origins 3/, male

2003.....	2.5	4.2	6.0	9.2	14.9	22.5	37.5	66.1	153.5
2002.....	2.3	3.5	5.5	8.2	12.7	22.2	35.9	64.8	153.4
2001.....	2.3	3.7	5.2	8.1	13.7	20.0	36.0	61.3	150.8
2000.....	2.3	3.8	5.4	9.1	11.5	20.6	34.5	62.9	142.4
1999.....	2.0	3.5	5.0	8.2	13.4	20.1	33.1	59.4	132.4

All origins 3/, female

2003.....	1.2	2.4	4.0	6.9	11.4	20.4	39.1	72.2	204.6
2002.....	1.3	2.3	3.7	6.5	11.0	19.5	34.2	67.9	192.1
2001.....	1.4	2.3	3.8	6.9	11.0	18.2	35.2	67.5	180.8
2000.....	1.3	2.2	3.5	6.7	11.5	17.9	33.4	63.0	171.3
1999.....	1.3	2.1	3.3	6.5	10.8	17.5	32.4	60.2	160.0

Hispanic, both sexes

2003.....	1.7	2.3	4.3	7.9	12.6	21.9	34.3	61.1	171.0
2002.....	*	1.4	4.4	8.7	10.0	22.3	34.7	68.2	148.6
2001.....	1.5	2.2	2.9	6.6	11.4	22.8	38.2	67.5	160.1
2000.....	*	2.7	3.5	10.0	10.8	20.3	36.1	63.5	148.0
1999.....	1.7	2.7	3.2	7.3	12.3	19.0	35.5	71.7	129.8

Hispanic, male

2003.....	2.4	3.3	4.5	11.6	11.0	21.4	33.4	63.4	133.9
2002.....	*	*	5.5	9.2	8.8	23.7	39.0	59.5	132.2
2001.....	2.2	*	*	5.7	10.6	24.7	39.6	72.7	129.5
2000.....	*	4.2	4.4	13.0	9.7	22.4	42.1	73.6	141.1
1999.....	*	*	*	8.6	16.2	21.4	43.7	67.6	131.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Hispanic, female

2003.....	*	*	4.1	4.7	13.9	22.3	34.9	59.5	189.8
2002.....	*	*	3.4	8.3	11.1	21.3	31.5	73.7	156.7
2001.....	*	*	*	7.3	12.1	21.3	37.3	64.3	175.0
2000.....	*	*	*	7.4	11.8	18.8	31.9	57.3	151.3
1999.....	*	*	*	6.2	9.1	17.2	29.7	74.1	128.9

Non-Hispanic 4/, both sexes

2003.....	1.8	3.4	5.0	8.0	13.0	21.1	38.6	70.1	189.4
2002.....	1.9	3.0	4.5	7.2	11.8	20.5	34.9	66.5	181.1
2001.....	1.9	3.0	4.5	7.5	12.3	18.7	35.2	65.0	172.0
2000.....	1.8	3.0	4.5	7.6	11.5	18.9	33.5	62.9	162.9
1999.....	1.7	2.8	4.2	7.3	11.9	18.6	32.5	59.1	152.3

Non-Hispanic 4/, male

2003.....	2.5	4.3	6.1	9.0	15.1	22.4	37.7	66.2	154.4
2002.....	2.4	3.7	5.4	8.1	12.8	22.0	35.6	64.8	153.9
2001.....	2.3	3.8	5.3	8.3	13.8	19.6	35.6	60.6	151.1
2000.....	2.3	3.8	5.4	8.7	11.5	20.4	33.9	62.4	142.0
1999.....	2.0	3.5	5.1	8.1	13.2	20.0	32.6	58.5	131.9

Non-Hispanic 4/, female

2003.....	1.2	2.5	4.0	7.1	11.2	20.1	39.2	72.6	204.7
2002.....	1.4	2.3	3.7	6.3	10.9	19.3	34.3	67.5	192.8
2001.....	1.4	2.3	3.8	6.8	10.9	18.0	35.0	67.6	180.7
2000.....	1.3	2.2	3.5	6.6	11.5	17.8	33.3	63.2	171.3
1999.....	1.3	2.1	3.2	6.5	10.9	17.4	32.5	59.5	160.5

Non-Hispanic White, both sexes

2003.....	1.1	2.2	3.1	5.6	9.5	16.1	32.9	64.1	181.0
2002.....	1.1	2.0	2.9	5.1	8.9	16.3	29.3	59.7	170.6
2001.....	1.0	1.8	2.9	5.2	8.9	14.7	30.3	58.1	162.8
2000.....	1.0	1.7	2.9	5.0	8.4	14.9	28.3	57.0	152.7
1999.....	1.0	1.5	2.7	4.8	8.8	14.9	27.1	52.2	141.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Non-Hispanic White, male

2003.....	1.5	2.8	4.0	6.1	11.6	17.6	32.7	60.0	145.4
2002.....	1.4	2.6	3.6	6.1	9.8	17.7	30.2	57.3	145.4
2001.....	1.3	2.2	3.5	5.9	10.2	15.6	30.8	53.4	142.1
2000.....	1.3	2.3	3.7	5.9	8.7	16.3	29.1	56.5	131.8
1999.....	1.3	1.9	3.6	5.6	10.0	16.2	27.5	51.2	119.8

Non-Hispanic White, female

2003.....	0.7	1.7	2.3	5.0	7.6	14.9	33.0	66.6	196.6
2002.....	0.7	1.4	2.2	4.2	8.0	15.1	28.7	61.1	181.4
2001.....	0.7	1.3	2.3	4.5	7.8	14.0	29.9	60.9	171.5
2000.....	0.6	1.1	2.2	4.2	8.0	13.8	27.8	57.2	161.1
1999.....	0.7	1.1	1.8	4.1	7.8	13.7	26.9	52.8	150.5

Non-Hispanic Black, both sexes

2003.....	7.2	12.3	21.1	29.6	43.7	69.6	103.4	147.2	304.3
2002.....	7.6	11.1	18.8	24.8	36.9	62.6	98.1	153.2	312.4
2001.....	7.9	12.9	18.3	27.6	41.2	57.4	90.7	150.1	291.5
2000.....	7.3	13.1	18.3	29.5	41.0	59.9	93.8	132.8	285.9
1999.....	6.5	13.2	17.3	27.3	38.7	58.3	96.0	140.3	272.2

Non-Hispanic Black, male

2003.....	9.9	16.8	25.0	36.9	50.6	75.5	103.3	149.4	280.2
2002.....	9.5	13.4	22.2	26.3	40.7	68.4	107.0	166.9	276.8
2001.....	10.2	16.7	21.9	30.9	49.5	63.4	95.8	161.1	277.2
2000.....	9.0	15.9	22.9	35.3	40.8	65.3	93.9	139.9	268.9
1999.....	8.1	17.1	19.9	30.9	43.8	63.4	98.6	156.8	268.6

Non-Hispanic Black, female

2003.....	4.8	8.5	17.8	24.0	38.5	65.7	103.5	146.1	313.6
2002.....	6.0	9.1	16.0	23.7	34.2	58.7	92.8	146.1	326.1
2001.....	5.9	9.6	15.4	25.0	35.0	53.4	87.7	144.5	296.9
2000.....	5.9	10.6	14.5	25.0	41.1	56.3	93.8	129.2	292.2
1999.....	5.1	9.8	15.2	24.4	34.8	54.9	94.4	132.1	273.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cerebrovascular diseases (I60-I69)

All origins 3/, both sexes

2003.....	11.8	18.6	27.8	45.9	79.7	150.7	294.1	571.1	1,370.1
2002.....	12.3	18.3	28.6	48.3	83.8	160.7	308.8	601.7	1,445.9
2001.....	12.1	18.5	28.8	49.6	85.7	164.3	320.2	622.4	1,500.2
2000.....	13.1	19.3	30.8	53.8	91.0	169.1	332.8	654.0	1,589.2
1999.....	12.5	18.4	31.7	51.4	91.1	173.5	340.6	664.0	1,614.8

All origins 3/, male

2003.....	12.9	21.1	31.3	53.5	91.4	170.8	322.5	595.6	1,236.0
2002.....	13.4	20.4	32.7	55.8	97.0	178.7	334.8	615.6	1,317.9
2001.....	13.2	20.5	32.8	57.2	98.5	187.9	349.0	654.6	1,380.2
2000.....	14.2	21.3	35.2	62.3	102.6	192.9	367.6	695.1	1,484.3
1999.....	13.3	20.7	35.7	60.0	102.8	197.7	376.9	706.8	1,512.8

All origins 3/, female

2003.....	10.8	16.3	24.6	38.9	69.5	134.5	273.9	556.1	1,429.4
2002.....	11.2	16.3	24.8	41.4	72.4	146.2	290.4	593.3	1,501.5
2001.....	10.9	16.5	25.1	42.7	74.7	145.6	300.0	603.2	1,550.4
2000.....	12.0	17.3	26.6	46.0	81.0	150.3	308.5	629.7	1,632.0
1999.....	11.7	16.1	28.0	43.7	81.1	154.5	315.3	639.0	1,656.0

Hispanic, both sexes

2003.....	11.6	19.3	29.1	43.9	77.5	124.8	252.4	421.4	880.5
2002.....	13.0	18.4	28.2	46.2	70.3	143.8	240.2	453.2	898.0
2001.....	12.4	20.3	28.6	44.5	74.7	145.5	271.5	463.4	1,042.9
2000.....	13.8	17.0	28.2	54.6	83.1	151.1	262.9	491.8	1,082.9
1999.....	13.6	18.3	31.4	45.6	90.5	150.2	265.7	498.7	1,077.6

Hispanic, male

2003.....	14.2	23.3	34.7	56.1	91.0	143.9	275.9	441.8	837.4
2002.....	16.2	22.0	36.0	57.7	86.7	173.8	269.4	462.1	856.7
2001.....	14.6	24.1	34.2	56.2	95.3	168.4	311.0	529.8	1,005.6
2000.....	16.2	20.5	37.7	63.3	99.6	183.8	310.8	556.5	1,029.9
1999.....	16.2	23.8	36.3	58.7	105.5	187.4	330.0	526.6	1,136.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)									
Hispanic, female									
2003.....	8.9	15.4	23.8	33.2	66.4	110.1	235.8	408.1	902.3
2002.....	9.7	14.9	21.1	36.2	57.0	120.7	219.4	447.5	918.5
2001.....	10.2	16.7	23.5	34.3	58.0	128.1	243.5	422.2	1,061.2
2000.....	11.4	13.8	19.6	47.1	69.8	126.3	229.0	452.4	1,108.9
1999.....	11.0	13.1	26.9	34.4	78.4	122.2	220.2	482.0	1,049.2
Non-Hispanic 4/, both sexes									
2003.....	11.8	18.5	27.6	45.9	79.7	152.0	296.0	577.2	1,388.4
2002.....	12.2	18.3	28.6	48.3	84.6	161.4	311.7	606.9	1,465.0
2001.....	12.0	18.2	28.8	49.8	86.3	165.1	321.7	627.7	1,514.0
2000.....	12.9	19.4	30.9	53.5	91.3	169.6	335.1	658.3	1,604.5
1999.....	12.3	18.3	31.6	51.6	90.9	174.2	343.0	668.9	1,630.4
Non-Hispanic 4/, male									
2003.....	12.7	20.8	30.9	53.0	91.2	172.1	324.6	602.2	1,252.9
2002.....	13.0	20.2	32.3	55.4	97.5	178.7	337.3	621.1	1,336.0
2001.....	13.1	20.2	32.5	56.9	98.5	188.7	350.0	658.1	1,392.9
2000.....	13.9	21.3	34.9	62.0	102.6	192.7	369.1	698.8	1,499.9
1999.....	12.9	20.4	35.5	59.7	102.3	197.6	377.8	712.2	1,524.8
Non-Hispanic 4/, female									
2003.....	10.9	16.3	24.5	39.4	69.6	135.8	275.7	562.0	1,448.0
2002.....	11.3	16.4	25.1	41.7	73.4	147.5	293.5	598.3	1,520.7
2001.....	10.9	16.4	25.2	43.2	75.8	146.3	302.0	609.6	1,564.4
2000.....	12.0	17.6	27.1	45.7	81.7	151.4	311.4	634.5	1,646.8
1999.....	11.7	16.3	28.0	44.2	81.1	155.9	318.9	643.6	1,672.7
Non-Hispanic White, both sexes									
2003.....	8.6	13.4	21.8	37.6	69.9	140.7	284.2	571.5	1,401.1
2002.....	9.0	13.4	23.3	40.4	74.2	149.9	300.9	600.4	1,479.2
2001.....	8.8	13.0	23.4	41.1	74.9	153.2	310.4	621.9	1,525.0
2000.....	9.2	15.1	25.0	45.6	81.4	158.8	323.1	651.1	1,616.8
1999.....	8.8	14.0	25.6	43.0	81.8	161.2	331.1	662.9	1,648.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)									
Non-Hispanic White, male									
2003.....	9.4	15.0	23.7	43.3	80.7	158.7	311.1	596.6	1,264.2
2002.....	9.8	14.6	25.9	45.8	84.7	166.1	325.1	615.2	1,345.9
2001.....	9.6	14.4	26.0	46.9	85.6	174.7	337.7	651.5	1,399.0
2000.....	10.0	16.4	27.5	52.8	90.8	179.5	355.4	689.2	1,505.9
1999.....	9.0	15.3	28.7	49.6	91.7	182.4	363.1	705.6	1,535.1
Non-Hispanic White, female									
2003.....	7.7	11.9	20.1	32.4	60.4	126.0	264.8	556.1	1,461.3
2002.....	8.1	12.2	20.7	35.3	64.9	136.6	283.5	591.4	1,536.7
2001.....	8.0	11.6	20.8	35.7	65.4	135.7	291.1	604.0	1,577.4
2000.....	8.3	13.8	22.5	38.8	73.2	142.2	300.4	628.4	1,661.6
1999.....	8.6	12.7	22.6	37.0	73.2	144.3	308.7	637.7	1,693.0
Non-Hispanic Black, both sexes									
2003.....	32.2	53.9	73.9	114.5	159.8	261.6	440.4	686.0	1,366.0
2002.....	32.7	52.4	72.4	112.1	172.6	276.8	452.2	721.0	1,398.0
2001.....	33.1	56.2	71.9	119.3	181.5	281.4	465.4	725.7	1,465.5
2000.....	38.0	52.4	78.6	115.6	178.0	284.5	479.0	775.9	1,545.8
1999.....	35.1	51.3	79.6	119.8	172.9	312.5	493.4	762.8	1,501.5
Non-Hispanic Black, male									
2003.....	35.0	64.3	92.5	143.7	187.0	314.8	511.2	737.3	1,200.7
2002.....	34.8	63.0	90.8	141.7	217.1	334.6	508.9	738.6	1,266.8
2001.....	37.3	65.4	89.2	145.9	222.5	340.9	530.4	776.0	1,359.4
2000.....	41.5	61.7	99.4	138.8	216.4	343.4	548.2	872.8	1,479.5
1999.....	39.8	62.3	94.3	151.1	207.7	372.2	565.0	824.5	1,455.9
Non-Hispanic Black, female									
2003.....	29.7	45.2	58.6	91.7	139.7	225.8	398.4	659.1	1,430.0
2002.....	30.9	43.4	57.4	88.9	139.7	238.1	418.3	711.8	1,448.1
2001.....	29.6	48.4	57.8	98.4	151.2	241.8	426.6	700.1	1,504.8
2000.....	34.9	44.5	61.7	97.3	149.6	245.5	437.3	727.1	1,570.4
1999.....	31.1	42.0	67.7	95.2	147.1	272.9	450.2	732.3	1,518.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)									
All origins 3/, both sexes									
2003.....	0.3	0.6	1.1	2.4	4.5	9.0	19.9	43.5	154.6
2002.....	0.3	0.6	1.4	2.4	5.2	9.3	20.5	47.8	168.1
2001.....	0.3	0.7	1.4	2.5	4.9	9.8	21.9	49.2	179.1
2000.....	0.4	0.6	1.3	2.6	5.2	10.5	22.7	52.4	188.8
1999.....	0.3	0.8	1.3	2.9	5.1	11.2	24.1	53.6	202.0
All origins 3/, male									
2003.....	0.5	0.8	1.5	3.2	5.4	10.6	23.9	48.3	138.6
2002.....	0.4	0.8	1.7	3.2	6.4	11.5	24.4	52.5	156.6
2001.....	0.4	0.8	1.7	3.2	6.2	12.1	25.0	53.2	167.2
2000.....	0.5	0.8	1.7	3.6	6.6	13.1	27.5	59.8	176.5
1999.....	0.4	1.0	1.8	3.8	6.3	13.3	28.1	60.0	189.0
All origins 3/, female									
2003.....	*	0.4	0.7	1.7	3.7	7.8	17.0	40.6	161.7
2002.....	*	0.5	1.0	1.7	4.1	7.5	17.8	45.0	173.1
2001.....	0.2	0.5	1.1	1.8	3.9	8.1	19.7	46.8	184.1
2000.....	0.2	0.4	0.9	1.7	4.0	8.4	19.4	48.1	193.8
1999.....	*	0.5	0.9	2.2	4.0	9.5	21.4	49.8	207.2
Hispanic, both sexes									
2003.....	*	*	*	*	3.1	6.5	11.9	30.3	86.0
2002.....	*	*	*	*	3.7	5.0	15.5	34.8	108.6
2001.....	*	*	*	*	*	6.3	17.0	33.8	130.7
2000.....	*	*	*	*	4.0	8.2	16.2	39.5	132.7
1999.....	*	*	*	*	*	8.5	19.4	41.1	129.1
Hispanic, male									
2003.....	*	*	*	*	*	9.4	18.8	33.3	95.4
2002.....	*	*	*	*	*	*	18.6	41.2	102.1
2001.....	*	*	*	*	*	*	21.1	39.0	127.6
2000.....	*	*	*	*	*	12.6	25.1	45.6	104.8
1999.....	*	*	*	*	*	11.2	23.4	47.2	151.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerosis (I70)

Hispanic, female

2003.....	*	*	*	*	*	*	*	28.4	81.3
2002.....	*	*	*	*	*	*	13.3	30.7	111.9
2001.....	*	*	*	*	*	*	14.0	30.6	132.2
2000.....	*	*	*	*	*	*	*	35.8	146.4
1999.....	*	*	*	*	*	*	16.5	37.5	118.5

Non-Hispanic 4/, both sexes

2003.....	0.4	0.6	1.1	2.5	4.6	9.2	20.2	43.9	157.1
2002.....	0.3	0.7	1.4	2.5	5.3	9.5	20.7	48.3	170.2
2001.....	0.3	0.6	1.4	2.6	5.1	10.0	22.0	49.6	180.5
2000.....	0.4	0.6	1.3	2.7	5.2	10.6	23.0	52.8	190.3
1999.....	0.3	0.8	1.3	3.0	5.2	11.3	24.3	53.8	203.9

Non-Hispanic 4/, male

2003.....	0.6	0.9	1.5	3.3	5.5	10.6	24.1	48.8	140.3
2002.....	0.4	0.9	1.8	3.4	6.5	11.8	24.6	52.9	158.7
2001.....	0.5	0.8	1.7	3.4	6.4	12.4	25.0	53.6	168.2
2000.....	0.5	0.8	1.8	3.7	6.7	13.1	27.5	60.2	178.7
1999.....	0.4	1.0	1.8	3.8	6.4	13.4	28.2	60.2	189.9

Non-Hispanic 4/, female

2003.....	*	0.4	0.7	1.8	3.7	8.0	17.5	40.9	164.5
2002.....	*	0.5	1.1	1.7	4.2	7.7	17.9	45.4	175.1
2001.....	0.2	0.5	1.1	1.9	4.0	8.2	19.9	47.3	185.5
2000.....	0.2	0.4	0.9	1.7	4.0	8.6	19.8	48.5	195.1
1999.....	*	0.5	0.9	2.2	4.1	9.7	21.5	50.1	209.6

Non-Hispanic White, both sexes

2003.....	0.4	0.6	1.0	2.5	4.4	9.1	20.6	44.5	162.2
2002.....	0.3	0.6	1.4	2.3	5.4	9.4	20.5	48.8	177.0
2001.....	0.3	0.6	1.4	2.4	5.1	10.0	22.4	50.8	187.4
2000.....	0.3	0.5	1.3	2.5	5.3	10.5	23.3	53.4	196.8
1999.....	0.3	0.7	1.2	2.9	5.1	11.4	24.3	54.9	209.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)									
Non-Hispanic White, male									
2003.....	0.5	0.8	1.4	3.2	5.4	10.3	24.0	49.0	145.2
2002.....	0.4	0.7	1.7	3.1	6.7	11.6	24.2	53.5	165.7
2001.....	0.4	0.7	1.7	3.2	6.4	12.4	25.1	54.4	175.2
2000.....	0.4	0.7	1.8	3.6	6.6	13.3	28.3	60.8	183.6
1999.....	0.4	1.0	1.6	3.7	6.3	13.3	27.9	61.0	194.6
Non-Hispanic White, female									
2003.....	*	0.4	0.7	1.8	3.6	8.1	18.1	41.7	169.6
2002.....	*	0.4	1.0	1.6	4.3	7.6	17.9	45.9	181.9
2001.....	*	0.5	1.1	1.6	4.0	8.1	20.4	48.6	192.6
2000.....	*	0.3	0.8	1.5	4.1	8.3	19.8	49.1	202.1
1999.....	*	0.4	0.9	2.1	4.0	9.8	21.9	51.4	215.9
Non-Hispanic Black, both sexes									
2003.....	*	1.3	2.0	3.7	6.6	12.3	20.9	45.6	122.9
2002.....	*	1.5	2.0	4.3	5.3	11.7	27.0	52.1	118.0
2001.....	*	1.0	2.0	5.2	6.3	12.1	23.2	42.5	126.1
2000.....	*	1.5	2.1	4.8	5.8	13.1	23.6	54.6	140.2
1999.....	*	1.3	2.0	4.8	7.0	12.2	27.6	47.0	155.8
Non-Hispanic Black, male									
2003.....	*	*	*	5.0	8.1	17.2	29.0	58.9	105.5
2002.....	*	2.4	*	6.5	6.4	15.4	35.7	60.1	109.4
2001.....	*	*	*	6.1	7.8	15.1	30.8	48.2	113.2
2000.....	*	*	*	6.2	8.8	14.0	25.0	67.8	150.3
1999.....	*	*	*	5.9	9.0	15.9	37.7	56.7	166.4
Non-Hispanic Black, female									
2003.....	*	*	*	*	5.4	9.1	16.1	38.7	129.6
2002.....	*	*	*	*	4.6	9.2	21.9	47.9	121.2
2001.....	*	*	*	4.4	5.2	10.1	18.7	39.6	130.9
2000.....	*	*	*	3.7	*	12.5	22.7	48.0	136.5
1999.....	*	*	*	3.9	5.5	9.6	21.5	42.1	151.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of circulatory system (I71-I78)									
All origins 3/, both sexes									
2003.....	2.4	3.5	5.7	11.3	20.5	34.2	56.6	82.7	146.0
2002.....	2.2	3.1	6.2	11.7	22.3	36.5	57.8	83.8	148.4
2001.....	2.0	3.3	6.0	12.1	22.3	36.6	60.6	88.1	158.2
2000.....	2.1	3.5	6.4	12.7	24.5	40.6	62.9	92.2	163.2
1999.....	2.1	3.3	6.7	12.5	23.5	40.1	63.5	94.3	165.2
All origins 3/, male									
2003.....	3.2	5.1	7.9	15.9	29.4	46.1	78.9	111.3	179.7
2002.....	3.2	4.4	8.6	16.3	31.7	49.2	78.3	113.7	187.4
2001.....	2.8	4.4	8.2	16.8	30.8	51.2	81.0	122.5	197.3
2000.....	2.8	4.9	9.3	18.0	34.6	57.6	88.6	126.6	214.9
1999.....	3.0	4.4	9.4	18.2	33.5	56.7	89.1	129.3	207.5
All origins 3/, female									
2003.....	1.6	2.1	3.7	7.2	12.8	24.6	40.7	65.2	131.1
2002.....	1.4	2.0	3.9	7.5	14.2	26.4	43.4	65.6	131.5
2001.....	1.2	2.2	3.9	7.8	15.0	25.1	46.3	67.4	141.8
2000.....	1.5	2.1	3.7	7.9	15.7	27.2	45.0	71.9	142.2
1999.....	1.3	2.1	4.1	7.3	15.0	27.2	45.7	73.7	148.2
Hispanic, both sexes									
2003.....	1.4	1.8	3.8	5.7	11.3	17.6	37.6	50.0	106.2
2002.....	1.4	2.1	4.2	8.2	11.1	19.5	34.7	51.7	92.6
2001.....	1.6	2.8	3.7	6.2	11.3	16.8	35.4	45.2	95.0
2000.....	*	2.7	4.7	7.2	14.7	27.0	39.5	47.9	102.8
1999.....	*	1.8	4.3	7.6	12.6	21.2	31.3	67.0	110.8
Hispanic, male									
2003.....	*	2.8	5.8	9.6	15.3	23.9	54.1	66.5	138.5
2002.....	*	2.8	4.8	10.3	14.9	26.3	44.8	77.8	102.1
2001.....	2.1	3.6	4.4	9.0	16.7	24.2	51.8	55.8	108.6
2000.....	*	3.6	6.1	10.4	19.0	34.5	59.8	66.3	131.0
1999.....	*	*	5.8	10.1	17.0	29.0	44.5	97.5	163.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

Hispanic, female

2003.....	*	*	*	*	8.0	12.8	25.9	39.2	89.8
2002.....	*	*	3.7	6.5	8.0	14.2	27.4	35.0	87.9
2001.....	*	*	*	*	6.9	11.2	23.7	38.6	88.4
2000.....	*	*	*	*	11.2	21.4	25.1	36.7	89.0
1999.....	*	*	*	5.4	9.1	15.3	22.0	48.8	85.3

Non-Hispanic 4/, both sexes

2003.....	2.5	3.7	5.9	11.7	21.2	35.1	57.5	84.1	147.5
2002.....	2.3	3.2	6.3	11.9	23.0	37.5	58.8	85.1	150.3
2001.....	2.1	3.3	6.2	12.5	23.0	37.7	61.7	89.6	160.1
2000.....	2.2	3.5	6.5	13.0	25.0	41.3	63.8	93.7	165.0
1999.....	2.2	3.4	6.9	12.8	24.1	41.1	64.8	95.0	166.8

Non-Hispanic 4/, male

2003.....	3.4	5.3	8.0	16.3	30.4	47.3	80.1	113.3	181.5
2002.....	3.3	4.5	8.9	16.8	32.8	50.5	79.9	115.1	190.7
2001.....	2.9	4.4	8.5	17.3	31.6	52.6	82.2	124.9	200.5
2000.....	2.9	5.0	9.5	18.5	35.6	58.7	89.6	128.8	217.9
1999.....	3.1	4.6	9.7	18.7	34.3	58.0	91.1	130.2	209.0

Non-Hispanic 4/, female

2003.....	1.7	2.2	3.8	7.6	13.1	25.3	41.4	66.2	132.6
2002.....	1.4	2.0	4.0	7.5	14.6	27.1	44.0	66.9	132.8
2001.....	1.2	2.2	3.9	8.1	15.6	25.9	47.3	68.4	143.3
2000.....	1.5	2.1	3.7	8.1	16.0	27.5	45.8	73.0	143.6
1999.....	1.4	2.2	4.2	7.5	15.3	27.8	46.6	74.5	149.9

Non-Hispanic White, both sexes

2003.....	2.1	3.2	5.6	11.4	21.4	35.8	58.5	84.3	148.3
2002.....	1.9	2.8	5.9	11.6	23.2	38.1	59.8	86.1	150.9
2001.....	1.8	2.9	5.7	12.1	23.3	38.0	61.8	90.0	160.1
2000.....	1.9	3.0	6.1	12.7	25.2	42.1	64.0	94.6	164.8
1999.....	1.9	3.0	6.5	12.9	24.4	41.8	65.5	95.6	166.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

Non-Hispanic White, male

2003.....	2.9	4.7	7.8	15.9	30.8	48.9	81.8	114.0	185.8
2002.....	2.8	3.9	8.2	16.5	33.7	51.2	82.2	117.6	194.6
2001.....	2.5	3.8	8.2	16.9	32.5	53.3	83.4	126.6	204.7
2000.....	2.7	4.4	9.0	18.4	36.6	60.3	90.5	132.0	222.7
1999.....	2.8	4.1	9.5	19.1	35.4	59.8	93.2	131.9	212.8

Non-Hispanic White, female

2003.....	1.3	1.8	3.5	7.3	13.0	25.1	41.7	66.0	131.8
2002.....	1.0	1.7	3.7	7.2	13.9	27.3	43.7	66.9	132.1
2001.....	1.1	1.9	3.4	7.7	15.3	25.5	46.5	67.9	141.6
2000.....	1.1	1.7	3.3	7.5	15.2	27.5	45.3	72.3	141.5
1999.....	1.1	2.0	3.6	7.2	14.8	27.5	46.1	74.1	148.3

Non-Hispanic Black, both sexes

2003.....	5.5	7.5	8.7	16.6	24.5	36.0	56.1	93.6	152.9
2002.....	5.4	6.8	11.3	16.6	26.2	38.6	59.0	84.6	157.6
2001.....	4.1	6.6	10.9	18.2	23.2	43.5	69.8	97.1	175.7
2000.....	4.6	6.9	11.3	18.6	27.8	40.1	69.9	93.1	180.0
1999.....	4.7	6.3	11.4	13.8	26.5	39.1	64.9	97.4	180.1

Non-Hispanic Black, male

2003.....	7.1	10.1	11.6	23.8	33.9	39.8	73.4	120.7	141.7
2002.....	6.9	9.9	16.6	23.4	31.2	49.9	65.7	100.6	168.5
2001.....	5.7	9.6	13.2	24.0	28.1	56.5	78.7	129.2	168.6
2000.....	5.6	10.2	15.2	23.0	33.6	52.3	91.5	106.4	179.7
1999.....	6.3	8.9	14.3	18.1	32.4	49.9	75.4	118.7	184.2

Non-Hispanic Black, female

2003.....	4.1	5.2	6.4	11.0	17.5	33.4	45.8	79.4	157.2
2002.....	4.2	4.2	7.0	11.3	22.4	31.0	54.9	76.3	153.4
2001.....	2.7	4.1	9.0	13.6	19.6	34.9	64.5	80.8	178.4
2000.....	3.8	4.2	8.0	15.1	23.6	32.0	56.9	86.4	180.1
1999.....	3.3	4.1	9.1	10.4	22.0	31.9	58.5	86.9	178.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)									
All origins 3/, both sexes									
2003.....	1.7	2.4	3.7	7.3	13.5	22.7	36.6	50.7	70.0
2002.....	1.5	2.0	4.0	7.4	14.7	23.9	37.5	51.5	69.4
2001.....	1.4	2.0	3.7	7.8	15.0	23.8	39.0	55.4	77.5
2000.....	1.3	2.2	4.3	8.4	16.3	27.8	41.0	56.5	79.2
1999.....	1.4	2.1	4.4	8.1	16.5	27.3	42.2	58.2	80.3
All origins 3/, male									
2003.....	2.5	3.7	5.8	11.4	20.2	33.0	53.8	74.2	105.5
2002.....	2.4	3.2	6.3	11.5	23.0	34.9	55.0	74.5	106.6
2001.....	2.1	3.2	5.8	12.0	22.2	36.0	56.3	82.5	114.8
2000.....	1.9	3.5	6.7	13.1	24.9	41.8	61.9	84.9	127.3
1999.....	2.2	3.3	6.8	13.0	25.3	41.0	63.3	88.6	125.3
All origins 3/, female									
2003.....	0.9	1.1	1.7	3.6	7.8	14.5	24.3	36.3	54.2
2002.....	0.7	1.0	1.8	3.7	7.5	15.1	25.1	37.6	53.3
2001.....	0.6	0.9	1.8	3.9	8.8	14.0	26.9	39.2	61.9
2000.....	0.8	1.0	2.1	4.1	8.9	16.7	26.5	39.7	59.6
1999.....	0.7	1.0	2.1	3.7	9.0	16.6	27.5	40.4	62.2
Hispanic, both sexes									
2003.....	*	*	2.5	3.9	5.7	11.7	25.0	29.5	49.2
2002.....	*	*	2.7	5.3	6.4	9.0	19.2	24.5	31.0
2001.....	*	1.6	1.9	3.4	7.1	9.5	21.2	25.8	31.9
2000.....	*	2.1	2.7	4.3	7.7	15.7	25.4	30.1	41.8
1999.....	*	*	2.6	4.3	7.3	11.6	18.4	34.1	40.7
Hispanic, male									
2003.....	*	*	4.1	6.6	8.4	17.9	37.1	40.5	80.0
2002.....	*	*	3.9	7.7	9.5	15.4	25.6	43.5	43.5
2001.....	*	*	*	5.7	12.4	16.5	35.5	33.8	53.3
2000.....	*	3.0	*	6.6	12.3	20.9	42.1	42.7	64.5
1999.....	*	*	*	7.1	11.6	18.8	24.2	53.4	79.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)									
Hispanic, female									
2003.....	*	*	*	*	*	6.9	16.4	22.3	33.6
2002.....	*	*	*	*	*	*	14.6	*	24.9
2001.....	*	*	*	*	*	*	11.1	20.9	21.4
2000.....	*	*	*	*	*	11.8	13.6	22.4	30.7
1999.....	*	*	*	*	*	*	14.3	22.5	21.8
Non-Hispanic 4/, both sexes									
2003.....	1.8	2.5	3.8	7.6	14.1	23.4	37.1	51.6	70.7
2002.....	1.6	2.1	4.1	7.6	15.2	24.8	38.3	52.6	70.8
2001.....	1.4	2.1	3.9	8.1	15.5	24.5	39.8	56.4	79.0
2000.....	1.4	2.2	4.5	8.7	16.8	28.4	41.6	57.3	80.3
1999.....	1.5	2.2	4.5	8.4	17.0	28.1	43.2	58.9	81.7
Non-Hispanic 4/, male									
2003.....	2.7	3.9	6.0	11.8	21.0	33.8	54.6	75.8	106.6
2002.....	2.5	3.3	6.5	11.7	23.8	36.1	56.4	75.7	109.1
2001.....	2.2	3.2	6.0	12.4	22.8	37.0	57.2	84.2	117.0
2000.....	2.0	3.5	6.9	13.5	25.7	42.9	62.6	86.5	129.6
1999.....	2.3	3.4	7.0	13.3	26.1	42.1	65.0	89.7	127.0
Non-Hispanic 4/, female									
2003.....	0.9	1.1	1.8	3.8	8.1	15.0	24.7	36.8	55.0
2002.....	0.8	1.0	1.8	3.8	7.7	15.8	25.5	38.6	54.3
2001.....	0.7	0.9	1.9	4.1	9.2	14.6	27.7	39.8	63.2
2000.....	0.8	1.0	2.2	4.3	9.2	16.9	27.0	40.2	60.4
1999.....	0.7	1.0	2.1	3.9	9.3	17.1	28.0	40.9	63.5
Non-Hispanic White, both sexes									
2003.....	1.5	2.2	3.7	7.6	14.7	24.4	38.7	53.1	73.2
2002.....	1.3	1.8	3.8	7.7	16.0	25.8	39.9	54.3	72.8
2001.....	1.2	1.7	3.7	8.1	16.1	25.5	40.7	58.1	81.3
2000.....	1.2	2.0	4.2	8.8	17.3	29.7	42.6	59.1	82.0
1999.....	1.2	1.9	4.3	8.8	17.7	29.2	44.7	60.7	83.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)									
Non-Hispanic White, male									
2003.....	2.2	3.5	5.9	11.8	22.0	35.6	57.1	78.1	111.2
2002.....	2.0	2.8	6.1	11.9	25.3	37.4	59.1	79.1	114.7
2001.....	1.9	2.8	5.9	12.3	23.9	38.5	58.7	87.4	122.9
2000.....	1.9	3.2	6.7	13.8	27.0	45.0	64.3	90.3	135.3
1999.....	2.0	3.0	7.0	14.1	27.4	44.2	68.3	92.7	131.8
Non-Hispanic White, female									
2003.....	0.7	0.9	1.5	3.7	8.2	15.1	25.4	37.8	56.5
2002.....	0.6	0.8	1.7	3.8	7.7	16.3	26.2	39.2	54.7
2001.....	0.5	0.7	1.6	4.1	9.2	14.9	27.8	40.5	64.0
2000.....	0.5	0.8	1.9	4.1	8.8	17.4	27.4	40.5	60.5
1999.....	0.5	0.9	1.8	3.8	9.2	17.3	28.1	41.8	64.5
Non-Hispanic Black, both sexes									
2003.....	4.0	4.8	5.9	9.2	12.0	17.8	26.3	36.6	47.3
2002.....	3.9	4.8	6.8	8.3	11.7	18.8	25.1	37.9	50.5
2001.....	2.8	4.5	6.2	9.5	11.6	18.6	34.5	43.3	55.9
2000.....	2.9	3.7	7.2	9.5	14.6	20.5	34.1	41.1	63.0
1999.....	3.6	4.1	6.4	6.7	14.2	20.0	29.8	40.8	61.1
Non-Hispanic Black, male									
2003.....	6.0	7.2	8.2	14.9	16.5	19.9	33.2	50.6	52.2
2002.....	5.8	7.7	11.2	13.2	14.8	25.3	31.0	41.3	49.2
2001.....	4.1	7.0	8.5	14.8	14.9	26.5	42.7	55.6	49.7
2000.....	3.6	6.1	10.3	13.2	18.1	28.4	47.7	46.3	70.5
1999.....	5.5	6.6	9.1	9.6	19.4	25.6	30.0	54.9	77.3
Non-Hispanic Black, female									
2003.....	2.2	2.8	3.9	4.8	8.6	16.4	22.2	29.2	45.4
2002.....	2.2	2.3	3.3	4.5	9.3	14.5	21.6	36.1	51.0
2001.....	1.7	2.3	4.4	5.3	9.2	13.3	29.7	37.1	58.2
2000.....	2.4	*	4.7	6.5	12.0	15.2	25.8	38.5	60.2
1999.....	2.0	*	4.3	4.4	10.4	16.3	29.7	33.8	55.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

All origins 3/, both sexes

2003.....	0.7	1.2	2.0	4.0	7.0	11.4	20.0	32.0	76.0
2002.....	0.7	1.1	2.2	4.3	7.7	12.6	20.4	32.3	79.0
2001.....	0.6	1.2	2.2	4.3	7.3	12.9	21.5	32.7	80.7
2000.....	0.8	1.3	2.1	4.3	8.2	12.8	21.8	35.8	84.0
1999.....	0.7	1.2	2.3	4.4	7.0	12.9	21.3	36.1	84.9

All origins 3/, male

2003.....	0.7	1.3	2.0	4.4	9.3	13.0	25.1	37.1	74.2
2002.....	0.8	1.2	2.3	4.9	8.7	14.2	23.4	39.2	80.8
2001.....	0.7	1.2	2.4	4.8	8.5	15.2	24.7	40.0	82.5
2000.....	0.8	1.4	2.6	4.9	9.7	15.7	26.6	41.7	87.6
1999.....	0.8	1.2	2.6	5.2	8.1	15.7	25.9	40.8	82.2

All origins 3/, female

2003.....	0.7	1.0	1.9	3.6	5.0	10.1	16.3	28.9	76.9
2002.....	0.6	1.0	2.1	3.7	6.7	11.3	18.3	28.1	78.2
2001.....	0.6	1.3	2.0	3.9	6.3	11.1	19.3	28.3	79.9
2000.....	0.7	1.1	1.6	3.7	6.9	10.5	18.5	32.2	82.6
1999.....	0.6	1.2	2.1	3.6	6.0	10.6	18.2	33.3	86.0

Hispanic, both sexes

2003.....	*	*	*	*	5.6	5.9	12.6	20.5	57.0
2002.....	*	*	*	3.0	4.7	10.5	15.5	27.2	61.5
2001.....	*	*	*	2.8	4.1	7.3	14.2	19.4	63.2
2000.....	*	*	*	2.9	7.0	11.3	14.1	17.8	61.0
1999.....	*	*	*	3.3	5.4	9.6	12.9	32.9	70.2

Hispanic, male

2003.....	*	*	*	*	6.8	*	17.0	26.0	58.5
2002.....	*	*	*	*	*	11.0	19.2	34.3	58.6
2001.....	*	*	*	*	*	*	16.4	*	55.2
2000.....	*	*	*	*	*	13.6	17.7	*	66.5
1999.....	*	*	*	*	*	10.2	20.3	44.0	84.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

Hispanic, female

2003.....	*	*	*	*	*	*	9.5	16.9	56.3
2002.....	*	*	*	*	*	10.1	12.8	22.6	63.0
2001.....	*	*	*	*	*	7.0	12.6	17.7	67.0
2000.....	*	*	*	*	7.2	9.6	11.5	*	58.4
1999.....	*	*	*	*	*	9.2	*	26.3	63.4

Non-Hispanic 4/, both sexes

2003.....	0.7	1.2	2.0	4.1	7.1	11.7	20.4	32.5	76.7
2002.....	0.7	1.1	2.3	4.3	7.8	12.7	20.6	32.5	79.5
2001.....	0.6	1.2	2.3	4.4	7.5	13.2	21.9	33.1	81.1
2000.....	0.8	1.3	2.1	4.4	8.2	12.9	22.2	36.3	84.7
1999.....	0.7	1.2	2.4	4.4	7.1	13.0	21.7	36.1	85.1

Non-Hispanic 4/, male

2003.....	0.7	1.4	2.0	4.5	9.4	13.5	25.5	37.6	74.9
2002.....	0.8	1.2	2.4	5.0	9.0	14.4	23.5	39.3	81.7
2001.....	0.7	1.2	2.5	4.9	8.8	15.6	25.0	40.7	83.5
2000.....	0.9	1.5	2.6	4.9	9.9	15.8	27.0	42.3	88.2
1999.....	0.8	1.3	2.7	5.3	8.3	15.9	26.1	40.5	82.0

Non-Hispanic 4/, female

2003.....	0.8	1.0	2.0	3.8	5.1	10.3	16.7	29.4	77.6
2002.....	0.6	1.0	2.1	3.7	6.9	11.3	18.5	28.3	78.5
2001.....	0.6	1.2	2.1	4.0	6.4	11.3	19.6	28.6	80.1
2000.....	0.7	1.1	1.6	3.8	6.8	10.6	18.8	32.8	83.3
1999.....	0.6	1.2	2.1	3.6	6.0	10.7	18.6	33.6	86.4

Non-Hispanic White, both sexes

2003.....	0.6	1.1	2.0	3.8	6.6	11.4	19.8	31.1	75.1
2002.....	0.6	1.0	2.1	3.9	7.2	12.3	19.8	31.8	78.2
2001.....	0.6	1.1	2.1	4.0	7.3	12.5	21.1	31.9	78.8
2000.....	0.7	1.0	1.9	3.9	7.8	12.4	21.3	35.5	82.8
1999.....	0.7	1.1	2.1	4.2	6.7	12.6	20.9	34.8	83.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic White, male

2003.....	0.7	1.2	2.0	4.1	8.8	13.2	24.7	35.9	74.6
2002.....	0.8	1.1	2.1	4.5	8.4	13.8	23.1	38.5	79.9
2001.....	0.6	1.0	2.3	4.5	8.6	14.8	24.6	39.2	81.8
2000.....	0.8	1.2	2.4	4.5	9.5	15.3	26.2	41.7	87.4
1999.....	0.9	1.2	2.5	5.1	8.0	15.6	25.0	39.2	81.0

Non-Hispanic White, female

2003.....	0.6	0.9	1.9	3.6	4.8	9.9	16.3	28.2	75.4
2002.....	0.5	1.0	2.0	3.4	6.2	11.0	17.5	27.7	77.4
2001.....	0.5	1.2	1.8	3.6	6.1	10.6	18.7	27.5	77.6
2000.....	0.6	0.9	1.3	3.4	6.4	10.1	17.9	31.8	81.0
1999.....	0.6	1.1	1.8	3.4	5.5	10.2	18.0	32.2	83.9

Non-Hispanic Black, both sexes

2003.....	1.5	2.6	2.9	7.4	12.5	18.2	29.8	57.0	105.6
2002.....	1.6	2.0	4.5	8.3	14.5	19.8	33.9	46.7	107.1
2001.....	1.3	2.2	4.6	8.7	11.6	24.9	35.3	53.8	119.9
2000.....	1.7	3.2	4.0	9.1	13.2	19.6	35.9	52.0	117.0
1999.....	1.0	2.2	5.0	7.1	12.2	19.1	35.1	56.6	119.0

Non-Hispanic Black, male

2003.....	*	2.9	3.3	8.9	17.5	19.9	40.2	70.2	89.5
2002.....	*	2.1	5.4	10.2	16.4	24.6	34.7	59.3	119.2
2001.....	*	2.6	4.7	9.2	13.3	30.0	36.0	73.6	119.0
2000.....	2.0	4.1	4.9	9.8	15.5	23.9	43.8	60.1	109.2
1999.....	*	*	5.2	8.5	13.0	24.2	45.4	63.8	107.0

Non-Hispanic Black, female

2003.....	1.9	2.4	2.5	6.2	8.8	17.0	23.6	50.2	111.8
2002.....	2.0	1.9	3.8	6.8	13.1	16.5	33.4	40.2	102.4
2001.....	*	*	4.6	8.4	10.4	21.6	34.8	43.7	120.2
2000.....	*	2.4	3.3	8.6	11.6	16.8	31.1	48.0	119.9
1999.....	*	*	4.8	6.0	11.6	15.6	28.8	53.1	123.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	19.5
2002.....	*	*	*	*	*	*	*	*	23.2
2001.....	*	*	*	*	*	*	*	*	33.5
2000.....	*	*	*	*	*	*	*	*	23.7
1999.....	*	*	*	*	*	*	*	*	23.9

Non-Hispanic 4/, both sexes

2003.....	1.3	1.5	1.7	2.5	3.3	5.1	7.0	12.1	24.3
2002.....	1.2	1.5	2.1	2.6	3.5	4.3	7.6	13.3	24.7
2001.....	1.1	1.3	2.0	2.7	3.8	5.4	7.8	11.9	25.8
2000.....	1.3	1.5	1.8	2.7	3.4	5.4	7.7	12.0	25.6
1999.....	1.0	1.0	1.6	2.2	3.3	4.2	6.9	12.7	23.3

Non-Hispanic 4/, male

2003.....	1.4	1.7	2.2	2.7	3.6	5.0	6.9	11.0	19.3
2002.....	1.4	1.6	2.4	2.6	3.6	4.3	7.1	13.5	21.1
2001.....	1.2	1.5	2.3	3.0	3.9	6.1	8.3	12.0	21.6
2000.....	1.5	1.7	2.0	2.7	3.9	5.7	8.0	12.3	22.1
1999.....	1.0	1.2	1.7	2.3	3.8	4.7	7.6	13.3	20.5

Non-Hispanic 4/, female

2003.....	1.2	1.3	1.3	2.3	3.1	5.2	7.0	12.7	26.5
2002.....	1.1	1.4	1.8	2.6	3.3	4.3	7.9	13.2	26.2
2001.....	0.9	1.0	1.8	2.4	3.8	4.9	7.4	11.8	27.6
2000.....	1.1	1.3	1.6	2.7	3.0	5.1	7.5	11.8	27.0
1999.....	1.0	0.9	1.5	2.1	2.9	3.8	6.5	12.4	24.4

Non-Hispanic White, both sexes

2003.....	1.1	1.4	1.6	2.3	3.1	4.8	6.5	11.8	23.8
2002.....	1.1	1.4	1.9	2.2	3.2	4.1	7.3	12.8	24.5
2001.....	0.9	1.1	1.8	2.4	3.5	5.1	7.3	11.8	25.7
2000.....	1.2	1.4	1.6	2.4	2.9	5.1	7.3	11.9	25.5
1999.....	0.8	0.9	1.6	2.1	2.9	4.0	6.6	12.4	23.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

Non-Hispanic White, male

2003.....	1.2	1.5	2.1	2.3	3.2	4.8	6.4	10.3	18.7
2002.....	1.2	1.5	2.1	2.3	3.3	4.0	6.8	12.5	20.5
2001.....	1.2	1.3	2.1	2.5	3.6	5.7	7.8	11.9	21.2
2000.....	1.4	1.6	1.7	2.4	3.4	5.6	7.8	12.1	21.8
1999.....	0.9	1.0	1.7	2.3	3.3	4.5	7.1	12.8	20.3

Non-Hispanic White, female

2003.....	1.1	1.2	1.0	2.3	2.9	4.9	6.5	12.7	26.1
2002.....	1.0	1.3	1.7	2.2	3.2	4.1	7.6	12.9	26.3
2001.....	0.7	1.0	1.6	2.3	3.5	4.7	6.9	11.8	27.6
2000.....	1.0	1.1	1.4	2.4	2.5	4.7	7.0	11.8	27.0
1999.....	0.8	0.8	1.5	1.9	2.6	3.6	6.3	12.2	24.1

Non-Hispanic Black, both sexes

2003.....	2.7	2.6	3.7	5.3	6.3	9.0	13.9	17.9	33.6
2002.....	2.4	2.9	4.3	5.9	6.4	7.4	13.0	22.7	29.9
2001.....	2.2	2.4	4.0	6.2	7.1	8.9	14.0	14.9	29.7
2000.....	2.5	3.2	4.0	5.3	8.7	9.3	14.5	14.4	29.9
1999.....	2.4	2.4	2.1	4.0	7.0	6.5	11.6	17.9	28.5

Non-Hispanic Black, male

2003.....	3.5	3.5	3.8	7.5	7.8	9.1	15.9	21.1	33.0
2002.....	3.1	2.9	5.5	5.7	7.7	7.9	12.2	28.9	31.7
2001.....	1.8	3.1	4.8	8.8	8.1	10.7	16.1	*	28.9
2000.....	3.2	3.8	4.9	5.7	9.3	8.9	13.0	*	30.5
1999.....	2.7	3.0	*	*	9.6	6.9	15.0	21.3	*

Non-Hispanic Black, female

2003.....	2.0	1.8	3.7	3.6	5.1	8.9	12.8	16.2	33.8
2002.....	1.8	3.0	3.3	6.0	5.5	7.1	13.5	19.5	29.3
2001.....	2.5	*	3.3	4.1	6.3	7.8	12.7	14.6	29.9
2000.....	2.0	2.7	3.3	4.9	8.3	9.5	15.4	14.7	29.7
1999.....	2.2	*	*	4.3	5.1	6.2	9.6	16.2	30.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Influenza and pneumonia (J10-J18)

All origins 3/, both sexes

2003.....	4.2	6.4	8.2	15.2	26.6	49.4	103.3	216.8	666.1
2002.....	4.2	5.5	8.5	14.8	25.3	50.9	106.5	227.3	696.6
2001.....	3.9	5.4	8.1	13.9	24.4	49.3	101.1	216.9	685.6
2000.....	3.9	5.6	8.9	15.6	26.6	52.6	106.7	240.8	744.1
1999.....	4.1	5.3	8.4	14.3	25.0	50.5	105.9	233.8	751.8

All origins 3/, male

2003.....	4.9	8.1	9.6	17.9	33.3	62.1	130.1	272.3	757.8
2002.....	4.9	6.7	10.2	18.0	31.2	63.9	133.9	287.7	805.8
2001.....	4.9	6.7	10.0	17.0	30.3	61.6	130.7	274.7	808.4
2000.....	4.7	7.0	10.9	18.9	32.9	65.7	134.1	308.8	892.0
1999.....	5.2	6.8	9.7	16.8	31.7	64.2	133.5	301.7	894.7

All origins 3/, female

2003.....	3.5	4.8	6.8	12.7	20.7	39.2	84.2	182.9	625.6
2002.....	3.4	4.3	6.8	11.8	20.1	40.6	87.2	190.6	649.2
2001.....	3.0	4.1	6.4	11.1	19.2	39.5	80.5	182.3	634.2
2000.....	3.1	4.3	7.0	12.5	21.2	42.2	87.6	200.7	683.9
1999.....	3.0	3.9	7.2	12.0	19.2	39.7	86.7	194.1	694.2

Hispanic, both sexes

2003.....	2.9	5.3	7.9	14.7	24.6	48.2	93.6	200.4	516.2
2002.....	2.9	4.0	8.3	13.4	22.8	47.5	102.6	209.4	566.0
2001.....	2.9	4.4	6.8	12.1	22.2	51.9	98.0	206.1	660.3
2000.....	3.0	4.8	8.5	14.5	23.0	55.9	95.8	210.0	643.6
1999.....	2.9	4.4	6.4	13.4	24.7	40.4	81.3	186.2	614.6

Hispanic, male

2003.....	3.5	7.6	9.1	15.3	30.9	59.8	111.8	257.8	577.2
2002.....	3.9	5.8	11.0	18.0	29.8	61.9	120.9	266.5	655.9
2001.....	4.9	5.3	10.0	13.3	25.2	60.8	122.8	270.1	765.6
2000.....	4.6	6.2	11.0	19.9	28.0	69.5	122.5	295.9	763.9
1999.....	3.8	6.2	7.4	14.3	31.7	53.5	100.6	243.6	720.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza and pneumonia (J10-J18)

Hispanic, female

2003.....	2.3	3.1	6.8	14.2	19.5	39.3	80.6	163.1	485.2
2002.....	1.9	*	5.9	9.4	17.1	36.5	89.6	173.0	521.4
2001.....	*	3.6	4.0	11.0	19.8	45.0	80.4	166.5	608.8
2000.....	*	3.6	6.4	9.9	19.0	45.7	76.9	157.7	584.6
1999.....	*	*	5.4	12.6	19.1	30.6	67.7	151.9	563.6

Non-Hispanic 4/, both sexes

2003.....	4.3	6.5	8.2	15.1	26.5	49.3	103.6	217.2	671.1
2002.....	4.3	5.6	8.5	14.9	25.3	51.0	106.3	227.3	700.2
2001.....	4.0	5.4	8.2	14.0	24.4	48.8	101.0	216.7	684.8
2000.....	3.9	5.6	8.8	15.6	26.6	52.2	106.7	241.0	745.7
1999.....	4.1	5.3	8.5	14.2	24.8	50.8	106.6	234.7	754.0

Non-Hispanic 4/, male

2003.....	5.0	8.0	9.6	17.9	33.2	62.0	130.8	272.6	764.6
2002.....	5.0	6.8	10.1	18.0	31.1	63.7	134.2	287.4	810.7
2001.....	4.8	6.7	9.9	17.2	30.5	61.1	130.6	274.2	808.2
2000.....	4.7	7.0	10.8	18.7	32.9	65.2	134.0	307.9	894.9
1999.....	5.2	6.7	9.7	16.7	31.5	64.5	134.4	302.7	897.9

Non-Hispanic 4/, female

2003.....	3.7	5.0	6.8	12.5	20.7	39.1	84.2	183.4	630.0
2002.....	3.5	4.5	6.9	12.0	20.4	40.8	86.7	190.7	652.6
2001.....	3.2	4.1	6.5	11.1	19.1	39.0	80.2	182.3	633.5
2000.....	3.3	4.4	7.0	12.7	21.2	41.9	87.7	201.6	685.4
1999.....	3.1	4.0	7.3	11.9	19.1	40.1	87.2	194.9	696.4

Non-Hispanic White, both sexes

2003.....	3.8	5.7	7.2	13.8	25.0	47.6	102.5	218.5	684.5
2002.....	3.9	5.0	7.6	13.8	24.5	49.8	105.0	227.7	712.7
2001.....	3.3	4.8	7.5	12.9	23.2	47.4	98.2	216.2	693.9
2000.....	3.3	4.9	7.9	14.2	24.4	49.7	105.5	240.7	759.8
1999.....	3.4	4.6	7.7	12.9	23.8	48.9	106.1	236.5	766.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza and pneumonia (J10-J18)

Non-Hispanic White, male

2003.....	4.3	7.1	8.3	15.5	30.5	58.8	127.6	271.3	774.1
2002.....	4.4	6.0	8.8	16.3	29.6	62.0	131.3	286.1	820.6
2001.....	4.0	5.7	8.9	15.3	28.3	58.5	125.1	271.6	813.9
2000.....	3.7	5.7	9.2	16.9	29.5	61.4	131.6	305.1	908.1
1999.....	4.2	5.5	8.6	15.0	29.7	61.0	132.9	302.9	909.3

Non-Hispanic White, female

2003.....	3.2	4.4	6.2	12.1	20.0	38.4	84.4	186.0	645.1
2002.....	3.4	4.1	6.3	11.5	20.0	39.8	86.1	192.0	666.1
2001.....	2.6	3.9	6.1	10.6	18.7	38.3	79.2	182.8	644.0
2000.....	2.9	4.0	6.7	11.7	20.0	40.2	87.2	202.5	700.0
1999.....	2.6	3.7	6.8	11.0	18.6	39.3	87.3	197.2	710.2

Non-Hispanic Black, both sexes

2003.....	8.7	12.7	16.1	27.6	42.1	71.1	123.9	207.9	528.3
2002.....	7.7	10.3	17.6	26.8	35.6	70.0	124.3	233.8	579.8
2001.....	9.2	10.8	15.4	23.7	34.9	66.2	133.9	224.9	583.1
2000.....	8.6	12.4	18.1	27.6	46.2	78.5	124.8	247.4	604.5
1999.....	9.9	11.9	16.3	26.6	37.0	76.4	119.8	220.4	634.7

Non-Hispanic Black, male

2003.....	10.5	16.8	22.0	40.5	62.0	101.8	179.4	287.5	642.5
2002.....	10.1	13.2	24.2	36.7	48.1	91.0	175.1	324.5	695.8
2001.....	11.6	15.9	20.8	34.3	50.8	95.4	203.4	309.9	718.4
2000.....	11.7	17.6	26.2	35.5	65.0	108.7	173.4	345.9	763.2
1999.....	13.6	17.4	21.8	34.2	52.6	110.8	162.4	311.7	803.4

Non-Hispanic Black, female

2003.....	7.1	9.2	11.3	17.6	27.4	50.4	91.0	166.3	484.1
2002.....	5.5	7.9	12.1	19.1	26.4	56.0	93.9	186.7	535.5
2001.....	7.1	6.6	11.1	15.4	23.1	46.8	92.5	181.6	533.0
2000.....	6.0	8.1	11.4	21.3	32.2	58.5	95.5	197.9	545.6
1999.....	6.7	7.3	11.8	20.7	25.4	53.5	94.1	175.2	571.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza (J10-J11)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.1	0.2	0.3	0.3	0.8	1.4	2.8	4.8	17.5
2002.....	*	*	*	*	*	0.4	0.8	2.3	9.0
2001.....	*	*	*	*	*	*	*	0.7	2.3
2000.....	0.1	0.1	0.3	0.5	0.8	1.5	2.6	6.2	20.5
1999.....	*	*	0.2	0.4	0.6	1.2	2.1	6.2	21.6

Non-Hispanic 4/, male

2003.....	*	*	0.4	*	0.8	1.4	3.0	4.7	17.5
2002.....	*	*	*	*	*	*	0.9	2.7	9.2
2001.....	*	*	*	*	*	*	*	*	2.3
2000.....	*	*	*	0.5	1.0	1.7	2.8	6.2	21.1
1999.....	*	*	*	0.5	0.6	1.2	2.4	7.5	20.7

Non-Hispanic 4/, female

2003.....	*	*	*	0.3	0.7	1.3	2.6	4.8	17.6
2002.....	*	*	*	*	*	*	0.7	2.1	8.9
2001.....	*	*	*	*	*	*	*	0.8	2.2
2000.....	*	*	*	0.6	0.7	1.3	2.5	6.2	20.3
1999.....	*	*	*	*	0.7	1.2	1.9	5.4	22.0

Non-Hispanic White, both sexes

2003.....	0.1	0.2	0.3	0.3	0.8	1.5	3.0	5.2	18.8
2002.....	*	*	*	*	*	0.5	0.9	2.5	9.6
2001.....	*	*	*	*	*	*	*	0.7	2.4
2000.....	*	0.2	0.3	0.6	0.9	1.5	2.8	6.5	21.7
1999.....	*	*	0.2	0.4	0.7	1.2	2.2	6.5	23.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)									
All origins 3/, both sexes									
2003.....	4.1	6.2	7.9	14.9	25.8	48.1	100.6	212.1	649.1
2002.....	4.1	5.4	8.4	14.6	25.1	50.5	105.7	225.1	687.8
2001.....	3.9	5.3	8.1	13.8	24.3	49.2	100.9	216.1	683.4
2000.....	3.8	5.5	8.6	15.1	25.8	51.2	104.1	234.7	723.9
1999.....	4.0	5.2	8.2	13.9	24.3	49.3	103.8	227.7	730.6
All origins 3/, male									
2003.....	4.8	7.9	9.3	17.6	32.4	60.8	127.2	267.5	740.9
2002.....	4.9	6.7	10.1	17.8	30.9	63.4	133.0	285.0	796.9
2001.....	4.8	6.6	9.9	16.9	30.3	61.4	130.5	273.9	806.2
2000.....	4.6	6.8	10.6	18.4	31.9	64.1	131.5	302.6	871.3
1999.....	5.1	6.7	9.5	16.3	31.1	63.0	131.1	294.5	874.7
All origins 3/, female									
2003.....	3.4	4.7	6.5	12.4	20.1	37.9	81.7	178.2	608.5
2002.....	3.4	4.3	6.8	11.8	20.0	40.2	86.5	188.6	640.5
2001.....	2.9	4.1	6.3	11.0	19.1	39.4	80.2	181.5	632.0
2000.....	3.0	4.2	6.7	12.0	20.5	41.0	85.1	194.6	663.9
1999.....	2.9	3.9	7.0	11.7	18.6	38.6	84.8	188.7	672.5
Hispanic, both sexes									
2003.....	2.9	5.3	7.7	14.3	24.2	48.2	92.6	197.2	511.0
2002.....	2.9	4.0	8.3	13.4	22.7	47.4	102.1	209.0	563.2
2001.....	2.9	4.4	6.8	12.1	22.2	51.7	97.1	205.1	660.3
2000.....	2.9	4.8	8.4	14.4	22.5	55.7	94.9	206.1	633.0
1999.....	2.9	4.2	6.1	13.0	24.0	39.6	79.7	183.8	607.6
Hispanic, male									
2003.....	3.5	7.6	8.9	15.0	29.9	59.8	111.2	251.6	571.1
2002.....	3.8	5.8	11.0	18.0	29.5	61.9	120.9	265.4	654.2
2001.....	4.8	5.3	10.0	13.3	25.2	60.8	122.8	267.5	765.6
2000.....	4.5	6.2	10.7	19.6	27.2	69.5	121.8	290.1	751.8
1999.....	3.8	5.9	6.9	13.7	30.1	51.9	99.9	242.1	718.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)									
Hispanic, female									
2003.....	2.2	3.0	6.5	13.7	19.5	39.3	79.3	161.7	480.5
2002.....	1.9	*	5.9	9.4	17.1	36.2	88.7	173.0	518.1
2001.....	*	3.6	4.0	11.0	19.8	44.7	78.9	166.5	608.8
2000.....	*	3.6	6.4	9.9	18.7	45.3	75.8	155.0	574.7
1999.....	*	*	5.4	12.3	19.1	30.3	65.5	149.1	554.2
Non-Hispanic 4/, both sexes									
2003.....	4.2	6.3	7.9	14.8	25.8	47.9	100.8	212.4	653.6
2002.....	4.2	5.5	8.4	14.7	25.1	50.5	105.5	225.0	691.2
2001.....	4.0	5.4	8.1	13.9	24.3	48.7	100.8	216.0	682.5
2000.....	3.8	5.5	8.5	15.1	25.8	50.7	104.1	234.8	725.2
1999.....	4.1	5.3	8.2	13.8	24.2	49.6	104.4	228.5	732.4
Non-Hispanic 4/, male									
2003.....	4.9	7.8	9.3	17.7	32.3	60.6	127.8	267.8	747.1
2002.....	4.9	6.7	10.0	17.7	30.8	63.1	133.3	284.7	801.5
2001.....	4.8	6.7	9.8	17.1	30.4	60.9	130.4	273.6	805.9
2000.....	4.5	6.8	10.5	18.2	31.9	63.5	131.3	301.6	873.8
1999.....	5.1	6.6	9.5	16.3	30.9	63.3	132.0	295.2	877.2
Non-Hispanic 4/, female									
2003.....	3.5	4.8	6.5	12.1	20.0	37.7	81.6	178.6	612.4
2002.....	3.5	4.4	6.8	11.9	20.2	40.4	86.0	188.7	643.7
2001.....	3.1	4.1	6.5	10.9	19.0	38.9	80.0	181.6	631.3
2000.....	3.1	4.3	6.8	12.2	20.5	40.6	85.2	195.4	665.2
1999.....	3.0	4.0	7.0	11.6	18.4	38.9	85.3	189.6	674.4
Non-Hispanic White, both sexes									
2003.....	3.6	5.6	6.9	13.4	24.2	46.1	99.5	213.4	665.7
2002.....	3.8	5.0	7.5	13.7	24.3	49.3	104.1	225.2	703.1
2001.....	3.3	4.7	7.4	12.8	23.1	47.3	98.0	215.5	691.5
2000.....	3.2	4.7	7.6	13.7	23.6	48.2	102.7	234.2	738.1
1999.....	3.3	4.5	7.5	12.5	23.1	47.7	103.9	230.0	743.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)									
Non-Hispanic White, male									
2003.....	4.2	6.8	8.0	15.3	29.6	57.3	124.4	266.2	755.2
2002.....	4.3	5.9	8.7	16.1	29.4	61.4	130.3	283.2	810.6
2001.....	4.0	5.6	8.8	15.3	28.3	58.4	124.9	270.9	811.5
2000.....	3.7	5.6	8.9	16.4	28.5	59.6	128.7	298.4	886.1
1999.....	4.1	5.5	8.4	14.5	29.1	59.8	130.3	295.2	886.8
Non-Hispanic White, female									
2003.....	3.1	4.3	6.0	11.7	19.3	36.9	81.5	180.9	626.4
2002.....	3.4	4.0	6.2	11.4	19.9	39.4	85.4	189.8	656.6
2001.....	2.6	3.9	6.0	10.5	18.6	38.2	78.9	182.1	641.7
2000.....	2.8	3.9	6.3	11.1	19.3	39.0	84.5	196.0	678.4
1999.....	2.5	3.6	6.6	10.6	17.8	38.1	85.3	191.5	686.6
Non-Hispanic Black, both sexes									
2003.....	8.4	12.5	15.8	27.5	41.5	70.6	123.1	206.1	522.4
2002.....	7.7	10.3	17.5	26.5	35.2	69.8	124.1	233.0	577.4
2001.....	9.2	10.8	15.4	23.5	34.6	66.2	133.9	223.8	581.5
2000.....	8.4	12.3	17.9	27.2	45.6	77.0	123.9	245.1	595.0
1999.....	9.9	11.8	16.0	26.4	36.7	75.0	118.8	217.5	628.9
Non-Hispanic Black, male									
2003.....	10.1	16.6	21.4	40.3	61.3	101.4	178.5	286.8	639.3
2002.....	10.1	13.2	24.1	36.3	47.4	91.0	175.1	322.9	693.6
2001.....	11.6	15.9	20.6	33.9	50.3	95.4	203.4	309.9	716.1
2000.....	11.4	17.4	25.9	35.3	64.5	107.6	171.5	345.0	750.3
1999.....	13.5	17.3	21.8	34.0	52.1	109.4	161.9	307.3	799.9
Non-Hispanic Black, female									
2003.....	6.9	9.0	11.2	17.5	26.8	49.8	90.2	163.9	477.1
2002.....	5.5	7.9	12.1	18.9	26.2	55.5	93.6	186.3	533.0
2001.....	7.1	6.6	11.1	15.4	23.1	46.8	92.5	179.9	531.7
2000.....	5.8	8.1	11.4	20.8	31.7	56.7	95.2	194.9	537.3
1999.....	6.7	7.2	11.4	20.5	25.4	52.2	92.9	173.0	565.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute lower respiratory infections (J20-J22)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	*	*	*	*	*	*	0.5	0.9	3.6
2002.....	*	*	*	*	*	0.3	0.3	0.9	3.4
2001.....	*	*	*	*	*	*	0.3	0.6	3.3
2000.....	*	*	*	*	*	0.3	0.6	0.8	4.5
1999.....	*	*	*	*	*	0.2	0.6	1.3	6.1

Non-Hispanic 4/, male

2003.....	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	2.9
2001.....	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	*	*	*	0.7	*	3.7
1999.....	*	*	*	*	*	*	0.8	1.7	4.1

Non-Hispanic 4/, female

2003.....	*	*	*	*	*	*	*	0.8	4.1
2002.....	*	*	*	*	*	*	*	0.9	3.6
2001.....	*	*	*	*	*	*	*	0.7	3.7
2000.....	*	*	*	*	*	*	0.5	0.9	4.8
1999.....	*	*	*	*	*	*	0.5	1.1	6.9

Non-Hispanic White, both sexes

2003.....	*	*	*	*	*	*	0.5	0.9	3.8
2002.....	*	*	*	*	*	*	*	0.9	3.5
2001.....	*	*	*	*	*	*	0.3	0.7	3.5
2000.....	*	*	*	*	*	0.3	0.6	0.9	4.7
1999.....	*	*	*	*	*	*	0.7	1.4	6.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute bronchitis and bronchiolitis (J20-J21)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	*	*	*	*	*	*	*	0.5	2.2
2002.....	*	*	*	*	*	*	*	0.5	2.1
2001.....	*	*	*	*	*	*	*	0.4	2.1
2000.....	*	*	*	*	*	*	0.4	0.5	2.6
1999.....	*	*	*	*	*	*	0.4	0.9	3.6

Non-Hispanic 4/, male

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	2.2
1999.....	*	*	*	*	*	*	*	*	2.6

Non-Hispanic 4/, female

2003.....	*	*	*	*	*	*	*	*	2.6
2002.....	*	*	*	*	*	*	*	*	2.2
2001.....	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	*	*	*	*	*	2.7
1999.....	*	*	*	*	*	*	*	0.7	4.0

Non-Hispanic White, both sexes

2003.....	*	*	*	*	*	*	*	0.5	2.3
2002.....	*	*	*	*	*	*	*	0.5	2.1
2001.....	*	*	*	*	*	*	*	0.4	2.2
2000.....	*	*	*	*	*	*	0.4	0.5	2.7
1999.....	*	*	*	*	*	*	0.4	0.9	3.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic lower respiratory diseases (J40-J47)

All origins 3/, both sexes

2003.....	5.9	11.9	27.5	63.9	121.1	210.8	330.0	455.9	635.1
2002.....	5.8	12.0	28.2	60.8	119.0	211.5	330.1	465.7	637.6
2001.....	5.4	11.9	29.4	62.9	124.2	215.4	329.2	452.9	644.7
2000.....	5.8	11.8	29.6	62.5	124.1	218.2	327.8	473.4	648.6
1999.....	5.5	12.0	31.9	66.6	130.4	227.6	340.2	484.5	646.0

All origins 3/, male

2003.....	6.3	12.3	29.4	69.2	132.6	244.0	395.8	569.1	865.9
2002.....	5.9	12.8	29.5	65.9	131.5	246.0	405.6	595.9	894.8
2001.....	5.8	12.3	31.4	67.0	137.1	252.9	406.7	583.5	916.9
2000.....	6.2	12.1	31.8	68.0	139.7	257.8	404.2	628.5	967.9
1999.....	5.7	12.1	34.3	74.6	151.0	278.3	432.4	648.3	996.7

All origins 3/, female

2003.....	5.4	11.5	25.7	59.0	111.2	184.2	283.1	386.7	533.0
2002.....	5.6	11.2	26.9	56.1	108.3	183.8	276.8	386.5	526.0
2001.....	5.1	11.5	27.5	59.1	113.1	185.6	275.0	374.6	530.8
2000.....	5.5	11.5	27.5	57.5	110.7	187.0	274.6	381.9	518.5
1999.....	5.3	11.9	29.7	59.3	112.8	188.0	276.1	388.7	504.6

Hispanic, both sexes

2003.....	2.8	4.4	9.4	17.4	36.9	74.0	129.9	237.7	460.2
2002.....	2.4	4.9	9.0	20.6	38.4	73.1	136.0	242.9	457.3
2001.....	2.4	5.2	9.5	19.9	42.0	73.1	145.3	212.1	480.2
2000.....	2.5	4.4	9.1	20.9	40.0	69.6	143.2	252.3	481.1
1999.....	3.1	4.9	10.4	19.9	45.3	84.6	155.9	265.5	536.0

Hispanic, male

2003.....	2.8	3.7	11.8	23.4	40.9	103.8	170.7	334.7	646.5
2002.....	2.7	5.4	11.7	25.7	51.5	92.2	199.7	321.4	602.4
2001.....	2.3	4.4	10.6	23.6	57.1	99.3	195.1	279.2	676.1
2000.....	2.4	4.8	8.8	30.5	50.7	100.6	191.9	366.6	671.1
1999.....	2.5	5.0	11.6	25.3	52.5	120.7	216.9	385.1	836.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)									
Hispanic, female									
2003.....	2.8	5.1	7.1	12.2	33.6	51.1	100.9	174.6	365.6
2002.....	2.1	4.4	6.4	16.1	27.6	58.5	90.5	192.7	385.5
2001.....	2.6	6.0	8.5	16.7	29.8	53.0	109.9	170.5	384.4
2000.....	2.7	4.0	9.3	12.7	31.4	46.0	108.8	182.8	387.8
1999.....	3.7	4.9	9.4	15.1	39.4	57.5	112.9	194.1	391.0
Non-Hispanic 4/, both sexes									
2003.....	6.2	12.5	28.9	67.2	127.1	219.5	340.6	465.1	641.2
2002.....	6.1	12.6	29.6	63.8	124.6	219.7	339.4	474.4	643.5
2001.....	5.7	12.4	30.9	65.9	129.7	223.3	337.5	461.7	649.4
2000.....	6.1	12.3	31.0	65.3	129.3	226.0	335.4	480.7	653.2
1999.....	5.7	12.5	33.5	69.7	135.5	234.6	347.3	491.4	648.0
Non-Hispanic 4/, male									
2003.....	6.7	13.0	30.7	72.3	138.8	252.6	407.6	579.0	874.6
2002.....	6.2	13.5	30.9	68.8	136.7	254.8	415.0	606.7	906.0
2001.....	6.1	12.9	32.9	69.9	142.2	261.1	416.4	594.6	924.8
2000.....	6.5	12.6	33.4	70.3	144.8	265.6	412.8	637.1	978.0
1999.....	5.9	12.6	35.9	77.7	156.5	285.5	440.8	656.5	1,000.2
Non-Hispanic 4/, female									
2003.....	5.7	12.0	27.2	62.5	116.9	192.9	293.0	395.7	538.6
2002.....	6.0	11.7	28.5	59.2	114.1	191.6	286.1	394.1	530.3
2001.....	5.3	12.0	29.0	62.3	118.9	193.3	282.4	382.2	534.8
2000.....	5.7	12.1	28.8	60.8	115.9	194.7	281.5	388.6	521.9
1999.....	5.5	12.4	31.2	62.4	117.4	194.8	282.3	394.8	507.0
Non-Hispanic White, both sexes									
2003.....	6.0	12.4	29.6	71.3	136.6	233.4	359.4	489.6	671.6
2002.....	5.8	12.4	30.6	67.2	133.4	233.0	357.7	496.4	673.1
2001.....	5.4	12.3	31.8	69.9	137.9	236.4	355.6	483.3	676.9
2000.....	5.5	12.0	31.7	68.8	136.5	238.6	351.8	503.7	680.8
1999.....	5.0	12.3	34.3	73.2	143.2	246.1	361.3	512.6	675.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic White, male

2003.....	6.5	12.8	30.9	75.1	146.5	262.0	419.8	596.6	903.6
2002.....	5.9	13.1	31.1	70.6	143.1	264.3	428.0	624.9	935.4
2001.....	5.9	12.5	33.4	72.5	147.9	270.1	429.6	611.8	951.5
2000.....	5.9	12.1	33.6	72.5	148.9	274.6	425.1	658.0	1,008.6
1999.....	5.2	12.1	36.1	79.7	162.0	293.9	450.4	673.4	1,033.5

Non-Hispanic White, female

2003.....	5.4	11.9	28.4	67.7	127.9	209.9	315.7	423.8	569.5
2002.....	5.8	11.7	30.2	64.0	124.8	207.4	307.4	417.7	559.8
2001.....	4.8	12.1	30.3	67.4	129.1	209.0	303.2	405.8	562.7
2000.....	5.1	11.9	29.9	65.3	125.7	209.7	300.2	412.0	548.7
1999.....	4.8	12.6	32.5	67.2	126.8	208.0	298.8	417.7	532.8

Non-Hispanic Black, both sexes

2003.....	9.4	17.1	32.0	53.7	88.4	149.3	214.6	274.0	354.9
2002.....	9.3	17.6	31.4	54.8	87.4	149.7	217.7	300.6	370.8
2001.....	9.3	16.8	32.3	52.7	95.9	150.7	208.7	279.2	379.3
2000.....	11.3	18.5	33.8	58.1	98.7	152.8	213.1	273.6	376.2
1999.....	11.5	17.5	36.9	58.6	101.6	164.5	245.5	297.3	377.0

Non-Hispanic Black, male

2003.....	10.0	18.2	38.9	68.7	114.4	216.0	331.7	475.4	608.4
2002.....	10.0	20.3	38.4	72.7	118.0	216.2	345.1	479.7	653.1
2001.....	9.1	19.0	37.8	66.7	129.2	224.1	333.2	475.9	682.6
2000.....	11.8	19.6	40.3	74.2	140.2	228.6	328.0	451.4	696.3
1999.....	12.3	20.0	44.3	80.5	141.6	244.1	391.5	524.3	707.2

Non-Hispanic Black, female

2003.....	8.8	16.2	26.4	42.1	69.1	104.3	145.1	168.6	256.7
2002.....	8.7	15.4	25.7	40.7	64.7	105.1	141.6	207.4	263.0
2001.....	9.5	14.9	27.8	41.7	71.4	101.9	134.4	179.1	266.9
2000.....	10.8	17.4	28.4	45.5	68.0	102.5	143.7	184.1	257.3
1999.....	10.9	15.3	31.0	41.5	72.1	111.7	157.5	184.8	253.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Bronchitis, chronic and unspecified (J40-J42)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	*	*	0.1	0.2	0.4	0.8	1.5	2.5	7.3
2002.....	*	*	0.2	0.3	0.5	1.0	1.7	2.9	8.0
2001.....	*	0.1	0.2	0.3	0.6	0.9	1.7	2.6	9.1
2000.....	*	*	*	0.5	0.7	1.0	2.1	3.5	11.8
1999.....	*	0.2	0.3	0.6	0.7	1.3	1.9	4.1	10.7

Non-Hispanic 4/, male

2003.....	*	*	*	*	*	0.8	1.9	2.6	6.9
2002.....	*	*	*	*	0.7	1.2	2.3	2.9	8.8
2001.....	*	*	*	0.4	0.8	1.1	2.1	2.6	9.3
2000.....	*	*	*	0.5	1.0	0.9	2.9	4.8	12.8
1999.....	*	*	0.4	0.6	0.8	1.5	2.5	4.7	11.9

Non-Hispanic 4/, female

2003.....	*	*	*	*	0.5	0.8	1.1	2.5	7.5
2002.....	*	*	*	*	*	0.9	1.4	2.9	7.6
2001.....	*	*	*	*	0.4	0.8	1.4	2.6	9.1
2000.....	*	*	*	0.5	0.5	1.1	1.7	2.8	11.4
1999.....	*	*	*	0.6	0.6	1.1	1.5	3.7	10.2

Non-Hispanic White, both sexes

2003.....	*	*	*	0.2	0.5	0.8	1.6	2.5	7.5
2002.....	*	*	*	0.3	0.6	1.1	1.9	2.9	8.4
2001.....	*	0.1	0.2	0.4	0.6	0.9	1.7	2.7	9.7
2000.....	*	*	*	0.5	0.7	1.0	2.2	3.6	12.6
1999.....	*	*	0.3	0.6	0.7	1.3	1.9	4.1	11.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)									
All origins 3/, both sexes									
2003.....	0.7	1.5	4.0	9.3	16.6	28.5	41.9	51.6	54.6
2002.....	0.7	1.6	4.3	9.1	17.4	31.3	43.2	54.7	59.3
2001.....	0.7	1.7	5.0	10.1	19.8	33.2	46.9	55.5	62.1
2000.....	0.8	1.8	5.0	10.4	20.7	35.5	47.5	60.4	63.3
1999.....	0.7	2.0	5.5	12.1	23.5	38.4	51.3	65.3	62.1
All origins 3/, male									
2003.....	1.0	1.9	4.4	10.3	18.3	34.4	51.9	67.5	79.2
2002.....	0.9	1.9	4.6	10.7	19.7	37.8	56.1	74.9	90.0
2001.....	0.9	2.0	5.8	10.8	22.4	40.3	59.2	74.9	101.2
2000.....	1.0	2.1	5.6	11.2	24.5	42.6	60.4	83.6	104.2
1999.....	0.9	2.2	6.2	13.6	28.0	48.9	67.3	91.5	105.4
All origins 3/, female									
2003.....	0.5	1.2	3.6	8.4	15.2	23.7	34.8	41.9	43.7
2002.....	0.6	1.3	4.0	7.7	15.4	26.1	34.0	42.4	46.0
2001.....	0.5	1.4	4.2	9.5	17.6	27.5	38.3	43.8	45.8
2000.....	0.6	1.5	4.5	9.7	17.5	29.9	38.5	46.8	46.7
1999.....	0.5	1.7	4.9	10.8	19.8	30.2	40.2	49.9	44.7
Hispanic, both sexes									
2003.....	*	*	*	*	3.5	9.1	13.6	30.3	34.7
2002.....	*	*	*	2.4	3.8	10.3	10.9	32.1	36.6
2001.....	*	*	*	*	5.2	8.3	17.3	25.8	39.4
2000.....	*	*	*	*	4.3	8.2	18.4	29.5	31.2
1999.....	*	*	*	*	4.8	10.1	18.1	33.5	44.2
Hispanic, male									
2003.....	*	*	*	*	*	13.2	21.9	41.6	53.9
2002.....	*	*	*	*	6.8	12.3	17.3	43.5	53.5
2001.....	*	*	*	*	*	12.4	28.0	45.4	60.9
2000.....	*	*	*	*	*	12.6	25.8	45.6	58.4
1999.....	*	*	*	*	*	16.8	26.5	55.0	77.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)									
Hispanic, female									
2003.....	*	*	*	*	*	*	*	23.0	25.0
2002.....	*	*	*	*	*	8.8	*	24.8	28.2
2001.....	*	*	*	*	*	*	9.7	*	28.9
2000.....	*	*	*	*	*	*	13.1	19.7	*
1999.....	*	*	*	*	*	*	12.1	20.6	28.1
Non-Hispanic 4/, both sexes									
2003.....	0.8	1.6	4.2	9.8	17.5	29.7	43.4	52.6	55.4
2002.....	0.8	1.7	4.6	9.6	18.3	32.6	44.8	55.6	60.1
2001.....	0.7	1.8	5.3	10.7	20.8	34.6	48.2	56.6	62.8
2000.....	0.8	1.9	5.4	11.0	21.8	36.9	48.7	61.5	64.3
1999.....	0.7	2.0	5.9	12.8	24.7	39.9	52.6	66.3	62.5
Non-Hispanic 4/, male									
2003.....	1.0	2.0	4.7	10.8	19.3	35.7	53.4	68.6	80.4
2002.....	1.0	2.0	4.9	11.3	20.6	39.3	58.0	76.3	91.3
2001.....	0.9	2.1	6.0	11.4	23.4	41.8	60.6	76.0	102.7
2000.....	1.1	2.2	6.0	11.7	25.6	44.0	61.9	85.1	105.9
1999.....	0.9	2.3	6.6	14.2	29.2	50.5	68.9	92.7	106.1
Non-Hispanic 4/, female									
2003.....	0.5	1.3	3.8	8.9	16.0	24.9	36.3	42.8	44.5
2002.....	0.6	1.4	4.3	8.2	16.4	27.2	35.5	43.1	46.6
2001.....	0.5	1.5	4.5	10.1	18.6	28.9	39.6	44.9	46.3
2000.....	0.6	1.6	4.8	10.3	18.5	31.3	39.5	47.7	47.5
1999.....	0.5	1.8	5.2	11.5	20.8	31.6	41.3	50.9	45.0
Non-Hispanic White, both sexes									
2003.....	0.8	1.8	4.5	10.5	19.2	31.9	46.0	55.2	57.9
2002.....	0.8	1.8	4.9	10.4	19.8	34.8	47.7	58.4	63.1
2001.....	0.8	1.8	5.5	11.6	22.7	36.9	51.4	59.9	65.9
2000.....	0.8	1.9	5.6	11.9	23.2	39.5	51.6	64.9	66.9
1999.....	0.7	2.1	6.3	13.8	26.8	42.6	55.3	69.9	65.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)									
Non-Hispanic White, male									
2003.....	1.0	2.0	4.8	11.2	20.7	37.4	54.9	70.8	81.6
2002.....	1.0	2.0	5.1	11.7	21.2	41.1	60.3	78.9	94.3
2001.....	1.0	2.0	6.1	12.0	25.1	43.3	63.4	79.2	106.6
2000.....	0.9	2.1	6.2	12.3	26.3	46.3	64.8	88.6	107.3
1999.....	0.8	2.1	6.8	14.9	30.8	52.4	71.4	95.7	110.0
Non-Hispanic White, female									
2003.....	0.5	1.5	4.2	9.8	17.8	27.4	39.6	45.7	47.4
2002.....	0.6	1.6	4.7	9.2	18.6	29.6	38.8	45.9	49.7
2001.....	0.5	1.5	4.9	11.3	20.6	31.7	42.9	48.3	48.9
2000.....	0.6	1.6	5.1	11.5	20.5	34.1	42.3	50.8	50.6
1999.....	0.5	2.0	5.7	12.9	23.2	34.7	43.9	54.8	47.3
Non-Hispanic Black, both sexes									
2003.....	1.1	1.3	3.7	7.4	10.5	16.7	25.4	31.1	31.2
2002.....	1.1	1.5	3.4	6.1	12.1	21.4	23.7	35.0	32.1
2001.....	*	2.4	5.0	6.3	11.7	20.4	21.8	26.5	35.9
2000.....	1.6	2.6	4.8	7.8	15.3	19.2	24.5	31.0	37.5
1999.....	1.4	2.5	4.7	8.2	14.2	21.1	31.8	31.7	37.8
Non-Hispanic Black, male									
2003.....	*	*	5.4	10.5	13.7	25.3	43.9	51.3	71.4
2002.....	*	2.4	4.6	9.6	20.7	31.5	42.3	61.6	65.6
2001.....	*	3.0	6.9	9.2	15.9	32.4	34.6	49.9	69.3
2000.....	2.5	3.6	6.1	10.8	25.9	29.4	34.7	53.2	89.2
1999.....	2.0	4.2	6.6	12.2	23.1	37.4	49.8	66.4	73.7
Non-Hispanic Black, female									
2003.....	*	*	2.4	4.9	8.1	10.9	14.4	20.5	15.7
2002.....	*	*	*	3.3	5.7	14.7	12.6	21.1	19.2
2001.....	*	*	3.4	4.1	8.7	12.4	14.1	14.6	23.5
2000.....	*	*	3.7	5.4	7.5	12.5	18.3	19.9	18.3
1999.....	*	*	3.1	5.1	7.7	10.3	21.0	14.5	24.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)									
Hispanic, female									
2003.....	*	*	*	*	5.6	*	*	*	17.2
2002.....	*	*	*	*	5.8	*	*	*	*
2001.....	*	2.8	*	*	*	*	*	*	18.6
2000.....	*	*	*	*	6.0	*	*	*	*
1999.....	*	*	*	*	6.9	*	11.0	*	*
Non-Hispanic 4/, both sexes									
2003.....	1.4	1.6	1.8	2.2	2.6	3.2	4.9	7.1	14.1
2002.....	1.4	1.7	2.0	2.1	2.6	3.7	5.4	7.7	14.7
2001.....	1.4	1.7	1.8	2.6	3.0	3.9	5.0	8.1	16.1
2000.....	1.6	1.7	2.0	2.3	3.3	4.5	5.5	8.7	16.6
1999.....	1.6	1.7	2.6	2.2	3.6	4.6	6.1	9.5	17.3
Non-Hispanic 4/, male									
2003.....	1.2	1.1	1.4	1.7	1.7	2.4	3.8	5.5	10.7
2002.....	1.0	1.3	1.5	1.6	2.0	2.8	4.1	6.2	11.1
2001.....	1.0	1.3	1.1	1.9	2.3	2.9	3.7	5.6	12.2
2000.....	1.3	1.2	1.6	1.6	2.6	3.2	3.8	6.7	12.7
1999.....	1.0	1.2	1.9	1.8	2.5	3.8	4.8	7.1	12.6
Non-Hispanic 4/, female									
2003.....	1.7	2.1	2.2	2.7	3.3	3.8	5.7	8.1	15.6
2002.....	1.8	2.0	2.4	2.5	3.1	4.5	6.2	8.5	16.3
2001.....	1.7	2.1	2.5	3.2	3.6	4.7	6.0	9.6	17.7
2000.....	1.9	2.1	2.4	3.0	3.9	5.5	6.7	9.8	18.2
1999.....	2.2	2.3	3.3	2.7	4.5	5.3	7.0	10.9	19.1
Non-Hispanic White, both sexes									
2003.....	1.0	1.1	1.4	1.7	2.1	2.6	4.4	6.6	14.1
2002.....	1.0	1.1	1.4	1.7	2.0	3.1	5.1	7.4	14.7
2001.....	0.9	1.3	1.3	1.8	2.6	3.3	4.5	7.7	15.7
2000.....	1.0	1.2	1.4	1.8	2.5	3.9	5.0	8.4	15.8
1999.....	1.1	1.2	2.0	1.7	2.9	3.9	5.6	9.2	16.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)									
Non-Hispanic White, male									
2003.....	0.8	0.7	1.0	1.2	1.5	1.9	3.0	5.1	10.6
2002.....	0.7	0.9	1.0	1.2	1.5	2.3	3.6	5.8	10.8
2001.....	0.7	0.9	0.8	1.2	1.9	2.3	3.2	5.0	11.4
2000.....	0.9	0.9	1.0	1.1	1.8	2.9	3.5	6.3	11.1
1999.....	0.6	0.8	1.4	1.3	1.9	2.9	4.2	6.9	10.6
Non-Hispanic White, female									
2003.....	1.3	1.4	1.7	2.3	2.8	3.1	5.4	7.5	15.7
2002.....	1.4	1.3	1.7	2.0	2.5	3.8	6.1	8.4	16.4
2001.....	1.2	1.7	1.8	2.4	3.2	4.1	5.5	9.3	17.5
2000.....	1.2	1.4	1.7	2.3	3.2	4.8	6.1	9.7	17.7
1999.....	1.6	1.6	2.5	2.1	3.8	4.6	6.7	10.5	18.8
Non-Hispanic Black, both sexes									
2003.....	4.2	5.8	5.3	5.1	6.8	7.7	8.4	10.6	11.9
2002.....	4.2	5.8	7.3	5.7	6.2	10.3	7.2	9.1	11.8
2001.....	4.4	5.0	6.1	7.9	6.3	9.5	10.3	11.3	16.5
2000.....	5.6	5.3	6.7	7.2	9.1	9.8	10.9	9.8	22.3
1999.....	5.0	5.9	7.5	6.3	8.6	11.2	10.0	11.4	20.7
Non-Hispanic Black, male									
2003.....	3.5	4.6	3.9	5.0	5.1	*	9.3	*	*
2002.....	3.4	4.7	6.1	5.5	6.4	8.9	*	*	*
2001.....	3.8	4.6	3.7	6.1	*	9.6	*	*	*
2000.....	4.1	3.2	6.3	6.0	8.8	6.8	*	*	24.7
1999.....	3.7	4.3	5.2	5.9	6.9	12.1	9.7	*	*
Non-Hispanic Black, female									
2003.....	4.8	6.9	6.4	5.3	8.1	8.9	7.8	11.8	13.6
2002.....	4.8	6.8	8.2	5.9	6.1	11.2	7.8	9.7	12.5
2001.....	5.0	5.3	8.0	9.4	7.5	9.4	11.6	12.1	17.1
2000.....	6.9	7.1	7.1	8.2	9.2	11.8	14.5	9.9	21.4
1999.....	6.0	7.2	9.4	6.7	9.8	10.6	10.2	14.0	21.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)									
All origins 3/, both sexes									
2003.....	3.6	8.6	21.5	52.2	101.4	178.5	281.8	394.6	558.9
2002.....	3.5	8.7	21.8	49.3	98.4	175.4	279.8	400.3	555.5
2001.....	3.3	8.4	22.4	49.8	100.7	177.4	275.4	386.8	557.4
2000.....	3.4	8.2	22.4	49.1	99.2	177.1	272.5	400.7	556.7
1999.....	3.1	8.2	23.5	51.6	102.5	183.2	280.8	405.5	555.8
All origins 3/, male									
2003.....	4.1	9.1	23.3	57.0	112.2	206.5	338.3	493.3	769.0
2002.....	3.8	9.5	23.3	53.2	109.1	204.3	343.0	511.9	785.1
2001.....	3.8	9.0	24.2	54.0	111.6	208.6	341.5	500.4	794.5
2000.....	3.8	8.7	24.4	54.5	111.6	210.9	337.1	533.3	838.0
1999.....	3.7	8.6	25.8	58.7	119.6	223.9	357.9	544.9	866.7
All origins 3/, female									
2003.....	3.2	8.0	19.8	47.8	92.1	156.0	241.5	334.2	466.0
2002.....	3.3	7.9	20.4	45.7	89.3	152.3	235.2	332.4	455.8
2001.....	2.8	7.8	20.7	46.1	91.4	152.7	229.1	318.6	458.2
2000.....	2.9	7.8	20.5	44.3	88.7	150.4	227.6	322.5	442.2
1999.....	2.6	7.8	21.4	45.2	87.9	151.4	227.2	324.0	430.5
Hispanic, both sexes									
2003.....	1.0	2.6	6.0	13.4	29.3	61.9	111.2	195.9	403.7
2002.....	1.1	3.3	6.9	15.1	30.4	59.0	117.3	198.7	397.4
2001.....	1.1	3.0	5.9	15.5	33.6	59.6	118.4	176.8	418.9
2000.....	1.2	2.2	6.6	14.1	29.7	54.5	116.0	209.4	422.7
1999.....	1.3	2.6	7.2	15.3	34.6	68.0	127.8	213.2	462.3
Hispanic, male									
2003.....	*	*	8.0	18.2	35.4	87.1	145.2	278.6	574.2
2002.....	*	3.7	9.9	19.8	43.0	77.2	175.3	268.8	535.4
2001.....	*	2.9	6.6	19.8	47.1	82.3	157.5	227.2	605.7
2000.....	*	*	7.0	22.2	38.8	79.7	159.5	307.7	584.5
1999.....	*	*	8.8	20.6	41.0	98.8	184.1	319.1	735.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)									
Hispanic, female									
2003.....	*	3.2	4.1	9.2	24.3	42.6	87.1	142.1	317.2
2002.....	*	2.9	4.2	11.0	20.2	45.0	75.9	154.0	329.1
2001.....	*	3.2	5.2	11.8	22.7	42.2	90.5	145.6	327.7
2000.....	*	*	6.2	7.2	22.3	35.3	85.2	149.6	343.3
1999.....	*	*	5.8	10.8	29.4	44.8	88.1	150.0	330.7
Non-Hispanic 4/, both sexes									
2003.....	3.9	9.1	22.7	55.0	106.5	185.9	290.8	403.0	564.3
2002.....	3.8	9.1	22.9	51.8	103.1	182.4	287.5	408.2	560.8
2001.....	3.5	8.8	23.6	52.3	105.2	184.0	282.5	394.4	561.3
2000.....	3.5	8.7	23.5	51.5	103.5	183.6	279.1	407.0	560.5
1999.....	3.3	8.6	24.7	54.0	106.6	188.8	286.7	411.5	557.6
Non-Hispanic 4/, male									
2003.....	4.4	9.8	24.4	59.6	117.3	213.8	348.5	502.4	776.7
2002.....	4.0	10.0	24.3	55.6	113.4	211.5	350.7	521.4	794.9
2001.....	4.0	9.4	25.5	56.2	115.6	215.3	350.0	510.4	800.6
2000.....	4.1	9.2	25.7	56.4	115.7	217.5	344.3	540.5	846.6
1999.....	3.9	9.0	27.0	61.1	124.1	229.6	364.7	552.0	869.6
Non-Hispanic 4/, female									
2003.....	3.4	8.4	21.0	50.8	97.1	163.4	249.8	342.3	471.0
2002.....	3.5	8.3	21.7	48.3	94.3	159.0	243.0	339.5	459.8
2001.....	3.0	8.2	21.9	48.7	96.2	159.0	235.3	325.1	461.7
2000.....	3.0	8.2	21.5	47.0	93.0	156.8	233.7	328.4	444.8
1999.....	2.7	8.2	22.5	47.5	91.6	156.9	232.5	329.4	432.7
Non-Hispanic White, both sexes									
2003.....	4.1	9.4	23.6	58.8	114.8	198.1	307.5	425.2	592.2
2002.....	3.9	9.4	24.2	54.9	111.0	194.0	303.0	427.6	586.9
2001.....	3.6	9.1	24.8	56.0	112.0	195.4	298.0	413.0	585.7
2000.....	3.6	8.9	24.5	54.6	110.1	194.1	292.9	426.8	585.5
1999.....	3.2	8.9	25.8	57.0	112.9	198.4	298.5	429.4	582.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)									
Non-Hispanic White, male									
2003.....	4.6	10.0	24.9	62.4	123.9	222.0	360.0	518.3	804.2
2002.....	4.1	10.1	24.9	57.4	119.7	219.6	361.6	537.3	821.0
2001.....	4.1	9.5	26.2	58.9	120.0	223.5	361.0	525.0	823.9
2000.....	4.0	9.0	26.3	58.6	119.8	224.5	353.8	558.0	876.5
1999.....	3.8	9.1	27.5	63.0	128.5	237.0	372.4	566.3	900.5
Non-Hispanic White, female									
2003.....	3.6	9.0	22.4	55.4	106.8	178.5	269.5	368.0	498.8
2002.....	3.8	8.8	23.6	52.6	103.3	173.0	261.1	360.4	485.7
2001.....	3.0	8.7	23.5	53.4	104.9	172.5	253.3	345.3	486.6
2000.....	3.2	8.7	22.9	51.0	101.5	169.7	250.1	348.8	468.2
1999.....	2.6	8.8	24.1	51.5	99.2	167.5	246.7	348.6	456.1
Non-Hispanic Black, both sexes									
2003.....	3.9	9.7	22.6	41.0	70.9	124.3	180.3	229.9	306.1
2002.....	3.9	10.2	20.5	42.8	68.7	117.7	186.1	254.1	322.7
2001.....	4.2	9.4	21.2	38.2	77.1	119.8	175.2	240.6	323.4
2000.....	4.0	10.3	22.2	42.9	73.5	122.6	175.9	229.9	312.6
1999.....	5.0	8.6	24.5	43.7	78.4	130.6	201.5	250.1	314.0
Non-Hispanic Black, male									
2003.....	4.6	11.4	29.4	53.0	95.7	184.0	277.1	412.8	526.3
2002.....	4.9	12.9	27.5	57.4	90.1	175.5	295.3	408.7	572.1
2001.....	4.5	11.3	27.2	51.0	107.6	180.8	288.7	414.6	592.5
2000.....	5.2	12.7	27.9	56.8	105.0	191.0	286.6	386.2	578.9
1999.....	6.2	11.0	32.1	62.0	111.3	192.5	329.2	442.8	610.9
Non-Hispanic Black, female									
2003.....	3.2	8.3	17.1	31.7	52.6	84.1	122.9	134.3	220.8
2002.....	3.1	7.9	14.8	31.3	52.8	78.9	120.8	173.7	227.4
2001.....	3.9	7.8	16.2	28.1	54.6	79.1	107.5	152.0	223.7
2000.....	2.9	8.3	17.5	31.9	50.3	77.1	109.2	151.2	213.7
1999.....	3.9	6.6	18.3	29.4	53.9	89.6	124.6	154.5	203.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumoconioses and chemical effects (J60-J66,J68)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	*	*	0.2	0.4	1.0	1.8	3.0	5.0	6.6
2002.....	*	*	0.2	0.4	0.8	1.7	3.6	4.3	6.9
2001.....	*	*	0.2	0.3	1.0	2.0	3.5	5.3	6.2
2000.....	*	*	0.2	0.4	0.9	2.3	3.5	4.7	7.6
1999.....	*	*	*	0.4	1.1	2.0	3.5	5.3	6.2

Non-Hispanic 4/, male

2003.....	*	*	0.3	0.7	2.1	3.6	7.1	12.7	20.7
2002.....	*	*	0.3	0.8	1.7	3.6	8.4	10.7	21.9
2001.....	*	*	0.4	0.7	2.2	4.5	8.1	13.6	20.5
2000.....	*	*	*	0.7	1.8	5.0	8.1	12.2	25.7
1999.....	*	*	*	0.8	2.3	4.5	8.4	14.0	21.1

Non-Hispanic 4/, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic White, both sexes

2003.....	*	*	0.2	0.4	1.1	1.9	3.2	5.3	7.1
2002.....	*	*	0.2	0.4	0.9	1.8	3.8	4.5	7.3
2001.....	*	*	0.2	0.3	1.1	2.2	3.8	5.5	6.6
2000.....	*	*	0.2	0.4	1.0	2.5	3.6	4.9	8.1
1999.....	*	*	*	0.3	1.2	2.1	3.8	5.7	6.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)									
All origins 3/, both sexes									
2003.....	0.9	1.1	2.0	3.8	6.3	13.3	30.1	65.4	176.2
2002.....	0.7	1.1	1.8	3.4	6.5	13.4	31.3	67.3	186.0
2001.....	0.7	1.3	1.8	3.6	6.2	14.1	30.4	68.0	189.4
2000.....	0.8	1.2	1.9	3.2	6.7	14.2	30.4	65.7	187.6
1999.....	0.7	0.9	2.0	3.1	6.1	13.2	28.2	60.7	175.6
All origins 3/, male									
2003.....	1.0	1.4	2.6	4.8	8.6	17.8	41.5	92.3	246.2
2002.....	0.9	1.5	2.2	4.2	8.6	18.2	43.4	95.1	268.5
2001.....	0.8	1.7	2.3	4.2	8.2	19.1	41.0	98.1	270.3
2000.....	1.0	1.5	2.4	3.9	9.2	20.2	42.4	95.2	276.0
1999.....	0.9	1.0	2.4	4.2	8.3	18.4	41.6	90.1	258.4
All origins 3/, female									
2003.....	0.7	0.8	1.5	2.9	4.3	9.7	22.1	49.0	145.3
2002.....	0.5	0.8	1.4	2.8	4.8	9.6	22.8	50.4	150.2
2001.....	0.6	0.9	1.4	3.0	4.4	10.2	23.0	49.9	155.6
2000.....	0.6	0.9	1.5	2.5	4.6	9.4	22.1	48.2	151.6
1999.....	0.5	0.8	1.6	2.0	4.2	9.2	18.8	43.5	142.2
Hispanic, both sexes									
2003.....	*	*	1.6	*	4.2	9.5	15.9	34.0	113.0
2002.....	*	*	*	2.7	3.7	7.3	19.5	45.0	131.4
2001.....	*	*	*	*	4.3	8.3	19.0	42.2	161.3
2000.....	*	*	*	*	3.5	9.0	20.8	40.7	137.4
1999.....	*	*	*	*	3.6	7.9	18.4	35.8	130.5
Hispanic, male									
2003.....	*	*	*	*	*	12.0	20.7	52.0	123.1
2002.....	*	*	*	*	*	*	26.9	72.1	167.3
2001.....	*	*	*	*	*	11.4	25.2	57.1	220.9
2000.....	*	*	*	*	*	12.2	32.5	63.3	169.3
1999.....	*	*	*	*	*	*	26.5	48.7	159.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)									
Hispanic, female									
2003.....	*	*	*	*	*	7.5	12.5	22.3	107.8
2002.....	*	*	*	*	*	6.8	14.2	27.7	113.6
2001.....	*	*	*	*	*	14.5	33.0	132.2	
2000.....	*	*	*	*	*	*	12.5	26.9	121.7
1999.....	*	*	*	*	*	*	12.7	28.1	116.5
Non-Hispanic 4/, both sexes									
2003.....	0.9	1.1	2.0	3.9	6.4	13.5	30.8	66.8	178.5
2002.....	0.7	1.2	1.9	3.5	6.7	13.7	31.9	68.1	187.8
2001.....	0.7	1.3	1.8	3.7	6.3	14.4	30.9	68.8	190.0
2000.....	0.8	1.2	1.9	3.2	7.0	14.5	30.8	66.4	189.0
1999.....	0.7	1.0	2.1	3.1	6.2	13.4	28.5	61.4	176.8
Non-Hispanic 4/, male									
2003.....	1.1	1.4	2.6	4.9	8.7	18.1	42.5	94.0	251.5
2002.....	0.9	1.6	2.2	4.2	8.8	18.7	44.2	95.8	272.1
2001.....	0.8	1.7	2.3	4.4	8.4	19.5	41.7	99.6	271.6
2000.....	1.0	1.5	2.3	4.0	9.6	20.6	42.6	96.0	279.7
1999.....	0.9	1.0	2.5	4.3	8.5	18.7	42.1	91.4	261.6
Non-Hispanic 4/, female									
2003.....	0.7	0.8	1.5	3.0	4.4	9.8	22.5	50.2	146.5
2002.....	0.5	0.8	1.5	2.9	4.9	9.7	23.2	51.4	151.4
2001.....	0.6	0.9	1.4	3.1	4.5	10.4	23.4	50.4	156.0
2000.....	0.6	1.0	1.5	2.5	4.7	9.6	22.6	48.9	152.3
1999.....	0.5	0.9	1.7	2.0	4.3	9.3	19.0	43.8	142.9
Non-Hispanic White, both sexes									
2003.....	0.9	1.2	2.0	3.5	6.0	13.3	30.4	66.3	181.0
2002.....	0.7	1.1	1.8	3.3	6.3	13.4	31.5	69.0	191.3
2001.....	0.7	1.2	1.8	3.6	6.1	13.9	30.7	69.0	191.2
2000.....	0.7	1.1	1.8	3.0	6.6	13.6	30.5	66.3	192.5
1999.....	0.6	0.9	1.9	3.0	5.9	12.7	27.9	60.9	179.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)									
Non-Hispanic White, male									
2003.....	1.0	1.5	2.5	4.3	8.4	17.5	41.6	93.4	256.5
2002.....	0.9	1.4	2.1	4.1	8.2	18.0	43.3	97.3	278.4
2001.....	0.9	1.6	2.3	4.1	8.1	18.8	41.3	100.4	276.9
2000.....	0.9	1.3	2.3	3.6	8.8	19.0	41.7	96.9	288.1
1999.....	0.8	0.9	2.2	4.0	7.9	17.4	40.9	90.2	267.4
Non-Hispanic White, female									
2003.....	0.7	0.8	1.5	2.7	4.0	9.8	22.3	49.7	147.8
2002.....	0.5	0.7	1.5	2.6	4.6	9.6	23.0	51.8	153.7
2001.....	0.5	0.9	1.4	3.1	4.3	9.8	23.2	50.1	155.5
2000.....	0.5	0.9	1.4	2.3	4.6	9.3	22.6	48.2	153.9
1999.....	0.4	0.8	1.5	2.0	4.2	8.9	18.8	43.7	144.2
Non-Hispanic Black, both sexes									
2003.....	1.3	1.3	2.7	8.6	11.1	17.4	40.7	77.5	165.7
2002.....	0.8	2.2	2.4	5.6	12.0	19.2	42.7	68.3	167.3
2001.....	1.0	2.5	2.5	6.1	9.5	22.7	39.2	76.2	187.3
2000.....	1.7	2.3	3.0	5.9	11.5	25.5	39.1	72.4	160.6
1999.....	1.3	1.8	4.0	5.2	10.2	24.0	40.9	71.6	168.5
Non-Hispanic Black, male									
2003.....	1.8	*	3.9	11.7	14.7	26.6	60.7	110.2	215.2
2002.....	*	3.2	3.7	6.5	17.1	29.1	63.4	92.0	237.4
2001.....	*	3.5	3.2	8.4	12.8	31.3	56.9	107.9	243.7
2000.....	2.4	3.2	3.4	7.4	17.9	42.0	62.1	97.8	210.2
1999.....	*	*	5.2	8.3	15.7	37.7	65.3	111.6	229.4
Non-Hispanic Black, female									
2003.....	*	*	*	6.2	8.4	11.1	28.9	60.4	146.5
2002.....	*	*	*	4.9	8.2	12.6	30.3	56.0	140.5
2001.....	*	*	*	4.3	7.1	17.0	28.6	60.0	166.4
2000.....	*	*	*	4.7	6.9	14.5	25.3	59.6	142.2
1999.....	*	*	3.0	*	6.1	14.9	26.2	51.8	145.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All origins 3/, both sexes

2003.....	2.6	4.0	6.6	11.6	21.0	36.2	57.3	83.8	138.1
2002.....	2.5	4.1	7.0	11.9	20.6	37.6	56.8	83.7	133.5
2001.....	2.6	4.4	7.3	12.6	22.8	38.0	61.8	89.6	142.7
2000.....	2.6	4.6	7.8	12.7	22.9	36.9	60.5	89.2	141.7
1999.....	2.7	4.1	6.9	12.7	22.5	36.8	59.5	87.6	147.6

All origins 3/, male

2003.....	2.8	4.3	7.5	13.8	25.9	43.8	71.8	105.5	165.0
2002.....	2.5	4.3	7.9	13.9	24.4	47.8	70.5	104.2	156.6
2001.....	2.9	4.9	7.7	15.0	27.1	47.0	76.2	110.0	169.1
2000.....	2.8	5.0	8.8	14.6	26.8	45.7	77.3	112.4	179.7
1999.....	2.8	4.4	7.4	14.7	27.3	46.4	74.9	115.0	182.3

All origins 3/, female

2003.....	2.5	3.7	5.7	9.6	16.8	30.0	47.0	70.5	126.2
2002.....	2.4	3.9	6.2	10.2	17.3	29.4	47.0	71.1	123.4
2001.....	2.3	4.0	6.8	10.5	19.1	30.8	51.7	77.3	131.6
2000.....	2.4	4.2	6.8	10.8	19.6	29.9	48.8	75.6	126.2
1999.....	2.6	3.8	6.5	10.9	18.4	29.3	48.8	71.6	133.6

Hispanic, both sexes

2003.....	2.0	3.0	7.5	10.8	18.5	34.3	48.9	73.4	118.7
2002.....	2.2	3.4	6.2	11.1	18.7	35.3	49.3	72.6	112.0
2001.....	2.2	4.3	7.9	10.3	19.8	32.1	52.7	87.9	134.4
2000.....	2.2	3.1	8.6	11.2	20.5	31.4	53.3	84.1	135.4
1999.....	2.3	4.3	5.8	14.3	22.6	30.2	56.2	78.1	134.0

Hispanic, male

2003.....	2.3	4.2	8.9	14.8	21.8	42.7	56.5	74.8	127.8
2002.....	2.2	3.2	7.9	12.6	24.4	47.8	49.9	89.2	108.8
2001.....	2.5	4.7	8.6	13.3	23.4	38.9	64.8	103.9	167.6
2000.....	2.5	3.6	9.6	15.0	22.7	37.9	67.9	107.5	133.0
1999.....	2.6	3.7	7.2	17.3	26.7	35.6	71.0	105.3	161.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Hispanic, female									
2003.....	*	*	6.2	7.3	15.7	27.9	43.5	72.4	114.1
2002.....	2.1	3.6	4.7	9.8	14.1	25.7	48.9	62.0	113.6
2001.....	*	4.0	7.2	7.8	16.9	26.9	44.1	78.0	118.2
2000.....	*	*	7.7	7.9	18.7	26.5	42.9	69.9	136.5
1999.....	*	4.9	4.6	11.8	19.4	26.0	45.7	61.9	120.6
Non-Hispanic 4/, both sexes									
2003.....	2.7	4.1	6.5	11.6	21.2	36.2	57.7	84.2	138.7
2002.....	2.5	4.1	7.1	11.9	20.6	37.7	57.1	84.0	134.1
2001.....	2.6	4.4	7.2	12.8	23.0	38.2	62.1	89.4	142.7
2000.....	2.7	4.7	7.6	12.7	23.0	37.0	60.7	89.1	141.6
1999.....	2.7	4.1	6.9	12.5	22.4	37.0	59.5	87.7	147.7
Non-Hispanic 4/, male									
2003.....	2.9	4.3	7.3	13.6	26.2	43.8	72.5	106.9	166.5
2002.....	2.5	4.3	7.8	13.9	24.3	47.7	71.6	104.7	158.5
2001.....	2.9	4.9	7.6	15.1	27.3	47.2	76.5	110.0	169.1
2000.....	2.9	5.1	8.7	14.6	27.0	46.0	77.5	112.3	181.3
1999.....	2.8	4.5	7.3	14.5	27.2	46.8	74.8	114.8	182.7
Non-Hispanic 4/, female									
2003.....	2.5	3.8	5.7	9.8	16.8	30.1	47.1	70.4	126.4
2002.....	2.4	3.9	6.4	10.1	17.4	29.6	46.9	71.4	123.6
2001.....	2.4	3.9	6.8	10.7	19.2	31.0	52.0	77.1	131.8
2000.....	2.4	4.3	6.6	11.1	19.6	30.0	49.0	75.5	125.5
1999.....	2.6	3.7	6.6	10.8	18.2	29.4	48.8	71.8	133.7
Non-Hispanic White, both sexes									
2003.....	2.3	3.7	6.1	11.5	21.6	37.3	58.9	87.9	143.7
2002.....	2.0	3.8	6.7	11.6	21.0	38.2	59.0	85.8	138.6
2001.....	2.2	3.8	6.9	12.5	23.2	39.0	63.8	92.2	145.9
2000.....	2.2	4.4	7.4	12.5	22.9	37.9	62.1	91.0	145.7
1999.....	2.3	3.7	6.4	12.4	22.5	37.2	60.6	89.6	150.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic White, male

2003.....	2.5	4.0	6.9	13.5	26.6	45.1	73.5	111.4	172.0
2002.....	2.1	4.0	7.5	13.4	25.2	48.1	73.6	107.1	164.1
2001.....	2.5	4.4	7.5	14.5	27.7	48.6	78.9	113.3	173.4
2000.....	2.4	4.7	8.4	14.2	27.0	46.6	79.0	114.9	185.9
1999.....	2.3	4.2	6.5	14.4	27.6	47.4	76.5	117.4	186.1

Non-Hispanic White, female

2003.....	2.0	3.5	5.4	9.7	17.2	31.0	48.3	73.4	131.2
2002.....	2.0	3.6	5.9	9.9	17.2	30.2	48.5	72.8	127.6
2001.....	2.0	3.3	6.3	10.6	19.1	31.2	53.1	79.4	134.5
2000.....	2.1	4.0	6.4	10.8	19.3	30.8	50.3	76.9	129.5
1999.....	2.2	3.3	6.3	10.6	18.0	29.1	49.4	73.2	136.6

Non-Hispanic Black, both sexes

2003.....	5.8	7.5	10.6	14.5	20.6	29.7	51.7	50.8	91.0
2002.....	5.6	6.9	11.1	17.2	21.2	34.5	41.4	68.9	90.7
2001.....	5.9	9.5	11.2	17.4	24.3	36.1	50.3	67.0	110.8
2000.....	6.1	7.9	11.8	15.9	28.2	33.5	50.2	71.8	99.8
1999.....	6.0	7.3	12.1	15.6	25.2	38.7	53.6	66.9	112.8

Non-Hispanic Black, male

2003.....	5.7	7.4	12.9	18.7	25.8	38.8	70.6	66.4	116.1
2002.....	5.8	7.5	12.3	21.8	22.8	45.5	54.5	88.1	102.8
2001.....	6.4	9.6	10.8	22.2	27.9	39.6	59.7	85.0	131.7
2000.....	7.4	8.7	13.7	18.1	32.2	41.3	68.9	83.3	135.0
1999.....	6.8	7.9	14.1	18.1	28.7	44.7	67.2	85.0	146.2

Non-Hispanic Black, female

2003.....	5.9	7.5	8.7	11.3	16.7	23.6	40.5	42.6	81.3
2002.....	5.4	6.3	10.2	13.5	20.1	27.1	33.6	58.8	86.1
2001.....	5.5	9.4	11.5	13.6	21.7	33.7	44.7	57.9	103.1
2000.....	5.1	7.2	10.3	14.3	25.2	28.4	38.9	66.1	86.8
1999.....	5.3	6.8	10.4	13.7	22.6	34.7	45.4	58.0	100.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Peptic ulcer (K25-K28)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	20.3
2002.....	*	*	*	*	*	*	*	*	17.4
2001.....	*	*	*	*	*	*	*	*	25.1
2000.....	*	*	*	*	*	*	*	22.4	29.7
1999.....	*	*	*	*	*	*	*	*	22.9

Non-Hispanic 4/, both sexes

2003.....	0.6	0.8	1.1	1.7	2.3	4.1	7.3	13.5	26.7
2002.....	0.6	0.8	1.4	1.7	2.6	4.8	8.0	13.2	28.6
2001.....	0.7	0.8	1.3	2.2	3.1	4.8	8.3	16.0	33.6
2000.....	0.7	1.0	1.5	2.2	3.2	5.1	8.8	15.5	35.7
1999.....	0.6	0.9	1.4	2.2	3.3	5.9	9.1	17.2	34.8

Non-Hispanic 4/, male

2003.....	0.7	1.1	1.4	2.3	3.0	5.2	8.8	16.1	29.6
2002.....	0.6	1.2	1.7	2.0	3.0	6.1	9.3	16.8	31.7
2001.....	0.9	1.1	1.8	2.8	4.1	5.6	10.2	16.8	35.5
2000.....	0.9	1.6	2.1	2.8	3.9	6.9	11.3	18.1	41.0
1999.....	0.7	1.2	1.7	3.0	4.5	8.0	10.9	21.1	37.7

Non-Hispanic 4/, female

2003.....	0.4	0.5	0.9	1.1	1.7	3.3	6.3	11.9	25.4
2002.....	0.5	0.5	1.0	1.4	2.2	3.7	7.1	11.0	27.3
2001.....	0.5	0.6	0.8	1.6	2.3	4.3	7.0	15.5	32.8
2000.....	0.5	0.5	1.0	1.7	2.5	3.6	7.0	14.0	33.5
1999.....	0.5	0.5	1.1	1.5	2.3	4.3	7.9	14.9	33.7

Non-Hispanic White, both sexes

2003.....	0.5	0.7	1.1	1.6	2.3	4.2	7.2	13.3	27.3
2002.....	0.5	0.8	1.2	1.6	2.5	4.8	7.9	13.7	29.4
2001.....	0.6	0.7	1.1	2.1	3.0	4.8	8.3	16.3	34.7
2000.....	0.6	0.9	1.4	2.2	3.1	5.1	8.7	15.9	36.3
1999.....	0.5	0.7	1.3	2.2	3.1	5.8	9.2	17.2	35.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Peptic ulcer (K25-K28)

Non-Hispanic White, male

2003.....	0.6	0.9	1.2	2.2	2.9	5.1	8.5	15.5	29.8
2002.....	0.6	1.0	1.5	1.9	2.9	6.0	9.2	17.3	32.6
2001.....	0.8	0.9	1.5	2.7	3.9	5.3	9.9	16.9	36.3
2000.....	0.8	1.3	2.0	2.8	3.8	6.7	11.0	18.1	41.0
1999.....	0.6	1.1	1.5	2.9	4.1	7.9	11.0	21.1	38.6

Non-Hispanic White, female

2003.....	0.3	0.5	0.9	1.1	1.8	3.4	6.2	12.0	26.2
2002.....	0.5	0.5	1.0	1.3	2.2	3.9	7.1	11.5	27.9
2001.....	0.5	0.6	0.8	1.5	2.3	4.5	7.1	15.9	34.0
2000.....	0.4	0.5	0.9	1.7	2.4	3.7	7.0	14.5	34.4
1999.....	0.4	0.4	1.1	1.6	2.2	4.3	7.9	14.9	34.4

Non-Hispanic Black, both sexes

2003.....	1.2	1.8	2.0	2.4	2.2	4.5	8.7	14.8	17.3
2002.....	0.9	1.6	2.4	2.9	3.3	5.4	8.8	6.7	18.8
2001.....	1.1	1.6	2.5	3.1	3.5	5.6	8.0	10.5	20.9
2000.....	1.4	2.1	2.6	2.7	4.2	5.7	9.4	11.2	26.7
1999.....	1.3	1.9	2.5	3.2	5.1	6.5	7.6	16.4	25.5

Non-Hispanic Black, male

2003.....	1.7	3.0	3.0	4.0	*	7.1	11.7	24.1	*
2002.....	*	2.9	3.7	4.3	*	8.2	11.7	*	*
2001.....	1.9	2.8	4.0	4.6	5.2	9.3	10.9	*	24.3
2000.....	*	3.4	3.6	*	*	10.3	12.5	*	31.7
1999.....	1.9	2.9	3.8	5.2	7.7	9.0	*	18.6	28.5

Non-Hispanic Black, female

2003.....	*	*	*	*	*	*	6.9	9.9	16.1
2002.....	*	*	*	*	*	*	7.0	*	18.4
2001.....	*	*	*	*	*	*	6.2	9.2	19.7
2000.....	*	*	*	*	*	*	7.6	8.6	24.9
1999.....	*	*	*	*	*	4.8	7.0	15.4	24.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of appendix (K35-K38)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.1	0.1	*	0.2	0.3	0.4	0.8	1.4	1.6
2002.....	0.1	0.2	0.2	0.3	0.3	0.7	0.8	1.1	2.1
2001.....	0.1	*	0.2	0.2	0.4	0.6	0.5	1.0	2.2
2000.....	*	0.2	0.2	0.3	0.3	0.5	0.7	1.4	1.7
1999.....	0.1	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.5

Non-Hispanic 4/, male

2003.....	*	*	*	*	*	0.7	0.9	2.3	2.2
2002.....	*	*	*	0.4	0.6	0.9	1.1	1.7	2.3
2001.....	*	*	*	*	0.5	0.8	0.7	1.5	3.0
2000.....	*	*	*	0.5	*	0.7	1.0	1.6	2.0
1999.....	*	*	*	*	*	*	0.9	*	3.0

Non-Hispanic 4/, female

2003.....	*	*	*	*	*	*	0.7	0.9	1.4
2002.....	*	*	*	*	*	0.5	0.6	0.8	2.0
2001.....	*	*	*	*	*	0.5	*	0.7	1.8
2000.....	*	*	*	*	*	*	0.6	1.2	1.5
1999.....	*	*	*	*	*	*	*	1.1	0.9

Non-Hispanic White, both sexes

2003.....	*	*	*	*	0.3	0.4	0.7	1.5	1.7
2002.....	*	0.2	0.2	0.3	0.3	0.6	0.8	1.2	2.1
2001.....	*	*	*	*	0.3	0.7	0.5	0.9	2.1
2000.....	*	*	*	0.3	0.3	0.4	0.6	1.3	1.8
1999.....	*	0.2	*	*	0.3	0.4	0.6	1.1	1.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hernia (K40-K46)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.2	0.3	0.4	0.7	1.0	1.6	2.7	4.8	12.1
2002.....	0.2	0.2	0.4	0.6	0.9	1.6	2.8	5.2	13.0
2001.....	0.1	0.3	0.4	0.6	0.9	1.5	2.4	5.0	12.5
2000.....	0.1	0.3	0.4	0.6	0.9	1.4	2.9	4.9	13.1
1999.....	*	0.2	0.3	0.7	0.8	1.5	2.5	5.4	12.7

Non-Hispanic 4/, male

2003.....	0.2	0.4	0.4	0.8	1.0	1.7	3.0	5.2	12.5
2002.....	*	*	0.4	0.5	0.9	1.7	2.7	5.1	11.9
2001.....	*	0.3	0.4	0.8	0.9	1.4	2.4	5.2	12.1
2000.....	*	0.4	*	0.6	0.9	1.3	2.9	5.6	12.8
1999.....	*	*	0.4	0.7	0.8	1.4	2.5	6.5	13.3

Non-Hispanic 4/, female

2003.....	*	0.2	0.5	0.7	1.0	1.6	2.4	4.5	12.0
2002.....	*	0.3	0.4	0.7	0.9	1.5	2.9	5.3	13.4
2001.....	*	0.3	0.5	0.6	0.8	1.5	2.5	4.8	12.7
2000.....	*	0.3	0.5	0.5	0.8	1.5	3.0	4.5	13.2
1999.....	*	*	*	0.6	0.8	1.6	2.5	4.7	12.4

Non-Hispanic White, both sexes

2003.....	0.2	0.3	0.4	0.7	1.0	1.6	2.8	4.7	12.4
2002.....	0.1	0.2	0.4	0.5	0.8	1.6	3.0	5.3	13.6
2001.....	*	0.3	0.4	0.6	0.9	1.5	2.4	5.0	12.8
2000.....	*	0.3	0.4	0.6	0.8	1.5	2.9	5.0	13.3
1999.....	*	0.2	0.3	0.7	0.7	1.5	2.5	5.5	13.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

All origins 3/, both sexes

2003.....	15.9	21.1	22.0	24.4	28.0	31.2	31.0	28.5	20.1
2002.....	16.0	20.3	21.7	24.5	27.9	31.2	32.5	29.9	21.4
2001.....	17.1	20.1	21.1	24.7	28.9	31.1	31.3	28.7	22.2
2000.....	16.8	18.7	21.9	26.2	28.9	30.8	31.9	29.6	23.1
1999.....	16.4	18.6	21.4	26.5	29.7	31.6	32.6	30.8	23.2

All origins 3/, male

2003.....	23.4	32.3	32.4	35.6	38.4	40.0	38.1	35.0	27.5
2002.....	23.3	31.0	32.0	34.5	37.2	40.0	39.4	36.9	28.4
2001.....	25.2	30.1	31.5	35.3	39.7	40.8	37.6	35.3	32.0
2000.....	25.4	28.6	31.7	37.7	39.3	40.7	40.7	34.6	34.3
1999.....	25.3	28.8	32.5	37.2	41.8	41.5	40.1	38.7	33.1

All origins 3/, female

2003.....	8.6	10.3	12.2	14.2	19.0	24.1	26.0	24.5	16.8
2002.....	8.9	10.1	12.1	15.3	19.8	24.1	27.6	25.6	18.4
2001.....	9.3	10.5	11.4	15.2	19.6	23.5	26.9	24.7	18.1
2000.....	8.4	9.1	12.7	15.7	19.9	23.1	25.8	26.6	18.5
1999.....	7.8	8.8	11.1	16.8	19.4	23.9	27.4	26.2	19.3

Hispanic, both sexes

2003.....	21.4	28.1	35.6	42.3	47.2	56.0	66.8	54.1	31.1
2002.....	23.1	31.9	37.1	40.1	50.5	63.2	52.8	54.8	45.5
2001.....	25.0	31.9	35.9	43.2	51.1	58.6	62.0	57.6	36.9
2000.....	24.8	30.4	39.6	43.7	55.6	61.8	65.8	60.2	48.4
1999.....	25.2	28.4	35.6	43.2	53.6	59.7	60.4	57.6	51.2

Hispanic, male

2003.....	35.4	46.6	56.7	70.1	59.8	63.2	79.6	57.2	35.4
2002.....	38.0	52.4	61.1	58.5	76.5	79.4	60.8	70.9	45.2
2001.....	41.9	52.5	58.9	70.6	78.0	75.9	81.8	64.9	41.9
2000.....	41.2	51.5	64.7	67.9	86.9	78.8	89.3	61.8	60.5
1999.....	42.3	46.9	59.4	66.7	76.5	82.5	72.6	67.6	66.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)									
Hispanic, female									
2003.....	7.0	10.0	16.2	18.0	36.8	50.5	57.8	52.1	28.9
2002.....	7.9	12.1	15.0	24.2	29.3	50.7	47.1	44.5	45.6
2001.....	8.0	12.2	15.0	19.5	29.3	45.4	47.9	53.1	34.4
2000.....	8.4	10.3	16.9	22.8	30.2	49.0	49.1	59.1	42.5
1999.....	8.2	10.8	14.2	22.8	35.0	42.5	51.8	51.6	43.7
Non-Hispanic 4/, both sexes									
2003.....	15.2	20.3	20.8	22.8	26.5	29.5	29.0	27.2	19.6
2002.....	15.1	19.1	20.3	23.1	26.1	29.0	31.3	28.7	20.4
2001.....	16.3	19.0	19.8	23.2	27.2	29.4	29.7	27.4	21.6
2000.....	15.9	17.5	20.4	24.7	26.9	29.0	30.3	28.3	22.1
1999.....	15.5	17.6	20.3	25.2	28.0	30.0	31.3	29.8	22.1
Non-Hispanic 4/, male									
2003.....	21.9	30.7	30.3	32.7	36.7	38.4	35.7	33.9	27.1
2002.....	21.5	28.8	29.4	32.4	34.3	37.4	38.1	35.3	27.7
2001.....	23.4	28.0	29.1	32.4	36.8	38.6	35.4	34.1	31.4
2000.....	23.7	26.4	29.1	35.3	36.0	38.5	38.4	33.4	33.0
1999.....	23.6	27.1	30.3	35.0	39.4	39.2	38.5	37.6	31.6
Non-Hispanic 4/, female									
2003.....	8.7	10.3	11.8	13.9	17.6	22.3	24.2	23.2	16.3
2002.....	9.0	9.9	11.8	14.6	19.0	22.3	26.6	24.7	17.3
2001.....	9.3	10.3	11.1	14.8	18.8	22.0	25.7	23.4	17.5
2000.....	8.4	9.0	12.3	15.1	19.1	21.5	24.7	25.3	17.7
1999.....	7.8	8.5	10.9	16.3	18.3	22.9	26.3	25.3	18.3
Non-Hispanic White, both sexes									
2003.....	15.3	20.0	20.5	22.8	27.1	30.8	30.7	28.5	20.3
2002.....	15.0	18.7	20.1	23.1	26.9	30.2	32.9	29.6	21.2
2001.....	16.0	18.1	19.1	23.5	27.7	30.2	30.9	28.6	22.5
2000.....	15.7	16.7	19.9	24.7	27.2	29.7	31.6	29.1	22.8
1999.....	14.6	17.2	19.4	24.7	28.3	30.8	32.2	30.9	22.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic White, male

2003.....	22.2	30.1	29.5	32.5	37.5	40.0	37.6	34.9	27.9
2002.....	21.6	28.2	29.1	32.3	34.8	38.7	40.2	36.3	28.7
2001.....	23.2	26.7	27.7	32.3	36.8	39.6	36.6	35.8	32.9
2000.....	23.8	25.1	28.5	35.4	36.4	39.2	39.7	34.7	34.1
1999.....	22.3	26.4	29.0	34.7	39.5	40.0	39.3	39.1	32.7

Non-Hispanic White, female

2003.....	8.6	10.0	11.7	13.8	18.0	23.3	25.7	24.5	17.0
2002.....	8.5	9.5	11.5	14.7	19.8	23.3	27.7	25.5	18.0
2001.....	9.0	9.7	10.8	15.3	19.6	22.6	26.8	24.3	18.2
2000.....	7.7	8.4	11.8	14.8	19.0	22.1	26.0	25.9	18.2
1999.....	7.0	8.1	10.3	15.5	18.5	23.5	27.2	26.1	18.6

Non-Hispanic Black, both sexes

2003.....	15.3	26.1	26.7	25.3	25.8	22.0	13.8	14.5	12.5
2002.....	17.8	25.3	24.9	25.8	23.2	19.9	19.7	16.3	12.1
2001.....	19.4	27.2	28.3	23.8	27.1	23.0	20.6	14.6	10.6
2000.....	19.8	26.3	26.0	26.3	27.8	25.0	18.3	17.8	14.0
1999.....	22.8	23.2	28.4	31.4	28.3	25.1	23.3	19.1	16.8

Non-Hispanic Black, male

2003.....	22.0	42.0	42.5	39.3	37.2	29.0	18.7	22.6	22.4
2002.....	23.9	38.9	37.3	39.3	35.1	28.7	22.1	21.8	*
2001.....	28.9	42.8	44.8	38.5	43.8	34.4	27.0	*	*
2000.....	27.2	41.2	38.9	37.8	38.4	37.9	27.9	19.7	*
1999.....	34.8	37.1	44.3	43.3	44.1	37.0	32.4	25.7	*

Non-Hispanic Black, female

2003.....	9.5	12.6	13.7	14.4	17.3	17.3	10.8	10.3	8.7
2002.....	12.6	13.9	14.7	15.3	14.4	14.0	18.2	13.4	10.5
2001.....	11.3	14.0	14.8	12.3	14.8	15.4	16.7	14.6	9.8
2000.....	13.3	13.6	15.4	17.3	20.1	16.3	12.5	16.9	11.8
1999.....	12.6	11.4	15.5	22.0	16.5	17.2	17.8	15.8	15.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)									
All origins 3/, both sexes									
2003.....	9.4	11.8	11.6	11.6	11.0	8.3	6.6	4.4	1.8
2002.....	9.4	11.4	11.4	11.8	10.7	9.5	6.7	4.4	2.1
2001.....	10.2	11.3	11.1	12.0	11.1	9.0	6.9	3.9	2.4
2000.....	10.0	10.7	11.7	12.9	11.4	9.1	7.2	4.4	2.2
1999.....	10.0	10.9	11.5	12.6	11.6	9.4	7.3	5.2	2.8
All origins 3/, male									
2003.....	14.2	18.3	18.1	18.6	17.7	14.2	11.3	8.2	3.6
2002.....	13.8	17.7	17.6	19.2	17.4	15.4	10.9	8.0	4.7
2001.....	15.3	17.0	18.0	19.0	17.9	15.2	12.1	7.0	5.5
2000.....	15.3	16.9	18.0	20.3	18.7	15.7	13.8	7.7	5.3
1999.....	15.5	17.2	18.7	20.0	19.7	16.2	12.1	9.7	6.1
All origins 3/, female									
2003.....	4.8	5.5	5.5	5.3	5.3	3.6	3.3	2.1	1.0
2002.....	5.1	5.4	5.6	5.1	4.9	4.8	3.7	2.2	1.0
2001.....	5.3	5.8	4.7	5.6	5.3	4.0	3.3	2.1	1.0
2000.....	4.8	4.8	5.8	6.2	5.1	3.9	2.6	2.4	1.0
1999.....	4.6	4.8	4.9	5.9	4.6	4.0	3.9	2.7	1.5
Hispanic, both sexes									
2003.....	13.2	16.0	21.0	22.0	20.1	14.7	16.1	10.7	*
2002.....	13.6	19.3	20.1	19.5	21.2	20.4	11.7	*	*
2001.....	16.0	18.6	20.9	22.2	22.2	19.8	19.3	12.9	*
2000.....	15.4	18.6	22.7	24.3	25.5	19.7	19.0	*	*
1999.....	16.1	17.8	21.5	22.1	21.4	19.5	16.5	*	*
Hispanic, male									
2003.....	23.0	28.8	36.8	40.6	31.8	25.2	30.4	*	*
2002.....	22.6	33.0	36.7	35.7	41.3	39.1	19.2	*	*
2001.....	27.5	31.7	37.8	40.5	43.9	39.3	34.1	*	*
2000.....	26.7	32.8	41.9	44.0	51.8	39.9	41.3	*	*
1999.....	27.7	31.0	40.2	39.6	39.8	39.2	29.6	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)									
Hispanic, female									
2003.....	3.2	3.6	6.3	5.8	10.4	6.6	*	*	*
2002.....	4.5	6.0	4.9	5.4	*	*	*	*	*
2001.....	4.5	6.0	5.6	6.3	*	*	*	*	*
2000.....	4.1	5.0	5.4	7.2	*	*	*	*	*
1999.....	4.5	5.3	4.6	6.9	6.6	*	*	*	*
Non-Hispanic 4/, both sexes									
2003.....	9.0	11.3	10.8	10.8	10.3	7.8	6.0	4.1	1.7
2002.....	8.9	10.6	10.6	11.2	9.8	8.8	6.4	4.2	1.9
2001.....	9.6	10.6	10.3	11.1	10.3	8.2	6.3	3.6	2.2
2000.....	9.4	10.0	10.8	12.0	10.4	8.5	6.6	4.2	2.2
1999.....	9.4	10.2	10.8	11.9	10.9	8.8	6.8	5.1	2.5
Non-Hispanic 4/, male									
2003.....	13.1	17.2	16.5	16.8	16.6	13.4	10.2	7.7	3.4
2002.....	12.7	16.2	15.9	17.8	15.5	13.9	10.3	7.6	4.3
2001.....	14.0	15.7	16.3	17.3	15.9	13.6	11.0	6.3	5.1
2000.....	14.0	15.5	16.1	18.4	16.4	14.3	12.4	7.3	5.0
1999.....	14.3	16.0	17.0	18.5	18.3	14.8	11.2	9.2	5.5
Non-Hispanic 4/, female									
2003.....	4.9	5.6	5.5	5.3	4.8	3.4	3.1	1.9	1.0
2002.....	5.1	5.3	5.6	5.1	4.9	4.7	3.6	2.2	0.9
2001.....	5.3	5.8	4.6	5.5	5.4	4.0	3.0	1.9	1.0
2000.....	4.8	4.8	5.8	6.1	5.2	3.9	2.6	2.4	1.0
1999.....	4.6	4.7	4.9	5.8	4.5	4.0	3.8	2.6	1.4
Non-Hispanic White, both sexes									
2003.....	9.1	11.2	10.8	10.6	10.4	8.1	6.4	4.2	1.7
2002.....	9.0	10.5	10.5	11.2	10.0	9.0	6.5	4.3	1.9
2001.....	9.7	10.1	9.9	11.3	10.1	8.4	6.4	3.6	2.1
2000.....	9.2	9.7	10.4	11.9	10.4	8.6	6.7	4.1	2.2
1999.....	9.0	10.0	10.3	11.5	10.7	8.8	6.7	5.0	2.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)									
Non-Hispanic White, male									
2003.....	13.2	17.1	16.4	16.5	16.7	13.7	10.6	7.6	3.5
2002.....	13.0	16.0	15.9	17.8	15.6	14.1	10.5	7.7	4.1
2001.....	14.3	15.1	15.6	17.4	15.4	13.6	11.2	6.4	4.8
2000.....	14.1	15.0	15.8	18.4	16.5	14.3	12.6	7.3	5.1
1999.....	13.9	15.7	16.2	18.3	17.8	14.7	10.9	9.3	5.1
Non-Hispanic White, female									
2003.....	5.1	5.5	5.4	5.0	4.8	3.4	3.3	2.1	1.0
2002.....	5.0	5.1	5.3	5.0	5.0	4.8	3.6	2.1	1.0
2001.....	5.2	5.3	4.5	5.6	5.4	4.1	3.0	1.9	1.0
2000.....	4.4	4.4	5.3	5.8	5.1	4.0	2.5	2.3	1.0
1999.....	4.2	4.4	4.6	5.3	4.5	4.0	3.7	2.5	1.3
Non-Hispanic Black, both sexes									
2003.....	7.7	12.8	12.3	13.1	11.3	7.1	*	*	*
2002.....	8.5	12.4	12.4	12.1	9.6	7.6	5.8	*	*
2001.....	9.1	14.9	14.4	11.6	13.1	8.0	5.7	*	*
2000.....	11.3	13.6	14.0	12.8	12.0	8.6	6.3	*	*
1999.....	12.8	12.9	15.0	15.9	13.2	10.2	8.2	*	*
Non-Hispanic Black, male									
2003.....	12.2	20.8	19.7	21.0	18.7	12.8	*	*	*
2002.....	11.9	19.1	17.2	20.4	16.4	13.3	9.9	*	*
2001.....	13.3	22.5	24.0	18.8	23.4	15.5	*	*	*
2000.....	15.5	21.0	20.5	18.5	19.5	17.1	11.6	*	*
1999.....	19.2	20.4	25.3	23.7	24.7	18.3	14.5	*	*
Non-Hispanic Black, female									
2003.....	3.7	6.2	6.3	7.0	5.8	*	*	*	*
2002.....	5.5	6.7	8.6	5.6	4.6	*	*	*	*
2001.....	5.6	8.4	6.5	5.9	5.4	*	*	*	*
2000.....	7.7	7.3	8.7	8.4	6.5	*	*	*	*
1999.....	7.4	6.6	6.7	9.7	4.7	4.8	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)									
All origins 3/, both sexes									
2003.....	6.5	9.3	10.4	12.7	17.0	22.9	24.5	24.1	18.2
2002.....	6.6	8.9	10.3	12.6	17.2	21.7	25.8	25.5	19.3
2001.....	6.9	8.8	10.0	12.8	17.7	22.2	24.4	24.7	19.9
2000.....	6.8	7.9	10.2	13.3	17.5	21.7	24.7	25.2	20.9
1999.....	6.4	7.7	9.9	13.9	18.1	22.3	25.4	25.6	20.4
All origins 3/, male									
2003.....	9.2	14.0	14.3	17.0	20.7	25.8	26.8	26.8	23.9
2002.....	9.5	13.3	14.4	15.3	19.8	24.6	28.6	28.9	23.7
2001.....	10.0	13.1	13.5	16.3	21.8	25.6	25.5	28.4	26.5
2000.....	10.2	11.6	13.7	17.4	20.6	25.0	27.0	26.8	29.0
1999.....	9.8	11.6	13.8	17.3	22.0	25.3	28.0	29.0	27.0
All origins 3/, female									
2003.....	3.8	4.8	6.6	8.9	13.8	20.5	22.8	22.5	15.8
2002.....	3.8	4.7	6.5	10.2	14.9	19.4	23.9	23.4	17.4
2001.....	4.0	4.7	6.7	9.6	14.2	19.5	23.6	22.6	17.1
2000.....	3.6	4.3	6.9	9.5	14.8	19.2	23.2	24.2	17.5
1999.....	3.2	4.0	6.2	10.8	14.8	19.9	23.5	23.6	17.7
Hispanic, both sexes									
2003.....	8.2	12.1	14.6	20.3	27.1	41.4	50.7	43.4	26.9
2002.....	9.5	12.6	17.0	20.7	29.4	42.8	41.1	46.3	38.8
2001.....	9.0	13.3	15.0	21.0	28.9	38.8	42.8	44.7	30.6
2000.....	9.4	11.8	16.9	19.5	30.0	42.1	46.8	51.2	44.5
1999.....	9.1	10.5	14.1	21.1	32.1	40.2	43.9	47.0	40.0
Hispanic, male									
2003.....	12.4	17.8	19.9	29.5	27.9	38.0	49.2	37.4	*
2002.....	15.4	19.4	24.4	22.8	35.2	40.4	41.6	54.9	*
2001.....	14.3	20.8	21.2	30.1	34.0	36.6	47.7	41.6	*
2000.....	14.4	18.7	22.8	23.9	35.1	38.9	48.0	42.7	50.4
1999.....	14.6	15.8	19.2	27.1	36.7	43.3	42.9	45.6	45.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)									
Hispanic, female									
2003.....	3.8	6.5	9.8	12.2	26.4	43.9	51.7	47.4	26.6
2002.....	3.4	6.1	10.1	18.8	24.6	44.6	40.7	40.9	42.3
2001.....	3.5	6.2	9.4	13.2	24.7	40.5	39.2	46.6	33.5
2000.....	4.4	5.3	11.5	15.6	26.0	44.6	46.0	56.4	41.5
1999.....	3.7	5.5	9.6	15.9	28.4	37.9	44.6	47.8	37.4
Non-Hispanic 4/, both sexes									
2003.....	6.3	9.0	10.0	12.1	16.2	21.6	22.9	23.1	17.9
2002.....	6.3	8.5	9.7	11.9	16.2	20.2	25.0	24.5	18.5
2001.....	6.7	8.4	9.6	12.1	16.9	21.1	23.4	23.8	19.4
2000.....	6.6	7.5	9.6	12.7	16.5	20.5	23.7	24.1	19.9
1999.....	6.1	7.4	9.5	13.3	17.2	21.3	24.5	24.8	19.6
Non-Hispanic 4/, male									
2003.....	8.8	13.5	13.8	15.9	20.1	25.0	25.6	26.2	23.7
2002.....	8.7	12.7	13.6	14.6	18.7	23.4	27.8	27.7	23.4
2001.....	9.5	12.4	12.9	15.1	20.8	25.0	24.4	27.7	26.4
2000.....	9.7	11.0	13.0	16.9	19.6	24.1	26.0	26.1	28.0
1999.....	9.3	11.1	13.3	16.5	21.1	24.4	27.3	28.4	26.2
Non-Hispanic 4/, female									
2003.....	3.8	4.7	6.4	8.6	12.7	18.9	21.1	21.3	15.3
2002.....	3.9	4.6	6.2	9.5	14.1	17.7	23.0	22.5	16.4
2001.....	4.0	4.6	6.5	9.3	13.5	18.0	22.7	21.5	16.5
2000.....	3.5	4.3	6.5	9.0	14.0	17.6	22.1	22.9	16.7
1999.....	3.1	3.9	5.9	10.5	13.8	18.9	22.5	22.6	17.0
Non-Hispanic White, both sexes									
2003.....	6.2	8.7	9.7	12.3	16.7	22.8	24.3	24.3	18.6
2002.....	6.0	8.2	9.7	11.9	16.9	21.2	26.5	25.3	19.3
2001.....	6.3	8.0	9.2	12.3	17.5	21.9	24.5	25.0	20.4
2000.....	6.5	7.0	9.5	12.8	16.7	21.1	25.0	25.0	20.6
1999.....	5.6	7.2	9.1	13.2	17.6	22.1	25.5	25.9	20.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)									
Non-Hispanic White, male									
2003.....	8.9	13.0	13.2	16.0	20.8	26.3	26.9	27.3	24.5
2002.....	8.6	12.2	13.2	14.4	19.2	24.6	29.7	28.6	24.7
2001.....	8.9	11.6	12.1	14.9	21.4	25.9	25.4	29.3	28.1
2000.....	9.7	10.1	12.7	16.9	20.0	24.8	27.1	27.4	29.0
1999.....	8.5	10.7	12.8	16.4	21.6	25.2	28.4	29.8	27.5
Non-Hispanic White, female									
2003.....	3.5	4.5	6.3	8.8	13.1	19.9	22.4	22.4	16.0
2002.....	3.5	4.4	6.3	9.6	14.8	18.5	24.1	23.4	17.0
2001.....	3.8	4.4	6.3	9.8	14.2	18.5	23.8	22.4	17.2
2000.....	3.3	4.0	6.5	9.0	13.8	18.1	23.5	23.6	17.2
1999.....	2.8	3.7	5.7	10.3	14.0	19.5	23.5	23.5	17.3
Non-Hispanic Black, both sexes									
2003.....	7.6	13.2	14.4	12.2	14.4	14.9	10.6	10.6	11.0
2002.....	9.4	13.0	12.4	13.8	13.6	12.4	13.9	12.3	9.7
2001.....	10.3	12.3	13.9	12.2	14.1	15.0	14.9	11.6	7.8
2000.....	8.5	12.7	12.0	13.5	15.8	16.4	12.0	13.5	12.4
1999.....	10.0	10.2	13.4	15.5	15.0	14.9	15.1	13.5	12.3
Non-Hispanic Black, male									
2003.....	9.8	21.3	22.9	18.3	18.5	16.2	13.5	*	*
2002.....	12.1	19.8	20.1	18.9	18.7	15.4	12.2	*	*
2001.....	15.6	20.2	20.8	19.6	20.3	18.9	18.0	*	*
2000.....	11.8	20.2	18.5	19.3	18.9	20.8	16.4	*	*
1999.....	15.6	16.6	19.0	19.6	19.4	18.7	17.9	*	*
Non-Hispanic Black, female									
2003.....	5.7	6.4	7.4	7.4	11.5	14.1	8.9	9.5	8.3
2002.....	7.0	7.2	6.2	9.7	9.9	10.3	14.9	11.0	9.6
2001.....	5.7	5.6	8.3	6.4	9.4	12.4	13.0	12.9	*
2000.....	5.6	6.2	6.7	8.9	13.6	13.4	9.3	14.3	11.3
1999.....	5.2	4.8	8.8	12.3	11.8	12.4	13.4	12.3	12.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	19.5
2002.....	*	*	*	*	*	*	*	*	23.2
2001.....	*	*	*	*	*	*	*	*	30.7
2000.....	*	*	*	*	*	*	*	*	31.7
1999.....	*	*	*	*	*	*	*	*	32.2

Non-Hispanic 4/, both sexes

2003.....	0.2	0.4	0.5	0.9	1.9	3.3	5.8	11.2	21.9
2002.....	0.2	0.4	0.6	1.0	2.0	3.5	5.5	10.7	22.8
2001.....	0.3	0.4	0.6	1.3	2.0	3.6	5.8	9.8	23.4
2000.....	0.2	0.3	0.6	1.0	1.8	2.9	5.5	10.1	24.6
1999.....	0.3	0.4	0.7	1.0	1.8	3.3	5.5	10.7	23.6

Non-Hispanic 4/, male

2003.....	*	0.4	0.6	1.1	2.1	4.2	7.4	13.5	23.0
2002.....	0.3	0.5	0.8	1.1	2.3	3.9	7.3	13.6	23.9
2001.....	0.4	0.4	0.8	1.5	2.1	4.2	7.5	11.9	26.6
2000.....	*	0.3	0.7	1.1	2.1	3.5	7.0	12.6	27.7
1999.....	0.3	0.3	0.9	1.2	2.0	4.2	6.6	13.1	27.5

Non-Hispanic 4/, female

2003.....	0.2	0.4	0.4	0.7	1.8	2.6	4.6	9.8	21.4
2002.....	*	0.3	0.5	0.9	1.7	3.2	4.3	8.9	22.4
2001.....	0.3	0.3	0.4	1.2	1.8	3.1	4.6	8.6	22.1
2000.....	*	0.4	0.5	0.9	1.5	2.5	4.5	8.6	23.4
1999.....	0.2	0.5	0.6	0.8	1.6	2.7	4.7	9.2	22.0

Non-Hispanic White, both sexes

2003.....	0.2	0.3	0.4	0.8	1.8	3.4	5.7	11.2	22.0
2002.....	0.2	0.3	0.6	0.9	2.0	3.4	5.3	10.6	23.4
2001.....	0.3	0.3	0.5	1.2	1.7	3.4	5.7	9.6	23.6
2000.....	0.1	0.3	0.5	0.9	1.7	2.6	5.5	9.9	25.3
1999.....	0.2	0.4	0.6	0.8	1.5	3.2	5.5	10.7	24.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All origins 3/, both sexes

2003.....	3.8	6.1	10.4	17.9	30.7	50.6	84.5	143.9	293.1
2002.....	3.7	5.9	10.0	16.8	29.4	50.0	85.4	142.2	288.6
2001.....	3.6	5.8	9.9	16.9	31.7	49.3	82.2	135.8	287.7
2000.....	3.4	5.5	9.4	16.9	29.5	47.2	79.5	132.8	277.8
1999.....	3.1	5.2	9.0	15.8	28.2	46.7	77.7	127.4	268.9

All origins 3/, male

2003.....	4.5	7.6	12.2	19.6	34.6	58.1	102.9	187.9	387.6
2002.....	4.3	6.9	11.0	18.6	33.6	59.8	105.7	182.7	385.5
2001.....	4.4	6.7	10.9	18.2	36.5	57.0	100.5	176.8	393.7
2000.....	4.4	6.4	10.7	18.6	32.8	54.3	99.8	173.0	380.7
1999.....	3.8	5.8	9.9	17.8	31.0	54.9	99.2	168.9	371.1

All origins 3/, female

2003.....	3.2	4.7	8.6	16.4	27.4	44.7	71.4	117.1	251.4
2002.....	3.1	4.9	9.2	15.2	25.8	42.1	71.1	117.6	246.6
2001.....	2.9	4.9	8.9	15.7	27.6	43.2	69.4	111.2	243.3
2000.....	2.5	4.6	8.2	15.5	26.6	41.6	65.3	109.0	235.9
1999.....	2.4	4.5	8.1	14.0	25.8	40.3	62.8	103.1	227.7

Hispanic, both sexes

2003.....	3.2	6.3	12.0	19.7	33.0	48.0	80.7	105.7	233.2
2002.....	2.7	6.3	10.5	17.7	32.9	40.3	74.9	111.8	201.2
2001.....	2.7	5.5	11.1	15.1	34.6	49.9	83.8	113.2	222.0
2000.....	2.5	4.7	9.1	19.3	32.5	43.8	71.9	108.1	229.6
1999.....	3.0	4.4	10.5	18.5	29.7	43.1	77.1	109.8	232.2

Hispanic, male

2003.....	3.6	8.4	12.8	20.9	36.1	50.8	91.8	134.1	297.1
2002.....	2.8	7.6	11.7	19.8	38.9	43.4	92.1	147.6	235.9
2001.....	2.8	5.1	12.6	18.7	35.4	53.5	102.3	131.2	281.9
2000.....	3.6	5.6	9.9	20.1	33.6	50.6	85.6	132.5	292.2
1999.....	3.6	5.4	13.4	23.5	32.5	47.9	103.8	124.2	336.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)									
Hispanic, female									
2003.....	2.9	4.3	11.3	18.7	30.4	45.9	72.9	87.3	200.8
2002.....	2.7	5.1	9.3	15.9	27.9	37.9	62.6	89.1	184.0
2001.....	2.5	5.8	9.8	12.0	33.9	47.1	70.7	102.1	192.7
2000.....	*	3.9	8.3	18.6	31.7	38.7	62.2	93.2	198.8
1999.....	2.4	3.5	7.9	14.1	27.5	39.5	58.4	101.3	182.0
Non-Hispanic 4/, both sexes									
2003.....	3.9	6.1	10.2	17.7	30.5	50.7	84.5	145.4	294.9
2002.....	3.8	5.8	9.9	16.7	29.1	50.3	85.7	143.2	291.2
2001.....	3.7	5.8	9.8	17.0	31.5	49.1	82.0	136.3	289.6
2000.....	3.5	5.5	9.4	16.6	29.2	47.2	79.6	133.4	279.0
1999.....	3.1	5.2	8.8	15.5	28.0	46.8	77.6	127.7	269.2
Non-Hispanic 4/, male									
2003.....	4.6	7.5	12.1	19.4	34.3	58.4	103.2	190.1	390.8
2002.....	4.5	6.8	10.9	18.4	33.1	60.5	106.0	183.7	391.0
2001.....	4.5	6.8	10.8	18.1	36.5	57.0	100.3	178.2	397.6
2000.....	4.5	6.4	10.8	18.3	32.7	54.3	100.3	174.1	384.0
1999.....	3.8	5.8	9.6	17.3	30.8	55.1	98.8	170.2	370.9
Non-Hispanic 4/, female									
2003.....	3.2	4.7	8.4	16.1	27.2	44.5	71.1	118.1	252.7
2002.....	3.1	4.9	9.1	15.0	25.6	42.2	71.3	118.6	248.2
2001.....	2.9	4.8	8.8	16.0	27.1	42.8	69.1	111.2	244.6
2000.....	2.6	4.7	8.2	15.1	26.2	41.6	65.2	109.4	236.6
1999.....	2.4	4.6	8.1	14.0	25.7	40.3	62.9	102.8	228.4
Non-Hispanic White, both sexes									
2003.....	2.5	4.0	7.4	13.5	24.4	42.9	77.0	137.2	285.9
2002.....	2.6	3.6	7.0	12.6	23.5	43.1	78.0	136.0	281.1
2001.....	2.3	3.9	6.8	12.5	25.0	42.4	74.4	128.0	279.7
2000.....	2.2	3.6	6.6	12.5	23.0	40.2	72.7	125.6	269.4
1999.....	1.8	3.4	6.2	11.6	22.4	40.1	70.5	118.6	255.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic White, male

2003.....	3.0	4.7	8.7	14.9	28.2	50.3	96.8	181.7	386.3
2002.....	3.0	4.2	7.6	13.8	27.5	53.2	99.0	178.5	382.6
2001.....	2.7	4.8	7.8	13.9	29.9	50.9	94.0	170.0	392.6
2000.....	2.8	4.2	7.7	14.3	27.3	48.4	93.5	167.6	378.9
1999.....	2.2	3.7	7.2	13.2	25.5	48.8	90.8	160.8	359.6

Non-Hispanic White, female

2003.....	2.0	3.3	6.0	12.2	21.0	36.9	62.7	109.8	241.7
2002.....	2.1	3.1	6.4	11.6	20.1	34.8	63.0	110.0	237.3
2001.....	1.8	3.1	5.9	11.2	20.6	35.5	60.5	102.6	232.7
2000.....	1.7	3.1	5.6	10.9	19.3	33.6	58.2	100.6	225.2
1999.....	1.5	3.0	5.3	10.3	19.8	33.1	56.2	93.7	213.8

Non-Hispanic Black, both sexes

2003.....	13.0	21.4	35.1	55.1	85.7	130.7	173.7	263.6	446.5
2002.....	12.0	22.1	34.4	51.4	80.3	128.1	179.7	244.8	444.0
2001.....	12.9	20.3	33.3	54.2	90.4	121.3	174.0	250.8	432.9
2000.....	13.1	20.1	33.0	51.3	84.6	121.2	167.2	245.1	410.2
1999.....	12.1	20.1	29.6	47.9	78.9	118.2	165.6	246.3	436.8

Non-Hispanic Black, male

2003.....	15.3	29.8	44.3	63.7	96.4	159.1	194.8	335.8	526.3
2002.....	15.0	27.4	41.0	62.3	90.4	150.9	208.4	284.7	568.9
2001.....	17.0	23.6	38.6	56.2	103.4	132.6	190.6	312.3	517.4
2000.....	18.0	24.3	38.3	54.9	87.9	128.5	201.4	292.7	496.7
1999.....	15.9	24.1	29.8	55.5	82.3	131.9	210.3	325.9	532.4

Non-Hispanic Black, female

2003.....	11.1	14.3	27.6	48.4	77.7	111.6	161.2	225.9	415.6
2002.....	9.4	17.5	28.9	42.8	72.9	112.9	162.6	224.0	396.3
2001.....	9.3	17.5	28.9	52.6	80.8	113.8	164.1	219.5	401.7
2000.....	8.9	16.4	28.7	48.5	82.2	116.4	146.6	221.2	378.1
1999.....	8.9	16.8	29.4	42.0	76.4	109.1	138.6	206.8	401.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	*	*	*	*	*	*	*	0.6	0.5
2002.....	*	*	*	*	*	*	*	0.6	0.9
2001.....	*	*	*	*	*	*	0.4	*	0.8
2000.....	*	*	*	*	*	*	*	0.7	1.2
1999.....	*	*	*	*	*	0.3	0.5	0.5	0.8

Non-Hispanic 4/, male

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	1.8
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, female

2003.....	*	*	*	*	*	*	*	*	0.6
2002.....	*	*	*	*	*	*	*	*	0.9
2001.....	*	*	*	*	*	*	*	*	0.7
2000.....	*	*	*	*	*	*	*	0.7	0.9
1999.....	*	*	*	*	*	*	0.5	*	0.7

Non-Hispanic White, both sexes

2003.....	*	*	*	*	*	*	*	0.5	*
2002.....	*	*	*	*	*	*	*	0.5	0.8
2001.....	*	*	*	*	*	*	0.4	*	0.8
2000.....	*	*	*	*	*	*	*	0.8	1.2
1999.....	*	*	*	*	*	0.3	0.5	0.5	0.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	*	0.2	*	0.2	0.4	0.7	1.0	1.6	2.8
2002.....	0.1	0.1	0.2	*	0.3	0.7	1.1	1.8	2.8
2001.....	*	0.1	*	0.2	0.5	0.7	0.9	1.5	2.8
2000.....	*	*	0.2	0.3	0.5	0.7	1.1	1.9	2.8
1999.....	*	0.2	0.2	0.3	0.4	0.7	1.5	2.0	3.2

Non-Hispanic 4/, male

2003.....	*	0.2	*	*	0.5	0.6	1.3	1.9	4.3
2002.....	*	*	*	*	*	1.0	1.5	2.1	3.8
2001.....	*	*	*	*	0.5	1.1	1.3	2.3	3.7
2000.....	*	*	*	*	0.6	0.8	1.5	2.1	4.2
1999.....	*	*	*	*	*	0.7	2.0	2.8	5.1

Non-Hispanic 4/, female

2003.....	*	*	*	*	*	0.8	0.8	1.4	2.2
2002.....	*	*	*	*	*	0.4	0.8	1.5	2.3
2001.....	*	*	*	*	0.5	0.5	0.7	1.1	2.4
2000.....	*	*	*	*	*	0.6	0.8	1.8	2.2
1999.....	*	*	*	*	*	0.7	1.1	1.5	2.4

Non-Hispanic White, both sexes

2003.....	*	*	*	*	0.4	0.6	0.9	1.5	2.7
2002.....	*	*	0.2	*	0.3	0.7	1.0	1.7	2.6
2001.....	*	*	*	*	0.4	0.7	0.9	1.5	2.8
2000.....	*	*	0.2	*	0.4	0.7	1.0	1.8	2.4
1999.....	*	*	*	0.2	0.3	0.6	1.4	1.9	2.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)									
All origins 3/, both sexes									
2003.....	3.7	5.9	10.2	17.6	30.2	49.7	83.2	141.7	289.7
2002.....	3.6	5.7	9.8	16.5	29.0	49.1	83.9	139.8	284.8
2001.....	3.5	5.7	9.7	16.6	31.1	48.4	80.9	133.8	284.0
2000.....	3.3	5.3	9.2	16.6	28.9	46.2	78.1	130.0	273.7
1999.....	3.0	4.9	8.7	15.3	27.7	45.7	75.7	124.8	264.8
All origins 3/, male									
2003.....	4.3	7.4	12.0	19.3	33.9	57.2	101.3	185.2	383.1
2002.....	4.2	6.7	10.6	18.4	33.1	58.8	103.9	180.0	380.7
2001.....	4.2	6.5	10.7	17.9	35.9	55.8	98.9	173.9	388.7
2000.....	4.2	6.1	10.4	18.1	32.1	53.3	97.9	170.1	374.2
1999.....	3.6	5.6	9.7	17.3	30.5	53.9	96.7	165.1	364.6
All origins 3/, female									
2003.....	3.2	4.6	8.5	16.0	27.0	43.6	70.2	115.0	248.4
2002.....	2.9	4.8	9.0	14.9	25.4	41.4	69.9	115.3	243.2
2001.....	2.8	4.8	8.8	15.5	27.0	42.5	68.3	109.8	240.1
2000.....	2.4	4.5	8.0	15.2	26.1	40.6	64.4	106.4	232.8
1999.....	2.3	4.3	7.9	13.5	25.3	39.3	61.1	101.2	224.5
Hispanic, both sexes									
2003.....	3.1	6.1	11.6	19.4	32.4	47.3	79.4	103.3	230.6
2002.....	2.7	6.3	10.1	17.3	32.3	40.1	73.0	110.1	198.4
2001.....	2.4	5.3	10.8	14.6	33.9	48.7	82.7	111.7	217.6
2000.....	2.4	4.4	8.9	18.9	32.0	43.0	70.7	105.3	225.6
1999.....	2.8	4.4	10.2	17.8	28.2	42.2	75.5	107.5	228.0
Hispanic, male									
2003.....	3.3	8.0	12.3	20.9	35.4	50.4	90.5	129.9	295.5
2002.....	2.8	7.6	11.0	19.2	38.6	43.4	88.9	145.3	232.6
2001.....	2.8	4.8	12.2	17.9	34.7	52.6	100.9	128.6	278.1
2000.....	3.4	5.3	9.6	19.3	33.6	49.6	84.2	129.6	280.1
1999.....	3.5	5.4	12.9	22.6	30.9	47.9	102.2	122.6	330.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)									
Hispanic, female									
2003.....	2.9	4.3	10.9	18.0	29.9	44.9	71.6	85.9	197.7
2002.....	2.5	5.0	9.3	15.7	27.1	37.5	61.7	87.6	181.5
2001.....	2.1	5.7	9.6	11.8	33.3	45.7	69.7	101.3	188.0
2000.....	*	3.6	8.1	18.6	30.8	37.9	61.2	90.5	198.8
1999.....	*	3.5	7.7	13.6	25.9	37.9	56.7	98.5	178.8
Non-Hispanic 4/, both sexes									
2003.....	3.8	5.9	10.0	17.4	29.9	49.8	83.1	143.1	291.4
2002.....	3.6	5.6	9.7	16.4	28.6	49.5	84.2	140.7	287.4
2001.....	3.6	5.6	9.6	16.7	30.8	48.2	80.6	134.3	285.9
2000.....	3.4	5.3	9.2	16.3	28.6	46.2	78.3	130.6	274.9
1999.....	3.0	5.0	8.6	15.1	27.6	45.7	75.6	125.0	265.0
Non-Hispanic 4/, male									
2003.....	4.4	7.3	12.0	19.1	33.6	57.6	101.7	187.5	386.2
2002.....	4.3	6.6	10.6	18.2	32.6	59.4	104.1	181.0	386.2
2001.....	4.4	6.6	10.6	17.8	35.9	55.8	98.7	175.3	392.7
2000.....	4.3	6.1	10.5	17.9	31.9	53.2	98.4	171.3	377.7
1999.....	3.6	5.5	9.4	16.9	30.3	54.0	96.3	166.3	364.4
Non-Hispanic 4/, female									
2003.....	3.2	4.6	8.2	15.8	26.7	43.5	69.9	116.1	249.7
2002.....	3.0	4.7	8.8	14.7	25.2	41.6	70.1	116.3	244.8
2001.....	2.8	4.7	8.6	15.8	26.5	42.1	68.0	109.7	241.5
2000.....	2.5	4.6	8.0	14.8	25.8	40.7	64.3	106.7	233.4
1999.....	2.4	4.4	7.8	13.5	25.2	39.2	61.2	100.9	225.2
Non-Hispanic White, both sexes									
2003.....	2.4	3.9	7.2	13.2	23.9	42.1	75.8	135.1	282.6
2002.....	2.4	3.5	6.8	12.4	23.1	42.2	76.7	133.7	277.6
2001.....	2.2	3.8	6.7	12.3	24.4	41.4	73.1	126.1	276.1
2000.....	2.1	3.6	6.4	12.2	22.6	39.2	71.5	122.8	265.7
1999.....	1.7	3.2	6.0	11.3	22.0	39.1	68.5	116.2	251.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)									
Non-Hispanic White, male									
2003.....	2.9	4.5	8.6	14.7	27.6	49.5	95.5	179.5	381.9
2002.....	2.9	4.1	7.3	13.5	26.9	52.1	97.4	175.9	377.8
2001.....	2.7	4.6	7.6	13.7	29.4	49.7	92.4	167.2	387.6
2000.....	2.6	4.1	7.4	14.0	26.6	47.3	91.7	164.6	373.4
1999.....	2.1	3.5	7.1	12.8	25.0	47.9	88.3	157.2	353.8
Non-Hispanic White, female									
2003.....	1.9	3.2	5.9	11.8	20.5	36.0	61.5	107.8	238.9
2002.....	2.0	2.9	6.2	11.3	19.6	34.0	61.8	107.9	234.3
2001.....	1.8	3.0	5.8	11.1	20.0	34.7	59.4	101.3	229.7
2000.....	1.6	3.0	5.4	10.6	19.0	32.7	57.3	98.0	222.2
1999.....	1.4	2.9	5.1	10.0	19.4	32.1	54.6	92.0	210.8
Non-Hispanic Black, both sexes									
2003.....	12.9	21.0	35.0	54.5	84.7	128.9	171.5	260.3	442.3
2002.....	11.6	21.5	33.9	51.1	79.9	127.3	177.1	241.6	436.5
2001.....	12.6	19.8	33.0	53.6	88.9	120.5	172.2	248.1	428.9
2000.....	12.7	19.2	32.6	50.6	83.2	120.1	164.9	241.7	404.2
1999.....	12.0	19.4	29.0	46.9	77.7	116.8	162.8	241.9	428.7
Non-Hispanic Black, male									
2003.....	14.9	29.2	44.3	62.9	94.9	157.0	191.1	330.5	521.0
2002.....	14.8	26.7	40.4	62.3	89.8	149.5	204.2	281.6	562.3
2001.....	16.6	22.8	38.2	55.4	101.4	131.5	189.1	309.9	514.0
2000.....	17.6	22.9	37.9	53.8	85.8	127.8	198.9	290.1	487.3
1999.....	15.7	23.3	29.3	54.4	81.3	130.2	206.9	319.7	519.4
Non-Hispanic Black, female									
2003.....	11.1	14.0	27.3	47.9	77.2	110.0	159.8	223.5	411.9
2002.....	8.8	17.0	28.7	42.3	72.5	112.4	160.9	220.8	388.4
2001.....	9.2	17.2	28.8	52.2	79.6	113.1	162.1	216.6	397.4
2000.....	8.5	16.0	28.3	48.1	81.2	115.0	144.3	217.3	373.3
1999.....	8.8	16.1	28.7	41.0	75.0	108.0	136.3	203.3	394.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Infections of kidney (N10-N12,N13.6,N15.1)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.2	0.2	0.2	0.2	0.6	0.5	1.2	2.8	6.0
2002.....	*	*	0.2	0.3	0.5	0.6	1.5	2.3	6.4
2001.....	*	0.1	0.2	0.2	0.3	0.6	1.7	2.4	6.1
2000.....	*	0.1	0.2	0.3	0.4	0.8	1.5	2.7	7.6
1999.....	*	0.2	0.2	0.3	0.5	0.9	1.4	2.6	7.5

Non-Hispanic 4/, male

2003.....	*	*	*	*	*	*	0.7	2.5	5.4
2002.....	*	*	*	*	*	0.7	1.3	1.5	5.3
2001.....	*	*	*	*	*	*	1.2	2.2	4.3
2000.....	*	*	*	*	*	*	1.2	2.1	5.8
1999.....	*	*	*	*	*	0.6	1.2	2.4	6.9

Non-Hispanic 4/, female

2003.....	*	*	*	*	0.7	0.6	1.6	3.1	6.2
2002.....	*	*	*	0.4	0.6	0.6	1.7	2.7	6.8
2001.....	*	*	*	*	*	0.6	2.1	2.5	6.9
2000.....	*	*	*	0.4	0.5	1.1	1.8	3.0	8.3
1999.....	*	0.3	*	*	0.7	1.1	1.6	2.8	7.7

Non-Hispanic White, both sexes

2003.....	0.1	0.2	*	*	0.5	0.6	1.2	2.7	6.0
2002.....	*	*	0.2	0.2	0.5	0.7	1.5	2.3	6.5
2001.....	*	*	*	*	0.3	0.6	1.8	2.2	6.2
2000.....	*	*	0.2	0.3	0.3	0.7	1.4	2.5	7.6
1999.....	*	0.2	0.2	*	0.5	0.8	1.4	2.7	7.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hyperplasia of prostate (N40)

Hispanic, female

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic 4/, both sexes

2003.....	*	*	*	*	*	0.4	0.7	2.0	5.3
2002.....	*	*	*	*	*	0.4	0.6	2.2	4.9
2001.....	*	*	*	*	*	0.3	0.8	1.6	5.7
2000.....	*	*	*	*	*	0.3	0.9	2.0	5.2
1999.....	*	*	*	*	*	0.4	0.8	2.1	5.1

Non-Hispanic 4/, male

2003.....	*	*	*	*	*	0.8	1.8	5.2	17.5
2002.....	*	*	*	*	*	0.8	1.5	5.7	16.2
2001.....	*	*	*	*	*	0.8	2.0	4.2	19.3
2000.....	*	*	*	*	*	0.7	2.1	5.4	18.0
1999.....	*	*	*	*	*	0.8	2.0	5.6	18.0

Non-Hispanic 4/, female

2003.....
2002.....
2001.....
2000.....
1999.....

Non-Hispanic White, both sexes

2003.....	*	*	*	*	*	0.4	0.7	2.0	5.6
2002.....	*	*	*	*	*	0.3	0.6	2.2	5.0
2001.....	*	*	*	*	*	0.3	0.8	1.6	5.9
2000.....	*	*	*	*	*	0.3	0.9	2.0	5.2
1999.....	*	*	*	*	*	0.3	0.9	2.0	5.2

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	1.6	1.9	2.4	2.4	2.4	2.2	3.0	4.6	6.7
2002.....	1.7	2.0	2.4	2.5	2.4	2.6	3.1	4.8	6.7
2001.....	1.5	1.9	2.2	2.6	2.3	2.7	3.2	4.3	7.2
2000.....	1.5	1.9	2.2	2.5	2.6	2.5	3.2	4.9	7.5
1999.....	1.7	1.8	2.4	2.4	2.6	2.4	3.4	4.9	7.9

Non-Hispanic 4/, male

2003.....	1.7	1.9	2.5	2.6	2.5	2.3	2.8	4.6	6.7
2002.....	1.9	2.1	2.4	2.8	2.5	2.9	3.4	4.6	7.0
2001.....	1.5	1.9	2.2	2.4	2.5	3.0	3.5	4.8	7.6
2000.....	1.6	2.2	2.0	2.5	2.8	2.6	3.1	5.2	10.4
1999.....	1.9	2.0	2.5	2.6	2.4	2.5	3.4	5.7	8.5

Non-Hispanic 4/, female

2003.....	1.4	1.8	2.4	2.3	2.3	2.2	3.2	4.5	6.8
2002.....	1.5	1.9	2.4	2.3	2.3	2.3	2.9	4.9	6.6
2001.....	1.5	1.8	2.2	2.8	2.1	2.4	3.0	4.1	7.0
2000.....	1.3	1.6	2.4	2.5	2.5	2.4	3.3	4.8	6.4
1999.....	1.6	1.7	2.2	2.2	2.8	2.3	3.5	4.5	7.6

Non-Hispanic White, both sexes

2003.....	1.5	2.0	2.6	2.5	2.4	2.3	3.1	4.7	6.9
2002.....	1.7	2.0	2.5	2.6	2.5	2.6	3.2	4.8	6.9
2001.....	1.4	2.0	2.3	2.8	2.4	2.7	3.3	4.4	7.4
2000.....	1.5	2.0	2.2	2.6	2.7	2.5	3.3	5.1	7.8
1999.....	1.8	1.9	2.5	2.5	2.7	2.4	3.4	5.0	8.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins 3/, both sexes

2003.....	7.3	6.8	6.8	8.2	10.6	17.3	26.7	55.7	224.1
2002.....	6.4	6.9	7.0	8.3	10.4	16.1	28.3	53.3	208.0
2001.....	9.1	8.8	8.4	9.9	12.7	17.8	29.8	54.3	206.4
2000.....	8.6	8.0	8.6	10.2	14.2	19.0	30.1	56.5	208.1
1999.....	5.9	5.8	6.9	8.9	11.9	17.6	26.5	50.5	187.7

All origins 3/, male

2003.....	9.4	8.9	8.9	10.6	13.7	21.3	31.2	58.0	177.9
2002.....	8.5	9.0	9.0	11.2	12.7	19.7	32.9	60.6	175.5
2001.....	11.9	11.8	11.0	13.2	15.5	20.9	33.2	60.6	175.0
2000.....	11.7	11.2	11.4	13.4	18.9	22.4	36.1	62.3	182.7
1999.....	8.2	7.9	9.1	11.9	15.1	21.8	31.5	54.4	166.3

All origins 3/, female

2003.....	5.1	4.8	4.9	6.0	7.9	14.2	23.5	54.3	244.6
2002.....	4.4	5.0	5.0	5.6	8.4	13.3	25.1	48.9	222.1
2001.....	6.3	5.9	5.9	6.9	10.2	15.4	27.4	50.4	219.5
2000.....	5.6	5.0	5.9	7.3	10.1	16.4	26.0	53.0	218.5
1999.....	3.7	3.9	4.9	6.2	9.2	14.2	23.0	48.3	196.4

Hispanic, both sexes

2003.....	3.5	4.0	3.5	3.6	5.3	7.2	14.6	24.6	110.9
2002.....	3.1	5.0	3.9	6.1	5.5	9.0	11.7	30.3	91.5
2001.....	9.0	7.4	6.7	7.1	8.7	11.3	14.4	27.8	108.2
2000.....	8.1	7.9	7.5	5.6	8.2	12.2	16.5	27.8	102.2
1999.....	3.8	2.3	3.2	4.7	6.2	8.1	13.9	29.4	87.0

Hispanic, male

2003.....	5.7	5.5	5.6	5.4	6.5	10.2	15.2	24.9	95.4
2002.....	3.8	5.8	4.8	8.5	7.8	11.8	15.4	37.7	88.7
2001.....	14.0	10.7	10.0	11.4	12.8	16.0	18.4	37.7	87.6
2000.....	12.4	12.3	9.0	8.9	13.1	16.0	19.9	36.8	106.8
1999.....	6.2	*	*	6.9	8.5	*	21.8	*	77.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Hispanic, female

2003.....	*	2.5	*	*	*	*	14.2	24.4	118.8
2002.....	2.4	4.2	*	*	*	6.8	9.1	25.5	92.8
2001.....	4.0	4.4	3.8	*	*	7.7	11.6	21.7	118.2
2000.....	3.8	3.6	6.2	*	*	9.2	14.1	22.4	99.9
1999.....	*	*	*	*	*	8.0	*	30.0	91.5

Non-Hispanic 4/, both sexes

2003.....	7.6	7.0	7.0	8.5	11.0	17.9	27.3	57.1	228.4
2002.....	6.7	7.0	7.2	8.3	10.7	16.6	29.1	54.1	211.9
2001.....	9.0	8.8	8.5	10.0	12.9	18.2	30.4	55.3	209.6
2000.....	8.6	7.9	8.5	10.5	14.5	19.3	30.6	57.4	211.6
1999.....	6.0	6.0	7.1	9.1	12.2	17.9	27.0	51.2	190.9

Non-Hispanic 4/, male

2003.....	9.7	9.1	9.0	10.9	14.1	21.8	32.0	59.4	181.1
2002.....	9.0	9.2	9.2	11.2	12.9	20.1	33.7	61.1	178.6
2001.....	11.6	11.8	11.1	13.2	15.7	21.1	33.8	61.6	178.2
2000.....	11.5	10.9	11.3	13.5	19.0	22.6	36.6	63.0	185.4
1999.....	8.3	8.1	9.3	12.1	15.4	22.3	31.8	55.2	169.5

Non-Hispanic 4/, female

2003.....	5.6	5.0	5.1	6.2	8.2	14.8	23.9	55.6	249.2
2002.....	4.5	5.0	5.2	5.7	8.8	13.7	25.9	49.8	226.3
2001.....	6.4	6.0	6.1	7.1	10.5	15.8	28.0	51.6	222.7
2000.....	5.7	5.0	5.8	7.7	10.5	16.7	26.5	54.1	222.3
1999.....	3.9	4.0	5.1	6.5	9.4	14.5	23.6	48.9	199.5

Non-Hispanic White, both sexes

2003.....	7.0	6.4	6.5	7.9	10.4	17.2	27.1	57.6	232.9
2002.....	6.1	6.1	6.6	7.9	10.1	16.3	28.7	54.1	215.8
2001.....	8.7	8.3	7.9	9.5	12.1	17.5	30.1	55.0	213.3
2000.....	8.0	7.1	7.7	9.7	13.2	18.7	30.0	57.0	215.8
1999.....	5.2	5.2	6.3	8.4	11.3	17.3	25.9	50.1	193.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic White, male									
2003.....	8.9	8.3	8.2	10.0	13.2	21.0	31.8	59.3	183.3
2002.....	8.1	7.8	8.3	10.6	12.1	19.7	32.3	60.6	181.5
2001.....	11.2	11.1	10.0	12.1	14.4	20.5	33.9	61.6	180.4
2000.....	10.9	9.7	10.0	12.5	17.1	21.6	35.9	62.6	189.6
1999.....	6.9	7.0	8.2	11.2	14.2	21.5	30.5	53.3	171.1
Non-Hispanic White, female									
2003.....	5.0	4.4	4.9	6.0	8.0	14.2	23.7	56.5	254.7
2002.....	4.1	4.4	4.9	5.4	8.3	13.6	26.1	50.2	230.6
2001.....	6.2	5.5	5.9	7.0	10.0	15.1	27.4	51.0	227.0
2000.....	5.2	4.5	5.6	7.1	9.7	16.3	25.9	53.6	226.4
1999.....	3.5	3.5	4.6	5.9	8.8	14.0	22.7	48.2	202.8
Non-Hispanic Black, both sexes									
2003.....	13.7	13.5	13.2	15.9	18.4	31.1	35.3	61.2	214.5
2002.....	12.2	15.9	13.9	13.8	18.0	23.3	39.7	65.4	198.1
2001.....	13.2	14.9	14.6	15.7	21.8	28.1	39.2	67.6	194.2
2000.....	13.9	16.0	15.8	19.1	28.3	30.3	44.7	74.1	189.2
1999.....	13.2	13.2	15.0	17.1	21.9	27.8	43.3	73.4	182.0
Non-Hispanic Black, male									
2003.....	17.8	17.9	18.8	23.2	26.6	39.8	42.1	70.2	197.1
2002.....	17.3	22.4	19.8	20.4	22.8	31.1	57.3	87.4	182.7
2001.....	17.4	19.2	22.2	25.9	29.7	32.0	41.2	71.1	186.0
2000.....	18.3	23.5	24.7	25.5	41.0	38.6	55.4	85.0	163.2
1999.....	20.0	18.7	22.3	23.9	30.0	36.4	54.6	89.4	180.7
Non-Hispanic Black, female									
2003.....	10.1	9.8	8.6	10.2	12.4	25.2	31.4	56.5	221.2
2002.....	7.9	10.4	9.1	8.8	14.4	18.1	29.2	54.0	204.0
2001.....	9.6	11.2	8.4	7.7	16.0	25.5	37.9	65.8	197.2
2000.....	10.1	9.7	8.5	14.1	18.9	24.7	38.3	68.7	198.9
1999.....	7.5	8.5	9.1	11.8	15.9	22.1	36.4	65.4	182.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

All origins 3/, both sexes

2003.....	33.3	45.3	56.6	80.9	114.5	178.3	303.7	559.8	1,508.4
2002.....	33.6	42.6	56.2	77.1	110.4	173.8	297.8	555.5	1,493.0
2001.....	32.1	40.9	53.2	76.4	112.0	172.3	295.5	539.3	1,476.7
2000.....	31.0	40.4	53.1	74.1	106.0	168.2	288.9	537.8	1,445.7
1999.....	29.9	38.4	51.0	72.1	107.7	167.8	281.3	510.8	1,410.7

All origins 3/, male

2003.....	40.3	55.2	65.6	91.0	125.7	191.0	316.5	563.7	1,258.1
2002.....	41.0	52.4	64.4	87.9	122.8	186.8	309.6	555.2	1,262.9
2001.....	38.8	49.1	59.9	85.8	124.3	186.5	307.6	539.6	1,271.6
2000.....	38.5	47.6	60.9	83.1	117.6	183.1	303.4	537.6	1,266.2
1999.....	36.3	45.4	59.5	80.9	117.1	184.3	298.1	528.1	1,225.4

All origins 3/, female

2003.....	26.5	36.0	48.1	71.7	104.8	168.0	294.7	557.4	1,619.1
2002.....	26.4	33.2	48.6	67.3	99.6	163.5	289.5	555.6	1,592.8
2001.....	25.6	33.0	46.9	67.9	101.4	161.0	287.0	539.1	1,562.6
2000.....	23.7	33.4	45.8	65.9	96.1	156.5	278.8	537.8	1,518.8
1999.....	23.7	31.7	43.1	64.2	99.5	154.8	269.7	500.7	1,485.3

Hispanic, both sexes

2003.....	25.3	36.9	46.3	63.5	87.0	132.0	219.9	389.4	887.2
2002.....	25.6	30.1	43.3	55.6	90.9	136.7	212.2	372.5	870.8
2001.....	22.8	30.7	42.2	59.7	83.9	136.4	209.8	396.3	991.0
2000.....	24.8	31.0	45.2	50.5	84.3	127.4	217.6	404.9	913.0
1999.....	26.0	31.2	43.3	61.6	88.6	135.5	202.4	374.7	1,039.7

Hispanic, male

2003.....	32.5	46.4	59.4	72.1	97.5	141.4	229.1	426.2	757.3
2002.....	34.0	37.5	51.9	62.6	109.0	147.9	233.6	391.2	808.2
2001.....	30.6	40.0	51.7	69.0	96.0	144.6	205.3	403.8	939.0
2000.....	33.3	38.7	59.2	55.8	105.9	138.1	221.5	378.4	870.7
1999.....	35.4	39.2	53.2	74.8	96.6	159.9	220.0	389.8	923.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

Hispanic, female

2003.....	17.8	27.6	34.2	56.0	78.4	124.9	213.4	365.4	953.1
2002.....	17.0	22.9	35.4	49.4	76.0	128.1	197.0	360.6	901.9
2001.....	14.9	21.9	33.5	51.7	74.0	130.2	213.0	391.7	1,016.5
2000.....	16.3	23.7	32.5	45.9	66.7	119.3	214.9	421.1	933.8
1999.....	16.7	23.5	34.4	50.3	82.2	117.2	190.0	365.7	1,096.0

Non-Hispanic 4/, both sexes

2003.....	34.0	46.0	57.3	82.0	116.2	181.0	307.7	567.1	1,531.7
2002.....	34.3	43.4	57.0	78.4	111.5	175.8	301.5	561.8	1,514.9
2001.....	32.9	41.5	53.8	77.5	113.6	174.0	299.1	543.9	1,491.3
2000.....	31.3	40.9	53.4	75.6	107.1	169.8	291.3	541.5	1,462.0
1999.....	30.0	38.8	51.3	72.6	108.5	169.1	284.1	514.7	1,420.1

Non-Hispanic 4/, male

2003.....	41.0	55.7	65.9	92.0	127.3	193.8	320.6	569.7	1,278.3
2002.....	41.5	53.4	65.0	89.3	123.3	188.7	312.9	561.0	1,280.6
2001.....	39.4	49.5	60.0	86.7	125.7	188.5	311.8	544.1	1,282.3
2000.....	38.7	48.0	60.8	84.6	117.8	184.8	306.1	542.6	1,279.8
1999.....	36.1	45.6	59.6	80.8	117.9	185.0	300.8	532.1	1,234.0

Non-Hispanic 4/, female

2003.....	27.3	36.6	49.1	72.8	106.6	170.7	298.6	565.5	1,643.1
2002.....	27.3	33.9	49.5	68.5	101.3	165.4	293.4	562.3	1,615.9
2001.....	26.6	33.9	47.9	69.1	103.1	162.4	290.1	543.7	1,578.3
2000.....	24.2	34.1	46.5	67.3	97.8	158.0	281.0	540.9	1,535.7
1999.....	24.2	32.3	43.6	65.1	100.4	156.6	272.4	504.6	1,494.7

Non-Hispanic White, both sexes

2003.....	31.2	41.8	53.8	77.8	113.4	178.0	305.7	569.1	1,558.9
2002.....	31.2	39.8	53.6	74.5	108.5	171.7	299.2	562.5	1,540.5
2001.....	29.2	37.5	49.8	73.3	110.2	170.7	296.0	545.5	1,512.9
2000.....	27.3	37.3	49.3	71.2	103.2	165.9	287.8	540.7	1,481.3
1999.....	26.0	34.8	47.4	68.6	103.6	164.6	279.7	516.0	1,443.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

Non-Hispanic White, male

2003.....	38.4	50.8	61.8	86.8	122.7	188.5	315.2	566.4	1,300.3
2002.....	38.3	49.3	60.8	84.6	118.5	182.6	307.9	558.6	1,293.4
2001.....	35.5	45.0	55.1	82.2	121.0	183.6	306.0	544.6	1,293.1
2000.....	34.7	44.1	56.1	78.5	112.4	178.8	300.6	538.2	1,288.3
1999.....	31.8	41.2	55.0	76.3	112.0	179.6	293.3	529.5	1,239.8

Non-Hispanic White, female

2003.....	24.1	32.9	46.1	69.4	105.1	169.4	298.8	570.8	1,672.7
2002.....	24.2	30.4	46.7	65.1	99.6	162.8	293.0	564.9	1,647.2
2001.....	23.0	30.2	44.7	65.1	100.7	160.1	288.9	546.0	1,604.4
2000.....	19.9	30.7	42.7	64.4	95.2	155.5	278.8	542.2	1,559.1
1999.....	20.3	28.5	40.1	61.5	96.2	152.6	270.1	508.1	1,525.2

Non-Hispanic Black, both sexes

2003.....	59.7	85.3	98.5	133.8	164.7	245.2	387.7	636.4	1,451.6
2002.....	61.9	79.5	97.9	126.0	158.0	246.8	373.5	648.9	1,443.4
2001.....	64.1	80.9	97.2	124.2	165.8	239.3	384.5	619.4	1,431.5
2000.....	65.4	77.2	97.4	124.6	159.8	237.4	375.9	624.5	1,403.3
1999.....	63.0	77.3	93.5	118.8	169.4	242.1	373.1	560.6	1,295.3

Non-Hispanic Black, male

2003.....	67.5	104.5	115.9	160.6	194.9	294.2	447.6	722.2	1,243.4
2002.....	72.8	96.4	118.3	148.7	188.1	292.8	421.6	699.6	1,331.3
2001.....	75.8	97.7	116.3	144.0	199.8	276.5	449.4	669.7	1,329.4
2000.....	75.8	90.8	116.7	153.5	191.1	284.0	432.1	698.6	1,340.9
1999.....	74.5	91.6	112.6	135.8	192.6	274.5	439.4	638.5	1,303.8

Non-Hispanic Black, female

2003.....	52.9	69.1	84.1	112.9	142.3	212.2	352.1	591.6	1,532.3
2002.....	52.5	65.3	81.1	108.4	135.7	215.9	344.8	622.5	1,486.2
2001.....	54.1	66.6	81.6	108.6	140.6	214.5	345.7	593.9	1,469.3
2000.....	56.4	65.7	81.6	101.8	136.6	206.4	342.1	587.2	1,426.5
1999.....	53.2	65.2	78.0	105.5	152.2	220.6	333.1	522.0	1,292.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All origins 3/, both sexes

2003.....	40.7	36.6	32.8	32.9	38.1	50.9	78.8	133.5	278.9
2002.....	38.7	34.3	30.4	32.6	38.3	50.8	80.0	131.1	275.4
2001.....	36.0	31.9	29.3	31.5	36.8	49.3	80.4	130.4	276.4
2000.....	34.2	30.8	29.9	32.2	35.4	48.8	75.5	124.6	273.5
1999.....	34.1	29.2	30.1	31.2	38.7	50.9	78.6	133.3	282.4

All origins 3/, male

2003.....	58.9	52.3	47.2	47.0	52.3	66.8	104.9	175.2	352.5
2002.....	56.6	49.6	44.8	45.7	53.4	68.9	106.4	170.1	346.6
2001.....	54.0	46.9	41.8	44.5	49.9	65.6	108.3	173.2	358.5
2000.....	51.6	46.0	43.1	45.3	48.1	64.6	99.8	164.8	359.7
1999.....	51.7	43.6	43.6	44.5	53.1	68.2	102.8	181.8	375.4

All origins 3/, female

2003.....	23.0	21.6	19.3	20.1	25.8	38.1	60.3	108.1	246.4
2002.....	21.3	19.6	16.8	20.7	25.3	36.3	61.4	107.3	244.6
2001.....	18.5	17.5	17.6	19.6	25.5	36.3	60.9	104.8	242.0
2000.....	17.3	16.1	17.5	20.4	24.5	36.4	58.5	100.9	238.4
1999.....	17.0	15.5	17.6	19.2	26.4	37.3	61.8	105.0	244.8

Hispanic, both sexes

2003.....	33.9	31.5	27.8	34.5	35.0	42.1	59.0	90.2	151.8
2002.....	33.0	30.5	27.3	30.5	34.7	45.4	69.9	90.0	153.0
2001.....	32.6	30.5	29.7	31.6	32.5	49.3	62.9	92.9	166.3
2000.....	30.5	30.2	28.3	29.3	31.2	48.4	59.1	95.8	184.5
1999.....	32.5	29.6	27.4	29.8	42.3	42.6	59.1	88.7	182.4

Hispanic, male

2003.....	53.3	46.9	42.1	50.0	49.7	58.1	88.7	125.8	195.5
2002.....	50.0	47.0	43.1	44.1	56.2	61.9	97.3	121.2	224.2
2001.....	50.0	47.3	42.6	48.3	46.1	66.3	92.8	128.6	198.1
2000.....	49.1	47.4	42.3	44.3	47.0	70.5	86.4	109.0	251.9
1999.....	52.9	46.6	42.3	45.0	62.6	64.2	78.0	128.9	258.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Hispanic, female									
2003.....	14.0	16.5	14.6	21.0	22.9	29.8	37.9	67.0	129.7
2002.....	15.8	14.6	12.8	18.6	17.1	32.8	50.3	70.1	117.7
2001.....	15.2	14.6	18.1	17.2	21.5	36.3	41.6	70.8	150.8
2000.....	12.0	13.8	15.7	16.4	18.4	31.7	39.7	87.8	151.3
1999.....	12.2	13.5	14.0	16.7	25.9	26.4	45.7	64.7	145.6
Non-Hispanic 4/, both sexes									
2003.....	41.3	36.8	33.1	32.6	38.1	51.2	79.7	135.3	283.7
2002.....	39.1	34.4	30.4	32.6	38.4	50.9	80.3	132.4	279.8
2001.....	36.0	31.8	29.0	31.2	36.9	49.1	81.1	131.5	279.8
2000.....	34.3	30.5	29.8	32.3	35.5	48.6	76.0	125.4	276.1
1999.....	33.9	28.9	30.0	31.1	38.3	51.2	79.2	134.3	285.0
Non-Hispanic 4/, male									
2003.....	59.2	52.4	47.3	46.6	52.1	67.1	105.4	177.2	358.9
2002.....	56.9	49.5	44.5	45.5	52.8	68.9	106.5	171.3	351.2
2001.....	54.0	46.5	41.4	43.9	50.0	65.2	108.8	174.4	364.5
2000.....	51.4	45.4	42.8	45.0	47.9	64.0	100.1	166.5	363.1
1999.....	51.1	42.9	43.1	44.1	52.2	68.2	103.5	182.5	378.7
Non-Hispanic 4/, female									
2003.....	23.8	22.0	19.7	19.9	25.9	38.5	61.4	109.7	250.6
2002.....	21.8	20.0	17.2	20.9	25.9	36.4	61.7	108.8	249.0
2001.....	18.7	17.8	17.4	19.7	25.6	36.2	61.8	105.8	244.6
2000.....	17.7	16.2	17.5	20.6	24.9	36.5	59.2	101.1	240.9
1999.....	17.4	15.6	17.7	19.4	26.3	37.9	62.2	106.1	247.5
Non-Hispanic White, both sexes									
2003.....	41.4	35.7	32.3	32.1	37.2	51.3	81.1	138.4	296.5
2002.....	38.8	33.5	29.4	32.0	37.3	51.0	81.2	136.1	290.9
2001.....	34.7	30.8	28.1	30.1	36.5	48.5	81.9	134.4	289.3
2000.....	33.0	28.7	28.6	31.7	34.5	47.8	76.2	128.1	285.6
1999.....	32.3	27.4	28.6	29.2	37.1	49.7	78.5	136.1	293.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic White, male

2003.....	58.6	50.4	45.5	45.3	50.4	67.1	106.1	179.6	372.9
2002.....	56.6	47.3	42.2	44.1	50.1	68.3	106.5	174.1	361.6
2001.....	51.3	44.3	39.2	41.8	48.8	64.2	108.7	176.3	374.7
2000.....	49.1	42.2	40.2	43.3	45.4	62.5	98.8	168.3	373.8
1999.....	47.9	40.1	40.3	41.3	49.9	65.4	101.3	184.2	387.1

Non-Hispanic White, female

2003.....	24.3	21.4	19.5	19.7	25.4	38.3	62.9	113.1	262.8
2002.....	21.1	20.1	17.2	20.8	25.9	36.8	63.2	112.8	260.3
2001.....	18.3	17.5	17.4	19.4	25.6	35.9	63.0	109.1	253.8
2000.....	17.1	15.5	17.6	20.9	24.9	35.9	60.3	104.3	250.0
1999.....	16.9	14.9	17.5	18.0	25.9	37.1	62.5	107.7	255.9

Non-Hispanic Black, both sexes

2003.....	49.0	49.8	44.6	40.8	48.6	54.1	68.4	105.0	161.8
2002.....	49.1	45.9	42.7	40.8	47.4	52.6	76.0	97.7	175.4
2001.....	50.8	45.7	41.2	43.9	45.9	58.0	79.2	105.7	183.9
2000.....	47.9	47.6	42.7	39.6	46.8	60.4	75.9	100.3	183.2
1999.....	51.3	44.5	44.0	46.4	51.3	67.5	90.7	125.0	208.9

Non-Hispanic Black, male

2003.....	75.6	76.0	71.3	65.7	75.4	74.1	106.1	152.4	228.0
2002.....	73.0	73.6	72.1	65.4	75.5	79.4	119.2	151.3	262.5
2001.....	83.0	73.2	67.0	69.8	69.8	81.6	121.4	167.6	256.4
2000.....	76.2	77.4	72.6	65.3	75.2	87.1	120.4	156.2	277.1
1999.....	84.1	73.2	72.8	70.5	78.9	103.5	136.3	178.0	318.5

Non-Hispanic Black, female

2003.....	25.8	27.6	22.6	21.3	28.7	40.7	46.1	80.2	136.2
2002.....	28.4	22.4	18.6	21.6	26.6	34.6	50.2	69.8	142.1
2001.....	23.0	22.3	20.0	23.5	28.3	42.2	54.0	74.1	157.0
2000.....	23.5	22.3	18.3	19.3	26.0	42.7	49.1	72.2	148.3
1999.....	23.0	20.2	20.5	27.5	30.9	43.7	63.2	98.8	167.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Transport accidents (V01-V99,Y85)

All origins 3/, both sexes

2003.....	16.8	16.3	15.9	16.0	16.2	19.0	23.9	29.4	29.7
2002.....	16.9	15.8	14.9	16.7	16.7	20.2	25.0	29.7	28.8
2001.....	16.3	15.6	15.2	15.4	16.4	19.5	26.2	30.4	28.9
2000.....	16.2	15.5	15.7	16.4	16.8	19.4	25.1	30.0	31.5
1999.....	15.9	14.9	16.3	15.7	18.1	20.9	25.9	32.1	32.0

All origins 3/, male

2003.....	24.9	23.6	23.2	22.7	22.5	25.0	32.9	42.1	52.0
2002.....	25.1	22.7	22.2	22.8	23.4	27.5	34.5	42.6	53.0
2001.....	24.0	22.5	21.4	22.0	22.1	26.8	36.2	45.5	52.7
2000.....	23.6	22.3	22.4	22.6	22.6	26.3	34.5	44.0	59.8
1999.....	23.1	21.5	23.0	21.8	24.8	27.5	36.2	48.3	63.5

All origins 3/, female

2003.....	9.0	9.2	9.0	9.9	10.7	14.2	17.5	21.7	19.9
2002.....	8.8	9.2	7.9	11.1	10.9	14.3	18.3	21.8	18.3
2001.....	8.8	8.9	9.4	9.3	11.6	13.8	19.2	21.3	18.9
2000.....	8.9	8.9	9.5	10.7	11.7	13.9	18.5	21.7	19.9
1999.....	8.9	8.6	10.0	10.2	12.4	15.7	18.8	22.6	19.3

Hispanic, both sexes

2003.....	15.2	15.0	14.6	17.4	17.7	19.7	20.9	27.9	24.9
2002.....	15.3	15.0	13.6	17.3	16.6	20.6	27.5	24.1	25.5
2001.....	16.3	16.5	17.1	19.0	18.1	22.8	28.0	33.8	23.8
2000.....	15.8	16.7	15.9	15.3	16.4	25.6	23.3	25.1	25.2
1999.....	14.6	15.5	16.2	14.5	22.8	21.0	24.5	25.8	25.3

Hispanic, male

2003.....	22.9	21.7	20.4	22.9	25.3	27.3	30.4	43.7	46.2
2002.....	22.1	20.9	20.9	23.9	27.8	29.4	40.3	32.0	45.2
2001.....	22.4	24.1	21.4	29.1	25.9	31.1	45.7	61.0	43.8
2000.....	23.0	25.4	23.5	23.6	23.5	37.4	39.1	30.9	44.3
1999.....	22.0	23.4	21.7	20.0	32.8	30.0	35.9	48.7	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)									
Hispanic, female									
2003.....	7.4	8.5	9.2	12.7	11.5	13.8	14.2	17.6	*
2002.....	8.4	9.2	6.9	11.6	7.5	13.9	18.3	19.0	*
2001.....	10.1	9.3	13.2	10.3	11.8	16.4	15.5	16.9	*
2000.....	8.7	8.3	9.1	8.2	10.6	16.6	12.0	21.5	*
1999.....	7.3	8.1	11.3	9.8	14.7	14.2	16.5	*	*
Non-Hispanic 4/, both sexes									
2003.....	16.9	16.3	15.9	15.8	16.0	18.8	24.0	29.4	29.8
2002.....	16.9	15.8	14.9	16.5	16.6	20.1	24.8	29.7	28.9
2001.....	16.2	15.4	15.0	15.0	16.3	19.2	26.0	30.1	29.0
2000.....	16.1	15.2	15.6	16.3	16.7	18.9	25.1	30.0	31.7
1999.....	15.9	14.8	16.2	15.7	17.7	20.8	25.9	32.0	32.2
Non-Hispanic 4/, male									
2003.....	25.0	23.6	23.3	22.5	22.1	24.7	33.0	42.0	52.0
2002.....	25.3	22.8	22.2	22.6	23.0	27.2	34.1	42.7	53.2
2001.....	24.0	22.3	21.3	21.3	21.8	26.4	35.7	44.6	52.9
2000.....	23.6	21.9	22.2	22.3	22.5	25.5	34.3	44.3	60.4
1999.....	23.1	21.3	22.9	21.8	24.2	27.2	36.1	47.7	64.3
Non-Hispanic 4/, female									
2003.....	9.1	9.3	8.9	9.6	10.6	14.1	17.7	21.8	20.1
2002.....	8.9	9.1	8.0	11.0	11.1	14.3	18.3	21.9	18.4
2001.....	8.6	8.9	9.0	9.3	11.5	13.5	19.3	21.4	19.1
2000.....	8.9	8.9	9.5	10.9	11.8	13.8	18.8	21.7	20.1
1999.....	9.0	8.6	9.9	10.2	12.2	15.7	18.8	22.9	19.3
Non-Hispanic White, both sexes									
2003.....	17.0	16.1	15.6	15.5	15.7	19.0	24.0	30.0	30.6
2002.....	16.9	15.8	14.8	16.5	16.3	20.3	25.3	30.4	29.7
2001.....	15.7	15.4	14.8	14.7	16.2	19.2	26.5	30.8	29.9
2000.....	15.9	14.8	15.4	16.4	16.5	18.7	25.3	30.9	32.4
1999.....	15.6	14.5	15.8	15.3	17.6	20.7	26.0	32.4	32.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)									
Non-Hispanic White, male									
2003.....	24.8	23.3	22.8	22.3	21.7	25.1	32.6	42.5	52.8
2002.....	25.4	22.5	21.7	22.4	22.1	27.0	33.9	42.9	54.0
2001.....	23.1	22.2	20.8	20.7	21.4	26.2	35.7	45.7	54.3
2000.....	23.2	21.1	21.4	21.9	21.6	24.9	34.0	44.8	62.3
1999.....	22.3	20.7	22.3	21.0	23.8	26.7	35.7	47.8	64.6
Non-Hispanic White, female									
2003.....	9.2	9.1	8.7	9.2	10.3	14.0	17.8	22.3	20.8
2002.....	8.6	9.2	8.1	10.9	11.2	14.8	19.1	22.8	19.2
2001.....	8.5	8.8	9.0	9.1	11.6	13.5	20.0	21.9	19.8
2000.....	8.7	8.5	9.6	11.3	12.0	13.7	19.2	22.6	20.4
1999.....	8.9	8.4	9.7	10.1	12.1	15.8	19.1	23.3	19.7
Non-Hispanic Black, both sexes									
2003.....	18.5	18.6	19.2	17.9	18.8	17.5	23.2	21.8	21.4
2002.....	19.0	16.8	17.2	17.4	17.2	17.3	20.5	23.8	18.8
2001.....	20.7	17.2	18.1	18.9	18.9	18.7	22.3	19.9	17.8
2000.....	19.8	19.6	18.4	16.4	18.6	21.5	19.4	20.7	21.9
1999.....	20.0	18.3	19.1	17.8	18.2	21.1	24.7	28.2	27.8
Non-Hispanic Black, male									
2003.....	30.0	29.4	30.8	28.0	28.9	23.3	38.3	37.7	43.7
2002.....	29.2	26.6	29.6	27.3	28.4	29.1	39.0	43.7	47.0
2001.....	34.5	26.1	29.3	31.3	28.9	25.5	39.8	31.9	35.8
2000.....	30.0	30.3	31.5	27.2	31.7	33.8	34.7	39.5	45.8
1999.....	31.8	29.2	29.8	28.7	27.4	32.2	40.6	48.7	68.9
Non-Hispanic Black, female									
2003.....	8.6	9.5	9.7	10.0	11.3	13.6	14.1	13.4	12.8
2002.....	10.2	8.4	7.0	9.7	8.9	9.4	9.5	13.4	*
2001.....	8.8	9.6	8.8	9.2	11.5	14.2	11.9	13.7	11.1
2000.....	11.0	10.5	7.8	7.9	8.8	13.4	10.2	11.2	13.1
1999.....	9.7	9.1	10.4	9.2	11.4	13.8	15.1	18.0	12.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All origins 3/, both sexes

2003.....	15.1	14.6	14.1	14.3	14.8	17.7	22.5	28.2	28.8
2002.....	15.2	14.4	13.4	15.2	15.3	18.9	23.8	28.4	28.0
2001.....	14.5	13.9	13.7	13.7	14.8	18.0	24.6	28.9	27.8
2000.....	14.6	13.8	14.2	14.7	15.1	18.1	23.8	28.5	30.4
1999.....	14.0	13.1	14.4	13.9	16.4	19.3	24.1	30.1	30.4

All origins 3/, male

2003.....	22.0	20.8	20.3	19.6	20.1	22.8	30.5	40.0	50.0
2002.....	22.2	20.3	19.6	20.2	20.8	25.0	32.3	39.9	51.7
2001.....	21.2	19.7	18.9	19.2	19.2	24.2	33.6	43.1	51.1
2000.....	21.1	19.6	19.9	19.7	19.8	23.9	32.2	41.4	57.5
1999.....	20.0	18.6	19.8	18.7	22.0	24.7	33.6	45.2	59.8

All origins 3/, female

2003.....	8.3	8.7	8.3	9.4	10.3	13.7	16.9	21.0	19.5
2002.....	8.4	8.7	7.5	10.6	10.6	14.1	17.7	21.4	17.7
2001.....	8.0	8.3	8.8	8.8	11.0	13.1	18.2	20.4	18.0
2000.....	8.3	8.2	9.0	10.2	11.0	13.5	17.9	20.9	19.3
1999.....	8.1	8.0	9.4	9.5	11.6	15.0	17.5	21.3	18.5

Hispanic, both sexes

2003.....	14.1	14.2	13.4	16.2	16.4	18.7	20.4	27.5	24.4
2002.....	14.2	13.6	12.7	16.6	16.1	20.0	25.6	22.3	24.9
2001.....	13.8	13.8	14.4	16.0	15.2	20.8	23.5	31.3	20.0
2000.....	14.8	15.4	14.7	14.0	14.2	23.7	21.7	23.4	22.6
1999.....	13.3	14.1	15.5	13.4	20.6	18.6	23.2	24.1	23.2

Hispanic, male

2003.....	21.1	20.3	18.8	21.2	23.4	26.5	29.2	42.6	46.2
2002.....	20.4	18.6	19.3	22.3	26.7	28.5	37.8	28.6	43.5
2001.....	19.9	20.5	18.6	24.7	21.6	28.4	38.2	55.8	*
2000.....	21.0	23.3	21.5	20.7	20.1	33.5	36.9	*	*
1999.....	19.7	20.6	20.1	17.6	29.0	25.0	32.8	45.6	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Hispanic, female

2003.....	7.0	8.3	8.4	11.8	10.7	12.8	14.2	17.6	*
2002.....	7.9	8.7	6.6	11.6	7.5	13.5	16.9	18.2	*
2001.....	7.7	7.4	10.7	8.5	10.0	15.0	13.1	16.1	*
2000.....	8.6	7.9	8.5	8.2	9.4	16.2	11.0	21.5	*
1999.....	7.0	7.9	11.3	9.8	13.8	13.8	16.5	*	*

Non-Hispanic 4/, both sexes

2003.....	15.1	14.6	14.1	14.0	14.6	17.5	22.6	28.2	28.9
2002.....	15.2	14.4	13.4	15.0	15.2	18.8	23.6	28.5	28.1
2001.....	14.5	13.8	13.6	13.5	14.7	17.7	24.6	28.7	28.0
2000.....	14.5	13.6	14.1	14.7	15.1	17.7	23.9	28.6	30.6
1999.....	14.0	13.0	14.2	13.9	16.1	19.2	24.1	30.1	30.5

Non-Hispanic 4/, male

2003.....	22.1	20.7	20.3	19.4	19.7	22.3	30.5	39.8	49.9
2002.....	22.2	20.3	19.5	19.9	20.3	24.6	31.9	40.0	51.9
2001.....	21.2	19.5	18.9	18.6	19.0	23.8	33.3	42.4	51.6
2000.....	21.1	19.2	19.6	19.5	19.7	23.2	32.0	41.7	58.3
1999.....	19.9	18.3	19.5	18.7	21.5	24.6	33.5	44.7	60.7

Non-Hispanic 4/, female

2003.....	8.4	8.8	8.3	9.2	10.2	13.7	17.0	21.1	19.6
2002.....	8.5	8.7	7.6	10.4	10.8	14.1	17.8	21.5	17.8
2001.....	8.0	8.3	8.6	8.8	11.0	12.9	18.4	20.4	18.2
2000.....	8.2	8.2	9.0	10.3	11.1	13.3	18.2	20.9	19.4
1999.....	8.2	7.9	9.2	9.5	11.4	15.1	17.5	21.6	18.5

Non-Hispanic White, both sexes

2003.....	15.2	14.3	13.7	13.7	14.3	17.6	22.6	28.7	29.6
2002.....	15.2	14.3	13.3	14.8	15.0	19.0	24.1	29.2	28.9
2001.....	14.1	13.8	13.4	13.1	14.7	17.6	25.0	29.4	29.0
2000.....	14.3	13.0	13.9	14.7	14.9	17.4	24.1	29.3	31.3
1999.....	13.6	12.7	13.8	13.5	15.9	19.1	24.2	30.6	30.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Non-Hispanic White, male

2003.....	21.9	20.3	19.7	19.0	19.4	22.6	30.2	40.4	50.7
2002.....	22.3	20.0	19.1	19.7	19.5	24.5	31.9	40.2	52.6
2001.....	20.3	19.4	18.5	17.8	18.7	23.5	33.3	43.5	53.1
2000.....	20.7	18.3	18.9	18.9	18.8	22.7	31.7	42.0	60.0
1999.....	19.1	17.7	18.8	18.0	21.1	24.2	33.2	44.9	61.2

Non-Hispanic White, female

2003.....	8.6	8.5	8.0	8.8	9.9	13.5	17.2	21.6	20.3
2002.....	8.2	8.7	7.7	10.3	11.0	14.5	18.5	22.4	18.6
2001.....	8.0	8.2	8.6	8.8	11.1	12.8	19.2	21.0	19.0
2000.....	8.0	7.8	9.1	10.7	11.4	13.2	18.7	21.8	19.8
1999.....	8.1	7.7	9.0	9.3	11.4	15.1	17.9	22.1	18.8

Non-Hispanic Black, both sexes

2003.....	16.6	17.2	17.9	16.4	17.2	17.0	21.6	20.5	21.1
2002.....	17.0	15.5	15.3	16.3	15.8	16.2	18.8	22.4	18.5
2001.....	18.5	15.7	16.2	17.2	16.8	18.0	20.4	19.0	16.9
2000.....	18.0	18.6	17.0	15.5	16.9	20.3	18.1	19.8	20.7
1999.....	18.2	16.4	17.4	15.8	16.7	19.7	22.9	25.8	25.5

Non-Hispanic Black, male

2003.....	26.6	26.5	28.1	24.8	25.6	22.2	36.0	34.7	42.6
2002.....	25.5	24.0	25.8	24.8	25.1	26.3	34.3	40.6	45.9
2001.....	30.6	23.7	25.3	28.8	24.5	23.8	37.0	31.1	35.8
2000.....	27.0	28.5	28.6	25.5	28.2	31.8	32.3	38.6	44.6
1999.....	28.8	25.8	26.7	24.8	25.0	29.1	39.2	45.2	61.8

Non-Hispanic Black, female

2003.....	8.0	9.3	9.5	9.9	11.1	13.4	13.0	13.0	12.8
2002.....	9.6	8.3	6.8	9.5	8.9	9.4	9.5	13.0	*
2001.....	8.1	8.9	8.7	8.1	11.2	14.2	10.5	12.9	9.8
2000.....	10.3	10.2	7.5	7.5	8.5	12.7	9.6	10.4	11.8
1999.....	9.0	8.5	9.8	8.7	10.6	13.6	13.1	16.2	12.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.6	0.5	0.5	0.5	0.4	0.4	0.6	0.7	0.5
2002.....	0.6	0.4	0.4	0.4	0.4	0.4	0.4	0.5	*
2001.....	0.6	0.5	0.4	0.6	0.7	0.6	0.7	0.7	0.6
2000.....	0.6	0.6	0.4	0.5	0.5	0.5	0.7	0.8	0.8
1999.....	0.9	0.7	0.8	0.7	0.7	0.7	1.3	1.3	1.3

Non-Hispanic 4/, male

2003.....	1.1	0.9	0.9	0.8	0.7	0.8	0.9	1.3	*
2002.....	1.0	0.8	0.7	0.7	0.6	0.8	0.8	1.1	*
2001.....	1.0	0.9	0.8	0.8	1.2	0.9	1.0	*	*
2000.....	0.9	0.8	0.7	0.9	0.8	0.9	1.1	1.2	*
1999.....	1.4	1.1	1.3	1.1	1.0	1.0	1.6	2.0	3.0

Non-Hispanic 4/, female

2003.....	0.2	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	0.3	0.2	*	*	*	*	*	0.6	*
2000.....	0.3	0.4	*	*	*	*	*	*	*
1999.....	0.4	0.3	0.4	0.4	0.5	0.4	1.0	1.0	*

Non-Hispanic White, both sexes

2003.....	0.6	0.5	0.5	0.5	0.4	0.4	0.6	0.7	0.5
2002.....	0.5	0.4	0.3	0.4	0.3	0.4	0.4	0.5	*
2001.....	0.6	0.5	0.4	0.5	0.6	0.6	0.7	0.7	0.5
2000.....	0.5	0.6	0.4	0.5	0.4	0.5	0.6	0.8	0.8
1999.....	0.9	0.7	0.8	0.7	0.7	0.7	1.2	1.3	1.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	1.1	1.2	1.2	1.3	1.0	0.8	0.8	0.5	*
2002.....	1.1	1.0	1.1	1.1	1.0	0.9	0.8	0.7	*
2001.....	1.1	1.1	1.0	1.0	0.9	0.9	0.8	0.7	*
2000.....	1.0	1.1	1.1	1.1	1.2	0.7	0.6	0.7	*
1999.....	1.0	1.1	1.2	1.1	0.9	0.9	0.5	0.6	*

Non-Hispanic 4/, male

2003.....	1.9	2.1	2.2	2.4	1.7	1.6	1.6	*	*
2002.....	2.1	1.7	2.0	2.0	2.1	1.8	1.4	1.6	*
2001.....	1.9	1.9	1.7	1.9	1.6	1.7	1.3	1.3	*
2000.....	1.7	1.9	1.9	1.9	2.1	1.4	1.2	1.3	*
1999.....	1.8	1.9	2.1	2.0	1.7	1.6	0.9	*	*

Non-Hispanic 4/, female

2003.....	0.4	0.3	0.3	*	*	*	*	*	*
2002.....	0.3	0.4	0.3	0.4	*	*	*	*	*
2001.....	0.3	0.3	*	*	*	*	*	*	*
2000.....	0.3	0.3	*	0.4	0.5	*	*	*	*
1999.....	0.3	0.4	*	*	*	*	*	*	*

Non-Hispanic White, both sexes

2003.....	1.2	1.3	1.3	1.3	0.9	0.9	0.8	0.5	*
2002.....	1.2	1.1	1.2	1.2	1.0	0.9	0.8	0.7	*
2001.....	1.1	1.1	0.9	1.1	0.9	0.9	0.8	0.7	*
2000.....	1.1	1.2	1.1	1.2	1.2	0.7	0.6	0.7	*
1999.....	1.1	1.1	1.2	1.1	1.0	0.9	0.5	0.5	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nontransport accidents (W00-X59,Y86)

All origins 3/, both sexes

2003.....	23.9	20.3	17.0	16.9	21.9	31.9	54.9	104.1	249.2
2002.....	21.8	18.4	15.5	16.0	21.7	30.6	55.0	101.4	246.7
2001.....	19.7	16.3	14.1	16.1	20.3	29.8	54.3	100.1	247.5
2000.....	18.0	15.3	14.2	15.9	18.6	29.4	50.4	94.6	242.0
1999.....	18.2	14.3	13.9	15.5	20.6	30.0	52.7	101.3	250.3

All origins 3/, male

2003.....	34.1	28.7	24.0	24.3	29.7	41.8	72.0	133.0	300.5
2002.....	31.5	26.8	22.6	22.9	30.0	41.4	71.9	127.5	293.6
2001.....	30.0	24.3	20.5	22.5	27.8	38.8	72.1	127.8	305.8
2000.....	28.0	23.7	20.7	22.7	25.5	38.4	65.3	120.8	299.8
1999.....	28.6	22.0	20.6	22.7	28.3	40.7	66.6	133.5	311.9

All origins 3/, female

2003.....	14.1	12.3	10.4	10.2	15.1	23.9	42.7	86.4	226.5
2002.....	12.5	10.4	8.9	9.7	14.5	21.9	43.1	85.5	226.3
2001.....	9.7	8.6	8.2	10.3	13.9	22.6	41.8	83.5	223.1
2000.....	8.4	7.3	8.0	9.7	12.8	22.4	40.0	79.2	218.4
1999.....	8.1	6.9	7.6	9.0	14.0	21.7	43.0	82.4	225.5

Hispanic, both sexes

2003.....	18.7	16.5	13.3	17.1	17.3	22.4	38.1	62.3	127.0
2002.....	17.7	15.6	13.7	13.1	18.1	24.8	42.4	65.9	127.5
2001.....	16.3	14.0	12.6	12.6	14.4	26.5	34.8	59.1	142.6
2000.....	14.7	13.5	12.4	14.0	14.8	22.8	35.8	70.7	159.2
1999.....	17.8	14.1	11.2	15.3	19.5	21.6	34.5	62.8	157.2

Hispanic, male

2003.....	30.4	25.2	21.7	27.1	24.4	30.7	58.3	82.1	149.3
2002.....	27.9	26.1	22.2	20.3	28.4	32.5	57.0	89.2	179.0
2001.....	27.5	23.1	21.2	19.3	20.2	35.2	47.1	67.5	154.3
2000.....	26.1	22.0	18.9	20.7	23.5	33.1	47.2	78.0	207.6
1999.....	30.9	23.2	20.6	25.0	29.8	34.1	42.1	80.2	217.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)									
Hispanic, female									
2003.....	6.6	8.0	5.4	8.4	11.5	16.1	23.7	49.4	115.6
2002.....	7.5	5.4	5.9	6.9	9.7	18.9	32.0	51.1	102.0
2001.....	5.1	5.3	4.9	6.8	9.8	19.9	26.1	53.9	136.8
2000.....	3.4	5.5	6.5	8.2	7.9	15.1	27.7	66.3	135.5
1999.....	4.9	5.5	*	6.9	11.3	12.3	29.2	52.5	127.9
Non-Hispanic 4/, both sexes									
2003.....	24.3	20.5	17.2	16.8	22.1	32.4	55.7	105.9	253.9
2002.....	22.1	18.6	15.5	16.1	21.8	30.8	55.4	102.7	250.9
2001.....	19.8	16.4	14.1	16.2	20.6	29.8	55.1	101.4	250.8
2000.....	18.1	15.2	14.2	15.9	18.8	29.7	50.8	95.3	244.4
1999.....	18.0	14.1	13.8	15.4	20.5	30.4	53.3	102.3	252.8
Non-Hispanic 4/, male									
2003.....	34.2	28.7	24.0	24.0	29.9	42.4	72.4	135.3	306.9
2002.....	31.7	26.6	22.3	22.9	29.8	41.7	72.4	128.6	298.0
2001.....	29.9	24.2	20.1	22.6	28.1	38.8	73.1	129.8	311.6
2000.....	27.8	23.5	20.6	22.7	25.4	38.5	65.9	122.2	302.7
1999.....	28.0	21.6	20.2	22.3	28.0	40.9	67.4	134.9	314.3
Non-Hispanic 4/, female									
2003.....	14.8	12.7	10.8	10.3	15.3	24.4	43.7	87.9	230.6
2002.....	12.9	10.8	9.1	9.9	14.8	22.1	43.5	86.9	230.6
2001.....	10.1	8.9	8.4	10.5	14.1	22.7	42.5	84.4	225.5
2000.....	8.8	7.3	8.1	9.7	13.1	22.7	40.4	79.5	220.8
1999.....	8.4	6.9	7.9	9.1	14.1	22.1	43.4	83.2	228.2
Non-Hispanic White, both sexes									
2003.....	24.4	19.6	16.6	16.5	21.5	32.3	57.0	108.4	265.9
2002.....	21.8	17.7	14.6	15.6	20.9	30.7	56.0	105.6	261.2
2001.....	19.0	15.3	13.3	15.5	20.3	29.4	55.4	103.6	259.4
2000.....	17.2	13.9	13.2	15.3	18.0	29.1	50.9	97.3	253.2
1999.....	16.7	12.9	12.8	13.9	19.5	29.0	52.5	103.7	260.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)									
Non-Hispanic White, male									
2003.....	33.8	27.1	22.7	23.0	28.7	42.0	73.5	137.2	320.0
2002.....	31.1	24.8	20.4	21.7	28.0	41.3	72.6	131.2	307.6
2001.....	28.2	22.0	18.4	21.1	27.5	38.0	73.0	130.7	320.4
2000.....	25.9	21.1	18.8	21.4	23.8	37.6	64.9	123.5	311.5
1999.....	25.6	19.4	18.0	20.3	26.0	38.7	65.5	136.4	322.6
Non-Hispanic White, female									
2003.....	15.1	12.3	10.8	10.5	15.1	24.4	45.1	90.8	242.0
2002.....	12.6	10.9	9.1	9.9	14.6	22.0	44.1	90.0	241.1
2001.....	9.8	8.7	8.4	10.3	14.0	22.3	43.0	87.2	234.0
2000.....	8.5	6.9	7.9	9.6	12.9	22.2	41.0	81.7	229.7
1999.....	8.0	6.5	7.8	8.0	13.9	21.3	43.4	84.3	236.2
Non-Hispanic Black, both sexes									
2003.....	30.4	31.1	25.3	22.9	29.9	36.6	45.3	83.2	140.4
2002.....	30.1	29.1	25.5	23.4	30.2	35.3	55.5	73.9	156.7
2001.....	30.1	28.5	23.1	24.9	27.0	39.2	56.9	85.8	166.1
2000.....	28.1	28.0	24.3	23.2	28.3	38.9	56.5	79.6	161.2
1999.....	31.3	26.2	24.9	28.6	33.1	46.4	66.0	96.9	181.1
Non-Hispanic Black, male									
2003.....	45.7	46.6	40.5	37.7	46.6	50.9	67.7	114.7	184.3
2002.....	43.8	47.1	42.5	38.1	47.1	50.3	80.3	107.6	215.5
2001.....	48.6	47.1	37.7	38.5	40.9	56.1	81.5	135.7	220.6
2000.....	46.2	47.1	41.1	38.1	43.4	53.3	85.7	116.7	231.3
1999.....	52.3	44.0	43.0	41.8	51.5	71.3	95.7	129.3	249.6
Non-Hispanic Black, female									
2003.....	17.2	18.1	12.8	11.3	17.5	27.0	31.9	66.7	123.4
2002.....	18.2	14.0	11.6	11.9	17.6	25.2	40.6	56.4	134.2
2001.....	14.2	12.7	11.2	14.3	16.7	28.0	42.2	60.4	145.9
2000.....	12.5	11.8	10.5	11.4	17.1	29.3	38.9	60.9	135.2
1999.....	13.2	11.1	10.1	18.3	19.5	29.9	48.1	80.8	155.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)									
All origins 3/, both sexes									
2003.....	2.3	2.8	3.7	5.3	8.1	14.7	28.7	57.5	135.9
2002.....	2.3	2.8	3.4	4.9	7.9	13.9	27.1	54.0	130.4
2001.....	2.5	2.7	3.3	4.9	7.8	12.4	25.2	49.9	121.5
2000.....	2.0	2.6	3.3	4.6	6.2	12.0	22.1	44.5	112.6
1999.....	2.3	2.2	3.0	4.6	6.9	11.3	22.3	45.5	110.2
All origins 3/, male									
2003.....	3.6	4.4	5.3	7.3	10.7	18.9	36.9	73.8	165.1
2002.....	3.6	4.3	5.2	7.1	10.7	18.9	35.1	67.9	158.9
2001.....	4.2	4.4	5.0	7.1	10.6	16.0	34.0	64.1	155.7
2000.....	3.4	4.2	5.0	6.9	8.6	15.7	29.4	57.3	145.4
1999.....	3.8	3.6	4.7	7.0	9.8	15.3	27.9	61.6	142.5
All origins 3/, female									
2003.....	1.1	1.3	2.1	3.5	5.8	11.3	22.8	47.5	122.9
2002.....	1.0	1.4	1.8	3.0	5.5	9.9	21.4	45.5	118.1
2001.....	0.9	1.2	1.6	2.9	5.4	9.5	19.0	41.3	107.2
2000.....	0.8	1.2	1.8	2.6	4.2	9.1	17.1	36.9	99.2
1999.....	0.8	0.9	1.5	2.4	4.5	8.1	18.4	36.0	97.2
Hispanic, both sexes									
2003.....	2.3	3.0	3.4	7.3	6.2	13.2	19.9	34.4	80.8
2002.....	2.1	2.9	4.0	5.0	7.0	13.7	21.3	42.3	71.5
2001.....	2.7	2.4	3.8	4.8	5.5	13.5	19.0	31.8	70.0
2000.....	2.2	2.8	3.6	5.2	6.7	9.6	18.4	41.2	81.0
1999.....	2.0	2.4	3.3	4.5	8.1	8.1	17.8	35.8	86.3
Hispanic, male									
2003.....	3.9	4.9	5.1	10.8	8.1	16.7	31.0	44.7	90.8
2002.....	3.4	4.5	6.6	7.4	11.2	17.1	25.6	57.2	97.0
2001.....	5.1	3.6	6.4	8.1	7.4	16.5	22.5	37.7	72.4
2000.....	3.9	5.1	5.9	7.5	10.4	15.6	25.1	41.2	104.8
1999.....	3.3	4.5	6.5	8.0	10.8	12.2	22.6	48.7	120.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)									
Hispanic, female									
2003.....	*	*	*	4.3	*	10.5	12.1	27.7	75.8
2002.....	*	*	*	*	*	11.2	18.3	32.8	58.9
2001.....	*	*	*	*	*	11.2	16.5	28.1	68.9
2000.....	*	*	*	*	*	*	13.6	41.2	69.2
1999.....	*	*	*	*	*	*	14.3	28.1	69.7
Non-Hispanic 4/, both sexes									
2003.....	2.3	2.8	3.7	5.1	8.2	14.7	29.1	58.4	137.9
2002.....	2.3	2.8	3.4	4.9	8.0	13.9	27.3	54.3	132.6
2001.....	2.4	2.8	3.2	4.8	7.9	12.3	25.5	50.4	123.1
2000.....	2.0	2.6	3.3	4.6	6.2	12.1	22.2	44.5	113.5
1999.....	2.3	2.2	3.0	4.5	6.8	11.4	22.3	45.6	110.7
Non-Hispanic 4/, male									
2003.....	3.6	4.3	5.3	7.0	10.9	19.0	37.1	75.1	168.3
2002.....	3.6	4.2	5.1	7.0	10.6	18.9	35.5	68.0	161.3
2001.....	4.0	4.5	4.9	6.9	10.7	15.8	34.6	64.9	159.0
2000.....	3.3	4.0	4.9	6.8	8.4	15.7	29.5	57.9	146.5
1999.....	3.8	3.5	4.5	6.8	9.7	15.4	27.9	61.7	142.9
Non-Hispanic 4/, female									
2003.....	1.1	1.3	2.1	3.4	5.9	11.3	23.3	48.3	124.6
2002.....	1.0	1.4	1.8	3.0	5.7	9.9	21.5	46.0	120.2
2001.....	0.9	1.1	1.6	2.9	5.5	9.4	19.2	41.7	108.1
2000.....	0.8	1.2	1.8	2.5	4.3	9.3	17.2	36.7	100.1
1999.....	0.8	1.0	1.6	2.4	4.4	8.3	18.5	36.2	97.8
Non-Hispanic White, both sexes									
2003.....	2.3	2.7	3.6	5.2	8.3	15.2	30.6	61.1	146.8
2002.....	2.3	2.8	3.3	5.0	8.1	14.3	28.5	57.5	140.4
2001.....	2.4	2.8	3.1	4.9	8.1	12.7	26.6	53.1	129.9
2000.....	2.1	2.5	3.2	4.5	6.1	12.5	23.2	46.7	120.2
1999.....	2.2	2.1	2.8	4.3	6.8	11.5	23.1	47.9	116.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)									
Non-Hispanic White, male									
2003.....	3.5	4.0	5.2	7.0	10.7	19.6	38.8	77.9	178.7
2002.....	3.6	4.1	4.9	7.1	10.6	19.3	36.7	71.3	169.2
2001.....	3.8	4.4	4.7	6.9	10.9	16.3	35.7	67.5	166.0
2000.....	3.4	3.7	4.6	6.8	8.2	16.2	30.4	60.3	154.1
1999.....	3.6	3.2	4.1	6.4	9.4	15.4	28.4	64.4	149.7
Non-Hispanic White, female									
2003.....	1.0	1.3	2.1	3.5	6.2	11.7	24.6	50.8	132.8
2002.....	1.0	1.5	1.8	3.2	5.9	10.2	22.6	49.1	128.0
2001.....	1.0	1.2	1.6	3.0	5.6	9.7	20.1	44.4	114.8
2000.....	0.8	1.2	1.8	2.5	4.3	9.5	18.2	38.6	106.5
1999.....	0.8	0.9	1.5	2.3	4.5	8.4	19.4	38.1	103.4
Non-Hispanic Black, both sexes									
2003.....	2.9	3.6	4.4	5.5	7.7	10.7	13.2	28.3	44.6
2002.....	2.6	3.2	3.9	4.8	6.1	10.8	14.9	18.4	49.3
2001.....	2.6	3.4	4.0	4.9	8.0	9.8	16.7	21.8	44.0
2000.....	2.0	3.6	4.6	5.1	7.1	9.7	11.8	20.7	39.1
1999.....	3.3	3.7	4.5	6.7	9.0	9.7	13.3	23.5	45.3
Non-Hispanic Black, male									
2003.....	4.7	5.8	6.7	7.9	12.9	12.1	17.8	35.5	56.5
2002.....	4.1	5.5	6.3	7.5	9.2	15.4	22.5	28.9	76.6
2001.....	4.9	6.1	6.4	7.5	12.0	13.4	24.6	36.8	68.1
2000.....	3.3	6.3	8.0	8.5	10.7	12.3	18.8	26.6	61.1
1999.....	5.7	6.1	7.5	10.2	15.1	14.9	19.8	31.9	58.2
Non-Hispanic Black, female									
2003.....	*	*	2.6	3.6	3.8	9.8	10.5	24.5	40.0
2002.....	*	*	*	*	3.8	7.8	10.4	13.0	38.9
2001.....	*	*	*	*	5.0	7.3	11.9	14.2	35.1
2000.....	*	*	*	*	4.5	7.9	7.6	17.7	31.0
1999.....	*	*	*	3.9	4.5	6.2	9.3	19.3	40.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental discharge of firearms (W32-W34)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.2	0.2	0.2	0.2	0.2	*	*	0.4	*
2002.....	0.3	0.2	0.3	0.2	*	*	*	*	*
2001.....	0.3	0.2	0.3	0.2	*	*	0.3	*	*
2000.....	0.2	0.2	0.3	*	0.3	*	*	*	*
1999.....	0.3	0.3	0.2	0.2	*	*	*	*	*

Non-Hispanic 4/, male

2003.....	0.4	0.4	0.5	0.4	*	*	*	*	*
2002.....	0.5	0.5	0.5	*	*	*	*	*	*
2001.....	0.5	0.4	0.5	0.4	*	*	0.7	*	*
2000.....	0.3	0.5	0.5	*	0.5	*	*	*	*
1999.....	0.5	0.4	0.4	*	*	*	*	*	*

Non-Hispanic 4/, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic White, both sexes

2003.....	0.2	0.3	0.3	0.2	0.3	*	*	*	*
2002.....	0.3	0.3	0.3	*	*	*	*	*	*
2001.....	0.3	0.3	0.3	0.3	*	*	0.3	*	*
2000.....	0.2	0.3	0.3	*	0.3	*	*	*	*
1999.....	0.3	0.3	0.2	0.2	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental drowning and submersion (W65-W74)									
Hispanic, female									
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes									
2003.....	1.0	1.0	1.0	0.9	1.0	1.2	1.3	1.3	1.8
2002.....	1.0	0.9	0.8	0.8	0.9	1.0	1.1	1.5	1.7
2001.....	1.0	0.8	0.8	0.9	0.8	1.0	1.2	1.4	1.7
2000.....	0.9	1.0	0.9	0.9	0.9	1.1	1.0	1.7	1.5
1999.....	1.0	1.0	1.0	0.9	0.9	1.2	1.2	1.5	1.7
Non-Hispanic 4/, male									
2003.....	1.6	1.6	1.6	1.5	1.6	2.0	1.9	2.3	3.4
2002.....	1.6	1.5	1.3	1.2	1.4	1.7	1.7	2.4	3.6
2001.....	1.8	1.4	1.3	1.4	1.4	1.7	1.8	2.5	3.4
2000.....	1.6	1.7	1.6	1.6	1.5	1.7	1.3	2.7	2.8
1999.....	1.7	1.7	1.6	1.5	1.5	1.6	2.0	2.8	2.4
Non-Hispanic 4/, female									
2003.....	0.4	0.4	0.4	0.4	0.5	0.6	0.9	*	1.1
2002.....	0.4	0.4	0.3	0.4	*	*	0.7	0.9	0.9
2001.....	0.3	0.3	0.4	0.5	*	*	0.7	0.8	1.0
2000.....	0.3	0.3	*	*	*	0.6	0.8	1.1	1.0
1999.....	0.3	0.3	0.4	0.4	0.4	0.8	0.6	0.7	1.4
Non-Hispanic White, both sexes									
2003.....	1.0	1.0	1.0	0.9	1.1	1.2	1.2	1.3	1.7
2002.....	0.9	0.8	0.8	0.8	0.8	0.9	1.1	1.4	1.6
2001.....	0.9	0.8	0.7	0.9	0.8	1.0	1.2	1.4	1.7
2000.....	0.9	0.9	0.9	0.9	0.9	1.0	0.9	1.8	1.4
1999.....	1.1	1.0	0.9	0.9	0.9	1.0	1.1	1.5	1.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental exposure to smoke, fire and flames (X00-X09)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	1.3	1.3	1.3	1.6	2.1	2.7	3.0	4.4	5.4
2002.....	1.0	1.3	1.2	1.4	1.8	2.5	2.9	4.2	4.8
2001.....	1.1	1.2	1.6	1.5	2.1	2.6	3.1	3.9	5.7
2000.....	1.2	1.3	1.4	1.7	2.0	2.4	3.4	4.3	6.3
1999.....	1.0	1.0	1.2	1.5	2.1	2.6	3.5	4.7	6.0

Non-Hispanic 4/, male

2003.....	1.8	1.6	1.8	2.1	2.5	3.5	4.0	5.1	7.3
2002.....	1.5	1.6	1.4	2.0	2.6	3.3	3.6	6.2	7.0
2001.....	1.5	1.6	2.1	1.8	2.6	3.1	3.5	5.2	8.3
2000.....	1.6	1.9	2.0	2.3	2.5	3.2	4.4	5.9	9.3
1999.....	1.5	1.4	1.6	1.8	2.5	3.5	4.2	6.6	9.2

Non-Hispanic 4/, female

2003.....	0.9	1.1	0.9	1.2	1.7	2.0	2.3	4.0	4.5
2002.....	0.6	1.0	0.9	0.9	1.1	1.8	2.4	2.9	3.8
2001.....	0.6	0.7	1.1	1.2	1.6	2.2	2.8	3.1	4.6
2000.....	0.7	0.8	0.9	1.2	1.7	1.7	2.8	3.4	5.0
1999.....	0.6	0.7	0.9	1.2	1.7	1.9	3.1	3.6	4.7

Non-Hispanic White, both sexes

2003.....	1.2	1.2	1.2	1.5	1.8	2.3	2.6	3.8	4.7
2002.....	1.0	1.1	1.0	1.1	1.6	2.1	2.5	3.7	3.9
2001.....	0.9	1.0	1.3	1.3	1.7	2.2	2.6	3.3	4.7
2000.....	1.0	1.2	1.1	1.4	1.7	2.0	2.8	3.5	5.1
1999.....	0.9	0.9	1.1	1.2	1.7	2.1	2.8	3.9	4.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)									
Non-Hispanic White, male									
2003.....	1.6	1.4	1.6	1.9	2.2	3.1	3.2	4.2	6.5
2002.....	1.4	1.4	1.3	1.7	2.2	2.7	3.1	5.8	5.7
2001.....	1.3	1.3	1.8	1.5	2.2	2.5	2.8	4.5	6.9
2000.....	1.4	1.7	1.6	1.8	2.0	2.6	3.6	4.9	8.1
1999.....	1.3	1.2	1.4	1.6	2.0	2.7	3.2	5.7	7.8
Non-Hispanic White, female									
2003.....	0.8	0.9	0.8	1.0	1.5	1.7	2.2	3.5	3.9
2002.....	0.6	0.9	0.8	0.7	1.0	1.6	2.0	2.4	3.2
2001.....	0.6	0.7	0.9	1.1	1.4	1.9	2.5	2.6	3.8
2000.....	0.6	0.7	0.7	1.0	1.5	1.4	2.3	2.6	3.9
1999.....	0.5	0.6	0.9	0.8	1.4	1.5	2.6	2.8	3.5
Non-Hispanic Black, both sexes									
2003.....	2.4	2.9	2.9	3.0	4.5	6.5	8.7	12.7	14.9
2002.....	1.6	2.8	2.6	4.1	3.9	6.5	7.4	10.7	16.0
2001.....	2.3	2.6	3.8	3.3	5.1	7.3	8.7	11.9	17.8
2000.....	2.2	2.5	3.9	4.6	5.1	7.2	11.1	16.1	20.7
1999.....	2.1	2.3	2.7	4.4	5.9	8.4	12.2	15.6	22.0
Non-Hispanic Black, male									
2003.....	3.5	3.2	4.8	4.0	6.6	9.1	15.4	17.4	*
2002.....	2.2	4.2	3.4	6.1	6.4	9.6	9.4	*	26.3
2001.....	3.6	4.4	5.2	5.0	7.3	11.4	12.3	*	26.6
2000.....	3.2	4.0	5.6	7.2	7.7	10.6	15.9	21.5	28.2
1999.....	3.1	3.4	4.2	4.4	8.0	13.5	17.4	20.4	29.7
Non-Hispanic Black, female									
2003.....	1.5	2.6	*	*	*	4.8	*	10.3	13.2
2002.....	*	*	*	*	*	*	6.2	8.9	12.1
2001.....	*	*	2.8	*	*	4.6	6.5	10.4	14.5
2000.....	*	*	*	*	*	5.0	8.1	13.4	17.9
1999.....	*	*	*	4.4	4.3	5.1	9.0	13.2	19.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)									
All origins 3/, both sexes									
2003.....	15.4	10.9	6.1	3.4	2.3	2.0	2.0	2.5	3.5
2002.....	13.8	9.2	5.0	3.0	2.3	1.9	2.1	2.7	3.9
2001.....	10.9	7.0	3.7	2.4	1.7	1.5	2.1	2.4	3.5
2000.....	9.9	6.1	3.3	2.2	1.5	1.5	1.6	2.6	4.3
1999.....	9.8	5.4	3.4	2.0	1.7	1.6	1.9	2.6	3.4
All origins 3/, male									
2003.....	20.8	14.7	7.6	4.5	3.0	2.2	2.3	3.5	4.0
2002.....	18.7	12.8	6.8	3.6	2.8	2.1	2.3	3.2	5.1
2001.....	15.7	9.9	5.0	3.0	2.0	2.1	2.7	2.8	3.8
2000.....	14.7	8.9	4.0	2.7	1.8	1.7	1.6	2.9	6.2
1999.....	15.1	7.9	4.7	3.0	2.2	1.9	2.1	2.9	4.4
All origins 3/, female									
2003.....	10.2	7.3	4.6	2.5	1.8	1.8	1.7	2.0	3.3
2002.....	9.1	5.8	3.4	2.4	1.8	1.7	1.9	2.3	3.4
2001.....	6.2	4.1	2.6	1.9	1.4	1.0	1.6	2.2	3.3
2000.....	5.3	3.4	2.7	1.8	1.3	1.3	1.6	2.4	3.6
1999.....	4.7	3.1	2.2	1.2	1.2	1.4	1.7	2.4	3.0
Hispanic, both sexes									
2003.....	11.9	8.6	5.2	3.2	*	*	*	*	*
2002.....	11.7	7.9	4.7	3.1	*	*	*	*	*
2001.....	8.6	6.6	4.7	*	*	*	*	*	*
2000.....	8.9	6.8	4.0	*	*	*	*	*	*
1999.....	11.4	7.6	4.1	2.9	*	*	*	*	*
Hispanic, male									
2003.....	18.8	13.2	8.6	5.7	*	*	*	*	*
2002.....	17.6	13.4	7.2	5.4	*	*	*	*	*
2001.....	14.0	11.5	8.0	*	*	*	*	*	*
2000.....	15.7	10.8	5.3	*	*	*	*	*	*
1999.....	19.7	12.0	7.6	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)									
Hispanic, female									
2003.....	4.9	4.2	*	*	*	*	*	*	*
2002.....	5.7	2.6	*	*	*	*	*	*	*
2001.....	3.1	*	*	*	*	*	*	*	*
2000.....	*	3.0	*	*	*	*	*	*	*
1999.....	3.1	3.5	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes									
2003.....	15.7	11.0	6.1	3.4	2.4	2.0	2.0	2.6	3.5
2002.....	14.0	9.3	5.0	2.9	2.2	1.9	2.0	2.7	4.1
2001.....	11.0	6.9	3.6	2.4	1.7	1.5	2.1	2.4	3.6
2000.....	9.9	5.9	3.2	2.2	1.5	1.5	1.6	2.6	4.4
1999.....	9.5	5.1	3.3	1.9	1.6	1.6	1.9	2.6	3.5
Non-Hispanic 4/, male									
2003.....	20.9	14.6	7.4	4.3	2.9	2.2	2.3	3.6	4.0
2002.....	18.7	12.7	6.6	3.4	2.7	2.0	2.2	3.2	5.3
2001.....	15.7	9.6	4.6	2.9	1.9	2.0	2.7	2.9	3.9
2000.....	14.4	8.5	3.8	2.7	1.7	1.7	1.7	2.9	6.2
1999.....	14.4	7.3	4.4	2.7	2.1	1.9	2.1	3.0	4.4
Non-Hispanic 4/, female									
2003.....	10.6	7.6	4.8	2.6	1.9	1.9	1.7	2.0	3.3
2002.....	9.4	6.1	3.5	2.5	1.8	1.7	2.0	2.3	3.5
2001.....	6.5	4.3	2.6	1.9	1.5	1.0	1.7	2.2	3.4
2000.....	5.5	3.3	2.6	1.8	1.4	1.3	1.6	2.4	3.6
1999.....	4.8	3.0	2.3	1.3	1.2	1.4	1.8	2.4	3.1
Non-Hispanic White, both sexes									
2003.....	15.9	10.6	5.9	3.4	2.0	2.0	2.0	2.8	3.5
2002.....	13.7	8.7	4.5	2.7	2.2	1.9	2.0	2.7	4.1
2001.....	10.5	6.0	3.2	2.1	1.6	1.4	2.0	2.3	3.4
2000.....	9.2	5.0	2.9	2.1	1.5	1.4	1.7	2.7	4.1
1999.....	8.7	4.4	3.0	1.6	1.5	1.5	2.0	2.7	3.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)									
Non-Hispanic White, male									
2003.....	20.9	13.7	6.8	4.1	2.1	2.1	2.3	3.7	3.6
2002.....	18.2	11.4	5.5	2.7	2.5	1.9	2.2	2.9	5.4
2001.....	14.7	8.0	3.8	2.3	1.8	1.9	2.4	2.7	3.8
2000.....	13.2	7.0	3.0	2.3	1.4	1.6	1.7	3.0	5.8
1999.....	12.9	5.9	3.6	2.0	1.9	1.7	2.2	3.1	4.6
Non-Hispanic White, female									
2003.....	11.0	7.6	5.0	2.7	1.8	1.9	1.8	2.2	3.5
2002.....	9.4	6.1	3.5	2.7	1.9	1.8	1.9	2.5	3.5
2001.....	6.3	4.0	2.7	1.9	1.5	0.9	1.8	2.0	3.2
2000.....	5.3	3.0	2.7	1.9	1.5	1.3	1.7	2.5	3.4
1999.....	4.5	2.9	2.4	1.2	1.2	1.2	1.8	2.4	2.8
Non-Hispanic Black, both sexes									
2003.....	19.2	17.1	9.7	4.9	5.5	*	*	*	*
2002.....	19.8	16.5	10.6	6.0	2.8	*	*	*	*
2001.....	18.1	15.6	7.3	5.2	2.5	*	*	*	6.6
2000.....	16.8	14.3	6.9	3.8	2.6	*	*	*	8.3
1999.....	18.2	12.2	6.5	5.4	2.8	3.3	*	*	*
Non-Hispanic Black, male									
2003.....	27.3	25.8	15.8	8.9	10.4	*	*	*	*
2002.....	27.9	26.2	18.3	11.2	5.1	*	*	*	*
2001.....	28.4	25.2	12.6	9.0	*	*	*	*	*
2000.....	27.0	23.2	11.7	6.8	*	*	*	*	*
1999.....	30.0	20.7	12.2	9.6	*	*	*	*	*
Non-Hispanic Black, female									
2003.....	12.3	9.8	4.6	*	*	*	*	*	*
2002.....	12.8	8.3	4.4	*	*	*	*	*	*
2001.....	9.2	7.5	3.0	*	*	*	*	*	*
2000.....	8.1	6.7	3.0	*	*	*	*	*	*
1999.....	8.0	4.9	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)									
All origins 3/, both sexes									
2003.....	3.8	4.0	4.7	5.5	8.1	11.2	19.8	38.1	102.5
2002.....	3.5	4.0	4.8	5.7	8.6	11.1	21.6	38.8	105.6
2001.....	3.9	4.5	4.6	6.3	7.8	12.3	22.5	42.2	115.0
2000.....	3.9	4.1	5.0	6.3	7.8	12.2	22.0	41.4	117.1
1999.....	3.8	4.4	5.1	6.3	8.9	13.3	23.6	46.9	128.9
All origins 3/, male									
2003.....	6.0	6.1	7.1	8.7	11.4	15.1	26.5	47.6	120.2
2002.....	5.8	6.2	7.4	8.7	12.0	15.0	28.6	47.2	118.6
2001.....	6.3	6.8	6.7	8.9	10.9	15.9	29.3	52.7	134.0
2000.....	6.5	6.7	7.7	9.1	10.8	15.6	28.1	51.6	135.2
1999.....	6.0	7.2	7.8	9.1	12.0	18.0	29.9	59.3	152.7
All origins 3/, female									
2003.....	1.6	2.1	2.5	2.7	5.3	8.2	15.0	32.3	94.7
2002.....	1.3	1.9	2.4	3.0	5.7	8.1	16.6	33.8	100.0
2001.....	1.6	2.2	2.5	3.8	5.2	9.5	17.7	35.9	107.0
2000.....	1.4	1.7	2.4	3.8	5.2	9.6	17.8	35.4	109.8
1999.....	1.7	1.7	2.5	3.7	6.3	9.6	19.3	39.6	119.3
Hispanic, both sexes									
2003.....	2.9	3.1	3.2	4.9	6.7	5.7	12.9	23.4	37.3
2002.....	2.4	3.3	3.0	3.9	6.4	5.9	14.9	16.0	48.2
2001.....	3.2	4.1	3.3	4.4	5.9	10.1	12.7	21.9	63.8
2000.....	2.8	2.6	3.2	4.9	5.3	7.3	13.5	22.3	70.3
1999.....	2.9	2.6	3.0	5.8	7.1	10.7	11.3	21.7	65.9
Hispanic, male									
2003.....	4.9	4.4	5.3	7.9	9.1	9.0	21.3	31.2	44.6
2002.....	4.3	5.4	5.3	5.9	9.1	*	21.1	*	70.3
2001.....	5.3	7.1	5.4	6.0	8.5	13.7	17.7	26.0	68.6
2000.....	4.8	4.1	5.3	8.3	7.8	9.7	16.2	*	88.7
1999.....	5.3	4.2	5.3	7.4	10.4	16.8	*	*	84.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)									
Hispanic, female									
2003.....	*	*	*	*	*	*	*	18.3	33.6
2002.....	*	*	*	*	*	*	10.5	*	37.3
2001.....	*	*	*	*	*	7.3	*	19.3	61.4
2000.....	*	*	*	*	*	*	11.5	19.7	61.3
1999.....	*	*	*	*	*	*	11.6	19.7	57.2
Non-Hispanic 4/, both sexes									
2003.....	3.8	4.1	4.9	5.6	8.2	11.6	20.1	38.7	105.1
2002.....	3.6	4.0	4.9	5.8	8.8	11.4	21.9	39.8	107.6
2001.....	4.0	4.5	4.6	6.4	7.9	12.4	22.9	42.9	116.6
2000.....	4.0	4.2	5.1	6.4	7.9	12.4	22.3	42.0	118.5
1999.....	3.9	4.5	5.1	6.3	9.0	13.4	24.1	47.7	130.7
Non-Hispanic 4/, male									
2003.....	6.0	6.2	7.2	8.7	11.6	15.5	26.6	48.2	123.5
2002.....	5.9	6.2	7.4	8.9	12.2	15.4	29.0	48.2	120.4
2001.....	6.4	6.8	6.7	9.1	11.0	15.9	29.8	53.6	136.5
2000.....	6.6	6.8	7.9	9.1	10.9	15.8	28.5	52.4	136.9
1999.....	6.1	7.3	7.8	9.1	12.0	18.0	30.7	60.3	154.8
Non-Hispanic 4/, female									
2003.....	1.7	2.1	2.6	2.7	5.3	8.5	15.5	32.9	97.0
2002.....	1.3	1.9	2.6	3.1	5.8	8.3	16.9	34.7	102.2
2001.....	1.6	2.3	2.6	3.9	5.2	9.6	18.1	36.5	108.4
2000.....	1.4	1.7	2.4	3.9	5.4	9.8	18.0	35.8	111.1
1999.....	1.8	1.8	2.7	3.7	6.4	9.8	19.5	40.3	121.1
Non-Hispanic White, both sexes									
2003.....	3.8	3.9	4.7	5.4	8.0	11.4	20.3	39.1	109.1
2002.....	3.6	4.0	4.7	5.7	8.0	11.3	21.6	40.1	110.9
2001.....	3.8	4.4	4.5	6.0	7.8	12.0	22.7	43.2	119.5
2000.....	3.8	4.1	4.9	6.2	7.5	12.0	21.9	42.5	122.0
1999.....	3.6	4.3	4.7	5.7	8.6	12.7	23.3	47.6	134.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)									
Non-Hispanic White, male									
2003.....	5.8	6.0	7.0	8.2	11.4	15.0	26.7	48.1	127.5
2002.....	6.1	6.0	7.0	8.7	10.9	15.2	28.4	48.2	123.1
2001.....	6.2	6.5	6.4	8.7	10.8	15.3	29.6	52.7	139.7
2000.....	6.1	6.5	7.4	8.6	10.2	15.3	27.5	52.0	139.7
1999.....	5.5	7.0	7.1	8.5	11.2	16.9	29.4	60.1	157.3
Non-Hispanic White, female									
2003.....	1.7	2.0	2.5	2.7	5.0	8.5	15.8	33.5	101.0
2002.....	1.2	2.0	2.6	2.9	5.3	8.0	16.8	35.1	105.7
2001.....	1.5	2.4	2.7	3.6	5.1	9.2	17.9	37.4	111.2
2000.....	1.5	1.7	2.4	4.0	5.2	9.4	18.0	36.9	114.9
1999.....	1.7	1.7	2.5	3.2	6.3	9.4	19.0	40.3	125.2
Non-Hispanic Black, both sexes									
2003.....	4.8	6.4	6.9	8.3	11.5	15.1	20.0	39.4	74.1
2002.....	4.4	5.2	7.3	7.4	16.3	14.7	28.4	39.8	84.4
2001.....	5.6	6.0	6.1	10.5	10.5	18.9	27.5	44.7	96.5
2000.....	5.8	5.8	8.1	8.8	12.6	18.1	30.8	39.4	90.9
1999.....	6.7	6.6	9.5	11.0	14.1	23.2	36.3	54.0	106.7
Non-Hispanic Black, male									
2003.....	8.0	9.5	10.7	14.3	15.7	22.9	29.9	55.8	92.7
2002.....	6.6	8.8	12.6	11.4	24.3	19.2	41.8	54.6	107.2
2001.....	8.9	9.7	10.1	14.8	15.9	25.1	35.6	73.6	115.5
2000.....	10.4	10.3	14.2	14.2	18.7	23.6	47.7	64.4	124.5
1999.....	11.3	11.0	15.9	16.1	21.3	35.3	51.2	70.0	155.7
Non-Hispanic Black, female									
2003.....	2.0	3.7	3.8	3.6	8.4	9.8	14.1	30.8	66.9
2002.....	2.6	2.1	3.0	4.3	10.4	11.7	20.5	32.1	75.7
2001.....	2.7	2.8	2.8	7.1	6.5	14.7	22.6	30.0	89.4
2000.....	1.8	*	3.0	4.5	8.1	14.5	20.6	26.8	78.5
1999.....	2.8	2.9	4.3	7.0	8.9	15.2	27.4	46.1	88.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All origins 3/, both sexes

2003.....	16.0	15.7	14.7	12.6	12.0	13.6	15.6	17.7	16.9
2002.....	16.3	15.1	14.6	12.3	12.5	14.6	16.6	19.3	18.0
2001 5/.....	15.7	14.6	14.0	12.0	12.7	13.9	16.4	18.9	17.5
2000.....	14.9	13.9	12.7	11.4	11.3	13.8	16.5	19.3	19.6
1999.....	14.3	13.4	12.8	11.4	12.4	14.5	17.4	19.2	19.3

All origins 3/, male

2003.....	24.4	24.4	23.1	21.1	21.2	26.0	31.8	40.1	47.8
2002.....	24.9	23.8	23.5	20.4	22.0	27.9	34.3	44.0	50.7
2001 5/.....	24.3	22.4	22.3	19.5	22.7	26.7	33.8	44.2	51.1
2000.....	23.3	21.4	20.1	18.7	19.8	26.1	34.9	44.8	57.5
1999.....	22.3	20.6	20.3	19.1	22.1	27.7	35.6	44.1	57.1

All origins 3/, female

2003.....	7.9	7.5	6.7	4.9	4.0	3.6	4.0	3.9	3.3
2002.....	8.0	6.7	6.2	5.0	4.3	3.9	4.1	4.4	3.8
2001 5/.....	7.4	7.0	6.2	5.2	4.1	3.7	4.3	3.8	3.4
2000.....	6.7	6.8	5.8	4.9	4.0	4.1	3.8	4.2	4.2
1999.....	6.7	6.5	5.8	4.4	4.2	4.1	4.8	4.6	4.1

Hispanic, both sexes

2003.....	7.4	6.9	7.2	6.5	6.4	6.7	7.3	9.0	10.9
2002.....	7.2	7.0	7.9	6.2	6.8	8.4	8.0	11.1	12.7
2001 5/.....	6.8	7.4	6.2	6.2	6.7	8.7	6.8	12.4	13.8
2000.....	7.9	7.6	6.5	5.3	7.2	8.6	9.8	15.0	*
1999.....	7.2	6.9	7.1	6.5	7.9	5.9	11.3	16.4	15.4

Hispanic, male

2003.....	11.8	11.1	13.4	12.6	12.3	14.1	15.8	21.8	*
2002.....	11.0	11.7	14.5	11.3	13.2	16.7	16.0	25.2	35.1
2001 5/.....	11.0	11.9	11.2	11.9	13.8	17.4	15.7	28.6	41.9
2000.....	12.6	12.3	11.8	10.1	14.5	17.0	19.9	38.3	*
1999.....	12.4	10.4	12.7	11.6	15.8	11.2	23.4	37.7	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Hispanic, female

2003.....	2.9	2.9	*	*	*	*	*	*	*
2002.....	3.3	2.5	*	*	*	*	*	*	*
2001 5/.....	2.5	3.0	*	*	*	*	*	*	*
2000.....	3.3	3.0	*	*	*	*	*	*	*
1999.....	*	3.6	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	16.9	16.5	15.2	13.1	12.3	14.0	16.0	18.1	17.1
2002.....	17.2	15.7	15.1	12.7	12.8	14.9	17.0	19.7	18.1
2001 5/.....	16.5	15.1	14.5	12.3	13.1	14.2	16.9	19.1	17.5
2000.....	15.4	14.4	13.1	11.8	11.5	14.0	16.8	19.4	19.9
1999.....	14.9	13.8	13.2	11.7	12.6	14.9	17.6	19.2	19.3

Non-Hispanic 4/, male

2003.....	25.7	25.6	23.8	21.7	21.7	26.8	32.6	41.0	48.6
2002.....	26.3	24.8	24.1	20.9	22.3	28.5	35.2	44.8	50.9
2001 5/.....	25.6	23.2	23.1	19.9	23.1	27.2	34.6	44.6	51.3
2000.....	24.2	22.0	20.6	19.1	20.0	26.5	35.5	44.9	58.4
1999.....	23.1	21.2	20.8	19.5	22.3	28.5	36.0	44.2	57.6

Non-Hispanic 4/, female

2003.....	8.4	7.9	7.1	5.2	4.1	3.8	4.1	4.1	3.3
2002.....	8.5	7.1	6.5	5.2	4.5	4.0	4.2	4.4	3.9
2001 5/.....	7.8	7.4	6.5	5.5	4.3	3.8	4.4	3.8	3.5
2000.....	7.0	7.1	6.1	5.2	4.1	4.1	3.8	4.4	4.3
1999.....	7.1	6.7	6.1	4.6	4.4	4.2	4.8	4.6	4.0

Non-Hispanic White, both sexes

2003.....	19.3	18.8	16.9	14.4	13.5	15.2	17.3	19.4	18.4
2002.....	19.7	17.7	16.7	14.1	14.1	16.2	18.1	21.3	19.2
2001 5/.....	18.8	17.0	16.0	13.6	14.2	15.2	18.2	20.3	18.7
2000.....	17.3	15.9	14.4	13.1	12.6	15.0	18.0	20.6	21.3
1999.....	16.6	15.3	14.5	12.7	13.6	16.1	18.8	20.4	20.8

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All origins 3/, both sexes

2003.....	7.8	8.3	8.4	8.2	8.5	10.1	11.9	13.1	11.9
2002.....	7.8	7.9	8.6	8.1	8.8	10.7	12.4	14.3	12.0
2001.....	7.5	8.0	8.4	8.0	9.0	10.3	12.4	14.1	12.2
2000.....	7.4	7.8	7.4	7.7	8.3	10.3	12.4	14.0	13.2
1999.....	7.4	7.2	8.1	7.5	8.7	10.9	12.9	13.5	12.6

All origins 3/, male

2003.....	13.2	14.5	14.6	15.3	16.5	20.7	26.3	32.4	36.6
2002.....	13.3	14.0	15.4	14.8	17.0	22.5	27.8	35.9	37.8
2001.....	12.8	13.8	14.9	14.2	17.3	21.4	27.7	35.5	38.9
2000.....	12.8	13.3	13.1	13.9	15.9	21.3	28.1	35.4	42.9
1999.....	12.6	12.4	14.0	14.0	16.9	22.4	29.1	34.6	41.6

All origins 3/, female

2003.....	2.6	2.3	2.5	1.7	1.6	1.6	1.6	1.3	0.9
2002.....	2.4	2.1	2.3	2.0	1.8	1.3	1.5	1.2	0.8
2001.....	2.3	2.4	2.4	2.3	1.9	1.4	1.7	1.3	1.0
2000.....	2.2	2.5	2.1	2.1	1.8	1.6	1.4	1.4	1.1
1999.....	2.4	2.1	2.6	1.6	1.7	1.8	1.6	1.2	0.9

Hispanic, both sexes

2003.....	3.1	2.8	2.6	2.9	*	3.7	*	*	*
2002.....	2.8	2.5	3.5	3.2	4.0	5.0	*	*	*
2001.....	2.8	3.3	3.0	3.4	3.8	4.2	*	*	*
2000.....	4.1	3.8	2.9	*	4.0	5.2	*	*	*
1999.....	3.8	2.5	3.2	*	4.1	*	7.7	*	*

Hispanic, male

2003.....	5.3	5.4	5.1	6.2	*	8.5	*	*	*
2002.....	4.9	4.8	6.8	6.4	8.1	11.0	*	*	*
2001.....	5.0	5.8	6.4	6.8	8.2	9.6	*	*	*
2000.....	6.9	6.8	5.9	*	8.6	11.2	*	*	*
1999.....	6.6	4.2	6.0	*	8.5	*	18.7	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	8.3	8.8	8.9	8.6	9.0	10.5	12.3	13.5	12.1
2002.....	8.2	8.4	9.0	8.4	9.1	11.1	12.8	14.7	12.1
2001.....	7.9	8.4	8.8	8.3	9.3	10.6	12.8	14.4	12.4
2000.....	7.7	8.1	7.8	8.0	8.6	10.5	12.7	14.3	13.4
1999.....	7.8	7.5	8.4	7.8	8.9	11.3	13.1	13.7	12.7

Non-Hispanic 4/, male

2003.....	14.1	15.3	15.4	15.9	17.2	21.5	27.2	33.5	37.5
2002.....	14.1	14.8	16.0	15.3	17.4	23.1	28.7	36.8	38.4
2001.....	13.6	14.5	15.4	14.6	17.8	22.0	28.6	36.1	39.7
2000.....	13.3	13.8	13.6	14.4	16.3	21.7	28.9	36.1	43.8
1999.....	13.2	13.1	14.5	14.5	17.3	23.2	29.4	35.1	42.2

Non-Hispanic 4/, female

2003.....	2.8	2.5	2.7	1.9	1.8	1.7	1.7	1.4	0.9
2002.....	2.5	2.3	2.5	2.2	1.9	1.4	1.5	1.3	0.8
2001.....	2.5	2.6	2.6	2.5	2.0	1.5	1.8	1.4	1.0
2000.....	2.3	2.6	2.3	2.2	1.9	1.6	1.4	1.5	1.1
1999.....	2.5	2.2	2.8	1.7	1.7	1.9	1.7	1.2	0.9

Non-Hispanic White, both sexes

2003.....	9.8	10.0	10.1	9.6	10.0	11.6	13.5	14.6	13.2
2002.....	9.5	9.5	10.2	9.5	10.3	12.2	13.8	16.0	13.1
2001.....	9.1	9.5	9.9	9.3	10.4	11.5	13.9	15.3	13.4
2000.....	8.7	9.1	8.6	9.0	9.5	11.4	13.8	15.5	14.5
1999.....	8.7	8.5	9.4	8.6	9.8	12.3	14.1	14.6	13.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All origins 3/, both sexes									
2003.....	8.2	7.5	6.2	4.5	3.4	3.5	3.7	4.6	5.1
2002.....	8.5	7.2	5.9	4.2	3.7	3.9	4.2	5.0	6.0
2001 5/.....	8.2	6.6	5.6	4.0	3.7	3.6	4.0	4.8	5.3
2000.....	7.4	6.2	5.2	3.7	3.0	3.5	4.2	5.3	6.5
1999.....	6.9	6.2	4.7	3.9	3.8	3.6	4.5	5.7	6.7
All origins 3/, male									
2003.....	11.2	9.9	8.5	5.8	4.7	5.4	5.5	7.7	11.1
2002.....	11.6	9.9	8.2	5.7	5.0	5.5	6.5	8.1	12.9
2001 5/.....	11.5	8.6	7.5	5.3	5.4	5.4	6.1	8.7	12.2
2000.....	10.6	8.1	6.9	4.8	3.9	4.8	6.8	9.4	14.6
1999.....	9.6	8.1	6.4	5.1	5.2	5.3	6.6	9.5	15.5
All origins 3/, female									
2003.....	5.3	5.1	4.1	3.2	2.4	2.0	2.4	2.6	2.4
2002.....	5.6	4.6	3.9	2.9	2.5	2.6	2.6	3.1	3.0
2001 5/.....	5.1	4.6	3.8	2.8	2.2	2.3	2.6	2.4	2.4
2000.....	4.4	4.3	3.7	2.8	2.1	2.5	2.4	2.8	3.2
1999.....	4.2	4.4	3.2	2.8	2.5	2.3	3.1	3.4	3.1
Hispanic, both sexes									
2003.....	4.3	4.1	4.6	3.7	3.8	*	*	*	*
2002.....	4.3	4.6	4.4	3.0	*	*	*	*	*
2001 5/.....	4.0	4.1	3.1	2.8	*	4.6	*	*	*
2000.....	3.9	3.7	3.5	2.8	*	*	*	*	*
1999.....	3.4	4.4	3.9	3.9	3.8	*	*	*	*
Hispanic, male									
2003.....	6.6	5.7	8.2	6.4	6.5	*	*	*	*
2002.....	6.1	6.9	7.7	*	*	*	*	*	*
2001 5/.....	6.0	6.1	4.8	*	*	*	*	*	*
2000.....	5.8	5.6	5.9	*	*	*	*	*	*
1999.....	5.9	6.2	6.7	6.0	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years and over

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hispanic, female									
2003.....	2.1	2.6	*	*	*	*	*	*	*
2002.....	2.5	*	*	*	*	*	*	*	*
2001 5/.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes									
2003.....	8.6	7.8	6.3	4.5	3.4	3.5	3.7	4.5	5.1
2002.....	9.0	7.4	6.0	4.3	3.7	3.9	4.2	5.0	5.9
2001 5/.....	8.6	6.7	5.7	4.1	3.7	3.6	4.1	4.7	5.1
2000.....	7.7	6.3	5.3	3.8	2.9	3.5	4.1	5.1	6.5
1999.....	7.2	6.3	4.8	3.9	3.7	3.6	4.6	5.5	6.6
Non-Hispanic 4/, male									
2003.....	11.6	10.3	8.4	5.8	4.5	5.3	5.4	7.5	11.2
2002.....	12.1	10.0	8.1	5.6	4.9	5.4	6.4	8.0	12.5
2001 5/.....	12.0	8.8	7.6	5.3	5.4	5.2	6.0	8.6	11.6
2000.....	10.9	8.2	6.9	4.7	3.7	4.7	6.7	8.8	14.6
1999.....	9.9	8.2	6.3	5.0	5.0	5.3	6.7	9.1	15.4
Non-Hispanic 4/, female									
2003.....	5.6	5.4	4.4	3.4	2.4	2.1	2.5	2.7	2.4
2002.....	5.9	4.8	4.0	3.0	2.6	2.6	2.6	3.2	3.1
2001 5/.....	5.4	4.8	3.9	3.0	2.3	2.3	2.7	2.4	2.5
2000.....	4.6	4.5	3.8	2.9	2.2	2.5	2.4	2.9	3.2
1999.....	4.5	4.5	3.3	2.8	2.6	2.3	3.1	3.5	3.1
Non-Hispanic White, both sexes									
2003.....	9.6	8.7	6.9	4.8	3.5	3.6	3.8	4.8	5.2
2002.....	10.1	8.2	6.6	4.6	3.8	4.0	4.3	5.3	6.1
2001 5/.....	9.6	7.4	6.2	4.3	3.8	3.7	4.3	5.0	5.3
2000.....	8.6	6.8	5.9	4.2	3.1	3.6	4.2	5.1	6.8
1999.....	7.9	6.8	5.1	4.1	3.8	3.8	4.7	5.8	7.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All origins 3/, both sexes									
2003.....	5.6	4.2	3.1	2.4	2.3	2.4	2.5	2.4	2.2
2002.....	5.4	4.0	3.5	2.7	2.4	2.2	2.3	2.4	2.1
2001 5/.....	7.0	5.5	4.4	3.5	3.3	2.5	2.5	2.5	2.4
2000.....	5.3	4.0	3.4	2.6	2.4	2.4	2.3	2.6	2.4
1999.....	5.1	4.0	3.2	2.8	2.7	2.4	2.5	2.4	2.4
All origins 3/, male									
2003.....	8.3	6.4	4.6	3.7	3.4	3.4	3.5	3.0	3.2
2002.....	8.0	6.0	5.3	4.2	3.4	3.2	3.3	2.9	3.0
2001 5/.....	10.6	8.3	6.7	5.0	4.6	3.3	3.2	3.0	3.2
2000.....	7.8	5.9	5.2	3.9	3.3	3.4	2.8	3.8	3.3
1999.....	7.7	5.8	5.1	4.3	4.2	3.1	3.4	3.4	3.4
All origins 3/, female									
2003.....	3.0	2.1	1.7	1.3	1.4	1.6	1.8	2.1	1.8
2002.....	2.9	2.2	1.8	1.4	1.5	1.4	1.6	2.1	1.7
2001 5/.....	3.6	2.7	2.3	2.2	2.1	1.9	1.9	2.2	2.0
2000.....	2.8	2.1	1.7	1.5	1.7	1.6	2.0	1.9	2.0
1999.....	2.5	2.2	1.4	1.5	1.5	1.9	1.9	1.9	2.0
Hispanic, both sexes									
2003.....	7.3	5.3	3.0	4.4	3.5	4.1	*	*	*
2002.....	6.2	5.5	4.9	3.7	*	*	*	*	*
2001 5/.....	8.1	6.2	4.4	4.5	4.9	*	*	*	*
2000.....	7.6	5.6	5.2	3.6	*	4.4	*	*	*
1999.....	7.9	5.0	5.2	3.0	3.5	*	*	*	*
Hispanic, male									
2003.....	11.2	8.7	4.3	8.4	6.5	*	*	*	*
2002.....	10.0	8.0	8.6	6.4	*	*	*	*	*
2001 5/.....	12.6	9.6	6.6	7.1	8.5	*	*	*	*
2000.....	11.1	9.0	8.8	*	*	*	*	*	*
1999.....	11.9	8.0	8.6	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Hispanic, female

2003.....	3.3	*	*	*	*	*	*	*	*
2002.....	2.4	3.1	*	*	*	*	*	*	*
2001 5/.....	3.7	3.0	*	*	*	*	*	*	*
2000.....	4.2	*	*	*	*	*	*	*	*
1999.....	3.9	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	5.4	4.1	3.1	2.3	2.2	2.2	2.4	2.3	2.1
2002.....	5.3	3.9	3.3	2.6	2.4	2.2	2.3	2.2	2.1
2001 5/.....	6.8	5.3	4.3	3.3	3.1	2.5	2.5	2.4	2.4
2000.....	5.0	3.8	3.2	2.5	2.4	2.3	2.3	2.6	2.3
1999.....	4.8	3.8	3.0	2.8	2.6	2.4	2.5	2.4	2.4

Non-Hispanic 4/, male

2003.....	7.9	6.1	4.5	3.4	3.2	3.0	3.4	2.9	3.0
2002.....	7.7	5.8	5.0	4.0	3.3	3.1	3.3	2.8	2.9
2001 5/.....	10.2	8.0	6.6	4.7	4.3	3.2	3.2	2.9	3.1
2000.....	7.4	5.6	4.9	3.7	3.2	3.2	2.7	3.8	3.1
1999.....	7.3	5.5	4.8	4.1	3.9	3.0	3.3	3.3	3.4

Non-Hispanic 4/, female

2003.....	2.9	2.1	1.7	1.3	1.4	1.6	1.7	2.0	1.8
2002.....	2.9	2.1	1.8	1.4	1.5	1.5	1.5	1.9	1.7
2001 5/.....	3.5	2.7	2.2	2.1	2.1	1.9	2.0	2.2	2.1
2000.....	2.6	2.0	1.6	1.4	1.6	1.5	2.0	1.9	2.0
1999.....	2.4	2.2	1.4	1.6	1.5	1.9	2.0	1.9	2.0

Non-Hispanic White, both sexes

2003.....	3.4	2.8	2.3	1.7	1.7	1.7	2.0	2.0	2.0
2002.....	3.4	2.6	2.6	2.0	1.9	1.9	1.8	1.9	1.8
2001 5/.....	5.1	4.1	3.7	2.8	2.5	2.1	2.1	2.1	2.2
2000.....	3.2	2.6	2.4	1.8	1.7	1.8	2.0	2.0	2.1
1999.....	3.1	2.7	2.3	2.1	2.2	2.0	2.0	2.1	2.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hispanic White, male									
2003.....	4.6	4.0	3.1	2.5	2.3	2.2	2.8	2.3	2.9
2002.....	4.3	3.5	3.6	2.9	2.5	2.4	2.4	2.1	2.7
2001 5/.....	7.4	6.0	5.3	3.8	3.2	2.5	2.6	2.4	2.8
2000.....	4.3	3.5	3.5	2.6	2.1	2.4	2.3	2.6	2.6
1999.....	4.5	3.9	3.4	3.0	3.1	2.3	2.5	2.8	2.6
Non-Hispanic White, female									
2003.....	2.3	1.7	1.6	1.0	1.2	1.3	1.5	1.8	1.6
2002.....	2.5	1.6	1.7	1.3	1.3	1.5	1.3	1.8	1.5
2001 5/.....	2.9	2.3	2.0	1.8	1.9	1.7	1.7	1.9	2.0
2000.....	2.0	1.7	1.4	1.2	1.4	1.3	1.8	1.6	1.9
1999.....	1.7	1.6	1.2	1.4	1.5	1.7	1.7	1.6	1.9
Non-Hispanic Black, both sexes									
2003.....	18.1	12.9	9.1	6.9	5.8	6.9	6.4	6.7	*
2002.....	17.9	13.2	9.0	7.7	6.0	5.5	8.2	5.9	*
2001 5/.....	18.2	13.9	10.0	7.5	7.6	6.6	7.1	6.3	*
2000.....	17.1	12.4	9.2	7.5	8.1	7.0	5.4	9.8	*
1999.....	16.4	11.2	9.0	7.6	6.7	6.1	8.4	6.2	*
Non-Hispanic Black, male									
2003.....	30.6	22.2	17.4	11.5	11.1	11.5	11.7	*	*
2002.....	31.4	22.8	17.2	13.8	10.2	10.9	15.0	*	*
2001 5/.....	30.1	23.6	17.6	11.7	13.0	12.1	11.9	*	*
2000.....	29.2	22.5	16.3	13.4	14.9	12.6	*	19.7	*
1999.....	27.7	18.5	17.2	13.1	13.0	10.4	14.0	*	*
Non-Hispanic Black, female									
2003.....	7.1	5.1	*	3.2	*	*	*	*	*
2002.....	6.3	5.0	*	*	*	*	*	*	*
2001 5/.....	7.8	5.7	3.8	4.1	*	*	*	*	*
2000.....	6.5	3.8	3.5	*	*	*	*	*	*
1999.....	6.6	5.1	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All origins 3/, both sexes

2003.....	3.1	2.2	1.6	1.2	0.9	0.9	0.7	0.6	0.5
2002.....	2.8	2.2	1.8	1.3	1.0	0.8	1.0	0.8	*
2001.....	2.7	2.1	1.5	1.4	1.2	0.9	0.8	0.7	*
2000.....	2.9	2.1	1.7	1.2	1.0	1.0	0.7	0.5	0.7
1999.....	2.8	2.0	1.6	1.6	1.1	1.0	0.8	0.7	0.6

All origins 3/, male

2003.....	5.0	3.5	2.4	1.8	1.4	1.4	0.9	*	*
2002.....	4.2	3.4	2.7	1.9	1.4	1.0	1.4	1.0	*
2001.....	4.2	3.2	2.3	2.0	1.8	1.1	1.2	*	*
2000.....	4.5	3.3	2.6	1.8	1.3	1.3	0.8	*	*
1999.....	4.4	3.0	2.6	2.5	1.5	1.4	1.1	*	*

All origins 3/, female

2003.....	1.4	1.0	0.8	0.6	0.4	0.5	0.6	0.7	*
2002.....	1.5	1.1	0.9	0.7	0.6	0.6	0.6	0.7	*
2001.....	1.1	1.0	0.7	0.8	0.7	0.7	0.5	0.7	*
2000.....	1.3	1.0	0.9	0.7	0.8	0.7	0.7	*	*
1999.....	1.3	1.1	0.6	0.7	0.7	0.7	0.6	0.6	*

Hispanic, both sexes

2003.....	4.3	2.7	*	*	*	*	*	*	*
2002.....	3.1	3.2	1.8	*	*	*	*	*	*
2001.....	3.4	2.4	*	*	*	*	*	*	*
2000.....	4.5	3.4	2.3	*	*	*	*	*	*
1999.....	4.8	2.5	2.5	*	*	*	*	*	*

Hispanic, male

2003.....	7.3	3.9	*	*	*	*	*	*	*
2002.....	5.1	4.9	*	*	*	*	*	*	*
2001.....	5.5	3.9	*	*	*	*	*	*	*
2000.....	7.0	5.6	*	*	*	*	*	*	*
1999.....	6.9	4.2	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	2.6	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	3.0	2.2	1.6	1.1	0.8	0.9	0.7	0.6	0.5
2002.....	2.8	2.1	1.7	1.3	1.0	0.8	1.0	0.8	*
2001.....	2.6	2.0	1.5	1.4	1.1	0.9	0.9	0.7	*
2000.....	2.7	2.0	1.7	1.2	1.0	0.9	0.7	0.6	0.7
1999.....	2.6	2.0	1.5	1.5	1.1	1.1	0.8	0.7	0.5

Non-Hispanic 4/, male

2003.....	4.7	3.5	2.5	1.7	1.3	1.3	0.8	*	*
2002.....	4.1	3.3	2.7	1.9	1.4	1.0	1.4	*	*
2001.....	4.1	3.2	2.3	2.0	1.6	1.1	1.3	*	*
2000.....	4.2	3.1	2.5	1.8	1.4	1.2	0.8	*	*
1999.....	4.1	2.9	2.5	2.5	1.5	1.5	1.0	*	*

Non-Hispanic 4/, female

2003.....	1.4	1.0	0.8	0.6	*	0.5	0.7	0.8	*
2002.....	1.5	1.0	0.9	0.7	0.6	0.6	0.7	0.7	*
2001.....	1.1	0.9	0.8	0.8	0.7	0.7	0.6	*	*
2000.....	1.3	1.0	0.9	0.6	0.7	0.7	0.7	*	*
1999.....	1.1	1.1	0.5	0.7	0.6	0.7	0.7	*	*

Non-Hispanic White, both sexes

2003.....	1.8	1.5	1.3	0.9	0.6	0.7	0.6	0.6	0.5
2002.....	1.8	1.4	1.4	0.9	0.7	0.7	0.9	0.7	*
2001.....	1.7	1.4	1.2	1.2	1.0	0.7	0.7	0.6	*
2000.....	1.7	1.4	1.3	0.9	0.8	0.7	0.7	*	0.6
1999.....	1.6	1.4	1.1	1.2	0.9	0.9	0.6	0.7	0.6

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Non-Hispanic White, male

2003.....	2.5	2.3	1.8	1.3	1.0	1.0	*	*	*
2002.....	2.2	2.1	2.0	1.3	0.9	0.8	1.2	*	*
2001.....	2.5	2.1	1.5	1.7	1.2	0.8	1.0	*	*
2000.....	2.5	1.9	1.9	1.2	1.0	0.9	0.8	*	*
1999.....	2.4	2.1	1.7	1.7	1.3	1.2	*	*	*

Non-Hispanic White, female

2003.....	1.2	0.8	0.8	0.5	*	*	0.6	0.8	*
2002.....	1.3	0.8	0.9	0.5	0.5	0.7	0.6	0.7	*
2001.....	0.9	0.8	0.8	0.8	0.8	0.6	0.5	*	*
2000.....	1.0	0.9	0.8	0.7	0.7	0.7	0.7	*	*
1999.....	0.8	0.7	0.5	0.7	0.7	0.7	0.6	*	*

Non-Hispanic Black, both sexes

2003.....	10.5	7.0	4.6	3.1	2.5	*	*	*	*
2002.....	9.7	7.2	4.2	4.1	2.8	*	*	*	*
2001.....	8.5	6.6	4.4	2.9	2.8	*	*	*	*
2000.....	9.7	6.9	4.3	2.7	2.8	2.7	*	*	*
1999.....	9.7	6.1	4.6	4.3	2.3	*	*	*	*

Non-Hispanic Black, male

2003.....	19.7	12.3	9.1	5.2	*	*	*	*	*
2002.....	17.6	12.3	8.4	6.7	5.4	*	*	*	*
2001.....	15.6	12.1	8.7	5.0	6.0	*	*	*	*
2000.....	17.1	13.1	7.3	5.5	5.3	*	*	*	*
1999.....	17.4	9.8	9.6	8.5	*	*	*	*	*

Non-Hispanic Black, female

2003.....	2.6	2.6	*	*	*	*	*	*	*
2002.....	2.8	2.8	*	*	*	*	*	*	*
2001.....	2.4	2.0	*	*	*	*	*	*	*
2000.....	3.3	*	*	*	*	*	*	*	*
1999.....	3.1	2.9	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1)

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All origins 3/, both sexes									
2003.....	2.5	1.9	1.5	1.3	1.5	1.5	1.8	1.8	1.7
2002.....	2.6	1.8	1.7	1.5	1.4	1.4	1.3	1.6	1.7
2001 5/.....	4.4	3.4	2.9	2.1	2.1	1.6	1.6	1.8	2.0
2000.....	2.4	1.8	1.6	1.4	1.4	1.4	1.6	2.1	1.7
1999.....	2.3	1.9	1.6	1.3	1.6	1.4	1.7	1.7	1.8
All origins 3/, male									
2003.....	3.3	2.9	2.2	1.9	2.1	2.0	2.7	2.6	2.5
2002.....	3.8	2.6	2.6	2.3	2.0	2.2	1.8	1.9	2.3
2001 5/.....	6.4	5.0	4.4	3.0	2.8	2.2	2.0	2.1	2.7
2000.....	3.3	2.6	2.6	2.0	2.0	2.1	2.0	3.0	2.5
1999.....	3.3	2.8	2.5	1.8	2.6	1.7	2.2	2.6	2.6
All origins 3/, female									
2003.....	1.6	1.0	0.9	0.7	1.0	1.0	1.1	1.3	1.4
2002.....	1.4	1.1	1.0	0.7	0.9	0.8	0.9	1.4	1.4
2001 5/.....	2.5	1.8	1.5	1.4	1.4	1.2	1.4	1.6	1.7
2000.....	1.5	1.0	0.8	0.8	0.9	0.9	1.3	1.5	1.4
1999.....	1.3	1.1	0.9	0.8	0.8	1.2	1.3	1.2	1.5
Hispanic, both sexes									
2003.....	2.9	2.7	2.2	2.5	*	*	*	*	*
2002.....	3.1	2.3	3.1	2.5	*	*	*	*	*
2001 5/.....	4.7	3.8	3.0	3.8	3.2	*	*	*	*
2000.....	3.1	2.2	2.9	*	*	*	*	*	*
1999.....	3.2	2.5	2.6	*	*	*	*	*	*
Hispanic, male									
2003.....	3.9	4.8	*	5.2	*	*	*	*	*
2002.....	4.9	3.1	5.3	*	*	*	*	*	*
2001 5/.....	7.1	5.7	4.4	6.0	*	*	*	*	*
2000.....	4.1	3.5	4.6	*	*	*	*	*	*
1999.....	5.0	3.8	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1)

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hispanic, female									
2003.....	2.0	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001 5/.....	2.3	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes									
2003.....	2.4	1.9	1.4	1.2	1.4	1.4	1.7	1.7	1.6
2002.....	2.5	1.7	1.6	1.4	1.4	1.4	1.3	1.5	1.7
2001 5/.....	4.2	3.3	2.8	2.0	1.9	1.6	1.6	1.7	2.1
2000.....	2.3	1.8	1.5	1.3	1.4	1.3	1.5	2.1	1.6
1999.....	2.2	1.8	1.5	1.2	1.6	1.3	1.7	1.7	1.8
Non-Hispanic 4/, male									
2003.....	3.2	2.7	2.1	1.7	1.9	1.7	2.6	2.5	2.3
2002.....	3.6	2.5	2.3	2.1	1.9	2.1	1.8	1.8	2.3
2001 5/.....	6.1	4.9	4.3	2.6	2.6	2.1	1.9	2.0	2.7
2000.....	3.2	2.5	2.4	1.9	1.9	2.0	1.9	3.1	2.4
1999.....	3.1	2.6	2.3	1.6	2.4	1.6	2.3	2.6	2.6
Non-Hispanic 4/, female									
2003.....	1.6	1.1	0.8	0.7	1.0	1.1	1.0	1.2	1.3
2002.....	1.4	1.0	1.0	0.7	0.9	0.8	0.9	1.2	1.4
2001 5/.....	2.4	1.7	1.4	1.3	1.4	1.2	1.4	1.6	1.8
2000.....	1.4	1.0	0.7	0.8	0.9	0.8	1.2	1.5	1.3
1999.....	1.2	1.1	0.8	0.8	0.9	1.1	1.3	1.2	1.5
Non-Hispanic White, both sexes									
2003.....	1.6	1.3	1.1	0.8	1.1	1.0	1.4	1.4	1.5
2002.....	1.6	1.1	1.2	1.1	1.1	1.1	0.9	1.2	1.4
2001 5/.....	3.4	2.7	2.5	1.6	1.5	1.3	1.3	1.5	1.9
2000.....	1.5	1.2	1.1	0.9	0.9	1.0	1.3	1.6	1.5
1999.....	1.5	1.4	1.2	1.0	1.3	1.1	1.4	1.4	1.5

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1)

Non-Hispanic White, male									
2003.....	2.0	1.7	1.4	1.2	1.3	1.2	2.1	2.0	2.2
2002.....	2.1	1.4	1.5	1.6	1.6	1.6	1.2	1.4	2.0
2001 5/.....	4.9	3.9	3.8	2.2	2.1	1.7	1.6	1.8	2.3
2000.....	1.9	1.6	1.6	1.3	1.1	1.5	1.5	2.2	1.9
1999.....	2.1	1.8	1.7	1.3	1.8	1.2	1.8	2.1	*
Non-Hispanic White, female									
2003.....	1.1	0.9	0.8	0.5	0.9	0.9	0.9	1.0	1.2
2002.....	1.1	0.8	0.9	0.7	0.7	0.8	0.7	1.1	1.2
2001 5/.....	2.0	1.5	1.2	1.0	1.1	1.1	1.2	1.2	1.7
2000.....	1.1	0.9	0.5	0.5	0.7	0.7	1.1	1.2	1.3
1999.....	0.9	0.9	0.7	0.7	0.8	1.0	1.1	1.0	1.4
Non-Hispanic Black, both sexes									
2003.....	7.5	5.9	4.5	3.7	3.3	4.3	4.7	5.2	*
2002.....	8.2	6.0	4.8	3.6	3.2	3.7	5.8	*	*
2001 5/.....	9.6	7.3	5.7	4.5	4.9	4.1	4.8	*	*
2000.....	7.4	5.5	5.0	4.8	5.3	4.2	4.2	6.9	*
1999.....	6.7	5.2	4.4	3.3	4.4	3.5	5.3	*	*
Non-Hispanic Black, male									
2003.....	11.0	9.9	8.2	6.4	6.3	6.7	*	*	*
2002.....	13.7	10.5	8.7	7.1	*	7.2	10.3	*	*
2001 5/.....	14.6	11.5	8.9	6.7	7.0	7.2	*	*	*
2000.....	12.2	9.5	9.0	7.9	9.6	6.8	*	*	*
1999.....	10.3	8.8	7.7	4.6	8.5	*	*	*	*
Non-Hispanic Black, female									
2003.....	4.6	2.6	*	*	*	*	*	*	*
2002.....	3.5	2.2	*	*	*	*	*	*	*
2001 5/.....	5.4	3.7	3.0	*	*	*	*	*	*
2000.....	3.3	2.0	*	*	*	*	*	*	*
1999.....	3.5	2.1	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Legal intervention (Y35,Y89.0)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.1	0.1	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	*	0.2	*	*	*	*	*	*	*

Non-Hispanic 4/, male

2003.....	0.3	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*
1999.....	*	0.3	*	*	*	*	*	*	*

Non-Hispanic 4/, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic White, both sexes

2003.....	0.1	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*
1999.....	*	0.2	*	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	3.7	2.7	1.6	1.0	0.8	0.6	0.6	0.9	1.1
2002.....	3.5	2.4	1.4	1.0	0.8	0.6	0.6	1.0	1.4
2001.....	3.1	2.0	1.1	0.6	0.6	0.5	0.7	1.1	0.9
2000.....	2.8	1.8	1.1	0.7	0.5	0.6	0.6	0.7	1.4
1999.....	2.8	1.6	1.0	0.8	0.8	0.5	0.7	0.9	1.4

Non-Hispanic 4/, male

2003.....	4.8	3.5	2.0	1.2	0.8	0.7	0.9	1.0	*
2002.....	4.2	3.0	1.8	1.2	1.0	0.9	0.8	1.1	2.1
2001.....	4.5	2.5	1.5	0.8	0.9	0.6	0.9	1.5	*
2000.....	4.0	2.4	1.6	0.9	0.8	0.8	1.0	*	1.7
1999.....	3.9	2.1	1.4	1.1	1.1	0.6	1.0	1.4	2.1

Non-Hispanic 4/, female

2003.....	2.7	1.9	1.2	0.9	0.8	0.6	*	0.9	1.0
2002.....	2.8	1.9	1.1	0.7	0.7	*	*	0.9	1.1
2001.....	1.8	1.5	0.7	0.5	*	*	0.6	0.8	0.8
2000.....	1.8	1.2	0.7	0.5	*	*	*	*	1.2
1999.....	1.8	1.1	0.7	0.6	0.5	*	0.5	0.7	1.1

Non-Hispanic White, both sexes

2003.....	3.8	2.6	1.5	0.9	0.7	0.6	0.6	0.9	1.2
2002.....	3.5	2.3	1.4	0.9	0.8	0.6	0.6	0.9	1.5
2001.....	3.1	1.9	1.1	0.6	0.5	0.5	0.7	1.1	1.0
2000.....	2.8	1.7	1.0	0.7	0.5	0.5	0.6	0.7	1.3
1999.....	2.6	1.4	1.0	0.8	0.8	0.4	0.7	1.0	1.4

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	3.7	2.6	1.5	1.0	0.8	0.6	0.6	0.8	1.1
2002.....	3.4	2.4	1.4	0.9	0.8	0.6	0.6	0.9	1.4
2001.....	3.0	1.9	1.0	0.6	0.5	0.4	0.6	1.0	0.9
2000.....	2.8	1.7	1.0	0.6	0.5	0.5	0.5	0.7	1.3
1999.....	2.7	1.5	1.0	0.7	0.7	0.4	0.6	0.8	1.3

Non-Hispanic 4/, male

2003.....	4.7	3.4	1.8	1.1	0.8	0.6	0.7	*	*
2002.....	4.1	2.9	1.8	1.2	0.9	0.8	0.7	1.0	2.1
2001.....	4.4	2.4	1.4	0.7	0.8	*	0.7	1.3	*
2000.....	3.9	2.2	1.4	0.9	0.7	0.7	0.9	*	*
1999.....	3.7	2.1	1.4	0.9	0.9	*	0.8	*	1.7

Non-Hispanic 4/, female

2003.....	2.7	1.9	1.2	0.8	0.7	0.6	*	0.8	1.0
2002.....	2.7	1.9	1.0	0.7	0.6	*	*	0.9	1.1
2001.....	1.8	1.5	0.7	0.5	*	*	0.6	0.8	0.8
2000.....	1.7	1.2	0.7	0.4	*	*	*	*	1.2
1999.....	1.8	1.0	0.6	0.6	0.5	*	*	0.7	1.1

Non-Hispanic White, both sexes

2003.....	3.7	2.5	1.4	0.9	0.7	0.5	0.5	0.8	1.2
2002.....	3.4	2.2	1.4	0.9	0.7	0.5	0.5	0.9	1.5
2001.....	3.0	1.8	1.0	0.6	0.5	0.4	0.6	1.0	0.9
2000.....	2.7	1.6	0.9	0.6	0.4	0.5	0.5	0.7	1.2
1999.....	2.6	1.4	0.9	0.7	0.7	0.4	0.6	0.8	1.3

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Complications of medical and surgical care (Y40-Y84,Y88)

Hispanic, female

2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	0.7	0.7	1.2	1.7	2.2	3.7	5.5	7.4	11.4
2002.....	0.6	0.8	1.1	1.6	2.3	3.9	5.2	7.7	11.6
2001.....	0.7	0.8	1.3	1.9	2.6	3.7	5.9	8.3	12.9
2000.....	0.7	1.0	1.2	2.1	2.7	4.4	5.8	8.1	13.7
1999.....	0.5	0.7	1.1	1.7	2.5	4.2	5.6	8.6	12.8

Non-Hispanic 4/, male

2003.....	0.7	0.7	1.4	1.7	2.3	4.2	5.7	9.0	13.2
2002.....	0.6	0.7	1.2	1.6	2.5	3.9	5.4	7.8	12.9
2001.....	0.7	0.8	1.4	2.0	2.6	4.0	6.3	9.9	14.5
2000.....	0.6	1.0	1.2	2.1	2.8	5.3	6.9	10.4	16.1
1999.....	0.4	0.7	1.2	1.7	2.4	4.8	6.5	10.3	16.8

Non-Hispanic 4/, female

2003.....	0.6	0.7	1.0	1.7	2.1	3.3	5.3	6.4	10.6
2002.....	0.5	0.9	1.0	1.7	2.2	4.0	5.0	7.7	11.1
2001.....	0.6	0.8	1.2	1.8	2.5	3.5	5.6	7.4	12.2
2000.....	0.7	0.9	1.2	2.1	2.6	3.7	5.0	6.8	12.7
1999.....	0.6	0.8	0.9	1.6	2.5	3.7	5.0	7.6	11.2

Non-Hispanic White, both sexes

2003.....	0.6	0.6	1.0	1.6	2.1	3.5	5.2	7.5	11.2
2002.....	0.5	0.7	1.0	1.5	2.1	3.9	5.0	7.6	11.5
2001.....	0.6	0.7	1.2	1.7	2.4	3.3	5.8	8.2	12.2
2000.....	0.6	0.9	1.1	1.8	2.5	4.1	5.5	8.0	13.2
1999.....	0.4	0.6	0.9	1.6	2.4	3.9	5.3	8.2	12.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic White, male

2003.....	0.6	0.6	1.2	1.6	2.3	4.1	5.6	9.3	13.1
2002.....	0.5	0.6	1.1	1.5	2.2	3.8	5.3	7.9	12.4
2001.....	0.6	0.7	1.2	1.7	2.5	3.3	6.2	10.1	13.5
2000.....	0.6	1.0	1.2	1.8	2.5	5.0	6.8	10.6	15.9
1999.....	0.3	0.6	1.0	1.6	2.3	4.5	6.3	9.9	15.8

Non-Hispanic White, female

2003.....	0.6	0.6	0.8	1.5	2.0	3.0	4.9	6.3	10.4
2002.....	0.4	0.7	0.9	1.5	2.1	4.0	4.9	7.4	11.1
2001.....	0.5	0.7	1.2	1.6	2.3	3.2	5.5	7.1	11.6
2000.....	0.6	0.9	1.0	1.8	2.5	3.5	4.5	6.5	12.0
1999.....	0.5	0.6	0.8	1.5	2.5	3.5	4.7	7.2	10.7

Non-Hispanic Black, both sexes

2003.....	1.3	1.6	2.5	3.0	3.7	5.7	9.7	7.8	15.2
2002.....	1.3	2.2	2.6	2.9	4.1	5.4	7.7	9.6	14.2
2001.....	1.6	1.6	2.8	4.2	4.6	8.5	8.1	11.0	23.7
2000.....	1.2	1.7	2.7	4.8	4.8	8.0	10.5	10.6	21.3
1999.....	0.9	1.8	2.7	2.6	3.5	7.3	9.4	12.3	20.4

Non-Hispanic Black, male

2003.....	*	2.2	2.9	*	*	*	*	*	*
2002.....	*	*	*	*	5.1	*	*	*	*
2001.....	1.8	*	3.5	5.2	5.2	11.4	*	*	30.0
2000.....	*	*	*	4.7	6.4	9.6	10.1	*	*
1999.....	*	*	*	*	*	9.3	10.2	*	23.8

Non-Hispanic Black, female

2003.....	*	*	*	3.1	3.9	6.4	10.8	7.9	15.3
2002.....	*	2.5	2.5	3.2	*	5.0	8.4	10.1	11.7
2001.....	*	*	*	3.5	4.2	6.7	8.2	11.2	21.4
2000.....	*	*	3.2	4.9	*	7.0	10.7	10.8	22.2
1999.....	*	*	*	*	*	6.0	9.0	11.0	19.1

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/

All origins 3/, both sexes									
2003.....	16.0	18.9	18.3	18.8	16.8	13.2	10.3	7.6	4.1
2002 7/.....	15.6	18.2	18.2	18.6	16.5	14.4	10.9	7.2	4.4
2001 7/.....	16.5	17.9	17.6	18.9	17.3	13.7	11.0	7.5	5.0
2000 7/.....	15.9	16.7	18.2	19.3	17.2	14.3	11.2	7.9	5.4
1999 7/.....	16.0	16.9	18.1	19.3	17.6	14.1	11.9	8.5	5.5
All origins 3/, male									
2003.....	24.5	29.9	29.1	30.6	27.6	22.8	18.0	15.0	7.8
2002 7/.....	23.7	29.2	29.2	31.0	27.9	23.8	18.7	13.4	9.7
2001 7/.....	25.4	28.0	28.7	31.0	28.5	23.8	19.4	13.8	11.6
2000 7/.....	25.0	26.9	28.8	31.4	28.9	25.0	21.1	14.9	11.9
1999 7/.....	25.2	27.4	29.7	31.5	29.9	24.5	21.2	16.6	11.8
All origins 3/, female									
2003.....	7.7	8.3	8.1	8.1	7.5	5.4	4.9	3.1	2.5
2002 7/.....	7.7	7.7	7.9	7.2	6.7	6.9	5.3	3.4	2.1
2001 7/.....	7.9	8.3	7.2	8.0	7.6	5.7	5.1	3.8	2.3
2000 7/.....	7.1	7.0	8.3	8.3	7.2	5.9	4.3	3.8	2.7
1999 7/.....	7.1	6.8	7.3	8.3	7.0	6.0	5.5	3.8	3.0
Hispanic, both sexes									
2003.....	18.3	22.8	27.7	27.4	26.1	19.1	19.7	14.8	*
2002 7/.....	18.7	24.5	26.9	25.7	28.0	25.8	16.3	11.6	*
2001 7/.....	21.5	23.9	27.3	29.5	27.7	24.0	22.4	16.9	*
2000 7/.....	20.6	24.5	30.0	28.8	32.5	24.3	22.0	13.4	*
1999 7/.....	21.3	24.2	28.0	29.6	28.5	23.2	20.7	15.3	14.0
Hispanic, male									
2003.....	31.3	40.5	50.3	50.7	44.2	35.4	37.7	30.1	*
2002 7/.....	31.7	42.4	49.4	47.5	54.5	50.5	28.8	22.9	*
2001 7/.....	36.9	41.2	49.7	54.6	55.6	48.0	41.6	31.2	*
2000 7/.....	35.9	43.4	55.9	53.3	66.0	49.1	48.0	30.9	*
1999 7/.....	36.9	42.6	51.3	54.8	54.1	47.4	37.4	33.0	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/

Hispanic, female

2003.....	4.9	5.6	6.8	7.1	11.2	6.6	*	*	*
2002 7/.....	5.4	7.1	6.2	6.7	6.4	6.8	*	*	*
2001 7/.....	6.1	7.4	7.1	7.8	*	*	*	*	*
2000 7/.....	5.3	6.5	6.5	7.7	*	*	*	*	*
1999 7/.....	5.8	6.7	6.9	8.0	7.8	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	15.6	18.4	17.4	18.0	16.0	12.7	9.8	7.3	4.1
2002 7/.....	15.1	17.5	17.4	17.9	15.5	13.6	10.5	7.0	4.1
2001 7/.....	15.9	17.2	16.6	18.0	16.4	13.0	10.4	7.2	5.0
2000 7/.....	15.3	15.9	17.2	18.5	16.1	13.7	10.6	7.7	5.3
1999 7/.....	15.4	16.1	17.3	18.4	16.8	13.5	11.5	8.3	5.2

Non-Hispanic 4/, male

2003.....	23.6	28.7	27.2	28.7	26.2	21.8	16.9	14.2	7.8
2002 7/.....	22.5	27.8	27.3	29.5	25.8	22.0	18.1	12.9	9.1
2001 7/.....	24.0	26.5	26.7	29.0	26.4	22.2	18.1	13.1	11.4
2000 7/.....	23.7	25.2	26.6	29.6	26.2	23.6	19.6	14.3	11.7
1999 7/.....	23.9	25.9	28.0	29.5	28.2	23.0	20.3	16.0	11.1

Non-Hispanic 4/, female

2003.....	7.9	8.5	8.2	8.2	7.2	5.3	4.8	3.1	2.5
2002 7/.....	7.9	7.7	8.0	7.3	6.7	6.9	5.2	3.4	1.9
2001 7/.....	8.0	8.3	7.1	8.0	7.7	5.6	4.9	3.6	2.3
2000 7/.....	7.2	7.0	8.5	8.3	7.3	5.9	4.4	3.8	2.6
1999 7/.....	7.2	6.7	7.3	8.3	6.9	6.0	5.4	3.7	2.9

Non-Hispanic White, both sexes

2003.....	15.8	18.0	17.1	17.2	16.0	12.7	10.0	7.2	4.1
2002 7/.....	15.2	17.2	16.6	17.1	15.4	13.3	10.5	6.8	4.0
2001 7/.....	15.7	16.1	15.5	17.5	15.6	12.9	10.2	7.2	4.7
2000 7/.....	14.7	15.1	16.1	17.6	15.5	13.4	10.5	7.6	5.2
1999 7/.....	14.4	15.2	16.0	17.3	16.0	12.9	11.0	8.1	5.0

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/

Non-Hispanic White, male

2003.....	23.5	27.8	26.5	26.9	26.0	21.6	17.0	13.7	7.7
2002 7/.....	22.5	27.3	26.1	28.0	24.9	20.9	18.0	12.5	8.7
2001 7/.....	23.7	24.8	24.5	27.8	24.5	21.7	17.8	13.2	10.8
2000 7/.....	22.9	23.8	24.9	28.0	24.9	22.8	19.1	14.2	11.4
1999 7/.....	22.4	24.4	25.4	27.9	26.3	21.7	19.3	15.8	10.5

Non-Hispanic White, female

2003.....	8.2	8.5	8.0	8.1	7.1	5.3	4.9	3.2	2.5
2002 7/.....	7.9	7.4	7.6	6.9	6.9	7.1	5.2	3.3	2.1
2001 7/.....	7.8	7.5	6.9	7.9	7.7	5.8	4.9	3.7	2.2
2000 7/.....	6.6	6.5	7.7	8.0	7.1	5.9	4.4	3.7	2.7
1999 7/.....	6.5	6.2	6.9	7.5	6.9	5.9	5.2	3.5	2.8

Non-Hispanic Black, both sexes

2003.....	15.3	22.8	22.7	26.9	19.2	14.5	9.2	8.8	*
2002 7/.....	16.3	22.2	25.6	25.3	18.7	18.0	12.3	9.1	*
2001 7/.....	18.6	27.8	27.2	24.8	25.5	14.7	13.6	7.4	*
2000 7/.....	21.9	24.6	27.1	26.6	24.0	17.9	13.6	8.3	6.4
1999 7/.....	23.1	24.4	29.8	29.3	25.3	21.1	16.7	11.7	7.8

Non-Hispanic Black, male

2003.....	25.2	39.5	37.5	49.2	33.2	28.0	17.8	21.9	*
2002 7/.....	25.1	36.0	42.3	45.4	36.3	35.6	23.0	19.5	*
2001 7/.....	29.5	44.5	48.6	44.5	48.7	29.6	26.5	*	*
2000 7/.....	33.5	40.5	44.2	45.3	43.2	36.6	28.4	17.2	*
1999 7/.....	36.1	40.5	53.7	49.0	49.1	41.2	31.9	23.0	*

Non-Hispanic Black, female

2003.....	6.8	8.6	10.6	9.4	8.8	5.5	*	*	*
2002 7/.....	8.7	10.4	11.8	9.5	5.7	6.2	5.9	*	*
2001 7/.....	9.2	13.6	9.6	9.4	8.3	4.8	5.9	*	*
2000 7/.....	11.8	11.0	13.0	11.9	9.8	5.4	*	*	*
1999 7/.....	11.9	10.9	10.4	13.8	7.7	7.8	7.6	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0-F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

All origins 3/, both sexes

2003.....	22.7	17.0	9.8	5.7	4.5	3.7	3.9	4.6	6.0
2002 7/.....	20.9	14.5	8.2	5.1	3.8	3.3	3.6	4.1	5.7
2001 7/.....	17.6	11.6	6.3	4.1	3.1	2.9	3.4	3.6	5.1
2000 7/.....	15.8	10.3	5.9	3.6	2.6	2.7	3.2	3.9	5.7
1999 7/.....	15.5	9.3	5.8	3.8	3.2	2.9	3.6	4.2	4.8

All origins 3/, male

2003.....	29.2	21.7	11.5	6.6	5.2	3.9	4.7	5.3	7.8
2002 7/.....	26.4	19.1	10.0	5.7	4.1	3.4	3.9	4.7	7.3
2001 7/.....	23.9	15.0	7.7	4.3	3.7	3.3	3.8	3.9	5.7
2000 7/.....	21.9	13.8	6.7	3.9	2.9	2.8	3.6	4.3	7.5
1999 7/.....	21.9	12.2	7.1	4.7	3.7	3.1	3.5	4.4	6.2

All origins 3/, female

2003.....	16.5	12.4	8.2	4.9	4.0	3.4	3.3	4.2	5.2
2002 7/.....	15.6	10.2	6.6	4.6	3.6	3.3	3.3	3.8	4.9
2001 7/.....	11.6	8.3	5.1	3.9	2.6	2.6	3.2	3.4	4.9
2000 7/.....	9.8	7.0	5.2	3.4	2.3	2.7	2.9	3.8	4.9
1999 7/.....	9.2	6.6	4.6	2.9	2.7	2.7	3.7	4.0	4.3

Hispanic, both sexes

2003.....	16.4	13.0	7.6	4.9	3.8	*	*	*	*
2002 7/.....	15.9	11.1	6.5	4.4	*	*	*	*	*
2001 7/.....	12.0	9.6	6.3	2.8	*	*	*	*	*
2000 7/.....	12.1	8.6	6.4	*	*	*	*	*	*
1999 7/.....	15.1	11.2	5.0	5.2	*	*	*	*	*

Hispanic, male

2003.....	24.3	19.1	12.3	7.9	*	*	*	*	*
2002 7/.....	23.6	17.7	9.9	6.9	*	*	*	*	*
2001 7/.....	18.5	15.7	10.6	*	*	*	*	*	*
2000 7/.....	20.1	13.1	9.0	*	*	*	*	*	*
1999 7/.....	25.9	16.5	9.2	8.6	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

Hispanic, female

2003.....	8.3	7.1	3.2	*	*	*	*	*	*
2002 7/.....	8.1	4.6	3.4	*	*	*	*	*	*
2001 7/.....	5.5	3.8	*	*	*	*	*	*	*
2000 7/.....	4.2	4.3	4.0	*	*	*	*	*	*
1999 7/.....	4.4	6.2	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	23.3	17.2	9.9	5.7	4.5	3.7	3.9	4.6	6.0
2002 7/.....	21.3	14.8	8.3	5.1	3.8	3.3	3.5	4.2	5.8
2001 7/.....	18.0	11.6	6.2	4.2	3.1	2.9	3.5	3.6	5.2
2000 7/.....	15.9	10.3	5.8	3.7	2.6	2.7	3.2	4.0	5.7
1999 7/.....	15.3	9.0	5.7	3.6	3.1	2.9	3.7	4.2	4.9

Non-Hispanic 4/, male

2003.....	29.5	21.7	11.3	6.4	5.0	3.8	4.7	5.1	7.8
2002 7/.....	26.5	19.1	9.9	5.6	4.0	3.3	3.7	4.7	7.5
2001 7/.....	24.1	14.8	7.3	4.2	3.6	3.3	3.8	4.0	5.8
2000 7/.....	21.8	13.6	6.3	3.8	2.9	2.8	3.7	4.4	7.5
1999 7/.....	21.3	11.6	6.8	4.2	3.5	3.1	3.6	4.6	6.3

Non-Hispanic 4/, female

2003.....	17.2	12.9	8.6	5.1	4.1	3.6	3.3	4.2	5.3
2002 7/.....	16.3	10.6	6.9	4.7	3.6	3.3	3.4	3.8	5.1
2001 7/.....	12.1	8.6	5.2	4.1	2.6	2.6	3.3	3.4	5.0
2000 7/.....	10.2	7.1	5.2	3.6	2.4	2.7	2.9	3.8	5.0
1999 7/.....	9.6	6.6	4.7	3.0	2.8	2.7	3.8	4.0	4.4

Non-Hispanic White, both sexes

2003.....	23.6	16.7	9.7	5.6	4.4	3.9	4.2	4.8	6.3
2002 7/.....	21.2	14.0	8.0	4.9	3.9	3.4	3.6	4.4	5.9
2001 7/.....	17.3	10.8	6.0	4.0	3.1	3.0	3.6	3.8	5.2
2000 7/.....	15.0	9.3	5.5	3.7	2.6	2.9	3.5	4.2	5.7
1999 7/.....	14.2	8.1	5.4	3.3	3.1	2.9	3.9	4.4	4.9

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

Non-Hispanic White, male

2003.....	29.4	20.2	10.4	5.9	4.5	3.9	5.0	5.4	7.8
2002 7/.....	25.9	17.2	8.7	4.7	3.7	3.4	3.8	4.6	7.8
2001 7/.....	22.4	12.8	6.5	3.4	3.5	3.2	3.6	4.2	5.7
2000 7/.....	20.0	11.7	5.5	3.5	2.6	2.7	3.9	4.5	7.2
1999 7/.....	19.1	9.8	5.7	3.5	3.3	3.2	3.6	4.8	6.6

Non-Hispanic White, female

2003.....	17.9	13.2	9.0	5.4	4.4	3.8	3.7	4.5	5.7
2002 7/.....	16.6	10.9	7.3	5.1	4.0	3.5	3.5	4.2	5.1
2001 7/.....	12.2	8.8	5.6	4.5	2.7	2.8	3.6	3.5	4.9
2000 7/.....	10.1	7.0	5.5	3.9	2.5	3.0	3.1	4.0	5.1
1999 7/.....	9.4	6.6	5.0	3.2	3.0	2.8	4.1	4.2	4.2

Non-Hispanic Black, both sexes

2003.....	28.9	25.8	14.8	7.6	6.5	3.0	*	*	*
2002 7/.....	28.9	25.3	13.9	8.9	3.6	*	*	*	*
2001 7/.....	28.6	21.9	10.1	6.9	3.4	2.9	*	*	6.2
2000 7/.....	26.5	21.2	9.8	4.6	3.6	*	*	*	7.6
1999 7/.....	28.1	18.4	9.6	6.4	3.8	2.9	*	*	*

Non-Hispanic Black, male

2003.....	40.3	38.9	23.4	12.9	11.6	*	*	*	*
2002 7/.....	39.7	40.5	23.3	16.1	6.7	*	*	*	*
2001 7/.....	44.1	35.1	16.9	12.1	5.2	*	*	*	*
2000 7/.....	41.5	33.9	16.3	7.4	5.9	*	*	*	*
1999 7/.....	44.6	30.1	17.6	11.8	6.4	*	*	*	*

Non-Hispanic Black, female

2003.....	19.0	14.8	7.7	3.6	*	*	*	*	*
2002 7/.....	19.5	12.5	6.2	3.3	*	*	*	*	*
2001 7/.....	15.1	10.7	4.5	*	*	*	*	*	*
2000 7/.....	13.5	10.4	4.6	*	*	*	*	*	*
1999 7/.....	13.8	8.6	3.1	*	*	*	*	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All origins 3/, both sexes									
2003.....	11.4	10.9	10.4	9.6	9.7	11.3	12.9	14.2	12.5
2002.....	11.1	10.5	10.7	9.6	10.1	11.8	13.6	15.5	12.5
2001.....	10.6	10.4	10.4	9.7	10.5	11.4	13.7	15.3	12.8
2000.....	10.7	10.3	9.6	9.1	9.7	11.5	13.4	14.7	14.2
1999.....	10.7	9.6	9.9	9.4	10.0	12.2	14.0	14.5	13.5
All origins 3/, male									
2003.....	18.9	18.6	17.8	17.6	18.5	22.5	27.8	33.8	37.8
2002.....	18.4	18.0	18.6	17.1	18.8	24.1	29.8	37.6	38.9
2001.....	17.9	17.6	17.9	16.7	19.6	22.9	29.9	37.3	40.2
2000.....	17.9	17.3	16.5	16.1	17.8	23.1	29.6	36.6	44.7
1999.....	17.8	16.1	17.1	17.1	18.9	24.5	30.8	36.2	43.3
All origins 3/, female									
2003.....	4.1	3.5	3.4	2.4	2.1	2.3	2.3	2.2	1.3
2002.....	4.0	3.2	3.3	2.8	2.5	2.0	2.2	2.0	1.1
2001.....	3.6	3.5	3.3	3.2	2.6	2.2	2.3	2.2	1.3
2000.....	3.7	3.5	3.1	2.8	2.7	2.4	2.2	1.9	1.7
1999.....	3.7	3.4	3.2	2.4	2.4	2.5	2.3	1.9	1.5
Hispanic, both sexes									
2003.....	7.7	5.6	3.5	4.9	4.2	5.4	5.3	*	*
2002.....	6.1	5.9	5.4	4.4	5.2	5.3	*	*	*
2001.....	6.6	6.0	4.5	4.4	5.7	4.8	*	*	*
2000.....	8.8	7.3	5.3	4.3	5.0	7.1	*	*	*
1999.....	9.0	5.2	5.9	4.5	6.2	*	8.7	*	*
Hispanic, male									
2003.....	12.9	9.6	6.8	9.8	9.1	11.1	*	*	*
2002.....	10.2	10.0	10.3	8.5	10.2	11.8	*	*	*
2001.....	11.0	10.3	8.8	8.4	11.7	10.1	*	*	*
2000.....	14.3	12.5	10.1	7.8	9.7	13.6	*	*	*
1999.....	14.1	8.8	10.6	8.9	12.0	*	21.1	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year

45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Hispanic, female

2003.....	2.2	*	*	*	*	*	*	*	*
2002.....	1.9	*	*	*	*	*	*	*	*
2001.....	2.2	*	*	*	*	*	*	*	*
2000.....	3.3	*	*	*	*	*	*	*	*
1999.....	3.9	*	*	*	*	*	*	*	*

Non-Hispanic 4/, both sexes

2003.....	11.8	11.4	10.9	10.0	10.1	11.6	13.3	14.7	12.7
2002.....	11.5	10.8	11.1	9.9	10.4	12.2	14.1	15.8	12.7
2001.....	11.0	10.8	10.8	10.0	10.7	11.7	14.1	15.5	12.9
2000.....	10.8	10.5	9.9	9.4	9.9	11.7	13.8	15.1	14.4
1999.....	10.8	9.9	10.2	9.7	10.2	12.6	14.2	14.7	13.6

Non-Hispanic 4/, male

2003.....	19.5	19.5	18.6	18.2	19.1	23.2	28.7	35.0	38.7
2002.....	19.1	18.6	19.3	17.6	19.3	24.9	30.8	38.3	39.5
2001.....	18.5	18.2	18.5	17.2	20.0	23.5	30.9	37.9	40.8
2000.....	18.1	17.6	16.9	16.5	18.2	23.5	30.4	37.4	45.8
1999.....	18.1	16.6	17.5	17.6	19.3	25.3	31.1	36.6	43.9

Non-Hispanic 4/, female

2003.....	4.3	3.6	3.6	2.5	2.2	2.3	2.4	2.3	1.3
2002.....	4.2	3.4	3.4	2.9	2.6	2.1	2.3	2.1	1.2
2001.....	3.7	3.6	3.5	3.4	2.7	2.3	2.4	2.2	1.4
2000.....	3.7	3.6	3.3	2.9	2.8	2.4	2.2	1.9	1.7
1999.....	3.7	3.5	3.4	2.5	2.4	2.7	2.4	1.8	1.5

Non-Hispanic White, both sexes

2003.....	12.0	12.0	11.7	10.8	11.0	12.6	14.4	15.7	13.8
2002.....	11.8	11.3	12.0	10.6	11.3	13.3	14.9	17.1	13.7
2001.....	11.4	11.4	11.5	10.9	11.7	12.5	15.1	16.4	14.0
2000.....	10.8	10.8	10.3	10.1	10.7	12.4	14.9	16.1	15.5
1999.....	10.8	10.3	10.7	10.2	11.0	13.5	15.0	15.6	14.7

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Non-Hispanic White, male

2003.....	19.5	20.4	19.8	19.5	20.7	24.9	30.9	37.4	42.1
2002.....	19.2	19.3	20.4	18.8	20.9	26.8	32.3	41.3	42.5
2001.....	19.0	19.0	19.3	18.5	21.4	24.8	32.9	39.9	44.0
2000.....	18.0	18.0	17.5	17.5	19.4	24.6	32.8	39.7	49.6
1999.....	17.8	17.3	18.0	18.2	20.6	26.7	32.9	38.7	47.5

Non-Hispanic White, female

2003.....	4.6	3.8	3.9	2.7	2.4	2.4	2.5	2.3	1.4
2002.....	4.5	3.4	3.9	3.0	2.9	2.2	2.4	2.3	1.2
2001.....	3.9	3.8	3.9	3.8	3.0	2.4	2.5	2.2	1.5
2000.....	3.7	3.8	3.5	3.2	3.0	2.6	2.3	2.0	1.8
1999.....	3.7	3.5	3.8	2.7	2.6	2.9	2.5	1.9	1.6

Non-Hispanic Black, both sexes

2003.....	12.8	10.4	7.4	6.3	6.0	6.4	5.7	6.5	*
2002.....	13.2	10.3	7.3	7.1	5.2	6.5	8.6	*	*
2001.....	11.4	9.9	8.1	5.4	5.9	7.6	6.9	8.3	*
2000.....	13.4	10.5	7.5	5.9	6.3	7.9	4.7	6.9	*
1999.....	13.7	8.8	8.7	8.0	6.6	7.0	8.5	6.2	*

Non-Hispanic Black, male

2003.....	24.1	18.6	13.9	11.9	11.6	12.5	12.6	15.1	*
2002.....	24.6	18.0	14.6	12.4	10.5	14.0	20.2	*	*
2001.....	20.5	17.9	16.3	10.2	12.0	16.2	16.1	20.4	*
2000.....	23.7	19.2	13.9	11.5	13.1	18.5	9.6	17.2	*
1999.....	24.7	15.5	17.8	15.7	13.0	15.9	19.8	*	*

Worktable Orig291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Non-Hispanic Black, female	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
2003.....	2.9	3.5	*	*	*	*	*	*	*
2002.....	3.3	3.7	*	*	*	*	*	*	*
2001.....	3.5	3.0	*	*	*	*	*	*	*
2000.....	4.5	3.1	*	*	*	*	*	*	*
1999.....	4.3	3.2	*	*	*	*	*	*	*

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Figures for origin not stated are included in "All origins" but not distributed among specified origins.

4/ Includes races other than white and black.

5/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

6/ Included in selected categories.

7/ Figures may differ from those previously shown as they are based on a modified list of causes; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003."

Source: CDC/NCHS, National Vital Statistics System, Mortality.