

**INTERACTION PROFILE FOR:
1,1,1-TRICHLOROETHANE, 1,1-DICHLOROETHANE,
TRICHLOROETHYLENE, AND TETRACHLOROETHYLENE**

**U.S. Department of Health and Human Services
Public Health Service
Agency for Toxic Substances and Disease Registry**

May 2004

PREFACE

The Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) mandates that the Agency for Toxic Substances and Disease Registry (ATSDR) shall assess whether adequate information on health effects is available for the priority hazardous substances. Where such information is not available or under development, ATSDR shall, in cooperation with the National Toxicology Program, initiate a program of research to determine these health effects. The Act further directs that where feasible, ATSDR shall develop methods to determine the health effects of substances in combination with other substances with which they are commonly found. The Food Quality Protection Act (FQPA) of 1996 requires that factors to be considered in establishing, modifying, or revoking tolerances for pesticide chemical residues shall include the available information concerning the cumulative effects of substances that have a common mechanism of toxicity, and combined exposure levels to the substance and other related substances. The FQPA requires that the Administrator of the Environmental Protection Agency (EPA) consult with the Secretary of the Department of Health and Human Services (which includes ATSDR) in implementing some of the provisions of the act.

To carry out these legislative mandates, ATSDR's Division of Toxicology (DT) has developed and coordinated a mixtures program that includes trend analysis to identify the mixtures most often found in environmental media, *in vivo* and *in vitro* toxicological testing of mixtures, quantitative modeling of joint action, and methodological development for assessment of joint toxicity. These efforts are interrelated. For example, the trend analysis suggests mixtures of concern for which assessments need to be conducted. If data are not available, further research is recommended. The data thus generated often contribute to the design, calibration or validation of the methodology. This pragmatic approach allows identification of pertinent issues and their resolution as well as enhancement of our understanding of the mechanisms of joint toxic action. All the information obtained is thus used to enhance existing or developing methods to assess the joint toxic action of environmental chemicals. Over a number of years, ATSDR scientists in collaboration with mixtures risk assessors and laboratory scientists have developed approaches for the assessment of the joint toxic action of chemical mixtures. As part of the mixtures program a series of documents, Interaction Profiles, are being developed for certain priority mixtures that are of special concern to ATSDR.

The purpose of an Interaction Profile is to evaluate data on the toxicology of the "whole" priority mixture (if available) and on the joint toxic action of the chemicals in the mixture in order to recommend approaches for the exposure-based assessment of the potential hazard to public health. Joint toxic action includes additivity and interactions. A weight-of-evidence approach is commonly used in these documents to evaluate the influence of interactions in the overall toxicity of the mixture. The weight-of-evidence evaluations are qualitative in nature, although ATSDR recognizes that observations of toxicological interactions depend greatly on exposure doses and that some interactions appear to have thresholds. Thus, the interactions are evaluated in a qualitative manner to provide a sense of what influence the interactions may have when they do occur.

CONTRIBUTORS

CHEMICAL MANAGER(S)/AUTHORS:

[Sharon Wilbur, M.A.](#)

ATSDR, Division of Toxicology, Atlanta, GA

Author:

[Peter McClure, Ph.D., D.A.B.T.](#)

Syracuse Research Corporation, North Syracuse, NY

PEER REVIEW

A peer review panel was assembled for this profile. The panel consisted of the following members:

1. Dr. James V. Bruckner
College of Pharmacy
University of Georgia
Athens, GA
2. Dr. Kannan Krishnan
Department of Occupational and Environmental Health
University of Montreal
Montreal, Quebec
3. Dr. Harihari Mehendale
College of Pharmacy
University of Louisiana at Monroe
Monroe, LA

All reviewers were selected in conformity with the conditions for peer review specified in Sect 104(I)(13) of the Comprehensive Environmental Response, Compensation, and Liability Act, as amended.

Scientists from the Agency for Toxic Substances and Disease Registry (ATSDR) have reviewed the peer reviewers' comments and determined which comments will be included in the profile. A listing of the peer reviewers' comments not incorporated in the profile, with a brief explanation of the rationale for their exclusion, exists as part of the administrative record for this compound. A list of databases reviewed and a list of unpublished documents cited are also included in the administrative record.

The citation of the peer review panel should not be understood to imply its approval of the profile's final content. The responsibility for the content of this profile lies with the ATSDR.

SUMMARY

1,1,1-Trichloroethane, 1,1-dichloroethane, trichloroethylene, and tetrachloroethylene frequently occur together in water samples collected from and in the vicinity of National Priorities List (NPL) hazardous waste sites and other sites assessed by the Agency for Toxic Substances and Disease Registry (ATSDR). These chemicals occur together more frequently than other volatile organic chemicals at the sites. In an unpublished survey of ATSDR Public Health Assessments for 210 NPL hazardous waste sites, this mixture of chemicals was found in groundwater samples from 95% of the sites, in soil samples from 23% of the sites, and in air samples from 12% of the sites. The purposes of this profile are (1) to evaluate data on the toxicology of mixtures of 1,1,1-trichloroethane, 1,1-dichloroethane, trichloroethylene, and tetrachloroethylene, (2) to evaluate data on the joint toxic actions (e.g., additive, less-than-additive, or greater-than-additive joint actions) of these chemicals in producing health hazards, and (3) to make recommendations for exposure-based assessments of the potential impact of joint toxic action of the mixture on public health.

There are no studies available that directly characterize health hazards and dose-response relationships for exposures to “whole” mixtures containing 1,1,1-trichloroethane, 1,1-dichloroethane, trichloroethylene, and tetrachloroethylene. Furthermore, physiologically based pharmacokinetic (PBPK) models have not been developed to predict dispositional and toxicological outcomes of joint action of mixtures of these four chemicals.

Exposure to each of the individual chemicals can produce neurological impairment via parent chemical-induced physical and chemical changes in neuronal membranes and cause non-carcinogenic and carcinogenic responses (via reactive metabolites) in the liver and kidney of animals. No studies are available that directly examine joint toxic actions of binary or trinary mixtures of these chemicals on the nervous system, but additive joint action is plausible. Limited studies of joint toxic action of binary or trinary mixtures of these chemicals on the liver and kidney provide no evidence of greater-than-additive joint toxic actions. Additive joint action on the liver and kidney is plausible for binary combinations of each of the components, with the exception of limited evidence that tetrachloroethylene may inhibit the toxic action of trichloroethylene on the liver and kidney (Goldsworthy and Popp 1987; Seiji et al. 1989).

A component-based hazard index approach that assumes additive joint toxic action and uses ATSDR Minimal Risk Levels (MRLs) based on neurological impairment is recommended for exposure-based assessments of possible health hazards from exposure to mixtures of 1,1,1-trichloroethane, 1,1-dichloroethane, trichloroethylene, and tetrachloroethylene. There is no evidence to indicate that greater-than-additive interactions would cause liver and kidney effects to occur at exposure levels lower than those influencing the nervous system.

CONTENTS

PREFACE	-ii-
CONTRIBUTORS	-iii-
PEER REVIEW	-v-
SUMMARY	-vii-
CONTENTS	-ix-
LIST OF FIGURES AND TABLES	-xi-
LIST OF ACRONYMS, ABBREVIATIONS, AND SYMBOLS	-xiii-
1. Introduction	1
2. Joint Toxic Action Data for the Mixture of Concern and Component Mixtures	3
2.1 Mixture of Concern	3
2.2 Component Mixtures	3
2.2.1 1,1,1-Trichloroethane, Trichloroethylene, and Tetrachloroethylene	3
2.2.2 1,1,1-Trichloroethane and 1,1-Dichloroethane	5
2.2.3 1,1,1-Trichloroethane and Trichloroethylene	8
2.2.4 1,1,1-Trichloroethane and Tetrachloroethylene	12
2.2.5 1,1-Dichloroethane and Trichloroethylene	15
2.2.6 1,1-Dichloroethane and Tetrachloroethylene	16
2.2.7 Trichloroethylene and Tetrachloroethylene	17
2.3 Relevance of the Joint Toxic Action Data and Approaches to Public Health	24
3. Recommendation for Exposure-Based Assessment of Joint Toxic Action of the Mixture	43
4. Conclusions	47
5. List of References	49
Appendix A: Background Information for 1,1,1-Trichloroethane	59
A.1 Toxicokinetics	59
A.2 Health Effects	61
A.3 Mechanisms of Action	62
A.4 Health Guidelines	64
Appendix B: Background Information for 1,1-Dichloroethane	67
B.1 Toxicokinetics	67
B.2 Health Effects	68
B.3 Mechanisms of Action	68
B.4 Health Guidelines	69

Appendix C: Background Information for Trichloroethylene	71
C.1 Toxicokinetics	71
C.2 Health Effects	73
C.3 Mechanisms of Action	74
C.4 Health Guidelines	75
Appendix D: Background Information for Tetrachloroethylene	77
D.1 Toxicokinetics	77
D.2 Health Effects	79
D.3 Mechanisms of Action	80
D.4 Health Guidelines	81
Appendix E: Chemical Structures of Mixture Components	85

LIST OF FIGURES AND TABLES

Table 1. Summary of Available Data on the Influence of 1,1,1-Trichloroethane on Toxicity/ Carcinogenicity of Trichloroethylene and the Influence of Trichloroethylene on Toxicity/Carcinogenicity after 1,1,1-Trichloroethane after Simultaneous Exposure	11
Table 2. Summary of Available Data on the Influence of 1,1,1-Trichloroethane on Toxicity/ Carcinogenicity of Tetrachloroethylene and the Influence of Tetrachloroethylene on Toxicity/Carcinogenicity of 1,1,1-Trichloroethane after Simultaneous Exposure	14
Table 3. Summary of Available Data on the Influence of Trichloroethylene on Toxicity/Carcinogenicity of Tetrachloroethylene and the Influence of Tetrachloroethylene on Toxicity/Carcinogenicity of Trichloroethylene after Simultaneous Exposure	22
Table 4. Health Effects Forming the Basis of ATSDR Inhalation and Oral MRLs for Chemicals of Concern	25
Figure 1. Binary Weight-of-Evidence Scheme for the Assessment of Chemical Interactions	28
Table 5. Effect of 1,1,1-Trichloroethane on 1,1-Dichloroethane	29
Table 6. Effect of 1,1-Dichloroethane on 1,1,1-Trichloroethane	30
Table 7. Effect of 1,1,1-Trichloroethane on Trichloroethylene	31
Table 8. Effect of Trichloroethylene on 1,1,1-Trichloroethane	32
Table 9. Effect of Tetrachloroethylene on 1,1,1-Trichloroethane	33
Table 10. Effect of 1,1,1-Trichloroethane on Tetrachloroethylene	34
Table 11. Effect of 1,1-Dichloroethane on Trichloroethylene	35
Table 12. Effect of Trichloroethylene on 1,1-Dichloroethane	36
Table 13. Effect of 1,1-Dichloroethane on Tetrachloroethylene	37
Table 14. Effect of Tetrachloroethylene on 1,1-Dichloroethane	38
Table 15. Effect of Trichloroethylene on Tetrachloroethylene	39
Table 16. Effect of Tetrachloroethylene on Trichloroethylene	40
Table 17. Matrix of BINWOE Determinations for Nervous System Effects from Simultaneous Exposure to Chemicals of Concern	41
Table 18. Matrix of BINWOE Determinations for Liver and Kidney Effects from Simultaneous Exposure to Chemicals of Concern	42

LIST OF ACRONYMS, ABBREVIATIONS, AND SYMBOLS

ALT	alanine aminotransferase
AST	aspartate aminotransferase
ATSDR	Agency for Toxic Substances and Disease Registry
BINWOE	binary weight of evidence
bw	body weight
CYP	cytochrome P-450
DCVC	S-(1,2-dichlorovinyl)-L-cysteine
DCVG	S-(1,2-dichlorovinyl)glutathione
EPA	Environmental Protection Agency
IRIS	Integrated Risk Information System
kg	kilogram
L	liter
LOAEL	lowest-observed-adverse-effect level
LSE	Levels of Significant Exposure
mg	milligram
mL	milliliter
mmol	millimole
MRL	Minimal Risk Level
NOAEL	no-observed-adverse-effect level
NPL	National Priorities List
PBPK	physiologically based pharmacokinetic
ppm	parts per million
RfC	Reference Concentration
RfD	Reference Dose
SDH	sorbitol dehydrogenase
U.S.	United States
WOE	weight of evidence
>	greater than
≥	greater than or equal to
=	equal to
<	less than
≤	less than or equal to