This is an Archived Document.

Table 2. Tuberculosis Cases, Percentages, and Case Rates per 100,000 Population by Hispanic Ethnicity and non-Hispanic Race: United States, 1993-2003

					Non-Hispanic																						
	Total	Hispanic or Latino ¹			Multiple Race ²			American Indian or Alaska Native			Asian³			Asian or Pacific Islander ⁴			Black or African American			Native Hawaiian or Other Pacific Islander ⁵			White			Unknown or Missing	
Year	Cases	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)
1993	25,287	5,194	(21)	20.6				274	(1)	14.5				3,680	(15)	45.0	8,951	(35)	29.1				6,922	(27)	3.6	266	(1)
1994	24,361	5,074	(21)	19.4				332	(1)	17.4				3,821	(16)	44.9	8,345	(34)	26.7				6,494	(27)	3.4	295	(1)
1995	22,860	4,847	(21)	17.9				319	(1)	16.5				3,997	(17)	45.2	7,555	(33)	23.9				5,989	(26)	3.1	153	(1)
1996	21,337	4,533	(21)	16.1				284	(1)	14.5				3,814	(18)	41.5	7,106	(33)	22.2				5,506	(26)	2.8	94	(0)
1997	19,851	4,228	(21)	14.5				264	(1)	13.3				3,833	(19)	40.2	6,610	(33)	20.4				4,872	(25)	2.5	44	(0)
1998	18,361	4,099	(22)	13.5				253	(1)	12.6				3,623	(20)	36.7	5,831	(32)	17.8				4,495	(24)	2.3	60	(0)
1999	17,531	3,875	(22)	12.4				240	(1)	11.8				3,591	(20)	35.3	5,552	(32)	16.8				4,224	(24)	2.2	49	(0)
2000	16,377	3,805	(23)	10.7				236	(1)	10.0				3,451	(21)	30.1	5,161	(32)	14.7				3,674	(22)	1.9	50	(0)
2001	15,989	4,001	(25)	10.7				233	(1)	9.8				3,552	(22)	29.8	4,796	(30)	13.4				3,357	(21)	1.7	50	(0)
2002	15,075	3,976	(26)	10.3				187	(1)	7.8				3,345	(22)	27.0	4,439	(29)	12.3				3,041	(20)	1.5	87	(1)
2003	14,874	4,115	(28)	10.3	37	(0)	1.0	176	(1)	8.1	3,425	(23)	29.3				4,145	(28)	11.6	85	(1)	21.8	2,790	(19)	1.4	101	(1)

¹Persons of Hispanic or Latino origin may be of any race or multiple race.

Note: Case counts for years 1993–2002 may include persons of multiple race. Previously published rates for 1993–1999 have been updated using U.S. Census midyear post-censal resident population files (http://eire.census.gov/popest/archives/national/nation3.php). Denominators for computing case rates were obtained as follows: For 2000, 2001, and 2002, NCHS bridged population estimates were used (http://www.cdc.gov/nchs/about/major/dvs/popbridge/popbridge.htm). Denominators for computing 2003 case rates were obtained from the 2003 U.S. Census Bureau midyear population estimates by sex, race, and Hispanic origin (www.eire.census.gov/popsest/data/national/tables/NC-EST2003-srh.pop). Case counts for race categories (American Indian or Alaska Native, Asian, Black or African American, Native Hawaiian or Other Pacific Islander, and White) do not include persons of Hispanic ethnicity or multiple race. See Technical Notes (page 9).

See Surveillance Slides #8 and #9.

Zero % (0) denotes < 1%.

²Indicates two or more races reported for a person. Category first reported in 2003. Does not include persons of Hispanic or Latino ethnicity.

³Asian race first reported in 2003.

⁴Asian or Pacific Islander race reported 1993–2002.

⁵Native Hawaiian or Other Pacific Islander race first reported in 2003.