

Table 5. Tuberculosis Cases by Case Verification Criterion and by Site of Disease: United States, 1990-2000

Year	Total Cases	Verification Criterion ¹								Site of Disease			
		Positive Culture		Positive Smear		Clinical Case Definition		Provider Diagnosis		Pulmonary ²		Extra-pulmonary	
		No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
1990	25,701	20,897	81	556	2	2,958	12	1,290	5	21,576	84	4,091	16
1991	26,283	21,417	81	388	1	2,992	11	1,486	6	21,937	83	4,327	16
1992	26,673	21,398	80	407	2	3,141	12	1,727	6	22,371	84	4,288	16
1993	25,287	20,081	79	309	1	2,994	12	1,903	8	21,255	84	3,995	16
1994	24,361	19,537	80	236	1	2,794	11	1,794	7	20,385	84	3,964	16
1995	22,860	18,292	80	220	1	2,664	12	1,684	7	18,991	83	3,860	17
1996	21,337	17,234	81	150	1	2,556	12	1,397	7	17,445	82	3,870	18
1997	19,851	16,015	81	177	1	2,355	12	1,304	7	16,285	82	3,554	18
1998	18,361	14,830	81	166	1	2,207	12	1,158	6	14,813	81	3,541	19
1999	17,531	13,997	80	176	1	2,058	12	1,300	7	14,083	80	3,438	20
2000	16,377	13,035	80	169	1	1,901	12	1,272	8	13,142	80	3,220	20

1. Based on the public health surveillance case definition for tuberculosis: CDC. Case definitions for infectious conditions under public health surveillance. MMWR 1997;46(No. RR-10):40-41. See Appendix B.

2. Includes cases of both pulmonary and extrapulmonary disease and cases of military TB.

Note: See Technical Notes (Appendix A) for a description of national TB surveillance.

Table 6. Pulmonary Tuberculosis Cases by Sputum Smear and Sputum Culture Results: United States, 1990-2000

Year	Total Pulmonary Cases ¹	Sputum Smear Results						Sputum Culture Results					
		Positive		Negative		Not Done or Unknown		Positive		Negative		Not Done or Unknown	
		No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
1990	21,576	9,391	44	6,865	32	5,320	25	14,816	69	2,124	10	4,636	21
1991	21,937	9,095	41	7,281	33	5,561	25	15,022	68	2,232	10	4,683	21
1992	22,371	8,975	40	7,413	33	5,983	27	15,124	68	2,476	11	4,771	21
1993	21,255	9,324	44	7,747	36	4,184	20	14,708	69	2,675	13	3,872	18
1994	20,385	8,845	43	7,770	38	3,770	18	14,080	69	2,618	13	3,687	18
1995	18,991	8,068	42	7,717	41	3,206	17	13,236	70	2,597	14	3,158	17
1996	17,445	7,449	43	7,337	42	2,659	15	12,232	70	2,507	14	2,706	16
1997	16,285	6,882	42	6,878	42	2,525	16	11,481	71	2,226	14	2,578	16
1998	14,813	6,630	45	6,016	41	2,167	15	10,472	71	2,101	14	2,240	15
1999	14,083	6,252	44	5,626	40	2,205	16	9,777	69	2,049	15	2,257	16
2000	13,142	5,865	45	5,332	41	1,945	15	9,214	70	1,912	15	2,016	15

1. Includes cases of both pulmonary and extrapulmonary disease and cases of military TB.