

Table 35. Tuberculosis Cases and Percentages by Initial Drug Regimen: Reporting Areas, 2007

Reporting Area	Total Cases	Cases in Persons Alive at Diagnosis	Cases with Information on Initial Drug Regimen		Percentage of Cases in Persons with Initial Drug Regimen ^{1,2}			
			No.	(%)	IR	IRZ	IRZ,E/S	IRZE ³
United States	13,299	12,999	12,954	(99.7)	(1.1)	(4.7)	(83.3)	(83.3)
Alabama	175	167	167	(100.0)	(0.6)	(4.2)	(85.6)	(85.6)
Alaska	51	51	50	(98.0)	(0.0)	(12.0)	(86.0)	(86.0)
Arizona	304	291	289	(99.3)	(1.4)	(5.2)	(79.6)	(79.2)
Arkansas	106	104	104	(100.0)	(7.7)	(38.5)	(39.4)	(39.4)
California	2,726	2,676	2,668	(99.7)	(0.9)	(2.4)	(87.6)	(87.5)
Colorado	111	103	102	(99.0)	(0.0)	(2.9)	(80.4)	(80.4)
Connecticut	108	106	105	(99.1)	(1.0)	(4.8)	(84.8)	(84.8)
Delaware	19	19	19	(100.0)	(0.0)	(5.3)	(89.5)	(89.5)
District of Columbia	60	57	56	(98.2)	(0.0)	(1.8)	(98.2)	(98.2)
Florida	989	968	968	(100.0)	(0.6)	(5.9)	(84.1)	(84.1)
Georgia	474	464	464	(100.0)	(0.4)	(5.2)	(62.5)	(62.5)
Hawaii	122	121	121	(100.0)	(2.5)	(9.9)	(79.3)	(79.3)
Idaho	9	8	8	(100.0)	(0.0)	(0.0)	(75.0)	(75.0)
Illinois	521	514	514	(100.0)	(1.0)	(3.5)	(83.1)	(83.1)
Indiana	128	123	123	(100.0)	(0.8)	(3.3)	(92.7)	(92.7)
Iowa	43	42	42	(100.0)	(0.0)	(9.5)	(90.5)	(90.5)
Kansas	59	57	57	(100.0)	(0.0)	(8.8)	(89.5)	(89.5)
Kentucky	120	118	118	(100.0)	(0.0)	(7.6)	(65.3)	(65.3)
Louisiana	218	214	213	(99.5)	(0.0)	(2.3)	(95.3)	(95.3)
Maine	19	19	19	(100.0)	(0.0)	(0.0)	(89.5)	(89.5)
Maryland	270	266	266	(100.0)	(1.1)	(4.5)	(90.6)	(90.6)
Massachusetts	224	222	221	(99.5)	(0.9)	(1.4)	(82.8)	(82.8)
Michigan	226	215	212	(98.6)	(3.3)	(18.9)	(74.1)	(74.1)
Minnesota	238	235	234	(99.6)	(0.9)	(7.7)	(84.6)	(84.6)
Mississippi	137	132	132	(100.0)	(3.0)	(8.3)	(80.3)	(80.3)
Missouri	119	112	112	(100.0)	(0.0)	(7.1)	(75.0)	(75.0)
Montana	11	11	11	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
Nebraska	25	25	24	(96.0)	(0.0)	(8.3)	(87.5)	(87.5)
Nevada	102	94	94	(100.0)	(0.0)	(0.0)	(97.9)	(97.9)
New Hampshire	11	9	9	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
New Jersey	467	459	458	(99.8)	(0.9)	(2.8)	(86.5)	(86.5)
New Mexico	51	51	51	(100.0)	(0.0)	(0.0)	(92.2)	(92.2)
New York State ⁴	261	252	252	(100.0)	(1.2)	(5.2)	(86.9)	(86.9)
New York City	914	897	897	(100.0)	(1.0)	(3.0)	(85.3)	(85.3)
North Carolina	345	337	337	(100.0)	(0.3)	(3.9)	(89.3)	(89.3)
North Dakota	7	7	6	(85.7)	(0.0)	(0.0)	(83.3)	(83.3)
Ohio	252	247	247	(100.0)	(0.0)	(5.7)	(83.0)	(83.0)
Oklahoma	149	145	145	(100.0)	(11.0)	(19.3)	(62.8)	(62.8)
Oregon	94	92	91	(98.9)	(0.0)	(2.2)	(93.4)	(93.4)
Pennsylvania	276	271	262	(96.7)	(0.4)	(2.3)	(42.4)	(42.4)
Rhode Island	45	45	44	(97.8)	(6.8)	(0.0)	(84.1)	(84.1)
South Carolina	218	213	213	(100.0)	(0.5)	(10.3)	(82.2)	(82.2)
South Dakota	13	13	13	(100.0)	(0.0)	(15.4)	(69.2)	(69.2)
Tennessee	234	231	231	(100.0)	(0.0)	(6.1)	(61.0)	(61.0)
Texas	1,510	1,471	1,464	(99.5)	(1.9)	(3.1)	(88.5)	(88.5)
Utah	39	39	39	(100.0)	(0.0)	(5.1)	(89.7)	(89.7)
Vermont	3	3	3	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
Virginia	309	303	299	(98.7)	(0.3)	(3.7)	(89.6)	(89.6)
Washington	291	288	288	(100.0)	(0.0)	(3.8)	(88.2)	(88.2)
West Virginia	24	21	21	(100.0)	(4.8)	(4.8)	(81.0)	(81.0)
Wisconsin	70	69	69	(100.0)	(0.0)	(5.8)	(85.5)	(85.5)
Wyoming	2	2	2	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
American Samoa ⁵	3	3	3	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
Fed. States of Micronesia ⁵	66	65	64	(98.5)	(0.0)	(18.8)	(79.7)	(79.7)
Guam ⁵	92	90	90	(100.0)	(0.0)	(20.0)	(75.6)	(75.6)
Marshall Islands ⁵	117	112	111	(99.1)	(0.0)	(19.8)	(78.4)	(78.4)
N. Mariana Islands ⁵	41	40	40	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)
Puerto Rico ⁵	98	93	93	(100.0)	(0.0)	(3.2)	(91.4)	(91.4)
Republic of Palau ⁵	11	11	11	(100.0)	(0.0)	(9.1)	(72.7)	(72.7)
U.S. Virgin Islands ⁵

¹Overall U.S. percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥75% of cases.

²I=isoniazid; R=rifampin; Z=pyrazinamide; E=ethambutol; S=streptomycin.

³Streptomycin is no longer considered a first-line drug. CDC. Treatment of Tuberculosis. American Thoracic Society, CDC, and Infectious Diseases Society of America. MMWR 2003;52(No. RR-11): 1-77.

⁴Excludes New York City.

⁵Not included in U.S. totals.

Note: Excluding cases with no information on drug regimen, 127 (0.98%) persons were not started on any drugs, 21 (<0.2%) were started on one drug, and 1,272 (9.8%) had an initial multidrug regimen other than IR, IRZ, or IRZ,E/S.

Ellipses indicate data not available.