Table 16. Tuberculosis Case Rates per 100,000 Population by Hispanic Ethnicity and Non-Hispanic Race, Sex, and Age Group: United States, 2006

Race/Ethnicity and Sex	Age Group						
	All Ages	Under 5	5–14	15–24	25–44	45–64	≥65
Total Rate	4.6	2.4	0.8	3.6	5.6	5.4	7.2
Male	5.8	2.4	0.8	4.0	6.6	7.7	10.4
Female	3.4	2.4	0.8	3.3	4.6	3.2	4.8
Hispanic or Latino[1]	9.2	5.5	1.6	9.2	10.7	13.1	20.5
Male	11.7	5.7	1.6	11.3	13.6	18.6	29.3
Female	6.5	5.3	1.6	6.9	7.3	7.7	13.9
American Indian or Alaska Native	7.4	2.3	1.4	2.7	8.6	12.6	16.0
Male	8.4	2.2	1.6	1.9	10.1	16.1	18.2
Female	6.4	2.3	1.1	3.5	7.2	9.5	14.3
Asian	25.6	5.2	2.5	19.1	26.1	30.9	65.2
Male	29.0	5.0	2.5	17.1	25.3	41.2	95.0
Female	22.3	5.5	2.5	21.2	26.8	22.0	42.7
Black/African American	10.2	4.5	2.0	6.7	12.3	15.7	16.5
Male	13.2	4.2	2.1	6.8	15.2	24.8	25.6
Female	7.4	4.9	2.0	6.6	9.6	8.2	10.8
Native Hawaiian or Pacific Islander	13.6	3.6	6.4	17.0	16.1	13.9	17.5
Male	14.4	0.0	6.2	11.0	21.6	16.4	15.5
Female	12.7	7.6	6.5	23.4	10.4	11.5	19.1
White	1.2	0.4	0.1	0.4	1.0	1.5	2.9
Male	1.6	0.4	0.1	0.4	1.2	2.3	4.2
Female	0.8	0.4	0.1	0.3	0.8	0.7	1.9
Multiple Race[2]	0.9	0.2	0.1	0.3	1.0	2.5	3.2
Male	1.0	0.3	0.0	0.3	1.3	3.1	4.3
Female	0.7	0.0	0.2	0.3	0.6	1.9	2.4

[1]Persons of Hispanic or Latino origin may be of any race or multiple race.
[2]Indicates two or more races reported for a person. Category first reported in 2003.
Note: Denominators for computing case rates were obtained from the U.S. Census Bureau Monthly Postcensal Resident Population, by single year of age, sex, race, and Hispanic origin: July 1, 2006 (http://www.census.gov/popest/national/asrh/files/NC-EST2006-ALLDATA-R-File14.dat) (Accessed May 21, 2007).
Case counts for race categories (American Indian or Alaska Native, Asian, Black or African American, Native Hawaiian or Other Pacific Islander, and White) are mutually exclusive and do not include persons of Hispanic ethnicity or multiple race. Multiple Race does not include persons of Hispanic ethnicity.
See Technical Notes (page 9).
See Surveillance Slide #10.