

Table 8. Tuberculosis Cases and Percentages by Case Verification Criterion and Site of Disease: United States, 1993-2003

Year	Total Cases	Verification Criterion ¹								Site of Disease			
		Positive Culture		Positive Smear		Clinical Case Definition		Provider Diagnosis		Pulmonary ²		Extra-pulmonary	
		No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	25,287	20,081	(79)	309	(1)	2,994	(12)	1,903	(8)	21,255	(84)	3,995	(16)
1994	24,361	19,537	(80)	236	(1)	2,794	(11)	1,794	(7)	20,385	(84)	3,964	(16)
1995	22,860	18,292	(80)	220	(1)	2,664	(12)	1,684	(7)	18,991	(83)	3,860	(17)
1996	21,337	17,234	(81)	150	(1)	2,556	(12)	1,397	(7)	17,445	(82)	3,870	(18)
1997	19,851	16,015	(81)	177	(1)	2,355	(12)	1,304	(7)	16,285	(82)	3,554	(18)
1998	18,361	14,830	(81)	166	(1)	2,207	(12)	1,158	(6)	14,813	(81)	3,541	(19)
1999	17,531	13,997	(80)	176	(1)	2,058	(12)	1,300	(7)	14,083	(80)	3,438	(20)
2000	16,377	13,035	(80)	169	(1)	1,901	(12)	1,272	(8)	13,142	(80)	3,220	(20)
2001	15,989	12,780	(80)	131	(1)	1,843	(12)	1,235	(8)	12,768	(80)	3,212	(20)
2002	15,075	11,993	(80)	108	(1)	1,774	(12)	1,200	(8)	11,912	(79)	3,143	(21)
2003	14,874	11,704	(79)	128	(1)	1,738	(12)	1,304	(9)	11,832	(80)	3,029	(20)

¹Based on the public health surveillance case definition for tuberculosis: CDC. Case definitions for infectious conditions under public health surveillance. *MMWR* 1997;46(No. RR-10):40-41. See Appendix A.

²Includes cases among persons with both pulmonary and extrapulmonary disease and cases of miliary TB.

Note: See Technical Notes (page 9) for a description of national TB surveillance.

Table 9. Pulmonary Tuberculosis Cases and Percentages by Sputum Smear and Sputum Culture Results: United States, 1993-2003

Year	Total Pulmonary Cases ¹	Sputum Smear Result						Sputum Culture Result					
		Positive		Negative		Not Done or Unknown		Positive		Negative		Not Done or Unknown	
		No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	21,255	9,324	(44)	7,747	(36)	4,184	(20)	14,708	(69)	2,675	(13)	3,872	(18)
1994	20,385	8,845	(43)	7,770	(38)	3,770	(18)	14,080	(69)	2,618	(13)	3,687	(18)
1995	18,991	8,068	(42)	7,717	(41)	3,206	(17)	13,236	(70)	2,597	(14)	3,158	(17)
1996	17,445	7,449	(43)	7,337	(42)	2,659	(15)	12,232	(70)	2,507	(14)	2,706	(16)
1997	16,285	6,882	(42)	6,878	(42)	2,525	(16)	11,481	(71)	2,226	(14)	2,578	(16)
1998	14,813	6,630	(45)	6,016	(41)	2,167	(15)	10,472	(71)	2,101	(14)	2,240	(15)
1999	14,083	6,252	(44)	5,626	(40)	2,205	(16)	9,777	(69)	2,049	(15)	2,257	(16)
2000	13,142	5,865	(45)	5,332	(41)	1,945	(15)	9,214	(70)	1,912	(15)	2,016	(15)
2001	12,768	5,600	(44)	5,311	(42)	1,857	(15)	8,855	(69)	1,937	(15)	1,976	(15)
2002	11,912	5,380	(45)	4,774	(40)	1,758	(15)	8,274	(69)	1,748	(15)	1,890	(16)
2003	11,832	5,303	(45)	4,819	(41)	1,710	(14)	8,143	(69)	1,850	(16)	1,839	(16)

¹Includes cases among persons with both pulmonary and extrapulmonary disease and cases of miliary TB.