

Table 36. Culture-Positive Tuberculosis Cases and Percentages with Drug- Susceptibility Results, by Resistance to INH or Multidrug Resistance: 60 Reporting Areas, 2004

Reporting Area	Total Culture Positive Cases	Cases with Initial Drug-Susceptibility Testing Performed		Resistance ¹			
				Isoniazid		Isoniazid and Rifampin	
		No.	(%)	No.	(%)	No.	(%)
United States	11,315	10,621	(93.9)	846	(8.0)	125	(1.2)
Alabama	173	169	(97.7)	7	(4.1)	0	(0.0)
Alaska	36	36	(100.0)	2	(5.6)	0	(0.0)
Arizona	206	204	(99.0)	20	(9.8)	2	(1.0)
Arkansas	91	91	(100.0)	2	(2.2)	0	(0.0)
California	2,311	2,220	(96.1)	250	(11.3)	37	(1.7)
Colorado	75	74	(98.7)	4	(5.4)	1	(1.4)
Connecticut	92	88	(95.7)	12	(13.6)	1	(1.1)
Delaware	28	27	(96.4)	0	(0.0)	0	(0.0)
District of Columbia	66	62	(93.9)	2	(3.2)	1	(1.6)
Florida	862	815	(94.5)	57	(7.0)	8	(1.0)
Georgia	435	425	(97.7)	35	(8.2)	2	(0.5)
Hawaii	77	74	(96.1)	9	(12.2)	1	(1.4)
Idaho	9	9	(100.0)	0	(0.0)	0	(0.0)
Illinois	430	398	(92.6)	21	(5.3)	5	(1.3)
Indiana	109	109	(100.0)	1	(0.9)	0	(0.0)
Iowa	35	34	(97.1)	1	(2.9)	0	(0.0)
Kansas	42	41	(97.6)	4	(9.8)	1	(2.5)
Kentucky	104	101	(97.1)	3	(3.0)	0	(0.0)
Louisiana	181	167	(92.3)	4	(2.4)	1	(0.6)
Maine	13	12	(92.3)	0	(0.0)	0	(0.0)
Maryland	225	224	(99.6)	16	(7.1)	3	(1.3)
Massachusetts	216	210	(97.2)	24	(11.4)	4	(1.9)
Michigan	217	214	(98.6)	14	(6.5)	3	(1.4)
Minnesota	139	139	(100.0)	18	(12.9)	5	(3.6)
Mississippi	98	91	(92.9)	8	(8.8)	0	(0.0)
Missouri	103	100	(97.1)	9	(9.0)	0	(0.0)
Montana	12	12	(100.0)	2	(16.7)	0	(0.0)
Nebraska	30	30	(100.0)	2	(6.7)	0	(0.0)
Nevada	76	65	(85.5)	10	(15.4)	1	(1.5)
New Hampshire	18	18	(100.0)	2	(11.1)	1	(5.6)
New Jersey	386	382	(99.0)	42	(11.0)	3	(0.8)
New Mexico	38	34	(89.5)	2	(5.9)	2	(5.9)
New York State ²	247	243	(98.4)	26	(10.7)	6	(2.5)
New York City	800	779	(97.4)	81	(10.4)	17	(2.2)
North Carolina	289	287	(99.3)	20	(7.0)	1	(0.3)
North Dakota	3	2	(66.7)	—	—	—	—
Ohio	181	179	(98.9)	12	(6.7)	1	(0.6)
Oklahoma	111	98	(88.3)	2	(2.0)	0	(0.0)
Oregon	93	93	(100.0)	8	(8.6)	0	(0.0)
Pennsylvania	272	250	(91.9)	18	(7.2)	1	(0.4)
Rhode Island	32	32	(100.0)	2	(6.3)	0	(0.0)
South Carolina	182	171	(94.0)	5	(2.9)	0	(0.0)
South Dakota	9	8	(88.9)	1	(12.5)	0	(0.0)
Tennessee	216	208	(96.3)	5	(2.4)	0	(0.0)
Texas	1,342	1,024	(76.3)	46	(4.5)	14	(1.4)
Utah	31	31	(100.0)	4	(12.9)	0	(0.0)
Vermont	4	4	(100.0)	0	(0.0)	0	(0.0)
Virginia	266	235	(88.3)	9	(3.8)	1	(0.4)
Washington	203	202	(99.5)	18	(8.9)	2	(1.0)
West Virginia	22	22	(100.0)	0	(0.0)	0	(0.0)
Wisconsin	76	75	(98.7)	6	(8.0)	0	(0.0)
Wyoming	3	3	(100.0)	0	(0.0)	0	(0.0)
American Samoa ³	3	3	(100.0)	0	(0.0)	0	(0.0)
Fed. States of Micronesia ³	7	7	(100.0)	3	(42.9)	2	(28.6)
Guam ³	28	23	(82.1)	2	(8.7)	0	(0.0)
Marshall Islands ³	32	31	(96.9)	2	(6.5)	2	(6.5)
N. Mariana Islands ³	27	26	(96.3)	5	(19.2)	1	(3.8)
Puerto Rico ³	114	99	(86.8)	4	(4.0)	0	(0.0)
Republic of Palau ³	1	0	(0.0)
U.S. Virgin Islands ³

¹Isolates may be resistant to other drugs. Overall U.S. percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥ 75% of cases.

²Excludes New York City.

³Not included in U.S. totals.

Note: Ellipses indicate data not available.