

Table 34. Tuberculosis Cases and Percentages by Excess Alcohol Use¹: 60 Reporting Areas, 2004

| Reporting Area | Total Cases | Cases with Information on Excess Alcohol Use | | Cases Reporting Excess Alcohol Use | |
|--|---------------|--|---------------|------------------------------------|---------------|
| | | No. | (%) | No. | (%) |
| United States | 14,517 | 14,133 | (97.4) | 2,007 | (14.2) |
| Alabama | 211 | 211 | (100.0) | 32 | (15.2) |
| Alaska | 43 | 40 | (93.0) | 11 | (27.5) |
| Arizona | 272 | 252 | (92.6) | 48 | (19.0) |
| Arkansas | 132 | 132 | (100.0) | 13 | (9.8) |
| California | 2,989 | 2,917 | (97.6) | 304 | (10.4) |
| Colorado | 127 | 127 | (100.0) | 13 | (10.2) |
| Connecticut | 101 | 94 | (93.1) | 10 | (10.6) |
| Delaware | 32 | 32 | (100.0) | 5 | (15.6) |
| District of Columbia | 81 | 81 | (100.0) | 13 | (16.0) |
| Florida | 1,076 | 1,056 | (98.1) | 248 | (23.5) |
| Georgia | 536 | 507 | (94.6) | 75 | (14.8) |
| Hawaii | 116 | 77 | (66.4) | — | — |
| Idaho | 11 | 8 | (72.7) | — | — |
| Illinois | 569 | 506 | (88.9) | 59 | (11.7) |
| Indiana | 128 | 128 | (100.0) | 35 | (27.3) |
| Iowa | 47 | 47 | (100.0) | 7 | (14.9) |
| Kansas | 62 | 62 | (100.0) | 9 | (14.5) |
| Kentucky | 127 | 127 | (100.0) | 21 | (16.5) |
| Louisiana | 249 | 235 | (94.4) | 42 | (17.9) |
| Maine | 20 | 20 | (100.0) | 1 | (5.0) |
| Maryland | 314 | 310 | (98.7) | 21 | (6.8) |
| Massachusetts | 283 | 275 | (97.2) | 32 | (11.6) |
| Michigan | 273 | 262 | (96.0) | 18 | (6.9) |
| Minnesota | 199 | 198 | (99.5) | 8 | (4.0) |
| Mississippi | 119 | 119 | (100.0) | 23 | (19.3) |
| Missouri | 127 | 120 | (94.5) | 12 | (10.0) |
| Montana | 15 | 15 | (100.0) | 6 | (40.0) |
| Nebraska | 39 | 39 | (100.0) | 2 | (5.1) |
| Nevada | 95 | 92 | (96.8) | 9 | (9.8) |
| New Hampshire | 24 | 24 | (100.0) | 2 | (8.3) |
| New Jersey | 482 | 482 | (100.0) | 27 | (5.6) |
| New Mexico | 42 | 27 | (64.3) | — | — |
| New York State ² | 324 | 318 | (98.1) | 24 | (7.5) |
| New York City | 1,039 | 1,016 | (97.8) | 167 | (16.4) |
| North Carolina | 382 | 379 | (99.2) | 68 | (17.9) |
| North Dakota | 4 | 4 | (100.0) | 1 | (25.0) |
| Ohio | 219 | 218 | (99.5) | 40 | (18.3) |
| Oklahoma | 178 | 178 | (100.0) | 21 | (11.8) |
| Oregon | 106 | 106 | (100.0) | 20 | (18.9) |
| Pennsylvania | 327 | 325 | (99.4) | 17 | (5.2) |
| Rhode Island | 51 | 51 | (100.0) | 3 | (5.9) |
| South Carolina | 234 | 231 | (98.7) | 69 | (29.9) |
| South Dakota | 11 | 11 | (100.0) | 3 | (27.3) |
| Tennessee | 279 | 277 | (99.3) | 61 | (22.0) |
| Texas | 1,683 | 1,668 | (99.1) | 314 | (18.8) |
| Utah | 36 | 36 | (100.0) | 7 | (19.4) |
| Vermont | 6 | 6 | (100.0) | 1 | (16.7) |
| Virginia | 329 | 327 | (99.4) | 2 | (0.6) |
| Washington | 244 | 239 | (98.0) | 29 | (12.1) |
| West Virginia | 24 | 23 | (95.8) | 7 | (30.4) |
| Wisconsin | 95 | 93 | (97.9) | 14 | (15.1) |
| Wyoming | 5 | 5 | (100.0) | 0 | (0.0) |
| American Samoa ³ | 3 | 3 | (100.0) | 0 | (0.0) |
| Fed. States of Micronesia ³ | 8 | 7 | (87.5) | 2 | (28.6) |
| Guam ³ | 51 | 50 | (98.0) | 0 | (0.0) |
| Marshall Islands ³ | 41 | 38 | (92.7) | 1 | (2.6) |
| N. Mariana Islands ³ | 55 | 55 | (100.0) | 5 | (9.1) |
| Puerto Rico ³ | 123 | 121 | (98.4) | 35 | (28.9) |
| Republic of Palau ³ | 5 | 5 | (100.0) | 1 | (20.0) |
| U.S. Virgin Islands ³ | ... | ... | ... | ... | ... |

¹Excess alcohol use within past 12 months of TB diagnosis. Percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥ 75% of cases.

²Excludes New York City.

³Not included in U.S. totals.

Note: Ellipses indicate data not available.