

Table 28. Tuberculosis Cases and Case Rates per 100,000 Population, Ranked and Grouped by Number of Cases: States and the District of Columbia, 2003 and 2004

| State | 2004 | | 2003 | | 2003–2004 % Change | | Overall Rank by 2004 Rate |
|------------------------------|---------------|------------|---------------|------------|--------------------|-------------|---------------------------|
| | No. | Rate | No. | Rate | No. | Rate | |
| >400 cases in 2004 | | | | | | | |
| California | 2,989 | 8.3 | 3220 | 9.1 | -7.2 | -8.3 | 3 |
| Texas | 1,683 | 7.5 | 1589 | 7.2 | 5.9 | 4.1 | 4 |
| New York ¹ | 1,363 | 7.1 | 1471 | 7.7 | -7.3 | -7.4 | 5 |
| Florida | 1,076 | 6.2 | 1041 | 6.1 | 3.4 | 1.0 | 7 |
| Illinois | 569 | 4.5 | 634 | 5.0 | -10.3 | -10.7 | 19 |
| Georgia | 536 | 6.1 | 532 | 6.1 | 0.8 | -1.0 | 8 |
| New Jersey | 482 | 5.5 | 495 | 5.7 | -2.6 | -3.3 | 11 |
| 100-399 cases in 2004 | | | | | | | |
| North Carolina | 382 | 4.5 | 374 | 4.4 | 2.1 | 0.7 | 19 |
| Virginia | 329 | 4.4 | 332 | 4.5 | -0.9 | -2.2 | 21 |
| Pennsylvania | 327 | 2.6 | 336 | 2.7 | -2.7 | -3.0 | 33 |
| Maryland | 314 | 5.6 | 268 | 4.9 | 17.2 | 16.2 | 9 |
| Massachusetts | 283 | 4.4 | 261 | 4.1 | 8.4 | 8.5 | 21 |
| Tennessee | 279 | 4.7 | 286 | 4.9 | -2.4 | -3.4 | 15 |
| Michigan | 273 | 2.7 | 243 | 2.4 | 12.3 | 12.0 | 32 |
| Arizona | 272 | 4.7 | 295 | 5.3 | -7.8 | -10.4 | 15 |
| Louisiana | 249 | 5.5 | 259 | 5.8 | -3.9 | -0.3 | 11 |
| Washington | 244 | 3.9 | 250 | 4.1 | -2.4 | -3.5 | 25 |
| South Carolina | 234 | 5.6 | 254 | 6.1 | -7.9 | -9.0 | 9 |
| Ohio | 219 | 1.9 | 229 | 2.0 | -4.4 | -4.5 | 39 |
| Alabama | 211 | 4.7 | 258 | 5.7 | -18.2 | -18.7 | 15 |
| Minnesota | 199 | 3.9 | 214 | 4.2 | -7.0 | -7.7 | 25 |
| Oklahoma | 178 | 5.1 | 163 | 4.6 | 9.2 | 8.7 | 13 |
| Arkansas | 132 | 4.8 | 128 | 4.7 | 3.1 | 2.2 | 14 |
| Indiana | 128 | 2.1 | 143 | 2.3 | -10.5 | -11.0 | 38 |
| Kentucky | 127 | 3.1 | 138 | 3.4 | -8.0 | -8.6 | 28 |
| Colorado | 127 | 2.8 | 111 | 2.4 | 14.4 | 13.1 | 31 |
| Missouri | 127 | 2.2 | 131 | 2.3 | -3.1 | -3.7 | 35 |
| Mississippi | 119 | 4.1 | 128 | 4.4 | -7.0 | -7.7 | 23 |
| Hawaii | 116 | 9.2 | 116 | 9.3 | 0.0 | -1.1 | 2 |
| Oregon | 106 | 2.9 | 106 | 3.0 | 0.0 | -0.8 | 29 |
| Connecticut | 101 | 2.9 | 111 | 3.2 | -9.0 | -9.4 | 29 |
| <100 cases in 2004 | | | | | | | |
| Nevada | 95 | 4.1 | 106 | 4.7 | -10.4 | -13.9 | 23 |
| Wisconsin | 95 | 1.7 | 66 | 1.2 | 43.9 | 43.0 | 41 |
| District of Columbia | 81 | 14.6 | 79 | 14.2 | 2.5 | 3.3 | 1 |
| Kansas | 62 | 2.3 | 75 | 2.8 | -17.3 | -17.7 | 34 |
| Rhode Island | 51 | 4.7 | 46 | 4.3 | 10.9 | 10.4 | 15 |
| Iowa | 47 | 1.6 | 40 | 1.4 | 17.5 | 17.0 | 42 |
| Alaska | 43 | 6.6 | 57 | 8.8 | -24.6 | -25.4 | 6 |
| New Mexico | 42 | 2.2 | 49 | 2.6 | -14.3 | -15.4 | 35 |
| Nebraska | 39 | 2.2 | 28 | 1.6 | 39.3 | 38.5 | 35 |
| Utah | 36 | 1.5 | 39 | 1.7 | -7.7 | -9.1 | 44 |
| Delaware | 32 | 3.9 | 32 | 3.9 | 0.0 | -1.5 | 25 |
| New Hampshire | 24 | 1.8 | 15 | 1.2 | 60.0 | 58.7 | 40 |
| West Virginia | 24 | 1.3 | 21 | 1.2 | 14.3 | 14.0 | 47 |
| Maine | 20 | 1.5 | 24 | 1.8 | -16.7 | -17.2 | 44 |
| Montana | 15 | 1.6 | 7 | 0.8 | 114.3 | 112.3 | 42 |
| South Dakota | 11 | 1.4 | 20 | 2.6 | -45.0 | -45.4 | 46 |
| Idaho | 11 | 0.8 | 13 | 1.0 | -15.4 | -17.0 | 50 |
| Vermont | 6 | 1.0 | 9 | 1.5 | -33.3 | -33.6 | 48 |
| Wyoming | 5 | 1.0 | 4 | 0.8 | 25.0 | 23.9 | 48 |
| North Dakota | 4 | 0.6 | 6 | 0.9 | -33.3 | -33.4 | 51 |
| Total | 14,517 | 4.9 | 14,852 | 5.1 | -2.3 | -3.2 | |

¹Includes New York City.

Note: Denominators for computing 2003 and 2004 rates for states and the District of Columbia were obtained from Annual Estimates of the Population for the United States and States, and for Puerto Rico: April 1, 2000–July 1, 2004 (NST-EST2004-01) (<http://www.census.gov/popest/states/tables/NST-EST2004-01.pdf>).

See Table 20, page 37, for ranking of states without the District of Columbia.