

TABLE 2. Percentage of students in high school (grades 9-12) who were current users* of any tobacco product, by product type, sex, and race/ethnicity -- National Youth Tobacco Survey, United States, 2002 and 2004

| Characteristic | Any tobacco† | | Cigarettes | | Cigars | | Smokeless tobacco | | Pipes | | Bidis | | Kreteks | |
|--------------------------|--------------|---------------|-------------|---------------|-------------|---------------|-------------------|---------------|------------|---------------|------------|---------------|------------|---------------|
| | % | (95% CI§) | % | (95% CI§) | % | (95% CI§) | % | (95% CI§) | % | (95% CI§) | % | (95% CI§) | % | (95% CI§) |
| High school, 2004 | | | | | | | | | | | | | | |
| Sex | | | | | | | | | | | | | | |
| Male | 30.7 | (±2.7) | 21.6 | (±2.4) | 18.4 | (±1.8) | 9.9 | (±1.9) | 4.8 | (±0.8) | 3.7 | (±0.7) | 3.4 | (±0.7) |
| Female | 24.1 | (±2.5) | 21.8 | (±2.5) | 7.6 | (±1.1) | 1.2 | (±0.4) | 1.5 | (±0.5) | 1.6 | (±0.4) | 1.6 | (±0.5) |
| Race/Ethnicity | | | | | | | | | | | | | | |
| White, non-Hispanic | 30.8 | (±3.2) | 24.8 | (±3.0) | 13.7 | (±1.7) | 6.9 | (±1.4) | 3.1 | (±0.7) | 2.4 | (±0.5) | 2.5 | (±0.6) |
| Black, non-Hispanic | 16.8¶ | (±2.5) | 10.9 | (±2.5) | 10.0 | (±1.6) | 1.4 | (±0.7) | 1.7¶ | (±0.7) | 2.1 | (±0.7) | 1.3 | (±0.6) |
| Hispanic | | (±2.3) | 20.5 | (±2.3) | 13.5¶ | (±1.6) | 3.2 | (±0.8) | 4.8 | (±0.9) | 4.8¶ | (±0.8) | 3.6 | (±0.8) |
| Asian | 13.3 | (±3.3) | 11.3 | (±2.6) | 6.2 | (±2.5) | 2.0 | (±1.3) | 2.1 | (±1.1) | 2.1 | (±1.1) | 1.3 | (±0.9) |
| Total | 27.4 | (±2.4) | 21.7 | (±2.2) | 12.9 | (±1.3) | 5.5 | (±1.1) | 3.2 | (±0.5) | 2.7 | (±0.4) | 2.5 | (±0.4) |
| High school, 2002 | | | | | | | | | | | | | | |
| Sex | | | | | | | | | | | | | | |
| Male | 25.7 | | 32.6 | (±2.3) | 23.9 | (±2.1) | 16.9 | (±1.4) | 10.5 | (±2.0) | 5.0 | (±0.9) | 3.7 | (±0.8) |
| Female | | | 23.7 | (±1.8) | 21.0 | (±1.9) | 6.2 | (±0.9) | 1.2 | (±0.3) | 1.4 | (±0.4) | 1.5 | (±0.4) |
| Race/Ethnicity | | | | | | | | | | | | | | |
| White, non-Hispanic | | | 30.9 | (±2.0) | 25.2 | (±1.8) | 11.8 | (±1.0) | 7.3 | (±1.4) | 2.8 | (±0.6) | 2.2 | (±0.5) |
| Black, non-Hispanic | | | 21.7 | (±2.9) | 13.8 | (±2.8) | 12.0 | (±1.9) | 1.8 | (±0.8) | 3.7 | (±1.2) | 3.4 | (±1.1) |
| Hispanic | | | | (±2.7) | 19.8 | (±2.5) | 10.8 | (±1.5) | 3.3 | (±1.1) | 4.6 | (±1.1) | 3.5 | (±0.9) |
| Asian | | | 14.6 | (±3.8) | 12.2 | (±3.4) | 5.4 | (±2.3) | 2.1 | (±1.5) | 2.7 | (±1.5) | 2.9 | (±1.6) |
| Total | | | 28.2 | (±1.7) | 22.5 | (±1.6) | 11.6 | (±0.9) | 5.9 | (±1.1) | 3.2 | (±0.6) | 2.6 | (±0.5) |

* Used tobacco on at least 1 day during the 30 days preceding the survey.

† Cigarettes, cigars, smokeless tobacco, pipes, bidis (leaf-wrapped, flavored cigarettes from India), or kreteks (clove cigarettes).

§ Confidence interval.

¶ Significant difference ($p < 0.05$), 2004 versus 2002.