

All pneumoconioses: Death rates (per million population) by race and sex, U.S. residents age 15 and over, 1995–2004

Year	Overall	White		Black		Other	
		Male	Female	Male	Female	Male	Female
Crude Death Rate							
1995	15.37	35.03	0.64	14.17	0.15	1.96	-
1996	15.01	33.76	0.91	14.26	0.15	3.02	0.21
1997	13.96	31.74	0.78	11.41	0.52	2.04	-
1998	13.15	30.13	0.68	10.36	0.37	1.96	0.79
1999	12.81	29.21	0.65	11.19	0.14	3.61	-
2000	12.90	29.30	0.83	11.34	0.21	2.89	-
2001	12.22	27.73	0.73	11.38	0.27	2.44	-
2002	11.94	27.12	0.85	10.24	0.34	1.68	0.16
2003	11.47	25.92	0.89	9.69	0.47	2.75	-
2004	10.87	24.54	0.75	10.94	0.20	1.41	0.15
1995–2004	12.72	28.94	0.76	11.25	0.28	2.20	0.12
Age-Adjusted Death Rate							
1995	15.80	42.82	0.53	27.72	0.21	3.67	-
1996	15.33	41.15	0.75	26.01	0.18	6.32	0.42
1997	14.22	38.57	0.64	21.18	0.70	3.82	-
1998	13.27	35.68	0.55	19.28	0.43	3.81	1.25
1999	12.88	34.41	0.52	20.91	0.16	6.27	-
2000	13.14	35.77	0.68	22.43	0.24	6.18	-
2001	12.40	33.09	0.59	22.27	0.30	5.34	-
2002	12.07	32.04	0.70	20.36	0.41	3.60	0.17
2003	11.53	30.38	0.73	18.38	0.55	5.03	-
2004	10.88	28.54	0.59	21.15	0.20	2.48	0.25
1995–2004	12.95	34.80	0.62	22.05	0.34	4.50	0.20

- indicates no deaths listed.

NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Census Bureau.