

**NORA healthcare and social assistance sector and all pneumoconioses: Most frequently recorded industries and occupations on death certificate, U.S. residents age 15 and over, selected states and years, 1990–1999**

Bureau of Census Industry/Occupation		Number of Deaths*	Percent
<b>CIC</b>	<b>Industry</b>		
831	Hospitals	35	67.3
840	Health services, n.e.c.	5	9.6
832	Nursing and personal care facilities	4	7.7
870	Residential care facilities, without nursing	3	5.8
812	Offices and clinics of physicians	2	3.8
871	Social services, n.e.c.	2	3.8
	All other industries within this sector	1	1.9
	<b>TOTAL</b>	<b>52</b>	<b>100.0</b>
<b>COC</b>	<b>Occupation</b>		
453	Janitors and cleaners	11	21.2
447	Nursing aides, orderlies, and attendants	3	5.8
678	Dental laboratory and medical appliance technicians	3	5.8
084	Physicians	2	3.8
095	Registered nurses	2	3.8
203	Clinical laboratory technologists and technicians	2	3.8
448	Supervisors, cleaning and building service workers	2	3.8
449	Maids and housemen	2	3.8
575	Electricians	2	3.8
585	Plumbers, pipefitters, and steamfitters	2	3.8
696	Stationary engineers	2	3.8
	All other occupations within this sector	19	36.5
	Occupation not reported within this sector	0	0.0
	<b>TOTAL</b>	<b>52</b>	<b>100.0</b>

NORA - National Occupational Research Agenda    CIC - Census Industry Code    COC - Census Occupation Code    n.e.c. - not elsewhere classified

\* The total number of all pneumoconioses deaths was 10,014 in these same selected states and years, and the comparable number of all pneumoconioses deaths in the entire United States for this same time period was 31,455.

NOTE: Percentages may not total to 100% due to rounding. See appendices for source description, methods, ICD codes, industry and occupation codes, industrial sector groupings, and list of selected states and years.

SOURCE: National Center for Health Statistics multiple cause-of-death data.