TABLE 1
Prevalence estimates of selected sociodemographic characteristics, access to health care, and selected health indicators among American Indian/Alaska Native and non-Hispanic White adults residing in CHSDA counties—Behavioral Risk Factor Surveillance System, 2000–2006<sup>a</sup>

	NHW Total <sup>a</sup>		AI/AN Total <sup>a</sup>		Northern Plains <sup>b</sup>		Alaska <sup>c</sup>		Southern Plains <sup>d</sup>		Pacific Coast <sup>e</sup>		East		Southwest <sup>g</sup>	
	% h	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Age group																
18–49	56.2	(55.9-56.4)	71.1	(69.3-72.9)	74.3	(71.5-76.9)	72.1	(69.7-74.4)	67.9	(65.7–70.0)	69.3	(64.5-73.7)	66.9	(61.6-71.8)	77.3	(75.2-79.4)
50+	43.8	(43.6-44.1)	28.9	(27.1-30.7)	25.7	(23.1-28.5)	27.9	(25.6-30.3)	32.1	(30.0-34.3)	30.7	(26.3-35.5)	33.1	(28.2-38.4)	22.7	(20.6-24.8)
Gender																
Male	48.7	(48.4-49.0)	51.1	(49.1-53.1)	47.3	(43.9-50.7)	47.6	(45.0-50.2)	49.0	(46.7-51.4)	55.3	(50.5-60.1)	55.4	(50.5-60.2)	46.1	(43.3-48.9)
Female	51.3	(51.0-51.6)	48.9	(46.9-50.9)	52.7	(49.3-56.1)	52.4	(49.8-55.0)	51.0	(48.6-53.3)	44.7	(39.9-49.5)	44.6	(39.8-49.5)	53.9	(51.1-56.7)
Marital status																
Married	61.1	(60.8-61.4)	48.9	(47.0-50.8)	43.9	(40.7-47.1)	47.4	(44.8-50.1)	55.2	(53.0-57.4)	49.7	(45.2-54.2)	46.3	(41.6-51.2)	48.8	(46.1-51.4)
Divorced, widowed, separated	17.8	(17.6-18.0)	25.0	(23.3-26.7)	25.3	(22.8-27.9)	21.9	(19.8-24.2)	25.6	(23.7–27.5)	25.4	(21.5-29.8)	30.4	(26.7-34.4)	20.0	(17.9-22.2)
Never married or a member of an unmarried couple	21.1	(20.9–21.4)	26.1	(24.7–27.5)	30.8	(28.0-33.9)	30.6	(28.7–32.6)	19.2	(17.6–21.0)	24.9	(21.6–28.4)	23.3	(19.5–27.5)	31.3	(29.3–33.3)
Education																
< High school	7.2	(7.1-7.4)	23.1	(21.3-24.9)	26.7	(23.9-29.8)	26.8	(24.5-29.2)	17.5	(15.6-19.6)	25.4	(21.2-30.1)	24.9	(20.7-29.6)	20.4	(18.3-22.8)
High school	28.8	(28.6–29.1)	36.3	(34.3-38.3)	34.4	(31.2-37.9)	43.6	(41.3–46.0)	36.7	(34.5-39.0)	36.6	(31.8-41.6)	34.5	(30.0-39.3)	35.1	(32.4–37.8)
Some college or technical school	30.8	(30.5-31.1)	27.7	(26.0-29.4)	25.5	(23.0-28.2)	20.4	(18.6–22.5)	29.3	(27.2-31.5)	26.8	(22.9-31.1)	28.7	(24.3-33.6)	30.0	(27.5-32.6)
College graduate	33.1	(32.9-33.4)	12.9	(11.8-14.1)	13.3	(11.5-15.4)	9.1	(7.3-11.2)	16.4	(14.9-18.1)	11.2	(8.9-14.1)	11.9	(9.3-15.2)	14.5	(12.5-16.8)
Income																
<\$15,000	7.1	(6.9-7.3)	16.8	(15.5-18.3)	16.9	(14.9-19.0)	17.3	(15.6–19.2)	14.9	(13.2-16.7)	18.6	(15.2-22.5)	15.4	(12.4-19.0)	15.3	(13.5-17.3)
\$15,000-\$34,999	25.0	(24.8–25.3)	40.3	(38.3-42.3)	40.5	(37.4-43.8)	35.5	(33.0-38.2)	43.6	(41.3-45.9)	36.6	(31.8-41.8)	44.5	(39.5-49.7)	41.5	(38.6-44.4)
\$35,000-\$74,999	34.5	(34.2-34.8)	24.8	(23.1-26.5)	22.3	(19.7-25.0)	21.0	(18.6-23.5)	23.3	(21.5-25.2)	28.0	(24.0-32.4)	23.6	(19.5-28.3)	24.9	(22.4–27.6)
≥ \$75K	22.0	(21.7–22.2)	6.9	(6.1–7.8)	6.3	(4.9-8.2)	10.0	(8.5–11.7)	7.4	(6.1–9.1)	7.4	(5.5–10.0)	5.9	(4.3-8.0)	6.4	(5.2–7.7)
Unknown	11.4	(11.2–11.6)	11.3	(10.0-12.7)	14.0	(11.0-17.6)	16.2	(14.5-18.0)	10.8	(9.5-12.2)	9.4	(6.8-12.8)	10.6	(8.2-13.5)	11.9	(10.1-14.1)
Employment																
Employed	63.4	(63.2-63.7)	55.0	(53.3-56.7)	51.8	(48.5-55.1)	54.1	(51.4-56.7)	57.4	(55.1-59.5)	53.5	(49.3-57.7)	58.3	(54.1-62.4)	57.6	(55.1-60.1)
Unemployed	8.1	(7.9–8.3)	18.1	(16.5–19.8)	21.2	(18.0-24.8)	24.3	(22.1–26.7)	15.3	(13.7–17.0)	18.5	(14.8–22.8)	17.3	(14.2-21.0)	15.4	(13.6–17.4)
Homemaker, student, retired	28.5	(28.2-28.7)	26.9	(25.3-28.6)	27.0	(24.6-29.6)	21.7	(19.6-23.9)	27.4	(25.6-29.2)	28.0	(24.2-32.2)	24.4	(21.1-27.9)	27.0	(25.0-29.0)
Health care																
Health care coverage	87.9	(87.7-88.1)	76.6	(75.1–78.1)	75.3	(72.6-77.7)	80.0	(77.9-81.8)	74.5	(72.3-76.7)	82.9	(78.9-86.3)	74.9	(70.5–78.8)	68.4	(66.0-70.8)
Have personal provider	81.2	(81.0-81.5)	70.0	(67.9–71.9)	71.9	(68.5–75.0)	66.6	(64.2–69.0)	76.3	(74.1–78.3)	70.3	(65.2–74.9)	72.2	(67.1–76.8)	62.0	(59.0-65.0)
Health																
Health status (good or better)	87.9	(87.7-88.0)	74.1	(72.2-75.9)	75.9	(73.2-78.4)	78.0	(75.7-80.1)	76.2	(74.2-78.1)	70.7	(66.2-74.8)	76.1	(71.8-80.0)	75.7	(73.0-78.2)
Ever told have diabetes <sup>j</sup>	5.7	(5.6–5.9)	14.4	(13.0–15.9)	17.0	(14.7–19.4)	5.7	(4.5–7.2)	14.3	(12.9–15.8)	14.9	(11.9–18.6)	11.4	(8.8–14.6)	13.3	(11.7–15.1)

CHSDA indicates Contract Health Service Delivery Areas of the Indian Health Service; NHW, non-Hispanic whites; Al/AN, American Indians/Alaska Natives; 95% CI, 95% confidence interval.

<sup>&</sup>lt;sup>j</sup>Does not include those who had diabetes during pregnancy.





<sup>&</sup>lt;sup>a</sup>Data from 34 US states (includes states listed in the footnotes below).

<sup>&</sup>lt;sup>b</sup>AI/AN in Indiana, Iowa, Michigan, Minnesota, Montana, Nebraska, North Dakota, South Dakota, Wisconsin, and Wyoming.

cAI/AN in Alaska.

<sup>&</sup>lt;sup>d</sup>AI/AN in Kansas, Oklahoma, and Texas.

<sup>&</sup>lt;sup>e</sup>AI/AN in California, Idaho, Oregon and Washington.

fAI/AN in Alabama, Connecticut, Florida, Louisiana, Maine, Massachusetts, Mississippi, New York, North Carolina, Rhode Island, and South Carolina.

<sup>&</sup>lt;sup>g</sup>AI/AN in Arizona, Colorado, Nevada, New Mexico, and Utah.

hAll prevalence estimates are weighted. Except for age group, estimates are age-adjusted to the 2000 US standard population. "Refused" and "don't know" responses are excluded. i2001–2006.