

**All pneumoconioses: Years of potential life lost to age 65 and to life expectancy by race and sex, U.S. residents age 15 and over, 1995–2004**

Year	White		Black		Other		Total
	Male	Female	Male	Female	Male	Female	
<b>Years of Potential Life Lost to Age 65</b>							
1995	2,365	40	165	15	50	-	2,635
1996	1,815	115	140	-	55	-	2,125
1997	1,810	105	130	-	-	-	2,045
1998	1,665	60	170	30	10	5	1,940
1999	1,340	60	260	25	30	-	1,715
2000	1,745	80	185	25	25	-	2,060
2001	1,610	145	220	35	30	-	2,040
2002	1,535	240	150	10	5	5	1,945
2003	1,735	195	175	55	55	-	2,215
2004	1,790	45	210	25	30	-	2,100
<b>TOTAL</b>	<b>17,410</b>	<b>1,085</b>	<b>1,805</b>	<b>220</b>	<b>290</b>	<b>10</b>	<b>20,820</b>
<b>Years of Potential Life Lost to Life Expectancy</b>							
1995	28,815	684	1,440	37	152	-	31,128
1996	27,385	1,004	1,566	28	195	8	30,186
1997	25,884	865	1,305	63	111	-	28,228
1998	24,977	746	1,213	95	112	51	27,194
1999	23,921	673	1,397	51	243	-	26,285
2000	24,784	960	1,405	58	198	-	27,405
2001	24,408	899	1,481	87	181	-	27,056
2002	24,048	1,155	1,294	66	105	22	26,690
2003	23,838	1,171	1,373	142	254	-	26,778
2004	23,127	813	1,582	75	146	7	25,750
<b>TOTAL</b>	<b>251,187</b>	<b>8,970</b>	<b>14,056</b>	<b>702</b>	<b>1,697</b>	<b>88</b>	<b>276,700</b>

- indicates no deaths listed.

NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data.