

**DRAFT
TOXICOLOGICAL PROFILE FOR
STYRENE**

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
Agency for Toxic Substances and Disease Registry

September 2007

DISCLAIMER

The use of company or product name(s) is for identification only and does not imply endorsement by the Agency for Toxic Substances and Disease Registry.

This information is distributed solely for the purpose of pre dissemination public comment under applicable information quality guidelines. It has not been formally disseminated by the Agency for Toxic Substances and Disease Registry. It does not represent and should not be construed to represent any agency determination or policy.

UPDATE STATEMENT

A Toxicological Profile for styrene was released in 1992. This present edition supersedes any previously released draft or final profile.

Toxicological profiles are revised and republished as necessary. For information regarding the update status of previously released profiles, contact ATSDR at:

Agency for Toxic Substances and Disease Registry
Division of Toxicology and Environmental Medicine/Applied Toxicology Branch
1600 Clifton Road NE
Mailstop F-32
Atlanta, Georgia 30333

This page is intentionally blank.

FOREWORD

This toxicological profile is prepared in accordance with guidelines developed by the Agency for Toxic Substances and Disease Registry (ATSDR) and the Environmental Protection Agency (EPA). The original guidelines were published in the *Federal Register* on April 17, 1987. Each profile will be revised and republished as necessary.

The ATSDR toxicological profile succinctly characterizes the toxicologic and adverse health effects information for the hazardous substance described therein. Each peer-reviewed profile identifies and reviews the key literature that describes a hazardous substance's toxicologic properties. Other pertinent literature is also presented, but is described in less detail than the key studies. The profile is not intended to be an exhaustive document; however, more comprehensive sources of specialty information are referenced.

The focus of the profiles is on health and toxicologic information; therefore, each toxicological profile begins with a public health statement that describes, in nontechnical language, a substance's relevant toxicological properties. Following the public health statement is information concerning levels of significant human exposure and, where known, significant health effects. The adequacy of information to determine a substance's health effects is described in a health effects summary. Data needs that are of significance to protection of public health are identified by ATSDR and EPA.

Each profile includes the following:

- (A) The examination, summary, and interpretation of available toxicologic information and epidemiologic evaluations on a hazardous substance to ascertain the levels of significant human exposure for the substance and the associated acute, subacute, and chronic health effects;
- (B) A determination of whether adequate information on the health effects of each substance is available or in the process of development to determine levels of exposure that present a significant risk to human health of acute, subacute, and chronic health effects; and
- (C) Where appropriate, identification of toxicologic testing needed to identify the types or levels of exposure that may present significant risk of adverse health effects in humans.

The principal audiences for the toxicological profiles are health professionals at the Federal, State, and local levels; interested private sector organizations and groups; and members of the public. We plan to revise these documents in response to public comments and as additional data become available. Therefore, we encourage comments that will make the toxicological profile series of the greatest use.

Comments should be sent to:

Agency for Toxic Substances and Disease Registry
Division of Toxicology and Environmental Medicine
1600 Clifton Road, N.E.
Mail Stop F-32
Atlanta, Georgia 30333


The toxicological profiles are developed in response to the Superfund Amendments and Reauthorization Act (SARA) of 1986 (Public Law 99-499) which amended the Comprehensive Environmental Response, Compensation, and Liability Act of 1980 (CERCLA or Superfund). This public law directed ATSDR to prepare toxicological profiles for hazardous substances most commonly found at facilities on the CERCLA National Priorities List and that pose the most significant potential threat to human health, as determined by ATSDR and the EPA. The availability of the revised priority list of 275 hazardous substances was announced in the *Federal Register* on December 7, 2005 (70 FR 72840). For prior versions of the list of substances, see *Federal Register* notices dated April 17, 1987 (52 FR 12866); October 20, 1988 (53 FR 41280); October 26, 1989 (54 FR 43619); October 17, 1990 (55 FR 42067); October 17, 1991 (56 FR 52166); October 28, 1992 (57 FR 48801); February 28, 1994 (59 FR 9486); April 29, 1996 (61 FR 18744); November 17, 1997 (62 FR 61332); October 21, 1999 (64 FR 56792); October 25, 2001 (66 FR 54014); and November 7, 2003 (68 FR 63098). Section 104(i)(3) of CERCLA, as amended, directs the Administrator of ATSDR to prepare a toxicological profile for each substance on the list.

This profile reflects ATSDR's assessment of all relevant toxicologic testing and information that has been peer-reviewed. Staff of the Centers for Disease Control and Prevention and other Federal scientists have also reviewed the profile. In addition, this profile has been peer-reviewed by a nongovernmental panel and is being made available for public review. Final responsibility for the contents and views expressed in this toxicological profile resides with ATSDR.



Howard Frumkin, M.D., Dr. P.H.
Director

National Center for Environmental Health/
Agency for Toxic Substances and
Disease Registry



Julie Louise Gerberding, M.D., M.P.H.
Administrator

Agency for Toxic Substances and
Disease Registry

Managing Hazardous Materials Incidents is a three-volume set of recommendations for on-scene (prehospital) and hospital medical management of patients exposed during a hazardous materials incident. Volumes I and II are planning guides to assist first responders and hospital emergency department personnel in planning for incidents that involve hazardous materials. Volume III—*Medical Management Guidelines for Acute Chemical Exposures*—is a guide for health care professionals treating patients exposed to hazardous materials.

Fact Sheets (ToxFAQs) provide answers to frequently asked questions about toxic substances.

Other Agencies and Organizations

The National Center for Environmental Health (NCEH) focuses on preventing or controlling disease, injury, and disability related to the interactions between people and their environment outside the workplace. Contact: NCEH, Mailstop F-29, 4770 Buford Highway, NE, Atlanta, GA 30341-3724 • Phone: 770-488-7000 • FAX: 770-488-7015.

The National Institute for Occupational Safety and Health (NIOSH) conducts research on occupational diseases and injuries, responds to requests for assistance by investigating problems of health and safety in the workplace, recommends standards to the Occupational Safety and Health Administration (OSHA) and the Mine Safety and Health Administration (MSHA), and trains professionals in occupational safety and health. Contact: NIOSH, 200 Independence Avenue, SW, Washington, DC 20201 • Phone: 800-356-4674 or NIOSH Technical Information Branch, Robert A. Taft Laboratory, Mailstop C-19, 4676 Columbia Parkway, Cincinnati, OH 45226-1998 • Phone: 800-35-NIOSH.

The National Institute of Environmental Health Sciences (NIEHS) is the principal federal agency for biomedical research on the effects of chemical, physical, and biologic environmental agents on human health and well-being. Contact: NIEHS, PO Box 12233, 104 T.W. Alexander Drive, Research Triangle Park, NC 27709 • Phone: 919-541-3212.

Referrals

The Association of Occupational and Environmental Clinics (AOEC) has developed a network of clinics in the United States to provide expertise in occupational and environmental issues. Contact: AOEC, 1010 Vermont Avenue, NW, #513, Washington, DC 20005 • Phone: 202-347-4976 • FAX: 202-347-4950 • e-mail: AOEC@AOEC.ORG • Web Page: <http://www.aoec.org/>.

The American College of Occupational and Environmental Medicine (ACOEM) is an association of physicians and other health care providers specializing in the field of occupational and environmental medicine. Contact: ACOEM, 25 Northwest Point Boulevard, Suite 700, Elk Grove Village, IL 60007-1030 • Phone: 847-818-1800 • FAX: 847-818-9266.

CONTRIBUTORS

CHEMICAL MANAGER(S)/AUTHOR(S):

Zemoria Rosemond, B.A.
Selene Chou, Ph.D.
Jewell Wilson, Ph.D.
Michael Schwartz, M.D.
Francisco Tomei-Torres, Ph.D.
ATSDR, Division of Toxicology and Environmental Medicine, Atlanta, GA

Lisa Ingerman, Ph.D., DABT
David B. Knaebel, Ph.D.
Mario Citra, Ph.D.
Syracuse Research Corporation, North Syracuse, NY

THE PROFILE HAS UNDERGONE THE FOLLOWING ATSDR INTERNAL REVIEWS:

1. Health Effects Review. The Health Effects Review Committee examines the health effects chapter of each profile for consistency and accuracy in interpreting health effects and classifying end points.
2. Minimal Risk Level Review. The Minimal Risk Level Workgroup considers issues relevant to substance-specific Minimal Risk Levels (MRLs), reviews the health effects database of each profile, and makes recommendations for derivation of MRLs.
3. Data Needs Review. The Applied Toxicology Branch reviews data needs sections to assure consistency across profiles and adherence to instructions in the Guidance.
4. Green Border Review. Green Border review assures the consistency with ATSDR policy.

This page is intentionally blank.

PEER REVIEW

A peer review panel was assembled for styrene. The panel consisted of the following members:

1. George Cruzan, Ph.D., DABT, ToxWorks, Bridgeton, New Jersey;
2. Teresa Leavens, Ph.D., The Hamner Institutes for Health Sciences, Research Triangle Park, North Carolina; and
3. Jean Rabovsky, Ph.D., Retired Toxicologist, El Cerrito, California.

These experts collectively have knowledge of styrene's physical and chemical properties, toxicokinetics, key health end points, mechanisms of action, human and animal exposure, and quantification of risk to humans. All reviewers were selected in conformity with the conditions for peer review specified in Section 104(I)(13) of the Comprehensive Environmental Response, Compensation, and Liability Act, as amended.

Scientists from the Agency for Toxic Substances and Disease Registry (ATSDR) have reviewed the peer reviewers' comments and determined which comments will be included in the profile. A listing of the peer reviewers' comments not incorporated in the profile, with a brief explanation of the rationale for their exclusion, exists as part of the administrative record for this compound.

The citation of the peer review panel should not be understood to imply its approval of the profile's final content. The responsibility for the content of this profile lies with the ATSDR.

This page is intentionally blank.

CONTENTS

DISCLAIMER	ii
UPDATE STATEMENT	iii
FOREWORD	v
QUICK REFERENCE FOR HEALTH CARE PROVIDERS.....	vii
CONTRIBUTORS	ix
PEER REVIEW	xi
CONTENTS.....	xiii
LIST OF FIGURES	xvii
LIST OF TABLES	xix
1. PUBLIC HEALTH STATEMENT.....	1
2. RELEVANCE TO PUBLIC HEALTH	9
2.1 BACKGROUND AND ENVIRONMENTAL EXPOSURES TO STYRENE IN THE UNITED STATES.....	9
2.2 SUMMARY OF HEALTH EFFECTS.....	9
2.3 MINIMAL RISK LEVELS (MRLs)	12
3. HEALTH EFFECTS.....	23
3.1 INTRODUCTION.....	23
3.2 DISCUSSION OF HEALTH EFFECTS BY ROUTE OF EXPOSURE	23
3.2.1 Inhalation Exposure	24
3.2.1.1 Death	24
3.2.1.2 Systemic Effects.....	25
3.2.1.3 Immunological and Lymphoreticular Effects.....	51
3.2.1.4 Neurological Effects.....	52
3.2.1.5 Reproductive Effects	66
3.2.1.6 Developmental Effects	68
3.2.1.7 Cancer.....	69
3.2.2 Oral Exposure.....	75
3.2.2.1 Death	75
3.2.2.2 Systemic Effects.....	76
3.2.2.3 Immunological and Lymphoreticular Effects.....	85
3.2.2.4 Neurological Effects.....	86
3.2.2.5 Reproductive Effects	86
3.2.2.6 Developmental Effects	87
3.2.2.7 Cancer.....	87
3.2.3 Dermal Exposure.....	88
3.2.3.1 Death	88
3.2.3.2 Systemic Effects.....	88
3.2.3.3 Immunological and Lymphoreticular Effects.....	89
3.2.3.4 Neurological Effects.....	89
3.2.3.5 Reproductive Effects	89
3.2.3.6 Developmental Effects	89
3.2.3.7 Cancer.....	89
3.3 GENOTOXICITY	89
3.4 TOXICOKINETICS	93
3.4.1 Absorption.....	93
3.4.1.1 Inhalation Exposure.....	93

3.4.1.2	Oral Exposure.....	95
3.4.1.3	Dermal Exposure.....	95
3.4.2	Distribution.....	96
3.4.2.1	Inhalation Exposure.....	96
3.4.2.2	Oral Exposure.....	97
3.4.2.3	Dermal Exposure.....	97
3.4.3	Metabolism.....	98
3.4.4	Elimination and Excretion.....	101
3.4.4.1	Inhalation Exposure.....	101
3.4.4.2	Oral Exposure.....	102
3.4.4.3	Dermal Exposure.....	103
3.4.5	Physiologically Based Pharmacokinetic (PBPK)/Pharmacodynamic (PD) Models.....	103
3.4.5.1	Summary of PBPK Models.....	106
3.4.5.2	Discussion of Model.....	106
3.5	MECHANISMS OF ACTION.....	108
3.5.1	Pharmacokinetic Mechanisms.....	108
3.5.2	Mechanisms of Toxicity.....	113
3.5.3	Animal-to-Human Extrapolations.....	114
3.6	TOXICITIES MEDIATED THROUGH THE NEUROENDOCRINE AXIS.....	115
3.7	CHILDREN'S SUSCEPTIBILITY.....	117
3.8	BIOMARKERS OF EXPOSURE AND EFFECT.....	119
3.8.1	Biomarkers Used to Identify or Quantify Exposure to Styrene.....	120
3.8.2	Biomarkers Used to Characterize Effects Caused by Styrene.....	121
3.9	INTERACTIONS WITH OTHER CHEMICALS.....	122
3.10	POPULATIONS THAT ARE UNUSUALLY SUSCEPTIBLE.....	123
3.11	METHODS FOR REDUCING TOXIC EFFECTS.....	123
3.11.1	Reducing Peak Absorption Following Exposure.....	123
3.11.2	Reducing Body Burden.....	124
3.11.3	Interfering with the Mechanism of Action for Toxic Effects.....	124
3.12	ADEQUACY OF THE DATABASE.....	125
3.12.1	Existing Information on Health Effects of Styrene.....	126
3.12.2	Identification of Data Needs.....	126
3.12.3	Ongoing Studies.....	135
4.	CHEMICAL AND PHYSICAL INFORMATION.....	137
4.1	CHEMICAL IDENTITY.....	137
4.2	PHYSICAL AND CHEMICAL PROPERTIES.....	137
5.	PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL.....	141
5.1	PRODUCTION.....	141
5.2	IMPORT/EXPORT.....	144
5.3	USE.....	144
5.4	DISPOSAL.....	145
6.	POTENTIAL FOR HUMAN EXPOSURE.....	147
6.1	OVERVIEW.....	147
6.2	RELEASES TO THE ENVIRONMENT.....	147
6.2.1	Air.....	149
6.2.2	Water.....	152
6.2.3	Soil.....	153
6.3	ENVIRONMENTAL FATE.....	153

6.3.1	Transport and Partitioning.....	153
6.3.2	Transformation and Degradation	155
6.3.2.1	Air.....	155
6.3.2.2	Water	156
6.3.2.3	Sediment and Soil.....	157
6.4	LEVELS MONITORED OR ESTIMATED IN THE ENVIRONMENT	158
6.4.1	Air	158
6.4.2	Water	158
6.4.3	Sediment and Soil	160
6.4.4	Other Environmental Media.....	161
6.5	GENERAL POPULATION AND OCCUPATIONAL EXPOSURE	163
6.6	EXPOSURES OF CHILDREN	164
6.7	POPULATIONS WITH POTENTIALLY HIGH EXPOSURES	165
6.8	ADEQUACY OF THE DATABASE.....	165
6.8.1	Identification of Data Needs	166
6.8.2	Ongoing Studies	169
7.	ANALYTICAL METHODS	171
7.1	BIOLOGICAL MATERIALS	171
7.2	ENVIRONMENTAL SAMPLES.....	172
7.3	ADEQUACY OF THE DATABASE.....	175
7.3.1	Identification of Data Needs	175
7.3.2	Ongoing Studies	180
8.	REGULATIONS AND ADVISORIES	181
9.	REFERENCES	187
10.	GLOSSARY	249
APPENDICES		
A.	ATSDR MINIMAL RISK LEVELS AND WORKSHEETS	A-1
B.	USER’S GUIDE.....	B-1
C.	ACRONYMS, ABBREVIATIONS, AND SYMBOLS.....	C-1
D.	INDEX	D-1

This page is intentionally blank.

LIST OF FIGURES

3-1. Levels of Significant Exposure to Styrene - Inhalation	42
3-2. Levels of Significant Exposure to Styrene - Oral	81
3-3. Scheme for Styrene Metabolism in Humans and Animals	99
3-4. Conceptual Representation of a Physiologically Based Pharmacokinetic (PBPK) Model for a Hypothetical Chemical Substance	105
3-5. Styrene and Styrene Oxide Models used in the Sarangapani PBPK Model	107
3-6. Existing Information on Health Effects of Styrene	127
6-1. Frequency of NPL Sites with Styrene Contamination	148

This page is intentionally blank.

LIST OF TABLES

2-1. Results of Selected Human Neurotoxicity Studies	17
3-1. Levels of Significant Exposure to Styrene - Inhalation	26
3-2. Summary of Occupational Exposure Studies Examining Neurological End Points.....	54
3-3. Levels of Significant Exposure to Styrene - Oral	77
3-4. Genotoxicity of Styrene <i>In Vivo</i>	90
3-5. Genotoxicity of Styrene <i>In Vitro</i>	94
3-6. Physiological and Flow Parameters Used in the Sarangapani PBPK Model.....	109
3-7. Respiratory Tract-Specific Physiological Parameters Used in the Sarangapani PBPK Model	110
3-8. Kinetic Parameters Used in the Sarangapani PBPK Model	111
3-9. Stereospecific Kinetic Parameters for Styrene and Styrene Oxide Metabolism in Rodents Used in the Sarangapani PBPK Model.....	112
4-1. Chemical Identity of Styrene	138
4-2. Physical and Chemical Properties of Styrene	139
5-1. Facilities that Produce, Process, or Use Styrene.....	142
6-1. Releases to the Environment from Facilities that Produce, Process, or Use Styrene	150
6-2. Styrene Concentrations in Representative Air Samples	159
6-3. Styrene Levels in Food Items	162
7-1. Analytical Methods for Determining Styrene in Biological Materials.....	173
7-2. Analytical Methods for Determining Styrene in Environmental Samples	176
8-1. Regulations and Guidelines Applicable to Styrene	182

This page is intentionally blank.

1. PUBLIC HEALTH STATEMENT

This public health statement tells you about styrene and the effects of exposure to it.

The Environmental Protection Agency (EPA) identifies the most serious hazardous waste sites in the nation. These sites are then placed on the National Priorities List (NPL) and are targeted for long-term federal clean-up activities. Styrene has been found in at least 31 of the 1,689 current or former NPL sites. Although the total number of NPL sites evaluated for this substance is not known, the possibility exists that the number of sites at which styrene is found may increase in the future as more sites are evaluated. This information is important because these sites may be sources of exposure, and exposure to this substance may harm you.

When a substance is released either from a large area, such as an industrial plant, or from a container, such as a drum or bottle, it enters the environment. Such a release does not always lead to exposure. You can be exposed to a substance only when you come in contact with it. You may be exposed by breathing, eating, or drinking the substance, or by skin contact.

If you are exposed to styrene, many factors will determine whether you will be harmed. These factors include the dose (how much), the duration (how long), and how you come in contact with it. You must also consider any other chemicals you are exposed to and your age, sex, diet, family traits, lifestyle, and state of health.

1. PUBLIC HEALTH STATEMENT

What is styrene?

Description	<p>Styrene is a colorless liquid that evaporates easily.</p> <p>In its pure form, styrene has a sweet smell. Manufactured styrene may contain aldehydes, which give it a sharp, unpleasant odor.</p>
<p>Uses</p> <p>•manufacturing</p> <p>•consumer products</p>	<p>Large amounts of styrene are produced in the United States. Small amounts are produced naturally by plants, bacteria, and fungi. Styrene is also present in combustion products such as cigarette smoke and automobile exhaust.</p> <p>Styrene is widely used to make plastics and rubber. Consumer products containing styrene include:</p> <ul style="list-style-type: none"> • packaging materials • insulation for electrical uses (i.e., wiring and appliances) • insulation for homes and other buildings • fiberglass, plastic pipes, automobile parts • drinking cups and other "food-use" items • carpet backing <p>These products mainly contain styrene linked together in long chains (polystyrene). However, most of these products also contain a small amount of unlinked styrene.</p>

For more information on the physical and chemical properties of styrene, and its production, disposal, and use, see Chapters 4 and 5.

What happens to styrene when it enters the environment?

Sources	<p>Styrene can be found in air, soil, and water after release from the manufacture, use, and disposal of styrene-based products.</p>
Break down	<p><i>Air:</i> Styrene is quickly broken down in the air, usually within 1–2 days.</p> <p><i>Water and soil:</i> Styrene evaporates from shallow soils and surface water. Styrene that remains in soil or water may be broken down by bacteria or other microorganisms.</p>

For more information on styrene in the environment, see Chapter 6.

1. PUBLIC HEALTH STATEMENT

How might I be exposed to styrene?

<p><i>Air</i></p>	<p>The primary way you can be exposed to styrene is by breathing air containing it. Releases of styrene into the air occur from:</p> <ul style="list-style-type: none"> • industries using or manufacturing styrene • automobile exhaust • cigarette smoke, and • use of photocopiers <p>Rural or suburban air generally contains lower concentrations of styrene than urban air. Indoor air often contains higher levels of styrene than outdoor air.</p> <ul style="list-style-type: none"> • 0.06–4.6 parts per billion (ppb) in outdoor air • 0.023–11.5 ppb in indoor air
<p><i>Water and soil</i></p>	<p>Styrene is occasionally detected in groundwater, drinking water, or soil samples. Drinking water containing styrene or bathing in water containing styrene may expose you to low levels of this chemical.</p>
<p><i>Workplace air</i></p>	<p>A large number of workers are potentially exposed to styrene. The highest potential exposure occurs in the reinforced-plastics industry, where workers may be exposed to high air concentrations and also have dermal exposure to liquid styrene or resins.</p> <p>Workers involved in styrene polymerization, rubber manufacturing, and styrene-polyester resin facilities and workers at photocopy centers may also be exposed to styrene.</p>
<p><i>Food</i></p>	<p>Low levels of styrene occur naturally in a variety of foods, such as fruits, vegetables, nuts, beverages, and meats. Small amounts of styrene can be transferred to food from styrene-based packaging material.</p>

For more information on human exposure to styrene, see Chapter 6.

1. PUBLIC HEALTH STATEMENT

How can styrene enter and leave my body?

<i>Enter your body</i>	<i>Inhalation:</i> When you breathe air containing styrene, most of the styrene will rapidly enter your body through your lungs. <i>Ingestion:</i> Styrene in food or water may also rapidly enter your body through the digestive tract. <i>Dermal contact:</i> A very small amount may enter through your skin when you come into contact with liquids containing styrene.
<i>Leave your body</i>	Once in your body, styrene is broken down into other chemicals. Most of these other chemicals leave your body in the urine within few days.

For more information on how styrene enters and leaves the body, see Chapter 3.

How can styrene affect my health?

Scientists use many tests to protect the public from harmful effects of toxic chemicals and to find ways for treating persons who have been harmed.

1. PUBLIC HEALTH STATEMENT

Health Effects	
Workers • <i>Inhalation</i>	<p>The most common health problems in workers exposed to styrene involve the nervous system. These health effects include changes in color vision, tiredness, feeling drunk, slowed reaction time, concentration problems, and balance problems.</p> <p>The styrene concentrations that cause these effects are more than 1,000 times higher than the levels normally found in the environment.</p>
Laboratory animals • <i>Inhalation</i>	<p>Hearing loss has been observed in animals exposed to very high concentrations of styrene.</p> <p>Animal studies have shown that inhalation of styrene can result in changes in the lining of the nose and damage to the liver. However, animals may be more sensitive than humans to the nose and liver effects.</p>
Laboratory animals • <i>Oral</i>	<p>Impaired learning has been observed in rats exposed to high doses of styrene.</p> <p>Sperm damage has also been observed in rats exposed to high doses of styrene.</p>
Cancer	<p>The International Agency for Research on Cancer has determined that styrene is a possible carcinogen.</p>

Further information on the health effects of styrene in humans and animals can be found in Chapters 2 and 3.

How can styrene affect children?

This section discusses potential health effects in humans from exposures during the period from conception to maturity at 18 years of age.

Effects in children	<p>There are no studies evaluating the effect of styrene exposure on children or immature animals. It is likely that children would have the same health effects as adults. We do not know whether children would be more sensitive than adults to the effects of styrene.</p>
Birth defects	<p>Studies in workers have examined whether styrene can cause birth defects or low birth weight; however, the results are inconclusive. No birth defects were observed in animal studies.</p>
Breast milk	<p>Nursing infants can be exposed to styrene from breast milk.</p>

1. PUBLIC HEALTH STATEMENT

How can families reduce the risk of exposure to styrene?

<i>Tobacco smoke</i>	Styrene is a component of tobacco smoke. Avoid smoking in enclosed spaces like inside the home or car in order to limit exposure to children and other family members.
<i>Copier</i>	Styrene is released during the use of home copiers. Families should use a copier only when needed and turn it off when finished. It is also important to keep the room with the copier well ventilated.

Is there a medical test to determine whether I have been exposed to styrene?

<i>Detecting exposure</i>	Styrene can be measured in blood, urine, and body tissues for a short time following exposure to moderate-to-high levels.
<i>Measuring exposure</i>	<p>The presence of styrene breakdown products (metabolites) in urine might indicate that you were exposed to styrene; however, these metabolites can also form when you are exposed to other substances.</p> <p>Measuring styrene metabolites in urine within 1 day of exposure allows medical personnel to estimate actual exposure level.</p> <p>The detection of these metabolites in your urine cannot be used to predict the kind of health effects that might develop from that exposure.</p>

Information about tests for detecting styrene in the body is given in Chapters 3 and 7.

What recommendations has the federal government made to protect human health?

The federal government develops regulations and recommendations to protect public health. Regulations *can* be enforced by law. The EPA, the Occupational Safety and Health Administration (OSHA), and the Food and Drug Administration (FDA) are some federal agencies that develop regulations for toxic substances. Recommendations provide valuable guidelines to protect public health, but *cannot* be enforced by law. The Agency for Toxic Substances and Disease Registry (ATSDR) and the National Institute for Occupational Safety and Health (NIOSH) are two federal organizations that develop recommendations for toxic substances.

Regulations and recommendations can be expressed as “not-to-exceed” levels. These are levels of a toxic substance in air, water, soil, or food that do not exceed a critical value. This critical value is usually based

1. PUBLIC HEALTH STATEMENT

on levels that affect animals; they are then adjusted to levels that will help protect humans. Sometimes these not-to-exceed levels differ among federal organizations because they used different exposure times (an 8-hour workday or a 24-hour day), different animal studies, or other factors.

Recommendations and regulations are also updated periodically as more information becomes available. For the most current information, check with the federal agency or organization that provides it.

Some regulations and recommendations for styrene include the following:

<i>Drinking water</i>	<p>The EPA has determined that exposure to styrene in drinking water at concentrations of 20 ppm for 1 day or 2 ppm for 10 days is not expected to cause any adverse effects in a child.</p> <p>The EPA has determined that lifetime exposure to 0.1 ppm styrene in drinking water is not expected to cause any adverse effects.</p>
<i>Bottled water</i>	<p>The FDA has determined that the styrene concentration in bottled drinking water should not exceed 0.1 ppm.</p>
<i>Workplace air</i>	<p>OSHA set a legal limit of 100 ppm styrene in air averaged over an 8-hour work day.</p>

For more information on regulations and advisories, see Chapter 8.

Where can I get more information?

If you have any more questions or concerns, please contact your community or state health or environmental quality department, or contact ATSDR at the address and phone number below.

ATSDR can also tell you the location of occupational and environmental health clinics. These clinics specialize in recognizing, evaluating, and treating illnesses that result from exposure to hazardous substances.

1. PUBLIC HEALTH STATEMENT

Toxicological profiles are also available on-line at www.atsdr.cdc.gov and on CD-ROM. You may request a copy of the ATSDR ToxProfiles™ CD-ROM by calling the toll-free information and technical assistance number at 1-800-CDCINFO (1-800-232-4636), by e-mail at cdcinfo@cdc.gov, or by writing to:

Agency for Toxic Substances and Disease Registry
Division of Toxicology and Environmental Medicine
1600 Clifton Road NE
Mailstop F-32
Atlanta, GA 30333
Fax: 1-770-488-4178

Organizations for-profit may request copies of final Toxicological Profiles from the following:

National Technical Information Service (NTIS)
5285 Port Royal Road
Springfield, VA 22161
Phone: 1-800-553-6847 or 1-703-605-6000
Web site: <http://www.ntis.gov/>

2. RELEVANCE TO PUBLIC HEALTH

2.1 BACKGROUND AND ENVIRONMENTAL EXPOSURES TO STYRENE IN THE UNITED STATES

Styrene is a high production chemical; over 13 billion pounds of styrene was produced in the United States in 2006. Small amounts of styrene are naturally present in foods such as legumes, beef, clams, eggs, nectarines, and spices. It can also be present in packaged foods by migration from polystyrene food containers and packaging materials. Styrene is a combustion product of cigarette smoke and automobile exhaust. Manufactured styrene is primarily used in the production of polystyrene plastics and resins used principally for insulation or in the fabrication of fiberglass boats; production of copolymers such as styrene-acrylonitrile and acrylonitrile-butadiene-styrene, which are used to manufacture piping, automotive components, and plastic drinking glasses; production of styrene-butadiene rubber used to manufacture car tires, hoses for industrial purposes, and shoes; or formulated with unsaturated polyester resins used as fiberglass reinforcement materials. Styrene copolymers are also frequently used in liquid toner for photocopiers and printers.

Median styrene concentrations in urban and rural/suburban air samples are 0.07–0.9 ppb and 0.06–0.08 ppb. The median styrene concentration in indoor air samples ranged from 0.09 to 2 ppb; the primary sources of styrene in indoor air are cigarette smoke and photocopiers. Styrene is rarely detected in drinking water samples and is rarely detected in soil samples.

General population exposure to styrene in air and food has been estimated to be 18–54 and 0.2–1.2 $\mu\text{g}/\text{person}/\text{day}$, respectively, with a total daily exposure of 18.2–55.2 $\mu\text{g}/\text{day}$ or 0.0003–0.0008 $\text{mg}/\text{kg}/\text{day}$ (assuming a 70-kg reference body weight).

2.2 SUMMARY OF HEALTH EFFECTS

Styrene-induced neurotoxicity has been reported in workers since the 1970s. Studies over the last 15 years have firmly established the central nervous system as the critical target of toxicity. Both short- and long-term exposures to styrene can result in neurological effects. Acute exposure data are limited to the finding of impaired performance on tests of vestibular function in test subjects exposed to 87–376 ppm for 1–3 hours. A variety of neurological effects have been observed in chronically exposed styrene workers; these effects include decreased color discrimination, vestibular effects, hearing impairment, symptoms of neurotoxicity, particularly "feeling drunk" and tiredness, delays in reaction

2. RELEVANCE TO PUBLIC HEALTH

time, impaired performance on tests measuring attention and memory, impaired nerve conduction velocity, and EEG alterations. The LOAELs for these effects range from about 10 ppm to 93 ppm. In most of the occupational exposure studies, neurological function tests were conducted in the morning before work, suggesting that the deficits were not acute effects. Results of a meta-analysis suggest that the severity of the some of the neurological symptoms increases with exposure duration. For example, 8, 15, 25, and 35% increases in reaction time were observed in workers exposed to 100 ppm for 2, 4, 6, and 8 work-years, respectively. However, this may also be reflective of higher exposure levels in the past rather than a duration-related increase in severity. The existing data are inadequate to determine whether chronic styrene exposure results in permanent damage. Mixed results have been found in studies examining workers before and after an extended period without styrene exposure. Neurotoxicity studies in animals have primarily focused on effects on hearing and damage to the organ of Corti.

Other effects that have been observed in animal studies include damage to the nasal olfactory epithelium and liver necrosis; testicular damage and developmental effects have also been reported, but the weight of evidence does not support concluding that these are sensitive targets. Damage to the nasal olfactory epithelium was observed in mice after 3 days of exposure. The severity of the lesion progressed from single cell necrosis to atrophy and respiratory metaplasia with increasing exposure duration. The lowest-observed-adverse-effect levels (LOAELs) for these lesions are 80, 50, and 20 ppm for acute, intermediate, and chronic exposure, respectively. Rats do not appear to be as sensitive as mice to the nasal olfactory epithelial damage; an intermediate-duration study identified a no-observed-adverse-effect level (NOAEL) and LOAEL of 500 and 1,000 ppm for focal hyperplasia and a chronic study identified a LOAEL of 50 ppm for atrophy and degeneration. The observed species differences may be due to differences in styrene metabolism in the nasal cavity. In particular, rats have a higher capacity to detoxify styrene oxide with epoxide hydrolases and glutathione S-transferase. Humans are not likely sensitive to the nasal toxicity of styrene because styrene oxide has not been detected and high levels of epoxide hydrolases have been found in *in vitro* assays of human nasal tissue.

Unlike the nasal lesions, the severity of hepatic lesions decreases with increased exposure durations. Severe hepatocellular necrosis was observed in mice exposed to 250 ppm for 3 days; however, continued exposure at this concentration resulted in focal necrosis and an increase in pigmented macrophages. Centrilobular aggregates of siderophages were observed in mice exposed to 200 ppm for 13 weeks; no liver effects were observed at 160 ppm after 2 years of exposure. Rats are less sensitive than mice to liver toxicity; no liver effects were observed in an intermediate-duration study in which rats were exposed to a styrene concentration 10-fold higher than the concentration eliciting hepatic effects in mice. No

2. RELEVANCE TO PUBLIC HEALTH

alterations in serum markers of liver damage were observed in styrene workers exposed to 40 ppm for approximately 5 years. Liver effects have not been observed in rats orally exposed to 35 mg/kg/day for 105 weeks. Some hepatic alterations (increases in liver weight and small areas of focal necrosis) have been reported in rats exposed to 400 mg/kg for an intermediate duration; however, the studies are poorly reported and lack statistical comparisons with controls. No studies examined systemic end points following acute exposure.

Occupational exposure studies have not found significant increases in the occurrence of stillbirth, infant death, malformations, or low birth weight. An increase in fetal deaths were observed in hamsters exposed to very high concentrations (1,000 ppm on gestation days 6–18) and in rats exposed to 300 ppm on gestation days 6–20. However, most single and multigeneration inhalation and oral exposure animal studies did not find significant alterations in fetus/pup survival, growth, or incidence of abnormalities in rats, mice, rabbits, and hamsters exposed to styrene. Two studies have examined neurodevelopmental effects in rats; one study found some minor effects (slight delays in some developmental landmarks). The other, higher-quality study did not find any significant alterations in a number of neurodevelopmental end points. The National Toxicology Program (NTP) Expert Panel examining the developmental potential of styrene (NTP 2006) concluded that the human data are not sufficient to evaluate the potential developmental toxicity of styrene in humans and that there was no convincing evidence of developmental toxicity in animals.

Although several epidemiology studies have examined potential reproductive effects in male and female styrene workers, adequate analysis of the data is limited by the lack of exposure information and concomitant exposure to other compounds. Mixed results have been found for increased occurrence of spontaneous abortions and oligomenorrhea. In male workers, sperm abnormalities have been reported, but not alterations in time-to-pregnancy or fertility rates. No adverse reproductive effects were observed in inhalation and oral multigeneration studies in rats. A series of studies found decreases in spermatozoa counts in rats exposed as adults, as neonates, and through lactation. However, as noted by the NTP Expert Panel, this finding is not consistent with the lack of reproductive effects found in the inhalation two-generation study. The NOAEL identified in the two-generation inhalation study was 500 ppm (6 hours/day), which is roughly equivalent to 230 mg/day using a reference inhalation rate of 0.42 m³/day. The LOAEL for spermatozoa effects in adult rats was 400 mg/kg (6 days/week), which is roughly equivalent to 158 mg/day using a reference body weight of 0.462 kg.

2. RELEVANCE TO PUBLIC HEALTH

There are several epidemiologic studies of workers at styrene manufacturing and polymerization facilities and reinforced plastics facilities that suggest an association between occupational exposure and an increased incidence of cancer of the lymphatic and hematopoietic tissues in styrene. However, the reported studies are inconclusive due to exposure to multiple chemicals (including benzene) and the small size of the cohorts. Other studies have reported negative results. More consistent results for increases in the risk of lymphatic and hematopoietic cancers have been observed among workers at styrene-butadiene manufacturing facilities. There is suggestive evidence that these increased risks may be due to exposure to 1,3-butadiene rather styrene exposure; however, it is difficult to separate the risks for styrene and 1,3-butadiene because the exposure is highly correlated. There are no reports of cancer resulting from styrene exposure by the oral or dermal routes in humans. Species differences in styrene carcinogenicity have been detected in animal studies. Inhalation and oral exposure studies in rats have not found significant increases in neoplastic lesions. However, increases in lung tumors have been found in mice following inhalation and oral exposure. The increased production of styrene 7,8-oxide in lung Clara cells and the higher ratio of styrene oxide R- to S-enantiomers likely resulted in the increased sensitivity of mice. Overall, human and animal studies suggest that styrene may be a weak human carcinogen. The International Agency for Research on Cancer (IARC) has assigned styrene to Group 2B, possibly carcinogenic to humans.

2.3 MINIMAL RISK LEVELS (MRLs)

Estimates of exposure levels posing minimal risk to humans (MRLs) have been made for styrene. An MRL is defined as an estimate of daily human exposure to a substance that is likely to be without an appreciable risk of adverse effects (noncarcinogenic) over a specified duration of exposure. MRLs are derived when reliable and sufficient data exist to identify the target organ(s) of effect or the most sensitive health effect(s) for a specific duration within a given route of exposure. MRLs are based on noncancerous health effects only and do not consider carcinogenic effects. MRLs can be derived for acute, intermediate, and chronic duration exposures for inhalation and oral routes. Appropriate methodology does not exist to develop MRLs for dermal exposure.

Although methods have been established to derive these levels (Barnes and Dourson 1988; EPA 1990), uncertainties are associated with these techniques. Furthermore, ATSDR acknowledges additional uncertainties inherent in the application of the procedures to derive less than lifetime MRLs. As an example, acute inhalation MRLs may not be protective for health effects that are delayed in development or are acquired following repeated acute insults, such as hypersensitivity reactions, asthma, or chronic

2. RELEVANCE TO PUBLIC HEALTH

bronchitis. As these kinds of health effects data become available and methods to assess levels of significant human exposure improve, these MRLs will be revised.

Inhalation MRLs***Acute-duration Inhalation MRL***

- An MRL of 2 ppm has been derived for acute-duration inhalation exposure (14 days or less) to styrene.

The acute-duration inhalation toxicity database for styrene consists of several human experimental studies primarily examining neurotoxicity (Ödkvist et al. 1982; Seeber et al. 2004; Stewart et al. 1968), systemic toxicity studies in mice (Cruzan et al. 1997, 2001; Morgan et al. 1993a, 1993b, 1993c), neurotoxicity studies in rats (Campo et al. 2001; Crofton et al. 1994; Lataye et al. 2003), mice (Cruzan et al. 1997; DeCeaurrez et al. 1983), and guinea pigs (Lataye et al. 2003), a reproductive toxicity study in mice (Salomaa et al. 1985), and developmental toxicity studies in rats (Murray et al. 1978), mice (Kankaanpää et al. 1980), hamsters (Kankaanpää et al. 1980), and rabbits (Murray et al. 1978). Eye irritation was reported in humans exposed to 99 ppm for 7 hours or 376 ppm for 1 hour (Stewart et al. 1968); nasal irritation was also reported at 376 ppm. A significant inhibition of the vestibular-oculomotor system was observed in subjects exposed to 87 ppm for 1 hour (Ödkvist et al. 1982). Studies by (Stewart et al. 1968) found alterations in tests of balance or coordination in subjects exposed to 376 ppm for 1 hour, but not after exposure to 99 ppm for 7 hours or 216 ppm for 1 hour; the test used in the Stewart et al. (1968) studies is probably less sensitive than those used by Ödkvist et al. (1982). No significant alterations in performance on tests of reaction time were observed in subjects exposed to 20 ppm for 3 hours (Seeber et al. 2004).

In mice, the most sensitive target of styrene toxicity appears to be the nasal olfactory epithelium; single cell necrosis was observed following exposure to 80 ppm 6 hours/day for 3 days (Cruzan et al. 2001). At 250 ppm, hepatocellular necrosis and degeneration have been observed (Cruzan et al. 1997; Morgan et al. 1993a, 1993b, 1993c). The severity of this lesion appears to be inversely related to the duration of exposure, with more severe damage observed in mice killed within 3 days of exposure (Morgan et al. 1993a, 1993b, 1993c) compared to animals killed after 2 weeks of exposure (Cruzan et al. 1997; Morgan et al. 1993a). Exposure to 250 ppm 6 hours/day, 5 days/week for 2 weeks also resulted in lethargy and unsteady gait in mice (Cruzan et al. 1997). Impaired performance on a swimming test was observed in mice exposed to 610 ppm for 4 hours, but not in animals exposed to 413 ppm (DeCeaurrez et al. 1983).

2. RELEVANCE TO PUBLIC HEALTH

Exposure of rats to high concentrations (1,000 or 1,600 ppm) 6–8 hours/day for 5–14 days resulted in auditory threshold shifts (indicative of hearing loss) and loss of outer hair cells (OHC) in the organ of Corti (Campo et al. 2001; Crofton et al. 1994; Lataye et al. 2003). No alterations in sperm morphology were observed in mice exposed to 300 ppm styrene 5 hours/day for 5 days (Salomaa et al. 1985) and no developmental effects were observed in rats or rabbits exposed to 600 ppm 7 hours/day on gestational days 6–15 or 6–18, respectively, (Murray et al. 1978) or mice exposed to 250 ppm 6 hours/day on gestational days 6–16 (Kankaanpää et al. 1980). An increase in fetal deaths or resorptions was observed in hamsters exposed to 1,000 ppm 6 hours/day on gestational days 6–18 (Kankaanpää et al. 1980).

These data suggest that the nervous system is the most sensitive target of styrene toxicity in humans following acute-duration inhalation exposure. The lowest LOAEL for a relevant end point in humans is 87 ppm for vestibular impairment in subjects exposed to styrene for 1 hour (Ödkvist et al. 1982). A similar LOAEL (80 ppm) was identified for nasal effects in mice exposed to styrene for 3 days (Cruzan et al. 2001). Although nasal irritation has been observed in humans exposed to 376 ppm for 1 hour (Stewart et al. 1968) and focal hyperplasia in the nasal olfactory epithelium was observed in rats exposed to 1,000 ppm (NOAEL of 500 ppm) for 13 weeks (Cruzan et al. 1997), mice appears to be unusually susceptible to this effect. As discussed in Section 2.2, mice appear to have a greater capacity than humans to generate the reactive metabolite, styrene oxide, in the nasal cavity and a lower capacity to detoxify styrene oxide (Green et al. 2001a). Thus, nasal lesions in mice were not considered suitable as the basis of an MRL. The identification of the nervous system as the critical target of toxicity for styrene is supported by a large number of occupational exposure studies. Delays in reaction time have been observed in workers exposed to 21.9–92 ppm (Cherry et al. 1980; Fallas et al. 1992; Gamberale et al. 1976; Jegaden et al. 1993; Mutti et al. 1984a; Tsai and Chen 1996) and vestibular effects have been observed at 18–36 ppm (Calabrese et al. 1996; Möller et al. 1990; Toppila et al. 2006).

The Ödkvist et al. (1982) study did not identify a NOAEL for vestibular effects; however, a NOAEL of 20 ppm for performance on several tests of reaction time and attention was identified by Seeber et al. (2004) in subjects exposed to styrene for 3 hours. Although there is some uncertainty whether deriving an MRL based on a 3-hour exposure study would be protective of continuous exposure to styrene for 2 weeks, the Seeber et al. (2004) study was selected as the basis of an acute duration inhalation MRL for styrene.

Groups of eight volunteers (gender not reported) were exposed to 0.5 or 20 ppm styrene for 3 hours (Seeber et al. 2004). The subjects were tested for simple reaction time, choice reaction time, and attention

2. RELEVANCE TO PUBLIC HEALTH

prior to exposure initiation, during the third hour of exposure, and 1.5 hours after exposure termination. The subjects were also asked to complete a symptom questionnaire before, during, and after exposure. The mean concentration of styrene in blood was 2.2 and 80 µg/L after 3 hours of exposure to 0.5 and 20 ppm, respectively. The blood levels of styrene were correlated with styrene levels in air ($r_{xy}=0.98$). No significant alterations in performance on neurobehavioral tests were found. An increase in the reporting of breathing problems was found; however, it was not statistically significant and the ranking of the severity of the breathing problem was very low (0.5 on a scale of 5).

The NOAEL of 20 ppm from the Seeber et al. (2004) study was divided by an uncertainty factor of 10 to account for human variability resulting in an acute-duration inhalation MRL of 2 ppm.

Intermediate-duration Inhalation MRL

No human intermediate-duration studies were identified. Animal studies examining systemic, neurological, reproductive, and developmental toxicity have identified the respiratory tract as the most sensitive target of toxicity. Atrophy of the olfactory epithelium, hypertrophy/hyperplasia of Bowman's gland has been observed in mice exposed to 50 ppm 6 hours/day, 5 days/week for 13 weeks (Cruzan et al. 1997), decreased nasal cilia activity has been observed in rats exposed to 150 ppm 4 hours/day, 5 days/week for 21 days (Ohashi et al. 1986), and focal hyperplasia has been observed in rats exposed to 1,000 ppm 6 hours/day, 5 days/week for 13 weeks (Cruzan et al. 1997). As discussed previously, the mouse does not appear to be a good model for nasal effects in humans due metabolic differences. Other systemic effects that have been observed include eye irritation in rats exposed to 200 ppm 6 hours/day, 5 days/week for 13 weeks (Cruzan et al. 1997) and centrilobular aggregates of siderophages in the livers of mice exposed to 200 ppm 6 hours/day, 5 days/week for 13 weeks (Cruzan et al. 1997).

A number of studies in rats have reported outer hair cell loss in the organ of Corti in rats exposed to 600–650 ppm for 4 weeks (Loquet et al. 2000; Makitie et al. 2002; Pouyatos et al. 2002) and hearing loss at 750–1,000 ppm for 3–4 weeks (Campo et al. 2001; Lataye et al. 2000, 2001; Loquet et al. 1999, 2000; Pouyatos et al. 2002). A NOAEL of hearing effects of 300 ppm was identified in rats exposed for 12 hours/day, 5 days/week for 4 weeks (Makitie et al. 2002). Other neurological effects include alterations in astroglial cells in rats continuously exposed to 320 ppm for 3 months (Rosengren and Haglid 1989) and decreased sensory nerve conduction velocity in rats exposed to 2,000 ppm 8 hours/day, 5 days/week for 32 weeks (Yamamoto et al. 1997). No reproductive, developmental, or neurodevelopmental effects were observed in a two-generation study (Cruzan et al. 2005a, 2005b); the NOAEL was

2. RELEVANCE TO PUBLIC HEALTH

500 ppm. In contrast, an increase in neonatal deaths, developmental landmark delays, and alterations in neurochemical levels were observed in the offspring of rats exposed 6 hours/day on gestational days 6–20 (Katakura et al. 1999, 2001).

Chronic-duration studies suggest that the most sensitive target of styrene toxicity is the nervous system. It is likely that this would also be the most sensitive effect following intermediate-duration exposure. In the absence of human neurotoxicity data, an intermediate-duration inhalation MRL is not recommended at this time.

Chronic-duration Inhalation MRL

- An MRL of 0.2 ppm has been derived for chronic-duration inhalation exposure (greater than 365 days) to styrene.

A large number of studies have examined the neurotoxicity of styrene in workers at reinforced plastic manufacturing facilities. These studies reported a variety of neurological effects, including decreased color discrimination, slowed reaction time, permanent hearing threshold shifts, vestibular effects, and increases in subjective symptoms. A summary of the results of studies for some of these neurological effects is presented in Table 2-1. The LOAELs for these effects range from 6 to 93 ppm.

Chronic-duration studies in animals identify the nasal olfactory epithelium as the most sensitive end point. Atrophic and/or degenerative changes were observed in rats exposed to 50 ppm styrene 6 hours/day, 5 days/week for 104 weeks (Cruzan et al. 1998) and respiratory metaplasia in the nasal olfactory epithelium has been observed in mice exposed to 20 ppm 6 hours/day, 5 days/week for 98–104 weeks (Cruzan et al. 2001). As noted previously, mice do not appear to be a good model for potential respiratory effects in humans.

Neurotoxicity observed in styrene workers was selected as the basis of the chronic-duration inhalation MRL for styrene. Two approaches to deriving the MRL were considered. In the first approach, the MRL is based on a single study identifying a sensitive LOAEL. In the study by Kishi et al. (2001), styrene workers were divided into three groups based on urinary mandelic acid excretion levels. A significant increase in color confusion index (CCI) was found in the workers exposed to equivalent styrene concentrations of 10 or 46 ppm, as compared to age-matched controls. No significant alterations were observed in workers exposed to 4 ppm. The second approach involves the use of a LOAEL estimated

2. RELEVANCE TO PUBLIC HEALTH

Table 2-1. Results of Selected Human Neurotoxicity Studies

Result	Reference	NOAEL ppm	LOAEL ppm
Decreased color discrimination	Chia et al. 1994		6
	Kishi et al. 2001	4	10
	Gong et al. 2002		10
	Gobba et al. 1991		16
	Iregren et al. 2005		22
	Fallas et al. 1992		24.3
	Campagna et al. 1996		26
	Eguchi et al. 1995	8	93
Neurological symptoms	Flodin et al. 1989	6	
	Edling et al. 1993	8.6	
	Checkoway et al. 1992	10.8	18.9
	Cherry et al. 1980		92
Vestibular effects	Möller et al. 1990		18
	Toppila et al. 2006		24.8
	Calabrese et al. 1996		36
Reaction time	Edling et al. 1993	8.6	
	Tsai and Chen 1996		21.9
	Jegaden et al. 1993		22.68
	Fallas et al. 1992		24.3
	Mutti et al. 1984a		25
	Gamberale et al. 1976		47
	Cherry et al. 1980		92
	Hearing	Morata et al. 2002	
Śliwińska-Kowalska et al. 2003			15.6
Morioka et al. 1999			16
Möller et al. 1990		18	
Calabrese et al. 1996		36	
Nerve conduction velocity		Seppäläinen and Härkönen 1976	30
	Štětkařová et al. 1993		50
	Triebig et al. 1985	100	

2. RELEVANCE TO PUBLIC HEALTH

from a meta-analysis of occupational exposure studies finding effects on color vision and reaction time (Benignus et al. 2005). Benignus et al. (2005) used data color vision impairment data from the Campagna et al. (1996), Eguchi et al. (1995), Gobba et al. (1991), Gong et al. (2002), and Kishi et al. (2001) studies and choice reaction time data from the Jegaden et al. (1993), Mutti et al. (1984a), Triebig et al. (1989), and Tsai and Chen (1996) studies. Average styrene exposure concentrations were estimated from individual data reported in the papers; for studies reporting individual data as urinary mandelic acid levels, Benignus et al. (2005) used standardized methods for converting to styrene exposure levels. Cumulative styrene exposure was estimated by multiplying exposure level by length of employment. A common metric of effect magnitude (percentage of baseline) was calculated for the different neurological effects. The analysis found a significant linear relationship between choice reaction time and cumulative styrene exposure; cumulative exposure accounted for 91% of the variance in reaction time. Similarly, a significant relationship between CCI and cumulative styrene exposure was found, with cumulative exposure accounting for 35% of the variance in CCI. Using the regression equations for these two effects, Benignus et al. (2005) estimated that exposure to 150 ppm for 8 work-years would result in a 50% increase in choice reaction time and a 17% increase in CCI score; exposure to 20 ppm for 8 work-years would result in a 6.5% increase in choice reaction time and a 2.23% increase in CCI score. As discussed in Benignus et al. (2005), a 7% decrease in reaction time would prevent 58,000–70,000 injuries per year from automobile accidents. The investigators also noted that CCI increases with age, the rate of increase is about 10% per 13 years of age; thus, a 2.23% decrease in color perception would be roughly equivalent to 2.9 additional years of age. Based on this analysis, 20 ppm is considered a LOAEL for neurological effects.

The LOAEL of 20 ppm is consistent with the LOAEL values identified in many of the individual studies. However, using the LOAEL identified from the Benignus et al. (2005) meta-analysis has several advantages over selecting a single study as the basis of the MRL. Because data were pooled from several studies, the relationships between styrene exposure and effects were examined in a large number of subjects (302 subjects for choice reaction time and 383 subjects for color vision). The use of standardized methods for estimating styrene exposure levels from urinary biomarker levels is also an advantage. Additionally, the biological relevance of the observed deficits in reaction time and color vision was estimated. The LOAEL of 20 ppm was divided by an uncertainty factor of 100 (10 for use of a LOAEL and 10 for human variability) resulting in a chronic-duration inhalation MRL of 0.2 ppm.

2. RELEVANCE TO PUBLIC HEALTH

Oral MRLs***Acute-duration Oral MRL***

- An MRL of 0.1 mg/kg/day has been derived for acute-duration oral exposure (14 days or less) to styrene.

A limited number of studies have examined the acute toxicity of orally administered styrene; these studies have examined potential neurotoxicity and developmental toxicity. No developmental effects were observed in rats administered a single dose of 300 mg/kg on gestational day 11 (Daston et al. 1991) or administered 300 mg/kg/day (administered as two daily doses of 150 mg/kg) on gestational days 6–15 (Murray et al. 1978). Impaired learning was observed in rats administered via gavage 100 or 200 mg/kg/day for 14 days; increases in serotonin levels were observed in the hypothalamus, hippocampus, and midbrain (Husain et al. 1985). In another study, increases in dopamine receptor binding was observed in rats administered a single gavage dose of 200 mg/kg (Agrawal et al. 1982).

Although a limited number of toxicity end points have been examined following acute-duration oral exposure, longer-term oral studies examining systemic and reproductive end points have identified LOAELs that were higher than the 100 mg/kg/day LOAEL identified for neurotoxicity in the Husain et al. (1985) study. The lowest LOAEL identified for a systemic effect is 400 mg/kg/day for Heinz body formation in dogs administered styrene by gavage for 561 days (Quast et al. 1979); the NOAEL was 200 mg/kg/day. Decreased spermatozoa counts were observed in adult rats administered 400 mg/kg 6 days/week for 60 days (Srivastava et al. 1989), young rats exposed via lactation on postnatal days 1–21 (maternal dose of 400 mg/kg/day) (Srivastava et al. 1992a), and young rats administered 200 mg/kg 6 days/week on postnatal days 1–61 (Srivastava et al. 1992b); the NOAELs identified in these three studies were 200, 200, and 100 mg/kg, respectively. Marked degeneration of the seminiferous tubules was also observed in the adult rats administered 400 mg/kg (Srivastava et al. 1989). Impaired learning observed in rats administered 500 mg/kg 5 days/week for 8 weeks (no NOAEL identified) (Bushnell 1994) also supports the identification of neurotoxicity as a sensitive end point following oral exposure. Additionally, the extensive inhalation toxicity database for styrene supports the selection of neurotoxicity as the most sensitive target of toxicity; both the acute- and chronic-duration inhalation MRLs are based on neurological effects in humans. Neurological effects observed in chronically exposed styrene workers include decreased color discrimination, slowed reaction time, increased prevalence of neurological symptoms, and ototoxicity (hearing and vestibular effects).

2. RELEVANCE TO PUBLIC HEALTH

The Husain et al. (1985) study was selected as the basis of the acute-duration oral MRL. In this study, groups of 15 male Wistar rats were administered by gavage 0, 100, or 200 mg/kg/day styrene in ground nut oil for 14 consecutive days. Spontaneous motor activity with or without amphetamine induction was observed 1 day after the last dose. Two days after exposure termination, the rats underwent acquisition training for 4 days. Learning was assessed by measuring the number of times the rat climbed the pole after the conditioned stimulus to avoid the foot-shock unconditioned stimulus. Noradrenaline, dopamine, and serotonin levels were measured in seven regions of the brain in six rats/group sacrificed after the acquisition training. No overt signs of toxicity were observed. No significant alterations in locomotor activity were observed with or with amphetamine induction. Significantly greater increases in percent avoidance response in the conditioned avoidance response test (indicative of impaired learning) were observed at 100 and 200 mg/kg/day; no difference was found between the two styrene groups. The effects were observed on test day 3 and 4. Significant increases in the level of serotonin in the hypothalamus (70%), hippocampus (51%), and midbrain (29%) were observed at 200 mg/kg/day. Styrene exposure did not affect brain noradrenaline and dopamine levels.

The LOAEL of 100 mg/kg/day was divided by an uncertainty factor of 1,000 (10 for use of a LOAEL, 10 for extrapolation from animals to humans, and 10 for human variability).

Intermediate-duration Oral MRL

The systemic toxicity of styrene has not been investigated in intermediate-duration oral exposure studies. Neurotoxicity studies have identified a LOAEL of 200 mg/kg/day for increased dopamine receptor binding in rats (Agrawal et al. 1982), a LOAEL of 500 mg/kg (5 days/week) for impaired learning in rats (Bushnell 1994), and a LOAEL of 906 mg/kg/day for alterations in serotonin and noradrenaline levels in rats (Husain et al. 1980); none of these studies identified a NOAEL for neurological effects. An increase in dopamine receptor binding was also observed in the offspring of rats administered 200 mg/kg/day during gestation, lactation, or both (Zaidi et al. 1985). Reproductive and immunological effects were reported in the other intermediate-duration oral studies. Decreases in spermatozoa counts were observed in rats exposed as 400 mg/kg (6 days/week) as adults, 200 mg/kg (6 days/week) as neonates, or during lactation (maternal dose of 400 mg/kg/day) (Srivastava et al. 1989, 1992a, 1992b). Impaired immune function was observed in mice exposed to 30 mg/kg/day and in rats exposed to 294 mg/kg/day (Dogra et al. 1992); the NOAELs were 23 and 196 mg/kg/day.

2. RELEVANCE TO PUBLIC HEALTH

Dogra et al. (1992) identified the lowest LOAEL following intermediate-duration oral exposure to styrene; however, there are limited data to support the identification of the immune system as a sensitive, relevant target for humans. Although, the sensitivity of the nervous system has been firmly established following inhalation and oral exposure, the LOAELs identified in the intermediate-duration studies are higher than the lowest LOAEL for neurotoxicity identified in an acute-duration study (Husain et al. 1985). An MRL based on the Dogra et al. (1992) study would be higher than the acute-duration oral MRL; thus, an intermediate-duration MRL is not recommended at this time.

Chronic-duration Oral MRL

The available data on the chronic toxicity of styrene comes from three systemic toxicity studies. No adverse effects were observed in rats exposed to 35 mg/kg/day styrene in drinking water for 2 years (Beliles et al. 1985) and no liver or kidney alterations were observed in rats administered 500 mg/kg 1 day/week for 120 weeks (Ponomarev and Tomatis 1978). Increase in Heinz body formation was observed in dogs administered 400 mg/kg/day for 561 days (Quast et al. 1979); the NOAEL for this effect is 200 mg/kg/day.

The chronic-duration inhalation database provides strong evidence that neurotoxicity is the most sensitive target of styrene toxicity. It is not known if this would also be true for chronic-duration oral exposure; the acute-toxicity oral database provides suggestive evidence that it would be a sensitive target. In the absence of a long-term oral study examining neurological end points, a chronic-duration oral MRL is not recommended.

2. RELEVANCE TO PUBLIC HEALTH

This page is intentionally blank.

3. HEALTH EFFECTS

3.1 INTRODUCTION

The primary purpose of this chapter is to provide public health officials, physicians, toxicologists, and other interested individuals and groups with an overall perspective on the toxicology of styrene. It contains descriptions and evaluations of toxicological studies and epidemiological investigations and provides conclusions, where possible, on the relevance of toxicity and toxicokinetic data to public health.

A glossary and list of acronyms, abbreviations, and symbols can be found at the end of this profile.

3.2 DISCUSSION OF HEALTH EFFECTS BY ROUTE OF EXPOSURE

To help public health professionals and others address the needs of persons living or working near hazardous waste sites, the information in this section is organized first by route of exposure (inhalation, oral, and dermal) and then by health effect (death, systemic, immunological, neurological, reproductive, developmental, genotoxic, and carcinogenic effects). These data are discussed in terms of three exposure periods: acute (14 days or less), intermediate (15–364 days), and chronic (365 days or more).

Levels of significant exposure for each route and duration are presented in tables and illustrated in figures. The points in the figures showing no-observed-adverse-effect levels (NOAELs) or lowest-observed-adverse-effect levels (LOAELs) reflect the actual doses (levels of exposure) used in the studies. LOAELs have been classified into "less serious" or "serious" effects. "Serious" effects are those that evoke failure in a biological system and can lead to morbidity or mortality (e.g., acute respiratory distress or death). "Less serious" effects are those that are not expected to cause significant dysfunction or death, or those whose significance to the organism is not entirely clear. ATSDR acknowledges that a considerable amount of judgment may be required in establishing whether an end point should be classified as a NOAEL, "less serious" LOAEL, or "serious" LOAEL, and that in some cases, there will be insufficient data to decide whether the effect is indicative of significant dysfunction. However, the Agency has established guidelines and policies that are used to classify these end points. ATSDR believes that there is sufficient merit in this approach to warrant an attempt at distinguishing between "less serious" and "serious" effects. The distinction between "less serious" effects and "serious" effects is considered to be important because it helps the users of the profiles to identify levels of exposure at which major health effects start to appear. LOAELs or NOAELs should also help in determining whether or not

3. HEALTH EFFECTS

the effects vary with dose and/or duration, and place into perspective the possible significance of these effects to human health.

The significance of the exposure levels shown in the Levels of Significant Exposure (LSE) tables and figures may differ depending on the user's perspective. Public health officials and others concerned with appropriate actions to take at hazardous waste sites may want information on levels of exposure associated with more subtle effects in humans or animals (LOAELs) or exposure levels below which no adverse effects (NOAELs) have been observed. Estimates of levels posing minimal risk to humans Minimal Risk Levels or (MRLs) may be of interest to health professionals and citizens alike.

A User's Guide has been provided at the end of this profile (see Appendix B). This guide should aid in the interpretation of the tables and figures for Levels of Significant Exposure and the MRLs.

3.2.1 Inhalation Exposure

Most information on the effects of inhalation exposure to styrene in humans comes from studies of workers exposed to styrene vapors in the production and use of plastics and resins, especially polyester resins dissolved in styrene. In most cases, the studies involve workplace exposures such as fiberglass boat building factories where the actual levels of styrene are reported as a range of styrene air concentrations. However, there are a few human clinical studies in which exposures are better quantified. A common limitation of many of the occupational exposure studies is the phenomenon of the healthy worker effect. The selection of healthy individuals for employment and the likelihood that more susceptible workers are more likely to leave the workforce can result in workers who are healthier than the general population. This type of bias typically affects comparisons with the general population and is less likely to influence comparisons with other groups of workers. Provided below are descriptions of the known effects of inhalation exposure of humans and animals to styrene.

3.2.1.1 Death

There have been no reports of deaths in humans directly associated with exposure to styrene in the workplace (EPA 1985a; Gosselin et al. 1984; NIOSH 1983).

In animals, inhalation studies indicate that the acute toxicity of styrene is low to moderate. An LC₅₀ of 2,770 ppm after 2 hours of exposure was reported in rats, and the LC₅₀ for mice after exposure for 4 hours was 4,940 ppm (Shugaev 1969). All rats and guinea pigs survived after exposure to 1,300 ppm styrene

3. HEALTH EFFECTS

for 30 hours and 16 hours, respectively (Spencer et al. 1942). However, all animals died after 40 hours of exposure. Gender differences in mortality were observed in repeated-exposure studies (6 hours/day, 5 days/week for 2 weeks) (Cruzan et al. 1997). Increases in mortality were observed in female CD-1 and B6C3F1 female mice exposed to 250 ppm; no deaths were observed at 500 ppm. In the CD-1 and B6C3F1 males, very few deaths were observed at 250 ppm, but increases in deaths were observed at 500 ppm. A similar finding was reported by Morgan et al. (1993a): increases in mortality were observed in female B6C3F1 mice exposed to 250 ppm and no deaths were observed at 500 ppm; in males, deaths were observed at 250 and 500 ppm. In contrast to these findings, no deaths were observed in Sprague Dawley rats exposed to concentrations as high as 1,500 ppm 6 hours/day, 5 days/week for 13 weeks (Cruzan et al. 1997).

All reliable LOAEL values and LC₅₀ values for lethality in each species and duration category are recorded in Table 3-1 and plotted in Figure 3-1.

3.2.1.2 Systemic Effects

No studies were located regarding dermal or metabolic effects in humans or animals after inhalation exposure to styrene.

For the following systemic effects resulting from inhalation exposure to styrene, the highest NOAEL values and all reliable LOAEL values in each species and duration category are recorded in Table 3-1 and plotted in Figure 3-1.

Respiratory Effects. Several human studies have examined the respiratory effects caused by inhalation exposure to styrene. The most commonly reported general symptom is mucous membrane irritation. Irritation of the upper respiratory tract (i.e., nose and throat) has been reported by volunteers (Carpenter et al. 1944; Stewart et al. 1968) and workers (NIOSH 1983). Throat irritation and increased nasal secretion occurred following exposure of two male subjects to 800 ppm for 4 hours (Carpenter et al. 1944). Nasal irritation was observed in all volunteers after exposure to 376 ppm styrene for 60 minutes (Stewart et al. 1968). Obstructive lung changes were observed in 4 of 21 workers exposed to styrene for about 10 years (Chmielewski and Renke 1975). However, exposure levels were not defined. No histological alterations were observed in nasal biopsies from styrene workers exposed to 50–60 ppm styrene for 7 years (Ödkvist et al. 1985).

3. HEALTH EFFECTS

3. HEALTH EFFECTS

3. HEALTH EFFECTS

2. RELEVANCE TO PUBLIC HEALTH

2.1 BACKGROUND AND ENVIRONMENTAL EXPOSURES TO STYRENE IN THE UNITED STATES

Styrene is a high production chemical; over 13 billion pounds of styrene was produced in the United States in 2006. Small amounts of styrene are naturally present in foods such as legumes, beef, clams, eggs, nectarines, and spices. It can also be present in packaged foods by migration from polystyrene food containers and packaging materials. Styrene is a combustion product of cigarette smoke and automobile exhaust. Manufactured styrene is primarily used in the production of polystyrene plastics and resins used principally for insulation or in the fabrication of fiberglass boats; production of copolymers such as styrene-acrylonitrile and acrylonitrile-butadiene-styrene, which are used to manufacture piping, automotive components, and plastic drinking glasses; production of styrene-butadiene rubber used to manufacture car tires, hoses for industrial purposes, and shoes; or formulated with unsaturated polyester resins used as fiberglass reinforcement materials. Styrene copolymers are also frequently used in liquid toner for photocopiers and printers.

Median styrene concentrations in urban and rural/suburban air samples are 0.07–0.9 ppb and 0.06–0.08 ppb. The median styrene concentration in indoor air samples ranged from 0.09 to 2 ppb; the primary sources of styrene in indoor air are cigarette smoke and photocopiers. Styrene is rarely detected in drinking water samples and is rarely detected in soil samples.

General population exposure to styrene in air and food has been estimated to be 18–54 and 0.2–1.2 $\mu\text{g}/\text{person}/\text{day}$, respectively, with a total daily exposure of 18.2–55.2 $\mu\text{g}/\text{day}$ or 0.0003–0.0008 $\text{mg}/\text{kg}/\text{day}$ (assuming a 70-kg reference body weight).

2.2 SUMMARY OF HEALTH EFFECTS

Styrene-induced neurotoxicity has been reported in workers since the 1970s. Studies over the last 15 years have firmly established the central nervous system as the critical target of toxicity. Both short- and long-term exposures to styrene can result in neurological effects. Acute exposure data are limited to the finding of impaired performance on tests of vestibular function in test subjects exposed to 87–376 ppm for 1–3 hours. A variety of neurological effects have been observed in chronically exposed styrene workers; these effects include decreased color discrimination, vestibular effects, hearing impairment, symptoms of neurotoxicity, particularly "feeling drunk" and tiredness, delays in reaction

2. RELEVANCE TO PUBLIC HEALTH

time, impaired performance on tests measuring attention and memory, impaired nerve conduction velocity, and EEG alterations. The LOAELs for these effects range from about 10 ppm to 93 ppm. In most of the occupational exposure studies, neurological function tests were conducted in the morning before work, suggesting that the deficits were not acute effects. Results of a meta-analysis suggest that the severity of the some of the neurological symptoms increases with exposure duration. For example, 8, 15, 25, and 35% increases in reaction time were observed in workers exposed to 100 ppm for 2, 4, 6, and 8 work-years, respectively. However, this may also be reflective of higher exposure levels in the past rather than a duration-related increase in severity. The existing data are inadequate to determine whether chronic styrene exposure results in permanent damage. Mixed results have been found in studies examining workers before and after an extended period without styrene exposure. Neurotoxicity studies in animals have primarily focused on effects on hearing and damage to the organ of Corti.

Other effects that have been observed in animal studies include damage to the nasal olfactory epithelium and liver necrosis; testicular damage and developmental effects have also been reported, but the weight of evidence does not support concluding that these are sensitive targets. Damage to the nasal olfactory epithelium was observed in mice after 3 days of exposure. The severity of the lesion progressed from single cell necrosis to atrophy and respiratory metaplasia with increasing exposure duration. The lowest-observed-adverse-effect levels (LOAELs) for these lesions are 80, 50, and 20 ppm for acute, intermediate, and chronic exposure, respectively. Rats do not appear to be as sensitive as mice to the nasal olfactory epithelial damage; an intermediate-duration study identified a no-observed-adverse-effect level (NOAEL) and LOAEL of 500 and 1,000 ppm for focal hyperplasia and a chronic study identified a LOAEL of 50 ppm for atrophy and degeneration. The observed species differences may be due to differences in styrene metabolism in the nasal cavity. In particular, rats have a higher capacity to detoxify styrene oxide with epoxide hydrolases and glutathione S-transferase. Humans are not likely sensitive to the nasal toxicity of styrene because styrene oxide has not been detected and high levels of epoxide hydrolases have been found in *in vitro* assays of human nasal tissue.

Unlike the nasal lesions, the severity of hepatic lesions decreases with increased exposure durations. Severe hepatocellular necrosis was observed in mice exposed to 250 ppm for 3 days; however, continued exposure at this concentration resulted in focal necrosis and an increase in pigmented macrophages. Centrilobular aggregates of siderophages were observed in mice exposed to 200 ppm for 13 weeks; no liver effects were observed at 160 ppm after 2 years of exposure. Rats are less sensitive than mice to liver toxicity; no liver effects were observed in an intermediate-duration study in which rats were exposed to a styrene concentration 10-fold higher than the concentration eliciting hepatic effects in mice. No

2. RELEVANCE TO PUBLIC HEALTH

alterations in serum markers of liver damage were observed in styrene workers exposed to 40 ppm for approximately 5 years. Liver effects have not been observed in rats orally exposed to 35 mg/kg/day for 105 weeks. Some hepatic alterations (increases in liver weight and small areas of focal necrosis) have been reported in rats exposed to 400 mg/kg for an intermediate duration; however, the studies are poorly reported and lack statistical comparisons with controls. No studies examined systemic end points following acute exposure.

Occupational exposure studies have not found significant increases in the occurrence of stillbirth, infant death, malformations, or low birth weight. An increase in fetal deaths were observed in hamsters exposed to very high concentrations (1,000 ppm on gestation days 6–18) and in rats exposed to 300 ppm on gestation days 6–20. However, most single and multigeneration inhalation and oral exposure animal studies did not find significant alterations in fetus/pup survival, growth, or incidence of abnormalities in rats, mice, rabbits, and hamsters exposed to styrene. Two studies have examined neurodevelopmental effects in rats; one study found some minor effects (slight delays in some developmental landmarks). The other, higher-quality study did not find any significant alterations in a number of neurodevelopmental end points. The National Toxicology Program (NTP) Expert Panel examining the developmental potential of styrene (NTP 2006) concluded that the human data are not sufficient to evaluate the potential developmental toxicity of styrene in humans and that there was no convincing evidence of developmental toxicity in animals.

Although several epidemiology studies have examined potential reproductive effects in male and female styrene workers, adequate analysis of the data is limited by the lack of exposure information and concomitant exposure to other compounds. Mixed results have been found for increased occurrence of spontaneous abortions and oligomenorrhea. In male workers, sperm abnormalities have been reported, but not alterations in time-to-pregnancy or fertility rates. No adverse reproductive effects were observed in inhalation and oral multigeneration studies in rats. A series of studies found decreases in spermatozoa counts in rats exposed as adults, as neonates, and through lactation. However, as noted by the NTP Expert Panel, this finding is not consistent with the lack of reproductive effects found in the inhalation two-generation study. The NOAEL identified in the two-generation inhalation study was 500 ppm (6 hours/day), which is roughly equivalent to 230 mg/day using a reference inhalation rate of 0.42 m³/day. The LOAEL for spermatozoa effects in adult rats was 400 mg/kg (6 days/week), which is roughly equivalent to 158 mg/day using a reference body weight of 0.462 kg.

2. RELEVANCE TO PUBLIC HEALTH

There are several epidemiologic studies of workers at styrene manufacturing and polymerization facilities and reinforced plastics facilities that suggest an association between occupational exposure and an increased incidence of cancer of the lymphatic and hematopoietic tissues in styrene. However, the reported studies are inconclusive due to exposure to multiple chemicals (including benzene) and the small size of the cohorts. Other studies have reported negative results. More consistent results for increases in the risk of lymphatic and hematopoietic cancers have been observed among workers at styrene-butadiene manufacturing facilities. There is suggestive evidence that these increased risks may be due to exposure to 1,3-butadiene rather styrene exposure; however, it is difficult to separate the risks for styrene and 1,3-butadiene because the exposure is highly correlated. There are no reports of cancer resulting from styrene exposure by the oral or dermal routes in humans. Species differences in styrene carcinogenicity have been detected in animal studies. Inhalation and oral exposure studies in rats have not found significant increases in neoplastic lesions. However, increases in lung tumors have been found in mice following inhalation and oral exposure. The increased production of styrene 7,8-oxide in lung Clara cells and the higher ratio of styrene oxide R- to S-enantiomers likely resulted in the increased sensitivity of mice. Overall, human and animal studies suggest that styrene may be a weak human carcinogen. The International Agency for Research on Cancer (IARC) has assigned styrene to Group 2B, possibly carcinogenic to humans.

2.3 MINIMAL RISK LEVELS (MRLs)

Estimates of exposure levels posing minimal risk to humans (MRLs) have been made for styrene. An MRL is defined as an estimate of daily human exposure to a substance that is likely to be without an appreciable risk of adverse effects (noncarcinogenic) over a specified duration of exposure. MRLs are derived when reliable and sufficient data exist to identify the target organ(s) of effect or the most sensitive health effect(s) for a specific duration within a given route of exposure. MRLs are based on noncancerous health effects only and do not consider carcinogenic effects. MRLs can be derived for acute, intermediate, and chronic duration exposures for inhalation and oral routes. Appropriate methodology does not exist to develop MRLs for dermal exposure.

Although methods have been established to derive these levels (Barnes and Dourson 1988; EPA 1990), uncertainties are associated with these techniques. Furthermore, ATSDR acknowledges additional uncertainties inherent in the application of the procedures to derive less than lifetime MRLs. As an example, acute inhalation MRLs may not be protective for health effects that are delayed in development or are acquired following repeated acute insults, such as hypersensitivity reactions, asthma, or chronic

2. RELEVANCE TO PUBLIC HEALTH

bronchitis. As these kinds of health effects data become available and methods to assess levels of significant human exposure improve, these MRLs will be revised.

Inhalation MRLs***Acute-duration Inhalation MRL***

- An MRL of 2 ppm has been derived for acute-duration inhalation exposure (14 days or less) to styrene.

The acute-duration inhalation toxicity database for styrene consists of several human experimental studies primarily examining neurotoxicity (Ödkvist et al. 1982; Seeber et al. 2004; Stewart et al. 1968), systemic toxicity studies in mice (Cruzan et al. 1997, 2001; Morgan et al. 1993a, 1993b, 1993c), neurotoxicity studies in rats (Campo et al. 2001; Crofton et al. 1994; Lataye et al. 2003), mice (Cruzan et al. 1997; DeCeaurrez et al. 1983), and guinea pigs (Lataye et al. 2003), a reproductive toxicity study in mice (Salomaa et al. 1985), and developmental toxicity studies in rats (Murray et al. 1978), mice (Kankaanpää et al. 1980), hamsters (Kankaanpää et al. 1980), and rabbits (Murray et al. 1978). Eye irritation was reported in humans exposed to 99 ppm for 7 hours or 376 ppm for 1 hour (Stewart et al. 1968); nasal irritation was also reported at 376 ppm. A significant inhibition of the vestibular-oculomotor system was observed in subjects exposed to 87 ppm for 1 hour (Ödkvist et al. 1982). Studies by (Stewart et al. 1968) found alterations in tests of balance or coordination in subjects exposed to 376 ppm for 1 hour, but not after exposure to 99 ppm for 7 hours or 216 ppm for 1 hour; the test used in the Stewart et al. (1968) studies is probably less sensitive than those used by Ödkvist et al. (1982). No significant alterations in performance on tests of reaction time were observed in subjects exposed to 20 ppm for 3 hours (Seeber et al. 2004).

In mice, the most sensitive target of styrene toxicity appears to be the nasal olfactory epithelium; single cell necrosis was observed following exposure to 80 ppm 6 hours/day for 3 days (Cruzan et al. 2001). At 250 ppm, hepatocellular necrosis and degeneration have been observed (Cruzan et al. 1997; Morgan et al. 1993a, 1993b, 1993c). The severity of this lesion appears to be inversely related to the duration of exposure, with more severe damage observed in mice killed within 3 days of exposure (Morgan et al. 1993a, 1993b, 1993c) compared to animals killed after 2 weeks of exposure (Cruzan et al. 1997; Morgan et al. 1993a). Exposure to 250 ppm 6 hours/day, 5 days/week for 2 weeks also resulted in lethargy and unsteady gait in mice (Cruzan et al. 1997). Impaired performance on a swimming test was observed in mice exposed to 610 ppm for 4 hours, but not in animals exposed to 413 ppm (DeCeaurrez et al. 1983).

2. RELEVANCE TO PUBLIC HEALTH

Exposure of rats to high concentrations (1,000 or 1,600 ppm) 6–8 hours/day for 5–14 days resulted in auditory threshold shifts (indicative of hearing loss) and loss of outer hair cells (OHC) in the organ of Corti (Campo et al. 2001; Crofton et al. 1994; Lataye et al. 2003). No alterations in sperm morphology were observed in mice exposed to 300 ppm styrene 5 hours/day for 5 days (Salomaa et al. 1985) and no developmental effects were observed in rats or rabbits exposed to 600 ppm 7 hours/day on gestational days 6–15 or 6–18, respectively, (Murray et al. 1978) or mice exposed to 250 ppm 6 hours/day on gestational days 6–16 (Kankaanpää et al. 1980). An increase in fetal deaths or resorptions was observed in hamsters exposed to 1,000 ppm 6 hours/day on gestational days 6–18 (Kankaanpää et al. 1980).

These data suggest that the nervous system is the most sensitive target of styrene toxicity in humans following acute-duration inhalation exposure. The lowest LOAEL for a relevant end point in humans is 87 ppm for vestibular impairment in subjects exposed to styrene for 1 hour (Ödkvist et al. 1982). A similar LOAEL (80 ppm) was identified for nasal effects in mice exposed to styrene for 3 days (Cruzan et al. 2001). Although nasal irritation has been observed in humans exposed to 376 ppm for 1 hour (Stewart et al. 1968) and focal hyperplasia in the nasal olfactory epithelium was observed in rats exposed to 1,000 ppm (NOAEL of 500 ppm) for 13 weeks (Cruzan et al. 1997), mice appears to be unusually susceptible to this effect. As discussed in Section 2.2, mice appear to have a greater capacity than humans to generate the reactive metabolite, styrene oxide, in the nasal cavity and a lower capacity to detoxify styrene oxide (Green et al. 2001a). Thus, nasal lesions in mice were not considered suitable as the basis of an MRL. The identification of the nervous system as the critical target of toxicity for styrene is supported by a large number of occupational exposure studies. Delays in reaction time have been observed in workers exposed to 21.9–92 ppm (Cherry et al. 1980; Fallas et al. 1992; Gamberale et al. 1976; Jegaden et al. 1993; Mutti et al. 1984a; Tsai and Chen 1996) and vestibular effects have been observed at 18–36 ppm (Calabrese et al. 1996; Möller et al. 1990; Toppila et al. 2006).

The Ödkvist et al. (1982) study did not identify a NOAEL for vestibular effects; however, a NOAEL of 20 ppm for performance on several tests of reaction time and attention was identified by Seeber et al. (2004) in subjects exposed to styrene for 3 hours. Although there is some uncertainty whether deriving an MRL based on a 3-hour exposure study would be protective of continuous exposure to styrene for 2 weeks, the Seeber et al. (2004) study was selected as the basis of an acute duration inhalation MRL for styrene.

Groups of eight volunteers (gender not reported) were exposed to 0.5 or 20 ppm styrene for 3 hours (Seeber et al. 2004). The subjects were tested for simple reaction time, choice reaction time, and attention

2. RELEVANCE TO PUBLIC HEALTH

prior to exposure initiation, during the third hour of exposure, and 1.5 hours after exposure termination. The subjects were also asked to complete a symptom questionnaire before, during, and after exposure. The mean concentration of styrene in blood was 2.2 and 80 µg/L after 3 hours of exposure to 0.5 and 20 ppm, respectively. The blood levels of styrene were correlated with styrene levels in air ($r_{xy}=0.98$). No significant alterations in performance on neurobehavioral tests were found. An increase in the reporting of breathing problems was found; however, it was not statistically significant and the ranking of the severity of the breathing problem was very low (0.5 on a scale of 5).

The NOAEL of 20 ppm from the Seeber et al. (2004) study was divided by an uncertainty factor of 10 to account for human variability resulting in an acute-duration inhalation MRL of 2 ppm.

Intermediate-duration Inhalation MRL

No human intermediate-duration studies were identified. Animal studies examining systemic, neurological, reproductive, and developmental toxicity have identified the respiratory tract as the most sensitive target of toxicity. Atrophy of the olfactory epithelium, hypertrophy/hyperplasia of Bowman's gland has been observed in mice exposed to 50 ppm 6 hours/day, 5 days/week for 13 weeks (Cruzan et al. 1997), decreased nasal cilia activity has been observed in rats exposed to 150 ppm 4 hours/day, 5 days/week for 21 days (Ohashi et al. 1986), and focal hyperplasia has been observed in rats exposed to 1,000 ppm 6 hours/day, 5 days/week for 13 weeks (Cruzan et al. 1997). As discussed previously, the mouse does not appear to be a good model for nasal effects in humans due metabolic differences. Other systemic effects that have been observed include eye irritation in rats exposed to 200 ppm 6 hours/day, 5 days/week for 13 weeks (Cruzan et al. 1997) and centrilobular aggregates of siderophages in the livers of mice exposed to 200 ppm 6 hours/day, 5 days/week for 13 weeks (Cruzan et al. 1997).

A number of studies in rats have reported outer hair cell loss in the organ of Corti in rats exposed to 600–650 ppm for 4 weeks (Loquet et al. 2000; Makitie et al. 2002; Pouyatos et al. 2002) and hearing loss at 750–1,000 ppm for 3–4 weeks (Campo et al. 2001; Lataye et al. 2000, 2001; Loquet et al. 1999, 2000; Pouyatos et al. 2002). A NOAEL of hearing effects of 300 ppm was identified in rats exposed for 12 hours/day, 5 days/week for 4 weeks (Makitie et al. 2002). Other neurological effects include alterations in astroglial cells in rats continuously exposed to 320 ppm for 3 months (Rosengren and Haglid 1989) and decreased sensory nerve conduction velocity in rats exposed to 2,000 ppm 8 hours/day, 5 days/week for 32 weeks (Yamamoto et al. 1997). No reproductive, developmental, or neurodevelopmental effects were observed in a two-generation study (Cruzan et al. 2005a, 2005b); the NOAEL was

2. RELEVANCE TO PUBLIC HEALTH

500 ppm. In contrast, an increase in neonatal deaths, developmental landmark delays, and alterations in neurochemical levels were observed in the offspring of rats exposed 6 hours/day on gestational days 6–20 (Katakura et al. 1999, 2001).

Chronic-duration studies suggest that the most sensitive target of styrene toxicity is the nervous system. It is likely that this would also be the most sensitive effect following intermediate-duration exposure. In the absence of human neurotoxicity data, an intermediate-duration inhalation MRL is not recommended at this time.

Chronic-duration Inhalation MRL

- An MRL of 0.2 ppm has been derived for chronic-duration inhalation exposure (greater than 365 days) to styrene.

A large number of studies have examined the neurotoxicity of styrene in workers at reinforced plastic manufacturing facilities. These studies reported a variety of neurological effects, including decreased color discrimination, slowed reaction time, permanent hearing threshold shifts, vestibular effects, and increases in subjective symptoms. A summary of the results of studies for some of these neurological effects is presented in Table 2-1. The LOAELs for these effects range from 6 to 93 ppm.

Chronic-duration studies in animals identify the nasal olfactory epithelium as the most sensitive end point. Atrophic and/or degenerative changes were observed in rats exposed to 50 ppm styrene 6 hours/day, 5 days/week for 104 weeks (Cruzan et al. 1998) and respiratory metaplasia in the nasal olfactory epithelium has been observed in mice exposed to 20 ppm 6 hours/day, 5 days/week for 98–104 weeks (Cruzan et al. 2001). As noted previously, mice do not appear to be a good model for potential respiratory effects in humans.

Neurotoxicity observed in styrene workers was selected as the basis of the chronic-duration inhalation MRL for styrene. Two approaches to deriving the MRL were considered. In the first approach, the MRL is based on a single study identifying a sensitive LOAEL. In the study by Kishi et al. (2001), styrene workers were divided into three groups based on urinary mandelic acid excretion levels. A significant increase in color confusion index (CCI) was found in the workers exposed to equivalent styrene concentrations of 10 or 46 ppm, as compared to age-matched controls. No significant alterations were observed in workers exposed to 4 ppm. The second approach involves the use of a LOAEL estimated

2. RELEVANCE TO PUBLIC HEALTH

Table 2-1. Results of Selected Human Neurotoxicity Studies

Result	Reference	NOAEL ppm	LOAEL ppm
Decreased color discrimination	Chia et al. 1994		6
	Kishi et al. 2001	4	10
	Gong et al. 2002		10
	Gobba et al. 1991		16
	Iregren et al. 2005		22
	Fallas et al. 1992		24.3
	Campagna et al. 1996		26
	Eguchi et al. 1995	8	93
Neurological symptoms	Flodin et al. 1989	6	
	Edling et al. 1993	8.6	
	Checkoway et al. 1992	10.8	18.9
	Cherry et al. 1980		92
Vestibular effects	Möller et al. 1990		18
	Toppila et al. 2006		24.8
	Calabrese et al. 1996		36
Reaction time	Edling et al. 1993	8.6	
	Tsai and Chen 1996		21.9
	Jegaden et al. 1993		22.68
	Fallas et al. 1992		24.3
	Mutti et al. 1984a		25
	Gamberale et al. 1976		47
	Cherry et al. 1980		92
	Hearing	Morata et al. 2002	
Śliwińska-Kowalska et al. 2003			15.6
Morioka et al. 1999			16
Möller et al. 1990		18	
Calabrese et al. 1996		36	
Nerve conduction velocity		Seppäläinen and Härkönen 1976	30
	Štětkařová et al. 1993		50
	Triebig et al. 1985	100	

2. RELEVANCE TO PUBLIC HEALTH

from a meta-analysis of occupational exposure studies finding effects on color vision and reaction time (Benignus et al. 2005). Benignus et al. (2005) used data color vision impairment data from the Campagna et al. (1996), Eguchi et al. (1995), Gobba et al. (1991), Gong et al. (2002), and Kishi et al. (2001) studies and choice reaction time data from the Jegaden et al. (1993), Mutti et al. (1984a), Triebig et al. (1989), and Tsai and Chen (1996) studies. Average styrene exposure concentrations were estimated from individual data reported in the papers; for studies reporting individual data as urinary mandelic acid levels, Benignus et al. (2005) used standardized methods for converting to styrene exposure levels. Cumulative styrene exposure was estimated by multiplying exposure level by length of employment. A common metric of effect magnitude (percentage of baseline) was calculated for the different neurological effects. The analysis found a significant linear relationship between choice reaction time and cumulative styrene exposure; cumulative exposure accounted for 91% of the variance in reaction time. Similarly, a significant relationship between CCI and cumulative styrene exposure was found, with cumulative exposure accounting for 35% of the variance in CCI. Using the regression equations for these two effects, Benignus et al. (2005) estimated that exposure to 150 ppm for 8 work-years would result in a 50% increase in choice reaction time and a 17% increase in CCI score; exposure to 20 ppm for 8 work-years would result in a 6.5% increase in choice reaction time and a 2.23% increase in CCI score. As discussed in Benignus et al. (2005), a 7% decrease in reaction time would prevent 58,000–70,000 injuries per year from automobile accidents. The investigators also noted that CCI increases with age, the rate of increase is about 10% per 13 years of age; thus, a 2.23% decrease in color perception would be roughly equivalent to 2.9 additional years of age. Based on this analysis, 20 ppm is considered a LOAEL for neurological effects.

The LOAEL of 20 ppm is consistent with the LOAEL values identified in many of the individual studies. However, using the LOAEL identified from the Benignus et al. (2005) meta-analysis has several advantages over selecting a single study as the basis of the MRL. Because data were pooled from several studies, the relationships between styrene exposure and effects were examined in a large number of subjects (302 subjects for choice reaction time and 383 subjects for color vision). The use of standardized methods for estimating styrene exposure levels from urinary biomarker levels is also an advantage. Additionally, the biological relevance of the observed deficits in reaction time and color vision was estimated. The LOAEL of 20 ppm was divided by an uncertainty factor of 100 (10 for use of a LOAEL and 10 for human variability) resulting in a chronic-duration inhalation MRL of 0.2 ppm.

2. RELEVANCE TO PUBLIC HEALTH

Oral MRLs***Acute-duration Oral MRL***

- An MRL of 0.1 mg/kg/day has been derived for acute-duration oral exposure (14 days or less) to styrene.

A limited number of studies have examined the acute toxicity of orally administered styrene; these studies have examined potential neurotoxicity and developmental toxicity. No developmental effects were observed in rats administered a single dose of 300 mg/kg on gestational day 11 (Daston et al. 1991) or administered 300 mg/kg/day (administered as two daily doses of 150 mg/kg) on gestational days 6–15 (Murray et al. 1978). Impaired learning was observed in rats administered via gavage 100 or 200 mg/kg/day for 14 days; increases in serotonin levels were observed in the hypothalamus, hippocampus, and midbrain (Husain et al. 1985). In another study, increases in dopamine receptor binding was observed in rats administered a single gavage dose of 200 mg/kg (Agrawal et al. 1982).

Although a limited number of toxicity end points have been examined following acute-duration oral exposure, longer-term oral studies examining systemic and reproductive end points have identified LOAELs that were higher than the 100 mg/kg/day LOAEL identified for neurotoxicity in the Husain et al. (1985) study. The lowest LOAEL identified for a systemic effect is 400 mg/kg/day for Heinz body formation in dogs administered styrene by gavage for 561 days (Quast et al. 1979); the NOAEL was 200 mg/kg/day. Decreased spermatozoa counts were observed in adult rats administered 400 mg/kg 6 days/week for 60 days (Srivastava et al. 1989), young rats exposed via lactation on postnatal days 1–21 (maternal dose of 400 mg/kg/day) (Srivastava et al. 1992a), and young rats administered 200 mg/kg 6 days/week on postnatal days 1–61 (Srivastava et al. 1992b); the NOAELs identified in these three studies were 200, 200, and 100 mg/kg, respectively. Marked degeneration of the seminiferous tubules was also observed in the adult rats administered 400 mg/kg (Srivastava et al. 1989). Impaired learning observed in rats administered 500 mg/kg 5 days/week for 8 weeks (no NOAEL identified) (Bushnell 1994) also supports the identification of neurotoxicity as a sensitive end point following oral exposure. Additionally, the extensive inhalation toxicity database for styrene supports the selection of neurotoxicity as the most sensitive target of toxicity; both the acute- and chronic-duration inhalation MRLs are based on neurological effects in humans. Neurological effects observed in chronically exposed styrene workers include decreased color discrimination, slowed reaction time, increased prevalence of neurological symptoms, and ototoxicity (hearing and vestibular effects).

2. RELEVANCE TO PUBLIC HEALTH

The Husain et al. (1985) study was selected as the basis of the acute-duration oral MRL. In this study, groups of 15 male Wistar rats were administered by gavage 0, 100, or 200 mg/kg/day styrene in ground nut oil for 14 consecutive days. Spontaneous motor activity with or without amphetamine induction was observed 1 day after the last dose. Two days after exposure termination, the rats underwent acquisition training for 4 days. Learning was assessed by measuring the number of times the rat climbed the pole after the conditioned stimulus to avoid the foot-shock unconditioned stimulus. Noradrenaline, dopamine, and serotonin levels were measured in seven regions of the brain in six rats/group sacrificed after the acquisition training. No overt signs of toxicity were observed. No significant alterations in locomotor activity were observed with or with amphetamine induction. Significantly greater increases in percent avoidance response in the conditioned avoidance response test (indicative of impaired learning) were observed at 100 and 200 mg/kg/day; no difference was found between the two styrene groups. The effects were observed on test day 3 and 4. Significant increases in the level of serotonin in the hypothalamus (70%), hippocampus (51%), and midbrain (29%) were observed at 200 mg/kg/day. Styrene exposure did not affect brain noradrenaline and dopamine levels.

The LOAEL of 100 mg/kg/day was divided by an uncertainty factor of 1,000 (10 for use of a LOAEL, 10 for extrapolation from animals to humans, and 10 for human variability).

Intermediate-duration Oral MRL

The systemic toxicity of styrene has not been investigated in intermediate-duration oral exposure studies. Neurotoxicity studies have identified a LOAEL of 200 mg/kg/day for increased dopamine receptor binding in rats (Agrawal et al. 1982), a LOAEL of 500 mg/kg (5 days/week) for impaired learning in rats (Bushnell 1994), and a LOAEL of 906 mg/kg/day for alterations in serotonin and noradrenaline levels in rats (Husain et al. 1980); none of these studies identified a NOAEL for neurological effects. An increase in dopamine receptor binding was also observed in the offspring of rats administered 200 mg/kg/day during gestation, lactation, or both (Zaidi et al. 1985). Reproductive and immunological effects were reported in the other intermediate-duration oral studies. Decreases in spermatozoa counts were observed in rats exposed as 400 mg/kg (6 days/week) as adults, 200 mg/kg (6 days/week) as neonates, or during lactation (maternal dose of 400 mg/kg/day) (Srivastava et al. 1989, 1992a, 1992b). Impaired immune function was observed in mice exposed to 30 mg/kg/day and in rats exposed to 294 mg/kg/day (Dogra et al. 1992); the NOAELs were 23 and 196 mg/kg/day.

2. RELEVANCE TO PUBLIC HEALTH

Dogra et al. (1992) identified the lowest LOAEL following intermediate-duration oral exposure to styrene; however, there are limited data to support the identification of the immune system as a sensitive, relevant target for humans. Although, the sensitivity of the nervous system has been firmly established following inhalation and oral exposure, the LOAELs identified in the intermediate-duration studies are higher than the lowest LOAEL for neurotoxicity identified in an acute-duration study (Husain et al. 1985). An MRL based on the Dogra et al. (1992) study would be higher than the acute-duration oral MRL; thus, an intermediate-duration MRL is not recommended at this time.

Chronic-duration Oral MRL

The available data on the chronic toxicity of styrene comes from three systemic toxicity studies. No adverse effects were observed in rats exposed to 35 mg/kg/day styrene in drinking water for 2 years (Beliles et al. 1985) and no liver or kidney alterations were observed in rats administered 500 mg/kg 1 day/week for 120 weeks (Ponomarkov and Tomatis 1978). Increase in Heinz body formation was observed in dogs administered 400 mg/kg/day for 561 days (Quast et al. 1979); the NOAEL for this effect is 200 mg/kg/day.

The chronic-duration inhalation database provides strong evidence that neurotoxicity is the most sensitive target of styrene toxicity. It is not known if this would also be true for chronic-duration oral exposure; the acute-toxicity oral database provides suggestive evidence that it would be a sensitive target. In the absence of a long-term oral study examining neurological end points, a chronic-duration oral MRL is not recommended.

2. RELEVANCE TO PUBLIC HEALTH

This page is intentionally blank.

3. HEALTH EFFECTS

3.1 INTRODUCTION

The primary purpose of this chapter is to provide public health officials, physicians, toxicologists, and other interested individuals and groups with an overall perspective on the toxicology of styrene. It contains descriptions and evaluations of toxicological studies and epidemiological investigations and provides conclusions, where possible, on the relevance of toxicity and toxicokinetic data to public health.

A glossary and list of acronyms, abbreviations, and symbols can be found at the end of this profile.

3.2 DISCUSSION OF HEALTH EFFECTS BY ROUTE OF EXPOSURE

To help public health professionals and others address the needs of persons living or working near hazardous waste sites, the information in this section is organized first by route of exposure (inhalation, oral, and dermal) and then by health effect (death, systemic, immunological, neurological, reproductive, developmental, genotoxic, and carcinogenic effects). These data are discussed in terms of three exposure periods: acute (14 days or less), intermediate (15–364 days), and chronic (365 days or more).

Levels of significant exposure for each route and duration are presented in tables and illustrated in figures. The points in the figures showing no-observed-adverse-effect levels (NOAELs) or lowest-observed-adverse-effect levels (LOAELs) reflect the actual doses (levels of exposure) used in the studies. LOAELs have been classified into "less serious" or "serious" effects. "Serious" effects are those that evoke failure in a biological system and can lead to morbidity or mortality (e.g., acute respiratory distress or death). "Less serious" effects are those that are not expected to cause significant dysfunction or death, or those whose significance to the organism is not entirely clear. ATSDR acknowledges that a considerable amount of judgment may be required in establishing whether an end point should be classified as a NOAEL, "less serious" LOAEL, or "serious" LOAEL, and that in some cases, there will be insufficient data to decide whether the effect is indicative of significant dysfunction. However, the Agency has established guidelines and policies that are used to classify these end points. ATSDR believes that there is sufficient merit in this approach to warrant an attempt at distinguishing between "less serious" and "serious" effects. The distinction between "less serious" effects and "serious" effects is considered to be important because it helps the users of the profiles to identify levels of exposure at which major health effects start to appear. LOAELs or NOAELs should also help in determining whether or not

3. HEALTH EFFECTS

the effects vary with dose and/or duration, and place into perspective the possible significance of these effects to human health.

The significance of the exposure levels shown in the Levels of Significant Exposure (LSE) tables and figures may differ depending on the user's perspective. Public health officials and others concerned with appropriate actions to take at hazardous waste sites may want information on levels of exposure associated with more subtle effects in humans or animals (LOAELs) or exposure levels below which no adverse effects (NOAELs) have been observed. Estimates of levels posing minimal risk to humans Minimal Risk Levels or (MRLs) may be of interest to health professionals and citizens alike.

A User's Guide has been provided at the end of this profile (see Appendix B). This guide should aid in the interpretation of the tables and figures for Levels of Significant Exposure and the MRLs.

3.2.1 Inhalation Exposure

Most information on the effects of inhalation exposure to styrene in humans comes from studies of workers exposed to styrene vapors in the production and use of plastics and resins, especially polyester resins dissolved in styrene. In most cases, the studies involve workplace exposures such as fiberglass boat building factories where the actual levels of styrene are reported as a range of styrene air concentrations. However, there are a few human clinical studies in which exposures are better quantified. A common limitation of many of the occupational exposure studies is the phenomenon of the healthy worker effect. The selection of healthy individuals for employment and the likelihood that more susceptible workers are more likely to leave the workforce can result in workers who are healthier than the general population. This type of bias typically affects comparisons with the general population and is less likely to influence comparisons with other groups of workers. Provided below are descriptions of the known effects of inhalation exposure of humans and animals to styrene.

3.2.1.1 Death

There have been no reports of deaths in humans directly associated with exposure to styrene in the workplace (EPA 1985a; Gosselin et al. 1984; NIOSH 1983).

In animals, inhalation studies indicate that the acute toxicity of styrene is low to moderate. An LC₅₀ of 2,770 ppm after 2 hours of exposure was reported in rats, and the LC₅₀ for mice after exposure for 4 hours was 4,940 ppm (Shugaev 1969). All rats and guinea pigs survived after exposure to 1,300 ppm styrene

3. HEALTH EFFECTS

for 30 hours and 16 hours, respectively (Spencer et al. 1942). However, all animals died after 40 hours of exposure. Gender differences in mortality were observed in repeated-exposure studies (6 hours/day, 5 days/week for 2 weeks) (Cruzan et al. 1997). Increases in mortality were observed in female CD-1 and B6C3F1 female mice exposed to 250 ppm; no deaths were observed at 500 ppm. In the CD-1 and B6C3F1 males, very few deaths were observed at 250 ppm, but increases in deaths were observed at 500 ppm. A similar finding was reported by Morgan et al. (1993a): increases in mortality were observed in female B6C3F1 mice exposed to 250 ppm and no deaths were observed at 500 ppm; in males, deaths were observed at 250 and 500 ppm. In contrast to these findings, no deaths were observed in Sprague Dawley rats exposed to concentrations as high as 1,500 ppm 6 hours/day, 5 days/week for 13 weeks (Cruzan et al. 1997).

All reliable LOAEL values and LC₅₀ values for lethality in each species and duration category are recorded in Table 3-1 and plotted in Figure 3-1.

3.2.1.2 Systemic Effects

No studies were located regarding dermal or metabolic effects in humans or animals after inhalation exposure to styrene.

For the following systemic effects resulting from inhalation exposure to styrene, the highest NOAEL values and all reliable LOAEL values in each species and duration category are recorded in Table 3-1 and plotted in Figure 3-1.

Respiratory Effects. Several human studies have examined the respiratory effects caused by inhalation exposure to styrene. The most commonly reported general symptom is mucous membrane irritation. Irritation of the upper respiratory tract (i.e., nose and throat) has been reported by volunteers (Carpenter et al. 1944; Stewart et al. 1968) and workers (NIOSH 1983). Throat irritation and increased nasal secretion occurred following exposure of two male subjects to 800 ppm for 4 hours (Carpenter et al. 1944). Nasal irritation was observed in all volunteers after exposure to 376 ppm styrene for 60 minutes (Stewart et al. 1968). Obstructive lung changes were observed in 4 of 21 workers exposed to styrene for about 10 years (Chmielewski and Renke 1975). However, exposure levels were not defined. No histological alterations were observed in nasal biopsies from styrene workers exposed to 50–60 ppm styrene for 7 years (Ödkvist et al. 1985).

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
ACUTE EXPOSURE								
Death								
1	Rat	3-40 hr				1300	(100% mortality after >40 hours exposure)	Spencer et al. 1942 Styrene
2	Mouse (CD-1 and B6C3F1)	6 hr/d 5 d/wk 2 wk				250	(increased mortality)	Cruzan et al. 1997 Styrene
3	Mouse (B6C3F1)	6 hr/d 14 d				250 M	(44% mortality)	Morgan et al. 1993a Styrene
4	Gn Pig	3-40 hr				1300	(100% mortality after 40 hours exposure)	Spencer et al. 1942 Styrene
Systemic								
5	Human	7 hr	Ocular		99 M (mild, transient eye irritation)			Stewart et al. 1968 Styrene
6	Human	1 or 2 hr	Resp	216 M	376 M (nasal irritation)			Stewart et al. 1968 Styrene
			Ocular	216 M	376 M (eye irritation)			

DRAFT FOR PUBLIC COMMENT

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
7	Mouse (CD-1 and B6C3F1)	6 hr/d 5 d/wk 2 wk	Resp		250	(shallow breathing)	Cruzan et al. 1997 Styrene	
			Hepatic	60	250	(increased liver weight and centrilobular hepatocyte necrosis)		
8	Mouse (CD-1)	6 hr/d 3 d	Resp		80 M	(single cell necrosis in nasal olfactory epithelium)	Cruzan et al. 2001 Styrene	
9	Mouse (CD-1)	6 hours/day 3 days	Resp	40 M	160 M	(moderate to marked degenerative changes in olfactory epithelium)	Green et al. 2001a Styrene	
10	Mouse (B6C3F1)	6 hr/d 14 d	Hepatic	125	250 M	(pigmented macrophages and focal necrosis)	Morgan et al. 1993a Styrene	
			Renal	500				
			Bd Wt	500				
11	Mouse (B6C3F1)	6 hr/d 1-3 d	Hepatic			250 M (mild to marked necrosis)	Morgan et al. 1993a Styrene	

DRAFT FOR PUBLIC COMMENT

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
12	Mouse (B6C3F1)	6 hr/d 3 d	Hepatic	125		250 (severe hepatocellular degeneration/necrosis)	Morgan et al. 1993b Styrene	
13	Mouse	6 hr/d 4 d	Hepatic	125		250 (marked degeneration and/or coagulative necrosis of centrilobular hepatocytes)	Morgan et al. 1993c Styrene	
Immuno/ Lymphoret								
14	Mouse (BALB/c)	6 hr/d 4 d			100 F (exacerbated inflammatory reaction after ovalbumin challenge)		Ban et al. 2006 Styrene	
Neurological								
15	Human	1 hr			87 (inhibition of vestibular-oculomotor system)		Odkvist et al. 1982 Styrene	
16	Human	3 or 4 hr		20 ^b			Seeber et al. 2004 Styrene	
17	Human	1 or 2 hr		216 M	376 M (impaired performance on balance and coordination tests)		Stewart et al. 1968 Styrene	

DRAFT FOR PUBLIC COMMENT

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
18	Human	7 hr		99 M			Stewart et al. 1968 Styrene	
19	Rat (Long- Evans)	6 hr/d 5 d/wk 1 or 2 wk				1000 M (hearing loss and loss of OHC)	Campo et al. 2001 Styrene	
20	Rat (Long- Evans)	8 hr/d 5 d				1600 M (hearing loss at 8 and 16 kHz)	Crofton et al. 1994 Styrene	
21	Rat (Long- Evans)	6 hr/d 5 d				1000 M (hearing loss, loss of OHC)	Lataye et al. 2003 Styrene	
22	Mouse (CD-1 and B6C3F1)	6 hr/d 5 d/wk 2 wk		60		250 (lethargy and unsteady gait)	Cruzan et al. 1997 Styrene	
23	Mouse (Swiss OF1)	4 hr		413 M	610 M (impaired performance on a swimming test)		De Ceaurriz et al. 1983 Styrene	
24	Gn Pig (NS)	6 hr/d 5 d		1000 M			Lataye et al. 2003 Styrene	
Reproductive 25	Mouse	5 d 5 hr/d		300 M			Salomaa et al. 1985 Styrene	

DRAFT FOR PUBLIC COMMENT

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure	Species ^a (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
Developmental								
26	Rat (Sprague-Dawley)	7 hr/d Gd 6-15		600 F			Murray et al. 1978 Styrene	
27	Mouse (BMR/T6T6)	6 hr/d Gd 6-16		250 F			Kankaanpaa et al. 1980 Styrene	
28	Hamster (Chinese)	6 hr/d Gd 6-18		750 F		1000 F (fetal deaths or resorptions)	Kankaanpaa et al. 1980 Styrene	
29	Rabbit (New Zealand)	7 hr/d Gd 6-18		600 F			Murray et al. 1978 Styrene	
INTERMEDIATE EXPOSURE								
Systemic								
30	Rat (Sprague-Dawley)	6 hr/d 5 d/wk 13 wk	Resp	500	1000	(focal hyperplasia in nasal olfactory epithelium)	Cruzan et al. 1997 Styrene	
			Hemato	1500				
			Hepatic	1500				
			Renal	1500				
			Ocular		200	(eye irritation)		
			Bd Wt	1000 M	1500 M	(10% decrease in body weight gain)		

DRAFT FOR PUBLIC COMMENT

STYRENE

3. HEALTH EFFECTS

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
31	Rat (Long- Evans)	6 hr/d 5 d/wk 4 wk	Hepatic	750 M			Loquet et al. 2000 Styrene	Urinary markers of renal toxicity and serum markers of liver toxicity.
			Renal	750 M				
32	Rat	21 d 5 d/wk 4 hr/d	Resp		150 M (decreased nasal cilia activity)	1000 M (disabled nasal cilia activity)	Ohashi et al. 1986 Styrene	
33	Rat (Sprague-Dawley)	13 wk 5 d/wk 7 hr/d	Renal	133			Viau et al. 1987 Styrene	
34	Rat (Wistar)	8 hr/d 5 d/wk 32 wk	Bd Wt	200 M	2000 M (>15% decrease in body weight gain)		Yamamoto et al. 1997 Styrene	
35	Rat (Fischer- 344)	14 hr/d 5 d/wk 3 wk	Bd Wt		800 M (10-13.5% decrease in body weight)		Yano et al. 1992 Styrene	

DRAFT FOR PUBLIC COMMENT

STYRENE

3. HEALTH EFFECTS

31

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure	Species ^a (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
36	Mouse (CD-1)	6 hr/d 5 d/wk 13 wk	Resp		50	(atrophy of olfactory epithelium, dilatation, hypertrophy, hyperplasia of Bowman's gland; decreased eosinophilia of bronchiolar epithelial cells)		Cruzan et al. 1997 Styrene
			Hemato	200				
			Hepatic	100 F	150 F	(centrilobular aggregates of siderophages)		
37	Pig	3 wk 5 d/wk 6 hr/d	Bd Wt	150 M	200 M	(decreased body weight gain)		Johnston et al. 1983 Styrene
			Hemato	360				
Neurological								
38	Rat (Long- Evans)	6 hr/d 5 d/wk 3 or 4 wk				1000 M (hearing loss, loss of OHC in organ of Corti)		Campo et al. 2001 Styrene
39	Rat (Long- Evans)	6 hr/d 5 d/wk 4 wk				750 M (hearing loss and loss of OHC)		Lataye et al. 2000 Styrene

DRAFT FOR PUBLIC COMMENT

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure	Species ^a (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
40	Rat (Long- Evans)	6 hr/d 5 d/wk 4 wk					1000 M (loss of OHC and spiral ganglion cell density in organ of Corti)	Lataye et al. 2001 Styrene
41	Rat (Long- Evans)	6 hr/d 5 d/wk 4 wk			650 M (OHC loss)		850 M (hearing loss and loss of OHC)	Loquet et al. 1999 Styrene
42	Rat (Long- Evans)	6 hr/d 5 d/wk 4 wk					750 M (hearing loss and loss of OHC)	Loquet et al. 2000 Styrene
43	Rat (Wistar)	12 hr/d 5 d/wk 4 wk		300 M	600 M (hearing impairment and loss of OHC)			Makitie et al. 2002 Styrene
44	Rat (Long- Evans)	6 hr/d 5 d/wk 4 wk			650 M (OHC loss)		750 M (hearing loss and OHC loss)	Pouyatos et al. 2002 Styrene
45	Rat (Fischer- 344)	3 wk 14 hr/d					800 M (hearing loss)	Pryor et al. 1987 Styrene
46	Rat (Sprague- Dawley)	3 mo continuous		90 M	320 M (astroglial alterations)			Rosengren and Haglid 1989 Styrene

DRAFT FOR PUBLIC COMMENT

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
47	Rat (Wistar)	8 hr/d 5 d/wk 32 wk		200 M	2000 M (decreased sensory nerve conduction velocity)		Yamamoto et al. 1997 Styrene	
48	Rat (Fischer- 344)	14 hr/d 5 d/wk 3 wk				800 M (hearing loss and loss of OHC in organ of Corti)	Yano et al. 1992 Styrene	
49	Mouse (CD-1)	6 hr/d 5 d/wk 13 wk		50	100 (atrophy of olfactory nerve fibers)	200 F (transient lethargy, cold to touch, and slow respiration)	Cruzan et al. 1997 Styrene	
Reproductive								
50	Rat (CD)	6 hr/d 70 pmd 14 d mating Gd 0-21, Ld 5-21		500			Cruzan et al. 2005a, 2005b Styrene	
Developmental								
51	Rat (CD)	6 hr/d 70 pmd 14 d mating Gd 0-21, Ld 5-21		500			Cruzan et al. 2005a, 2005b Styrene	
52	Rat (Wistar)	6 hr/d Gd 6-20				300 (increased neonatal deaths, delays in righting reflex and incisor eruption, alterations in neurochemical levels)	Katakura et al. 1999, 2001 Styrene	

DRAFT FOR PUBLIC COMMENT

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
CHRONIC EXPOSURE								
Systemic								
53	Human	5.1 yr (Occup)	Hepatic	40 F			Harkonen et al. 1984 Styrene	
54	Human	occupational exposure 7 years	Resp	46 M			Odkvist et al. 1985 Styrene	
55	Human	12.6 yr (Occup)	Renal	26			Verplanke et al. 1998 Styrene	
56	Rat (Sprague-Dawley)	6 hr/d 5 d/wk 104 wk	Resp		50	(atrophic and/or degenerative changes in nasal olfactory epithelium)	Cruzan et al. 1998 Styrene	
			Cardio	1000				
			Gastro	1000				
			Hemato	1000				
			Hepatic	1000				
			Renal	1000				
			Ocular	1000				
			Bd Wt	200 F	500 F	(decreased body weight gain)		

DRAFT FOR PUBLIC COMMENT

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
57	Mouse (CD-1)	6 hr/d 5 d/wk 98-104 wk	Resp		20	(respiratory metaplasia in nasal olfactory epithelium, bronchiolar epithelial hyperplasia)		Cruzan et al. 2001 Styrene
			Cardio	160				
			Gastro	160				
			Hemato	160				
			Hepatic	160				
			Renal	160				
			Ocular	160				
			Bd Wt	80 M	160 M (11% decrease in body weight gain)			
Immuno/ Lymphoret								
58	Human	7 yr (Occup)			30	(alterations in lymphocyte subsets)		Bergamaschi et al. 1995b Styrene
59	Human	13 yr (Occup)			26	(impaired immune response to concanavalin A)		Tulinska et al. 2000 Styrene
Neurological								
60	Human	(Occup)			20 ^c	(decreased color discrimination and reaction time)		Benignus et al. 2005 Styrene

DRAFT FOR PUBLIC COMMENT

STYRENE

3. HEALTH EFFECTS

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
61	Human	7.6 yr (Occup)			36	(altered performance vestibular tests)	Calabrese et al. 1996 Styrene	
62	Human	62.5 or 79.3 mo (Occup)			26	(decreased color discrimination)	Campagna et al. 1996 Styrene	
63	Human	4-6.4 yr (Occup)		10.8	18.9	(increased prevalence of neurological symptoms)	Checkoway et al. 1992 Styrene	
64	Human	(Occup)			92 M	(tiredness, slow reaction times, mood changes)	Cherry et al. 1980 Styrene	
65	Human	18.8 yr (Occup)			6 M	(decreased color discrimination and performance on neurobehavioral tests)	Chia et al. 1994 Styrene	
66	Human	12.5 yr (Occup)		24.6			Dalton et al. 2003 Styrene	
67	Human	9 yr (Occup)			8.6 M	(increased symptoms of neurotoxicity)	Edling et al. 1993 Styrene	

DRAFT FOR PUBLIC COMMENT

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
68	Human	7.0 yr (Occup)		8 M	93 M (decreased color discrimination)		Eguchi et al. 1995 Styrene	
69	Human	6.5 yr (Occup)			24.3 M (decreased color discrimination)		Fallas et al. 1992 Styrene	
70	Human	2.7 yr (Occup)			47 M (slowed reaction time)		Gamberale et al. 1976 Styrene	
71	Human	(Occup)			16 (decreased color discrimination)		Gobba et al. 1991 Styrene	
72	Human	76.6 mo (Occup)			10 M (decreased color discrimination)		Gong et al. 2002 Styrene	
73	Human	12.9-17.8 yr (Occup)			22 M (decreased color discrimination)		Iregren et al. 2005 Styrene	
74	Human	5 yr (Occup)			22.68 (impaired performance on neurobehavioral tests of visual reaction and memory)		Jegaden et al. 1993 Styrene	

DRAFT FOR PUBLIC COMMENT

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
75	Human	6.2 yr (Occup)		4	10	(decreased color discrimination)	Kishi et al. 2001 Styrene	
76	Human	4.9 yr (Occup)			75 M	(impaired performance on visuomotor accuracy and psychomotor performance tests)	Lindstrom et al. 1976 Styrene	
77	Human	10.8 yr (Occup)			18 M	(impaired vestibular function)	Moller et al. 1990 Styrene	
78	Human	17 yr (Occup)		3.68			Morata et al. 2002 Styrene	
79	Human	9.4 yr (Occup)			16 M	(reduction in upper limit of hearing)	Morioka et al. 1999 Styrene	
80	Human	5 yr (Occup)			22 M	(slowed distribution of nerve conduction velocities and ECG R-R interval)	Murata et al. 1991 Styrene	
81	Human	8.6 yr (Occup)			25	(decreased verbal learning skills)	Mutti et al. 1984a Styrene	

DRAFT FOR PUBLIC COMMENT

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
82	Human	5 yr (Occup)			30 M (EEG abnormalities)		Seppalainen and Harkonen 1976 Styrene	
83	Human	(Occup)			15.6 (hearing loss)		Sliwinska-Kowalska et al. 2003 Styrene	
84	Human	11 yr (Occup)			50 F (decreased peripheral nerve conduction velocity and prolonged latency of somatosensory evoked potentials)		Stetkarova et al. 1993 Styrene	
85	Human	(Occup)			24.8 M (impaired postural stability)		Toppila et al. 2006 Styrene	
86	Human	4 yr		100 M			Triebig et al. 1985 Styrene	Measured nerve conduction velocity.
87	Human	8.3 yr (Occup)			21.9 (impaired performance on neurobehavioral tests)		Tsai and Chen 1996 Styrene	
Developmental 88	Human	>1 yr 7 d/wk 8 hr/d		82 F			Lemasters et al. 1989 Styrene	

DRAFT FOR PUBLIC COMMENT

STYRENE

3. HEALTH EFFECTS

Table 3-1 Levels of Significant Exposure to Styrene - Inhalation

(continued)

Key to Figure	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (ppm)	LOAEL		Reference Chemical Form	Comments
					Less Serious (ppm)	Serious (ppm)		
Cancer								
89	Mouse (CD-1)	6 hr/d 5 d/wk 98-104 wk					160 F (CEL: bronchioloalveolar carcinoma)	Cruzan et al. 2001 Styrene

a The number corresponds to entries in Figure 3-1.


b The acute-duration inhalation MRL of 2 ppm; concentration divided by an uncertainty factor of 10 to account for human variability.

c The chronic-duration inhalation MRL of 0.2 ppm was calculated from a LOAEL of 20 ppm identified in meta-analysis (Benignus et al. 2005) of occupational exposure studies reporting significant alterations in color vision and choice reaction time and an uncertainty factor of 100 (10 for the use of a LOAEL and 10 for human variability).

Bd Wt = body weight; Cardio = cardiovascular; CEL = cancer effect level; d = day(s); ECG = electrocardiographic; F = Female; Gastro = gastrointestinal; Gd = gestational day; Gn pig = guinea pig; Hemato = hematological; hr = hour(s); kHz = kilohertz; LOAEL = lowest-observed-adverse-effect level; M = male; min = minute(s); mo = month(s); NOAEL = no-observed-adverse-effect level; NS = not specified; occup = occupational; OHC = outer hair cell(s); pmd = pre-mating day; ppm = parts per million; Resp = respiratory; x = time(s); wk = week(s); yr = year(s)

DRAFT FOR PUBLIC COMMENT

Figure 3-1 Levels of Significant Exposure to Styrene - Inhalation
Acute (≤14 days)



DRAFT FOR PUBLIC COMMENT

Figure 3-1 Levels of Significant Exposure to Styrene - Inhalation (Continued)
Intermediate (15-364 days)

DRAFT FOR PUBLIC COMMENT




Figure 3-1 Levels of Significant Exposure to Styrene - Inhalation (Continued)

Chronic (≥365 days)

DRAFT FOR PUBLIC COMMENT





Figure 3-1 Levels of Significant Exposure to Styrene - Inhalation (Continued)
 Chronic (≥365 days)

DRAFT FOR PUBLIC COMMENT



3. HEALTH EFFECTS

In rats exposed to 150 ppm styrene 4 hours/day, 5 days/week for 3 weeks a decrease in nasal ciliar activity was observed; at 1,000 ppm, the nasal cilia activity was considered disabled (Ohashi et al. 1986). Electron microscopic examination of the nasal cavity of rats exposed to 1,000 ppm revealed very few ciliated cells and severe degeneration with marked vacuolization. Decreases in cilia activity were also observed in the trachea at 150 and 1,000 ppm. After a 12-week recovery period, nasal and tracheal cilia activity in the 150 ppm group was similar to controls; decreases in nasal cilia activity in the 1,000 ppm group was still lower than controls but was increased compared to rats killed at the end of the exposure period (Ohashi et al. 1986). In a longer-term study, focal hyperplasia was observed in the nasal olfactory epithelium of rats exposed to 1,000 ppm for 13 weeks (Cruzan et al. 1997); at 500 ppm, no histological alterations were observed in the respiratory tract. Chronic exposure to 50 ppm resulted in atrophic and/or degenerative changes in the nasal olfactory epithelium (Cruzan et al. 1998).

Mice appear to be more sensitive than rats to the respiratory toxicity of styrene. Exposure to 50 ppm styrene for 13 weeks resulted in atrophy of the nasal olfactory epithelium and dilatation, hypertrophy and hyperplasia of Bowman's gland (Cruzan et al. 1997). At 100 ppm, atrophy of the nasal olfactory nerve fibers was observed; focal crowding of nonciliated epithelial cells in the bronchioles were observed at 150 ppm. Chronic exposure resulted in respiratory metaplasia of the nasal olfactory epithelium and dilatation, respiratory metaplasia, epithelial hyperplasia of the Bowman's gland in mice exposed to 20 ppm and higher for 2 years (Cruzan et al. 2001). Decreased eosinophilia of epithelial cells and bronchiolar epithelial hyperplasia were observed in the lungs of mice exposed to 20 ppm and higher.

A study by Spencer et al. (1942) also provides some information on species differences. Rats and guinea pigs exposed 1,300 ppm for 7–8 hours/day, 5 days/week for 6 months showed nasal irritation, but rabbits and monkeys did not (Spencer et al. 1942). Histopathological examinations revealed no changes between test and control rats, but pronounced lung irritation was seen in guinea pigs that died after a few exposures. The irritation, which included congestion, hemorrhages, edema, exudation, and a general acute inflammatory reaction, was not seen in the guinea pigs, rabbits, and monkeys that survived the 6-month exposure period (Spencer et al. 1942).

Green et al. (2001a) suggest that the observed species differences between mice and rats are due to differences in styrene metabolism in the nasal epithelium. The rates of metabolism of styrene by cytochromes P-450 CYP2E1 and CYP2F2 to styrene oxide was similar for the two species. However, styrene oxide is more efficiently metabolized by epoxide hydrolases and glutathione S-transferases in rats than in mice. Thus, the higher levels of the reactive epoxide styrene oxide in mice is the likely cause of

3. HEALTH EFFECTS

the increased sensitivity in this species. In *in vitro* assays in fresh human nasal tissues, styrene oxide was not detected and high levels of epoxide hydrolases were detected, suggesting that humans have limited capacity to metabolize styrene in the nasal cavity and a high potential to detoxify styrene oxide. These data suggest that rodents may not be a good model for nasal toxicity in humans.

These well-conducted human and animal studies demonstrate the characteristic irritant properties of styrene on the upper respiratory tract.

Cardiovascular Effects. No studies were located regarding cardiovascular effects in humans after inhalation exposure to styrene.

No cardiovascular effects were observed in rats or mice exposed to concentrations as high as 1,000 ppm or 160 ppm, respectively, for 2 years (Cruzan et al. 1998, 2001).

Gastrointestinal Effects. Nausea was observed in humans exposed to 376 ppm styrene after 1 hour of exposure (Stewart et al. 1968). This effect is probably secondary to effects on the central nervous system, although mucociliary transport of styrene aerosol droplets from the upper respiratory tract to the gastrointestinal tract might also contribute to gastrointestinal irritation. A Russian study (Basirov 1975) reviewed by the World Health Organization (WHO 1983) investigated the effects of styrene on digestive function by testing the secretory, excretory, motor, and pepsinogen-generating functions of the stomach in 20 unexposed and 80 exposed workers. The authors reported that some workers in the styrene-butadiene synthetic rubber manufacture exposed to 60–130 mg/m³ (14–31 ppm) styrene had decreased digestive function and decreased stomach acidity.

No histological alterations were observed in the stomach or intestines of rats exposed to 1,000 ppm (Cruzan et al. 1998) or mice exposed to 160 ppm (Cruzan et al. 2001) styrene for 2 years.

Hematological Effects. Several studies indicate that inhalation exposure of humans to styrene cause mild or no effects on the blood. In one study, the incidence of abnormal values for hematological parameters including erythrocyte, leukocyte, and platelet counts, and hemoglobin levels for 84 styrene workers generally exposed to <1 ppm styrene for 1–36 years was investigated. However, these workers were also exposed to intermittent high levels of styrene as well as to other chemicals. The percentages of the exposed group with abnormally low hemoglobin and erythrocyte values or abnormally high leukocyte values were less than those percentages in the 62-person control group. There were no abnormal

3. HEALTH EFFECTS

thrombocyte values reported in either the exposed or control groups (Thiess and Friedheim 1978). Findings from a group of 93 workers engaged in the manufacture of styrene polymers and exposed to generally <1 ppm styrene for 1–38 years were also presented in this study; only the incidence of abnormally low erythrocyte counts (in the group exposed to styrene) was found to be statistically significant ($p \leq 0.05$). However, because exposures could not be determined accurately and because there were concomitant exposures to other chemicals, the results of these studies are difficult to interpret.

Lowered erythrocyte counts, hemoglobin, platelets, and neutrophils and slightly higher mean corpuscular red cell volumes and neutrophil band counts were observed in workers in a styrene-butadiene rubber manufacturing plant (Checkoway and Williams 1982). The highest mean styrene level was 13.67 ppm. However, interpretation of this study is limited because multiple-chemical exposures were involved and exposure and clinical signs were measured at the same time and only once. An earlier study of styrene workers showed no definite pattern of hematological changes (Lorimer et al. 1978). In these studies, exposure levels were uncertain and multiple chemicals were involved.

In rats exposed to 49 ppm styrene, erythrocyte-aminolevulinate dehydratase (ALA-D) was depressed markedly. The decrease in enzyme activity was accompanied by a decrease in the enzyme content in bone marrow cells (Fujita et al. 1987). The investigators suggested that the changes may have been a result of styrene oxide reducing the enzyme protein is based on *in vitro* data. No hematological alterations were observed in 2-year studies in rats (Cruzan et al. 1998) and mice (Cruzan et al. 2001) exposed to concentrations as high as 1,000 or 160 ppm, respectively.

The well-conducted Thiess and Friedheim (1978) study as well as the more limited studies indicate that few adverse hematological effects occurred in styrene-exposed workers. However, the full meaning of the findings is not clear because of poor characterization of the exposure level and concurrent exposures to other chemicals.

Musculoskeletal Effects. No studies were located regarding musculoskeletal effects in humans after inhalation exposure to styrene.

No histological alterations were observed in skeletal muscle or bone of rats exposed to 1,000 ppm (Cruzan et al. 1998) or mice exposed to 160 ppm (Cruzan et al. 2001) for 2 years.

3. HEALTH EFFECTS

Hepatic Effects. Human studies on the hepatic effects of styrene inhalation frequently used serum levels of enzymes as indicators of liver dysfunction. In general, human studies have resulted in negative or equivocal results (Härkönen et al. 1984; Hotz et al. 1980; Lorimer et al. 1978; Thiess and Friedheim 1978). No significant alterations in alanine aminotransferase, aspartate aminotransferase, or γ -glutamyl transferase levels were observed in workers exposed to generally <1 ppm for 1–36 years (Thiess and Friedheim 1978) or 50–120 ppm for 5.1 years (Härkönen et al. 1984). A significant increase in γ -glutamyl transferase levels was observed in workers exposed to 5–20 ppm for up to 20 years; however, no alterations in alanine aminotransferase, aspartate aminotransferase, or alkaline phosphatase levels were observed (Lorimer et al. 1978). Another study of workers (Hotz et al. 1980) found significant correlations between the exposure level (as measured by styrene metabolite concentrations in morning urine) and ornithine carbamoyl transferase, alanine aminotransferase, and γ -glutamyl transferase levels. Among workers exposed to 50–100 ppm, the increases in these enzymes were modest, 67.8, 55, and 64.9% of reference levels.

Animal studies provide evidence that the liver is a target tissue for styrene; however, the hepatotoxicity of styrene in mice is inversely related to the duration of exposure. Hepatic effects have been observed following acute- and intermediate-duration exposure, but not after chronic exposure and the severity of the effects decreases with continuing exposure. Exposure to 250 or 500 ppm for 1–4 days resulted in marked to severe hepatocellular necrosis and degeneration in mice (Morgan et al. 1993a, 1993b, 1993c). The necrosis was characterized as centrilobular coagulative necrosis and was accompanied by pooling of erythrocytes in dilated sinusoids (Morgan et al. 1993a). The necrosis was often observed after a single exposure to 500 ppm or a 2-day exposure to 250 ppm and the severity did not increase with increasing duration (Morgan et al. 1993a). However, continued exposure resulted in regeneration and repair of the initial hepatic damage. After 14 days of exposure, minimal to mild focal necrosis was observed in female mice exposed to 250 ppm and no hepatic effects were observed in male mice exposed to 250 ppm or male and female mice exposed to 500 ppm (Morgan et al. 1993a). Similarly, a 13-week exposure to 200 ppm resulted in focal loss of hepatocytes with siderosis and centrilobular aggregates of siderophages in female mice (Cruzan et al. 1997). No histological alterations were observed in the livers of mice exposed to 160 ppm for 2 years (Cruzan et al. 2001). Strain differences have also been detected in mice. Morgan et al. (1993c) found that B6C3F1 and C57BL/6 mice were more sensitive than DBA/2 mice, which were more sensitive than Swiss mice. The severity scores for hepatocellular degeneration/necrosis following a 4-day exposure to 250 ppm were 3.2–3.5 in B6C3F1 mice, 3.6 in C57BL/6 mice, 2.4–2.9 in DBA/2 mice, and 2.0 in Swiss mice.

3. HEALTH EFFECTS

Rats appear to be less sensitive than mice to styrene-induced hepatotoxicity. No histological alterations were observed in the livers of Sprague Dawley rats exposed to 1,000 ppm styrene for 13 weeks (Cruzan et al. 1997) or 2 years (Cruzan et al. 1998; Jersey et al. 1978). Parenchymal hydropic degeneration, steatosis, and congestion were observed in rats exposed to 300 ppm for 2 weeks (Vainio et al. 1979); the lack of incidence data limits the interpretation of these results.

Renal Effects. Based on the results of occupational exposure studies and animal toxicity studies, the kidney does not appear to be a sensitive target of styrene toxicity. Occupational exposure studies of workers exposed to 24 ppm (Viau et al. 1987), 53 ppm (Vyskocil et al. 1989), or 26 ppm styrene (Verplanke and Herber 1998) did not find significant alterations in urinary levels of β -microglobulin (not examined in Verplanke and Herber 1998 study), retinol-binding protein, or albumin. The Vyskocil et al. (1989) study also found no significant alterations in total protein, glucose, lysozyme, lactate dehydrogenase, or β -N-acetyl-D-glucosaminidase levels and Verplanke and Herber (1998) did not find alterations in β -galactosidase, N-acetyl- β -D-glucosaminidase, or alanine aminopeptidase. No histological alterations were observed in the kidneys following acute exposure of rats to 300 ppm (Vainio et al. 1979) or mice to 500 ppm (Morgan et al. 1993a), intermediate exposure of rats to 133–1,500 ppm (Cruzan et al. 1997; Spencer et al. 1942; Viau et al. 1987), or chronic exposure of rats to 1,000 ppm (Cruzan et al. 1998) or mice to 160 ppm (Cruzan et al. 2001). Additionally, no alterations in urinary levels of N-acetyl-D-glucosaminidase, γ -glutamyl transpeptidase, protein, or urea were observed in rats exposed to 500 ppm for 4 weeks (Loquet et al. 2000).

Endocrine Effects. Several occupational studies have examined potential endocrine effects in reinforced plastics industry workers exposed to styrene. Significant increases in serum prolactin levels were observed in male and female workers (Bergamaschi et al. 1996, 1997; Luderer et al. 2004; Mutti et al. 1984b). The serum prolactin levels significantly correlated with urinary metabolite (mandelic acid plus phenylglyoxylic acid) levels (Mutti et al. 1984b) and blood styrene levels (Luderer et al. 2004). Based on a logistic regression model, Luderer et al. (2004) estimated that workers exposed to styrene exposures >20 ppm would be more likely to have elevated serum prolactin levels than workers exposed to lower levels; a 10-fold increase in blood styrene concentrations would result in a 2.06-fold increase in serum prolactin levels. Similarly, Arfini et al. (1987) found that female styrene workers had an abnormal response to an intravenous dose of thyrotropin-releasing hormone; the levels of serum prolactin were significantly higher following exposure to thyrotropin-releasing hormone, as compared to referents. Two of these workers were re-examined after a 3-month period without styrene exposure; the serum prolactin response following thyrotrophin-releasing hormone exposure was similar to that in the referent group. No

3. HEALTH EFFECTS

significant alterations in the levels of thyroid stimulating hormone, follicle stimulating hormone, or luteinizing hormone were found in this study (Arfini et al. 1987). However, thyroid stimulating hormone levels were significantly correlated with urinary metabolite (mandelic acid plus phenylglyoxylic acid) levels.

Significant increases in serum prolactin levels have also been observed in female rats exposed to 150 ppm, 8 hours/day for 10 days (Umemura et al. 2005). No significant alterations in serum prolactin levels were observed in similarly exposed male rats (Umemura et al. 2005) or in male rats exposed to approximately 150, 500, or 1,500 ppm 6 hours/day for 5 days (Jarry et al. 2002). Acute exposure to styrene did not result in significant alterations in thyroid stimulating hormone levels in male or female rats (Umemura et al. 2005).

Ocular Effects. Eye irritation in humans has been reported at high styrene concentrations (Carpenter et al. 1944; Stewart et al. 1968). Immediate eye irritation was reported in two human subjects exposed to 800 ppm styrene for 4 hours (Carpenter et al. 1944). Eye irritation was also noted by Stewart et al. (1968) in two of five volunteers exposed to 376 ppm styrene for 1 hour. Also, 345 styrene-exposed workers (98% male) were evaluated for ocular toxicity due to exposure to styrene (5–200 ppm) for 7–20 years. No evidence of optic neuritis, central retinal vein occlusion, or retrobulbar neuritis was found. Conjunctival irritation was a complaint of 22% of the 345 workers exposed to styrene levels above 50 ppm (Kohn 1978).

Eye and nasal irritation was observed in rats and guinea pigs exposed to 1,300 or 2,000 ppm styrene, 7–8 hours/day, 5 days/week for durations ranging from 21 to 30 weeks (Wolf et al. 1956). Rabbits and monkeys were exposed for up to 360 days with no effects.

3.2.1.3 Immunological and Lymphoreticular Effects

Two occupational studies have found significant alterations in lymphocyte subsets in styrene workers. Increase in percentage of CD4+/CD3CD4 T-lymphocytes and decrease in CD35+CD+ peripheral lymphocytes were observed in oil industry workers exposed to styrene, as compared to unexposed controls (Biró et al. 2002). However, these results should be interpreted cautiously because there were marked differences in smoking habits between the styrene workers and controls (80% styrene workers smoked compared to 20% of controls) and the two groups were not matched for gender. Bergamaschi et al. (1995b) found an altered distribution of lymphocyte subsets in styrene workers exposed to an 8-hour

3. HEALTH EFFECTS

time-weighted average (TWA) of 10–50 ppm. Some of these alterations, particularly the reduction in total T-lymphocytes (CD3+) and T-helper cells (CD4+), may be indicative of reduced cell-mediated immunity. This is supported by the finding of an impaired response to concanavalin in styrene workers exposed to a median styrene concentration of 26 ppm (Tulinska et al. 2000), 187–256 ppm (Somorovská et al. 1999) or 54–56 ppm (Somorovská et al. 1999). No alterations in the response to pokeweed mitogen were observed (Somorovská et al. 1999; Tulinska et al. 2000).

In patch-testing studies of cross-reactors to styrene, styrene 7,8-oxide was more sensitizing than styrene itself (Sjöborg et al. 1984). The authors interpreted this as evidence that styrene requires metabolism by skin aryl hydrocarbon hydroxylase to styrene epoxide for its sensitizing activity.

In animals, styrene exacerbated the inflammatory reaction in mice challenged with ovalbumin (Ban et al. 2006). Styrene-only exposure resulted in slight increases in Th2 cytokine (IL-4, IL-5, IL-13) and Th1 cytokine (interferon- γ) levels; however, the statistical significance of these alterations were not reported.

3.2.1.4 Neurological Effects

The available human data suggest that the nervous system is the most sensitive target following chronic-duration inhalation exposure. It is likely the most sensitive target following shorter-term durations, but this has not been as extensively investigated. In studies examining the acute neurotoxicity of styrene, impairment of the vestibular-oculomotor system was observed in experimental subjects exposed to 87 ppm for 1 hour (Ödkvist et al. 1982) or 376 ppm for 1 hour (Stewart et al. 1968). No alterations in the performance of balance tests were observed at 216 ppm for 1 hour (Stewart et al. 1968), 117 ppm for 2 hours (Stewart et al. 1968), or 99 ppm for 7 hours (Stewart et al. 1968). Although these NOAELs are higher than the LOAEL identified in the Ödkvist et al. (1982) study, the studies are not comparable. The Ödkvist et al. (1982) study used sensitive tests of vestibular-oculomotor function compared to the modified Romberg test (subjects stand on one foot with eyes closed, walk heel to toe, touch finger to nose) used in the Stewart et al. (1968) studies. An increase in the reporting of “feeling inebriated” was found in subjects exposed to 376 ppm for 1 hour (Stewart et al. 1968); no increases in subjective symptoms were observed in subjects exposed to 20 ppm for 3–4 hours (Seeber et al. 2004) or 216 ppm for 1 hour (Stewart et al. 1968). Additionally, no alterations in reaction time were observed in subjects exposed to 20 ppm for 3 or 4 hours (Seeber et al. 2004). No human studies examined neurotoxicity following intermediate-duration exposure.

3. HEALTH EFFECTS

In an international cohort of styrene workers, a significant association between mortality from central nervous system disease and cumulative styrene exposure was found (Welp et al. 1996c). The rate ratio was 3.29 (95% confidence interval [CI] of 0.48–22.65) for workers exposed to 25–49 ppm-years and 16.32 (95% CI 3.47–76.73) for those exposed for 200–349 ppm-years. A similar relationship was found for shorter durations of styrene exposure. The rate ratio was 2.33 (95% CI 0.40–13.56) for workers exposed for 6–11 months and 8.80 (95% CI 1.87–41.33) for workers exposed for 7–9 months. A significant association between mortality from epilepsy and duration of styrene exposure was found; the rate ratio in workers exposed for ≥ 10 years was 28.4 (95% CI 2.11–381.5). Time since first exposure was also significantly associated with mortality from epilepsy. Significant associations between mental disorders and duration of exposure and between suicide and duration of exposure were also found; however, for both of these causes of death, the rate ratio decreased with increasing duration of exposure and the investigators noted that lifestyle factors, rather than a direct effect of styrene, appear to be the most likely cause of the higher mortality.

A variety of neurological effects have been reported in workers chronically exposed to styrene including altered vestibular function, impaired hearing, decreased color discrimination, altered performance on neurobehavioral tests, and increased clinical symptoms. In general, these occupational exposure studies have several limitations. In most cases, the exposure levels reflect current exposure conditions and do not take into consideration past exposure to higher styrene levels that may have resulted in permanent damage. Some workers, particularly laminators, wore respiratory masks with or without canisters; many investigators estimated exposure based on biomarker levels, particularly urinary mandelic acid levels, while others did not. As discussed in greater detail in Sections 3.4.3, 3.4.3.1, and 3.8.1, urinary levels of styrene metabolites mandelic acid and phenoxyglycolic acid have been shown to correlate with time-weighted average styrene exposure levels and may be a more reliable biological indicator of styrene exposure in workplaces with highly variable styrene exposure levels. Significant differences between workers and referents were reported as LOAELs; however, the magnitude of the alteration may have been subclinical. A summary of the neurological effects observed in styrene workers is presented in Table 3-2.

Color vision appears to be one of the more sensitive targets of styrene toxicity, with many studies reporting alterations. Color vision was typically measured using the Lanthony desaturated panel D-15 test in which the subjects were asked to arrange 15 painted caps in a line with definite chromatic

3. HEALTH EFFECTS

Table 3-2. Summary of Occupational Exposure Studies Examining Neurological End Points

Reference	Number of workers	Mean duration of exposure (years)	NOAEL (ppm)	LOAEL (ppm)	Comments
Decreased color discrimination					
Chia et al. 1994	21	18.8 (range: 5–23)		6	Styrene exposure was estimated from mean urinary levels of mandelic acid (84.0 mg/g creatinine; range of 1.3–504.1 mg/g creatinine). Significant decreases in color discrimination, as expressed as total color difference score; no concentration-response relationship was found.
Kishi et al. 2001	21–42	6.2	4	10	Workers divided into three groups based on urinary mandelic acid levels. Significant differences in CCI, compared to age-matched controls, were found in the two highest groups. Mean CCI in 4, 10, and 46 ppm groups (CCI levels in age-paired controls): 1.21 (1.17), 1.23 (1.12), and 1.27 (1.13). Significant difference in CCI also found in analysis using 87 age-matched workers/controls.
Gong et al. 2002	43	6.4		10	Workers divided into two groups based on combined urinary mandelic acid and phenylglyoxylic acid level dividing line of 0.24 g/g creatinine (approximately 10 ppm); both groups were significantly different from controls. Mean CCI (mean of right and left eyes) in controls, low exposure, and high exposure groups: 1.02, 1.09, and 1.14.
Gobba et al. 1991	41			16	Significant differences found when compared to age-matched controls (41 workers/controls) and in older workers (≥ 40 years of age). CCI was significantly different in workers exposed to ≥ 50 ppm compared to workers exposed to < 50 ppm.
Iregren et al. 2005	53–55	12.9–17.9 (range: 2–39)		22	Lifetime weighted average exposure calculated for each worker using historical exposure data. Total error score was significantly different from workers in the low exposure group (9 ppm).

3. HEALTH EFFECTS

Table 3-2. Summary of Occupational Exposure Studies Examining Neurological End Points

Reference	Number of workers	Mean duration of exposure (years)	NOAEL (ppm)	LOAEL (ppm)	Comments
Fallas et al. 1992	60	6.5 (range: >1–29)		24.3	Significant difference in the number of subjects with error axis in the red-green or blue-yellow ranges; no significant difference in calculated error scores.
Campagna et al. 1996	118	5.2 or 6.6		26	Mathematical threshold of impaired CCI was 4 ppm; the upper limit of the CI was 25.7 ppm.
Eguchi et al. 1995	57	7.0 (range: 0.2–26.8)	8	93	Significant difference in CCI between age-matched workers and controls. Workers divided into two groups—significant difference in CCI between high-concentration workers (urinary mandelic acid level of 1.06 g/L [range: 0.46–3.98 g/L] equivalent to 93 ppm) and age-matched controls. No difference in low concentration workers (urinary mandelic acid levels—mean of 0.02 g/L; range of 0.04–0.41 g/L; equivalent to 8 ppm). CCI scores in low and high exposure groups (CCI in age-matched controls) of 1.173 (1.118) and 1.332 (1.125).
Reaction time					
Edling et al. 1993	20	9 (range: 1–25)	8.6		Simple and choice reaction time (measured before and after work).
Tsai and Chen 1996	45	8.3		21.9	Directly exposed workers compared to workers without direct styrene exposure (range of styrene levels was 0–6.4 ppm). Complex reaction time (continuous performance test): 532.8 and 495.6 ms in directly and indirectly exposed groups.
Jegaden et al. 1993	30	5		22.68	Reaction times (measured in morning before work) in workers and controls were 0.29 and 0.27 seconds for simple reaction time and 0.37 and 0.32 seconds for complex reaction time.
Fallas et al. 1992	60	6.5 (range: >1–29)	24.3		No significant difference in simple reaction time (23.7 seconds versus 22.7 seconds in controls).

3. HEALTH EFFECTS

Table 3-2. Summary of Occupational Exposure Studies Examining Neurological End Points

Reference	Number of workers	Mean duration of exposure (years)	NOAEL (ppm)	LOAEL (ppm)	Comments
Mutti et al. 1984a	50			25	Workers divided into four groups based on combined levels of urinary mandelic acid and phenylglyoxylic acid of <150, 150–299, 300–350, and >450 mmole/mole creatinine; 150 mmole/mole creatinine equivalent to 25 ppm. Complex reaction time in all workers and controls: 623.6 and 488.3 ms; the percent variations from matched controls were 120 and 165% in the 25 and 50 ppm groups.
Gamberale et al. 1976	106	2.7 (range: 0.1–11)		47	Mean styrene levels in each location were 16.6, 59.3, 41.6, and 101.4 ppm for resin applicators and 13.6 and 49.3 ppm for assemblers. The average styrene level for the six sites was 47 ppm. Simple reaction times (measured in morning before work) were 274 ms in workers and 260 ms in controls.
Cherry et al. 1980	27	NR		92	Simple reaction times (measured in morning before work) were 252 and 230 ms in workers and controls. A slower reaction time was also observed during the workshift.
Other neurobehavioral tests					
Chia et al. 1994	21	18.8 (range: 5–23)		6	Styrene exposure was estimated from mean urinary levels of mandelic acid (84.0 mg/g creatinine; range of 1.3–504.1 mg/g creatinine). Several tests of memory—Benton Visual Retention test (score 6.0 in workers vs. 7.7 in controls), digit symbol test (26.3 in workers vs. 38.0 in controls), and digit span (11.7 in workers vs. 15.6 in controls) were significantly affected. No significant relations between test score and urinary mandelic acid or phenylglyoxylic acid levels. No alteration in Santa Ana dexterity test or pursuit aiming.
Edling et al. 1993	20	9 (range: 1–25)	8.6		No effect on performance of symbol digit test.
Jegaden et al. 1993	30	5		22.68	Significant impairment in performance on digit span test.

3. HEALTH EFFECTS

Table 3-2. Summary of Occupational Exposure Studies Examining Neurological End Points

Reference	Number of workers	Mean duration of exposure (years) (range: >1–29)	NOAEL (ppm)	LOAEL (ppm)	Comments
Fallas et al. 1992	60	6.5 (range: >1–29)		24.3	Significant alteration in aiming test. No significant alteration in Santa Ana test of dexterity, digit span, digit symbol, or Benton visual retention tests. However, there were significantly more workers with digit span test scores differing by more than one standard deviation from the mean, particularly in workers exposed to styrene for >10 years.
Mutti et al. 1984a	50			25	Workers divided into four groups based on combined levels of urinary mandelic acid and phenylglyoxylic acid of <150, 150–299, 300–350, and >450 mmole/mole creatinine; 150 mmole/mole creatinine equivalent to 25 ppm. Impaired performance on verbal learning test; no significant relationship with test scores in duration of exposure or exposure level.
Clinical symptoms					
Flodin et al. 1989	8 or 9	11.6 (range: 6–21)		6	High prevalence of clinical symptoms—abnormal tiredness (7/8 subjects) and short memory (8/8 subjects) in workers exposed to 6 ppm styrene; high prevalences of problems concentrating (7/9 subjects) and irritation (8/9 subjects) were also observed in workers exposed to 12 ppm. Prevalence was not compared to a referent group.
Edling et al. 1993	20	9 (range: 1–25)		8.6	Reported more acute symptoms than controls (mean of 2.9 in workers versus 1.8 in controls). More frequently responded positively to the following questions: are you abnormally tired, do you often have painful tingling in some parts of your body, do you have a headache at least once a week.
Checkoway et al. 1992	16–27	3.4–4.1	10.8	18.9	Higher prevalence of headache, dizziness, light headedness, fatigue, irritability, feeling "drunk" at work, and memory loss.
Cherry et al. 1980	27	NR		92	At the end of the workshift, changes in self-reported physical and mental tiredness and general health scores were correlated with blood styrene concentrations.

3. HEALTH EFFECTS

Table 3-2. Summary of Occupational Exposure Studies Examining Neurological End Points

Reference	Number of workers	Mean duration of exposure (years)	NOAEL (ppm)	LOAEL (ppm)	Comments
Hearing					
Morata et al. 2002	65	17 (range: 1–39)		3.68	Noise level 82 dBA. A higher prevalence of high frequency hearing loss (47%) compared to controls (33%), but difference was not statistically significant. Significantly poorer thresholds were observed at 2, 3, 4, 6, and 8 kHz in the styrene group. The OR of 1.19 (95% CI, 1.11–1.28) times greater for each 1 year of age was calculated and 2.44 times greater (95% CI, 1.01–5.89) for each increment of 1 mmol mandelic acid/g creatinine in urine.
Śliwińska-Kowalska et al. 2003	194	NR		15.6	Current styrene concentrations ranged from 0.05 to 46 ppm; average worklife mean styrene concentration was 15.6 ppm. Average noise level was 80.3 dBA. Abnormal audiograms were found in 63.3% of styrene workers, compared to 33.8% in the unexposed controls. The OR of hearing loss was 5.2 (95% CI, 2.9–8.9). A positive linear relationship between styrene working life exposure levels and hearing thresholds.
Morioka et al. 1999	93	9.4		16	Sound levels in the workplace ranged from 53 to 95 dBA. Significant correlation between individual percentiles of the upper limit of hearing and styrene concentrations in workers exposed for ≥5 years; the prevalence rates below the 75th percentile were significantly higher than 25% in workers exposed to >16 ppm.
Möller et al. 1990	18	10.8 (range: 6–15)	18		No styrene-related alterations in the pure-tone audiometry and speech discrimination tests were found. Seven workers displayed abnormal results in distorted speech and/or cortical response audiometry tests.
Calabrese et al. 1996	20	7.6 (range: 2–23)	36		All workers had normal hearing thresholds and no abnormalities of stapedial reflex were found. No significant effect on ABRs (as compared to 10 control subjects) were observed; additionally, when nine subjects were re-tested after 3 weeks without exposure, no significant difference between pre- and postrecovery values were found.

3. HEALTH EFFECTS

Table 3-2. Summary of Occupational Exposure Studies Examining Neurological End Points

Reference	Number of workers	Mean duration of exposure (years)	NOAEL (ppm)	LOAEL (ppm)	Comments
Vestibular					
Möller et al. 1990	18	10.8 (range: 6–15)		18	Significantly higher sway with eyes open or closed in the static posturography test, increased latency in saccade test, a phase lag and depressed gain in unpredictable and predictable stimulation in the smooth eye pursuit test, and impaired ability to suppress vestibuloocular reflex in sinusoidal and pseudorandomized tests.
Toppila et al. 2006	88	NR		24.8	Nonlaminators used as the comparison group; 88 pairs of age-matched workers were used for postural stability tests; the mean styrene and mandelic acid phenylglyoxylic acid concentrations for these pairs were 24.8 ppm and 1.4 mmol/L for the laminators and 4.8 ppm and 0.3 mmol/L for the nonlaminators. Poorer performance in dynamic tests on the static platform. In the tilting platform and virtual reality tests, sway velocity was greater and workers became unstable and displayed large correctional movements.
Calabrese et al. 1996	20	7.6 (range: 2–23)		36	Abnormal results in the vestibuloocular tests, these alterations persisted after the 3-week recovery period. No alterations in visual suppression test or postural performance were found.
Nerve conduction velocity					
Seppäläinen and Härkönen 1976	96	5	30		Styrene exposure was estimated using end-of-shift urinary mandelic acid levels collected weekly for 5 consecutive weeks. No significant differences in median, ulnar, deep peroneal, or posterior tibial nerve motor or sensory conduction velocities were observed, as compared to a referent group with a similar age distribution.
Štětkařová et al. 1993	15	11		50	Decreased peripheral conduction velocities in median and tibial nerves were observed in female styrene workers; prolonged latencies of peripheral and cortical somatosensory evoked potentials were also observed in the female workers exposed to styrene.

3. HEALTH EFFECTS

Table 3-2. Summary of Occupational Exposure Studies Examining Neurological End Points

Reference	Number of workers	Mean duration of exposure (years)	NOAEL (ppm)	LOAEL (ppm)	Comments
Triebig et al. 1985	11	4	100		No significant alterations in maximum conduction velocity in the ulnar nerve or distal conduction velocity of the sensory fibers of the ulnar and median nerves.

ABR = auditory brainstem response; CCI = color confusion index; CI = confidence interval; LOAEL = lowest-observed-adverse-effect level; MS = millisecond; NOAEL = no-observed-adverse-effect level; NR = not reported; OR = odds ratio

3. HEALTH EFFECTS

sequence; the color confusion index (CCI) quantifies the number of types of mistake. A significant correlation between CCI and urinary mandelic acid concentration (after correction for age) was observed in workers at fiberglass reinforced plastic facilities (Kishi et al. 2001). When workers were divided into three groups based on end-of-shift urinary mandelic acid levels, there were significant differences between CCI in workers with a mean a mandelic acid level of 0.14 or 0.65 g/L and age-matched referents; no difference was found for the third group with a mean mandelic acid level of 0.05 g/L. The investigators estimated that these urinary mandelic acid levels were equivalent to styrene exposure levels of 4, 10, and 46 ppm. Thus, this study identifies a NOAEL of 4 ppm and a LOAEL of 10 ppm for decreased color discrimination. Similarly, Gong et al. (2002) found significantly higher CCI values in workers at a fiberglass reinforced plastic boat facility with end-of-shift urinary mandelic acid and phenylglyoxylic acid levels of ≥ 0.24 g/g creatinine or < 0.24 g/g creatinine; a mandelic acid plus phenylglyoxylic acid urine level of 0.24 g/g creatinine is equivalent to a styrene exposure level of 10 ppm. A significant increase in CCI was also observed in workers at fiberglass reinforced plastic facilities exposed to a geometric mean concentration of 16 ppm, as compared to age-matched controls (Gobba et al. 1991). In contrast to other studies, Gobba et al. (1991) did not find a significant relationship between end-of-shift urinary mandelic acid levels and CCI; however, urinary styrene levels correlated with CCI values. Significantly higher CCI values were observed in fiberglass reinforced workers with a mean urinary mandelic acid levels of 1.06 g/L, which is roughly equivalent to a styrene exposure level of 93 ppm (Eguchi et al. 1995). This study did not find significant alteration in workers with a mean urinary mandelic acid level of 0.02 g/L, equivalent to 8 ppm. Another study of fiberglass reinforced plastic workers (some of this cohort was examined by Gobba et al. 1991 and Campagna et al. 1995) found a significant association between CCI and styrene exposure levels (Campagna et al. 1996). The investigators concluded that color vision impairment could be detected at styrene levels of 4 ppm with a 95% upper confidence limit of 26 ppm. Two other occupational exposure studies using different measures of color vision impairment also found significant alterations. Chia et al. (1994) found significantly poorer color discrimination, after adjusting for age, education, and alcohol consumption, in 21 workers at a fiber-reinforced plastic boat manufacturing facility; the styrene exposure level of 6 ppm was estimated from a mean end-of-shift urinary mandelic acid level of 84.0 mg/g creatinine. No relationship between the total color difference score and the urinary mandelic acid level was found. In 60 workers in the shipbuilding industry with a mean styrene exposure level of 24.3 ppm, a significantly higher incidence of workers with errors in the blue-yellow or red-green ranges, compared to a referent group, was found (Fallas et al. 1992). Total error score was significantly different in workers, with a lifetime weighted average exposure level of 22 ppm styrene, as compared to workers in a low exposure group (9 ppm) (Iregren et al. 2005). Several studies found improvements in color vision following an

3. HEALTH EFFECTS

extended period of no styrene exposure or lower exposure. Triebig et al. (2001) reported a significant improvement in CCI scores following a 4-week period with no styrene exposure; in contrast, no improvement in CCI scores was found in another group of styrene workers following a 1-month period without styrene exposure (Gobba et al. 1991). Two studies found significant improvements in color vision (age-adjusted color confusion score or CCI score) were observed in styrene workers following a decrease in styrene air level (Castillo et al. 2001; Triebig et al. 2001). However, one study found no change in age-adjusted near visual contrast sensitivity following a decrease in styrene exposure levels (Castillo et al. 2001).

A number of studies have found significant alterations in performance on a variety of neurobehavioral tests; among these studies, reaction time appears to be the most frequently examined end point. Significant increases in simple reaction time have been observed in styrene workers exposed to concentrations of 21.9, 22.68, 47, or 92 ppm (Cherry et al. 1980; Gamberale et al. 1976; Jegaden et al. 1993; Tsai and Chen 1996); tests for reaction time were measured in the morning before the work shift, suggesting that the effect was not due to acute exposure to styrene. The reaction times were 4–10% slower in the styrene workers as compared to the referent groups. No significant alterations in simple reaction time were observed in workers exposed to 8.6 ppm (Edling et al. 1993) or 24.3 ppm (Fallas et al. 1992). Similarly, complex reaction time was significantly increased among styrene workers exposed to 21.9, 22.68, or 25 ppm (Fallas et al. 1992; Jegaden et al. 1993; Mutti et al. 1984a); the variance from controls ranged from 7.5 to 20%. No alterations in complex reaction time were observed in workers exposed to 8.6 ppm (Edling et al. 1993). Impaired performance on the digit span test, which measures attention/concentration, was observed in workers exposed to 6 ppm (Chia et al. 1994), 22.68 ppm (Jegaden et al. 1993), or 24.3 ppm (Fallas et al. 1992). Other neurobehavioral performance tests that may be altered by chronic exposure to styrene included digit symbol or symbol digit tests at 6 ppm (Chia et al. 1994), visuomotor at 50 ppm (Mutti et al. 1984a) or 75 ppm (Lindstrom et al. 1976), and memory at 25 ppm (Mutti et al. 1984a). However, other studies have not found significant alterations in digit symbol at 8.6 ppm (Edling et al. 1993), 24.3 ppm (Fallas et al. 1992), or 25 ppm (Mutti et al. 1984a), or memory at 75 ppm (Lindstrom et al. 1976). By pooling response data for several tests of neurobehavioral performance, Mutti et al. (1984a) were able to analyze exposure-response relationships. When workers were divided into four groups based on morning urinary mandelic acid and phenylglyoxylic acid levels, the number of subjects with abnormal scores on greater than one, two, or three tests increased with increasing exposure concentration. Additional analyses demonstrated that the exposure intensity and duration of exposure affected a worker's performance on neurobehavioral tests and the duration of exposure appeared to affect performance more than exposure intensity.

3. HEALTH EFFECTS

Clinical symptoms of neurotoxicity have been reported by styrene workers; commonly reported symptoms included headaches, dizziness, impaired memory, and feeling “drunk”. At 6 or 12 ppm, abnormal tiredness and short memory were reported by most of the styrene workers examined by Flodin et al. (1989); problems concentrating and irritation were also reported by most workers exposed to 12 ppm. After a 7-month period without styrene exposure, there was a marked improvement in symptoms and the mean number of symptoms reported was 1.9, compared to 5.3 reported 7 months earlier. Fiberglass reinforced plastic industry workers exposed to 18.9 or 50.0 ppm reported a higher prevalence of headaches, dizziness, light headedness, fatigue, irritability, feeling “drunk”, and memory loss (Checkoway et al. 1992); the prevalence of clinical signs was not significantly increased in workers exposed to 10.8 ppm. Increases in the incidence of headache, memory disturbances, forgetfulness, dizziness, and sensory symptoms in the upper and lower extremities were observed in workers with high exposure to styrene compared to those with low styrene exposure (Matikainen et al. 1993a); exposure levels were not reported. A significantly higher incidence of subjective symptoms (nausea, feeling of drunkenness, dizziness, and disturbance) was observed in styrene workers exposed to 4–164 ppm, as compared to controls (Geuskens et al. 1992). No significant difference in the incidence of symptoms related to cognitive motor disturbances (lack of concentration, understanding, trouble with movements) was found. Fiberglass boat manufacturing workers exposed to 92 ppm reported a higher prevalence of physical and mental tiredness at the end of the work shift than controls (Cherry et al. 1980). No alterations in the reporting of clinical symptoms were observed in plastic industry workers exposed to a mean concentration of 8.6 ppm (Edling et al. 1993). One study examined styrene’s potential to affect olfactory function. No alterations in three clinical tests of olfactory function were observed in styrene workers exposed to 24.6 ppm styrene for minimum of 4 years (Dalton et al. 2003).

Styrene-induced damage to hearing and the vestibular system have been examined in a number of studies of chronically exposed workers. Several studies have reported significant associations between styrene exposure and hearing impairment; however, interpretation of the findings is limited by confounding exposure to noise or other solvents. Noise studies have found that exposure to >85 dB for over 10 years can result in a 10% hearing loss (Prince et al. 1997). Morioka et al. (1999) found an increased prevalence of workers with a urinary mandelic acid level of >0.3 g/L (approximately 16 ppm) with an upper frequency of hearing below the 75th percentile for normal. However, interpretation of the results is limited by confounding exposure to noise and exposure to other solvents, particularly toluene, which has been shown to be ototoxic. The noise levels ranged from 53.0 to 95.0 dBA with 14% of the measurements exceeding 85 dBA. Another study (Muijsers et al. 1988) of styrene workers found a

3. HEALTH EFFECTS

significant difference in hearing threshold at 8 kHz between indirectly exposed workers (mean styrene level of 14 ppm) and directly exposed workers (mean styrene level of 32 ppm); however, no differences were found in comparisons of indirectly and directly exposed workers with referent workers. The noise level for both groups of styrene workers was 80–85 dBA for most of the day. Śliwińska-Kowalska et al. (2003) found a significantly elevated risks of hearing loss among styrene workers exposed to a mean styrene concentration of 15.6 ppm (average noise level of 80.3 dBA). The odds ratio (adjusted for noise and gender) in workers only exposed to styrene was 5.2 (95% CI 2.9–8.9). The hearing losses were found within the range of 2–8 kHz. Morata et al. (2002) found significant decreases in hearing thresholds at 2, 3, 4, and 6 kHz in workers exposed to 0.05–22 ppm (mean of 4 ppm); no difference in the prevalence of high frequency hearing loss, as compared to referent workers, was found. The fairly wide range of exposure levels adds a great deal of uncertainty to estimating the LOAEL from this study; although the mean exposure is reported as the LOAEL in Figure 3-1, this value may be an overly conservative estimate of the true LOAEL. Other studies have not found significant alterations in hearing. Möller et al. (1990) found no indications of hearing loss in workers exposed to 18 ppm styrene. Sass-Korstak et al. (1995) did not find significant relationships between lifetime styrene exposure and hearing loss in workers at fiber-reinforced plastics manufacturing facilities. The cumulative styrene exposure level was calculated using data for current exposure (25 ppm for directly exposed workers and 8 ppm for indirectly exposed workers), length of time in each job category, and a downward adjustment for self-reported respirator use. The average noise levels (L_{eq}) were 88.1 and 89.2 dBA for the directly and indirectly exposed workers, for nonexposed workers, a sound level of 80 dBA was assumed. In another study of fiberglass workers (Calabrese et al. 1996), no significant alterations in audiometric tests or auditory brainstem response were observed in workers exposed to a mean styrene level of 36 ppm. Additionally, a 3-week recovery period did not result in any significant changes in auditory brainstem responses (pre- and post-recovery) in nine of the workers.

Other studies have examined workers for styrene-induced vestibular effects. Significant alteration in tests of central vestibulocular and optocular motor movements (i.e., static posturography, smooth eye pursuit, saccade, and vestibulocular reflex tests) were observed in workers at a plastic boat manufacturing facility exposed to a TWA styrene concentration of 18 ppm (Möller et al. 1990). No indications of labyrinthine or peripheral vestibular lesions were observed. Toppila et al. (2006) also found significant alterations in postural stability in workers at fiberglass-reinforced plastic boat manufacturing facilities exposed to 25 ppm styrene. In contrast, Calabrese et al. (1996) did not find significant alterations in visual suppression tests or postural stability in fiberglass plant workers exposed to 36 ppm styrene. Significant

3. HEALTH EFFECTS

alterations in vestibuloocular reflex were found. A 3-week recovery period did not result in significant changes in the test results.

Workers exposed to styrene in several industries at mean concentrations of 5–125 ppm had mild sensory neuropathy characterized by decreased sensory conduction amplitude and increased duration, but there were too few people to define a NOAEL (Rosen et al. 1978). Peripheral neuropathy and reduced nerve conduction velocity was also reported in an individual following a 2-day exposure to an unknown amount of styrene (and other chemicals) (Fung and Clark 1999). Leg weakness, leg muscle cramps, and paresthesia were also reported in two styrene workers (Gobba et al. 1995). Moderate sensorimotor neuropathy of the demyelinating type was diagnosed in both cases based on the clinical symptoms and the decreased motor nerve conduction velocity in the peroneal nerve and decreased sensory nerve conduction velocity in the sural and median nerves. Alterations in nerve conduction velocity have also been observed in styrene workers. Decreased peripheral conduction velocities in the median and tibial nerves and prolonged latencies of peripheral and cortical somatosensory evoked potentials were observed in female styrene workers exposed to 30–130 ppm (midpoint of the range is 50 ppm) (Štětkářová et al. 1993). Significant decreases in ulnar and peroneal maximum conduction velocities and increased peroneal motor distal latencies were observed in fiber reinforced workers exposed with urinary mandelic acid levels (end of shift) of ≥ 250 mg/L, as compared to referent workers. Motor distal latencies in the workers with urinary mandelic acid levels ≥ 250 mg/L were also significantly lower than in workers with urinary mandelic acid levels < 250 mg/L (Yuasa et al. 1996). In contrast, no alterations in motor or sensory nerve conduction velocity in the ulnar, median, deep peroneal, or posterior tibial nerve were observed in workers exposed to a TWA styrene concentration of 30 ppm (based on urinary mandelic acid excretion) (Seppäläinen and Härkönen 1976), and no alteration in motor or sensory nerve conduction velocity was observed in workers exposed to approximately 100 ppm for a median of 4 years (Triebig et al. 1985). Although Seppäläinen and Härkönen (1976) did not find alterations in nerve conduction velocity, they found abnormal EEGs in 24% of the styrene workers, as compared to reported values for the normal population. The mean urinary mandelic acid level (975 mg/dm^3) was higher in workers with abnormal EEGs compared to those with normal readings (750 mg/dm^3). Similarly, a significantly higher absolute EEG power in alpha band in the fronto-temporal region of the brain was found in workers with high styrene exposures, as compared to workers with low-level exposure (Matikainen et al. 1993a).

The majority of the available animal neurotoxicity studies have focused on hearing impairment. Hearing loss and a loss of outer hair cells (OHC) in the organ of Corti were observed in rats acutely exposed to 1,000 ppm (Campo et al. 2001; Lataye et al. 2003) or 1,600 ppm (Crofton et al. 1994). In contrast, acute

3. HEALTH EFFECTS

exposure of guinea pigs to 1,000 ppm did not result in hearing loss or OHC damage (Lataye et al. 2003). Intermediate-duration exposure studies have consistently found hearing loss and loss of OHC in rats exposed to ≥ 750 ppm styrene (Campo et al. 2001; Lataye et al. 2000, 2001; Loquet et al. 2000; Pouyatos et al. 2002; Pryor et al. 1987; Yano et al. 1992). Exposure to 600–650 ppm resulted in OHC losses but no alterations in hearing threshold (Loquet et al. 1999; Makitie et al. 2002; Pouyatos et al. 2002). A NOAEL of 300 ppm was identified by Makitie et al. (2002).

Other neurological effects that have been observed in animal studies include lethargy and unsteady gait in mice exposed to 250 ppm for 2 weeks (Cruzan et al. 1997), an increase in astroglial alterations at 320 ppm (Rosengren and Haglid 1989), a decrease in nerve conduction velocity in rats exposed to 2,000 ppm, but not 200 ppm, for 32 weeks (Yamamoto et al. 1997), and concentration-related alterations in nystagmus elicited by optokinetic, vestibular, simultaneous optokinetic-vestibular, and saccadic stimulation in rats exposed to 830–4,000 ppm styrene for at least 60 minutes (actual duration of exposure was not reported) (Niklasson et al. 1993).

The highest NOAEL values and all reliable LOAEL values for neurological effects in each species and duration category are recorded in Table 3-1 and plotted in Figure 3-1.

3.2.1.5 Reproductive Effects

Information on the reproductive effects of styrene in humans is available from epidemiological studies of the reproductive outcomes of females employed in the various industrial operations in which styrene is used. However, exposures to styrene were not adequately quantified in any of the studies cited. In one study, spontaneous abortions among 9,000 Finnish chemical workers from 1973 to 1976 were analyzed (Hemminki et al. 1980). The risk of spontaneous abortion expressed as number of abortions per 100 pregnancies) was significantly higher in women employed in styrene production compared to all women in Finland 15.0 vs. 5.5). However, this increase was not detected in a follow-up study of the same workers (Hemminki et al. 1984). An increase in the occurrence of spontaneous abortions was also observed in a study of 76 women involved in processing polystyrene plastics (McDonald et al. 1988); the ratio of observed to expected abortions was 1.58 (95% CI 1.02–2.35). The possible embryotoxic effects of styrene on 67 female lamination workers compared to 67 age-matched controls were evaluated in a second study (Härkönen and Holmberg 1982). The number of births was significantly lower among the workers exposed to styrene. This result was explained in part by a greater number of induced abortions in the styrene-exposed group. The number of spontaneous abortions was not elevated in the exposed

3. HEALTH EFFECTS

women. No increased risk of spontaneous abortions among workers processing polymerized plastics or heated plastics made of vinyl chloride or styrene was reported (Lindbohm et al. 1985). The authors reported that the statistical power of the study was low due to the small study population. These studies are not conclusive since the workers were exposed to chemicals other than styrene in the workplace and the concentrations of styrene were not adequately reported. Two studies have examined the potential of styrene to induce menstrual disturbances. A significant increase in the incidence of oligomenorrhea was observed in petrochemical industry workers; the adjusted odds ratio was 1.65 (95% CI 1.05–2.55) (Cho et al. 2001). Although the odds ratio includes an adjustment for exposure to other aromatic chemicals, there was potential for exposure to other chemicals; only three workers were exposed to styrene only and none of these women reported oligomenorrhea. In contrast, no significant alterations were observed in women working at reinforced plastics facilities with a mean styrene exposure level of 52 ppm for women directly exposed to styrene and 13 ppm for those indirectly exposed (Lemasters et al. 1985). Several studies have examined levels of prolactin, follicle stimulating hormone, and luteinizing hormone levels in female styrene workers; the results of these studies are discussed in Section 3.2.1.2, Endocrine Effects.

Several studies have also examined styrene's potential to induce male reproductive effects. A significant decrease in sperm concentration, total sperm count, percentage of normal sperm, and percentage of nonvital sperm and an increase in sperm velocity were observed in 23 workers employed at a styrene manufacturing facility for approximately 6 months, as compared to levels during the first week of employment (Kolstad et al. 1999a; results are also presented in Kolstad et al. 1999b). No significant relationships between urinary mandelic acid level and sperm density, total sperm count, or proportion of sperm with normal morphology were observed. A positive correlation between mandelic acid levels and the percentage of nonvital sperm was found, but the trend test was not statistically significant. In a large multinational study of male styrene workers, no significant alterations in time-to-pregnancy were found; the odds ratio, adjusted for maternal age, use of oral contraceptives, maternal and paternal smoking habits, time-to-pregnancy starting year, length of employment, and country, was 0.79 (95% CI 0.59–1.05) (Kolstad et al. 2000). Additionally, no significant alterations in fertility rates were found when workers were divided into groups based on length of exposure, period of attempting pregnancy, or exposure group (monitoring data from a subset of facilities were used to divide workers into different exposure groups) (preliminary data from this study was reported by Kolstad et al. 1999c). Similarly, no significant alterations in time to pregnancy among styrene workers exposed to high or intermediate/low exposure levels (based on urinary mandelic acid levels) (Sallmén et al. 1998).

3. HEALTH EFFECTS

Two animal studies have examined the reproductive toxicity of styrene following inhalation exposure. No statistically significant alterations in the frequency of abnormal sperm heads were observed in mice 3–5 weeks after exposure to 300 ppm for 5 days (Salomaa et al. 1985). In a two-generation study in rats (Cruzan et al. 2005b), no significant alterations in reproductive performance, estrous cycle length, spermatogenic parameters, or histological alterations in reproductive tissues were observed at concentrations up to 500 ppm.

3.2.1.6 Developmental Effects

Limited information concerning developmental effects of styrene in humans is available from studies of delivery outcome of women employed in the plastics industry (processing styrene or polyurethane plastics). Case-control studies performed in Sweden and Norway did not detect an increase in the odds ratio for developmental effects (stillbirth, infant death, malformations, low birth weight) in women who worked in the plastic industry (Ahlborg et al. 1987). However, actual levels of styrene exposure were not known for either group of workers. Another study did not find significant increases in the occurrence of congenital malformations in children of men or women working at reinforced plastics facilities (Härkönen et al. 1984). A <4% lower birthweight were observed in the children of women who worked in the reinforced plastics industry in areas with elevated levels of styrene (mean concentration of approximately 82 ppm) during pregnancy (Lemasters et al. 1989). However, this decrease was not statistically significant ($p=0.08$). These studies suggest that developmental effects in exposed workers are not of major concern, but the data are not adequate to exclude this effect. Moreover, interpretation of the results is complicated due to exposure of the workers to other chemicals in the workplace such as toluene, xylene, acetone, methylene chloride, and methyl ketone (Lemasters et al. 1989), as well as thermal degradation products of styrene polymers (Ahlborg et al. 1987). Workers may also be exposed to aerosols containing aldehydes, ketones, alcohols, esters, acids, and anhydrides. An expert panel convened by the National Toxicology Program (NTP 2006) concluded that the human data are not sufficient to evaluate the potential developmental toxicity of styrene in humans.

The developmental toxicity of styrene has been examined in several animal studies. The average fetal crown-rump length was significantly reduced in rats exposed to 300 ppm on gestational days 6–15, but was not affected in rats exposed to 600 ppm (Murray et al. 1978); the investigators concluded that this effect was not treatment-related. A few skeletal variants such as lumbar spurs and delayed ossification of sternbrae occurred in the styrene-exposed litters at a higher incidence than the control litters; however, the occurrence of this effect was similar to historical controls. No developmental effects were observed

3. HEALTH EFFECTS

in rabbits exposed to 600 ppm styrene on days 6–18 of gestation (Murray et al. 1978). Although there was a significant increase in the incidence of unossified sternebrae in the 600 ppm group, it did not exceed that found in historical control data. No significant alterations in the number of live fetuses, dead/resorbed fetuses, or malformed fetuses were observed in the offspring of mice exposed to 250 ppm styrene on days 6–16 of gestation (Kankaanpää et al. 1980). In the same study, exposure of hamsters to 1,000 ppm of styrene on days 6–18 of gestation resulted in a significant increase in the number of dead or resorbed fetuses; no other alterations were observed (Kankaanpää et al. 1980). No effects were observed at 750 ppm. Similarly, no developmental effects were observed in a two-generation study in which rats were exposed to 500 ppm prior to mating, during mating, and during gestation and lactation (Cruzan et al. 2005b). In contrast, Katakura et al. (1999, 2001) reported a significant increase in neonatal deaths in the offspring of rats exposed to 300 ppm on gestational days 6–20.

Two studies have examined potential neurodevelopmental effects. Some minor alterations in tests of developmental milestones (incisor eruption), functional observational battery tests (forelimb grip strength), and swimming maze test were observed in the F2 offspring of rats exposed to 500 ppm in a two-generation study; these alterations were attributed to a lower body weight rather than a neurodevelopmental effect of styrene (Cruzan et al. 2005a). No alterations in locomotor activity, acoustic startle response, or brain morphology and weights were observed in this study. Another study found delays in righting reflex and incisor eruption in the offspring of rats exposed to 300 ppm on gestational days 6–20 (Katakura et al. 2001). This study (Katakura et al. 1999, 2001) also found alterations in homovanillic acid levels in the cerebrum and 5-hydroxyindoleacetic acid levels in the hippocampus of the offspring in the 300 ppm group.

The highest NOAEL values and all reliable LOAEL values for developmental effects in each species and duration category are recorded in Table 3-1 and plotted in Figure 3-1.

3.2.1.7 Cancer

A number of studies have examined the carcinogenic potential of styrene in workers at styrene manufacturing and polymerization facilities, reinforced plastics facilities, and styrene-butadiene manufacturing facilities and among community members exposed to elevated styrene workers. Although there are several epidemiologic studies which suggest there may be an association between styrene exposure and an increased risk of leukemia and lymphoma, the evidence is generally inconclusive due to

3. HEALTH EFFECTS

multiple chemical exposures and inadequate documentation of the levels and durations of exposure to styrene.

Of the industries examined, workers employed at glass-reinforced plastics manufacturing facilities are likely to be exposed to higher levels of styrene and have lower potential for exposure to other carcinogenic agents. Some studies of glass-reinforced plastic workers have found suggestive evidence of increased cancer risks, particularly in workers with longer exposures to higher levels of styrene. No alterations in the number of deaths from cancer were observed in workers with high styrene exposure (mean levels at two facilities were 42.5 and 71.5 ppm) (Okun et al. 1985). In a follow-up study of these workers (Ruder et al. 2004), a significant increase in the number of deaths from urinary tract cancer (standardized mortality ratio [SMR] 3.44; 95% CI 1.26–7.50) was observed among workers with high styrene exposure; a trend for increasing SMRs for urinary tract cancer with increasing duration of exposure was also observed. The SMRs were not significantly elevated for other cancer types. In a very large epidemiological study of nearly 16,000 workers in the styrene plastic industry, the death rate from leukemia was twice as high in areas of high exposure as in areas of low exposure (Wong 1990); however, there were no statistically significant differences. In a follow-up study conducted 12 years later (Wong et al. 1994), significant increases in deaths from all cancers (SMR 115.5, 95% CI 104.8–127.1), cancer of the esophagus (SMR 191.7; 95% CI 104.8–321.7), bronchus, trachea, or lung (SMR 140.6; 95% CI 119.3–164.0), cervix or uteri (SMR 283.5; 95% CI 135.9–521.3), and female genital organs (SMR 201.6; 95% CI 107.4–344.8) were observed. However, no relationships between styrene exposure (exposure level or duration of exposure) and deaths from these cancer types were found. No significant increases in the incidence of non-Hodgkin's lymphoma, Hodgkin's disease, multiple myeloma, leukemia, or all lymphohematopoietic malignancies were observed in workers at Danish reinforced plastics facilities in which 50–100% of the workers were involved in reinforced plastics production (Kolstad et al. 1993, 1994). However, when workers were divided by first year of employment, there was a significant increase in leukemia (standard incidence ratio [SIR] 1.69, 95% CI 1.09–2.49) among workers with a latency of ≥ 10 years and first year of employment of 1964–1970; when the data were analyzed by the length of employment, the incidence of leukemia was only significantly elevated among workers employed for <1 year. Significant increases in incidence were also observed for pancreatic cancer in workers with a high probability of styrene exposure (incidence rate ratio of 2.2; 95% CI 1.1–4.5) and urinary bladder cancer in workers with the highest probability of exposure and employed for >1 year (incidence rate ratio of 2.1; 95% CI 1.1–4.1) (Kolstad et al. 1995). No significant alterations in the incidence of leukemia, lymphoma, or other cancers were observed in styrene workers at eight British reinforced plastic manufacturing facilities (Coggon et al. 1987). In a large international cohort of workers

3. HEALTH EFFECTS

employed in the reinforced plastics industry (this cohort included the British cohort examined by Coggon et al. 1987 and the Danish cohort examined by Kolstad et al. 1993, 1994, 1995), no significant alterations in no excess in mortality from all cancer or cancer of the lymphatic and hematopoietic tissues were observed (Kogevinas et al. 1993, 1994). However, significant increases in the incidence of lymphatic and hematopoietic neoplasms were observed in workers with a latency of at least 10 years (relative risk [RR] 2.90; 95% CI 1.29–6.48 in workers with a latency of 10–19 years and RR 3.97; 95% CI 1.30–12.13 for workers with a latency of ≥ 20 years) and in workers exposed to ≥ 100 ppm styrene (RR 3.11; 95% CI 1.07–9.06 for workers exposed to 100–119 ppm; RR 3.08; 95% CI 1.04–9.08 for workers exposed to 120–199 ppm; RR 3.59; 95% CI 0.98–13.14 for workers exposed to ≥ 200 ppm), as compared to unexposed workers. An increase in the number deaths from malignant lymphomas was also observed in workers exposed to 120–199 ppm styrene (RR 7.15; 95% CI 1.21–42.11).

An increase in lymphatic leukemia (4 observed deaths versus 0.5 expected) in workers exposed to polymer extrusion fumes, solvents, and colorants, but was not found to be related to duration or level of exposure (Ott et al. 1980). In a follow-up to this study, which followed the workers for another 11 years (Bond et al. 1992), a nonsignificant increase in the number of deaths from lymphatic and hematopoietic tissue cancers (SMR 144; 95% CI 95–208) was observed. Statistically significant increases in the number of deaths from lymphatic and hematopoietic cancer were observed in workers exposed to 1–4 ppm styrene and a 15-year minimum latency (SMR 160; 95% CI 102–238); however, significant alterations were not found in workers exposed to ≥ 5 ppm or with longer latency periods. In another study of workers involved in the styrene production, polymerization, or processing, a statistically significant excess of lymphoma deaths (3 deaths observed versus 0.56 expected) was reported; 2 of the 3 deaths occurred in men < 40 years of age who had been exposed for at least a year (Hodgson and Jones 1985). However, the lack of association with actual exposure levels or specific durations and the small number of observed deaths requires cautious interpretation. No significant alterations in the number of deaths from cancer were observed in workers (1960 subjects) exposed to styrene in a production and polymerization facility (Frentzel-Beyme et al. 1978). In a study of workers at a styrene-polystyrene manufacturing facility who had at least 5 years of exposure, there were no significant increases in cause-specific mortality (Nicholson et al. 1978). However, when workers employed for < 5 years were included in the analysis, there was an apparent increase in the number of deaths from lymphoma or leukemia (statistical analysis not conducted).

A number of older studies provide suggestive evidence of increased risk of lymphatic and hematopoietic cancers in workers at styrene-butadiene rubber manufacturing facilities (Matanoski and Schwartz 1987;

3. HEALTH EFFECTS

Matanoski et al. 1990; McMichael et al. 1976; Meinhardt et al. 1982); however, these studies provided limited exposure data and did not adjust for contribution of 1,3-butadiene to the overall cancer risk. A case-control study (Matanoski et al. 1993, 1997; Santos-Burgoa et al. 1992) provides suggestive evidence that the increase in leukemia was due to exposure to 1,3-butadiene rather than to styrene exposure. However, increases in the risk of lymphosarcoma and myeloma were associated with styrene exposure (Matanoski et al. 1997). More recent studies of styrene-butadiene workers include adjustments for 1,3-butadiene exposure. A cohort mortality study conducted by Delzell and associates (Delzell et al. 1996; Macaluso et al. 1996) examined workers at many of the same styrene-butadiene rubber manufacturing facilities examined by Matanoski and associates and Meinhardt and associates. In this examination of 15,649 male synthetic rubber workers employed for at least 1 year at one of eight styrene-butadiene rubber manufacturing facilities in the United States or Canada, significant increases in deaths from leukemia were observed among hourly employees (SMR 143; 95% CI 104–191), particularly among workers employed for ≥ 10 and ≥ 20 years since hire (SMR 224; 95% CI 149–323) (Delzell et al. 1996). When workers were divided by year of hire and age at death, leukemia deaths were elevated in workers who were hired between 1950 and 1959 (SMR 200; 95% CI 122–310) and who were < 55 years of age at the time of death (SMR 179; 95% CI 104–287). Using calculated estimates of exposure levels to 1,3-butadiene, styrene, and benzene, Macaluso et al. (1996) found that 75% of the cohort was exposed to 1,3-butadiene with a median cumulative exposure of 11.2 ppm-years, 83% of the cohort was exposed to styrene with a median cumulative exposure of 7.4 ppm-years, and 25% of the cohort was exposed to benzene with a cumulative exposure of 2.9 ppm-years. Among workers with leukemia, 86% had 1,3-butadiene exposure and 90% had styrene exposure; median cumulative exposure levels of 1,3-butadiene and styrene were about 3 times higher than the rest of the cohort. Workers with a cumulative exposure of 20–79 ppm 1,3-butadiene had a relative risk of leukemia mortality (after adjustment by race, age, and cumulative styrene exposure) that was 50% higher than workers with a cumulative exposure of 0.1–19 ppm and workers with a cumulative exposure of > 80 ppm had a 70% higher relative risk than the low exposure group; the progressive of relative risk with increasing cumulative exposure was statistically significant. Although a similar progression was observed for cumulative styrene exposure, the trend was not statistically significant. A follow-up to the Delzell et al. (1996) study, which tracked deaths for an additional 7 years (Sathiakumar et al. 2005), found similar results. A significant increase in deaths from leukemia was observed among hourly workers employed for > 10 years and hired 20–29 years earlier (SMR 258; 95% CI 156–403). Increases in deaths for colorectal cancer among workers employed for > 10 years and hired 20–29 years earlier (SMR 147; 95% CI 103–205) and deaths from prostate cancer among workers employed for < 10 years and hired > 30 years earlier (SMR 155; 95% CI 113–206). Significant increases deaths from leukemia were observed in

3. HEALTH EFFECTS

workers involved in polymerization, coagulation, and finishing processes, maintenance workers, and laboratory workers; these workers had the highest potential exposure to 1,3-butadiene, styrene, and possibly dimethyldithiocarbamate. Subsequent analysis of these data using updated exposure assessments (Cheng et al. 2007; Delzell et al. 2001; Graff et al. 2005) found that the increased risk of leukemia was positively associated with 1,3-butadiene exposure. Positive associations between cumulative 1,3-butadiene exposure (ppm-years) and leukemia and between cumulative styrene exposure and leukemia were observed; the associations were only statistically significant at the highest cumulative exposure levels for 1,3-butadiene (≥ 362.2 ppm-years) or styrene (≥ 60.4 ppm) (Delzell et al. 2001). However, when the relative risks were adjusted for 1,3-butadiene and dimethyldithiocarbamate cumulative exposure, cumulative styrene exposure was no longer significantly associated with leukemia (Delzell et al. 2001; Graff et al. 2005). Because styrene, 1,3-butadiene, and dimethyldithiocarbamate exposure were correlated, it is difficult to separate the risks for each individual compound.

Several population-based studies have examined the possible carcinogenicity of styrene. A case-control study found a significant increase in prostate cancer (odds ratio of 5.5; 95% CI 1.4–21.8) and rectal cancer (odds ratio of 5.1; 95% CI 1.4–19.4) among workers with medium to high exposure to styrene (Gerin et al. 1998). Workers in the following professions were considered to have medium to high styrene exposure: motor vehicle painters, motor vehicle repairers, firemen, and plastic mouldmakers. Another study found a significant increase in the incidence of rectal cancer (SIR 3.11; 95% CI 1.14–6.77) among individuals with occupational exposure to styrene (Antilla et al. 1998). A limitation of both of these studies is the lack of exposure information, including levels of styrene and confounding exposure to other chemicals; thus, it is difficult to ascribe the increased cancer risks to styrene exposure. Loughlin et al. (1999) examined former students who attended a high school adjacent to synthetic styrene-butadiene rubber production facilities between 1963 and 1993 and found no significant alterations in deaths from lymphatic and hematopoietic cancer. Two studies have examined the possible association between styrene exposure and breast cancer. A case-control study by Cantor et al. (1995) found significant elevations in the risk of breast cancer among women possibly exposed to styrene in the workplace. Coyle et al. (2005) found a significant higher incidence of age-adjusted breast cancer rate in men and women, women, and women ≥ 50 years of age and living in counties with EPA toxics release inventory (TRI) facilities with on-site releases of styrene. As with the other population-based studies, these studies did not monitor styrene levels or exposure to other potentially carcinogenic chemicals and thus provided limited information on the carcinogenic potential of styrene.

3. HEALTH EFFECTS

The carcinogenicity of styrene has been examined in three studies in rats (Conti et al. 1988; Cruzan et al. 1998; Jersey et al. 1978; Maltoni et al. 1982) and one study in mice (Cruzan et al. 2001). No significant increases in the incidence neoplastic lesions were observed in rats exposed to styrene concentrations as high as 1,000 ppm 6 hours/day, 5 days/week for 2 years (Cruzan et al. 1998). Similarly, exposure of female rats to 600 or 1,000 ppm styrene 6 hours/day, 5 days/week for 21 months did not result in styrene-related increases in the incidence neoplastic tumors (Jersey et al. 1978); a high incidence of chronic murine pneumonia in the control and 1,000 ppm male rats precludes the use of the male data for assessing the carcinogenic potential of styrene. There was a significant trend (Cochran-Armitage test conducted by ATSDR) for increased incidence of malignant mammary tumors in female rats exposed to styrene 4 hours/day, 5 days/week for 52 weeks (Conti et al. 1988); the incidences were 6/60, 6/30, 4/30, 9/30, 12/30, and 9/30 in the 0, 25, 50, 100, 200, and 300 ppm groups, respectively. No other significant increases in specific tumors were observed in this study (Conti et al. 1988; Maltoni et al. 1982). The findings of the Conti et al. (1988) study conflict with those of Cruzan et al. (1998), who found a concentration-related decrease in mammary tumors in female rats exposed to similar or higher styrene concentrations for a longer duration (20/60, 13/44, 9/43, 5/38, and 2/59 in female rats exposed to 0, 50, 200, 500, or 1,000 ppm, respectively). The decrease in body weight observed in the female rats exposed to ≥ 200 ppm may have influenced the occurrence of mammary tumors. In contrast to the results in rat studies, significant increases in the incidence of bronchioalveolar carcinoma were observed in female mice exposed to 160 ppm 6 hours/day, 5 days/week for approximately 2 years (Cruzan et al. 2001). The incidences of bronchioalveolar carcinoma were 0/50, 0/50, 2/50, 0/50, and 7/50 in the 0, 20, 40, 80, and 160 ppm female mice, respectively). Significant trends for increasing incidences of bronchioalveolar adenoma were also observed for the male and female mice; the respective incidence of adenomas was 15/50, 21/50, 35/50, 30/50, and 33/50 in males and 6/50, 16/50, 16/50, 11/50, and 24/50 in females. The incidence of adenoma was significantly higher than controls in males exposed to 40, 80, or 160 ppm and in females exposed to 20, 40, or 160 ppm.

As discussed by IARC (2002) and Cruzan et al. (2002), the lung tumors observed in the mice are likely due to the *in situ* formation of styrene 7,8-oxide resulting in cytotoxicity and increased cell proliferation. The relevance of these tumors to humans has been questioned due to species differences in the metabolism of styrene in the lungs. In rats and mice, Clara cells have the ability to metabolize styrene to styrene 7,8-oxide in the lung, whereas humans have limited ability to metabolize styrene to styrene 7,8-oxide in the lung. A physiologically based pharmacokinetic (PBPK) model predicted that the total amount of styrene oxide in the terminal bronchioles in mice is 10 times higher than in rats and 100-fold higher than in humans. In addition to these quantitative differences in the generation of styrene 7,8-oxide

3. HEALTH EFFECTS

between rats and mice, there are qualitative differences in styrene metabolism. Mice produce higher levels of the R-enantiomer of styrene oxide, as compared to rats; the R-enantiomer has been shown to be more potent pneumotoxic than the S-enantiomer. The ratio of R- to S-enantiomers ranges from 2.2 to 2.87 in mice exposed to 20–160 ppm styrene and from 0.7 to 0.73 in rats exposed to 50–1,000 ppm. Thus, mice appear to be very sensitive to the induction of lung tumors and the mechanism of inducing lung tumors is not likely to be relevant to humans. Although the mechanism involved in the development of lung tumors in mice may not be applicable to humans, other mechanisms of styrene carcinogenicity may be relevant for humans.

3.2.2 Oral Exposure

No studies were located regarding health effects in humans after oral ingestion of styrene. Based on the animal data that follow, the oral toxicity of styrene in humans would be expected to be low to moderate.

3.2.2.1 Death

No deaths in humans from ingesting styrene have been reported in the evaluations of case studies (EPA 1989c; Gosselin et al. 1984; NIOSH 1983).

The approximate reported oral LD₅₀ for male and female rats was 5,000 mg/kg (Wolf et al. 1956). A 100% survival rate and 100% mortality rate were reported in rats exposed to single oral doses of styrene (observation period 2 weeks) at 1,600 and 8,000 mg/kg, respectively (Spencer et al. 1942). Death in this study was mainly due to pronounced irritation of the esophagus and stomach. In another study, female mice were given a single oral dose of 1,350 mg/kg styrene on the 17th day of pregnancy (Ponomarkov and Tomatis 1978). After weaning, the progeny received the same dose once per week. The treatment was suspended after 16 weeks due to high mortality among the progeny (including both males and females). Fifty percent of the males and 20% of the females had died after 20 weeks, despite the suspension of treatment at week 16. The cause of death was liver necrosis and lung congestion. A high mortality rate was reported in 40 female rats exposed to 250 mg/kg/day styrene for 52 weeks (Conti et al. 1988). Mortality was significantly elevated in male and female rats administered styrene by gavage at a dosage level of 2,000 mg/kg/day for 78 weeks (NCI 1979b). In this study, mortality was unaffected at dosage levels of 500 and 1,000 mg/kg/day in male and female rats. Male mice administered styrene at doses of 150 or 300 mg/kg/day for 78 weeks showed increased mortality; however, the female mice did not.

3. HEALTH EFFECTS

The highest reliable LOAEL values and LD₅₀s values in each species and duration category are recorded in Table 3-3 and plotted in Figure 3-2.

3.2.2.2 Systemic Effects

No studies were located regarding endocrine, metabolic, musculoskeletal, or dermal/ocular effects in humans or animals after oral exposure to styrene.

For the following systemic effects resulting from oral exposure to styrene, the highest NOAEL values and all reliable LOAEL values for each species and duration category are recorded in Table 3-3 and plotted in Figure 3-2.

Respiratory Effects. No studies were located regarding respiratory effects in humans after oral exposure to styrene.

Severe lung congestion was observed in mice that were the offspring of dams given a single oral dose of styrene at 1,350 mg/kg on the 17th day of gestation and that continued to receive the same dose once per week after weaning (Ponomarkov and Tomatis 1978). The lung congestion was noted following 16 weeks of styrene administration. No respiratory effects were observed in rats exposed to 35 mg/kg/day styrene in drinking water for 105 weeks (Beliles et al. 1985).

Cardiovascular Effects. No studies were located regarding cardiovascular effects in humans following oral exposure to styrene.

No cardiovascular effects were observed in rats chronically exposed to 35 mg/kg/day in drinking water (Beliles et al. 1985).

Gastrointestinal Effects. Abdominal pain was reported by 11% of the residents of two apartment buildings exposed to elevated levels of styrene in drinking water for 3 days (Arnedo-Pena et al. 2003). The concentration of styrene in the water was 900 µg/L and the dose was estimated to be 0.026 mg/kg/day. Based on other symptoms and the higher prevalence of symptoms among residents living near the contaminated water tank, it is likely that some of the observed effects were due to inhalation exposure of styrene vapors from the repair of a firewater tank adjacent to the drinking water tank.

Table 3-3 Levels of Significant Exposure to Styrene - Oral

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
ACUTE EXPOSURE								
Death								
1	Rat	1 d (GO)				5000 (LD50)	Wolf et al. 1956 Styrene	
Neurological								
2	Rat	once (GO)			200 M (increased dopamine receptor binding)		Agrawal et al. 1982 Styrene	
3	Rat (Wistar)	14 d 1 x/d (GO)			100 M ^b (impaired learning)		Husain et al. 1985 Styrene	
Developmental								
4	Rat (Sprague-Dawley)	Gd 11 (GO)		300			Daston et al. 1991 Styrene	
5	Rat (Sprague-Dawley)	2 x/d Gd 6-15 (GW)		300 F			Murray et al. 1978 Styrene	
INTERMEDIATE EXPOSURE								
Immuno/ Lymphoret								
6	Rat (UF)	5 d/wk 4 wk (GO)		196 M	294 M (impaired immune response)		Dogra et al. 1992 Styrene	
7	Mouse (Swiss)	5 d/wk 4 wk (GO)		23 M	30 M (impaired immune response)		Dogra et al. 1992 Styrene	

DRAFT FOR PUBLIC COMMENT

Table 3-3 Levels of Significant Exposure to Styrene - Oral

(continued)

Key to Figure	Species ^a (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
Neurological								
8	Rat	90 d 1 x/d (GO)			200 M (increased dopamine receptor binding)		Agrawal et al. 1982 Styrene	
9	Rat (Long- Evans)	5 d/wk 8 wk (GO)			500 M (impaired learning)		Bushnell 1994 Styrene	
10	Rat	15 d 1 x/d (G)			906 M (increased serotonin and noradrenaline and decreased monoamine oxidase levels)		Husain et al. 1980 Styrene	
11	Rat (Wistar)	15 days (GO)		250 F			Khanna et al. 1994 Styrene	
Reproductive								
12	Rat	90 d (continuous) (W)		35			Beliles et al. 1985 Styrene	
13	Rat	60 d 6 d/wk 1 x/d (GO)		200 M		400 M (marked degeneration of seminiferous tubules, decreased spermatozoa)	Srivastava et al. 1989 Styrene	
Developmental								
14	Rat	Ld 1-21 (GO)		200 M	400 M (altered testicular enzyme levels and decreased spermatozoa counts)		Srivastava et al. 1992a Styrene	

DRAFT FOR PUBLIC COMMENT

STYRENE

3. HEALTH EFFECTS

78

Table 3-3 Levels of Significant Exposure to Styrene - Oral

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	NOAEL (mg/kg/day)	LOAEL		Reference Chemical Form	Comments
					Less Serious (mg/kg/day)	Serious (mg/kg/day)		
15	Rat (Wistar)	6 d/wk pnd 1-61 (GO)		100 M	200 M (decreased testes weight and spermatozoa counts)		Srivastava et al. 1992b Styrene	
16	Rat (NS)	Gd 1-21, Gd 1- Ld 14-21, or LD 1- Ld 14-21 (GO)			200 (increased dopamine receptor binding)		Zaidi et al. 1985 Styrene	
Cancer								
17	Mouse	16 wk 1 d/wk (GO)				1350 (CEL: lung tumors)	Ponomarev and Tomatis 1978 Styrene	
CHRONIC EXPOSURE								
Death								
18	Rat	78 wk 5 d/wk 1 x/d (GO)				2000 (decreased survival in males and females)	NCI 1979b Styrene	

DRAFT FOR PUBLIC COMMENT

Table 3-3 Levels of Significant Exposure to Styrene - Oral

(continued)

Key to Figure ^a	Species (Strain)	Exposure/Duration/Frequency (Route)	System	LOAEL		Reference Chemical Form	Comments
				NOAEL (mg/kg/day)	Less Serious (mg/kg/day)		
Systemic							
19	Rat	105 wk 7 d/wk (W)	Resp	35			Beliles et al. 1985 Styrene
			Cardio	35			
			Gastro	35			
			Hemato	35			
			Musc/skel	35			
			Hepatic	35			
			Renal	35			
			Dermal	35			
			Other	35			
20	Rat	120 wk 1 d/wk 1 x/d (GO)	Hepatic	500			Ponomarkov and Tomatis 1978 Styrene
			Renal	500			
21	Dog	561 d 1 x/d (GO)	Hemato	200	400 (Heinz body formation)		Quast et al. 1979 Styrene
Cancer							
22	Mouse	78-103 wk 5 d/wk 1 x/d (GO)				300 (CEL: lung tumors)	NCI 1979b Styrene

a The number corresponds to entries in Figure 3-2.

b The acute-duration oral MRL of 0.1 mg/kg/day; dose divided by an uncertainty factor of 1000 (10 for use of a LOAEL, 10 for extrapolation from animals to humans and 10 human variability).

Cardio = cardiovascular; CEL = cancer effect level; d = day(s); F = Female; (G) = gavage; Gastro = gastrointestinal; Gd = gestational day; (GO) = gavage in oil; (GW) = gavage in water; Hemato = hematological; LD50 = lethal dose, 50% kill; LOAEL = lowest-observed-adverse-effect level; M = male; min = minute(s); mo = month(s); Musc/skel = musculoskeletal; NOAEL = no-observed-adverse-effect level; NS = not specified; pnd = post-natal day; Resp = respiratory; x = time(s); (W) = drinking water; wk = week(s)

DRAFT FOR PUBLIC COMMENT

STYRENE

3. HEALTH EFFECTS

Figure 3-2 Levels of Significant Exposure to Styrene - Oral
Acute (≤14 days)




Figure 3-2 Levels of Significant Exposure to Styrene - Oral (Continued)
Intermediate (15-364 days)





Figure 3-2 Levels of Significant Exposure to Styrene - Oral (Continued)
Chronic (≥365 days)



DRAFT FOR PUBLIC COMMENT

c-Cat	-Humans	f-Ferret	n-Mink	◆ Cancer Effect Level-Animals	▼ Cancer Effect Level-Humans	■ LD50/LC50
d-Dog	k-Monkey	j-Pigeon	o-Other	● LOAEL, More Serious-Animals	▲ LOAEL, More Serious-Humans	⋮ Minimal Risk Level
r-Rat	m-Mouse	e-Gerbil		◐ LOAEL, Less Serious-Animals	△ LOAEL, Less Serious-Humans	⋮ for effects
p-Pig	h-Rabbit	s-Hamster		○ NOAEL - Animals	△ NOAEL - Humans	⋮ other than
q-Cow	a-Sheep	g-Guinea Pig				⋮ Cancer

3. HEALTH EFFECTS

No gastrointestinal effects were observed in rats chronically exposed to 35 mg/kg/day styrene in drinking water (Beliles et al. 1985).

Hematological Effects. No studies were located regarding hematological effects in humans after oral exposure to styrene.

Intra-erythrocytic Heinz bodies were regularly detected in a dose-related manner in male and female dogs chronically exposed to 400 or 600 mg/kg/day groups and sporadically in females in the 200 mg/kg/day group (Quast et al. 1979). There were occasional decreased red blood cell counts, hemoglobin levels, and erythrocyte sedimentation rates in males and females in the 600 mg/kg/day groups. Increased hemosiderin deposits and intranuclear inclusions in liver were noted in animals dosed with 600 mg/kg/day. This was probably secondary to the effects on the red blood cells. The formation of intra-erythrocytic Heinz bodies was readily reversible upon discontinuing the administration of styrene in the 600 mg/kg/day group after 470 days of exposure. No hematological effects were observed in rats chronically exposed to 35 mg/kg/day (Beliles et al. 1985).

Hepatic Effects. No studies were located regarding hepatic effects in humans after oral exposure to styrene.

Some animal studies have reported hepatic effects; however, the inconsistency of the findings and poor reporting of the data preclude drawing conclusion on the hepatotoxicity of orally administered styrene. Small areas of focal necrosis was observed in the livers of rats administered 400 mg/kg styrene in groundnut oil 6 days/week for 100 days (Srivastava et al. 1982). Because the incidence or statistical analysis data were not reported, it is not possible to determine whether 400 mg/kg is an adverse effect level. This study also found alteration in mitochondrial and microsomal enzymes at 200 and 400 mg/kg; the significance of these alterations in the absence of histological damage is not known. An increase in liver weights was observed in rats administered 400 or 677 mg/kg via gavage 5 days/week for 6 months (Wolf et al. 1956); no histological alterations were observed. Although the alterations in the liver weight were considered slight at 400 mg/kg and moderate at 677 mg/kg, the magnitude of the change and statistical significance is not known; slight and moderate alterations in body weight were also observed at these dose levels. Hepatic glutathione content was reduced in rats orally administered 900 mg/kg styrene for 7 consecutive days (Das et al. 1981); the toxicological significance of this effect is not known. As noted above, increased numbers of hemosiderin deposits and intranuclear crystalline inclusions were

3. HEALTH EFFECTS

reported in the hepatocytes of dogs orally administered 600 mg/kg/day of styrene by gavage for 316 days (Quast et al. 1979). This was presumably secondary to Heinz body formation, and no other hepatic histological effects were in this study. No hepatic effects were observed in rats exposed to 35 mg/kg/day styrene in drinking water for 105 weeks (Beliles et al. 1985) or in rats administered 500 mg/kg styrene 1 day/week for 120 weeks (Ponomarkov and Tomatis 1978).

Renal Effects. No studies were located regarding renal effects in humans after oral exposure to styrene.

A decrease in renal glutathione content and decreased glutathione-S-transferase activity was noted in rats orally administered 900 mg/kg styrene for 7 days (Das et al. 1983). Growth depression and slightly increased kidney weight were reported in female rats administered 400 and 667 mg/kg, 5 days/week for 6 months (Wolf et al. 1956); the magnitude and statistical significance of the effect were not reported. Histopathological examination of kidney tissue showed no abnormalities; thus, the changes in organ weight were not considered adverse. In another study, female rats and mice were exposed to 1,350 mg/kg/day of styrene on the 17th day of gestation; the offspring were also administered styrene, by gavage, 1 day/week for 120 weeks. No statistically significant increases in the incidence of kidney lesions were observed in rats exposed to 500 mg/kg (Ponomarkov and Tomatis 1978). No histological alterations were observed in the kidneys of rats chronically exposed to 35 mg/kg/day styrene in drinking water (Beliles et al. 1985).

3.2.2.3 Immunological and Lymphoreticular Effects

No studies were located regarding immunological effects in humans after oral exposure to styrene.

The World Health Organization (WHO 1983) reviewed a Russian study conducted by Sinitskij in which styrene was fed to 36 rabbits at doses of 250 mg/kg for 58 days, 5 mg/kg for 216 days, and 0.5 mg/kg for 202 days. Impairment of the immunological defense system was indicated by a nearly total suppression of leukocyte phagocytic activity. Although no statistical analysis was provided, the data showed a dose-response relationship for both the severity of the effect and the time of onset. Similarly, impaired host resistance was observed in mice exposed to 30 mg/kg/day and infected with encephalomyocarditis or a rodent strain of malaria and rats exposed to 294 mg/kg/day and infected with a rodent hookworm parasite (Dogra et al. 1992).

3. HEALTH EFFECTS

3.2.2.4 Neurological Effects

No studies were located regarding neurological effects in humans following oral exposure to styrene.

Neurobehavioral effects and alterations in neurochemicals have been observed in animal studies. Significant learning impairment was observed in an operant behavioral test in rats administered 500 mg/kg styrene in corn oil, 5 days/week for 8 weeks (Bushnell 1994). A reversal of the effect was not observed 1 year after exposure termination. Another study found significantly increased mean percent avoidance response, indicative of impaired learning, in rats administered 100 or 200 mg/kg/day styrene for 14 days (Husain et al. 1985). No alterations in foot shock-induced aggressive behavior or amphetamine-induced motor activity were observed in young rats administered via gavage 250 mg/kg/day styrene for 15 days (Khanna et al. 1994). However, significant alterations were observed in similarly exposed rats maintained on a low protein diet (8% casein versus 20% in normal diet).

Significant increases in serotonin levels in the hypothalamus, hippocampus, and midbrain were observed in rats administered 200 mg/kg/day for 14 days (Husain et al. 1985); no alterations in dopamine or noradrenaline levels were observed. Exposure to a higher dose (906 mg/kg/day) for 15 days resulted in increases in serotonin and noradrenalin in brain tissue (Husain et al. 1980). Neither study found significant alterations in brain dopamine levels. Another study found a significant increase in dopamine receptor binding, as assessed using labeled spiroperidol binding, in rats administered 200 or 400 mg/kg/day for 1 day or 90 days (Agrawal et al. 1982).

The highest NOAEL and LOAEL values for neurological effects in each species and duration category are recorded in Table 3-3 and plotted in Figure 3-2.

3.2.2.5 Reproductive Effects

No studies were located regarding reproductive effects in humans after oral exposure to styrene.

Marked degeneration in the seminiferous tubules and decreased spermatozoa were observed in rats administered 400 mg/kg styrene via gavage 6 days/week for 60 days (Srivastava et al. 1989). No adverse reproductive effects were observed in a three-generation reproduction study in which rats were exposed to 35 mg/kg/day in drinking water (Beliles et al. 1985). In another study, marked degeneration of seminiferous tubules and decreased spermatozoa counts were observed in adult rats administered 400 mg/kg styrene via gavage 6 days/week for 60 days (Srivastava et al. 1989). Significant decreases in

3. HEALTH EFFECTS

sorbitol dehydrogenase and acid phosphatase levels and increases in lactate dehydrogenase, glucose-6-phosphate dehydrogenase, β -glucuronidase, and γ -glutamyl transpeptidase levels were also observed at this dose level; the investigators considered these enzymes to be markers for testicular function.

The highest NOAEL values and all reliable LOAEL values for reproductive effects in rats in the acute and intermediate duration categories are recorded in Table 3-3 and plotted in Figure 3-2.

3.2.2.6 Developmental Effects

No studies were located regarding developmental effects in humans after oral exposure to styrene.

No developmental effects were observed in the offspring of rats receiving a single gavage dose of 300 mg/kg on gestational day 11 (Daston et al. 1991) or in rats administered 300 mg/kg/day (150 mg/kg administered twice daily) on gestational days 6–15 (Murray et al. 1978).

Other developmental studies have focused on the impaired development of the reproductive system or neurodevelopmental effects. Decreases in spermatozoa counts (measured on postnatal days 61 and 91) were observed in offspring of rats administered to 400 mg/kg styrene on lactation days 0–21; no effects were observed at 200 mg/kg/day (Srivastava et al. 1992a). However, decreases in spermatozoa counts were observed in rats exposed to 200 mg/kg 6 days/week on postnatal days 1–61 (Srivastava et al. 1992b). Both studies found significant alterations in testicular enzyme levels at the same dose level as spermatozoa effects; these enzymes were considered markers of testicular function. A decrease in dopamine receptor binding, as assessed using labeled spiroperidol, was observed in the offspring of rats administered to 200 mg/kg/day styrene throughout gestation and lactation or only during lactation, but was not observed in rat pups only observed during gestation (Zaidi et al. 1985). This decrease in receptor binding was attributed to an increase in the number of dopamine receptors rather than an alteration in binding affinity. Impaired amphetamine-induced locomotor activity and apomorphine-induced stereotypy were also observed in the pups exposed during gestation and lactation.

The highest NOAEL value for developmental effects is recorded in Table 3-3 and plotted in Figure 3-2.

3.2.2.7 Cancer

No studies were located regarding cancer effects in humans after oral exposure to styrene.

3. HEALTH EFFECTS

Investigations of the carcinogenic potential of styrene in animals after oral exposure have yielded variable results. No significant alterations in the incidence of neoplastic tumors were observed in rats exposed to gavage doses as high as 250 mg/kg 4–5 days/week for 52 weeks (Conti et al. 1988; Maltoni et al. 1982) or 2,000 mg/kg 5 days/week for 78–103 weeks (NCI 1979b) or in rats exposed to 35 mg/kg/day styrene in drinking water for 2 years (Beliles et al. 1985). In contrast, significant increases in the incidence of lung tumors were observed in mice receiving gavage doses of 300 mg/kg 5 days/week for 78–103 weeks (NCI 1979b). The incidences of bronchiolo-alveolar carcinoma in male mice were 0/20, 3/44, and 5/43 in mice exposed to 0, 150, or 300 mg/kg, respectively, and the respective combined incidences of bronchiolo-alveolar carcinoma and adenoma in male mice were 0/20, 6/44, and 9/43. The incidence of bronchiolo-alveolar carcinoma in the 300 mg/kg group was similar to the incidence in untreated historical controls (12%), but lower than the incidence in historical vehicle controls 0/40; however, the National Cancer Institute (NCI 1979b) noted that the incidence in historical vehicle controls is based on too small a number of animals for meaningful use of historical control data. Two studies conducted by Ponomarkov and Tomatis (1978) examined the carcinogenicity of styrene following gestation and postnatal exposure. In the offspring of mice administered 1,350 mg/kg on gestation day 17 with continued exposure of the weanling mice to this dose level (1 day/week) for 16 weeks, a significant increase in lung tumors was observed during the 100-week observation period; this dose was also associated with treatment-related toxicity and mortality. In the second study, the mice were exposed to 300 mg/kg on gestation day 17 followed by exposure to 300 mg/kg for 120 weeks (1 day/week) beginning at weaning. No significant alterations in tumor incidence were observed.

3.2.3 Dermal Exposure

No studies were located regarding health effects in humans after dermal exposure to styrene.

3.2.3.1 Death

No studies were located regarding lethality in humans or animals after dermal exposure to styrene.

3.2.3.2 Systemic Effects

No studies were located regarding respiratory, cardiovascular, gastrointestinal, hematological, musculoskeletal, hepatic, endocrine, or renal effects in humans or animals after dermal exposure to styrene.

3. HEALTH EFFECTS

Dermal Effects. Marked irritation with denaturation of the skin was noted when styrene was applied in small amounts over a 4-week period to the shaved abdomen of rabbits at 20,000 mg/kg total dose) (Spencer et al. 1942).

Ocular Effects. Moderate conjunctival irritation and transient corneal injury of the eyes were observed when undiluted styrene was tested in rabbit eyes (Wolf et al. 1956). The effects were produced immediately (within 3 minutes) by a single administration of two drops (about 0.1 mL) and persisted throughout the 7-day observation period.

No studies were located regarding the following health effects in humans or animals after dermal exposure to styrene:

3.2.3.3 Immunological and Lymphoreticular Effects

3.2.3.4 Neurological Effects

3.2.3.5 Reproductive Effects

3.2.3.6 Developmental Effects

3.2.3.7 Cancer

3.3 GENOTOXICITY

The genotoxicity of styrene has been examined in numerous *in vivo* studies of workers and laboratory animals; these data are summarized in Table 3-4. Chromosomal damage, DNA strand breaks, and mutagenic effects have frequently been studied in workers exposed to styrene in the production of reinforced plastic products and styrene/polystyrene production. In general, these studies are limited by the fact that workers in these industries are often exposed to chemicals other than styrene, such as methylene chloride and epoxide resins, and many studies did not control for potential confounding factors such as age, sex, and smoking status. Chromosomal aberrations have been reported in numerous studies of workers exposed to styrene for 1–25 years in reinforced plastic operations (Anwar and Shamy 1995; Artuso et al. 1995; Hogstedt et al. 1979; Mäki-Paakkanen et al. 1991; Meretoja et al. 1977, 1978; Somorovská et al. 1999; Tomanin et al. 1992); however, other studies have not found significant increases in chromosomal aberrations (Andersson et al. 1980; Hansteen et al. 1984; Jablonicka et al. 1988; Mäki-Paakkanen 1987; Nordenson and Beckman 1984; Thiess et al. 1980; Vodicka et al. 2004; Watanabe et al. 1981). The results of the Artuso et al. (1995) and Tomanin et al. (1992) studies provide suggestive evidence that the occurrence of chromosomal aberrations is concentrated-related. Significant increases in

3. HEALTH EFFECTS

Table 3-4. Genotoxicity of Styrene *In Vivo*

Species (test system)	End point	Results	Reference
Human Studies			
Human lymphocytes	Chromosomal aberrations	+	Meretoja et al. 1977
Human lymphocytes	Chromosomal aberrations	+	Meretoja et al. 1978
Human lymphocytes	Chromosomal aberrations	+	Hogstedt et al. 1979
Human lymphocytes	Chromosomal aberrations	+	Anwar and Shamy 1995
Human lymphocytes	Chromosomal aberrations	+	Artuso et al. 1995
Human lymphocytes	Chromosomal aberrations	+	Mäki-Paakkanen et al. 1991
Human lymphocytes	Chromosomal aberrations	+	Tomanin et al. 1992
Human mononuclear leukocytes	Chromosomal aberrations	+	Somorovská et al. 1999
Human lymphocytes	Chromosomal aberrations	–	Thiess et al. 1980
Human lymphocytes	Chromosomal aberrations	–	Andersson et al. 1980
Human lymphocytes	Chromosomal aberrations	–	Watanabe et al. 1981
Human lymphocytes	Chromosomal aberrations	–	Nordenson and Beckman 1984
Human lymphocytes	Chromosomal aberrations	–	Hansteen et al. 1984
Human lymphocytes	Chromosomal aberrations	–	Maki-Paakkanen 1987
Human lymphocytes	Chromosomal aberrations	–	Jablonicka et al. 1988
Human lymphocytes	Chromosomal aberrations	–	Vodicka et al. 2004
Human lymphocytes	Sister chromatid exchange	+	Yager et al. 1993
Human lymphocytes	Sister chromatid exchange	+	Laffon et al. 2002
Human lymphocytes	Sister chromatid exchange	+	Karakaya et al. 1997
Human lymphocytes	Sister chromatid exchange	+	Artuso et al. 1995
Human lymphocytes	Sister chromatid exchange	+	Hallier et al. 1994
Human lymphocytes	Sister chromatid exchange	+	Andersson et al. 1980
Human lymphocytes	Sister chromatid exchange	–	Mäki-Paakkanen et al. 1991
Human lymphocytes	Sister chromatid exchange	–	Hansteen et al. 1984
Human lymphocytes	Sister chromatid exchange	–	Watanabe et al. 1981
Human lymphocytes	Sister chromatid exchange	–	Maki-Paakkanen 1987
Human lymphocytes	Micronuclei	+	Nordenson and Beckman 1984
Human lymphocytes	Micronuclei	+	Hogstedt et al. 1983
Human lymphocytes	Micronuclei	+	Laffon et al. 2002
Human lymphocytes	Micronuclei	–	Karakaya et al. 1997

3. HEALTH EFFECTS

Table 3-4. Genotoxicity of Styrene *In Vivo*

Species (test system)	End point	Results	Reference
Human lymphocytes	Micronuclei	–	Anwar and Shamy 1995
Human lymphocytes	Micronuclei	–	Mäki-Paakkanen et al. 1991
Human lymphocytes	Micronuclei	–	Tomanin et al. 1992
Human lymphocytes	Micronuclei	–	Vodicka et al. 2004
Human erythrocytes	Mutations in glycophorin A	+	Bigbee et al. 1996
Human erythrocytes	Mutations in glycophorin A	+	Compton-Quintana et al. 1993
Human lymphocytes	HPRT mutations	±	Vodicka et al. 1995
Human mononuclear leukocytes	DNA Single strand breaks	+	Somorovská et al. 1999
Human lymphocytes	DNA Single strand breaks	+	Shamy et al. 2002
Human lymphocytes	DNA Single strand breaks	+	Mäki-Paakkanen et al. 1991
Human leukocytes	DNA Single strand breaks	+	Walles et al. 1993
Human lymphocytes	DNA Single strand breaks	+	Vodicka et al. 1995
Human lymphocytes	DNA Single strand breaks	–	Vodicka et al. 2004
Human lymphocytes	Unscheduled DNA synthesis	+	Pero et al. 1982
Laboratory animal studies			
Mouse bone marrow	Chromosomal aberrations	–	Sbrana et al. 1983
Rat bone marrow	Chromosomal aberrations	–	Sinha et al. 1983
Mouse bone marrow	Sister chromatid exchange	±	Simula and Priestly 1992
Mouse spleen, lung, erythrocytes	Sister chromatid exchange	+	Kligerman et al. 1992, 1993
Mouse bone marrow, liver cells, and alveolar macrophages	Sister chromatid exchange	+	Conner et al. 1980
Mouse bone marrow	Sister chromatid exchange	±	Simula and Priestly 1992
Rat lymphocyte	Sister chromatid exchange	+	Kligerman et al. 1992, 1993
Mouse bone marrow and polychromatic erythrocytes	Micronuclei	±	Norppa 1981
Mouse bone marrow	Micronuclei	±	Simula and Priestly 1992
Mouse bone marrow	Micronuclei	–	Engelhardt et al. 2003
Mouse spleen, lung, erythrocytes	Micronuclei	–	Kligerman et al. 1992, 1993
Mouse bone marrow	Micronuclei	–	Simula and Priestly 1992
Rat lymphocyte	Micronuclei	–	Kligerman et al. 1992, 1993
Mouse bone marrow and lymphocytes	DNA single strand breaks	–	Vodicka et al. 2001a
Mouse kidney, liver, lung, testes, and brain	DNA	+	Walles and Orsen 1983

3. HEALTH EFFECTS

Table 3-4. Genotoxicity of Styrene *In Vivo*

Species (test system)	End point	Results	Reference
Mouse liver	Unscheduled DNA synthesis	–	Clay 2004

– = negative result; + = positive result; ± = weakly positive result; DNA = deoxyribonucleic acid

3. HEALTH EFFECTS

chromosomal aberrations alterations were observed in high exposure groups (20–326 or 27–104 ppm), but not in workers exposed to lower exposure levels (0.5–24 ppm). As with the occurrence of chromosomal aberrations, mixed results have been observed in studies of sister chromatid exchange (Andersson et al. 1980; Artuso et al. 1995; Hallier et al. 1994; Hansteen et al. 1984; Karakaya et al. 1997; Laffon et al. 2002; Mäki-Paakkanen 1987; Mäki-Paakkanen et al. 1991; Watanabe et al. 1981; Yager et al. 1993) and micronuclei formation (Anwar and Shamy 1995; Hogstedt et al. 1983; Karakaya et al. 1997; Laffon et al. 2002; Mäki-Paakkanen et al. 1991; Nordenson and Beckman 1984; Tomanin et al. 1993; Vodicka et al. 2004) in styrene workers. Other genotoxicity studies in styrene workers have found significant increases in glycophorin A mutations (Bigbee et al. 1996; Compton-Quintana et al. 1993), DNA single strand breaks (Mäki-Paakkanen et al. 1991; Shamy et al. 2002; Somorovská et al. 1999; Walles et al. 1993; Vodicka et al. 1995), and increases in unscheduled DNA synthesis (Pero et al. 1982).

Studies in laboratory animals have found significant increases in the occurrence of sister chromatid exchange (Conner et al. 1980; Kligerman et al. 1992, 1993; Simula and Priestly 1992). However, the results for chromosomal aberrations (Sbrana et al. 1983; Sinha et al. 1983), micronuclei formation (Engelhardt et al. 2003; Kligerman et al. 1992, 1993; Norppa 1981; Simula and Priestly 1992), DNA single strand breaks (Vodicka et al. 2001a), and unscheduled DNA synthesis (Clay 2004) have been weakly positive or negative.

Styrene has been tested for genotoxic potential in a variety of *in vitro* systems, as summarized in Table 3-5. In the absence of metabolic activation, styrene has not produced gene mutations in *Salmonella typhimurium* (DeMeester et al. 1981; Dunkel et al. 1985; Vainio et al. 1976) or *Escherichia coli* (Dunkel et al. 1985); mixed results have been found in the presence of metabolic activation. Increases in chromosomal aberrations (Jantunen et al. 1986) and sister chromatid exchange (Norppa et al. 1983) have been found in human lymphocytes in the absence of metabolic activation.

3.4 TOXICOKINETICS

3.4.1 Absorption

3.4.1.1 Inhalation Exposure

The uptake of styrene following inhalation exposure in humans and animals is rapid (Ramsey and Andersen 1984; Ramsey and Young 1978; Ramsey et al. 1980; Withey and Collins 1979; Withey and Karpinski 1985). Pulmonary retention of inhaled styrene in humans is approximately 2/3 of the

3. HEALTH EFFECTS

Table 3-5. Genotoxicity of Styrene *In Vitro*

Species (test system)	End point	Results		Reference
		With activation	Without activation	
Prokaryotic organisms:				
<i>Salmonella typhimurium</i> (two strains, plate incorporation method)	Gene mutation	+	–	Vainio et al. 1976
<i>S. typhimurium</i> (three strains, plate incorporation method)	Gene mutation	–	–	Vainio et al. 1976
<i>S. typhimurium</i> (three strains, vapor exposure – disiccator test)	Gene mutation	+	–	DeMeester et al. 1981
<i>S. typhimurium</i> (four strains, vapor exposure – disiccator test)	Gene mutation	–	–	DeMeester et al. 1981
<i>S. typhimurium</i> (five strains, preincubation method)	Gene mutation	–	–	Dunkel et al. 1985
<i>Escherichia coli</i> (one strain, preincubation method)	Gene mutation	–	–	Dunkel et al. 1985
Mammalian cells:				
Human lymphocytes	Sister chromatid exchange	No data	+	Norppa et al. 1983
Human lymphocytes	Chromosomal aberrations	No data	+	Jantunen et al. 1986

– = negative result; + = positive result

3. HEALTH EFFECTS

administered concentrations (Engstrom et al. 1978a, 1978b). For example, male human subjects were exposed to styrene in inspired air during 30-minute rest and three 30-minute work periods on a bicycle ergometer. The mean uptake was approximately 63% (range was 59–70%) of the amount of inspired styrene. In exercising volunteers exposed to 50 ppm styrene for 2 hours, an average of 66.5% of the inhaled styrene was absorbed (Johanson et al. 2000). Another study in volunteers exposed to 50 ppm styrene for 2 hours during exercise calculated that 64% of the styrene was absorbed (Norstöm et al. 1992). Exposures of rats to styrene concentrations of 50–2,000 ppm for 5 hours yielded blood uptakes that showed a continued and increasing rapid absorption, proportional to the styrene air level (Withey and Collins 1979). Plateau levels of styrene in rats' blood were reached within 6–8 hours during exposures ranging from 80 to 1,200 ppm styrene for up to 24 hours (Ramsey and Young 1978).

3.4.1.2 Oral Exposure

No studies were located regarding absorption in humans after oral exposure to styrene.

The absorption of styrene from the gastrointestinal tract was rapid and complete in rats deprived of food overnight and administered, via gavage, 9.3 mg/kg styrene in aqueous solution. A peak blood level of 6 µg/mL was reached in a few minutes. There was a much slower uptake of the styrene administered in vegetable oil (Withey 1976). Styrene administered in vegetable oil at a total dose of 32.61 mg produced a peak level of 12 µg/mL. This was reached at about 100 minutes (Withey 1976).

3.4.1.3 Dermal Exposure

Limited data indicate that absorption of styrene via the dermal route is probably low compared to absorption via other routes. When liquid styrene was applied to the forearms of male subjects, the absorption rate was estimated to be 9–15 mg/cm²/hour (Dutkiewicz and Tyras 1968). By contrast, the rate of absorption through human skin was very low (1±0.5 µg/cm²/minute) in subjects who dipped one hand into liquid styrene (Berode et al. 1985). The higher absorption rate reported by Dutkiewicz and Tyras (1968) likely included the disappearance rate of the solvent from the surface of the skin (Guillemin and Berode 1988). Riihimaki and Pfaffli (1978) demonstrated that in humans, dermal exposure to moderate concentrations of styrene vapor (300 and 600 ppm) resulted in percutaneous penetration corresponding to approximately 0.1–2% of the amount estimated to be absorbed from the respiratory tract. Similarly, Limasset et al. (1999) did not find significant differences in the levels of urinary metabolites in workers wearing total protective equipment (insulating suit and respiratory mask) and those wearing a respiratory mask only.

3. HEALTH EFFECTS

Although absorption of styrene applied to the abdomen of rabbits was reported, there was no information on absorption rates (Spencer et al. 1942). Dermal exposure of rats to a neat solution of styrene resulted in peak blood levels of 5.3 µg/mL within 1 hour of exposure (Morgan et al. 1991).

3.4.2 Distribution

A blood:air partition coefficient of 40.2 and fat:air partition coefficient of 3,476 were calculated for rats (Gargas et al. 1989). Fisher et al. (1997) calculated a human blood:air partition coefficient of 69.74 and a breast milk:blood partition coefficient of 2.17.

In a study of 81 adults without occupational exposure to styrene, average blood styrene levels were 0.221 µg/L; in comparison, blood styrene levels in reinforced plastics industry workers were 1,068–1,590 µg/L at the end of workshift and 60–119 µg/L in the morning after exposure (Brugnone et al. 1993).

3.4.2.1 Inhalation Exposure

Inhalation studies in both humans and animals resulted in the widespread distribution of styrene with the highest concentration in adipose tissue.

Three humans were exposed to 8–20 ppm styrene which resulted in a mean daily uptake of 193–558 mg styrene (Engstrom et al. 1978b). The concentration of styrene in adipose tissue was 2.8–8.1 mg/kg at the beginning of the week and 4.7–11.6 mg/kg at the end of the week. The authors estimated the half-life of styrene in the subcutaneous fat of humans to be about 72 hours. Subsequent studies by this author confirmed this estimate and reported the half-life of styrene in adipose tissue to be 2–5 days (Engstrom et al. 1978a).

Fiberglass factory workers exposed to >50 ppm of styrene for 8-hour work shifts had blood styrene levels which ranged from 120 to 684 µg/L at the end of the shift (Apostoli et al. 1983). The concentrations of urinary MA and phenylglyoxilic acid (PGA) were 133–2,100 and 107–685 mg/L, respectively. These levels were also determined at the end of the work-shift. Distribution of styrene was also studied in adult men exposed to about 70 ppm of styrene for 2 hours during light physical exercise (Wigaeus et al. 1983). Blood styrene reached a level of approximately 2,000 µg/L after 75 minutes. The concentration of styrene in adipose tissue was about 5,000 µg/kg after 30–90 minutes of exposure.

3. HEALTH EFFECTS

Rats were exposed for 5 hours to styrene at concentrations ranging from 50 to 2,000 ppm (Withey and Collins 1979). Tissue concentrations of styrene in the heart, liver, lung, kidney, spleen, brain, and perirenal fat demonstrated different patterns of distribution as the dose increased. The styrene concentration in perirenal fat was 10 times greater than in other organs. The largest amounts of styrene were found in the subcutaneous fat of male rats exposed to about 45 ppm of radioactively labeled styrene in the inspired air for 1–8 hours (Carlsson 1981). The concentration increased steadily during the first 4 hours of exposure. Styrene concentrations in brain tissue and muscles were about 70% of the arterial blood value. Other investigators (Ramsey and Andersen 1984; Ramsey and Young 1978; Savolainen and Pfaffli 1978; Withey 1976) demonstrated that higher levels of styrene in adipose tissue increase with higher exposures to styrene. Styrene was found to distribute to the fetuses of pregnant rats after inhalation exposure, but at concentrations much lower than those measured in maternal organs and tissues (Withey and Karpinski 1985).

3.4.2.2 Oral Exposure

No studies were located regarding distribution in humans after oral exposure to styrene.

An oral dose of 20 mg/kg of ¹⁴C styrene was administered to male and female rats (Plotnick and Weigel 1979). Tissue levels peaked at 4 hours or earlier after dosing. Less than 10% of the administered dose was found in the stomach, small intestine, and large intestine 8 hours after dosing. The kidney had the highest concentration of radioactivity at all time intervals, with decreasing amounts in the liver and pancreas. Fat tissue showed increased levels after 2 hours. All tissue levels were below 1 µg/g at 24 hours and at 48 and 72 hours were below the limit of detection. Excretion data from the Plotnick and Weigel (1979) study are presented in Section 3.4.4.2.

3.4.2.3 Dermal Exposure

No studies were located regarding distribution in humans after dermal exposure to styrene.

Immersion of rats' tails in pure liquid styrene for 1 hour resulted in styrene levels in the liver and brain that were estimated to be between 50 and 70% of the concentrations found in the same organs after 4-hour inhalation exposure to a vapor concentration of 11.8 g/m³ (Shugaev 1969). A skin:air partition coefficient of 91.9 was calculated for rat skin (Mattie et al. 1994).

3. HEALTH EFFECTS

3.4.3 Metabolism


There have been numerous *in vivo* studies, conducted primarily via inhalation, and *in vitro* studies that address the metabolism of styrene in humans and animals. There are several metabolic pathways for styrene (Cruzan et al. 2002; IARC 2002; Sumner and Fennel 1994), as illustrated in Figure 3-3. The primary pathway is oxidation of the side chain by cytochrome P450 to form styrene 7,8-oxide. Styrene oxide is predominantly metabolized by epoxide hydrolase to form styrene glycol; the styrene glycol is subsequently converted to mandelic acid, phenylglyoxylic acid, and hippuric acid. Styrene 7,8-oxide can also be conjugated with glutathione to ultimately form phenylhydroxyethylmercapturic acids. A minor pathway of styrene metabolism involves the formation of phenylacetaldehyde from styrene 7,8-oxide or cytochrome P450 conversion of styrene to phenylethanol and subsequent metabolism to phenylacetic acid. An alternative minor pathway involves ring oxidation resulting in the production of styrene 3,4-oxide, which is further metabolized to 4-vinylphenol.

As summarized by Cruzan et al. (2002), over 95% of the styrene urinary metabolites excreted by humans are derived from styrene glycol (mandelic acid, phenylglyoxylic acid, hippuric acid) compared to 49–59% in mice and 68–72% in rats. In mice and rats, 25–35% of the metabolites are derived from glutathione conjugation (mercapturic acids). The remaining metabolites derive from phenylacetic acid production (12–22% in mice and 3–5% in rats) and ring oxidation (4–8% in mice and <1% in rats). Trace amounts of mercapturic acids (DePalma et al. 2001) and 4-vinylphenol (Manini et al. 2002, 2003) have also been detected in humans; both metabolites each account for <1% of the total styrene metabolites.

The liver is the primary site of styrene metabolism and the source of styrene oxide in the blood (Cruzan et al. 2005). However, styrene is metabolized in other tissues, particularly the lung and nasal cavity following inhalation exposure, and it is this localized metabolism that results in the observed toxicity and/or carcinogenicity in these tissues. Studies in humans, mice, and rats indicate that styrene metabolism is concentration-dependent. At air concentrations of <200–300 ppm, most of the inhaled styrene is metabolized, only small amounts are exhaled unchanged, and there is little accumulation (Filser et al. 1993; Ramsey and Andersen 1984). At concentrations >300 ppm, metabolism was progressively limited by metabolic capacity and was saturated (V_{\max}) at 700 ppm in rats and at 800 ppm in mice (Filser et al. 1993). Löf and Johanson (1993) estimated that metabolic saturation occurs at 100–200 ppm in humans; however, a subsequent analysis by this group (Jonsson and Johanson 2002) found that the V_{\max} was 40% higher.

3. HEALTH EFFECTS

Figure 3-3. Scheme for Styrene Metabolism in Humans and Animals



GSH = glutathione

Source: Adapted from IARC 2002; Manini et al. 2002; Sumner and Fennel 1994

3. HEALTH EFFECTS

Comparisons of cytochrome P450, epoxide hydrolase, and glutathione S-transferase activity in the liver and lungs of rats and mice demonstrated 2–15-fold higher activities in the liver (Mendrala et al. 1993). In both tissues, previous exposure to styrene did not result in a dose-related increase in enzyme activity. In contrast, cytochrome P450 from human lung has a very limited ability to metabolize styrene to styrene oxide (Carlson et al. 2000; Nakajima et al. 1994); the capacity was 100-fold lower than in rat lung microsomes (Cruzan et al. 2002).

A number of cytochrome P450 isozymes have the capacity to catalyze styrene to styrene oxide. In human livers, CYP2B6 was the most active isoform; the activities of CYP1A2 and CYP2E1 were about half that of CYP2B6 (Nakajima et al. 1993). Kim et al. (1997) found that CYP2E1 was the main isoform at low styrene concentrations and CYP2B6 at high styrene concentrations. In mice and rats, the CYP1A1 and CYP2B1, respectively, were the most active isoforms in the liver (Nakajima et al. 1993). In the lungs, CYP2F1 was the most active isoform in human (Nakajima et al. 1994) lung microsomes. In mice and rat lung and nasal cavity microsomes, CYP2F2 and CYP2E1 are the predominant isoforms (Hynes et al. 1999).

Studies by Mendrala et al. (1993) compared the kinetic constants of cytochrome P450 from the livers of humans, mice, and rats. The affinity of cytochrome P450 for styrene (based on K_m values) was similar for the three species. However, the mouse had the greatest capacity to form styrene 7,8-oxide from styrene based on the V_{max} values and relative liver and body size, and humans had the lowest capacity. In contrast, marked differences in the K_m values for epoxide hydrolase were found between species; the K_m values were 0.01, 0.74, and 0.13–0.23 mmol for humans, mice, and rats, respectively. These results suggest that humans have a greater affinity to metabolize styrene 7,8-oxide and is more efficient at low levels of styrene oxide, as compared to rodent species.

Styrene oxide exists as two enantiomeric forms (R) and (S). As with other aspects of styrene metabolism, species differences in the ratio of R enantiomer and S enantiomer have been detected. In human liver samples, the R:S ratio was 0.15 at low styrene concentrations (0.016 mM) and 1.4 at high styrene concentrations (1.1 mM) (Wenker et al. 2001). In mouse and rat liver microsomes (incubated with 2 mM styrene), the R:S ratio was 0.57 and 1.18, respectively (Hynes et al. 1999). In the lungs, the R:S ratio was 2.4 and 0.52 for mice and rats, respectively.

3. HEALTH EFFECTS

3.4.4 Elimination and Excretion**3.4.4.1 Inhalation Exposure**

Several studies have demonstrated that styrene is almost totally excreted as urinary metabolites in humans, and at higher doses, the elimination profile indicates saturation of metabolic excretion or processes (Ramsey and Young 1978; Ramsey et al. 1980). Most of the inhaled styrene is excreted in urine as MA and PGA. In a study of the excretion of styrene and its metabolites resulting from a 100-ppm/8-hour inhalation exposure, 2.6% of the total uptake was excreted as unchanged styrene in exhaled air (Guillemin and Berode 1988). The metabolites MA, PGA, and hippuric acid were excreted in the urine at 56.9, 33, and 7.5% of the absorbed dose, respectively. In exercising volunteers exposed to 50 ppm styrene for 2 hours, 0.7–2.2% of the retained dose was exhaled as unchanged styrene (Johanson et al. 2000). Peak levels of styrene in the urine were measured immediately after exposure termination, whereas urinary excretion of MA and PGA peaked at 2 hours after exposure termination. MA excretion accounted for 6–29% of the estimated retained dose and PGA excretion accounted for 4–6%; the half-time excretion rates of MA and PGA were 2.2–4.2 and 3.5–13.9 hours, respectively. Phenylacetic acid and hippuric acid was also detected in the urine samples collected 2 hours after exposure termination. At this time point, MA account for 73% of the total excreted metabolites, PGA 18%, phenylacetic acid 4.5%, and hippuric acid 5.7%. In styrene workers exposed to 29–42 ppm styrene, both R-mandelic acid and S-mandelic acid were detected in the urine (Hallier et al. 1995). The ratio of R- to S-mandelic acid ranged from 0.7 to 1.2 in 19 of the 20 workers; in the last worker, the ratio was 2.2.

An alternative pathway for the metabolism of styrene 7,8-oxide is conjugation with glutathione, resulting in the excretion of mercapturic acids. Low levels of mercapturic acids have been detected in workers exposed to an unspecified amount of styrene (Maestri et al. 1997a). The mean concentrations of styrene metabolites were 580 mg/g creatinine mandelic acid, 174 mg/g creatinine phenylglyoxylic acid, 1.517 mg/g N-acetyl-S-(1-phenyl-2-hydroxyethyl)-cysteine S-enantiomer, 0.0637 mg/g N-acetyl-S-(1-phenyl-2-hydroxyethyl)-cysteine R- enantiomer, and 1.519 mg/g N-acetyl-S-(2-phenyl-2-hydroxyethyl)-cysteine. Another study of styrene workers (exposure level of 29–41 ppm) only detected styrene-specific mercapturic acid in 1 of 20 workers (Hallier et al. 1995). Similarly, in volunteers exposed to 50 ppm for 2 hours during exercise, N-acetyl-S-(2-phenyl-2-hydroxyethyl)-cysteine was not detected in urine samples collected up to 5 hours after exposure termination (Norström et al. 1992).

In volunteers exposed to 80 ppm styrene, styrene is cleared from the blood in a biphasic manner, indicating a two-compartment pharmacokinetic model. The half-lives for the rapid and slow clearance

3. HEALTH EFFECTS

phases are 0.58 and 13.0 hours, respectively. The half-life of styrene in subcutaneous adipose tissue of humans is 2–5 days (Engstrom et al. 1978a). The quantities of the major metabolites of styrene in urine compared with the quantity of styrene eliminated unchanged in expired air indicated that approximately 97% is cleared by the metabolic route (Ramsey et al. 1980).

Another human inhalation study determined that between 59 and 66% of inhaled styrene (50–200 ppm) was retained after a 4–8-hour exposure (Guillemin and Bauer 1979). Urinary elimination of MA was biphasic with a half-life for the first phase of 4 hours and for the second phase, 25 hours. These findings were comparable to those reported by Engstrom et al. (1976). The half-life of urinary elimination of PGA was determined to be 11 hours. This was regarded by the authors as being the first phase of elimination since MA is a precursor of PGA.

A lactational transfer pharmacokinetic model developed by Fisher et al. (1997) predicted that exposure to 50 ppm styrene would result in 0.650 mg styrene being ingested by a nursing infant over a 24-hour period.

Styrene is almost totally excreted as urinary metabolites in animals. The blood elimination curve for rats is biphasic exponential at 80 and 200 ppm styrene over 6 hours. For exposures >600 ppm exposure levels for 6 hours duration), a nonlinear blood elimination curve following Michaelis-Menten kinetics was observed. In going from 80 to 1,200 ppm (a 15-fold increase) the area under the blood concentration curves increases by 112-fold (Ramsey and Young 1978; Young et al. 1979). Rats exposed to 50–2,000 ppm styrene by inhalation for 5 hours exhibited a dose dependent biphasic pattern of elimination (Withey and Collins 1979).

3.4.4.2 Oral Exposure

No studies were located regarding excretion in humans after oral exposure to styrene.

Styrene was rapidly excreted in the urine of male and female rats administered 20 mg/kg of ¹⁴C styrene with 90% of the dose detected in the urine within 24 hours of administration (Plotnick and Weigel 1979). Less than 2% of the dose was found in the feces. Detectable tissue levels were not found 48 and 72 hours after administration. In mice administered 200 mg/kg for 70 days, 26.4, 13.3, and 19.0% of urinary metabolites excreted on day 70 were mandelic acid, phenylglyoxylic acid, and hippuric acid (Sbrana et al. 1983). Approximately 80% of the dose was excreted in the first 24 hours.

3. HEALTH EFFECTS

3.4.4.3 Dermal Exposure

In a study of the absorption of liquid styrene applied to the forearms of male volunteers, about 13% of the absorbed dose was excreted as MA (Dutkiewicz and Tyras 1968).

No studies were located regarding excretion in animals after dermal exposure to styrene.

3.4.5 Physiologically Based Pharmacokinetic (PBPK)/Pharmacodynamic (PD) Models

Physiologically based pharmacokinetic (PBPK) models use mathematical descriptions of the uptake and disposition of chemical substances to quantitatively describe the relationships among critical biological processes (Krishnan et al. 1994). PBPK models are also called biologically based tissue dosimetry models. PBPK models are increasingly used in risk assessments, primarily to predict the concentration of potentially toxic moieties of a chemical that will be delivered to any given target tissue following various combinations of route, dose level, and test species (Clewell and Andersen 1985). Physiologically based pharmacodynamic (PBPD) models use mathematical descriptions of the dose-response function to quantitatively describe the relationship between target tissue dose and toxic end points.

PBPK/PD models refine our understanding of complex quantitative dose behaviors by helping to delineate and characterize the relationships between: (1) the external/exposure concentration and target tissue dose of the toxic moiety, and (2) the target tissue dose and observed responses (Andersen and Krishnan 1994; Andersen et al. 1987a). These models are biologically and mechanistically based and can be used to extrapolate the pharmacokinetic behavior of chemical substances from high to low dose, from route to route, between species, and between subpopulations within a species. The biological basis of PBPK models results in more meaningful extrapolations than those generated with the more conventional use of uncertainty factors.

The PBPK model for a chemical substance is developed in four interconnected steps: (1) model representation, (2) model parameterization, (3) model simulation, and (4) model validation (Krishnan and Andersen 1994). In the early 1990s, validated PBPK models were developed for a number of toxicologically important chemical substances, both volatile and nonvolatile (Krishnan and Andersen 1994; Leung 1993). PBPK models for a particular substance require estimates of the chemical-specific physicochemical parameters, and species-specific physiological and biological parameters. The numerical estimates of these model parameters are incorporated within a set of differential and algebraic

3. HEALTH EFFECTS

equations that describe the pharmacokinetic processes. Solving these differential and algebraic equations provides the predictions of tissue dose. Computers then provide process simulations based on these solutions.

The structure and mathematical expressions used in PBPK models significantly simplify the true complexities of biological systems. If the uptake and disposition of the chemical substance(s) are adequately described, however, this simplification is desirable because data are often unavailable for many biological processes. A simplified scheme reduces the magnitude of cumulative uncertainty. The adequacy of the model is, therefore, of great importance, and model validation is essential to the use of PBPK models in risk assessment.


PBPK models improve the pharmacokinetic extrapolations used in risk assessments that identify the maximal (i.e., the safe) levels for human exposure to chemical substances (Andersen and Krishnan 1994). PBPK models provide a scientifically sound means to predict the target tissue dose of chemicals in humans who are exposed to environmental levels for example, levels that might occur at hazardous waste sites) based on the results of studies where doses were higher or were administered in different species. Figure 3-4 shows a conceptualized representation of a PBPK model.

If PBPK models for styrene exist, the overall results and individual models are discussed in this section in terms of their use in risk assessment, tissue dosimetry, and dose, route, and species extrapolations.

Several investigators have developed toxicokinetic models for styrene (Csanády et al. 1994, 2003; Jonsson and Johanson 2002; Leavens and Bond 1996). The Csanády et al. (1994, 2003) model is useful for evaluating the carcinogenic risk associated with inhalation exposure to styrene. As discussed in Section 3.2.1.7, species differences exist in the metabolism of styrene in the lungs of rats, mice, and humans; these differences result in increased sensitivity of mice. Jonsson and Johanson (2002) developed a population-based PBPK model for styrene, which decreased the intraindividual variability for estimating the metabolic capacity for styrene in humans. Leavens and Bond (1996) described initial work on developing a model for co-exposure to 1,3-butadiene and styrene in mice. Some of these models provide strong support for the observed differences in styrene toxicity between rats, mice, and humans. As discussed further in Section 3.5.3, some have primarily focused on the species differences in the metabolism of styrene and metabolic differences between rats, mice, and humans.

3. HEALTH EFFECTS

Figure 3-4. Conceptual Representation of a Physiologically Based Pharmacokinetic (PBPK) Model for a Hypothetical Chemical Substance



Note: This is a conceptual representation of a physiologically based pharmacokinetic (PBPK) model for a hypothetical chemical substance. The chemical substance is shown to be absorbed via the skin, by inhalation, or by ingestion, metabolized in the liver, and excreted in the urine or by exhalation.

Source: adapted from Krishnan and Andersen 1994

3. HEALTH EFFECTS

3.4.5.1 Summary of PBPK Models

A number of investigators have developed toxicokinetic models for styrene. The earliest model was developed by Ramsey and Anderson (1984) to relate styrene exposure concentrations quantitatively to blood concentrations. A model developed by Csanády et al. (1994, 2003) is useful in describing the blood/tissue time course of styrene and styrene oxide in rats, mice, and humans following multiple routes of administration. A model developed by Sarangapani et al. (2002) expands these models and adds a description of styrene and styrene oxide levels in multiple compartments of the respiratory tract; this model is described below.

Additionally, Jonsson and Johanson (2002) developed a population-based PBPK model for styrene, which decreased the intraindividual variability for estimating the metabolic capacity for styrene in humans. Leavens and Bond (1996) described initial work on developing a model for co-exposure to 1,3-butadiene and styrene in mice.

3.4.5.2 Discussion of Model


Risk assessment. The Sarangapani model quantifies the observed differences in the occurrence of lung tumors between rats, mice, and humans.

Description of the model. The Sarangapani model is a mode of action based PBPK model developed to predict blood, liver, and respiratory tract tissue (particularly the terminal bronchioles) levels of styrene and styrene oxide and allow for interspecies extrapolations. The model has a nested architecture with a model for styrene and a linked submodel for styrene oxide. Both models consist of four respiratory tract tissue compartments (upper respiratory tract, conducting airways, terminal bronchioles, and pulmonary) and systemic tissue compartments for liver, fat, richly perfused tissue, and poorly perfused tissue. The metabolism of styrene to styrene 7,8-oxide and the detoxification of styrene 7,8-oxide by epoxide hydrolase and glutathione S-transferase was modeled to occur in the liver and selected regions of the lung. The styrene oxide formed in the parent model in any tissue compartment was passed to the corresponding tissue compartment in the metabolite submodel; the schematic of the PBPK model is presented in Figure 3-5. Inhalation was the only exposure route considered in this model.

Mass balance equations were used to account for the transport of styrene across the lumen and tissue subcompartments in the upper, conducting, and transitional airways and in the liver. Additionally mass balance equations were used to describe the production of styrene 7,8-oxide, elimination of styrene oxide,

3. HEALTH EFFECTS

Figure 3-5. Styrene and Styrene Oxide Models used in the Sarangapani PBPK Model



EH = epoxide hydrase; GST = glutathione S-transferase; PBPK = physiologically based pharmacokinetic; T. Bronchioles = transitional bronchioles; URT = upper respiratory tract; V_a = alveolar ventilation; V_m = minute ventilation or pulmonary ventilation;

Source: Sarangapani et al. 2002

3. HEALTH EFFECTS

and kinetics of cytosolic glutathione in the liver and lung. Most of the physiological and flow parameters (Table 3-6) and respiratory tract-specific physiological parameters (Table 3-7) used in the model were obtained from the literature. Tissue volumes for the respiratory tract compartments were estimated by multiplying the appropriate surface areas with the tissue thickness. The kinetic constants used for hepatic and lung cytochrome P450, epoxide hydrolase, and glutathione S-transferase are presented in Table 3-8. Stereospecific kinetic parameters (Table 3-9) and steady-state concentrations of R-styrene oxide and S-styrene oxide were used to account for species differences in the stereospecific metabolism of styrene to styrene oxide.

Validation of the model. Ten independent data sets ranging from closed chamber data to concentration measurements of styrene and styrene oxide in multiple tissues following multiple routes of exposure were used to validate various dose metrics in the mouse, rat, and human. The model provided good fit across species and at multiple target sites, including the whole lung for both styrene and styrene oxide. The model was not validated for styrene and styrene oxide levels in the transitional bronchioles due to the lack of experimental data.

Target tissues. The model was used to predict steady-state styrene oxide levels in arterial blood and the terminal bronchioles.

Species extrapolation. The model predicted that the levels of styrene oxide in the transitional bronchioles are approximately 10- and 100-fold lower in rats and humans, as compared to mice, thus suggesting that humans would be 100-fold less sensitive than mice to styrene-induced lung tumors.

Interroute extrapolation. Interroute extrapolation was not attempted in this model.

3.5 MECHANISMS OF ACTION

3.5.1 Pharmacokinetic Mechanisms

Styrene is rapidly absorbed through the respiratory tract (Ramsey and Andersen 1984; Ramsey and Young 1978; Ramsey et al. 1980; Withey and Collins 1979; Withey and Karpinski 1985) with a mean uptake of approximately 60–70% in humans (Johanson et al. 2000; Norström et al. 1992). A concentration-dependent uptake efficiency was found in the upper respiratory tract of rats and mice (Morris 2000). In rats, the uptake efficiency was 23.7% at 5 ppm and 10.1% at 200 ppm; in mice, uptake efficiency decreased from 41.7% at 5 ppm to 9.6% at 200 ppm. Based on the decreased uptake efficiency

3. HEALTH EFFECTS

Table 3-6. Physiological and Flow Parameters Used in the Sarangapani PBPK Model

Parameter	Mouse	Rat	Human
Body weight (g)	25	250	70,000
Tissue volume as fraction of body weight			
Liver	5.5	3.66	2.6
Richly perfused tissues	10	12.34	8.5
Poorly perfused tissues	70	70	60
Fat	7	6.5	21.4
Blood	7.5	7.5	7.5
Minute ventilation (mL/minute)	24	150	15,000
Pulmonary ventilation (mL/minute)	12–1	75–110	10,500
Cardiac output (mL/minute)	14	110	5,200
Tissue blood flow as fraction of cardiac output			
Liver	15–30	15–30	22.7
Richly perfused tissues	48	28.7	43
Fat	5.9	7	5.2
Upper airways	1.0	1.0	0.25
Conducting airways	0.5	2.1	0.75
Transitional airways	0.1	0.15	0.67
Partition coefficients for styrene			
Blood:air	40	40	48
Liver:blood	2	2	2
Fat:blood	87	87	50
Tissue:blood	1.3	1.3	1.3
Partition coefficients for styrene 7,8-oxide			
Blood:air	2,000	2,000	2,000
Liver:blood	1	1	1
Fat:blood	14	14	14
Tissue:blood	0.6	0.6	0.6
Styrene:styrene oxide tissue-phase diffusivity (cm ² /minute)	0.0002	0.0002	0.0002
Styrene:styrene oxide air-phase diffusivity (cm ² /minute)	6	6	6

Source: Sarangapani et al. 2002

3. HEALTH EFFECTS

Table 3-7. Respiratory Tract-Specific Physiological Parameters Used in the Sarangapani PBPK Model

Parameter	Mouse	Rat	Human
Tissue thickness (cm)			
Upper airway epithelium	0.005	0.005	0.005
Conducting airway epithelium	0.0025	0.0025	0.0025
Transitional airway epithelium	0.001	0.001	0.001
Pulmonary airway epithelium	0.0003	0.00025	0.0005
Mucus	0.0005	0.001	0.001
Upper airway submucosa	0.01	0.01	0.01
Conducting airway submucosa	0.005	0.005	0.005
Transitional airway submucosa	0.002	0.002	0.002
Surface area (cm ²)			
Upper airway compartment	2.7	13.2	138
Conducting airway compartment	8.87	48.3	2,000
Transitional airway compartment	0.48	5.5	,6220
Pulmonary airway compartment	500	3,400	540,000
Mass transfer coefficient (cm/min)			
Upper airway air-phase	7,200	7,200	1,980
Conducting airway air-phase	312	228	181
Transitional airway air-phase	1,136	481	158
Tissue liquid phase	32	16	19.2
Intracompartment clearance (cm ³ /minute)	10	40	400
Lung microsomal protein (mg/mL)	3.8	3.8	3.8
Liver microsomal protein (mg/mL)	13	11	23
Lung cytosolic protein (mg/mL)	68	60	43
Liver cytosolic protein (mg/mL)	94	90	45

Source: Sarangapani et al. 2002

3. HEALTH EFFECTS

Table 3-8. Kinetic Parameters Used in the Sarangapani PBPK Model

Parameter	Mouse	Rat	Human
V_{\max} Cytochrome P450 (nmol/minute/mL)			
Liver	200	50–150	50
Upper airway	183	98	50
Transitional airway	362	46.4	1.7
K_m Cytochrome P450 (nmol/mL)	10	10	10
V_{\max} Epoxide hydrolase (nmol/minute/mL)			
Liver	200	250	900
Upper airway	250	250	500
Transitional airway	250	250	500
K_m Epoxide hydrolase (nmol/mL)	100	100	100
V_{\max} Glutathione S-transferase (nmol/minute/mL)			
Liver	11,000	6,300	1,400
Upper airway	1,000	1,000	300
Transitional airway	1,000	1,000	300
K_m Glutathione S-transferase for styrene (nmol/mL)	2,500	2,500	2,500
K_m Glutathione S-transferase for styrene oxide (nmol/mL)	700	700	500
Liver glutathione basal concentration (nmol/mL)	8,300	6,300	6,000
Upper airway glutathione basal concentration (nmol/mL)	1,000	2,500	1,000
Transitional airway glutathione basal concentration (nmol/mL)	1,000	1,000	1,000
Glutathione production rate (per minute)	0.012	0.012	0.012

Source: Sarangapani et al. 2002

3. HEALTH EFFECTS

Table 3-9. Stereospecific Kinetic Parameters for Styrene and Styrene Oxide Metabolism in Rodents Used in the Sarangapani PBPK Model

Parameter	Mouse		Rat	
	R	S	R	S
V_{max} Cytochrome P450-liver (nmol/minute/mL)	108.3	91.7	33.8	59.2
V_{max} Cytochrome P450-terminal bronchioles (nmol/minute/mL)	211.8	88.2	130	250
K_m Cytochrome P450 (nmol/mL)	10	10	10	10
K_m Epoxide hydrolase, liver (nmol/mL)	66.7	133.3	570	151
K_m Epoxide hydrolase, terminal bronchioles (nmol/mL)	125	125	200	50
K_m Epoxide hydrolase (nmol/mL)	29	155	29	155
V_{max} Glutathione S-transferase-liver (nmol/minute/mL)	4,400	6,600	2,400	3,600
V_{max} Glutathione S-transferase-terminal bronchioles (nmol/minute/mL)	400	600	400	600
K_m Glutathione S-transferase for styrene oxide (nmol/mL)	700	2,000	700	2,000
K_m Glutathione S-transferase for glutathione (nmol/mL)	2,500	2,500	2,500	2,500

Source: Sarangapani et al. 2002

3. HEALTH EFFECTS

observed in rats and mice following exposure to the cytochrome P450 inhibitor, metyrapone, Morris (2000) suggested that styrene was metabolized *in situ* and this metabolism enhanced styrene uptake. In humans, blood styrene levels reached steady state after 75 minutes of exposure to 70 ppm (Wigaeus et al. 1983). The elimination of styrene from blood was biphasic, with a half-time of 1 minute for the rapid distribution phase and 40.8 minutes for the elimination phase. Styrene is rapidly distributed throughout the body with the highest concentrations found in adipose tissue. In rats, the styrene concentration in the adipose tissue was approximately 50-fold higher than in muscle; the biological half-time was 6.3 hours in adipose tissue and 2.4–2.0 hours in the blood, liver, kidney, spleen, muscle, and brain (Teramoto and Horiguchi 1979). In humans, styrene is primarily excreted in the urine as mandelic acid and phenylglyoxylic acid. The half-times of mandelic acid and phenylglyoxylic acid in the urine were 3.6 and 8.8 hours, respectively, in humans exposed to 70 ppm for 2 hours (Wigaeus et al. 1983); another study reported elimination half-times of 2.2–4.2 hours for mandelic acid and 3.5–13.9 hours for phenylglyoxylic acid following a 2-hour exposure to 50 ppm styrene (Johanson et al. 2000).

3.5.2 Mechanisms of Toxicity

A large number of studies have investigated the mechanism of styrene carcinogenic activity, particularly the increased susceptibility of mice. Increases in malignant lung tumors have been observed in mice exposed to 160 ppm 6 hours/day, 5 days/week for approximately 2 years (Cruzan et al. 2001) and following gavage exposure to 300 mg/kg/day administered 5 days/week (NCI 1979b); however, neoplastic tumors have not been observed in rats exposed to concentrations as high as 1,000 ppm 6 hours/day, 5 days/week for 2 years (Cruzan et al. 1998) or 2,000 mg/kg/day 5 days/week for 2 years (NCI 1979b), suggesting that mice are particularly sensitive. As reviewed by IARC (2002), Cohen et al. (2002), and Cruzan et al. (2002), genotoxic and nongenotoxic modes of action have been proposed. Although styrene itself does not appear to be DNA reactive, styrene 7,8-oxide is DNA reactive and has been shown to form stable N₂ and O₆ adducts of deoxyguanosine. Styrene oxide, DNA adducts, and genotoxic effects have been detected in humans, rats, and mice. Styrene (styrene 7,8-oxide is the likely causative agent) has been shown to be mutagenic in bacteria, and exposure can result in increased frequency of sister chromatid exchange, chromosomal aberrations, micronucleated cells, and DNA strand breaks. However, elevated levels of blood styrene oxide do not explain the species differences in tumor formation. In humans, styrene 7,8-oxide is rapidly hydrolyzed by epoxide hydrolase as evidenced by the high levels of mandelic acid, phenylglyoxylic acid, and hippuric acid detected in the urine. Styrene 7,8-oxide is relatively stable in rats and mice, and elevated levels have been detected in blood. The blood

3. HEALTH EFFECTS

levels of styrene oxide in rats exposed to 1,000 ppm is 100-fold higher than the levels in mouse exposed to 20–40 ppm; however, tumors have not been detected in rats.

Although the nongenotoxic potential mode of action also implicates styrene 7,8-oxide as the causative agent, it involves cytotoxic damage at the target tissue, the lung. In the mouse lung, styrene is primarily metabolized by cytochrome P450, particularly the CYP2F2 isoform, in the Clara cell. The continued exposure to styrene 7,8-oxide results in Clara cell cytotoxicity, increased cell proliferation, bronchiolar epithelial hyperplasia, and eventually lung tumors. Several species differences account for the increased sensitivity of mice, compared to rats and likely humans. Humans appear to have a lower capacity to metabolize styrene in the lung compared to rats and a much lower capacity compared to mice and humans and rats have fewer Clara cells than mice. Mouse Clara cells metabolize higher levels of styrene than rat Clara cells and produce a higher proportion of R-styrene oxide than S-styrene oxide, as compared to rats. It has been estimated that mice produce 15 times more R-styrene oxide than rats; in humans, the S enantiomer also predominates. This is particularly important since R-styrene oxide is a more potent pneumotoxicant than S-styrene oxide. In mice and rats, a portion of the styrene oxide generated is metabolized via glutathione conjugation. Mice appear to be more susceptible to glutathione depletion than rats, and glutathione depletion has been observed in mouse lung tissue at exposure concentrations of 80–300 ppm.

IARC (2002) concluded that the proposed mechanism involving the metabolism of styrene to styrene 7,8-oxide in the mouse Clara cell is the likely mode of action resulting in lung tumors in mice. This mode of action is not likely to be relevant to humans to a biologically significant extent. However, this mechanism and a genotoxic mode of action have not been excluded for humans, and styrene is considered a possible human carcinogen.

3.5.3 Animal-to-Human Extrapolations

Species differences exist in the metabolism of styrene in humans, rats, and mice; these differences are discussed in greater detail in Section 3.4.3. Although all three species predominantly metabolize styrene to styrene 7,8-oxide, there are species differences in the subsequent metabolism of styrene 7,8-oxide. As discussed in the metabolism section, styrene 7,8-oxide is primarily hydrolyzed to mandelic acid via epoxide hydrolase in humans. In rats and mice, styrene 7,8-oxide is also conjugated to form mercapturic acids and styrene is metabolized to phenylacetic acid and/or 4-vinylphenol. In rats, 68–72% of the styrene metabolites in urine are generated from the epoxide hydrolase pathway and 26–35% are from the

3. HEALTH EFFECTS

glutathione transferase pathway; in mice, 48–59 and 20–35% arise from the epoxide hydrolase and glutathione transferase pathways, respectively (Cruzan et al. 2002). In contrast, 95–100% of the styrene 7,8-oxide is metabolized via the epoxide hydrolase pathway; only trace amounts of mercapturic acids have been detected in human urine. The difference in metabolism could result in significant increases in styrene 7,8-oxide levels in the body following exposure to high levels of styrene which may result in depletion of glutathione. Additionally, a small percentage of styrene can undergo ring oxidation resulting in the formation of 4-vinylphenol. Ring-opened compounds account for 4–8% of the urinary metabolites in mice, less than 1% in rats, and were not detected in humans. The production of 4-vinylphenol is potentially significant mode of action because it is considered to be more toxic to the liver and lung than styrene or styrene oxide (Cruzan et al. 2005b).

However, these differences in the hepatic metabolism of styrene do not account for all of the observed species differences in styrene toxicity. As discussed in Section 3.2, mice appear to be especially sensitive to styrene toxicity in the liver, nasal olfactory, and lung. In the respiratory tract, the species differences between rats and mice are due to local metabolism of styrene to R-styrene oxide and/or other oxidized metabolites. The higher rate of metabolism in mice and higher production of the more reactive enantiomer likely result in increased susceptibility. The fact that humans have a more limited ability to metabolize styrene in the respiratory tract and possibly a higher potential to detoxify styrene oxide suggests that mice are not a good model for end points in which styrene oxide is the causative agent.

3.6 TOXICITIES MEDIATED THROUGH THE NEUROENDOCRINE AXIS

Recently, attention has focused on the potential hazardous effects of certain chemicals on the endocrine system because of the ability of these chemicals to mimic or block endogenous hormones. Chemicals with this type of activity are most commonly referred to as *endocrine disruptors*. However, appropriate terminology to describe such effects remains controversial. The terminology *endocrine disruptors*, initially used by (Thomas and Colborn 1992), was also used in 1996 when Congress mandated the EPA to develop a screening program for “...certain substances [which] may have an effect produced by a naturally occurring estrogen, or other such endocrine effect[s]...”. To meet this mandate, EPA convened a panel called the Endocrine Disruptors Screening and Testing Advisory Committee (EDSTAC), and in 1998, the EDSTAC completed its deliberations and made recommendations to EPA concerning *endocrine disruptors*. In 1999, the National Academy of Sciences released a report that referred to these same types of chemicals as *hormonally active agents*. The terminology *endocrine modulators* has also been used to convey the fact that effects caused by such chemicals may not necessarily be adverse. Many scientists

3. HEALTH EFFECTS

agree that chemicals with the ability to disrupt or modulate the endocrine system are a potential threat to the health of humans, aquatic animals, and wildlife. However, others think that endocrine-active chemicals do not pose a significant health risk, particularly in view of the fact that hormone mimics exist in the natural environment. Examples of natural hormone mimics are the isoflavonoid phytoestrogens (Adlercreutz 1995; Livingston 1978; Mayr et al. 1992). These chemicals are derived from plants and are similar in structure and action to endogenous estrogen. Although the public health significance and descriptive terminology of substances capable of affecting the endocrine system remains controversial, scientists agree that these chemicals may affect the synthesis, secretion, transport, binding, action, or elimination of natural hormones in the body responsible for maintaining homeostasis, reproduction, development, and/or behavior (EPA 1997). Stated differently, such compounds may cause toxicities that are mediated through the neuroendocrine axis. As a result, these chemicals may play a role in altering, for example, metabolic, sexual, immune, and neurobehavioral function. Such chemicals are also thought to be involved in inducing breast, testicular, and prostate cancers, as well as endometriosis (Berger 1994; Giwercman et al. 1993; Hoel et al. 1992).

No *in vitro* studies were located regarding endocrine disruption of styrene.

There is some evidence in styrene workers and in female rats that inhalation exposure to styrene may disrupt the tuberoinfundibular dopaminergic system. Significant alterations in serum prolactin levels have been observed in male and female workers exposed to air concentrations as low as 50 ppm (Bergamaschi et al. 1996, 1997; Luderer et al. 2004; Mutti et al. 1984b); a regression model predicts that exposure to ≥ 20 ppm would result in significant elevations in serum prolactin levels (Luderer et al. 2004). Elevated serum prolactin levels have also been observed in female rats acutely exposed to 150 ppm (Umemura et al. 2005); alterations have not been observed in male rats exposed to concentrations as high as 1,500 ppm (Jarry et al. 2002; Umemura et al. 2005). As noted by NTP (2006), the clinical significance of the increased serum prolactin levels, in the absence of other reproductive effects, is not known. Styrene exposure does not appear to adversely affect thyroid stimulating hormone levels in humans (Arfini et al. 1987) or rats (Umemura et al. 2005) or follicle stimulating hormone or luteinizing hormone in humans (Arfini et al. 1987).

No significant alterations were observed in a gonadal sex differentiation assay using genetic male frogs exposed to styrene (Ohtani et al. 2001).

3. HEALTH EFFECTS

3.7 CHILDREN'S SUSCEPTIBILITY

This section discusses potential health effects from exposures during the period from conception to maturity at 18 years of age in humans, when all biological systems will have fully developed. Potential effects on offspring resulting from exposures of parental germ cells are considered, as well as any indirect effects on the fetus and neonate resulting from maternal exposure during gestation and lactation.

Relevant animal and *in vitro* models are also discussed.

Children are not small adults. They differ from adults in their exposures and may differ in their susceptibility to hazardous chemicals. Children's unique physiology and behavior can influence the extent of their exposure. Exposures of children are discussed in Section 6.6, Exposures of Children.

Children sometimes differ from adults in their susceptibility to hazardous chemicals, but whether there is a difference depends on the chemical (Guzelian et al. 1992; NRC 1993). Children may be more or less susceptible than adults to health effects, and the relationship may change with developmental age (Guzelian et al. 1992; NRC 1993). Vulnerability often depends on developmental stage. There are critical periods of structural and functional development during both prenatal and postnatal life, and a particular structure or function will be most sensitive to disruption during its critical period(s). Damage may not be evident until a later stage of development. There are often differences in pharmacokinetics and metabolism between children and adults. For example, absorption may be different in neonates because of the immaturity of their gastrointestinal tract and their larger skin surface area in proportion to body weight (Morselli et al. 1980; NRC 1993); the gastrointestinal absorption of lead is greatest in infants and young children (Ziegler et al. 1978). Distribution of xenobiotics may be different; for example, infants have a larger proportion of their bodies as extracellular water, and their brains and livers are proportionately larger (Altman and Dittmer 1974; Fomon 1966; Fomon et al. 1982; Owen and Brozek 1966; Widdowson and Dickerson 1964). The infant also has an immature blood-brain barrier (Adinolfi 1985; Johanson 1980) and probably an immature blood-testis barrier (Setchell and Waites 1975). Many xenobiotic metabolizing enzymes have distinctive developmental patterns. At various stages of growth and development, levels of particular enzymes may be higher or lower than those of adults, and sometimes unique enzymes may exist at particular developmental stages (Komori et al. 1990; Leeder and Kearns 1997; NRC 1993; Vieira et al. 1996). Whether differences in xenobiotic metabolism make the child more or less susceptible also depends on whether the relevant enzymes are involved in activation of the parent compound to its toxic form or in detoxification. There may also be differences in excretion, particularly in newborns who all have a low glomerular filtration rate and have not developed efficient

3. HEALTH EFFECTS

tubular secretion and resorption capacities (Altman and Dittmer 1974; NRC 1993; West et al. 1948). Children and adults may differ in their capacity to repair damage from chemical insults. Children also have a longer remaining lifetime in which to express damage from chemicals; this potential is particularly relevant to cancer.

Certain characteristics of the developing human may increase exposure or susceptibility, whereas others may decrease susceptibility to the same chemical. For example, although infants breathe more air per kilogram of body weight than adults breathe, this difference might be somewhat counterbalanced by their alveoli being less developed, which results in a disproportionately smaller surface area for alveolar absorption (NRC 1993).

No studies were identified that examined the toxicity of styrene in children. Several occupational exposure studies have examined the developmental toxicity of styrene; these studies did not find statistically significant alterations in the occurrence of stillbirths, infant deaths, malformations, or birth weight (Ahlborg et al. 1987; Härkönen et al. 1984; Lemasters et al. 1989). However, an expert panel (NTP 2006) evaluating these data concluded that the human studies were not sufficient to evaluate developmental toxicity due to the low statistical power of the studies and the lack of adequate information on exposure.

In general, animal studies have not found styrene-related developmental effects following inhalation exposure in rats (Murray et al. 1978), mice (Kankaanpää et al. 1980), rabbits (Murray et al. 1978), or hamsters (Kankaanpää et al. 1980) or oral exposure in rats (Daston et al. 1991; Murray et al. 1978); additionally, no developmental effects were observed in a rat two-generation study (Cruzan et al. 2005b). An expert panel determined that there was sufficient animal data to conclude that styrene does not cause developmental toxicity in rats following inhalation or oral exposure or in rabbits following inhalation exposure. No studies examined styrene toxicity following exposure of young laboratory animals.

Studies in adults, particularly reinforced plastics industry workers, have identified the nervous system as the most sensitive target of styrene toxicity. Inconsistent results have been found in animal neurodevelopmental toxicity studies. Minor alterations in forelimb grip strength and swimming ability were observed in F2 offspring of rats exposed to 500 ppm styrene; however, the investigators (Cruzan et al. 2005a) attributed these alterations to a lower body weight rather than a neurodevelopmental effect of styrene. Another inhalation study found impaired righting reflex in the offspring of rats exposed to 300 ppm during gestation (Katakura et al. 2001). Similarly, impaired amphetamine-induced locomotor

3. HEALTH EFFECTS

activity and apomorphine-induced stereotypy were observed in the offspring of rats orally administered 200 mg/kg/day styrene during gestation and lactation (Zaidi et al. 1985).

No human or animal data were located on the toxicokinetic properties of styrene in children or immature animals or possible age-related differences in the toxicokinetics of styrene. A lactational toxicokinetic model predicted that styrene can be transferred via maternal milk (Fisher et al. 1997). Subsequent sections of this chapter (Sections 3.8, 3.10, and 3.11) discuss the available information on biomarkers, interactions, and methods for reducing toxic effects. The available information is from adults and mature animals; no child-specific information was identified. It is likely that this information will also be applicable to children.

3.8 BIOMARKERS OF EXPOSURE AND EFFECT

Biomarkers are broadly defined as indicators signaling events in biologic systems or samples. They have been classified as markers of exposure, markers of effect, and markers of susceptibility (NAS/NRC 1989).

Due to a nascent understanding of the use and interpretation of biomarkers, implementation of biomarkers as tools of exposure in the general population is very limited. A biomarker of exposure is a xenobiotic substance or its metabolite(s) or the product of an interaction between a xenobiotic agent and some target molecule(s) or cell(s) that is measured within a compartment of an organism (NAS/NRC 1989). The preferred biomarkers of exposure are generally the substance itself, substance-specific metabolites in readily obtainable body fluid(s), or excreta. However, several factors can confound the use and interpretation of biomarkers of exposure. The body burden of a substance may be the result of exposures from more than one source. The substance being measured may be a metabolite of another xenobiotic substance (e.g., high urinary levels of phenol can result from exposure to several different aromatic compounds). Depending on the properties of the substance (e.g., biologic half-life) and environmental conditions (e.g., duration and route of exposure), the substance and all of its metabolites may have left the body by the time samples can be taken. Identification of individuals exposed to hazardous substances that are commonly found in body tissues and fluids (e.g., essential mineral nutrients such as copper, zinc, and selenium) may be difficult. Biomarkers of exposure to styrene are discussed in Section 3.8.1.

Biomarkers of effect are defined as any measurable biochemical, physiologic, or other alteration within an organism that, depending on magnitude, can be recognized as an established or potential health

3. HEALTH EFFECTS

impairment or disease (NAS/NRC 1989). This definition encompasses biochemical or cellular signals of tissue dysfunction e.g., increased liver enzyme activity or pathologic changes in female genital epithelial cells), as well as physiologic signs of dysfunction such as increased blood pressure or decreased lung capacity. Note that these markers are not often substance specific. They also may not be directly adverse, but can indicate potential health impairment (e.g., DNA adducts). Biomarkers of effects caused by styrene are discussed in Section 3.8.2.

A biomarker of susceptibility is an indicator of an inherent or acquired limitation of an organism's ability to respond to the challenge of exposure to a specific xenobiotic substance. It can be an intrinsic genetic or other characteristic or a preexisting disease that results in an increase in absorbed dose, a decrease in the biologically effective dose, or a target tissue response. If biomarkers of susceptibility exist, they are discussed in Section 3.10, Populations That Are Unusually Susceptible.

3.8.1 Biomarkers Used to Identify or Quantify Exposure to Styrene

The elimination of styrene via expired air may be used to identify exposure to styrene (Guillemin and Berode 1988; Stewart et al. 1968). Only a small percentage of unchanged styrene is expired after cessation of exposure. There are no adequate studies correlating post-exposure exhaled styrene with previous exposure levels. Assessment of occupational exposure involving measurement of unchanged styrene in urine has been reported (Dolara et al. 1984). In this study of workers, the styrene air concentrations were 3.8–14 ppm and the urinary concentrations of styrene were 0.7–4.1 µg/L. Urinary mutagenic activity was also evaluated in this study and was not a good indication of exposure to styrene. Only a small fraction of unchanged styrene is recovered in the urine. However, measurement of styrene in urine is a reliable indicator of styrene exposure if the exposure is recent (Dolara et al. 1984; Gobba et al. 1993; Guillemin and Berode 1988; Pezzagno et al. 1985).

Analysis of unchanged styrene in blood may be used as a qualitative indicator of styrene exposure (Antoine et al. 1986). In one study, styrene was detected in the blood of humans exposed to 80 ppm (Ramsey et al. 1980). The maximum blood concentration at the end of exposure was 0.92 ± 0.26 µg/mL. The half-life values for rapid and slow clearance curves were 0.58 and 13 hours, respectively. In another study, the concentration of styrene in blood (0.2–3.7 mg/L) increased with the level and duration of styrene exposure (Baselt 1988a).

3. HEALTH EFFECTS

The presence of styrene in adipose tissue is also an indicator of exposure. The concentration of styrene in the adipose tissue of two workers exposed to 7.5–20 ppm of styrene during a work week suggested a half-life of 5.2 days for one worker and 2.8 days for the other worker. The elimination time was estimated to be 5 weeks (Engstrom et al. 1978b).

Levels of occupational exposure to styrene may also be estimated by measurement of styrene metabolites such as MA and PGA in urine (Bartolucci et al. 1986; Elia et al. 1980; Engstrom et al. 1976; Sedivec et al. 1984; Sollenberg et al. 1988). However, large intra-individual differences in MA and PGA urinary concentrations have been reported. A study of the inter- and intra-individual differences found that PGA levels were less variable than MA levels (Symanski et al. 2001) and variability was higher in post-shift urine samples compared to pre-shift urine samples. Expressing MA and PGA levels in units of mg per gram creatinine decreased the source intra-individual variability. Some studies found a good correlation between the time-weighted styrene exposure and urinary MA concentrations (Chua et al. 1993; Engstrom et al. 1976; Härkönen et al. 1974), while other studies found a better correlation with the sum of urinary MA and PGA at the end of the work period (Elia et al. 1980; Ong et al. 1994; Sollenberg et al. 1988). A good correlation between environmental styrene levels and urinary PGA levels has also been found (Chua et al. 1993). Total MA and PGA measured the morning after exposure may be a more reliable biological indicator of styrene exposure in factories where there is high variability in the environmental styrene concentration (Bartolucci et al. 1986).

Reference levels of styrene urinary metabolites likely to be observed in workers exposed to the time-weighted average concentrations by inhalation have been reported. The American Conference of Governmental Industrial Hygienists (ACGIH 2006) recommends a biological exposure index of 400 mg/g creatinine for the sum of MA and PGA in urine.

3.8.2 Biomarkers Used to Characterize Effects Caused by Styrene

The nervous system is the most sensitive target of styrene toxicity in humans. Styrene affects both sensory (color vision, hearing, vestibular) and motor (nerve conduction velocity) function. Impaired performance on neurobehavioral tests and diminished color vision may be indicative of styrene exposure; however, these effects are not specific to styrene and have been observed following exposure to other solvents such as toluene. Several studies have found significant associations between performance on tests of color discrimination (Eguchi et al. 1995; Kishi et al. 2001) or reaction time (Mutti et al. 1984a) and levels of styrene urinary metabolites (mandelic acid and/or phenylglyoxylic acid).

3. HEALTH EFFECTS

Other investigators have proposed the use of genotoxicity biomarkers. Cytogenetic monitoring of peripheral lymphocytes as a biomarker of effect has been proposed (DeJong et al. 1988; Pero et al. 1982). Future biomarkers may include hemoglobin adducts. Using unscheduled DNA synthesis (UDS) as an indicator of DNA damage, the lymphocytes of 38 individuals occupationally exposed to styrene were evaluated. The induced UDS was significantly increased for the group exposed to 1–40 ppm styrene (Pero et al. 1982). Measurement of chromosome aberration in peripheral blood lymphocyte has been used for many years to monitor the biologic effects of genotoxic chemicals. However, due to high background levels of chromosomal aberration and exposures to other genotoxic workplace chemicals, the sensitivity of this biomarker for the effects of styrene is probably not adequate (DeJong et al. 1988). The role of hepatic glutathione in the toxicity of styrene has been proposed as inhibiting the covalent binding of styrene. This has been confirmed in animal studies by decreased glutathione in styrene-exposed animals (Parkki 1978). However, its use as a biomarker of effect in humans remains to be demonstrated since data on the adverse effects of styrene on the human liver are insufficient.

3.9 INTERACTIONS WITH OTHER CHEMICALS

Styrene metabolism is known to be inhibited by the presence of other chemicals such as toluene, trichloromethylene, and ethyl benzene. The biotransformation of styrene in rats to PGA, MA, and hippuric acid was suppressed by co-administration of toluene (Ikeda et al. 1972). This may be due to competitive inhibition of oxidative mechanisms. Similar results were reported by Ikeda and Hirayama (1978) in rats when styrene metabolism was inhibited by the administration of trichloroethylene. Urinary metabolites of styrene may be markedly reduced when humans or animals are concurrently exposed to organic solvents that inhibit styrene metabolism.

In numerous polymer industries, workers are exposed to styrene and 1,3-butadiene, and several animal studies have found that styrene affects the metabolism and toxicity of 1,3-butadiene (Laib et al. 1992; Leavens and Bond 1996; Leavens et al. 1997) and 1,3-butadiene affects the metabolism of styrene (Leavens et al. 1996); however, the affect of 1,3-butadiene on styrene toxicity has not been well examined. In other industries, workers are co-exposed to styrene and acrylonitrile; in rats receiving an intraperitoneal dose of styrene and gavage dose of acrylonitrile, increases in serum creatinine and asparate aminotransferase levels were observed, as compared to styrene-only exposure (Normandeau et al. 1984).

3. HEALTH EFFECTS

3.10 POPULATIONS THAT ARE UNUSUALLY SUSCEPTIBLE

A susceptible population will exhibit a different or enhanced response to styrene than will most persons exposed to the same level of styrene in the environment. Reasons may include genetic makeup, age, health and nutritional status, and exposure to other toxic substances (e.g., cigarette smoke). These parameters result in reduced detoxification or excretion of styrene, or compromised function of organs affected by styrene. Populations who are at greater risk due to their unusually high exposure to styrene are discussed in Section 6.7, Populations with Potentially High Exposures.

Styrene is a hazardous substance found in the workplace with much lower levels found in the environment. Therefore, the populations at risk are workers in industries making polystyrene plastics, coating, polyester resins, and other products. Although no populations of unusually susceptible individuals have been identified for styrene, based on the targets of styrene toxicity, an assumption can be made that persons with pre-existing respiratory or neurological problems would be at risk for the irritant action and central nervous system effects of styrene, respectively.

3.11 METHODS FOR REDUCING TOXIC EFFECTS

This section will describe clinical practice and research concerning methods for reducing toxic effects of exposure to styrene. However, because some of the treatments discussed may be experimental and unproven, this section should not be used as a guide for treatment of exposures to styrene. When specific exposures have occurred, poison control centers and medical toxicologists should be consulted for medical advice. The following texts provide specific information about treatment following exposures to styrene:

Bronstein AC, Currence PL. 1988. Emergency care for hazardous materials exposure. St. Louis, MO: The C.V. Mosby Company, 221-222.

Ellenhorn MJ, Barceloux DG. 1988. Medical toxicology: Diagnosis and treatment of human poisoning. New York, NY: Elsevier, 956-959.

Haddad LM, Winchester JF. 1990. Clinical management of poisoning and drug overdose. 2nd ed. Philadelphia, PA: W.B. Saunders Company, 1226-1228.

3.11.1 Reducing Peak Absorption Following Exposure

Human exposure to styrene may occur by inhalation, ingestion, or dermal contact. General recommendations for reducing absorption of styrene following exposure include removing the exposed

3. HEALTH EFFECTS

individual from the contaminated area and removing the contaminated clothing. If the eyes and skin were exposed, they should be flushed with water. Since aspiration of styrene into the lung can cause pulmonary edema and hemorrhage, some authors advise against the use of emetics, but recommend administration of water for dilution of gastric lavage (Bronstein and Currance 1988; Haddad and Winchester 1990). Following acute inhalation exposure, administration of oxygen and use of mechanical ventilation to support respiration have been suggested (Bronstein and Currance 1988; Ellenhorn and Barceloux 1988; Haddad and Winchester 1990). Administration of aminophylline and inhaled bronchodilators may be required to treat bronchospasm (Ellenhorn and Barceloux 1988). Furthermore, cardiac monitoring has been suggested. Supportive treatment may be needed for neurological effects of styrene exposure (Haddad and Winchester 1990).

3.11.2 Reducing Body Burden

Styrene is metabolized by the body, and most styrene that is absorbed is excreted in the urine as metabolites of the parent compound. Styrene is cleared rapidly from the human body. Its half-life is several hours in the blood and about 2–4 days in subcutaneous adipose tissue (see Section 3.4). No method is commonly used to enhance the elimination of the absorbed dose of styrene.

3.11.3 Interfering with the Mechanism of Action for Toxic Effects

In humans, central nervous system depression and upper respiratory tract irritation were reported following acute exposure to higher styrene concentrations (see Section 3.2). Studies in animals indicate that chronic styrene exposure causes liver and kidney effects and may induce cancer. Styrene oxide was found to be the active mutagenic metabolite of styrene in several studies (de Raat 1978; Donner et al. 1979; Norppa et al. 1979, 1980a, 1980b, 1981, 1984, 1988; Pohlova et al. 1985; Vainio et al. 1976). Based on these studies, it can be concluded that styrene is a typical indirect mutagen that needs metabolic activation to be able to bind covalently to macromolecules (e.g., nucleic acids). In one of the possible metabolic pathways, styrene oxide is further metabolized to hydroxyphenylethyl mercapturic acid. The reaction utilizes glutathione (Bond 1989). The mutagenic activity of styrene oxide was decreased in the presence of glutathione in *S. typhimurium* TA100 (Yoshikawa et al. 1980). This experiment, therefore, suggests that glutathione may reduce the mutagenic effects of styrene oxide.

The formation of styrene oxide may also contribute to other effects following styrene exposure. Glutathione decreases the cytotoxicity of many reactive chemicals by acting as a scavenger of toxic metabolites. Exposure of rodents to high levels of styrene caused depletion of glutathione content in the

3. HEALTH EFFECTS

liver cells of these animals (Das et al. 1983; Vainio et al. 1979). Glutathione has been suggested to decrease the hepatotoxicity by preventing styrene oxide reaction with other endogenous macromolecules. Similarly, depletion of glutathione was found in all regions of rat brain following exposure to styrene oxide (Dixit et al. 1982; Trenga et al. 1991). The authors speculated that the depletion of brain glutathione may lead to an increased concentration of free styrene oxide with increased binding to cellular nucleophiles. This process would contribute to oxidative injury to neuronal and glial cells and may be a part of styrene-induced neurotoxicity. However, that styrene itself, being a lipophilic compound, may disrupt the nerve membrane function in a manner similar to anesthetic agents.

Although results from *in vitro* studies in bacteria and *in vivo* animal studies demonstrate that exogenous glutathione precursors may decrease the effects of styrene toxicity, the benefit of this treatment is not known in humans. For low-level exposure cases, the endogenous glutathione levels are not likely to be decreased to a significant extent. Therefore, exogenous glutathione precursors such as N-acetylcysteine are not likely to be effective in mitigating the toxic effects of styrene. Exogenous doses of reducing agents may be useful following acute high dose exposure to styrene. In this case, a significant depletion of glutathione may occur as a result of the presence of high levels of styrene oxide. However, there are no clinical data available to date that support the use of this treatment.

3.12 ADEQUACY OF THE DATABASE

Section 104(I)(5) of CERCLA, as amended, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of styrene is available. Where adequate information is not available, ATSDR, in conjunction with the National Toxicology Program (NTP), is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of styrene.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that if met would reduce the uncertainties of human health assessment. This definition should not be interpreted to mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

3. HEALTH EFFECTS

3.12.1 Existing Information on Health Effects of Styrene

The existing data on health effects of inhalation, oral, and dermal exposure of humans and animals to styrene are summarized in Figure 3-6. The purpose of this figure is to illustrate the existing information concerning the health effects of styrene. Each dot in the figure indicates that one or more studies provide information associated with that particular effect. The dot does not necessarily imply anything about the quality of the study or studies, nor should missing information in this figure be interpreted as a “data need”. A data need, as defined in ATSDR’s *Decision Guide for Identifying Substance-Specific Data Needs Related to Toxicological Profiles* (Agency for Toxic Substances and Disease Registry 1989), is substance-specific information necessary to conduct comprehensive public health assessments. Generally, ATSDR defines a data gap more broadly as any substance-specific information missing from the scientific literature.

There is information on most categories of human toxicity via the inhalation route from occupational studies. However, there are limited data on humans exposed to styrene by the oral or dermal routes. Data from animal studies are more extensive, with studies available for most areas of toxicity resulting from exposure via the oral and inhalation routes. Little is known about the effects of dermal exposure to styrene in animals.

3.12.2 Identification of Data Needs

Acute-Duration Exposure. The possibility for brief human exposure to high concentrations of styrene exists in occupational settings, and might also exist near major spills. Exposure of the general public to episodic high concentrations of styrene at hazardous waste sites, in the home, or in the general environment is unlikely. The respiratory tract and central nervous system are the likely target organ systems for inhaled styrene (Alarie 1973; Carpenter et al. 1944; DeCeuriz et al. 1983; Kankaanpää et al. 1980; Murray et al. 1978; Seeber et al. 2004; Spencer et al. 1942; Stewart et al. 1968). Animal studies have reported hepatic (Cruzan et al. 1997, 2001; Morgan et al. 1993a, 1993b, 1993c; Vainio et al. 1979) and nasal (Cruzan et al. 2001) effects and hearing impairments (Campo et al. 2001; Crofton et al. 1994; Lataye et al. 2003). Available toxicokinetic data suggest that the mouse may be more sensitive to the hepatic and nasal toxicity of styrene than humans; thus, these data are not suitable for derivation of an acute-duration inhalation MRL. Studies have also examined potential reproductive (Salomaa et al. 1985) and developmental (Kankaanpää et al. 1980; Murray et al. 1978) effects; the highest doses tested in these studies were NOAELs. An acute-duration inhalation MRL based on a NOAEL for neurological effects in

3. HEALTH EFFECTS

Figure 3-6. Existing Information on Health Effects of Styrene

		Systemic									
		Death	Acute	Intermediate	Chronic	Immunologic/Lymphoretic	Neurologic	Reproductive	Developmental	Genotoxic	Cancer
Inhalation		●	●	●	●	●	●	●	●	●	●
Oral											
Dermal											

Human

		Systemic									
		Death	Acute	Intermediate	Chronic	Immunologic/Lymphoretic	Neurologic	Reproductive	Developmental	Genotoxic	Cancer
Inhalation		●	●	●	●		●	●	●	●	●
Oral		●	●	●	●	●	●	●	●	●	●
Dermal			●								

Animal

● Existing Studies

3. HEALTH EFFECTS

humans (Seeber et al. 2004) was derived. Episodic high-level exposures to styrene from contaminated food or water are unlikely. Few studies have examined the toxicity of styrene following exposure to an acute oral dose. Abdominal discomfort was observed in residents exposed to elevated levels of styrene in drinking water (Arnedo-Pena et al. 2003); concomitant inhalation exposure to styrene limits the utilization of this study for MRL derivation. A study in rats identified a LOAEL for neurotoxicity (Husain et al. 1985) and another rat study examined potential developmental effects, but found no adverse effects (Daston et al. 1991). An acute-duration oral MRL was derived using the neurotoxicity study conducted by Husain et al. (1985). Although data on the toxicity of styrene following acute inhalation or oral exposure were considered adequate for the derivation of MRLs, the databases are limited to a few studies; additional studies confirming the dose-response relationships would increase the confidence in these MRLs. Dermal exposure to styrene at significant levels is unlikely except in the case of workplace spills and dermal absorption is probably low based on limited human studies. However, the almost complete lack of dermal toxicity data in animals and humans creates a degree of uncertainty on this issue. Therefore, single-dose dermal studies would be useful in determining target organs and thresholds for dermal exposure. In designing these types of studies, precautions should be taken to avoid concomitant inhalation exposure.

Intermediate-Duration Exposure. Information on the toxicity of styrene in humans following intermediate-duration inhalation exposure is limited to a study examining potential reproductive effects in workers (Lindbohm et al. 1985); however, exposure information was not provided. Inhalation studies in animals have reported damage to the nasal olfactory epithelium in rats (Cruzan et al. 1997, 2005a, 2005b; Ohashi et al. 1986) and mice (Cruzan et al. 1997, 2001), liver damage in mice (Cruzan et al. 1997), eye irritation in rats (Cruzan et al. 1997) and guinea pigs (Spencer et al. 1942), ototoxicity in rats (Campo et al. 2001; Lataye et al. 2000; Loquet et al. 1999, 2000; Makitie et al. 2002; Pouyatos et al. 2002; Pryor et al. 1987; Yano et al. 1992), and impaired nerve conduction velocity (Yamamoto et al. 1997). A two-generation study in rats did not find reproductive, developmental, or neurodevelopmental effects (Cruzan et al. 2005a, 2005b), but another study did find neurodevelopmental effects (Katakura et al. 1999, 2001). However, additional studies are needed, as the data are not considered sufficient to derive an intermediate-duration inhalation MRL. Chronic exposure studies provide strong evidence that the nervous system is the most sensitive target of styrene toxicity; studies examining neurological function of workers exposed to styrene for <1 year would provide valuable data for deriving an intermediate-duration inhalation MRL. Oral exposure studies of intermediate-duration are limited to a small number of animal studies and no human data; observed effects include impaired learning in rats (Bushnell 1994) and decreases in spermatozoa in rats (Srivastava et al. 1989, 1992a, 1992b). The results of the Srivastava

3. HEALTH EFFECTS

studies have been questioned by the NTP Expert Panel (NTP 2006) because the findings are inconsistent with the lack of reproductive effects found in an inhalation two-generation study conducted by Cruzan et al. (2005b). The LOAELs identified in these studies were higher than the lowest LOAEL identified in an acute-duration neurotoxicity study (Husain et al. 1985). The intermediate-duration database was considered inadequate for derivation of an MRL because an intermediate-duration oral MRL based on the LOAELs identified in the intermediate-duration oral studies would be higher than the acute-duration oral MRL. Studies in animals are needed to establish dose-response relationships for neurotoxicity, the presumed sensitive end point. Additionally, more studies examining the potential reproductive toxicity of ingested styrene are needed to confirm the results of the Srivastava studies. One study examined the dermal toxicity of styrene in rabbits (Spencer et al. 1942); basic information on the adverse effects of intermediate-duration dermal exposure to styrene in animals is also needed due to the sparsity of available data.

Chronic-Duration Exposure and Cancer. A large number of occupational exposure studies have examined the chronic toxicity of styrene. Systemic toxicity studies have examined endocrine (Bergamaschi et al. 1997; Mutti et al. 1984b), hematological (Checkoway and Williams 1982; Thiess and Friedheim 1978), hepatic (Hotz et al. 1980; Lorimer et al. 1978), or renal (Verplanke and Herber 1998; Viau et al. 1987; Vyskocil et al. 1989) end points; most studies relied on biomarkers of toxicity. The most widely examined end point is neurotoxicity and the available data suggest that this is the most sensitive end point. Examined neurological end points included color vision (Campagna et al. 1995, 1996; Chia et al. 1994; Eguchi et al. 1995; Fallas et al. 1992; Gobba et al. 1991; Gong et al. 2002; Kishi et al. 2001; Mutti et al. 1984a), vestibular effects (Calabrese et al. 1996; Möller et al. 1990), hearing impairment (Morata et al. 2002; Morioka et al. 1999; Muijser et al. 1988; Śliwińska-Kowalska et al. 2003), symptoms of neurotoxicity (Checkoway et al. 1992; Cherry et al. 1980; Edling et al. 1993; Viaene et al. 1998, 2001), performance on neurobehavioral tests (Cherry et al. 1980; Edling et al. 1993; Gamberale et al. 1976; Jegaden et al. 1993; Lindstrom et al. 1976; Mutti et al. 1984a; Tsai and Chen 1996; Viaene et al. 1998, 2001), nerve conduction velocity (Behari et al. 1986; Murata et al. 1991; Rosen et al. 1978), olfactory alterations (Dalton et al. 2003), and EEG alterations (Härkönen et al. 1984; Seppäläinen and Härkönen 1976). Other human studies have examined reproductive (Härkönen and Holmberg 1982; Hemminki et al. 1980) and developmental (Ahlborg et al. 1987; Lemasters et al. 1989) end points. The chronic toxicity of styrene has also been examined in rat (Cruzan et al. 1998; Jersey et al. 1978) and mouse (Cruzan et al. 2001) studies. The occupational exposure studies were considered adequate for derivation of a chronic-duration inhalation MRL for styrene. Further research to define the dose-response curve more fully and to identify a chronic inhalation NOAEL for neurological effects

3. HEALTH EFFECTS

would be valuable and would help to reduce uncertainty in the MRL. Data on chronic oral exposure to styrene is only available through animal studies (Beliles et al. 1985; Conti et al. 1988; NCI 1979b; Quast et al. 1979). In these studies, the most sensitive indicator of toxicity appears to be Heinz body formation in red blood cells in dogs (Quast et al. 1979) and the EPA has calculated a chronic oral RfD based on this study (IRIS 2007). However, there is some doubt regarding the chronic oral NOAEL, and whether hematological effects are really more sensitive than neurological effects. Moreover, decreased survival has been noted in rats at exposure levels only slightly higher than the no-effect level for hematological effects (Conti et al. 1988). Therefore, no chronic oral MRL has been derived. Further studies on the effects of oral exposure, with special emphasis on neurological or neurobehavioral effects, would be valuable. Although chronic dermal exposure by the general public is not likely, there may be some potential for dermal contact with soil at hazardous waste sites. Therefore, data on long-term effects of dermal contact with styrene would be useful.

Taken together, the animal and human data indicate that styrene may possibly be a weak human carcinogen. Although data from epidemiological studies are limited due to concurrent chemical exposures and small cohorts, the data are suggestive of some carcinogenic potential in humans (Antilla et al. 1998; Bond et al. 1992; Cheng et al. 2007; Coggon et al. 1987; Delzell et al. 1996, 2001; Frentzel-Beyme et al. 1978; Gerin et al. 1998; Graff et al. 2005; Hodgson and Jones 1985; Kogevinas et al. 1993, 1994; Kolstad et al. 1993, 1994, 1995; Macaluso et al. 1996; Matanoski and Schwartz 1987; Matanoski et al. 1990; McMichael et al. 1976; Meinhardt et al. 1982; Nicholson et al. 1978; Okun et al. 1985; Ott et al. 1980; Sathiakumar et al. 2005; Wong 1990; Wong et al. 1994). Inhalation and oral exposure studies in rats have not found significant increases in the incidence of neoplastic tumors (Beliles et al. 1985; Conti et al. 1988; Cruzan et al. 1998; Jersey et al. 1978; Maltoni et al. 1979; NCI 1979b). However, inhalation and oral studies in mice have found significant increases in the incidence of neoplastic lung tumors (Cruzan et al. 2001; NCI 1979b). The available data suggest that toxicokinetic differences between rats, mice, and humans result in an increased sensitivity of mice. Clarification of the data is needed in several areas. Almost all of the available epidemiological studies involve concurrent exposures to other chemicals. The role of the metabolism of styrene in humans and animals needs to be clarified and the carcinogenic mechanisms needed to be further elucidated. Additional studies that account for these issues would be valuable.

Genotoxicity. The results of genotoxicity tests for styrene both *in vivo* and *in vitro* are frequently conflicting, and the genotoxic potential of styrene is not clear (Andersson et al. 1980; Beliles et al. 1985; Hogstedt et al. 1979; Meretoja et al. 1977, 1978; Watanabe et al. 1981). The reasons for the mixed or

3. HEALTH EFFECTS

conflicting genotoxicity results may be differences in the metabolism or detoxification of styrene in the various test systems employed. The role of the metabolite styrene oxide in genotoxicity assays on styrene should be fully evaluated, preferably in mammalian *in vivo* systems. Toxicokinetic studies evaluating the presence, level, and activity of styrene oxide in humans will influence the interpretation of genotoxicity studies on styrene and their relevance to public health.

Reproductive Toxicity. Occupational exposure studies have examined male and female styrene workers to evaluate potential reproductive effects; however, most of these studies did not quantify styrene exposure or exposure to other compounds, thus, interpretation of results is difficult. Inconsistent results have been reported for spontaneous abortions with some studies reporting significant increases (Härkönen and Holmberg 1982; Hemminki et al. 1980; McDonald et al. 1988) and others reporting no effect (Härkönen and Holmberg 1982; Hemminki et al. 1980, 1984; Lindbohm et al. 1985). Oligomenorrhea was observed in one study of workers (Cho et al. 2001), but not in another study (Lemasters et al. 1985). Studies in male workers have found alterations in sperm parameters (Kolstad et al. 1999a), but no alterations in time-to-pregnancy (Kolstad et al. 2000; Sallmén et al. 1998) or fertility rates (Kolstad et al. 1999c). A two-generation inhalation study (Cruzan et al. 2005b) and three-generation oral study (Beliles et al. 1985) in rats showed no styrene-related reproductive effects. However, testicular effects have been observed in an oral exposure study (Srivastava et al. 1989), but not in two inhalation studies (Cruzan et al. 2005b; Salomaa et al. 1985). Additional reproductive data on occupationally-exposed males would be useful in evaluating the existing animal data that indicates altered testicular function and studies in females would be useful in evaluating the inconsistent findings in the existing studies.

Developmental Toxicity. Data on the developmental effects of inhalation exposure to styrene are available in humans and animals. Occupational exposure studies (Ahlborg et al. 1987; Härkönen et al. 1984; Lemasters et al. 1989) have not found increases in the occurrence of birth defects or decreases in birth weight. However, interpretation of the results are complicated by exposure to other chemicals and lack of information on exposure levels. Additional occupational studies are needed to adequately assess this end point. Developmental studies in animals via inhalation (Cruzan et al. 2005b; Kankaanpää et al. 1980; Murray et al. 1978) or oral (Beliles et al. 1985) exposure have not found effects on fetal outcome, birth weight, or incidence of abnormalities. However, several studies have reported neurodevelopmental (Katakura et al. 1999, 2001; Zaidi et al. 1985) or reproductive (Srivastava et al. 1992a, 1992b) effects. Additional studies are needed to examine the potential effects on the nervous and reproductive systems of developing organisms. No studies examined the developmental toxicity of styrene following dermal exposure.

3. HEALTH EFFECTS

Immunotoxicity. Occupational exposure studies have found alterations in lymphocyte subsets (Bergamaschi et al. 1995b; Biró et al. 2002), which may be indicative of reduced cell-mediated immunity and an impaired immune response to concanavalin (Tulinska et al. 2000). Limited data in animals indicate that inhalation (Ban et al. 2006) and oral (Sinitskij discussed in WHO 1983) exposure can also result in impaired immune response. No dermal exposure studies examining immunotoxicity were identified. Human and animal studies provide suggestive evidence that the immune system is a target; additional studies would be useful to further investigate the effect of styrene on immune function.

Neurotoxicity. The neurotoxicity of styrene in workers in the reinforced plastic industry has been extensively examined (Behari et al. 1986; Calabrese et al. 1996; Campagna et al. 1995, 1996; Castillo et al. 2001; Checkoway et al. 1992; Cherry et al. 1980; Chia et al. 1994; Dalton et al. 2003; Edling et al. 1993; Eguchi et al. 1995; Fallas et al. 1992; Fung and Clark 1999; Gamberale et al. 1976; Gobba et al. 1991, 1995; Gong et al. 2002; Härkönen et al. 1984; Iregren et al. 2005; Jegaden et al. 1993; Kishi et al. 2001; Lindstrom et al. 1976; Matikainen et al. 1993a, 1993b; Möller et al. 1990; Morata et al. 2002; Morioka et al. 1999; Muijser et al. 1988; Murata et al. 1991; Mutti et al. 1984a; Niklasson et al. 1993; Rosen et al. 1978; Seppäläinen and Härkönen 1976; Śliwińska-Kowalska et al. 2003; Štětkařová et al. 1993; Triebig et al. 1985, 2001; Tsai and Chen 1996; Viaene et al. 1998, 2001; Yuasa et al. 1996). A variety of neurological effects have been observed in these studies including decreased color discrimination, slowed reaction time, altered performance on neurobehavioral tests of memory and learning, altered vestibular function, altered hearing, reduced nerve conduction velocity, and increased clinical symptoms such as dizziness, tiredness, memory loss, and feeling drunk. Additionally, several experimental studies have examined the effects of acute exposure on vestibular function (Ödkvist et al. 1982; Stewart et al. 1968), clinical symptoms (Seeber et al. 2004; Stewart et al. 1968), and performance on neurobehavioral tests (Seeber et al. 2004). Animal studies have primarily focused on the damage to the organ of Corti and hearing loss (Campo et al. 2001; Crofton et al. 1994; Lataye et al. 2003; Loquet et al. 1999, 2000; Makitie et al. 2002; Pouyatos et al. 2002; Pryor et al. 1987; Yano et al. 1992), although nerve conduction velocity has also been examined (Kulig 1988; Yamamoto et al. 1997). The potential for neurotoxicity has not been examined in humans orally exposed to styrene and a limited number of end points have been examined in animals (Agrawal et al. 1982; Bushnell 1994; Husain et al. 1980, 1985; Khanna et al. 1994). The neurological effects observed in styrene workers were used as the basis of a chronic-duration inhalation MRL. Since this is based on a LOAEL, further studies which define the chronic NOAEL, as well as acute- and intermediate-duration NOAELs, would be valuable especially at levels of styrene causing problems with coordination and psychological function. These and other

3. HEALTH EFFECTS

neurological effects may play a role in the rate of workplace accidents and the level of performance. Additional studies in mammalian animal models are needed to determine if styrene causes chronic damage to the central and/or peripheral nervous systems and to determine the associated mechanism of toxicity. Also, information is needed to determine if neurotoxicity is a sensitive end point from exposure to styrene via the oral route.

Epidemiological and Human Dosimetry Studies. Numerous studies have examined the toxicity of styrene in workers, as discussed in other sections, most of these studies have focused on neurotoxicity and potential carcinogenicity of styrene. A common limitation of these studies is the poor characterization of exposure levels and possible exposure to other chemicals. Some studies provided no data on styrene exposure levels and other studies provide current exposure levels with limited or no data on past exposure levels. Occupational exposure and experimental studies also provide suggestive evidence of acute upper respiratory tract irritation and eye irritation (Carpenter et al. 1944; NIOSH 1983; Stewart et al. 1968) and possible endocrine effects (elevated levels of serum prolactin) (Arfini et al. 1987; Bergamaschi et al. 1996, 1997; Luderer et al. 2004; Mutti et al. 1984b); additional studies are needed to confirm the results of these studies and to establish dose-response relationships. Additionally, there are suggestive findings that styrene has the potential to induce reproductive effects (Cho et al. 2001; Härkönen and Holmberg 1982; Hemminki et al. 1980; Kolstad et al. 1999c; McDonald et al. 1988); however, poor characterization of styrene exposure, possible exposure to other compounds (particularly for the Cho et al. (2001) study), the low statistical power of the studies, and the lack of positive associations in the follow-up study (Hemminki et al. 1984) to the Hemminki et al. (1980) study limit the usefulness of the studies. Studies of males and female styrene workers examining a variety of reproductive end points and adequately characterized exposure would be useful.

Biomarkers of Exposure and Effect.

Exposure. Available studies indicate that there are good quantitative relationships between styrene metabolites (MA and PGA) in the urine and styrene exposure levels in humans (Bartolucci et al. 1986; Chua et al. 1993; Elia et al. 1980; Engstrom et al. 1976; Härkönen et al. 1978; Ong et al. 1994; Sedivec et al. 1984; Sollenberg et al. 1988; Symanski et al. 2001). Levels of styrene in blood have also been used as a biomarker of exposure (Antoine et al. 1986; Baselt et al. 1988a; Ramsey et al. 1980).

Effect. There are currently no biomarkers specific for the effects of styrene that are not also typical of other central nervous system depressants. Further research is needed to evaluate potential biomarkers of

3. HEALTH EFFECTS

effect in the areas of chromosome aberrations, psychomotor decrement, hepatic glutathione depletion, and adipose tissue retention of styrene. These potential biomarkers should be evaluated in terms of long-term or chronic exposure periods, and their specificity for exposure to styrene.

Absorption, Distribution, Metabolism, and Excretion. Styrene oxide (styrene epoxide) has been identified as an intermediate metabolite of styrene (Drummond et al. 1989; Engstrom et al. 1976; Korn et al. 1984, 1987; Leibman 1975; Lof et al. 1983; Withey and Collins 1979; Young et al. 1979). However, styrene oxide has only been found in minute amounts in human studies (Lof et al. 1986a). The presence of styrene oxide, a mutagen and carcinogen, may account for some conflicting results and/or interspecies variation in mutagenicity tests and cancer bioassays. The role, if any, of styrene oxide in the overall toxicity of styrene needs to be evaluated by additional metabolism studies to confirm its presence, level, and duration in human tissues. The toxicokinetics of styrene exposure via inhalation are reasonably well defined. However, oral and dermal exposure data are needed to better characterize absorption rates and the elimination ratios of the metabolites (MA and PGA).

Comparative Toxicokinetics. Interspecies variations in styrene metabolism have been established. Differences in the relative proportion of urinary metabolites, which is indicative of different metabolic pathways have been found in humans, mice, and rats. Additionally, there are differences in the kinetic constants for cytochrome P450 and epoxide hydrolase, which result in higher levels of reactive metabolites in the liver, lungs, and nasal epithelium. Also, mice appear to generate a higher proportion of R-styrene oxide than S-styrene oxide; the R-enantiomer is believed to be more cytotoxic. These metabolic differences are believed to result in mice being more sensitive than rats or humans to liver, lung, and nasal toxicity. Potential species differences in the neurotoxicity, the most sensitive end point in humans, have not been examined; if the neurological effects are due to styrene rather than one of its metabolites, the observed species differences may not be relevant. Efforts should continue to identify which animal model best approximates human metabolism of styrene.

Methods for Reducing Toxic Effects. Recommended methods for the mitigation of acute effects of styrene intoxication include mechanical ventilatory support, administration of oxygen, and drug therapy for bronchospasm, if exposure is by inhalation (Bronstein and Currance 1988; Ellenhorn and Barceloux 1988). Thorough washing or flushing with water is recommended for dermal/ocular exposure. Supportive treatment is indicated for neurological effects of styrene exposure (Haddad and Winchester 1990). No information was located concerning mitigation of effects of lower-level or longer-term exposure to styrene. Further information on techniques to mitigate such effects would be useful in

3. HEALTH EFFECTS

determining the safety and effectiveness of possible methods for treating styrene-exposed populations in the vicinity of hazardous waste sites. This includes further studies on the mechanism(s) of styrene toxicity, so that methods may be developed to interfere with or block styrene's toxic actions in the body.

Children's Susceptibility. Data needs relating to both prenatal and childhood exposures, and developmental effects expressed either prenatally or during childhood, are discussed in detail in the Developmental Toxicity subsection above.

No studies were identified that examined the toxicity of styrene in children or young laboratory animals. No consistently observed developmental effects have been reported in occupational exposure studies (Ahlborg et al. 1987; Härkönen et al. 1984; Lemasters et al. 1989) or in animal studies (Cruzan et al. 2005b; Daston et al. 1991; Kankaanpää et al. 1980; Murray et al. 1978). The nervous system is the most sensitive target of styrene toxicity in adults. No adverse styrene-related effects were observed in neurobehavioral function tests in rats exposed to styrene during gestation and lactation (Cruzan et al. 2005a); however, neurological effects have been observed in another inhalation study of rats exposed during gestation (Katakura et al. 2001) and in an oral gestation and lactation study (Zaidi et al. 1985). Possible neurological effects have not been assessed following post-weaning exposure; these data would be useful in evaluating whether growing children are more susceptible than adults to styrene-induced neurotoxicity.

Child health data needs relating to exposure are discussed in Section 6.8.1, Identification of Data Needs: Exposures of Children.

3.12.3 Ongoing Studies

No ongoing studies on styrene were identified in Federal Research in Progress database (FEDRIP 2007).

3. HEALTH EFFECTS

This page is intentionally blank.

4. CHEMICAL AND PHYSICAL INFORMATION

4.1 CHEMICAL IDENTITY


Information regarding the chemical identity of styrene is located in Table 4-1.

4.2 PHYSICAL AND CHEMICAL PROPERTIES

Information regarding the physical and chemical properties of styrene is located in Table 4-2.

4. CHEMICAL AND PHYSICAL INFORMATION

Table 4-1. Chemical Identity of Styrene

Characteristic	Information	Reference
Chemical name	Styrene	Verschuereen 1983
Synonym(s)	cinnamene; cinnamol; ethenylbenzene; phenylethylene; styrol; vinylbenzene;	Verschuereen 1983; HSDB 2007
Registered trade name(s)	No data	
Chemical formula	C ₈ H ₈	Windholz 1983
Chemical structure		IARC 1994
Identification numbers:		
CAS registry	100-42-5	Sax and Lewis 1987
NIOSH RTECS	WL3675000	HSDB 2007
EPA hazardous waste	No data	
EINICS	202-851-5	ESIS 2007
OHM/TADS	7216911	HSDB 2007
DOT/UN/NA/IMDG shipping	IMDG 3.3	HSDB 2007
	UN 2055	NLM 1989
HSDB	171	HSDB 2007
NCI	C0220	NLM 1989

CAS = Chemical Abstracts Service; DOT/UN/NA/IMDG = Department of Transportation/United Nations/North America/International Maritime Dangerous Goods Code; EPA = Environmental Protection Agency; ESIS = European chemical Substances Information System HSDB = Hazardous Substances Data Bank; NCI = National Cancer Institute; NIOSH = National Institute for Occupational Safety and Health; OHM/TADS = Oil and Hazardous Materials/Technical Assistance Data System; RTECS = Registry of Toxic Effects of Chemical Substances

4. CHEMICAL AND PHYSICAL INFORMATION

Table 4-2. Physical and Chemical Properties of Styrene

Property	Information	Reference
Molecular weight	104.15	O'Neil et al. 2001 Lide 2005
Color	Colorless to yellowish	Windholz 1983
Physical state	Liquid	Sax and Lewis 1987
Melting point	-30.6 °C	O'Neil et al. 2001
Boiling point	145.2 °C	Verschueren 2001; Weast 1985
Density at 20 °C	0.9059	O'Neil et al. 2001
Odor	If pure, sweet and pleasant; commonly contains aldehydes which provide it with a penetrating, sharp, and unpleasant smell	Verschueren 2001
Odor threshold:		
Water	0.73 mg/L 0.011 mg/L	HSDB 2007 Amoore and Hautala 1983
Air	1.36 mg/m ³	Amoore and Hautala 1983
Solubility:		
Water at 15 °C	280 mg/L	Verschueren 2001
Water at 20 °C	300 mg/L	
Water at 40 °C	400 mg/L	
Organic solvents	Soluble in alcohol, ether, acetone, carbon disulfide	Windholz 1983
Partition coefficients:		
Log K _{ow}	2.95	Hansch et al. 1995; EPA 1984a
Log K _{oc}	2.96	Sabljić et al. 1995
Vapor pressure at 20 °C	5 mmHg	Verschueren 2001
Henry's law constant at 25 °C	2.61x10 ⁻³ atm-m ³ /mol (calculated)	EPA 1981
Autoignition temperature	914 °F (490 °C)	Sax and Lewis 1987
Flashpoint	87 °F (31 °C) (closed cup) 34.4°C (Tag open cup)	O'Neil et al. 2001; Kirk-Othmer 2001
Flammability limits	No data 0.9 (lower); 6.8 (higher) 1.1 (lower); 6.1 (higher)	CEFIC 2002; Kirk-Othmer 2001
Conversion factors	1 mg/m ³ =0.23 ppm; 1 ppm=4.33 mg/m ³	Verschueren 2001

4. CHEMICAL AND PHYSICAL INFORMATION

This page is intentionally blank.

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

5.1 PRODUCTION

In the United States, styrene is produced principally by the catalytic dehydrogenation of ethylbenzene. Hence, ethylbenzene is a common contaminant. Styrene is also produced by oxidation of ethylbenzene to ethylbenzene hydroperoxide, which is then reacted with propylene to produce propylene oxide and α -methylphenyl carbinol. The carbinol is then further dehydrated to produce styrene (Dickson et al. 1973; HSDB 2007; IARC 1979). The first route of manufacture (dehydrogenation of ethylbenzene) represents 90% of styrene production. The other described method is the second most commonly used route of styrene synthesis. Other methods of styrene production are rarely used.

Styrene has been manufactured in the United States since 1938, with production increasing dramatically over the last 30 years. Since 1977, U.S. total styrene production has more than doubled. Production increased 16% in the decade between 1977 and 1987, but production increased >32% between 1987 and 1999, and rose again by another 28% between the years 1999 and 2006 (HSDB 2007; SRI 2006). Specifically, U.S. production of styrene in 1978 was 6.8 billion pounds, and then in 1987, production was approximately 8 billion pounds (USITC 1987, 1988). In 1999, U.S. styrene production was over 10 billion pounds, and in 2006, the United States produced >13 billion pounds (HSDB 2007; SRI 2006).

Information regarding the locations of the numerous styrene production facilities and the amounts of styrene that may be present on-site is presented in Table 5-1. Current domestic producers of styrene include Chevron Phillips Chemical Company LP (Aromatics and Styrenics Business Unit), St. James, Louisiana; Cos-Mar Company, Carville, Louisiana; the Dow Chemical Company, Freeport, Texas; INEOS Americas, LLC, Texas City, Texas; Lyondell Chemical Company, Channelview, Texas; NOVA Chemicals Corp, Bayport, Texas; Sterling Chemicals, Inc., Texas City, Texas; and Westlake Styrene Corporation, Sulphur, Louisiana (SRI 2006). The production of styrene at these facilities is directed primarily for captive processes (on-site conversion to other materials) or for merchant sales to other entities (these include export). The information presented in Table 5-1 reflects the locations of these production plants, where it can be noted that the greatest production capacity occurs primarily in Texas and Louisiana.

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

Table 5-1. Facilities that Produce, Process, or Use Styrene

State ^a	Number of facilities	Minimum amount on site in pounds ^b	Maximum amount on site in pounds ^b	Activities and uses ^c
AL	84	0	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
AR	57	100	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 14
AZ	54	0	999,999	1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12
CA	186	0	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
CO	24	100	9,999,999	2, 3, 6, 7, 8, 9, 11, 12, 13
CT	24	0	49,999,999	3, 5, 6, 7, 8, 9, 11, 12
DE	21	100	999,999	1, 2, 3, 4, 6, 7, 8, 11, 12, 14
FL	160	0	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
GA	100	0	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13
GU	1	100	999	9
HI	3	100	9,999	2, 3, 6, 7, 8, 10, 11
IA	48	0	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12
ID	15	0	99,999	1, 2, 3, 6, 7, 8, 11, 12
IL	106	0	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13
IN	163	0	99,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
KS	52	0	99,999,999	1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
KY	66	0	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
LA	119	0	10,000,000,000	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
MA	34	0	499,999,999	2, 3, 4, 6, 7, 8, 9, 12, 13
MD	39	100	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11
ME	15	100	99,999	2, 3, 6, 7, 8, 10
MI	99	0	99,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
MN	72	100	9,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13
MO	81	0	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12
MS	45	0	99,999,999	1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12
MT	9	100	9,999,999	3, 6, 7, 8, 9, 10, 13
NC	110	0	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12
ND	17	1,000	499,999,999	2, 3, 6, 7, 8, 9, 10, 11, 13
NE	38	0	999,999	1, 2, 3, 6, 7, 8, 10, 11, 12, 13
NH	26	100	499,999,999	2, 3, 5, 6, 7, 8, 10, 11, 12
NJ	52	100	99,999,999	2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13
NM	19	100	49,999,999	1, 2, 3, 6, 7, 8, 9, 11, 12
NV	11	1,000	99,999,999	2, 3, 6, 7, 8, 10, 11
NY	56	0	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
OH	198	0	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
OK	55	100	49,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
OR	55	100	49,999,999	1, 2, 3, 6, 7, 8, 9, 10, 11, 12
PA	123	0	99,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13
PR	29	0	9,999,999	1, 2, 3, 6, 7, 8, 10, 11, 12

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

Table 5-1. Facilities that Produce, Process, or Use Styrene

State ^a	Number of facilities	Minimum amount on site in pounds ^b	Maximum amount on site in pounds ^b	Activities and uses ^c
RI	29	100	999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12
SC	84	100	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14
SD	7	1,000	99,999	1, 2, 3, 6, 7, 8, 11
TN	103	0	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
TX	259	0	10,000,000,000	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
UT	42	0	99,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14
VA	63	0	49,999,999	1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12
VT	5	1,000	99,999	6, 7, 8
WA	89	0	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13
WI	92	0	499,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14
WV	40	100	999,999,999	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13
WY	2	1,000	9,999	6, 8, 9

^aPost office state abbreviations used

^bAmounts on site reported by facilities in each state

^cActivities/Uses:

- | | | |
|--------------------------|--------------------------|-----------------------------|
| 1. Produce | 6. Impurity | 11. Chemical Processing Aid |
| 2. Import | 7. Reactant | 12. Manufacturing Aid |
| 3. Onsite use/processing | 8. Formulation Component | 13. Ancillary/Other Uses |
| 4. Sale/Distribution | 9. Article Component | 14. Process Impurity |
| 5. Byproduct | 10. Repackaging | |

Source: TRI05 2007 (Data are from 2005)

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

5.2 IMPORT/EXPORT

Imports of styrene have generally been <1% of U.S. domestic production volumes, with imported styrene amounts decreasing over the last decades, and exported amounts increasing during the same time period. Styrene imports were reported to be 26.4 million pounds for 1976, 320 million pounds in 1986 (Dickson et al. 1973; IARC 1979), but only 1 million pounds in 1999 (HSDB 2007). These trends indicate a higher capacity for domestic producers to meet industry needs. Styrene exports were <1 billion pounds in 1978, but had exceeded 1 billion pounds by 1983. Exports have slowly increased such that recent export data indicate that the U.S. exports >2 billion pounds of styrene annually (HSDB 2007), also indicating that domestic production is more than capable to serve domestic needs.

5.3 USE

Styrene is used predominantly (65% of total product) in the production of polystyrene plastics and resins (James and Castor 2005). In addition, fiberglass products used for boats are also made from polyester resins dissolved in styrene. Styrene is also used as an intermediate in the synthesis of materials used for ion exchange resins and to produce copolymers such as styrene-acrylonitrile (SAN) and acrylonitrile-butadiene-styrene (ABS), both representing approximately 9% of styrene use, and styrene-butadiene rubber (SBR), representing approximately 6% of styrene use. SBR is used for such products as car tires, hoses used for industrial applications, and shoes. A related polymer, styrene-butadiene latex (approximately 7%), is used in making carpet, coatings for paper, and as part of latex paints. SAN and ABS are used for materials such as piping, automotive components, refrigerator liners, plastic drinking glasses, and car battery enclosures. An additional 7% of styrene is formulated with unsaturated polyester resins in such things as boat hulls (fiberglass reinforcement materials). The remaining amounts of styrene produced are used for several types of applications, including less common thermoplastics and even for laboratory and water purification uses (ion-exchange resins) and glues and adhesives (James and Castor 2005). Styrene copolymers are also frequently used in liquid toner for photocopiers and printers (HSDB 2007).

The Food and Drug Administration (FDA) permits styrene to be used as a direct additive for synthetic flavoring and an indirect additive in polyester resins, ion-exchange membranes, and in rubber articles (5% by weight maximum) intended for use with foods (HSDB 2007; IARC 1979; NIOSH 1983).

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

5.4 DISPOSAL

Typical means of styrene disposal include absorption on vermiculite or similar material, followed by disposal in an EPA-permitted landfill. Incineration is also a useful disposal method, but this must be carefully controlled since pure styrene is highly flammable (HSDB 2007). No data were located regarding the quantities of styrene disposed by these means on a national level, but the state of Massachusetts reported that most styrene disposal occurred via incineration (95.5%), followed by smaller amounts being disposed of in landfills (0.5%), a slightly greater amount being subjected to solvent recovery (0.7%), and slightly more being transferred to waste/energy brokers (3.3%) (Keenan and Harriman 1993). The total amounts represented were ~250,000 pounds. Whether the data reported for Massachusetts are representative of the proportions disposed of by these means in other states is not known.

5. PRODUCTION, IMPORT/EXPORT, USE, AND DISPOSAL

This page is intentionally blank.

6. POTENTIAL FOR HUMAN EXPOSURE

6.1 OVERVIEW

Styrene has been identified in at least 31 of the 1,689 hazardous waste sites that have been proposed for inclusion on the EPA National Priorities List (NPL) (HazDat 2007). However, the number of sites evaluated for styrene is not known. The frequency of these sites can be seen in Figure 6-1. Of these sites, 30 are located within the United States and one is located in the Commonwealth of Puerto Rico (not shown).

Styrene is a widely used industrial chemical with reported atmospheric emissions of >51 million pounds annually in the United States (TRI05 2007). Styrene photodegrades in the atmosphere, with a half-life ranging between 7 and 16 hours (which are the degradation half-lives catalyzed by reactions with hydroxyl radical and ozone, respectively). Styrene is moderately mobile in soil and volatilizes from water to the atmosphere. Styrene will undergo biodegradation in most top soils and aquatic environments, but degradation will be much slower in environments that are anaerobic. Bioconcentration does not appear to be significant.


The principal route of styrene exposure for the general population is probably by inhalation of contaminated indoor air. Mean indoor air levels of styrene have been reported in the range of 0.1–50 $\mu\text{g}/\text{m}^3$, and can be attributed to emissions from building materials, consumer products, and tobacco smoke. It should be pointed out that the workplace or home office may have substantially higher levels of airborne styrene, due to emissions from laser printers and photocopiers. General workplace styrene concentrations ranged from 89 to $1.5 \times 10^6 \mu\text{g}/\text{m}^3$ ($20\text{--}3.4 \times 10^5$ ppmv). The most significant exposure route in these settings is also likely by inhalation. The industries with the highest potential exposure are probably the reinforced plastics factories, boatbuilding facilities, and polystyrene factories. Exposure may also be high in areas near major spills. Exposure to styrene from hazardous waste sites is potentially important, but the magnitude of the problem is unknown. The potential for outdoor exposure to styrene is lower than indoor exposure, with reported mean air levels ranging from 0.28 to 20 $\mu\text{g}/\text{m}^3$ (0.064–4.6 ppmv).

6.2 RELEASES TO THE ENVIRONMENT

The Toxics Release Inventory (TRI) data should be used with caution because only certain types of facilities are required to report (EPA 2005). This is not an exhaustive list. Manufacturing and processing

6. POTENTIAL FOR HUMAN EXPOSURE

Figure 6-1. Frequency of NPL Sites with Styrene Contamination



Derived from HazDat 2007

6. POTENTIAL FOR HUMAN EXPOSURE

facilities are required to report information to the TRI only if they employ 10 or more full-time employees; if their facility is included in Standard Industrial Classification (SIC) Codes 10 (except 1011, 1081, and 1094), 12 (except 1241), 20–39, 4911 (limited to facilities that combust coal and/or oil for the purpose of generating electricity for distribution in commerce), 4931 (limited to facilities that combust coal and/or oil for the purpose of generating electricity for distribution in commerce), 4939 (limited to facilities that combust coal and/or oil for the purpose of generating electricity for distribution in commerce), 4953 (limited to facilities regulated under RCRA Subtitle C, 42 U.S.C. section 6921 et seq.), 5169, 5171, and 7389 (limited S.C. section 6921 et seq.), 5169, 5171, and 7389 (limited to facilities primarily engaged in solvents recovery services on a contract or fee basis); and if their facility produces, imports, or processes $\geq 25,000$ pounds of any TRI chemical or otherwise uses $>10,000$ pounds of a TRI chemical in a calendar year (EPA 2005).

6.2.1 Air

Estimated releases of more than 51 million pounds (>23 thousand metric tons) of styrene to the atmosphere from 1,563 domestic manufacturing and processing facilities in 2005, accounted for $>93\%$ of the estimated total environmental releases from facilities required to report to the TRI (TRI05 2007). These releases are summarized in Table 6-1.

Styrene may be emitted to the atmosphere from industrial production and usage processes, motor vehicle operation, combustion processes, building materials, and consumer products. Estimated atmospheric industrial styrene emissions reported to EPA for the 2005 TRI totaled >51 million pounds, with >41 million pounds released from point sources and >10 million pounds released as fugitive emissions (TRI05 2007). Styrene ranked 16th among air emissions for reported chemicals and chemical group compounds in the United States in 2005. Since EPA regulations that require reporting of toxic chemical emissions apply only to selected facilities producing and/or using substantial quantities of the chemical (EPA 1988a), the total air emissions of styrene are probably greater than those reported. Typical sources of industrial styrene emissions are those facilities producing styrene, polystyrene, other plastics, synthetic rubber, and resins (EPA 1975, 1987d; Graedel 1978; IARC 1979; NIOSH 1983). The number of facilities reporting styrene emissions to the TRI are listed in Table 6-1, along with number of reporting facilities in each state and Puerto Rico, and the primary routes of styrene release from those facilities.

Styrene has been identified as a component of motor vehicle emissions from both gasoline- and diesel-powered engines (Hampton et al. 1982, 1983). Styrene emission rates ranging from 6.2 to 7.0 mg/km

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-1. Releases to the Environment from Facilities that Produce, Process, or Use Styrene^a

State ^c	RF ^d	Reported amounts released in pounds per year ^b							
		Air ^e	Water ^f	UI ^g	Land ^h	Other ⁱ	Total release		On- and off-site
							On-site ^j	Off-site ^k	
AL	35	1,415,407	251	0	2,303	0	1,415,658	2,303	1,417,961
AR	25	624,028	0	0	33,520	0	624,028	33,520	657,548
AZ	26	495,751	0	0	60,884	0	495,751	60,884	556,635
CA	108	2,686,333	0	0	11,821	1,048	2,688,515	10,687	2,699,202
CO	10	101,053	No data	0	5,600	0	101,053	5,600	106,653
CT	6	52,488	1	0	52	0	52,489	52	52,541
DE	6	39,835	0	0	0	317	39,835	317	40,152
FL	119	4,439,463	1	0	753	62,382	4,439,967	62,632	4,502,599
GA	57	3,489,799	5	0	378	0	3,489,804	378	3,490,182
HI	1	3,488	5	0	5	0	3,498	0	3,498
IA	22	777,501	No data	0	5,824	480	777,501	6,304	783,805
ID	2	73,935	No data	0	0	0	73,935	0	73,935
IL	54	1,277,897	11	0	365,681	12,256	1,346,152	309,693	1,655,845
IN	96	5,490,355	170	0	73,269	46,527	5,491,275	119,046	5,610,321
KS	23	792,279	No data	0	517	0	792,279	517	792,796
KY	32	343,856	60	0	21,441	700	343,916	22,141	366,057
LA	42	673,578	138	0	51,517	0	673,716	51,517	725,233
MA	14	52,972	0	0	26,865	369	52,972	27,234	80,206
MD	10	587,546	0	0	0	212	587,546	212	587,758
ME	6	65,579	No data	0	0	0	65,579	0	65,579
MI	41	1,832,832	57	0	12,194	2,250	1,839,698	7,635	1,847,332
MN	31	1,185,353	0	0	0	1,015	1,185,353	1,015	1,186,368
MO	30	1,861,601	0	0	750	2,662	1,861,601	3,412	1,865,014
MS	14	561,660	10	0	157,603	0	561,670	157,603	719,272
MT	3	1	No data	0	0	0	1	0	1
NC	68	1,809,027	0	0	10,602	96,536	1,811,816	104,349	1,916,165
ND	4	210,811	5	0	0	0	210,816	0	210,816
NE	10	162,436	No data	0	8,811	0	162,461	8,786	171,247
NH	6	62,918	No data	0	0	0	62,918	0	62,918
NJ	24	192,588	3	0	2	0	192,593	0	192,593
NM	2	21,009	No data	0	0	0	21,009	0	21,009
NV	7	99,820	0	0	2,272	0	99,820	2,272	102,092
NY	18	89,190	26	0	8,999	96	89,216	9,095	98,311
OH	124	1,673,076	12	305	457,492	3,776	1,673,088	461,573	2,134,661
OK	21	482,448	0	0	118	0	482,448	118	482,566
OR	21	978,715	0	0	4,707	4,238	978,720	8,940	987,660

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-1. Releases to the Environment from Facilities that Produce, Process, or Use Styrene^a

State ^c	RF ^d	Reported amounts released in pounds per year ^b							
		Air ^e	Water ^f	UI ^g	Land ^h	Other ⁱ	Total release		
							On-site ^j	Off-site ^k	On- and off-site
PA	61	1,255,183	4	0	68,645	2,643	1,255,187	71,288	1,326,475
PR	6	45,994	0	0	45,517	9,817	45,994	55,334	101,328
RI	4	86,013	No data	0	0	0	86,013	0	86,013
SC	46	1,758,070	0	12	476,030	60	1,758,070	476,102	2,234,172
SD	3	78,181	No data	0	0	0	78,181	0	78,181
TN	47	5,425,447	255	0	55,904	0	5,425,702	55,904	5,481,606
TX	165	4,325,419	3,797	980,163	239,749	30,092	5,314,576	264,644	5,579,220
UT	9	87,271	No data	0	0	0	87,271	0	87,271
VA	19	734,299	No data	0	0	0	734,299	0	734,299
VT	1	10,546	No data	0	0	0	10,546	0	10,546
WA	26	1,252,647	43	0	1	2,668	1,252,691	2,668	1,255,359
WI	45	914,643	No data	0	27,261	10,143	914,643	37,404	952,046
WV	12	423,589	5	0	22,889	800	423,704	23,579	447,283
WY	1	11,660	No data	0	0	0	11,660	0	11,660
Total	1,563	51,115,590	4,858	980,480	2,259,976	291,087	52,187,233	2,464,757	54,651,991

^aThe TRI data should be used with caution since only certain types of facilities are required to report. This is not an exhaustive list. Data are rounded to nearest whole number.

^bData in TRI are maximum amounts released by each facility.

^cPost office state abbreviations are used.

^dNumber of reporting facilities.

^eThe sum of fugitive and point source releases are included in releases to air by a given facility.

^fSurface water discharges, waste water treatment-(metals only), and publicly owned treatment works (POTWs) (metal and metal compounds).

^gClass I wells, Class II-V wells, and underground injection.

^hResource Conservation and Recovery Act (RCRA) subtitle C landfills; other on-site landfills, land treatment, surface impoundments, other land disposal, other landfills.

ⁱStorage only, solidification/stabilization (metals only), other off-site management, transfers to waste broker for disposal, unknown

^jThe sum of all releases of the chemical to air, land, water, and underground injection wells.

^kTotal amount of chemical transferred off-site, including to POTWs.

RF = reporting facilities; UI = underground injection

Source: TRI05 2007 (Data are from 2005)

6. POTENTIAL FOR HUMAN EXPOSURE

distance for gasoline-powered vehicles and 1.4–2.1 mg/km for diesel trucks have been reported (Hampton et al. 1983). Styrene may also be emitted into the air by other combustion processes. Styrene has been identified in the stack emissions from waste incineration (Junk and Ford 1980), and Kleindienst et al. (1986) reported the presence of styrene in wood smoke emissions, but no quantitative data were reported.

Emissions of styrene from building materials (carpets, floor tiles, insulation), office copiers, and consumer products (disinfectants, plastics, paint, cigarettes) may contribute significantly to indoor air pollution (Crump 1995). A styrene emission rate from glued carpet of 98 ng/minute/m² was calculated by Wallace et al. (1987b), and Girman et al. (1986) identified styrene as a major emittant from adhesives used in the constructing and finishing of buildings. Hodgson et al. (1993) determined an average styrene emission rate from new carpets of 410 ng/minute/m² over a 24-hour time period, but this was reduced to 30 ng/minute/m² when emissions were measured over 168 hours. Carpet cushioning material showed higher styrene emission rates of 2,300 ng/minute/m² when measured over 6 hours, but this material also showed significantly lower emission rates of 83 ng/minute/m² when measured over a longer span of 96 hours (Schaeffer et al. 1996). Polystyrene products such as packaging materials, toys, housewares, and appliances that may contain small amounts of the monomer also contribute to air levels. The workplace or home office may have substantial levels of airborne styrene due to emissions from laser printers and photocopiers. In the case of laser printers, styrene concentrations measured in test chambers during printer operation were reported to be as high as 380 µg/m³ (87 ppmv) (Kagi et al. 2007). For photocopiers, emission rates from four different copiers averaged 3,300 µg/hour, but one copier had an emission rate of 12,000 µg/hour (Leovic et al. 1996). General workplace styrene concentrations ranged from 89 to 1.5x10⁶ µg/m³ (20–3.4x10⁵ ppmv). Styrene has also been detected in sidestream smoke emitted from cigarettes but concentrations were not reported (IARC 1979).

6.2.2 Water

Estimated releases of 4,858 pounds (~2.2 metric tons) of styrene to surface water from 1,563 domestic manufacturing and processing facilities in 2005, accounted for about 0.01% of the estimated total environmental releases from facilities required to report to the TRI (TRI05 2007). No styrene was reported to have been released to publicly owned treatment works (POTWs) (TRI05 2007). These releases are summarized in Table 6-1.

The principal sources of styrene releases to water are industrial effluents. Styrene has been detected in effluents from chemical, textile, latex, and coal gasification plants (EPA 1976; Pellizzari et al. 1979).

6. POTENTIAL FOR HUMAN EXPOSURE

Styrene was also identified in one of 63 industrial effluents at a concentration of <10 µg/L (EPA 1979b). Styrene occurred at concentrations up to 83 µg/L in coal gasification effluents (Pellizzari et al. 1979), and King and Sherbin (1986) reported styrene concentrations up to 970 µg/L in chemical plant effluents. The daily styrene loading from a single chemical plant into the St. Clair River (just south of Lake Huron on the Michigan/Ontario border) was estimated at 133 kg (King and Sherbin 1986). Styrene was detected (but not quantified) in the leachate from an industrial landfill in a study of 58 municipal and industrial landfill leachates (Brown and Donnelly 1988). Styrene has also been detected at trace concentrations in the River Elbe at two different sampling locations, with concentrations ranging from 6.1 to 46 ng/L (Gotz et al. 1998).

6.2.3 Soil

Estimated releases of more than two million pounds (~1,000 metric tons) of styrene to soils from 1,563 domestic manufacturing and processing facilities in 2005, accounted for about 4.1% of the estimated total environmental releases from facilities required to report to the TRI (TRI05 2007). An additional 0.980 million pounds (~444 metric tons), constituting about 1.8% of the total environmental emissions, were released via underground injection (TRI05 2007). These releases are summarized in Table 6-1.

Soil and sediments may become contaminated with styrene by chemical spills, landfill disposal of styrene-containing wastes, or discharge of styrene-contaminated water. A small amount of styrene is produced naturally through the activities of microorganisms, and some plants also produce styrene that may be released to soil. The amounts released to soil through these processes, however, are not expected to be significant in comparison to human activities that generate and release styrene to soil.

6.3 ENVIRONMENTAL FATE**6.3.1 Transport and Partitioning**

Should styrene be released to the environment, its vapor pressure indicates that it will partition to the atmosphere. In the atmosphere, styrene exists as a vapor. Styrene is an oily liquid that is slightly volatile; its vapor pressure has been determined to be approximately 5 mm Hg at 20 °C (Verschueren 2001). A small fraction of the styrene released to the atmosphere may dissolve into condensed water vapor such as clouds and raindrops. A Henry's law constant (H) is a measure of the tendency of a chemical to partition between its gas phase and water. A value for H has not been experimentally measured for styrene, but it

6. POTENTIAL FOR HUMAN EXPOSURE

may be estimated by dividing the vapor pressure of styrene by its solubility in water at the same temperature (EPA 1982a). In this case, the value of H is approximately 2.61×10^{-3} atm-m³/mole at 25 °C. Analogous air-water partition coefficients were measured at 37 °C, yielding a value of approximately 5.4×10^{-1} atm-m³/mole (Sato and Nakajima 1979). The magnitude of these values suggests that only a small fraction of vapor-phase styrene would dissolve into atmospheric water droplets. Physical processes such as precipitation and dry deposition would not be significant mechanisms for removing styrene from the atmosphere because of its high photochemical reactivity (EPA 1984b).

Should styrene be released to lakes, rivers, streams, or other waterways, the magnitude of the estimated Henry's law constant (2.61×10^{-3} atm-m³/mole, assuming a water solubility of 300 mg/L at 20 °C [Verschueren 2001]) and its water solubility suggest that a large fraction of the chemical dissolved in the water will volatilize into the atmosphere (depending on temperature gradients, relative humidity, air currents, and the extent of mixing of the solution). From a modeling perspective, it should be pointed out that the dependence of this process depends on an accurate determination of the chemical's water solubility. In this case, while styrene is only sparingly soluble in water, the reported values are not precise, and range from 160 mg/L at 23 °C to 310 mg/L at 20 °C to 400 mg/L at 40 °C (Banerjee et al. 1980; Valvani et al. 1981; Verschueren 2001). Therefore, estimated rates of volatilization have some uncertainty associated with them. Volatilization of styrene, however, has been measured in surface lake and distilled water samples such that different loadings of styrene were examined. When 2–10 mg of styrene was added to 1 L of lake water, 50% was lost to the atmosphere in 1–3 hours. For the distilled water, volatilization occurred more slowly, with 50% loss occurring within 6–7 hours (Fu and Alexander 1992). The volatilization half-life of styrene in moving water that is 1 meter deep (assuming a solubility of 300 mg/L) may be on the order of 6 hours, based on the empirical relationship reported by Dilling (1977) for the volatilization of chlorinated hydrocarbons from water. The half-life of styrene in the Rhine River was estimated from field measurements at about 14 hours, but it was not known if the loss was due to volatilization, biodegradation, or photodegradation (Zoeteman et al. 1980). Volatilization from ponds and lakes is estimated to be slower, with half-life estimates ranging from 3 to 13 days (EPA 1984b).

Styrene released to soils or sediments will also likely volatilize to the atmosphere, but the rate of this process depends on the characteristics of the soil or sediment. The extent of adsorption of sparingly water-soluble compounds such as styrene is often correlated with the organic carbon content of the adsorbent (i.e., the soil or sediment; Hassett et al. 1983). When adsorption is expressed as a function of organic-carbon content, an organic carbon/water partition coefficient (K_{oc}) is generated, and may be used to rank the relative mobility of the chemical in soil. A K_{oc} value for styrene has not been experimentally

6. POTENTIAL FOR HUMAN EXPOSURE

measured, but may be estimated from its solubility in water, using the empirical regression of Hassett et al. (1983). Assuming that the solubility of styrene is 300 mg/L, a calculated K_{oc} value for styrene is 260. The magnitude of this estimated K_{oc} suggests that styrene is "moderately mobile" in soil (Roy and Griffin 1985). In surface soils, where the amount of organic carbon will be highest, the movement of styrene will therefore be retarded by adsorption. In deeper subsurface environments where the amount of organic carbon may be lower, adsorption may not be as significant. Based on field measurements, the rate of movement of styrene in an aquifer was about 80 times slower than that of the groundwater (Roberts et al. 1980). The slower rate of movement was attributed to adsorption. No information was located to corroborate the estimated K_{oc} value, and apparently, there are no studies in which the adsorption-desorption characteristics of styrene by soils and sediments have been measured.

The octanol/water partition coefficient (K_{ow}) reflects the partitioning of a chemical between octanol and water and is believed to be a good indication of the tendency for a chemical to accumulate in the fatty structures in plants and animal tissues (Kenaga and Goring 1980). The K_{ow} of styrene has been measured to be 1,445 (Banerjee et al. 1980), 891 ($\log K_{ow}=2.95$) (Hansch et al. 1995), and 891 (Valvani et al. 1981), suggesting that styrene will partition to fat tissues. This is shown to be the case by the work of Engstrom et al. (1978a) and EPA (1986d).

Even though styrene does tend to partition into fat, it does not tend to bioaccumulate to high levels, mainly because of its metabolism and excretion. A bioconcentration factor (BCF) relates the concentration of a chemical in an organism to the concentration of the chemical in the medium in which it is exposed. Based on the empirical regression of Kenaga (1980), the BCF for styrene is about 25. An experimentally-measured BCF for goldfish was 13.5 (Ogata et al. 1984). These low BCFs suggest that bioconcentration is not a significant fate of styrene released into the environment (EPA 1984b). No other measured BCFs were located to corroborate these reported values.

6.3.2 Transformation and Degradation

6.3.2.1 Air

The major fate of atmospheric styrene is determined by the rate of photooxidation. Styrene may be transformed by direct photolysis, but the half-life of this process may be on the order of 50 years (EPA 1984b). Kopczynski et al. (1972) found that styrene was not degraded by direct photolysis after 6 hours of exposure.

6. POTENTIAL FOR HUMAN EXPOSURE

Styrene is more quickly photooxidized by ozone and hydroxyl radicals. The rate constant for the reaction of styrene with ozone at ambient temperatures (about 25 °C) has been measured and is approximately $0.17\text{--}2.16 \times 10^{-19}$ cm³/molecule-second (Atkinson et al. 1982; EPA 1979a). Assuming that the mean concentration of ozone in the troposphere is 10×10^{12} molecules/cm³ (EPA 1980), the half-life of styrene would be approximately 13 hours. The rate constant for the reaction of styrene with hydroxyl radicals has been measured as 5.3×10^{-11} cm³/molecule-second (Bignozzi et al. 1981). Assuming that the concentration of tropospheric hydroxyl radicals varies from 3×10^5 to 1×10^7 molecules/cm³ (Mac Leod et al. 1984), it follows that the atmospheric half-life of styrene would be between 0.5 and 17 hours. More recent studies provide a similar rate constant of 5.9×10^{-11} cm³/molecule-second (Bunce and Dryfhout 1992) and 5.8×10^{-12} cm³/molecule-second (EPA 1993a), with a corresponding half-life of ~2.2 hours. Consequently, it is not expected that styrene will persist in the atmosphere, due to the combined and rapid effects of ozone- and hydroxyl radical-initiated atmospheric degradation processes. Transformation products resulting from such degradation processes include primarily oxygen-containing hydrocarbons such as phenol, phenylacetaldehyde, and phenoxy radical (Sloane and Brudzynski 1979) or other compounds (Atkinson and Arey 2003), as well as other aromatic hydrocarbons such as the benzyl radical and other unsaturated hydrocarbons (Sloane and Brudzynski 1979).

6.3.2.2 Water

Little is known about abiotic transformations of styrene in water. The reaction of styrene with peroxy radicals appears to be too slow to be significant (EPA 1984b), and no relevant information regarding photochemical reactions in water was located. There is no information that styrene will hydrolyze in water, nor would its chemical structure suggest such potential.

While little is known about the abiotic degradation potential of styrene in water, it has been shown to be biologically degraded in several types of aquatic systems including sewage treatment facilities, biofilm reactors, groundwater, and lakes. In sewage samples, styrene showed a range of biodegradability. In one study where styrene was added at a concentration of 1 mg/L, 20% of the added styrene was completely degraded (i.e., mineralized to CO₂) within 3 days, with >60% degraded within 30 days (Fu and Alexander 1992). In another sewage fate study, however, styrene was only slightly biodegraded; in five different evaluations, conducted for either 17 or 36 days, styrene degradation ranged from 6 to 23% (Pahren and Bloodgood 1961). Bridie et al. (1979) found that 42% of the styrene initially present degraded in 5 days when unadapted sewage was used as the source of microorganisms; when an adapted sewage was used, 80% degradation was observed. Very low concentrations of styrene (<10 µg/L) were completely

6. POTENTIAL FOR HUMAN EXPOSURE

degraded within 20 minutes in an aerobic biofilm reactor after acclimation, but the chemical was degraded much more slowly (~8% after 2 days) under methanogenic biofilm column conditions (Bouwer and McCarty 1984). When studied in groundwater, styrene biodegradation ranged between 4 and 12% per week (when added at 600–800 µg/L; Wilson et al. 1983) and a more recent study showed similar rates of degradation, with over 40% mineralization after 30 days (Fu and Alexander 1992). Styrene had a similar degradation potential in lake water, with ~35% degradation after 30 days (Fu and Alexander 1992).

6.3.2.3 Sediment and Soil

Styrene is rapidly degraded in most soils when incubated under aerobic conditions, but it persists when soil conditions are anaerobic (e.g., waterlogged). Styrene was rapidly degraded when added to either a low organic-matter content landfill soil or a high organic-matter content loamy soil. When added at 2 g/kg, >87% of the styrene was degraded in the landfill soil, and >95% was degraded in the loamy soil (measured over a period of 16 weeks; Sielicki et al. 1978). When styrene was added to the soils at higher concentrations (5 g/kg), degradation was slower and less was degraded, with ~60% degraded in both soils after a 16-week incubation. Fu and Alexander (1992) showed that styrene was biodegraded in loamy soil (when added at 2 mg/kg soil), with >50% degradation occurring over a 30-day incubation period, and the same soil in a later study showed similar amounts of degradation (40% degradation in 50 days; Fu et al. 1994). In contrast to the findings in aerobic soils, this research group showed that styrene persisted when soil conditions were waterlogged and anaerobic (Fu and Alexander 1996). Other researchers, however, have been able to demonstrate degradation of styrene under anaerobic conditions by consortia of microbial cultures (Grbic-Galic et al. 1990). The relevance of these consortia studies to actual environmental samples is unknown.

Several bacterial and fungal species have been isolated from soils that are capable of using styrene as a sole-carbon source (Braun-Lüllemann et al. 1997; Burbach and Perry 1993; Hartmans 1995; Sielicki et al. 1978; Warhurst and Fewson 1994), and these organisms degrade styrene by either side chain oxidation or aromatic ring attack. Initial biodegradation products included styrene oxide, 1-phenylethanol, 2-phenylethanol, and then phenylacetaldehyde, acetophenone, and phenylethanedol, and then phenylacetic acid, with degradation proceeding towards normal metabolic intermediates such as acetaldehyde and pyruvate (Warhurst and Fewson 1994).

6. POTENTIAL FOR HUMAN EXPOSURE

6.4 LEVELS MONITORED OR ESTIMATED IN THE ENVIRONMENT

Reliable evaluation of the potential for human exposure to styrene depends in part on the reliability of supporting analytical data from environmental samples and biological specimens. Concentrations of styrene in unpolluted atmospheres and in pristine surface waters are often so low as to be near the limits of current analytical methods. In reviewing data on styrene levels monitored or estimated in the environment, it should also be noted that the amount of chemical identified analytically is not necessarily equivalent to the amount that is bioavailable. The analytical methods available for monitoring styrene in a variety of environmental media are detailed in Chapter 7.

6.4.1 Air

Styrene is a common contaminant of ambient urban air. Concentrations of styrene greater than rural air concentrations have been identified in urban and industrial source areas, near hazardous waste sites, in motor vehicle tunnels, in indoor air, and in workplace environments. A summary of monitoring data for these locations is presented in Table 6-2. The data suggest that indoor air concentrations of styrene can be considerably higher than outdoor concentrations. Cigarette smoke has also been implicated as a significant source of styrene in indoor air (EPA 1987e; Wallace et al. 1986a), as has the operation of photocopying machinery (Stefaniak et al. 2000; Leovic et al. 1996, 1998) and laser printers (Kagi et al. 2007).

Monitoring studies in Minnesota detected styrene in over 1,400 air samples collected from a total of 2,507 samples (there were 1,004 samples where styrene was below the detection limits) over an 8-year period. The average concentration detected was $0.1 \mu\text{g}/\text{m}^3$ (0.02 ppmv); the median concentration detected was $0.08 \mu\text{g}/\text{m}^3$ (0.02 ppmv); and the maximum detected amount was $1.49 \mu\text{g}/\text{m}^3$ (0.343 ppmv) (Pratt et al. 2000). Styrene monitoring in ambient air conducted in Chiba City, Japan over an 8-week period showed slightly higher mean concentrations, ranging from 0.11 to $0.36 \mu\text{g}/\text{m}^3$ (0.025–0.083 ppmv) (Uchiyama and Hasegawa 2000).

6.4.2 Water

Styrene is not frequently found in U.S. water supplies. Styrene was not detected in any of the >1,000 samples of drinking water analyzed during three federal surveys (EPA 1985a), but had been reported occasionally in drinking water supplies in several states (Coleman et al. 1984; EPA 1975, 1976; Kleopfer and Fairless 1972; Kool et al. 1982; Sanjivamurthy 1978) well water (EPA 1985b; Krill and

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-2. Styrene Concentrations in Representative Air Samples

Location	Concentration ($\mu\text{g}/\text{m}^3$)		References
	Maximum	Mean	
Indoor			
Indoors, unspecified	6,500	0.4–8.9 0.3–50	EPA 1987e, 1988c; Fishbein 1992; Sakaguchi and Akabayashi 2003; Shields and Weschler 1992; Wallace et al. 1986a
New homes	14.3	0.9–2.6	Hodgson et al. 2000
Remodeled buildings	167	0.1–10.1	Rothweiler et al. 1992; Zabiegala 1999
Workplaces	4.5×10^6	$<1-1.5 \times 10^6$	Bartolucci et al. 1986; Cocheo et al. 1983; Correa et al. 2004; Fishbein 1992; Kagi et al. 2007; NIOSH 1983
Photocopy centers	12,000 ^a ; 220 ^b	7,000 ^a ; 89 ^b	Stefaniak et al. 2000; Leovic et al. 1996, 1998
Outdoor			
Rural/suburban	53	0.28–0.34 ^c ; 0.43	EPA 1988c; Graedel 1978; Islam and Stancheva 1999; Kinney et al. 2002
Urban	2,500	0.29–3.8; 20	EPA 1987e, 1988c; Fishbein 1992; Grosjean and Fung 1984; Grosjean et al. 1998; Harkov et al. 1985; Wallace et al. 1986b
Industrial source areas	25	1.3 ^c –2.1	EPA 1978, 1983, 1988c
Municipal waste sites	6,100	No data	Assmuth and Kalevi 1992; Eitzer 1995
Hazardous waste sites	65	1.1–6.4	Harkov et al. 1985; La Regina and Bozzelli 1986
Tunnels	46	1.1–6.6 ^d	Hampton et al. 1983; Zielinska et al. 1996

^aValue provided is the emission rate, as $\mu\text{g}/\text{hour}$; the emission rate was measured during operation of copier in test chambers.

^bValue provided is the emission rate, as $\mu\text{g}/\text{hour}$; the emission rate was measured during the idle phase of copier operation in test chambers.

^cMedian value

^dRange of values, no mean given.

6. POTENTIAL FOR HUMAN EXPOSURE

Sonzogni 1986), river water (EPA 1976; Sheldon and Hites 1978), and Lake Erie (Konasewich et al. 1978). Quantitative data were not available in these reports. A more recent survey provided by the EPA National Contaminant Occurrence Database (EPA 2006) noted that styrene was detected rarely in groundwater, where it was detected only 295 times out of over 174,000 analyses (<0.2% of the samples); detected concentrations ranged from <10 to 40 µg/L. Styrene concentrations in raw and treated waters ranged from 0.1 to ≥1.0 µg/L in an evaluation of organic compounds in Canadian water supplies at nine municipalities along the Great Lakes (Otson 1987). Styrene was rarely detected in aquifer materials in a large evaluation of U.S. groundwater and wells conducted by the U.S. Geological Survey (USGS). For >3,400 test evaluations of aquifer materials, there was a very low frequency of detections, with the median concentration being 0.015 µg/L. It was detected less frequently and also at low concentrations in domestic wells (0.014 µg/L), and at slightly higher concentrations in public wells (median=0.13 µg/L) (USGS 2006).

Other bodies of water—those potentially highly contaminated and those that are used for direct ingestion—likewise do not usually contain styrene. Styrene is not commonly detected in groundwater even near superfund sites, and was not found in a drinking water evaluation of public water sources in Torino, Italy (Canter and Sabatini 1994; Zelano et al. 1998). In a study of wells near a superfund site in Florida, styrene was detected at a very low concentration (the maximum detected concentration was 6.3 µg/L) near the site, but it was not detected in any treated water effluents (Canter and Sabatini 1994). Squillace et al. (1999) surveyed a vast number of drinking water wells in the United States over a 10-year period. Styrene was detected in <1% of the 2,900 surveyed urban and suburban wells between 1985 and 1995, and the concentrations detected were >2 orders of magnitude lower than the health advisory level. Styrene was only detected in rural wells, not in urban ones. Finally, Zelano et al. (1998) did not report the detection of any styrene in an evaluation of 21 public drinking water fountains in Torino, Italy.

6.4.3 Sediment and Soil

Limited data were located regarding estimation of styrene in sediments and soils (see Section 6.2.3). Water and sediment samples from the Lower Tennessee River were evaluated for styrene, and low concentrations (4.2 ppb) were found (Goodley and Gordon 1976).

6. POTENTIAL FOR HUMAN EXPOSURE

6.4.4 Other Environmental Media

Styrene is a natural component of many foods; however, most styrene associated with food is the result of packaging of the food material in polystyrene containers. Styrene has been found as a natural component of roasted filberts, dried legumes, fried chicken, cooked pork, roasted beef, mussels, clams, eggs, nectarines, and Beaufort cheese (Dumont and Adda 1978; Kinlin et al. 1972; Lovegren et al. 1979; Takeoko et al. 1988; Tang et al. 1983, 2000), but detected concentrations were often very low (Tang et al. 2000), except for turkey sausage, where detected levels were 100 ppb, and in some cheeses, where concentrations detected were up to 5,000 ppb (Tang et al. 2000). In contrast, styrene is a natural component of cinnamon, with concentrations up to 40,000 ppb (Tang et al. 2000). Data on styrene levels from an FDA monitoring study (1996–2000) of volatile organic compounds in food items are presented in Table 6-3 (Fleming-Jones and Smith 2003).

Styrene may enter packaged foods by migration from polystyrene food containers and packaging materials, with concentrations ranging from <100 to >3,000 ppm, but common levels being much lower (5–30 ppb) (EPA 1985a; Tang et al. 2000). Concentrations of styrene measured in yogurt packaged in polystyrene containers ranged from 5.5 to 150 ppb (Withey 1976). Mean levels of styrene in foods packaged in plastic in the United Kingdom ranged from <1 to 180 ppb (Gilbert and Startin 1983). Similar concentrations of styrene were detected in other dairy products packaged in polystyrene containers (IARC 1979). The rate of styrene migration into food is mainly a function of the diffusion coefficient of the monomer in the polymer and of the lipophilicity of the food (Till et al. 1987). For example, 4–6% of the free monomer in polystyrene packaging migrated into corn oil or sunflower oil within 10 days, while only 0.3–0.6% migrated into milk, beef, or water. Similarly, migration of styrene from foam cups into liquids such as water, tea, or coffee was about 8 ng/cm², while migration into 8% ethanol (as might be encountered in wine or other alcoholic drinks) was 36 ng/cm² (Varner and Breder 1981). However, Withey and Collins (1978) found no clear relationship between the styrene monomer content of packaging material (which varied widely) and the amount leached into food after comparable residence times. Styrene was detected, but not quantified, in samples of mother's milk from four urban areas (Pellizzari et al. 1982).

Styrene has been identified as a component of cigarette smoke (EPA 1984b) and has been detected in concentrations of 18 µg/cigarette in the smoke of cigarettes made in the United States (IARC 1979). Indoor air concentrations of styrene may be significantly higher in homes of smokers than nonsmokers (EPA 1987e).

6. POTENTIAL FOR HUMAN EXPOSURE

Table 6-3. Styrene Levels in Food Items

Food	Number of detections	Minimum (ppb)	Maximum (ppb)
American cheese	4	2	11
Cheddar cheese	3	4	70
Mixed cuts	14	21	104
Ground beef	6	4	13
Pork bacon	7	6	85
Cream cheese	2	2	3
Frankfurters, beef	8	4	77
Chocolate cake with icing	12	7	57
Tuna canned in oil	1	2	2
Fruit flavored cereal	3	2	10
Eggs, scrambled	6	5	10
Peanut butter	13	16	38
Avocado, raw	8	3	550
Popcorn, popped in oil	3	2	2
Blueberry muffin	10	8	141
Strawberries, raw	9	12	350
Orange, raw	2	2	3
Coleslaw with dressing	1	2	2
Sweet roll/Danish	13	13	91
Potato chops	6	2	16
Popsicle	3	4	11
Quarter pound hamburger, cooked	6	4	27
Margarine	10	9	20
Sandwich cookies	14	15	165
Butter	12	11	28
Chocolate chip cookies	12	15	111
Sour cream	3	5	30
Apple pie, fresh/frozen	9	10	40
Chicken nuggets	12	10	66
Graham crackers	6	4	21
French fries	12	8	68
Cheeseburger, quarter pound	5	5	22
Cheese pizza	6	3	23
Bologna	7	2	78
Cheese and pepperoni pizza	7	8	20
Olive/safflower oil	11	3	54
Sugar cookies	14	24	142
Cake doughnuts with icing	10	6	45

Source: Fleming-Jones and Smith (2003).

6. POTENTIAL FOR HUMAN EXPOSURE

6.5 GENERAL POPULATION AND OCCUPATIONAL EXPOSURE

Exposure to styrene may occur by inhalation, ingestion, or dermal absorption. The most likely mode of exposure of the general population to styrene is by inhalation of indoor air (EPA 1985a). Based on the EPA (1989e) estimate that the average person spends 20.4 hours/day indoors (inhaling about 17 m³ of air during that time based on an air inhalation rate of 20 m³/day) and the range of mean indoor air concentrations presented in Table 6-2, typical indoor exposure levels to styrene may range from 1.7 to 850 µg/day. Additional exposures may occur from inhalation of outdoor air and ingestion of food that was stored in polystyrene containers. Based on estimated food consumption rates, Tang et al. (2000) reported an estimated annual general population exposure to styrene ranging from 0.8 to 4.5 mg/person from food. Outdoor air concentrations are likely to be lower in rural than urban areas and are likely to be small compared to indoor air concentrations. Exposure from municipal drinking water is probably insignificant. However, groundwater at hazardous waste sites where styrene has been detected may provide significant exposure to styrene if used as a local water supply.

The exposure of the population to styrene varies significantly from the typical to the worst-case scenario. The daily general population exposure to styrene via food has been estimated at 0.2–1.2 µg/person and the exposure via inhalation has been estimated at 18–54 µg/person, with a total estimated exposure ranging from 18.2 to 55.2 µg/day. This is equivalent to 6.7–20.2 mg/year. Therefore, the primary route of exposure for the general population is via inhalation (Tang et al. 2000). Worst-case exposure estimates, on the other hand, are 0–0.5 µg/day from drinking water, 30 µg/day from food, and 65,000 µg/day from air (EPA 1985a). These estimates are based on the highest levels estimated or monitored and, therefore, reflect the highest potential exposure rather than typical exposure for the general population.

Exposure of the general population to styrene is confirmed by human monitoring data. Styrene has been identified in adipose tissue at concentrations of 8–350 ng/g (EPA 1986d), in blood at a mean concentration of 0.4 µg/L (Antoine et al. 1986), and in exhaled breath at mean concentrations of 0.7–1.6 µg/m³ (EPA 1987e).

A large number of workers are potentially exposed to styrene. NIOSH estimates that approximately 300,000 workers at 22,000 facilities may be exposed to styrene (NIOSH 1990); about 30,000 of these on a full-time basis (NIOSH 1983) and about 86,000 are females. The highest potential exposure occurs in the reinforced-plastics industry, where workers may be exposed to high air concentrations and also have dermal exposure to liquid styrene or resins (Lemasters et al. 1985; NIOSH 1983). Hemminki and Vianio

6. POTENTIAL FOR HUMAN EXPOSURE

(1984) estimated that heavily exposed workers in this industry in Finland might be exposed to up to 3 g of styrene per day. Significant occupational exposures may also occur in other industrial settings, including styrene polymerization, rubber manufacturing, and styrene-polyester resin facilities (Engstrom et al. 1978b; NIOSH 1983; Rappaport and Fraser 1977) as well as in photocopy centers or facilities (Leovic et al. 1996, 1998; Stefaniak et al. 2000).

6.6 EXPOSURES OF CHILDREN

This section focuses on exposures from conception to maturity at 18 years in humans. Differences from adults in susceptibility to hazardous substances are discussed in Section 3.7, Children's Susceptibility.

Children are not small adults. A child's exposure may differ from an adult's exposure in many ways. Children drink more fluids, eat more food, breathe more air per kilogram of body weight, and have a larger skin surface in proportion to their body volume. A child's diet often differs from that of adults. The developing human's source of nutrition changes with age: from placental nourishment to breast milk or formula to the diet of older children who eat more of certain types of foods than adults. A child's behavior and lifestyle also influence exposure. Children crawl on the floor, put things in their mouths, sometimes eat inappropriate things (such as dirt or paint chips), and spend more time outdoors. Children also are closer to the ground, and they do not use the judgment of adults to avoid hazards (NRC 1993).

Children can be exposed to styrene at home by inhalation of contaminated air and by food consumption. Inhalation-based exposures may occur in both urban and rural home environments, both of which may be contaminated by vehicular and industrial emissions. In addition, exposure to tobacco smoke may provide another route of styrene exposure, especially in homes where one or both parents, any siblings, or other relatives smoke. Children may be also be exposed to higher levels of styrene indoors at home during painting of indoor rooms, especially during winter months (such as over winter school vacations) when the child stays indoors more and during which time, windows may not be opened.

From a food-based exposure perspective, infants may be exposed to styrene from consuming food items such as those listed in Table 6-3. In addition, it is possible that exposure may also result from consumption of infant formula or from nursing practices. In the 5-year FDA study on volatile organic compounds in foods (Fleming-Jones and Smith 2003), soy- and milk-based infant formula was included in the study; however, the results for styrene were not reported. Baby foods and infant formula are often stored in polystyrene containers and the migration of low levels of non polymerized styrene into food

6. POTENTIAL FOR HUMAN EXPOSURE

items from polystyrene containers has been demonstrated (EPA 1985a; Tang et al. 2000). In a study on chemicals in mother's milk, styrene was identified, but not quantified, in 8 out of 12 samples of mother's milk samples collected from mothers living in four U.S. urban areas (Pellizzari 1982). Although children are exposed to styrene from these oral routes, it has been estimated that >90% of human exposure to styrene arises due to inhalation routes (Fleming-Jones and Smith 2003; Tang et al. 2000).

Aside from food-related intake, children's exposure to styrene may differ from exposures to adults, especially during school, home, or play activities that may expose the children to styrene sources. For example, for elementary aged children (grades 2, 3, 4, and 5) attending inner city schools in Minneapolis, it was found that the lowest exposure to styrene occurred either outdoors or in school, and the highest exposure occurred at home. The latter can be substantially influenced (increased) if smoking occurs in the home. Exposures to styrene while outside, in either winter or spring, were very low (winter: $0.0 \mu\text{g}/\text{m}^3$; spring: $0.1 \mu\text{g}/\text{m}^3$), whereas exposures were much higher at school (winter: $31.3 \mu\text{g}/\text{m}^3$; spring: $39.7 \mu\text{g}/\text{m}^3$), but were almost three times higher at home (winter: $91.9 \mu\text{g}/\text{m}^3$; spring: $91.9 \mu\text{g}/\text{m}^3$) (Adgate et al. 2004). These exposures led to blood level concentrations of styrene that were generally twice as high as the general population (Sexton et al. 2005).

6.7 POPULATIONS WITH POTENTIALLY HIGH EXPOSURES

People working in various styrene industries are likely to have the highest exposures to styrene. Lower levels may be encountered near industrial facilities or hazardous waste sites emitting styrene to outdoor air. High indoor styrene concentrations in the home may be due to emissions from building materials, consumer products, tobacco smoke, photocopiers and laser printers. Smokers and those eating a high proportion of foods packaged in polystyrene may also have above average exposure to styrene, with the amounts estimated by smoking ($100 \mu\text{g}$ from 20 cigarettes) more than doubling the normal estimated exposure to styrene (Tang et al. 2000). In addition, workers with long-term employment at photocopy centers may also be exposed to high concentrations of styrene.

6.8 ADEQUACY OF THE DATABASE

Section 104(i)(5) of CERCLA, as amended, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of styrene is available. Where adequate information is not available, ATSDR, in conjunction with NTP, is required to assure the initiation of a program of research

6. POTENTIAL FOR HUMAN EXPOSURE

designed to determine the health effects (and techniques for developing methods to determine such health effects) of styrene.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that if met would reduce the uncertainties of human health assessment. This definition should not be interpreted to mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

6.8.1 Identification of Data Needs

Physical and Chemical Properties. The solubility of an organic compound in water is indicative of how that chemical will partition between water, soil, and organisms (Banerjee et al. 1980; Hassett et al. 1983; Valvani et al. 1981). Clarification of the exact solubility of styrene in water would be helpful because a range of values is currently reported (Table 4-2). The Henry's law constant and K_{oc} value for styrene need to be verified experimentally to provide more accurate predictions of air-water and soil-water partitioning.

Production, Import/Export, Use, Release, and Disposal. According to the Emergency Planning and Community Right-to-Know Act of 1986, 42 U.S.C. Section 11023, industries are required to submit substance release and off-site transfer information to the EPA. The TRI, which contains this information for 2005, became available in May of 2007. This database is updated yearly and should provide a list of industrial production facilities and emissions.

Substantial quantities of styrene are currently produced and used in the United States (Heylin 1989; HSDB 2007; SRI 1989; USITC 1988). Production and import quantities, producers, and uses are well documented, with 4.7×10^{12} g produced in 2000 (HSDB 2007), representing a slight increase in U.S. production since 1997 (3.12×10^{12} g). The United States has imported less styrene over the last several decades, with amounts decreasing more than an order of magnitude, from 1.4×10^{10} g imported in 1978 to 5.7×10^8 g being imported in 2001 (HSDB 2007). Interestingly, styrene exports increased from 3.6×10^{10} g in 1978 (representing 1% of total U.S. production) to $>1.2 \times 10^{12}$ g exported in 2001 (representing >26% of total U.S. production) (HSDB 2007).

6. POTENTIAL FOR HUMAN EXPOSURE

Quantities of styrene disposed of by various disposal methods, other than those reported to the TRI, are not known. Styrene releases into water are regulated by EPA, but styrene is not listed as a hazardous waste constituent and, therefore, land disposal restrictions do not apply to this compound. Additional information on disposal methods used for styrene and styrene-containing products and the quantities disposed of by each method would help to better characterize the potential for human exposure to this compound from disposal at waste sites or other locations.

Environmental Fate. Styrene will partition among the environmental media, with a tendency to volatilize from water to air and to adsorb to soils (EPA 1984b; Roberts et al. 1980; Sato and Nakajima 1979). However, data on styrene volatilization from water and confirmation of the estimated K_{oc} value by adsorption/desorption data would be useful to estimate more accurately the tendency of styrene to partition to air and soil. Confirmation of the K_{oc} would also provide a more reliable basis for estimating the mobility of styrene in the various types of soil.

Although the reaction mechanisms of styrene transformations in the atmosphere are fairly well understood (Atkinson et al. 1982; Bigozzi et al. 1981; EPA 1979a; Sloane and Brudzynski 1979), more information regarding the environmental fates of the transformation products would allow a more accurate prediction of the atmospheric fate of this compound. Biodegradation data are available for styrene under both aerobic and anaerobic conditions (Bridie et al. 1979; Grbic-Galic et al. 1990).

Bioavailability from Environmental Media. Styrene is known to be absorbed following inhalation, oral, and dermal contact (Dutkiewicz and Tyras 1968; Engstrom et al. 1978a, 1978b; Ramsey and Andersen 1984; Ramsey and Young 1978; Withey 1976; Withey and Collins 1979). Absorption rates via inhalation are known (Withey and Collins 1978). Additional data are needed to evaluate absorption rates following oral and dermal exposure. It is believed that absorption of styrene from the gut is believed to be generally rapid and therefore, contact with styrene contaminated food, soil, or water will probably also result in significant absorption. However, this may depend on the medium in which it is contained.

Food Chain Bioaccumulation. Bioconcentration of styrene in aquatic organisms is not likely to be significant, based on both a measured BCF for a single goldfish species (BCF=13.5; Ogata et al. 1984) and an estimated BCF (EPA 1984b; Kenaga 1980; Ogata et al. 1984). No data on biomagnification of styrene in the food chain were located. Since significant bioaccumulation is unlikely, this lack of data may not be a major limitation. No data needs are identified at this time.

6. POTENTIAL FOR HUMAN EXPOSURE

Exposure Levels in Environmental Media. Reliable monitoring data for the levels of styrene in contaminated media at hazardous waste sites are needed so that the information obtained on levels of styrene in the environment can be used in combination with the known body burden of styrene to assess the potential risk of adverse health effects in populations living in the vicinity of hazardous waste sites.

Monitoring data for styrene in environmental compartments are extensive, and historical and recent data are available for air (EPA 1987e, 1988c; Uchiyama and Hasegawa 2000), groundwater (Canter and Sabatini 1994; Squillace et al. 1999), and rivers (Gotz et al. 1998; Rathbun 2000). Additional data on styrene levels in water and soil, especially in the vicinity of hazardous waste sites, would be useful in assessing the potential for human exposure. Estimates of human intake from food, air, water, and soil have been made (EPA 1985a; Tang et al. 2000) and will undoubtedly be revised as additional data become available.

Exposure Levels in Humans. Styrene has been detected in human blood, breath, milk, and adipose tissue of the general population (Antoine et al. 1986; EPA 1986d, 1987e; Pellizzari et al. 1982) and metabolites of styrene have been detected in urine of workers exposed to styrene (Elia et al. 1980; Sollenberg et al. 1988). However, data generated by biological monitoring of populations in the vicinity of waste sites with the most sensitive methods (Section 7.1) would be useful in assessing the magnitude of human exposures from this source.

This information is necessary for assessing the need to conduct health studies on these populations.

Exposures of Children. Additional studies on the exposure of children to styrene would improve understanding of any mechanisms and pathways for styrene exposure and subsequent effects on this subpopulation. The exposure study report by Adgate et al. (2004) and the corresponding blood level study by Sexton et al. (2005) provide a good understanding for styrene exposures and uptake in one metropolitan area in the Midwestern United States, but additional reports investigating exposures and uptake, for example, in other climates, would be beneficial.

Child health data needs relating to susceptibility are discussed in Section 3.12.2, Identification of Data Needs: Children's Susceptibility.

Exposure Registries. No exposure registries for styrene were located. This substance is not currently one of the compounds for which a sub-registry has been established in the National Exposure

6. POTENTIAL FOR HUMAN EXPOSURE

Registry. The substance will be considered in the future when chemical selection is made for sub-registries to be established. The information that is amassed in the National Exposure Registry facilitates the epidemiological research needed to assess adverse health outcomes that may be related to exposure to this substance.

6.8.2 Ongoing Studies

No current ongoing studies located in the Federal Research in Progress (FEDRIP) database were related to the potential for human exposure. The Computer Retrieval of Information on Scientific Projects (CRISP 2007) National Institutes of Health database identified one study. Scott Ensign of Utah State University proposed to investigate microbial pathways of short-chain hydrocarbon metabolism and the properties of the enzymes, cofactors, and regulatory elements associated with these pathways. The work will be sponsored by the National Institutes of Health/National Institute of General Medical Sciences and will investigate the central roles of carboxylation and novel cofactors/enzymes in bacterial hydrocarbon/epoxide/ketone metabolism of these chemicals, including styrene, by pursuing the biochemical, structural, and genetic characterization of these processes (CRISP 2007).

As part of the Third National Health and Nutrition Evaluation Survey (NHANES III), the Environmental Health Laboratory Sciences Division of the National Center for Environmental Health, Centers for Disease Control and Prevention, will be analyzing human blood samples for styrene and other volatile organic compounds. These data will give an indication of the frequency of occurrence and background levels of these compounds in the general population.

6. POTENTIAL FOR HUMAN EXPOSURE

This page is intentionally blank.

7. ANALYTICAL METHODS

The purpose of this chapter is to describe the analytical methods that are available for detecting, measuring, and/or monitoring styrene, its metabolites, and other biomarkers of exposure and effect to styrene. The intent is not to provide an exhaustive list of analytical methods. Rather, the intention is to identify well-established methods that are used as the standard methods of analysis. Many of the analytical methods used for environmental samples are the methods approved by federal agencies and organizations such as EPA and the National Institute for Occupational Safety and Health (NIOSH). Other methods presented in this chapter are those that are approved by groups such as the Association of Official Analytical Chemists (AOAC) and the American Public Health Association (APHA). Additionally, analytical methods are included that modify previously used methods to obtain lower detection limits and/or to improve accuracy and precision.

As a volatile material, styrene is readily determined by gas chromatographic (GC) analysis. As a hydrocarbon, styrene is detected very sensitively by flame ionization detection (FID); its aromatic nature enables some selectivity by photoionization detection (PID); and it can be specifically identified by mass spectrometry (MS). Styrene is usually collected from the gas phase or from vapor evolved from the sample matrix on a column of solid sorbent, such as Tenax®. Cryogenic (low temperature) collection and sorption in organic liquids are also possible.

Capillary gas chromatography, also known broadly as high-resolution gas chromatography (HRGC), has greatly facilitated the analysis of compounds such as styrene that can be measured by gas chromatography and has resulted in vast improvements in resolution and sensitivity. It has made the choice of a stationary phase less important than is the case with the use of packed columns. The instrumental capability to separate volatile analytes by HRGC is, for the most part, no longer the limiting factor in their analysis.

The specific analytical methods used to quantify styrene in biological and environmental media samples are summarized below.

7.1 BIOLOGICAL MATERIALS

Methods have been described for the determination of styrene in expired air (Kneip and Crable 1988a; Stewart et al. 1968; Wallace and Pellizzari 1995; Wallace et al. 1996), blood (Antoine et al. 1986; Ashley et al. 1992; Bartolucci et al. 1986; Guillemain and Berode 1988; Withey and Collins 1977), urine (Dolara et al. 1984; Ghittori et al. 1987; Pezzagno et al. 1985), adipose tissue (Engstrom et al. 1978a), and other

7. ANALYTICAL METHODS

tissues (heart, lungs, liver, spleen, kidney, brain) (Withey and Collins 1977) (Table 7-1). These methods generally require styrene release from the sample matrix and collection on a column of solid sorbent or collection as headspace gas; more recent methods permit the exhaled breath to be delivered directly, via a breath interface, to the mass spectrometer (Wallace et al. 1996). Cryogenic collection is also possible (Romieu et al. 1999). Of the available methods for detecting styrene, FID is the most sensitive, and MS is the most specific, the latter of which can be made more sensitive by tandem MS approaches (MS/MS) (Wallace et al. 1996). Levels of detection are in the low parts per billion range for breath samples (Stewart et al. 1968; Wallace et al. 1996) and parts per trillion range for blood samples (Ashley et al. 1992).

The major metabolites of styrene in humans are mandelic acid (MA) and phenylglyoxylic acid (PGA). Detection of these metabolites in urine is the most commonly performed procedure as an indicator of exposure to styrene. Procedures have been described for their measurement in urine (Baselt 1988a; Dolara et al. 1984; Engstrom et al. 1976; Kneip and Crable 1988b, 1988c; Korn et al. 1984; Pezzagno et al. 1985; Sedivec et al. 1984; Sollenberg et al. 1988). Generally, these styrene metabolites are converted to volatile derivatives and measured gas chromatographically or determined directly by high performance liquid chromatography (HPLC). Two other styrene metabolites that may result from exposure to styrene are 4-vinylphenol (Pfaffli et al. 1981) and styrene glycol (phenyl ethylene glycol) (Guillemin and Berode 1988), but methods for the detection of these metabolites in biological materials have not been worked out in detail. Sensitive methods are also available for measuring styrene oxide in blood (Kessler et al. 1990; Langvardt and Nolan 1991), although these techniques are probably more useful in research on styrene toxicity than in detecting or quantifying styrene exposure.

Methods for detection of styrene and its metabolites in biological materials are summarized in Table 7-1.

7.2 ENVIRONMENTAL SAMPLES

Styrene determined in environmental samples is usually collected on solid sorbents (from air; Zielinska et al. 1996) or on solid sorbents after purging in a gas stream (water, soil, solid waste samples; Miermans et al. 2000). Styrene from such samples is measured very sensitively by GC/FID and very specifically by GC/MS. Methods for the determination of styrene in environmental samples have been standardized by the American Society for Testing and Materials (ASTM 1988a, 1988b), EPA (1986a, 1986b, 1989c, 1989d, 2003), and NIOSH (1984). Relatively low detection limits can be achieved for the determination of styrene in environmental samples and the accuracy appears to be acceptable for those limited cases in

7. ANALYTICAL METHODS

Table 7-1. Analytical Methods for Determining Styrene in Biological Materials

Sample matrix	Preparation method	Analytical method	Sample detection limit	Percent recovery	Reference
Styrene analyte					
Adipose tissue	Evaporation into nitrogen, collection as vapor	GC	No data	No data	Engstrom et al. 1978a
Breath ^a	Collection in Saran bag	GC/FID	0.05 ppm	No data	Stewart et al. 1968
Breath	Sorption onto silicagel, desorption into headspace	GC	0.1 ppm	No data	Kneip and Crable 1988a
Breath	Collection in Tedlar bag, followed by sorption of VOCs (including styrene) onto Tenax	GC/MS	Low ppb	No data	Wallace et al. 1996
Breath	Collection onto personal monitoring badge, followed by solvent extraction	GC/FID	0.9 ppb ^b	No data	Romieu et al. 1999
Blood	Purge at 40–50 °C with helium, collection on Tenax-GC/silica	GC/MS	No data	CV<5%	Antoine et al. 1986
Blood	Headspace analysis	GC/FID	No data	No data	Bartolucci et al. 1986
Blood	Collection in vacutainer with EDTA as anticoagulant, headspace analysis	GC	No data	No data	Guillemin and Berode 1988
Blood	Headspace analysis	GC	0.02 µg/mL	No data	Withey and Collins 1977
Blood	Vacutainer collection, followed by purge and trap	GC/MS	9 ppt (95% CI: 4–21 ppt)	95–121%	Ashley et al. 1992
Heart, lungs, liver, spleen, kidney, brain	Hemogenate prepared for headspace analysis	GC	0.01 µg/g	No data	Withey and Collins 1977
Urine	Headspace from sample maintained and 37 °C for 2 hours	GC/MS	No data	No data	Ghitori et al. 1987
Urine	Sorption on XAD-2, elution with n-hexane	HPLC/UV	<0.7 µg/L	72±10%	Dolara et al. 1984
Urine	Headspace analysis	GS/MS	No data	No data	Pezzagno et al. 1985
Styrene metabolite analyte					
Urine for MA	Extraction with ethyl acetate, derivatization to isopropyl ester	GC/FID	No data	No data	Korn et al. 1984

7. ANALYTICAL METHODS

Table 7-1. Analytical Methods for Determining Styrene in Biological Materials

Sample matrix	Preparation method	Analytical method	Sample detection limit	Percent recovery	Reference
Urine for MA	Extraction with diethyl ether, silylation	GC	No data	No data	Engstrom et al. 1976
Urine for MA and PGA metabolites	Extraction and derivatization	GC/FID	0.05 ppm	94% MA, 98% PGA	Bartolucci et al. 1986
Urine for MA metabolite	Acidification, extraction, derivatization	HRGC/FID	10 mg/L	No data	Kneip and Crable 1988b
Urine for PGA metabolite	Reduction, acidification, extraction, derivatization	HRGC/FID	10 mg/L	No data	Kneip and Crable 1988c
Urine for MA and PGA metabolites	Extraction with ethyl acetate, derivatization to methyl esters with diazomethane	GC	10 mg/L	No data	Sedivec et al. 1984
Urine for MA and PGA metabolites	Extraction with ethyl acetate, evaporation derivatization	GC/FID	No data	97–99% relative recovery	Baselt 1988a
Urine for MA, PGA, and hippuric acid metabolites	Direct injection	HPLC/UV	<1 µg/mL ^c	<3% deviation from true value at 5 µg/mL	Regnaud et al. 1987
Urine for MA and PGA (stereo-selective)	Extraction and derivatization	HRGC/FID	No data	No data	Korn et al. 1987
Blood for styrene oxide	Extraction with n-hexanone, concentration by evaporation	GC/FID	1 ng/mL	72±8%	Kessler et al. 1990
Blood for styrene oxide	Extraction with benzene	GC/MS	10 ng/g	92±21%	Langvardt and Nolan 1991

^aUnless otherwise designated, analyses are for styrene.

^bValues were reported for benzene; authors indicated that other VOCs had similar levels of detection

^cDetection limits were 0.63 µg/mL for mandelic acid, 0.78 µg/mL for phenylglyoxylic acid, and 0.52 µg/mL for hippuric acid.

CV = coefficient of variation; CI = confidence interval; EDTA = ethylene diamine tetra acetic acid; FID = flame ionization detector; GC = gas chromatography; HPLC = high-performance liquid chromatography; HRGC = high-resolution gas chromatography; MA = mandelic acid; MS = mass spectrometry; PGA = phenylglyoxylic acid; UV = ultraviolet; VOC = volatile organic compound

7. ANALYTICAL METHODS

which accuracy data are available. For example, the most sensitive for styrene detection limits were 0.002 µg/L in water, 4 µg/kg in soil, and 500 µg/kg in solid waste. No significant reports were found pertaining to styrene degradation products in environmental samples. Information on methods for the determination of styrene in environmental samples is summarized in Table 7-2.

7.3 ADEQUACY OF THE DATABASE

Section 104(i)(5) of CERCLA, as amended, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of styrene is available. Where adequate information is not available, ATSDR, in conjunction with NTP, is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of styrene.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that if met would reduce the uncertainties of human health assessment. This definition should not be interpreted to mean that all data needs discussed in this section must be filled. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

7.3.1 Identification of Data Needs

Methods for Determining Biomarkers of Exposure and Effect. Styrene and its primary metabolites, MA and PGA, can be detected in several human tissues (blood, urine, adipose tissue, and in several organs [Table 7-1]), as well as in exhaled breath. The detection limits range in the parts per billion to parts per trillion range (Ashley et al. 1992). Approaches have been developed to provide more efficient and rapid assessment of exposure, such as where the subject breathes for short periods of time into different types of sample collectors (tedlar bags, evacuated canisters, even directly into a mass spectrometer interface [Wallace et al. 1996]), allowing samples to be collected efficiently for subsequent or immediate analysis. Personal monitoring badges containing charcoal have also been developed for longer-term assessment of exposure to styrene and other volatiles, which have been calibrated for comparison to levels detected in the subjects' blood (Romieu et al. 1999). There are no data needs at this time.

7. ANALYTICAL METHODS

Table 7-2. Analytical Methods for Determining Styrene in Environmental Samples

Sample matrix	Preparation method	Analytical method	Sample detection limit	Percent recovery	Reference
Food	Homogenization, headspace sampling	GC/MS	1 µg/kg	No data	Gilbert and Startin 1983
Air	Retention by activated carbon	GC ^a	No data	No data	ASTM 1988a
Air	Retention by activated carbon, elution with carbon disulfide	GC ^b	No data	No data	ASTM 1988b
Air	Retention by activated carbon, elution with carbon disulfide	HRGC/FID	0.01 mg/sample	No data	NIOSH 1984
Wastewater	Stable isotope dilution method, followed by solvent extraction	GC/MS	10 µg/L	No data	EPA 2001
Water	Purge by helium, collection on activated charcoal/silica gel/Tenax®	GC/PID	0.01 µg/L	96–104%	EPA 1989f
Water	Purge by helium, collection on activated charcoal/silica gel/Tenax®	GC/PID	0.008 µg/L	No data	EPA 1989g
Water	Purge by helium, collection on activated charcoal/silica gel/Tenax®	HRGC/MS	0.20 µg/L	120% (at 1 µg/L)	EPA 1989h
Water	Purge by helium, collection on activated charcoal/silica gel/Tenax®	HRGC/MS	0.04 µg/L	102%	EPA 1989i
Water	Purge and trap with nitrogen or helium	GC/PID	0.01 µg/L	104%	EPA 1995
Water	Purge and trap	GC/PID	0.01 µg/L	104%	EPA 1996
Water	Purge by helium, and trap at low temperature	GC/FID	0.002 µg/L	No data	Miermans et al. 2000
Water	Purge and trap on VOCARB or equivalent	GC/MS	0.039 µg/L	94–110%	USGS 1998
Soil, low level	Purge by helium, collection on solid, thermal desorption	GC/MS	4 µg/kg	No data	EPA 1986c
Solid waste, nonwater miscible	Purge by helium, collection on solid, thermal desorption	GC/MS	500 µg/kg ^c	No data	EPA 1986c

7. ANALYTICAL METHODS

Table 7-2. Analytical Methods for Determining Styrene in Environmental Samples

Sample matrix	Preparation method	Analytical method	Sample detection limit	Percent recovery	Reference
Solid waste	Purge by helium, collection on solid, thermal desorption	GC/MS	500 µg/kg ^c	No data	EPA 1986b

^aAbsorption characteristics for sampling atmospheric vapor with activated carbon for subsequent analysis by GC.

^bGeneric method for the determination of organics.

^cEstimation from detection limits in water.

FID = flame ionization detector; GC = gas chromatography; HRGC = high-resolution gas chromatography; MS = mass spectrometry; PID = photoionization detection; UV = ultraviolet

7. ANALYTICAL METHODS

Methods for Determining Biomarkers of Exposure and Effect. Biological monitoring of styrene exposure has been reviewed (Guillemin and Berode 1988). Sensitive and selective methods are available for the qualitative and quantitative measurement of styrene and its two major metabolites, MA and PGA, in samples from exposed individuals after the analytes are separated from their biological sample matrix. The concentration of these metabolites in urine has been found to correlate with average exposure levels in air (Härkönen et al. 1978), and so may be used as a biomarker of exposure. However, measurements of MA and PGA are not specific for this purpose (Bartolucci et al. 1986) and these metabolites can result from the metabolism of other organic substances, particularly ethylbenzene (Baselt 1988b). As noted in Chapter 6 for studies of the general population, styrene has been identified in adipose tissue at concentrations of 8–350 ng/g (EPA 1986d), in blood at a mean concentration of 0.4 µg/L (Antoine et al. 1986) and in exhaled breath at mean concentrations of 0.7–1.6 µg/L. Levels of MA and PGA in biological samples from the general population probably are below the detection limits of methods that are currently used (Baselt 1988a). However, it is likely that normal background levels of these metabolites in unexposed individuals are too low to be of any significance. Although new and improved methods for the determination of styrene and its metabolites in biological samples need not have a high priority, additional work on standardization of these methods for use in biological samples accompanied by additional studies involving interlaboratory comparisons of recovery, accuracy, and precision data would be useful.

Clinical means have been proposed to indicate exposure to styrene. In general, these are not sufficiently sensitive, specific, or well characterized. The most common symptom of exposure, impairment of central nervous system function, is not unique to styrene. Neither cytogenetic monitoring of peripheral lymphocytes nor unscheduled DNA synthesis have been sufficiently well characterized as biomarkers of exposure to styrene.

There is currently some information that can be used to correlate levels of biomarkers of exposure to styrene in biological media with adverse health effects. Central nervous system depression has been correlated with a urinary MA concentration of ≥ 800 mg/L and a decrement in psychomotor performance in association with a concentration of $\geq 1,200$ mg/L (Härkönen et al. 1978). The styrene concentrations in air producing these effects and urinary MA levels were relatively high. Studies to determine if effects at lower levels of exposure could also be correlated to metabolite levels in urine would be valuable. However, the design of studies involving controlled inhalation exposures in humans is precluded by the potential carcinogenicity of styrene.

7. ANALYTICAL METHODS

Methods for Determining Parent Compounds and Degradation Products in Environmental

Media. In an occupational setting, the medium that is of most concern for human exposure to styrene is air, although at Superfund sites, contaminated groundwater may pose a danger. Methods are well developed for the determination of styrene in water and air with excellent selectivity and sensitivity (ASTM 1988a, 1988b; EPA 1989f, 1989g, 1989i, 1995, 1996; NIOSH 1984; USGS 1998). Methods for the determination of styrene in soil and waste samples are not as well developed and may require additional testing and validation (EPA 1986b, 1986c, 2001).

The detection limits for styrene in environmental media cited in Table 7-2 (0.01 mg/sample, typically 10 L, NIOSH 1984; 0.002 µg/L in water, Miermans et al. 2000; and 4 µg/kg in soil, EPA 1986c) are low enough to enable the determination of styrene in any environmental medium likely to pose a hazard to health based upon information currently available in the literature. These detection limits are probably below most ambient background levels of styrene.

Sampling methodologies for compounds such as styrene pose typical collection problems that include the collection of samples that are nonrepresentative, may be of insufficient sample volume, may contain interfering materials that result in low sample recovery, or may contain interfering contaminating chemicals. Other sampling methods may be labor-intensive, or require tedious extraction and purification procedures (Green and Le Pape 1987; Miermans et al. 2000). Methods that measure organic compounds such as styrene *in situ* in water and other environmental media without the need for sampling and extraction procedures to isolate the analyte prior to analysis are desirable. One such method has been patented, but no commercial products have been identified (see below).

In regard to methods for determining parent styrene and degradation products in environmental media, the following conclusions may be drawn: Because styrene can be detected instrumentally and determined in air and normal water samples with totally adequate selectivity and sensitivity, no additional data are needed at this time. A moderate need exists to improve methodologies to determine styrene in soil, sludges, and solid wastes. Styrene degradation products are a different matter in that little information is available on their determination in environmental samples. In air, these compounds should consist predominantly of photochemical oxidation products, whereas in water and soil samples, they are expected to be biodegradation products. Additional research is needed on the determination of these materials.

7. ANALYTICAL METHODS

7.3.2 Ongoing Studies

No current ongoing studies were found in the Federal Research in Progress database (FEDRIP 2007) regarding the development of analytical methods for the detection of styrene or its metabolites.

The Environmental Health Laboratory Sciences Division of the National Center for Environmental Health, Centers for Disease Control and Prevention, is developing methods for the analysis of styrene and other volatile organic compounds in blood. These methods use purge and trap methodology, high-resolution gas chromatography, and magnetic sector mass spectrometry, which give detection limits in the low parts per trillion (ppt) range.

In addition to the detection and identification of styrene using chromatographic means, where current detection limits approach analytical thresholds, other technologies may offer additional routes for detection of styrene in samples, especially for aquatic environmental samples. Immunoassay-based methods of analysis offer rapid analysis times, high sensitivity, and often high selectivity for various organic pollutants and other chemicals of concern. While styrene and its metabolites are considered important chemicals to monitor using such techniques, no known assays have been commercialized, even though one patent has been issued for a method that claims to detect styrene and other VOCs by immunoassay (patent 5,358,851, issued in 1994 [Peck 1994]). No additional information was found regarding the continued development (or use or application) of this technology.

8. REGULATIONS AND ADVISORIES

The international and national regulations and guidelines regarding styrene in air, water, and other media are summarized in Table 8-1.

ATSDR has derived an acute-duration inhalation MRL of 2 ppm for styrene. This MRL is based on a NOAEL of 20 ppm for alterations in tests of reaction time and attention in subjects exposed to styrene for 3 hours (Seeber et al. 2004) and an uncertainty factor of 10 to account for human variability.

ATSDR has derived a chronic-duration inhalation MRL of 0.2 ppm for styrene. This MRL is based on a LOAEL of 20 ppm estimated from a meta-analysis of data (Benignus et al. 2005) from occupational exposure studies demonstrating alterations in choice reaction time and color discrimination in styrene workers and an uncertainty factor of 100 (10 for use of a LOAEL and 10 for human variability).

EPA (IRIS 2007) has derived an inhalation reference concentration (RfC) of 1 mg/m³ (0.2 ppm) based on a NOAEL_{HEC} value of 34 mg/m³ in a cross-sectional study in workers finding central nervous system effects in workers exposed to >22 ppm (Mutti et al. 1984a) and an uncertainty factor of 30 (3 for the use less than chronic study, 3 for human variability, and 3 for database inadequacies).

ATSDR has derived an acute-duration oral MRL of 0.1 mg/kg/day for styrene. This MRL is based on a LOAEL of 100 mg/kg/day for impaired learning in rats exposed to styrene for 14 days (Husain et al. 1985) and an uncertainty factor of 1,000 (10 for use of a LOAEL, 10 for extrapolation from rats to humans, and 10 for human variability).

EPA (IRIS 2007) has established an oral reference dose (RfD) of 0.2 mg/kg/day based on a NOAEL of 200 mg/kg/day from a subchronic oral toxicity study in beagle dogs in which red blood cell and liver effects was observed (Quast et al. 1979). The uncertainty factor used in this assessment was 1,000 (10 for interspecies extrapolation, 10 for human variability variation, and 10 for extrapolation of subchronic effects to chronic effects).

The International Agency for Research on Cancer (IARC) has classified styrene as a Group 2B carcinogen (possibly carcinogenic to humans) (IARC 2006). The Department of Health and Human Services and the EPA has not classified styrene for human carcinogenicity (IRIS 2007; NTP 2005).

Table 8-1. Regulations and Guidelines Applicable to Styrene

Agency	Description	Information	Reference
<u>INTERNATIONAL</u>			
Guidelines:			
IARC	Carcinogenicity classification	Group 2B ^a	IARC 2006
WHO	Air quality guidelines		WHO 2000
	TWA based on effects other than cancer or odor/annoyance using an averaging time of 1 week	0.26 mg/m ³	
	Based on sensory effects or annoyance reactions, using an averaging time of 30 minutes		
	Detection threshold	0.07 mg/m ³	
	Recognition threshold	0.21–0.28 mg/m ³	
	Guideline value	0.07 mg/m ³	
	Drinking water quality guidelines	0.02 mg/L ^b	WHO 2004
<u>NATIONAL</u>			
Regulations and Guidelines:			
a. Air			
ACGIH	TLV (8-hour TWA)	85 mg/m ³	ACGIH 2006
	STEL (15-minute TWA)	170 mg/m ³	
AIHA	ERPG-1 ^c	213 mg/m ³	AIHA 1995
	ERPG-2 ^c	1,065 mg/m ³	
	ERPG-3 ^c	4,260 mg/m ³	
EPA	AEGL-1 ^d		EPA 2007a
	10 minutes	85 mg/m ³	
	30 minutes	85 mg/m ³	
	60 minutes	85 mg/m ³	
	4 hours	85 mg/m ³	
	8 hours	85 mg/m ³	
	AEGL-2 ^d		
	10 minutes	980 mg/m ³	
	30 minutes	682 mg/m ³	
	60 minutes	554 mg/m ³	
4 hours	554 mg/m ³		
8 hours	554 mg/m ³		

Table 8-1. Regulations and Guidelines Applicable to Styrene

Agency	Description	Information	Reference
NATIONAL (cont.)			
EPA	AEGL-3 ^d		EPA 2007a
	10 minutes ^e	8,094 mg/m ³ (1,862 ppm)	
	30 minutes ^e	8,094 mg/m ³ (1,862 ppm)	
	60 minutes ^e	4,686 mg/m ³ (1,078 ppm)	
	4 hours	1,448 mg/m ³ (333 ppm)	
	8 hours	1,448 mg/m ³ (333 ppm)	
	Level of distinct odor awareness	2.3 mg/m ³ (0.5 ppm)	
EPA	Hazardous air pollutant	Yes	EPA 2007c 42 USC 7412
NIOSH	REL (10-hour TWA)	50 ppm	NIOSH 2005
	STEL (15-minute TWA)	100 ppm	
	IDLH	2,982 mg/m ³ (686 ppm)	
OSHA	PEL (8-hour TWA) for general industry	100 ppm	OSHA 2006c 29 CFR 1910.1000, Table Z-2
	Acceptable ceiling concentration	200 ppm	
	Acceptable maximum peak above the acceptable ceiling concentration for an 8-hour shift for a maximum duration of 5 minutes in any 3 hours	600 ppm	
	PEL (8-hour TWA) for shipyard industry	100 ppm	
	PEL (8-hour TWA) for construction industry	100 ppm	
b. Water			
EPA	Designated as hazardous substances in accordance with Section 311(b)(2)(A) of the Clean Water Act	Yes	EPA 2007b 40 CFR 116.4

Table 8-1. Regulations and Guidelines Applicable to Styrene

Agency	Description	Information	Reference
NATIONAL (cont.)			
EPA	Drinking water standards and health advisories		EPA 2006
	1-day health advisory for a 10-kg child	20 mg/L	
	10-day health advisory for a 10-kg child	2 mg/L	
	DWEL	7 mg/L	
	Lifetime	0.1 mg/L	
	10 ⁻⁴ Cancer risk	No data	
	Master Testing List	Yes ^f	EPA 2007d
	National primary drinking water standards		EPA 2003
	MCLG	0.1 mg/L	
	MCL	0.1 mg/L	
	Public health goal	0.1 mg/L	
	Reportable quantities of hazardous substances designated pursuant to Section 311 of the Clean Water Act	1,000 pounds	EPA 2007e 40 CFR 117.3
c. Food			
FDA	Bottled drinking water	0.1 mg/L	FDA 2006a 21 CFR 165.110
	Food additives permitted for direct addition to food for human consumption	Yes	FDA 2007b 21 CFR 172
	Indirect food additives: adhesives and components of coatings	Yes	FDA 2006b 21 CFR 175
	EAFUS	Yes ^h	FDA 2007a
d. Other			
ACGIH	Carcinogenicity classification	A4 ⁱ	ACGIH 2006
	Biological exposure indices (end of shift at end of workweek)		
	Sum of mandelic acid and phenyl glyoxylic acid in urine	400 mg/g creatinine	
	Styrene in venous blood	0.2 mg/L	
EPA	Carcinogenicity classification	No data	IRIS 2007
	RfC	1 mg/m ³ (0,2 ppm)	
	RfD	0.2 mg/kg/day	
	Superfund, emergency planning, and community right-to-know		
	Designated CERCLA hazardous substance	Yes ^j	EPA 2007f 40 CFR 302.4
	Reportable quantity	1,000 pounds	

Table 8-1. Regulations and Guidelines Applicable to Styrene

Agency	Description	Information	Reference
NATIONAL (cont.)			
EPA	Effective date of toxic chemical release reporting	01/01/2000	EPA 2007g 40 CFR 372.65
NTP	Carcinogenicity classification	No data	NTP 2005

^aGroup 2B: possibly carcinogenic to humans

^bConcentrations of the substance at or below the health-based guideline value may affect the appearance, taste, or odor of the water, leading to consumer complaints (WHO 2004).

^cERPG-1 is the maximum airborne concentration below which nearly all individuals could be exposed for up to 1 hour without experiencing other than mild, transient health effects; ERPG-2 is the maximum airborne concentration below which nearly all individuals could be exposed for up to 1 hour without experiencing irreversible or other serious adverse effects; and ERPG-3 is the maximum airborne concentration below which nearly all individuals could be exposed for up to 1 hour without life-threatening health effects (AIHA 1995).

^dAEGL-1 is the airborne concentration of a substance above which it is predicted that the general population, including susceptible individuals, could experience notable discomfort, irritation, or certain asymptomatic nonsensory effects, however, the effects are not disabling and are transient and reversible upon cessation of exposure; AEGL-2 is the airborne concentration of a substance above which it is predicted that the general population, including susceptible individuals, could experience irreversible or other serious, long-lasting adverse health effects or an impaired ability to escape; and AEGL-3 is the airborne concentration of a substance above which it is predicted that the general population, including susceptible individuals, could experience life-threatening health effects or death (EPA 2007a).

^eSafety considerations against the hazard of explosion must be taken into account.

^fStyrene was recommended to the MTL by the U.S. EPA's Office of Pollution Prevention and Toxics on the basis of the SIDS. Styrene was added to the MTL in 1993 and the chemical testing program is currently underway by way of a VTA. The testing needs include health effects, environmental effects, and environmental fate and exposure.

^gPotential health effects from exposure above the MCL include liver, kidney, or circulatory system problems. Common sources of contaminant in drinking water include discharges from rubber and plastic factories and leaching from landfills (EPA 2003).

^hThe EAFUS list of substances contains ingredients added directly to food that FDA has either approved as food additives or listed or affirmed as GRAS.

ⁱA4: not classifiable as a human carcinogen

^jDesignated CERCLA hazardous substance pursuant to Section 311(b)(2) of the Clean Water Act and Section 112 of the Clean Air Act.

ACGIH = American Conference of Governmental Industrial Hygienists; AEGL = acute exposure guideline levels; AIHA = American Industrial Hygiene Association; CERCLA = Comprehensive Environmental Response, Compensation, and Liability Act; CFR = Code of Federal Regulations; DWEL = drinking water equivalent level; EAFUS = Everything Added to Food in the United States; EPA = Environmental Protection Agency; ERPG = emergency response planning guidelines; FDA = Food and Drug Administration; GRAS = Generally Recognized As Safe; IARC = International Agency for Research on Cancer; IDLH = immediately dangerous to life or health; IRIS = Integrated Risk Information System; MCL = maximum contaminant level; MCLG = maximum contaminant level goal; MTL = Master Testing List; NIOSH = National Institute for Occupational Safety and Health; NTP = National Toxicology Program; OSHA = Occupational Safety and Health Administration; PEL = permissible exposure limit; REL = recommended exposure limit; RfC = inhalation reference concentration; RfD = oral reference dose; SIDS = Screening Information Data Sets; STEL = short-term exposure limit; TLV = threshold limit values; TWA = time-weighted average; USC = United States Code; VTA = Voluntary Testing Agreement; WHO = World Health Organization

This page is intentionally blank.

9. REFERENCES

- ACGIH. 1988. Documentation of the threshold limit values and biological exposure indices. 5th ed. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- ACGIH. 1988-1989. Threshold limit values and biological exposure indices for 1988-1989. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- *ACGIH. 2006. Styrene. Threshold limit values for chemical substances and physical agents and biological exposure indices. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- *Adgate JL, Church TR, Ryan AD, et al. 2004. Outdoor, indoor, and personal exposure to VOCs in children. *Environ Health Perspect* 112:1386-1392.
- *Adinolfi M. 1985. The development of the human blood-CSF-brain barrier. *Dev Med Child Neurol* 27(4):532-537.
- *Adlercreutz H. 1995. Phytoestrogens: Epidemiology and a possible role in cancer protection. *Environ Health Perspect Suppl* 103(7):103-112.
- *Agency for Toxic Substances and Disease Registry. 1989. Decision guide for identifying substance-specific data needs related to toxicological profiles; Notice. Agency for Toxic Substances and Disease Registry, Division of Toxicology. *Fed Regist* 54(174):37618-37634.
- Agency for Toxic Substances and Disease Registry. 1990a. Biomarkers of organ damage or dysfunction for the renal, hepatobiliary, and immune systems. Subcommittee on Biomarkers of Organ Damage and Dysfunction. Atlanta, GA: Agency for Toxic Substances and Disease Registry.
- Agency for Toxic Substances and Disease Registry. 1990b. Toxicological profile for manganese and compounds. Atlanta, GA: Agency for Toxic Substances and Disease Registry.
- +*Agrawal AK, Srivastava SP, Seth PK. 1982. Effect of styrene on dopamine receptors. *Bull Environ Contam Toxicol* 39:400-403.
- +*Ahlborg G, Bjerkedal T, Egenaes J. 1987. Delivery outcome among women employed in the plastics industry in Sweden and Norway. *Am J Ind Med* 12:507-517.
- *AIHA. 1995. Styrene. Emergency Response Planning Guidelines (ERPG). Fairfax, VA: American Industrial Hygiene Association.
- Aitio A. 1993. Letters to the editor and authors' reply Re: "Urinary styrene in the biological monitoring of styrene exposure" by Gobba et al. *Scand J Work Environ Health* 19:432-435.
- +*Alarie Y. 1973. Sensory irritation of the upper airways by airborne chemicals. *Toxicol Appl Pharmacol* 24:279-297.

*Cited in text

+Cited in supplemental document

9. REFERENCES

- *Ali SF, Hong JS, Wilson WE, et al. 1983. Effect of acrylamide on neurotransmitter metabolism and neuropeptide levels in several brain regions and upon circulating hormones. *Arch Toxicol* 52:35-43.
- +*Aliberti LM, Severini G. 1987. Urinary enzyme excretion in subjects exposed to styrene. *Ann Clin Biochem* 24:114.
- *Altman PL, Dittmer DS. 1974. *Biological handbooks: Biology data book*. Vol. III. 2nd ed. Bethesda, MD: Federation of American Societies for Experimental Biology, 1987-2008, 2041.
- *Amoore JE, Hautala E. 1983. Odor as an aid to chemical safety: Odor thresholds compared with threshold limit values and volatilities for 214 industrial chemicals in air and water dilution. *J Appl Toxicol* 3:272-290.
- *Andersen ME, Krishnan K. 1994. Relating *in vitro* to *in vivo* exposures with physiologically based tissue dosimetry and tissue response models. In: Salem H, ed. *Animal test alternatives: Refinement, reduction, replacement*. New York, NY: Marcel Dekker, Inc., 9-25.
- *Andersen ME, Clewell HJ, Gargas ML, et al. 1987a. Physiologically based pharmacokinetics and the risk assessment process for methylene chloride. *Toxicol Appl Pharmacol* 87(2):185-205.
- Andersen ME, Gargas ML, Ramsey JC. 1984. Inhalation pharmacokinetics: Evaluating systemic extraction, total *in vivo* metabolism, and the time course of enzyme induction for inhaled styrene in rats based on arterial blood: Inhaled air concentration ratios. *Toxicol Appl Pharmacol* 73:176-187.
- Andersen ME, MacNaughton MG, Clewell HJ, et al. 1987b. Adjusting exposure limits for long and short exposure periods using a physiological pharmacokinetics model. *Am Ind Hyg Assoc J* 48:335-343.
- Anderson KE. 1986. Styrene exposure during the manufacturing of reinforced fiberglass pipe. *Diss Abstr Int* 47:979-B.
- *Andersson HC, Tranberg EA, Uggla AH, et al. 1980. Chromosomal aberrations and sister-chromatid exchanges in lymphocytes of men occupationally exposed to styrene in a plastic-boat factory. *Mutat Res* 73:387-401.
- *Antilla A, Pukkala E, Riala R, et al. 1998. Cancer incidence among Finnish workers exposed to aromatic hydrocarbons. *Int Arch Occup Environ Health* 71:187-193.
- *Antoine SR, DeLeon IR, O'Dell-Smith RM. 1986. Environmentally significant volatile organic pollutants in human blood. *Bull Environ Contam Toxicol* 36:364-371.
- *Anwar WA, Shamy MY. 1995. Chromosomal aberrations and micronuclei in reinforced plastics workers exposed to styrene. *Mutat Res* 327:41-47.
- *Apostoli P, Brugnone F, Perbellini L, et al. 1983. Occupational styrene exposure: Environmental and biological monitoring. *Am J Ind Med* 4:741-754.
- *Arfini G, Mutti A, Vescovi P, et al. 1987. Impaired dopaminergic modulation of pituitary secretion in workers occupationally exposed to styrene: Further evidence from PRL response to TRH stimulation. *J Occup Med* 29:826-830.

9. REFERENCES

- Aringer L, Lof A, Elinder CG. 1991. The applicability of the measurement of urinary thioethers. A study of humans exposed to styrene during diet standardization. *Int Arch Occup Environ Health* 63:341-346.
- +*Arnedo-Pena A, Bellido-Blasco J, Villamarin-Vazquez JL, et al. 2003. Acute health effects after accidental exposure to styrene from drinking water in Spain. *Environ Health* 2(6):1-9.
- *Artuso M, Angotzi G, Bonassi S, et al. 1995. Cytogenetic biomonitoring of styrene-exposed plastic boat builders. *Arch Environ Contam Toxicol* 29:270-274.
- Artuso M, DeFerrari M, Bonatti S, et al. 1992. Cytogenetic biomonitoring of workers exposed to styrene. *Mutat Res* 271(2):199.
- Ashby J, Richardson CR. 1985. Tabulation and assessment of 113 human surveillance cytogenetic studies conducted between 1965 and 1984. *Mutat Res* 154:111-113.
- *Ashley DL, Bonin MA, Cardinali FL, et al. 1992. Determining volatile organic compounds in human blood from a large sample population by using purge and trap gas chromatography/mass spectrometry. *Anal Chem* 64:1021-1029.
- Ashley DL, Bonin MA, Cardinali FL, et al. 1994. Blood concentrations of volatile organic compounds in a nonoccupationally exposed U.S. population and in groups with suspected exposure. *Clin Chem* 40(7):1401-1404.
- Ashley DL, Bonin MA, Cardinali FL, et al. 1996. Measurement of volatile organic compounds in human blood. *Environ Health Perspect* 104:871-877.
- *Assmuth T, Kalevi K. 1992. Concentrations and toxicological significance of trace organic compounds in municipal solid waste landfill gas. *Chemosphere* 24(9):1207-1216.
- *ASTM. 1988a. D 3686-84. Standard practice for sampling atmospheres to collect organic compound vapors. (Activated charcoal tube adsorption method). In: Storer RA, Cornillot JL, Richardson D, et al., eds. 1988 Annual book of ASTM standards. Vol. 11.03. Section 11: Water and environmental technology. Philadelphia, PA: American Society for Testing and Materials, 234-240.
- *ASTM. 1988b. D3687-84. Standard practice for analysis of organic compound vapors collected by the activated charcoal tube absorption method. In: Storer RA, Cornillot JL, Richardson D, et al., eds. 1988 Annual book of ASTM standards. Vol. 11.03. Section 11. Water and environmental technology. Philadelphia, PA: American Society for Testing and Materials, 241-246.
- Atkinson R. 1994. Gas-phase tropospheric chemistry of organic compounds. *J Phys Chem Ref Data* 1:231.
- *Atkinson R, Arey J. 2003. Atmospheric degradation of volatile organic compounds. *Chem Rev* 103:4605-4638.
- *Atkinson R, Aschmann SM, Fitz DR, et al. 1982. Rate constants for the gas-phase reactions of O₃ with selected organics at 296K. *Int J Chem Kinet* 14:13-18.
- +Axelson O, Gustavson J. 1978. Some hygienic and clinical observations on styrene exposure. *Scand J Work Environ Health* 4:215-219.

9. REFERENCES

- Badings HT, De Jong C, Dooper RP. 1985. Automatic system for rapid analysis of volatile compounds by purge-and-cold trapping/capillary gas chromatography. *J High Resolut Chromatogr Commun* 8:755-763.
- Baehr AL, Stackelberg PE, Baker RJ. 1999. Evaluation of the atmosphere as a source of volatile organic compounds in shallow groundwater. *Water Resour Res* 35:127-136.
- +*Ban M, Langonne I, Huguet N, et al. 2006. Inhaled chemicals may enhance allergic airway inflammation in ovalbumin-sensitised mice. *Toxicology* 226:161-171.
- *Banerjee S, Yalkowsky SH, Valvani SC. 1980. Water solubility and octanol/water partition coefficients of organics. Limitations of the solubility-partition coefficient correlation. *Environ Sci Technol* 14:1227-1229.
- Barale R. 1991. The genetic toxicology of styrene and styrene oxide. *Mutat Res* 257:107-126.
- *Barnes DG, Dourson M. 1988. Reference dose (RfD): Description and use in health risk assessments. *Regul Toxicol Pharmacol* 8(4):471-486.
- *Bartolucci GB, De Rosa E, Gori GP, et al. 1986. Biomonitoring of occupational exposure to styrene. *Appl Ind Hyg* 1:125-131.
- *Baselt RC. 1988a. Styrene. In: *Biological monitoring methods for industrial chemicals*. 2nd ed. Littleton, MA: PSG Publishing, Inc., 265-267.
- *Baselt RC. 1988b. Ethylbenzene. In: *Biological monitoring methods for industrial chemicals*. 2nd ed. Littleton, MA: PSG Publishing, Inc., 146-148.
- *Basirov AA. 1975. [Biochemical indexes of the gastric juice in the early diagnosis of stomach illness under the effect of toxic substances (1,3-butadiene and styrene).] *Azerb Med Zh* 52:60-66. (Russian)
- +*Behari M, Choudhary C, Roy S, et al. 1986. Styrene-induced peripheral neuropathy: A case report. *Eur Neurol* 25:424-427.
- +*Beliles RP, Butala JH, Stack CR, et al. 1985. Chronic toxicity and three-generation reproduction study of styrene monomer in the drinking water of rats. *Fundam Appl Toxicol* 5:855-868.
- *Belvedere G, Tursi F. 1981. Styrene oxidation to styrene oxide in human blood erythrocytes and lymphocytes. *Res Commun Chem Pathol Pharmacol* 33:273-282.
- +*Benignus VA, Geller AM, Boyes WK, et al. 2005. Human neurobehavioral effects of long-term exposure to styrene: A meta-analysis. *Environ Health Perspect* 113:532-538.
- Bergamaschi E, Mutti A, Cavazzini S, et al. 1995a. Peripheral markers of neurochemical effects among styrene-exposed workers. *Neurotoxicology* 16(3):545.
- *Bergamaschi E, Mutti A, Cavazzini S, et al. 1996. Peripheral markers of neurochemical effects among styrene-exposed workers. *Neurotoxicology* 17(3-4):753-759.

9. REFERENCES

- +Bergamaschi E, Smargiassi A, Mutti A, et al. 1995b. Immunological changes among workers occupationally exposed to styrene. *Int Arch Occup Environ Health* 67:165-171.
- +*Bergamaschi E, Smargiassi A, Mutti A, et al. 1997. Peripheral markers of catecholaminergic dysfunction and symptoms of neurotoxicity among styrene-exposed workers. *Int Arch Occup Environ Health* 69:209-214.
- *Berger GS, ed. 1994. Epidemiology of endometriosis. In: *Endometriosis: Modern surgical management of endometriosis*. New York, NY: Springer-Verlag, 3-7.
- Bernardini S, Hirvonen A, Jarventaus H, et al. 2002. Influence of GSTM1 and GSTT1 genotypes on sister chromatid exchange induction by styrene in cultured human lymphocytes. *Carcinogenesis* 23(5):893-897.
- *Berode M, Droz PO, Guillemin M. 1985. Human exposure to styrene: VI. Percutaneous absorption in human volunteers. *Int Arch Occup Environ Health* 55:331-336.
- *BfR. 2006. High daily intakes of cinnamon: Health risks cannot be ruled out. Bundesinstitut für Risikobewertung. BfR Health Assessment No. 044/2006. http://www.bfr.bund.de/cm/245/high_daily_intakes_of_cinnamon_health_risk_cannot_be_ruled_out.pdf. August 17, 2007.
- *Bigbee WL, Grant SG, Langlois RG, et al. 1996. Glycophorin a somatic cell mutation frequencies in Finnish reinforced plastics workers exposed to styrene. *Cancer Epidemiol Biomarkers Prev* 5:801-810.
- *Bigozzi CA, Maldotti A, Chiorboli C, et al. 1981. Kinetics and mechanisms of reactions between aromatic olefins and hydroxyl radicals. *Int J Chem Kinet* 13:1235-1242.
- *Biró A, Pallinger E, Major J, et al. 2002. Lymphocyte phenotype analysis and chromosome aberration frequency of workers occupationally exposed to styrene, benzene, polycyclic aromatic hydrocarbons or mixed solvents. *Immunol Lett* 81:133-140.
- Boffetta P, Sali D, Kolstad H, et al. 1998. Mortality of short-term workers in two international cohorts. *J Occup Environ Med* 40(12):1120-1126.
- Bolt HM. 1993. Interspecies differences in metabolism and kinetics of 1,3-butadiene, isobutene and styrene. *IARC Sci Publ* (127):37-44.
- Bommaraju TV, Luke B, O'Brien TF, et al. 2004. Chlorine. In: *Kirk-Othmer encyclopedia of chemical technology*. Vol. 6. John Wiley & Sons, Inc. <http://www.mrw.interscience.wiley.com/emrw/9780471238966/kirk/article/chlocurl.a01/current/pdf>. March 29, 2007.
- *Bond GG, Bodner KM, Olsen GW, et al. 1992. Mortality among workers engaged in the development or manufacture of styrene-based products-an update. *Scand J Work Environ Health* 18:145-154.
- *Bond JA. 1989. Review of toxicology of styrene. *CRC Crit Rev Toxicol* 19:227-249.
- Bonde JP. 1999. Occupational risk to male reproduction. *Int Arch Occup Environ Health* 72:133-134.

9. REFERENCES

- Boogaard PJ, de Kloe KP, Sumner SCJ, et al. 2000a. Disposition of [ring-U-¹⁴C]styrene in rats and mice exposed by recirculating nose-only inhalation. *Toxicol Sci* 58:161-172.
- Boogaard PJ, de Kloe KP, Wong BA, et al. 2000b. Quantification of DNA adducts formed in liver, lungs, and isolated lung cells of rats and mice exposed to ¹⁴C-styrene by nose-only inhalation. *Toxicol Sci* 57:203-216.
- Bouhamra WS, BuHamra SS, Thomson MS. 1997. Determination of volatile organic compounds in indoor and ambient air of residences in Kuwait. *Environ Int* 23(1):197-204.
- *Bouwer EJ, McCarty PL. 1984. Modeling of trace organics biotransformation in the subsurface. *Ground Water* 22:433-440.
- Brandorff NP, Flyvhokm M-A, Beck ID, et al. 1995. National survey on the use of chemicals in the working environment: Estimated exposure events. *Occup Environ Med* 52:454-463.
- Brantley AS, Townsend TG. 1999. Leaching of pollutants from reclaimed asphalt pavement. *Environ Eng Sci* 16(2):105-116.
- *Braun-Lüllemann A, Majcherczyk A, Huttermann A. 1997. Degradation of styrene by white-rot fungus. *Appl Microbiol Biotechnol* 47:150-155.
- Brenner DD, Jeffrey AM, Latriano L, et al. 1991. Biomarkers in styrene-exposed boatbuilders. *Mutat Res* 261(3):225-236.
- *Bridie AL, Wolff CJ, Winter M. 1979. BOD and COD of some petrochemicals. *Water Res* 13:627-630.
- Brinkmann E, Kaul G, Zeller H-J. 2000. Neuropsychiatric symptoms at low-level exposure to styrene [Abstract]. *Neurotoxicology* 21(5):897-898.
- Brodkin CA, Moon J-D, Camp J, et al. 2001. Serum hepatic biochemical activity in two populations of workers exposed to styrene. *Occup Environ Med* 58:95-102.
- Brondeau MT, Hesbert A, Beausoleil C, et al. 1999. To what extent are biomonitoring data available in chemical risk assessment? *Hum Exp Toxicol* 18:322-326.
- *Bronstein AC, Currence PL. 1988. Emergency care for hazardous materials exposure. St. Louis, MO: The C.V. Mosby Company, 221-222.
- Brosseau J, Heitz M. 1994. Trace gas compound emissions from municipal landfill sanitary sites. *Atmos Environ* 28:285-293.
- *Brown KW, Donnelly KC. 1988. An estimation of the risk associated with the organic constituents of hazardous and municipal waste landfill leachates. *Haz Waste Haz Mater* 5:1-30.
- Brown NA. 1991. Reproductive and developmental toxicity of styrene. *Reprod Toxicol* 5:3-29.
- Brown NA, Lamb JC, Brown SM, et al. 2000. A review of the developmental and reproductive toxicity of styrene. *Regul Toxicol Pharmacol* 32:228-247.

9. REFERENCES

- Brown SK, Sim MR, Abramson MJ, et al. 1994. Concentrations of volatile organic compounds in indoor air-A review. *Indoor Air* 4:123-134.
- Brown-Woodman PDC, Webster WS, Picker K, et al. 1994. *In vitro* assessment of individual and interactive effects of aromatic hydrocarbons on embryonic development of the rat. *Reprod Toxicol* 8:121-135.
- *Brugnone F, Perbellini L, Wang GZ, et al. 1993. Blood styrene concentrations in a "normal" population and in exposed workers 16 hours after the end of the workshift. *Int Arch Occup Environ Health* 65:125-130.
- *Bunce NJ, Dryfhout HG. 1992. Diurnal and seasonal modeling of the tropospheric half-lives of polycyclic aromatic hydrocarbons. *Can J Chem* 70:1966-1970.
- *Burbach BL, Perry JJ. 1993. Biodegradation and biotransformation of groundwater pollutant mixtures by *Mycobacterium vaccae*. *Appl Environ Microbiol* 59(4):1025-1029.
- Burns CJ, Cruzan G, Tsai SP. 2006. Styrene and breast cancer incidence in Texas: A comment on an ecological association. *Breast Cancer Res Treat* 97:339-340.
- Buschini A, De Palma G, Poli P, et al. 2003. Genetic polymorphism of drug-metabolizing enzymes and styrene-induced DNA damage. *Environ Mol Mutagen* 41:243-252.
- +*Bushnell PJ. 1994. Styrene impairs serial spatial reversal learning in rats. *J Am Coll Toxicol* 13(4):279-300.
- Bushnell PJ, Levin ED. 1993. Effects of dopaminergic drugs on working and reference memory in rats. *Pharmacol Biochem Behav* 45:765-776.
- +*Calabrese G, Martini A, Sessa G, et al. 1996. Otoneurological study in workers exposed to styrene in the fiberglass industry. *Int Arch Occup Environ Health* 68(4):219-223.
- +*Campagna D, Gobba F, Mergler D, et al. 1996. Color vision loss among styrene-exposed workers neurotoxicological threshold assessment. *Neurotoxicology* 17(2):367-373.
- +*Campagna D, Mergler D, Huel G, et al. 1995. Visual dysfunction among styrene-exposed workers. *Scand J Work Environ Health* 21:382-390.
- Campbell ME, Benson BA, Muir MA. 1995. Urban air quality and human health: A Toronto perspective. *Can J Public Health* 86(5):351-357.
- +*Campo P, Lataye R, Loquet G, et al. 2001. Styrene-induced hearing loss: A membrane insult. *Hear Res* 154(1-2):170-180.
- Campo P, Loquet G, Blachere V, et al. 1999. Toluene and styrene intoxication route in the rat cochlea. *Neurotoxicol Teratol* 21(4):427-434.
- *Canter LW, Sabatini DA. 1994. Contamination of public ground water supplies by Superfund sites. *Int J Environ Stud* 46:35-57.

9. REFERENCES

- *Cantoni L, Salmona M, Facchinetti T, et al. 1978. Hepatic and extrahepatic formation and hydration of styrene oxide *in vitro* in animals of different species and sex. *Toxicol Lett* 2:179-186.
- Cantor KP, Stewart PA, Brinton LA, et al. 1995. Occupational exposures and female breast cancer mortality in the United States. *J Occup Environ Med* 37(3):336-348.
- Cantoreggi S, Lutz WK. 1993. Covalent binding of styrene to DNA in rat and mouse. *Carcinogenesis* 14(3):355-360.
- Carlson GP. 1997a. Comparison of mouse strains for susceptibility to styrene-induced hepatotoxicity and pneumotoxicity. *J Toxicol Environ Health* 51:177-187.
- Carlson GP. 1997b. Effects of inducers and inhibitors on the microsomal metabolism of styrene to styrene oxide in mice. *J Toxicol Environ Health* 51:477-488.
- Carlson GP. 2003. *In vitro* metabolism of styrene to styrene oxide in liver and lung of CYP2E1 knockout mice. *J Toxicol Environ Health A* 66:861-869.
- Carlson GP. 2004a. Comparison of the susceptibility of wild-type and CYP2E1 knockout mice to the hepatotoxic and pneumotoxic effects of styrene and styrene oxide. *Toxicol Lett* 150:335-339.
- Carlson GP. 2004b. Influence of selected inhibitors on the metabolism of the styrene metabolite 4-vinylphenol in wild-type and CYP2E1 knockout mice. *J Toxicol Environ Health* 67A:905-909.
- *Carlson EM, Erskine MG. 1973. Styrene. *Chemical economics handbook*. Menlo Park, CA: Standford Research Institute.
- Carlson GP, Hynes DE, Mantick NA. 1998. Effects of inhibitors of CYP1A and CYP2B on styrene metabolism in mouse liver and lung microsomes. *Toxicol Lett* 98:131-137.
- *Carlson GP, Mantick NA, Powley MW. 2000. Metabolism of styrene by human liver and lung. *J Toxicol Environ Health A* 59:591-595.
- Carlson GP, Perez Rivera AA, Mantick NA. 2001. Metabolism of the styrene metabolite 4-vinylphenol by rat and mouse liver and lung. *J Toxicol Environ Health A* 63:541-551.
- Carlson GP, Turner M, Mantick NA. 2006. Effects of styrene and styrene oxide on glutathione-related antioxidant enzymes. *Toxicology* 227(3):217-226.
- *Carlsson A. 1981. Distribution and elimination of 14-C styrene in rat. *Scand J Work Environ Health* 7:45-50.
- +*Carpenter CP, Shaffer CB, Weil CS, et al. 1944. Studies on the inhalation of 1,3-butadiene; with a comparison of its narcotic effect with benzol, toluol, and styrene, and a note on the elimination of styrene by the human. *J Ind Hyg Toxicol* 26:69-78.
- *Castillo L, Baldwin M, Sassine MP, et al. 2001. Cumulative exposure to styrene and visual functions. *Am J Ind Med* 39:351-360.

9. REFERENCES

- *CEFIC. 2002. Styrene monomer: Environmental, health and safety guidelines. Conseil Européen de L'Industrie Chimique.
<http://www.styrenemonomer.org/Styrene%20Monomer%20Environmental%20Health%20Safety%20and%20Distribution%20Guidelines.pdf>. May 30, 2007.
- Chakrabarti SK. 1999. Styrene and styrene oxide affect the transport of dopamine in purified rat striatal synaptic vesicles. *Biochem Biophys Res Commun* 255:70-74.
- Chakrabarti SK, Duhr M-A, Senecal-Quevillon M, et al. 1993. Dose-dependent genotoxic effects of styrene on human blood lymphocytes and the relationship to its oxidative and metabolic effects. *Environ Mol Mutagen* 22:85-92.
- +Chakrabarti SK, Labelle L, Tuchweber B. 1987. Studies on the subchronic nephrotoxic potential of styrene in Sprague-Dawley rats. *Toxicology* 44:355-365.
- Chamkhia N, Sakly M, Rhouma KB. 2006. Male reproductive impacts of styrene in rat. *Toxicol Ind Health* 22(8):349-355.
- Chao J, Lin CT, Chung TH. 1983. Vapor pressure of coal chemicals. *J Phys Chem Ref Data* 12:1033, 1054.
- +*Checkoway H, Williams TM. 1982. A hematology survey of workers at a styrene-butadiene synthetic rubber manufacturing plant. *Am Ind Hyg Assoc J* 43:164-169.
- +*Checkoway H, Costa LG, Camp J, et al. 1992. Peripheral markers of neurochemical function among workers exposed to styrene. *Br J Ind Med* 49:560-565.
- *Cheng H, Sathiakumar N, Graff J, et al. 2007. 1,3-Butadiene and leukemia among synthetic rubber industry workers: Exposure-response relationships. *Chem Biol Interact* 166:15-24.
- Cherry N, Rodgers B, Venables H, et al. 1981. Acute behavioral effects of styrene exposure: A further analysis. *Br J Ind Med* 38:346-350.
- +*Cherry N, Waldron HA, Wells GG, et al. 1980. An investigation of the acute behavioral effects of styrene on factory workers. *Br J Ind Med* 37:234-240.
- +*Chia SE, Jeyaratnam J, Ong CN, et al. 1994. Impairment of color vision among workers exposed to low concentrations of styrene. *Am J Ind Med* 26:481-488.
- *Chmielewski J, Renke W. 1975. Clinical and experimental studies on the pathogenesis of toxic effects of styrene. II. The effect of styrene on the respiratory system. *Bull Inst Marit Trop Med Gdynia* 26:299-302.
- *Cho SI, Damokosh AI, Ryan LM, et al. 2001. Effects of exposure to organic solvents on menstrual cycle length. *J Occup Environ Med* 43(6):565-575.
- Christakopoulos A, Bergmark E, Zorcec V, et al. 1993. Monitoring occupational exposure to styrene from hemoglobin adducts and metabolites in blood. *Scand J Work Environ Health* 19:255-263.
- Chu KC, Cueto C, Ward JM. 1981. Factors in the evaluation of 200 National Cancer Institute carcinogen bioassays. *J Toxicol Environ Health* 8:251-280.

9. REFERENCES

- *Chua SC, Lee BL, Liau LS, et al. 1993. Determination of mandelic acid and phenylglyoxylic acid in the urine and its use in monitoring of styrene exposure. *J Anal Toxicol* 17(3):129-132.
- Chung JK, Yuan W, Zheng J. 2006. Investigation of bioactivation and toxicity of styrene in CYP2E1 transgenic cells. *Toxicology* 226(2-3):99-106.
- *Clay P. 2004. Styrene monomer does not induce unscheduled DNA synthesis in the mouse liver following inhalation exposure. *Mutagenesis* 19(6):489-492.
- *Clewell HJ, Andersen ME. 1985. Risk assessment extrapolations and physiological modeling. *Toxicol Ind Health* 1(4):111-131.
- Coccini T, Di Nucci A, Tonini M, et al. 1996. Effects of ethanol administration on cerebral non-protein sulfhydryl content in rats exposed to styrene vapour. *Toxicology* 106(1-3):115-122.
- Coccini T, Fenoglio C, Maestri L, et al. 1998. Effect of subchronic ethanol ingestion on styrene-induced damage to the tracheal and pulmonary epithelium of the rat. *J Appl Toxicol* 18:349-356.
- Coccini T, Fenoglio C, Nano R, et al. 1997. Styrene-induced alterations in the respiratory tract of rats treated by inhalation or intraperitoneally. *J Toxicol Environ Health* 52:63-77.
- Coccini T, Nappi R, Randine G, et al. 2000. The neuroendocrine system as a target for stressors: Interaction of styrene with the oestrus cycle in rat [Abstract]. *Eur J Neurosci* 12(Suppl 11):477.
- Coccini T, Randine G, Li B, et al. 1999. Effect of styrene on monoamine oxidase B activity in rat brain. *J Toxicol Environ Health* 56:59-68.
- *Cocheo V, Bellomo ML, Bombi GG. 1983. Rubber manufacture: Sampling and identification of volatile pollutants. *Am Ind Hyg Assoc J* 44:521-527.
- Coggon D. 1994. Epidemiological studies of styrene-exposed populations. *Crit Rev Toxicol* 24(S1):S107-S115.
- +*Coggon D, Osmond C, Pannett B, et al. 1987. Mortality of workers exposed to styrene in the manufacture of glass-reinforced plastics. *Scand J Work Environ Health* 13:94-99.
- Cohen JT, Carlson G, Charnley G, et al. 2002. A comprehensive evaluation of the potential health risks associated with occupational and environmental exposure to styrene. *J Toxicol Environ Health B Crit Rev* 5:1-263.
- *Coleman WE, Munch JW, Streicher RP, et al. 1984. The identification and measurement of components in gasoline, kerosene, and No. 2 fuel oil that partition into the aqueous phase after mixing. *Arch Environ Contam Toxicol* 13:171-178.
- *Compton-Quintana PJ, Jensen RH, Bigbee WL, et al. 1993. Use of the glycophorin A human mutation assay to study workers exposed to styrene. *Environ Health Perspect* 99:297-301.
- Conner MK, Alarie Y, Dombroske RL. 1979. Sister chromatid exchange in regenerating liver and bone marrow cells of mice exposed to styrene. *Toxicol Appl Pharmacol* 50:365-367.

9. REFERENCES

- *Conner MK, Alarie Y, Dombroske RL. 1980. Sister chromatid exchange in Murine alveolar macrophages, bone marrow, and regenerating liver cells induced by styrene inhalation. *Toxicol Appl Pharmacol* 55:37-42.
- +*Conti B, Maltoni C, Perino G, et al. 1988. Long-term carcinogenicity bioassays on styrene administered by inhalation, ingestion and injection and styrene oxide administered by ingestion in Sprague-Dawley rats, and *para*-methylstyrene administered by ingestion in Sprague-Dawley rats and Swiss mice. *Ann NY Acad Sci* 534:203-234.
- Contos DA, Holdren MW, Smith DL. 1995. Sampling and analysis of volatile organic compounds evolved during thermal processing of acrylonitrile butadiene styrene composite resins. *J Air Waste Manage Assoc* 45:686-694.
- *Correa SM, Torres AR, Arbilla G. 2004. Aromatic volatile organic compounds emissions in a tire recapping unit. *Bull Environ Contam Toxicol* 72:255-260.
- Coyle YM, Hynan LS, Euhus DM, et al. 2005. An ecological study of the association of environmental chemicals on breast cancer incidence in Texas. *Breast Cancer Res Treat* 92:107-114.
- *CRISP. 2007. Styrene: Computer Retrieval of Information on Scientific Projects Database. Bethesda, MD: National Institutes of Health, Division of Research Grants.
- +*Crofton KM, Lassiter TL, Rebert CS. 1994. Solvent-induced ototoxicity in rats: An atypical selective mid-frequency hearing deficit. *Hear Res* 80:25-30.
- *Crump D. 1995. Volatile organic compounds in indoor air. *Issues Environ Sci Technol* 4:109-124.
- *Cruzan G, Carlson GP, Johnson KA. 2002. Styrene respiratory tract toxicity and mouse lung tumors are mediated by CYP2F-generated metabolites. *Regul Toxicol Pharmacol* 35:308-319.
- +*Cruzan G, Carlson GP, Turner M, et al. 2005b. Ring-oxidized metabolites of styrene contribute to styrene-induced Clara-cell toxicity in mice. *J Toxicol Environ Health* 68A:229-237.
- +*Cruzan G, Cushman JR, Andrews LS, et al. 1997. Subchronic inhalation studies of styrene in CD rats and CD-1 mice. *Fundam Appl Toxicol* 35(2):152-165.
- +*Cruzan G, Cushman JR, Andrews LS, et al. 1998. Chronic toxicity/oncogenicity study of styrene in CD rats by inhalation exposure for 104 weeks. *Toxicol Sci* 46:266-281.
- +*Cruzan G, Cushman JR, Andrews LS, et al. 2001. Chronic toxicity/oncogenicity study of styrene in CD-1 mice by inhalation exposure for 104 weeks. *J Appl Toxicol* 21:185-198.
- Cruzan G, Faber WD, Johnson KA, et al. 2005c. Two generation reproduction study of styrene by inhalation in Crl-CD rats. *Birth Defects Res B Dev Reprod Toxicol* 74(3):211-220.
- +*Cruzan G, Faber WD, Johnson KA, et al. 2005a. Developmental neurotoxicity study of styrene by inhalation in Crl-CD rats. *Birth Defects Res B Dev Reprod Toxicol* 74(3):221-232.
- *Csanády GA, Kessler W, Hoffmann HD, et al. 2003. A toxicokinetic model for styrene and its metabolite styrene-7,8-oxide in mouse, rat, and human with special emphasis on the lung. *Toxicol Lett* 138:75-102.

9. REFERENCES

- *Csanády GA, Mendrala AL, Noland RJ, et al. 1994. A physiologic pharmacokinetic model for styrene and styrene-7,8-oxide in mouse, rat and man. *Arch Toxicol* 68:143-157.
- +*Dalton P, Cowart B, Dilks D, et al. 2003. Olfactory function in workers exposed to styrene in the reinforced-plastics industry. *Am J Ind Med* 44(1):1-11.
- Danishefsky I, Willhite M. 1954. The metabolism of styrene in the rat. *J Biol Chem* 211:544-553.
- +*Das M, Dixit R, Mushtaq M, et al. 1981. Effect of styrene on hepatic mixed function oxidases, glutathione content and glutathione-S-transferase activity in rats. *Drug Chem Toxicol* 4:219-227.
- +*Das M, Srivastava SP, Seth PK. 1983. Effect of styrene on glutathione content and some xenobiotic metabolizing enzymes of rat kidney. *Acta Pharmacol Toxicol (Copenh)* 52:47-50.
- +*Daston GP, Overmann GJ, Taubeneck MW, et al. 1991. The role of metallothionein induction and altered zinc status in maternally mediated developmental toxicity: Comparison of the effects of urethane and styrene in rats. *Toxicol Appl Pharmacol* 110:450-463.
- +*DeCeaurriz J, Desiles JP, Bonnet P, et al. 1983. Concentration-dependent behavioral changes in mice following short-term inhalation exposure to various industrial solvents. *Toxicol Appl Pharmacol* 67:383-389.
- *DeJong G, van Sittert NJ, Natarajan AT. 1988. Cytogenetic monitoring of industrial populations potentially exposed to genotoxic chemicals and of control populations. *Mutat Res* 204:451-464.
- Delbressine LP, Van Bladeren PJ, Smeets FL, et al. 1981. Stereoselective oxidation of styrene to styrene oxide in rats as measured by mercapturic acid excretion. *Xenobiotica* 11:589-594.
- *Delzell E, Macalluso M, Sathiakumar N, et al. 2001. Leukemia and exposure to 1,3-butadiene, styrene and dimethyldithiocarbamate among workers in the synthetic rubber industry. *Chem Biol Interact* 135-136:515-534.
- *Delzell E, Sathiakumar N, Hovinga M, et al. 1996. A follow-up study of synthetic rubber workers. *Toxicology* 113:182-189.
- *DeMeester C, Duverger-van Bogaert M, Lambotte-Vandepaer M, et al. 1981. Mutagenicity of styrene in the *Salmonella typhimurium* test system. *Mutat Res* 90:443-450.
- DeMeester C, Poncelet F, Roberfroid M, et al. 1977. Mutagenicity of styrene and styrene oxide. *Mutat Res* 56:147-152.
- DePalma G, Manini P, Mozzoni P, et al. 2001. Polymorphism of xenobiotic-metabolizing enzymes and excretion of styrene-specific mercapturic acids. *Chem Res Toxicol* 14:1393-1400.
- *de Raat VK. 1978. Induction of sister chromatid exchanges by styrene and its presumed metabolite styrene oxide in the presence of rat liver homogenate. *Chem Biol Interact* 20:163-170.
- De Rosa E, Cellini M, Sessa G, et al. 1993. Biological monitoring of workers exposed to styrene and acetone. *Int Arch Occup Environ Health* 65(Suppl 1):S107-S110.

9. REFERENCES

- *Dilling WL. 1977. Interphase transfer processes. II. Evaporation rates of chloro methanes, ethanes, ethylenes, propanes, and propylenes from dilute aqueous solutions. Comparisons with theoretical predictions. *Environ Sci Technol* 11:405-409.
- Diodovich C, Bowe G, Gribaldo L. 2004. Response of human cord blood cells to styrene exposure: Evaluation of its effects on apoptosis and gene expression by genomic technology [Abstract]. *Toxicol Appl Pharmacol* 197(3):276-277.
- +*Dixit R, Das M, Mushtaq M, et al. 1982. Depletion of glutathione content and inhibition of glutathione-S-transferase and aryl hydrocarbon hydroxylase activity of rat brain following exposure to styrene. *Neurotoxicology* 3:142-145.
- +*Dogra RKS, Chandra K, Chandra S, et al. 1992. Host resistance assays as predictive models in styrene immunomodulation. *Int J Immunopharmacol* 14(6):1003-1009.
- Dogra RKS, Khanna S, Srivastava SN, et al. 1993. Immunomodulation due to coexposure to styrene and dioctyl phthalate in mice. *Immunopharmacol Immunotoxicol* 15(4):491-514.
- *Dolara P, Caderni G, Lodovici M, et al. 1984. Determination of styrene in the urine of workers manufacturing polystyrene plastics. *Ann Occup Hyg* 28:195-199.
- *Donner M, Sorsa M, Vainio H. 1979. Recessive lethals induced by styrene and styrene oxide in *Drosophila melanogaster*. *Mutat Res* 67:373-376.
- Doskey PV, Yoshiko F, Sultan M. 1999. Source profiles for nonmethane organic compounds in the atmosphere of Cairo, Egypt. *J Air Waste Manage Assoc* 49:814-822.
- *Drummond L, Caldwell J, Wilson HK. 1989. The metabolism of ethylbenzene and styrene to mandelic acid: Stereochemical considerations. *Xenobiotica* 19:199-207.
- *Dumont JP, Adda J. 1978. Occurrence of sesquiterpenes in mountain cheese volatiles. *J Agric Food Chem* 26:364-367.
- Dunbar RC. 1989. Time-resolved unimolecular dissociation of styrene ion. Rats and activation parameters. *J Am Chem Soc* 111:5572-5576.
- *Dunkel VC, Zeiger E, Brusick D, et al. 1985. Reproducibility of microbial mutagenicity assays: II. Testing of carcinogens and noncarcinogens in *Salmonella typhimurium* and *Escherichia coli*. *Environ Mut* 7(Suppl 5):1-248.
- Durkal SK, Biswas P, Shi J, Bernhard MJ. 1998. Characterization of polycyclic aromatic hydrocarbon particulate and gaseous emissions from polystyrene combustion. *Environ Sci Technol* 32:2301-2307.
- *Dutkiewicz T, Tyras H. 1968. Skin absorption of toluene, styrene, and xylene by man. *Br J Ind Med* 25:243.
- ECIS. 2007. Styrene. European chemical Substances Information. <http://ecb.jrc.it.esis/>. May 30, 2007.
- Edling C, Ekberg K. 1985. No acute behavioural effects of exposure to styrene: A safe level of exposure? *Br J Ind Med* 42:301-304.

9. REFERENCES

- +*Edling C, Anundi H, Johanson G, et al. 1993. Increase in neuropsychiatric symptoms after occupational exposure to low levels of styrene (Comment in: *Occup Environ Med* 51(4):286-287). *Br J Ind Med* 50(9):843-850.
- +*Eguchi T, Kishi R, Harabuchi I, et al. 1995. Impaired colour discrimination among workers exposed to styrene: Relevance of a urinary metabolite. *Occup Environ Med* 52:534-538.
- *Eitzer BD. 1995. Emissions of volatile organic chemicals from municipal solid waste composting facilities. *Environ Sci Technol* 29:896-902.
- *Elia VJ, Anderson LA, MacDonald TJ, et al. 1980. Determination of urinary mandelic and phenylglyoxylic acids in styrene exposed workers and a control population. *Am Ind Hyg Assoc J* 41:922-926.
- *Ellenhorn MJ, Barceloux DG. 1988. Styrene. In: *Medical toxicology: Diagnosis and treatment of human poisoning*. New York, NY: Elsevier, 956-959.
- *Engelhardt G, Gamer A, Vodicka P, et al. 2003. A re-assessment of styrene-induced clastogenicity in mice in a subacute inhalation study. *Arch Toxicol* 77:56-61.
- *Engstrom J, Astrand I, Wigaeus E. 1978b. Exposure to styrene in a polymerization plant: Uptake in the organism and concentration in subcutaneous adipose tissue. *Scand J Work Environ Health* 4:324-329.
- *Engstrom J, Bjurstrom R, Astrand I, et al. 1978a. Uptake, distribution and elimination of styrene in man: Concentration in subcutaneous adipose tissue. *Scand J Work Environ Health* 4:315-323.
- *Engstrom K, Harkonen H, Kalliokoski P, et al. 1976. Urinary mandelic acid concentration after occupational exposure to styrene and its use as a biological exposure test. *Scand J Work Environ Health* 2:21-26.
- *EPA. 1975. Identification of organic compounds in effluents from industrial sources. Washington, DC: U.S. Environmental Protection Agency, Office of Toxic Substances. EPA560375002.
- *EPA. 1976. Frequency of organic compounds identified in water. Athens, GA: U.S. Environmental Protection Agency, Office of Research and Development. EPA600476062. PB265470.
- *EPA. 1978. Environmental monitoring near industrial sites: Brominated chemicals. Part I. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Toxic Substances. PB286484.
- *EPA. 1979a. Atmospheric reaction products of organic compounds. Washington, DC: U.S. Environmental Protection Agency, Office of Toxic Substances. EPA5601279001. PB301384.
- *EPA. 1979b. Identification of organic compounds in industrial effluent discharges. Athens, GA: U.S. Environmental Protection Agency, Office of Research and Development. EPA600479016. PB294794.
- *EPA. 1980. Fate of toxic and hazardous materials in the air environment. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development. EPA600380084. PB80221948.
- *EPA. 1981. Treatability manual. Vol. I. Treatability data. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development. EPA600282001A.

9. REFERENCES

- *EPA. 1982a. Aquatic fate process data for organic priority pollutants. Washington, DC: U.S. Environmental Protection Agency, Office of Water Regulations and Standards. EPA440481014. PB87169090.
- EPA. 1982b. Chemical information rules; manufacturers reporting; preliminary assessment information. U.S. Environmental Protection Agency. Fed Regist 47:26992, 27007.
- *EPA. 1983. Volatile organic chemicals in the atmosphere: An assessment of available data. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development. EPA600383027A.
- *EPA. 1984a. Drinking water criteria document for styrene. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Criteria and Assessment Office. EPA600X84195.
- *EPA. 1984b. Health and environmental effects profile for styrene. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Criteria and Assessment Office. EPA600X84325.
- *EPA. 1985a. Drinking water criteria document for styrene. Final draft. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment. ECAOCIN409.
- *EPA. 1985b. Synthetic organic compound sampling survey of public water supplies. Washington, DC: U.S. Environmental Protection Agency. PB85214427.
- *EPA. 1986a. National primary drinking water regulations; synthetic organic chemicals, inorganic chemicals and microorganisms; correction. U.S. Environmental Protection Agency. Fed Regist 51(25):4618.
- *EPA. 1986b. Gas chromatography/mass spectrometry for volatile organics, method 8240. In: Test methods for evaluating solid waste. Vol. IA: Laboratory manual, physical/chemical methods. 3rd ed. SW-846. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response.
- *EPA. 1986c. Method 8020. Aromatic volatile organics. In: Test methods for evaluating solid waste. 3rd ed. SW-846. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response.
- *EPA. 1986d. Broad scan analysis of the FY82 national human adipose tissue survey specimens. Vol. I. Executive summary. Washington, DC: U.S. Environmental Protection Agency, Office of Toxic Substances. EPA560586035.
- EPA. 1986e. Report on the interim data base for state and local air toxic volatile organic chemical measurements. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards. EPA450486012. PB87168779.
- EPA. 1987a. Styrene: Health advisory. Washington, DC: U.S. Environmental Protection Agency, Office of Drinking Water.
- EPA. 1987b. National primary drinking water regulations; synthetic organic chemicals, monitoring for unregulated contaminants. Part II. U.S. Environmental Protection Agency. Fed Regist 52:25710.

9. REFERENCES

EPA. 1987c. List (phase 1) of hazardous constituents for ground water monitoring, final rule. U.S. Environmental Protection Agency: Fed Regist 52(131):25942-25952.

*EPA. 1987d. Toxic air pollutant/source crosswalk: A screening tool for locating possible sources emitting toxic air pollutants. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards. EPA450487023a. PB88161146.

*EPA. 1987e. The Total Exposure Assessment Methodology (TEAM) study: Summary and analysis. Vol. I. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development. EPA600687002a. PB088100060.

*EPA. 1988a. Toxic chemical release reporting: Community right to know. U.S. Environmental Protection Agency. Fed Regist 53:4500-4501.

*EPA. 1988b. Drinking water criteria document for styrene. Final draft. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment. ECAOCIN409.

*EPA. 1988c. National ambient volatile organic compounds (VOCs) database update. Research Triangle Park, NC: U.S. Environmental Protection Agency. EPA600338010a. PB88195631.

EPA. 1989a. List of hazardous substances and reportable quantities. U.S. Environmental Protection Agency. Fed Regist 54(155):33425-33484.

EPA. 1989b. National primary and secondary drinking water regulations. U.S. Environmental Protection Agency. Fed Regist 54:22064, 22090.

*EPA. 1989c. The toxics-release inventory: A national perspective. Washington, DC: U.S. Environmental Protection Agency, Office of Toxic Substances. EPA560489005.

*EPA. 1989d. Interim methods for development of inhalation reference doses. Washington, DC: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment. EPA600888066F.

*EPA. 1989e. Exposure factors handbook. Washington, DC: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment. EPA600889043.

*EPA. 1989f. Method 503.1 volatile aromatic and unsaturated organic compounds in water by purge and trap gas chromatography. U.S. Environmental Protection Agency, Office of Research and Development.

*EPA. 1989g. Methods 508, 501.1, 502.2. Methods for the determination of organic compounds in drinking water. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory. EPA600488039.

*EPA. 1989h. Method 524.1. Measurement of purgeable organic compounds in water by packed column gas chromatography/mass spectrometry. Methods for the determination of organic compounds in drinking water. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Research and Development. EPA600488039.

*EPA. 1989i. Method 524.2. Measurement of purgeable organic compounds in water by capillary column gas chromatography/mass spectrometry. In: Methods for the determination of organic compounds in drinking water. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Research and Development. EPA600488039.

9. REFERENCES

- *EPA. 1990. Interim methods for development of inhalation reference concentrations. Washington, DC: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment, Office of Research and Development, Environmental Criteria and Assessment Office. EPA600890066A. PB90238890.
- *EPA. 1993a. Determination of rates of reaction in the gas-phase in the troposphere. Theory and practice. 5. Rate of indirect photoreaction: Evaluation of the atmospheric oxidation computer program of Syracuse Research Corporation for estimating the second-order rate constant for the reaction of an organic chemical with hydroxyl radicals. Washington, DC: U.S. Environmental Protection Agency. EPA744R93001.
- EPA. 1993b. Final report on literature review of atmospheric transformation products of Clean Air Act Title III hazardous air pollutants. Research Triangle Park: U.S. Environmental Protection Agency. EPA600R94088.
- *EPA. 1995. Method 502.2. Volatile organic compounds in water by purge and trap capillary column gas chromatography with photoionization and electrolytic conductivity detectors in series. Cincinnati, OH: U.S. Environmental Protection Agency.
- *EPA. 1996. Method 8021B: Aromatic and halogenated volatiles by gas chromatography using photoionization and/or electrolytic conductivity detectors. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste.
- *EPA. 1997. Special report on environmental endocrine disruption: An effects assessment and analysis. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. EPA630R96012.
- *EPA. 2001. Method 1625. Semivolatile organic compounds by isotope dilution GCMS. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR Part 136, App. A.
- *EPA. 2003. National primary drinking water regulations. Washington, DC: U.S. Environmental Protection Agency, Office of Ground Water and Drinking Water. EPA816F03016. <http://www.epa.gov/safewater/mcl.html>. March 07, 2006.
- *EPA. 2005. Toxic chemical release inventory reporting forms and instructions: Revised 2004 version. Section 313 of the Emergency Planning and Community Right-to-Know Act (Title III of the Superfund Amendments and Reauthorization Act of 1986). U.S. Environmental Protection Agency. Office of Environmental Information. EPA260B05001.
- *EPA. 2006. Drinking water standards and health advisories. Washington, DC: U.S. Environmental Protection Agency, Office of Water. EPA822R04005. <http://epa.gov/waterscience/criteria/drinking/>. April 11, 2007.
- *EPA. 2007a. Acute exposure guideline levels (AEGLs) Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics.
- *EPA. 2007b. Designated as hazardous substances in accordance with Section 311(b)(2)(A) of the Clean Water Act. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 116.4. <http://www.epa.gov/epacfr40/chapt-I.info/chi-toc.htm>. April 11, 2007.

9. REFERENCES

- *EPA. 2007c. Hazardous air pollutants. Clean Air Act. U.S. Environmental Protection Agency. United States Code. 42 USC 7412. <http://www.epa.gov/ttn/atw/orig189.html>. April 11, 2007.
- *EPA. 2007d. Master testing list. Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics. <http://www.epa.gov/opptintr/chemtest/pubs/mtl.htm>. May 10, 2007.
- *EPA. 2007e. Reportable quantities of hazardous substances designated pursuant to Section 311 of the Clean Water Act. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 117.3. <http://www.epa.gov/epacfr40/chapt-I.info/chi-toc.htm>. April 11, 2007.
- *EPA. 2007f. Superfund, emergency planning, and community right-to-know programs. Designation, reportable quantities, and notifications. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 302.4. <http://www.epa.gov/epacfr40/chapt-I.info/chi-toc.htm>. April 11, 2007.
- *EPA. 2007g. Superfund, emergency planning, and community right-to-know programs. Toxic chemical release reporting. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 372.65. <http://www.epa.gov/epacfr40/chapt-I.info/chi-toc.htm>. April 11, 2007.
- Evans GF, Lumpkin TA, Smith DL, et al. 1992. Measurements of VOCs from the Tams network. *J Air Waste Manage Assoc* 42:1319-1323.
- +*Fallas C, Fallas J, Maslard P, et al. 1992. Subclinical impairment of colour vision among workers exposed to styrene. *Br J Ind Med* 49(10):679-682.
- Fawell JK, Fielding M. 1985. Identification and assessment of hazardous compounds in drinking water. *Sci Total Environ* 47:317-341.
- *FDA. 2006a. Beverages. Bottled water. U.S. Food and Drug Administration. Code of Federal Regulations. 21 CFR 165.110. <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm>. April 11, 2007.
- *FDA. 2006b. Indirect food additives: Adhesives and components of coatings. U.S. Food and Drug Administration. Code of Federal Regulations. 21 CFR 175. <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm>. May 10, 2007.
- *FDA. 2007a. Everything added to food in the United States (EAFUS). Washington, DC: U.S. Food and Drug Administration. <http://vm.cfsan.fda.gov/~dms/eafus.html>. April 11, 2007.
- *FDA. 2007b. Food additives permitted for direct addition to food for human. Code of Federal Regulations. 21 CFR 172. Washington, DC: U.S. Food and Drug Administration. <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm>. May 15, 2007.
- *FEDRIP. 2007. Styrene. Federal Research in Progress database. Springfield, VA: National Technical Information Service.
- Fechter LD. 1993. Effects of acute styrene and simultaneous noise exposure on auditory function in the guinea pig. *Neurotoxicol Teratol* 15(3):151-155.
- Fellin P, Otson R. 1994. Assessment of the influence of climatic factors on concentration levels of volatile organic compounds (VOC's) in Canadian homes. *Atmos Environ* 28:3581-3586.

9. REFERENCES

- Fernandez-Nieto M, Quirce S, Fraj J, et al. 2006. Airway inflammation in occupational asthma caused by styrene. *J Allergy Clin Immunol* 117(4):948-950.
- Filser JG, Johanson G, Kessler W, et al. 1993. A pharmacokinetic model to describe toxicokinetic interactions between 1,3-butadiene and styrene in rats: Predictions for human exposure. *IARC Sci Publ* (127):65-78.
- Filser JG, Kessler W, Csanady GA. 2002. Estimation of a possible tumorigenic risk of styrene from daily intake via food and ambient air. *Toxicol Lett* 126:1-18.
- Filser JG, Schwegler U, Csanady GA, et al. 1993. Species-specific pharmacokinetics of styrene in rat and mouse. *Arch Toxicol* 67(8):517-530.
- *Fishbein L. 1992. Exposure from occupational versus other sources. *Scand J Work Environ Health* 18(1):5-16.
- *Fisher J, Mahle D, Bankston L, et al. 1997. Lactational transfer of volatile chemicals in breast milk. *Am Ind Hyg Assoc J* 58(6):425-431.
- Fjeldstad PE, Woldbaek T. 1992. A national exposure database. Special Publication of the Royal Chemical Society 108:303-310.
- Fjeldstad PE, Thorud S, Wannag A. 1979. Styrene oxide in the manufacture of reinforced polyester plastics. *Scand J Work Environ Health* 5:162-163.
- *Fleming-Jones ME, Smith RE. 2003. Volatile organic compounds in foods: A five-year study. *J Agric Food Chem* 51(27):8120-8127.
- +*Flodin U, Ekberg K, Andersson L. 1989. Neuropsychiatric effects of low exposure to styrene. *Br J Ind Med* 46:805-808.
- *Fomon SJ. 1966. Body composition of the infant: Part 1: The male reference infant. In: Faulkner F, eds. *Human development*. Philadelphia, PA: WB Saunders, 239-246.
- *Fomon SJ, Haschke F, Ziegler EE, et al. 1982. Body composition of reference children from birth to age 10 years. *Am J Clin Nutr* 35((Suppl 5)):1169-1175.
- Font R, Esperanza M, Garcia AN. 2003. Toxic by-products from the combustion of Kraft lignin. *Chemosphere* 52:1047-1058.
- Fouremant GL, Harris C, Guengerich FP, et al. 1989. Stereoselectivity of styrene oxidation in microsomes and in purified cytochrome P-450 enzymes from rat liver. *J Pharmacol Exp Ther* 248:492-497.
- *Frentzel-Beyme R, Thiess AM, Wieland R. 1978. Survey of mortality among employees engaged in manufacture of styrene and polystyrene at the BASF Ludwigshafen works. *Scand J Work Environ Health* 4:231-239.
- FSTRAC. 1988. Summary of state and federal drinking water standards and guidelines. Washington, DC: Federal-State Toxicology and Regulatory Alliance Committee.

9. REFERENCES

- *Fu MH, Alexander M. 1992. Biodegradation of styrene in samples of natural environments. *Environ Sci Technol* 26:1540-1544.
- *Fu MH, Alexander M. 1996. Biodegradation of styrene in waterlogged soils and aquifer solids. *Soil Sci* 161(12):846-851.
- *Fu MH, Mayton H, Alexander M. 1994. Desorption and biodegradation of sorbed styrene in soil and aquifer solids. *Environ Toxicol Chem* 13(5):749-753.
- +*Fujita H, Koizumi A, Furusawa T, et al. 1987. Decreased erythrocyte β -aminolevulinic acid dehydratase activity after styrene exposure. *Biochem Pharmacol* 36:711-716.
- *Fung F, Clark RF. 1999. Styrene-induced peripheral neuropathy. *Clin Toxicol* 37(1):91-97.
- Gagnaire F, Chalansonnet M, Carabin N, et al. 2006. Effects of subchronic exposure to styrene on the extracellular and tissue levels of dopamine, serotonin and their metabolites in rat brain. *Arch Toxicol* 80(10):703-712.
- Gamberale F, Hultengren M. 1974. Exposure to styrene. II. Psychological function. *Work Environ Health* 11:86-93.
- +*Gamberale F, Lisper HO, Olson BA. 1976. The effect of styrene vapour on the reaction time of workers in the plastic boat industry. In: Horvath M, ed. *Adverse effects of environmental chemicals and psychomotor drugs. Neurophysiological and behavioral tests.* London, England: Elsevier Scientific Pub., 135-148.
- Gamer AO, Leibold E, Deckardt K, et al. 2004. The effects of styrene on lung cells in female mice and rats. *Food Chem Toxicol* 42(10):1655-1667.
- *Gargas ML, Burgess RJ, Voisard DE, et al. 1989. Partition coefficients of low-molecular-weight volatile chemicals in various liquids and tissues. *Toxicol Appl Pharmacol* 98:87-99.
- *Gerin M, Siemiatycki J, Desy M, et al. 1998. Associations between several sites of cancer and occupational exposure to benzene, toluene, xylene, and styrene: Results of a case-control study in Montreal. *Am J Ind Med* 34:144-156.
- *Geuskens RBM, van der Klaauw MM, van der Tuin J, et al. 1992. Exposure to styrene and health complaints in the Dutch glass-reinforced plastics industry. *Ann Occup Hyg* 36:47-57.
- *Ghittori S, Imbriani M, Pezzagno G, et al. 1987. The urinary concentration of solvents as a biological indicator of exposure: Proposal for the biological equivalent exposure limit for nine solvents. *Am Ind Hyg Assoc J* 48:786-790.
- Ghittori S, Maestri L, Imbriani M, et al. 1997. Urinary excretion of specific mercapturic acids in workers exposed to styrene. *Am J Ind Med* 31(5):636-644.
- *Gilbert J, Startin JR. 1983. A survey of styrene monomer levels in foods and plastic packaging by coupled mass spectrometry-automatic headspace gas chromatography. *J Sci Food Agric* 34:647-652.
- Girman JR, Hogson AT, Newton AS. 1986. Emissions of volatile organic compounds from adhesives with indoor applications. *Environ Int* 12:317-321.

9. REFERENCES

- *Giwercman A, Carlsen E, Keiding N, et al. 1993. Evidence for increasing incidence of abnormalities of the human testis: A review. *Environ Health Perspect Suppl* 101(2):65-71.
- Gobba F. 2000. Color vision: A sensitive indicator of exposure to neurotoxins. *Neurotoxicology* 21(5):857-862.
- Gobba F, Cavalleri A. 1993. Kinetics of urinary excretion and effects on colour vision after exposure to styrene. *IARC Sci Publ* (127):79-88.
- *Gobba F, Cavalleri F, Bontadi D, et al. 1995. Peripheral neuropathy in styrene-exposed workers. *Scand J Work Environ Health* 21:517-520.
- *Gobba F, Galassi C, Ghittori S, et al. 1993. Urinary styrene in the biological monitoring of styrene exposure. *Scand J Work Environ Health* 19:175-182.
- +*Gobba F, Galassi C, Imbriani M, et al. 1991. Acquired dyschromatopsia among styrene-exposed workers (Comment in: *J Occup Med* 34(5):563-564). *J Occup Med* 33(7):761-765.
- Godderis L, De Boeck M, Haufroid V, et al. 2004. Influence of genetic polymorphisms on biomarkers of exposure and genotoxic effects in styrene-exposed workers. *Environ Mol Mutagen* 44:293-303.
- +*Gong YY, Kishi R, Katakura Y, et al. 2002. Relation between colour vision loss and occupational styrene exposure level. *Occup Environ Med* 59(12):824-829.
- *Goodley PC, Gordon M. 1976. Characterization of industrial organic compounds in water. *Trans KY Acad Sci* 37(1-2):11-15.
- *Gosselin RE, Smith RP, Hodge HC, et al. 1984. *Clinical toxicology of commercial products*. 5th ed. Baltimore, MD: Williams and Wilkins, II-152.
- Gotell P, Axelson O, Lindelof B. 1972. Field studies on human styrene exposure. *Work Environ Health* 9:76-83.
- *Gotz R, Bauer OR, Friesel P, et al. 1998. Organic trace compounds in the water of the River Elbe near Hamburg. Part II. *Chemosphere* 36:2103-2118.
- *Graedel TE. 1978. Aromatic compounds: Benzene and derivatives. In: *Chemical compounds in the atmosphere*. New York, NY: Academic Press, 105-113.
- *Graff JJ, Sathiakumar N, Macaluso M, et al. 2005. Chemical exposures in the synthetic rubber industry and lymphohematopoietic cancer mortality. *J Occup Environ Med* 47(9):916-932.
- *Grbic-Galic D, Churchman-Eisel N, Mrakovic I. 1990. Microbial transformation of styrene by anaerobic consortia. *J Appl Bacteriol* 69:247-260.
- *Green DR, Le Pape D. 1987. Stability of hydrocarbon samples on solid-phase extraction columns. *Anal Chem* 59:699-703.
- +*Green T, Lee R, Toghil A, et al. 2001a. The toxicity of styrene to the nasal epithelium of mice and rats: Studies on the mode of action and relevance to humans. *Chem Biol Interact* 137:185-202.

9. REFERENCES

- Green T, Toghil A, Foster JR. 2001b. The role of cytochromes P-450 in styrene induced pulmonary toxicity and carcinogenicity. *Toxicology* 169(2):107-117.
- *Grosjean D, Fung K. 1984. Hydrocarbons and carbonyls in Los Angeles air. *J Air Pollut Control Assoc* 34:537-543.
- *Grosjean E, Grosjean D, Rasmussen RA. 1998. Ambient concentrations, sources, emission rates, and photochemical reactivity of C2-C10 hydrocarbons in Porto Alegre, Brazil. *Environ Sci Technol* 32(14):2061-2069.
- *Guillemin MP, Bauer D. 1979. Human exposure to styrene. III. Elimination kinetics of urinary mandelic and phenylglyoxylic acids after single experimental exposure. *Int Arch Occup Environ Health* 44:249-263.
- *Guillemin MP, Berode M. 1988. Biological monitoring of styrene: A review. *Am Ind Hyg Assoc J* 49:497-505.
- *Guzelian PS, Henry CJ, Olin SS, eds. 1992. Similarities and differences between children and adults: implications for risk assessment. Washington DC: International Life Sciences and Press Institute Press.
- *Haddad LM, Winchester JF. 1990. Clinical management of poisoning and drug overdose. 2nd ed. Philadelphia, PA: W.B. Saunders Company, 1226-1228.
- *Hallier E, Goergens HW, Hallier K, et al. 1994. Intervention study on the influence of reduction of occupational exposure to styrene on sister chromatid exchanges in lymphocytes. *Int Arch Occup Environ Health* 66(3):167-172.
- *Hallier E, Goergens HW, Karels H, et al. 1995. A note on individual differences in the urinary excretion of optical enantiomers of styrene metabolites and of styrene-derived mercapturic acids in humans. *Arch Toxicol* 69(5):300-305.
- *Hampton CV, Pierson WR, Harvey TM, et al. 1982. Hydrocarbon gases emitted from vehicles on the road. 1. A qualitative gas chromatography/mass spectrometry survey. *Environ Sci Technol* 16:287-298.
- *Hampton CV, Pierson WR, Schuetz D, et al. 1983. Hydrocarbon gases emitted from vehicles on the road. 2. Determination of emission rates from diesel and spark-ignition vehicles. *Environ Sci Technol* 17:699-708.
- *Hansch C, Leo A, Hoekman D. 1995. Exploring QSAR. Hydrophobic, electronic, and steric constants. Washington, DC: American Chemical Society, 40.
- *Hansteen IL, Jelmert O, Torgrimsen T, et al. 1984. Low human exposure to styrene in relation to chromosome breaks, gaps and sister-chromatid exchanges. *Hereditas* 100:87-91.
- +*Härkönen H, Holmberg PC. 1982. Obstetric histories of women occupationally exposed to styrene. *Scand J Work Environ Health* 8:74-77.
- +*Härkönen H, Lehtniemi A, Aitio A. 1984. Styrene exposure and the liver. *Scand J Work Environ Health* 10:59-61.

9. REFERENCES

- *Härkönen H, Kalliokoski P, Hietala S, et al. 1974. Concentrations of mandelic and phenylglyoxylic acid in urine as indicators of styrene exposure. *Work Environ Health* 11:162-169.
- *Härkönen H, Lindstrom K, Seppäläinen AM, et al. 1978. Exposure-response relationship between styrene exposure and central nervous functions. *Scand J Work Environ Health* 4:53-59.
- *Harkov R, Gianti SJ, Bozzelli JW, et al. 1985. Monitoring volatile organic compounds at hazardous and sanitary landfills in New Jersey. *J Environ Sci Health A20*:491-501.
- Harkov R, Kebbekus B, Bozzelli JW, et al. 1983. Measurement of selected volatile organic compounds at three locations in New Jersey during the summer season. *J Air Pollut Control Assoc* 33:1177-1183.
- Harkov R, Kebbekus B, Bozzelli JW, et al. 1984. Comparison of selected volatile organic compounds during the summer and winter at urban sites in New Jersey. *Sci Total Environ* 38:259-274.
- Harkov R, Kebbekus B, Bozzelli JW, et al. 1987. Volatile organic compounds at urban sites in New Jersey. In: Lioy P, Daisey J, eds. *Toxic air pollutants*. Chelsea, MI: Lewis Publishers, Inc., 69-85.
- Harley RA, Cass GR. 1994. Modeling the concentrations of gas-phase toxic organic air pollutants: Direct emissions and atmospheric formation. *Environ Sci Technol* 28:88-98.
- Harley RA, Cass GR. 1995. Modeling the atmospheric concentrations of individual volatile organic compounds. *Atmos Environ* 29(8):905-922.
- *Hartmans S. 1995. Microbial degradation of styrene. In: Singh VP, ed. *Biotransformations: Microbial degradation of health risk compounds*. Amsterdam: Elsevier Science, 227-238.
- Hartmans S, Smits JP, VanderWerf MJ, et al. 1989. Metabolism of styrene oxide and 2-phenylethanol in the styrene-degrading xanthobacter strain 124x. *Appl Environ Microbiol* 55(11):2850-2855.
- Harvilchuck JA, Carlson GP. 2006. Comparison of styrene and its metabolites styrene oxide and 4-vinylphenol on cytotoxicity and glutathione depletion in Clara cells of mice and rats. *Toxicology* 227(1-2):165-172.
- *Hassett JJ, Banwart WL, Griffin RA. 1983. Correlation of compound properties with sorption characteristics of nonpolar compounds by soils and sediments: Concepts and limitations. In: Francis CW, Auerbach SI, eds. *Environmental and solid wastes: Characterization, treatment, and disposal*. Boston, MA: Butterworths, 161-176.
- Haufroid V, Jakubowski M, Janasik B, et al. 2002. Interest of genotyping and phenotyping of drug-metabolizing enzymes for the interpretation of biological monitoring of exposure to styrene. *Pharmacogenetics* 12(9):691-702.
- Hayes JP, Lambourn L, Hopkirk JAC, et al. 1991. Occupational asthma due to styrene. *Thorax* 46(5):396-397.
- Hays MD, Geron CD, Linna KJ, et al. 2002. Speciation of gas-phase and fine particle emissions from burning of foliar fuels. *Environ Sci Technol* 36(11):2281-2295.

9. REFERENCES

- *HazDat. 2007. Styrene. HazDat Database: ATSDR's Hazardous Substance Release and Health Effects Database. Atlanta, GA: Agency for Toxic Substances and Disease Registry. <http://www.atsdr.cdc.gov/hazdat.html>. May 21, 2007.
- Heavner DL, Morgan WT, Ogden MW. 1995. Determination of volatile organic compounds and its apportionment in 49 homes. *Environ Int* 21:3-21.
- Heavner DL, Morgan WT, Ogden MW. 1996. Determination of volatile organic compounds and respirable suspended particulate matter in New Jersey and Pennsylvania homes and workplaces. *Environ Int* 22:159-183.
- Hemminki K, Vainio H. 1984. Human exposure to potentially carcinogenic compounds. *IARC Sci Publ* (59):37-45.
- +*Hemminki K, Franssila E, Vainio H. 1980. Spontaneous abortions among female chemical workers in Finland. *Int Arch Occup Environ Health* 45:123-126.
- *Hemminki K, Lindbohm ML, Hemminki T, et al. 1984. Reproductive hazards and plastics industry. In: Jarvisalo J, Pfaffli P, Vainio H, eds. *Industrial hazards of plastics and synthetic elastomers*. New York, NY: Alan R. Liss, Inc., 79-87.
- Henderson LM, Speit G. 2005. Review of the genotoxicity of styrene in humans. *Mutat Res* 589(3):158-191.
- *Heylin M, ed. 1989. Production by the U.S. chemical industry: Growth maintains a strong pace. *Chem Eng News* 67:38-45.
- Hirasawa F, Kawagoe M, Arany S, et al. 2005. Styrene monomer primarily induces CYP2B1 mRNA in rat liver. *Xenobiotica* 35(12):1089-1099.
- *Hodgson AT, Rudd AF, Beal D, et al. 2000. Volatile organic compound concentrations and emission rates in new manufactures and site-built houses. *Indoor Air* 10:178-192.
- Hodgson AT, Wooley JD, Daisey JM. 1993. Emissions of volatile organic compounds from new carpet measured in a large scale environmental chamber. *J Air Waste Manage Assoc* 43:316-324.
- +*Hodgson JT, Jones RD. 1985. Mortality of styrene production, polymerization and processing workers at a site in northwest England. *Scand J Work Environ Health* 11:347-352.
- *Hoel DG, Davis DL, Miller AB, et al. 1992. Trends in cancer mortality in 15 industrialized countries, 1969-1986. *J Natl Cancer Inst* 84(5):313-320.
- Hoffmann K, Krause C, Seifert B, et al. 2000. The German Environmental Survey 1990/92 (GERES II). Sources of personal exposure to volatile organic compounds. *J Expo Anal Environ Epidemiol* 10:115-125.
- Hofmann C, Putz C, Semder B, et al. 2006. Styrene-7,8-oxide burden in ventilated, perfused lungs of mice and rats exposed to vaporous styrene. *Toxicol Sci* 90(1):39-48.

9. REFERENCES

- *Hogstedt B, Akesson B, Axell K, et al. 1983. Increased frequency of lymphocyte micronuclei in workers producing reinforced polyester resin with low exposure to styrene. *Scand J Work Environ Health* 9:241-246.
- *Hogstedt B, Hender K, Mark-Vendel E, et al. 1979. Increased frequency of chromosome aberrations in workers exposed to styrene. *Scand J Work Environ Health* 5:333-335.
- Holmberg PC. 1977. Central nervous defects in two children of mothers exposed to chemicals in the reinforced plastics industry. *Scand J Work Environ Health* 3:212-214.
- Horvath AL. 1982. Halogenated hydrocarbons: Solubility - miscibility with water. New York, NY: Marcel Dekker, Inc., 324-325.
- Horvath E, Pongracz K, Rappaport S, et al. 1994. ³²P-Post-labeling detection of DNA adducts in mononuclear cells of workers occupationally exposed to styrene. *Carcinogenesis* 15(7):1309-1315.
- +*Hotz P, Guillemin MP, Lob M. 1980. Study of some hepatic effects (induction and toxicity) caused by occupational exposure to styrene in the polyester industry. *Scand J Work Environ Health* 6:206-215.
- Hou CT, Patel R, Laskin AI, et al. 1983. Epoxidation of short-chain alkenes by resting-cell suspensions of propane-grown bacteria. *Appl Environ Microbiol* 46:171-177.
- *HSDB. 2007. Styrene. Hazardous Substances Data Bank. National Library of Medicine. <http://toxnet.nlm.nih.gov>. August 30, 2007.
- Hsieh CC, Tsai JH. 2003. VOC concentration characteristics in Southern Taiwan. *Chemosphere* 50:545-556.
- +*Husain R, Srivastava SP, Mushtaq M, et al. 1980. Effect of styrene on levels of serotonin, noradrenaline, dopamine and activity of acetyl cholinesterase and monoamine oxidase in rat brain. *Toxicol Lett* 7:47-50.
- +*Husain R, Srivastava SP, Seth PK. 1985. Some behavioral effects of early styrene intoxication in experimental animals. *Arch Toxicol* 57:53-55.
- *Hynes DE, DeNicola DB, Carlson GP. 1999. Metabolism of styrene by mouse and rat isolated lung cells. *Toxicol Sci* 51:195-201.
- *IARC. 1979. Styrene, polystyrene and styrene-butadiene copolymers. IARC monographs on the evaluation of the carcinogenic risk of chemicals to humans. Vol. 19. Some monomers, plastics and synthetic elastomers, and acrolein. Geneva, Switzerland: International Agency for Research on Cancer, World Health Organization, 231-274.
- IARC. 1985. IARC monographs on the evaluation of the carcinogenic risk of chemicals to humans. Vol. 36. Allyl compounds, aldehydes, epoxies and peroxides. Lyon, France: World Health Organization.
- IARC. 1987. IARC monographs on the evaluation of carcinogenic risks to humans. Supplement No. 7. Lyon, France: World Health Organization, 72, 345-347.
- IARC. 1988. IARC monographs on the evaluation of carcinogenic risk of chemicals to humans. Vol. 42. Silica and some silicates. Lyon, France: World Health Organization.

9. REFERENCES

- *IARC. 1994. IARC monographs on the evaluation of the carcinogenic risk of chemicals to humans. Lyon, France: International Agency for Research on Cancer, 233-320.
- *IARC. 2002. IARC monographs on the evaluation of carcinogenic risk to humans. Vol. 82. Some traditional herbal medicines, some mycotoxins, naphthalene and styrene. Lyon, France: International Agency for Research on Cancer, 437-550.
- *IARC. 2006. Agents Reviewed by the IARC Monographs. Volumes 1–96. Lyon, France: International Agency for Research on Cancer. <http://monographs.iarc.fr/ENG/Classification/index.php>. April 11, 2007.
- *Ikeda M, Hirayama T. 1978. Possible metabolic interaction of styrene with organic solvents. *Scand J Work Environ Health* 4:41-46.
- *Ikeda M, Ohtsuji H, Imamura T. 1972. *In vivo* suppression of benzene and styrene oxidation by co-administered toluene in rats and effects of phenobarbital. *Xenobiotica* 2:101-106.
- Imbriani M, Maestri L, Marraccini P, et al. 2002. Urinary determination of N-acetyl-S-(N-methylcarbamoyl)cysteine and N-methylformamide in workers exposed to N,N-dimethylformamide. *Int Arch Occup Environ Health* 75:445-452.
- +*Iregren A, Johnson AC, Nylen P. 2005. Low-level styrene exposure and color vision in Swedish styrene workers. *Environ Toxicol Pharmacol* 20(3):506.
- IRIS. 1991. Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency.
- *IRIS. 2007. Styrene. Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/index.html>. May 15, 2007.
- Ishidate M, Yoshikawa K. 1980. Chromosome aberration tests with Chinese hamster cells *in vitro* with and without metabolic activation - a comparative study on mutagens and carcinogens. *Arch Toxicol Suppl* 4:41-44.
- *Islam MA, Stancheva E. 1999. Volatile organic compounds in the ambient air of Rousse, Bulgaria. *Water Air Soil Pollut* 115:309-320.
- *Jablonicka A, Karellova J, Polakova H, et al. 1988. Analysis of chromosomes in peripheral blood lymphocytes of styrene-exposed workers. *Mutat Res* 206:167-169.
- Jaeger RJ, Conolly RB, Murphy SD. 1974. Toxicity and biochemical changes in rats after inhalation exposure to 1,1-dichloroethylene, bromobenzene, styrene, acrylonitrile, or 2-chlorobutadiene [Abstract]. *Toxicol Appl Pharmacol* 29:81.
- *James DH, Castor WM. 2005. Styrene. In: Ullman's encyclopedia of industrial chemistry. http://www.mrw.interscience.wiley.com/emrw/9783527306732/ueic/article/a25_329/current/pdf. May 30, 2007.

9. REFERENCES

- Jang JY, Droz PO. 1997. Ethnic differences in biological monitoring of several organic solvents II. A simulation study with a physiologically based pharmacokinetic model. *Int Arch Occup Environ Health* 70:41-50.
- Jang JY, Droz PO, Berode M. 1997. Ethnic differences in biological monitoring of several organic solvents. I. Human exposure experiment. *Int Arch Occup Environ Health* 69:343-349.
- *Jantunen K, Maki-Paakkanen J, Norppa H. 1986. Induction of chromosome aberrations by styrene and vinylacetate in cultured human lymphocytes: Dependence on erythrocytes. *Mutat Res* 159:109-116.
- *Jarry H, Metten M, Gamer AO, et al. 2002. Effects of 5-day styrene inhalation on serum prolactin and dopamine levels and on hypothalamic and striatal catecholamine concentrations in male rats. *Arch Toxicol* 76:657-663.
- +*Jegaden D, Amann D, Simon JF, et al. 1993. Study of the neurobehavioural toxicity of styrene at low levels of exposure. *Int Arch Occup Environ Health* 64(7):527-531.
- +*Jersey GM, Balmer J, Quast J, et al. 1978. Two-year chronic inhalation toxicity and carcinogenicity study on monomeric styrene in rats - final report. Dow Chemical Company. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8E. OTS8EHQ-0692-4867.
- *Johanson CE. 1980. Permeability and vascularity of the developing brain: Cerebellum vs. cerebral cortex. *Brain Res* 190(1):3-16.
- Johanson G, Naslund PH. 1988. Spreadsheet programming-a new approach to physiologically based modelling of solvent toxicokinetics. *Toxicol Lett* 41:115-127.
- *Johanson G, Ernstgard L, Gullstrand E, et al. 2000. Styrene oxide in blood, hemoglobin adducts, and urinary metabolites in human volunteers exposed to ¹³C₈-styrene vapors. *Toxicol Appl Pharmacol* 168:36-49.
- *Jonsson F, Johanson G. 2002. Physiologically based modeling of the inhalation kinetics of styrene in humans using a Bayesian population approach. *Toxicol Appl Pharmacol* 179:35-49.
- Jung J, Sofer S, Lakhwala F. 1993. A comparison study on biodegradation of selected monomers: Effect of molecular structure. *Haz Waste Haz Mater* 10(1):3-11.
- *Junk GA, Ford CS. 1980. A review of organic emissions from selected combustion processes. *Chemosphere* 9:187-230.
- *Kagi N, Fujii S, Horiba Y, et al. 2007. Indoor air quality for chemical and ultrafine particle contaminants from printers. *Build Environ* 42(5):1949-1954.
- +*Kankaanpää JT, Eovaara E, Hemminki K, et al. 1980. The effect of maternally inhaled styrene on embryonal and foetal development in mice and Chinese hamsters. *Acta Pharmacol Toxicol* 47:127-129.
- *Karakaya AE, Karahalil B, Yilmazer M, et al. 1997. Evaluation of genotoxic potential of styrene in furniture workers using unsaturated polyester resins. *Mutat Res* 392:261-268.
- +*Katakura Y, Kishi R, Ikeda T, et al. 1999. Effects of prenatal exposure to styrene on neurochemical levels in rat brain. *Toxicol Lett* 105:239-249.

9. REFERENCES

- +*Katakura Y, Kishi R, Ikeda T, et al. 2001. Effects of prenatal styrene exposure on postnatal development and brain serotonin and catecholamine levels in rats. *Environ Res* 85(1):41-47.
- +Kato T, Higashi K, Inoue N. 1989. Subchronic effects of styrene and styrene oxide on lipid peroxidation and the metabolism of glutathione in rat liver and brain. *J Toxicol Sci* 14:1-9.
- Kaufmann W, Mellert W, Van Ravenzwaay B, et al. 2005. Effects of styrene and its metabolites on different lung compartments of the mouse-cell proliferation and histomorphology. *Regul Toxicol Pharmacol* 42:24-36.
- Kawai T, Mizunuma K, Yasugi T, et al. 1995. Effects of methanol on styrene metabolism among workers occupationally exposed at low concentrations. *Arch Environ Contam Toxicol* 28:543-546.
- Kawata K, Tanabe A, Saito S, et al. 1997. Screening of volatile organic compounds in river sediment. *Bull Environ Contam Toxicol* 58:893-900.
- *Keenan C, Harriman E. 1993. Styrene use in Massachusetts. Lowell, MA: University of Massachusetts. Toxic Use Reduction Institute.
- Kelly TJ, Mukund R, Spicer CW, et al. 1994. Concentrations and transformations of hazardous air pollutants. *Environ Sci Technol* 28:378A-387A.
- *Kenaga EE. 1980. Predicted bioconcentration factors and soil sorption coefficients of pesticides and other chemicals. *Ecotoxicol Environ Safety* 4:26-38.
- *Kenaga EE, Goring CAI. 1980. Relationship between water solubility, soil sorption, octanol-water partitioning, and concentration of chemicals in biota. *ASTM Spec Tech Publ* 707:78-115.
- *Kessler W, Jiang X, Filser JG. 1990. Direct determination of styrene-7,8-oxide in blood by gas chromatography with flame ionization detector. *J Chromatog* 534:67-75.
- Khanna VK, Husain R, Hanig JP, et al. 1991. Increased neurobehavioral toxicity of styrene in protein-malnourished rats. *Neurobehav Toxicol Teratol* 13:153-159.
- +*Khanna VK, Husain R, Seth PK. 1994. Effect of protein malnutrition on the neurobehavioural toxicity of styrene in young rats. *J Appl Toxicol* 14(5):351-356.
- *Kim H, Wang RS, Elovaara E, et al. 1997. Cytochrome P450 isozymes responsible for the metabolism of toluene and styrene in human liver microsomes. *Xenobiotica* 27(7):657-665.
- Kim YM, Harrad S, Harrison RM. 2001. Concentrations and sources of VOCs in urban domestic and public microenvironments. *Environ Sci Technol* 35:997-1004.
- Kim YM, Harrad S, Harrison RM. 2002. Levels and sources of personal inhalation exposure to volatile organic compounds. *Environ Sci Technol* 36:5405-5410.
- King JW. 1989. Fundamentals and applications of supercritical fluid extraction in chromatographic science. *J Chromatogr Sci* 27:355-364.

9. REFERENCES

- *King L, Sherbin G. 1986. Point sources of toxic organics to the upper St. Clair River. *Water Poll Res J Canada* 21:433-446.
- *Kinlin TE, Muralidhara R, Pittet AO, et al. 1972. Volatile components of roasted filberts. *J Agric Food Chem* 20:1021-1028.
- *Kinney PL, Chillrud SN, Ramstrom S, et al. 2002. Exposures to multiple air toxics in New York City. *Environ Health Perspect* 110(Suppl 4):539-546.
- *Kirk-Othmer. 2001. Styrene. In: Kirk-Othmer encyclopedia of chemical technology. John Wiley & Sons, Inc.
<http://www.mrw.interscience.wiley.com/emrw/9780471238966/kirk/article/styrchen.a01/current/pdf>.
May 30, 2007.
- +Kishi R, Chen BQ, Katakura Y, et al. 1995. Effect of prenatal exposure to styrene on the neurobehavioral development, activity, motor coordination, and learning behavior of rats. *Neurobehav Toxicol Teratol* 17(2):121-130.
- ++Kishi R, Eguchi T, Yuasa J, et al. 2001. Effects of low-level occupational exposure to styrene on color vision: Dose relation with a urinary metabolite. *Environ Res* 85(1):25-30.
- Kishi R, Harabuchi I, Katakura Y, et al. 1993. Neurobehavioral effects of chronic occupational exposure to organic solvents among Japanese industrial painters. *Environ Res* 62:303-313.
- +Kishi R, Katakura Y, Ikeda T, et al. 1992. Neurochemical effects in rats following gestational exposure to styrene. *Toxicol Lett* 63:141-146.
- Kishi R, Sata F, Katakura Y, et al. 2005. Effects of pregnancy, age and sex in the metabolism of styrene in rat liver in relation to the regulation of cytochrome P450 enzymes. *J Occup Health* 47:49-55.
- Kishi R, Tozaki S, Gong YY. 2000a. Impairment of neurobehavioral function and color vision loss among workers exposed to low concentration of styrene: A review of literatures. *Ind Health* 38(2):120-126.
- Kishi R, Tozaki S, Katakura Y, et al. 2000b. Contrast vision dysfunction among styrene-exposed workers [Abstract]. *Neurotoxicology* 21(5):890.
- Kivisto H, Pekari K, Aitio A. 1993. Analysis and stability of phenylglyoxylic and mandelic acids in the urine of styrene-exposed people. *Int Arch Occup Environ Health* 64:399-403.
- Kleindienst TE, Shepson PB, Edney EO. 1986. Wood smoke: Measurement of the mutagenic activities of its gas- and particulate-phase photooxidation products. *Environ Sci Technol* 20:493-501.
- *Kleopfer RD, Fairless BJ. 1972. Characterization of organic components in a municipal water supply. *Environ Sci Technol* 6:1036-1037.
- *Kligerman AD, Allen JW, Bryant MF, et al. 1992. Cytogenetic studies of mice exposed to styrene by inhalation. *Mutat Res* 280:35-43.
- Kligerman AD, Allen JW, Erexson GL, et al. 1993. Cytogenetic studies of rodents exposed to styrene by inhalation. *IARC Sci Publ* (127):217-224.

9. REFERENCES

- *Kneip TJ, Crable JV. 1988a. Styrene in expired air. In: *Methods for biological monitoring*. Washington, DC: American Public Health Association, 457-462.
- *Kneip TJ, Crable JV. 1988b. Mandelic acid in urine. In: *Methods for biological monitoring*. Washington, DC: American Public Health Association, 387-391.
- *Kneip TJ, Crable JV. 1988c. Phenylglyoxylic acid in urine. In: *Methods for biological monitoring*. Washington, DC: American Public Health Association, 393-399.
- Knudsen LE, Sorsa M. 1993. Human biological monitoring of occupational genotoxic exposures. *Pharmacol Toxicol* 72(1):86-92.
- *Kogevinas M, Ferro G, Andersen A, et al. 1994. Cancer mortality in a historical cohort study of workers exposed to styrene. *Scand J Work Environ Health* 50:251-261.
- *Kogevinas M, Ferro G, Saracci R, et al. 1993. Cancer mortality in an international cohort of workers exposed to styrene. *IARC Sci Publ* (127):289-300.
- +*Kohn AN. 1978. Ocular toxicity of styrene. *Am J Ophthalmol* 85:569-570.
- Kohn J, Minotti S, Durham H. 1995. Assessment of the neurotoxicity of styrene, styrene oxide, and styrene glycol in primary cultures of motor and sensory neurons. *Toxicol Lett* 75:29-37.
- *Kolstad HA, Bisanti L, Roeleveld N, et al. 1999c. Time to pregnancy for men occupationally exposed to styrene in several European reinforced plastics companies. *Scand J Work Environ Health* 25:66-69.
- *Kolstad HA, Bisanti L, Roeleveld N, et al. 2000. Time to pregnancy among male workers of the reinforced plastics industry in Denmark, Italy and the Netherlands. *Scand J Work Environ Health* 26(4):353-358.
- *Kolstad HA, Bonde JP, Spano M, et al. 1999a. Change in semen quality and sperm chromatin structure following occupational styrene exposure. *Int Arch Occup Environ Health* 72:135-141.
- *Kolstad HA, Bonde JPE, Spano M, et al. 1999b. Sperm chromatin structure and semen quality following occupational styrene exposure. *Scand J Work Environ Health* 25:70-73.
- *Kolstad HA, Juel K, Olsen J, et al. 1995. Exposure to styrene and chronic health effects: Mortality and incidence of solid cancers in the Danish reinforced plastics industry. *Occup Environ Med* 52:320-327.
- *Kolstad HA, Lynge E, Olsen J. 1993. Cancer incidence in the Danish reinforced plastics industry. *IARC Sci Publ* (127):301-308.
- *Kolstad HA, Lynge E, Olsen J, et al. 1994. Incidence of lymphohematopoietic malignancies among styrene-exposed workers of the reinforced plastics industry. *Scand J Work Environ Health* 20:272-278.
- Kolstad HA, Pedersen B, Olsen J, et al. 1996. Clonal chromosome aberrations in myeloid leukemia. *Scand J Work Environ Health* 22(1):58-61.
- *Komori M, Nishio K, Kitada M, et al. 1990. Fetus-specific expression of a form of cytochrome P-450 in human livers. *Biochemistry* 29(18):4430-4433.

9. REFERENCES

- *Konasewich D, Traversy W, Zar H. 1978. Status report on organic and heavy metal contaminants in the Lakes Erie, Michigan, Huron and Superior basins. Great Lakes Water Quality Board.
- *Kool HJ, van Kreijl CF, Zoeteman BC. 1982. Toxicology assessment of organic compounds in drinking water. *Crit Rev Environ Control* 12:307-357.
- Kopczynski SL, Lonneman WA, Sutterfield FD, et al. 1972. Photochemistry of atmospheric samples in Los Angeles. *Environ Sci Technol* 6:342-347.
- Korn M, Gfrorer W, Filser JG, et al. 1994a. Styrene-7,8-oxide in blood of workers exposed to styrene (Erratum in: *Arch Toxicol* 69(1):72). *Arch Toxicol* 68(8):524-527.
- Korn M, Gfrorer W, Filser JG, et al. 1994b. Erratum: Styrene-7,8-oxide in blood of workers exposed to styrene (Erratum to: *Arch Toxicol* 68:524-527). *Arch Toxicol* 69(1):72.
- *Korn M, Wodarz R, Drysch K, et al. 1987. Stereometabolism of styrene in man. Urinary excretion of chiral styrene metabolites. *Arch Toxicol* 60:86-88.
- *Korn M, Wodarz R, Schoknecht W, et al. 1984. Styrene metabolism in man: Gas chromatographic separation of mandelic acid enantiomers in the urine of exposed persons. *Arch Toxicol* 55:59-63.
- Koskinen M, Vodicka P, Hemminki K. 2001. Identification of 1-adenine DNA adducts in workers occupationally exposed to styrene. *J Occup Environ Med* 43(8):694-700.
- *Krill RM, Sonzogni WC. 1986. Chemical monitoring of Wisconsin's groundwater. *J Am Water Works Assoc* 78:70-75.
- *Krishnan K, Andersen ME. 1994. Physiologically based pharmacokinetic modeling in toxicology. In: Hayes AW, eds. *Principles and methods of toxicology*. 3rd ed. New York, NY: Raven Press, Ltd., 149-188.
- *Krishnan K, Anderson ME, Clewell HJ, et al. 1994. Physiologically based pharmacokinetic modeling of chemical mixtures. In: Yang RSH, eds. *Toxicology of chemical mixtures*. New York, NY: Academic Press, 399-437.
- Kubo T, Urano K, Utsumi H. 2002. Mutagenicity characteristics of 255 environmental chemicals. *J Health Sci* 48(6):545-554.
- +*Kulig BM. 1988. The neurobehavioral effects of chronic styrene exposure in the rat. *Neurotoxicol Teratol* 10:511-517.
- Kurppa K, Holmberg PC, Hernberg S, et al. 1983. Screening for occupational exposures and congenital malformations: Preliminary results from a nationwide case-referent study. *Scand J Work Environ Health* 9:89-93.
- Laffon B, Lema M, Mendez J. 2001. Simultaneous high-performance liquid chromatographic determination of urinary mandelic and phenylglyoxylic acids as indirect evaluation of styrene exposure. *J Chromatogr B Biomed Sci Appl* 753(2):385-393.

9. REFERENCES

- *Laffon B, Pasaro E, Mendez J. 2002. Evaluation of genotoxic effects in a group of workers exposed to low levels of styrene. *Toxicology* 171(2-3):175-186.
- Laffon B, Perez-Cadahia B, Pasaro E, et al. 2003. Individual sensitivity to DNA damage induced by styrene *in vitro*: Influence of cytochrome P450, epoxide hydrolase and glutathione *S*-transferase genotypes. *Toxicology* 186(1-2):131-141.
- *Laib RJ, Tucholski M, Filser JG, et al. 1992. Pharmacokinetic interaction between 1,3-butadiene and styrene in Sprague-Dawley rats. *Arch Toxicol* 66(5):310-314.
- Landrigan PJ. 1993. Critical assessment of epidemiological studies on the carcinogenicity of 1,3-butadiene and styrene. *IARC Sci Publ* (127):375-388.
- *Langvardt PW, Nolan RJ. 1991. Determination of styrene 7,8-oxide in whole rat blood by gas chromatography-mass spectrometry. *J Chromatog* 567:93-103.
- LaRegina J, Bozzelli JW. 1986. Volatile organic compounds at hazardous waste sites and a sanitary landfill in New Jersey: An up-to-date review of the present situation. *Environ Prog* 5:18-27.
- +*Lataye R, Campo P, Barthelemy C, et al. 2001. Cochlear pathology induced by styrene. *Neurotoxicol Teratol* 23(1):71-79.
- +*Lataye R, Campo P, Loquet G. 2000. Combined effects of noise and styrene exposure on hearing function in the rat. *Hear Res* 139:86-96.
- +*Lataye R, Campo P, Pouyatos B, et al. 2003. Solvent ototoxicity in the rat and guinea pig. *Neurotoxicol Teratol* 25(1):39-50.
- Latriano L, Wazneh L, Dong Z, et al. 1991. Exposure to styrene: Comparison of DNA and haemoglobin adducts as biomarkers. In: Garner RC, Farmer PB, Steel GT, et al., eds. *Human carcinogen exposure*. Oxford: IRL Press at Oxford University Press, 305-314.
- Lauwerys RR. 1986. Occupational toxicology. In: Klaassen CD, Amdur MO, Doull J, eds. *Casarett and Doull's toxicology. The basic science of poisons*. New York, NY: Macmillan Publishing Company, 902-916.
- Lawton BW, Hoffmann J, Triebig G. 2006. The ototoxicity of styrene: A review of occupational investigations. *Int Arch Occup Environ Health* 79:93-102.
- *Leavens TL, Bond JA. 1996. Pharmacokinetic model describing the disposition of butadiene and styrene in mice. *Toxicology* 113(1-3):310-313.
- *Leavens TL, Farris GM, James RA, et al. 1997. Genotoxicity and cytotoxicity in male B6C3F1 mice following exposure to mixtures of 1,3-butadiene and styrene. *Environ Mol Mutagen* 29:335-345.
- Leavens TL, Moss OR, Turner MJ, et al. 1996. Metabolic interactions of 1,3-butadiene and styrene in male B6C3F1 mice. *Toxicol Appl Pharmacol* 141(2):628-636.
- Lee SH, Norppa H. 1995. Effects of indomethacin and arachidonic acid on sister chromatid exchange induction by styrene and styrene-7,8-oxide. *Mutat Res* 348(4):175-181.

9. REFERENCES

- Lee YK, Kim MC, Whang KJ. 1989. Simultaneous determination of natural and styrene-butadiene rubber tire tread particles in atmospheric dusts by pyrolysis-gas chromatography. *J Anal Appl Pyrolysis* 16:49-56.
- *Leeder JS, Kearns GL. 1997. Pharmacogenetics in pediatrics: Implications for practice. *Pediatr Clin North Am* 44(1):55-77.
- Lees PS, Stefaniak A, Emmett EA, et al. 2003. Exposure assessment for study of olfactory function in workers exposed to styrene in the reinforced-plastics industry. *Am J Ind Med* 44(1):12-23.
- Legraverend C, Elovaara E, Zitting A, et al. 1984. DNA binding of [¹⁴C] styrene in isolated rat hepatocytes. *Chem Biol Interact* 51:91-102.
- Lehmann E, Weidlich U, Gmehling J. 1986. Survey on organic solvents in various products and methods for estimating workplace exposures. *Prog Clin Biol Res* 220:31-41.
- *Leibman KC. 1975. Metabolism and toxicity of styrene. *Environ Health Perspect* 11:115-119.
- *Lemasters GK, Carson A, Samuels SJ. 1985. Occupational styrene exposure for twelve product categories in the reinforced-plastics industry. *Am Ind Hyg Assoc J* 46:434-441.
- +*Lemasters GK, Samuels SJ, Morrison JA, et al. 1989. Reproductive outcomes of pregnant workers employed at 36 reinforced plastics companies. II. Lowered birth weight. *J Occup Med* 31:115-120.
- Lenvik K, Osvoll PO, Woldbaek T. 1999. Occupational exposure to styrene in Norway, 1972-1996. *Appl Occup Environ Hyg* 14:165-170.
- *Leovic KW, Sheldon LS, Whitaker DA, et al. 1996. Measurement of indoor air emissions from dry-process photocopier machines. *J Air Waste Manage Assoc* 46:821-829.
- *Leovic KW, Whitaker D, Norheim C, et al. 1998. Evaluation of a test method for measuring indoor air emissions from dry-process photocopiers. *J Air Waste Manage Assoc* 48:915-923.
- Lesage S, McBride RA, Cureton PM, et al. 1993. Fate of organic solvents in landfill leachates under simulated field conditions and in anaerobic microcosms. *Waste Manage Res* 11:215-226.
- *Leung HW. 1993. Physiologically-based pharmacokinetic modelling. In: Ballantyne B, Marro T, Turner P, eds. *General and applied toxicology*. Vol. 1.1 New York, NY: Stockton Press, 153-164.
- *Lide CR. 2005. Styrene. In: *CRC Handbook of chemistry and physics*. 86th ed. Boca Raton, FL: Taylor & Francis, 3-464.
- Lijinsky W. 1986. Rat and mouse forestomach tumors induced by chronic oral administration of styrene oxide. *J Natl Cancer Inst* 77:471-476.
- Limasset JC, Simon P, Poirot P, et al. 1999. Estimation of the percutaneous absorption of styrene in an industrial situation. *Int Arch Occup Environ Health* 72:46-51.
- Lindbohm ML. 1993. Effects of styrene on the reproductive health of women: A review. *IARC Sci Publ* (127):163-169.

9. REFERENCES

Lindbohm ML. 1995. Effects of parental exposure to solvents on pregnancy outcome. *J Occup Environ Med* 17(8):908-914.

+*Lindbohm ML, Hemminki K, Kyyronen P. 1985. Spontaneous abortions among women employed in the plastics industry. *Am J Ind Med* 8:579-586.

+*Lindstrom K, Harkonen H, Hernberg S. 1976. Disturbances in psychological functions of workers occupationally exposed to styrene. *Scand J Work Environ Health* 3:129-139.

Lindstrom K, Harkonen H, Mantere P. 1978. Alcohol consumption and tolerance of workers exposed to styrene in relation to level of exposure and psychological symptoms and signs. *Scand J Work Environ Health* 4:196-199.

Linhart I, Gut I, Smejkal J, et al. 2000. Biotransformation of styrene in mice. Stereochemical aspects. *Chem Res Toxicol* 13:36-44.

Linhart I, Smejkal J, Mladkova I. 1998. Stereochemical aspects of styrene biotransformation. *Toxicol Lett* 94:127-135.

Linhart I, Weidenhoffer Z, Mraz J, et al. 1997. The evidence for conjugated mandelic and phenylglyoxylic acids in the urine of rats dosed with styrene. *Toxicol Lett* 90:199-205.

Lioy PJ, Daisey JM, Atherholt T, et al. 1983. The New Jersey project on Airborne Toxic Elements and Organic Substances (ATEOS): A summary of the 1981 summer and 1982 winter studies. *J Air Pollut Control Assoc* 33:649-657.

Little JC, Hodgson AT, Gadgil AJ. 1994. Modeling emissions of volatile organic compounds from new carpets. *Atmos Environ* 28:227-234.

*Livingston AL. 1978. Forage plant estrogens. *J Toxicol Environ Health* 4(2-3):301-324.

*Lof A, Johanson G. 1993. Dose-dependent kinetics of inhaled styrene in man. *IARC Sci Publ* (127):89-99.

*Lof A, Gullstrand E, Byfalt Nordqvist M. 1983. Tissue distribution of styrene, styrene glycol and more polar styrene metabolites in the mouse. *Scand J Work Environ Health* 9:419-430.

*Lof A, Gullstrand E, Lundgren E, et al. 1984. Occurrence of styrene-7,8-oxide and styrene glycol in mouse after the administration of styrene. *Scand J Work Environ Health* 10:179-187.

Lof A, Lundgren E, Byfalt Nordqvist M. 1986b. Kinetics of styrene in workers from a plastics industry after controlled exposure: A comparison with subjects not previously exposed. *Br J Ind Med* 43:537-543.

*Lof A, Lundgren E, Nydahl EM, et al. 1986a. Biological monitoring of styrene metabolites in blood. *Scand J W Environ Health* 12:70-74.

+*Loquet G, Campo P, Lataye R. 1999. Comparison of toluene-induced and styrene-induced hearing losses. *Neurotoxicology* 21(6):689-697.

9. REFERENCES

- +*Loquet G, Campo P, Lataye R, et al. 2000. Combined effects of exposure to styrene and ethanol on the auditory function in the rat. *Hear Res* 148(1-2):173-180.
- +*Lorimer WV, Lilis R, Fischbein A, et al. 1978. Health status of styrene-polystyrene polymerization workers. *Scand J Work Environ Health* 4:220-226.
- Lorimer WV, Lilis R, Nicholson WJ, et al. 1976. Clinical studies of styrene workers: Initial findings. *Environ Health Perspect* 17:171-181.
- Loughlin JE, Rothman KJ, Dreyer NA. 1999. Lymphatic and haematopoietic cancer mortality in a population attending school adjacent to styrene-butadiene facilities, 1963-1999. *J Epidemiol Community Health* 53:283-287.
- *Lovegren NV, Fisher GS, Legendre MG, et al. 1979. Volatile constituents of dried legumes. *J Agric Food Chem* 27:851-853.
- Luderer U, Collins TF, Daston GP, et al. 2006a. NTP-CERHR Expert Panel Report on the reproductive and developmental toxicity of styrene (Erratum in: *Birth Defects Res B Dev Reprod Toxicol* 77:250). *Birth Defects Res B Dev Reprod Toxicol* 77(2):110-193.
- Luderer U, Collins TF, Daston GP, et al. 2006b. Corrigendum to NTP-CERHR expert panel report on the reproductive and developmental toxicity of styrene (Erratum to: *Birth Defects Res B Dev Reprod Toxicol* 77:110-193). *Birth Defects Res B Dev Reprod Toxicol* 77(3):250.
- *Luderer U, Tornero-Velez R, Shay T, et al. 2004. Temporal association between serum prolactin concentration and exposure to styrene. *Occup Environ Med* 61(4):325-333.
- Ma M, Umemura T, Mori Y, et al. 2005. Influence of genetic polymorphisms of styrene-metabolizing enzymes and smoking habits on levels of urinary metabolites after occupational exposure to styrene. *Toxicol Lett* 160:84-91.
- *Macaluso M, Larson R, Delzell E, et al. 1996. Leukemia and cumulative exposure to butadiene, styrene and benzene among workers in the synthetic rubber industry. *Toxicology* 113:190-202.
- *Mackay CJ, Kelman GR. 1986. Choice reaction time in workers exposed to styrene vapour. *Human Toxicol* 5:85-89.
- *MacLeod H, Jourdain JL, Poulet G, et al. 1984. Kinetic study of reactions of some organic sulfur compounds with OH radicals. *Atmos Environ* 18:2621-2626.
- Madden S, Roper C, Greenough R. 2004. *In vitro* assessment of styrene absorption through human skin [Abstract]. *Toxicologist* 78(1-5):330-331.
- *Maestri L, Ghittori S, Imbriani M. 1997a. Determination of specific mercapturic acids as an index of exposure to environmental benzene, toluene, and styrene. *Ind Health* 35:489-501.
- Maestri L, Imbriani M, Ghittori S, et al. 1997b. Excretion of *N*-acetyl-S-(1-phenyl-2-hydroxyethyl)-cysteine and *N*-acetyl-S-(2-phenyl-2-hydroxyethyl)-cysteine in workers exposed to styrene. *Sci Total Environ* 199:13-22.

9. REFERENCES

- Mahler JF, Price HC, O'Connor RW, et al. 1999. Characterization of hepatocellular resistance and susceptibility to styrene toxicity in B6C3F1 mice. *Toxicol Sci* 48:123-133.
- *Maki-Paakkanen J. 1987. Chromosome aberrations, micronuclei and sister-chromatid exchanges in blood lymphocytes after occupational exposure to low levels of styrene. *Mutat Res* 189:39-406.
- *Maki-Paakkanen J, Walles S, Osterman-Golkar S, et al. 1991. Single-strand breaks, chromosome aberrations, sister-chromatid exchanges, and micronuclei in blood lymphocytes of workers exposed to styrene during the production of reinforced plastics. *Environ Mol Mutagen* 17(1):27-31.
- +*Makitie A, Pirvola U, Pyykko I, et al. 2002. Functional and morphological effects of styrene on the auditory system of the rat. *Arch Toxicol* 76:40-47.
- Makitie A, Pirvola U, Pyykko I, et al. 2003. The ototoxic interaction of styrene and noise. *Hear Res* 179(1-2):9-20.
- Male R, Lillehaug JR, Djurhuus R, et al. 1985. *In vitro* transformation and tumor promotion studies of styrene and styrene oxide. *Carcinogenesis* 6:1367-1370.
- Malorni W, Formisano G, Iosi F. 1988. *In vitro* effects of styrene on cytoskeletal apparatus: An immunocytochemical study. *Drug Chem Toxicol* 11:419-431.
- +*Maltoni C, Ciliberti A, Carretti D. 1982. Experimental contributions in identifying brain potential carcinogens in the petrochemical industry. *Ann NY Acad Sci* 381:216-249.
- *Maltoni C, Failla G, Kassapidis G. 1979. First experimental demonstration of the carcinogenic effects of styrene oxide. *Med Lav* 5:358-362.
- Manini P, Andreoli R, Poll D, et al. 2002. Liquid chromatography/electrospray tandem mass spectrometry characterization of styrene metabolism in man and in rat. *Rapid Commun Mass Spectrom* 16:2239-2248.
- Manini P, Buzio L, Andreoli R, et al. 2003. Assessment of biotransformation of the arene moiety of styrene in volunteers and occupationally exposed workers. *Toxicol Appl Pharmacol* 189:160-169.
- Manini P, De Palma G, Mozzoni P, et al. 2002. GSTM1 polymorphism and styrene metabolism: Insights from an acute accidental exposure. *Toxicol Lett* 134:201-208.
- Marczynski B, Peel M, Bauer X. 2000. New aspects in genotoxic risk assessment of styrene exposure - a working hypothesis. *Med Hypotheses* 54(4):619-623.
- Marczynski B, Rozynek P, Elliehausen H, et al. 1996. DNA-fragmentation changes in white blood cells of workers occupationally exposed to styrene. *Naunyn-Schmiedeberg's Arch Pharmacol* 353:R136.
- Marczynski B, Rozynek P, Elliehausen H, et al. 1997. Detection of 8-hydroxydeoxyguanosine, a marker of oxidative DNA damage, in white blood cells of workers occupationally exposed to styrene. *Arch Toxicol* 71:496-500.
- Marhuenda D, Prieto MJ, Periago JF, et al. 1997. Biological monitoring of styrene exposure and possible interference of acetone co-exposure. *Int Arch Occup Environ Health* 69:455-460.

9. REFERENCES

- Mastrangelo G, Fedeli U, Priolo G, et al. 2005. Lung cancer risk in the vinyl chloride industry. Letter to editor. *Cancer Causes Control* 16:189-190.
- +*Matanoski G, Schwartz L. 1987. Mortality of workers in styrene-butadiene polymer production. *J Occup Med* 29:675-680.
- Matanoski G, Tao XG. 2003. Styrene exposure and ischemic heart disease: A case-cohort study. *Am J Epidemiol* 158(10):988-995.
- *Matanoski G, Elliott E, Tao X, et al. 1997. Lymphohematopoietic cancers and butadiene and styrene exposure in synthetic rubber manufacture. *Ann N Y Acad Sci* 837:157-169.
- *Matanoski G, Francis M, Correa-Villasenor A, et al. 1993. Cancer epidemiology among styrene-butadiene rubber workers. *IARC Sci Publ* (127):363-374.
- *Matanoski G, Santos-Burgoa C, Schwartz L. 1990. Mortality of a cohort of workers in the styrene-butadiene polymer manufacturing industry. *Environ Health Perspect* 86:107-117.
- *Matikainen E, Forsman-Gronholm L, Pfaffli P, et al. 1993a. Nervous system effects of occupational exposure to styrene: A clinical and neurophysiological study. *Environ Res* 61(1):84-92.
- *Matikainen E, Forsman-Gronholm L, Pfaffli P, et al. 1993b. Neurotoxicity in workers exposed to styrene. *IARC Sci Publ* (127):153-161.
- Matsuoka A, Hayashi M, Ishidate M. 1979. Chromosomal aberration tests on 29 chemicals combined with 59 mix *in vitro*. *Mutat Res* 66:277-290.
- Matthiessen P, Thani JE, Law RJ, et al. 1993. Attempts to assess the environmental hazard posed by complex mixtures of organic chemicals in UK estuaries. *Mar Pollut Bull* 26(2):90-95.
- *Mattie DR, Bates GD, Jepson GW, et al. 1994. Determination of skin: Air partition coefficients for volatile chemicals: Experimental method and applications. *Fundam Appl Toxicol* 22:51-57.
- *Mayr U, Butsch A, Schneider S. 1992. Validation of two *in vitro* test systems for estrogenic activities with zearalenone, phytoestrogens and cereal extracts. *Toxicology* 74(2-3):135-149.
- McConnell EE, Swenberg JA. 1993. Styrene and styrene oxide: Results of studies on carcinogenicity in experimental animals. *IARC Sci Publ* (127):323-333.
- McConnell EE, Swenberg JA. 1994. Review of styrene and styrene oxide long-term animal studies. *Crit Rev Toxicol* 24(S1):S49-S55.
- *McDonald AD, Lavoie J, Côté R. 1988. Spontaneous abortion in women employed in plastics manufacture. *Am J Ind Med* 14:9-14.
- McMichael AJ. 1976. Standardized mortality ratios and the "healthy worker effect": Scratching beneath the surface. *J Occup Med* 18:165-168.
- *McMichael AJ, Spirtas R, Gamble JF, et al. 1976. Mortality among rubber workers: relationship to specific jobs. *J Occup Med* 18:178-185.

9. REFERENCES

- +*Meinhardt TJ, Lemen RA, Crandall MS, et al. 1982. Environmental epidemiologic investigation of the styrene-butadiene rubber industry: Mortality patterns with discussion of the hematopoietic and lymphatic malignancies. *Scand J Work Environ Health* 8:250-259.
- *Mendrala AL, Langvardt PW, Nitschke KD, et al. 1991. *In vitro* metabolism of styrene and styrene oxide in human and animal tissues. Midland, MI: Dow Chemical Company, 1-63.
- *Meretoja T, Jarventaus H, Sorsa M, et al. 1978. Chromosome aberrations in lymphocytes of workers exposed to styrene. *Scand J Work Environ Health* 4:259-264.
- *Meretoja T, Vainio H, Sorsa M, et al. 1977. Occupational styrene exposure and chromosomal aberrations. *Mutat Res* 56:193-197.
- Mergler D, Huel G, Belanger S, et al. 1996. Surveillance of early neurotoxic dysfunction. *Neurotoxicology* 17(3-4):803-812.
- Michael LC, Pellizzari ED, Wiseman RW. 1988. Development and evaluation of a procedure for determining volatile organics in water. *Environ Sci Technol* 22:565-570.
- Mierauskiene JR, Lekevicius RK. 1985. Cytogenic studies of workers occupationally exposed to phenol, styrene and formaldehyde. *Mutat Res* 147:308-309.
- *Miermans CJH, van der Velde LE, Frintrop PCM. 2000. Analysis of volatile organic compounds, using the purge and trap injector coupled to a gas chromatograph/ion-trap mass spectrometer: Review of the results in Dutch surface water of the Rhine, Meuse, Northern Delta area and Westerscheldt over the period 1992-1997. *Chemosphere* 40:39-48.
- Migliore L, Naccarati A, Scarpato R. 2001. Effect of occupational exposure to styrene in somatic and germ cells of male workers. *Mutat Res* 483(Suppl 1):S174.
- Migliore L, Naccarati A, Zanello A, et al. 2002. Assessment of sperm DNA integrity in workers exposed to styrene. *Hum Reprod* 17(11):2912-2918.
- Miller CA, Srivastava RK, Ryan JV. 1994. Emissions of organic hazardous air pollutants from the combustion of pulverized coal in a small-scale combustor. *Environ Sci Technol* 28:1150-1158.
- Misumi J, Nagano M, Zhao W, et al. 2000. Neurophysiological changes in rats subchronically treated with styrene or its metabolites. *J Occup Health* 42:328-335.
- Mizunuma K, Yasugi T, Kawai T, et al. 1993. Exposure-excretion relationship of styrene and acetone in factory workers: A comparison of a lipophilic solvent and a hydrophilic solvent. *Arch Environ Contam Toxicol* 25(1):129-133.
- Mizutani T, Irie Y, Nakanishi K. 1994. Styrene-induced hepatotoxicity in mice depleted of glutathione. *Res Commun Mol Pathol Pharmacol* 86(3):361-374.
- Moen BE, Hollund BE. 2000. Exposure to organic solvents among car painters in Bergen, Norway. *Ann Occup Hyg* 44(3):185-189.
- Mohamed MF, Kang D, Aneja VP. 2002. Volatile organic compounds in some urban locations in United States. *Chemosphere* 47:863-882.

9. REFERENCES

- +*Möller C, Ödkvist L, Larsby B, et al. 1990. Otoneurological findings in workers exposed to styrene. *Scand J Work Environ Health* 16:189-194.
- +*Morata TC, Johnson AC, Nysten P, et al. 2002. Audiometric findings in workers exposed to low levels of styrene and noise. *J Occup Environ Med* 44(9):806-814.
- *Morgan DL, Cooper SW, Carlock DL, et al. 1991. Dermal absorption of neat and aqueous volatile organic chemicals in the Fischer 344 rat. *Environ Res* 55:51-63.
- +*Morgan DL, Mahler JF, Dill JA, et al. 1993a. Styrene inhalation toxicity studies in mice: II. Sex differences in susceptibility of B6C3F1 mice. *Fundam Appl Toxicol* 21:317-325.
- +*Morgan DL, Mahler JF, Dill JA, et al. 1993b. Styrene inhalation toxicity studies in mice: III. Strain differences in susceptibility. *Fundam Appl Toxicol* 21:326-333.
- Morgan DL, Mahler JF, Moorman MP, et al. 1995. Comparison of styrene hepatotoxicity in B6C3F1 and Swiss mice. *Fundam Appl Toxicol* 27(2):217-222.
- +*Morgan DL, Mahler JF, O'Connor RW, et al. 1993c. Styrene inhalation toxicity studies in mice: I. Hepatotoxicity in B6C3F1 mice. *Fundam Appl Toxicol* 20:325-335.
- Morgan DL, Mahler JF, Wilson RE, et al. 1997. Effects of various pretreatments on the hepatotoxicity of inhaled styrene in the B6C3F1 mouse. *Xenobiotica* 27(4):401-411.
- +*Morioka I, Kuroda M, Miyashita K, et al. 1999. Evaluation of organic solvent ototoxicity by the upper limit of hearing. *Arch Environ Health* 54(5):341-346.
- *Morris JB. 2000. Uptake of styrene in the upper respiratory tract of the CD mouse and Sprague-Dawley rat. *Toxicol Sci* 54:222-228.
- *Morselli PL, Franco-Morselli R, Bossi L. 1980. Clinical pharmacokinetics in newborns and infants: Age-related differences and therapeutic implications. *Clin Pharmacokinet* 5(6):485-527.
- Moscato G, Biscaldi GP, Brunetti G, et al. 1984. Occupational asthma caused by styrene. *J Occup Med* 26:552.
- Moscato G, Biscaldi G, Cottica D, et al. 1987. Occupational asthma due to styrene: Two case reports. *J Occup Med* 29:957-960.
- Mueller M, Klein W. 1993. Fugacity calculations using estimated physico-chemical properties. *SAR QSAR Environ Res* 1:235-262.
- +*Muijser H, Hoogendijk EM, Hooisma J. 1988. The effects of occupational exposure to styrene on high-frequency hearing thresholds. *Toxicology* 49:331-340.
- +*Murata K, Araki S, Yokoyama K. 1991. Assessment of the peripheral, central, and autonomic nervous system function in styrene workers. *Am J Ind Med* 20:775-784.

9. REFERENCES

- Murer AJL, Christensen JM, Midtgaard T, et al. 1994. Determination of the urinary metabolites of styrene: Estimation of the method evaluation function and evaluation of reference values in Danish subjects. *Int Arch Occup Environ Health* 65:313-318.
- +*Murray FJ, John JA, Balmer MF, et al. 1978. Teratologic evaluation of styrene given to rats and rabbits by inhalation or by gavage. *Toxicology* 11:335-343.
- Mutti A, Smargiassi A. 1998. Selective vulnerability of dopaminergic systems to industrial chemicals: Risk assessment of related neuroendocrine changes. *Toxicol Ind Health* 14(1-2):311-323.
- Mutti A, Coccini T, Alinovi R, et al. 1999. Exposure to hydrocarbons and renal disease: An experimental animal model. *Ren Fail* 21(3&4):369-385.
- Mutti A, Falzoi M, Romanelli A, et al. 1984c. Regional alterations of brain catecholamine by styrene exposure in rabbits. *Arch Toxicol* 55:173-177.
- +*Mutti A, Mazzucchi A, Rustichelli P, et al. 1984a. Exposure-effect and exposure-response relationships between occupational exposure to styrene and neuropsychological functions. *Am J Ind Med* 5:275-286.
- Mutti A, Romanelli A, Falzoi M, et al. 1985. Styrene metabolism and striatal dopamine depletion in rabbits. *Arch Toxicol Suppl* 8:447-450.
- +*Mutti A, Vescovi PP, Falzoi M, et al. 1984b. Neuroendocrine effects of styrene on occupationally exposed workers. *Scand J Work Environ Health* 10:225-228.
- Muttray A, Jung D, Konietzko J. 1992. Acquired dyschromatopsia: Some comments concerning the procedure of testing and interpretation [including response from original authors Gallassi et al.]. (Comment on: *J Occup Med* 33(7):761-765). *J Occup Med* 34(5):563-564.
- *Nakajima T, Elovaara E, Gonzalez FJ, et al. 1993. Characterization of the human cytochrome P450 isozymes responsible for styrene metabolism. *IARC Sci Publ* (127):101-108.
- *Nakajima T, Wang R-S, Elovaara E, et al. 1994. CYP2C11 and CYP2B1 are major cytochrome P450 forms involved in styrene oxidation in liver and lung microsomes from untreated rats, respectively. *Biochem Pharmacol* 48(4):637-642.
- Nano R, Rossi A, Fenoglio C, et al. 2000. Evaluation of a possible styrene-induced damage to the haematopoietic tissues in the rat. *Anticancer Res* 20(3A):1615-1619.
- *NAS/NRC. 1989. Report of the oversight committee. Biologic markers in reproductive toxicology. Washington, DC: National Academy of Sciences, National Research Council, National Academy Press, 15-35.
- Nasterlack M, Triebig G. 1994. Increase in neuropsychiatric symptoms after occupational exposure to low levels of styrene (Comment on: *Br J Ind Med* 50(9):843-850). *Occup Environ Med* 51(4):286-287.
- NATICH. 1989. National Air-Toxics Information Clearinghouse: NATICH data base report on state, local and EPA air toxics activities. July 1989. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards. EPA45038929.

9. REFERENCES

- NCHS. 1988. National Health and Nutrition Examination Survey III. Appendix C. Examination content. Hyattsville, MD: U.S. Department of Health and Human Services, National Center for Health Statistics.
- NCI. 1979a. Bioassay of a solution of β -nitrostyrene and styrene for possible carcinogenicity. Bethesda, MD: National Cancer Institute, Division of Cancer Cause and Prevention. DHEW Publication No. (NIH) 79-1726.
- +*NCI. 1979b. Bioassay of styrene for possible carcinogenicity. Bethesda, MD: National Cancer Institute, Division of Cancer Cause and Prevention. Technical Report Series No. 185. NCI-CG-TR-185. NIH 79-1741. PB300977.
- Negri S, Maestri L, Andreoli R, et al. 2006. Mercapturic acids of styrene in man: Comparability of the results obtained by LC/MS/MS and by HPLC-fluorimeter, and stability of samples under different storage conditions. *Toxicol Lett* 162:225-233.
- Nestmann ER, Lynch BSW, Ratpan F. 2003. Evaluation of the genotoxic potential of styrene [Abstract]. *Environ Mol Mutagen* 41(3):195.
- Nestmann ER, Lynch BS, Ratpan F. 2005. Perspectives on the genotoxic risk of styrene. *J Toxicol Environ Health B Crit Rev* 8:95-107.
- Newhook R, Caldwell I. 1993. Exposure to styrene in the general Canadian population. *IARC Sci Publ* (127):27-33.
- Nicholson WJ, Tarr D. 1984. Occupational hazards in production and processing of styrene polymers-epidemiologic findings. *Prog Clin Biol Res* 141:263-277.
- +*Nicholson WJ, Selikoff IJ, Seidman H. 1978. Mortality experience of styrene-polystyrene polymerization workers: Initial findings. *Scand J Work Environ Health* 4:247-252.
- *Niklasson M, Tham R, Larsby B, et al. 1993. Effects of toluene, styrene, trichloroethylene, and trichloroethane on the vestibulo- and opto-oculo motor system in rats. *Neurotoxicol Teratol* 15:327-334.
- *NIOSH. 1983. Criteria for a recommended standard: Occupational exposure to styrene. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health. PB84148295.
- *NIOSH. 1984. Method 1501: Hydrocarbons, aromatic. In: NIOSH manual of analytical methods. Vol. 2. Cincinnati, OH: U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health.
- NIOSH. 1985. Pocket guide to chemical hazards. Washington, DC: U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health. 85-114.
- NIOSH. 1988. NIOSH recommendations for occupational safety and health standards. National Institute for Occupational Safety and Health. *MMWR* 37:S-7.

9. REFERENCES

- *NIOSH. 1990. Styrene. Estimated numbers of employees potentially exposed to specific agents by 2-digit Standard Industrial Classification (SIC). National Occupational Exposure Survey conducted from 1981-1983. Centers for Disease Control. National Institute for Occupational Safety and Health. <http://www.cdc.gov/noes/>. August 09, 2007.
- *NIOSH. 2005. NIOSH pocket guide to chemical hazards. Atlanta, GA: National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention. <http://www.cdc.gov/niosh/npg/>. May 15, 2007.
- *NLM. 1989. Chemline. Bethesda, MD: National Library of Medicine. September 5, 1989.
- NOHS. 1989. National Occupational Hazard Survey. Cincinnati, OH: National Institute for Occupational Safety and Health. October 18, 1989.
- *Nordenson I, Beckman L. 1984. Chromosomal aberrations in lymphocytes of workers exposed to low levels of styrene. *Hum Hered* 34:178-182.
- Nordqvist MB, Lof A, Osteman-Golkar S, et al. 1985. Covalent binding of styrene and styrene-7,8-oxide to plasma proteins, hemoglobin and DNA in the mouse. *Chem Biol Interact* 55:63-73.
- *Normandeau J, Chakrabarti S, Brodeur J. 1984. Influence of simultaneous exposure to acrylonitrile and styrene on the toxicity and metabolism of styrene in rats. *Toxicol Appl Pharmacol* 75:346-349.
- *Norppa H. 1981. Styrene and vinyltoluene induce micronuclei in mouse bone marrow. *Toxicol Lett* 8:247-251.
- Norppa H, Sorsa M. 1993. Genetic toxicity of 1,3-butadiene and styrene. *IARC Sci Publ* (127):185-193.
- Norppa H, Jaerventausta H, Kubiak R, et al. 1991. Chromosome aberrations, sister-chromatid exchanges, and micronuclei in blood lymphocytes of Finnish reinforced plastics workers exposed to styrene. *Environ Mol Mutagen* 17:56.
- *Norppa H, Maki-Paakkanen J, Jantunen K, et al. 1988. Mutagenicity studies on styrene and vinyl acetate. *Ann NY Acad Sci* 534:671-678.
- *Norppa H, Sorsa M, Pfaffli P, et al. 1979. Effects of styrene oxide on chromosome aberrations, sister chromatid exchange and hepatic drug biotransformation in Chinese hamsters *in vivo*. *Chem Biol Interact* 26:305-315.
- *Norppa H, Sorsa M, Pfaffli P, et al. 1980a. Styrene and styrene oxide induce SCEs and are metabolized in human lymphocyte cultures. *Carcinogenesis* 1:357-361.
- *Norppa H, Sorsa M, Vainio H. 1980b. Chromosomal aberrations in bone marrow of Chinese hamsters exposed to styrene and ethanol. *Toxicol Lett* 5:241-244.
- *Norppa H, Vainio H, Sorsa M. 1981. Chromosome aberrations in lymphocytes of workers exposed to styrene. *Am J Ind Med* 2:299-304.
- *Norppa H, Vainio H, Sorsa M. 1983. Metabolic activation of styrene by erythrocytes detected as increased sister-chromatid exchanges in cultured human lymphocytes. *Cancer Res* 43:3579-3582.

9. REFERENCES

- *Norppa H, Vainio H, Sorsa M, et al. 1984. Erythrocyte-dependent metabolic activation of styrene and induction of sister-chromatid exchange in cultured human lymphocytes. *Arch Toxicol Suppl* 7:286-290.
- *Norström A, Lof A, Aringer L, et al. 1992. Determination of N-acetyl-S-(2-phenyl-2-hydroxyethyl) cysteine in human urine after experimental exposure to styrene. *Chemosphere* 24(11):1553-1561.
- *NRC. 1993. Pesticides in the diets of infants and children. Washington, DC: National Research Council. National Academy Press. PB93216091.
- *NTP. 2005. Introduction. Report on carcinogens. 11th ed. Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service, National Toxicology Program. <http://ntp.niehs.nih.gov/ntp/roc/eleventh/intro.pdf>. May 23, 2007.
- *NTP. 2006. NTP-CERHR monograph on the potential human reproductive and developmental effects of styrene. National Toxicology Program. NIH Publication No. 06-4475.
- Nunez CM, Ramsey GH, Bahner MA, Clayton CA. 1999. An empirical model to predict styrene emissions from fiber-reinforced plastics fabrication processes. *J Air Waste Manage Assoc* 49:1168-1178.
- Oberheitmann B, Frentzel-Beyme R, Hoffmann W. 2001. An application of the challenge assay in boat builders exposed to low levels of styrene- a feasibility study of a possible biomarker for acquired susceptibility. *Int J Hyg Environ Health* 204(1):23-29.
- +*Ödkvist LM, Edling C, Hellquist H, et al. 1985. Influence of vapours on the nasal mucosa among industry workers. *Rhinology* 23:121-127.
- +*Ödkvist LM, Larsby B, Tham R, et al. 1982. Vestibulo-oculomotor disturbances in humans exposed to styrene. *Acta Otolaryngol* 94:487-493.
- Ogata M, Taguchi T. 1987. Quantitation of urinary metabolites of toluene, xylene, styrene, ethylbenzene, benzene and phenol by automated high performance liquid chromatography. *Int Arch Occup Environ Health* 59:263-272.
- Ogata M, Taguchi T. 1988. Simultaneous determination of urinary creatinine and metabolites of toluene, xylene, styrene, ethylbenzene, benzene and phenol by automated high performance liquid chromatography. *Int Arch Occup Environ Health* 61:131-140.
- *Ogata M, Fujisawa K, Ogino Y, et al. 1984. Partition coefficients as a measure of bioconcentration potential of crude oil compounds in fish and shellfish. *Bull Environ Contam Toxicol* 33:561-567.
- +*Ohashi Y, Nakai Y, Ikeoka H, et al. 1986. Degeneration and regeneration of respiratory mucosa of rats after exposure to styrene. *J Appl Toxicol* 6:405-412.
- *Ohtani H, Ichikawa Y, Iwamoto E, et al. 2001. Effects of styrene monomer and trimer on gonadal sex differentiation of genetic males of the frog *Rana rugosa*. *Environ Res* 87(3):175-180.
- Ohtsuji M, Ikeda J. 1971. Metabolism of styrene in the rat and the stimulatory effect of phenobarbital. *Toxicol Appl Pharmacol* 18:321-328.
- +*Okun AH, Beaumont JJ, Meinhardt TJ, et al. 1985. Mortality patterns among styrene-exposed boatbuilders. *Am J Ind Med* 8:193-205.

9. REFERENCES

- *O'Neil MJ, Smith A, Heckelman PE, et al., eds. 2001. Styrene. In: The Merck index. An encyclopedia of chemicals, drugs, and biologicals. 13th ed. Whitehouse Station, NJ: Merck & Co., Inc., 1578.
- Oner F, Mungan D, Numanoglu N, et al. 2004. Occupational asthma in the furniture industry: Is it due to styrene? *Respiration* 71(4):336-341.
- *Ong CN, Shi CY, Chia SE, et al. 1994. Biological monitoring of exposure to low concentrations of styrene. *Am J Ind Med* 25(5):719-730.
- ORS. 1989. Massachusetts drinking water standards and guidelines. Boston, MA: Executive Office of Environmental Affairs, Department of Environmental Quality Engineering, Office of Research and Standards.
- OSHA. 1989. Occupational Safety and Health Administration: Part III. *Federal Register* 54:2952.
- *OSHA. 2006a. Air contaminants. Occupational safety and health standards for shipyard employment. Occupational Safety and Health Administration. Code of Federal Regulations. 29 CFR 1915.1000. <http://www.osha.gov/comp-links.html>. April 11, 2007.
- *OSHA. 2006b. Gases, vapors, fumes, dusts, and mists. Safety and health regulations for construction. Occupational Safety and Health Administration. Code of Federal Regulations. 29 CFR 1926.55, Appendix A. <http://www.osha.gov/comp-links.html>. April 11, 2007.
- *OSHA. 2006c. Toxic and Hazardous Substances. Occupational safety and health standards. Occupational Safety and Health Administration. Code of Federal Regulations. 29 CFR 1910.1000, Table Z-2. <http://www.osha.gov/comp-links.html>. May 15, 2007.
- Osswald P, Baveye P, Block J. 1996. Bacterial influence on partitioning rate during the biodegradation of styrene in a biphasic aqueous-organic system. *Biodegradation* 7:297-302.
- Osterman-Golkar S, Christakopoulos A, Zorcec V, et al. 1995. Dosimetry of styrene 7,8-oxide in styrene- and styrene oxide-exposed mice and rats by quantification of haemoglobin. *Chem Biol Interact* 95:79-87.
- OTA. 1990. Neurotoxicology: Identifying and controlling poisons of the nervous system. Washington, DC: Office of Technology Assessment. OTABA438.
- *Otson R. 1987. Purgeable organics in Great Lakes raw and treated water. *Int J Environ Anal Chem* 31:41-53.
- Otson R. 1996. Vapor-phase organic compounds in Canadian residences. In: Wang W, Schnoor J, Doi J, eds. Volatile organic compounds in the environment. American Society for Testing and Materials, 66-76.
- Otson R, Chan C. 1987. Sample handling and analysis for 51 volatile organics by an adapted purge and trap GC-MS technique. *Int J Environ Anal Chem* 30:275-287.
- Otson R, Fellin P. 1992. Volatile organics in the indoor environment: Sources and occurrence. In: Nriagu JO, ed. Gaseous pollutants: Characterization and cycling. John Wiley & Sons, Inc., 335-421.

9. REFERENCES

- Ott WR, Roberts JW. 1998. Everyday exposure to toxic pollutants. *Sci Am* 278:86-91.
- +*Ott MG, Kolesar RC, Schamweber HC, et al. 1980. A mortality survey of employees engaged in the development or manufacture of styrene-based products. *J Occup Med* 22:445-460.
- Otteneder M, Lutz U, Lutz WK. 2002. DNA adducts of styrene-7,8-oxide in target and non-target organs for tumor induction in rat and mouse after repeated inhalation exposure to styrene. *Mutat Res* 500(1-2):111-116.
- *Owen GM, Brozek J. 1966. Influence of age, sex and nutrition on body composition during childhood and adolescence. In: Falkner F, ed., eds. *Human development*. Philadelphia, PA: WB Saunders, 222-238.
- Pacifici GM, Guthenberg C, Warholm M, et al. 1988. Conjugation of styrene oxide by the basic and acidic forms of glutathione transferase in the human fetal liver. *Dev Pharmacol Ther* 11:243-251.
- Pacifici GM, Warholm M, Guthenberg C, et al. 1987. Detoxification of styrene oxide by human liver glutathione transferase. *Hum Toxicol* 6:483-490.
- *Pahren HR, Bloodgood DE. 1961. Biological oxidation of several vinyl compounds. *J Water Pollut Control Fed* 33:233-238.
- Pankow JF, Rosen ME. 1988. Determination of volatile compounds in water by purging directly to a capillary column with whole column cryotrapping. *Environ Sci Technol* 22:398-405.
- Paramei GV, Meyer-Baron M, Seeber A. 2004. Impairments of colour vision induced by organic solvents: A meta-analysis study. *Neurotoxicology* 25(5):803-816.
- *Parkki MG. 1978. The role of glutathione in the toxicity of styrene. *Scand J Work Environ Health* 4(Suppl 2):53-59.
- Pauwels W, Severi M, Veulemans H. 1996. Determination of macromolecular adducts as biomarkers for styrene exposure [Abstract]. *Mutat Res* 360(3):245.
- Pauwels W, Vodiceka P, Severi M, et al. 1996. Adduction formation on DNA and haemoglobin in mice intraperitoneally administered with styrene. *Carcinogenesis* 17(12):2673-2680.
- *Peck DP. 1994. Immunoassay for aromatic ring containing compounds. United States Patent no. 5,358,851. Date of patent June 06, 1991. <http://patft.uspto.gov/netacgi/nph-Parser?Sect1=PTO2&Sect2=HITOFF&p=1&u=%2Fmetahtml%2FPTO%2Fsearch-bool.html&r=7&f=G&l=50&co1=AND&d=PTXT&s1=5358851&OS=5358851&RS=5358851>. May 30, 2007.
- Pekari K, Nylander-French L, Pfäffli P, et al. Biological monitoring of exposure to styrene-Assessment of different approaches. *J Occup Med Toxicol* 2(2):115-126.
- *Pellizzari ED, Castillo NP, Willis S, et al. 1979. Identification of organic components in aqueous effluents from energy-related processes. *ASTM Spec Tech Publ* 686:256-274.

9. REFERENCES

- *Pellizzari ED, Hartwell TD, Harris BS, et al. 1982. Purgeable organic compounds in mother's milk. *Bull Environ Contam Toxicol* 28:322-328.
- Perbellini L, Mozzo P, Turri PV, et al. 1988. Biological exposure index of styrene suggested by a physiologico-mathematical model. *Int Arch Occup Environ Health* 60:187-193.
- *Pero RW, Bryngelsson T, Hogstedt B, et al. 1982. Occupational and *in vitro* exposure to styrene assessed by unscheduled DNA synthesis in resting human lymphocytes. *Carcinogenesis* 3:681-685.
- *Pezzagno G, Ghittori S, Imbriani M, et al. 1985. Urinary elimination of styrene in experimental and occupational exposure. *Scand J Work Environ Health* 11:371-379.
- Pfaffli P, Saamanen A. 1993. The occupational scene of styrene. *IARC Sci Publ* (127):15-26.
- *Pfaffli P, Hesso A, Vainio H, et al. 1981. 4-Vinylphenol excretion suggestive of arene oxide formation in workers occupationally exposed to styrene. *Toxicol Appl Pharmacol* 60:85-90.
- Phillips DH, Farmer PB. 1994. Evidence for DNA and protein binding by styrene and styrene oxide. *Crit Rev Toxicol* 24(Suppl 1):S35-S46.
- Piao M, Chu S, Zheng M, et al. 1999. Characterization of the combustion products of polyethylene. *Chemosphere* 39:1497-1512.
- Pierce CH, Tozer TN. 1992. Styrene in adipose tissue of nonoccupationally exposed persons. *Environ Res* 58:230-235.
- Pierce CH, Becker CE, Tozer TN, et al. 1998. Modeling the acute neurotoxicity of styrene. *J Occup Environ Med* 40(3):230-240.
- Pierce CH, Dills RL, Silvey GW, et al. 1996. Partition coefficients between human blood or adipose tissue and air for aromatic solvents. *Scand J Work Environ Health* 22:112-118.
- Pitzer KS, Guttman L, Westrum EF. 1946. The heat capacity, heats of fusion and vaporization, vapor pressure, entropy, vibration frequencies and barrier to internal rotation of styrene. *J Am Chem Soc* 68:2209-2212.
- *Plotnick HB, Weigel WW. 1979. Tissue distribution and excretion of ¹⁴C-styrene in male and female rats. *Res Commun Chem Pathol Pharmacol* 24:515-524.
- *Pohlova H, Rossner P, Sram RJ. 1985. Cytogenetic analysis of human peripheral blood lymphocytes in culture exposed *in vitro* to styrene and styrene oxide. *J Hyg Epidem Microbiol Immunol* 29:269-274.
- Poliakova OV, Lebedev AT, Hanninen O. 2000. Organic pollutants in snow of urban and rural Russia and Finland. *Toxicol Environ Chem* 75:181-194.
- +*Ponomarkov V, Tomatis L. 1978. Effects of long-term oral administration of styrene to mice and rats. *Scand J Work Environ Health* 4(Suppl 2):127-135.
- Ponomarkov V, Cabral JRP, Wahrendorf J, et al. 1984. A carcinogenicity study of styrene-7,8-oxide in rats. *Cancer Lett* 24:95-101.

9. REFERENCES

+*Pouyatos B, Campo P, Lataye R. 2002. Use of DPOAEs for assessing hearing loss caused by styrene in the rat. *Hear Res* 165(1-2):156-164.

Pouyatos B, Morel G, Lambert-Xolin AM, et al. 2004. Consequences of noise- or styrene-induced cochlear damages on glutamate decarboxylase levels in the rat inferior colliculus. *Hear Res* 189(1-2):83-91.

Pratt GC, McCourtney M, Wu CY, et al. 1998. Measurement and source apportionment of human exposures to toxic air pollutants in the Minneapolis-St. Paul metropolitan area. Measurement of toxic and related air pollutants. Sewickley, PA: Air and Waste Management Association, 64-72.

*Pratt GC, Palmer K, Wu CY, et al. 2000. An assessment of air toxics in Minnesota. *Environ Health Perspect* 108(9):815-825.

Preston RJ, Abernethy DJ. 1993. Studies of the induction of chromosomal aberration and sister chromatid exchange in rats exposed to styrene by inhalation. *IARC Sci Publ* (127):225-233.

*Prince MM, Stayner LT, Smith RJ, et al. 1997. A re-examination of risk estimates from the NIOSH Occupational Noise and Hearing Survey (OHNS). *J Acoust Soc Am* 101(2):950-963.

Prito MJ, Marhuenda D, Cardona A. 2002. Analysis of styrene and its metabolites in blood and urine of workers exposed to both styrene and acetone. *J Anal Toxicol* 26:23-28.

Proctor NH, Hughes JP, Fischman ML. 1988. Chemical hazards of the workplace. 2nd ed. Philadelphia, PA: J.B. Lippincott Company, 448-449.

+*Pryor GT, Rebert CS, Howd RA. 1987. Hearing loss in rats caused by inhalation of mixed xylenes and styrene. *J Appl Toxicol* 7:55-61.

+*Quast JF, Humiston CG, Kalnins RV, et al. 1979. Results of a toxicity study of monomeric styrene administered to beagle dogs by oral intubation for 19 months. Midland, MI: Toxicology Research Laboratory, 199.

Rabovsky J, Fowles J, Hill MD, et al. 2001. A health risk benchmark for the neurologic effects of styrene: Comparison with NOAEL/LOAEL approach. *Risk Anal* 21(1):117-126.

*Ramsey JC, Andersen ME. 1984. A physiologically based description of the inhalation pharmacokinetics of styrene in rats and humans. *Toxicol Appl Pharmacol* 73:159-175.

*Ramsey JC, Young JD. 1978. Pharmacokinetics of inhaled styrene in rats and humans. *Scand J Work Environ Health* 4(Suppl 2):84-91.

*Ramsey JC, Young JD, Karbowski RJ, et al. 1980. Pharmacokinetics of inhaled styrene in human volunteers. *Toxicol Appl Pharmacol* 53:54-63.

*Rappaport SM, Fraser DA. 1977. Air sampling and analysis in a rubber vulcanization area. *Am Ind Hyg Assoc J* 38:205-210.

Rappaport SM, Yager JW. 1996. Correspondence: Re: Papers published in *Mutation Research* [including response from original authors Scott and Preston] (Comment on: *Mutat Res* 318(3):175-203). *Mutat Res* 340(2-3):183-185.

9. REFERENCES

- Rappaport SM, Yeowell-O'Connor K. 1999. Protein adducts as dosimeters of human exposure to styrene, styrene-7,8-oxide, and benzene. *Toxicol Lett* 108:117-126.
- Rappaport SM, Dionne L, Woodlee J, et al. 1992. The relationship between exposure to styrene biomarkers of exposure and cytogenetic response. *Rev Epidemiol Sante Publique* 40(Suppl 1):S80-S81.
- Rappaport SM, Symanski E, Yager JW, et al. 1995. The relationship between environmental monitoring and biological markers in exposure assessment. *Environ Health Perspect* 103:49-53.
- Rappaport SM, Yeowell-O'Connell K, Bodell W, et al. 1996. An investigation of multiple biomarkers among workers exposed to styrene and styrene-7,8-oxide. *Cancer Res* 56:5410-5416.
- *Rathbun Re. 2000. Transport, behavior and fate of volatile organic compounds in streams. *Crit Rev Environ Sci* 30:129-295.
- Rebert CS, Hall TA. 1994. The neuroepidemiology of styrene: A critical review of representative literature. *Crit Rev Toxicol* 24(S1):S57-S106.
- Rebert CS, Schwartz RW, Svendsgaard DJ, et al. 1995. Combined effect of paired solvents on the rat's auditory system. *Toxicology* 105:345-354.
- *Regnaud L, Sirois G, Colin P, et al. 1987. Simultaneous ion-pairing liquid chromatographic determination of the major metabolites of styrene and carbamazepine and of unchanged carbamazepine in urine. *J Liq Chromatogr* 10(11):2369-2382.
- Rhouma KB, Chamkhia N, Sakly M. 2005. Toxicity of styrene in rat liver [Abstract]. *FASEB J* 19(4):A760.
- +*Riihimaki V, Pfaffli P. 1978. Percutaneous absorption of solvent vapors in man. *Scand J Work Environ Health* 4:73-85.
- *Roberts PV, McCarty PL, Reinhard M, et al. 1980. Organic contaminant behavior during groundwater recharge. *J Water Pollut Control Fed* 52(1):161-172.
- Roe FJC. 1994. Styrene: Toxicity studies-What do they show? *Crit Rev Toxicol* 24(S1):S117-S125.
- *Romieu I, Ramirez M, Meneses F, et al. 1999. Environmental exposure to volatile organic compounds among workers in Mexico City assessed by personal monitors and blood concentrations. *Environ Health Perspect* 107:511-515.
- Rorije E, Mueller M, Peijnenburg WJGM. 1997. Prediction of environmental degradation rates for high production volume chemicals (HPVC) using quantitative structure-activity relationships. Bilthoven: National Institute of Public Health and the Environment. RIVM report no: 719101030.
- +*Rosen I, Haeger-Aronsen B, Rehnstrom S, et al. 1978. Neurophysiological observations after chronic styrene exposure. *Scand J Work Environ Health* 4(Suppl. 2):184-194.
- Rosenbaum AS, Axelrod DA, Woodruff TJ, et al. 1999. National estimates of outdoor air toxics concentrations. *J Air Waste Manage Assoc* 49:1138-1152.

9. REFERENCES

- +*Rosengren LE, Haglid KG. 1989. Long term neurotoxicity of styrene. A quantitative study of glial fibrillary acidic protein (GFA) and S-100. *Br J Ind Med* 46:316-320.
- Rothberg M, Heloma A, Svinhufvud J, et al. 1998. Measurement and analysis of nicotine and other VOCs in indoor air as an indicator of passive smoking. *Ann Occup Hyg* 42(2):129-134.
- *Rothweiler H, Wager PA, Schlatter C. 1992. Volatile organic compounds and some very volatile organic compounds in new and recently renovated buildings in Switzerland. *Atmos Environ* 26(12):2219-2225.
- *Roy WR, Griffin RA. 1985. Mobility of organic solvents in water-saturated soil materials. *Environ Geol Water Sci* 7:241-247.
- Roycroft JH, TJ Mast, HA Ragan, et al. 1992. Toxicological effects of inhalation exposure to styrene in rats and mice. *Toxicologist* 12(1):397.
- *Ruder AM, Ward EM, Dong M, et al. 2004. Mortality patterns among workers exposed to styrene in the reinforced plastic boatbuilding industry: An update. *Am J Ind Med* 45(2):165-176.
- *Sabljić A, Gusten H, Hermens J. 1995. QSAR modelling of soil sorption. Improvements and systematics of log K_{oc} vs. log K_{ow} correlations. *Chemosphere* 31:4489-4514.
- Sack TM, Steele DH, Hammerstrom K. 1992. A survey of household products for volatile organic compounds. *Atmos Environ* 26A(6):1063-1070.
- *Sakaguchi J, Akabayashi S. 2003. Field survey of indoor air quality in detached houses in Niigata Prefecture. *Indoor Air* 13:42-49.
- *Sallmén M, Lindbohm M-L, Anttila A, et al. 1998. Time to pregnancy among the wives of men exposed to organic solvents. *Occup Environ Med* 55:24-30.
- +*Salomaa S, Donner M, Norppa H. 1985. Inactivity of styrene in the mouse sperm morphology test. *Toxicol Lett* 24:151-155.
- Salthammer T. 1997. Emission of volatile organic compounds from furniture coatings. *Indoor Air* 7:189-197.
- Samimi B, Falbo L. 1982. Monitoring of workers exposure to low levels of airborne monomers in a polystyrene production plant. *Am Ind Hyg Assoc J* 43:858-862.
- *Sanjivamurthy VA. 1978. Analysis of organics in Cleveland water supply. *Water Res* 12:31-33.
- Santos-Burgoa C, Matanoski GM. 1993. RE: "Lymphohematopoietic cancer in styrene-butadiene polymerization workers" (Comment on: *Am J Epidemiol* 36:843-854). *Am J Epidemiol* 138(9):765-768.
- *Santos-Burgoa C, Matanoski GM, Zeger S, et al. 1992. Lymphohematopoietic cancer in styrene-butadiene polymerization workers. *Am J Epidemiol* 136(7):843-854.
- *Sarangapani R, Teeguarden JG, Cruzan G, et al. 2002. Physiologically based pharmacokinetic modeling of styrene and styrene oxide respiratory-tract dosimetry in rodents and humans. *Inhal Toxicol* 14:789-834.

9. REFERENCES

- Sass-Korstak AM, Corey PN, Robertson JM. 1995. An investigation of the association between exposure to styrene and hearing loss. *Ann Epidemiol* 5(1):15-24.
- *Sathiakumar N, Graff J, Macaluso M, et al. 2005. An updated study of mortality among North American synthetic rubber industry workers. *Occup Environ Med* 62(12):822-829.
- *Sato A, Nakajima T. 1979. Partition coefficients of some aromatic hydrocarbons and ketones in water, blood and oil. *Br J Ind Med* 36:231-234.
- Savela K, Hemminki K. 1986. Reaction products of styrene oxide with deoxynucleosides and DNA *in vitro*. *Arch Toxicol Suppl* 9:281-285.
- *Savolainen H, Pfaffli P. 1978. Accumulation of styrene monomer and neurochemical effects of long-term inhalation exposure in rats. *Scand J Work Environ Health* 4:78-83.
- Savolainen H, Vaino H. 1977. Organ distribution and nervous system binding of styrene and styrene oxide. *Toxicology* 8:135-141.
- Sawyer RF. 1993. Trends in auto emissions and gasoline composition. *Environ Health Perspect* 101(Suppl 6):5-12.
- *Sax NI, Lewis RJ. 1987. *Hawley's condensed chemical dictionary*. 11th ed. New York: Van Nostrand Reinhold Company, 1101.
- *Sbrana I, Lascialfari D, Rossi AM, et al. 1983. Bone marrow cell chromosomal aberrations and styrene biotransformation in mice given styrene on a repeated oral schedule. *Chem Biol Interact* 45:349-357.
- Scapellato ML, Marcuzzo G, Mastrangelo G, et al. 1998. Environmental and biological monitoring of styrene exposure: Urinary excretion of D-glucaric acid compared with exposure indices. *J Occup Health* 40:313-318.
- Scelo G, Constantinescu V, Csiki I, et al. 2004. Occupational exposure to vinyl chloride, acrylonitrile and styrene and lung cancer risk (Europe). *Cancer Causes Control* 15(5):445-452.
- *Schaeffer VH, Bhooshan B, Chen S-B, et al. 1996. Characterization of volatile organic chemical emissions from carpet cushions. *J Air Waste Manage Assoc* 46:813-820.
- Schubert P, Rosenberg E, Grasserbauer M. 1999. Measurement of volatile organic compound (VOC) concentration profiles in the atmosphere of a remote forest region. *Int J Environ Anal Chem* 75(1-2):127-147.
- Scott D. 1993. Butadiene and styrene: Assessment of health hazards. *IARC Sci Publ* (127):275-286.
- Scott D, Preston RJ. 1993. A re-evaluation of the cytogenetic effects of styrene (Comment in: *Mutat Res* 340(2-3):183-185). *Mutat Res* 318(3):175-203.
- Scott D, Preston RJ. 1994. A critical review of the cytogenetic effects of styrene with an emphasis on human population monitoring: A synopsis. *Crit Rev Toxicol* 24(S1):S47-S48.

9. REFERENCES

- Scow KM, Johnson CR. 1997. Effect of sorption on biodegradation of soil pollutants. *Adv Agron* 58:1-56.
- *Sedivec V, Flek J, Mraz M. 1984. Urinary excretion of mandelic and phenylglyoxylic acids after human exposure to styrene vapour. *J Hyg Epid Microbiol Immunol* 28:241-255.
- +*Seeber A, Blaszkewicz M, Golka K, et al. 2004. Neurobehavioral effects of experimental exposures to low levels of styrene. *Toxicol Lett* 151:183-192.
- Seppäläinen AM. 1978. Neurotoxicity of styrene in occupational and experimental exposure. *Scand J Work Environ Health* 4(Suppl 2):181-183.
- +*Seppäläinen AM, Härkönen H. 1976. Neurophysiological findings among workers occupationally exposed to styrene. *Scand J Work Environ Health* 3:140-146.
- *Setchell BP, Waites GMH. 1975. The blood testis barrier. In: Creep RO, Astwood EB, Geiger SR, eds. *Handbook of physiology: Endocrinology V*. Washington, DC: American Physiological Society, 143-172.
- Severi M, Pauwels W, VanHummeien P, et al. 1994. Urinary mandelic acid and hemoglobin adducts in fiberglass-reinforced plastics workers exposed to styrene. *Scand J Work Environ Health* 20:451-458.
- *Sexton K, Adgate JL, Church TR, et al. 2005. Children's exposure to volatile organic compounds as determined by longitudinal measurements in blood. *Environ Health Perspect* 113(3):342-349.
- Sexton K, Adgate JL, Mongin SJ, et al. 2004. Evaluating differences between measured personal exposures to volatile organic compounds and concentrations in outdoor and indoor air. *Environ Sci Technol* 38:2593-2602.
- *Shamy MY, Osman HH, Kandeel KM, et al. 2002. DNA single strand breaks induced by low levels of occupational exposure to styrene: The gap between standards and reality. *J Environ Pathol Toxicol Oncol* 21(1):57-61.
- *Sheldon LS, Hites RA. 1978. Organic compounds in the Delaware River. *Environ Sci Tech* 12:1188-1194.
- Shen TT. 1982. Air quality assessment for land disposal of industrial wastes. *Environ Manage* 6:297-305.
- Shi CY, Chua SC, Lee BL, et al. 1994. Kinetics of styrene urinary metabolites: A study in a low-level occupational exposure setting in Singapore. *Int Arch Occup Environ Health* 65:319-323.
- *Shields HC, Weschler CJ. 1992. Volatile organic compounds measured at a telephone switching center form 5/30/85-12/6/88: A detailed case study. *J Air Waste Manage Assoc* 42:792-804.
- Shimkin MB, Stoner GD. 1975. Lung tumors in mice: Application to carcinogenesis bioassay.
- Shiu WY, Ma KC. 2000. Temperature dependence of physical-chemical properties of selected chemicals of environmental interest. I. Mononuclear and polynuclear aromatic hydrocarbons. *J Phys Chem Ref Data* 29(1):41-130.

9. REFERENCES

- +*Shugaev BB. 1969. Concentrations of hydrocarbons in tissues as a measure of toxicity. Arch Environ Health 18:878-882.
- *Sielicki M, Focht DD, Martin JP. 1978. Microbial transformations of styrene and [¹⁴C] styrene in soil and enrichment cultures. Appl Environ Microbiol 35:124-128.
- Sielken RL, Valdez-Flores C. 2001. Dose-response implications of the University of Alabama study of lymphohematopoietic cancer among workers exposed to 1,3-butadiene and styrene in the synthetic rubber industry. Chem Biol Interact 135-136:637-651.
- *Simula AP, Priestly BG. 1992. Species differences in the genotoxicity of cyclophosphamide and styrene in three *in vivo* assays. Mutat Res 271:49-58.
- Sinatra MG, Baiardi P, Manzo L. 2000. Effects of styrene on the nervous and immune systems in occupational low dose exposure [Abstract]. Neurotoxicology 21(5):898.
- Singh AP, Srivastava SP. 1986. Inhibitory characteristics of styrene and styrene oxide on Na⁺, K⁺ activated adenosine triphosphatase. J Environ Pathol Toxicol Oncol 6:29-36.
- +*Sinha AK, Jersey GC, Linscombe VA, et al. 1983. Cytogenetic evaluation of bone marrow cells from rats exposed to styrene vapor for one year. Fundam Appl Toxicol 3:95-98.
- +*Sjöborg S, Fregert S, Trulsson L. 1984. Contact allergy to styrene and related chemicals. Contact Dermatitis 10:94-96.
- +*Śliwińska-Kowalska M, Zamysłowska-Szmytke E, Szymczak W, et al. 2003. Ototoxic effects of occupational exposure to styrene and co-exposure to styrene and noise. J Occup Environ Med 45(1):15-24.
- *Sloane TM, Brudzynski RJ. 1979. Competition between reactive sites in the reactions of oxygen atoms and hydroxyl radicals with phenylacetylene and styrene. J Am Chem Soc 101 (March 14):1495-1499.
- Smith RM. 1988. Supercritical fluid chromatography. Anal Chem 60(24):1394A.
- Snyder RC, Breder OV. 1985. New FDA migration cell used to study migration of styrene from polystyrene into various solvents. J Assoc Off Anal Chem 68:770-775.
- *Sollenberg J, Bjurström R, Wrangskag K, et al. 1988. Biological exposure limits estimated from relations between occupational styrene exposure during a workweek and excretion of mandelic and phenylglyoxylic acids in urine. Int Arch Occup Environ Health 60:365-370.
- Solveig Walles SA, Orsen I. 1983. Single-strand breaks in DNA of various organs of mice induced by styrene and styrene oxide. Cancer Lett 21:9-15.
- *Somorovská M, Jahnova E, Tulinska J, et al. 1999. Biomonitoring of occupational exposure to styrene in a plastic lamination plant. Mutat Res 428(1-2):255-269.
- Speit G, Henderson L. 2004. Review of the *in vivo* genotoxicity tests performed with styrene. Mutat Res 589(1):67-79.

9. REFERENCES

- +*Spencer HC, Irish DD, Adams EM, et al. 1942. The response of laboratory animals to monomeric styrene. *J Ind Hyg Toxicol* 24:295-301.
- *Squillace PJ, Moran MJ, Lapman WW, et al. 1999. Volatile organic compounds in untreated ambient groundwater of the United States, 1985-1995. *Environ Sci Technol* 33(23):4176-4187.
- SRI. 1988a. Directory of chemical producers: United States of America. Menlo Park, CA: SRI International, 961-962.
- SRI. 1988b. Directory of chemical producers: United States of America. Supplement II. Menlo Park, CA: SRI International, 56.
- *SRI. 1989. Directory of chemical producers: United States of America. Menlo Park, CA: SRI International, 968.
- *SRI. 2006. Directory of chemical producers. United States of America. Menlo Park, CA: SRI Consulting, 894.
- +*Srivastava S, Das M, Mushtaq M, et al. 1982. Hepatic effects of orally administered styrene in rats. *J Appl Toxicol* 2:219-222.
- +*Srivastava S, Seth PK, Srivastava SP. 1989. Effect of styrene administration on rat testis. *Arch Toxicol* 63:43-46.
- +*Srivastava S, Seth PK, Srivastava SP, et al. 1992a. Biochemical and morphological studies in testes of rat offspring of mothers exposed to styrene during lactation. *Pharmacol Toxicol* 70:314-316.
- +*Srivastava S, Seth PK, Srivastava SP. 1992b. Effect of styrene on testicular enzymes of growing rat. *Ind J Exp Biol* 30:399-401.
- Staudinger J, Roberts PV. 1996. A critical review of Henry's Law constants for environmental applications. *Crit Rev Environ Sci* 26:205-297.
- *Stefaniak AB, Breyse PN, Murray MPM, et al. 2000. An evaluation of employee exposure to volatile organic compounds in three photocopy centers. *Environ Res* 83:162-173.
- +*Štětkářová I, Urban P, Prochazka B, et al. 1993. Somatosensory evoked potentials in workers exposed to toluene and styrene. *Br J Ind Med* 50(6):520-527.
- +*Stewart RD, Dodd HC, Baretta ED, et al. 1968. Human exposure to styrene vapor. *Arch Environ Health* 16:656-662.
- Stoltz DR, Withey RJ. 1977. Mutagenicity testing of styrene and styrene epoxide in *Salmonella typhimurium*. *Bull Environ Contam Toxicol* 17:739-742.
- Stoner GD, Shimkin MB. 1982. Strain A mouse lung tumor bioassay. *J Am Coll Toxicol* 1:145-169.
- Sumner SC, Cattley RC, Asgharian B, et al. 1997. Evaluation of the metabolism and hepatotoxicity of styrene in F344 rats, B6C3F1 mice, and CD-1 mice following single and repeated inhalation exposures. *Chem Biol Interact* 106(1):47-65.

9. REFERENCES

- *Sumner SJ, Fennell TR. 1994. Review of the metabolic fate of styrene. *Crit Rev Toxicol* 24(Suppl 1):S11-S33.
- Sweet CW, Vermette SJ. 1992. Toxic volatile organic compounds in urban air in Illinois. *Environ Sci Technol* 26:165-173.
- *Symanski E, Bergamaschi E, Mutti A. 2001. Inter-and intra-individual sources of variation in levels of urinary styrene metabolites. *Int Arch Occup Environ Health* 74:336-344.
- *Takeoko GR, Flath RA, Guntert M, et al. 1988. Nectarine volatiles: Vacuum steam distillation versus headspace. *J Agric Food Chem* 36:553-560.
- *Tang J, Jin QZ, Shen GH, et al. 1983. Isolation and identification of volatile compounds from fried chicken. *J Agric Food Chem* 31:1287-1292.
- *Tang W, Hemm I, Eisenbrand G. 2000. Estimation of human exposure to styrene and ethylbenzene. *Toxicology*. *Toxicology* 144(1-3):39-50.
- Teixeira J, Gaspar J, Silva S, et al. 2004a. Haemoglobin adducts as biomarkers of occupational exposure to styrene [Abstract] *Toxicol Appl Pharmacol* 197(3):190.
- Teixeira JP, Gaspar J, Silva S, et al. 2004b. Occupational exposure to styrene: Modulation of cytogenetic damage and levels of urinary metabolites of styrene by polymorphisms in genes CYP2E1, EPHX1, GSTM1, GSTT1 and GSTP1. *Toxicology* 195(2-3):231-242.
- *Teramoto K, Horiguchi S. 1979. Absorption, distribution and elimination of styrene in man and experimental animals. *Arh Hig Rada Toksikol* 30:431-437.
- +*Thiess AM, Friedheim M. 1978. Morbidity among persons employed in styrene production, polymerization and processing plants. *Scand J Work Environ Health* 4 (Suppl 2):203-214.
- *Thiess AM, Schwegler H, Fleig I. 1980. Chromosome investigations in lymphocytes of workers employed in areas in which styrene-containing unsaturated polyester resins are manufactured. *Am J Ind Med* 1:205-210.
- Thomas AO, Chiampo A. 1998. Characterisation of natural attenuation in an industrial site contaminated with phenols and alkylbenzenes. Contaminated soil '98. Proceedings of the sixth international FZK/TNO conference on contaminated soil. London: Thomas Telford Publishing, 181-190.
- *Thomas K, Colborn T. 1992. Organochlorine endocrine disruptors in human tissue. In: Colborn T, Clement C, eds. Chemically induced alterations in sexual and functional development: The wildlife/human connection. Princeton, NJ: Princeton Scientific Publishing, 365-394.
- *Till D, Schwöpe AD, Ehntholt DJ, et al. 1987. Indirect food additive migration from polymeric food packaging materials. *CRC Crit Rev Toxicol* 18:215-243.
- Ting D, Smith MT, Doane-Stezer P, et al. 1991. Measurement of styrene-oxide cysteine adducts in hemoglobin by selective catalytic reduction. *Adv Exp Med Biol* 283:837-841.

9. REFERENCES

- *Tomanin R, Ballarin C, Bartolucci GB, et al. 1992. Chromosome aberrations and micronuclei in lymphocytes of workers exposed to low and medium levels of styrene. *Int Arch Occup Environ Health* 64:209-215.
- +*Toppila E, Forsman P, Pyykko I, et al. 2006. Effect of styrene on postural stability among reinforced plastic boat plant workers in Finland. *J Occup Environ Med* 48(2):175-180.
- Tossavainen A, Jaakkola J. 1994. Occupational exposure to chemical agents in Finland. *Appl Occup Environ Hyg* 9(1):28-31.
- *Trenga CA, Kunkel DD, Eaton DL, et al. 1991. Effects of styrene oxide on rat brain glutathione. *Neurotoxicity* 12:165-178.
- *TRI05. 2007. TRI explorer: Providing access to EPA's toxics release inventory data. Washington, DC: Office of Information Analysis and Access. Office of Environmental Information. U.S. Environmental Protection Agency. Toxics Release Inventory. <http://www.epa.gov/triexplorer/>. March 08, 2007.
- *Triebig G, Lehl S, Weltle D, et al. 1989. Clinical and neurobehavioral study of the acute and chronic neurotoxicity of styrene. *Br J Ind Med* 46:799-804.
- *Triebig G, Schaller KH, Valentin H. 1985. Investigations on neurotoxicity of chemical substances at the workplace. *Int Arch Occup Environ Health* 56:239-247.
- +*Triebig G, Stark T, Ihrig A, et al. 2001. Intervention study on acquired color vision deficiencies in styrene-exposed workers. *J Occup Environ Med* 43(5):494-500.
- +*Tsai SP, Chen JD. 1996. Neurobehavioural effects of occupational exposure to low-level styrene. *Neurotoxicol Teratol* 18(4):463-469.
- +*Tulinska J, Dusinska M, Jahnova E, et al. 2000. Changes in cellular immunity among workers occupationally exposed to styrene in a plastics lamination plant. *Am J Ind Med* 38(5):576-583.
- Tursi F, Samaia M, Salmona M, et al. 1983. Styrene oxidation to styrene oxide in human erythrocytes is catalyzed by oxyhemoglobin. *Experientia* 39:593-594.
- *Uchiyama S, Hasegawa S. 2000. Investigation of a long-term sampling period for monitoring volatile organic compounds in ambient air. *Environ Sci Technol* 34:4656-4661.
- *Umemura T, Kurahashi N, Kondo T, et al. 2005. Acute effects of styrene inhalation on the neuroendocrinological system of rats and the different effects in male and female rats. *Arch Toxicol* 79(11):653-659.
- *USGS. 1998. Methods of analysis by the U.S. Geological Survey National Water Quality Laboratory: Determination of 86 volatile organic compounds in water by gas chromatography/mass spectrometry, including detections less than reporting limits. Denver, CO: U.S. Geological Survey, 97-829.
- *USGS. 2006. Volatile organic compounds in the nation's ground water and drinking water supply wells. Reston, VA: U.S. Geological Survey.

9. REFERENCES

- *USITC. 1987. Synthetic organic chemicals: United States production and sales, 1986. Washington, DC: U.S. International Trade Commission. USITC Publication 2009, 26, 39.
- *USITC. 1988. Synthetic organic chemicals: United States production and sales, 1987. Washington, DC: U.S. International Trade Commission. USITC Publication 2118, 3-2, 3-14.
- Uziel M, Munro NB, Katz DS, et al. 1992. DNA adduct formation by 12 chemicals with populations potentially suitable for molecular epidemiological studies. *Mutat Res* 277(1):35-90.
- Vaghef H, Hellman B. 1998. Detection of styrene and styrene oxide-induced DNA damage in various organs of mice using the Comet assay. *Pharmacol Toxicol* 83:69-74.
- +*Vainio H, Jarvisalo J, Taskinen E. 1979. Adaptive changes caused by intermittent styrene inhalation on xenobiotic biotransformation. *Toxicol Appl Pharmacol* 49:7-14.
- Vainio H, Norppa H, Belvedere G. 1984. Metabolism and mutagenicity of styrene and styrene oxide. In: *Industrial hazards of plastics and synthetic elastomers*. New York, NY: Alan R. Liss, Inc., 215-225.
- Vainio H, Norppa H, Hemminki K, et al. 1981. Metabolism and genotoxicity of styrene. *Adv Exp Med Biol* 136(P&A):257-274.
- *Vainio H, Paakkonen R, Ronnholm K, et al. 1976. A study on the mutagenic activity of styrene and styrene oxide. *Scand J Work Environ Health* 3:147-151.
- *Valvani SC, Yalkowsky SH, Roseman TJ. 1981. Solubility and partitioning IV: Aqueous solubility and octanol-water partition coefficients of liquid nonelectrolytes. *J Pharm Sci* 70:502-507.
- van Agteren MH, Keuning S, Janssen DB. 1998. Styrene. *Handbook on biodegradation and biological treatment of hazardous organic compounds*. Dordrecht: Kluwer Academic Publishers, 235-242.
- Van Duuren BL, Nelson N, Orris L, et al. 1963. Carcinogenicity of epoxies, lactones, and peroxy compounds. *J Natl Cancer Inst* 31:41-55.
- Van Hummelen P, Severi M, Pauwels W, et al. 1994. Cytogenetic analysis of lymphocytes from fiberglass-reinforced plastics workers occupationally exposed to styrene. *Mutat Res* 310(1):157-165.
- *Varner SL, Breder CV. 1981. Headspace sampling and gas chromatographic determination of styrene migration from food-contact polystyrene cups into beverages and food stimulants. *J Assoc Off Anal Chem* 64:1122-1130.
- +*Verplanke AJW, Herber RFM. 1998. Effects on the kidney of occupational exposure to styrene. *Int Arch Occup Environ Health* 71:47-52.
- *Verschuere K. 1983. *Handbook of environmental data on organic chemicals*. 2nd ed. New York: Van Nostrand Reinhold Company, 1055-1057.
- *Verschuere K. 2001. *Handbook of environmental data on organic chemicals*. Vol. 2. 4th ed. New York, NY: John Wiley & Sons, Inc., 1899-1902.
- Vettori MV, Corradi D, Coccini T, et al. 2000. Styrene-induced changes in amacrine retinal cells: An experimental study in the rat. *Neurotoxicology* 21(4):607-614.

9. REFERENCES

- Viaene MK. 2003. Relation between colour vision loss and occupational styrene exposure (Comment on: *Occup Environ Med* 59:824-829). *Occup Environ Med* 60(3):222.
- +*Viaene MK, Pauwels W, Veulemans H, et al. 2001. Neurobehavioural changes and persistence of complaints in workers exposed to styrene in a polyester boat building plant: Influence of exposure. *Occup Environ Med* 58(2):103-112.
- +*Viaene M, Veulemans H, Masschelein R. 1998. Experience with a vocabulary test for workers previously and still exposed to styrene. *Scand J Work Environ Health* 24(4):308-311.
- +*Viau C, Bernard A, De Russis R, et al. 1987. Evaluation of the nephrotoxic potential of styrene in man and in rat. *J Appl Toxicol* 7(5):313-316.
- *Vieira I, Sonnier M, Cresteil T. 1996. Developmental expression of CYP2E1 in the human liver: Hypermethylation control of gene expression during the neonatal period. *Eur J Biochem* 238(2):476-483.
- VIEW Database. 1989. Chemicals in ATSDR VIEW database. Atlanta, GA: Agency for Toxic Substances and Disease Registry, Office of External Affairs, Exposure and Disease Branch.
- *Vodicka P, Bastlova T, Vodickova L, et al. 1995. Biomarkers of styrene exposure in lamination workers: Levels of {IO}{+6}-guanine DNA adducts, DNA strand breaks and mutant frequencies in the hypoxanthine guanine phosphoribosyltransferase gene in T-lymphocytes. *Carcinogenesis* 16(7):1473-1481.
- Vodicka P, Koskinen M, Arand M, et al. 2002. Spectrum of styrene-induced DNA adducts: The relationship to other biomarkers and prospects in human biomonitoring. *Mutat Res* 511(3):239-254.
- Vodicka P, Koskinen M, Stetina R, et al. 2003. The role of various biomarkers in the evaluation of styrene genotoxicity. *Cancer Detect Prev* 27:275-284.
- *Vodicka P, Koskinen M, Vodickova L, et al. 2001a. DNA adducts, strand breaks and micronuclei in mice exposed to styrene by inhalation. *Chem Biol Interact* 137:213-227.
- Vodicka PE, Linhart I, Novak J, et al. 2006. 7-Alkylguanine adduct levels in urine, lungs and liver of mice exposed to styrene by inhalation. *Toxicol Appl Pharmacol* 210:1-8.
- Vodicka P, Soucek P, Tates AD, et al. 2001b. Association between genetic polymorphisms and biomarkers in styrene-exposed workers. *Mutat Res* 482(1-2):89-103.
- Vodicka P, Stetina R, Koskinen M, et al. 2002. New aspects in biomonitoring of occupational exposure to styrene. *Int Arch Occup Environ Health* 75(Suppl):S75-S85.
- *Vodicka P, Tuimala J, Stetina R, et al. 2004. Cytogenetic markers, DNA single-strand breaks, urinary metabolites, and DNA repair rates in styrene-exposed lamination workers. *Environ Health Perspect* 112:867-871.
- Vodicka P, Tvrdik T, Osterman-Golkar S, et al. 1999. An evaluation of styrene genotoxicity using several biomarkers in a 3-year follow-up study of hand-lamination workers. *Mutat Res* 445:205-224.

9. REFERENCES

Vodicka P, Vodickova L, Hemminki K. 1993. ³²P-postlabeling of DNA adducts of styrene-exposed lamination workers. *Carcinogenesis* 14(10):2059-2061.

Vodicka P, Vodickova L, Trejbalovaa K, et al. 1994. Persistence of O6-guanine DNA adducts in styrene-exposed lamination workers determined by ³²P-postlabelling. *Carcinogenesis* 15(9):1949-1953.

+*Vyskocil A, Emminger S, Malir F, et al. 1989. Lack of nephrotoxicity of styrene at current TLV level (50 ppm). *Int Arch Occup Environ Health* 61:409-411.

*Wallace L, Buckley T, Pellizzari E, Gordon S. 1996. Breath measurements as volatile organic compounds biomarkers. *Environ Health Perspect* 104:861-869.

Wallace LA. 1986. Personal exposures, indoor and outdoor air concentrations, and exhaled breath concentrations of selected volatile organic compounds measured for 600 residents of New Jersey, North Dakota, North Carolina and California. *Toxicol Environ Chem* 12:215-236.

*Wallace LA, Pellizzari ED. 1995. Recent advances in measuring exhaled breath and estimating exposure and body burden for volatile organic compounds. *Environ Health Perspect* 103:95-98.

*Wallace LA, Pellizzari ED, Hartwell TD, et al. 1986a. Total exposure assessment methodology (TEAM) study: Personal exposures, indoor-outdoor relationships, and breath levels of volatile organic compounds in New Jersey. *Environ Int* 12:369-387.

Wallace LA, Pellizzari ED, Hartwell TD, et al. 1987a. The TEAM study: Personal exposures to toxic substances in air, drinking water, and breath of 400 residents of New Jersey, North Carolina and North Dakota. *Environ Res* 43:290-307.

*Wallace LA, Pellizzari ED, Leaderer B, et al. 1987b. Emissions of volatile organic compounds from building materials and consumer products. *Atmos Environ* 21:385-393.

*Wallace LA, Pellizzari ED, Sheldon L, et al. 1986b. The total exposure assessment methodology (TEAM) study: Direct measurement of personal exposures through air and water for 600 residents of several U.S. cities. In: Cohen Y, ed. *Pollutants in a multimedia environment*. New York, NY: Plenum Publishing Corporation, 289-315.

*Walles SAS, Orsen I. 1983. Single-strand breaks in DNA of various organs of mice induced by styrene and styrene oxide. *Cancer Lett* 21:9-15.

*Walles SAS, Edling C, Anundi H, et al. 1993. Exposure dependent increase in DNA single strand breaks in leucocytes from workers exposed to low concentrations of styrene. *Br J Ind Med* 50:570-574.

Wang JL, Chew C, Chen SW, et al. 2000. Concentration variability of anthropogenic halocarbons and applications as internal reference in volatile organic compound measurements. *Environ Sci Technol* 34:2243-2248.

Wang Y, Kupper LL, Lof A, et al. 1996. Comparison of average estimated metabolic rates for styrene in previously exposed and unexposed groups with pharmacokinetics modeling. *Occup Environ Med* 53(9):601-605.

*Warhurst AM, Fewson CA. 1994. Microbial metabolism and biotransformation of styrene. *J Appl Bacteriol* 77:597-606.

9. REFERENCES

Watabe T, Hiratsuka A, Aizawa T, et al. 1982. Studies on metabolism and toxicity of styrene: IV. 1-vinylbenzene 3,4-oxide, a potent mutagen formed as a possible intermediate in the metabolism *in vivo* of styrene to 4-vinylphenol. *Mutat Res* 93:45-55.

Watabe T, Ozawa N, Hiratsuka A. 1983. Studies on metabolism and toxicity of styrene - VI. Regioselectivity in glutathione S-conjugation and hydrolysis of racemic, R- and S-phenyloxiranes in rat liver. *Biochem Pharmacol* 32:777-785.

Watanabe T, Endo A, Kumai M, et al. 1983. Chromosome aberrations and sister chromatid exchanges in styrene-exposed workers with reference to their smoking habits. *Environ Mut* 5:299-309.

*Watanabe T, Endo A, Sato K, et al. 1981. Mutagenic potential of styrene in man. *Ind Health* 19:37-45.

*Weast RC, ed. 1985. CRC handbook of chemistry and physics. Boca Raton, FL: CRC Press, Inc., C-498.

Welke B, Ettlenger K, Riederer M. 1998. Sorption of volatile organic chemicals in plant surfaces. *Environ Sci Technol* 32:1099-1104.

*Welp E, Kogevinas M, Andersen A, et al. 1996c. Exposure to styrene and mortality from nervous system diseases and mental disorders. *Am J Epidemiol* 144(7):623-633.

Welp E, Partanen T, Kogevinas M, et al. 1996a. Exposure to styrene and mortality from non-malignant respiratory diseases. *Occup Environ Med* 53(7):499-501.

Welp E, Partanen T, Kogevinas M, et al. 1996b. Exposure to styrene and mortality from nonmalignant diseases of the genitourinary system. *Scand J Work Environ Health* 22(3):223-226.

*Wenker MAM, Kezic S, Monster AC, et al. 2001. Stereochemical metabolism of styrene in volunteers. *Int Arch Occup Environ Health* 74:359-365.

Weschler CJ, Hodgson AT, Wooley JD. 1992. Indoor chemistry: Ozone, volatile organic compounds, and carpets. *Environ Sci Technol* 26:2371-2377.

*West JR, Smith HW, Chasis H. 1948. Glomerular filtration rate, effective renal blood flow, and maximal tubular excretory capacity in infancy. *J Pediatr* 32:10-18.

*WHO. 1983. Environmental health criteria 26: Styrene. Geneva: World Health Organization. <http://www.inchem.org/documents/ehc/ehc/ehc26.htm>. April 05, 2007.

*WHO. 2000. Air quality guidelines. 2nd ed. Geneva, Switzerland: World Health Organization. <http://www.euro.who.int/Document/AIQ/AirQualRepMtg.pdf>. March 08, 2006.

*WHO. 2004. Guidelines for drinking-water quality. Vol. 1. Recommendations. 3rd ed. Geneva, Switzerland: World Health Organization. http://www.who.int/water_sanitation_health/dwq/gdwq3/en/. March 08, 2006.

*Widdowson EM, Dickerson JWT. 1964. Chemical composition of the body. In: Comar CL, Bronner F, eds. Mineral metabolism: An advanced treatise. Volume II: The elements Part A. New York, NY: Academic Press.

9. REFERENCES

- Wieboldt RC, Adams GE, Later DW. 1988. Sensitivity improvement in infrared detection for supercritical fluid chromatography. *Anal Chem* 60(1):2422-2427.
- Wieczorek H. 1985. Evaluation of low exposure to styrene: II. Dermal absorption of styrene vapours in humans under experimental conditions. *Int Arch Occup Environ Health* 57:71-75.
- Wieczorek H, Piotrowski JK. 1985. Evaluation of low exposure to styrene: I. Absorption of styrene vapours by inhalation under experimental conditions. *Int Arch Occup Environ Health* 57:57-59.
- Wieczorek H, Piotrowski JK. 1988. Kinetic interpretation of the exposure test for styrene. *Int Arch Occup Environ Health* 61:107-113.
- *Wigaeus E, Lof A, Bjurström R, et al. 1983. Exposure to styrene: Uptake, distribution, metabolism and elimination in man. *Scand J Work Environ Health* 9:479-488.
- Wilkins K, Larsen K. 1996. Volatile organic compounds from garden waste. *Chemosphere* 32(10):2049-2055.
- Wilson RH. 1944. Health hazards encountered in the manufacture of synthetic rubber. *J Am Med Assoc* 124:701-703.
- *Wilson JT, McNabb JF, Wilson BH, et al. 1983. Biotransformation of selected organic pollutants in ground water. *Developments in Industrial Microbiology* 24:225-233.
- *Windholz, M, ed. 1983. Styrene. *The Merck index*. 10th ed. Rahway, NJ: Merck and Co., Inc., 1270.
- *Wintrobe MM, Thorn GW, Adams RD, et al., eds. 1970. *Harrison's principles of internal medicine*. Vol. 2. 6th ed. New York, NY: McGraw-Hill, Inc., 1614-1615.
- *Withey JR. 1976. Quantitative analysis of styrene monomer in polystyrene and foods including some preliminary studies of the uptake and pharmacodynamics of the monomer in rats. *Environ Health Perspect* 17:125-133.
- *Withey JR, Collins PG. 1977. Pharmacokinetics and distribution of styrene monomer in rats after intravenous administration. *J Toxicol Environ Health* 3:1011-1020.
- *Withey JR, Collins PG. 1978. Styrene monomer in foods: A limited Canadian survey. *Bull Environ Contam Toxicol* 19:86-94.
- *Withey JR, Collins PG. 1979. The distribution and pharmacokinetics of styrene monomer in rats by the pulmonary route. *J Environ Pathol Toxicol* 2:1329-1342.
- *Withey JR, Karpinski K. 1985. Fetal distribution of styrene in rats after vapor phase exposures. *Biol Res Pregnancy* 6:59-64.
- +*Wolf MA, Rowe VK, McCollister DD, et al. 1956. Toxicological studies of certain alkylated benzenes and benzene: Experiments on laboratory animals. *AMA Arch Ind Health* 14:387-398.
- +*Wong O. 1990. A cohort mortality study and a case-control study of workers potentially exposed to styrene in the reinforced plastics and composites industry. *Br J Ind Med* 47:753-762.

9. REFERENCES

- Wong O, Trent LS. 1999. Mortality from nonmalignant diseases of the respiratory, genitourinary and nervous systems among workers exposed to styrene in the reinforced plastics and composites industry. *Scand J Work Environ Health* 25(4):317-325.
- *Wong O, Trent LS, Whorton MD. 1994. An updated cohort mortality study of workers exposed to styrene in the reinforced plastics and composites industry. *Occup Environ Med* 51:386-396.
- Wrangskog K, Sollenberg J, Soderman E. 1996. Application of a single-compartment model for estimation of styrene uptake from measurements of urinary excretion of mandelic and phenylglyoxylic acids after occupational exposure. *Int Arch Occup Environ Health* 68(5):337-341.
- *Yager JW, Paradisin WM, Rappaport SM. 1993. Sister-chromatid exchanges in lymphocytes are increased in relation to longitudinally measured occupational exposure to low concentrations of styrene. *Mutat Res* 319:155-165.
- Yager JW, Paradisin WM, Symanski E, et al. 1990. Sister chromatid exchanges induced in peripheral lymphocytes of workers exposed to low concentrations of styrene. In: Mendelsohn ML, Albertini RJ, eds. *Mutation and the environment. Part C: Somatic and heritable mutation, adduction, and epidemiology*. New York, NY: Wiley-Liss, 347-356.
- +*Yamamoto T, Teramoto K, Horiguchi S. 1997. Effects of styrene on peripheral nerve conduction velocities in rats. *J Occup Health* 39:319-324.
- +*Yano BL, Dittenber DA, Albee RR, et al. 1992. Abnormal auditory brainstem responses and cochlear pathology in rats induced by an exaggerated styrene exposure regimen. *Toxicol Pathol* 20(1):1-6.
- Yeowell-O'Connell K, Jin Z, Rappaport SM. 1996. Determination of albumin and hemoglobin adducts in workers exposed to styrene and styrene oxide. *Cancer Epidemiol Biomarkers Prev* 5:205-215.
- *Yoshikawa K, Isobe M, Watabe T, et al. 1980. Studies on metabolism and toxicity of styrene: III. The effect of metabolic inactivation by rat-liver S9 on the mutagenicity of phenylloxirane toward *Salmonella typhimurium*. *Mutat Res* 78:219-226.
- *Young JD, Ramsey JC, Blau GE, et al. 1979. Pharmacokinetics of inhaled or intraperitoneally administered styrene in rats. In: *Toxicology and Occupational Medicine*. New York, NY: Elsevier/North-Holland, 297-310.
- *Yuasa J, Kishi R, Eguchi T, et al. 1996. Study of urinary mandelic acid concentration and peripheral nerve conduction among styrene workers. *Am J Ind Med* 30:41-47.
- Zabiegala B, Namiesnik J, Przyk E, et al. 1999. Changes in concentration levels of selected VOCs in newly erected and remodeled buildings in Gdansk. *Chemosphere* 39(12):2035-2046.
- +*Zaidi NF, Agrawal AK, Srivastava SP, et al. 1985. Effect of gestational and neonatal styrene exposure on dopamine receptors. *Neurobehav Toxicol Teratol* 7:23-28.
- *Zelano V, Rizzi S, Gastaldi D, et al. 1998. Volatile organic compounds in drinking water. *Ann Chim* 88:449-460.

9. REFERENCES

- Zetterberg G. 1982. Styrene-a widespread mutagen: Conclusions from the results of testing. Proceedings of the 3rd Environmental Mutagens and Carcinogens International Conference, Tokyo, Japan. New York, NY: Alan R. Liss, Inc., 315-322.
- Zhang J, Wilson WE, Liou PJ. 1994. Indoor air chemistry: Formation of organic acids and aldehydes. Environ Sci Technol 28:1975-1982.
- Zhang XX, Chakrabarti S, Malick AM, et al. 1993. Effects of different styrene metabolites on cytotoxicity, sister-chromatid exchanges and cell-cycle kinetics in human whole blood lymphocytes *in vitro*. Mutat Res 302:213-218.
- Zheng J, Chung J, Yuan W. 2004. Mechanistic investigation on cytotoxicity of styrene. [Abstract] Toxicologist 78(1-5):137.
- *Ziegler EE, Edwards BB, Jensen RL, et al. 1978. Absorption and retention of lead by infants. Pediatr Res 12(1):29-34.
- Zielinska B, Fung KK. 1994. The composition and concentration of hydrocarbons in the range of C₂ to C₁₈ emitted from motor vehicles. Sci Total Environ 146/147:281-288.
- *Zielinska B, Sagebiel JC, Harshfield G, et al. 1996. Volatile organic compounds up to C₂₀ emitted from motor vehicles; measurement methods. Atmos Environ 30:2269-2286.
- *Zoeteman BC, Harmsen K, Linders JB, et al. 1980. Persistent organic pollutants in river water and groundwater of the Netherlands. Chemosphere 9:231-241.

10. GLOSSARY

Absorption—The taking up of liquids by solids, or of gases by solids or liquids.

Acute Exposure—Exposure to a chemical for a duration of 14 days or less, as specified in the Toxicological Profiles.

Adsorption—The adhesion in an extremely thin layer of molecules (as of gases, solutes, or liquids) to the surfaces of solid bodies or liquids with which they are in contact.

Adsorption Coefficient (K_{oc})—The ratio of the amount of a chemical adsorbed per unit weight of organic carbon in the soil or sediment to the concentration of the chemical in solution at equilibrium.

Adsorption Ratio (K_d)—The amount of a chemical adsorbed by sediment or soil (i.e., the solid phase) divided by the amount of chemical in the solution phase, which is in equilibrium with the solid phase, at a fixed solid/solution ratio. It is generally expressed in micrograms of chemical sorbed per gram of soil or sediment.

Benchmark Dose (BMD)—Usually defined as the lower confidence limit on the dose that produces a specified magnitude of changes in a specified adverse response. For example, a BMD_{10} would be the dose at the 95% lower confidence limit on a 10% response, and the benchmark response (BMR) would be 10%. The BMD is determined by modeling the dose response curve in the region of the dose response relationship where biologically observable data are feasible.

Benchmark Dose Model—A statistical dose-response model applied to either experimental toxicological or epidemiological data to calculate a BMD.

Bioconcentration Factor (BCF)—The quotient of the concentration of a chemical in aquatic organisms at a specific time or during a discrete time period of exposure divided by the concentration in the surrounding water at the same time or during the same period.

Biomarkers—Broadly defined as indicators signaling events in biologic systems or samples. They have been classified as markers of exposure, markers of effect, and markers of susceptibility.

Cancer Effect Level (CEL)—The lowest dose of chemical in a study, or group of studies, that produces significant increases in the incidence of cancer (or tumors) between the exposed population and its appropriate control.

Carcinogen—A chemical capable of inducing cancer.

Case-Control Study—A type of epidemiological study that examines the relationship between a particular outcome (disease or condition) and a variety of potential causative agents (such as toxic chemicals). In a case-controlled study, a group of people with a specified and well-defined outcome is identified and compared to a similar group of people without outcome.

Case Report—Describes a single individual with a particular disease or exposure. These may suggest some potential topics for scientific research, but are not actual research studies.

Case Series—Describes the experience of a small number of individuals with the same disease or exposure. These may suggest potential topics for scientific research, but are not actual research studies.

10. GLOSSARY

Ceiling Value—A concentration of a substance that should not be exceeded, even instantaneously.

Chronic Exposure—Exposure to a chemical for 365 days or more, as specified in the Toxicological Profiles.

Cohort Study—A type of epidemiological study of a specific group or groups of people who have had a common insult (e.g., exposure to an agent suspected of causing disease or a common disease) and are followed forward from exposure to outcome. At least one exposed group is compared to one unexposed group.

Cross-sectional Study—A type of epidemiological study of a group or groups of people that examines the relationship between exposure and outcome to a chemical or to chemicals at one point in time.

Data Needs—Substance-specific informational needs that if met would reduce the uncertainties of human health assessment.

Developmental Toxicity—The occurrence of adverse effects on the developing organism that may result from exposure to a chemical prior to conception (either parent), during prenatal development, or postnatally to the time of sexual maturation. Adverse developmental effects may be detected at any point in the life span of the organism.

Dose-Response Relationship—The quantitative relationship between the amount of exposure to a toxicant and the incidence of the adverse effects.

Embryotoxicity and Fetotoxicity—Any toxic effect on the conceptus as a result of prenatal exposure to a chemical; the distinguishing feature between the two terms is the stage of development during which the insult occurs. The terms, as used here, include malformations and variations, altered growth, and *in utero* death.

Environmental Protection Agency (EPA) Health Advisory—An estimate of acceptable drinking water levels for a chemical substance based on health effects information. A health advisory is not a legally enforceable federal standard, but serves as technical guidance to assist federal, state, and local officials.

Epidemiology—Refers to the investigation of factors that determine the frequency and distribution of disease or other health-related conditions within a defined human population during a specified period.

Genotoxicity—A specific adverse effect on the genome of living cells that, upon the duplication of affected cells, can be expressed as a mutagenic, clastogenic, or carcinogenic event because of specific alteration of the molecular structure of the genome.

Half-life—A measure of rate for the time required to eliminate one half of a quantity of a chemical from the body or environmental media.

Immediately Dangerous to Life or Health (IDLH)—The maximum environmental concentration of a contaminant from which one could escape within 30 minutes without any escape-impairing symptoms or irreversible health effects.

Immunologic Toxicity—The occurrence of adverse effects on the immune system that may result from exposure to environmental agents such as chemicals.

10. GLOSSARY

Immunological Effects—Functional changes in the immune response.

Incidence—The ratio of individuals in a population who develop a specified condition to the total number of individuals in that population who could have developed that condition in a specified time period.

Intermediate Exposure—Exposure to a chemical for a duration of 15–364 days, as specified in the Toxicological Profiles.

In Vitro—Isolated from the living organism and artificially maintained, as in a test tube.

In Vivo—Occurring within the living organism.

Lethal Concentration_(LO) (LC_{LO})—The lowest concentration of a chemical in air that has been reported to have caused death in humans or animals.

Lethal Concentration₍₅₀₎ (LC₅₀)—A calculated concentration of a chemical in air to which exposure for a specific length of time is expected to cause death in 50% of a defined experimental animal population.

Lethal Dose_(LO) (LD_{LO})—The lowest dose of a chemical introduced by a route other than inhalation that has been reported to have caused death in humans or animals.

Lethal Dose₍₅₀₎ (LD₅₀)—The dose of a chemical that has been calculated to cause death in 50% of a defined experimental animal population.

Lethal Time₍₅₀₎ (LT₅₀)—A calculated period of time within which a specific concentration of a chemical is expected to cause death in 50% of a defined experimental animal population.

Lowest-Observed-Adverse-Effect Level (LOAEL)—The lowest exposure level of chemical in a study, or group of studies, that produces statistically or biologically significant increases in frequency or severity of adverse effects between the exposed population and its appropriate control.

Lymphoreticular Effects—Represent morphological effects involving lymphatic tissues such as the lymph nodes, spleen, and thymus.

Malformations—Permanent structural changes that may adversely affect survival, development, or function.

Minimal Risk Level (MRL)—An estimate of daily human exposure to a hazardous substance that is likely to be without an appreciable risk of adverse noncancer health effects over a specified route and duration of exposure.

Modifying Factor (MF)—A value (greater than zero) that is applied to the derivation of a Minimal Risk Level (MRL) to reflect additional concerns about the database that are not covered by the uncertainty factors. The default value for a MF is 1.

Morbidity—State of being diseased; morbidity rate is the incidence or prevalence of disease in a specific population.

Mortality—Death; mortality rate is a measure of the number of deaths in a population during a specified interval of time.

10. GLOSSARY

Mutagen—A substance that causes mutations. A mutation is a change in the DNA sequence of a cell's DNA. Mutations can lead to birth defects, miscarriages, or cancer.

Necropsy—The gross examination of the organs and tissues of a dead body to determine the cause of death or pathological conditions.

Neurotoxicity—The occurrence of adverse effects on the nervous system following exposure to a chemical.

No-Observed-Adverse-Effect Level (NOAEL)—The dose of a chemical at which there were no statistically or biologically significant increases in frequency or severity of adverse effects seen between the exposed population and its appropriate control. Effects may be produced at this dose, but they are not considered to be adverse.

Octanol-Water Partition Coefficient (K_{ow})—The equilibrium ratio of the concentrations of a chemical in *n*-octanol and water, in dilute solution.

Odds Ratio (OR)—A means of measuring the association between an exposure (such as toxic substances and a disease or condition) that represents the best estimate of relative risk (risk as a ratio of the incidence among subjects exposed to a particular risk factor divided by the incidence among subjects who were not exposed to the risk factor). An OR of greater than 1 is considered to indicate greater risk of disease in the exposed group compared to the unexposed group.

Organophosphate or Organophosphorus Compound—A phosphorus-containing organic compound and especially a pesticide that acts by inhibiting cholinesterase.

Permissible Exposure Limit (PEL)—An Occupational Safety and Health Administration (OSHA) allowable exposure level in workplace air averaged over an 8-hour shift of a 40-hour workweek.

Pesticide—General classification of chemicals specifically developed and produced for use in the control of agricultural and public health pests.

Pharmacokinetics—The dynamic behavior of a material in the body, used to predict the fate (disposition) of an exogenous substance in an organism. Utilizing computational techniques, it provides the means of studying the absorption, distribution, metabolism, and excretion of chemicals by the body.

Pharmacokinetic Model—A set of equations that can be used to describe the time course of a parent chemical or metabolite in an animal system. There are two types of pharmacokinetic models: data-based and physiologically-based. A data-based model divides the animal system into a series of compartments, which, in general, do not represent real, identifiable anatomic regions of the body, whereas the physiologically-based model compartments represent real anatomic regions of the body.

Physiologically Based Pharmacodynamic (PBPD) Model—A type of physiologically based dose-response model that quantitatively describes the relationship between target tissue dose and toxic end points. These models advance the importance of physiologically based models in that they clearly describe the biological effect (response) produced by the system following exposure to an exogenous substance.

Physiologically Based Pharmacokinetic (PBPK) Model—Comprised of a series of compartments representing organs or tissue groups with realistic weights and blood flows. These models require a

10. GLOSSARY

variety of physiological information: tissue volumes, blood flow rates to tissues, cardiac output, alveolar ventilation rates, and possibly membrane permeabilities. The models also utilize biochemical information, such as air/blood partition coefficients, and metabolic parameters. PBPK models are also called biologically based tissue dosimetry models.

Prevalence—The number of cases of a disease or condition in a population at one point in time.

Prospective Study—A type of cohort study in which the pertinent observations are made on events occurring after the start of the study. A group is followed over time.

q_1^* —The upper-bound estimate of the low-dose slope of the dose-response curve as determined by the multistage procedure. The q_1^* can be used to calculate an estimate of carcinogenic potency, the incremental excess cancer risk per unit of exposure (usually $\mu\text{g/L}$ for water, mg/kg/day for food, and $\mu\text{g/m}^3$ for air).

Recommended Exposure Limit (REL)—A National Institute for Occupational Safety and Health (NIOSH) time-weighted average (TWA) concentration for up to a 10-hour workday during a 40-hour workweek.

Reference Concentration (RfC)—An estimate (with uncertainty spanning perhaps an order of magnitude) of a continuous inhalation exposure to the human population (including sensitive subgroups) that is likely to be without an appreciable risk of deleterious noncancer health effects during a lifetime. The inhalation reference concentration is for continuous inhalation exposures and is appropriately expressed in units of mg/m^3 or ppm.

Reference Dose (RfD)—An estimate (with uncertainty spanning perhaps an order of magnitude) of the daily exposure of the human population to a potential hazard that is likely to be without risk of deleterious effects during a lifetime. The RfD is operationally derived from the no-observed-adverse-effect level (NOAEL, from animal and human studies) by a consistent application of uncertainty factors that reflect various types of data used to estimate RfDs and an additional modifying factor, which is based on a professional judgment of the entire database on the chemical. The RfDs are not applicable to nonthreshold effects such as cancer.

Reportable Quantity (RQ)—The quantity of a hazardous substance that is considered reportable under the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA). Reportable quantities are (1) 1 pound or greater or (2) for selected substances, an amount established by regulation either under CERCLA or under Section 311 of the Clean Water Act. Quantities are measured over a 24-hour period.

Reproductive Toxicity—The occurrence of adverse effects on the reproductive system that may result from exposure to a chemical. The toxicity may be directed to the reproductive organs and/or the related endocrine system. The manifestation of such toxicity may be noted as alterations in sexual behavior, fertility, pregnancy outcomes, or modifications in other functions that are dependent on the integrity of this system.

Retrospective Study—A type of cohort study based on a group of persons known to have been exposed at some time in the past. Data are collected from routinely recorded events, up to the time the study is undertaken. Retrospective studies are limited to causal factors that can be ascertained from existing records and/or examining survivors of the cohort.

Risk—The possibility or chance that some adverse effect will result from a given exposure to a chemical.

10. GLOSSARY

Risk Factor—An aspect of personal behavior or lifestyle, an environmental exposure, or an inborn or inherited characteristic that is associated with an increased occurrence of disease or other health-related event or condition.

Risk Ratio—The ratio of the risk among persons with specific risk factors compared to the risk among persons without risk factors. A risk ratio greater than 1 indicates greater risk of disease in the exposed group compared to the unexposed group.

Short-Term Exposure Limit (STEL)—The American Conference of Governmental Industrial Hygienists (ACGIH) maximum concentration to which workers can be exposed for up to 15 minutes continually. No more than four excursions are allowed per day, and there must be at least 60 minutes between exposure periods. The daily Threshold Limit Value-Time Weighted Average (TLV-TWA) may not be exceeded.

Standardized Mortality Ratio (SMR)—A ratio of the observed number of deaths and the expected number of deaths in a specific standard population.

Target Organ Toxicity—This term covers a broad range of adverse effects on target organs or physiological systems (e.g., renal, cardiovascular) extending from those arising through a single limited exposure to those assumed over a lifetime of exposure to a chemical.

Teratogen—A chemical that causes structural defects that affect the development of an organism.

Threshold Limit Value (TLV)—An American Conference of Governmental Industrial Hygienists (ACGIH) concentration of a substance to which most workers can be exposed without adverse effect. The TLV may be expressed as a Time Weighted Average (TWA), as a Short-Term Exposure Limit (STEL), or as a ceiling limit (CL).

Time-Weighted Average (TWA)—An allowable exposure concentration averaged over a normal 8-hour workday or 40-hour workweek.

Toxic Dose₍₅₀₎ (TD₅₀)—A calculated dose of a chemical, introduced by a route other than inhalation, which is expected to cause a specific toxic effect in 50% of a defined experimental animal population.

Toxicokinetic—The absorption, distribution, and elimination of toxic compounds in the living organism.

Uncertainty Factor (UF)—A factor used in operationally deriving the Minimal Risk Level (MRL) or Reference Dose (RfD) or Reference Concentration (RfC) from experimental data. UFs are intended to account for (1) the variation in sensitivity among the members of the human population, (2) the uncertainty in extrapolating animal data to the case of human, (3) the uncertainty in extrapolating from data obtained in a study that is of less than lifetime exposure, and (4) the uncertainty in using lowest-observed-adverse-effect level (LOAEL) data rather than no-observed-adverse-effect level (NOAEL) data. A default for each individual UF is 10; if complete certainty in data exists, a value of 1 can be used; however, a reduced UF of 3 may be used on a case-by-case basis, 3 being the approximate logarithmic average of 10 and 1.

Xenobiotic—Any chemical that is foreign to the biological system.

APPENDIX A. ATSDR MINIMAL RISK LEVELS AND WORKSHEETS

The Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) [42 U.S.C. 9601 et seq.], as amended by the Superfund Amendments and Reauthorization Act (SARA) [Pub. L. 99–499], requires that the Agency for Toxic Substances and Disease Registry (ATSDR) develop jointly with the U.S. Environmental Protection Agency (EPA), in order of priority, a list of hazardous substances most commonly found at facilities on the CERCLA National Priorities List (NPL); prepare toxicological profiles for each substance included on the priority list of hazardous substances; and assure the initiation of a research program to fill identified data needs associated with the substances.

The toxicological profiles include an examination, summary, and interpretation of available toxicological information and epidemiologic evaluations of a hazardous substance. During the development of toxicological profiles, Minimal Risk Levels (MRLs) are derived when reliable and sufficient data exist to identify the target organ(s) of effect or the most sensitive health effect(s) for a specific duration for a given route of exposure. An MRL is an estimate of the daily human exposure to a hazardous substance that is likely to be without appreciable risk of adverse noncancer health effects over a specified duration of exposure. MRLs are based on noncancer health effects only and are not based on a consideration of cancer effects. These substance-specific estimates, which are intended to serve as screening levels, are used by ATSDR health assessors to identify contaminants and potential health effects that may be of concern at hazardous waste sites. It is important to note that MRLs are not intended to define clean-up or action levels.

MRLs are derived for hazardous substances using the no-observed-adverse-effect level/uncertainty factor approach. They are below levels that might cause adverse health effects in the people most sensitive to such chemical-induced effects. MRLs are derived for acute (1–14 days), intermediate (15–364 days), and chronic (365 days and longer) durations and for the oral and inhalation routes of exposure. Currently, MRLs for the dermal route of exposure are not derived because ATSDR has not yet identified a method suitable for this route of exposure. MRLs are generally based on the most sensitive chemical-induced end point considered to be of relevance to humans. Serious health effects (such as irreparable damage to the liver or kidneys, or birth defects) are not used as a basis for establishing MRLs. Exposure to a level above the MRL does not mean that adverse health effects will occur.

MRLs are intended only to serve as a screening tool to help public health professionals decide where to look more closely. They may also be viewed as a mechanism to identify those hazardous waste sites that are not expected to cause adverse health effects. Most MRLs contain a degree of uncertainty because of

APPENDIX A

the lack of precise toxicological information on the people who might be most sensitive (e.g., infants, elderly, nutritionally or immunologically compromised) to the effects of hazardous substances. ATSDR uses a conservative (i.e., protective) approach to address this uncertainty consistent with the public health principle of prevention. Although human data are preferred, MRLs often must be based on animal studies because relevant human studies are lacking. In the absence of evidence to the contrary, ATSDR assumes that humans are more sensitive to the effects of hazardous substance than animals and that certain persons may be particularly sensitive. Thus, the resulting MRL may be as much as 100-fold below levels that have been shown to be nontoxic in laboratory animals.

Proposed MRLs undergo a rigorous review process: Health Effects/MRL Workgroup reviews within the Division of Toxicology and Environmental Medicine, expert panel peer reviews, and agency-wide MRL Workgroup reviews, with participation from other federal agencies and comments from the public. They are subject to change as new information becomes available concomitant with updating the toxicological profiles. Thus, MRLs in the most recent toxicological profiles supersede previously published levels. For additional information regarding MRLs, please contact the Division of Toxicology and Environmental Medicine, Agency for Toxic Substances and Disease Registry, 1600 Clifton Road NE, Mailstop F-32, Atlanta, Georgia 30333.

APPENDIX A

MINIMAL RISK LEVEL (MRL) WORKSHEET

Chemical Name: Styrene
CAS Numbers: 100-42-5
Date: August 2007
Profile Status: Pre-Public Comment, Final Draft
Route: Inhalation Oral
Duration: Acute Intermediate Chronic
Graph Key: 16
Species: Human

Minimal Risk Level: 2 mg/kg/day ppm

Reference: Seeber A, Blaszkewicz M, Golka K, et al. 2004. Neurobehavioral effects of experimental exposures to low levels of styrene. Toxicol Lett 151:183-192.

Experimental design: Groups of eight volunteers (gender not reported) were exposed to 0.5 or 20 ppm styrene for 3 hours. Prior to exposure, during the third hour of exposure, and 1.5 hours after exposure termination, the subjects were tested for simple reaction time, choice reaction time, and attention. The subjects were also asked to complete a symptom questionnaire before, during, and after exposure.

Effect noted in study and corresponding doses: The mean concentrations of styrene in blood were 2.2 and 80 µg/L after 3 hours of exposure to 0.5 and 20 ppm, respectively. The blood levels of styrene were correlated with styrene levels in air ($r_{xy}=0.98$). No significant alterations in performance on neurobehavioral tests were found. An increase in the reporting of breathing problems was found; however, it was not significantly significant and the ranking of the severity of the breathing problem was very low (0.5 on a scale of 5).

Dose and end point used for MRL derivation:

NOAEL LOAEL

A NOAEL of 20 ppm for the lack of alterations in tests of simple reaction time, choice reaction time, or attention.

Uncertainty Factors used in MRL derivation:

- 10 for use of a LOAEL
- 10 for extrapolation from animals to humans
- 10 for human variability

Was a conversion factor used from ppm in food or water to a mg/body weight dose? Not applicable.

If an inhalation study in animals, list conversion factors used in determining human equivalent dose: Not applicable.

Was a conversion used from intermittent to continuous exposure? No, because the subjects were only exposed once for 3 hours.

Other additional studies or pertinent information that lend support to this MRL: Exposure to 99 ppm for 7 hours or 376 ppm for 1 hour (Stewart et al. 1968) resulted in eye irritation in experimental subjects;

APPENDIX A

nasal irritation was also reported at 376 ppm. A significant inhibition of the vestibular-oculomotor system was observed in subjects exposed to 87 ppm for 1 hour (Ödkvist et al. 1982). Studies by Stewart et al. (1968) found alterations in tests of balance or coordination in subjects exposed to 376 ppm for 1 hour, but not after exposure to 99 ppm for 7 hours or 216 ppm for 1 hour; the test used in the Stewart et al. (1968) studies is probably less sensitive than those used by Ödkvist et al. (1982). No significant alterations in performance on neurobehavioral tests were observed in subjects exposed to 20 ppm for 3 hours (Seeber et al. 2004).

The available data suggest that the nervous system is the most sensitive target of styrene toxicity following acute-duration inhalation exposure. The lowest LOAEL for a relevant end point in humans is 87 ppm for vestibular impairment in subjects exposed to styrene for 1 hour (Ödkvist et al. 1982). A similar LOAEL (80 ppm) was identified for nasal effects in mice exposed to styrene for 3 days (Cruzan et al. 2001); this effect was not considered suitable as the basis of an MRL. As stated previously, mice appear to have a greater capacity than humans to generate the reactive metabolite, styrene oxide, in the nasal cavity and a lower capacity to detoxify styrene oxide (Green et al. 2001a). The identification of the nervous system as the critical target of toxicity for styrene is supported by a large number of occupational exposure studies. Delays in reaction time have been observed in workers exposed to 21.9–92 ppm (Cherry et al. 1980; Fallas et al. 1992; Gamberale et al. 1976; Jegaden et al. 1993; Mutti et al. 1984a; Tsai and Chen 1996) and vestibular effects have been observed at 18–36 ppm (Calabrese et al. 1996; Möller et al. 1990; Toppila et al. 2006).

Agency Contacts (Chemical Managers): Zemoria Rosemond, Jewell Wilson, Selene Chou

APPENDIX A

MINIMAL RISK LEVEL (MRL) WORKSHEET

Chemical Name: Styrene
CAS Numbers: 100-42-5
Date: August 2007
Profile Status: Pre-Public Comment, Final Draft
Route: Inhalation Oral
Duration: Acute Intermediate Chronic
Graph Key: 60
Species: Human

Minimal Risk Level: 0.2 mg/kg/day ppm

Reference: Benignus VA, Geller AM, Boyes WK, et al. 2005. Human neurobehavioral effects of long-term exposure to styrene: a meta-analysis. Environ Health Perspect 113:532-538.

Experimental design: Benignus et al. (2005) used data from occupational exposure studies examining color vision impairment (Campagna et al. 1996; Eguchi et al. 1995; Gobba et al. 1991; Gong et al. 2002; Kishi et al. 2001) and delays in choice reaction time (Jegaden et al. 1993; Mutti et al. 1984a; Triebig et al. 1989; Tsai and Chen 1996). Average styrene exposure concentrations for each study were estimated from individual data reported in the papers; for studies reporting individual data as urinary mandelic acid levels, standardized methods for converting to styrene exposure levels were used. Cumulative styrene exposure was estimated by multiplying exposure level by length of employment. A common metric of effect magnitude (percentage of baseline) was calculated for the different neurological effects.

Effect noted in study and corresponding doses: A significant linear relationship between choice reaction time and cumulative styrene exposure was found; cumulative exposure accounted for 91% of the variance in reaction time. Similarly, a significant relationship between color confusion index (CCI) and cumulative styrene exposure was found, with cumulative exposure accounting for 35% of the variance in CCI. Using the regression equations for these two effects, the investigators estimated that exposure to 150 ppm for 8 work-years would result in a 50% increase in choice reaction time and a 17% increase in CCI score; exposure to 20 ppm for 8 work-years would result in a 6.5% increase in choice reaction time and a 2.23% increase in CCI score. As discussed in Benignus et al. (2005), a 7% decrease in reaction time would prevent 58,000–70,000 injuries per year from automobile accidents. The investigators also noted that CCI increases with age, the rate of increase is about 10% per 13 years of age; thus, a 2.23% decrease in color perception would be roughly equivalent to 2.9 additional years of age. Based on this analysis, 20 ppm is considered a LOAEL for neurological effects.

Dose and end point used for MRL derivation:

NOAEL LOAEL

A LOAEL of 20 ppm was identified for biologically significant increases in choice reaction time and decrease in color perception in workers exposed to styrene for 8 work years.

Uncertainty Factors used in MRL derivation: 100

- 10 for use of a LOAEL
- 10 for extrapolation from animals to humans
- 10 for human variability

APPENDIX A

Was a conversion factor used from ppm in food or water to a mg/body weight dose? Not applicable.

If an inhalation study in animals, list conversion factors used in determining human equivalent dose: Not applicable.

Was a conversion used from intermittent to continuous exposure? No.

Other additional studies or pertinent information that lend support to this MRL: A large number of occupational exposure studies have examined the neurotoxicity of styrene in reinforced plastic workers, particularly laminators. A summary of the studies examining color vision, vestibular function, reaction time, and neurological symptoms is presented in Table A-1; more details regarding these studies and other studies of neurological effects in styrene workers are presented in Table 3-2.

Table A-1. Results of Selected Human Neurotoxicity Studies

Result	Reference	NOAEL (ppm)	LOAEL (ppm)
Decreased color discrimination	Chia et al. (1994)		6
	Kishi et al. (2001)	4	10
	Gong et al. (2002)		105
	Gobba et al. (1991)		16
	Iregren et al. (2005)		22
	Fallas et al. (1992)		24.3
	Campagna et al. (1996)		265
	Eguchi et al. (1995)	8	93
Neurological symptoms	Flodin et al. (1989)	6	
	Edling et al. (1993)	8.6	
	Checkoway et al. (1992)	10.8	18.9
	Cherry et al. 1980		92
Vestibular effects	Möller et al. (1990)		18
	Toppila et al. (2006)		24.8
	Calabrese et al. (1996)		36
Reaction time	Edling et al. (1993)	8.6	21.9
	Tsai and Chen (1996)		21.9
	Jegaden et al. (1993)		22.68
	Fallas et al. (1992)		24.3
	Mutti et al. (1984a)		25
	Gamberale et al. (1976)		4,755
	Cherry et al. (1980)		92

Agency Contacts (Chemical Managers): Zemoria Rosemond, Jewell Wilson, Selene Chou

APPENDIX A

MINIMAL RISK LEVEL (MRL) WORKSHEET

Chemical Name: Styrene
CAS Numbers: 100-42-5
Date: August 2007
Profile Status: Pre-Public Comment, Final Draft
Route: Inhalation Oral
Duration: Acute Intermediate Chronic
Graph Key: 3
Species: Rat

Minimal Risk Level: 0.1 mg/kg/day ppm

Reference: Husain R, Srivastava SP, Seth PK. 1985. Some behavioral effects of early styrene intoxication in experimental animals. Arch Toxicol 57:53-55.

Experimental design: Groups of 15 male Wistar rats were administered by gavage 0, 100, or 200 mg/kg/day styrene in ground nut oil for 14 consecutive days. Spontaneous motor activity for a period of 10 minutes was measured 24 hours after the last dose. After baseline activity was measured, the rats received an intraperitoneal injection of 2.5 mg/kg amphetamine, and amphetamine-induced spontaneous motor activity was measured. Two days after exposure termination, acquisition training was initiated using a Cook's pole climbing apparatus. Learning was assessed by measuring the number of times the rat climbed the pole after the conditioned stimulus to avoid the foot-shock unconditioned stimulus. The rats were tested for 4 days. Noradrenaline, dopamine, and serotonin levels were measured in seven regions of the brain in six rats/group killed after the acquisition training.

Effect noted in study and corresponding doses: No overt signs of toxicity were observed. No significant alterations in locomotor activity were observed with or without amphetamine induction. Significantly greater increases in percent avoidance response (indicative of impaired learning) were observed at 100 and 200 mg/kg/day; no difference was found between the two styrene groups. The effects were observed on test days 3 and 4. Significant increases in the level of serotonin in the hypothalamus (70%), hippocampus (51%), and midbrain (29%) were observed at 200 mg/kg/day. Styrene did not affect brain noradrenaline and dopamine levels.

Dose and end point used for MRL derivation:

NOAEL LOAEL

A LOAEL of 100 mg/kg/day for impaired learning

Uncertainty Factors used in MRL derivation:

- 10 for use of a LOAEL
- 10 for extrapolation from animals to humans
- 10 for human variability

Was a conversion factor used from ppm in food or water to a mg/body weight dose? Not applicable.

If an inhalation study in animals, list conversion factors used in determining human equivalent dose: Not applicable.

APPENDIX A

Was a conversion used from intermittent to continuous exposure? Not applicable.

Other additional studies or pertinent information that lend support to this MRL: A limited number of studies have examined the acute toxicity of orally-administered styrene; these studies have examined potential neurotoxicity and developmental toxicity. No developmental effects were observed in rats administered a single dose of 300 mg/kg on gestational day 11 (Daston et al. 1991) or administered 300 mg/kg/day (administered as two daily doses of 150 mg/kg) on gestational days 6–15 (Murray et al. 1978). Impaired learning was observed in rats administered via gavage 100 or 200 mg/kg/day for 14 days (Husain et al. 1985); increases in serotonin levels in the hypothalamus, hippocampus, and midbrain were also observed at 200 mg/kg/day. In another study, increases in dopamine receptor binding was observed in rats administered a single gavage dose of 200 mg/kg (Agrawal et al. 1982).

Although there is a limited acute toxicity database, longer-term oral studies support the selection of neurotoxicity as the principal effect. The lowest LOAEL identified for a systemic effect is 400 mg/kg/day for Heinz body formation in dogs administered styrene by gavage for 561 days (Quast et al. 1979); the identified NOAEL was 200 mg/kg/day. Decreased spermatozoa counts were observed in adult rats administered 400 mg/kg 6 days/week for 60 days (Srivastava et al. 1989), young rats exposed via lactation on postnatal days 1–21 (maternal dose of 400 mg/kg/day) (Srivastava et al. 1992a), and young rats administered 200 mg/kg 6 days/week on postnatal days 1–61 (Srivastava et al. 1992b); the NOAELs identified in these three studies were 200, 200, and 100 mg/kg, respectively. Marked degeneration of the seminiferous tubules was also observed in the adult rats administered 400 mg/kg (Srivastava et al. 1989). Impaired learning was also observed in rats administered 500 mg/kg 5 days/week for 8 weeks (Bushnell 1994); a NOAEL was not identified in this study. The extensive inhalation toxicity database for styrene also supports the selection of neurotoxicity as the most sensitive target of toxicity; both the acute- and chronic-duration inhalation MRLs are based on neurological effects in humans. Neurological effects observed in chronically exposed styrene workers include decreased color discrimination, slowed reaction time, increased prevalence of neurological symptoms, and ototoxicity (hearing and vestibular effects).

Agency Contacts (Chemical Managers): Zemoria Rosemond, Jewell Wilson, Selene Chou

APPENDIX B. USER'S GUIDE

Chapter 1

Public Health Statement

This chapter of the profile is a health effects summary written in non-technical language. Its intended audience is the general public, especially people living in the vicinity of a hazardous waste site or chemical release. If the Public Health Statement were removed from the rest of the document, it would still communicate to the lay public essential information about the chemical.

The major headings in the Public Health Statement are useful to find specific topics of concern. The topics are written in a question and answer format. The answer to each question includes a sentence that will direct the reader to chapters in the profile that will provide more information on the given topic.

Chapter 2

Relevance to Public Health

This chapter provides a health effects summary based on evaluations of existing toxicologic, epidemiologic, and toxicokinetic information. This summary is designed to present interpretive, weight-of-evidence discussions for human health end points by addressing the following questions:

1. What effects are known to occur in humans?
2. What effects observed in animals are likely to be of concern to humans?
3. What exposure conditions are likely to be of concern to humans, especially around hazardous waste sites?

The chapter covers end points in the same order that they appear within the Discussion of Health Effects by Route of Exposure section, by route (inhalation, oral, and dermal) and within route by effect. Human data are presented first, then animal data. Both are organized by duration (acute, intermediate, chronic). *In vitro* data and data from parenteral routes (intramuscular, intravenous, subcutaneous, etc.) are also considered in this chapter.

The carcinogenic potential of the profiled substance is qualitatively evaluated, when appropriate, using existing toxicokinetic, genotoxic, and carcinogenic data. ATSDR does not currently assess cancer potency or perform cancer risk assessments. Minimal Risk Levels (MRLs) for noncancer end points (if derived) and the end points from which they were derived are indicated and discussed.

Limitations to existing scientific literature that prevent a satisfactory evaluation of the relevance to public health are identified in the Chapter 3 Data Needs section.

Interpretation of Minimal Risk Levels

Where sufficient toxicologic information is available, ATSDR has derived MRLs for inhalation and oral routes of entry at each duration of exposure (acute, intermediate, and chronic). These MRLs are not meant to support regulatory action, but to acquaint health professionals with exposure levels at which adverse health effects are not expected to occur in humans.

APPENDIX B

MRLs should help physicians and public health officials determine the safety of a community living near a chemical emission, given the concentration of a contaminant in air or the estimated daily dose in water. MRLs are based largely on toxicological studies in animals and on reports of human occupational exposure.

MRL users should be familiar with the toxicologic information on which the number is based. Chapter 2, "Relevance to Public Health," contains basic information known about the substance. Other sections such as Chapter 3 Section 3.9, "Interactions with Other Substances," and Section 3.10, "Populations that are Unusually Susceptible" provide important supplemental information.

MRL users should also understand the MRL derivation methodology. MRLs are derived using a modified version of the risk assessment methodology that the Environmental Protection Agency (EPA) provides (Barnes and Dourson 1988) to determine reference doses (RfDs) for lifetime exposure.

To derive an MRL, ATSDR generally selects the most sensitive end point which, in its best judgement, represents the most sensitive human health effect for a given exposure route and duration. ATSDR cannot make this judgement or derive an MRL unless information (quantitative or qualitative) is available for all potential systemic, neurological, and developmental effects. If this information and reliable quantitative data on the chosen end point are available, ATSDR derives an MRL using the most sensitive species (when information from multiple species is available) with the highest no-observed-adverse-effect level (NOAEL) that does not exceed any adverse effect levels. When a NOAEL is not available, a lowest-observed-adverse-effect level (LOAEL) can be used to derive an MRL, and an uncertainty factor (UF) of 10 must be employed. Additional uncertainty factors of 10 must be used both for human variability to protect sensitive subpopulations (people who are most susceptible to the health effects caused by the substance) and for interspecies variability (extrapolation from animals to humans). In deriving an MRL, these individual uncertainty factors are multiplied together. The product is then divided into the inhalation concentration or oral dosage selected from the study. Uncertainty factors used in developing a substance-specific MRL are provided in the footnotes of the levels of significant exposure (LSE) tables.

Chapter 3

Health Effects

Tables and Figures for Levels of Significant Exposure (LSE)

Tables and figures are used to summarize health effects and illustrate graphically levels of exposure associated with those effects. These levels cover health effects observed at increasing dose concentrations and durations, differences in response by species, MRLs to humans for noncancer end points, and EPA's estimated range associated with an upper-bound individual lifetime cancer risk of 1 in 10,000 to 1 in 10,000,000. Use the LSE tables and figures for a quick review of the health effects and to locate data for a specific exposure scenario. The LSE tables and figures should always be used in conjunction with the text. All entries in these tables and figures represent studies that provide reliable, quantitative estimates of NOAELs, LOAELs, or Cancer Effect Levels (CELs).

The legends presented below demonstrate the application of these tables and figures. Representative examples of LSE Table 3-1 and Figure 3-1 are shown. The numbers in the left column of the legends correspond to the numbers in the example table and figure.

APPENDIX B

LEGEND**See Sample LSE Table 3-1 (page B-6)**

- (1) **Route of Exposure.** One of the first considerations when reviewing the toxicity of a substance using these tables and figures should be the relevant and appropriate route of exposure. Typically when sufficient data exist, three LSE tables and two LSE figures are presented in the document. The three LSE tables present data on the three principal routes of exposure, i.e., inhalation, oral, and dermal (LSE Tables 3-1, 3-2, and 3-3, respectively). LSE figures are limited to the inhalation (LSE Figure 3-1) and oral (LSE Figure 3-2) routes. Not all substances will have data on each route of exposure and will not, therefore, have all five of the tables and figures.
- (2) **Exposure Period.** Three exposure periods—acute (less than 15 days), intermediate (15–364 days), and chronic (365 days or more)—are presented within each relevant route of exposure. In this example, an inhalation study of intermediate exposure duration is reported. For quick reference to health effects occurring from a known length of exposure, locate the applicable exposure period within the LSE table and figure.
- (3) **Health Effect.** The major categories of health effects included in LSE tables and figures are death, systemic, immunological, neurological, developmental, reproductive, and cancer. NOAELs and LOAELs can be reported in the tables and figures for all effects but cancer. Systemic effects are further defined in the "System" column of the LSE table (see key number 18).
- (4) **Key to Figure.** Each key number in the LSE table links study information to one or more data points using the same key number in the corresponding LSE figure. In this example, the study represented by key number 18 has been used to derive a NOAEL and a Less Serious LOAEL (also see the two "18r" data points in sample Figure 3-1).
- (5) **Species.** The test species, whether animal or human, are identified in this column. Chapter 2, "Relevance to Public Health," covers the relevance of animal data to human toxicity and Section 3.4, "Toxicokinetics," contains any available information on comparative toxicokinetics. Although NOAELs and LOAELs are species specific, the levels are extrapolated to equivalent human doses to derive an MRL.
- (6) **Exposure Frequency/Duration.** The duration of the study and the weekly and daily exposure regimens are provided in this column. This permits comparison of NOAELs and LOAELs from different studies. In this case (key number 18), rats were exposed to "Chemical x" via inhalation for 6 hours/day, 5 days/week, for 13 weeks. For a more complete review of the dosing regimen, refer to the appropriate sections of the text or the original reference paper (i.e., Nitschke et al. 1981).
- (7) **System.** This column further defines the systemic effects. These systems include respiratory, cardiovascular, gastrointestinal, hematological, musculoskeletal, hepatic, renal, and dermal/ocular. "Other" refers to any systemic effect (e.g., a decrease in body weight) not covered in these systems. In the example of key number 18, one systemic effect (respiratory) was investigated.
- (8) **NOAEL.** A NOAEL is the highest exposure level at which no harmful effects were seen in the organ system studied. Key number 18 reports a NOAEL of 3 ppm for the respiratory system, which was used to derive an intermediate exposure, inhalation MRL of 0.005 ppm (see footnote "b").

APPENDIX B

- (9) LOAEL. A LOAEL is the lowest dose used in the study that caused a harmful health effect. LOAELs have been classified into "Less Serious" and "Serious" effects. These distinctions help readers identify the levels of exposure at which adverse health effects first appear and the gradation of effects with increasing dose. A brief description of the specific end point used to quantify the adverse effect accompanies the LOAEL. The respiratory effect reported in key number 18 (hyperplasia) is a Less Serious LOAEL of 10 ppm. MRLs are not derived from Serious LOAELs.
- (10) Reference. The complete reference citation is given in Chapter 9 of the profile.
- (11) CEL. A CEL is the lowest exposure level associated with the onset of carcinogenesis in experimental or epidemiologic studies. CELs are always considered serious effects. The LSE tables and figures do not contain NOAELs for cancer, but the text may report doses not causing measurable cancer increases.
- (12) Footnotes. Explanations of abbreviations or reference notes for data in the LSE tables are found in the footnotes. Footnote "b" indicates that the NOAEL of 3 ppm in key number 18 was used to derive an MRL of 0.005 ppm.

LEGEND**See Sample Figure 3-1 (page B-7)**

LSE figures graphically illustrate the data presented in the corresponding LSE tables. Figures help the reader quickly compare health effects according to exposure concentrations for particular exposure periods.

- (13) Exposure Period. The same exposure periods appear as in the LSE table. In this example, health effects observed within the acute and intermediate exposure periods are illustrated.
- (14) Health Effect. These are the categories of health effects for which reliable quantitative data exists. The same health effects appear in the LSE table.
- (15) Levels of Exposure. Concentrations or doses for each health effect in the LSE tables are graphically displayed in the LSE figures. Exposure concentration or dose is measured on the log scale "y" axis. Inhalation exposure is reported in mg/m³ or ppm and oral exposure is reported in mg/kg/day.
- (16) NOAEL. In this example, the open circle designated 18r identifies a NOAEL critical end point in the rat upon which an intermediate inhalation exposure MRL is based. The key number 18 corresponds to the entry in the LSE table. The dashed descending arrow indicates the extrapolation from the exposure level of 3 ppm (see entry 18 in the table) to the MRL of 0.005 ppm (see footnote "b" in the LSE table).
- (17) CEL. Key number 38m is one of three studies for which CELs were derived. The diamond symbol refers to a CEL for the test species-mouse. The number 38 corresponds to the entry in the LSE table.

APPENDIX B

- (18) Estimated Upper-Bound Human Cancer Risk Levels. This is the range associated with the upper-bound for lifetime cancer risk of 1 in 10,000 to 1 in 10,000,000. These risk levels are derived from the EPA's Human Health Assessment Group's upper-bound estimates of the slope of the cancer dose response curve at low dose levels (q_1^*).
- (19) Key to LSE Figure. The Key explains the abbreviations and symbols used in the figure.

SAMPLE

1 →

Table 3-1. Levels of Significant Exposure to [Chemical x] – Inhalation

Key to figure ^a	Species	Exposure frequency/ duration	System	NOAEL (ppm)	LOAEL (effect)		Reference
					Less serious (ppm)	Serious (ppm)	
INTERMEDIATE EXPOSURE							
3 →	Systemic	↓	↓	↓	↓	↓	↓
4 →	18	Rat	13 wk 5 d/wk 6 hr/d	Resp	3 ^b	10 (hyperplasia)	Nitschke et al. 1981
CHRONIC EXPOSURE							
	Cancer					11 ↓	
	38	Rat	18 mo 5 d/wk 7 hr/d			20	(CEL, multiple organs) Wong et al. 1982
	39	Rat	89–104 wk 5 d/wk 6 hr/d			10	(CEL, lung tumors, nasal tumors) NTP 1982
	40	Mouse	79–103 wk 5 d/wk 6 hr/d			10	(CEL, lung tumors, hemangiosarcomas) NTP 1982

12 →

^a The number corresponds to entries in Figure 3-1.


^b Used to derive an intermediate inhalation Minimal Risk Level (MRL) of 5×10^{-3} ppm; dose adjusted for intermittent exposure and divided by an uncertainty factor of 100 (10 for extrapolation from animal to humans, 10 for human variability).

DRAFT FOR PUBLIC COMMENT

APPENDIX B

SAMPLE

Figure 3-1. Levels of Significant Exposure to [Chemical X] - Inhalation



DRAFT FOR PUBLIC COMMENT

APPENDIX B

APPENDIX B

This page is intentionally blank.

APPENDIX C. ACRONYMS, ABBREVIATIONS, AND SYMBOLS

ACGIH	American Conference of Governmental Industrial Hygienists
ACOEM	American College of Occupational and Environmental Medicine
ADI	acceptable daily intake
ADME	absorption, distribution, metabolism, and excretion
AED	atomic emission detection
AFID	alkali flame ionization detector
AFOSH	Air Force Office of Safety and Health
ALT	alanine aminotransferase
AML	acute myeloid leukemia
AOAC	Association of Official Analytical Chemists
AOEC	Association of Occupational and Environmental Clinics
AP	alkaline phosphatase
APHA	American Public Health Association
AST	aspartate aminotransferase
atm	atmosphere
ATSDR	Agency for Toxic Substances and Disease Registry
AWQC	Ambient Water Quality Criteria
BAT	best available technology
BCF	bioconcentration factor
BEI	Biological Exposure Index
BMD	benchmark dose
BMR	benchmark response
BSC	Board of Scientific Counselors
C	centigrade
CAA	Clean Air Act
CAG	Cancer Assessment Group of the U.S. Environmental Protection Agency
CAS	Chemical Abstract Services
CDC	Centers for Disease Control and Prevention
CEL	cancer effect level
CELDS	Computer-Environmental Legislative Data System
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
CFR	Code of Federal Regulations
Ci	curie
CI	confidence interval
CL	ceiling limit value
CLP	Contract Laboratory Program
cm	centimeter
CML	chronic myeloid leukemia
CPSC	Consumer Products Safety Commission
CWA	Clean Water Act
DHEW	Department of Health, Education, and Welfare
DHHS	Department of Health and Human Services
DNA	deoxyribonucleic acid
DOD	Department of Defense
DOE	Department of Energy
DOL	Department of Labor
DOT	Department of Transportation
DOT/UN/ NA/IMCO	Department of Transportation/United Nations/ North America/Intergovernmental Maritime Dangerous Goods Code

APPENDIX C

DWEL	drinking water exposure level
ECD	electron capture detection
ECG/EKG	electrocardiogram
EEG	electroencephalogram
EEGL	Emergency Exposure Guidance Level
EPA	Environmental Protection Agency
F	Fahrenheit
F ₁	first-filial generation
FAO	Food and Agricultural Organization of the United Nations
FDA	Food and Drug Administration
FEMA	Federal Emergency Management Agency
FIFRA	Federal Insecticide, Fungicide, and Rodenticide Act
FPD	flame photometric detection
fpm	feet per minute
FR	Federal Register
FSH	follicle stimulating hormone
g	gram
GC	gas chromatography
gd	gestational day
GLC	gas liquid chromatography
GPC	gel permeation chromatography
HPLC	high-performance liquid chromatography
HRGC	high resolution gas chromatography
HSDB	Hazardous Substance Data Bank
IARC	International Agency for Research on Cancer
IDLH	immediately dangerous to life and health
ILO	International Labor Organization
IRIS	Integrated Risk Information System
K _d	adsorption ratio
kg	kilogram
kgg	metric ton
K _{oc}	organic carbon partition coefficient
K _{ow}	octanol-water partition coefficient
L	liter
LC	liquid chromatography
LC ₅₀	lethal concentration, 50% kill
LC _{Lo}	lethal concentration, low
LD ₅₀	lethal dose, 50% kill
LD _{Lo}	lethal dose, low
LDH	lactic dehydrogenase
LH	lutinizing hormone
LOAEL	lowest-observed-adverse-effect level
LSE	Levels of Significant Exposure
LT ₅₀	lethal time, 50% kill
m	meter
MA	mandelic acid
MAL	maximum allowable level
mCi	millicurie
MCL	maximum contaminant level
MCLG	maximum contaminant level goal
MF	modifying factor

APPENDIX C

MFO	mixed function oxidase
mg	milligram
mL	milliliter
mm	millimeter
mmHg	millimeters of mercury
mmol	millimole
mppcf	millions of particles per cubic foot
MRL	Minimal Risk Level
MS	mass spectrometry
NAAQS	National Ambient Air Quality Standard
NAS	National Academy of Science
NATICH	National Air Toxics Information Clearinghouse
NATO	North Atlantic Treaty Organization
NCE	normochromatic erythrocytes
NCEH	National Center for Environmental Health
NCI	National Cancer Institute
ND	not detected
NFPA	National Fire Protection Association
ng	nanogram
NHANES	National Health and Nutrition Examination Survey
NIEHS	National Institute of Environmental Health Sciences
NIOSH	National Institute for Occupational Safety and Health
NIOSHTIC	NIOSH's Computerized Information Retrieval System
NLM	National Library of Medicine
nm	nanometer
nmol	nanomole
NOAEL	no-observed-adverse-effect level
NOES	National Occupational Exposure Survey
NOHS	National Occupational Hazard Survey
NPD	nitrogen phosphorus detection
NPDES	National Pollutant Discharge Elimination System
NPL	National Priorities List
NR	not reported
NRC	National Research Council
NS	not specified
NSPS	New Source Performance Standards
NTIS	National Technical Information Service
NTP	National Toxicology Program
ODW	Office of Drinking Water, EPA
OERR	Office of Emergency and Remedial Response, EPA
OHM/TADS	Oil and Hazardous Materials/Technical Assistance Data System
OPP	Office of Pesticide Programs, EPA
OPPT	Office of Pollution Prevention and Toxics, EPA
OPPTS	Office of Prevention, Pesticides and Toxic Substances, EPA
OR	odds ratio
OSHA	Occupational Safety and Health Administration
OSW	Office of Solid Waste, EPA
OTS	Office of Toxic Substances
OW	Office of Water
OWRS	Office of Water Regulations and Standards, EPA
PAH	polycyclic aromatic hydrocarbon

APPENDIX C

PBPD	physiologically based pharmacodynamic
PBPK	physiologically based pharmacokinetic
PCE	polychromatic erythrocytes
PEL	permissible exposure limit
pg	picogram
PGA	phenylglyoxylic acid
PHS	Public Health Service
PID	photo ionization detector
pmol	picomole
PMR	proportionate mortality ratio
ppb	parts per billion
ppbv	parts per billion volume per volume
ppm	parts per million
ppmv	parts per million volume per volume
ppt	parts per trillion
PSNS	pretreatment standards for new sources
RBC	red blood cell
REL	recommended exposure level/limit
RfC	reference concentration
RfD	reference dose
RNA	ribonucleic acid
RQ	reportable quantity
RTECS	Registry of Toxic Effects of Chemical Substances
SARA	Superfund Amendments and Reauthorization Act
SCE	sister chromatid exchange
SGOT	serum glutamic oxaloacetic transaminase
SGPT	serum glutamic pyruvic transaminase
SIC	standard industrial classification
SIM	selected ion monitoring
SMCL	secondary maximum contaminant level
SMR	standardized mortality ratio
SNARL	suggested no adverse response level
SPEGL	Short-Term Public Emergency Guidance Level
STEL	short term exposure limit
STORET	Storage and Retrieval
TD ₅₀	toxic dose, 50% specific toxic effect
TLV	threshold limit value
TOC	total organic carbon
TPQ	threshold planning quantity
TRI	Toxics Release Inventory
TSCA	Toxic Substances Control Act
TWA	time-weighted average
UF	uncertainty factor
U.S.	United States
USDA	United States Department of Agriculture
USGS	United States Geological Survey
VOC	volatile organic compound
WBC	white blood cell
WHO	World Health Organization

APPENDIX C

>	greater than
≥	greater than or equal to
=	equal to
<	less than
≤	less than or equal to
%	percent
α	alpha
β	beta
γ	gamma
δ	delta
μm	micrometer
μg	microgram
q ₁ *	cancer slope factor
-	negative
+	positive
(+)	weakly positive result
(-)	weakly negative result

APPENDIX C

This page is intentionally blank.

APPENDIX D. INDEX

absorbed dose.....	101, 103, 120, 124
adipose tissue.....	96, 97, 102, 113, 121, 124, 134, 163, 168, 171, 175, 178
adsorption.....	154, 167
aerobic.....	157, 167
alanine aminotransferase.....	49
ambient air.....	158
anaerobic.....	147, 157, 167
aspartate aminotransferase.....	49
bioaccumulation.....	167
bioconcentration factor.....	155
biodegradation.....	147, 154, 157, 179
biological half-time.....	113
biomarker.....	18, 53, 119, 120, 122, 133, 178
biomarkers.....	119, 120, 122, 129, 133, 171, 178
blood cell count.....	84
breast milk.....	5, 96, 164
cancer.....	5, 12, 70, 71, 72, 73, 87, 118, 124, 134, 182
carcinogen.....	12, 114, 130, 134, 181, 185
carcinogenic.....	12, 23, 69, 70, 73, 74, 88, 104, 113, 130, 181, 185
carcinogenicity.....	12, 73, 74, 75, 88, 98, 133, 178, 181
carcinoma.....	74, 88
cardiovascular.....	47, 76, 88
cardiovascular effects.....	47, 76
CCI (see color confusion index).....	16, 18, 54, 55, 60, 61
chromosomal aberrations.....	89, 93, 113
clearance.....	101, 110, 120
color confusion index (see CCI).....	16, 60, 61
color discrimination.....	9, 16, 17, 19, 53, 54, 61, 121, 132, 181
death.....	11, 23, 53, 68, 70, 72, 75, 185
deoxyribonucleic acid (see DNA).....	89
dermal effects.....	89
developmental effects.....	10, 14, 19, 68, 69, 87, 89, 118, 128, 131, 135
DNA (see deoxyribonucleic acid).....	89, 91, 92, 93, 113, 120, 122, 178
dopamine.....	19, 20, 86, 87
elimination half-time.....	113
enantiomer.....	75, 100, 101, 114, 115, 134
endocrine.....	50, 76, 88, 115, 116, 129, 133
endocrine effects.....	50, 67, 133
fetus.....	11, 117
follicle stimulating hormone.....	51, 67, 116
gastrointestinal effects.....	47, 76, 84
general population.....	24, 119, 147, 163, 165, 168, 169, 178, 185
genotoxic.....	23, 93, 113, 114, 122, 130
genotoxicity.....	89, 93, 122, 130
groundwater.....	3, 155, 156, 160, 163, 168, 179
half-life.....	96, 102, 119, 120, 121, 124, 147, 154, 155, 156
hematological effects.....	47, 48, 84, 130
hematopoietic.....	12, 71, 73
hepatic effects.....	10, 49, 84

APPENDIX D

hydroxyl radical 147, 156

immune system 21, 132

immunological 20, 23, 85

immunological effects 20, 85

K_{ow} 139, 155

LD₅₀ 75

leukemia 69, 70, 71, 72

lymphatic 12, 71, 73

menstrual 67

metabolic effects 25

micronuclei 93

milk 119, 161, 164, 168

mucociliary 47

musculoskeletal effects 48

neonatal 16, 69

neoplastic 12, 74, 88, 113, 130

neurobehavioral 15, 53, 56, 62, 116, 121, 129, 132, 135

neurochemical 16

neurodevelopmental 11, 15, 69, 87, 118, 128, 131

neurological effects 9, 15, 16, 18, 19, 20, 52, 53, 66, 86, 89, 124, 126, 129, 132, 134, 135

non-Hodgkin’s lymphoma 70

ocular effects 51, 76, 89

odds ratio 60, 64, 67, 68, 73

partition coefficients 154

pharmacodynamic 103

pharmacokinetic 74, 101, 102, 103, 104, 105, 117

photolysis 155

rate constant 156

renal effects 50, 85, 88

reproductive effects 11, 66, 67, 86, 87, 89, 116, 128, 131, 133

respiratory effects 16, 25, 76

retention 57, 93, 134

solubility 154, 155, 166

spermatozoa 11, 19, 20, 86, 87, 128

systemic effects 15, 25, 76, 88

thyroid 51, 116

thyroid stimulating hormone 51, 116

toxicokinetic 23, 104, 106, 119, 126, 130, 134

tumors 12, 74, 88, 106, 108, 113, 114, 130

vapor pressure 153

volatilization 154, 167

weaning 88

