

**Coal workers' pneumoconiosis: Number of deaths, death rates (per million population), and years of potential life lost (YPLL) by state, U.S. residents age 15 and over, 1995–2004**

State	No. of Deaths Rank		Death Rate				YPLL to Life Expectancy			
			Crude	Rank	Age-Adjusted	Rank	Total	Rank	YPLL/death	Rank
Alabama	106	10	3.00	11	3.01	12	1,055	11	10.0	25
Alaska	1	45	0.21	43	0.73	25	6	49	6.3	49
Arizona	34	24	0.86	23	0.87	23	281	28	8.3	41
Arkansas	41	20	1.94	15	1.77	15	442	21	10.8	19
California	78	14	0.30	37	0.34	36	932	13	11.9	11
Colorado	83	12	2.45	13	3.25	11	789	14	9.5	32
Connecticut	7	35	0.26	40	0.23	43	73	35	10.4	21
Delaware	3	39	0.48	30	0.50	32	30	40	9.9	26
District of Columbia	2	43	0.42	32	0.46	33	17	44	8.7	39
Florida	166	8	1.29	17	0.97	20	1,642	9	9.9	26
Georgia	27	25	0.42	32	0.55	31	303	25	11.2	15
Hawaii	-	-	-	-	-	-	-	-	-	-
Idaho	6	37	0.60	27	0.64	28	84	34	14.0	9
Illinois	228	6	2.34	14	2.39	14	2,294	7	10.1	24
Indiana	141	9	2.95	12	3.01	12	1,369	10	9.7	29
Iowa	35	23	1.49	16	1.16	19	285	27	8.2	44
Kansas	17	32	0.80	24	0.79	24	296	26	17.4	5
Kentucky	863	4	26.49	3	27.70	3	10,088	4	11.7	12
Louisiana	24	27	0.69	25	0.72	26	582	18	24.2	2
Maine	1	45	0.10	48	0.08	49	7	48	6.8	48
Maryland	54	18	1.29	17	1.47	16	518	19	9.6	30
Massachusetts	1	45	0.02	50	0.02	50	14	45	14.3	8
Michigan	68	16	0.87	22	0.90	22	705	16	10.4	21
Minnesota	4	38	0.10	48	0.11	47	67	37	16.8	6
Mississippi	96	11	4.33	9	4.43	8	3,170	6	33.0	1
Missouri	20	30	0.45	31	0.42	35	223	31	11.1	16
Montana	9	34	1.24	20	1.17	18	67	37	7.5	47
Nebraska	3	39	0.22	42	0.20	45	24	42	7.9	45
Nevada	3	39	0.19	45	0.26	41	30	40	9.9	26
New Hampshire	3	39	0.31	36	0.33	37	24	42	7.9	45
New Jersey	40	22	0.60	27	0.57	29	364	23	9.1	34
New Mexico	47	19	3.34	10	3.77	10	391	22	8.3	41
New York	41	20	0.27	39	0.27	39	465	20	11.3	14
North Carolina	80	13	1.25	19	1.39	17	760	15	9.5	32
North Dakota	1	45	0.19	45	0.17	46	9	46	8.5	40
Ohio	406	5	4.52	8	4.36	9	3,660	5	9.0	37
Oklahoma	26	26	0.95	21	0.93	21	326	24	12.5	10
Oregon	7	35	0.26	40	0.25	42	73	35	10.4	21
Pennsylvania	4,234	1	42.81	2	34.79	2	38,400	1	9.1	34
Rhode Island	1	45	0.12	47	0.09	48	6	49	6.1	50
South Carolina	21	29	0.65	26	0.71	27	234	29	11.1	16
South Dakota	2	43	0.34	35	0.33	37	48	39	23.9	3
Tennessee	208	7	4.58	7	4.84	7	2,232	8	10.7	20
Texas	58	17	0.36	34	0.43	34	1,022	12	17.6	4
Utah	76	15	4.59	6	6.53	5	685	17	9.0	37
Vermont	1	45	0.20	44	0.21	44	8	47	8.3	41
Virginia	973	3	17.24	4	19.82	4	11,191	3	11.5	13
Washington	24	27	0.52	29	0.56	30	231	30	9.6	30
West Virginia	2,005	2	135.40	1	118.37	1	21,865	2	10.9	18
Wisconsin	12	33	0.28	38	0.27	39	181	32	15.1	7
Wyoming	19	31	4.83	5	5.40	6	173	33	9.1	34

- indicates no deaths listed.

NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Census Bureau.