

2005Table 2-1. Coal workers' pneumoconiosis: Number of deaths by sex, race, and age, and median age at death, U.S. residents age 15 and over, 1993–2002

Year	No. of Deaths	Underlying Cause (%)	Sex		Race			Age Group (yrs)							Median Age (yrs)	
			Male	Female	White	Black	Other	15–24	25–34	35–44	45–54	55–64	65–74	75–84		85+
1993	1,631	35.1	1,616	15	1,567	62	2	-	2	4	24	111	404	777	309	78.0
1994	1,478	33.3	1,465	13	1,434	42	2	-	-	6	15	78	345	741	293	79.0
1995	1,413	37.7	1,407	6	1,372	41	-	-	3	4	28	75	354	614	335	79.0
1996	1,417	37.8	1,407	10	1,375	39	3	-	-	4	22	51	327	673	340	79.0
1997	1,297	37.5	1,283	14	1,267	30	-	-	2	7	24	54	266	623	321	80.0
1998	1,103	38.2	1,093	10	1,066	35	2	-	1	4	15	51	235	503	294	80.0
1999	1,003	40.9	998	5	971	31	1	-	1	3	10	52	174	459	304	81.0
2000	950	41.1	940	10	924	25	1	2	1	7	18	42	171	382	327	81.0
2001	889	41.4	878	11	865	23	1	2	-	8	19	41	159	374	286	81.0
2002	858	41.3	847	11	837	20	1	1	3	5	18	43	124	370	294	81.0
TOTAL	12,039	37.9	11,934	105	11,678	348	13	5	13	52	193	598	2,559	5,516	3,103	79.0

- indicates no deaths listed.

NOTE: Slight variations in numbers may exist when compared to previous WoRLD Surveillance Reports. See appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data.