

Table 6. Estimated numbers of AIDS cases in adult and adolescent Hispanics, by transmission category and place of birth, 2006—United States and dependent areas

Transmission category	Place of birth											
	United States		Central/South America		Cuba		Mexico		Puerto Rico		Total ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact	1,172	45	402	49	89	62	790	59	249	18	3,337	43
Injection drug use	569	22	83	10	13	9	133	10	533	40	1,650	21
Male-to-male sexual contact and injection drug use	158	6	25	3	6	4	59	4	63	5	357	5
High-risk heterosexual contact ^b	661	25	296	36	35	24	342	26	487	36	2,282	30
Other ^c	47	2	8	1	0	0	9	1	14	1	103	1
Total^d	2,608	100	814	100	145	100	1,334	100	1,346	100	7,728	100

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

^a Includes 222 persons whose place of birth is not among those listed and 1,260 persons whose place of birth is unknown.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^d Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.