

**Table 9. Estimated numbers of persons living with HIV/AIDS at the end of 2006, by race/ethnicity, sex, and transmission category—33 states with confidential name-based HIV infection reporting**

Transmission category	White, not Hispanic		Black, not Hispanic		Hispanic		Asian/Pacific Islander		American Indian/Alaska Native		Total <sup>a</sup>	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>												
Male-to-male sexual contact	108,259	77	70,242	49	36,043	57	1,669	68	938	62	218,676	62
Injection drug use	12,328	9	31,339	22	14,472	23	264	11	216	14	59,077	17
Male-to-male sexual contact and injection drug use	11,283	8	9,587	7	3,720	6	93	4	205	13	25,085	7
High-risk heterosexual contact <sup>b</sup>	6,667	5	31,140	22	8,788	14	395	16	143	9	47,562	13
Other <sup>c</sup>	1,458	1	1,492	1	404	1	30	1	17	1	3,424	1
Subtotal	139,994	100	143,799	100	63,427	100	2,450	100	1,519	100	353,825	100
<b>Female adult or adolescent</b>												
Injection drug use	8,262	33	19,136	23	5,528	28	106	15	175	30	33,470	26
High-risk heterosexual contact <sup>b</sup>	16,338	65	63,204	75	14,132	71	573	82	392	68	95,403	73
Other <sup>c</sup>	450	2	1,469	2	344	2	24	3	12	2	2,321	2
Subtotal	25,050	100	83,809	100	20,004	100	703	100	578	100	131,195	100
<b>Child (&lt;13 yrs at diagnosis)</b>												
Perinatal	801	84	4,064	93	1,183	92	28	82	20	95	6,143	92
Other <sup>d</sup>	154	16	283	7	107	8	6	18	1	5	560	8
Subtotal	955	100	4,347	100	1,290	100	34	100	21	100	6,703	100
<b>Total<sup>e</sup></b>	<b>166,000</b>		<b>231,957</b>		<b>84,720</b>		<b>3,187</b>		<b>2,119</b>		<b>491,727</b>	

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

Data include persons with a diagnosis of HIV infection (not AIDS), a diagnosis of HIV infection and a later diagnosis of AIDS, or concurrent diagnoses of HIV infection and AIDS.

See Technical Notes for the list of 33 states that have had laws or regulations requiring confidential name-based HIV infection reporting since at least 2003.

<sup>a</sup> Includes 3,745 persons of unknown race or multiple races.

<sup>b</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>c</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>d</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

<sup>e</sup> Includes 5 persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.