

**2005Table 7-5. Malignant mesothelioma: Number of deaths, mortality rates (per million population), and years of potential life lost (YPLL) by state, U.S. residents age 15 and over, 1999–2002**

State	No. of Deaths		Crude Death Rate		Age-Adjusted Death Rate		YPLL to Life Expectancy			
	Deaths	Rank	Rate	Rank	Rate	Rank	Total	Rank	YPLL/death	Rank
Alabama	125	24	8.84	41	8.60	43	1,909	24	15.3	7
Alaska	20	50	11.37	26	25.28	1	251	50	12.5	49
Arizona	152	22	9.51	36	9.16	41	1,946	23	12.8	47
Arkansas	50	40	5.89	50	5.47	51	854	39	17.1	2
California	1,001	1	9.43	37	10.75	30	14,030	1	14.0	30
Colorado	107	28	7.73	46	9.61	38	1,580	28	14.8	11
Connecticut	123	25	11.35	27	10.45	32	1,670	26	13.6	38
Delaware	42	44	16.50	5	16.45	4	547	44	13.0	44
District of Columbia	12	51	6.30	49	6.67	50	209	51	17.4	1
Florida	685	2	12.98	18	10.12	34	8,984	2	13.1	43
Georgia	145	23	5.56	51	7.01	49	2,092	22	14.4	18
Hawaii	31	45	7.91	44	7.55	47	433	45	14.0	30
Idaho	49	41	12.15	23	12.83	21	706	41	14.4	18
Illinois	507	5	12.95	19	13.44	17	7,287	6	14.4	18
Indiana	211	15	11.00	30	11.12	28	3,105	14	14.7	12
Iowa	105	29	11.19	28	9.99	35	1,518	29	14.5	17
Kansas	102	30	12.07	24	11.52	26	1,274	32	12.5	49
Kentucky	119	27	9.09	40	9.28	39	1,802	25	15.1	8
Louisiana	178	19	12.82	20	13.51	16	2,765	18	15.5	6
Maine	96	32	22.86	1	20.83	2	1,380	31	14.4	18
Maryland	208	16	12.28	22	13.35	18	2,859	17	13.7	35
Massachusetts	330	10	15.98	8	15.25	9	4,128	11	12.5	49
Michigan	362	9	11.54	25	11.72	25	5,213	9	14.4	18
Minnesota	214	14	13.63	14	14.11	14	3,050	15	14.3	23
Mississippi	65	36	7.31	48	7.52	48	955	35	14.7	12
Missouri	192	18	10.75	31	10.28	33	2,631	19	13.7	35
Montana	48	42	16.97	2	15.73	7	703	42	14.7	12
Nebraska	67	34	12.40	21	11.79	24	985	34	14.7	12
Nevada	66	35	10.20	33	11.51	27	940	36	14.2	26
New Hampshire	57	38	14.19	12	14.97	10	794	40	13.9	33
New Jersey	449	8	16.69	3	16.02	6	5,746	8	12.8	47
New Mexico	57	38	10.55	32	10.70	31	914	37	16.0	3
New York	612	4	10.09	34	9.97	36	8,286	3	13.5	40
North Carolina	203	17	7.86	45	8.17	44	3,050	15	15.0	10
North Dakota	27	46	13.40	15	12.25	23	407	46	15.1	8
Ohio	468	7	13.01	17	12.45	22	6,508	7	13.9	33
Oklahoma	88	33	8.36	43	7.99	46	1,203	33	13.7	35
Oregon	171	20	15.57	9	15.28	8	2,438	21	14.3	23
Pennsylvania	637	3	16.06	7	13.59	15	8,231	4	12.9	46
Rhode Island	48	42	14.04	13	12.88	20	681	43	14.2	26
South Carolina	122	26	9.43	37	9.76	37	1,642	27	13.5	40
South Dakota	21	49	9.11	39	8.07	45	293	49	14.0	30
Tennessee	158	21	8.63	42	8.75	42	2,483	20	15.7	5
Texas	500	6	7.68	47	9.23	40	7,291	5	14.6	16
Utah	65	36	9.73	35	13.16	19	883	38	13.6	38
Vermont	22	48	11.06	29	10.76	29	347	47	15.8	4
Virginia	301	12	13.16	16	14.56	11	4,296	10	14.3	23
Washington	309	11	16.43	6	17.89	3	4,002	12	13.0	44
West Virginia	98	31	16.59	4	14.44	12	1,383	30	14.1	29
Wisconsin	248	13	14.49	11	14.18	13	3,293	13	13.3	42
Wyoming	24	47	15.38	10	16.03	5	342	48	14.2	26

– indicates no deaths listed.

NOTE: Slight variations in numbers may exist when compared to previous WoRLD Surveillance Reports. See appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Bureau of the Census.