

**2005Table 3-5. Silicosis: Number of deaths, death rates (per million population), and years of potential life lost (YPLL) by state, U.S. residents age 15 and over, 1993–2002**

State	No. of Deaths		Crude Death Rate		Age-Adjusted Death Rate		YPLL to Life Expectancy			
	Deaths	Rank	Rate	Rank	Rate	Rank	Total	Rank	YPLL/death	Rank
Alabama	30	23	0.87	23	0.88	23	424	22	14.1	13
Alaska	3	45	0.66	34	1.84	9	29	47	9.6	46
Arizona	37	16	1.03	16	1.04	17	423	23	11.4	36
Arkansas	15	33	0.75	27	0.71	31	333	25	22.2	1
California	85	6	0.34	46	0.39	45	1,073	5	12.6	23
Colorado	70	8	2.25	5	2.81	3	799	11	11.4	36
Connecticut	25	27	0.96	19	0.87	24	306	28	12.2	27
Delaware	5	44	0.84	24	0.90	20	49	45	9.9	45
District of Columbia	3	45	0.69	29	0.71	31	60	44	19.8	3
Florida	50	12	0.42	45	0.32	47	637	12	12.7	21
Georgia	28	25	0.47	42	0.61	37	308	27	11.0	41
Hawaii	1	49	0.11	51	0.11	51	22	48	22.0	2
Idaho	14	35	1.48	10	1.62	11	132	38	9.4	48
Illinois	64	10	0.68	32	0.69	34	883	8	13.8	16
Indiana	41	13	0.88	22	0.90	20	575	13	14.0	14
Iowa	22	30	0.97	18	0.82	25	252	32	11.5	34
Kansas	9	40	0.44	44	0.43	43	143	35	15.8	6
Kentucky	40	15	1.28	11	1.33	12	558	14	14.0	14
Louisiana	31	21	0.91	20	0.96	19	517	15	16.7	5
Maine	7	42	0.69	29	0.66	36	108	40	15.4	8
Maryland	27	26	0.67	33	0.75	29	310	26	11.5	34
Massachusetts	24	28	0.49	41	0.46	42	278	31	11.6	31
Michigan	86	5	1.12	14	1.15	14	972	7	11.3	38
Minnesota	41	13	1.11	15	1.08	15	419	24	10.2	43
Mississippi	12	36	0.56	39	0.57	39	234	33	19.5	4
Missouri	36	19	0.84	24	0.79	26	481	16	13.4	18
Montana	15	33	2.14	6	2.04	7	134	36	8.9	49
Nebraska	2	48	0.15	50	0.13	50	15	49	7.3	50
Nevada	12	36	0.89	21	1.04	17	122	39	10.1	44
New Hampshire	3	45	0.32	47	0.33	46	35	46	11.6	31
New Jersey	37	16	0.57	38	0.55	41	433	20	11.7	30
New Mexico	31	21	2.35	4	2.57	5	427	21	13.8	16
New York	89	4	0.62	36	0.60	38	1,123	4	12.6	23
North Carolina	68	9	1.15	13	1.19	13	839	10	12.3	26
North Dakota	1	49	0.20	49	0.14	49	6	51	6.1	51
Ohio	175	2	1.97	8	1.91	8	2,209	2	12.6	23
Oklahoma	12	36	0.46	43	0.43	43	134	36	11.2	40
Oregon	20	32	0.77	26	0.74	30	227	34	11.3	38
Pennsylvania	318	1	3.30	2	2.75	4	3,452	1	10.9	42
Rhode Island	8	41	1.01	17	0.89	22	93	42	11.6	31
South Carolina	22	30	0.72	28	0.78	27	281	29	12.8	20
South Dakota	7	42	1.22	12	1.06	16	86	43	12.2	27
Tennessee	30	23	0.69	29	0.71	31	444	19	14.8	10
Texas	99	3	0.66	34	0.77	28	1,569	3	15.8	6
Utah	37	16	2.43	3	3.23	2	471	17	12.7	21
Vermont	10	39	2.10	7	2.22	6	96	41	9.6	46
Virginia	33	20	0.61	37	0.69	34	471	17	14.3	11
Washington	23	29	0.51	40	0.57	39	280	30	12.2	27
West Virginia	58	11	3.90	1	3.48	1	862	9	14.9	9
Wisconsin	77	7	1.87	9	1.82	10	1,006	6	13.1	19
Wyoming	1	49	0.26	48	0.27	48	14	50	14.3	11

– indicates no deaths listed.

NOTE: Slight variations in numbers may exist when compared to previous WoRLD Surveillance Reports. See appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Bureau of the Census.