

The Northwest Corner of the Atlantic and Rapid Climate Change Bibliography

Matthew W. Hecht¹, Frank O. Bryan², Mathew Maltrud¹ and Richard D. Smith³

¹Los Alamos National Laboratory, Los Alamos, NM

²National Center for Atmospheric Research, Boulder CO

³Los Alamos, NM

Many of the North Atlantic regional modeling results are from [3]. Partial bottom cells are discussed in [1]. The benefit of smoothing of topography is discussed in [2], [9] and [4]. Observed position of the Gulf Stream is from [20], observed velocities off the Grand Banks at around 700m depth are from [15]. Version 3 of the Community Climate System Model is documented in the June 1, 2006 volume of *J. Clim.*. Others papers referred to in the poster include [5], [6], [7], [8], [10], [12], [11], [13], [14], [17], [16], [18] and [19].

References

- [1] A. Adcroft, C. Hill, and J. Marshall. Representation of topography by shaved cells in a height coordinate ocean model. *Mon. Weather Rev.*, 125:2293–2315, 1997.
- [2] Barnier Bernard, Gurvan Madec, Thierry Penduff, Jean-Marc Molines, Anne-Marie Treguier, Aike Beckmann, Arne Biastoch, Claus Böning, Joachim Dengg, Sergei Gulev, Julien Le Sommer, Elizabeth Remy, Claude Talandier, Sébastien Theetten, Mathew Maltrud, Julie McClean, and Beverly De Cuevas. Recent progress in modelling the global ocean circulation at eddy permitting resolution. *Ocean Dynamics*, pages submitted August 31, 2005, 2006.
- [3] F. O. Bryan, M. W. Hecht, and R. D. Smith. Resolution convergence and sensitivity studies with North Atlantic circulation models. Part I: The western boundary current system. *Ocean Modelling*, *in press*, 2006.
- [4] W. K. Dewar. Topography and barotropic transport control by bottom friction. *J. Marine Res.*, 56(2):295 – 328, 1998.
- [5] Scott C. Doney and Matthew W. Hecht. Antarctic bottom water formation and deep water chlorofluorocarbon distributions in a global ocean climate model. *J. Phys. Oceanogr.*, 32:1642–1666, 2002.
- [6] C. O. Iselin. A study of the circulation of the western North Atlantic. *Pap. Phys. Oceanogr. Meteorol.*, 4:101, 1936.
- [7] Mathew E. Maltrud and Julie L. McClean. An eddy resolving $1/10^\circ$ ocean simulation. *Ocean Modelling*, 8(1–2):31–54, 2005.
- [8] Tamay M. Özgökmen, Eric P. Chassignet, and Afonso M. Paiva. Impact of wind forcing, bottom topography, and inertia on midlatitude jet separation in a quasigeostrophic model. *J. Phys. Oceanogr.*, 27:2460–2476, 1997.

- [9] Thierry Penduff, Bernard Barnier, Marie-Aurélie Kerbiriou, and Jacques Verron. How topographic smoothing contributes to differences between the eddy flows simulated by sigma- and geopotential-coordinate models. *J. Phys. Oceanogr.*, 32(1):122–137, 2002.
- [10] S. Rahmstorf. Ocean circulation and climate during the past 120,000 years. *Nature*, 419(6903):207 – 14, 2002.
- [11] M. Roberts and D. Marshall. Do we require adiabatic dissipation schemes in eddy-resolving ocean models? *Journal of physical oceanography*, 28(10):2050 – 63, 1998.
- [12] S. G. Robinson, M. A. Maslin, and I. N. McCave. Magnetic-susceptibility variations in Upper Pleistocene deep-sea sediments of the NE Atlantic - Implications for ice rafting and paleocirculation at the last glacial maximum. *Paleoceanography*, 10(2):221 – 250, 1995.
- [13] T. Rossby. The North Atlantic Current and surrounding waters: At the crossroads. *Reviews of Geophysics*, 34(4):463–481, 1996.
- [14] T. Rossby, C. N. Flagg, and K. Donohue. Interannual variations in upper-ocean transport by the Gulf Stream and adjacent waters between New Jersey and Bermuda. *J. Marine Res.*, 63(1):203–226, 2005.
- [15] Friedrich A. Schott, Rainer Zantopp, Lothar Stramma, Marcus Dengler, Jürgen Fischer, and Mathieu Wibaux. Circulation and deep-water export at the western exit of the subpolar North Atlantic. *J. Phys. Oceanogr.*, 34(4):817–843, 2004.
- [16] Richard D. Smith and Peter R. Gent. Anisotropic GM parameterization for ocean models. *J. Phys. Oceanogr.*, 34:2541–2564, 2004.
- [17] Richard D. Smith, Mathew E. Maltrud, Frank O. Bryan, and Matthew W. Hecht. Numerical simulation of the North Atlantic Ocean at $\frac{1}{10}^\circ$. *J. Phys. Oceanogr.*, 30:1532–1561, 2000.
- [18] R. J. Stouffer, J. Yin, J. M. Gregory, K. W. Dixon, M. J. Spelman, W. Hurlin, A. J. Weaver, M. Eby, G. M. Flato, H. Hasumi, A. Hu, J. H. Jungclaus, I. V. Kamenkovich, A. Levermann, M. Montoya, S. Murakami, S. Nawrath, A. Oka, W. R. Peltier, D. Y. Robitaille, A. Sokolov, G. Vettoretti, and S. L. Weber. Investigating the cause of the response of the thermohaline circulation to past and future climate changes. *Journal of climate*, 19(8):1365 – 87, 2006.
- [19] J. D. Thompson and W. J. Schmitz. A limited-area model of the Gulf Stream: Design, initial experiments, and model-data intercomparison. *Journal of physical oceanography*, 19(6):791 – 814, 1989.
- [20] D. R. Watts, K. L. Tracey, J. M. Bane, and T. J. Shay. Gulf Stream path and thermocline structure near 74°W and 68°W. *J. Geophys. Res.*, 100:18291–18312, 1995.