

Hepatitis B Virus

- Is found in an infected person's blood, semen, and vaginal fluids.
- Can be spread by unprotected sex.
- Can be spread by sharing injection drug needles or equipment.
- Symptoms can include nausea, fatigue, yellowing skin (jaundice), and stomach or muscle pain.



Hepatitis B Virus

- Usually just lasts a few months, but sometimes turns into a long-term infection.
- Can be prevented
 - ✓ With a vaccine.
 - ✓ By practicing safer sex.
 - ✓ By not sharing drug needles or equipment.



www.hepatitis.va.gov
June 2005

Hepatitis B Virus

- Is found in an infected person's blood, semen, and vaginal fluids.
- Can be spread by unprotected sex.
- Can be spread by sharing injection drug needles or equipment.
- Symptoms can include nausea, fatigue, yellowing skin (jaundice), and stomach or muscle pain.



Hepatitis B Virus

- Usually just lasts a few months, but sometimes turns into a long-term infection.
- Can be prevented
 - ✓ With a vaccine.
 - ✓ By practicing safer sex.
 - ✓ By not sharing drug needles or equipment.



www.hepatitis.va.gov
June 2005

Hepatitis B Virus

- Is found in an infected person's blood, semen, and vaginal fluids.
- Can be spread by unprotected sex.
- Can be spread by sharing injection drug needles or equipment.
- Symptoms can include nausea, fatigue, yellowing skin (jaundice), and stomach or muscle pain.



Hepatitis B Virus

- Usually just lasts a few months, but sometimes turns into a long-term infection.
- Can be prevented
 - ✓ With a vaccine.
 - ✓ By practicing safer sex.
 - ✓ By not sharing drug needles or equipment.



www.hepatitis.va.gov
June 2005

Hepatitis B Virus

- Is found in an infected person's blood, semen, and vaginal fluids.
- Can be spread by unprotected sex.
- Can be spread by sharing injection drug needles or equipment.
- Symptoms can include nausea, fatigue, yellowing skin (jaundice), and stomach or muscle pain.



Hepatitis B Virus

- Usually just lasts a few months, but sometimes turns into a long-term infection.
- Can be prevented
 - ✓ With a vaccine.
 - ✓ By practicing safer sex.
 - ✓ By not sharing drug needles or equipment.



www.hepatitis.va.gov
June 2005

Hepatitis B Virus

- Is found in an infected person's blood, semen, and vaginal fluids.
- Can be spread by unprotected sex.
- Can be spread by sharing injection drug needles or equipment.
- Symptoms can include nausea, fatigue, yellowing skin (jaundice), and stomach or muscle pain.



Hepatitis B Virus

- Usually just lasts a few months, but sometimes turns into a long-term infection.
- Can be prevented
 - ✓ With a vaccine.
 - ✓ By practicing safer sex.
 - ✓ By not sharing drug needles or equipment.



www.hepatitis.va.gov
June 2005