



Soil Quality
National
Technology
Development
Team

September 2008

Soil Quality - Agronomy Technical Note No. ____

DRAFT

Soil Quality: Managing Cool, Wet Soils

Draft Issued September 2008

Soil Quality - Agronomy Technical Note No. ____ was written by Ray Archuleta, Soil Quality National Technology Development Team, East National Technology Support Center, Natural Resources Conservation Service (NRCS), Greensboro, North Carolina. Contributing authors and reviewers include Jon Stika, North Dakota NRCS; Chris Lawrence, Virginia NRCS; and Susan Andrews, Charles Kome and Holli Kuykendall, Soil Quality National Technology Development Team.

For further information, contact the Soil Quality National Technology Development Team at (336) 370-3331.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, SW., Washington, DC 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Soil Quality:

Managing Cool, Wet Soils

Dealing with cool, wet soils

When dealing with cool and wet soils, the first thing is to realize that you are dealing with a complex system. Any alteration of the individual properties (physical, chemical, biological) impacts the whole soil ecosystem thus diminishing function. This technical note will show that by mimicking nature, through limiting physical disturbance and increasing crop diversity, you can better manage cool and wet soils.

Before beginning to optimally manage cool and wet soils, we must first understand a few basic principles that will lay the foundation for best management practices.

One major soil function is to permit water infiltration. Soil infiltration is often reduced by tillage. Tillage destroys soil aggregates. Destruction of soil aggregates negatively impacts soil porosity (air spaces in the soil). When soil porosity is reduced, infiltration (the rate water travels in the soil) is reduced and the subsequent runoff is increased furthermore, free exchanges of gases between soil and atmosphere are hampered. Roots need a porous medium for gas exchange and water availability to achieve optimum growth. Water infiltration into cool, wet soils is important for several reasons, particularly its effects on the freeze/thaw cycle and soil structure.

This technical note will illustrate that cool, wet soils can be managed without disrupting the soil ecosystem.

Understanding the natural system


Understanding the natural system is crucial if we want to understand the problems associated with cold, wet soils. Snow cover is one of the most important climatic features of cold, wet soils because of its influence on energy and moisture budgets. Snow is highly reflective. Without snow cover the ground absorbs about four to six times more of the sun's energy.

In areas where snow cover is disappearing earlier in the spring due to climate change, large amounts of energy that would have melted the snow can now directly warm the soil. Snow cover accounts for the large

differences between summer and winter land surface albedo. The Albedo of an object is the extent to which it diffusely reflects light from the sun. Most land areas have an albedo of 10 to 40 percent. The average albedo on earth is about 30 percent. Dark wet soils have an albedo roughly of 6 to 15% and dry soils have an albedo of 22 to 34% (fig. 1) (Grobe 2006). Crops have an albedo of 16 to 25%. Dark soils have low albedo which means they absorb more sunlight (electromagnetic radiation) and reflect very little, thus heating up quicker than lighter colored soils. A daily life example of the albedo effect is: people that wear dark cloths in the summer put themselves at greater risk of heatstroke than those who wear light colored clothes.

Researchers and farmers both find that it takes a longer time for wet soils to warm up. The reason wet soils take longer to warm up is the high specific gravity of water, which means it takes a lot of energy to heat the soil water. Even though color and texture impact how quickly soils warm up, it is the soil water in cold climates that plays the largest role in the soil warming process in the early spring. This is why it is important to have good infiltration in your soils.

Figure 1 Percentage of diffusely reflected sun light in relation to various surface conditions of the earth (Grobe 2006)



If you have a light colored soil with a high albedo that is saturated with water, the light will be reflected, the soil will not absorb the heat from the light and the high water content will delay the soil from warming. In other words, dark soils heat up faster than light ones. Dark color is a reasonable indicator of soil carbon. Therefore, in theory, increasing soil carbon will darken the soil's color and decrease its albedo, potentially leading to warmer spring soil temperatures. But in reality that does not always occur. Soils with high organic matter content hold more water. More water means you need more energy to warm the soil profile. So the best way to warm your soils is to increase infiltration and to increase infiltration you must decrease tillage.

Snow cover impacts soils

The effects of snow and frost on soil water dynamics are extremely complex. Partitioning of snowmelt into soil surface infiltration and runoff is influenced by soil drainage characteristics, soil frost conditions, infiltration, soil water storage capacity, and snowmelt rates. Zhao and Gray (1999) report that numerous studies found seasonal infiltration is inversely related to the total moisture content of a soil at the time of melt.

Research has shown that soil freezing can alter the dynamics of water flow and nutrient cycling during snow melt (Brooks and Williams 1999). Seasonal snow packs in the northern latitudes are well known to reduce soil freezing by insulating the soil surface; frost depth generally varies inversely with snow depth. Lack of snow or a late snow pack accumulation results in soil freezes that are deeper and of longer duration than when the snow pack is established in early winter (Shanley and Chalmers 1999).

A study of drainage in agricultural land indicated that high water content soils at the onset of freezing reduced infiltration of snow melt. Freezing influences the potential for interaction between snow pack nutrients and soil microbes, which are critical regulators of nutrient cycling and retention during these periods (Schimel and Kieland 1996).

Concrete frost, which is partially saturated soil that freezes so pores are filled with ice, has very low permeability that greatly reduces infiltration of snow melt or rain and promotes overland flow. Soils in no-till systems have less concrete frost because no-till soils have larger soil pores compared with the same soil under conventional tillage. Soils in conventional tilled systems have a smaller soil pore which creates more concrete frost.

When soil aggregates are destroyed, pore spaces between soil aggregates are decreased. With decreased pore space, it's easier for ice to bridge the gap between the remaining aggregates, which creates an impermeable surface like concrete, thus the term "concrete frost." Concrete frost inhibits infiltration, water flow, and nutrient cycling by soil microbes (fig. 2).

Granular frost, occurs when unsaturated soils freeze (fig. 3). This results in a more permeable soil, allowing more infiltration and less overland flow than concrete frost.

Figure 2 Conceptual diagram of concrete frost





Figure 3 Conceptual diagram of granular frost



Increasing Granular frost

During the winter months soils function best in a **granular frost condition**. This occurs when permeable soils freeze under unsaturated conditions. Although the soil is frozen, infiltration still occurs and overland flow is minimized. Good soil structure with intact soil aggregates will increase infiltration rates, thus decreasing the water content in the soil. Granular frost in soils reduces the opportunity for ice to bridge the pore spaces between the aggregates, maintaining permeability.

How can the soil system be managed to increase soil function and productivity?

The better the soil is functioning (e.g. infiltrating), the fewer problems you will have with concrete frost and other associated water issues (Johnson and Lundin 1991). Functioning soils capture needed rain throughout the growing season. During the winter months soils will infiltrate and drain water throughout the soil profile as quickly as possible-which reduces water accumulation on the surface. Figure 4 shows what happens when soil pores are reduced in size.

Soil quality principles that increase soil function in cool and wet soils:

1) Reduce physical and chemical soil disturbance (tillage and pesticides):

Managing a cool and wet soil with tillage is like fixing a Swiss watch with a sledge hammer. Using the wrong tool can create a worse problem. Physical disturbance

like tillage depletes organic matter, increases soil crusting which reduces infiltration, increases concrete frost by reducing average pore space, and disturbs microbial function. All these factors impact water infiltration and nutrient availability throughout the growing season. Chemical disturbances, like pesticides, negatively impact soil microarthropods and earthworms, which are critical players for increasing infiltration and nutrient cycling. Pesticides and tillage also directly harm soil fungi and bacteria, which make soil “glues”, like glomalin and other polysaccharides, that help form soil aggregates. Tillage in cool, wet soils can create a string of events that can cause more harm than good.

Most wet areas in a field are wet because they receive additional water from higher places on the landscape by surface or subsurface flow and do not have an outlet for the excess water. Runoff is increased because tillage destroys the surface porosity needed for water to soak into the soil. One pathway water takes to infiltrate the soil is between soil aggregates. Tillage shears soil aggregates, leaving fewer pathways between aggregates and clogging the spaces that remain with the smaller pieces of smashed aggregates (fig. 5) (Laws and Evans 1949). Another way tillage reduces water infiltration into the soil pores; the pores created by roots and earthworms are destroyed. Pores that are not continuous with the soil surface do not conduct water down into the soil very well.

The net result is more water in low areas with no place to go but up (evaporation). When the weather and soil are cool in the spring, evaporation is a slow process. The best way to warm up the soil in these conditions is to use a pin wheel which is placed in front of a no-till drill. The pin wheel will push away the residue from the seedling thus allowing the soil warm next to the seedling without disturbing the soil. In sod or pasture applications, ripple coulters throw less soil and work well in heavy soils.

2) Use cover crops:

Cover crops are a good way of solving water problems in the winter. Cold tolerant cover crops such as winter wheat and rye, which survive through the winter, have the ability to utilize excess moisture and increase soil strength to ensure an earlier planting date. Ongoing research by Osborne and others (2002) in Brookings, South Dakota indicates that soil bearing strength and trafficability is increased with cover crops. The study revealed that conventional tilled fields and fields that grow no cover crops have a lower bearing strength versus no-till fields with cover crops (fig. 6). Soils with cover crops have more root and worm channels. The

Figure 4 Reduced pore size caused by tillage



roots in cover crops act like stabilizers in the soil, similar to rebar in concrete. Improved bearing strength may be related to the above biomass characteristics and the root system. In general terms the experiment illustrated the ability of cover crops to utilize excess soil moisture and increase soil strength compared to conventional tillage or no cover crop (Osborne et al. 2002).

Some other benefits of cover crops are the following:

- Cover crops increase nutrient cycling
- Cover crops enhance soil habitat for all types of soil microbes
- Cover crops improve water cycling in the soil profile
- Cover crops increase soil biodiversity

3) Reduce compaction:

Nature has built-in processes that will reduce soil compaction; cycles of wetting and drying, freezing and thawing. In the last 30 to 40 years, farming practices have changed drastically, creating situations where natural rejuvenation of the soil environment by wet-dry and freeze-thaw cycles is inadequate to maintain optimum conditions for crops.


Performing field operations on wet soils, using multiple field operations for crop production, eliminating perennial crops from crop rotations, and using heavy equipment contribute to more extensive and deeper compaction.

Management strategies for eliminating compaction:

- **Stay off wet soil.** Soil is most susceptible to compaction when soil water in the three- to six-inch soil depth is near field capacity or wetter. Under such moisture conditions, the potential for compaction increases as soil clay content increases and soil organic matter decreases.
- **Reduce tillage.** Tilled soils are more susceptible to compaction than no-till soils. Tillage contributes to the breakdown of soil structure by compressing and breaking soil aggregates, which are necessary for good air and water movement and good root growth. Tillage also results in the loss of soil organic matter which is important to soil aggregate stability. Using a pin wheel in front of the no-till drill will help remove the residue away from seedling allowing the soil to warm up quicker.

Figure 5 Effect of 50 years of continuous cropping on soil porosity (Adapted from Laws and Evans 1949)

Virgin Prairie (5.6% SOM)



Cropped 50 Years (2.9% SOM)





Figure 6 Bearing strength of soil (Osborne et al. 2002)



Build organic matter in soil. Organic matter promotes the development of good soil structure and decreases soil bulk density. It helps bind soil particles together as aggregates so they are not as easily cracked, split, or compressed by tillage or wheel traffic. Reducing disturbance and increasing diversity will increase organic matter.

- **Increase diversity of rotations with perennial crops or deep rooted crops.** When crop rotations include alfalfa, clover, or grass, soils usually are less compact than soils in fields without these rotations. The deeper rooting depth of alfalfa and clovers helps keep the soil more porous and overall produce more organic matter.

Conclusion

Nature is very complex. Trying to alter the way the soil ecosystem functions is usually expensive, ineffective, and detrimental in the long run. As we mimic the natural soil ecosystem by using cover crops and minimizing chemical/physical disturbance, the effectiveness of our efforts to manage cool and wet soils is accelerated.

References

- Brooks PD and Williams MW. 1999. Snowpack controls on nitrogen cycling and export in seasonally snow-covered catchments. *Hydrological Process* 13:2177-2190.
- Fahey TJ and Lang GE. 1975. Concrete frost along an elevational in New Hampshire. *Canadian Forest Research* 5:700-705.
- Grobe H. 2006. Percentage of diffusely reflected sunlight to various surface conditions of the earth. Creative Commons. Category: Climatology, Radiation and Albedo. Online: http://en.wikipedia.org/wiki/Image:Albedo-e_hg.svg, verified August 16, 2008.
- Hardy P, Groffman PM, Fitzhugh RD, Henry KS, Welman AT, Demers JD, Fahey TJ, Dricscoll CT, Tierney GL, and Nolan S. 2001. Snow depth manipulation and its influence on soil frost and water dynamics in a northern hardwood forest. *Biogeochemistry* 56:151-174
- Johnson H and Lundin LC. 1991. Surface runoff and soil water percolation as affected by snow and soil frost. *Journal of Hydrology* 122:141-159
- Laws WD and Evans DD. 1949. The effects of long-time cultivation on some physical and chemical properties of two rendzina soils. *Soil Science Society Periodical* 13:15-19.
- Osborne SL, Riedell WE, Schumacher TE, and Hamburg DS. 2002. Use of cover crops to increase corn emergence and field trafficability. Progress Report #SOIL PR 02-39. Ag. Exp. Stn., Plt. Sci. SDSU. Brookings SD.
- Schimel JP, Kieland K, and Chapin FS. 1996 Nutrient availability and uptake by tundra plants. In: JD Tenhunen ed., *Landscape Function: Implications for Ecosystem Response to Disturbance: A case study of Arctic Tundra*. Springer-Verlag, New York. pp.201-221.
- Shanley JB and Chalmers A. 1999. The effect of frozen soil on snow melt runoff at Sleepers River, Vermont. *Hydrological Process* 13:1843-1857
- Zhao L and Gray DM. 1999. Estimating snowmelt infiltration into frozen soils. *Hydrological Process* 13: 1827-1842.