

Innermost row of bracts enlarged with a rigid, incurved hooked, point



Pappus lacking



Flowering heads rayless



Scientific Name: *Stylocline filaginea* A. Gray

Bibliographic Reference: Gray, 1873. Proc. Amer. Acad. Arts 8: 652.

Common Name: Stylocline

Family (Common Name): Asteraceae or Compositae (Sunflower Family)

Synonyms: *Ancistrocarphus filagineus* A. Gray

Idaho Native Plant Society Category: State Monitor

Natural Heritage Program Rank: S2

Distribution: Blaine, Camas, Elmore, Gooding, Owyhee, and Washington counties, Idaho; Baja California; Jefferson and Baker counties, Oregon; Lander and w Elko counties, Nevada.

Habitat: Open, dry or vernal moist habitats in the valleys and foothills on shallow stony basalt with cindery graveled surface. Commonly associated with alkali sage, Owyhee sage or stiff sage, from 600-1900 m elevation.


Phenology: (April) May through June.

Look-alikes: Possibly confused with other species of the genus *Stylocline* but distinguished from them by the enlarged and conspicuous innermost row of receptacular bracts, each with a rigid, incurved hooked point.



Closeup of *Stylocline filaginea*

Ann Debolt



Species
Distribution