SHRUB PERFORMANCE ON ORGANIC SOILS

Sheet

them died.

612-5



United States
Department of
Agriculture

Conservation

Soil



(Plant Materials - Forestry)

INTRODUCTION: The USDA Soil Conservation Service Rose Lake Plant Materials Center established a combination field evaluation/demonstration windbreak in 1987 on Michigan State University's Muck Research Farm. The soil type was a Houghton Muck--level, deep, organic soil with moderate permeability and neutral pH. Classification was a Typic Medisaprists. The purpose of this planting was to evaluate plant performance of various shrub species commonly used as windbreaks on organic soils. Evaluations were carried out for four years (1987-1991).

PROCEDURE: Eleven species were selected for this study:
(1) 'Roselow' Sargent crabapple, Malus sargentii Rehder; (2) Eastern ninebark, Physocarpus opulifolius (L.) Maxium; (3) spirea, Spirea x vanhouttei (Broit) Zabe; (4) arrowwood, Viburnum dentatum L;
(5) 'Indigo' silky dogwood, Cornus amomum Miller; (6) highbush cranberry, Viburnum trilobum Marshall; (7) 'Cling-Red' Amur honeysuckle, Lonicera maackii Maxim; (8) nannyberry, Viburnum lentago L.; (9) tall hedge Rhamnus, Frangula var. columnaris L.; (10) Cheyenne privet, Ligustrum vulgare L.; and (11) 'Magenta, hybrid crabapple, Malus sp.

Twenty bare-root plants of each species were planted in a single row with 180 cm (6 ft.) spacing between plants. Weed control, both chemical and mechanical, was provided for the first four years. Annual application of fertilizer was applied according to soil tests. Supplemental application of manganese was important for good growth of shrubs on organic soils.

DISCUSSION: Some observations made on the planting are: Survival and growth rate for the planting was very good. This can be attributed to good weed control during the establishment period.

It was observed that those species with a fiberous root system, such as silky dogwood, were better able to resist frost heave. Those species with a non-fiberous root system, such as the crabapples, had portions of their roots exposed.

Discussion of the individual species, in order of their effectiveness, follows: (1) 'Indigo, gilky dogwood developed into a solid, uniform hedge. Survival was excellent. The plants were healthy and much tillering was evident; (2) Chevenne privet had excellent survi-The plants were large and healthy; (3) Arrowwood had excellent val. The plants were slightly smaller than the top performers; survival. (4) <u>'Cling Red, Amur-honeysuckle</u> grew fast but suffered die-back in the upper portion of its branches each year; (5) <u>Eastern</u> demonstrated good growth but survival was not good; (6) Tall hedge had excellent survival and good growth but was too compact to form a barrier at the selected spacing; (7) <u>Nannyberry</u> had excellent survival but initial growth was irregular. Growth that occurred later in the study was more uniform; (8) was slow to establish and growth rates were irregular; (9) American cranberrybush survival and plant vigor was low; (10) 'Roselow are abapple growth was irregular. Individual plant health was not uniform; (11) Spire: was A drought in 1988 stressed the plants and the majority of a failure.

DATA SUMMARY

. .

1991

SPECIES	\$ SUR.	AVE. HT. (Cm.)		UNI- FORMITY	EFFEC- TIVENESS	REL. RANK
'Indigo' silky Dogwood	100	±.250	320	2.6.2	2	1
'Cheyenne' Privet	100	273	273		1 74 3 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	** 2
Arrowwood	100	210	210	3	3	3
'Cling-Red' Amur Honeysuckle		263	297	5 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	en ja roj b estoj Kojska jost sal Prijaj koje k	44. 45. 38.54
Eastern, Ninebark	85	197	225	* 6	5	5
Tall Hedge	100	256⊗	120	5.00	ir karangan salah sa Arangan salah s	6
Nannyberry	100	232	209	5	1 , 42 5 7 7 1 7 1	7
'Magenta' Crabapple	88	212	141	7	7	8
American Cranberrybush	90	177	175	5	· · · · · · · · · · · · · · · · · · ·	9
'Roselow' Sargent Crabapple	95	144	115	6	8	10
Spirea	20	88	106	**** 8 ***	9.4.4.4	11

_1/

2/

4.3 Cont. (1.2 East)

Uniformity of size and shape. 1 = Identical - 10 = Extremely variable

Occular estimate of effectiveness as a windbreak based on size, density and uniformity. 1 = Excellent - 10 = Totally ineffective.