

# Rose Lake Plant Materials Center



## 2007 Technical Report

---

East Lansing, Michigan

## TABLE OF CONTENTS

	Page
Personnel.....	5
Introduction.....	5
Studies.....	6
<b>Partnership and Program Development</b> .....	8
Memorandum of Understanding with Indiana Department of Natural Resources.....	9
National Ash Tree ( <i>Fraxinus spp.</i> ) Seed Collection Initiative.....	11
Memorandum of Understanding for Ft. Custer Vegetative Restoration.....	12
Arrest Erosion Threatening Raspberry and Outer Island Light Stations.....	14
<b>Plant Material Collections</b> .....	15
Evaluation and Great Lakes Release of American Plum ( <i>Prunus americana</i> Marsh.).....	15
Evaluation and Release of Broomsedge Bluestem ( <i>Andropogon virginicus</i> L.).....	15
Development of a Great Lakes Release of Buttonbush ( <i>Cephalanthus occidentalis</i> L.).....	15
<b>Stabilization of Streambanks and Shorelines</b> .....	16
Common Elderberry ( <i>Sambucus canadensis</i> ) for Streambank Stabilization .....	17
Shrub Rooting Study.....	17
<b>Stabilization and Restoration of Dune Areas</b> .....	21
Testing of Prairie Sandreed ( <i>Calamovilfa longifolia</i> ) Potential Release.....	21
Evaluation of Dune Willow ( <i>Salix cordata</i> ).....	25
<b>Stabilize Cropped Areas of Concentrated Water Flow</b> .....	26
Vegetative Hedges for Controlling Erosion in Areas of Concentrated Flow .....	26
Intercenter Component of Evaluation and Release of Miscanthus Grass ( <i>Miscanthus sinensis</i> ).....	28
<b>Cover Crops</b> .....	29
Evaluation of Hairy Vetch Populations for Winter Hardiness.....	29
<b>Bioenergy</b> .....	30
Tall Wheatgrass Biofeedstock for Biofuel.....	30
<b>Plants for Wetland Restoration</b> .....	31
Evaluation and Release of Canada Bluejoint ( <i>Calamagrostis canadensis</i> (Michx.) Beauv.).....	31
<b>Restoration or Reclamation of Disturbed Areas</b> .....	34
Great Lakes Composite of Virginia Wildrye ( <i>Elymus virginicus</i> ).....	34
Increase and Release of Riverbank Wildrye ( <i>Elymus riparius</i> ).....	36
Great Lakes Composite of Bottlebrush Grass ( <i>Elymus hystrix</i> ).....	36

Evaluation and Release of Poverty Oatgrass ( <i>Danthonia spicata</i> (L.) Beauv. ex Roemer & J.A. Schultes).....	38
<b>Improved Warm-season Forage Grass</b> .....	41
Eastern Gamagrass ( <i>Tripsacum dactyloides</i> ) for Forage Production.....	42
Relative Maturity of Eastern Gamagrass ( <i>Tripsacum dactyloides</i> ) Accessions.....	43
Big Bluestem ( <i>Andropogon gerardii</i> ) for Forage Use.....	45
<b>Forest Improvement and Windbreak Technology</b> .....	47
Windbreaks in Muck Soils.....	47
Direct Seeding Tree and Shrub Establishment.....	48
Direct Seeding Northern Red Oak ( <i>Quercus rubra</i> ).....	50
<b>Native Plant Species to Enhance Wildlife Habitat</b> .....	53
Tick-trefoil ( <i>Desmodium spp.</i> ) for Wildlife Food Plots.....	53
<b>Appendix A: Released Material</b> .....	55
<b>Appendix B: Customer Assistance Summary</b> .....	60
<b>Appendix C: Weather and Climate</b> .....	61

## ROSE LAKE PLANT MATERIALS CENTER

7472 Stoll Road

E. LANSING, MI 48823-9420

Phone: 517-641-6300

Fax: 517-641-4421

Website: [www.ashseed.org](http://www.ashseed.org)

### PERSONNEL

John Leif..... Manager  
John Durling.....Agronomist  
Sergio Pérez.....Biological Sciences Technician  
Elaine Gerona.....Office Automation Assistant  
Gail Bischoff.....Biological Aide  
David Bricker.....Biological Aide

### Plant Materials Specialist

David Burgdorf

Phone: 517-641-7831

Fax: 517-641-4421

Cell: 517-281-5939

### State Conservationist Advisory Committee

Garry Lee, Chair.....Michigan  
Jane Hardisty.....Indiana  
Terry Cosby.....Ohio  
Patricia Leavenworth.....Wisconsin  
Bill Gradle.....Illinois  
William Hunt.....Minnesota  
Ron Alvarado.....New York  
Craig Derickson.....Pennsylvania

### Plant Materials Committee Members

#### Indiana

Ken Collins, Ch.  
Kelley Barkell  
Dave Burgdorf  
Barry Fisher  
Adam Heichelbech  
Susannah Hinds  
Jo Hodgett  
Brian Ingmire  
Ellen Jacquart  
Doris Scully  
Dave Stratman  
Jenny Vogel  
Shannon Zezula

#### Wisconsin

Greg Rebman, Ch.  
Tom Cogger  
Randy Gilbertson  
Pattie Haack  
Dexter Porter  
Victor Price  
John Pingry  
Robert Weihrouch  
Michele Wilkinson  
Pat Murphy  
Merrie Schamberger  
Robert Weihrouch

#### Michigan

Tom Ward, Ch.  
Vicki Anderson, Adv.  
Dave Burgdorf, Adv.  
Karen Blair  
Jerry Davis  
Demarys Mortenson  
James Scott  
Thomas Tauzer  
Jennifer Taylor  
Sally VanLieu  
Jim Williams

#### Ohio

Shawn Anderson, Ch.  
Clifford Thornton, Adv.  
Mary Ann Core  
Doug Deardorff  
Mark DeBrock  
Rick Ellsmore  
Erika Harold  
Mark Scarpitti  
Mark Smith  
Tammy Willis

## INTRODUCTION

**History** The Rose Lake Plant Materials Center (PMC), as part of the USDA Natural Resources Conservation Service (NRCS) Plant Materials Program, was established in 1958 on a 40-acre site at the Michigan Department of Natural Resources, Rose Lake Wildlife Research Station (42.8° N. Lat., 84.4° W. Long, Elevation 875 ft), and located eight miles northeast of Lansing, Michigan. Soil association is Boyer-Marlette-Houghton with well drained and moderately well drained, gently sloping to steep loamy sands to loams on moraines and very poorly drained muck in depressions.

**Responsibilities** The NRCS Plant Materials Program develops, tests, and transfers effective state-of-the-art plant science technology to meet customer and resource needs. The Plant Materials Program:

- Focuses on using plants as a natural way to solve conservation issues and re-establish ecosystem function.
- Collects, selects, and releases grasses, legumes, wildflowers, trees, and shrubs.
- Cooperates with public, private, commercial, and tribal partners and land managers to apply new conservation methods using plants.
- Provides plant materials and new applied technologies for national initiatives, like the Farm Bill.
- Offers plant solutions to battle invasive species, heal lands damaged by natural disasters, reduce the effects of drought, promote air and water quality, and produce alternative energy.

It is the responsibility of the Rose Lake PMC to:

- assemble, test, and release plant materials for conservation use;
- determine techniques for the successful use and management of conservation plant species;
- facilitate the commercial increase of conservation plant species; and
- provide for the development and transfer of state-of-the-art applied science technology.

**Long Range Plan** PMC operations are guided by a Long Range Plan which is a compilation of Plant Materials LRPs from Indiana, Ohio, Michigan, and Wisconsin. The PMC LRP is consistent with goals and objectives identified in the NRCS Strategic Plan.

## STUDIES

<u>Study Number</u>	<u>Study Name</u>	<u>Purpose</u>	<u>Year Initiated</u>	<u>Report Page</u>
26A097F	Testing of Prairie Sandreed ( <i>Calamovilfa longifolia</i> ) for Potential Release	Release	1989	21
26C126V	Vegetative Hedges for Controlling Erosion in Areas of Concentrated Flow	Technology	1996	26
26I080J	Tick-trefoil ( <i>Desmodium</i> spp.) for Wildlife Food Plots	Release	1987	53
26I101G	Big Bluestem ( <i>Andropogon gerardii</i> ) for Forage Use	Release	1992	45
26I124L	Common Elderberry ( <i>Sambucus canadensis</i> ) for Streambank Stabilization	Release	1997	17
MIPMC-P-0201-CR	Development of Great Lakes Composite of Virginia Wildrye ( <i>Elymus virginicus</i> )	Release	2002	34
MIPMC-P-0204-CR	Development of Great Lakes Composite of Bottlebrush Grass ( <i>Elymus hystrix</i> )	Release	2002	36
MIPMC-P-0207-CR	Increase and Release of Riverbank Wildrye ( <i>Elymus riparius</i> )	Release	2002	36
MIPMC-P-0208-CR	Evaluation and Release of Miscanthus Grass ( <i>Miscanthus sinensis</i> )	Release	2002	28
MIPMC-P-0209-CR	Evaluation of Dune Willow ( <i>Salix cordata</i> )	Release	2002	25
MIPMC-P-0503-WL	Evaluation and Great Lakes Release of American Plum ( <i>Prunus americana</i> Marsh.)	Release	2005	15
MIPMC-P-0504-WE	Development of a Great Lakes Release of Buttonbush ( <i>Cephalanthus occidentalis</i> L.)	Release	2005	15
MIPMC-P-0601-CR	Evaluation and Release of Broomsedge Bluestem ( <i>Andropogon virginicus</i> L.)	Release	2006	15
MIPMC-P-0602-CR	Evaluation and Release of Poverty Oatgrass ( <i>Danthonia spicata</i> (L.) Beauv. ex Roemer & J.A. Schultes)	Release	2006	38
MIPMC-P-0603-WE	Evaluation and Release of Canada Bluejoint ( <i>Calamagrostis canadensis</i> (Michx.) Beauv.)	Release	2006	31
MIPMC-T-0004-CR	Arrest Erosion Threatening Raspberry and Outer Island Light Stations	Technology	2000	14
MIPMC-T-0301-WO	Direct Seeding Tree and Shrub Establishment	Technology	2003	48
MIPMC-T-0302-PA	Eastern Gamagrass ( <i>Tripsacum dactyloides</i> ) Cultivar Evaluation	Technology	2003	43

MIPMC-T-0303-PA	Eastern Gamagrass ( <i>Tripsacum dactyloides</i> ) Production Evaluation	Technology	2003	42
MIPMC-T-0402-WO	Direct Seeding of Northern Red Oak ( <i>Quercus ruba</i> )	Technology	2004	50
MIPMC-T-0404-BU	Windbreaks in Muck Soils	Technology	2004	47
MIPMC-T-0604-CP	Evaluation of Hairy Vetch ( <i>Vicia villosa</i> ) Populations for Winter Hardiness	Technology	2004	29
MIPMC-T-0701-TE	National Ash Tree Seed Collection Initiative	Technology	2007	11
MIPMC-T-0702-RI	Shrub Rooting Study	Technology	2007	17
MIPMC-T-0709-CP	Tall Wheatgrass Biofeedstock Study for Potential Use in Biofuel (Liquid, Thermal, and Thermochemical) Applications in NE, Mid-Atlantic, and Upper Midwestern States	Technology	2007	30

## PARTNERSHIP AND PROGRAM DEVELOPMENT

The Rose Lake PMC partners with other federal, state, and local agencies; tribes; and private entities to achieve its missions and goals. The PMC is always looking to create new partnerships while strengthening its many existing partnerships. Partnering has two advantages: First, partnering allows expedient and effective land conservation by garnering the skills, knowledge, and abilities of each agency or group involved. Second, it broadens the PMC's range of plant and conservation skills thereby widening the customer base and demand. Often partnerships are established as *reimbursables* which help fund staff and equipment needs.

**Problems** Many partners lack either the skills or facilities to produce plants or complete research on various, adopted conservation projects.

**Needs** Plants, planting skills, and/or facilities to grow plants for conservation projects on public lands are needed.

### **Memorandum of Understanding with Indiana Department of Natural Resources**

**Background** The Indiana Department of Natural Resources (INDNR) Division of Fish and Wildlife and NRCS have mutual interests and goals concerning wildlife habitat management and conservation, specifically related to native plants from Indiana. The NRCS is the lead Federal agency for providing conservation assistance on private lands. The NRCS Plant Material Center's mission is to develop, test, and transfer effective state-of-the-art plant science technology, providing timely and effective vegetative solutions for identified customer and resource needs. The INDNR's mission is to professionally manage Indiana's fish and wildlife for present and future generations, balancing ecological, recreational, and economic benefits.

The Natural Resources Conservation Service and INDNR are cooperating to meet the demands of both public and private lands for establishing high quality wildlife habitats with native Indiana seed. Both agencies recognize the concern of introduced plant species (native and non-native) and the potential for these species to alter the genetics of existing native plant communities. Moreover, both agencies have recognized the need for testing and selecting plants native to Indiana for use in implementing their individual and mutual programs.

**Memorandum of Understanding** Through a Memorandum of Understanding (MOU) the INDNR and Rose Lake Plant Materials Center have agreed to cooperatively promote and propagate native plants from Indiana genotype sources for the expansion and implementation of both agencies' missions and programs. Specifically, they will cooperatively:

- Identify a list of plants that each agency deems important and set priorities for which species to work on first.



- Harvest seed from established fields or from remnant stands.
- Cooperatively review their MOU to determine any necessary revisions.
- Cooperatively release plant materials through the NRCS Plant Materials Program and publicize to commercial growers for establishment and increase for availability to the public.
- Secure Indiana Native Seed Certification of all seed.
- Promote the use of the released native plant material.

**Releases** Seed was collected from native stands (as designated by the Indiana Department of Natural Resources, Wildlife Division) of big bluestem (*Andropogon gerardii*), little bluestem (*Schizachyrium scoparium*), and Indiangrass (*Sorghastrum nutans*) within the State of Indiana in the early to mid 1990s. Seeds from these populations were planted into single crossing blocks for each species. Resultant seeds were planted in nine field plantings in central Indiana. Field plantings were evaluated for adaptability, potential for invasiveness, and usefulness in conservation planting.

Based on (1) the excellent field performance of these big bluestem, Indiangrass, and little bluestem composites over several years and locations; (2) the limited availability of plant material; and (3) the need for these products in ecosystem restoration and enhancement, seed stock of Prairie View Germplasm of these composites was released in 2005.

Prairie View Indiana Germplasm big bluestem is a perennial, warm-season bunch grass with short, scaly rhizomes. Big bluestem attracts insects and provides seeds that are used as food sources by songbirds, game birds, and small mammals. Many wildlife species also utilize big bluestem for nesting, escape, and winter cover. Other anticipated uses of Prairie View big bluestem include increasing species diversity, controlling erosion, and restoring native plant environments.

Prairie View Indiana Germplasm little bluestem is another perennial, warm-season bunch grass. Its anticipated uses also include providing food/cover for wildlife, increasing species diversity, controlling erosion, and restoring native plant environments. At 2 – 4 ft in height at maturity, it is the shortest of the Prairie View grasses.

Prairie View Indiana Germplasm Indiangrass, like the other Prairie Views, is a perennial, warm-season bunch grass. A distinguishing characteristic is a narrow, plume-like seed head that turns golden at maturity. Its anticipated uses also include providing food/cover for wildlife, increasing species diversity, controlling erosion, and restoring native plant environments.

Quantities of pure live seed produced in 2007 under this Memo of Understanding at the Jasper-Pulaski Fish & Wildlife Area and processed by the Rose Lake PMC were approximately 3 lbs bulk of big bluestem and 270 lbs bulk (212 PLS) of Indiangrass. This seed will be distributed to growers and demonstration plantings. Seed requests may be submitted to:

David Burgdorf, Plant Materials Specialist  
Rose Lake plant Materials Center  
7472 Stoll Road  
E. Lansing, MI 48823-9420  
517-641-7831  
517-281-5939 cell  
517-641-4421 FAX

[dave.burgdorf@mi.usda.gov](mailto:dave.burgdorf@mi.usda.gov)

**Study No. MIPMC-T-0701-TE**  
**National Ash Tree Seed Collection Initiative**

**Background** Ash trees in the PMC Service Area are being attacked and destroyed by Emerald Ash Borer (*Agrilus planipennis* (Fairmaire)). This introduced pest, first detected in summer 2002 in Southeast Michigan, has already spread to Illinois, Indiana, Maryland, Ohio, Ontario, Pennsylvania, and Maryland.

With this impending disaster (already more than 20 million trees have been killed) Rose Lake PMC entered into a non-funded cooperative agreement with the USDA Agriculture Research Service (ARS). The agreement states that the PMC will initiate an ash tree seed collection and that the ARS National Center for Genetic Resource Preservation Center in Ft. Collins, Colorado will store samples of all the seed collected in their long term seed storage facility. Tribes are specifically mentioned in the agreement because of the PMC's previous assistance to tribes with black ash and other culturally significant plants.

Voluntary seed collections are being sought. The PMC receives, processes, labels, and sends seed to the Preservation Center. Tribal seed remains tribal property in storage with tribes retaining authority over sharing and distribution.

**Presentations** Plant Materials Specialist David Burgdorf and Plant Materials Center staff have made numerous presentations on the ash collection effort to government agencies including tribes, partners, and the general public. Posters and brochures are being distributed to NRCS Field Offices and other partners. The effort has also received popular press and television coverage.

**Collection** Ash seed samples have been collected in California, Illinois, Indiana, Maryland, Michigan, Minnesota, New York, Ohio, Pennsylvania, Utah, West Virginia, and Wisconsin on tribal and non-tribal lands by NRCS and District personnel and the general public. PMC staff check seeds for fill and obvious pests. Staff clean, label, and forward seed to the USDA Forest Service National Seed Laboratory in Dry Branch, Georgia for X-ray analysis (nondestructive viability test). Seed is then shipped to the ARS National Center for Genetic Resource Preservation Center for long-term storage. After culling, six samples, 117 samples, and 12 samples of 2005, 2006, and 2007 growing season seed, respectively, were sent by the PMC to the National Center for Genetic Resource Preservation.

The collection continues. Collection information is available on the web at:

[www.ashseed.org](http://www.ashseed.org)

**MSU/PMC Collaborative Research** Dr. Deborah G. McCullough, Professor of Entomology and Professor of Forestry at Michigan State University, and her graduate students including Andrew Roy Tluczek are conducting ash-borer related research at the PMC where a 200-300 ash tree plantation had been established in the 1980s. Research questions being pursued at the PMC relate to factors affecting development of Emerald Ash Borer larvae.

## Memorandum of Understanding for Ft. Custer Vegetative Restoration

**Background** Fort Custer Training Center (FCTC) in Southwest Michigan is 7500 acres of military tactical training area used by the Michigan National Guard and other branches of the armed forces. While FCTC is an important training facility, it is also home to a wide variety of natural resources, e.g., wildlife, forests, wetlands, surface water, and various rare plant and animal species. Moreover, historic and cultural resources are located on the property. The facility is federally-owned and operated by the Michigan Department of Military and Veterans Affairs (MDMVA).

**Memorandum of Understanding** A memorandum of understanding between the NRCS Rose Lake PMC and the MDMVA was developed for restoration of native vegetation and habitats at Fort Custer. PMC staff will provide consultation and on-the-ground assistance for collection, propagation, establishment, and maintenance of native vegetation. PMC assistance with research and selection of proper native vegetation that meets the needs of the firing ranges, ammo bunkers, and convoy reaction course areas will mutually benefit MDMVA, FCTC, and NRCS.

**Deliverables** The following are being realized from the cooperative agreement between MDMVA and NRCS Rose Lake PMC:

- Identification and selection of specific native plant species for collection, increase, testing, and evaluation for conservation uses at FCTC, including grasses to reduce mowing needs on ammunition bunkers (see below) and firing ranges and native grasses for roadsides.
- Established production plots/fields of selected native plants (e.g., little bluestem, big bluestem, and Indiangrass) to provide seed to commercial growers and for use at FCTC for prairie plantings on post. Approximately 17,000 Indiangrass and 17,000 big bluestem plants were grown in the PMC greenhouse and transplanted into FCTC production areas in 2007.
- Technical expertise on plan development for establishing native prairie plants on the convoy reaction course.
- Consultation and hands on implementation of weed control, planting, plot/field production, and management.
- Protocols for harvesting, cleaning, and handling of seed. Approximately 2 lbs PLS of big bluestem seed and 80 lbs PLS of little bluestem seed from FCTC were cleaned by the PMC in 2007.

**Ammunition Bunkers** Demonstration plots were established on an ammunition bunker so that vegetation species and establishment methods could be evaluated (Table 1). Treatment strips were 30 ft up-and-down the slope and 3 or 7 ft wide. Replicates were established to face northeast, east, and southeast, respectively. Evaluation will be conducted in collaboration with FCTC environmental staff. Criteria include ease of establishment, mowing and maintenance requirements, soil stabilization on the 30-40° slopes, fire hazard, etc.

Table 1. Vegetation and establishment methods for Ft. Custer ammunition bunker trials.					
Common Name		Scientific Name	Establishment Method	Approximate Seed or Plant Population	Observations
Bermudagrass		<i>Cynodon dactylon</i> (L.) Pers.	Plugged into coir biodegradable mattress (with potting soil); established off-site; and transported to bunker in fall 2005	6 plants/ft <sup>2</sup>	Greenup was slow in spring. Sod was green and well formed by end-of-season.
Periwinkle		<i>Vinca minor</i> L.	Plugged into coir biodegradable mattress (with potting soil); established off-site; and transported to bunker in fall 2005	Actual planting population not calculated due to stoloniferous nature of Periwinkle	Didn't provide adequate groundcover during 2006 growing season.
Sedum	Two row stonecrop	<i>Sedum spurium</i> Bieb. 'Tricolor'	Plugged into soil overlaid with 0.75-in X 0.75-in mesh coir fiber, biodegradable mats covering in summer 2005	3 plants/ft <sup>2</sup>	Established well but vegetative growth was slow. Didn't provide adequate ground cover in 2006 growing season.
	Stone- crop	<i>Sedum acre</i> L.	Same as above	Same as above	
	Orange stonecrop	<i>Sedum kamtschaticum</i> Fisch. & C.A. Mey.	Same as above	Same as above	
	Orange stonecrop	<i>Sedum spurium</i> Bieb. 'Fudlaglut'	Same as above	Same as above	
Red fescue		<i>Festuca rubra</i> L.	Direct seeded on-site at bunker into soil overlaid with 0.5-in X 0.4-in mesh coir fiber, biodegradable mats in fall 2005	4 lbs/1000 ft <sup>2</sup>	Good germination in fall 2005 and good sod formation during 2006.
			Established off-site by direct seeding into coir biodegradable mattress (with potting soil); overlaid with 0.5-in X 0.4-in mesh coir fiber, biodegradable mats and transported to bunker in fall 2005	4 lbs/1000 ft <sup>2</sup>	Good growth and sod formation in mats off site and good establishment on bunkers in 2006.
Buffalograss		<i>Buchloe dactyloides</i> (Nutt.) Engelm. 'Top Gun'	Direct seeded on-site in summer 2006	3 lbs/1000 ft <sup>2</sup>	Very poor germination.
Pink		<i>Dianthus</i> sp.	Transplanted into biodegradable mattress in summer 2006	≈1 plant/ft <sup>2</sup>	Good stand was observed in 2007.

**Study No. MIPMC-T-0004-CR**  
**Arrest Erosion Threatening Raspberry and Outer Island Light Stations**

**Introduction** The Apostle Islands National Lakeshore is comprised of Raspberry Island, Outer Island, and others in Lake Superior near Bayfield, Wisconsin. Several of these islands have historic lighthouses that once guided mariners through the rough waters of Lake Superior. Continuous erosion of steep slopes has jeopardized these historic facilities. This project was initiated in 2000 to produce native plant stock for stabilizing slopes, preventing erosion, preserving native plant resources, and revegetating at Apostle Islands National Lakeshore.

The National Park Service entered into several reimbursable agreements with NRCS and the Rose Lake PMC to provide technical assistance and to assist with collecting and growing plants. Under a Memorandum of Agreement grass, forb, and shrub species were selected for propagation based on the materials' availability, viability, and site adaptability for the intended use.

**Accomplishments** The NRCS Great Lakes Region Plant Materials Specialist has provided training to the Apostle Islands National Lakeshore employees on several slope stabilization techniques including guidance on the installation of a vegetative crib wall and slope grid system for the Outer Island. Park staff have constructed their first vegetative crib wall on Raspberry Island under the direction of the Plant Materials Specialist.

**Technology Development** The NRCS Great Lakes Region Plant Materials Specialist has provided training to the Apostle Islands National Lakeshore employees on several soil bioengineering slope stabilization techniques including guidance on the installation of vegetative crib walls and slope grids for Outer Island. Park staff constructed several vegetative crib walls on Outer and Raspberry Islands.

**PLANT MATERIAL COLLECTIONS**

**Study No. MIPMC-P-0503-WL  
Evaluation and Great Lakes Release of American Plum (*Prunus americana* Marsh.)**

**Study No. MIPMC-P-0601-CR  
Evaluation and Release of Broomsedge Bluestem (*Andropogon virginicus* L.)**

**Study No. MIPMC-P-0504-WE  
Development of a Great Lakes Release of Buttonbush (*Cephalanthus occidentalis* L.)**

The collection, assembly, selection, and release of new plant varieties and/or germplasm are integral to the Plant Materials Program. Every release starts with a collection. To this end the Rose Lake PMC is in the process of collecting American plum, broomsedge bluestem, and buttonbush plant materials from throughout the service area. Details are provided in Table 2 below.

Table 2. Plant materials collections on-going at Rose Lake PMC in 2007.			
Common name	Scientific name	Description	Collection Location (number of collections received)
American plum  NOTE: cover photo of this Technical Report shows native American plum blooming in Hillsdale County, MI	<i>Prunus americana</i> Marsh.	American plum is a native, small tree of the PMC service area. It is found in thickets, borders of woods, streambanks, floodplains, and fencerows. Flowers are white and 5-petaled. The one inch drupe is stone-seeded and red to yellow. The fruit is palatable to wildlife and humans. <i>Prunus americana</i> is closely related to <i>P. nigra</i> , Canadian Plum.	Hillsdale County, MI (1); Clinton County, MI (1); Ashland County, WI (4); Bayfield County, WI (1); Dane County, WI (2); Fon du Lac County, WI (1); Sheboygan County, WI (3)
Broomsedge bluestem	<i>Andropogon virginicus</i> L.	Broomsedge bluestem is native to the Rose Lake MI, Alderson WV, and Elsberry MO PMC service areas. It is a warm-season, herbaceous, perennial bunch grass that begins its growth when the average daytime temperature is between 60° to 65° F. It benefits wildlife, is ornamental, and has potential conservation application. It is found on low fertility soils and prevents erosion where similar warm season grasses can not be found. It is grazed readily by cattle in spring and early summer, but decreases rapidly in forage quality and palatability with advancing maturity.	
Buttonbush	<i>Cephalanthus occidentalis</i> L.	Common Buttonbush is a native, multi-stemmed wetland shrub. Mature plant is up to 12 ft tall. White flowers in spherical clusters form one-inch diameter button-like seed heads. Dark green, glossy, deciduous foliage is opposite or whorled.	Washington County, IN (1); LaCrosse County, WI (1)

## STABILIZATION OF STREAMBANKS AND SHORELINES

Soil bioengineering is the art and applied science that uses living plant material as a main structural component to control erosion, sedimentation, and flooding. It is a unique technology offering a responsible, attractive, and distinct approach to land stabilization and habitat restoration. Soil bioengineering systems are intended to form a positive interaction with the complex relationships that connect our natural resources.

**Problems** Reduced water quality from sedimentation and nutrient accumulation in our lakes and waterways is a concern in the Great Lakes and Midwest states. Poor urban, agricultural, and forestry land management practices resulting in excessive erosion and nutrient runoff have contributed to this situation. Nonpoint source pollution is responsible for sediment, nitrates, and phosphates deposition and increased biological oxygen demand in our lakes and streams.

**Needs** Although many other best management practices treat the cause of water quality degradation, soil bioengineering focuses on restoration. With emphasis placed on native species, developing acceptable plant materials and innovative soil bioengineering techniques has been pushed further to the front. Focus in the Great Lakes and Midwest States has been placed on native shrub species that:

- propagate vegetatively
- establish rapidly
- grow vigorously
- exhibit excellent erosion control qualities or features
- provide food, shelter, or nesting for wildlife
- are low maintenance and wear resistant
- possess aesthetic qualities



**Study No. 26I124L**  
**Common Elderberry (*Sambucus canadensis*) for Streambank Stabilization**

**Background** Elderberry is an upright, native shrub that can grow to 10-ft tall. The thick, yellowish-brown to light brown bark roughens and furrows with age. Stems and twigs are commonly covered with numerous, small, wart-like bumps (lenticels). Leaves are compound opposite with 5 to 11 coarse-toothed, elliptical leaflets. Common elderberry has white pith. Flowers are borne in white, flat-topped clusters. Fruit is purple to black. It is often found on wet or moist sites, such as along drainage ditches and wet fields.

**Description of Study** Assemble, select, and release native ecotypes of common elderberry for use in (a) soil bioengineering practices as locally-adapted plant material to stabilize streambanks, (b) native plant restoration projects, and (c) wildlife plantings as a food and shelter source. Additionally, this species has potential as an income source from berry production for human consumption. Exploring such uses in production agriculture would necessitate advanced testing to insure that quality and performance standards would be met.

**Procedure** Thirty-seven collections of common elderberry cuttings were processed and planted in the greenhouse in late 1997 through early 1998. Cuttings were transplanted into three initial study areas located in Wisconsin, Indiana, and the Center. Data was recorded on fruit abundance, plant height, canopy width, canopy density, and vegetative spread at the Center site from 1998 through 2000 but adverse weather forced the termination of studies in Wisconsin and Indiana. PMC-harvested material was sectioned and used to grow cuttings for new field trials in Indiana, Wisconsin, and Ohio, but these were terminated after 2003 due to poor long-term survival. Germination protocol development began in 2007 in the PMC greenhouse and is continuing. Technical information on germination will be made available to commercial growers when the release is made.

**Anticipated Release** Data collected from the Rose Lake Plant Materials Center trial indicates that Accession 9084126 has excellent growth characteristics for height, canopy density, spread, fruit abundance, and re-growth after pruning. Further evaluation is ongoing at the Rose Lake and National PMCs and in Wisconsin and is anticipated in several MLRAs and plant hardiness zones in the Rose Lake PMC service area. A new production block was established at the Rose Lake PMC in 2005 and a release is expected.

**Study No. MIPMC-T-0702-RI**  
**Shrub Rooting Study**

**Background and Objective** Stream bank restoration and bioengineering commonly use cuttings from various shrub and tree species. This study will demonstrate the growth and rooting ability of dormant woody plant materials. The objective of the study is to photographically document the growth and root formation of the tested species when grown under uniform conditions.

Dormant cuttings of many woody plant species have been evaluated for their effectiveness in soil bioengineering. A compiled list of those species and their rated effectiveness is available (Burgdorf, D.W. et al. 2007. Plant Species with Rooting Ability from Live Hardwood Materials for Use in Soil Bioengineering Techniques. Plant Materials Technical Note – No. 1. USDA-NRCS.) However, a technical document with good quality photographs of these materials showing root formation from dormant cuttings is not available.

**Materials and Methods** NRCS Regional Plant Materials Specialists coordinated the collection of dormant hardwood cuttings from the plant lists on Plant Materials Technical Note – No. 1 with their respective Plant Materials Centers. Cuttings were sent to Rose Lake PMC and grown under greenhouse conditions. Entries received for testing in 2006-07 are listed in Table 3.

Table 3. Shrub rooting study entries. Rose Lake PMC. Winter 2006-07.	
Scientific name	Common name
<i>Alnus incana</i>	Speckled Alder
<i>Amorpha fruitensa</i>	False Indigo
<i>Baccharis salicifolia</i>	Seep Willow
<i>Cephalanthus occidentalis</i>	Buttonbush
<i>Cephalanthus occidentalis</i>	'Keystone' Buttonbush
<i>Cornus amomum</i>	'Indigo' Silky Dogwood
<i>Cornus sericea</i>	Redosier Dogwood
<i>Cornus sericea ssp. occidentalis</i>	'Mason' Western Redosier Dogwood
<i>Cornus sericea ssp.sericea</i>	'Ruby' Redosier Dogwood
<i>Holodiscus dumosus</i>	Rock Spirea
<i>Lonicera invoucrata</i>	Bearberry Honeysuckle
<i>Morus sp.</i>	Mulberry
<i>Philadelphus microphyllus</i>	Mock Orange
<i>Philadelphus lewisii</i>	Colfax Germplasm Lewis Mockorange
<i>Physocarpus capitatus</i>	Pacific Ninebark
<i>Physocarpus monogynus</i>	Mountain Ninebark
<i>Physocarpus sp.</i>	Ninebark
<i>Populus balsamifera ssp. trichocarpa</i>	Black Cottonwood
<i>Populus Canadensis</i>	'Imperial' Carolina Poplar
<i>Populus deltoids</i>	Native Cottonwood
<i>Populus deltoides ssp. monilifera</i>	Plains Cottonwood
<i>Populus deltoides ssp.wislizeni</i>	Rio Grande Cottonwood
<i>Populus deltoides X nigra</i>	'Spike' Hybrid Poplar
<i>Rubus spectabilis</i>	Salmonberry
<i>Salix alba</i>	White Willow
<i>Salix amygdaloides</i>	Peachleaf Willow

<i>Salix cottetti</i>	'Bankers' Dwarf Willow
<i>Salix discolor</i>	Pussy Willow
<i>Salix drummondiana</i>	'Curlew' Drummond Willow
<i>Salix eriocephala</i>	Missouri River Willow
<i>Salix eriocephala ssp. ligulifolia</i>	'Placer' Erect Willow
<i>Salix exigua</i>	'Silvar' Coyote Willow
<i>Salix exigua</i>	Coyote Willow
<i>Salix hindsiana</i>	Sandbar Willow
<i>Salix hookeriana</i>	'Clatsop' Hooker Willow
<i>Salix humilis</i>	Prairie Willow
<i>Salix interior</i>	'Greenbank' Sandbar Willow
<i>Salix interior</i>	Sandbar Willow
<i>Salix lasiolepis</i>	'Rogue' Arroyo Willow
<i>Salix lasiolepis</i>	Arroyo Willow
<i>Salix lemmonii</i>	'Palouse' Lemmon's Willow
<i>Salix lucida</i>	Shining Willow
<i>Salix matsudana X alba</i>	Austree Willow
<i>Salix nigra</i>	Western Black Willow
<i>Salix nigra</i>	Black Willow
<i>Salix pentandra</i>	Laurel Willow
<i>Salix prolixa</i>	'Rivar' Mackenzie Willow
<i>Salix purpurea</i>	'Streamco' Purpleosier Willow
<i>Salix purpurea</i>	Purpleosier Willow
<i>Salix sericea</i>	Riverbend Germplasm Silky Willow
<i>Salix sericea</i>	Silky Willow
<i>Salix sessilifolia</i>	'Multnomah' NW Sandbar Willow
<i>Sambucus nigra ssp. canadensis</i>	Common Elderberry
<i>Sambucus nigra ssp. cerulea</i>	Blue Elderberry
<i>Sambucus nigra ssp. canadensis</i>	American Elderberry
<i>Sambucus racemosa</i>	Red Elderberry
<i>Spirea douglasii</i>	'Bashaw' Douglas Spirea
<i>Symphoricarpos albus</i>	Snowberry

Three cuttings of each entry were trimmed to 2-ft long and placed horizontally (fascines) into growing media. Light (sun supplemented with artificial), temperature, and humidity were maintained to optimize growing conditions. Soil was maintained at field capacity and optimum nutrient level with overhead fertigation.

Another three cuttings of each entry were trimmed to 1-ft long and placed vertically into potted growing media. Growing conditions were maintained as for fascines.

Days to new shoot emergence and days to 6-, 12-, and 18- inch shoot height were recorded. Photographs were taken on or about day 70.

Initial greenhouse study was conducted in winter and spring 2006-07. The study is being repeated in 2007-08 for entries showing unexpected results and for entries not included in the first year.

**Results** Vegetative and root growth photographs are taken at the end of the growing period. The cuttings are removed from the planting media, washed, and photographed to photographically document the growth and rooting characteristics. The photography protocol was developed with the assistance of the MI Public Affairs Specialist. Results will be published as a supplement to Plant Materials Technical Note – No. 1.

**Acknowledgement** The Rose Lake PMC gratefully acknowledges receipt of plant specimens from PMCs in Alaska, California, Colorado, Florida, Idaho, Kansas, Mississippi, New Mexico, New York, North Dakota, Oregon, Texas, Washington, and West Virginia and the State University of New York at Syracuse which were used in the greenhouse 2006-07 season. Additional plant specimens were provided in the 2007-08 greenhouse season.

## STABILIZATION AND RESTORATION OF DUNE AREAS

A diversity of landforms and plant and animal communities make up the Great Lakes shoreline, including those of the dune system. The greatest dunes of the Great Lakes occur along the east coast of Lake Michigan. This is principally due to the massive amounts of sand and sediment that eroded into Lake Michigan as glaciers retreated northward, and the prevailing westerly winds that gather energy traveling across this uninterrupted expanse of water. Wind velocity and direction, water levels, direction of water current, topography, and existing vegetation determine the rate of erosion and deposition.

**Problems** Development and other human activities along these dune systems alter protective dunes and wetlands, remove stabilizing vegetation, and generally reduce the shoreline's ability to combat strong winds and waves.

**Needs** The Great Lakes region needs commercially available plant varieties, the technology to establish them, and information on dune systems. Released plant material should:

- be native to the dune systems
- have good vigor and establish readily
- exhibit erosion control qualities

### **Testing of Prairie Sandreed (*Calamovilfa longifolia*) Potential Release (Accession 9086408) from Study No. 26A097F**

**Background** *Calamovilfa longifolia* is a tall, coarse, perennial, sand-binding grass with two distinct varieties: *magna* and *longifolia*. Literature indicates var. *longifolia* occurs in many dry prairies on the interior plains of Canada and the United States whereas var. *magna* characteristically occurs on active and semi-established dunes fringing Lake Huron and Lake Michigan. Studies have shown var. *magna* is a dominant dune builder on sites with slower sand deposition or windward blowout slopes, but has limited representation on rapidly depositing surfaces.

Prairie sandreed is a C4 species that reproduces vegetatively by rhizomes and sexually by seed. *Calamovilfa longifolia* expands into adjacent territory by producing short rhizomes on the periphery of a clump. The species follows a conservative growth form of slow radial spread. Such a growth form consolidates a local patch of resources and is resistant to invasion by other plant species. A 1985 study suggested prairie sandreed does not establish vegetatively in nature because the rhizomes are woody, do not bear any dormant buds, and do not fragment by wave action.

In response to sand accretion, the tillers and rhizomes adopt an erect habit of growth, emerge from sand, and produce tillers at the new sand surface. The growth form with a

high density of stout tillers within a localized area of clumps is exceptionally adept in trapping windblown sand.

Recurrent phenotypic selection breeding procedures were employed to develop a superior release of *Calamovilfa longifolia* var. *magna* with improved seed production, drought tolerance, and seedling vigor for the Great Lakes states using four accessions previously selected for the desired characteristics.

**Objective** Develop release (Koch) and compare growth to parental lines and standards.

**Release Development** Koch Prairie sandreed (*Calamovilfa longifolia* var. *magna*) Accession 9086408 was derived from the four most promising accessions from Rose Lake PMC Project 26I057F (Table 4). Koch was developed through three phenotypic mass selection cycles as described below.

Table 4. Parents of potential Koch Prairie Sandreed ( <i>Calamovilfa longifolia</i> var. <i>magna</i> ).						
Accession	Material	Comparative Strengths	Collection Location		Year	Collector
477007	Seed	Leafy foliage and later-maturing seed heads	Manistee National Forest Recreation Area	Mason County, MI	1971	Dorian Carroll
9004939	Vegetative material	Excellent vigor and abundant foliage	Ottawa County, MI		1978	Ellis Humphrey
9004944	Vegetative material	Abundant uniform foliate and good vigor	Huron county, MI		1978	Ellis Humphrey
9004959	Vegetative material	Good vigor and abundant foliage	Porter County, IN		1978	D.V. Wilson

**Cycle #1** of the phenotypic mass selection process began in 1989. A grid system was used to select fifty healthy, greenhouse-grown plants of each accession (477007, 9004939, 9004944, and 9004959). These plants were established in a breeding block and F1 seed was harvested in 1991. The plumpest 3,000 seeds were selected and planted three seeds per cone-tainer in the greenhouse during late winter and early spring 1992. Plants were thinned to one per cone-tainer. In mid-summer the seedlings were transplanted to a field selection nursery. These plants represented the F1 progeny and were assigned accession number 9055449.

In **Cycle #2** the 100 best and most upright F1 plants in the field selection nursery were selected and retransplanted into a polycross nursery in 1993. Three thousand F2 seeds were collected from the polycross nursery. These 3,000 seeds were grown three per container in 1,000 containers, thinned, and transplanted into a selection nursery.

In **Cycle #3** one hundred F2 plants were selected and retransplanted into a polycross, breeder nursery. Seed from this nursery was designated F3.

A foundation seed block was established with F3 plants grown from F3 seed. F4 seed from this foundation seed block is the seed that will be released as the Koch germplasm prairie sandreed, Accession 9086408.

**Materials, Methods, and Experimental Design for Progeny Test** Plugs were established in the greenhouse in 2003 or earlier and planted in 2004 in two-row plots. Design was a randomized complete block with four replicates (or fewer for limited-inventory accessions).

**Results** No significant differences were observed among Koch and its progenitors when vigor, disease, insect damage, lodging, and height were observed in early July and again in late August when disease was observed (Table 5). Both 9055449 and Koch flowered later than the one parental accession (477007). Leaf rust (*Puccinia sp.*) and anthracnose (*Colletotrichum graminicola*) infected all plot entries by late August.

Compared to Pronghorn, Koch was flowered earlier.

Compared to Goshen, Koch was more vigorous, less diseased at the later date, taller, and earlier flowering.

**Conclusion and Recommendation** Koch's equal or superior performance when compared to its progenitors and standards Pronghorn and Goshen support its release by the Rose Lake Plant Materials Program. Koch Germplasm Prairie Sandreed was released as a selected class ecotype by NRCS-USDA and the Michigan Association of Conservation Districts on 5 Sept 2007. A ceremony to recognize the contribution of past Plant Materials Program personnel including the late Philip Koch was held at the PMC.

Table 5. Prairie sandreed growth at Rose Lake PMC, 2007.							
<u>Accession</u>	<u>Early Vigor</u> <sup>1,2,4</sup>	<u>Early Disease</u> <sup>1,2</sup>	<u>Late Disease</u> <sup>1,3,4</sup>	<u>E. Ins. Damage</u> <sup>1,2</sup>	<u>Early Lodging</u> <sup>1,2</sup>	<u>Height</u> <sup>2,4</sup> (ft)	<u>Flowering Stage</u> <sup>2</sup>
Koch	1.0 B	1.0	3.3 BC	1	1	3.5A	Early flowering
9055449 (F1 of Parental Crossing Block)	1.0 B	1.0	4.3AB	1	1	3.5A	Early flowering
477007 (One of 4 Parents)	1.0 B	1.0	4.0AB	1	1	3.3AB	Flowering
Pronghorn (Standard)	1.3AB	1.0	2.3 C	1	1	3.6A	Boot
Goshen (Standard)	2.0A	1.0	4.8A	1	1	2.6 B	Boot
LSD <sub>(0.05)</sub>				n.s.	n.s.		
<sup>1</sup> Vigor, disease, insect damage, and lodging: 1=excellent vigor; no disease, insect damage, or lodging; 9=poor vigor; severe disease, insect damage, or lodging.							
<sup>2</sup> Evaluated 7 July 2007							
<sup>3</sup> Evaluated 30 Aug 2007							
<sup>4</sup> Means followed by same letter in same column are not different at LSD <sub>(0.05)</sub>							



**Study No. MIPMC-P-0209CR**  
**Evaluation of Dune Willow (*Salix cordata*)**

**Background** *Salix cordata* propagates efficiently from dormant cuttings. It is found growing on dry dune areas. The plant has potential conservation and landscaping uses on dune areas. It has the potential to trap sand or sediment on dunes and stabilize dune areas that are vulnerable to erosion.

One concern with using this species in dune restoration is that a flea beetle (*Altica subplicata*) uses the plant as a food source. The flea beetle can nearly defoliate a dune willow plant by September or October. Multiple years of exposure of dune willow to the flea beetle causes significant plant loss, and subsequent loss of dunescape erosion protection.

**Description of Study** A dune willow population will be established at the Rose Lake Plant Materials Center and evaluated for three to five years for vigor, plant growth, and incidence of disease or insects throughout the growing season.

**Procedure** A Sand Dune willow was collected at Warren Dunes by Michigan DNR personnel in 2001. Plants were propagated by planting dormant cuttings in greenhouse containers and transplanting the rooted cuttings in a field trial in 2002. A total of 58 plants were transplanted into the field trial.

**Summary** No new data are available from 2007 field studies.

## STABILIZE CROPPED AREAS OF CONCENTRATED WATER FLOW

Controlling sediment and nutrient runoff from cropland is a concern in the Rose Lake PMC service area. Developing methods that control erosion, while keeping the most land in production is a continuing effort.

**Problems** Grassed waterways are often utilized to address erosion and runoff in areas of concentrated flow within cropped fields. In small watershed areas where the water flow is less aggressive this practice is often dismissed in lieu of periodic filling and reshaping of the eroded area. This allows more land to remain in production and eliminates any hindrance the waterway may pose to farming operations. Erosion and runoff, however, continue to be a problem.

**Needs** Vegetative barriers should:

- not include potentially invasive or otherwise environmentally undesirable plants
- exhibit excellent erosion control qualities
- not impede normal farming operations
- be easy to maintain
- remove as little land from production as possible

### Study No. 26C126V

#### Vegetative Hedges for Controlling Erosion in Areas of Concentrated Flow

**Background** Pre-established strips of perennial grasses (vegetative barriers) transplanted within the crop rows and designed to impede water flow may have application in erosion control. The concept is that where runoff concentrates in rills or ephemeral gullies, grass hedges will pond water upslope causing a large portion of the sediment load to settle and fill the eroding areas. This will create small, benched terraces that diffuse and slow runoff, thus limiting further erosion and increasing water absorption. This study was initiated for miscanthus (*Miscanthus sinensis*) at Michigan State University Kellogg Biological Station (KBS) near Battle Creek, MI in collaboration with NRCS Field Office Staff in 1993. In 1996 the study was expanded to include native warm-season grasses. Additional field plantings, including miscanthus or eastern gamagrass (*Tripsacum dactyloides*), were established in Michigan, Indiana, and Ohio in 2001-03.

**Description of Study** Vegetative barriers of miscanthus or eastern gamagrass were evaluated for their effectiveness to control erosion in cropped fields.

**Procedure** Whole plants from miscanthus or eastern gamagrass stands on the MI PMC were harvested and divided into sprigs. A line of sprigs was planted 5 to 10 inches deep, spaced 3 inches apart, in rows 3 inches apart in soil-filled troughs of varying lengths. Troughs were formed with rolled plastic material to contain root growth, and lined with coconut fiber to maintain trough integrity during installation. Material was grown for several months in the greenhouse prior to installation insuring good root development. Troughs were lifted by the coconut fiber liner and placed into excavated strips of the crop

field perpendicular to the path of concentrated flow, and within the row so as not to interfere with farming operations.

**Summary** The miscanthus plantings established well and were healthy. There was no evidence of miscanthus spreading from where it was planted, and no seeds were evident. Soil sediment and crop residue deposits were evident on the upslope side of each barrier at KBS, but heavy rains caused some breaks in the row. Survey stakes are present in the barriers and sediment buildup measurements are planned.

The eastern gamagrass planting in Indiana showed good growth. Visual evaluation of the gamagrass and miscanthus plots in Indiana showed no down slope erosion. The plots will be maintained to evaluate sediment or residue entrapment and to further monitor the development of rills or gullies.

**Study No. MIPMC-P-0208-CR**  
**Intercenter Component of**  
**Evaluation and Release of Miscanthus Grass (*Miscanthus sinensis*)**

**Background** The Rose Lake PMC has been working for about 15 years with miscanthus grass as a planting for vegetative barriers in concentrate flow areas. Miscanthus has performed extremely well in the field, but concerns have been raised (e.g. <http://horticulture.coafes.umn.edu/miscanthus/>) about the possibility of its becoming an invasive species. The PMC has had field plantings for about ten years and has not identified any spread of this specific collection by seed.

**Procedure** Plants from the Rose Lake PMC were divided, repotted in 2.5-in diameter plastic cones, and sent to seven other PMCs in 2002. Cooperating PMCs were requested to record plant growth data and collect seeds for viability testing.

**Results** Miscanthus growth and development were excellent at all reporting PMCs and other cooperating locations. In all the years of intercenter testing, no viable seed was reported from laboratory germination tests (Table 6). However, Missouri conducted a greenhouse germination test that did show germination. Dr. Suleiman Bughrara at Michigan State University reported very low pollen viability in preliminary testing done by his laboratory on pollen produced from miscanthus plants grown at the PMC in 2007. These results suggest the need for further work to determine if and where the plant will develop viable seed.

Table 6. Miscanthus seed analysis as conducted by Michigan Department of Agriculture Laboratory Division.		
<b>Growing Location</b>	<b>Seed Production Year</b>	<b>Laboratory Results</b>
Big Flats, NY	2004	No TZ test due to absence of seed. No fertile florets detected.
Big Flats, NY	2005	0% germination. All florets appeared to be sterile.
Elsberry, MO	2005	0% germination. No fertile florets detected.
Big Flats, NY	2006	0% germination. Dead seed present.

## COVER CROPS

Cover crops are legumes, grasses, or other herbaceous plants established for seasonal cover and other conservation purposes. Cover crops protect the soil from erosion during vulnerable times, e.g., after harvest in the fall through establishment of crop in the spring.

Organic and sustainable farming systems are especially reliant on cover crops for soil fertility and pest management. Legume cover crops capture nitrogen from the air. Cover crops capture nitrogen and release it as the cover crop decays. Weeds are suppressed and disease cycles are broken with appropriate timing. Cover crops may also provide supplemental forage for grazing or harvesting.

### Study No. MIPMC-T-0604-CP

#### Evaluation of Hairy Vetch (*Vicia villosa*) Populations for Winter Hardiness

**Background and Objective** USDA ARS, Beltsville, Maryland is evaluating vetch (*Vicia villosa*) as a cover crop for increasing soil fertility and suppressing weeds in organic systems. Five hairy vetch populations (K-12, B-35, AUEC, Groff's selection, and Nebraska common) with potential for organic farming systems have been identified. Until winter hardiness has been documented, their use cannot be recommended. Objective of this study is to determine winter hardiness of five hairy vetch populations at multiple locations in the northern regions of the US.

**Approach** NRCS Plant Materials Centers at Rose Lake, MI; Bismark, ND; Manhattan, KS; Bridger, MT; Alderson, WV; Big Flats, NY; and Elsberry, MO were identified as testing sites. Phosphorus and K were brought to a medium or high level according to soil test recommendations. Hairy vetch entries were direct seeded in fall 2006 and again in fall 2007 according to a three replicate, randomized complete block design provided by ARS. Seed was inoculated with appropriate *Rhizobium sp.* for hairy vetch. Planting dates were based on recommendations for cool season legumes in the respective regions. Herbicides and hand rouging were used to control weeds.

**Data Collection** Number of hairy vetch seedlings in the late fall and again in spring when plants begin active growth are being counted. Survival will be calculated as a percent of the plants that survived the winter. Flowering dates will be recorded in spring 2008. Data from various PMCs is being compiled and analyzed at ARS.

## BIOENERGY

With the increased concern over dwindling reserves of known fossil fuels and possible global climate change has come commensurate interest in plant biomass as an energy source. Interest in bioenergy plants for their role in carbon sequestration is also increasing.

The National Plant Materials Program Strategic Plan calls for increased alternative uses and specialized uses of conservation plant releases to meet emerging needs (Objective 2.3, 2006-2011 Plan). Further, it suggests collaboration by the Plant Materials Program with external agencies to develop bioenergy technology. The following meets Strategic Plan objectives in both its research application and collaborative approach.

### Study No. MIPMC-T-0709-CP

#### **Tall Wheatgrass Biofeedstock Study for Potential Use in Biofuel (Liquid, Thermal, And Thermochemical) Applications in NE, Mid-Atlantic, and Upper Midwestern States**

**Background and Objective** Efforts are increasing to develop grass biofeedstocks for various biofuel applications. A tall wheatgrass cultivar from Hungary, ‘Szarvasi-1’ Energygrass [*Thinopyrum ponticum* (Podp.) Z.-W. Liu & R.-C. Wang], has been suggested as a biofeedstock and alternative to switchgrass and other warm-season grasses in the Northeast and other moist and cool environments. The objective of this study is to field test the performance of ‘Szarvasi-1’ Energygrass as a biofuel stock at multiple locations in the Northeast.

**Materials and Methods** Three commercially available plant releases of tall wheatgrass from the US will be compared ‘Szarvasi-1’ which has been used as a bioenergy grass. The grasses will be tested in replicated studies at two seeding rates at participating Plant Materials Centers. One cultivar of reed canarygrass (*Phalaris arundinacea*) at one seeding rate will be used as a standard.

Plots were established 12 Sept 2007 at the Rose Lake PMC with personnel and equipment assistance from the Michigan State University Agronomy Farm.

**Data Collection** Stand density and longevity and plant response to environmental factors including disease and insects will be observed throughout the life of the study. Above ground biomass production and forage quality for biofuel production will be determined.

All data will be submitted to NY Plant Materials Specialist who will develop a final report for inclusion in all participating PMC annual reports and for publication.

## PLANTS FOR WETLAND RESTORATION

Wetlands are transitional areas between terrestrial and fully aquatic ecosystems. They include swamps, marshes, and bogs. Wetlands perform many important functions: They improve water quality by trapping nutrients such as nitrogen and phosphorous, toxic substances, and disease-causing microorganisms. They slow and intercept runoff, protect shorelines and banks from erosion, and protect upland areas from floods. Wetlands also provide valuable habitat for wildlife.

**Problem** Historically, humans have made large-scale efforts to drain wetlands for development or to flood them for use as recreational lakes. Since the 1700s the US has lost 50% of its wetlands. With this loss and degradation of wetlands comes:

- decreased water quality
- lost wildlife habitat
- increased flooding
- less groundwater recharge
- diminished recreational opportunities and aesthetics

**Need** In order for wetlands to be restored and for new wetlands to be created, information related to plant species establishment and management is required and plant material must be made available. The Rose Lake PMC has assembled and is testing Canada bluejoint (*Calamagrostis canadensis* (Michx.) Beauv.) to address this need.

### **Study No. MIPMC-P-0603-WE Evaluation and Release of Canada Bluejoint (*Calamagrostis canadensis* (Michx.) Beauv.)**

**Background** Canada bluejoint grows in dense tufts 1.5 – 5 ft tall, typically in meadows, bogs, and wet thickets or as an understory plant in open, wet woods. Seed heads are open and nodding during flowering, becoming somewhat contracted as the seed ripens. Each flower has a small tuft of hair at the base. The foliage and stem are comparatively soft and lax. This grass gets its name from the bluish-purple stem nodes and slender, bluish leaves. Leaf blades have a rough surface. The roots have numerous creeping rhizomes. The seed heads are purplish-green during flowering. Mature seed are smooth, yellow-brown, ellipsoid, and about 1/16 inch long. Bluejoint is perennial and native to much of the US except the Gulf Coast states. *C. canadensis* is a widely variable species.

**Description** A collection of Canada bluejoint was assembled from native stands. This material will be evaluated and selected for potential release.

**Materials and Methods** Native Canada bluejoint from Indiana, Michigan, and Wisconsin was assembled and accessioned. Plants were established from seed in the greenhouse and transplanted into the field. Field plots were established in a wet area in 2006 in LaPorte County, Indiana and at the PMC in Michigan with accessions listed in

Table 7. Field plots are being evaluated for survival, vigor, disease and insect damage, height, lodging, seed production, and germination over the duration of the study plan.

Table 7. Canada bluejoint accessions established in 2006 in evaluation plots in Michigan and Indiana.		
Accession	Origin	Entry Location
9084188	Chippewa County, Michigan	MI
9084330	Pulaski County, Indiana	IN, MI
9084331	White County, Indiana	IN, MI
9084357	Polk County, Wisconsin	IN, MI
9084529	Starke County, Indiana	IN, MI
9086445	Roscommon County, Michigan	MI
9086446	Roscommon County, Michigan	IN, MI
9086529	Ontonagon County, Michigan	IN, MI
9086623	Ontonagon County, Michigan	IN, MI

**2007 Results** Plant growth data from the Rose Lake PMC plots is presented in Table 8. In accessions where plants survived in only 2 or 3 replicates, missing values were supplied by Statistix®'s missing value function and analysis of variance was performed with degrees of freedom adjusted. Significant differences were observed among accessions in vigor, disease, and height.

Plant growth data from the Indiana plot is presented in Table 9. Statistix®'s missing value function was used to supply missing values for accessions where plants survived in only 3 of 4 replicates. The accession where only 1 replicate survived was not included in the analysis of variance. Of accessions included in the analysis of variance, significant differences were observed in height. Vigor, disease, insect damage, and lodging observations were not recorded.



Table 8. Canada bluejoint plant growth data, Rose Lake PMC site. 2007. <sup>1</sup>							
Accession	n <sup>2</sup>	Vigor <sup>3,4</sup>	Disease <sup>3,4</sup>	Insect Damage <sup>3,4</sup>	Lodging <sup>3,4</sup>	Height (ft) <sup>4</sup>	Growth Stage
9084188	4	2.3AB	2.3AB	1A	1.0A	2.0AB	vegetative
9084330	4	2.3AB	2.0AB	1A	1.3A	2.0AB	vegetative
9084331	3	2.2AB	2.2AB	1A	0.9A	1.9AB	vegetative
9084357	4	2.3AB	2.5AB	1A	1.3A	2.0AB	vegetative
9084529	4	2.8A	2.8A	1A	1.3A	1.8 B	vegetative-early flowering
9086445	2	2.3AB	2.2AB	1A	0.9A	1.8AB	late vegetative-early flowering
9086446	4	1.5 B	1.8 B	1A	1.0A	2.4A	flowering
9086623	4	2.3AB	2.5AB	1A	1.3A	1.6 B	vegetative
9086529	No plants survived. Not included in ANOVA.						
<sup>1</sup> Evaluated on 16 July 2007.							
<sup>2</sup> Number of replicates included in ANOVA.							
<sup>3</sup> Vigor, disease, insect damage, and lodging: 1 = excellent vigor; no disease, insect damage, or lodging; 9 = poor vigor; severe disease, insect damage, or lodging							
<sup>4</sup> Means with the same letter within the same column do not differ significantly at LSD <sub>(0.05)</sub>							

Table 9. Canada bluejoint plant growth data, LaPorte County, Indiana. 2007. <sup>1</sup>			
Accession	n <sup>2</sup>	Height (ft) <sup>3</sup>	Growth Stage
9084330	4	2.9A	flowering
9084331	3	1.7 BC	vegetative
9086623	3	0.9 C	vegetative
9084529	4	2.5AB	late vegetative-early flowering
9084357	3	1.8 BC	vegetative
9086446	4	2.6AB	flowering
9086529	Survived in only 1 of 4 reps. Not included in ANOVA.		
<sup>1</sup> Evaluated on 25 July 2007.			
<sup>2</sup> Number of replicates included in ANOVA.			
<sup>3</sup> Means with the same letter within the same column do not differ significantly at LSD <sub>(0.05)</sub>			

Data will be collected from the Michigan and Indiana plots in 2008.

## RESTORATION OR RECLAMATION OF DISTURBED AREAS

Numerous human activities have the potential to dramatically alter the natural resources in an area. Construction of roads, travel lanes or utility corridors, agricultural or mining operations, and municipal or recreational development often results in a drastic disturbance of the natural communities. Successful revegetation of these disturbed areas requires plant materials and the technology to use them.

**Problems** With the heightened interest and promotion of native species, the availability of regionally-sources and adapted ecotypes has not kept pace with demand. This is especially true with native grass species for use in Conservation Reserve Program plantings and restoration work. Some non-native species traditionally utilized for conservation activities have come under scrutiny due to their aggressive nature. This has generated an even greater need for native plant material and the corresponding research to replace these conventional conservation species.

**Needs** The states served by the Rose Lake PMC need commercially available quantities of regionally native, or adapted, non-aggressive plant species, and the technology to use them. Selected species should have proven capabilities for one or more conservation concerns.

### **Study No. MIPMC-P-0201-CR Great Lakes Composite of Virginia Wildrye (*Elymus virginicus*)**

**Background** Virginia wildrye is a native, cool-season perennial bunchgrass with erect stems that reach to 4-ft high. Leaves are flat, up to 0.5-in wide, and rough on both sides and the margins. Spikes are stiff and up to 5-in long. The lower portion of the spike is often enclosed by the sheath. Lemmas have awns that reach 1.5 inches. Auricles are claw-like and clasping. Virginia wildrye is found in moist woods, meadows, and prairies throughout the United States east of the Rockies. It has good tolerance to flooding and moderate tolerance to drought. There are approximately 96000 seeds/lb.

**Description of Study** A collection of Virginia wildrye was assembled from native stands in Michigan, Indiana, Ohio, and Wisconsin. Material is being evaluated and composite selected for restoration or revegetation potential as conservation cover or streambank protection in the Great Lakes and Midwest states.

**Procedures** Virginia wildrye was collected from native stands by field staff and partners and accessioned. Plants from each of 19 accessions were started in the greenhouse and transplanted into field plots. Field plots in Michigan were established in 2002 and in Indiana and Ohio in 2003. Each plot was evaluated for survival, vigor, plant density, height, lodging resistance, disease and insect damage, seed production, and germination. A randomized complete block advanced field trial of “finalists” with three replications was established at the PMC. Plants were established in cone-tainers and transplanted to the field on 23 June 2005. Trial entries were:

9084531  
 9084344  
 9084514  
 Omaha (standard)

**Summary and Future Direction** Data recorded in 2007 from the advanced trial established at the PMC in 2005 is summarized in Table 10. Significant differences were observed in vigor, lodging, and height.

The advanced trial will be continued in 2008. Forage production will be evaluated. Selection decisions may be made at the end of the 2008 season.

Table 10. Virginia wildrye plant growth data, Rose Lake PMC advanced test site. 2007. <sup>1</sup>						
Accession	Vigor <sup>2</sup>	Disease <sup>2</sup>	Insect Damage <sup>2</sup>	Lodging <sup>2</sup>	Height (ft)	Growth Stage
9084351	1.0	2.0	1.0	1.7	3.0	Flowering
Omaha	2.7	2.0	1.0	1.0	1.8	Flowering
9084344	1.3	2.0	1.0	1.0	2.8	Flowering
9084514	1.3	2.0	1.0	1.0	2.5	Flowering
LSD <sub>(0.05)</sub>	0.7	n.s.	n.s.	0.6	1.0	n.s.
<sup>1</sup> Evaluated on 16 July 2007.						
<sup>2</sup> Vigor, disease, insect damage, and lodging: 1 = excellent vigor; no disease, insect damage, or lodging; 9 = poor vigor; severe disease, insect damage, or lodging						

**Study No. MIPMC-P-0207-CR**  
**Increase and Release of Riverbank Wildrye (*Elymus riparius*)**

**Background** The Wisconsin State Plant Materials Committee (WIPMC) suggested the need for native cool-season grasses for use in conservation activities, noting also the lack or unavailability of native cool season grasses from commercial growers. The WIPMC identified riverbank wildrye as a candidate for collection, study, and possible release as a conservation plant. Riverbank wildrye is an erect plant with relatively short awns compared to the awns of Canada wildrye.

**Study Description** A source collection was made by the WIPMC and was sent to Rose Lake PMC for study and evaluation.

**Procedure** Seed from the source collection was evaluated in the greenhouse for germination and early vigor during August of 2002. Germination was very high in greenhouse plantings. The remainder of the seed was planted into a seed increase field at the PMC in September of 2002. Plants grown in the greenhouse for the germination evaluation were transplanted into the same seed increase field.

Growth resumed in spring of 2003 and seed was produced from most plants in the field during that growing season. Seed from that field was harvested with a combine and cleaned with a fanning mill at the PMC. Seed cleaning was relatively easy, compared to Canada wildrye, because of the short awns on the riverbank wildrye seed.

Seed was provided to PM committee members in 2004 to plant in critical area treatments to determine if riverbank wildrye can be used effectively as a conservation plant.

**Results** Observations in 2006 from the Wisconsin planting were “has potential” and “good growth” even where precipitation was unfavorable and reed canary grass was invading.

**Study No. MIPMC-P-0204-CR**  
**Great Lakes Composite of Bottlebrush Grass (*Elymus hystrix*)**

**Background** Bottlebrush grass is a cool-season, perennial bunchgrass. It may be found growing in moist to dry woods from Nova Scotia to Manitoba in Canada and south to Arkansas and Georgia. Bottlebrush-shaped spikes may be up to 10-in long with 1-4 spikelets per node. Spikelets spread horizontally as they mature, often becoming nearly perpendicular to the rachis. Leaf sheaths may be smooth or hairy. Leaf blades may be up to ½-in wide with rough texture. Lemmas have rough, straight awns that reach 1½ -in long. Plants grow to 4 ft.

**Description of Study** A collection of bottlebrush grass was assembled from native stands in Michigan, Indiana, Ohio and Wisconsin. Materials are being evaluated for restoration or revegetation potential as conservation cover or streambank protection in the Great Lakes and Midwest states. Seed from the breeder field will be made available to growers as a selected or tested class release.

**Procedures** Collections were made from native stands by field staff and partners. Each collection was accessioned and cleaned. Cone-tainerized plants from each accession were established in the greenhouse and transplanted to field sites at the PMC and in Indiana and Ohio in 2003. Plots were placed in a randomized complete block design with 3 replications and evaluated for survival, vigor, plant density, height, lodging resistance, disease and insect damage, seed production, and germination.

Based on evaluations of plots established in 2003, these “Finalists” were identified and placed in a randomized complete block field plot with 3 replicates at the Rose Lake PMC in June 2005 after having been established in greenhouse cone-tainers:

9086418

9084535

9084191

9084533

9084360

9084186

**Summary** Data recorded in 2007 at the PMC is summarized in Table 11. Differences among entries were not observed, except in height where accession 9084533 was significantly shorter than 9084360. This height difference was consistent with 2006 observations when 9084533 was shorter than four other accessions including 9084360.

Table 11. Bottlebrush grass plant growth data, Rose Lake PMC advanced test site. 2007. <sup>1</sup>							
Accession	Vigor <sup>2</sup>	Disease <sup>2</sup>	Insect Damage <sup>2</sup>	Lodging <sup>2</sup>	Height (ft)	Median Flowering Stage	Harvested Seedheads/Plot
9086418	2.7	2.0	1.0	1.0	2.2	Flowering	27
9084535	2.3	2.0	1.0	1.0	2.3	Flowering	37
9084191	2.3	2.0	1.0	1.0	2.3	Flowering	36
9084533	2.7	2.0	1.0	1.0	1.8	Flowering	29
9084360	2.0	2.0	1.0	1.0	2.5	Flowering	23
9084186	2.7	2.0	1.0	1.0	2.3	Flowering	23
Grand Mean	2.4	2.0	1.0	1.0	2.3		29
LSD <sub>(0.05)</sub>	n.s.	n.s.	n.s.	n.s.	0.6		n.s.
<sup>1</sup> Established 24 June 2005. Evaluated 16 July 2007.							
<sup>2</sup> Vigor, disease, insect damage, and lodging: 1 = excellent vigor; no disease, insect damage, or lodging; 9 = poor vigor; severe disease, insect damage, or lodging							

**Study No. MIPMC-P-0602-CR**  
**Evaluation and Release of Poverty Oatgrass (*Danthonia spicata* (L.) Beauv. ex Roemer & J.A. Schultes)**

**Background** Poverty oatgrass is native to the Rose Lake PMC service area. This perennial, bunch-type, cool-season grass is typically 1-2 feet tall (smaller and fewer leaves on poor soil) with curly basal leaves that straighten in damp weather. Gray's Manual of Botany describes poverty oats as "A polymorphous and very plastic species sadly in need of critical study."

Poverty oats have two types of florets: (1) florets that cross-fertilize (chasmogamous) located on the aerial panicle and containing more pollen grains and (2) the unopened, self-fertilized florets (cleistogamous) that are located inside one or more of the leaf sheaths throughout their development. This trait was observed at the PMC plots and corroborated by a literature review.

**Description** A collection of poverty oatgrass was assembled from native stands in Indiana and Michigan in 1998-2005. This material will be evaluated and selected.

**Materials and Methods** Native poverty oatgrass from the PMC service area was assembled and accessioned. Plants were established from seed in the greenhouse and

transplanted into the field. Field plots in LaPorte County, Indiana and at the Rose Lake PMC were established with the accessions listed in Table 12. Field plots are being evaluated for survival, vigor, disease and insect damage, height, lodging, seed production, and germination over the duration of the study.

**2007 Results** Plant growth data from the Rose Lake PMC is presented in Table 13. Four accessions did not survive in any of the replicates and two accessions only survived in one replicate. These accessions were not included in the analysis of variance. Of the remaining accessions significant differences were observed in height and number of florets.

When Indiana plots were evaluated on 25 July 2007 surviving poverty oatgrass could be identified in only about 20% of plots. Dry soil conditions at time of transplanting, weather, and weed pressure were likely contributors to the stand failure. Surviving accessions that could be identified and number of replicates are shown in Table 14.

Table 12. Poverty oatgrass accessions established in 2006 in evaluation plots in Michigan and Indiana.		
Accession	Origin	Entry Location
9084165	Gibson County, Indiana	IN, MI
9084166	Pike County, Indiana	IN, MI
9084175	Wexford County, Michigan	IN, MI
9084189	Chippewa County, Michigan	IN, MI
9084311	Pike County, Indiana	IN, MI
9084314	Gibson County, Indiana	IN, MI
9086433	Dubois County, Indiana	IN, MI
9086434	Dubois County, Indiana	IN, MI
9086620	Ontonagon County, Michigan	MI
9086507	Kalamazoo County, Michigan	MI

Table 13. Poverty oatgrass plant growth data, Rose Lake PMC site. 2007.							
Accession	Vigor <sup>1,2</sup>	Disease <sup>1,2</sup>	Insect Damage <sup>1,2</sup>	Lodging <sup>1,2</sup>	Height (ft) <sup>1</sup>	Growth Stage <sup>1</sup>	Number of Florets <sup>3</sup>
9084175	1.0	1.0	1.0	1.0	2.0	flowering	166
9084189	1.0	1.0	1.0	1.0	1.8	flowering	215
9084314	1.5	1.0	1.0	1.5	1.0	flowering	118
9086507	1.0	1.0	1.0	1.0	1.8	flowering	141
LSD <sub>(0.05)</sub>	n.s.	n.s.	n.s.	n.s.	0.5		78
9084165	No plants survived. Not included in ANOVA.						
9084166	No plants survived. Not included in ANOVA.						
9084311	Plants survived in only 1 of 4 reps. Not included in ANOVA.						
9086433	No plants survived. Not included in ANOVA.						
9086434	No plants survived. Not included in ANOVA.						
9086620	Plants survived in only 1 of 4 reps. Not included in ANOVA.						
<sup>1</sup> Evaluated on 16 July 2007.							
<sup>2</sup> Vigor, disease, insect damage, and lodging: 1 = excellent vigor; no disease, insect damage, or lodging; 9 = poor vigor; severe disease, insect damage, or lodging							
<sup>3</sup> Evaluated 31 July – 1 Aug 2007							

Table 14. Poverty oatgrass survival, LaPorte County, Indiana. 2007. <sup>1</sup>	
Accession	Replicates with at Least One Surviving Plant
9084175	1
9084311	1
9084314	1
9086433	3
<sup>1</sup> Evaluated on 25 July 2007.	

Observations at both plot sites are planned for 2008 with continued data collection planned for the Michigan plots.



## IMPROVED WARM-SEASON FORAGE GRASS

Warm and cool-season grasses have contrasting patterns of yield distribution. Warm season grasses produce more than 60% of their yield in mid-summer, while cool-season grasses have their greatest production in spring and fall. Cool and warm-season grasses can best be used in grazing systems that utilize separate pastures for each grass type. Including warm-season grasses in a grazing system permits resting cool-season grasses in mid-summer which improves their vigor and enhances forage production in the late summer and early fall. Cool-season grasses can be grazed in spring and fall and warm-season grasses during mid-summer.

**Problems** Currently, there are no commercial native warm season forage grass varieties originating from the area covered by the Rose Lake PMC. Concerns are growing over the loss of native germplasm.

**Needs** Native warm-season grasses are needed that:

- originate from the Great Lakes/Midwest States
- have good forage yields
- have good palatability and nutrition for livestock
- persist under grazing
- establish rapidly
- have good seed production

**Study No. MIPMC-T-0303-PA**  
**Eastern Gamagrass (*Tripsacum dactyloides*) for Forage Production**

**Background** Eastern gamagrass is a native warm-season perennial that can be found throughout the eastern half of the US. It is a highly productive grass that is best adapted to wet habitats. Remnant colonies are commonly found in flood plains along stream banks. Eastern gamagrass is in the same taxonomic family (Poaceae) as field corn (*Zea mays*) and is characterized by numerous short, well-developed rhizomes. Individual grass clumps can reach a diameter of 4 ft with seed heads growing on culms 3- to 9-ft tall.

Eastern gamagrass produces the majority of its growth from late spring through late September. It begins growing earlier in the spring than do other native grasses such as big bluestem (*Andropogon gerardii*) and switchgrass (*Panicum virgatum*). The distribution of eastern gamagrass yield throughout the summer makes this grass an excellent source of forage during the period of the year when cool-season grasses are relatively dormant.

Several eastern gamagrass varieties are well adapted to the central and southern United States. The Big Flats Plant Materials Center, near Corning, NY, is evaluating several eastern gamagrass accessions to determine adaptability in the northern regions of the United States. Little research has been done in Michigan to demonstrate adaptability of any eastern gamagrass releases or experimental accessions in the Great Lakes area.

**Forage Production Study Procedures** A forage production study was initiated by planting stratified Pete eastern gamagrass seed on 17 June 2003 in plots having row spacing of 15 inches or 30 inches between rows. Plots with 15-in row spacing were planted with a seeding rate of 2 seeds/row ft. Plots with 30-in row spacing were planted with a seeding rate of 4 seeds/row ft. Seeds were planted 1 inch deep. Each plot was divided in 2004 by superimposing two cutting heights (simulated grazing). The study was conducted in a randomized complete block experimental design with four replicates per treatment. The site was irrigated to insure good survival after planting. All plots were fertilized and herbicides were applied as needed. Data is being taken from the center of each plot throughout the course of the study.

**Forage Study Results and Discussion** Significant differences were not observed in forage yield in 2007 (Table 15) nor were they observed in 2006. Whether or not to continue in 2008 with yield data collection depends upon which of the following explanations are adopted for why significant yield differences were not observed:

- there were no differences in plant responses to treatments,
- different plant responses to treatments may only be observed in higher yield environments,
- forage sampling technique is inadequate for detecting yield differences, or
- differences in plant responses to treatments can only be observed later in the life of the stand.

Table 15. Effect of row width and cutting height on gamagrass forage yield and quality. Rose Lake PMC. 2007 <sup>1</sup> .	
	Yield (lbs/plot)
Wide, high	4687
Wide, low	6311
Narrow, high	5201
Narrow, low	7312
Mean	5878
LSD <sub>(0.05)</sub> within column	n.s.
<sup>1</sup> Harvested 7 Jun and 18 Jul 2007	

**Relative Maturity of Eastern Gamagrass (*Tripsacum dactyloides*) Accessions –  
Modification of Study No. MIPMC-T-0302-PA**

**Accession Study Procedures** In June 2003 seedlings of Highlander and Pete eastern gamagrass were received from the Jamie L. Whitten PMC at Coffeeville, MS and accessions 9086456 (NY Diploid) and Meadowcrest were received from the Big Flats PMC at Corning, NY. Seedlings were transplanted into a field study site at Rose Lake PMC. Each plot contained three rows of five plants. Plants were spaced 36 inches apart within the row and rows were spaced 30 inches apart. The study was conducted in a randomized complete block with three replicates per accession, although one replication was subsequently abandoned due to competition from surrounding vegetation. The site was irrigated to insure good survival after transplanting. All plots were fertilized and herbicides were applied as needed. Yield (2004-2006) and quality (2006) data were taken from the center row of each plot throughout the course of the study.

**Yield and Quality Results** Three years of yield data from this trial did not show yield differences among accessions that were consistent over the years and one year of quality data did not show differences among accessions. This lack of differentiation was perhaps due to the loss of one replicate as mentioned above and the compromised power of statistical tests based on just two remaining replicates. Rose Lake researchers therefore decided for the use this plot to comparing relative maturity of the four accessions in the 2007 season.

**Maturity Results** Accessions were cut when they reached a predetermined early flowering growth stage. Second cuttings were taken on accessions that achieved that same flowering stage a second time during the season. Growing degree day accumulations were calculated from daily low and high temperatures using 50° F minimum and 86°F maximum starting May 1. Relative maturities from earliest to latest were Pete, NY Diploid, Highlander, and Meadowcrest as shown in Table 16 below.

Table 16. Growing Degree Day (calendar day) accumulations to reach early flowering at Rose Lake PMC. 2007.		
	First Cutting	Second Cutting
Pete	501 (44)	1106 (60)
NY Diploid	613 (49)	1185 (66)
Highlander	776 (59)	Did not reach second early flowering stage
Meadowcrest	880 (66)	Did not reach second early flowering stage

**Application Summary** Graziers typically need to provide quality pasture for animals throughout the grazing season. This may be accomplished using warm and cool-season forages with differing relative maturities and growth characteristics. Relative maturity information as provided above may be useful in such a pasture establishment and management decision-making context.

**Study No. 26I101G**  
**Big Bluestem (*Andropogon gerardii*) for Forage Use**

**Background** Big bluestem is a native, perennial, warm-season grass reaching heights of 8 feet at maturity. It has long white hairs on the upper leaf surface near the base of the blade. Lower leaf sheaths and blades are sometimes hairy. Big bluestem is bluish in color during most of the summer, often becoming reddish purple when mature. Seedheads consist of two or three racemes that arise from a common joint on the stem and resemble a turkey's foot. Prior to European settlement it was a major constituent of the tall grass prairies and savannas across the Great Plains, Midwest, and northeastern United States.

Warm-season forage grasses such as big bluestem are being integrated into grazing systems to increase beef production during the summer months when cool-season forage production declines. Unlike cool-season grasses that have their greatest growth during cooler temperatures, warm-season grass production peaks at higher temperatures. Utilizing these contrasting patterns of yield distribution helps to ensure adequate feed throughout the summer months and enhance cool-season forage production in the late summer and fall.

Varieties of big bluestem vary in origin and maturity and should be selected for adaptability. Varieties adapted to southern regions are later maturing, taller, and more productive; however, they may not produce mature seed. There is also the potential for winter injury to occur under defoliation stress if grown and harvested by grazing or haying in northern regions. Varieties adapted to the north that are grown in the south mature earlier, are shorter, have lowered forage production, and are more susceptible to leaf and stem diseases.

**Description of Study** From assembled regional material select and release a big bluestem cultivar with improved forage characteristics. Other potential uses for this release will include wildlife habitat, erosion control, native habitat restoration, and biomass production.

**Procedure** More than 100 vegetative collections were assembled from Michigan, Indiana, Wisconsin, and Ohio in 1992. Each accession was started in the greenhouse, grouped into region of origin (northern, central, or southern), then transplanted to an initial evaluation site with 4 replicates. Data was collected on this material through the 1990s. Ten accessions were selected for advanced testing from each region based on vigor, foliage, biomass production, height, tillering, lodging, disease, and insect resistance. In 1998 advanced test sites were established in Wisconsin, Michigan, and Indiana for northern, central, and southern accessions, respectively. Each study site was established as a RCB design with 3 replicates. Two regionally-adapted, commercially-available varieties were also included at each test site for comparison. Data collected included survival count, vigor, foliage rating, heading date, height, lodging resistance,

and forage yield. A forage analysis for nitrogen, protein, phosphorus, and acid detergent fiber (ADF) was also completed.

**Summary and Plans – Michigan Plots** Five of 10 accessions from the Michigan advanced testing study (Table 17) were selected for further evaluation and establishment of seed increase plots based on plant growth and forage quality characteristics. All five showed better forage production characteristics than the standards Bonella and Champ. In summer 2007 vegetative material from each of the five accessions was separated, potted, and grown in the greenhouse in anticipation of additional testing and/or crossing in 2008.

Table 17. Big bluestem accessions selected from advanced testing and growing in isolation plots around Rose Lake Wildlife Area.	
Accession	Origin
9070139	Porter County, Indiana
9070149	Jasper County, Indiana
9070162	Porter County, Indiana
9070163	Jasper County, Indiana
9070197	Washtenaw County, Michigan

## FOREST IMPROVEMENT AND WINDBREAK TECHNOLOGY

Forestland accounts for up to 30 percent of the agricultural land in Michigan. Considerable emphasis has been placed on establishing, maintaining, and improving forestland in Michigan. Several NRCS conservation programs encourage these practices.

In addition to forestland, trees and shrubs are used in windbreaks, riparian areas, filter strips, and wildlife corridors. Windbreak planting is practiced on almost all soil types within the PMC service area.

### Problems

- Tree and shrub transplanting can be expensive and labor intensive.
- Direct seeding of tree and shrub species is not well understood in the PMC service area.
- Soil type influences tree or shrub species selection for windbreaks and other woody species plantings. Species establishment in muck soils is not as well understood as establishment in upland soils.

### Study No. MIPMC-T-0404-BU Windbreaks in Muck Soils

**Background** Huron County in North Central Ohio has a significant area of muck soil. Called Celeryville Marsh, this area is known for its vegetable production. However, wind erosion threatens the soils and their productivity. In the past farmers removed nearly all the trees to expand their fields because the trees exhibited undesirable characteristics. Now windbreaks are needed to protect this valuable soil. Renewed interest in establishing windbreaks has created a demand for information on which species will perform best and be accepted by farmers. Desirable characteristics include rapid growth, sufficient height for wind erosion protection, longevity, little breakage, low maintenance, and limited spread.

**Description of Study** Plant species were selected and evaluated for establishment and survival in windbreaks on muck soils. A series of “tall” species and a series of “short” species were planted in replicated plots to determine the effectiveness of these species as windbreak components.

**Materials and Methods** Twelve tree or shrub species were evaluated for effectiveness as windbreaks. “Tall” plants included arborvitae, hybrid crabapple, bald cypress, pin oak, and austree willow. “Short” plants included privet, silky willow, silky dogwood, highbush cranberry, Sargent’s crabapple, black chokeberry, and lilac.

The Rose Lake Plant Materials Center provided ‘Affinity’ arborvitae, Riverbend Germplasm silky willow, ‘Magenta’ hybrid crabapple, ‘Indigo’ silky dogwood, and

Leelanau Germplasm highbush cranberry. Ohio DNR and OARDC provided the other species needed for the study.

OARDC planted the species in randomized, replicated field plots during spring 2004. The species will be evaluated annually for growth characteristics and general windbreak effectiveness.

**Results** Data were not reported in 2007.

### **Study No. MIPMC-T-0301-WO Direct Seeding Tree and Shrub Establishment**

**Background** Direct seeding of hardwoods offers an alternative to the high cost of establishing new or re-stocking existing stands of hardwood species in Michigan. Direct seeding of hardwoods is successful in Indiana, Ohio, Wisconsin, Illinois, and Missouri. The practice is generally recognized as an effective alternative to planting hardwood tree seedlings. Direct seeding of hardwoods is currently a cost-shared component in of USDA conservation programs in some states. There is little current research or written information specific to Michigan for establishing or enhancing hardwood stands by direct seeding.

**Description of Study** A selection of hardwood tree and shrub species was planted by direct seeding in 2003. A second study was established in 2003 and 2004. Plots are being evaluated for emergence, survival, and plant growth characteristics for up to five years.

**Procedure** Seeds of seven heavy mast and seven light mast hardwood tree and shrub species were planted in field studies at the Rose Lake Plant Materials Center. (Species are listed in Table 18.) Planting dates were May 2003 and November 2003 in the first study. Planting density was approximately 4500 seeds/acre for each species. Soil in the test area is Boyer sandy loam or loamy sand. Each species was:

- 1) broadcast on tilled soil, followed by dragging and cultipacking;
- 2) drilled in rows (rows 1.5 ft apart, seeds 8-in spacing within rows) in tilled soil, followed by cultipacking; and
- 3) drilled in rows (rows 1.5 ft apart, seeds 8-in spacing within rows) in non-tilled soil, followed by cultipacking

Separate heavy mast and light mast studies were established each with three replicates in a randomized complete block design. Non-tilled plots were sprayed with glyphosate herbicide to control all emerged vegetation before seeds were planted. A rodent control product was placed in several locations across the trial. No fertilizer or irrigation was applied. Stands were evaluated in 2003, 2004, and 2005. Plants were counted in rows in the drilled plots and by a transect method in the broadcast plots.



A second study was established with plantings in fall 2003 and spring 2004 and with seeds drilled into tilled and non-tilled soil. Each treatment combination was planted in rows 1.5 ft apart with 8-in seed spacing within rows (43560 seeds per acre). There were three replicates in a randomized complete block design. Species were pignut hickory (*Carya glabra*), shellbark hickory (*Carya laciniosa*), bitternut hickory (*Carya cordiformis*), common winterberry (*Ilex verticillata*), and sugar maple (*Acer saccharum*).

**Results** Data collected in 2005 and reported in the 2005 RLPMC Technical Report suggest that fall plantings and heavier, larger mast seeds fare better than spring plantings and lighter, smaller mast seeds. Data were not collected in 2006 and 2007. However, data collection will resume in 2008 following which collection a final report will be written.

Results of the second study corroborated findings of the first study: Fall-plantings and heavier, larger mast seeds fare better than spring-plantings and lighter, smaller mast seeds.

**Discussion** Droughty soil and growing season conditions (summer of 2003) may have reduced the emergence of seedlings, especially for light-seeded species which were planted at a shallow, but appropriate depth. Fall planting dates give seeds the advantage of the earliest possible start to the growing season and an opportunity to extend roots deeper into the soil before summer heat and drought begin.

Drilled seed has the advantage of relatively accurate and consistent seed depth. Broadcast seeding produces a more natural plantation but a random seed depth; some will be deeper than ideal depth and some will be on the soil surface and subject to drying and consumption by seed-eating mammals and birds.

The trend in forest regeneration is toward increasing rates of trees planted per acre. A comparison of normal tree seedling planting practice shows that direct seeding can result in a higher number of seedlings per acre compared to seedling planting. A common seedling tree planting rate is 436/acre (10-ft X 10-ft spacing). Acceptable survival for many purposes is 80% or about 350 seedlings/acre. Several direct seeded species and techniques equaled or exceeded the typical seedling tree planting rates and survival expectations.

Although an economic analysis was not included in this study, it would appear that direct seeding could be accomplished at lower costs/acre than with seeding tree planting. Because direct seeding can result in higher numbers of seedlings per acre the cost per established seedling may be much lower with direct seeding than with seedling tree planting.

Table 18. Species planted in first direct seeding tree and shrub study in 2003.	
Common Name	Scientific Name
<b>Heavy Mast Species</b>	
Northern Red Oak	<i>Quercus rubra</i>
White Oak	<i>Quercus alba</i>
Scarlet Oak	<i>Quercus coccinea</i>
Bur Oak	<i>Quercus macrocarpa</i>
Black Walnut	<i>Juglans nigra</i>
Black Cherry	<i>Prunus serotina</i>
Shagbark Hickory	<i>Carya ovata</i>
<b>Light Mast Species</b>	
Arrowwood / Highbush Cranberry*	<i>Viburnum dentatum</i> / <i>Viburnum trilobum</i>
Staghorn Sumac	<i>Rhus typhina</i>
Green Ash	<i>Fraxinus pennsylvanica</i>
White Ash	<i>Fraxinus Americana</i>
Red Maple	<i>Acer rubrum</i>
Silver Maple	<i>Acer saccharinum</i>
White Birch	<i>Betula papyrifera</i>
* Arrowwood was used in the spring 2003 planting and highbush cranberry was used in the fall 2003 planting.	

**Study No. MIPMC-T-0402-WO**  
**Direct Seeding of Northern Red Oak (*Quercus rubra*)**

**Background** Beech bark disease causes significant defect and mortality in American beech (*Fagus grandifolia*). The disease results when bark, attacked and altered by the beech scale (*Cryptococcus fagisuga*) is invaded and killed by fungi, primarily *Nectria coccinea* var. *faginata*, and sometimes *N. galligena*.

Beech bark disease is becoming a serious pest in the Eastern Upper Peninsula (UP) and in the West Central Lower Peninsula of Michigan, threatening to destroy the beech component of these forests and spread throughout Michigan. Though not a prized wood for timber, American beech is an important heavy mast crop for wildlife. In some areas of Michigan, particularly the Eastern UP, it is the only heavy mast crop available to bear, turkey, deer, and grouse. Much interest has been generated in replacing beech as it dies off with red oak, since it is the hardiest and most northerly growing of the black oak group. There has been some discussion of whether direct seeding might be a reasonable alternative to transplanting.

**Materials and Methods** Side-by-side comparisons of direct-seeded and transplanted Northern red oak (*Quercus rubra*) were established in the Eastern UP in 2004. Site selection was based on proximity of broken canopy and open sunlight so oak seedling performance could be evaluated in both environments. Acorns and seedlings were supplied by the Rose Lake PMC to cooperators representing nine sites in the Upper

Peninsula. Suggested field design was a row of 100 red oak acorns to be planted next to a row of 25 red oak seedlings with three rows of each treatment per test site. Broken canopy and open sunlight sites were to be paired and in close proximity. Acorns were to be planted 1.5-in deep (2 times the diameter of the acorn) approximately 1 ft apart within a row with rows spaced 5 ft apart. Seedlings were to be spaced 4 ft apart within a row with rows spaced 5 ft apart.

Planting date was in spring 2004. Data were collected in 2004, 2005, 2006, and 2007 on oak survival and average plant height. Data were also to be collected on other species that establish in the test site, including species name, density, and height.

**Results and Discussion** Survival and height data are presented in Table 19. Survival and height generally appeared greater in transplants than direct-seeded plants. Differences in response to broken canopy vs. open sunlight were inconsistent. Little or no increase in plant height was observed from 2004 to 2006, perhaps due to extremely dry summers.

Design differences among the eight locations rendered them not amenable to an across-all-sites statistical analysis.

Table 19. Performance of direct seeded and transplanted northern red oak in Michigan's Upper Peninsula.										
Location	Site Description	Treatment	Survival (%)				Height (inches)			
			2004	2005	2006	2007	2004	2005	2006	2007
Luce County (site 1)	Broken canopy opening, northern hardwood stand	Direct Seeding	91	91	77	62	4	4	5	7
		Transplant	100	100	92	100	8.6	9.3	10.3	14
Luce County (site 2)	Edge open meadow, east side	Direct Seeding	40	40	36	37	4	4	5	7
		Transplant	90	86	76	76	7.2	8	7	10
Luce County (site 3)	Edge open meadow, west side	Direct Seeding	20	25	21	21	3	3	5.5	10
Luce County (site 4)	Edge open meadow, lower elevation	Direct Seeding	67	60	55	63	4	4	6	9
		Transplant	100	95	95	89	7.6	8.5	11.8	18
Luce County (site 5)	Broken canopy opening, aspen clear cut	Direct Seeding	71	82	32	46	3	3	3	3
		Transplant	67	67	22	56	6.5	7.5	5.5	7
Luce County (site 6)	Open meadow	Direct Seeding	60	40	23	18	4	4	4.3	5
		Transplant	83	83	77	70	7.2	8.2	6.6	9
Luce County (site 7)	Broken canopy opening, aspen	Direct Seeding	37	38	42	14	4	4	4	6
Mackinac County (Northern Timberland Ventures Inc., site)	old pasture reverting back to forest. sandy loam	Direct Seeding	71	31	9	No data available in 2007	3.7	4	3	No data available in 2007
		Transplant	95	77	16	No data available in 2007	8.6	9	9.3	No data available in 2007

## NATIVE PLANT SPECIES TO ENHANCE WILDLIFE HABITAT

With controversy surrounding non-native species, particularly in undomesticated settings, interest in native ecotypes has risen sharply. Some non-native species historically used in wildlife plantings have been labeled aggressive or invasive or less beneficial to wildlife than many native species.

**Problem** Availability of Great-Lakes area native plant species for wildlife use is limited.

**Needs** Native species are needed for wildlife food plots, shelter, nesting, and brood rearing cover that:

- originate from the Great Lakes/Midwest States
- have good survival, vigor, and seed and foliage production
- have documented wildlife benefit
- meet the criteria for non-invasive plants

### Study No. 26I080J

#### Tick-trefoil (*Desmodium spp.*) for Wildlife Food Plots

**Background** In the United States there are 30 species of *Desmodium*. All are native, perennial legumes with trifoliolate (or rarely 1 to 5 foliate) leaves, purple flowers, and flat, deeply lobed or jointed pods. The joints of the pods easily separate and attach to clothing or animals by means of small hooked hairs, hence the common names tickclover, tick trefoil, and beggar's lice. These species are well distributed throughout most of the eastern and central states with several also in the southwest. Most inhabit dry, sandy, open woods or slightly shaded areas. Seeds of *Desmodium* have been found in the stomachs of masked bobwhite, lesser scaup duck, eastern ruffed grouse, slate-colored junco, ring-necked pheasant, willow ptarmigan, Gambel quail, Mearns quail, red-eyed towhee, Virginia opossum, and Bangs flying squirrel. Their seeds are also said to be eaten by the greater prairie chicken and sharp-tailed grouse in Minnesota and eastern turkey in Missouri.

**Description of Study** This study plans calls for assembly and evaluation of *Desmodium spp.* and selection of a superior accession for use in establishment of wildlife food plots. The development of harvesting, cleaning, and seeding procedures for seed and plant increase will be included.

**Procedure** In 1988, seed from 49 accessions of various *Desmodium* species was field collected from eight states and 16 MLRAs, and assembled at the Rose Lake Plant Materials Center. Each accession was grown in the greenhouse for preliminary observation the following year. In 1990, forty accessions were transplanted into field plots arranged in a randomized, complete block for an initial 2-year evaluation period. Five accessions were selected for advanced testing based on survival, vigor, foliage,

flower and seed production, and maturity date. The advanced trial was completed in 1992.

**Releases** Three accessions were selected for increase and potential release (Table 20). Two accession were issued as tested releases for the Great Lakes and upper Midwest regions and named after their county of origin. Generation 1 plants will be maintained at the Rose Lake Plant Materials Center. Foundation seed increase seed plots were established for each potential release in 2003 with additional plants added to those fields in 2004 and 2005.

**Table 20. *Desmodium* species released or selected for potential release.**

Accession No	Release Name	Scientific Name	State of Origin	Maturity Period	Release Date
9005087	Marion Germplasm Dillenius' tick-trefoil	<i>Desmodium glabellum</i>	IL	Mid-season	pending
9055415	Alcona Germplasm Dillenius' tick-trefoil	<i>Desmodium glabellum</i>	MI	Early-season	2006
9055428	Grant Germplasm Panicleleaf tick-trefoil	<i>Desmodium paniculatum</i>	WI	Mid-season	2006

## APPENDIX A

### RELEASED MATERIAL

'Affinity' (*Thuja occidentalis* L.) Northern White Cedar

Released: 1993 (FY1993)

Accession Number: 9005060 (PI Number: 477011)

Release Type: cultivar

Plant Origin: native

Collection Location: Pulaski Co., IN

Plant Type: tree

Plant Duration: perennial

Propagation: seed

Uses: field and farmstead windbreaks, screen or border planting in urban situations, and winter browse

Alcona Germplasm (*Desmodium glabellum* (Michx.) DC.) Dillenius' Tick-Trefoil

Released: 2006 (FY2006)

Accession Number: 9055415

Release Type: tested germplasm

Plant Origin: native

Collection Location: Alcona County, MI

Plant Type: legume

Plant Duration: perennial

Propagation: seed

Uses: wildlife food plots as an alternative to introduced plant species

Grant Germplasm (*Desmodium paniculatum* (L.) DC.) Panicleleaf Tick-Trefoil

Released: 2006 (FY2006)

Accession Number: 9055428

Release Type: tested germplasm

Plant Origin: native

Collection Location: Grant County, WI

Plant Type: legume

Plant Duration: perennial

Propagation: seed

Uses: wildlife food plots as an alternative to introduced plant species

Icy Blue Germplasm (*Elymus canadensis* L.) Canada Wildrye

Released: 2004 (FY2004)

Accession Number: 9084347 (PI Number 641962)

Release Type: tested germplasm

Plant Origin: native

Collection Location: LaPorte County, IN

Plant Type: cool-season grass

Plant Duration: perennial

Propagation: seed

Uses: restoration, wildlife cover, and erosion control

Registration Document: Durling, J.C., J.W. Leif, and D.W. Burgdorf. 2006. Registration of Icy Blue Canada Wildrye Germplasm. Crop Sci. 46:2330-2331.

'Imperial' (*Populus canadensis* Moench var. *eugenei* (Simon-Louis) Schelle) Carolina Poplar  
Released: 1979 (FY1979)  
Accession Number: Mich-88 (PI Number: 432347)  
Release Type: cultivar  
Plant Origin: introduced  
Collection Location: Rice Co., MN  
Plant Type: tree  
Plant Duration: perennial  
Propagation: vegetative  
Uses: windbreaks (especially around orchards) and pulpwood

'Indigo' (*Cornus amomum* P. Mill.) Silky Dogwood  
Released: 1982 (FY1982)  
Accession Number: 9031863 (PI Number 468117)  
Release Type: cultivar  
Plant Origin: native  
Collection Location: Clinton Co., MI  
Plant Type: shrub  
Plant Duration: perennial  
Propagation: seed or vegetative  
Uses: single row windbreak under center pivot irrigation, field and farmstead windbreak, soil bioengineering, and wildlife food

Koch Germplasm [*Calamovilfa longifolia* (Hook.) Scribn. var. *magna* Scribn. & Merr.] Prairie Sandreed  
Released: 2007 (FY2007)  
Accession Number: 9086408  
Release Type: selected  
Plant Origin: native  
Collection Location: costal zones along Lakes Michigan and Huron  
Plant Type: warm-season grass  
Plant Duration: perennial  
Propagation: seed or vegetative  
Uses: wind erosion control, dune stabilization, and water quality improvement

'Lancer' (*Lathyrus latifolius* L.) Perennial Pea  
Released: 1984 (FY1984)  
Accession Number: 477009 (PI Number 477009)  
Release Type: cultivar  
Plant Origin: naturalized  
Collection Location: MI  
Plant Type: legume  
Plant Duration: perennial  
Propagation: seed  
Uses: erosion control plant, wildlife cover plant, land reclamation, brush management, roadside seeding mixtures, critical area planting where objective includes beautification



Leelanau Germplasm (*Viburnum opulus* L. var. *americanum* Ait.) Highbush Cranberry

Released: 1999 (FY1999)

Accession Number: 9031863

Release Type: selected

Plant Origin: native

Collection Location: Leelanau Co., MI

Plant Type: shrub

Plant Duration: perennial

Propagation: vegetative

Uses: windbreaks (especially on wet or organic soils) and wildlife habitat

'Magenta' (*Malus sp.*) Hybrid Crabapple

Released: 1990 (FY1990)

Accession Number: 9005032

Release Type: cultivar

Plant Origin: introduced

Collection Location: Clinton Co., MI

Plant Type: tree

Plant Duration: perennial

Propagation: seed

Uses: small tree for single row windbreaks & beautification

Prairie View Indiana Germplasm (*Andropogon gerardii* Vitman) Big Bluestem

Released: 2005 (FY2005)

Accession number: 9086588

Release Type: Selected

Plant Origin: Native

Collection Location: Indiana

Plant Type: warm-season grass

Plant Duration: perennial

Propagation: seed

Uses: wildlife food/cover, erosion control, increased species diversity, and native environment restoration

Prairie View Indiana Germplasm (*Schizachyrium scoparium* (Michx.)) Little Bluestem

Released: 2005 (FY2005)

Accession number: 9086577

Release Type: Selected

Plant Origin: Native

Collection Location: Indiana

Plant Type: warm-season grass

Plant Duration: perennial

Propagation: seed

Uses: wildlife food/cover, erosion control, increased species diversity, and native environment restoration

Prairie View Indiana Germplasm (*Sorghastrum nutans* (L.) Nash) Indiangrass  
Released: 2005 (FY2005)  
Accession number: 9086566  
Release Type: Selected  
Plant Origin: Native  
Collection Location: Indiana  
Plant Type: warm-season grass  
Plant Duration: perennial  
Propagation: seed  
Uses: wildlife food/cover, erosion control, increased species diversity, and native environment restoration

Riverbend Germplasm (*Salix sericea* Marsh.) Silky Willow  
Released: 2003 (FY2003)  
Accession Number: 9069052  
Release Type: tested  
Plant Origin: native  
Collection Location: Daviess County, IN  
Plant Type: shrub  
Plant Duration: perennial  
Propagation: vegetative  
Uses: streambank/shoreline restoration and riparian corridors

'Roselow' (*Malus sargentii* Rehder) Sargent's Crabapple  
Released: 1978 (FY1978)  
Accession Number: 9005026 (PI Number: 477986)  
Release Type: cultivar  
Plant Origin: introduced  
Collection Location: Japan  
Plant Type: tree  
Plant Duration: perennial  
Propagation: seed  
Uses: farm and field windbreaks

Southlow Michigan Germplasm (*Andropogon gerardii* Vitman) Big Bluestem  
Released: 2001 (FY2001)  
Accession number: 9084510 (PI Number: 642398)  
Release Type: Source Identified  
Plant Origin: Native  
Collection Location: Southern Lower Michigan  
Plant Type: warm-season grass  
Plant Duration: perennial  
Propagation: seed  
Uses: wildlife cover filter strips  
Registration Document: Durling, J.C., J.W. Leif, and D.W. Burgdorf. 2007. Registration of Southlow Michigan Big Bluestem Germplasm. *Crop Sci.* 47:455.

Southlow Michigan Germplasm (*Panicum virgatum* L.) Switchgrass  
Released: 2001 (FY2001)  
Accession number: 9084512 (PI Number: 642395)  
Release Type: Source Identified  
Plant Origin: Native  
Collection Location: Southern Lower Michigan  
Plant Type: warm-season grass  
Plant Duration: perennial  
Propagation: seed  
Uses: wildlife food/cover, erosion control, increased species diversity, and native environment restoration  
Registration Document: Durling, J.C., J.W. Leif, and D.W. Burgdorf. 2008. Registration of Southlow Michigan Germplasm Switchgrass. *Journal of Plant Registrations* 2:60.

Southlow Michigan Germplasm (*Schizachyrium scoparium* (Michx.) Nash) Little Bluestem  
Released: 2001 (FY2001)  
Accession number: 9084511 (PI Number: 642397)  
Release Type: Source Identified  
Plant Origin: Native  
Collection Location: Southern Lower Michigan  
Plant Type: warm-season grass  
Plant Duration: perennial  
Propagation: seed  
Uses: wildlife food/cover, erosion control, increased species diversity, and native environment restoration  
Registration Document: Durling, J.C., J.W. Leif, and D.W. Burgdorf. 2007. Registration of Southlow Michigan Little Bluestem Germplasm. *Journal of Plant Registrations*. 1:134.

Southlow Michigan Germplasm (*Sorghastrum nutans* L. Nash) Indiangrass  
Released: 2001 (FY2001)  
Accession number: 9084513 (PI Number: 642396)  
Release Type: Source Identified  
Plant Origin: Native  
Collection Location: Southern Lower Michigan  
Plant Type: warm-season grass  
Plant Duration: perennial  
Propagation: seed  
Uses: wildlife food/cover, erosion control, increased species diversity, and native environment restoration  
Registration Document: Durling, J.C., J.W. Leif, and D.W. Burgdorf. 2008. Registration of Southlow Michigan Germplasm Indiangrass. *Journal of Plant Registrations*. 2:56.

## **APPENDIX B**

### **CUSTOMER ASSISTANCE SUMMARY**

The Rose Lake Plant Materials Center recorded approximately 250 customer assists during FY 2007. Time spent per assist ranged from 5 minutes to 6 hours. Assistance was provided to individuals and groups. Partners and government agencies receiving assistance included:

Boy Scouts of America  
Cleveland (OH) Police Department  
Congressman Mike Rogers (R-MI)  
Ft. Custer Military Training Center  
Gun Lake, Huron, and Pokagon Band of Potawatomi Indians  
Indiana DNR  
Jasper-Pulaski Fish & Wildlife Center  
Michigan DNR  
MI Dept. of Agriculture  
Michigan Food and Farming Systems  
Michigan State University  
National Park Service  
Parks Canada  
Sleeping Bear Dunes National Lakeshore  
US Fish and Wildlife Service  
US Forest Service  
USDA-APHIS  
USDA-ARS  
WKAR Radio

The Rose Lake Center made approximately 25 presentations and created approximately 25 publications during 2007. Several presentations involved Hispanic outreach. Publications included the following:

Durling, J.C., J.W. Leif, and D.W. Burgdorf. 2007. Registration of Southlow Michigan Big Bluestem Germplasm. *Crop Sci.* 47:455.

Durling, J.C., J.W. Leif, and D.W. Burgdorf. 2007. Registration of Southlow Michigan Little Bluestem Germplasm. *Journal of Plant Registrations.* 1:134.

## APPENDIX C

### WEATHER AND CLIMATE

Table 21. Temperature and precipitation data (deviation from long-term average) for Rose Lake PMC for 2007 growing season. <sup>1</sup>					
	May	June	July	Aug	Sept
Average Daily Maximum (°F)	73.3 (+3.3)	80.9 (+0.9)	81.8 (-2.5)	80.9 (-1.5)	76.5 (+1.6)
Average Daily Minimum (°F)	44.6 (-0.4)	52.4 (-3.0)	52.2 (-7.0)	56.6 (-0.7)	48.7 (-2.0)
Precipitation (in)	3.62 (+0.62)	3.91 (+0.31)	0.80 (-2.01)	6.18 (+2.93)	1.98 (-0.76)
<sup>1</sup> Rose Lake PMC data based on records from Michigan State University Muck Research Farm in Laingsburg, MI, about 3 miles from Rose Lake PMC. Data available on-line at <a href="http://www.agweather.geo.msu.edu/mawn/">http://www.agweather.geo.msu.edu/mawn/</a> Long-term average was calculated as average from Clinton and Shiawassee Counties as recorded in respective soil surveys.					

**2007 Weather Summary** Severe drought conditions in July were preceded (May-June) and followed (August) by periods when precipitation exceeded long-term averages.

**Climate** The Rose Lake PMC is inland approximately 100 miles from Lakes Michigan, Huron, and Erie. These Great Lakes influence the climate toward modified maritime but the inland location of the PMC influences the climate toward continental. Prevailing winds are westerly. Annual precipitation is approximately 30 inches, distributed throughout the year. The average frost-free growing season is 143 days, from mid-May to early October. January and July are the coldest and hottest months, respectively.

"The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer."