



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Physics, Computer Science & Mathematics Division

RECEIVED
LAWRENCE
BERKELEY LABORATORY
JUL 21 1983
LIBRARY AND
DOCUMENTS SECTION

Submitted to Physical Review Letters


LOW FREQUENCY MEASUREMENT OF THE SPECTRUM OF
THE COSMIC BACKGROUND RADIATION

G.F. Smoot, G. De Amici, S.D. Friedman, C. Witebsky,
N. Mandolesi, R.B. Partridge, G. Sironi,
L. Danese, and G. De Zotti

June 1983

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782.*



LBL-16181
e.2

Low Frequency Measurement of the Spectrum
of the Cosmic Background Radiation

George F. Smoot, Giovanni De Amici, Scott D. Friedman, Chris Witebsky
Space Sciences Laboratory & Lawrence Berkeley Laboratory
University of California, Berkeley, California 94720

Nazzareno Mandolesi
Istituto TESRE-CNR, Bologna, Italy

R. Bruce Partridge
Haverford College, Haverford, PA 19041

Giorgio Sironi
Istituto di Fisica Cosmica-CNR, Milano, Italy

Luigi Danese, Gianfranco De Zotti
Osservatorio Astronomico, University of Padova, Padova, Italy

ABSTRACT

We have made measurements of the cosmic background radiation spectrum at 5 wavelengths (0.33, 0.9, 3, 6.3, and 12 cm) using radiometers with wavelength-scaled corrugated horn antennas having very low sidelobes. A single large-mouth (0.7 m diameter) liquid-helium-cooled absolute reference load was used for all five radiometers. The results of the observations are consistent with previous measurements and represent a significant improvement in accuracy.

PACS numbers: 98.70.Vc, 98.80.Bp

A careful measurement of the shape of the cosmic background radiation spectrum is crucial to the understanding of processes that occurred in the early universe. Several of these processes are thought to be capable of causing distortions from a blackbody distribution, particularly in the Rayleigh-Jeans region. However, previous measurements at wavelengths greater than 0.35 cm, all made before 1968, typically have errors of 15-20%¹. A significant distortion therefore may have gone undetected. At infrared wavelengths a distortion has been reported by Woody and Richards². We report here an experiment to measure, in a systematic way, the intensity of the cosmic background radiation at five wavelengths (0.33, 0.91, 3.0, 6.3, and 12 cm) and the atmospheric emission at an additional wavelength (3.2 cm). This paper describes the experiment and summarizes the results. The companion papers³⁻⁷ which follow discuss each instrument and the observations in more detail.

In the standard Big Bang model the cosmic background radiation is predicted to have a Planckian (blackbody) spectrum to very high accuracy. However, a more complete model must include perturbations in the early universe which result in the observed structure containing galaxies and clusters of galaxies. The extra energy associated with these perturbations produce distortions in the cosmic radiation spectrum, with the fractional distortion approximately equal to the ratio of the energy released to the energy in the cosmic radiation field^{8,9}.

A number of early processes could have caused a large energy release: dissipation of primeval turbulence, dissipation of sound and shock waves associated with adiabatic density perturbations or with gravity waves, isotropization of an anisotropic universe, matter-antimatter annihilation, and formation of helium and heavier elements. After the lepton era ($t > 1$ hour, corresponding to a redshift $z < 10^8$), processes which would relax a distorted spectrum to a Planckian form were, in general, so weak that the presently observed spectrum should still retain most of its original distortions, thus providing information about the initial energy release⁹.

The primary interaction between matter and radiation after the lepton era was Compton scattering, which conserves the photon number. Energy released after the lepton era would result in the cosmic background radiation assuming a Bose-Einstein distribution, with a consequent depletion of photons at low frequency compared to high frequencies¹⁰. Radiative processes would be unable to produce sufficient photons to reestablish a Planckian distribution except at wavelengths longer than 15 cm. The result would be a distortion in the Rayleigh-Jeans region. After the universe had expanded for roughly one hundred years ($z = 10^4$) there were insufficient Compton scatterings to establish a Bose-Einstein spectrum⁹. As shown by Sunyaev and Zel'dovich¹⁰, energy release at $z < 10^4$ which heats the intergalactic (or primordial) plasma would produce, by Compton scatterings, a distortion of the spectrum, depleting photons throughout the Rayleigh-Jeans region and creating an excess in the Wien region. The detection and measurement of distortions in the background radiation spectrum would provide data on important cosmological processes.

The goal of this experiment is to measure the low-frequency spectrum of the cosmic background radiation at several frequencies with small systematic errors.

The concept of the measurement is simple: compare the power received by an antenna directed upward at the sky with that received when the antenna is looking downward into an absolute reference cold load. The power received looking at the sky is the sum of power from the cosmic background radiation, galactic emission, atmospheric emission, diffracted ground emission, and miscellaneous other sources. The difference in power received (sky minus cold load) plus the known power of the absolute reference cold load is the power from the sky. By carefully accounting for all radiation entering the antenna, one can determine the power received from the cosmic background radiation.

The experimental design reduces the extraneous sources of radiation as much as is practical and then determines the residual values. Atmospheric emission is reduced by roughly a factor of three compared to sea level by going to a high, dry site (University of California's White Mountain Research Station at 3800 meters, with only 0.3 cm of precipitable H_2O). The

remaining atmospheric signal is then determined by zenith scans. The galactic background is minimized by taking data at high galactic latitudes where the emission is low and is estimated by scans and modeling. Other extraneous sources of radiation are greatly reduced through the use of low-sidelobe antennas¹¹ and ground shields.

We used an ambient-pressure liquid-helium-cooled target (Figure 1) to provide a cold load temperature near 3 K, which prevents the uncertainty in the measured gain of the instruments from contributing significantly to the error in the measured cosmic background signal. The liquid helium dewar is a large cylinder with an open mouth diameter of 70 cm. The interior radiometric walls of the cold load are made of aluminum-coated mylar. The aluminum is 13 microns thick, or seven skin depths at 12 cm wavelength. Completely covering the floor of the cold load is a 20-cm-thick circular slab of Eccosorb (Emerson and Cuming VHP-8), whose microwave emissivity is greater than 0.999 at all of our wavelengths. During operation, the Eccosorb is completely submerged in liquid helium. We maintained at all times 100-200 liters of the LHe in the cold load.

Two 18-micron-thick polyethylene windows, spaced about 15 cm apart, were installed near the top of the dewar. No other objects were between the antennas and the liquid helium. Boiled off helium gas was warmed and then passed through the space between the two windows to prevent condensation on the top window. The cold load was operated at a very slight (2mm Hg) pressure above ambient in order to maintain positive flow through the system without bursting the very thin windows.

The temperature of the LHe was 3.77 ± 0.01 K during the entire measurement period as determined by measuring the ambient air pressure. The cold load has a measured reflection coefficient less than 10^{-3} (typically 2×10^{-4}). We estimate that more than 99% of the emitted power came from the cooled Eccosorb target.

The instruments used to measure the cosmic background radiation are all differential, Dicke-switched radiometers, each composed of two wavelength-scaled corrugated-horn antennas and superheterodyne receivers. Table I summarizes the properties of the various radiometers.

All five radiometers were mounted on carts which rolled on a 20-meter-long set of rails and thus could be positioned over the cold load, which was in a hole and suspended below the middle of the rails.

In a set of measurements each radiometer in turn was positioned above the cold load and made a series of observations of the cold load and the vertical sky, together with zenith scans, gain calibrations, and related measurements. During this period (typically one hour) the other radiometers made zenith scans or galactic background measurements.

The most significant background we face is atmospheric emission; errors in determining the atmospheric signal translate directly into errors in the antenna temperature of the cosmic background radiation.

A sixth radiometer, operating at 3.2 cm (9.4 Ghz), provided automated measurements of the atmospheric emission continuously during all the cosmic background spectrum observations. The other radiometers, particularly the 0.33 and 0.91 cm instruments, were often operated to make atmospheric measurements. Atmospheric emission at 0.33 cm is particularly sensitive to water vapor and thus provided a good monitor of the atmospheric water vapor content. We have made models of atmospheric emission to tie together the observations at different frequencies. At this time the model and measurements are limited to an accuracy of about 0.1 K^5 at the three largest wavelengths, and worse at 0.91 and 0.33 cm³⁻⁷.

There are a number of other possible systematic errors. Perhaps the most important of these is the difference in signal level that may result from mechanical stresses or changing alignments when the radiometer is pointed up at the sky and then down into the cold load or during zenith scans. We have tested for these effects. The results and errors are discussed separately for each of the radiometers in the succeeding papers.

Measurements were made with liquid nitrogen in the cold load on July 4 and 7, 1982 and with liquid helium on July 5 and 6, 1982. The data were digitized and recorded both on magnetic tape and by hand. The results of the measurements are summarized in Table II. They are also plotted in Figure 2 together with the results of previous measurements. As can be seen in Figure 2 our data are in good agreement with previous results but represent an improvement in the low frequency region. This agreement is underscored when one compares the weighted average of our measurements, 2.79 ± 0.10 K, with that of the previous data at these frequencies, which have a weighted average of 2.74 ± 0.09 K¹. Our results are also in agreement with the Woody and Richards¹² infrared measurement value of 2.96 ± 0.08 K.

What conclusions can we draw from our results? First, let us suppose that the results of Woody and Richards¹² are high by ~25% because of an error in calibration, as suggested by Weiss¹. Then from 0.07 to 12 cm, the observed spectrum of the CBR is consistent with a blackbody of $T = 2.74 \pm 0.07$ K. However, if the Woody and Richards calibration is correct, their data and ours may provide evidence for a spectral distortion, with the temperature of the CBR smaller in the Rayleigh-Jeans region than near the peak. This distortion is consistent either with a Bose-Einstein spectrum with temperature (2.92 ± 0.03) K and chemical potential term $\mu = (5 \pm 3) \times 10^{-3}$ for $\Omega = 0.1$ or $\mu = (1.4 \pm 0.9) \times 10^{-2}$ for $\Omega = 1$, or with spectra produced by an early generation of stars¹³. A value of $\mu = 1.4 \times 10^{-2}$ precludes the possibility that turbulence had the major role in initiating the formation of galaxies¹⁴. If the universe contains large amounts of antimatter, the matter and antimatter must be segregated into regions larger than galactic masses; otherwise annihilation would have produced a distortion larger than we observe. We can also put a limit on the amplitude of adiabatic perturbations: $\Delta\rho/\rho < 0.1$ for $M > 10^5 M_\odot$ ¹⁰.

The succeeding papers provide a more detailed explanation of the measurements at each of the frequencies. We plan to repeat these measurements in the summer of 1983. There will be a number of improvements in the equipment to provide more complete and more accurate data. In particular, we have placed emphasis on better measurements of the atmospheric emission and on testing our atmospheric model. Another major objective is to add another radiometer capable of measurements at several additional frequencies, in order to add to the present spectral coverage.

We thank the staff of the White Mountain Research Station of the University of California, and the following individuals for their help: Alan Benner, Hal Dougherty, Bruno Falconi, John Gibson, Nancy Gusack, Enrico Mattaini, Faye Mitschang, and Stefano Silvestri.

This research was supported by NSF Grants PHY80-15694 and AST 800737, and by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098. This work was also supported by CNR Fellowship 203.2.13.

REFERENCES

1. Weiss, R. *Annual Review of Astronomy and Astrophysics* 18, 489 (1980).
2. Woody, D.P. and Richards, P.L. *Physics Review Letters* 42, 925 (1979).
3. Sironi, G. et al., companion paper (1983).
4. Mandolesi, N. et al., companion paper (1983).
5. Partridge, R.B., et al., companion paper, (1983).
6. Friedman, S.D., et al., companion paper (1983).
7. Witebsky, C., et al. companion paper (1983).
8. Chan, K.L. and Jones, B.J.T. *The Astrophysical Journal* 200, 454 (1975).
9. Danese, L. and De Zotti, G. *Riv. Nuovo Cimento* 7, 277 (1977), and *Astron. Astrophys.* 107, 39 (1982).
10. Illarionov, A.F. and Sunyaev, R.A. *Sov. Astron.* 16, 45 (1972), and Sunyaev, R.A. and Zel'dovich, Ya.B. *Ann. Rev. of Astron. and Astrophys.* 18, 537 (1982).
11. Bielli, P., Pagana, E., and Sironi, G. *Proc. ICAP, Vol. 1*, 509, (1983).
12. Woody, D.P. and Richards, P.L. *The Astrophysical Journal* 248, 18 (1982).
13. Negroponte, J., Rowan-Robinson, M., and Silk, J., *Ap. J.* 248, 38 (1981).
14. Jones, B.J.T. *Reviews of Modern Physics* 48, 1 (1976).

Table I: Radiometer Properties

Wavelength (cm)	Frequency (GHz)	Antenna			T_{RMS} K/Hz ^{1/2}
		RF Bandwidth	Beamwidth (HPBW)		
12.0	2.5	160 MHz	12.5°	0.2	
6.3	4.75	160 MHz	12.5°	0.1	
3.2	9.4	1 GHz	12.6°	0.1	
3.0	10.0	910 MHz	12.5°	0.05	
0.9	33.0	1 GHz	7.5°	0.08	
0.33	90.0	2 GHz	7.5°	0.11	

Table II: Summary of Results

Wavelength (cm)	Frequency (GHz)	T_{CBR} (K)	T_{ATM} (K)	T_{GAL} (K)	T_{GND} (K)
12.0	2.5	2.62 ± 0.25	0.95 ± 0.05	0.3	0.2
6.3	4.75	2.71 ± 0.2	1.0 ± 0.1	0.1	0.05
3.2	9.4	---	1.03 ± 0.03	0.05	0.03
3.0	10.0	2.91 ± 0.19	0.93 ± 0.16	0.03	0.01
0.9	33.0	2.87 ± 0.21	5.0 ± 0.1	0.01	0.01
0.33	90.0	2.4 ± 1.0	12.3 ± 0.8	0.01	0.01

where

T_{CBR} is the measured thermodynamic temperature of the cosmic background radiation;


T_{ATM} is the antenna temperature of the atmosphere observed vertically;

T_{GAL} is the maximum antenna temperature estimated for galactic backgrounds during our observations; and

T_{GND} is the estimated antenna temperature from the earth.

FIGURE CAPTIONS

- Figure 1 Schematic representation of the absolute reference cold load. Only two 18-micron-thick windows are between the antenna and the liquid-helium-immersed Eccosorb target.
- Figure 2 Plot of these results together with previous measurements of the temperature of the cosmic background radiation.



XBL 833-91

Fig. 1




Fig. 2