



For More Information

Visit our Plant Materials website at

<http://Plant-Materials.nrcs.usda.gov> to find more information on using plants to solve conservation problems.

USDA / NRCS

Rose Lake Plant Materials Center

7472 Stoll Road

East Lansing, MI 48823

Phone: (517) 641-6300

Fax: (517) 641-4421

Email Contacts

Plant Materials Specialist

dave.burgdorf@mi.usda.gov

PMC Manager

john.leif@mi.usda.gov



The United States Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation and marital or familiar status. (Not all prohibited bases apply to all programs). Persons with disability who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET center at (202) 720-2600 (Voice or TDD).

To file a complaint of discrimination, write USDA, Director, and Office of Civil Rights, Room 326 W. Whitten Building, 14th and Independence Avenues, SW, Washington DC 20250 or call (202) 720-5964 (Voice or TDD). USDA is an equal opportunity provider and/employer.

June, 2004

United States Department of Agriculture

NRCS Natural Resources
Conservation Service



'INDIGO' SILKY DOGWOOD



Released by
Rose Lake Plant Materials
Center

East Lansing, Michigan

'INDIGO' SILKY DOGWOOD

'Indigo' silky dogwood (*Cornus amomum* Mill.) is a dense, multi-stemmed woody plant. 'Indigo' is especially useful for single-row windbreaks in fields with center pivot irrigation systems because it grows low enough that the spray bar can pass over it.

'Indigo' can also be used in multi-row field windbreaks and farmstead windbreaks. Stems and leaves are desirable deer browse, and the fruit is eaten by upland game birds and many songbirds. Silky dogwood is also useful for beautification and landscaping.

Description

'Indigo' is a spreading shrub that can grow 10 feet tall. The stems are red when young but turn reddish brown and gray as they mature. 'Indigo' has yellowish-white flowers that bloom in mid-June. It produces an abundant crop of fruit, which ripens in September. The fruit is pale to dark blue and about 1/4 inch in diameter.

Silky dogwood can be distinguished from other blue-fruited species of dogwood by the dark brown pith in one and two year old stems. In other blue-fruited dogwoods the pith is white.



Adaptation

'Indigo' silky dogwood seed was first collected in 1961 from plants at the Rose Lake Wildlife Research Station in Clinton County, Michigan. 'Indigo' has been evaluated in 20 states, particularly in the Great Lake States, where it has performed well. Its natural range of climatic adaptation is shown on the map.

'Indigo' is adapted to a wide range of soils and soil textures, but it grows best in moist soil. It is adapted to soils that are somewhat poorly drained to well drained and have a pH of 5.0 to 7.0. It is moderately shade tolerant but is not drought tolerant.

Establishment

For help in planning windbreaks of 'Indigo', contact your local Natural Resources Conservation office.

Establish plantings of 'Indigo' from seedlings. Two year old seedlings are suitable for mechanical planting. One year old seedlings can be planted by hand, although survival may be low because of their small root systems.

Follow the planting dates and spacing recommendation for shrubs in your area; space the plants no more than six feet apart. Plant 'Indigo' seedlings in the spring as soon as the ground thaws. If you plant them much later than the average frost free date, their establishment may be poor unless a moist site is chosen or supplemental water is applied.

To ensure establishment, control weeds mechanically for the first few years. Once 'Indigo' is established, you can use approved herbicides for effective weed control. Several effective herbicides are available, be sure to follow the manufacturer's directions carefully. Consult Extension personnel for help in selecting herbicides. Silky dogwood is susceptible to infestations of scurfy scale and occasional webworm attacks. Also excessive deer browsing may retard growth.

