Hawaii, 2005

Primary and secondary syphilis

- CASES: Hawaii reported 11 cases in 2005.
- RATES: Hawaii ranked 41 among 50 states + DC + 3 territories: 0.9 cases per 100,000 population compared to the U.S. rate of 3.0 cases per 100,000 population.

Male-female rates of primary and secondary syphilis

- In Hawaii, the rate among males was 1.7 per 100,000 population compared to the U.S. male rate of 5.1 per 100,000.
- The rate among females was 0.0 per 100,000 compared to the U.S. female rate of 0.9 per 100,000.

Racial disparities of primary and secondary syphllis


In Hawaii, the race/ethnicity adjusted rates per 100,000 population were 2.1 among whites, 3.0 among blacks, 0.0 among Hispanics, 0.4 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.

• The rate among blacks was 1.4 times that of whites.

Congenital syphilis

- CASES: Hawaii reported 0 cases in 2005.
- RATES: Hawaii had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.0 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Hawaii, 1996-2005



Total P&S Syphilis Rates by County, Hawaii, 2005

