

Iowa, 2005

Primary and secondary syphilis

- CASES: Iowa reported 9 cases in 2005.
- RATES: Iowa ranked 47 among 50 states + DC + 3 territories: 0.3 cases per 100,000 population compared to the U.S. rate of 3.0 cases per 100,000 population.

Male-female rates of primary and secondary syphilis

- In Iowa, the rate among males was 0.6 per 100,000 population compared to the U.S. male rate of 5.1 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the U.S. female rate of 0.9 per 100,000.

Racial disparities of primary and secondary syphilis


In Iowa, the race/ethnicity adjusted rates per 100,000 population were 0.2 among whites, 4.1 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.

- The rate among blacks was 20.5 times that of whites.


Congenital syphilis

- CASES: Iowa reported 0 cases in 2005.
- RATES: Iowa had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.0 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Iowa, 1996-2005



Total P&S Syphilis Rates by County, Iowa, 2005



Rate (per 100,000 population)

