Oklahoma, 2005

Primary and secondary syphilis

- CASES: Oklahoma reported 44 cases in 2005.
- RATES: Oklahoma ranked 34 among 50 states + DC + 3 territories: 1.2 cases per 100,000 population compared to the U.S. rate of 3.0 cases per 100,000 population.

Male-female rates of primary and secondary syphilis

- In Oklahoma, the rate among males was 1.6 per 100,000 population compared to the U.S. male rate of 5.1 per 100,000.
- The rate among females was 1.0 per 100,000 compared to the U.S. female rate of 0.9 per 100,000.

Racial disparities of primary and secondary syphilis


In Oklahoma, the race/ethnicity adjusted rates per 100,000 population were 0.7 among whites, 8.8 among blacks, 0.0 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.3 among American Indians/Alaska Natives.

• The rate among blacks was 12.6 times that of whites.


Congenital syphilis

- CASES: Oklahoma reported 1 case in 2005.
- RATES: Oklahoma ranked 25 among 28 states + 2 territories reporting at least one congenital syphilis case: 2.0 cases per 100,000 live births compared to the U.S. rate of 8.0 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Oklahoma, 1996-2005



Total P&S Syphilis Rates by County, Oklahoma, 2005



Rate (per 100,000 population)

