# Hawaii, 2006

#### Primary and secondary syphilis

- CASES: Hawaii reported 18 cases in 2006.
- RATES: Hawaii ranked 36 among 50 states, DC, and 3 territories with 1.4 cases per 100,000 population compared to the total rate of 3.3 cases per 100,000 population.

### Rates of primary and secondary syphilis, by sex

- In Hawaii, the rate among males was 2.7 per 100,000 population compared to the total male rate of 5.7 per 100,000.
- The rate among females was 0.2 per 100,000 compared to the total female rate of 1.0 per 100,000.

# Rates of primary and secondary syphilis, by race/ethnicity


In Hawaii, the race/ethnicity adjusted rates per 100,000 population were 2.1 among whites, 8.8 among blacks, 1.0 among Hispanics, 0.9 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.

• The rate among blacks was 4.2 times that of whites.


## Congenital syphilis

- CASES: Hawaii reported 0 cases in 2006.
- RATES: Hawaii had a rate of 0.0 cases per 100,000 live births compared to the total rate of 8.7 cases per 100,000.

### Syphilis: P&S Rates and Congenital Cases, Hawaii, 1997-2006



Total P&S Syphilis Rates by County, Hawaii, 2006



Rate (per 100,000 population)

