Washington, 2006

Primary and secondary syphilis

- CASES: Washington reported 182 cases in 2006.
- RATES: Washington ranked 17 among 50 states, DC, and 3 territories with 2.9 cases per 100,000 population compared to the total rate of 3.3 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Washington, the rate among males was 5.7 per 100,000 population compared to the total male rate of 5.7 per 100,000.
- The rate among females was 0.1 per 100,000 compared to the total female rate of 1.0 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity


In Washington, the race/ethnicity adjusted rates per 100,000 population were 2.7 among whites, 9.8 among blacks, 3.4 among Hispanics, 1.3 among Asian/Pacific Islanders, and 1.0 among American Indians/Alaska Natives.

- The rate among blacks was 3.6 times that of whites.
- The rate among Hispanics was 1.3 times that of whites.

Congenital syphilis

- CASES: Washington reported 0 cases in 2006.
- RATES: Washington had a rate of 0.0 cases per 100,000 live births compared to the total rate of 8.7 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Washington, 1997-2006



Total P&S Syphilis Rates by County, Washington, 2006

