

**Table 4-2. Byssinosis: Death rates (per million population) by race and sex, U.S. residents age 15 and over, 1993–2002**

Year	White		Black		Other		
	Overall	Male	Female	Male	Female	Male	Female
<b>Crude Death Rate</b>							
1993	0.08	0.09	0.06	0.37	0.08	—	—
1994	0.08	0.11	0.07	0.18	—	—	—
1995	0.07	0.13	0.03	—	—	—	—
1996	0.04	0.05	0.06	—	—	—	—
1997	0.05	0.09	0.01	0.09	—	—	—
1998	0.03	0.06	0.01	—	—	—	—
1999	0.03	0.03	0.04	—	—	—	—
2000	0.05	0.09	0.02	—	—	—	—
2001	0.04	0.07	0.04	—	—	—	—
2002	0.04	0.05	0.04	—	—	—	—
<b>1993–2002</b>	<b>0.05</b>	<b>0.08</b>	<b>0.04</b>	<b>0.06</b>	<b>0.01</b>	—	—
<b>Age-Adjusted Death Rate</b>							
1993	0.09	0.10	0.05	0.47	0.11	—	—
1994	0.09	0.13	0.06	0.39	—	—	—
1995	0.07	0.14	0.03	—	—	—	—
1996	0.04	0.05	0.05	—	—	—	—
1997	0.05	0.11	0.01	0.18	—	—	—
1998	0.03	0.07	0.01	—	—	—	—
1999	0.03	0.03	0.03	—	—	—	—
2000	0.05	0.11	0.02	—	—	—	—
2001	0.05	0.08	0.03	—	—	—	—
2002	0.04	0.06	0.03	—	—	—	—
<b>1993–2002</b>	<b>0.05</b>	<b>0.09</b>	<b>0.03</b>	<b>0.09</b>	<b>0.01</b>	—	—

— indicates no deaths listed.

NOTE: Slight variations in numbers may exist when compared to previous WoRLD Surveillance Reports. See appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Bureau of the Census.