Investigation of Heavy Metals Cerro Grande Fire Los Alamos, New Mexico May 2000

CDC Epi-Aid 2000-40

Mitchell Wolfe¹, Josh Mott¹, Ron Voorhees⁴, C. Mack Sewell⁴, C.M. Wood³, Dan Paschal², Stephen Redd¹

- 1 -Air Pollution and Respiratory Health Branch, EHHE, NCEH, CDC
- 2 Toxicology Branch, Division of Laboratory Sciences, NCEH, CDC
- 3 Radiation Branch, EHHE, NCEH, CDC
- 4 New Mexico Department of Health





Investigation of Heavy Metals Cerro Grande Fire Los Alamos, New Mexico May 2000

CDC Epi-Aid 2000-40

Mitchell Wolfe¹, Josh Mott¹, Ron Voorhees⁴, C. Mack Sewell⁴, C.M. Wood2, Dan Paschal2, Stephen Redd1

- I Air Pullation and Respiratory Health Branck, EIRIE, NCER, CDC 7 Tournings Branck, Division of Laboratory Sciences, NCER, CDC 3 Radiation Branch, ERHE, NCER, CDC

- 4 New Mexico Department of Health





Health Effects of Forest Fires

Smoke exposure associated with:

- · increased respiratory symptoms
- · increased emergency room visits for respiratory diseases
- · increased hospitalizations for respiratory diseases (asthma, chronic obstructive pulmonary disease), heart disease

Health effects associated with:

- · Being in the area of fires
- · Elevated particulate matter







Background Cerro Grande Fire

- . May 4: Controlled burn by Nat'l Park Service begins in Bandelier National Monument adjacent to Los Alamos National Lab (LANL).
- . May 5: Declared wildland fire.
- May 10&11: 239 houses burned; 25,000 evacuated.
 - Mandatory: Los Alamos, White Rock
 - Voluntary: Española
- May 18: 100% contained, 47,650 acres
- . May 18: NMDOH invited CDC to assist:
 - Mitchell Wolfe, Josh Mott, and C.M. Wood departed May 18th









Background

- . Los Alamos National Laboratory
 - Mesas and canyons: elevation 6,200-7,800 ft.
 - 235,000 persons live within 50-mi radius
 - 27,500 acres total: fire burned 7,500 acres of LANL property
 - Fire burned part or all of 112 structures, mostly office trailers and storage sheds.



10.0

Objectives

- Assess environmental monitoring data
- Determine necessity of evaluating human exposures to potential contaminants from wildfire smoke
- If needed, conduct a study

11.3

Objective 1: Assess environmental monitoring data Rautine environmental monitoring


- · LANL: annual environmental surveillance report
 - Air, water, sediments, soils, animals, food
 - Testing for radiation, metals, PCBs, pesticides
- Results: "acceptable"/background levels of metals in most samples
- · Particulate matter (NMED, LANL, Porblus)
 - Part of fire smoke
 - Routine monitoring at various sates (Santa Fe, Taos, Bernalillo, LANL)
 - Not in Española
- · Radiation (LANL/DOE/NMED, EPA)
 - Large network of testing for gamma (real-time munitoring) plus routine
 - airborne particulate samples for gross alpha, beta, gamma or radionuclides
- Newner: http://newnet.lanl.gov/stabytot.asp
 Airnet: http://www.air-quality.lanl.gov/AirCone_CerroGrandeFire.htm



Environmental monitoring in response to the fire

erene.


1173



Environmental monitoring in response to the Cerro Grande Fire

- · Particulate Matter (NMED, EPA)
 - Additional sites and intervals in area
 - Española began May 13
 - Results: low except elevated on LANL (TA-54) May 12-13.






Environmental monitoring in response to the Cerro Grande Fire

- · Asbestos
 - NMED air/wipe samples in Los Alamos town
 - Results
 - · Air. low (10 times below occupational standards)
 - + Wipe 11 houses (including 1 school), negative

STATES

100



Environmental monitoring in response to the Cerro Grande Fire

- Radionuclides
 - Many agencies, individually and through coordinated testing
 - Results
 - Some samples contained small amounts of radioactive material
 - · Concentrations well below regulatory levels for safety
 - · Radioactive material determined to be from natural sources

MAIN

300

Environmental monitoring in response to the Cerro Grande Fire

- · Metals and chemicals (EPA)
 - 6 monitoring sites around LANL, May 12-17
 - VOCs (toluene, benzene), PAHs (pyrene), pesticides, and metals
 - Only metals in Española, May 14.
 - Results: very low VOC, PAH, and metals.

CTIL

€1.8

Assessment and Recommendations

- · Asbestos
 - No human testing recommended
- · Radiation
 - No human testing recommended
- · Metals and chemicals
 - Human testing recommended for heavy metals

17111

400

Objective 2: Determine necessity of evaluating human exposures to potential contaminants from wildfire smoke

Why did we test for metals?

- Metals would still have been present in people when we performed our testing
- Incomplete air monitoring data for metals during greatest potential for human exposure.
- Low levels of metals detected in testing during the fire and in previous routine environmental testing (before the fire).



17.5

Why did we only test some people who were exposed to smoke from the fire?

- Looked at the "warst-case scenario" people from groups most exposed to smoke were to be representatives of the exposed population to make sure we didn't miss detection of metals if they were there.
- Method of sampling was necessary because of our goal to perform testing as quickly as possible on a sample large enough to represent the population.





Objective 3: Conduct study

Research Questions:

- 1 "Was exposure to smoke from the Cerro Grande Fire associated with elevated levels of metals in people in the area of the smoke from the fire?"
- "Were metal levels detected in people high enough to have negative health effects or warrant further testing in more people?"



1

Potential human exposure

- · 1,600 firefighters
 - 1.400 (88%) during May 10-15, when most of LANL burned
- · Several hundred National Guard, City and State Police
 - Evacuations
 - Roadblocks
 - Traffic control
- · Residents of Española (pop. 9,000) and environs
 - including Tribal Lands, e.g. San Ildefonso and Santa Clara Pueblos





Screening

- · Questionnaire and urine sample to exposed and unexposed:
 - Firefighters
 - Community
 - People who were outside a great deal of the time during fire
 - National Guard, City Police, Postal, Health Department
- Also tested unexposed in case certain occupations or cities might give you higher metal levels
- Definition of "exposure":
 - Firefighters: fought fires on LANL during Cerro Grande Fire
- Community: were in Los Alamos or Española May 10 or 11




100

Screening (cont.)


- · Lab analysis
 - 16 metals based on air monitoring, previous LANL surveillance
 - Took into account diluted or concentrated urine
 - + For example, drinking a lot of water
 - Reference for expected metal levels: survey of general US population
 - Above reference: top 5% of samples in the national survey
 - If 100 people, top 5 are "above reference"
 - Recently obtained more stringent reference for some study metals





TOP











How many people were tested, and who were they?

Firefighters (92 persons tested)

Exposer tested)

Laposer tested

Laposer te

Firefighters (92 persons tested)

Exposer category Total

Exposer (heappeant (heappeant

How many people were tested, and who were they?

Research Question 1: "Was exposure to smoke from the Cerro Grande Fire associated with elevated metals"

Analysis

- Remove the effect of other things that can affect metal level other than smoke (age, gender, smoking, city)
- · Computer statistical techniques (regression analysis)

N/A/TE

103

For which metals was there a significant association between smoke exposure and metal level?

Section Section

Menal	Number of relievations of trees.	Accept take in openial prints	Arrange have be sentiment promi-
No.		**	8
(admin	6		ai

The shadow

(False)	Number of reliant observables and course	Arresta relación (species person-	fromprishe to secuped process
Mwae		44	**
Links	+3	7.000	44

MF BILL

TIE

Research Question 1: "Was exposure to smoke from the Cerro Grande Fire associated with elevated metals"

Answer: No positive association of metals with smoke exposure

- Only exception is cadmium in National Guard, but there were only 2 cadmium level above reference, and the difference in levels between exposed and unexposed people was small.
- Some negative associations, meaning higher levels in unexposed persons, which would be due to something other than smoke

er date

HE HOR

113

113

Now let's examine the number of people with metal levels above those found in most people in the general US population

Menul	Or Tana	Manufact organization in the otherwise	und allurium
Arrest	111	. 11	1
Byrise	317	11	
No. of Street	107	11.	, A.
Contractor or	817	E #	F-
Salest .	412	1	
Oktob loss	307	10	tr.
Lima	117	0	Ja .
liamen.	111	0.	
H-t-i-d-sum	(4)		
Miller	107	10	100
Limi	127	111	1
Piercete	111	11.	
Artister	100	16.	
Theire	101	- 11	
Craces	tr*	- 11	Lier
Tangent	200		

111117

700

Which metals have more than the expected number of people with levels above those found in most of the general population? (remember, these metals were not associated with smoke exposure)

May	Or Total	Property to be obstored	Sent of the Sent
Areset	137	11	- 1
Name	237	- 11	2.4
Beryline	191	- 11	1
Cadelen	127	-11	:4
اليطو (391	.00	
Chrombia	101	- 11	- 10
Comme	117	-11	- 11
hirmen	10+	.11	1
Niet horses	381		
Swiut	207	. 11	He .
Lord	21"	-11	
Platera	317		
Antonio	346		
Theiltea	217	744	
Graphen .	my	. 0	1er
Tangerra	286	10	


Was there an association between smoke exposure and nickel, chromism, cesium or granium level?

Free Parking and Section 1 Section 1

No association, so where might these levels have come from?

NE SETTE

POSTER





Cesium isotope testing

- Tested sample with highest cesium to determine cesium isotopes
- · Isotopes identified as naturally occurring
 - all Cs 133, no Cs 137

THE PERSON

777

Discussion

- · Some metal levels in people were above normal
- Of the 16 metals tested, cesium, uranium, chromium, and nickel have more than expected number of people with values above most in the general population
- No association of elevated metals in people with wildfire smoke exposure



11.15

Study issues

- · Sampling
 - needed to act quickly because in time interval from fire to testing, some metal levels may decrease
- . Urine testing
 - could test more people with single urine test than a 24-hour urine collection.
- · Classification of exposure
 - No biomarker to measure exact amount of exposure, so relied on questionnaire. Since fire smoke was so widespread, questionnaire was likely a good measure of exposure



TE

Summary

- We sought to evaluate human exposures to potential contaminants in wildfire smoke
- · Screening for heavy metals was deemed appropriate
- . We tested people from groups we expected to be most exposed
- Data don't show that metals detected in people were associated with wildfire smoke exposure
- But, we found more people than expected had metal levels of uranium, cesium, chromium, and nickel that were greater than those found in most of the general US population



Acknowledgments

New Meason Environment Depty Devid Datum SMDOH Los Alamos Fire Direct Randy Merker Cleria Johnson Santa Fe Fire Dep 3 Edith United Cathy Pawers Altuquetque For Dec't Sandra Lity Merthe Tenu Judy Knutt Senta Fe National Guard Dennis McQuillan Elica Kenney Retta Prophet Albuqueture National Grant Tabus Maria Jeffery Marques Therea L. Smith Lapubola City Police Dong Frye Bit Schmidt Santa Fe City Police CDC Dave Manning Irone Votel Labalisis Post Office Martha Marting Santa Fe Pen Office Ken Falter Albuquerque Pest Office Maria Lacers Charles Dodos Victoria Baro Ligateds Health Department Larry Neodbara Sarenne Utbern Santa Fe Health Department 47.6 WE KIND

Further Discussion

- Research Question 2: "Were metal levels detected in people high enough to cause negative health effects, or to warrant further testing in more people?"
 - Issues with specific metals
 - · possible sources of nickel, cessum, and chromium
 - Retesting?
 - Clinical follow-up?



Expected and Observed Number of Elevated Metal Values

Cerro Grande Fire

	Number	Number	Number of
Metal	Tests	elevated	observed
Arsenic	227	11	2
Barium	227	11	00
Beryllium	227	11	2
Cadmium	227	11	m
Cobalt	212	11	0
Chromium	227	11	23
Cesium	227	11	20
Mercury	227	11	ıv
Molybdenum	181	o	00
Nickel	227	11	116
Lead	227	11	0
Platinum	227	11	0
Antimony	208	10	0
Thallium	227	11	0
Uranium	227	11	105
Tungsten	208	10	0

Summary of Selected Final Regression Models

-										
	Metal	Final Model	Beta1	p2	Number of elevated values	GM ³ exposed	GM ³ unexposed	Adj ⁴ R²	12	C ₂ (6)
ard	Arsenic	Exp city cityxexp	17.6	00.	0	4.6	7.5	.21	000	.15
	Cadmium	Exp age gender	1.7	00.	7	.39	.33	.26	.00	.08
1	Initial mode	Initial model: exposure age gender smoking city occupational group cityxexp (interaction term)	smoking c	ity occup	pational group cityxe	xp (interactio	n term)			
hters	Arsenic	Exp city smoke cityxexp	-13.4	.05	ī	3.9	6'9	80.	.03	.04
	Cesium	Exp city age gender smoke cityxexp	-7.5	.03	7	3.9	4.5	.22	00.	.04

Nickel Exp city gender 1.1 .89 48 5.8 3.8 .08	Chromium Exp gender smoke -1.1 .27 7 .16 .20 .68	Initial model: exposure age gender smoking city occupational group cityxexp (interaction term)	Uranium Exp gender age 4.3 .45 24 .03 .02 .12 .05			Cesium Exp age -1.3 .06 5 3.6 5.5 .18 .00	Metal Final Model Beta ¹ P ² elevated values exposed unexposed R ² F ⁵	.24 04 06 .10 .01	.00 .01 .01 .05 .03 .03	.18 .17 .13 .08	GM3 unexposed 5.5 .19 .02 .02 .20 .3.8	GM3 exposed 3.6 .16 .03 xp (interactio	Number of elevated values 5 6 27 24 7 7 7 7 7 7 7 8 8 8 8 8 8 8 9 9 9 9 9 9	.06 .06 .08 .45 .ty occup	-1.3 1.0 1.7 4.3 smoking ci	Exp age gender Exp age gender Exp age Exp age City cityxexp City cityxexp Exp gender age City cityxexp Exp gender smoke Exp gender smoke	Metal Cesium Chromium Nickel Uranium Initial model
	Nickel Exp city gender 1.1 .89 48 5.8 3.8 .08	Exp gender smoke -1.1 .27 7 .16 .20 .68 Exp city gender 1.1 .89 48 5.8 3.8 .08	Initial model: exposure age gender smoking city occupational group cityxexp (interaction term) Chromium Exp gender smoke -1.1 .27 7 .16 .20 .68 Nickel Exp city gender 1.1 .89 48 5.8 3.8 .08	Uranium Exp gender age city cityxexp 4.3 .45 24 .03 .02 .12 Initial model: exposure age gender smoking city occupational group cityxexp (interaction term) Chromium Exp gender smoke -1.1 .27 7 .16 .20 .68 Nickel Exp city gender 1.1 .89 48 5.8 3.8 .08	Birckel Exp age 1.7 .08 27 6.4 2.8 .13 Uranium Exp gender age city cityxexp 4.3 .45 24 .03 .02 .12 Initial model: exposure age gender smoking city occupational group cityxexp (interaction term) 7 .16 .20 .68 Chromium Exp gender smoke -1.1 .27 7 .16 .20 .68 Nickel Exp city gender 1.1 .89 48 5.8 3.8 .08	Chromium Exp age gender 1.0 .77 6 .16 .19 .17 Nickel Exp age 1.7 .08 27 6.4 2.8 .13 Uranium Exp gender age city cityxexp 4.3 .45 24 .03 .02 .12 Initial model: exposure age gender smoking city occupational group cityxexp (interaction term) 7 .16 .20 .68 Nickel Exp city gender 1.1 .89 48 5.8 3.8 .08	Cesium Exp age -1.3 .06 5 3.6 5.5 .18 Chromium Exp age gender 1.0 .77 6 .16 .19 .17 Uranium Exp gender age 4.3 .45 24 .03 .02 .12 Initial model: exposure age gender smoking city occupational group cityxexp (interaction term) .11 .27 7 .16 .20 .68 Nickel Exp city gender 1.1 .89 48 5.8 3.8 .08	_	32	00	18	0.0	24	32		Fxn	Ilradium

Theta for exposure variable

2-value for exposure variable

3GM = geometric mean

4djusted R-squared of finat model

F = p-value for ANOVA F-test for entire model

6C-squared = part correlation of "exposure variable (i.e. the increase in R due to "exposure" value)

Occupational Group by City and Exposure Cerro Grande Fire

	Exposure	Exposure category	i
Occupational Group	Exposed (n=83)	Unexposed (n=52)	Total
National Guard			
Albuquerque	14	10	24
Santa Fe	24	4	28
Police			
Española	23	0	23
Santa Fe	7	19	21
Postal workers			
Española	o	0	0
Albuquerque	0	2	7
Santa Fe	1	7	00
Health department			
Española	10	0	10
Santa Fe	0	10	10
Firefighters			
Los Alamos	42	0	42
Santa FEW	20	-	21
Other (mainly Albuq.)	4	25	29

Naturally-occurring Uranium Concentrations, USA, 1993



Uring Uranium Tsotonic Analysis, Cerro Grande Fire (n=9)

or amunit assume Abundance		ricasulements	sales and deposits a ad an added	and the second second		
	Analysis Date: 6-26-00		Instrument ELAN 6000 Quadrupole ICPMS	Quadrupole ICP	MS	
A see	And the second second second	441.41.91	U-234	U-235	U-236	U-238
Natural Uranium Isotope Ab	n Isotope Abundance is	ce is	0.0055%	0.72%	N.	99.27%
		the order or the Co.		Observed Isotope	Observed Isotope Abundances (%)	39 39
1	Observed U Concer	Concentration (ppb)	U-234	U-235	U-236	U-238
Samples from Los Alamos Fi	os Alamos Fire Stu	re Study (case # 00-0033)	33)			
9200	2.3 ppb	and the same	0.00	0.76	-0.01%	99.21%
4		std dev	0,00	0.12%	0.05%	0.14%
0090	1.4 ppb		0.00	0.82%	-0.02%	99.21%
10000	100	std dev	0.00	0.13%	0.09%	0.21%
0095	0.8 ppb	die de	0.00	0.85%	-0404%	99.13%
- Constitution	A Section of	std dev	0.00	0.22%	0.10%	0.32%
0100	0.9ppb		00.00	0.80%	0.01%	99.16%
	0.000	std dev	0.00	0.11%	0.13%	0.205
0103	1.0 ppb	2	0.00	0.77%	0.04%	99.15%
		std dev	0.00	0.24%	0.09%	0.21%
0.104	3.4 ppb		00.0	0.72%	-0.01%	99.28%
		std dev	0.00	0.07%	0.02%	0.08%
0.111	0.8 ppb		0.00	0.77%	0.01%	99.25%
The state of		std dev	0.00	0.27%	0.23%	0.33%
1000	1.4 ppb		0.00	0.80%	-0.01%	99.21%
1000	The second second	std dev	00.00	0.13%	0.05%	0.12%
6000	0.7 ppb	Second Co.	0.00	0.83%	-0.05%	99.21%
		std dev	0.00	0.35%	0.05%	0.04%

Investigation of Heavy Metals Cerro Grande Fire Los Alamos, New Mexico May 2000

CDC Epi-Aid 2000-40

Mitchell Wolfe¹, Josh Mott¹, Ron Voorhees⁴, C. Mack Sewell⁴, C.M. Wood³, Dan Paschal², Stephen Redd¹

- 1-Air Pollution and Respiratory Health Branch, EHHE, NCEH, CDC
- 2 Toxicology Branch, Division of Laboratory Sciences, NCEH, CDC
- 3 Radiation Branch, EHHE, NCEH, CDC
- 4 New Mexico Department of Health

Background

- May 4: Controlled burn by National Park Service begins in Bandelier National Monument adjacent to Los Alamos National Lab (LANL).
 - May 5: Declared wildland fire.
 - May 10&11: 239 houses burned; 25,000 evacuated.
 - Mandatory: Los Alamos, White Rock
 - Voluntary: Española
- May 18: 100% contained, 47,650 acres
- May 18: NMDOH invited CDC to assist:
- Mitchell Wolfe, Josh Mott, and C.M. Wood departed May 18th

Los Alamos National Laboratory

- Mesas and canyons: elevation 6,200-7,800 ft.
- 27,500 acres total: fire burned 7,500 acres of LANL property
- Fire burned part or all of 112 structures

Objectives

Assess environmental monitoring data

Determine necessity of evaluating human exposures to potential contaminants from wildfire smoke

If needed, conduct a study

Objective 1: Assess environmental monitoring data Routine environmental monitoring

LANL: annual environmental surveillance report

- Air, water, sediments, soils, animals, food
- Testing for radiation, metals, poly chlorinated biphenyls (PCBs), pesticides
- Results: "acceptable"/background levels of metals in most samples

Particulate matter (New Mexico Environmental Department—NMED, LANL, Pueblos)

- Part of fire smoke
- Routine monitoring at various sites (Santa Fe, Taos, Bernalillo, LANL)
- Not in Española

Radiation (LANL/DOE/NMED, Environmental Protection Agency-EPA)

- Large network of testing for gamma (real-time monitoring) plus routine airborne particulate samples for gross alpha, beta, gamma or radionuclides
- Newnet: http://newnet.lanl.gov/stabyloc.asp
- Airnet: http://www.air-quality.lanl.gov/AirConc_CerroGrandeFire.htm

Environmental monitoring in response to the fire

Particulate Matter (NMED, EPA)

- Additional sites and intervals in area
- Española began May 13
- Results: low except elevated on LANL (TA-54) May 12-13.

Asbestos control

Environmental monitoring in response to the Cerro Grande Fire

Asbestos

- NMED air/wipe samples in Los Alamos town
- Results
 - Air: low (10 times below occupational standards)
 - Wipe: 11 houses (including 1 school), negative

Radiation and metal monitoring, Los Alamos area Environmental monitoring in response to the Radionuclides

- Many agencies, individually and through coordinated testing

- Results:
 - · Some samples contained small amounts of radioactive material
 - · Concentrations well below regulatory levels for safety
 - · Radioactive material determined to be from natural sources

Environmental monitoring in response to the Metals and chemicals (EPA)

- 6 monitoring sites around LANL, May 12-17
- Volatile Organic Compounds--VOCs (toluene, benzene), Polyaromatic Hydrocarbons--PAHs (pyrene), pesticides, and metals
- Only metals in Española, May 14.
- Results: very low VOC, PAH, and metals

Assessment and Recommendations

Asbestos

- No human testing recommended

Radiation

- No human testing recommended

Metals and chemicals

- Human testing recommended for heavy metals

Objective 2: Determine necessity of evaluating human exposures to potential contaminants from wildfire smoke

Why did we test for metals?

- · Metals would still have been present in people when we performed our testing.
- Incomplete air monitoring data for metals during greatest potential for human exposure.
- Low levels of metals detected in testing during the fire and in previous routine environmental testing (before the fire).

Why did we only test some people who were exposed to smoke from the fire?

- Looked at the "worst-case scenario" people from groups most exposed to smoke were to be representatives of the exposed population to make sure we didn't miss detection of metals if they were there.
- Method of sampling was necessary because of our goal to perform testing as quickly as possible on a sample large enough to represent the population.

Objective 3: Conduct study

Research Questions:

- 1. "Was exposure to smoke from the Cerro Grande Fire associated with elevated levels of metals in people in the area of the smoke from the fire?"
- 2. "Were metal levels detected in people high enough to have negative health effects or warrant further testing in more people?"

Potential human exposure

1,600 firefighters

- 1,400 (88%) during May 10-15, when most of LANL burned
- Several hundred National Guard, City and State Police
- Evacuations
- Roadblocks
- Traffic control

Residents of Española (pop. 9,000) and environs

- including Tribal Lands, e.g. San Ildefonso and Santa Clara Pueblos

Screening

Questionnaire and urine sample to exposed and unexposed:

- Firefighters
- Community
 - People who were outside a great deal of the time during fire
 - National Guard, City Police, Postal, Health Department
- Also tested unexposed in case certain occupations or cities might give you higher metal levels

Occupational Group by City and Exposure Cerro Grande Fire

	Exposure	e Category	
Occupational Group	Exposed n=83	Unexposed n=52	Total
National Guard			
Albuquerque	14	10	24
Santa Fe	24	4	28
Police			
Espanola	23	0	23
Santa Fe	2	19	21
Postal Workers			
Espanola	9	0	9
Albuquerque	0	2	2
Santa Fe	1	7	8
Health Department			
Espanola	10	0	10
Santa Fe	0	10	10
Fire Fighters			
Los Alamos	42	0	42
Santa Fe	20	1	21
Other (Mainly Albuquerque)	4	25	29

` Definition of "exposure":

- Firefighters: fought fires on LANL during Cerro Grande Fire
- Community: were in Los Alamos or Española May 10 or 11

Lab analysis

- 16 metals based on air monitoring, previous LANL surveillance
- Took into account diluted or concentrated urine
 - · For example, drinking a lot of water

What does "above reference" mean?

- Reference for expected metal levels: survey of general US population
- Above reference: top 5% of samples in the national survey
 - If 100 people, top 5 are "above reference"
 - Recently obtained more stringent reference for some study metals

Research Question 1: "Was exposure to smoke from the Cerro Grande Fire associated with elevated metals"

Answer: No positive association of metals with smoke exposure

- Only exception is cadmium in National Guard, but there were only 2 cadmium level above reference, and the difference in levels between exposed and unexposed people was small
- Some negative associations, meaning higher levels in unexposed persons, which would be due to something *other than* smoke.

Now let's examine the number of people with metal levels above those found in most people in the general US population.

Which metals have more than the expected number of people with levels above those found in most of the general population?

(remember, these metals were not associated with smoke exposure)

Naturally-occurring uranium concentrations, USA High levels previously shown in Northern New Mexico water

Cesium isotope testing

- Tested sample with highest cesium to determine cesium isotopes Isotopes identified as naturally occurring
 - all Cs 133 (naturally occurring)
 - no Cs 137

Discussion

- Some metal levels in people were above normal
- Of the 16 metals tested, cesium, uranium, chromium, and nickel have more than expected number of people with values above most in the general population
- No association of elevated metals in people with wildfire smoke exposure

Study issues

Sampling

 needed to act quickly because in time interval from fire to testing, some metal levels may decrease

Urine testing

- could test more people with single urine test than a 24-hour urine collection

Classification of exposure

No biomarker to measure exact amount of exposure, so relied on questionnaire.
 Since fire smoke was so widespread, questionnaire was likely a good measure of exposure

Summary

- We sought to evaluate human exposures to potential contaminants in wildfire smoke Screening for heavy metals was deemed appropriate
- We tested people from groups we expected to be most exposed
- Data don't show that metals detected in people were associated with wildfire smoke exposure
- But, we found more people than expected had metal levels of uranium, cesium, chromium, and nickel that were greater than those found in most of the general US population

Further Discussion

Research Question 2: "Were metal levels detected in people high enough to cause negative health effects, or to warrant further testing in more people?"

- Issues with specific metals
 - · possible sources of nickel, cesium, and chromium
- Re-testing?
- Clinical follow-up?