

Lethality of Commercial Beef Jerky Manufacturing Processes

1 Lethality of Commercial Whole-Muscle Beef Jerky Manufacturing Processes

2 Against *Salmonella* serovars and *Escherichia coli* O157:H7

3

4 Dennis R. Buege<sup>1</sup>, Gina Searls<sup>1</sup>, and Steven C. Ingham<sup>2\*</sup>

5

6 University of Wisconsin-Madison

7

8

9

10 <sup>1</sup>Department of Animal Sciences, <sup>2</sup>Department of Food Science

11

12 Short Title: Lethality of Commercial Beef Jerky Manufacturing Processes

13 Key Words: beef jerky, *Salmonella*, *Escherichia coli* O157:H7

14

15

16

17 \*Corresponding author

18 University of Wisconsin-Madison, Department of Food Science

19 1605 Linden Drive

20 Madison, WI 53706

21 Phone 608-265-4801, fax 608-262-6872, e-mail [scingham@wisc.edu](mailto:scingham@wisc.edu)

## Lethality of Commercial Beef Jerky Manufacturing Processes

### 1 ABSTRACT

2 Thermal processing used in making whole-muscle beef jerky also involves drying. This  
3 drying may cause enhanced pathogen thermotolerance and evaporative cooling that  
4 reduce process lethality. Several salmonellosis outbreaks have been associated with beef  
5 jerky. In this study, a standardized process was used to inoculate beef strips with 5-strain  
6 cocktails of either *Salmonella* serovars or *Escherichia coli* O157:H7, marinate the strips  
7 in pH 5.3 marinade for 22-24 h at 5°C, and then convert the strips to jerky using various  
8 heating/drying regimes. Numbers of surviving organisms were determined during and  
9 after the heating/drying. In some trials, a commercial lactic acid starter culture was also  
10 evaluated as a potential surrogate for the pathogens. The 5-log *Salmonella* reduction  
11 mandated by the United States Department of Agriculture (USDA), along with a 5-log  
12 reduction in *E. coli* O157:H7, was best achieved by ensuring that high wet-bulb  
13 temperatures were reached and maintained early in the process (51.7°C or 54.4°C for 60  
14 minutes, 57.2°C for 30 minutes, or 60°C for 10 minutes) followed by drying at 76.7°C  
15 (dry-bulb temperature). Processes that met the USDA guideline with smaller safety  
16 margins were 1) heating and drying at 76.7°C (dry-bulb) within 90 minutes of beginning  
17 the process, 2) heating for successive hourly intervals at 48.9, 54.4, 60, and 76.7°C (dry-  
18 bulb), or 3) heating at 51.7°C (dry-bulb), followed by 76.7°C (dry-bulb) drying started  
19 before product  $a_w$  was  $< 0.86$ . Achieving a  $\geq 3.0$  log reduction in the starter culture is a  
20 possible standard for validating process lethality.

1

2 Beef jerky processing, using whole muscle or restructured ground meat, is unique  
3 compared to the processing of other ready-to-eat meat products because heat processing  
4 is intended to attain considerable drying and desired texture and shelf-stability. This  
5 drying may reduce the process lethality against pathogenic bacteria in beef, and outbreaks  
6 of salmonellosis have been linked to the consumption of beef jerky (6). Previous  
7 research has suggested that sub-lethal drying conditions may lead to increased heat-  
8 resistance in pathogens such as *Salmonella* serovars (7). Furthermore, evaporative  
9 cooling during drying may lessen the effective temperature to which pathogens are  
10 exposed. A possible decrease in lethality related to evaporative cooling on the surface of  
11 cooked beef was noted previously in studies of *Salmonella* spp. survival during cooking  
12 of beef (2, 8). In fact, Blankenship et al., (1) recommended introduction of steam into the  
13 oven when cooking beef roasts to ensure that adequate lethality against salmonellae was  
14 attained on the roast surface. Evaporative cooling was also noted as a factor contributing  
15 to insufficient thermal lethality in making jerky associated with an outbreak of  
16 salmonellosis in New Mexico in 2003 (11). All of these factors have resulted in  
17 heightened scrutiny of the lethality of the heating and drying steps typically used during  
18 jerky processing.

19 United States Department of Agriculture (USDA) officials have issued a  
20 compliance guideline for jerky processors (14) that stresses the importance of  
21 maintaining high humidity during thermal processing in order to ensure sufficient  
22 destruction of *Salmonella* spp. and *Escherichia coli* O157:H7. However, processors have  
23 had difficulty either complying with USDA guidance or developing and validating

## Lethality of Commercial Beef Jerky Manufacturing Processes

1 adequate alternative processes while still obtaining desired finished product  
2 characteristics. Previously suggested techniques for ensuring adequate lethality included  
3 boiling beef strips in marinade prior to cooking and drying, and oven-heating beef jerky  
4 strips after drying (9, 10). Commercial processors have not widely adopted these  
5 methods, either because of perceived adverse effects on product sensory characteristics or  
6 because of economic or efficiency concerns. Development of validated heating/drying  
7 guidelines for processors of whole-muscle jerky has been further complicated by  
8 variables such as thickness of jerky strips, whether the strips have been marinated and, if  
9 so, the composition and conditions of marination, the type of smokehouse or oven used  
10 for heating/drying, and the weather and altitude at the processing plant.

11 Presently, the USDA has indicated that a jerky-making process has sufficient  
12 lethality if it results in a 5-log reduction of *Salmonella* spp. (written communication from  
13 Dr. Paul Uhler, USDA – FSIS, 2005). Furthermore, USDA officials have stated that the  
14 lethality of non-thermal steps in jerky-making, such as marination, can be counted toward  
15 meeting the overall process lethality requirement (Dan Englejohn, USDA – FSIS,  
16 personal communication, 2005). Pre-existing USDA guidance for certain other beef  
17 products specified a 6.5 log reduction in *Salmonella* spp. (15). The newer 5-log  
18 reduction standard resulted from a USDA draft risk assessment of the impact of different  
19 lethality standards for ready-to-eat meat and poultry products on the incidence of  
20 salmonellosis (13). This risk assessment determined that decreasing the pathogen  
21 reduction standard from 6.5 logs to 5 logs for products such as jerky that do not support  
22 the growth of salmonellae, would have little effect on the incidence of salmonellosis.

## Lethality of Commercial Beef Jerky Manufacturing Processes

1           The first objective of this study was to develop and validate sufficiently lethal  
2 processes for use by commercial jerky manufacturers in heating and drying whole-muscle  
3 beef jerky. In working towards this objective, marination and beef strip thickness were  
4 standardized based on typical industry practice. A standardized smokehouse loading  
5 procedure was used, and ambient weather conditions were noted.

6           A second objective of this work was to develop a simple method for commercial  
7 processors of whole-muscle beef jerky to evaluate the lethality of new processes.  
8 Because in-plant challenge studies involving pathogenic bacteria are not recommended  
9 for commercial meat processing establishments for safety reasons, and because  
10 laboratory-based challenge studies are neither practical nor affordable for most  
11 processors, we investigated the use of a commercially available lactic acid bacterial  
12 starter culture as a surrogate for *Salmonella* serovars and *E. coli* O157:H7.

13

### 14 MATERIALS AND METHODS

15           **Preparation of inoculum.** Five-strain cocktails of *Salmonella* serovars and *E.*  
16 *coli* O157:H7 were used to inoculate beef strips prior to jerky processing. *E. coli*  
17 O157:H7 strains ATCC 43894, 51657, and 51658 were clinical isolates and strain ATCC  
18 43895 was originally from ground beef implicated in an outbreak of food-borne illness;  
19 each of these strains was obtained from the American Type Culture Collection  
20 (Manassas, VA). Strain USDA-FSIS-380-94 was originally from salami implicated in an  
21 illness outbreak and was obtained from the laboratory of Dr. John Luchansky at the Food  
22 Research Institute, University of Wisconsin-Madison. The *Salmonella* serovars were all  
23 obtained from the laboratory of Dr. Eric Johnson at the Food Research Institute,

## Lethality of Commercial Beef Jerky Manufacturing Processes

1 University of Wisconsin-Madison. To obtain a working culture, each strain was cultured  
2 twice successively (from a previously frozen culture) at 35°C for 18-24 hours in Brain  
3 Heart Infusion Broth (BHIB; Difco, Becton-Dickenson, Sparks, MD), streaked to Brain  
4 Heart Infusion Agar (BHIA; Difco), incubated at 35°C for 18-24 hours, examined for  
5 purity, and then stored at 5°C. According to the work of Calicioglu et al. (3 - 5), acid-  
6 adaptation is unlikely to increase pathogen resistance to hurdles involved in beef jerky  
7 processing. Therefore, cultures for inoculation were grown in Brain Heart Infusion broth  
8 (BHIB; Difco, Becton Dickinson, Sparks, MD), a medium containing only a small  
9 amount of glucose that could be metabolized to produce organic acids. To achieve  
10 stationary-phase inoculum cultures, an isolated colony of each strain was transferred from  
11 its working culture plate to 9 mL of BHIB, and incubated at 35°C for 24 hours. To  
12 prepare a 5-strain inoculum cocktail of *Salmonella* serovars or *E. coli* O157:H7, the  
13 BHIB culture of each strain was combined into one 50-ml sterile plastic centrifuge tube,  
14 and centrifuged for 12 minutes at 5,000 x g. The supernatant in each tube was decanted  
15 and the pellets were re-suspended with approximately 20 ml of Butterfield's phosphate  
16 diluent (BPD; Nelson Jameson, Marshfield, WI). A commercial lactic acid bacteria  
17 starter culture, intended for making fermented meat products, was evaluated in several  
18 trials as a surrogate for the pathogens. The starter culture (Formula 100; Trumark,  
19 Linden, NJ) was stored at -20°C. Three different lots of this starter culture were tested  
20 during the study. Preliminary experiments showed that the starter culture was  
21 considerably more thermotolerant than the *Salmonella* and *E. coli* O157:H7 strains used,  
22 so a lower inoculum level was used for it. To prepare the starter culture for inoculation

## Lethality of Commercial Beef Jerky Manufacturing Processes

1 of beef strips, 0.5 g of culture was added to 99 ml BPD, mixed well, and then diluted  
2 another 100-fold in BPD.

3       **Inoculation of beef strips.** Frozen vacuum-packaged beef strips were obtained  
4 from a commercial jerky processor and thawed at 4°C or under running tap water prior to  
5 inoculation. The individual beef strips (5 – 7 mm thick) were placed in a biosafety hood  
6 on aluminum foil that had been previously sanitized with 70% (v/v) ethanol. To  
7 inoculate each strip, 0.4 ml of the undiluted pathogen cocktail or 0.4 ml of the diluted  
8 starter culture was pipetted onto the product surface and distributed as evenly as possible  
9 using a sterile plastic spreader. Aluminum foil was placed over the strips in a tented  
10 manner to minimize the amount of drying during microbial attachment (30 min), after  
11 which strips were turned over and the inoculation/attachment process was repeated on the  
12 other side. For each trial, one group of 9 - 12 beef strips was inoculated with *Salmonella*  
13 spp., another group of 9 - 12 beef strips was inoculated with *E. coli* O157:H7, and a third  
14 group of 4 - 6 uninoculated beef strips was used to monitor yield and water activity  
15 throughout thermal processing. In many trials, four additional strips were inoculated with  
16 the starter culture. Initial pathogen levels on inoculated beef strips were approximately  
17 10<sup>8</sup> CFU per beef strip and initial starter culture levels were about 10<sup>4</sup> CFU per beef strip.

18       **Jerky processing.** Each group of beef strips was tumbled manually in a closed  
19 zip-lock plastic bag for approximately 5 minutes in a non-acidic (pH 5.3) spice-  
20 containing marinade applied at a level of 15% (w/w; 9.7% water and 5.3% dry  
21 ingredients) intended to result in a pre-processing level of 2% (w/w) sodium chloride, 2%  
22 (w/w) sucrose, and 156 ppm sodium nitrite (w/w) in the meat. Following marination,  
23 strips were held for 22-24 h at 5°C. The next day, strips were arranged on racks placed in

## Lethality of Commercial Beef Jerky Manufacturing Processes

1 the center of a commercial one-truck smokehouse (Model TR2, Vortron, Beloit, WI) for  
2 processing. Pans of water were placed on the lowest rack in the smokehouse and a low  
3 fan speed was used to simulate as much as possible a drying rate consistent with a  
4 smokehouse containing several racks filled with product. The smokehouse dry-bulb and  
5 wet-bulb temperatures were monitored using thermocouples (L#113-1055 P/M,  
6 ThermoWorks, Alpine, UT) and a data logger (Model 92000-00, Barnant Co.,  
7 Barrington, IL). Percent relative humidity (%RH) was calculated from the wet-bulb and  
8 dry-bulb temperatures using a slide rule (Alkar, Lodi, WI). In all trials, the product-  
9 internal temperature was measured by inserting a thermocouple probe into the geometric  
10 center of a beef strip. Because insertion of the probe in this location is relatively difficult,  
11 a surrogate product-internal temperature was also obtained by tightly folding a beef strip  
12 once over a thermocouple probe in the majority of the trials. The latter temperature  
13 measurement method is considerably easier, but it was not known at first whether it could  
14 be considered an accurate surrogate for internal beef strip temperature. Smoke was not  
15 applied to the beef strips during processing. Several types of heating/drying processes  
16 were tested (summarized in Table 1). In Type 1-A processes, the dry-bulb temperature  
17 controller was set at 62.8°C (145°F) in the first 15 minutes and then at 76.7°C (170°F)  
18 during the next 15 minutes, with no added humidity. This two-step increase in dry-bulb  
19 temperature was done to simulate the beginning stages of heating a full smokehouse of  
20 moist, ambient-temperature beef strips. Next, humidity (steam or water) was introduced  
21 into the smokehouse via the wet-bulb temperature controller to obtain targeted increases  
22 in wet-bulb temperatures, referred to as “wet-bulb spikes”. The process lethality was  
23 determined for a series of trials conducted using early-process wet-bulb spikes of 51.7°C



## Lethality of Commercial Beef Jerky Manufacturing Processes

1 (125°F) for 60 min, 54.4°C (130°F) for 60 min., 57.2°C (135°F) for 30 min., and 60°C  
2 (140°F) for 10 min. Following completion of wet-bulb spikes, no further humidity was  
3 introduced into the smokehouse chamber as the product was further dried at a dry-bulb  
4 temperature of 76.7°C (170°F). To investigate the possible protective or lethal effects of  
5 marinade ingredients, a selected Type 1 process (wet bulb spike of 54.4°C for 60  
6 minutes) was done to products that were marinated either only in water (9.7% initial  
7 product weight gain) or only in the dry ingredients (5.3% initial product weight gain). In  
8 four trials, a Type 1-B process was used (wet-bulb spike of 54.4°C for 60 minutes) in  
9 which the dry-bulb temperature was held at either 65.5°C (150°F) or 87.8°C (190°F)  
10 throughout a 15-minute equilibration period before the wet-bulb spike, the wet-bulb spike  
11 itself, and final drying. Type 2 and Type 3 processes involved rapid (15 minutes) and  
12 slow (90 minutes) increases in dry-bulb temperature to 76.7 °C (170°F) followed by final  
13 drying at a dry-bulb temperature of 76.7°C (170°F). In Type 4 processes, the dry-bulb  
14 temperature was held constant at 51.7°C (125°F) until a desired approximate jerky water  
15 activity was attained, whereupon the dry-bulb temperature was increased to 76.7°C  
16 (170°F) for final drying (no humidity added during the process). In Type 5 processes, the  
17 dry-bulb temperature was held constant at 60, 71.1, or 82.2°C (140, 160, or 180°F) and  
18 no attempt was made to control wet-bulb temperature. Type 6 and 7 processes both  
19 involved sequential 1 hour exposures to dry-bulb temperatures of 48.9, 54.4, and 60°C  
20 (120, 130, and 140°F); in the Type 7 process beef strips were exposed an additional hour  
21 to a dry-bulb temperature of 76.7°C (170°F). In addition to trials in the commercial-scale  
22 smokehouse, a consumer-scale smokehouse (Pragotrade Model TS160, Cabela's, Sidney,  
23 NE) was used for additional trials testing the relationship between thermally induced

## Lethality of Commercial Beef Jerky Manufacturing Processes

1 death of the starter culture surrogate and inoculated pathogens (“consumer-type  
2 process”). Inoculated and uninoculated beef strips were placed on two racks in this  
3 smokehouse, two large (15 cm diameter) Petri dishes containing water were placed on the  
4 floor of the smokehouse, and the dry-bulb temperature was set at either 60 or 70.1°C (140  
5 or 160°F). These trials were done to compare pathogen and starter culture surrogate  
6 survival over a broad range of conditions. In all trials, uninoculated strips were evaluated  
7 at the intermediate sampling point for water activity (measured on-site using an AquaLab  
8 Series 3TE water activity meter, Decagon Devices, Inc., Pullman, WA). In all trials,  
9 additional uninoculated finished beef jerky strips were sent to a commercial testing  
10 laboratory for pH, water activity, % water (forced air oven method, AOAC method  
11 950.46Bb), % protein (Kjeldahl method, AOAC method 991.20.I), and % salt  
12 (potentiometric method, AOAC method 980.25). From these analyses, Moisture: Protein  
13 ratio (MPR), and % water-phase salt values were calculated.

14

15 **Enumeration of inoculum organisms.** The numbers of *Salmonella* spp. and *E.*  
16 *coli* O157:H7 on beef strips were determined prior to marination, after marination (in 21  
17 trials), after the early-process wet-bulb spike in the smokehouse (Type 1 process) or some  
18 other intermediate time (other process types), and following drying when the beef strips  
19 had reached a yield level predetermined to correlate with an average water activity of  $\leq$   
20 0.90. One beef strip comprised a sample and three samples were analyzed for pathogen  
21 numbers at each sampling time. In several trials, two strips each were analyzed for  
22 starter culture numbers after inoculation and at the end of drying. Each sample was  
23 placed in a whirl pak filter bag (Nasco, Fort Atkinson, WI), BPD (99 ml) was added, and

## Lethality of Commercial Beef Jerky Manufacturing Processes

1 the bag contents were stomached for 2 minutes at medium speed (Stomacher 400  
2 Circulator lab blender; Seward) or samples were manually massaged for 1 minute and  
3 shaken for 1 minute. This initial dilution was arbitrarily defined as  $10^{-1}$ . Serial decimal  
4 dilutions were made in BPD as needed. From the initial dilution, 1.0 ml was distributed  
5 for spread-plating among three plates of BHIA. From the original dilution and each  
6 subsequent dilution 0.1 ml was spread on one BHIA plate per dilution. Plates were  
7 incubated at 35°C for 1 h to allow for repair of injured cells, and then overlaid with  
8 MacConkey Sorbitol agar (SMAC; Difco), XLD agar (Difco), or mEnterococcus agar  
9 (mE; Difco) for selective/differential enumeration of *E. coli* O157:H7, *Salmonella*  
10 serovars, and starter culture surrogate, respectively. After 20-24 h (*E. coli* O157:H7,  
11 *Salmonella serovars*) or 72 h (starter culture surrogate) incubation at 35°C, plates were  
12 examined for typical colonies. For each sampling time, one presumptive colony each of  
13 *E. coli* O157:H7 and *Salmonella* was transferred to BHIA, incubated at 35°C for 20-24 h,  
14 and then tested to confirm colony identity. A single plate containing presumptive starter  
15 culture colonies was retained for confirmation tests at each sampling time. Confirmation  
16 tests for the presumptive pathogens were Gram reaction, cellular morphology, oxidase  
17 activity, and biochemical characteristics (API 20E kit, bioMerieux, Hazelwood, MO) for  
18 the pathogens, with an additional O157 latex agglutination test (Oxoid, Ogdensburg, NY)  
19 done to confirm *E. coli* O157:H7 isolates. Presumptive starter culture surrogate colonies  
20 were evaluated for Gram reaction, cellular morphology, and catalase activity. The log  
21 CFU for a given inoculum organism was calculated for each sample with a mean log  
22 CFU calculated for each sampling time. A value of 9 CFU (0.95 log CFU) was assigned  
23 when no colonies were present for the least dilute plating.

1           **Weather data.** Because outdoor weather conditions, particularly temperature and  
2 relative humidity, were believed to have a potentially important effect on jerky  
3 processing lethality, the dewpoint at noon in Madison, WI, on the day of each trial was  
4 obtained from meteorological archives  
5 (<http://www.channel3000.com/weather/index.html>).

6  
7

## 8 RESULTS AND DISCUSSION

9 Finished jerky made in the present study had a pH of 5.6 – 6.1, with 22 of 30 samples  
10 having pH of 5.7 – 5.9. The water activity, MPR, and % water-phase salt varied widely,  
11 as expected, given the range of heating and drying conditions evaluated. When the  
12 normal marination and a Type 1 process were used, the ranges of finished product water  
13 activity, MPR, and % water-phase salt were 0.78 - 0.93, 0.52 – 0.95, and 7.9 – 16.6,  
14 respectively. In contrast, the Type 1 product made with no spices had water activity,  
15 MPR, and % water-phase salt of 0.94 – 0.96, 0.63 – 0.65, and 0.60 – 0.70, respectively.  
16 Type 1 product marinated only with spices had a water activity, MPR, and % water-phase  
17 salt of 0.86 – 0.87, 0.67- 0.68, and 11.1 – 11.5, respectively. Products made using the  
18 normal marination procedure and any of the Type 2 - 7 processes had compositional  
19 values similar to those for Type 1 processes with the exception of product from one Type  
20 2 and three Type 3 processes that had water activity, MPR, and % water-phase salt ranges  
21 of 0.64 – 0.75, 0.41 – 0.66, and 11.2 – 17.1, respectively. It should be noted that in many  
22 trials achieving sufficient lethality, the finished product water activity and MPR were  
23 higher (and the % water-phase salt was lower) than desired for optimum consumer

## Lethality of Commercial Beef Jerky Manufacturing Processes

1 acceptance and required in USDA labeling standards (12). However, processors using  
2 the same heating/drying processes could simply extend the drying period to obtain  
3 desired product characteristics. This extended drying would have no adverse effect on  
4 lethality and might, in some situations, increase it.

5 In 21 trials, pathogen numbers were determined following the 22-24-hour post-  
6 marination refrigerated storage step. Numbers of *Salmonella* serovars and *E. coli*  
7 O157:H7 fell by 0.04 – 0.43 and 0.04 – 0.34 log CFU, respectively. We concluded that  
8 this marination step had little lethality, and discontinued post-marination sampling.

9 Throughout the study, consistent trends in dry-bulb, wet-bulb, product-internal,  
10 and surrogate product-internal (beef strip folded over probe) temperatures were observed.  
11 As shown for a typical Type 1 process (Figure 1), wet-bulb temperature was initially well  
12 below dry-bulb temperature. Product-internal temperature was always similar to the wet-  
13 bulb temperature early in the process and could effectively serve as an estimate of wet-  
14 bulb temperature until later in the process. At some time, though, evaporative cooling of  
15 the strips diminished and the product-internal temperature rose toward the dry-bulb  
16 temperature. It is important to note that throughout the jerky heating/drying process,  
17 product-internal temperature was always close to (within 1°C) or higher than the chamber  
18 wet-bulb temperature. Thus, maintaining the chamber wet-bulb temperature (and thereby  
19 the product temperature) high enough to cause pathogen destruction (ca. 51.7°C/125°F  
20 and higher) can strongly influence process lethality. Early in the heating/drying process,  
21 the surrogate product-internal temperature, measured with a jerky strip folded over the  
22 thermocouple, was often lower than the product-internal temperature because the applied  
23 heat had to pass through twice the thickness of meat in the folded strip. Later in the

1 process, the surrogate product-internal temperature rose above the product-internal  
2 temperature, presumably because the greater meat thickness with the folded strip  
3 diminished evaporative cooling near the thermocouple. By the end of the process, when  
4 little evaporative cooling still was occurring, the two temperatures were the same. The  
5 divergence of the two temperatures by a variable amount during much of the process calls  
6 into question the practice of using the surrogate product-internal temperature for overall  
7 process control or evaluation. However, the surrogate product-internal temperature was  
8 very close to the product-internal temperature early in processing and during the wet-bulb  
9 temperature spikes in Type 1 processes, and could be useful in early-process control or  
10 evaluation.

11 Earlier research has established the fact that sub-lethal drying can make pathogens  
12 such as *Salmonella* serovars more resistant to heat (7). This phenomenon was likewise  
13 observed in several early jerky-making trials (data not shown). Therefore, several early-  
14 process wet-bulb temperature spikes were applied to determine the extent of elevated-  
15 humidity heating conditions necessary to achieve desired lethality while maintaining  
16 product quality (Table 2). During the 54.4, 57.2, and 60°C (130, 135, and 140°F) wet-  
17 bulb spikes with concurrent 76.7°C (170°F) dry-bulb temperature, the product-internal  
18 temperature was generally quite similar to the wet-bulb temperature. However, during  
19 the 54.4°C (130°F) wet-bulb spike with a concurrent 87.8°C (190°F) dry-bulb  
20 temperature, the product-internal temperature rose faster than it did during the wet-bulb  
21 spike treatments conducted with 76.7°C (170°F) dry-bulb temperature, and the product-  
22 internal temperature reached at least 5°C higher (Figure 2) than in the latter process  
23 (Figure 1). This more rapid increase in product temperature resulted from faster jerky

## Lethality of Commercial Beef Jerky Manufacturing Processes

1 drying at the lower %RH under 87.8°C (190°F) dry-bulb/54.4°C (130°F) wet-bulb  
2 temperature conditions (see % RH values in Table 2). Similarly, when a wet-bulb  
3 temperature spike of 51.7°C (125°F) was applied concurrently with 76.7°C (170°F) dry-  
4 bulb temperature, the %RH was lower than under 54.4°C (130°F) wet-bulb/76.7°C  
5 (170°F) temperature conditions. This lower %RH led to a faster increase in product-  
6 internal temperature and a more rapid achievement of a  $\geq 5.0$  log reduction in  
7 salmonellae numbers (see first four lines of Table 2).

8 Presently, the USDA has indicated that a jerky-making process has sufficient  
9 lethality if it results in a 5-log reduction of *Salmonella* serovars. However, USDA  
10 guidance for certain other beef products specified a 6.5 log reduction in *Salmonella*  
11 serovars. All tested Type 1 processes (early-process wet-bulb temperature spike  
12 following the standard marination process) resulted in a  $\geq 6.4$  log reduction in both  
13 pathogens by the end of the complete process (including final drying; Table 2). Three  
14 Type 1 processes achieved a  $\geq 5.2$  log reduction in *Salmonella* serovars but caused  
15 smaller decreases in *E. coli* O157:H7 numbers at the end of the wet-bulb temperature  
16 spike (Table 1). These treatments were 51.7°C (125°F) for 60 min, 57.2°C (135°F) for  
17 30 min, and 60°C (140°F) for 10 min. However, a  $\geq 6.4$  log reduction of both *Salmonella*  
18 serovars and *E. coli* O157:H7 was achieved by the end of drying after each of these  
19 processes. One Type 1 process did not cause a  $> 5.0$  log reduction in either pathogen by  
20 the end of the web-bulb spike but subsequent drying resulted in sufficient overall  
21 lethality. This treatment, wet-bulb temperature of 54.4°C (130°F) for 60 minutes,  
22 resulted in decreases of 3.2 – 3.9 and 2.0 – 2.1 log CFU for *Salmonella* serovars and *E.*  
23 *coli* O157:H7, respectively. By the end of drying after these treatments, reductions of  $\geq$

1 6.9 logs had occurred for both pathogen species. The overall effectiveness of these Type  
2 1 processes appeared to result from the fact that product temperature was controllable via  
3 control of wet-bulb temperature and was increased rapidly to levels at which pathogens  
4 were killed while the beef strips were still moist enough to achieve high lethality. It is of  
5 interest to note that the time for which the wet-bulb temperature was elevated, i.e. the  
6 duration of the wet-bulb temperature spike, was generally shorter than the corresponding  
7 times for the same wet-bulb temperatures listed in USDA guidance (15) for the cooking  
8 of beef. The latter times were 112 minutes at 54.4°C (130°F), 36 minutes at 57.2°C  
9 (135°F), and 12 minutes at 60°C (140°F). By comparison, times used in the present  
10 study were 60, 30, and 10 minutes, respectively.

11         The dry-bulb temperature during a Type 1 process was found to have a major  
12 effect on process lethality. The application of a wet-bulb temperature spike of 54.4°C  
13 (130°F) for 60 minutes with a concurrent dry-bulb temperature of 65.5C (150°F; Figure  
14 3) reduced *Salmonella* serovar populations by 4.9 – 5.0 log CFU, but only resulted in 3.2  
15 – 3.6 log reductions in numbers of *E. coli* O157:7 (Table 2) without further drying.  
16 Subsequent drying at 65.5°C (150°F), however, did lead to overall  $\geq 6.7$  log CFU  
17 reductions for both pathogens. These final-sampling results were not noticeably different  
18 than results obtained with the same wet-bulb temperature applied with a concurrent dry-  
19 bulb temperature of 76.7°C (170°F), perhaps because pathogen populations following  
20 both types of process had fallen below the detection limit. When the concurrent dry-bulb  
21 temperature was increased to 87.8°C (190°F), though, the wet-bulb temperature spike  
22 caused a  $\geq 6.5$  log CFU reduction of both pathogens before final drying was even begun  
23 (Table 2).



## Lethality of Commercial Beef Jerky Manufacturing Processes

1           Although USDA guidance (14) recommended 90% relative humidity (RH) during  
2 the early heating period in jerky processing, it also stated that such a high humidity may  
3 not be necessary if alternative procedures are validated. In the Type 1-A processes  
4 conducted here using a dry-bulb temperature of 76.7°C (170°F), the calculated RH for  
5 early-process wet-bulb temperature spikes was 27 - 43% RH. When followed by drying  
6 at 76.7°C (170°F), these processes were sufficient to provide  $\geq 6.4$  log reduction in  
7 numbers of *Salmonella* serovars and *E. coli* O157:H7. Taking into account current USDA  
8 expectations for jerky processing lethality, processors using the Type 1 process  
9 conditions employed in this study [achieving 76.7°C (170°F) dry-bulb temperature within  
10 30 minutes and maintaining this temperature throughout processing] could employ any of  
11 the early-process wet-bulb spike treatments listed in Table 1 followed by drying at  
12 76.7°C (170°F) as scientifically validated processes for making safe whole-muscle beef  
13 jerky.

14           Lethality was compared for several different Type 1 processes at a  
15 common wet-bulb temperature spike time of 60 minutes. The relevant processes were  
16 51.7°C (125°F) or 54.4°C (130°F) wet-bulb temperature spikes with concurrent 76.7°C  
17 (170°F) dry-bulb temperature, and 54.4°C (130°F) wet-bulb temperature spikes with  
18 65.5°C (150°F) or 87.8°C (190°F) dry-bulb temperatures. The 60-minute lethality for  
19 *Salmonella* serovars for these processes averaged 5.4, 3.5, 4.9, and 6.6 logs, respectively  
20 (Table 2). Corresponding values for *E. coli* O157:H7 were 4.7, 2.1, 3.4, and 7.1 logs,  
21 respectively. The reason for the highest lethality during the 51.7°C (125°F) wet-bulb  
22 spike (76.7°C dry-bulb temperature) and the 54.4°C (130°F) wet-bulb temperature spike  
23 with 87.8°C dry-bulb temperature (Figure 2) is that both these processes resulted in a

## Lethality of Commercial Beef Jerky Manufacturing Processes

1 lower environmental relative humidity, and a more rapid increase in product temperature  
2 to lethal levels early in the process while there was still sufficient product moisture for  
3 enhanced lethality.

4 Separate trials were conducted with a Type 1-A process to evaluate the relative  
5 importance of the marinade components in achieving desired process lethality. Results  
6 showed that omission of either water or dry ingredients from the marinade led to  
7 somewhat greater reductions in pathogen numbers after a 54.4°C (130°F)/60 minute wet-  
8 bulb temperature spike (Table 2) but comparable lethality after the subsequent drying.  
9 Whole-muscle beef jerky prepared without the addition of salt, however, had a much  
10 higher water activity when trials were completed. We concluded that the choice of water  
11 level in the jerky marinade used in this study (and hence the amount of marinade pick-up)  
12 or the addition of only dry marinade ingredients was not a critical factor in attaining  
13 desired process lethality.

14 The Type 2 process involving a rapid (15 min) increase in dry-bulb temperature to  
15 76.7°C (170°F) achieved greater pathogen destruction than the Type 3 process in which  
16 dry-bulb temperature did not increase to 76.7°C (170°F) until after 90 minutes (Table 3).  
17 However, both of these processes did result in pathogen reductions exceeding 5.0 log  
18 CFU after drying was completed. An alternative approach that some processors may  
19 elect to use is to initially heat beef strips at the relatively moderate dry-bulb temperature  
20 of 51.7°C (125°F) for a short period of time followed by heating/drying of the strips at a  
21 the relatively high dry-bulb temperature of 76.7°C (170°F). The rationale for this  
22 approach (Type 4 process) is that the short initial heating imparts desirable product  
23 characteristics without increasing pathogen thermotolerance via sublethal stress. As seen

## Lethality of Commercial Beef Jerky Manufacturing Processes

1 in Table 4, the success of a Type 4 process depends on beginning the high-temperature  
2 drying when the product water activity is still relatively high. When drying was begun at  
3 a product water activity of 0.72 or 0.81, the reduction in *Salmonella* serovars was 4.7 log  
4 CFU. Reductions in numbers of *E. coli* O157:H7 in these trials were 5.4 log CFU,  
5 however. When drying was begun at product water activity of 0.86, 0.87, 0.95, or 0.96,  
6 the reduction in both pathogens was 4.9 – 6.7 log CFU. Although such a process clearly  
7 provided less of a safety margin than using a Type 1 process (wet-bulb temperature  
8 spike), it is likely that short-term heating at a low dry-bulb temperature such as 51.7°C  
9 followed by 76.7°C (dry-bulb temperature) drying could achieve sufficient lethality if the  
10 drying is begun when product water activity is  $\geq 0.86$ .

11 Heating whole-muscle beef strips at a constant dry-bulb temperature of either 60  
12 or 71.1°C (140 or 160°F) without the addition of humidity (Type 5 processes) did not  
13 achieve USDA-mandated lethality even when product was dried to water activity levels  
14 typical of commercial beef jerky (Table 5). These processes resulted in decreases in  
15 *Salmonella* serovars and *E. coli* O157:H7 of 3.8 – 4.7 and 3.9 – 4.0 logs, respectively.  
16 Heating at a constant dry-bulb temperature of 82.2°C (180°F) did result in a reduction of  
17 just over 5 log CFU for both pathogens. The cause of the lower lethality in Type 5  
18 processes can be induced from Figure 4. As can be seen, the wet-bulb temperature and  
19 product-internal temperature remained at sub-lethal levels for long periods of time,  
20 allowing pathogen survival during the drying that took place. The surviving cells  
21 apparently had enhanced thermotolerance during subsequent heating, as previously  
22 described (7). Three Type 5 process trials were conducted on winter days with very low  
23 noon dewpoint temperatures (-11, -12, -14°C) and one trial (60°C/140°F dry-bulb

1 temperature) was done in mid-July (noon dewpoint temperature of 20°C). Although it is  
2 possible that using a constant dry-bulb temperature process could attain more lethality  
3 than in the present study if it was employed in very humid weather, we observed no clear  
4 relationship between outdoor dew point and process lethality. We may have mitigated  
5 any weather effects, though, by adjusting process drying times to attain an acceptable  
6 reduction in water activity. Furthermore, our trials were conducted in a climate-  
7 controlled building. Processors with a lower degree of humidity control may need to  
8 adjust process parameters to account for weather extremes.

9         The Type 6 process that had a slow increase in dry-bulb temperature to a  
10 maximum of 60°C (140°F) did not cause more than a 3 log CFU reduction in pathogens.  
11 In contrast, the Type 7 process which had final 1-hour exposure to dry-bulb temperature  
12 of 76.7°C (170°F) caused > 5 log CFU decreases in pathogen numbers (Table 3).

13         The starter culture tested in 19 trials as a pathogen surrogate survived the jerky-  
14 making process considerably better than either of the tested pathogens (Table 6). When  
15 the starter culture population was reduced by at least 3.0 log CFU (nine trials), the  
16 populations of both pathogens decreased by at least 5.0 log CFU in eight of the trials. In  
17 the one exception, the *E. coli* O157:H7 population decreased by 5.0 log CFU and the  
18 population of *Salmonella* serovars was reduced by 4.7 log CFU. In contrast, when the  
19 starter culture population was reduced by < 3.0 log CFU (10 trials), pathogen populations  
20 were considerably less likely to decrease by at least 5.0 log CFU. *Salmonella* serovar and  
21 *E. coli* O157:H7 levels decreased by < 5.0 log CFU in four and three of these trials,  
22 respectively. We conclude that use of this starter culture surrogate with a target lethality

1 of at least 3.0 log CFU could be a useful tool for processors validating their whole-  
2 muscle beef jerky processes.

3         On the basis of our results, we conclude that of the two pathogens studied, *E. coli*  
4 O157:H7 is better able to survive the heating and drying steps used in making whole-  
5 muscle beef jerky. However, foodborne illness outbreaks linked to beef jerky have  
6 primarily involved *Salmonella* serovars, so it is prudent for any validation of a jerky-  
7 making process to involve both pathogens. Because our results clearly show the  
8 importance of wet-bulb temperature in achieving mandated lethality, we strongly  
9 recommend that processors buy or make a wet-bulb thermometer for use in processing, or  
10 use a hygrometer to monitor humidity and then use a commercially available slide rule to  
11 determine wet-bulb temperature from known dry-bulb temperature and %RH values.

12         Reductions in *Salmonella* serovars and *E. coli* O157:H7 of  $\geq 5.0$  log CFU can be  
13 achieved in the production of whole-muscle beef jerky by ensuring that high enough wet-  
14 bulb temperatures are reached and maintained early in the process (Type 1 processes) or  
15 that high dry-bulb temperature heating and drying is done before the beef strip water  
16 activity has fallen below 0.86 (Type 2, 3, 4, and 7 processes). Alternatively, guidance  
17 from USDA (14) has indicated that internal temperatures listed in USDA-accepted  
18 “Appendix A” time/temperature combinations (15) are effectively wet-bulb temperatures.  
19 Processors could consider a process valid in which the smokehouse wet-bulb temperature  
20 and, therefore the product-internal temperature, was at or above a designated level for a  
21 time at least as long as specified in Appendix A.

22

23

1 ACKNOWLEDGEMENTS

2 The authors gratefully acknowledge the laboratory assistance of Ryan Algino, Greg  
3 Burnham, Rebecca Engel, Melody Fanslau, Amy Haen, Erica Ready, Erica Schoeller,  
4 and Melissa Talbot. The authors also acknowledge the smokehouse assistance of Ruben  
5 Zarraga. This work was supported by a grant from the United States Department of  
6 Agriculture, Food Safety & Inspection Service.

7

8

9

10 REFERENCES

- 11 1. Blankenship, L.C., C.E. Davis, and G.J. Magner. 1980. Cooking methods for  
12 elimination of *Salmonella typhimurium* experimental surface contaminant from  
13 rare dry-roasted beef roasts. *J. Food Science* 45: 270-273.
- 14 2. Blankenship, L.C. 1978. Survival of *Salmonella typhimurium* experimental  
15 contaminant during cooking of beef roasts. *Appl. Environ. Microbiol.* 35: 1160-  
16 1165.
- 17 3. Calicioglu, M., J.N. Sofos, J. Samelis, P.A. Kendall, and G.C. Smith. 2003.  
18 Effect of acid adaptation on inactivation of *Salmonella* during drying and storage  
19 of beef jerky treated with marinades. *Int. J. Food Microbiol.* 89: 51-65.
- 20 4. Calicioglu, M., J.N. Sofos, J. Samelis, P.A. Kendall, and G.C. Smith. 2002a.  
21 Inactivation of acid-adapted and nonadapted *Escherichia coli* O157:H7 during  
22 drying and storage of beef jerky treated with different marinades. *J. Food Prot.*  
23 65: 1394-1405.

## Lethality of Commercial Beef Jerky Manufacturing Processes

- 1        5. Calicioglu, M., J.N. Sofos, J. Samelis, P.A. Kendall, and G.C. Smith. 2002b.  
2            Destruction of acid-and non-adapted *Listeria monocytogenes* during drying and  
3            storage of beef jerky. *Food Microbiol.* 19: 545-559.
- 4        6. Eidson, M., C.M. Sewell, G. Graves, and R. Olson. 2000. Beef jerky  
5            gastroenteritis outbreaks. *J. Environ. Health* 62: 9-13.
- 6        7. Goepfert, J.M., I.K. Iskander, and C.H. Amundson. 1970. Relation of the heat  
7            resistance of salmonellae to the water activity of the environment. *Appl.*  
8            *Microbiol.* 19: 429-433.
- 9        8. Goodfellow, S.J. and W.L. Brown. 1978. Fate of *Salmonella* inoculated into beef  
10            for cooking. *J. Food Prot.* 41: 598-605.
- 11       9. Harrison, J.A., M.A. Harrison, R.A. Rose-Morrow, and R.L. Shewfelt. 2001.  
12            Home-style beef jerky: effect of four preparation methods on consumer  
13            acceptability and pathogen inactivation. *J. Food Prot.* 64: 1194-1198.
- 14       10. Nummer, B.A., J.A. Harrison, M.A. Harrison, P. Kendall, J.N. Sofos, and E.L.  
15            Andress. 2004. Effects of preparation methods on the microbiological safety of  
16            home-dried meat jerky. *J. Food Prot.* 67: 2337-2341.
- 17       11. ProMED-mail. 2003. *Salmonella* Kiambu, beef jerky – USA (New Mexico).  
18            International Society for Infectious Diseases. Archive no. 20031001.2471.  
19            Available at <http://www.promedmail.org>. Accessed 21 July 2005.
- 20       12. United States Department of Agriculture, Food Safety & Inspection Service.  
21            2005a. Food Standards and Labeling Policy Book. Available at  
22            [http://www.fsis.usda.gov/OPPDE/larc/Policies/Labeling\\_Policy\\_Book\\_082005.pdf](http://www.fsis.usda.gov/OPPDE/larc/Policies/Labeling_Policy_Book_082005.pdf)  
23            f. Accessed 11 November 2005.

## Lethality of Commercial Beef Jerky Manufacturing Processes

- 1        13. United States Department of Agriculture, Food Safety & Inspection Service.  
2            2005b. Risk assessment of the impact of lethality standards on salmonellosis  
3            from ready-to-eat meat and poultry products; draft for public review and  
4            comment.  
5            [http://www.fsis.usda.gov/PDF/Risk\\_Assessment\\_RTE\\_Salm\\_Leth\\_Main.pdf](http://www.fsis.usda.gov/PDF/Risk_Assessment_RTE_Salm_Leth_Main.pdf)  
6            [accessed 21 July 2005](#)
- 7        14. United States Department of Agriculture, Food Safety & Inspection Service.  
8            2004. Compliance guideline for meat and poultry jerky produced by small and  
9            very small plants.  
10           [http://www.fsis.usda.gov/PDF/Compliance\\_Guideline\\_Jerky.pdf](http://www.fsis.usda.gov/PDF/Compliance_Guideline_Jerky.pdf). Accessed 4  
11           May 2005.
- 12        15. United States Department of Agriculture Food Safety and Inspection Service.  
13            1999. Compliance guidelines for meeting lethality performance standards for  
14            certain meat and poultry products, Appendix A. Available at:  
15            <http://www.fsis.usda.gov/oa/fr/95033f-a.htm>. Accessed 24 May 2005.



1

2 FIGURE LEGENDS

3 1. Wet-bulb, dry-bulb, and product-internal temperatures during the manufacture of  
4 whole-muscle beef jerky by a Type 1-A process with a 54.4°C / 130°F, 60-minute  
5 wet-bulb temperature spike.

6 2. Wet-bulb, dry-bulb, and product-internal temperatures during the manufacture of  
7 whole-muscle beef jerky by a Type 1-B process with a dry-bulb temperature  
8 setting of 87.8°C / 190°F.

9 3. Wet-bulb, dry-bulb, and product-internal temperatures during the manufacture of  
10 whole-muscle beef jerky by a Type 1-B process with a dry-bulb temperature  
11 setting of 65.5°C / 150°F.

12 4. Wet-bulb, dry-bulb, and product-internal temperatures during the manufacture of  
13 whole-muscle beef jerky by a Type 5 process with a dry-bulb temperature setting  
14 of 60.0°C / 140°F.

15

16

1 Table 1. Summary of heating/drying processes used to make whole-muscle beef jerky.

2

3 Process                      Dry-Bulb (Controlled)                      Wet-Bulb (Controlled)                      Cumulative  
 4 Type                      Temperature (°C/°F)    Time (min)                      Temperature (°C/°F)                      Time (min)                      Time (min)

5	<u>1-A</u>	62.8 / 145	15	Not Controlled = NC	---	15
6		76.7 / 170	15	NC	---	30
7	then	76.7 / 170	60	51.7 / 125	60	90
8	OR	76.7 / 170	60	54.4 / 130	60	90
9	OR	76.7 / 170	30	57.2 / 135	30	60
10	OR	76.7 / 170	10	60 / 140	10	40

11 Always followed by 76.7 / 170                      varied (to targeted    NC                      ----                      varied  
 12                      final product dryness)

13

14	1 – B	62.8 / 145	15	NC	---	15
15	then	65.5 / 150	15	NC		
16		65.5 / 150	60	54.4 / 130	60	90
17		65.5 / 150	varied (to targeted	NC	---	varied
18			final product dryness)			
19	OR	87.8 / 190	15	NC		
20		87.8 / 190	60	54.4 / 130	60	90
21		87.8 / 190	varied (to targeted	NC	----	varied

Lethality of Commercial Beef Jerky Manufacturing Processes

1			final product dryness)			
2						
3	2	62.8 / 145	15	NC	---	15
4		76.7 / 170	varied (to targeted	NC	---	varied
5			final product dryness)			
6						
7	3	62.8 / 145	90	NC	---	15
8		76.7 / 170	varied (to targeted	NC	---	varied
9			final product dryness)			
10						
11	4	51.7 / 125	varied (to targeted a <sub>w</sub> )	NC	---	varied
12		76.7 / 170	varied (to targeted	NC	---	varied
13			final product dryness)			
14						
15	5	60 / 140	varied (to targeted	NC	----	varied
16			final product dryness)			
17	OR	71.1 / 160	varied (to targeted	NC	---	varied
18			final product dryness)			
19	OR	82.2 / 180	varied (to targeted	NC	---	varied
20			final product dryness)			
21						

Lethality of Commercial Beef Jerky Manufacturing Processes

1	6	48.9 / 120	60	NC	---	60
2		54.4 / 130	60	NC	---	120
3		60 / 140	60	NC	---	180
4						
5	7	48.9 / 120	60	NC	---	60
6		54.4 / 130	60	NC	---	120
7		60 / 140	60	NC	---	180
8		76.7 / 170	60	NC	---	240
9						

Lethality of Commercial Beef Jerky Manufacturing Processes

1

2 Table 2. Process lethality<sup>a</sup> against *Salmonella* serovars (S) and *Escherichia coli* O157:H7 (EC) during Type 1 - A and 1 - B processing  
3 (wet-bulb temperature spike = WBS; followed by post-spike drying at 76.7°C = PSD) of whole muscle beef jerky.

4

5	WBS		PSD		Lethality as determined after				Water activity		Product-internal	
	6 Temp. (°C)	WBS Time (min)	%RH <sup>b</sup>	PSD time (min)	WBS		PSD		after		temperature after	
7	(°C)	(min)		(min)	S	EC	S	EC	WBS	PSD	WBS	PSD
8	Type 1 – A	Strips marinated with water and dry ingredients				Dry-bulb temperature at 76.7°C during wet-bulb spike.						
9	51.7	60	27	60	5.6	5.6	6.5	6.7	0.87	0.81	60.5	70.0
10	51.7	60	27	60	5.2	3.8	6.4	7.1	0.89	0.78	56.1	65.0
11	54.4	60	32	120	3.2	2.0	6.9	7.1	0.93	0.88	54.4	63.9
12	54.4	60	32	120	3.9	2.1	6.9	7.0	0.92	0.87	55.0	64.4
13	57.2	30	37	120	6.4	2.7	7.0	7.1	ND	0.86	58.3	63.9
14	57.2	30	37	90	5.3	3.1	7.0	7.1	0.93	0.90	57.8	66.1
15	60	10	43	120	6.2	3.8	7.0	7.2	0.96	0.90	58.9	67.8
16	60	10	43	120	6.7	2.2	6.8	7.0	0.96	0.84	60.0	62.8
17	Type 1 – B	Dry-bulb temperature at 65.5°C during wet-bulb spike										
18	54.4	60	56	90	4.9	3.2	6.7	7.1	0.95	0.85	55.6	57.2
19	54.4	60	56	150	5.0	3.6	6.9	6.9	0.97	0.91	55.0	57.8
20	Type 1 – B	Dry-bulb temperature at 87.8°C during wet-bulb spike										
21	54.4	60	19	30	6.7	7.3	7.0	7.4	0.93	0.89	66.7	71.1

Lethality of Commercial Beef Jerky Manufacturing Processes

1	54.4	60	19	45	6.5	6.9	7.2	7.1	0.95	0.89	65.0	73.9
2	Type 1 – A		Strips marinated only with water				Dry-bulb temperature at 76.7°C during wet-bulb spike					
3	54.4	60	32	60	5.6	4.9	6.6	6.9	0.98	0.94	55.6	63.9
4	54.4	60	32	60	5.4	4.5	6.1	7.0	0.99	0.96	55.6	64.4
5	Type 1 – A		Strips marinated only with dry ingredients				Dry-bulb temperature at 76.7°C during wet-bulb spike					
6	54.4	60	32	90	5.9	3.9	7.1	7.0	0.95	0.87	57.2	66.7
7	<u>54.4</u>	<u>60</u>	<u>32</u>	<u>60</u>	<u>5.6</u>	<u>6.9</u>	<u>6.6</u>	<u>7.1</u>	<u>0.93</u>	<u>0.86</u>	<u>58.9</u>	<u>68.3</u>

8 <sup>a</sup> Reduction in log CFU per sample relative to initial pathogen load prior to marination.

9 <sup>b</sup> Percent relative humidity during wet-bulb temperature spike calculated from wet-bulb and dry-bulb temperature settings using a  
10 slide rule.

11 ND = Not Determined

12

13

14

## Lethality of Commercial Beef Jerky Manufacturing Processes

1 Table 3. Process lethality against *Salmonella* serovars (S) and *Escherichia coli* O157:H7 (EC) during the type 4, 5, 6, and 7 processes  
 2 for heating and drying of whole-muscle beef jerky. Processes were 2 = fast come-up [15 minutes to reach dry-bulb temperature of  
 3 76.7°C (170°F), followed by drying at 76.7°C], 3 = slow come-up [90 minutes to reach dry-bulb temperature of 76.7°C (170°F),  
 4 followed by drying at 76.7°C], 6 = 1 hour each at dry-bulb temperatures of 48.9, 54.4, and 60°C (120, 130, and 140°F), or 7 = 1 hour  
 5 each at dry-bulb temperatures of 48.9, 54.4, 60, and 76.7°C (120, 130, 140, and 170°F). No humidity was added to the smokehouse  
 6 chamber during processing.

7

8 Process Lethality (reduction in log CFU)<sup>a</sup> and product characteristics at

9 Intermediate Time End

10 Product – internal Product - internal

11 Treatment RH<sup>b</sup> (min) S EC a<sub>w</sub><sup>c</sup> Temperature (°C) S EC a<sub>w</sub> Temperature (°C)

12 2 31 - 17 60 3.3 2.7 0.95 48.9 6.1 5.6 0.86 62.8

13 2 27 - 21 60 2.6 1.8 0.96 52.8 6.4 6.4 0.91 67.8

14 3 41 - 21 90 2.0 1.5 0.97 46.7 5.5 5.6 0.87 67.8

15 6 41 - 24 120 1.6 1.7 0.92 40.5 2.9 2.7 0.84 51.1

16 7 43 - 15 180 3.5 2.5 0.94 49.4 6.0 5.6 0.89 67.2

17

18 <sup>a</sup> Reduction in log CFU per sample relative to initial pathogen load prior to marination.

## Lethality of Commercial Beef Jerky Manufacturing Processes

- 1 <sup>b</sup> Percent relative humidity, change during process (initial value – final value). Values calculated from wet-bulb and dry-bulb
- 2 temperatures using a slide rule.
- 3 <sup>c</sup> Water activity.



Lethality of Commercial Beef Jerky Manufacturing Processes

1

2 Table 4. Process lethality against *Salmonella* serovars (S) and *Escherichia coli* O157:H7 (EC) during the Type 4 process of heating of  
 3 whole-muscle beef jerky at a constant dry-bulb temperature of 51.7°C (125°F) to attain a desired water activity followed by drying at  
 4 76.7°C (170°F). No humidity was added to the smokehouse chamber during processing.

5

6 Heating Lethality (reduction in log CFU)<sup>a</sup> and product characteristics at

7 Time	End of heating						End of drying				
	8 (min)	RH <sup>b</sup>	Product - internal		S	EC	Cumul. Time	Product - internal			
a <sub>w</sub> <sup>c</sup>			Temperature (°C)	(min)				a <sub>w</sub>	Temperature (°C)	S	EC
10 240	ND	0.72	47.2	3.3	3.1	300	0.65	74.4	4.7	5.4	
11 240	32 – 33	0.81	42.8	3.3	2.7	300	0.75	70.0	4.7	5.4	
12 180	30 - 36	0.86	42.8	3.2	2.2	240	0.67	68.3	5.8	5.0	
13 270	ND	0.87	45.6	4.3	2.9	330	0.82	71.7	5.8	6.7	
14 120	33 - 38	0.95	37.8	1.8	1.7	240	0.83	67.8	4.9	5.7	
15 120	35 - 38	0.96	38.9	2.1	1.9	225	0.82	70.0	5.6	5.9	

16

17 <sup>a</sup> Reduction in log CFU per sample relative to initial pathogen load prior to marination.

18 <sup>b</sup> Percent relative humidity, change during process (initial value – final value). Values calculated from wet-bulb and dry-bulb  
 19 temperatures using a slide rule.

20 <sup>c</sup> Water activity.

## Lethality of Commercial Beef Jerky Manufacturing Processes

- 1 ND = Not Determined.

Lethality of Commercial Beef Jerky Manufacturing Processes

1 Table 5. Process lethality against *Salmonella* serovars (S) and *Escherichia coli* O157:H7 (EC) during the Type 5 process of heating  
 2 and drying of whole-muscle beef jerky at a constant dry-bulb temperature of 60, 65.5, or 82.2°C (140, 160, or 180°F). No humidity  
 3 was added to the chamber during processing.

4

5 Dry-Bulb Lethality (reduction in log CFU)<sup>a</sup> and product characteristics at

6 Temperature Intermediate Time Final Time

7 Product – internal Product - internal

8 (°C)	RH <sup>b</sup>	(min)	S	EC	a <sub>w</sub> <sup>c</sup>	temperature (°C)	(min)	S	EC	a <sub>w</sub>	temperature (°C)
9 60	32 – 24	90	4.3	3.8	0.73	46.7	120	4.2	3.9	0.64	53.9
10 60	28 - 26	90	1.9	1.6	0.96	42.2	120	3.8	3.9	0.79	52.8
11 71.1	34 - 18	60	4.0	3.3	0.87	50.6	75	4.7	4.0	0.80	59.4
12 82.2	29 - 15	60	5.2	4.6	0.72	64.4	75	5.1	5.6	0.65	72.2

13

14 <sup>a</sup> Reduction in log CFU per sample relative to initial pathogen load prior to marination.

15 <sup>b</sup> Percent relative humidity, change during process (initial value – final value). Values calculated from wet-bulb and dry-bulb  
 16 temperatures using a slide rule.

17 <sup>c</sup> Water activity.

## Lethality of Commercial Beef Jerky Manufacturing Processes

1 Table 6. Comparison of *Salmonella* serovar (S), *Escherichia coli* O157:H7 (EC), and starter culture surrogate (SC) death during  
 2 various processes for making whole-muscle beef jerky.

3	4 Process	Lethality (reduction in log CFU) <sup>a</sup> for Finished Sample		
5	<u>Treatment</u>	<u>S</u>	<u>EC</u>	<u>SC</u>
6	Consumer	4.4	2.8	2.5
7	Consumer	4.2	3.0	1.8
8	1-A	7.0	7.2	3.5
9	1-B;87.8°C dry-bulb	7.0	7.4	3.2
10	1-B; 87.8°C dry-bulb	7.2	7.1	3.2
11	1-B; 65.5°C dry-bulb	6.7	7.1	3.2
12	1-B; 65.5°C dry-bulb	6.9	6.9	2.8
13	1-A; spices only	7.1	7.1	3.3
14	1-A; spices only	6.6	7.1	2.8
15	1-A; water only	6.6	6.9	2.7
16	1-A; water only	6.1	7.0	3.0
17	2	6.4	6.4	3.4
18	2	6.1	5.6	2.3
19	4	4.7	5.4	2.9
20	4	5.8	5.0	3.2
21	4	5.8	6.7	2.8

## Lethality of Commercial Beef Jerky Manufacturing Processes


1	4	4.7	5.4	3.3
2	5	3.8	3.9	1.9
3	6	6.0	5.6	2.6

---

4

5 <sup>a</sup> Reduction in log CFU per sample relative to initial pathogen load prior to marination.


# Lethality of Commercial Beef Jerky Manufacturing Processes



1

Lethality of Commercial Beef Jerky Manufacturing Processes


2.



1

Lethality of Commercial Beef Jerky Manufacturing Processes


3.



1



4.



1