

**COLORADO RIVER SYSTEM
CONSUMPTIVE USES
AND
LOSSES REPORT
1981-1985**



Mission: As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally-owned public lands and natural and cultural resources. This includes fostering wise use of our land and water resources, protecting our fish and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to assure that their development is in the best interests of all our people. The Department also promotes the goals of the Take Pride in America campaign by encouraging stewardship and citizen responsibility for the public lands and promoting citizen participation in their care. The Department also has a major responsibility for American Indian reservation communities and for people who live in Island Territories under U.S. Administration.

**COLORADO RIVER SYSTEM
CONSUMPTIVE USES
AND
LOSSES REPORT
1981-1985**

June 1991

**United States Department of the Interior
Bureau of Reclamation
Upper Colorado Region
Lower Colorado Region**


FOREWORD

This report was prepared pursuant to the Colorado River Basin Project Act of 1968, Public Law 90-537. The act directs the Secretary of the Interior to "make reports as to the annual consumptive uses and losses of water from the Colorado River System after each successive 5-year period, beginning with the 5-year period starting October 1, 1970 Such reports shall be prepared in consultation with the States of the Lower Basin individually and with the Upper Colorado River Commission and shall be transmitted to the President, the Congress, and to the Governors of each State signatory to the Colorado River Compact."


This report reflects the Department of the Interior's best estimate of actual consumptive uses and losses within the Colorado River Basin. The reliability of the estimate is affected by the availability of data and the current capabilities of data evaluation.

REPORTING AREA

1. Green River
2. Upper Main Stem
3. San Juan-Colorado
4. Muddy River
5. Virgin River
6. Mainstream and Remaining Tributary Area
7. Bill Williams River
8. Gila River
9. Little Colorado River



UNITED STATES
DEPARTMENT OF THE INTERIOR
BUREAU OF RECLAMATION
COLORADO RIVER SYSTEM
CONSUMPTIVE USES & LOSSES REPORT
PUBLIC LAW 90-537



SUMMARY

This report presents estimates of the consumptive uses and losses from the Colorado River System for each water year from 1981 through 1985. It includes a breakdown of the beneficial consumptive use by major types of use, by major tributary streams, and, where possible, by individual States.

The Colorado River rises in the Rocky Mountains of Colorado, flows southwesterly about 1,400 miles and terminates in the Gulf of California. Its drainage area of 242,000 square miles in this country represents one-fifteenth the area of the United States. Its water is used for irrigation, municipal and industrial purposes, electric power generation, mineral activities, livestock, fish and wildlife, and recreation. Large amounts are exported from the system to adjoining areas. The following table summarizes annual water use from the system by basins and States, including water use supplied by ground-water overdraft. Distribution of water use by types of use from the various reporting areas is contained within the body of the report.

Summary.--Colorado River System Uses and Losses Report, Public Law 90-537
Water use by states, basins, and tributaries¹ (1981-1985)
(Units: 1,000 acre-feet)

State and basin of use	Water year					Average 1981-85
	1981	1982	1983	1984	1985	
Arizona	6,167	5,143	4,237	5,041	4,838	5,085
Upper Basin	(42)	(40)	(42)	(44)	(44)	(42)
Lower Basin main stem	(1,282)	(1,127)	(910)	(999)	(1,032)	(1,070)
Lower Basin tributaries	(4,843)	(3,976)	(3,285)	(3,998)	(3,762)	(3,973)
California	4,839	4,349	3,953	4,679	4,710	4,506
Lower Basin main stem	(4,839)	(4,349)	(3,953)	(4,679)	(4,710)	(4,506)
Colorado	2,086	2,106	1,920	1,865	1,994	1,994
Upper Basin	(2,086)	(2,106)	(1,920)	(1,865)	(1,994)	(1,994)
Nevada	350	352	339	355	373	354
Lower Basin main stem	(109)	(106)	(92)	(99)	(103)	(102)
Lower Basin tributaries	(241)	(246)	(247)	(256)	(270)	(252)
New Mexico	342	425	425	417	401	402
Upper Basin	(314)	(399)	(400)	(394)	(375)	(377)
Lower Basin tributaries	(28)	(26)	(25)	(23)	(26)	(25)
Utah	782	746	718	762	879	777
Upper Basin	(666)	(633)	(596)	(638)	(755)	(657)
Lower Basin tributaries	(116)	(113)	(122)	(124)	(124)	(120)
Wyoming	341	330	346	307	336	332
Upper Basin	(341)	(330)	(346)	(307)	(336)	(332)
Other ²	1,598	1,403	1,896	1,197	1,783	1,575
Upper Basin Colorado River Storage Project reservoir evaporation	(541)	(501)	(614)	(659)	(645)	(592)
Lower Basin main stem reservoir evaporation and channel loss	(1,057)	(902)	(1,282)	(538)	(1,138)	(983)
Total--Colorado River System						
Upper Basin	3,449	3,508	3,304	3,248	3,504	3,402
Lower Basin main stem	6,230	5,582	4,955	5,777	5,845	5,678
Lower Basin tributaries	5,228	4,361	3,679	4,401	4,182	4,370
Other--Reservoir evaporation and channel loss	<u>1,598</u>	<u>1,403</u>	<u>1,896</u>	<u>1,197</u>	<u>1,783</u>	<u>1,575</u>
	16,505	14,854	13,834	14,623	15,314	15,025
Water passing to Mexico	3,997	1,817	9,782	16,992	13,396	9,197
Treaty	(1,751)	(1,495)	(1,646)	(1,694)	(1,671)	(1,652)
Minutes 218, 241, and 242	(131)	(146)	(166)	(138)	(131)	(142)
Excess release	(2,115)	(176)	(7,970)	(15,160)	(11,594)	(7,403)
Total--Colorado River System and water passing to Mexico	20,502	16,671	23,616	31,615	28,710	24,222

¹ Onsite consumptive uses and losses; includes water uses satisfied by ground-water overdraft (from tables C-2 through C-6 and LC-3).

² Represents main stem reservoir evaporation in the Upper Basin and main stem reservoir evaporation and channel loss below Lee Ferry in the Lower Basin.

CONTENTS

	Page
Foreword	i
General Location Map	ii
Summary	iii
Water Use by States, Basins, and Tributaries Summary	iv
Contents	v
Introduction	1
Authority	1
Plan of Study	1
Study Reporting Areas	1
Upper Colorado River Basin	2
Green River, Wyoming-Colorado-Utah	2
Upper Main Stem, Colorado-Utah	3
San Juan-Colorado, Colorado-New Mexico-Utah-Arizona	3
Lower Colorado River Basin	4
Main Stem Below Lee Ferry, Arizona-California-Nevada	4
Remaining Areas in Arizona-Nevada-Utah	5
Little Colorado River, Arizona-New Mexico	5
Virgin River, Arizona-Utah	5
Muddy River, Nevada	6
Bill Williams River, Arizona	6
Gila River, Arizona-New Mexico	7
Terminology	7
Methodology and Data Adequacy	9
Colorado River Basin Tributaries	9
Agriculture	9
Reservoir Evaporation	11
Ground Water	11
Stockpond Evaporation and Livestock	12
Mineral Resources	12
Thermal Electric Power	12
Municipal and Industrial	13
Transbasin Diversions	13
Lower Colorado River Main Stem	13
Beneficial Consumptive Uses and Losses	14
Upper Colorado River Tributaries	14
Lower Colorado River Main Stem	15
Lower Colorado River Tributaries	15

TABLES

Colorado River System

Table	Page
Summary - Water Use by States, Basins, and Tributaries, 1981-1985	iv
C-1 Drainage Area of the Colorado River System, Area Within Each State and Mexico by Major Tributary Streams	17
C-2 Summary of Estimated Water Use by States and Types of Use, 1981	18
C-3 Summary of Estimated Water Use by States and Types of Use, 1982	19
C-4 Summary of Estimated Water Use by States and Types of Use, 1983	20
C-5 Summary of Estimated Water Use by States and Types of Use, 1984	21
C-6 Summary of Estimated Water Use by States and Types of Use, 1985	22

Upper Colorado River

UC-1 Estimated Main Stem Reservoir Evaporation, 1981-1985	23
UC-2 Estimated Water Use Within States, by Major Tributaries, and Types of Use, 1981	24
UC-3 Estimated Water Use Within States, by Major Tributaries, and Types of Use, 1982	25
UC-4 Estimated Water Use Within States, by Major Tributaries, and Types of Use, 1983	26
UC-5 Estimated Water Use Within States, by Major Tributaries, and Types of Use, 1984	27
UC-6 Estimated Water Use Within States, by Major Tributaries, and Types of Use, 1985	28
UC-7 Irrigated Acreage, 1981-1985	29
UC-8 Population Estimates, 1981-1985	30

Lower Colorado River

LC-1 Colorado River Main Stem Estimated Reservoir Evaporation and Channel Loss, 1981-1985	31
LC-2 Estimated Water Use Including Colorado Main Stem by States and Types of Use, 1981-1985	32
LC-3 Colorado Main Stem Water Use and Exports Within States and Mexico, 1981-1985	33
LC-4 Estimated Water Use Within States, by Major Tributaries, and Types of Use, 1981	34
LC-5 Estimated Water Use Within States, by Major Tributaries, and Types of Use, 1982	35

TABLES (Continued)

LC-6	Estimated Water Use Within States, by Major Tributaries, and Types of Use, 1983	36
LC-7	Estimated Water Use Within States, by Major Tributaries, and Types of Use, 1984	37
LC-8	Estimated Water Use Within States, by Major Tributaries, and Types of Use, 1985	38
LC-9	Irrigated Acreage, 1981-1985	39
LC-10	Population Estimates, 1981-1985	40

COLORADO RIVER SYSTEM

CONSUMPTIVE USES AND LOSSES REPORT

1981-1985

Consumptive Uses and Losses

INTRODUCTION

The Colorado River System is composed of portions of seven States--Arizona, California, Colorado, Nevada, New Mexico, Utah, and Wyoming. It has a drainage area of about 242,000 square miles and represents about one-fifteenth the area of the United States.

This report incorporates annual estimates of consumptive uses and losses of water from the system from 1981 through 1985. Wherever available, water use reports prepared in accordance with legal requirements concerning the operation of the Colorado River were utilized. Base data needed to estimate onsite consumptive uses were taken largely from existing reports and studies and from ongoing programs. Where current data were not available, estimated values were developed by various techniques and reasoned judgment.

In general, methodology followed the techniques normally used within the system for estimating water use.

Nothing in this report is intended to interpret the provisions of the Colorado River Compact (45 Stat. 1057), the Upper Colorado River Basin Compact (63 Stat. 31), the Water Treaty of 1944 with the United Mexican States (Treaty Series 994; 59 Stat. 1219), the decree entered by the Supreme Court of the United States in *Arizona vs. California, et al.* (376 U.S. 340), the Boulder Canyon Project Act (45 Stat. 1057), the Boulder Canyon Project Adjustment Act (54 Stat. 774; 43 U.S.C. 618a), the Colorado River Storage Project Act, (70 Stat. 105; 43 U.S.C. 620), or the Colorado River Basin Project Act (82 Stat. 885; 43 U.S.C. 1501).

AUTHORITY

The authority for this report is contained in Public Law 90-537, the Colorado River Basin Project Act of 1968. Title VI, Section 601(b)(1) of the act reads as follows:

(b) The Secretary is directed to:

(1) Make reports as to the annual consumptive uses and losses of water from the Colorado River System after each successive 5-year period, beginning with the 5-year period starting October 1, 1970. Such reports shall include a detailed break down of the beneficial consumptive use of water on a State-by-State basis. Specific figures on quantities consumptively used from the major tributary streams flowing into the Colorado River shall also be included on a State-by-State basis. Such reports shall be prepared in consultation with the States of the Lower Basin individually and with the Upper Colorado River Commission, and shall be transmitted to the President, the Congress, and to the Governors of each State signatory to the Colorado River Compact.

PLAN OF STUDY

No plan of study was submitted for this report. The same procedures as were outlined in the previous 5-year plan of study were used in preparing this report.

STUDY REPORTING AREAS

The estimated drainage area of the Colorado River System in the United States is about 242,000 square miles.

The river originates in the Rocky Mountains of Colorado and Wyoming, flows southwest about 1,400 miles, and terminates in the Gulf of California. The system consists of portions of seven States: Arizona, California, Colorado, New Mexico, Nevada, Utah, and Wyoming. The drainage area was divided into nine subbasins for the purpose of this report.

The Colorado River Compact, signed November 24, 1922, was established because the Upper Basin States were concerned that any storage on the river would be put to use more rapidly by the Lower Basin States, thus allowing them to claim appropriative rights. The Upper Basin States wanted provisions for their future development.

The term "Upper Basin States" refers to the States of Colorado, New Mexico, Utah, and Wyoming. "Lower Basin States" refers to the States of Arizona, California, and Nevada. However, the Upper Colorado River Basin refers to the hydrologic boundaries. Lee Ferry is the division point between the Upper Colorado River Basin and the Lower Colorado River Basin. Therefore, the hydrologic boundaries include portions of Arizona in the Upper Colorado River Basins and portions of Utah and New Mexico in the Lower Colorado River Basin. Hydrologic boundaries are shown on the map on page ii.

The major tributary streams selected as reporting areas in the Upper Colorado River Basin are: Green River (Wyoming, Colorado, Utah); Upper Main Stem (Colorado, Utah); and San Juan-Colorado (Colorado, New Mexico, Utah, Arizona).

Five tributary areas in addition to the main stem were selected in the Lower Colorado River Basin: Little Colorado River (Arizona, New Mexico); Virgin River (Utah, Arizona); Muddy River (Nevada); Bill Williams River (Arizona); Gila River (Arizona, New Mexico); and remaining areas in Arizona, Nevada, and Utah. The outflow point and drainage area for each is shown in table C-1. The boundaries of the reporting areas are shown on the map on page ii. A brief description of each reporting area follows.

Upper Colorado River Basin

Green River, Wyoming-Colorado-Utah

The Green River reporting area comprises about 44,800 square miles in southwestern Wyoming, northwestern Colorado, and northeastern and east-central Utah.

Principal tributaries of the Green River are Blacks Fork, New Fork, and Big Sandy Creek in southwestern Wyoming; Yampa and White Rivers on the western slope of the Continental Divide in northwestern Colorado; and the Price, Duchesne, and San Rafael Rivers in eastern Utah. These streams are fed by numerous headwater lakes.

The largest towns in the reporting area are Rock Springs and Green River in Wyoming; Vernal and Price in Utah; and Craig, Steamboat Springs, and Meeker in Colorado.

Mineral production is the major industry. Oil and natural gas are of primary importance, as are coal,

gilsonite, asphalt, and trona (soda ash). Thermal electric power production is becoming an increasingly important industry.

Agriculture ranks near mineral production in importance to the local economy. Agricultural development is centered around livestock production, primarily beef cattle and sheep. Because of a short growing season, crop production is limited largely to small grain, hay, and pasture. These crops are used as winter livestock feed and complement the vast areas of public grazing lands.

Irrigation consumptive use accounts for about 75 percent of the total water use in the Green River reporting area exclusive of any share of main stem evaporation. Nearly 580,000 acres of land are irrigated in an average year. Large exports of water are made to the Great Basin in Utah.

Upper Main Stem, Colorado-Utah

The Upper Main Stem reporting area is drained by the Colorado River and its tributaries above the mouth of the Green River. Principal tributaries are the Roaring Fork, Gunnison, and the Dolores Rivers. The Upper Main Stem reporting area consists of 26,200 square miles, with about 85 percent of the area in Colorado and the remainder in Utah.

Grand Junction, Montrose, and Glenwood Springs are the principal towns in the Colorado portion of the upper main stem of the Colorado River. Moab is the only major community in the Utah portion of the upper main stem of the Colorado River.

Mineral production is the predominant industry. This area is the Nation's chief source of molybdenum and is a major source of vanadium, uranium, lead, zinc, coal, and gilsonite. On the Upper Main Stem reporting area, as in that of the Green River, agriculture centers around production of livestock which feeds on irrigated lands to complement the large areas of rangeland. Somewhat more diversification of crops occur in the Upper Main Stem, however, with some major land areas devoted to corn, beans, potatoes, table vegetables, and fruit. This diversification is made possible by climatic and topographic conditions that create favorable air drainage and minimize frost damage.

Irrigation consumptive use accounts for about 58 percent of the water use in the Upper Main Stem reporting area exclusive of any share of main stem evaporation. In an average year about 555,000 acres of land are irrigated.

A considerable amount (29 percent) of water is exported to serve agricultural and municipal needs on the eastern slope of the Continental Divide in Colorado.

San Juan-Colorado, Colorado-New Mexico-Utah- Arizona

The San Juan reporting area is drained by the Colorado River and its tributaries below the mouth of the Green River and above Lee Ferry, Arizona. The largest of the tributary stream is the San Juan River which heads on the western slope of the Continental Divide in southwestern Colorado. Principal tributaries of the San Juan River are the Navajo,

Piedra, Los Pinos, Animas, and La Plata Rivers. The other main tributaries in the basin are the Dirty Devil, Escalante, and Paria Rivers, which drain a portion of the eastern slope of the Wasatch Plateau in Utah. The reporting area includes about 38,600 square miles in portions of Utah, New Mexico, Arizona, and Colorado.

The largest towns are Durango and Cortez in Colorado; Monticello and Blanding in Utah; Farmington in New Mexico; and Page in Arizona.

Mining and agriculture form the economic base for the San Juan-Colorado reporting area. The agricultural development is similar to that of the Upper Main Stem with most of the cropland devoted to livestock feeds but with production of diversified market crops on lands with favorable air drainage. The main market crops are fruit, vegetables, and dry beans. Oil, natural gas, and coal are the most important minerals produced. Thermal electric power production is increasingly important to the economy of the area.

Irrigation accounts for the largest use of water, about 78 percent of the San Juan reporting area use exclusive of any share of main stem evaporation. About 275,000 acres of land are irrigated in an average year.

Lower Colorado River Basin

Main Stem Below Lee Ferry, Arizona-California-Nevada

The Colorado River has a length of over 700 miles and a drainage area of

132,300 square miles within the Lower Colorado River System in the United States. The dividing point between the Upper and Lower Basin is Lee Ferry. Diversions are made at Lake Mead to the rapidly expanding North Las Vegas-Las Vegas-Henderson-Boulder City area for municipal and industrial purposes. The river below Lake Mead courses through canyons and broad alluvial valleys interspersed with bordering groups of mountains. Lakes Mohave and Havasu provide flood control and regulatory storage below Lake Mead. Lake Havasu also provides a forebay for pumped diversion to the Central Arizona Project (CAP) in Arizona and export to the Metropolitan Water District of Southern California. Lake Mohave reregulates Hoover Dam releases for power production and for deliveries to Mexico. Lesser structures downstream include Senator Wash, Laguna, Headgate Rock, Palo Verde, Imperial, and Morelos Dams. Senator Wash and Laguna Dams provide very limited amounts of reregulation capacity while the others are used principally for diversion.

Diversions below Lake Mead for agriculture, municipal and industrial, power, export, and other purposes are of the magnitude of 6 million acre-feet annually. A portion of these diversions is satisfied from upstream return flows. Yuma and Lake Havasu City in Arizona and Needles and Blythe in California are the major cities along the main stem below Lake Mead. Current irrigated lands adjacent to the main stem are estimated to cover approximately 275,000 acres.

Remaining Areas in Arizona-Nevada-Utah

Development away from the Colorado River main stem is limited by the availability of water and the rugged terrain.

Most of the irrigated lands in this area are located in the lower reach of the Virgin River and Las Vegas Valley in Nevada, on Kanab Creek in Arizona and Utah, and the lower portions of the Gila and Bill Williams Rivers in Arizona. Reporting period irrigated land averaged approximately 160,000 acres.

North Las Vegas, Las Vegas, Henderson, and Boulder City in Nevada, and Kingman and Williams in Arizona, are the leading cities.

Both the remaining areas in Arizona-Nevada-Utah and the main stem below Lee Ferry, Arizona-California-Nevada are combined into one subbasin, subbasin 6 on the frontispiece map, but reported separately.

Little Colorado River, Arizona-New Mexico

The Little Colorado River drainage area occupies a large part of northern Arizona and a portion of west-central New Mexico. It originates on the north slopes of the White Mountains about 20 miles above Springerville, Arizona. The river has a main stem length of about 356 miles and joins the Colorado River on the east boundary of Grand Canyon National Park about 78 miles downstream from Glen Canyon Dam.

A series of saline springs near the mouth of the Little Colorado River produces an estimated 160,000 acre-feet of water annually. The U.S. Geological Survey (USGS) gauging station near Cameron, Arizona, is located on the Navajo Indian Reservation about 45 miles upstream from the mouth. Streamflow is undependable and erratic and is subject to flash floods of considerable magnitude. Flow at the gauging station during the 1981-85 period varied from 41,150 acre-feet in 1981 to 323,300 acre-feet in 1983. Only minor development of the ground water has occurred because of low yields and poor quality. Excessive erosion and sediment deposition plague the area.

Agriculture is concentrated along the main stem of the Little Colorado River in the upper reaches of the river, on Silver Creek, a southern tributary, and on the Zuni River in New Mexico. Current irrigated lands in the basin are estimated to average approximately 32,500 acres. Irrigated acreage in the basin is subject to variation because of frequent water shortages and inadequate storage facilities. Population is predominately rural with a relatively large Indian segment. Principal cities include Flagstaff, Winslow, and Holbrook in Arizona, and Gallup, Zuni, and Pueblo in New Mexico. Leading industries include tourism, recreation, manufacturing, mining, and forest management.

Virgin River, Arizona-Utah

The Virgin River originates in western Kane County, Utah. It flows south-westerly through the southwest corner of

Utah and the northwestern corner of Arizona and empties into the northern extremity of the Overton Arm of Lake Mead in Nevada. The selected outflow point is the long-term USGS gauging station at Littlefield, Arizona, which is about 36 miles upstream from Lake Mead and about 10 miles above the Arizona-Nevada State line. The river is fed chiefly from tributaries heading in the southern high plateaus and mountains in Utah. Several springs contribute water to the river at a relatively uniform rate. The two most significant of these springs are located near LaVerkin, Utah, and Littlefield, Arizona, and both are highly saline. Agricultural and municipal developments in Nevada below the selected outflow point are included in "remaining areas," as shown on the frontispiece map.

The major irrigated areas are located in the LaVerkin-Hurricane-St. George-Santa Clara areas of Washington County, Utah, and in the Littlefield area of Mohave County, Arizona. Small irrigated areas are scattered throughout. Irrigated lands were estimated to average approximately 25,000 acres. Ground water has been developed to a limited degree. Population is predominately rural with St. George, Utah, being the principal city in the basin.

Muddy River, Nevada

The Muddy River, a tributary of the Virgin River prior to the existence of Lake Mead, originates from warm springs in northern Clark County, Nevada, about 10 miles northwest of Glendale. The river flows southeasterly

for about 30 miles and terminates at the northwestern extremity of the Overton Arm of Lake Mead near Overton, Nevada. Meadow Valley Wash, the major tributary of Muddy River, originates in northeastern Lincoln County and flows south to join the parent stream at Glendale. The USGS gauging station near Glendale is about 2.4 miles downstream from Meadow Valley Wash. Outflow varies little from year to year. Meadow Valley Wash, although perennial in the vicinity of Caliente, is normally dry in the last 50-mile reach above Glendale. Irrigated lands averaged approximately 10,800 acres. The entire basin is sparsely populated.

Bill Williams River, Arizona

The Bill Williams River is formed by the merger of the Big Sandy and Santa Maria Rivers about 7.5 miles above Alamo Dam. The river above Alamo Dam drains an area of about 4,700 square miles from small, rough mountain ranges and intervening valleys in parts of Mohave, La Paz, and Yavapai Counties. Alamo Dam and Reservoir, primarily a flood control structure completed in 1968, was built to protect downstream development along the Colorado River. A minimum pool is maintained for recreation and game management purposes. Releases up to a maximum of 2,000 ft³/s from the allocated conservation pool above the minimum pool are coordinated with releases from main stem reservoirs. Releases from Alamo Dam and runoff from the intervening area flow westerly and enter at the lower end of Lake Havasu just above Parker Dam.

Irrigated lands are estimated to average approximately 10,000 acres. The limited development in the basin is dominated by copper mining at the unincorporated town of Bagdad. A large portion of the water supply in the basin is obtained from ground-water pumpage. Releases from Alamo Dam and Reservoir during the 1981-85 period varied from 17,350 acre-feet in 1982 to 644,500 acre-feet in 1985.

Gila River, Arizona-New Mexico

The Gila River is the largest tributary to the Colorado River in the Lower Colorado River System. The drainage area extends from the Continental Divide in New Mexico to the river's mouth near Yuma, Arizona. Elevations in the basin range from nearly 12,000 feet in the eastern mountains to about 150 feet at the mouth. The selected outflow point for the basin is at Painted Rock Dam, a flood control structure located about 20 miles west of Gila Bend, Arizona. The drainage area above Painted Rock Dam is about 50,900 square miles, of which 5,600 square miles are in New Mexico, 44,200 square miles are in Arizona, and 1,100 square miles in Mexico. The dam was constructed to protect agricultural and urban developments downstream.

Nearly three-fourths of the population of the Lower Colorado River System reside in the Gila River Basin in the metropolitan Phoenix and Tucson areas. Industry and recreation play a large part in the economy. About two-thirds of the agricultural development in the Lower Colorado River System is located in the Gila River Basin. This development is

concentrated in the central area of Maricopa, Pinal, and Pima Counties and is supported to a large degree by a long-term overdraft of the ground-water resources. Estimated irrigated lands ranged between 755,000 and 1,228,000 acres for the reporting period and averaged 965,000 acres. Nearly all of the surface water resources in the basin have been developed for decades.

TERMINOLOGY

The Colorado River is not only one of the most highly controlled rivers in the world, but is also one of the most institutionally encompassed. A multitude of legal documents, known collectively as the "Law of the River," affect and dictate its management and operation. Major documents include:

- Colorado River Compact--1922
- Boulder Canyon Project Act--1928
- California Limitation Act--1929
- California Seven Party Agreement--1931
- Mexican Water Treaty--1944
- Upper Colorado River Basin Compact--1948
- Colorado River Storage Project Act--1956
- United States Supreme Court Decree in Arizona vs. California--1964
- Colorado River Basin Project Act--1968
- Minute 242 of the International Boundary and Water Commission, United States and Mexico--1973
- Colorado River Basin Salinity Control Act--1974, amended 1984

The Colorado River System is defined in the Colorado River Compact of 1922 as "...that portion of the Colorado River and its tributaries within the United States," whereas the Colorado River Basin is defined as "...all of the drainage area of

the Colorado River System and all other territory within the United States of America to which waters of the Colorado River System shall be beneficially applied." The compact divided the Colorado River Basin into two subbasins--the "Upper Basin" and the "Lower Basin," with Lee Ferry as the division point on the river. Lee Ferry, located in Arizona, is a point in the main stem 1 mile below the mouth of the Paria River. For the purpose of this report, the Great Divide Basin, a closed basin in Wyoming, and the White River, also a closed basin, in Nevada have not been considered as part of the Colorado River System since flows from these basins never reach the Colorado River. Diversions from the system to areas outside its drainage area are considered herein as exports and have not been classified by types of use.

Beneficial consumptive use is normally construed to mean the consumption of water brought about by human endeavors and in this report includes use of water for municipal, industrial, agricultural, power generation, export, recreation, fish and wildlife, and other purposes, along with the associated losses incidental to these uses.

The storage of water and water in transit may also act as losses on the system although normally such water is recoverable in time. Qualitatively, what constitutes beneficial consumptive use is fairly well understood; however, an inability to exactly quantify these uses has led to various differences of opinion. The practical necessity of administering the various water rights, apportionments, etc., of the Colorado River has led

to definitions of consumptive use or depletions generally in terms of "how it shall be measured." The Upper Colorado River Basin Compact provides that the Upper Colorado River Commission is to determine the apportionment made to each State by "...the inflow-outflow method in terms of manmade depletions of the virgin flow at Lee Ferry...." There is further provision that the measurement method can be changed by unanimous action of the Commission. In contrast, article 1(A) of the decree of the Supreme Court of the United States in *Arizona vs. California* defines, for the purpose of the decree, "Consumptive use means diversions from the stream less such return flows thereto as are available for consumptive use in the United States or in satisfaction of the Mexican Treaty obligation." Nearly all the water exported from the Upper Colorado River System is measured; however, the remaining beneficial consumptive use, for the most part, must be estimated using theoretical methods and techniques. In the Lower Colorado River System tributaries to the main stem, similar methods must be employed to determine the amount of water consumptively used.

Reservoir evaporation loss is a consumptive use associated with the beneficial use of water for other purposes. For the purpose of this report, main stem reservoir evaporation is carried as a separate item for the Upper and Lower Basins.

Channel losses within the system are normally construed to be the consumptive use by riparian vegetation along the stream channel (or conveyance route) and the evaporation from the stream's

water surface and wetted materials. Seepage from the stream normally appears again downstream or reaches a ground-water aquifer where it may be usable again. A decided lack of data and acceptable methodology along with the intermittent flow characteristics of many Southwest streams combine to make a reasonable determination of channel loss difficult. Channel losses have not been estimated for this report within the Upper Basin nor on the tributaries of the Lower Colorado River main stem. Channel losses on the main stem below Lee Ferry and below Hoover Dam were estimated.

METHODOLOGY AND DATA ADEQUACY

This report is based almost entirely on data obtained from ongoing programs and current reports. Quantitative measurements of water use were used wherever available, but the majority of the basin water use was theoretically calculated. The following sections describe these calculations for both the Lower Colorado River Main Stem and the Upper and Lower Colorado River Basin tributaries.

Colorado River Basin Tributaries

In the tributary areas of the basin, records of diversions and return flows are not complete enough to allow direct calculation of consumptive water use. Theoretical and indirect methods of estimating consumptive use must then be relied upon. In the New Mexico portion of the Gila River Basin, the

annual consumptive use of water is reported by the New Mexico Interstate Stream Commission, pursuant to article VII of the March 9, 1964, decree of the United States Supreme Court in *Arizona vs. California, et al.*

Agriculture

The percentages of irrigation consumptive use range between 64 and 69 percent for the Upper Basin tributaries and between 73 and 81 percent for the Lower Basin tributaries. Both percent ranges exclude main stem evaporation. The annual irrigated acreage of most crops grown within each reporting area was estimated from information published in the yearly State Agriculture Statistics. Irrigated pasture and some minor crops not reported by the State statistics were estimated from information in the 1978 and 1983 National Census of Agriculture with supporting information from the comprehensive framework study and various other local reports. The total irrigated acreage values for the Upper and Lower Basins are shown in tables [UC-7](#) and [LC-9](#), respectively.

Since most of these data were presented on a county basis, it was necessary to separate them into smaller reporting areas for computational purposes. This was accomplished using land inventory maps and relationships developed for the comprehensive framework study.

These subbasins generally follow tributary stream basin and State boundaries. A representative climatic station was selected for each subbasin. Using historical records of temperature,

precipitation, and frost dates, a consumptive use rate was computed for each major crop in each of the reporting years. For the purpose of this report, the consumptive use rates were computed using the modified Blaney-Criddle evapotranspiration formula in the version described in the Soil Conservation Service Technical Release No. 21, "Irrigation Water Requirements," revised September 1970. Irrigation consumptive rates were determined by subtracting the effective precipitation from the consumptive use rates. Effective precipitation for the Upper Basin was computed using the Soil Conservation Service method. This method is referenced in "SCS Technical Release No. 21." (It should be noted that this method estimates less effective precipitation than the Reclamation method. Previous reports used the Reclamation method of computing effective precipitation.) The values of irrigation consumptive use rates were applied to the estimates of irrigated acreage to yield the final values of irrigation consumptive use.

An exception to this procedure occurred in the Lower Basin in the "low desert" regions of Arizona and Nevada where the Blaney-Criddle formula was used to estimate the crop consumptive use. This departure was based on the research results of Leonard Erie, et al. Seasonal crop consumptive use factors ("K") for the lower elevation desert areas were selected from Conservation Research Report Number 29, "Consumptive Use of Water by Major Crops in the Southwestern United States," issued May 1982 by the United States Department of Agriculture. Effective precipitation was derived from criteria

developed for the area by Wayne D. Criddle, former Utah State Engineer.

These theoretical consumptive use calculations were based on the assumption of full water supply during the crop growing season. However, it is estimated that in an average year, about 37 percent of the irrigated lands in the Upper Basin receive less than a full supply of water, either due to lack of distribution facilities or junior water rights. The degree to which these lands suffer shortages varies widely from year to year, depending in large part on the magnitude of runoff. For this study, an estimate of the short supply service lands was made for each subbasin, primarily on the basis of reports and investigations collected for the comprehensive framework study. A streamflow gauging station was selected within each subbasin and the magnitude of the recessional portion of the annual hydrograph was used as an index to select the date at which consumptive use calculations should be terminated for the short supply lands.

Comprehensive framework studies of the incidental consumptive use of water associated with irrigation indicated that this use varied between 5 and 29 percent of the irrigation consumptive use, depending upon the location of the study area within the Colorado Basin. These percentages were used in the Upper Basin and an average value of 15 percent was used in the Lower Basin to adjust the calculated consumptive use.

The agricultural data is generally adequate for use in this report. With the exception of Utah, each state prepared annual county irrigated acreage

estimates of the harvested crops during the reporting period. These statistics are assumed to be reliable. The irrigated pasture values were based largely on the 1978 and 1983 National Census of Agriculture since the State statistics do not include pastureland. Because of the length of time between reporting dates, this item needs to be considerably strengthened. In this regard, Wyoming and New Mexico have initiated aerial photographic mapping of their State's irrigated acreage to verify the annual statistical sampling. Other areas of agricultural data collection that need to be updated and verified are (1) the consumptive water use of lands that receive less than a full seasonal supply of irrigation water and the areal extent of these lands, and (2) the amount of incidental seepage and phreatophytic losses associated with irrigation.

Reservoir Evaporation

A comprehensive listing was developed of all reservoirs in the Colorado River Basin which included the latitude, elevation, and surface area at total capacity for each reservoir.

Monthly content records were obtained for those reservoirs for which records are available. The average annual water-surface area was determined for each year of the reporting period. For those reservoirs lacking records, a "fullness factor" was estimated on the basis of reservoir use and historical hydrologic conditions. These "fullness factors" were then used to obtain estimates of average annual water-surface area for the unreported reservoirs. For the majority of the

basin, historical evaporation rates were used to determine reservoir evaporation.

In the Upper Basin, a multiple regression equation relating gross annual evaporation to elevation and latitude was developed. Adjustments to computed evaporation were made based on climatic subareas. In the Lower Basin, evaporation values were calculated for each climatic subarea. An account was taken of precipitation and runoff salvage to determine net evaporation rates. The net evaporation rates were applied to the estimates of average annual water-surface area to yield the values of annual reservoir evaporation.

An exception to this procedure was the determination of evaporation from what are called the main stem reservoirs shown in table UC-1. Predetermined average evaporation rates were applied to historical surface areas to yield values of evaporation on a monthly basis.

Ground Water

Currently, all ground-water pumpage is counted as consumptive use charged against the Colorado River Basin. Obviously, this is not necessarily true. Depending on the location and depth of the well and what types of soils are present in the area, it is possible that little or none of the water pumped would have contributed to the Colorado River System for hundreds or even thousands of years. It has recently been proposed that an interagency study team be put together consisting of personnel from various State Engineers Offices, Bureau

of Reclamation, and any other pertinent agencies. This study team would establish guidelines for computing what amounts of ground water pumped should be charged against the Colorado River Basin. These guidelines will need to be established on an area by area basis rather than one set percentage for the entire basin. Results of this study will be incorporated in future Consumptive Uses and Losses Reports. However, until these guidelines are established, the Consumptive Uses and Losses Reports will continue to report all ground-water pumpage as depletion from the system.

Currently, the Arizona portion of the Upper Basin is the only part of the basin that reports ground-water pumpage as consumptive use. Although significant ground-water usage occurs in Arizona, Nevada, and New Mexico, for purposes of this report ground-water overdraft has not been taken into account in the computation of tributary consumptive use. It should be noted that present ground-water overdraft in Arizona has been estimated to be approximately 2.2 million acre-feet per year.

Stockpond Evaporation and Livestock

Stockpond surface areas were estimated from the May 1975 Soil Conservation Service (SCS) publication, "Livestock Water Use." The subbasin stockpond areas were subdivided by State and basin using the livestock population distribution. The same procedure used to calculate the unmeasured reservoir evaporation was used to estimate the stockpond evaporation.

Livestock population data were taken from annual State Agriculture Statistics and the 1978 and 1983 Census of Agriculture. Livestock population data included cattle, sheep, horses, and hogs. Consumption rates for the various livestock were derived from various reports, including the SCS publication, "Livestock Water Use," May 1975.

Stockpond and livestock data are adequate to prepare an estimate of this use. Considering the small amount of water use, any refining effort would be best spent on the irrigation or evaporation categories.

Mineral Resources

Arizona leads the nation in the production of copper and the net water use for its production represents about 90 percent of the total water use for mineral resources in the Lower Basin. The Upper Basin uses water in the production of numerous minerals in addition to energy-related materials such as oil and natural gas.

Estimates of the water consumptively used were based largely on phone surveys conducted by the U.S. Geological Survey in 1985, that quantified water use in the basin. Intermediate years were interpolated between 1980 and 1985. In some cases where, for privacy reasons, companies were unwilling to supply information, information was obtained from the U.S. Bureau of Mines.

Thermal Electric Power

The net use of water for the production of thermal electric energy from the

tributaries of the Colorado River Basin was estimated from records obtained from the various power companies in the Basin. These records were complete and were judged to be accurate.

Municipal and Industrial

The basis for estimating municipal and industrial uses was the urban and rural population within the reporting areas. Preparation of annual population estimates was guided by the 1970 and 1980 censuses, various State and county statistical reviews, and reports that included population estimates for local areas. The yearly population estimates for the Upper and Lower Basins are shown in tables [UC-8](#) and [LC-10](#), respectively. Water withdrawal rates for urban and rural uses in the various reporting areas were derived from available studies in the metropolitan areas, State Water Plan reports, and Bureau of Reclamation technical guidelines. These withdrawals were then converted to depletions using average basin consumptive use factors.

The population of the Colorado River System, estimated at nearly 4.5 million in 1985, has increased at an annual rate slightly in excess of 3 percent during this reporting period. A large portion of the population resides within Maricopa and Pima Counties, Arizona, and in Clark County, Nevada. Sixty percent of the Upper Basin and about 20 percent of the Lower Basin population were classified as rural with a significantly smaller per capita use of water. Both the urban and rural areas have the mutual problem of providing an adequate current and future water supply for a growing population in a water-short area. As a

result of almost continuous studies concerning these problems, adequate production and effluent records are usually available to adequately assess water use.

Transbasin Diversions

Nearly all the transbasin diversions both out of and into the Colorado River System were measured and reported by the Geological Survey, or local water commissioners and users. The remainder were estimated on the basis of past records and capacity of facilities. Due to the high degree of measurement, this area of basin consumptive use is considered to be quite accurately determined.

Lower Colorado River Main Stem

The annual consumptive use of water from the Colorado River main stem by the States and exports from the system were taken from the Bureau annual report entitled "Compilation of Records in Accordance with Article V of the Decree of the United States in Arizona vs. California." The estimated Colorado River component of the combined surface and subsurface return flows accruing to Las Vegas Wash and discharging into Lake Mead, as taken from the report, is credited to Nevada's municipal and industrial water uses.

Unmeasured subsurface return flows were estimated below Hoover Dam, based partially on preliminary information supplied by the Task Force on Ground-Water Return Flows. All unmeasured subsurface return flows

were credited to the irrigation water use taken from the Article V report, and were divided between California and Arizona based on their respective irrigation diversions.

Gross evaporation from Lake Mead is estimated by the USGS and published in its annual Water Resources Data reports. Net evaporation for Lake Mead is estimated by subtracting precipitation at nearby Boulder City, Nevada, from the gross evaporation. Net evaporation from Lakes Mohave and Havasu and Senator Wash Reservoir was derived incorporating Lake Mead gross evaporation and local precipitation records and operating data.

Annual channel losses were estimated as the inflow or outflow necessary to balance a simplified water budget for the Lee Ferry to Hoover Dam and Hoover Dam to International Boundary reaches. Channel losses include evaporation, seepage, phreatophyte consumptive use, and bank storage.

The accuracy of flow measurements of the Colorado River mainstream for use in determining the channel losses values in table [LC-1](#) is in question. The gauge error of an "excellent" USGS flow gauge is 5 percent. Actual flow at Lee Ferry, Hoover Dam, or to Mexico may therefore vary by 90,000 acre-feet for the low flow entering Mexico in 1982 to over 1.1 million acre-feet for 1984 flow below Hoover. Such inaccuracies, though a very small percent of total flow, will have dramatic effects on apparent channel loss computations, as reflected by channel gains.

The annual land use, water supply, and water use information being gathered for

the operation, maintenance, and administration of the Colorado River main stem below Lee Ferry is believed to be generally adequate in quantity, quality, and extent. These data are under constant review and are being continually upgraded. Studies and programs are in progress to remedy a lack of data on return flows from main stem diversions.

BENEFICIAL CONSUMPTIVE USES AND LOSSES

A summary table of the Colorado River System total annual water uses, 1981 through 1985, by states and water flowing to Mexico is shown on page iv. Tables [C-2](#) through [C-6](#) show on a yearly basis the same information broken down by State, basin, and type of use. Water use within the selected reporting areas is discussed below.

Upper Colorado River Tributaries

Summaries of estimated annual consumptive uses and losses in the Upper Colorado River Basin for each of the reporting years, broken down by State, reporting area, and type of use are shown in tables [UC-2](#) through [UC-6](#). Estimated main stem reservoir evaporation is shown in table [UC-1](#). Technically, these are not all main stem reservoirs but are reservoirs that participate in the Colorado River Storage Project (CRSP). The Basin Commission designates which reservoirs in the CRSP have evaporation losses charged to the State and which have losses charged to

the basin as a whole. Reservoirs listed in table UC-1 are those to be charged to the basin as a whole. These reservoir evaporation losses amount to about 15 percent of all Upper Basin losses.

Upper Basin consumptive use varied between 3.9 million to 4.2 million and averaged 4.0 million acre-feet per year for the reporting period, 1981 through 1985. Agricultural uses accounted for about 57 percent of the total Upper Basin consumptive uses and losses. Irrigated acreages fluctuated very little during this period, ranging between 1.37 million acres to 1.45 million acres, and averaged 1.41 million acres per year. Variation in consumptive use during the reporting period was largely due to year-to-year changes in climatic conditions.

Transbasin exports, the second largest Upper Basin use, on the average accounted for 17 percent of basin use, showed year by year variation during the reporting period ranging from a high of 820,000 acre-feet in 1982 to a low of 577,000 acre-feet in 1983. Exports were reduced during 1983 and 1984 due to record amounts of winter precipitation in the Rocky Mountain region, which produced abundant water supplies on the Front Range of Colorado, and in the Great Basin of Utah.

Water uses for thermal electric power generation remained fairly constant, averaging about 120,000 acre-feet per year, which represents about 3 percent of consumptive use in the Upper Basin. Increases noted in past reports were primarily due to new powerplants coming on line, whereas there have been no new power projects constructed during this reporting period.

Lower Colorado River Main Stem

Table LC-1 shows main stem reservoir evaporation and apparent channel losses, and table LC-3 shows water uses along the lower Colorado River main stem and flood plain including water passing to Mexico. Water passing to Mexico is made up of deliveries in satisfaction of the Treaty, deliveries made pursuant to Minute No. 242, Gila River flood releases, regulatory waste and anticipatory flood control releases from the main stem. The latter three are combined as excess releases in table LC-3 and for this reporting period totaled slightly more than 37 million acre-feet. The previous 5-year reporting period had identified excess releases to Mexico totaling about 5.4 million acre-feet, with nearly 80 percent of this total occurring in 1980 and corresponding to completion of initial filling of the Upper Basin's Lake Powell.

Annual average main stem reservoir evaporation consumed approximately 1.0 million acre-feet.

Transbasin diversions continued to be the single highest consumer. For the current reporting period, as for the proceeding reporting period, transbasin diversions continued to account for approximately 72 percent of the Lower Colorado River main stem depletions.

Lower Colorado River Tributaries

Tables LC-4 through LC-8 show annual water uses within states by tributary and type of use. Lower Basin consumptive use was estimated to be about 5.2 million acre-feet in 1981,

3.7 million acre-feet in 1983, and 4.2 million acre-feet in 1985. The average for 1981-85 was about 4.4 million acre-feet.

Significant ground-water usage occurs in Arizona, Nevada, and New Mexico. For the purpose of this report, ground-water overdraft has not been taken into account in the computation of tributary consumptive use. Also, tributary channel loss and salvage were not evaluated. However, it should be noted that present ground-water overdraft in Arizona has been estimated to be approximately 2.2 million acre-feet per year.

Consumptive use for the irrigation of crops represents between 73 and 81 percent of the total water use in the Lower Colorado tributary areas. Estimated annual consumptive use for the Lower Basin during the 5-year period average about 3.2 acre-feet per acre, varying from approximately 1.1 acre-feet per acre in parts of New Mexico to over 5 acre-feet in the western portion of the basin. Estimated crop

consumptive use varied considerably from year to year on the basis of climatic conditions, and acreage fluctuations, from a high of 4.3 million acre-feet in 1981 to a low of 2.7 million acre-feet in 1983. Irrigated lands amounted to about 1.47 million acres in 1981 but had decreased to about 980,000 acres in 1983. The decrease in irrigated acreage occurred primarily in the Gila River tributary, according to State agriculture statistics.

The consumptive use of water for municipal and industrial purposes is estimated to have increased by about 44 percent for 1985 when compared with the ending year of the previous reporting period. Nearly 60 percent of the increase occurred in 1981.

Water supply conditions were characterized by near normal runoff in 1981, exceptionally poor runoff in 1982, and extremely large runoff in 1983, 1984, and 1985. In addition, the runoff in 1985 produced an outflow below Painted Rock Dam of 2,385,000 acre-feet.

**Table C-1.--Drainage area of the Colorado River System,
Area within each State and Mexico by major tributary streams**

Units = 1,000 square miles

Major tributary streams and their selected outflow points	Wyoming	Colorado	Utah	New Mexico	Arizona	Nevada	California	Total	Mexico
Green River at Colorado River confluence, Utah	17.1	10.6	17.1	-	-	-	-	44.8	-
Upper Main Stem at Green River confluence, Utah	-	22.2	4.0	-	-	-	-	26.2	-
San Juan-Colorado at Lee Ferry, Arizona	-	5.8	16.2	9.7	6.9	-	-	38.6	-
Little Colorado River near Cameron, Arizona	-	-	-	5.3	21.2	-	-	26.5	-
Virgin River at Little- field, Arizona	-	-	3.0	-	1.9	0.2	-	5.1	-
Muddy River near Glendale, Nevada	-	-	-	-	-	6.8	-	6.8	-
Bill Williams River below Alamo Dam, Arizona	-	-	-	-	4.7	-	-	4.7	-
Gila River below Painted Rock Dam, Arizona	-	-	-	5.6	44.2	-	-	49.8	(1.1)
Main stem and remaining areas in Lower Basin	-	-	0.6	-	28.3	6.9	3.6	39.4	(0.1)
Colorado River System at Southerly International Boundary	17.1	38.6	40.9	20.6	107.2	13.9	3.6	241.9	(1.2)
Colorado River System above Lee Ferry	17.1	38.6	37.3	9.7	6.9	-	-	109.6	-
Colorado River System below Lee Ferry	-	-	3.6	10.9	100.3	13.9	3.6	132.3	(1.2)

**Table C-2.--Summary of estimated water use by States and types of use
1981**

(1,000 acre-feet)

State	Estimated beneficial consumptive uses and losses ¹							Total
	Reservoir evaporation	Irrigated agriculture ²	Municipal and industrial ³	Fish and wildlife recreation	Export outside system	Export within system	Unmeasured return flow	
Arizona	279.7	5,452.6	513.2	53.9	0.0	3.0	-135	6,167.4
California	0.0	641.9	3.0	0.0	4,258.7	0.0	-65	4,838.6
Colorado	82.7	1,418.7	46.7	0.0	538.0	0.0	0	2,086.1
Nevada	4.2	96.8	294.5	1.2	0.0	-3.0	-43.6	350.1
New Mexico	44.3	190.5	52.0	0.9	54.6	0.0	0	342.3
Utah	65.0	577.0	40.7	0.0	99.5	0.0	0	782.2
Wyoming	41.6	241.7	50.7	0.0	7.0	0.0	0	341.0
Other ⁴	<u>1,598.4</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>3,996.4</u>	<u>0.0</u>	<u>0</u>	<u>5,594.8</u>
Colorado River System total	2,115.9	8,619.2	1,000.8	56.0	8,954.2	0.0	-243.6	20,502.5

¹ From tables UC-1, UC-2, LC-1, and LC-2.

² Includes livestock water use and stockpond evaporation.

³ Includes water uses for thermal electric power generation and mineral resources.

⁴ Reservoir evaporation represents main stem reservoir evaporation in the Upper Basin and Lower Basin and main stem channel losses for Lower Basin. Exports outside system represent water passing to Mexico.

**Table C-3.--Summary of estimated water use by States and types of use
1982**

(1,000 acre-feet)

State	Estimated beneficial consumptive uses and losses ¹							Total
	Reservoir evaporation	Irrigated agriculture ²	Municipal and industrial ³	Fish and wildlife recreation	Export outside system	Export within system	Unmeasured return flow	
Arizona	212.1	4,480.9	532.6	49.5	0.0	3.0	-135.4	5,142.7
California	0.0	572.2	3.1	0.0	3,838.4	0.0	-64.6	4,349.1
Colorado	78.4	1,378.5	44.5	0.0	604.7	0.0	0	2,106.1
Nevada	3.9	97.4	301.9	1.0	0.0	-3.0	-49.2	352.0
New Mexico	42.3	195.2	54.5	0.9	131.8	0.0	0	424.7
Utah	59.4	569.6	41.2	0.0	75.5	0.0	0	745.7
Wyoming	39.2	228.7	50.8	0.0	11.1	0.0	0	329.8
Other ⁴	<u>1,402.9</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>1,816.8</u>	<u>0.0</u>	<u>0</u>	<u>3,219.7</u>
Colorado River System total	1,838.2	7,522.5	1,028.6	51.4	6,478.3	0.0	-249.2	16,669.8

¹ From tables UC-1, UC-3, LC-1, and LC-2.

² Includes livestock water use and stockpond evaporation.

³ Includes water uses for thermal electric power generation and mineral resources.

⁴ Reservoir evaporation represents main stem reservoir evaporation in the Upper Basin and Lower Basin and main stem channel losses for Lower Basin. Exports outside the system represents water passing to Mexico.

**Table C-4.--Summary of estimated water use by States and types of use
1983**

(1,000 acre-feet)

State	Estimated beneficial consumptive uses and losses ¹							Total
	Reservoir evaporation	Irrigated agriculture ²	Municipal and industrial ³	Fish and wildlife recreation	Export outside system	Export within system	Unmeasured return flow	
Arizona	280.4	3,548.2	517.2	30.6	0.0	3.0	-142.2	4,237.2
California	0.0	409.4	2.5	0.0	3,598.5	0.0	-57.8	3,952.6
Colorado	82.3	1,373.7	40.1	0.0	423.4	0.0	0	1,919.5
Nevada	4.6	91.2	311.6	1.1	0.0	-3.0	-66.2	339.3
New Mexico	39.0	198.8	55.6	0.9	131.2	0.0	0	425.5
Utah	63.0	587.1	43.1	0.0	24.9	0.0	0	718.1
Wyoming	36.7	257.6	45.2	0.0	6.6	0.0	0	346.1
Other ⁴	<u>1,896.2</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>9,781.8</u>	<u>0.0</u>	<u>0</u>	<u>11,678.0</u>
Colorado River System total	2,402.2	6,466.0	1,015.3	32.6	13,966.4	0.0	-266.2	23,616.3

¹ From tables UC-1, UC-4, LC-1, and LC-2.

² Includes livestock water use and stockpond evaporation.

³ Includes water uses for thermal electric power generation and mineral resources.

⁴ Reservoir evaporation represents main stem reservoir evaporation in the Upper Basin and Lower Basin and main stem channel losses for Lower Basin. Exports outside the system represents water passing to Mexico.

**Table C-5.--Summary of estimated water use by States and types of use
1984**

(1,000 acre-feet)

State	Estimated beneficial consumptive uses and losses ¹							Total
	Reservoir evaporation	Irrigated agriculture ²	Municipal and industrial ³	Fish and wildlife recreation	Export outside system	Export within system	Unmeasured return flow	
Arizona	274.5	4,344.9	555.7	5.8	0.0	3.0	-143	5,040.9
California	0.0	446.3	3.0	0.0	4,286.5	0.0	-57	4,678.8
Colorado	84.2	1,298.0	38.8	0.0	443.7	0.0	0	1,864.7
Nevada	5.1	95.4	323.6	1.0	0.0	-3.0	-66.9	355.2
New Mexico	44.9	199.7	58.0	0.9	114.1	0.0	0	417.6
Utah	64.7	630.0	44.0	0.0	23.3	0.0	0	762.0
Wyoming	34.5	229.4	39.4	0.0	4.0	0.0	0	307.3
Other ⁴	<u>1,197.1</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>16,991.3</u>	<u>0.0</u>	<u>0</u>	<u>18,188.4</u>
Colorado River System total	1,705.0	7,243.7	1,062.5	7.7	21,862.9	0.0	-266.9	31,614.9

¹ From tables UC-1, UC-5, LC-1, and LC-2.

² Includes livestock water use and stockpond evaporation.

³ Includes water uses for thermal electric power generation and mineral resources.

⁴ Reservoir evaporation represents main stem reservoir evaporation in the Upper Basin and Lower Basin and main stem channel losses for Lower Basin. Exports outside the system represents water passing to Mexico.

**Table C-6.--Summary of estimated water use by States and types of use
1985**

(1,000 acre-feet)

State	Estimated beneficial consumptive uses and losses ¹							Total
	Reservoir evaporation	Irrigated agriculture ²	Municipal and industrial ³	Fish and wildlife recreation	Export outside system	Export within system	Unmeasured return flow	
Arizona	266.2	4,120.0	585.2	6.0	0.0	3.0	-142.3	4,838.1
California	0.0	455.9	3.3	0.0	4,308.6	0.0	-57.7	4,710.1
Colorado	86.3	1,359.4	38.5	0.0	509.6	0.0	0.0	1,993.8
Nevada	4.7	102.6	336.0	1.0	0.0	-3.0	-69.2	372.1
New Mexico	38.4	211.0	58.1	0.9	92.8	0.0	0	401.2
Utah	69.0	705.5	45.3	0.0	59.6	0.0	0	879.4
Wyoming	30.5	250.4	44.2	0.0	11.3	0.0	0	336.4
Other ⁴	<u>1,782.9</u>	<u>0.0</u>	<u>0.0</u>	<u>0.0</u>	<u>13,395.9</u>	<u>0.0</u>	<u>0</u>	<u>15,178.8</u>
Colorado River System total	2,278.0	7,204.8	1,110.6	7.9	18,377.8	0.0	-269.2	28,709.9

¹ From tables UC-1, UC-6, LC-1, and LC-2.

² Includes livestock water use and stockpond evaporation.

³ Includes water uses for thermal electric power generation and mineral resources.

⁴ Reservoir evaporation represents main stem reservoir evaporation in the Upper Basin and Lower Basin and main stem channel losses for Lower Basin. Exports outside the system represents water passing to Mexico.

**Table UC-1.--Upper Colorado River Basin
Estimated main stem reservoir evaporation¹**

1981-1985

Reservoir	(1,000 acre-feet)					
	Evaporation					Average
	1981	1982	1983	1984	1985	
Flaming Gorge	74.2	72.5	81.1	81.4	82.2	78.3
Blue Mesa	8.2	9.0	12.4	9.1	9.7	9.7
Morrow Point	1.1	1.0	1.2	0.9	1.0	1.0
Lake Powell	<u>457.9</u>	<u>418.4</u>	<u>519.5</u>	<u>567.7</u>	<u>552.0</u>	<u>503.1</u>
Total	541.4	500.9	614.2	659.1	644.9	592.1

¹ Undistributed by States. Evaporation determined using average historical evaporation rates.

**Table UC-2.--Upper Colorado River Basin
Estimated water use within States, by major tributaries, and types of use
1981**

(1,000 acre-feet)												
State	Tributary	Reservoir ¹ evaporation	Agriculture			Municipal & industrial				Export		Totals
			Irri- gation	Stockpond evaporation & livestock	Total	Mineral resources	Thermal electric power	Urban rural & industrial	Total	Outside system	Within system	
Arizona	San Juan-Colo.	5.6	4.6	3.5	8.1	0.4	21.1	6.6	28.1	0.0	0.0	41.8
Colorado	Green River	6.5	154.9	3.7	158.6	4.0	12.7	1.9	18.6	0.0	0.0	183.7
	Upper Main Stem	67.4	1,059.4	7.4	1,066.8	9.9	0.1	13.1	23.1	535.2	120.9	1,813.4
	San Juan-Colo.	<u>8.8</u>	<u>187.3</u>	<u>6.0</u>	<u>193.3</u>	<u>2.0</u>	<u>0.0</u>	<u>3.0</u>	<u>5.0</u>	<u>2.8</u>	<u>-120.9</u>	<u>89.0</u>
	Subtotal	82.7	1,401.6	17.1	1,418.7	15.9	12.8	18.0	46.7	538.0	0.0	2,086.1
New Mexico	San Juan-Colo.	39.3	175.7	3.9	179.6	4.2	28.1	9.5	41.8	53.9	0.0	314.6
Utah	Green River	51.3	409.1	4.8	413.9	6.8	20.1	5.6	32.5	103.8	0.0	601.5
	Upper Main Stem	0.4	9.1	0.1	9.2	1.6	0.0	0.9	2.5	0.0	0.0	12.1
	San Juan-Colo.	<u>8.7</u>	<u>42.9</u>	<u>4.3</u>	<u>47.2</u>	<u>1.6</u>	<u>0.0</u>	<u>1.1</u>	<u>2.7</u>	<u>-5.7</u>	<u>0.0</u>	<u>52.9</u>
	Subtotal	60.4	461.1	9.2	470.3	10.0	20.1	7.6	37.7	98.1	0.0	666.5
Wyoming	Green River	41.6	237.1	4.6	241.7	16.4	28.6	5.7	50.7	7.0	0.0	341.0
Upper Basin	Green River	99.4	801.1	13.1	814.2	27.2	61.4	13.2	101.8	110.8	0.0	1,126.3
Totals	Upper Main Stem	67.8	1,068.5	7.5	1,076.0	11.5	0.1	14.0	25.6	535.2	120.9	1,825.5
	San Juan-Colo.	<u>62.4</u>	<u>410.5</u>	<u>17.7</u>	<u>428.2</u>	<u>8.2</u>	<u>49.2</u>	<u>20.2</u>	<u>77.6</u>	<u>51.0</u>	<u>-120.9</u>	<u>498.3</u>
	Subtotal	229.6	2,280.1	38.3	2,318.4	46.9	110.7	47.4	205.0	697.0	0.0	3,450.0

¹ Excludes reservoir evaporation from Colorado River main stem reservoirs listed on table UC-1.

**Table UC-3.--Upper Colorado River Basin
Estimated water use within States, by major tributaries, and types of use
1982**

(1,000 acre-feet)

State	Tributary	Reservoir ¹ evaporation	Agriculture			Municipal & industrial				Export		Totals
			Irri- gation	Stockpond evaporation & livestock	Total	Mineral resources	Thermal electric power	Urban rural & industrial	Total	Outside system	Within system	
Arizona	San Juan-Colo.	4.3	5.7	3.4	9.1	0.4	19.4	6.6	26.4	0.0	0.0	39.8
Colorado	Green River	6.5	140.8	3.6	144.4	3.1	13.2	1.9	18.2	0.0	0.0	169.1
	Upper Main Stem	63.2	1,016.9	7.2	1,024.1	7.7	0.2	13.8	21.7	601.7	156.1	1,866.8
	San Juan-Colo.	<u>8.7</u>	<u>204.1</u>	<u>5.9</u>	<u>210.0</u>	<u>1.5</u>	<u>0.0</u>	<u>3.1</u>	<u>4.6</u>	<u>3.0</u>	<u>-156.1</u>	<u>70.2</u>
	Subtotal	78.4	1,361.8	16.7	1,378.5	12.3	13.4	18.8	44.5	604.7	0.0	2,106.1
New Mexico	San Juan-Colo.	37.6	179.4	4.0	183.4	3.9	34.2	9.1	47.2	131.0	0.0	399.2
Utah	Green River	46.6	403.5	4.5	408.0	6.0	20.0	6.1	32.1	78.7	0.0	565.4
	Upper Main Stem	0.4	7.8	0.1	7.9	1.7	0.0	1.0	2.7	0.0	0.0	11.0
	San Juan-Colo.	<u>8.1</u>	<u>46.8</u>	<u>4.2</u>	<u>51.0</u>	<u>2.0</u>	<u>0.0</u>	<u>1.2</u>	<u>3.2</u>	<u>-6.0</u>	<u>0.0</u>	<u>56.3</u>
	Subtotal	55.1	458.1	8.8	466.9	9.7	20.0	8.3	38.0	72.7	0.0	632.7
Wyoming	Green River	39.2	224.3	4.4	228.7	13.9	31.0	5.9	50.8	11.1	0.0	329.8
Upper Basin	Green River	92.3	768.6	12.5	781.1	23.0	64.2	13.9	101.1	89.8	0.0	1,064.3
Totals	Upper Main Stem	63.6	1,024.7	7.3	1,032.0	9.4	0.2	14.8	24.4	601.7	156.1	1,877.8
	San Juan-Colo.	<u>58.7</u>	<u>436.0</u>	<u>17.5</u>	<u>453.5</u>	<u>7.8</u>	<u>53.6</u>	<u>20.0</u>	<u>81.4</u>	<u>128.0</u>	<u>-156.1</u>	<u>565.5</u>
	Subtotal	214.6	2,229.3	37.3	2,266.6	40.2	118.0	48.7	206.9	819.5	0.0	3,507.6

¹ Excludes reservoir evaporation from Colorado River main stem reservoirs listed on table UC-1.

**Table UC-4.--Upper Colorado River Basin
Estimated water use within States, by major tributaries, and types of use
1983**

(1,000 acre-feet)												
State	Tributary	Reservoir ¹ Evaporation	Agriculture			Municipal & Industrial				Export		Totals
			Irri- gation	Stockpond Evaporation & Livestock	Total	Mineral Resources	Thermal Electric Power	Urban Rural & Industrial	Total	Outside Within		
										System	System	
Arizona	San Juan-Colo.	4.0	6.5	3.5	10.0	0.4	20.2	7.0	27.6	0.0	0.0	41.6
Colorado	Green River	6.5	138.2	3.6	141.8	2.2	11.8	1.9	15.9	0.0	0.0	164.2
	Upper Main Stem	67.2	1,009.5	7.2	1,016.7	5.2	0.2	14.6	20.0	420.0	123.5	1,647.4
	San Juan-Colo.	<u>8.6</u>	<u>209.3</u>	<u>5.9</u>	<u>215.2</u>	<u>1.0</u>	<u>0.0</u>	<u>3.2</u>	<u>4.2</u>	<u>3.4</u>	<u>-123.5</u>	<u>107.9</u>
	Subtotal	82.3	1,357.0	16.7	1,373.7	8.4	12.0	19.7	40.1	423.4	0.0	1,919.5
New Mexico	San Juan-Colo.	34.3	185.2	4.0	189.2	5.4	33.2	8.5	47.1	130.3	0.0	400.9
Utah	Green River	49.2	407.3	4.5	411.8	5.2	21.4	6.6	33.2	20.0	0.0	514.2
	Upper Main Stem	0.4	9.1	0.1	9.2	1.8	0.0	1.1	2.9	0.0	0.0	12.5
	San Juan-Colo.	<u>8.3</u>	<u>58.8</u>	<u>4.2</u>	<u>63.0</u>	<u>2.4</u>	<u>0.0</u>	<u>1.2</u>	<u>3.6</u>	<u>-5.1</u>	<u>0.0</u>	<u>69.8</u>
	Subtotal	57.9	475.2	8.8	484.0	9.4	21.4	8.9	39.7	14.9	0.0	596.5
Wyoming	Green River	36.7	253.2	4.4	257.6	11.3	27.7	6.2	45.2	6.6	0.0	346.1
Upper Basin Totals	Green River	92.4	798.7	12.5	811.2	18.7	60.9	14.7	94.3	26.6	0.0	1,024.5
	Upper Main Stem	67.6	1,018.6	7.3	1,025.9	7.0	0.2	15.7	22.9	420.0	123.5	1,659.9
	San Juan-Colo.	<u>55.2</u>	<u>459.8</u>	<u>17.6</u>	<u>477.4</u>	<u>9.2</u>	<u>53.4</u>	<u>19.9</u>	<u>82.5</u>	<u>128.6</u>	<u>-123.5</u>	<u>620.2</u>
	Subtotal	215.2	2,277.1	37.4	2,314.5	34.9	114.5	50.3	199.7	575.2	0.0	3,304.6

¹ Excludes reservoir evaporation from Colorado River main stem reservoirs listed on table UC-1.

**Table UC-5.--Upper Colorado River Basin
Estimated water use within States, by major tributaries, and types of use
1984**

(1,000 acre-feet)												
State	Tributary	Reservoir ¹ evaporation	Agriculture			Municipal & industrial				Export		Totals
			Irri- gation	Stockpond evaporation & livestock	Total	Mineral resources	Thermal electric power	Urban rural & industrial	Total	Outside system	Within system	
Arizona	San Juan-Colo.	4.9	5.8	3.3	9.1	0.4	22.1	7.5	30.0	0.0	0.0	44.0
Colorado	Green River	6.5	134.7	3.6	138.3	1.3	13.1	1.9	16.3	0.0	0.0	161.1
	Upper Main Stem	69.0	922.5	7.2	929.7	3.2	0.0	15.4	18.6	440.0	137.7	1,595.0
	San Juan-Colo.	<u>8.7</u>	<u>224.1</u>	<u>5.9</u>	<u>230.0</u>	<u>0.6</u>	<u>0.0</u>	<u>3.3</u>	<u>3.9</u>	<u>3.7</u>	<u>-137.7</u>	<u>108.6</u>
	Subtotal	84.2	1,281.3	16.7	1,298.0	5.1	13.1	20.6	38.8	443.7	0.0	1,864.7
New Mexico	San Juan-Colo.	40.1	186.7	4.0	190.7	4.2	37.7	8.2	50.1	113.6	0.0	394.5
Utah	Green River	50.3	438.7	4.4	443.1	4.4	21.9	7.0	33.3	26.8	0.0	553.5
	Upper Main Stem	0.4	10.1	0.1	10.2	1.8	0.0	1.2	3.0	0.0	0.0	13.6
	San Juan-Colo.	<u>8.4</u>	<u>60.3</u>	<u>4.3</u>	<u>64.6</u>	<u>2.8</u>	<u>0.0</u>	<u>1.3</u>	<u>4.1</u>	<u>-6.1</u>	<u>0.0</u>	<u>71.0</u>
	Subtotal	59.1	509.1	8.8	517.9	9.0	21.9	9.5	40.4	20.7	0.0	638.1
Wyoming	Green River	34.5	225.2	4.2	229.4	8.7	24.3	6.4	39.4	4.0	0.0	307.3
Upper Basin	Green River	91.3	798.6	12.2	810.8	14.4	59.3	15.3	89.0	30.8	0.0	1,021.9
Totals	Upper Main Stem	69.4	932.6	7.3	939.9	5.0	0.0	16.6	21.6	440.0	137.7	1,608.6
	San Juan-Colo.	<u>62.1</u>	<u>476.9</u>	<u>17.5</u>	<u>494.4</u>	<u>8.0</u>	<u>59.8</u>	<u>20.3</u>	<u>88.1</u>	<u>111.2</u>	<u>-137.7</u>	<u>618.1</u>
	Subtotal	222.8	2,208.1	37.0	2,245.1	27.4	119.1	52.2	198.7	582.0	0.0	3,248.6

¹ Excludes reservoir evaporation from Colorado River main stem reservoirs listed on table UC-1.

**Table UC-6.--Upper Colorado River Basin
Estimated water use within States, by major tributaries, and types of use
1985**

(1,000 acre-feet)

State	Tributary	Reservoir ¹ evaporation	Agriculture			Municipal & industrial			Export		Totals	
			Irri- gation	Stockpond evaporation & livestock	Total	Mineral resources	Thermal electric power	Urban rural & industrial	Total	Outside system		Within system
Arizona	San Juan-Colo.	4.9	5.8	2.5	8.3	0.4	23.2	7.2	30.8	0.0	0.0	44.0
Colorado	Green River	6.5	129.8	3.5	133.3	0.3	15.6	1.9	17.8	0.0	0.0	157.6
	Upper Main Stem	71.1	1,011.3	7.0	1,018.3	0.9	0.0	16.2	17.1	505.3	140.7	1,752.5
	San Juan-Colo.	<u>8.7</u>	<u>202.0</u>	<u>5.8</u>	<u>207.8</u>	<u>0.2</u>	<u>0.0</u>	<u>3.4</u>	<u>3.6</u>	<u>4.3</u>	<u>-140.7</u>	<u>83.7</u>
	Subtotal	86.3	1,343.1	16.3	1,359.4	1.4	15.6	21.5	38.5	509.6	0.0	1,993.8
New Mexico	San Juan-Colo.	33.2	198.6	4.0	202.6	5.6	34.0	7.9	47.5	91.8	0.0	375.1
Utah	Green River	55.3	510.7	4.4	515.1	3.5	22.6	7.5	33.6	63.6	0.0	667.6
	Upper Main Stem	0.4	9.6	0.1	9.7	1.9	0.0	1.3	3.2	0.0	0.0	13.3
	San Juan-Colo.	<u>8.1</u>	<u>63.5</u>	<u>4.0</u>	<u>67.5</u>	<u>3.3</u>	<u>0.0</u>	<u>1.3</u>	<u>4.6</u>	<u>-6.1</u>	<u>0.0</u>	<u>74.1</u>
	Subtotal	63.8	583.8	8.5	592.3	8.7	22.6	10.1	41.4	57.5	0.0	755.0
Wyoming	Green River	30.5	245.9	4.5	250.4	6.2	31.3	6.7	44.2	11.3	0.0	336.4
Upper Basin Totals	Green River	92.3	886.4	12.4	898.8	10.0	69.5	16.1	95.6	74.9	0.0	1,161.6
	Upper Main Stem	71.5	1,020.9	7.1	1,028.0	2.8	0.0	17.5	20.3	505.3	140.7	1,765.8
	San Juan-Colo.	<u>54.9</u>	<u>469.9</u>	<u>16.3</u>	<u>486.2</u>	<u>9.5</u>	<u>57.2</u>	<u>19.8</u>	<u>86.5</u>	<u>90.0</u>	<u>-140.7</u>	<u>576.9</u>
	Subtotal	218.7	2,377.2	35.8	2,413.0	22.3	126.7	53.4	202.4	670.2	0.0	3,504.3

¹ Excludes reservoir evaporation from Colorado River main stem reservoirs listed on table UC-1.

**Table UC-7.--Upper Colorado River Basin
Irrigated acreage
1981-1985**

		(1,000 acres)				
State	Tributary	Irrigated acreage				
		1981	1982	1983	1984	1985
Arizona	San Juan-Colo.	5.4	5.4	5.4	3.5	3.6
Colorado	Green River	118.7	107.5	109.3	107.5	101.2
	Upper Main Stem	575.6	556.5	563.5	527.7	528.9
	San Juan-Colo.	<u>142.4</u>	<u>147.3</u>	<u>145.9</u>	<u>141.0</u>	<u>137.6</u>
	State total	836.1	811.3	818.7	776.2	767.7
New Mexico	San Juan-Colo.	93.8	95.4	93.5	89.9	92.9
Utah	Green River	210.9	209.1	215.5	224.0	235.6
	Upper Main Stem	4.4	4.4	4.4	4.5	5.0
	San Juan-Colo.	<u>34.8</u>	<u>32.5</u>	<u>33.7</u>	<u>35.9</u>	<u>36.4</u>
	State total	250.1	246.0	253.6	264.4	277.0
Wyoming	Green River	250.0	234.1	278.7	236.6	246.7
Upper Basin	Green River	579.0	550.7	603.5	568.1	583.5
	Upper Main Stem	580.0	560.9	567.9	532.2	533.9
	San Juan-Colo.	<u>276.4</u>	<u>280.6</u>	<u>278.5</u>	<u>270.3</u>	<u>270.5</u>
	Total	1,435	1,392	1,450	1,371	1,388

**Table UC-8.--Upper Colorado River Basin
Population estimates
1981-1985**

		(1,000's)				
State	Tributary	Estimated population				
		1981	1982	1983	1984	1985
Arizona	San Juan-Colo.	41.3	42.3	43.2	44.2	45.1
Colorado	Green River	32.8	32.9	33.0	33.1	33.2
	Upper Main Stem	213.6	219.3	225.0	230.7	236.4
	San Juan-Colo.	<u>49.4</u>	<u>50.6</u>	<u>51.8</u>	<u>53.1</u>	<u>54.3</u>
	State total	295.8	302.8	309.8	316.9	323.9
New Mexico	San Juan River	99.6	102.5	105.4	108.3	108.1
Utah	Green River	68.7	70.1	71.5	73.0	74.4
	Upper Main Stem	8.0	7.8	7.5	7.3	7.1
	San Juan-Colo.	<u>16.5</u>	<u>16.6</u>	<u>16.7</u>	<u>16.9</u>	<u>17.0</u>
	State total	93.2	94.5	95.7	97.2	98.5
Wyoming	Green River	58.7	61.4	64.1	66.8	69.4
Upper Basin	Green River	160.2	164.4	168.6	172.9	177.0
	Upper Main Stem	221.6	227.1	232.5	238.0	243.5
	San Juan-Colo.	<u>206.8</u>	<u>212.0</u>	<u>217.1</u>	<u>222.5</u>	<u>224.5</u>
	Total	588.6	603.5	618.2	633.4	645.0

**Table LC-1.--Lower Colorado River Basin
Colorado River main stem estimated reservoir evaporation and channel loss¹
1981-1985**

(1,000 acre-feet)										
Water year	Reservoir evaporation						Apparent channel losses ³			Total reservoir evaporation and channel loss
	Lake Mead	Lake Mohave	Lake Havasu	Senator Wash Reservoir	Other ²	Total	Lee Ferry to Hoover Dam	Hoover Dam to IB	Total	
1981	722	139	106	2	38	1,007	130	(-80)	50	1,057
1982	716	131	99	2	34	982	(-50)	(-30)	(-80)	902
1983	833	133	100	2	34	1,102	(-100)	280	180	1,282
1984	734	128	99	2	35	998	(-350)	(-110)	(-460)	538
1985	<u>793</u>	<u>138</u>	<u>100</u>	<u>2</u>	<u>35</u>	<u>1,068</u>	<u>(-10)</u>	<u>80</u>	<u>70</u>	<u>1,138</u>
Average	759	134	101	2	35	1,031	(-76)	28	(-48)	983

¹ Undistributed by States.

² "Other" impoundments include Palo Verde, Headgate Rock, Imperial, Laguna, and Morelos Diversion Dams.

³ Losses include channel evaporation, seepage, and phreatophyte consumptive use. Note that gauge error of a USGS "excellent" gauge is 5 percent, or approximately 400,000 acre-feet per year for the gauges at Lee Ferry, Hoover Dam, and the International Border. Such gauge errors may result in the computation of large apparent channel losses.

**Table LC-2.--Lower Colorado River Basin
Estimated water use including Colorado main stem by States and types of use¹
1981-1985**

(1,000 acre-feet)

Water year	State	Reservoir evaporation	Agriculture			Municipal and industrial				Fish, wild- life, and recreation	Export		Total
			Irrigation	Stockpond evaporation and livestock	Total	Mineral resources	Thermal electric power	Other ²	Total		Outside system	Inside system	
1981	Arizona	274.1	5,395.9	48.6	5,444.5	61.4	38.4	385.3	485.1	53.9	0.0	3.0	6,260.6
	Nevada	4.2	95.1	1.7	96.8	3.5	22.7	268.3	294.5	1.2	0.0	-3.0	393.7
	New Mexico	5.0	8.9	2.9	11.8	7.6	0.0	2.6	10.2	0.0	0.7	0.0	27.7
	Utah	4.6	105.4	1.3	106.7	0.0	0.0	3.0	3.0	0.0	1.4	0.0	115.7
	California	<u>0.0</u>	<u>641.9</u>	<u>0.0</u>	<u>641.9</u>	<u>0.0</u>	<u>0.0</u>	<u>3.0</u>	<u>3.0</u>	<u>0.0</u>	<u>4,258.7</u>	<u>0.0</u>	<u>4,903.6</u>
	Total	287.9	6,247.2	54.5	6,301.7	72.5	61.1	662.2	795.8	55.1	4,260.8	0.0	11,701.3
1982	Arizona	207.8	4,429.5	42.3	4,471.8	79.3	32.2	394.7	506.2	49.5	0.0	3.0	5,238.3
	Nevada	3.9	95.8	1.6	97.4	2.4	28.0	271.5	301.9	1.0	0.0	-3.0	401.2
	New Mexico	4.7	9.5	3.2	12.7	4.7	0.0	2.6	7.3	0.0	0.8	0.0	25.5
	Utah	4.3	101.5	1.2	102.7	0.0	0.0	3.2	3.2	0.0	2.8	0.0	113.0
	California	<u>0.0</u>	<u>572.2</u>	<u>0.0</u>	<u>572.2</u>	<u>0.0</u>	<u>0.0</u>	<u>3.1</u>	<u>3.1</u>	<u>0.0</u>	<u>3,838.4</u>	<u>0.0</u>	<u>4,413.7</u>
	Total	220.7	5,208.5	48.3	5,256.8	86.4	60.2	675.1	821.7	50.5	3,842.0	0.0	10,191.7
1983	Arizona	276.4	3,498.7	39.5	3,538.2	54.1	25.8	409.7	489.6	30.6	0.0	3.0	4,337.9
	Nevada	4.6	89.5	1.7	91.2	2.3	25.7	283.6	311.6	1.1	0.0	-3.0	405.5
	New Mexico	4.7	7.4	3.1	10.5	5.8	0.0	2.7	8.5	0.0	0.9	0.0	24.6
	Utah	5.1	101.8	1.3	103.1	0.0	0.0	3.4	3.4	0.0	10.0	0.0	121.6
	California	<u>0.0</u>	<u>409.4</u>	<u>0.0</u>	<u>409.4</u>	<u>0.0</u>	<u>0.0</u>	<u>2.5</u>	<u>2.5</u>	<u>0.0</u>	<u>3,598.5</u>	<u>0.0</u>	<u>4,010.4</u>
	Total	290.8	4,106.8	45.6	4,152.4	62.3	51.5	700.8	814.6	31.7	3,609.4	0.0	8,898.9
1984	Arizona	269.6	4,297.8	38.0	4,335.8	59.9	33.0	432.8	525.7	5.8	0.0	3.0	5,139.9
	Nevada	5.1	93.6	1.8	95.4	2.3	26.3	295.0	323.6	1.0	0.0	-3.0	422.1
	New Mexico	4.8	6.7	3.2	9.9	5.3	0.0	2.6	7.9	0.0	0.5	0.0	23.1
	Utah	5.6	110.8	1.3	112.1	0.0	0.0	3.6	3.6	0.0	2.6	0.0	123.9
	California	<u>0.0</u>	<u>446.3</u>	<u>0.0</u>	<u>446.3</u>	<u>0.0</u>	<u>0.0</u>	<u>3.0</u>	<u>3.0</u>	<u>0.0</u>	<u>4,286.5</u>	<u>0.0</u>	<u>4,735.8</u>
	Total	285.1	4,955.2	44.3	4,999.5	67.5	59.3	737.0	863.8	6.8	4,289.6	0.0	10,444.8
1985	Arizona	261.3	4,065.7	46.0	4,111.7	62.9	34.7	456.8	554.4	6.0	0.0	3.0	4,936.4
	Nevada	4.7	100.9	1.7	102.6	2.3	19.8	313.9	336.0	1.0	0.0	-3.0	441.3
	New Mexico	5.2	6.4	2.9	9.3	7.8	0.0	2.8	10.6	0.0	1.0	0.0	26.1
	Utah	5.2	111.9	1.3	113.2	0.0	0.0	3.9	3.9	0.0	2.1	0.0	124.4
	California	<u>0.0</u>	<u>455.9</u>	<u>0.0</u>	<u>455.9</u>	<u>0.0</u>	<u>0.0</u>	<u>3.3</u>	<u>3.3</u>	<u>0.0</u>	<u>4,308.6</u>	<u>0.0</u>	<u>4,767.8</u>
	Total	276.4	4,740.8	51.9	4,792.7	73.0	54.5	780.7	908.2	7.0	4,311.7	0.0	10,296.0

¹ A portion of the consumptive uses shown herein is satisfied by ground-water overdraft. Table does not include Colorado main stem unmeasured return flow.

² Includes rural, urban, and other industrial uses.

**Table LC-3.--Lower Colorado River Basin
Colorado main stem water use and exports within States and Mexico¹
1981-1985**

(1,000 acre-feet)

Water year	State	Estimated consumptive use ¹					Total	Unmeasured Return flow ³	Adjusted total	Water passing to Mexico ²			
		Irrigated agriculture	Municipal and industrial	Thermal electric power	Fish wildlife & rec.	Export				Treaty	Minute 242	Excess releases	Total
1981	Nevada	0.0	139.7	12.1	1.2	0.0	153.0	43.6	109.4	-	-	-	-
	Arizona	1,348.1	14.6	0.8	53.9	0.0	1,417.4	135.0	1,282.4	-	-	-	-
	California	<u>641.9</u>	<u>3.0</u>	<u>0.0</u>	<u>0.0</u>	<u>4,258.7</u>	<u>4,903.6</u>	<u>65.0</u>	<u>4,838.6</u>	-	-	-	-
	Total	1,990.0	157.3	12.9	55.1	4,258.7	6,474.0	243.6	6,230.4	1,750.8	130.8	2,114.8	3,996.4
1982	Nevada	0.0	137.3	16.8	1.0	0.0	155.1	49.2	105.9	-	-	-	-
	Arizona	1,198.9	13.4	0.6	49.5	0.0	1,262.4	135.4	1,127.0	-	-	-	-
	California	<u>572.2</u>	<u>3.1</u>	<u>0.0</u>	<u>0.0</u>	<u>3,838.4</u>	<u>4,413.7</u>	<u>64.6</u>	<u>4,349.1</u>	-	-	-	-
	Total	1,771.1	153.8	17.4	50.5	3,838.4	5,831.2	249.2	5,582.0	1,494.8	146.2	175.8	1,816.8
1983	Nevada	0.0	141.7	15.5	1.1	0.0	158.3	66.2	92.1	-	-	-	-
	Arizona	1,007.4	13.1	0.7	30.6	0.0	1,051.8	142.2	909.6	-	-	-	-
	California	<u>409.4</u>	<u>2.5</u>	<u>0.0</u>	<u>0.0</u>	<u>3,598.5</u>	<u>4,010.4</u>	<u>57.8</u>	<u>3,952.6</u>	-	-	-	-
	Total	1,416.8	157.3	16.2	31.7	3,598.5	5,220.5	266.2	4,954.3	1,646.3	166.0	7,969.5	9,781.8
1984	Nevada	0.0	148.9	16.1	1.0	0.0	166.0	66.9	99.1	-	-	-	-
	Arizona	1,120.8	14.3	0.7	5.8	0.0	1,141.6	143.0	998.6	-	-	-	-
	California	<u>446.3</u>	<u>3.0</u>	<u>0.0</u>	<u>0.0</u>	<u>4,286.5</u>	<u>4,735.8</u>	<u>57.0</u>	<u>4,678.8</u>	-	-	-	-
	Total	1,567.1	166.2	16.8	6.8	4,286.5	6,043.4	266.9	5,776.5	1,693.6	137.8	15,159.9	16,991.3
1985	Nevada	0.0	163.3	7.4	1.0	0.0	171.7	69.2	102.5	-	-	-	-
	Arizona	1,123.8	14.9	0.5	6.0	⁴ 29.3	1,174.5	142.3	1,032.2	-	-	-	-
	California	<u>455.9</u>	<u>3.3</u>	<u>0.0</u>	<u>0.0</u>	<u>4,308.6</u>	<u>4,767.8</u>	<u>57.7</u>	<u>4,710.1</u>	-	-	-	-
	Total	1,579.7	181.5	7.9	7.0	4,337.9	6,114.0	269.2	5,844.8	1,670.9	130.8	11,594.2	13,395.9

¹ From the Bureau of Reclamation calendar year reports "Compilation of Records in Accordance with Article V of the Decree of the Supreme Court of the United States in Arizona vs. California dated March 9, 1965." Exports to California and water passing to Mexico are demands on system water and consumption is outside the system.

² Source: International Boundary and Water Commission.

³ Estimates of unmeasured return flows are for the Colorado River diversions portions of Las Vegas Wash (Nevada) surface water discharge to Lake Mead, as found in decree accounting. Total unmeasured return flows for Arizona and California are estimated to be 200,000 acre-feet which is proportioned on the basis of irrigated agriculture diversions.

⁴ Central Arizona Project.

**Table LC-4.--Lower Colorado River Basin
Estimated water use within States, by major tributaries, and types of use¹
1981**

		(1,000 acre-feet)										
State	Tributary	Reservoir ⁴ evaporation	Agriculture			Municipal and industrial				Export		Total
			Stockpond evaporation	Irrigation and livestock	Total	Mineral resources	Thermal electric power	Other ²	Total	Outside system	Inside system	
Arizona	Little Colorado	14.9	103.2	5.5	108.7	3.1	20.9	8.5	32.5	0.0	+11.1	167.2
	Virgin	0.3	23.4	0.3	23.7	0.0	0.0	0.5	0.5	0.0	0.0	24.5
	Bill Williams	22.0	37.5	1.2	38.7	1.7	0.0	1.7	3.4	0.0	0.0	64.1
	Gila	236.3	3,723.0	32.5	3,755.5	54.8	16.7	336.0	407.5	0.0	-8.1	4,391.2
	Remaining area	<u>0.6</u>	<u>160.7</u>	<u>9.1</u>	<u>169.8</u>	<u>1.8</u>	<u>0.0</u>	<u>24.0</u>	<u>25.8</u>	<u>0.0</u>	<u>0.0</u>	<u>196.2</u>
	Total	274.1	4,047.8	48.6	4,096.4	61.4	37.6	370.7	469.7	0.0	+3.0	4,843.2
Nevada	Muddy	4.2	36.0	0.7	36.7	0.6	4.4	0.2	5.2	0.0	0.0	46.1
	Remaining area	<u>0.0</u>	<u>59.1</u>	<u>1.0</u>	<u>60.1</u>	<u>2.9</u>	<u>6.2</u>	<u>128.4</u>	<u>137.5</u>	<u>0.0</u>	<u>-3.0</u>	<u>194.6</u>
	Total	4.2	95.1	1.7	96.8	3.5	10.6	128.6	142.7	0.0	-3.0	240.7
N. Mex.	Little Colorado	4.4	1.7	0.7	2.4	1.2	0.0	2.4	3.6	0.0	0.0	10.4
	Gila	<u>0.6</u>	<u>7.2</u>	<u>2.2</u>	<u>9.4</u>	<u>3.4</u>	<u>0.0</u>	<u>0.2</u>	<u>6.6</u>	<u>0.7</u>	<u>0.0</u>	<u>17.3</u>
	Total	5.0	8.9	2.9	11.8	7.6	0.0	2.6	10.2	0.7	0.0	27.7
Utah	Virgin	4.5	95.3	1.1	96.4	0.0	0.0	2.9	2.9	1.4	0.0	105.2
	Remaining area	<u>0.1</u>	<u>10.1</u>	<u>0.2</u>	<u>10.3</u>	<u>0.0</u>	<u>0.0</u>	<u>0.1</u>	<u>0.1</u>	<u>0.0</u>	<u>0.0</u>	<u>10.5</u>
	Total	4.6	105.4	1.3	106.7	0.0	0.0	3.0	3.0	1.4	0.0	115.7
Lower Basin	Little Colorado	19.3	104.9	6.2	111.1	4.3	20.9	10.9	36.1	0.0	11.1	177.6
	Virgin	4.8	118.7	1.4	120.1	0.0	0.0	3.4	3.4	1.4	0.0	129.7
	Muddy	4.2	36.0	0.7	36.7	0.6	4.4	0.2	5.2	0.0	0.0	46.1
	Bill Williams	22.0	37.5	1.2	38.7	1.7	0.0	1.7	3.4	0.0	0.0	64.1
	Gila	236.9	3,730.2	34.7	3,764.9	61.2	16.7	336.2	414.1	0.7	-8.1	4,408.5
	Remaining area	<u>0.7</u>	<u>229.9</u>	<u>10.3</u>	<u>240.2</u>	<u>4.7</u>	<u>6.2</u>	<u>152.5</u>	<u>163.4</u>	<u>0.0</u>	<u>-3.0</u>	<u>401.3</u>
	Total	287.9	4,257.2	54.5	4,311.7	72.5	48.2	504.9	625.6	2.1	0.0	5,227.3

¹ Excludes Colorado River main stem and flood plain. A portion of the consumptive uses shown herein are satisfied by ground-water overdraft.

² Includes rural, urban, and other industrial uses.

³ This value is included in the "Other" Municipal and Industrial Uses category, as listed in the annual report by the New Mexico Interstate Stream Commission as required by the Supreme Court decree in Arizona vs. California.

⁴ Excludes reservoir evaporation from Colorado River main stem reservoirs listed on table LC-1.

**Table LC-5.--Lower Colorado River Basin
Estimated water use within States, by major tributaries, and types of use¹
1982**

(1,000 acre-feet)

State	Tributary	Reservoir ⁴ evaporation	Agriculture			Municipal and industrial				Export		Total
			Irrigation	Stockpond evap. and livestock	Total	Mineral resources	Thermal electric power	Other ²	Total	Outside system	Inside system	
Arizona	Little Colorado	11.2	93.2	4.4	97.6	2.2	22.3	8.6	33.1	0.0	+14.2	156.1
	Virgin	0.2	22.0	0.3	22.3	0.0	0.0	0.5	0.5	0.0	0.0	23.0
	Bill Williams	18.3	29.7	1.0	30.7	2.3	0.0	1.7	4.0	0.0	0.0	53.0
	Gila	177.6	2,952.8	28.4	2,981.2	72.6	9.3	346.5	428.4	0.0	-11.2	3,576.0
	Remaining area	<u>0.5</u>	<u>132.9</u>	<u>8.2</u>	<u>141.1</u>	<u>2.2</u>	<u>0.0</u>	<u>24.0</u>	<u>26.2</u>	<u>0.0</u>	<u>0.0</u>	<u>167.8</u>
	Total	207.8	3,230.6	42.3	3,272.9	79.3	31.6	381.3	492.2	0.0	3.0	3,975.9
Nevada	Muddy	3.9	32.0	0.6	32.6	0.6	7.0	0.1	7.7	0.0	0.0	44.2
	Remaining area	<u>0.0</u>	<u>63.8</u>	<u>1.0</u>	<u>64.8</u>	<u>1.8</u>	<u>4.2</u>	<u>134.1</u>	<u>140.1</u>	<u>0.0</u>	<u>-3.0</u>	<u>201.9</u>
	Total	3.9	95.8	1.6	97.4	2.4	11.2	134.2	147.8	0.0	-3.0	246.1
New Mexico	Little Colorado	4.2	2.3	0.8	3.1	1.2	0.0	2.4	3.6	0.0	0.0	10.9
	Gila	<u>0.5</u>	<u>7.2</u>	<u>2.4</u>	<u>9.6</u>	<u>3.5</u>	<u>0.0</u>	<u>0.2</u>	<u>3.7</u>	<u>0.8</u>	<u>0.0</u>	<u>14.6</u>
	Total	4.7	9.5	3.2	12.7	4.7	0.0	2.6	7.3	0.8	0.0	25.5
Utah	Virgin	4.2	92.4	1.0	93.4	0.0	0.0	3.1	3.1	2.8	0.0	103.5
	Remaining area	<u>0.1</u>	<u>9.1</u>	<u>0.2</u>	<u>9.3</u>	<u>0.0</u>	<u>0.0</u>	<u>0.1</u>	<u>0.1</u>	<u>0.0</u>	<u>0.0</u>	<u>9.5</u>
	Total	4.3	101.5	1.2	102.7	0.0	0.0	3.2	3.2	2.8	0.0	113.0
Lower Basin	Little Colorado	15.4	95.5	5.2	100.7	3.4	22.3	11.0	36.7	0.0	14.2	167.0
	Virgin	4.4	114.4	1.3	115.7	0.0	0.0	3.6	3.6	2.8	0.0	126.5
	Muddy	3.9	32.0	0.6	32.6	0.6	7.0	0.1	7.7	0.0	0.0	44.2
	Bill Williams	18.3	29.7	1.0	30.7	2.3	0.0	1.7	4.0	0.0	0.0	53.0
	Gila	178.1	2,960.0	30.8	2,990.8	76.1	9.3	346.7	432.1	0.8	-11.2	3590.6
	Remaining area	<u>0.6</u>	<u>205.8</u>	<u>9.4</u>	<u>215.2</u>	<u>4.0</u>	<u>4.2</u>	<u>158.2</u>	<u>166.4</u>	<u>0.0</u>	<u>-3.0</u>	<u>379.2</u>
	Total	220.7	3,437.4	48.3	3,485.7	86.4	42.8	521.3	650.5	3.6	0.0	4,360.5

¹ Excludes Colorado River main stem and flood plain. A portion of the consumptive uses shown herein are satisfied by ground-water overdraft.

² Includes rural, urban, and other industrial uses.

³ This value is included in the "Other" Municipal and Industrial Uses category, as listed in the annual report by the New Mexico Interstate Stream Commission as required by the Supreme Court decree in Arizona vs. California.

⁴ Excludes reservoir evaporation from Colorado River main stem reservoirs listed on table LC-1.

**Table LC-6.--Lower Colorado River Basin
Estimated water use within States, by major tributaries, and types of use¹
1983**

(1,000 acre-feet)

State	Tributary	Reservoir ⁴ evaporation	Agriculture			Municipal and industrial				Export		
			Irrigation	Stockpond evap. and livestock	Total	Mineral resources	Thermal electric power	Other ²	Total	Outside system	Inside system	Total
Arizona	Little Colorado	11.5	89.1	4.4	93.5	2.7	19.0	9.0	30.7	0.0	16.5	152.2
	Virgin	0.3	20.5	0.3	20.8	0.0	0.0	0.5	0.5	0.0	0.0	21.6
	Bill Williams	23.7	27.9	1.0	28.9	1.5	0.0	1.8	3.3	0.0	0.0	55.9
	Gila	240.4	2,240.8	25.8	2,266.6	48.2	6.1	361.2	415.5	0.0	-13.5	2,909.0
	Remaining area	<u>0.5</u>	<u>113.0</u>	<u>8.0</u>	<u>121.0</u>	<u>1.7</u>	<u>0.0</u>	<u>24.1</u>	<u>25.8</u>	<u>0.0</u>	<u>0.0</u>	<u>147.3</u>
	Total	276.4	2,491.3	39.5	2,530.8	54.1	25.1	396.6	475.8	0.0	3.0	3,286.0
Nevada	Muddy	4.6	32.7	0.7	33.4	0.6	6.8	0.1	7.5	0.0	0.0	45.5
	Remaining area	<u>0.0</u>	<u>56.8</u>	<u>1.0</u>	<u>57.8</u>	<u>1.7</u>	<u>3.4</u>	<u>141.8</u>	<u>146.9</u>	<u>0.0</u>	<u>-3.0</u>	<u>201.7</u>
	Total	4.6	89.5	1.7	91.2	2.3	10.2	141.9	154.4	0.0	-3.0	247.2
New Mexico	Little Colorado	4.2	1.7	0.8	2.5	1.2	0.0	2.4	3.6	0.0	0.0	10.3
	Gila	<u>0.5</u>	<u>5.7</u>	<u>2.3</u>	<u>8.0</u>	<u>3.4</u>	<u>0.0</u>	<u>0.3</u>	<u>4.9</u>	<u>0.9</u>	<u>0.0</u>	<u>14.3</u>
	Total	4.7	7.4	3.1	10.5	5.8	0.0	2.7	8.5	0.9	0.0	24.6
Utah	Virgin	4.9	92.8	1.1	93.9	0.0	0.0	3.3	3.3	10.0	0.0	112.1
	Remaining area	<u>0.2</u>	<u>9.0</u>	<u>0.2</u>	<u>9.2</u>	<u>0.0</u>	<u>0.0</u>	<u>0.1</u>	<u>0.1</u>	<u>0.0</u>	<u>0.0</u>	<u>9.5</u>
	Total	5.1	101.8	1.3	103.1	0.0	0.0	3.4	3.4	10.0	0.0	121.6
Lower Basin	Little Colorado	15.7	90.8	5.2	96.0	3.9	19.0	11.4	34.3	0.0	16.5	162.5
	Virgin	5.2	113.3	1.4	114.7	0.0	0.0	3.8	3.8	10.0	0.0	133.7
	Muddy	4.6	32.7	0.7	33.4	0.6	6.8	0.1	7.5	0.0	0.0	45.5
	Bill Williams	23.7	27.9	1.0	28.9	1.5	0.0	1.8	3.3	0.0	0.0	55.9
	Gila	240.9	2,246.5	28.1	2,274.6	52.8	6.1	361.5	420.4	0.9	-13.5	2,923.3
	Remaining area	<u>0.7</u>	<u>178.8</u>	<u>9.2</u>	<u>188.0</u>	<u>3.4</u>	<u>3.4</u>	<u>166.0</u>	<u>172.8</u>	<u>0.0</u>	<u>-3.0</u>	<u>358.5</u>
	Total	290.8	2,690.0	45.6	2,735.6	62.2	35.3	544.6	642.1	10.9	0.0	3,679.4

¹ A portion of the consumptive uses shown herein are satisfied by ground-water overdraft.

² Includes rural, urban, and other industrial uses.

³ This value is included in the "Other" Municipal and Industrial Uses category, as listed in the annual report by the New Mexico Interstate Stream Commission as required by the Supreme Court decree in Arizona vs. California.

⁴ Excludes reservoir evaporation from Colorado River main stem reservoirs listed on table LC-1.

**Table LC-7.--Lower Colorado River Basin
Estimated water use within States, by major tributaries, and types of use¹
1984**

(1,000 acre-feet)												
State	Tributary	Reservoir ⁴ evaporation	Agriculture			Municipal and industrial				Export		Total
			Irrigation	Stockpond evap. and livestock	Total	Mineral resources	Thermal electric power	Other ²	Total	Outside system	Inside system	
Arizona	Little Colorado	10.4	99.5	4.1	103.6	3.2	21.9	9.3	34.4	0.0	17.6	166.0
	Virgin	0.2	21.8	0.3	22.1	0.0	0.0	0.5	0.5	0.0	0.0	22.8
	Bill Williams	24.1	27.1	1.1	28.2	1.6	0.0	1.6	3.2	0.0	0.0	55.5
	Gila	234.5	2,901.8	24.2	2,926.0	53.3	10.4	381.9	445.6	0.0	-14.6	3,591.5
	Remaining area	<u>0.4</u>	<u>126.8</u>	<u>8.3</u>	<u>135.1</u>	<u>1.8</u>	<u>0.0</u>	<u>25.2</u>	<u>27.0</u>	<u>0.0</u>	<u>0.0</u>	<u>162.5</u>
	Total	269.6	3,177.0	38.0	3,215.0	59.9	32.3	418.5	510.7	0.0	3.0	3,998.3
Nevada	Muddy	5.1	35.4	0.8	36.2	0.6	6.8	0.3	7.7	0.0	0.0	49.0
	Remaining area	<u>0.0</u>	<u>58.2</u>	<u>1.0</u>	<u>59.2</u>	<u>1.7</u>	<u>3.4</u>	<u>145.8</u>	<u>150.9</u>	<u>0.0</u>	<u>-3.0</u>	<u>207.1</u>
	Total	5.1	93.6	1.8	95.4	2.3	10.2	146.1	158.6	0.0	-3.0	256.1
New Mexico	Little Colorado	4.3	1.6	0.8	2.4	1.2	0.0	2.4	3.6	0.0	0.0	10.3
	Gila	<u>0.5</u>	<u>5.1</u>	<u>2.4</u>	<u>7.5</u>	<u>3.4</u>	<u>0.0</u>	<u>0.2</u>	<u>4.3</u>	<u>0.5</u>	<u>0.0</u>	<u>12.8</u>
	Total	4.8	6.7	3.2	9.9	5.3	0.0	2.6	7.9	0.5	0.0	23.1
Utah	Virgin	5.4	101.4	1.1	102.5	0.0	0.0	3.5	3.5	2.6	0.0	114.0
	Remaining area	<u>0.2</u>	<u>9.4</u>	<u>0.2</u>	<u>9.6</u>	<u>0.0</u>	<u>0.0</u>	<u>0.1</u>	<u>0.1</u>	<u>0.0</u>	<u>0.0</u>	<u>9.9</u>
	Total	5.6	110.8	1.3	112.1	0.0	0.0	3.6	3.6	2.6	0.0	123.9
Lower Basin	Little Colorado	14.7	101.1	4.9	106.0	4.4	21.9	11.7	38.0	0.0	17.6	176.3
	Virgin	5.6	123.2	1.4	124.6	0.0	0.0	4.0	4.0	2.6	0.0	136.8
	Muddy	5.1	35.4	0.8	36.2	0.6	6.8	0.3	7.7	0.0	0.0	49.0
	Bill Williams	24.1	27.1	1.1	28.2	1.6	0.0	1.6	3.2	0.0	0.0	55.5
	Gila	235.0	2,906.9	26.6	2,933.5	57.4	10.4	382.1	449.9	0.5	-14.6	3,604.3
	Remaining area	<u>0.6</u>	<u>194.4</u>	<u>9.5</u>	<u>203.9</u>	<u>3.5</u>	<u>3.4</u>	<u>171.1</u>	<u>178.0</u>	<u>0.0</u>	<u>-3.0</u>	<u>379.5</u>
	Total	285.1	3,388.1	44.3	3,432.4	67.5	42.5	570.8	680.8	3.1	0.0	4,401.4

¹ A portion of the consumptive uses shown herein are satisfied by ground-water overdraft.

² Includes rural, urban, and other industrial uses.

³ This value is included in the "Other" Municipal and Industrial Uses category, as listed in the annual report by the New Mexico Interstate Stream Commission as required by the Supreme Court decree in Arizona vs. California.

⁴ Excludes reservoir evaporation from Colorado River main stem reservoirs listed on table LC-1.

**Table LC-8.--Lower Colorado River Basin
Estimated water use within States, by major tributaries, and types of use¹
1985**

(1,000 acre-feet)												
State	Tributary	Reservoir ⁴ evaporation	Agriculture			Municipal and industrial				Export		Total
			Irrigation	Stockpond evap. and livestock	Total	Mineral resources	Thermal electric power	Other ²	Total	Outside system	Inside system	
Arizona	Little Colorado	13.6	88.0	5.2	93.2	2.4	22.5	9.8	34.7	0.0	11.4	152.9
	Virgin	0.3	19.9	0.3	20.2	0.0	0.0	0.5	0.5	0.0	0.0	21.0
	Bill Williams	25.8	24.2	1.1	25.3	1.7	0.0	1.8	3.5	0.0	0.0	54.6
	Gila	221.0	2,686.6	30.8	2,717.4	56.8	11.7	404.4	472.9	0.0	-37.7	3,373.6
	Remaining area	<u>0.6</u>	<u>123.2</u>	<u>8.6</u>	<u>131.8</u>	<u>2.0</u>	<u>0.0</u>	<u>25.4</u>	<u>27.4</u>	<u>0.0</u>	<u>0.0</u>	<u>159.8</u>
	Total	261.3	2,941.9	46.0	2,987.9	62.9	34.2	441.9	539.0	0.0	-26.3	3,761.9
Nevada	Muddy	4.7	35.4	0.7	36.1	0.6	8.9	0.3	9.8	0.0	0.0	50.6
	Remaining area	<u>0.0</u>	<u>65.5</u>	<u>1.0</u>	<u>66.5</u>	<u>1.7</u>	<u>3.5</u>	<u>150.3</u>	<u>155.5</u>	<u>0.0</u>	<u>-3.0</u>	<u>219.0</u>
	Total	4.7	100.9	1.7	102.6	2.3	12.4	150.6	165.3	0.0	-3.0	269.6
New Mexico	Little Colorado	4.8	2.2	0.7	2.9	1.2	0.0	2.5	3.7	0.0	0.0	11.4
	Gila	<u>0.4</u>	<u>4.2</u>	<u>2.2</u>	<u>6.4</u>	<u>6.6</u> ³	<u>0.0</u>	<u>0.3</u>	<u>6.9</u>	<u>1.0</u>	<u>0.0</u>	<u>14.7</u>
	Total	5.2	6.4	2.9	9.3	7.8	0.0	2.8	10.6	1.0	0.0	26.1
Utah	Virgin	5.0	102.2	1.1	103.3	0.0	0.0	3.8	3.8	2.1	0.0	114.2
	Remaining area	<u>0.2</u>	<u>9.7</u>	<u>0.2</u>	<u>9.9</u>	<u>0.0</u>	<u>0.0</u>	<u>0.1</u>	<u>0.1</u>	<u>0.0</u>	<u>0.0</u>	<u>10.2</u>
	Total	5.2	111.9	1.3	113.2	0.0	0.0	3.9	3.9	2.1	0.0	124.4
Lower Basin	Little Colorado	18.4	90.2	5.9	96.1	3.6	22.5	12.3	38.4	0.0	11.4	164.3
	Virgin	5.3	122.1	1.4	123.5	0.0	0.0	4.3	4.3	2.1	0.0	135.2
	Muddy	4.7	35.4	0.7	36.1	0.6	8.9	0.3	9.8	0.0	0.0	50.6
	Bill Williams	25.8	24.2	1.1	25.3	1.7	0.0	1.8	3.5	0.0	0.0	54.6
	Gila	221.4	2,690.8	33.0	2,723.8	63.4	11.7	404.7	479.8	1.0	-37.7	3,388.3
	Remaining area	<u>0.8</u>	<u>198.4</u>	<u>9.8</u>	<u>208.2</u>	<u>3.7</u>	<u>3.5</u>	<u>175.8</u>	<u>183.0</u>	<u>0.0</u>	<u>-3.0</u>	<u>389.0</u>
	Total	276.4	3,161.1	51.9	3,213.0	73.0	46.6	599.2	718.8	3.1	⁵ -29.3	4,182.0

¹ A portion of the consumptive uses shown herein are satisfied by ground-water overdraft.

² Includes rural, urban, and other industrial uses.

³ This value is included in the "Other" Municipal and Industrial Uses category, as listed in the annual report by the New Mexico Interstate Stream Commission as required by the Supreme Court decree in Arizona vs. California.

⁴ Excludes reservoir evaporation from Colorado River main stem reservoirs listed on table LC-1.

⁵ Central Arizona Project.

**Table LC-9.--Lower Colorado River Basin
Irrigated acreage³
1981-1985**

		(1,000 acres)				
State	Tributary	Irrigated acreage ¹				
		1981	1982	1983	1984	1985
Arizona	Gila ²	1,223.7	990.4	751.1	956.2	883.7
	Little Colorado	32.6	31.5	30.5	31.6	27.6
	Bill Williams	12.0	10.7	9.8	9.5	8.6
	Virgin	3.4	3.1	3.1	2.8	2.6
	Remaining area	<u>31.8</u>	<u>29.0</u>	<u>35.6</u>	<u>39.5</u>	<u>38.0</u>
	Total	1,303.5	1,064.7	830.1	1,039.6	960.5
Nevada	Muddy	10.8	10.9	11.2	10.7	10.5
	Remaining area	<u>8.2</u>	<u>9.3</u>	<u>8.3</u>	<u>8.3</u>	<u>9.2</u>
	Total	19.0	20.2	19.5	19.0	19.7
New Mexico	Gila	4.5	4.4	4.1	3.4	2.3
	Little Colorado	<u>2.5</u>	<u>2.1</u>	<u>1.2</u>	<u>1.4</u>	<u>1.6</u>
	Total	7.0	6.5	5.3	4.8	3.9
Utah	Virgin	21.3	21.7	21.6	22.4	22.4
	Remaining area	<u>3.3</u>	<u>3.4</u>	<u>3.4</u>	<u>3.2</u>	<u>3.2</u>
	Total	24.6	25.1	25.0	25.6	25.6
California	Remaining area	<u>116.5</u>	<u>116.4</u>	<u>103.4</u>	<u>113.9</u>	<u>110.5</u>
	Total	116.5	116.4	103.4	113.9	110.5
<hr/>						
Lower Basin	Gila	1,228.2	994.8	755.2	959.6	886.0
	Little Colorado	35.1	33.6	31.7	33.0	29.2
	Bill Williams	12.0	10.7	9.8	9.5	8.6
	Virgin	24.7	24.8	24.7	25.2	25.0
	Muddy	10.8	10.9	11.2	10.7	10.5
	Remaining area	<u>159.8</u>	<u>158.1</u>	<u>150.7</u>	<u>164.9</u>	<u>160.9</u>
	Total	1,470.6	1,232.9	983.3	1,202.9	1,120.2

¹ Irrigated acreage includes all irrigated croplands harvested as well as irrigated pasture. Double-cropping is accounted.

² According to Arizona crop census reports, the fluctuation in irrigated acreage did occur.

³ Does not include main stem irrigated areas.

**Table LC-10.--Lower Colorado River Basin
Population estimates
1981-1985**

		(1,000's)				
State	Tributary	Estimated population				
		1981	1982	1983	1984	1985
Arizona	Lower Main Stem*	166.4	153.3	160.7	167.1	170.9
	Gila	2,348.1	2,419.0	2,505.7	2,633.7	2,752.6
	Little Colorado	131.9	134.9	140.1	142.0	149.1
	Bill Williams	17.0	17.7	17.5	19.4	19.6
	Virgin	<u>3.0</u>	<u>3.1</u>	<u>3.3</u>	<u>3.4</u>	<u>3.4</u>
	Total	2,666.4	2,728.0	2,827.3	2,965.6	3,095.6
California	Lower Main Stem	23.8	21.9	22.9	23.8	24.3
Nevada	Lower Main Stem*	502.2	527.1	557.4	572.6	590.8
	Muddy	<u>4.0</u>	<u>4.2</u>	<u>4.5</u>	<u>4.6</u>	<u>4.9</u>
	Total	506.2	531.3	561.9	577.2	595.7
New Mexico	Gila	9.0	9.3	9.6	9.6	9.3
	Little Colorado	<u>45.5</u>	<u>45.9</u>	<u>46.3</u>	<u>46.7</u>	<u>47.2</u>
	Total	54.5	55.2	55.9	56.3	56.5
Utah	Lower Main Stem*	0.8	0.8	0.9	0.9	0.9
	Virgin	<u>29.3</u>	<u>31.1</u>	<u>32.4</u>	<u>34.7</u>	<u>37.6</u>
	Total	30.1	31.9	33.3	35.6	38.5
Lower Basin	Lower Main Stem	693.2	703.1	741.9	764.4	786.9
	Gila	2,357.1	2,428.3	2,515.3	2,643.3	2,761.9
	Little Colorado	177.4	180.8	186.4	188.7	196.3
	Bill Williams	17.0	17.7	17.5	19.4	19.6
	Virgin	32.3	34.2	35.7	38.1	41.0
	Muddy	<u>4.0</u>	<u>4.2</u>	<u>4.5</u>	<u>4.6</u>	<u>4.9</u>
	Total	3,281.0	3,368.3	3,501.3	3,658.5	3,810.6

* Includes remaining area population.