

**2005Table 5-5. Unspecified and other pneumoconioses: Number of deaths, death rates (per million population), and years of potential life lost (YPLL) by state, U.S. residents age 15 and over, 1993–2002**

State	No. of Deaths		Crude Death Rate		Age-Adjusted Death Rate		YPLL to Life Expectancy			
	Deaths	Rank	Rate	Rank	Rate	Rank	Total	Rank	YPLL/death	Rank
Alabama	46	12	1.33	12	1.38	12	431	15	9.4	43
Alaska	2	43	0.44	22	0.99	15	23	45	11.5	21
Arizona	12	26	0.33	31	0.31	31	141	26	11.8	19
Arkansas	5	36	0.25	36	0.23	39	68	36	13.6	11
California	61	8	0.24	37	0.28	35	756	7	12.4	15
Colorado	48	11	1.54	9	2.00	9	429	16	8.9	45
Connecticut	11	28	0.42	25	0.38	27	155	25	14.0	9
Delaware	12	26	2.02	8	2.17	8	123	27	10.2	36
District of Columbia	1	46	0.23	40	0.24	38	38	41	38.4	1
Florida	63	7	0.53	19	0.39	25	662	8	10.5	34
Georgia	22	21	0.37	28	0.47	21	289	20	13.1	13
Hawaii	1	46	0.11	47	0.10	49	14	48	14.1	8
Idaho	2	43	0.21	41	0.22	41	30	42	15.2	4
Illinois	115	6	1.23	13	1.24	13	1,106	6	9.6	40
Indiana	39	16	0.84	15	0.86	16	424	17	10.9	29
Iowa	13	25	0.57	18	0.45	23	113	29	8.7	47
Kansas	7	31	0.34	30	0.31	31	71	35	10.2	36
Kentucky	245	3	7.84	4	8.16	4	2,734	3	11.2	26
Louisiana	8	30	0.24	37	0.25	37	113	29	14.2	6
Maine	2	43	0.20	42	0.19	42	23	45	11.3	24
Maryland	30	19	0.74	16	0.83	17	345	18	11.5	21
Massachusetts	18	23	0.37	28	0.33	30	183	23	10.2	36
Michigan	41	14	0.53	19	0.55	19	460	14	11.2	26
Minnesota	4	38	0.11	47	0.11	48	56	39	14.0	9
Mississippi	6	34	0.28	33	0.29	34	65	37	10.9	29
Missouri	17	24	0.40	26	0.38	27	216	22	12.7	14
Montana	3	41	0.43	24	0.41	24	29	43	9.5	42
Nebraska	–	–	–	–	–	–	–	–	–	–
Nevada	6	34	0.44	22	0.48	20	102	31	17.0	3
New Hampshire	3	41	0.32	32	0.34	29	27	44	8.9	45
New Jersey	40	15	0.62	17	0.60	18	566	11	14.2	6
New Mexico	19	22	1.44	11	1.63	10	183	23	9.6	40
New York	58	9	0.40	26	0.39	25	655	9	11.3	24
North Carolina	27	20	0.45	21	0.47	21	335	19	12.4	15
North Dakota	1	46	0.20	42	0.17	43	9	49	8.5	48
Ohio	198	5	2.23	7	2.18	7	2,107	5	10.6	33
Oklahoma	7	31	0.27	34	0.26	36	75	33	10.8	32
Oregon	4	38	0.15	45	0.15	45	48	40	12.1	18
Pennsylvania	967	1	10.03	2	8.32	3	8,959	1	9.3	44
Rhode Island	–	–	–	–	–	–	–	–	–	–
South Carolina	4	38	0.13	46	0.13	46	83	32	20.9	2
South Dakota	1	46	0.17	44	0.16	44	15	47	14.6	5
Tennessee	45	13	1.04	14	1.08	14	530	12	11.8	19
Texas	39	16	0.26	35	0.31	31	521	13	13.4	12
Utah	57	10	3.74	6	5.10	6	587	10	10.3	35
Vermont	7	31	1.47	10	1.54	11	72	34	10.2	36
Virginia	242	4	4.46	5	5.24	5	2,642	4	10.9	29
Washington	5	36	0.11	47	0.12	47	57	38	11.4	23
West Virginia	588	2	39.56	1	35.18	1	6,538	2	11.1	28
Wisconsin	10	29	0.24	37	0.23	39	122	28	12.2	17
Wyoming	35	18	9.26	3	10.90	2	271	21	7.7	49

– indicates no deaths listed.

NOTE: Slight variations in numbers may exist when compared to previous WoRLD Surveillance Reports. See appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Bureau of the Census.