

TABLE 1
THE 20 MOST FREQUENTLY REPORTED SALMONELLA SEROTYPES
FROM HUMAN SOURCES REPORTED TO CDC IN 2000 AND FROM
CLINICAL AND NON-CLINICAL NONHUMAN SOURCES REPORTED TO CDC AND NVSL IN 2000

HUMAN 2000				CLINICAL NONHUMAN 2000				NON-CLINICAL NONHUMAN 2000			
RANK	SEROTYPE	NUMBER	PERCENT	RANK	SEROTYPE	NUMBER	PERCENT	RANK	SEROTYPE	NUMBER	PERCENT
1	TYPHIMURIUM *	7085	22.1	1	TYPHIMURIUM *	2197	31.3	1	HEIDELBERG	3059	28.2
2	ENTERITIS	6224	19.4	2	HEIDELBERG	606	8.6	2	TYPHIMURIUM *	1384	12.8
3	NEWPORT	2975	9.3	3	NEWPORT	462	6.6	3	KENTUCKY	807	7.4
4	HEIDELBERG	1661	5.2	4	AGONA	315	4.5	4	BERTA	545	5.0
5	JAVIANA	1167	3.6	5	DERBY	264	3.8	5	SENFTENBERG	504	4.6
6	MONTEVIDEO	799	2.5	6	MONTEVIDEO	196	2.8	6	MONTEVIDEO	485	4.5
7	MUENCHEN	605	1.9	7	CHOLERAESUIS **	181	2.6	7	ANATUM	335	3.1
8	INFANTIS	577	1.8	8	MUENSTER	175	2.5	8	AGONA	315	2.9
9	THOMPSON	569	1.8	9	WORTHINGTON	174	2.5	9	HADAR	290	2.7
10	ORANIENBURG	538	1.7	10	ANATUM	168	2.4	10	ENTERITIS	236	2.2
11	SAINTPAUL	522	1.6	11	BREDENEY	144	2.0	11	READING	214	2.0
12	BRAENDERUP	507	1.6	12	DUBLIN	130	1.9	12	MUENSTER	209	1.9
13	JAVA	434	1.4	13	SENFTENBERG	117	1.7	13	DERBY	194	1.8
14	AGONA	382	1.2	14	KENTUCKY	91	1.3	14	SCHWARZENGRUND	182	1.7
15	TYPHI	368	1.1	15	INFANTS	89	1.3	15	MBANDAKA	173	1.6
16	HADAR	330	1.0	16	UGANDA	89	1.3	16	INFANTIS	156	1.4
17	POONA	322	1.0	17	CERRO	79	1.1	17	NEWPORT	146	1.3
18	BERTA	295	0.9	18	MBANDAKA	71	1.0	18	CERRO	145	1.3
19	MISSISSIPPI	285	0.9	19	MELEGRIDIS	66	0.9	19	OHTO	115	1.1
20	STANLEY	233	0.7	20	SAINTPAUL	62	0.9	20	BRAENDERUP	109	1.0
SUB TOTAL				SUB TOTAL				SUB TOTAL			
TOTAL				TOTAL				TOTAL			

* TYPHIMURIUM INCLUDES VAR. COPENHAGEN
** CHOLERAESUIS INCLUDES VAR. KUNZENDORF