

TABLE 1**The 20 most frequently reported *Salmonella* serotypes from human sources reported to CDC in 2006**

Human 2006			
Rank	Serotype	Reported	Percent
1	Typhimurium *	6872	16.9
2	Enteritidis	6740	16.6
3	Newport	3373	8.3
4	Heidelberg	1495	3.7
5	Javiana	1433	3.5
6	I 4,[5],12:i:-	1200	3.0
7	Montevideo	1061	2.6
8	Muenchen	753	1.9
9	Oranienburg	719	1.8
10	Mississippi	604	1.5
11	Saintpaul	588	1.4
12	Braenderup	561	1.4
13	Agona	538	1.3
14	Infantis	491	1.2
15	Thompson	447	1.1
16	Paratyphi B var. L(+) tartrate+	417	1.0
17	Typhi	413	1.0
18	Stanley	315	0.8
19	Tennessee	312	0.8
20	Hadar	275	0.7
	Sub Total	28607	70.3
	All Other Serotyped	6459	15.9
	Unknown	4042	9.9
	Partially serotyped	1448	3.6
	Rough, mucoid, and/or nonmotile isolates	110	0.3
	Sub Total	12059	29.7
	Total	40666	100
NOTE: ----- * Typhimurium includes var. 5- (Formerly var. Copenhagen)			

TABLE 1a

The 20 most frequently reported *Salmonella* serotypes from clinical and non-clinical nonhuman sources reported to CDC and NVSL in 2006

Clinical Nonhuman 2006			
Rank	Serotype	Reported	Percent
1	Typhimurium *	1592	19.9
2	Newport	648	8.1
3	Agona	455	5.7
4	Orion var. 15+,34+	365	4.6
5	Dublin	358	4.5
6	Montevideo	356	4.4
7	Anatum	315	3.9
8	Derby	302	3.8
9	Heidelberg	264	3.3
10	Senftenberg	232	2.9
11	Kentucky	213	2.7
12	Muenster	189	2.4
13	Mbandaka	172	2.1
14	I 4,5,12:i:-	160	2.0
15	Cerro	159	2.0
16	Infantis	159	2.0
17	Choleraesuis **	157	2.0
18	Reading	104	1.3
19	Meleagridis	98	1.2
20	Uganda	82	1.0
	Sub Total	6380	79.6
	All Other Serotyped	1635	20.4
	Sub Total	1635	20.4
	Total	8015	100

NOTE:

 * Typhimurium includes var. 5- (Formerly var. Copenhagen)
 ** Choleraesuis includes var. Decatur and Kunzendorf

Non-Clinical Nonhuman 2006			
Rank	Serotype	Reported	Percent
1	Heidelberg	774	11.5
2	Kentucky	732	10.9
3	Typhimurium *	695	10.4
4	Senftenberg	555	8.3
5	Hadar	438	6.5
6	Cerro	239	3.6
7	Enteritidis	222	3.3
8	Agona	204	3.0
9	Anatum	179	2.7
10	Montevideo	160	2.4
11	Newport	158	2.4
12	Mbandaka	137	2.0
13	Derby	129	1.9
14	Schwarzengrund	122	1.8
15	Oranienburg	120	1.8
16	I 4,5,12:i:-	118	1.8
17	Give	106	1.6
18	Worthington	105	1.6
19	Muenchen	95	1.4
20	Saintpaul	93	1.4
	Sub Total	5381	80.2
	All Other Serotyped	1329	19.8
	Sub Total	1329	19.8
	Total	6710	100

NOTE:

 * Typhimurium includes var. 5- (Formerly var. Copenhagen)

TABLE 1b**The 20 most frequently reported *Salmonella* serotypes from human sources
Percent change in reported isolates**

Rank			Serotype	Reported Isolates			Percent Change		
1996	2001	2006		1996	2001	2006	1996-2001	2001-2006	1996-2006
2	1	1	Typhimurium *	9500	7040	6872	-26	-2	-28
1	2	2	Enteritidis	9570	5634	6740	-41	20	-30
4	3	3	Newport	1985	3168	3373	60	6	70
3	4	4	Heidelberg	1998	1895	1495	-5	-21	-25
6	5	5	Javiana	749	1068	1433	43	34	91
35	20	6	I 4,[5],12:i:-	96	260	1200	171	362	1150
5	6	7	Montevideo	1227	630	1061	-49	68	-14
10	8	8	Muenchen	595	586	753	-2	28	27
7	7	9	Oranienburg	690	598	719	-13	20	4
24	16	10	Mississippi	180	336	604	87	80	236
12	10	11	Saintpaul	562	471	588	-16	25	5
13	13	12	Braenderup	531	396	561	-25	42	6
9	14	13	Agona	606	372	538	-39	45	-11
14	12	14	Infantis	503	441	491	-12	11	-2
11	9	15	Thompson	586	514	447	-12	-13	-24
18	11	16	Paratyphi B var. L(+) tartrate+	289	467	417	62	-11	44
15	15	17	Typhi	440	344	413	-22	20	-6
22	24	18	Stanley	200	173	315	-14	82	58
37	61	19	Tennessee	96	33	312	-66	845	225
8	19	20	Hadar	658	307	275	-53	-10	-58

NOTE:

* Typhimurium includes var. 5- (Formerly var. Copenhagen)