

**Table 3-12. Silicosis: Number of cases by ascertainment source and state, 1989-1998**

| Source                          | Michigan   |              | New Jersey |              | Ohio       |              | Total        |              |
|---------------------------------|------------|--------------|------------|--------------|------------|--------------|--------------|--------------|
|                                 | No.        | %            | No.        | %            | No.        | %            | No.          | %            |
| Health care professional report | 123        | 20.9         | 18         | 9.4          | 22         | 5.5          | 163          | 13.8         |
| Hospital discharge data         | 387        | 65.7         | 164        | 85.9         | 327        | 81.8         | 878          | 74.4         |
| Death certificate data          | 23         | 3.9          | 5          | 2.6          | 19         | 4.8          | 47           | 4.0          |
| Workers' compensation files     | 52         | 8.8          | -          | -            | 21         | 5.3          | 73           | 6.2          |
| Other                           | 4          | 0.7          | 4          | 2.1          | 11         | 2.8          | 19           | 1.6          |
| <b>TOTAL</b>                    | <b>589</b> | <b>100.0</b> | <b>191</b> | <b>100.0</b> | <b>400</b> | <b>100.0</b> | <b>1,180</b> | <b>100.0</b> |

- indicates no cases reported.

NOTE: Percentages may not sum to 100% due to rounding. See appendices for source description.

SOURCE: Provisional SENSOR surveillance data as of September 2002, aggregated by reporting source years, and reported by K Rosenman, MJ Reilly, and D Kalinowski (Michigan); D Valiante and D Schill (New Jersey); and E Socie and A Miglioizzi (Ohio).