

**2005Table 6-5. All pneumoconioses: Number of deaths, death rates (per million population), and years of potential life lost (YPLL) by state, U.S. residents age 15 and over, 1993–2002**

State	No. of Deaths		Crude Death Rate		Age-Adjusted Death Rate		YPLL to Life Expectancy			
	Deaths	Rank	Rate	Rank	Rate	Rank	Total	Rank	YPLL/death	Rank
Alabama	619	13	17.90	8	17.96	9	7,018	12	11.3	25
Alaska	29	48	6.38	34	16.50	10	317	49	10.9	35
Arizona	228	27	6.32	35	6.20	36	2,446	29	10.7	37
Arkansas	148	33	7.39	28	6.72	31	1,810	32	12.2	9
California	1,331	4	5.30	42	6.03	37	15,271	4	11.5	22
Colorado	311	20	10.00	19	12.68	15	3,278	22	10.5	41
Connecticut	191	31	7.32	29	6.61	33	2,220	31	11.6	18
Delaware	149	32	25.08	5	26.17	6	1,647	33	11.1	29
District of Columbia	12	51	2.76	50	2.74	50	175	51	14.6	2
Florida	1,013	8	8.48	24	6.24	35	11,206	8	11.1	29
Georgia	249	25	4.18	47	5.33	41	2,998	25	12.0	10
Hawaii	27	49	2.86	49	2.85	49	348	48	12.9	6
Idaho	69	42	7.31	30	7.92	26	730	43	10.6	39
Illinois	676	10	7.21	32	7.29	29	7,459	11	11.0	32
Indiana	309	21	6.65	33	6.80	30	3,410	21	11.0	32
Iowa	136	35	5.97	37	4.92	43	1,359	37	10.0	47
Kansas	106	38	5.15	44	4.90	44	1,409	36	13.3	5
Kentucky	1,267	5	40.53	3	42.14	3	14,695	5	11.6	18
Louisiana	305	22	8.98	21	9.75	22	4,101	20	13.4	4
Maine	133	36	13.20	14	12.50	16	1,505	34	11.3	25
Maryland	563	14	13.88	13	15.55	12	6,602	15	11.7	14
Massachusetts	419	18	8.51	23	7.86	27	4,462	19	10.6	39
Michigan	445	17	5.77	39	5.94	38	5,228	18	11.7	14
Minnesota	215	30	5.82	38	5.87	39	2,497	28	11.6	18
Mississippi	397	19	18.61	7	19.30	8	6,446	16	16.2	1
Missouri	224	28	5.23	43	4.88	45	2,630	27	11.7	14
Montana	76	40	10.85	16	10.36	20	758	40	10.0	47
Nebraska	50	45	3.85	48	3.49	48	495	45	9.9	49
Nevada	76	40	5.62	40	6.41	34	932	39	12.3	7
New Hampshire	68	43	7.25	31	7.82	28	749	41	11.0	32
New Jersey	1,005	9	15.61	9	14.83	13	10,788	9	10.7	37
New Mexico	138	34	10.45	18	11.75	17	1,434	35	10.4	42
New York	667	12	4.64	45	4.48	46	7,601	10	11.4	23
North Carolina	562	15	9.46	20	9.77	21	6,633	14	11.8	13
North Dakota	31	47	6.11	36	5.45	40	354	47	11.4	23
Ohio	1,247	6	14.06	12	13.68	14	13,571	7	10.9	35
Oklahoma	111	37	4.22	46	4.02	47	1,299	38	11.7	14
Oregon	297	23	11.41	15	10.99	19	3,087	24	10.4	42
Pennsylvania	7,282	1	75.52	2	62.45	2	69,668	1	9.6	50
Rhode Island	62	44	7.85	25	6.63	32	694	44	11.2	28
South Carolina	269	24	8.84	22	9.47	23	3,227	23	12.0	10
South Dakota	14	50	2.43	51	2.18	51	193	50	13.8	3
Tennessee	465	16	10.74	17	11.28	18	5,270	17	11.3	25
Texas	1,162	7	7.71	26	9.40	24	14,258	6	12.3	7
Utah	230	26	15.10	10	20.53	7	2,365	30	10.3	44
Vermont	36	46	7.56	27	8.11	25	371	46	10.3	44
Virginia	1,836	3	33.86	4	39.34	4	21,230	3	11.6	18
Washington	668	11	14.93	11	16.43	11	6,784	13	10.2	46
West Virginia	3,192	2	214.78	1	191.21	1	35,530	2	11.1	29
Wisconsin	222	29	5.39	41	5.21	42	2,655	26	12.0	10
Wyoming	87	39	23.01	6	26.80	5	735	42	8.5	51

– indicates no deaths listed.

NOTE: Slight variations in numbers may exist when compared to previous WoRLD Surveillance Reports. See appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Bureau of the Census.