

2005Table 6-6. All pneumoconioses: Percent of deaths by condition and state, U.S. residents age 15 and over, 1993–2002

State	Asbestosis	Silicosis	Coal Workers' Pneumoconiosis	Byssinosis	Unspecified/Other Pneumoconioses
Alabama	68.3	4.8	20.4	0.5	7.4
Alaska	79.3	10.3	6.9	–	6.9
Arizona	61.4	16.2	18.0	–	5.3
Arkansas	56.1	10.1	30.4	1.4	3.4
California	79.5	6.4	9.6	0.5	4.6
Colorado	31.8	22.5	30.2	1.0	15.4
Connecticut	74.3	13.1	7.3	–	5.8
Delaware	84.6	3.4	4.7	–	8.1
District of Columbia	58.3	25.0	8.3	–	8.3
Florida	72.3	4.9	17.6	0.4	6.2
Georgia	61.8	11.2	14.5	4.4	8.8
Hawaii	92.6	3.7	–	–	3.7
Idaho	66.7	20.3	10.1	–	2.9
Illinois	37.7	9.5	38.3	0.1	17.0
Indiana	24.6	13.3	50.5	–	12.6
Iowa	47.8	16.2	28.7	0.7	9.6
Kansas	65.1	8.5	21.7	–	6.6
Kentucky	7.3	3.2	70.8	0.1	19.3
Louisiana	81.0	10.2	6.6	–	2.6
Maine	93.2	5.3	0.8	–	1.5
Maryland	80.1	4.8	11.5	0.2	5.3
Massachusetts	90.2	5.7	0.2	0.2	4.3
Michigan	54.4	19.3	18.9	0.7	9.2
Minnesota	78.1	19.1	0.9	0.9	1.9
Mississippi	77.8	3.0	17.9	–	1.5
Missouri	64.3	16.1	12.9	0.4	7.6
Montana	63.2	19.7	11.8	1.3	3.9
Nebraska	90.0	4.0	6.0	–	–
Nevada	72.4	15.8	5.3	–	7.9
New Hampshire	88.2	4.4	5.9	–	4.4
New Jersey	89.1	3.7	4.4	–	4.0
New Mexico	29.0	22.5	34.8	–	13.8
New York	71.1	13.3	8.5	0.3	8.7
North Carolina	62.5	12.1	14.6	7.3	4.8
North Dakota	87.1	3.2	6.5	–	3.2
Ohio	33.7	14.0	38.8	–	15.9
Oklahoma	56.8	10.8	25.2	0.9	6.3
Oregon	90.6	6.7	2.4	–	1.3
Pennsylvania	13.9	4.4	70.0	<0.1	13.3
Rhode Island	83.9	12.9	3.2	1.6	–
South Carolina	77.7	8.2	10.8	2.6	1.5
South Dakota	35.7	50.0	7.1	–	7.1
Tennessee	36.1	6.5	49.9	–	9.7
Texas	85.0	8.5	4.3	0.3	3.4
Utah	22.2	16.1	38.7	–	24.8
Vermont	50.0	27.8	2.8	2.8	19.4
Virginia	27.2	1.8	58.2	0.2	13.2
Washington	91.2	3.4	5.1	–	0.7
West Virginia	9.4	1.8	72.0	<0.1	18.4
Wisconsin	55.9	34.7	5.0	1.4	4.5
Wyoming	25.3	1.1	33.3	–	40.2
TOTAL	42.4	6.8	40.9	0.4	10.9

– indicates no deaths listed. NOTE: Percentages may total more than 100% due to deaths with multiple pneumoconioses. Slight variations in numbers may exist when compared to previous WORLD Surveillance Reports. See appendices for source description, methods, and ICD codes. SOURCE: National Center for Health Statistics multiple cause-of-death data.