## Johnson's Seagrass (Halophila johnsonii) Critical Habitat Southeast Florida (50 CFR 226.213) Sebastian Inlet 100 150 200 Miles Holmes/ nta Rosa Okaloosa Nassau Walton Washington Gadsden Jefferson Hamilton{ **Calhouก์** <sup>/</sup>Madison Bay Duval SuwanneeColumbia Baker Ft. Pierce Inlet Wakulla **Taylor** Bradford Clay Lafayette (Franklip GilchristAlachua Putnam Dixie Critical habitat is designated St. Lucie Inlet Flagler to include substrate and water Levy in the following ten portions of the Marion Indian River Lagoon and Biscayne Volusia Bay, Florida, within the current range Citrus` of Johnson's seagrass. Seminole lernando-Orange (a) North of Sebastian Inlet Channel. Hobe Sound Pasco & Jupiter Inlet Sceola Brevard (b) South of Sebastian Inlet Channel. Hillsborough Polk Lake Worth (c) Fort Pierce Inlet. Indian Riv Lagoon Manatee Hardee 6 keechobee<sub>St</sub> (d) North of St. Lucie Inlet. lighlands (e) Hobe Sound. **DeSoto** arasota Martin Glades **Charlotte** (f) South side of Jupiter Inlet. Palm Beach Hendry (g) A portion of Lake Worth Lagoon north of Bingham Island. Lake Wyman (h) A portion of Lake Worth Lagoon, located just north of the Boynton Inlet. Broward Collier (i) A portion of northeast Lake Wyman, Boca Raton. Biscayne Bay Miami-Dade (j) A portion of Northern Biscayne Bay. **V**onroe See 50 CFR 226.213 for geographic coordinates. mentally .

82°W

84°W

80°W

78°W