

S U D A A N

Software for the Statistical Analysis of Correlated Data
Copyright Research Triangle Institute February

2005

Release 9.0.1

Number of observations read	:	19759	Weighted count	:278652243
Observations in subpopulation	:	9471	Weighted count:	198419259
Denominator degrees of freedom	:	29		

Variance Estimation Method: Taylor Series (WR)

For Subpopulation: RIDAGEYR >= 20

Mean systolic blood pressure: NHANES 1999-2002

by: Variable, Gender.

Variable	Sample	Mean	SE Mean
Gender	Size		

SBP average reported			
to examinee			
Total	9056	123.10	0.43
Male	4301	123.81	0.46
Female	4755	122.45	0.53

S U D A A N

Software for the Statistical Analysis of Correlated Data
Copyright Research Triangle Institute February

2005

Release 9.0.1

Number of observations read	:	19759	Weighted count	:278652243
Observations in subpopulation	:	9471	Weighted count:	198419259
Denominator degrees of freedom	:	29		

Variance Estimation Method: Taylor Series (WR)

For Subpopulation: RIDAGEYR >= 20

Significance test for difference between mean systolic blood pressure
for males and females

NHANES 1999-2002

by: Variable, One, Contrast.

for: Variable = SBP average reported to examinee.

One			P-value
Contrast		T-Test	T-Test
	Sample	Cont.Mean-	Cont.
	Size	=0	Mean=0

Total			
Males vs. Females	9056	2.64	0.0132
1			
Males vs. Females	9056	2.64	0.0132
