

### **3. CHEMICAL AND PHYSICAL INFORMATION**

#### **3.1 CHEMICAL IDENTITY**

Information regarding the chemical identity of propylene glycol is located in Table 3-1.

#### **3.2 PHYSICAL AND CHEMICAL PROPERTIES**

Information regarding the physical and chemical properties of propylene glycol is located in Table 3-2.

## 3. CHEMICAL AND PHYSICAL INFORMATION

**Table 3-1. Chemical Identity of Propylene Glycol<sup>a</sup>**

Characteristic	Information
Chemical name	Propylene glycol
Synonyms and trade names	1,2-Dihydroxypropane; 1,2-propanediol; 1,2-propylene glycol; 2,3-propanediol; hydroxy-propanol; alpha-propylene glycol; methyl glycol; methylethyl glycol; monopropylene glycol; trimethyl glycol.
Registered trade name(s)	PG-12; Sirlene
Chemical formula	C <sub>3</sub> H <sub>8</sub> O <sub>2</sub>
Chemical structure <sup>b</sup>	$  \begin{array}{c}  \text{CH}_3 \\    \\  \text{CH}_2 - \text{OH} \\    \\  \text{C} - \text{OH} \\    \\  \text{H}_2  \end{array}  $
Identification numbers:	
CAS registry	57-55-6
NIOSH RTECS	TY2000000
EPA hazardous waste	No data
OHM/TADS	7216877
DOT/UN/NA/IMDG shipping	No data
HSDB	174
NCI	No data

<sup>a</sup>All information obtained from HSDB 1994b, except where noted.

<sup>b</sup>EPA 1987a

CAS = Chemical Abstracts Service; DOT/UN/NA/IMDG = Department of Transportation/United Nations/North America/International Maritime Dangerous Goods Code; EPA = Environmental Protection Agency; HSDB = Hazardous Substances Data Bank; NCI = National Cancer Institute; NIOSH = National Institute for Occupational Safety and Health; OHM/TADS = Oil and Hazardous Materials/Technical Assistance Data System; RTECS = Registry of Toxic Effects of Chemical Substances

## 3. CHEMICAL AND PHYSICAL INFORMATION

**Table 3-2. Physical and Chemical Properties of Propylene Glycol<sup>a</sup>**

Property	Propylene glycol
Molecular weight	76.11 <sup>b</sup>
Color	Colorless <sup>d</sup>
Physical state	Liquid <sup>b</sup>
Melting point	-60 °C <sup>a,b</sup> (forms glass)
Boiling point	187.6; 188.2 °C <sup>b</sup>
Density:	
at 20 °C (g/cm <sup>3</sup> )	1.0361 <sup>c</sup>
Odor	Odorless
Odor threshold	No data
Solubility:	
Water at 20 °C	Miscible with water
Organic solvent(s)	Soluble in alcohol, ether, benzene, soluble in acetone, chloroform <sup>b</sup> .
Partition coefficients:	
Log K <sub>ow</sub>	-0.92 <sup>f,g</sup>
Log K <sub>oc</sub>	0.88 <sup>f</sup> , 0.76 <sup>g</sup>
Vapor pressure at 20 °C	0.07 mm Hg <sup>also d</sup>
Henry's law constant at 25 °C	1.2x10 <sup>-8</sup> atm-m <sup>3</sup> /mole; 1.7x10 <sup>-8</sup> atm-m <sup>3</sup> /mole <sup>g</sup>
Autoignition temperature	421.26 °C <sup>h</sup> ; 371 °C <sup>i</sup>
Flashpoint	99.04 °C <sup>h,i</sup>
Flammability limits	2.6–12.5% <sup>h,i</sup>
Conversion factors	1 ppm = 3.11 mg/m <sup>3j</sup> 1 mg/L = 321.6 ppm <sup>i</sup>
Explosive limits	No data

<sup>a</sup>All information obtained from HSDB 1995b, except where noted.

<sup>b</sup>Merck 1989

<sup>c</sup>Weast 1988

<sup>d</sup>Lewis 1993

<sup>e</sup>Daubert and Danner 1980

<sup>f</sup>EPA 1987a

<sup>g</sup>ASTER 1995

<sup>h</sup>Daubert and Danner 1989

<sup>i</sup>NFPA 1994

<sup>j</sup>Rowe and Wolf 1982